

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All causes													
All races, both sexes													
2007.....	2,423,712	29,138	4,703	6,147	33,982	42,572	79,606	184,686	287,110	389,238	652,682	713,647	201
2006.....	2,426,264	28,527	4,631	6,149	34,887	42,952	83,043	185,031	281,401	390,093	667,338	701,992	220
2005.....	2,448,017	28,440	4,756	6,602	34,234	41,925	84,785	183,530	275,301	398,355	686,665	703,169	255
2004.....	2,397,615	27,936	4,785	6,834	33,421	40,868	85,362	177,697	264,697	399,666	684,230	671,773	346
2003.....	2,448,288	28,025	4,965	6,954	33,568	41,300	89,461	176,781	262,519	413,497	703,024	687,852	342
2002.....	2,443,387	28,034	4,858	7,150	33,046	41,355	91,140	172,385	253,342	422,990	707,654	681,076	357
2001.....	2,416,425	27,568	5,107	7,095	32,252	41,683	91,674	168,065	244,139	430,960	701,929	665,531	422
2000.....	2,403,351	28,035	4,979	7,413	31,307	40,451	89,798	160,341	240,846	441,209	700,445	658,171	356
1999.....	2,391,399	27,937	5,249	7,595	30,656	41,066	89,256	152,974	238,979	452,600	698,590	646,141	356
All races, male													
2007.....	1,203,968	16,293	2,634	3,585	25,316	29,792	50,105	114,456	173,618	218,344	320,803	248,866	156
2006.....	1,201,942	15,980	2,541	3,625	26,070	30,188	52,150	115,172	168,987	218,165	327,315	241,578	171
2005.....	1,207,675	16,018	2,765	3,853	25,509	29,283	53,309	114,472	165,429	222,807	335,235	238,796	199
2004.....	1,181,668	15,718	2,649	3,999	24,587	28,359	53,677	111,163	158,032	222,891	333,669	226,650	274
2003.....	1,201,964	15,902	2,826	4,150	24,670	28,602	56,435	110,682	156,461	231,421	342,332	228,212	271
2002.....	1,199,264	15,717	2,806	4,198	24,416	28,736	57,593	107,722	151,363	237,021	343,504	225,906	282
2001.....	1,183,421	15,477	2,899	4,168	23,963	28,757	58,164	104,848	144,958	241,581	340,742	217,533	331
2000.....	1,177,578	15,718	2,824	4,401	23,071	27,890	57,297	100,398	143,321	247,408	340,219	214,742	289
1999.....	1,175,460	15,646	2,975	4,492	22,414	28,276	57,118	95,659	142,724	254,920	340,970	209,989	277

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All causes													
All races, female													
2007.....	1,219,744	12,845	2,069	2,562	8,666	12,780	29,501	70,230	113,492	170,894	331,879	464,781	45
2006.....	1,224,322	12,547	2,090	2,524	8,817	12,764	30,893	69,859	112,414	171,928	340,023	460,414	49
2005.....	1,240,342	12,422	1,991	2,749	8,725	12,642	31,476	69,058	109,872	175,548	351,430	464,373	56
2004.....	1,215,947	12,218	2,136	2,835	8,834	12,509	31,685	66,534	106,665	176,775	350,561	445,123	72
2003.....	1,246,324	12,123	2,139	2,804	8,898	12,698	33,026	66,099	106,058	182,076	360,692	459,640	71
2002.....	1,244,123	12,317	2,052	2,952	8,630	12,619	33,547	64,663	101,979	185,969	364,150	455,170	75
2001.....	1,233,004	12,091	2,208	2,927	8,289	12,926	33,510	63,217	99,181	189,379	361,187	447,998	91
2000.....	1,225,773	12,317	2,155	3,012	8,236	12,561	32,501	59,943	97,525	193,801	360,226	443,429	67
1999.....	1,215,939	12,291	2,274	3,103	8,242	12,790	32,138	57,315	96,255	197,680	357,620	436,152	79
White, both sexes													
2007 1/.....	2,074,151	18,807	3,287	4,480	25,137	31,181	59,571	142,233	231,287	327,377	579,122	651,521	148
2006 1/.....	2,077,549	18,403	3,233	4,460	25,871	31,127	62,060	141,969	227,108	328,578	593,016	641,545	179
2005 1/.....	2,098,097	18,514	3,408	4,710	25,410	30,103	63,584	140,224	222,218	336,362	610,820	642,539	205
2004 1/.....	2,056,643	18,231	3,363	4,900	25,013	29,275	64,040	135,719	214,171	338,262	609,660	613,740	269
2003 1/.....	2,103,714	18,440	3,509	5,031	24,976	29,689	66,748	134,512	212,821	350,549	627,452	629,695	292
2002.....	2,102,589	18,369	3,406	5,138	24,641	29,571	68,093	131,816	205,414	360,047	632,353	623,455	286
2001.....	2,079,691	17,955	3,670	5,170	23,748	29,994	68,137	128,699	197,853	367,128	627,752	609,265	320
2000.....	2,071,287	18,144	3,494	5,450	23,018	28,719	66,496	122,674	195,541	376,986	627,729	602,761	275
1999.....	2,061,348	18,067	3,690	5,493	22,352	29,400	65,607	117,011	193,582	387,437	626,489	591,949	271

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All causes													
White, male													
2007 1/.....	1,023,951	10,540	1,847	2,590	18,513	21,998	38,121	89,556	140,785	184,828	286,824	228,227	122
2006 1/.....	1,022,328	10,345	1,781	2,639	19,175	22,083	39,770	89,681	137,161	184,925	293,004	221,622	142
2005 1/.....	1,028,152	10,471	1,996	2,749	18,792	21,232	40,824	88,882	134,272	189,550	300,228	218,990	166
2004 1/.....	1,007,266	10,265	1,878	2,905	18,306	20,584	41,215	86,266	128,575	190,123	299,342	207,589	218
2003 1/.....	1,025,650	10,500	1,981	3,008	18,219	20,814	43,083	85,587	127,752	197,853	307,050	209,575	228
2002.....	1,025,196	10,433	1,958	3,005	18,082	20,798	43,955	83,651	123,551	203,735	308,654	207,144	230
2001.....	1,011,218	10,099	2,094	3,014	17,529	20,978	44,243	81,403	118,353	207,793	306,278	199,179	255
2000.....	1,007,191	10,177	2,004	3,255	16,868	20,140	43,378	77,866	117,114	213,379	306,370	196,409	231
1999.....	1,005,335	10,197	2,084	3,228	16,203	20,549	42,914	73,908	116,537	220,442	307,450	191,610	213
White, female													
2007 1/.....	1,050,200	8,267	1,440	1,890	6,624	9,183	21,450	52,677	90,502	142,549	292,298	423,294	26
2006 1/.....	1,055,221	8,058	1,452	1,821	6,696	9,044	22,290	52,288	89,947	143,653	300,012	419,923	37
2005 1/.....	1,069,945	8,043	1,412	1,961	6,618	8,871	22,760	51,342	87,946	146,812	310,592	423,549	39
2004 1/.....	1,049,377	7,966	1,485	1,995	6,707	8,691	22,825	49,453	85,596	148,139	310,318	406,151	51
2003 1/.....	1,078,064	7,940	1,528	2,023	6,757	8,875	23,665	48,925	85,069	152,696	320,402	420,120	64
2002.....	1,077,393	7,936	1,448	2,133	6,559	8,773	24,138	48,165	81,863	156,312	323,699	416,311	56
2001.....	1,068,473	7,856	1,576	2,156	6,219	9,016	23,894	47,296	79,500	159,335	321,474	410,086	65
2000.....	1,064,096	7,967	1,490	2,195	6,150	8,579	23,118	44,808	78,427	163,607	321,359	406,352	44
1999.....	1,056,013	7,870	1,606	2,265	6,149	8,851	22,693	43,103	77,045	166,995	319,039	400,339	58

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All causes													
Black, both sexes													
2007 1/.....	289,585	8,944	1,137	1,363	7,380	9,662	16,990	36,818	47,682	51,384	58,986	49,194	45
2006 1/.....	289,971	8,858	1,144	1,394	7,483	10,045	17,930	37,419	46,396	51,126	60,002	48,142	32
2005 1/.....	292,808	8,695	1,097	1,543	7,371	10,036	18,186	37,749	45,416	51,718	61,998	48,952	47
2004 1/.....	287,315	8,494	1,164	1,584	7,049	9,925	18,474	36,710	43,490	51,682	61,379	47,300	64
2003 1/.....	291,300	8,402	1,199	1,551	7,145	9,942	19,784	37,006	42,823	53,128	62,372	47,908	40
2002.....	290,051	8,524	1,196	1,666	7,045	10,171	20,235	35,623	41,336	53,314	62,862	48,022	57
2001.....	287,709	8,498	1,181	1,587	7,147	10,028	20,687	34,518	39,976	54,379	62,233	47,387	88
2000.....	285,826	8,771	1,248	1,643	7,049	10,239	20,557	33,289	39,281	55,195	61,444	47,038	72
1999.....	285,064	8,822	1,311	1,789	7,066	10,115	21,041	31,759	39,476	56,122	61,201	46,285	77
Black, male													
2007 1/.....	148,309	4,975	621	816	5,751	6,648	10,106	21,495	28,127	27,672	26,815	15,255	28
2006 1/.....	148,602	4,886	633	831	5,793	6,919	10,521	22,112	27,198	27,587	27,322	14,778	22
2005 1/.....	149,108	4,867	623	909	5,709	6,843	10,626	22,248	26,711	27,616	28,069	14,856	31
2004 1/.....	145,970	4,769	641	888	5,336	6,684	10,700	21,756	25,352	27,555	27,695	14,548	46
2003 1/.....	148,022	4,740	699	922	5,448	6,693	11,544	21,986	24,770	28,291	28,600	14,294	35
2002.....	146,835	4,652	702	998	5,364	6,848	11,849	21,122	24,000	28,198	28,478	14,584	40
2001.....	145,908	4,764	672	960	5,479	6,673	12,135	20,594	22,905	28,775	28,395	14,489	67
2000.....	145,184	4,901	692	974	5,318	6,750	12,251	19,913	22,678	29,071	28,026	14,560	50
1999.....	145,703	4,897	746	1,096	5,348	6,723	12,567	19,298	22,769	29,594	27,839	14,768	58

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

All causes

Black, female

2007 1/.....	141,276	3,969	516	547	1,629	3,014	6,884	15,323	19,555	23,712	32,171	33,939	17
2006 1/.....	141,369	3,972	511	563	1,690	3,126	7,409	15,307	19,198	23,539	32,680	33,364	10
2005 1/.....	143,700	3,828	474	634	1,662	3,193	7,560	15,501	18,705	24,102	33,929	34,096	16
2004 1/.....	141,345	3,725	523	696	1,713	3,241	7,774	14,954	18,138	24,127	33,684	32,752	18
2003 1/.....	143,278	3,662	500	629	1,697	3,249	8,240	15,020	18,053	24,837	33,772	33,614	5
2002.....	143,216	3,872	494	668	1,681	3,323	8,386	14,501	17,336	25,116	34,384	33,438	17
2001.....	141,801	3,734	509	627	1,668	3,355	8,552	13,924	17,071	25,604	33,838	32,898	21
2000.....	140,642	3,870	556	669	1,731	3,489	8,306	13,376	16,603	26,124	33,418	32,478	22
1999.....	139,361	3,925	565	693	1,718	3,392	8,474	12,461	16,707	26,528	33,362	31,517	19

American Indian 2/, both sexes

2007 1/.....	14,367	432	97	94	622	731	1,213	1,958	2,367	2,536	2,546	1,771	-
2006 1/.....	14,037	393	94	91	664	708	1,252	1,937	2,226	2,452	2,542	1,676	2
2005 1/.....	13,918	357	100	110	635	720	1,223	1,927	2,240	2,487	2,433	1,684	2
2004 1/.....	13,124	389	97	127	593	695	1,194	1,739	2,073	2,329	2,375	1,503	10
2003 1/.....	13,147	335	95	137	621	643	1,244	1,812	2,042	2,256	2,363	1,594	5
2002.....	12,415	343	90	126	583	652	1,169	1,602	1,886	2,291	2,215	1,451	7
2001.....	11,977	373	103	129	557	596	1,132	1,559	1,768	2,206	2,146	1,402	6
2000.....	11,363	323	91	113	519	596	1,081	1,354	1,698	2,141	2,062	1,382	3
1999.....	11,312	343	82	105	540	645	1,027	1,328	1,674	2,164	2,064	1,339	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

All causes

American Indian 2/, male

2007 1/.....	7,885	241	57	61	441	514	768	1,193	1,372	1,357	1,214	667	-
2006 1/.....	7,630	242	51	47	478	492	784	1,199	1,294	1,295	1,156	591	1
2005 1/.....	7,607	196	62	64	439	512	777	1,160	1,341	1,297	1,144	614	1
2004 1/.....	7,134	237	50	70	409	461	747	1,045	1,212	1,225	1,146	524	8
2003 1/.....	7,106	195	55	86	450	444	784	1,090	1,136	1,191	1,092	578	5
2002.....	6,750	191	49	65	416	457	749	978	1,074	1,201	1,048	516	6
2001.....	6,466	211	52	75	400	417	722	940	1,004	1,165	983	492	5
2000.....	6,185	193	49	61	369	410	675	858	964	1,113	989	501	3
1999.....	6,092	180	48	55	396	418	645	845	920	1,148	987	449	1

American Indian 2/, female

2007 1/.....	6,482	191	40	33	181	217	445	765	995	1,179	1,332	1,104	-
2006 1/.....	6,407	151	43	44	186	216	468	738	932	1,157	1,386	1,085	1
2005 1/.....	6,311	161	38	46	196	208	446	767	899	1,190	1,289	1,070	1
2004 1/.....	5,990	152	47	57	184	234	447	694	861	1,104	1,229	979	2
2003 1/.....	6,041	140	40	51	171	199	460	722	906	1,065	1,271	1,016	-
2002.....	5,665	152	41	61	167	195	420	624	812	1,090	1,167	935	1
2001.....	5,511	162	51	54	157	179	410	619	764	1,041	1,163	910	1
2000.....	5,178	130	42	52	150	186	406	496	734	1,028	1,073	881	-
1999.....	5,220	163	34	50	144	227	382	483	754	1,016	1,077	890	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All causes													
Asian or Pacific Islander 3/, both sexes													
2007 1/.....	45,609	955	182	210	843	998	1,832	3,677	5,774	7,941	12,028	11,161	8
2006 1/.....	44,707	873	160	204	869	1,072	1,801	3,706	5,671	7,937	11,778	10,629	7
2005 1/.....	43,194	874	151	239	818	1,066	1,792	3,630	5,427	7,788	11,414	9,994	1
2004 1/.....	40,533	822	161	223	766	973	1,654	3,529	4,963	7,393	10,816	9,230	3
2003 1/.....	40,127	848	162	235	826	1,026	1,685	3,451	4,833	7,564	10,837	8,655	5
2002.....	38,332	798	166	220	777	961	1,643	3,344	4,706	7,338	10,224	8,148	7
2001.....	37,048	742	153	209	800	1,065	1,718	3,289	4,542	7,247	9,798	7,477	8
2000.....	34,875	797	146	207	721	897	1,664	3,024	4,326	6,887	9,210	6,990	6
1999.....	33,675	705	166	208	698	906	1,581	2,876	4,247	6,877	8,836	6,568	7
Asian or Pacific Islander 3/, male													
2007 1/.....	23,823	537	109	118	611	632	1,110	2,212	3,334	4,487	5,950	4,717	6
2006 1/.....	23,382	507	76	108	624	694	1,075	2,180	3,334	4,358	5,833	4,587	6
2005 1/.....	22,808	484	84	131	569	696	1,082	2,182	3,105	4,344	5,794	4,336	1
2004 1/.....	21,298	447	80	136	536	630	1,015	2,096	2,893	3,988	5,486	3,989	2
2003 1/.....	21,186	467	91	134	553	651	1,024	2,019	2,803	4,086	5,590	3,765	3
2002.....	20,483	441	97	130	554	633	1,040	1,971	2,738	3,887	5,324	3,662	6
2001.....	19,829	403	81	119	555	689	1,064	1,911	2,696	3,848	5,086	3,373	4
2000.....	19,018	447	79	111	516	590	993	1,761	2,565	3,845	4,834	3,272	5
1999.....	18,330	372	97	113	467	586	992	1,608	2,498	3,736	4,694	3,162	5

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

All causes

Asian or Pacific Islander 3/, female

2007 1/.....	21,786	418	73	92	232	366	722	1,465	2,440	3,454	6,078	6,444	2
2006 1/.....	21,325	366	84	96	245	378	726	1,526	2,337	3,579	5,945	6,042	1
2005 1/.....	20,386	390	67	108	249	370	710	1,448	2,322	3,444	5,620	5,658	-
2004 1/.....	19,235	375	81	87	230	343	639	1,433	2,070	3,405	5,330	5,241	1
2003 1/.....	18,941	381	71	101	273	375	661	1,432	2,030	3,478	5,247	4,890	2
2002.....	17,849	357	69	90	223	328	603	1,373	1,968	3,451	4,900	4,486	1
2001.....	17,219	339	72	90	245	376	654	1,378	1,846	3,399	4,712	4,104	4
2000.....	15,857	350	67	96	205	307	671	1,263	1,761	3,042	4,376	3,718	1
1999.....	15,345	333	69	95	231	320	589	1,268	1,749	3,141	4,142	3,406	2

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Salmonella infections (A01-A02)													
All races, both sexes													
2007.....	30	2	2	-	-	2	-	3	2	4	6	9	-
2006.....	34	3	1	-	1	-	1	2	4	7	9	6	-
2005.....	30	-	-	-	1	2	2	2	6	3	9	5	-
2004.....	30	1	-	-	-	1	1	3	4	7	7	6	-
2003.....	43	3	-	1	-	1	3	2	6	8	10	9	-
2002.....	21	3	-	-	-	-	2	3	-	4	8	1	-
2001.....	41	2	-	-	-	2	1	3	3	7	14	9	-
2000.....	28	3	2	-	1	1	-	4	2	7	5	3	-
1999.....	38	3	-	-	2	-	2	2	3	7	12	7	-
All races, male													
2007.....	14	-	2	-	-	1	-	2	-	2	2	5	-
2006.....	24	2	-	-	-	-	-	2	3	7	7	3	-
2005.....	11	-	-	-	-	-	1	1	5	1	2	1	-
2004.....	22	1	-	-	-	-	1	3	4	3	6	4	-
2003.....	23	1	-	-	-	-	2	2	4	4	4	6	-
2002.....	10	1	-	-	-	-	-	1	-	2	5	1	-
2001.....	21	-	-	-	-	1	1	3	2	6	7	1	-
2000.....	15	2	-	-	1	1	-	2	2	5	1	1	-
1999.....	21	2	-	-	2	-	1	2	2	5	5	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Salmonella infections (A01-A02)													
All races, female													
2007.....	16	2	-	-	-	1	-	1	2	2	4	4	-
2006.....	10	1	1	-	1	-	1	-	1	-	2	3	-
2005.....	19	-	-	-	1	2	1	1	1	2	7	4	-
2004.....	8	-	-	-	-	1	-	-	-	4	1	2	-
2003.....	20	2	-	1	-	1	1	-	2	4	6	3	-
2002.....	11	2	-	-	-	-	2	2	-	2	3	-	-
2001.....	20	2	-	-	-	1	-	-	1	1	7	8	-
2000.....	13	1	2	-	-	-	-	2	-	2	4	2	-
1999.....	17	1	-	-	-	-	1	-	1	2	7	5	-
White, both sexes													
2007 1/.....	24	2	1	-	-	1	-	2	1	4	5	8	-
2006 1/.....	25	1	1	-	-	-	1	1	4	5	7	5	-
2005 1/.....	24	-	-	-	-	1	2	1	5	3	8	4	-
2004 1/.....	20	1	-	-	-	1	-	2	2	3	6	5	-
2003 1/.....	31	1	-	1	-	-	1	2	5	6	7	8	-
2002.....	18	2	-	-	-	-	2	3	-	3	7	1	-
2001.....	33	1	-	-	-	2	-	3	3	4	12	8	-
2000.....	25	2	2	-	-	1	-	4	2	6	5	3	-
1999.....	32	3	-	-	1	-	2	2	2	5	10	7	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Salmonella infections (A01-A02)													
White, male													
2007 1/.....	11	-	1	-	-	-	-	1	-	2	2	5	-
2006 1/.....	18	1	-	-	-	-	-	1	3	5	6	2	-
2005 1/.....	9	-	-	-	-	-	1	1	4	1	1	1	-
2004 1/.....	14	1	-	-	-	-	-	2	2	1	5	3	-
2003 1/.....	17	-	-	-	-	-	1	2	3	4	2	5	-
2002.....	9	1	-	-	-	-	-	1	-	1	5	1	-
2001.....	18	-	-	-	-	1	-	3	2	4	7	1	-
2000.....	13	1	-	-	-	1	-	2	2	5	1	1	-
1999.....	16	2	-	-	1	-	1	2	1	3	4	2	-
White, female													
2007 1/.....	13	2	-	-	-	1	-	1	1	2	3	3	-
2006 1/.....	7	-	1	-	-	-	1	-	1	-	1	3	-
2005 1/.....	15	-	-	-	-	1	1	-	1	2	7	3	-
2004 1/.....	6	-	-	-	-	1	-	-	-	2	1	2	-
2003 1/.....	14	1	-	1	-	-	-	-	2	2	5	3	-
2002.....	9	1	-	-	-	-	2	2	-	2	2	-	-
2001.....	15	1	-	-	-	1	-	-	1	-	5	7	-
2000.....	12	1	2	-	-	-	-	2	-	1	4	2	-
1999.....	16	1	-	-	-	-	1	-	1	2	6	5	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Salmonella infections (A01-A02)

Black, both sexes

2007 1/.....	3	-	-	-	-	-	-	1	1	-	1	-	-
2006 1/.....	8	2	-	-	1	-	-	1	-	1	2	1	-
2005 1/.....	5	-	-	-	-	1	-	1	1	-	1	1	-
2004 1/.....	4	-	-	-	-	-	-	1	1	2	-	-	-
2003 1/.....	8	2	-	-	-	1	1	-	-	1	3	-	-
2002.....	3	1	-	-	-	-	-	-	-	1	1	-	-
2001.....	5	1	-	-	-	-	-	-	-	1	2	1	-
2000.....	2	1	-	-	1	-	-	-	-	-	-	-	-
1999.....	4	-	-	-	1	-	-	-	-	1	2	-	-

Black, male

2007 1/.....	1	-	-	-	-	-	-	1	-	-	-	-	-
2006 1/.....	5	1	-	-	-	-	-	1	-	1	1	1	-
2005 1/.....	2	-	-	-	-	-	-	-	1	-	1	-	-
2004 1/.....	3	-	-	-	-	-	-	1	1	1	-	-	-
2003 1/.....	3	1	-	-	-	-	-	-	-	-	2	-	-
2002.....	1	-	-	-	-	-	-	-	-	1	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	1	-	-	-
2000.....	2	1	-	-	1	-	-	-	-	-	-	-	-
1999.....	3	-	-	-	1	-	-	-	-	1	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Salmonella infections (A01-A02)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	2	-	1	-	-	-	-	-	-	-	-	1	-
2006 1/.....	1	-	-	-	-	-	-	-	-	1	-	-	-
2005 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 1/.....	5	-	-	-	-	-	-	1	2	1	1	-	-
2003 1/.....	3	-	-	-	-	-	1	-	1	-	-	1	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	3	-	-	-	-	-	1	-	-	2	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	1	-	-	-
1999.....	2	-	-	-	-	-	-	-	1	1	-	-	-

Asian or Pacific Islander 3/, male

2007 1/.....	1	-	1	-	-	-	-	-	-	-	-	-	-
2006 1/.....	1	-	-	-	-	-	-	-	-	1	-	-	-
2005 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 1/.....	4	-	-	-	-	-	-	-	1	1	1	1	-
2003 1/.....	3	-	-	-	-	-	1	-	1	-	-	1	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	2	-	-	-	-	-	1	-	-	1	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	-	-	-	-	-	-	-	1	1	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Shigellosis and amebiasis (A03,A06)

All races, both sexes

2007.....	4	-	-	2	-	-	1	-	-	-	1	-	-
2006.....	6	-	3	-	-	-	-	1	1	-	1	-	-
2005.....	10	-	1	2	-	-	-	1	2	2	2	-	-
2004.....	3	-	-	1	1	-	-	-	-	1	-	-	-
2003.....	7	-	-	-	-	-	3	-	2	1	1	-	-
2002.....	8	-	1	-	1	-	1	2	1	-	1	1	-
2001.....	7	1	1	1	-	-	-	3	-	-	1	-	-
2000.....	12	-	-	-	-	-	1	1	1	1	7	1	-
1999.....	10	-	1	1	-	-	1	-	1	1	1	4	-

All races, male

2007.....	2	-	-	1	-	-	1	-	-	-	-	-	-
2006.....	2	-	1	-	-	-	-	1	-	-	-	-	-
2005.....	3	-	-	1	-	-	-	1	1	-	-	-	-
2004.....	2	-	-	1	-	-	-	-	-	1	-	-	-
2003.....	5	-	-	-	-	-	2	-	1	1	1	-	-
2002.....	4	-	-	-	1	-	1	2	-	-	-	-	-
2001.....	6	1	1	1	-	-	-	3	-	-	-	-	-
2000.....	5	-	-	-	-	-	1	-	1	-	3	-	-
1999.....	6	-	1	1	-	-	1	-	1	1	-	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Shigellosis and amebiasis (A03,A06)

All races, female

2007.....	2	-	-	1	-	-	-	-	-	-	1	-	-
2006.....	4	-	2	-	-	-	-	-	1	-	1	-	-
2005.....	7	-	1	1	-	-	-	-	1	2	2	-	-
2004.....	1	-	-	-	1	-	-	-	-	-	-	-	-
2003.....	2	-	-	-	-	-	1	-	1	-	-	-	-
2002.....	4	-	1	-	-	-	-	-	1	-	1	1	-
2001.....	1	-	-	-	-	-	-	-	-	-	1	-	-
2000.....	7	-	-	-	-	-	-	1	-	1	4	1	-
1999.....	4	-	-	-	-	-	-	-	-	-	1	3	-

White, both sexes

2007 1/.....	3	-	-	1	-	-	1	-	-	-	1	-	-
2006 1/.....	5	-	2	-	-	-	-	1	1	-	1	-	-
2005 1/.....	10	-	1	2	-	-	-	1	2	2	2	-	-
2004 1/.....	3	-	-	1	1	-	-	-	-	1	-	-	-
2003 1/.....	6	-	-	-	-	-	2	-	2	1	1	-	-
2002.....	7	-	-	-	1	-	1	2	1	-	1	1	-
2001.....	7	1	1	1	-	-	-	3	-	-	1	-	-
2000.....	10	-	-	-	-	-	-	1	1	-	7	1	-
1999.....	10	-	1	1	-	-	1	-	1	1	1	4	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Shigellosis and amebiasis (A03,A06)

White, male

2007 1/.....	2	-	-	1	-	-	1	-	-	-	-	-	-
2006 1/.....	2	-	1	-	-	-	-	1	-	-	-	-	-
2005 1/.....	3	-	-	1	-	-	-	1	1	-	-	-	-
2004 1/.....	2	-	-	1	-	-	-	-	-	1	-	-	-
2003 1/.....	4	-	-	-	-	-	1	-	1	1	1	-	-
2002.....	4	-	-	-	1	-	1	2	-	-	-	-	-
2001.....	6	1	1	1	-	-	-	3	-	-	-	-	-
2000.....	4	-	-	-	-	-	-	-	1	-	3	-	-
1999.....	6	-	1	1	-	-	1	-	1	1	-	1	-

White, female

2007 1/.....	1	-	-	-	-	-	-	-	-	-	1	-	-
2006 1/.....	3	-	1	-	-	-	-	-	1	-	1	-	-
2005 1/.....	7	-	1	1	-	-	-	-	1	2	2	-	-
2004 1/.....	1	-	-	-	1	-	-	-	-	-	-	-	-
2003 1/.....	2	-	-	-	-	-	1	-	1	-	-	-	-
2002.....	3	-	-	-	-	-	-	-	1	-	1	1	-
2001.....	1	-	-	-	-	-	-	-	-	-	1	-	-
2000.....	6	-	-	-	-	-	-	1	-	-	4	1	-
1999.....	4	-	-	-	-	-	-	-	-	-	1	3	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Certain other intestinal infections (A04,A07-A09)

All races, both sexes

2007.....	6,758	11	14	5	5	15	36	122	325	915	2,446	2,864	-
2006.....	6,639	13	9	4	10	11	33	124	363	911	2,484	2,677	-
2005.....	5,667	10	12	10	3	11	32	104	253	829	2,070	2,333	-
2004.....	4,357	13	15	1	2	12	31	94	252	608	1,662	1,667	-
2003.....	3,090	10	4	7	4	8	14	64	176	444	1,192	1,167	-
2002.....	2,465	13	8	3	6	4	18	56	142	362	889	964	-
2001.....	1,537	10	18	4	-	4	12	42	81	235	566	565	-
2000.....	1,328	13	10	-	1	3	19	29	70	222	462	499	-
1999.....	1,108	12	8	3	5	6	7	23	49	176	390	429	-

All races, male

2007.....	2,541	8	8	2	4	4	16	53	174	407	970	895	-
2006.....	2,550	5	6	2	4	8	21	64	193	420	983	844	-
2005.....	2,209	7	4	5	3	6	17	53	132	377	865	740	-
2004.....	1,622	8	5	-	1	5	19	48	125	264	665	482	-
2003.....	1,109	6	4	2	2	4	5	32	82	185	478	309	-
2002.....	947	3	5	1	3	1	8	29	71	185	358	283	-
2001.....	572	7	11	2	-	2	8	15	35	108	227	157	-
2000.....	524	10	6	-	-	3	10	16	39	105	195	140	-
1999.....	420	8	4	3	3	5	4	11	26	74	169	113	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Certain other intestinal infections (A04,A07-A09)

All races, female

2007.....	4,217	3	6	3	1	11	20	69	151	508	1,476	1,969	-
2006.....	4,089	8	3	2	6	3	12	60	170	491	1,501	1,833	-
2005.....	3,458	3	8	5	-	5	15	51	121	452	1,205	1,593	-
2004.....	2,735	5	10	1	1	7	12	46	127	344	997	1,185	-
2003.....	1,981	4	-	5	2	4	9	32	94	259	714	858	-
2002.....	1,518	10	3	2	3	3	10	27	71	177	531	681	-
2001.....	965	3	7	2	-	2	4	27	46	127	339	408	-
2000.....	804	3	4	-	1	-	9	13	31	117	267	359	-
1999.....	688	4	4	-	2	1	3	12	23	102	221	316	-

White, both sexes

2007 1/.....	6,252	5	9	4	4	9	31	98	275	818	2,284	2,715	-
2006 1/.....	6,165	8	4	3	9	8	24	88	302	819	2,347	2,553	-
2005 1/.....	5,271	6	9	8	2	9	23	85	208	749	1,945	2,227	-
2004 1/.....	4,075	10	12	1	2	9	22	76	223	554	1,563	1,603	-
2003 1/.....	2,877	5	3	6	3	6	11	50	150	413	1,120	1,110	-
2002.....	2,287	7	6	3	6	2	15	49	111	329	845	914	-
2001.....	1,417	6	15	3	-	2	10	37	66	211	527	540	-
2000.....	1,224	8	7	-	1	2	11	22	62	202	441	468	-
1999.....	1,026	8	7	1	4	6	4	18	41	162	364	411	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Certain other intestinal infections (A04,A07-A09)

White, male

2007 1/.....	2,345	4	5	2	3	3	14	46	148	360	905	855	-
2006 1/.....	2,353	4	3	2	4	7	15	45	162	380	924	807	-
2005 1/.....	2,044	4	2	3	2	5	11	43	110	347	804	713	-
2004 1/.....	1,502	6	3	-	1	4	13	37	108	239	629	462	-
2003 1/.....	1,036	2	3	2	2	3	5	26	68	173	455	297	-
2002.....	872	1	4	1	3	1	6	27	56	164	340	269	-
2001.....	518	4	9	1	-	2	6	12	31	96	208	149	-
2000.....	478	6	5	-	-	2	7	12	34	96	184	132	-
1999.....	389	6	3	1	3	5	3	7	22	72	160	107	-

White, female

2007 1/.....	3,907	1	4	2	1	6	17	52	127	458	1,379	1,860	-
2006 1/.....	3,812	4	1	1	5	1	9	43	140	439	1,423	1,746	-
2005 1/.....	3,227	2	7	5	-	4	12	42	98	402	1,141	1,514	-
2004 1/.....	2,573	4	9	1	1	5	9	39	115	315	934	1,141	-
2003 1/.....	1,841	3	-	4	1	3	6	24	82	240	665	813	-
2002.....	1,415	6	2	2	3	1	9	22	55	165	505	645	-
2001.....	899	2	6	2	-	-	4	25	35	115	319	391	-
2000.....	746	2	2	-	1	-	4	10	28	106	257	336	-
1999.....	637	2	4	-	1	1	1	11	19	90	204	304	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Certain other intestinal infections (A04,A07-A09)

Black, both sexes

2007 1/.....	425	4	4	1	-	5	4	19	43	83	135	127	-
2006 1/.....	379	3	3	1	-	3	8	32	53	78	106	92	-
2005 1/.....	324	3	3	2	1	1	8	17	38	66	98	87	-
2004 1/.....	237	1	2	-	-	3	7	16	26	45	81	56	-
2003 1/.....	186	5	1	1	1	1	2	11	23	25	64	52	-
2002.....	156	5	1	-	-	2	3	5	29	31	36	44	-
2001.....	101	3	3	1	-	2	2	5	13	23	30	19	-
2000.....	84	4	2	-	-	1	6	7	6	15	17	26	-
1999.....	74	4	-	2	1	-	3	3	6	14	25	16	-

Black, male

2007 1/.....	161	3	2	-	-	1	2	6	22	38	54	33	-
2006 1/.....	145	-	2	-	-	1	5	15	27	32	41	22	-
2005 1/.....	131	2	2	2	1	1	5	9	17	26	47	19	-
2004 1/.....	96	1	1	-	-	1	4	10	16	21	26	16	-
2003 1/.....	66	4	1	-	-	1	-	5	12	9	23	11	-
2002.....	66	1	1	-	-	-	2	2	15	20	13	12	-
2001.....	43	2	2	1	-	-	2	3	4	11	12	6	-
2000.....	37	3	1	-	-	1	2	4	4	6	9	7	-
1999.....	25	2	-	2	-	-	1	2	2	2	9	5	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Certain other intestinal infections (A04,A07-A09)

Black, female

2007 1/.....	264	1	2	1	-	4	2	13	21	45	81	94	-
2006 1/.....	234	3	1	1	-	2	3	17	26	46	65	70	-
2005 1/.....	193	1	1	-	-	-	3	8	21	40	51	68	-
2004 1/.....	141	-	1	-	-	2	3	6	10	24	55	40	-
2003 1/.....	120	1	-	1	1	-	2	6	11	16	41	41	-
2002.....	90	4	-	-	-	2	1	3	14	11	23	32	-
2001.....	58	1	1	-	-	2	-	2	9	12	18	13	-
2000.....	47	1	1	-	-	-	4	3	2	9	8	19	-
1999.....	49	2	-	-	1	-	2	1	4	12	16	11	-

American Indian 2/, both sexes

2007 1/.....	31	1	1	-	-	1	-	4	3	7	10	4	-
2006 1/.....	30	1	-	-	-	-	1	3	3	5	8	9	-
2005 1/.....	22	-	-	-	-	-	-	2	4	7	7	2	-
2004 1/.....	17	1	-	-	-	-	1	1	2	6	6	-	-
2003 1/.....	8	-	-	-	-	-	-	1	2	3	2	-	-
2002.....	8	-	1	-	-	-	-	1	1	1	2	2	-
2001.....	6	-	-	-	-	-	-	-	1	-	4	1	-
2000.....	6	-	-	-	-	-	2	-	2	1	-	1	-
1999.....	3	-	-	-	-	-	-	1	1	-	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Certain other intestinal infections (A04,A07-A09)

American Indian 2/, male

2007 1/.....	13	1	1	-	-	-	-	-	1	4	5	1	-
2006 1/.....	16	1	-	-	-	-	1	3	1	4	3	3	-
2005 1/.....	5	-	-	-	-	-	-	1	2	-	1	1	-
2004 1/.....	10	1	-	-	-	-	1	1	-	2	5	-	-
2003 1/.....	2	-	-	-	-	-	-	-	1	1	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	1	-	-
2001.....	3	-	-	-	-	-	-	-	-	-	3	-	-
2000.....	3	-	-	-	-	-	1	-	1	1	-	-	-
1999.....	2	-	-	-	-	-	-	1	1	-	-	-	-

American Indian 2/, female

2007 1/.....	18	-	-	-	-	1	-	4	2	3	5	3	-
2006 1/.....	14	-	-	-	-	-	-	-	2	1	5	6	-
2005 1/.....	17	-	-	-	-	-	-	1	2	7	6	1	-
2004 1/.....	7	-	-	-	-	-	-	-	2	4	1	-	-
2003 1/.....	6	-	-	-	-	-	-	1	1	2	2	-	-
2002.....	7	-	1	-	-	-	-	1	1	1	1	2	-
2001.....	3	-	-	-	-	-	-	-	1	-	1	1	-
2000.....	3	-	-	-	-	-	1	-	1	-	-	1	-
1999.....	1	-	-	-	-	-	-	-	-	-	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Certain other intestinal infections (A04,A07-A09)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	50	1	-	-	1	-	1	1	4	7	17	18	-
2006 1/.....	65	1	2	-	1	-	-	1	5	9	23	23	-
2005 1/.....	50	1	-	-	-	1	1	-	3	7	20	17	-
2004 1/.....	28	1	1	-	-	-	1	1	1	3	12	8	-
2003 1/.....	19	-	-	-	-	1	1	2	1	3	6	5	-
2002.....	14	1	-	-	-	-	-	1	1	1	6	4	-
2001.....	13	1	-	-	-	-	-	-	1	1	5	5	-
2000.....	14	1	1	-	-	-	-	-	-	4	4	4	-
1999.....	5	-	1	-	-	-	-	1	1	-	-	2	-

Asian or Pacific Islander 3/, male

2007 1/.....	22	-	-	-	1	-	-	1	3	5	6	6	-
2006 1/.....	36	-	1	-	-	-	-	1	3	4	15	12	-
2005 1/.....	29	1	-	-	-	-	1	-	3	4	13	7	-
2004 1/.....	14	-	1	-	-	-	1	-	1	2	5	4	-
2003 1/.....	5	-	-	-	-	-	-	1	1	2	-	1	-
2002.....	8	1	-	-	-	-	-	-	-	1	4	2	-
2001.....	8	1	-	-	-	-	-	-	-	1	4	2	-
2000.....	6	1	-	-	-	-	-	-	-	2	2	1	-
1999.....	4	-	1	-	-	-	-	1	1	-	-	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Tuberculosis (A16-A19)													
All races, both sexes													
2007.....	554	2	-	1	6	14	32	73	88	97	151	90	-
2006.....	652	-	4	1	4	18	31	97	101	117	168	111	-
2005.....	648	2	3	1	18	21	38	79	91	116	165	114	-
2004.....	657	1	1	2	7	16	53	80	79	137	165	116	-
2003.....	711	-	1	2	10	18	35	105	102	114	217	107	-
2002.....	784	1	-	-	9	18	52	81	107	161	234	121	-
2001.....	764	2	1	-	12	25	40	84	116	152	206	125	1
2000.....	776	1	6	1	11	20	42	91	81	155	252	115	1
1999.....	930	2	2	3	11	19	57	106	130	173	253	173	1
All races, male													
2007.....	342	1	-	-	4	7	23	57	66	53	82	49	-
2006.....	425	-	1	-	2	11	26	77	81	77	98	52	-
2005.....	392	2	2	-	12	12	30	54	64	74	89	53	-
2004.....	406	-	1	1	4	12	40	53	60	93	84	58	-
2003.....	427	-	1	2	3	13	25	79	71	72	116	45	-
2002.....	462	-	-	-	5	11	38	55	75	102	121	55	-
2001.....	453	-	-	-	8	14	27	63	85	98	100	57	1
2000.....	468	-	1	-	7	14	28	76	59	87	139	56	1
1999.....	584	-	2	2	4	11	43	87	96	116	137	85	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Tuberculosis (A16-A19)													
All races, female													
2007.....	212	1	-	1	2	7	9	16	22	44	69	41	-
2006.....	227	-	3	1	2	7	5	20	20	40	70	59	-
2005.....	256	-	1	1	6	9	8	25	27	42	76	61	-
2004.....	251	1	-	1	3	4	13	27	19	44	81	58	-
2003.....	284	-	-	-	7	5	10	26	31	42	101	62	-
2002.....	322	1	-	-	4	7	14	26	32	59	113	66	-
2001.....	311	2	1	-	4	11	13	21	31	54	106	68	-
2000.....	308	1	5	1	4	6	14	15	22	68	113	59	-
1999.....	346	2	-	1	7	8	14	19	34	57	116	88	-
White, both sexes													
2007 1/.....	343	1	-	1	4	8	16	41	48	54	94	76	-
2006 1/.....	388	-	2	1	4	9	17	45	55	65	112	78	-
2005 1/.....	376	2	2	1	9	10	25	44	46	63	96	78	-
2004 1/.....	390	1	1	1	3	8	29	38	42	78	113	76	-
2003 1/.....	437	-	1	2	6	8	11	45	60	76	150	78	-
2002.....	461	-	-	-	3	6	22	36	57	88	163	86	-
2001.....	469	1	1	-	8	10	22	42	67	90	140	87	1
2000.....	439	-	4	1	6	11	13	44	33	87	155	85	-
1999.....	564	1	-	1	5	5	25	52	73	108	172	122	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Tuberculosis (A16-A19)													
White, male													
2007 1/.....	209	1	-	-	3	6	10	31	38	30	48	42	-
2006 1/.....	237	-	-	-	2	5	15	36	43	41	63	32	-
2005 1/.....	226	2	2	-	6	7	19	35	34	38	48	35	-
2004 1/.....	237	-	1	1	2	5	19	26	34	53	58	38	-
2003 1/.....	255	-	1	2	3	5	9	37	43	46	77	32	-
2002.....	272	-	-	-	2	3	17	26	37	64	83	40	-
2001.....	271	-	-	-	6	8	16	30	51	53	69	37	1
2000.....	253	-	-	-	4	7	11	34	24	53	78	42	-
1999.....	348	-	-	1	4	4	19	46	52	73	91	58	-
White, female													
2007 1/.....	134	-	-	1	1	2	6	10	10	24	46	34	-
2006 1/.....	151	-	2	1	2	4	2	9	12	24	49	46	-
2005 1/.....	150	-	-	1	3	3	6	9	12	25	48	43	-
2004 1/.....	153	1	-	-	1	3	10	12	8	25	55	38	-
2003 1/.....	182	-	-	-	3	3	2	8	17	30	73	46	-
2002.....	189	-	-	-	1	3	5	10	20	24	80	46	-
2001.....	198	1	1	-	2	2	6	12	16	37	71	50	-
2000.....	186	-	4	1	2	4	2	10	9	34	77	43	-
1999.....	216	1	-	-	1	1	6	6	21	35	81	64	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Tuberculosis (A16-A19)													
Black, both sexes													
2007 1/.....	122	1	-	-	1	3	13	22	27	24	26	5	-
2006 1/.....	160	-	1	-	-	6	11	36	34	36	26	10	-
2005 1/.....	153	-	1	-	6	7	10	24	28	32	29	16	-
2004 1/.....	189	-	-	1	3	6	21	32	26	41	32	27	-
2003 1/.....	172	-	-	-	4	6	16	47	31	20	33	15	-
2002.....	204	-	-	-	4	8	24	35	34	43	37	19	-
2001.....	203	-	-	-	1	10	16	37	38	44	35	22	-
2000.....	244	-	2	-	2	7	25	42	34	52	60	19	1
1999.....	245	1	2	1	4	9	25	42	40	43	50	27	1
Black, male													
2007 1/.....	83	-	-	-	-	-	11	20	18	17	14	3	-
2006 1/.....	115	-	-	-	-	4	9	27	29	24	16	6	-
2005 1/.....	94	-	-	-	4	2	9	16	19	25	13	6	-
2004 1/.....	118	-	-	-	2	5	18	18	20	26	17	12	-
2003 1/.....	114	-	-	-	-	5	11	35	21	15	20	7	-
2002.....	127	-	-	-	2	5	17	24	25	25	22	7	-
2001.....	127	-	-	-	-	4	10	30	27	30	14	12	-
2000.....	161	-	1	-	1	5	14	37	26	28	40	8	1
1999.....	159	-	2	-	-	4	19	34	32	29	25	13	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Tuberculosis (A16-A19)

Black, female

2007 1/.....	39	1	-	-	1	3	2	2	9	7	12	2	-
2006 1/.....	45	-	1	-	-	2	2	9	5	12	10	4	-
2005 1/.....	59	-	1	-	2	5	1	8	9	7	16	10	-
2004 1/.....	71	-	-	1	1	1	3	14	6	15	15	15	-
2003 1/.....	58	-	-	-	4	1	5	12	10	5	13	8	-
2002.....	77	-	-	-	2	3	7	11	9	18	15	12	-
2001.....	76	-	-	-	1	6	6	7	11	14	21	10	-
2000.....	83	-	1	-	1	2	11	5	8	24	20	11	-
1999.....	86	1	-	1	4	5	6	8	8	14	25	14	-

American Indian 2/, both sexes

2007 1/.....	12	-	-	-	-	-	1	-	4	2	5	-	-
2006 1/.....	19	-	-	-	-	2	1	7	2	2	4	1	-
2005 1/.....	15	-	-	-	-	1	2	1	3	4	1	3	-
2004 1/.....	13	-	-	-	-	-	-	-	2	5	5	1	-
2003 1/.....	15	-	-	-	-	-	2	6	-	1	4	2	-
2002.....	27	-	-	-	-	1	2	3	5	7	7	2	-
2001.....	13	-	-	-	-	1	-	2	2	2	4	2	-
2000.....	20	1	-	-	-	-	-	2	6	5	5	1	-
1999.....	20	-	-	-	-	1	1	4	2	6	1	5	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Tuberculosis (A16-A19)

American Indian 2/, male

2007 1/.....	5	-	-	-	-	-	1	-	2	-	2	-	-
2006 1/.....	11	-	-	-	-	1	-	6	1	2	1	-	-
2005 1/.....	11	-	-	-	-	1	2	1	2	1	1	3	-
2004 1/.....	9	-	-	-	-	-	-	-	2	4	3	-	-
2003 1/.....	6	-	-	-	-	-	1	3	-	-	1	1	-
2002.....	14	-	-	-	-	1	-	1	3	3	4	2	-
2001.....	5	-	-	-	-	-	-	1	1	1	2	-	-
2000.....	12	-	-	-	-	-	-	2	4	3	3	-	-
1999.....	11	-	-	-	-	-	1	3	1	3	-	3	-

American Indian 2/, female

2007 1/.....	7	-	-	-	-	-	-	-	2	2	3	-	-
2006 1/.....	8	-	-	-	-	1	1	1	1	-	3	1	-
2005 1/.....	4	-	-	-	-	-	-	-	1	3	-	-	-
2004 1/.....	4	-	-	-	-	-	-	-	-	1	2	1	-
2003 1/.....	9	-	-	-	-	-	1	3	-	1	3	1	-
2002.....	13	-	-	-	-	-	2	2	2	4	3	-	-
2001.....	8	-	-	-	-	1	-	1	1	1	2	2	-
2000.....	8	1	-	-	-	-	-	-	2	2	2	1	-
1999.....	9	-	-	-	-	1	-	1	1	3	1	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Tuberculosis (A16-A19)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	77	-	-	-	1	3	2	10	9	17	26	9	-
2006 1/.....	85	-	1	-	-	1	2	9	10	14	26	22	-
2005 1/.....	104	-	-	-	3	3	1	10	14	17	39	17	-
2004 1/.....	65	-	-	-	1	2	3	10	9	13	15	12	-
2003 1/.....	87	-	-	-	-	4	6	7	11	17	30	12	-
2002.....	92	1	-	-	2	3	4	7	11	23	27	14	-
2001.....	79	1	-	-	3	4	2	3	9	16	27	14	-
2000.....	73	-	-	-	3	2	4	3	8	11	32	10	-
1999.....	101	-	-	1	2	4	6	8	15	16	30	19	-

Asian or Pacific Islander 3/, male

2007 1/.....	45	-	-	-	1	1	1	6	8	6	18	4	-
2006 1/.....	62	-	1	-	-	1	2	8	8	10	18	14	-
2005 1/.....	61	-	-	-	2	2	-	2	9	10	27	9	-
2004 1/.....	42	-	-	-	-	2	3	9	4	10	6	8	-
2003 1/.....	52	-	-	-	-	3	4	4	7	11	18	5	-
2002.....	49	-	-	-	1	2	4	4	10	10	12	6	-
2001.....	50	-	-	-	2	2	1	2	6	14	15	8	-
2000.....	42	-	-	-	2	2	3	3	5	3	18	6	-
1999.....	66	-	-	1	-	3	4	4	11	11	21	11	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Tuberculosis (A16-A19)

Asian or Pacific Islander 3/, female

2007 1/.....	32	-	-	-	-	2	1	4	1	11	8	5	-
2006 1/.....	23	-	-	-	-	-	-	1	2	4	8	8	-
2005 1/.....	43	-	-	-	1	1	1	8	5	7	12	8	-
2004 1/.....	23	-	-	-	1	-	-	1	5	3	9	4	-
2003 1/.....	35	-	-	-	-	1	2	3	4	6	12	7	-
2002.....	43	1	-	-	1	1	-	3	1	13	15	8	-
2001.....	29	1	-	-	1	2	1	1	3	2	12	6	-
2000.....	31	-	-	-	1	-	1	-	3	8	14	4	-
1999.....	35	-	-	-	2	1	2	4	4	5	9	8	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Respiratory tuberculosis (A16)													
All races, both sexes													
2007.....	424	2	-	1	3	10	27	53	62	73	119	74	-
2006.....	490	-	-	1	2	14	22	67	83	84	132	85	-
2005.....	480	2	1	-	9	11	28	62	62	81	129	95	-
2004.....	504	1	1	1	4	11	36	58	58	108	120	106	-
2003.....	562	-	1	2	9	6	26	86	81	88	177	86	-
2002.....	626	-	-	-	6	14	40	64	84	120	192	106	-
2001.....	600	2	1	-	7	17	28	61	96	115	166	106	1
2000.....	615	-	3	-	7	11	35	72	60	119	209	98	1
1999.....	764	1	1	1	8	14	45	76	109	139	216	153	1
All races, male													
2007.....	266	1	-	-	2	4	18	42	50	41	66	42	-
2006.....	335	-	-	-	1	8	20	54	70	59	82	41	-
2005.....	291	2	1	-	4	3	23	43	46	57	68	44	-
2004.....	316	-	1	-	2	8	28	40	43	74	64	56	-
2003.....	344	-	1	2	3	6	17	64	57	61	97	36	-
2002.....	372	-	-	-	4	9	31	43	60	80	96	49	-
2001.....	371	-	-	-	5	10	18	49	78	76	84	50	1
2000.....	393	-	1	-	4	8	26	64	50	73	116	50	1
1999.....	486	-	1	1	2	8	33	64	83	97	120	76	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Respiratory tuberculosis (A16)													
All races, female													
2007.....	158	1	-	1	1	6	9	11	12	32	53	32	-
2006.....	155	-	-	1	1	6	2	13	13	25	50	44	-
2005.....	189	-	-	-	5	8	5	19	16	24	61	51	-
2004.....	188	1	-	1	2	3	8	18	15	34	56	50	-
2003.....	218	-	-	-	6	-	9	22	24	27	80	50	-
2002.....	254	-	-	-	2	5	9	21	24	40	96	57	-
2001.....	229	2	1	-	2	7	10	12	18	39	82	56	-
2000.....	222	-	2	-	3	3	9	8	10	46	93	48	-
1999.....	278	1	-	-	6	6	12	12	26	42	96	77	-
White, both sexes													
2007 1/.....	264	1	-	1	2	5	15	28	35	39	76	62	-
2006 1/.....	287	-	-	1	2	5	13	32	44	45	88	57	-
2005 1/.....	273	2	1	-	5	4	18	33	31	44	74	61	-
2004 1/.....	300	1	1	-	1	4	20	27	32	63	82	69	-
2003 1/.....	347	-	1	2	6	4	9	35	48	57	125	60	-
2002.....	374	-	-	-	2	5	20	29	43	70	129	76	-
2001.....	368	1	1	-	4	7	14	30	57	69	114	70	1
2000.....	351	-	1	-	4	5	11	35	26	66	131	72	-
1999.....	472	-	-	-	3	3	20	40	61	87	145	113	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Respiratory tuberculosis (A16)													
White, male													
2007 1/.....	165	1	-	-	1	4	9	21	31	23	39	36	-
2006 1/.....	182	-	-	-	1	2	12	26	35	30	53	23	-
2005 1/.....	160	2	1	-	2	1	14	27	24	28	34	27	-
2004 1/.....	185	-	1	-	-	2	15	17	26	43	45	36	-
2003 1/.....	208	-	1	2	3	4	7	29	35	39	64	24	-
2002.....	223	-	-	-	2	3	16	20	29	54	62	37	-
2001.....	221	-	-	-	3	5	10	23	48	42	59	30	1
2000.....	216	-	-	-	3	4	11	30	20	43	67	38	-
1999.....	296	-	-	-	2	2	15	36	45	61	79	56	-
White, female													
2007 1/.....	99	-	-	1	1	1	6	7	4	16	37	26	-
2006 1/.....	105	-	-	1	1	3	1	6	9	15	35	34	-
2005 1/.....	113	-	-	-	3	3	4	6	7	16	40	34	-
2004 1/.....	115	1	-	-	1	2	5	10	6	20	37	33	-
2003 1/.....	139	-	-	-	3	-	2	6	13	18	61	36	-
2002.....	151	-	-	-	-	2	4	9	14	16	67	39	-
2001.....	147	1	1	-	1	2	4	7	9	27	55	40	-
2000.....	135	-	1	-	1	1	-	5	6	23	64	34	-
1999.....	176	-	-	-	1	1	5	4	16	26	66	57	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Respiratory tuberculosis (A16)													
Black, both sexes													
2007 1/.....	89	1	-	-	-	3	9	17	19	17	19	4	-
2006 1/.....	121	-	-	-	-	6	8	24	30	26	19	8	-
2005 1/.....	115	-	-	-	3	5	10	20	19	23	20	15	-
2004 1/.....	141	-	-	1	2	5	15	22	17	31	23	25	-
2003 1/.....	138	-	-	-	3	2	11	41	25	17	26	13	-
2002.....	164	-	-	-	2	8	15	29	29	32	34	15	-
2001.....	160	-	-	-	1	7	12	29	31	32	28	20	-
2000.....	192	-	2	-	1	6	20	32	22	43	48	17	1
1999.....	199	1	1	-	4	8	20	29	36	33	43	23	1
Black, male													
2007 1/.....	61	-	-	-	-	-	7	16	13	12	11	2	-
2006 1/.....	91	-	-	-	-	4	7	17	27	19	13	4	-
2005 1/.....	73	-	-	-	2	1	9	14	13	20	9	5	-
2004 1/.....	88	-	-	-	2	4	12	15	13	19	11	12	-
2003 1/.....	93	-	-	-	-	2	7	29	18	13	17	7	-
2002.....	104	-	-	-	1	5	11	20	22	19	21	5	-
2001.....	106	-	-	-	-	3	7	25	24	23	12	12	-
2000.....	134	-	1	-	-	4	12	29	22	26	32	7	1
1999.....	126	-	1	-	-	3	14	24	29	22	22	10	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Respiratory tuberculosis (A16)

Black, female

2007 1/.....	28	1	-	-	-	3	2	1	6	5	8	2	-
2006 1/.....	30	-	-	-	-	2	1	7	3	7	6	4	-
2005 1/.....	42	-	-	-	1	4	1	6	6	3	11	10	-
2004 1/.....	53	-	-	1	-	1	3	7	4	12	12	13	-
2003 1/.....	45	-	-	-	3	-	4	12	7	4	9	6	-
2002.....	60	-	-	-	1	3	4	9	7	13	13	10	-
2001.....	54	-	-	-	1	4	5	4	7	9	16	8	-
2000.....	58	-	1	-	1	2	8	3	-	17	16	10	-
1999.....	73	1	-	-	4	5	6	5	7	11	21	13	-

American Indian 2/, both sexes

2007 1/.....	8	-	-	-	-	-	1	-	1	2	4	-	-
2006 1/.....	11	-	-	-	-	2	-	4	-	2	3	-	-
2005 1/.....	9	-	-	-	-	1	-	1	2	2	-	3	-
2004 1/.....	9	-	-	-	-	-	-	-	1	2	5	1	-
2003 1/.....	10	-	-	-	-	-	1	5	-	-	3	1	-
2002.....	18	-	-	-	-	-	1	2	4	3	6	2	-
2001.....	8	-	-	-	-	-	-	-	1	1	4	2	-
2000.....	12	-	-	-	-	-	-	2	5	2	3	-	-
1999.....	10	-	-	-	-	-	-	1	1	5	1	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Respiratory tuberculosis (A16)													
American Indian 2/, male													
2007 1/.....	3	-	-	-	-	-	1	-	-	-	2	-	-
2006 1/.....	7	-	-	-	-	1	-	4	-	2	-	-	-
2005 1/.....	7	-	-	-	-	1	-	1	1	1	-	3	-
2004 1/.....	6	-	-	-	-	-	-	-	1	2	3	-	-
2003 1/.....	4	-	-	-	-	-	-	3	-	-	1	-	-
2002.....	9	-	-	-	-	-	-	1	2	1	3	2	-
2001.....	3	-	-	-	-	-	-	-	1	-	2	-	-
2000.....	8	-	-	-	-	-	-	2	4	1	1	-	-
1999.....	5	-	-	-	-	-	-	1	-	3	-	1	-
American Indian 2/, female													
2007 1/.....	5	-	-	-	-	-	-	-	1	2	2	-	-
2006 1/.....	4	-	-	-	-	1	-	-	-	-	3	-	-
2005 1/.....	2	-	-	-	-	-	-	-	1	1	-	-	-
2004 1/.....	3	-	-	-	-	-	-	-	-	-	2	1	-
2003 1/.....	6	-	-	-	-	-	1	2	-	-	2	1	-
2002.....	9	-	-	-	-	-	1	1	2	2	3	-	-
2001.....	5	-	-	-	-	-	-	-	-	1	2	2	-
2000.....	4	-	-	-	-	-	-	-	1	1	2	-	-
1999.....	5	-	-	-	-	-	-	-	1	2	1	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Respiratory tuberculosis (A16)													
Asian or Pacific Islander 3/, both sexes													
2007 1/.....	63	-	-	-	1	2	2	8	7	15	20	8	-
2006 1/.....	71	-	-	-	-	1	1	7	9	11	22	20	-
2005 1/.....	83	-	-	-	1	1	-	8	10	12	35	16	-
2004 1/.....	54	-	-	-	1	2	1	9	8	12	10	11	-
2003 1/.....	67	-	-	-	-	-	5	5	8	14	23	12	-
2002.....	70	-	-	-	2	1	4	4	8	15	23	13	-
2001.....	64	1	-	-	2	3	2	2	7	13	20	14	-
2000.....	60	-	-	-	2	-	4	3	7	8	27	9	-
1999.....	83	-	-	1	1	3	5	6	11	14	27	15	-
Asian or Pacific Islander 3/, male													
2007 1/.....	37	-	-	-	1	-	1	5	6	6	14	4	-
2006 1/.....	55	-	-	-	-	1	1	7	8	8	16	14	-
2005 1/.....	51	-	-	-	-	-	-	1	8	8	25	9	-
2004 1/.....	37	-	-	-	-	2	1	8	3	10	5	8	-
2003 1/.....	39	-	-	-	-	-	3	3	4	9	15	5	-
2002.....	36	-	-	-	1	1	4	2	7	6	10	5	-
2001.....	41	-	-	-	2	2	1	1	5	11	11	8	-
2000.....	35	-	-	-	1	-	3	3	4	3	16	5	-
1999.....	59	-	-	1	-	3	4	3	9	11	19	9	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Respiratory tuberculosis (A16)

Asian or Pacific Islander 3/, female

2007 1/.....	26	-	-	-	-	2	1	3	1	9	6	4	-
2006 1/.....	16	-	-	-	-	-	-	-	1	3	6	6	-
2005 1/.....	32	-	-	-	1	1	-	7	2	4	10	7	-
2004 1/.....	17	-	-	-	1	-	-	1	5	2	5	3	-
2003 1/.....	28	-	-	-	-	-	2	2	4	5	8	7	-
2002.....	34	-	-	-	1	-	-	2	1	9	13	8	-
2001.....	23	1	-	-	-	1	1	1	2	2	9	6	-
2000.....	25	-	-	-	1	-	1	-	3	5	11	4	-
1999.....	24	-	-	-	1	-	1	3	2	3	8	6	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other tuberculosis (A17-A19)													
All races, both sexes													
2007.....	130	-	-	-	3	4	5	20	26	24	32	16	-
2006.....	162	-	4	-	2	4	9	30	18	33	36	26	-
2005.....	168	-	2	1	9	10	10	17	29	35	36	19	-
2004.....	153	-	-	1	3	5	17	22	21	29	45	10	-
2003.....	149	-	-	-	1	12	9	19	21	26	40	21	-
2002.....	158	1	-	-	3	4	12	17	23	41	42	15	-
2001.....	164	-	-	-	5	8	12	23	20	37	40	19	-
2000.....	161	1	3	1	4	9	7	19	21	36	43	17	-
1999.....	166	1	1	2	3	5	12	30	21	34	37	20	-
All races, male													
2007.....	76	-	-	-	2	3	5	15	16	12	16	7	-
2006.....	90	-	1	-	1	3	6	23	11	18	16	11	-
2005.....	101	-	1	-	8	9	7	11	18	17	21	9	-
2004.....	90	-	-	1	2	4	12	13	17	19	20	2	-
2003.....	83	-	-	-	-	7	8	15	14	11	19	9	-
2002.....	90	-	-	-	1	2	7	12	15	22	25	6	-
2001.....	82	-	-	-	3	4	9	14	7	22	16	7	-
2000.....	75	-	-	-	3	6	2	12	9	14	23	6	-
1999.....	98	-	1	1	2	3	10	23	13	19	17	9	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other tuberculosis (A17-A19)													
All races, female													
2007.....	54	-	-	-	1	1	-	5	10	12	16	9	-
2006.....	72	-	3	-	1	1	3	7	7	15	20	15	-
2005.....	67	-	1	1	1	1	3	6	11	18	15	10	-
2004.....	63	-	-	-	1	1	5	9	4	10	25	8	-
2003.....	66	-	-	-	1	5	1	4	7	15	21	12	-
2002.....	68	1	-	-	2	2	5	5	8	19	17	9	-
2001.....	82	-	-	-	2	4	3	9	13	15	24	12	-
2000.....	86	1	3	1	1	3	5	7	12	22	20	11	-
1999.....	68	1	-	1	1	2	2	7	8	15	20	11	-
White, both sexes													
2007 1/.....	79	-	-	-	2	3	1	13	13	15	18	14	-
2006 1/.....	101	-	2	-	2	4	4	13	11	20	24	21	-
2005 1/.....	103	-	1	1	4	6	7	11	15	19	22	17	-
2004 1/.....	90	-	-	1	2	4	9	11	10	15	31	7	-
2003 1/.....	90	-	-	-	-	4	2	10	12	19	25	18	-
2002.....	87	-	-	-	1	1	2	7	14	18	34	10	-
2001.....	101	-	-	-	4	3	8	12	10	21	26	17	-
2000.....	88	-	3	1	2	6	2	9	7	21	24	13	-
1999.....	92	1	-	1	2	2	5	12	12	21	27	9	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other tuberculosis (A17-A19)													
White, male													
2007 1/.....	44	-	-	-	2	2	1	10	7	7	9	6	-
2006 1/.....	55	-	-	-	1	3	3	10	8	11	10	9	-
2005 1/.....	66	-	1	-	4	6	5	8	10	10	14	8	-
2004 1/.....	52	-	-	1	2	3	4	9	8	10	13	2	-
2003 1/.....	47	-	-	-	-	1	2	8	8	7	13	8	-
2002.....	49	-	-	-	-	-	1	6	8	10	21	3	-
2001.....	50	-	-	-	3	3	6	7	3	11	10	7	-
2000.....	37	-	-	-	1	3	-	4	4	10	11	4	-
1999.....	52	-	-	1	2	2	4	10	7	12	12	2	-
White, female													
2007 1/.....	35	-	-	-	-	1	-	3	6	8	9	8	-
2006 1/.....	46	-	2	-	1	1	1	3	3	9	14	12	-
2005 1/.....	37	-	-	1	-	-	2	3	5	9	8	9	-
2004 1/.....	38	-	-	-	-	1	5	2	2	5	18	5	-
2003 1/.....	43	-	-	-	-	3	-	2	4	12	12	10	-
2002.....	38	-	-	-	1	1	1	1	6	8	13	7	-
2001.....	51	-	-	-	1	-	2	5	7	10	16	10	-
2000.....	51	-	3	1	1	3	2	5	3	11	13	9	-
1999.....	40	1	-	-	-	-	1	2	5	9	15	7	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other tuberculosis (A17-A19)													
Black, both sexes													
2007 1/.....	33	-	-	-	1	-	4	5	8	7	7	1	-
2006 1/.....	39	-	1	-	-	-	3	12	4	10	7	2	-
2005 1/.....	38	-	1	-	3	2	-	4	9	9	9	1	-
2004 1/.....	48	-	-	-	1	1	6	10	9	10	9	2	-
2003 1/.....	34	-	-	-	1	4	5	6	6	3	7	2	-
2002.....	40	-	-	-	2	-	9	6	5	11	3	4	-
2001.....	43	-	-	-	-	3	4	8	7	12	7	2	-
2000.....	52	-	-	-	1	1	5	10	12	9	12	2	-
1999.....	46	-	1	1	-	1	5	13	4	10	7	4	-
Black, male													
2007 1/.....	22	-	-	-	-	-	4	4	5	5	3	1	-
2006 1/.....	24	-	-	-	-	-	2	10	2	5	3	2	-
2005 1/.....	21	-	-	-	2	1	-	2	6	5	4	1	-
2004 1/.....	30	-	-	-	-	1	6	3	7	7	6	-	-
2003 1/.....	21	-	-	-	-	3	4	6	3	2	3	-	-
2002.....	23	-	-	-	1	-	6	4	3	6	1	2	-
2001.....	21	-	-	-	-	1	3	5	3	7	2	-	-
2000.....	27	-	-	-	1	1	2	8	4	2	8	1	-
1999.....	33	-	1	-	-	1	5	10	3	7	3	3	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other tuberculosis (A17-A19)													
Black, female													
2007 1/.....	11	-	-	-	1	-	-	1	3	2	4	-	-
2006 1/.....	15	-	1	-	-	-	1	2	2	5	4	-	-
2005 1/.....	17	-	1	-	1	1	-	2	3	4	5	-	-
2004 1/.....	18	-	-	-	1	-	-	7	2	3	3	2	-
2003 1/.....	13	-	-	-	1	1	1	-	3	1	4	2	-
2002.....	17	-	-	-	1	-	3	2	2	5	2	2	-
2001.....	22	-	-	-	-	2	1	3	4	5	5	2	-
2000.....	25	-	-	-	-	-	3	2	8	7	4	1	-
1999.....	13	-	-	1	-	-	-	3	1	3	4	1	-
American Indian 2/, both sexes													
2007 1/.....	4	-	-	-	-	-	-	-	3	-	1	-	-
2006 1/.....	8	-	-	-	-	-	1	3	2	-	1	1	-
2005 1/.....	6	-	-	-	-	-	2	-	1	2	1	-	-
2004 1/.....	4	-	-	-	-	-	-	-	1	3	-	-	-
2003 1/.....	5	-	-	-	-	-	1	1	-	1	1	1	-
2002.....	9	-	-	-	-	1	1	1	1	4	1	-	-
2001.....	5	-	-	-	-	1	-	2	1	1	-	-	-
2000.....	8	1	-	-	-	-	-	-	1	3	2	1	-
1999.....	10	-	-	-	-	1	1	3	1	1	-	3	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other tuberculosis (A17-A19)													
American Indian 2/, male													
2007 1/.....	2	-	-	-	-	-	-	-	2	-	-	-	-
2006 1/.....	4	-	-	-	-	-	-	2	1	-	1	-	-
2005 1/.....	4	-	-	-	-	-	2	-	1	-	1	-	-
2004 1/.....	3	-	-	-	-	-	-	-	1	2	-	-	-
2003 1/.....	2	-	-	-	-	-	1	-	-	-	-	1	-
2002.....	5	-	-	-	-	1	-	-	1	2	1	-	-
2001.....	2	-	-	-	-	-	-	1	-	1	-	-	-
2000.....	4	-	-	-	-	-	-	-	-	2	2	-	-
1999.....	6	-	-	-	-	-	1	2	1	-	-	2	-
American Indian 2/, female													
2007 1/.....	2	-	-	-	-	-	-	-	1	-	1	-	-
2006 1/.....	4	-	-	-	-	-	1	1	1	-	-	1	-
2005 1/.....	2	-	-	-	-	-	-	-	-	2	-	-	-
2004 1/.....	1	-	-	-	-	-	-	-	-	1	-	-	-
2003 1/.....	3	-	-	-	-	-	-	1	-	1	1	-	-
2002.....	4	-	-	-	-	-	1	1	-	2	-	-	-
2001.....	3	-	-	-	-	1	-	1	-	-	-	-	-
2000.....	4	1	-	-	-	-	-	-	1	1	-	1	-
1999.....	4	-	-	-	-	1	-	1	-	1	-	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other tuberculosis (A17-A19)													
Asian or Pacific Islander 3/, both sexes													
2007 1/.....	14	-	-	-	-	1	-	2	2	2	6	1	-
2006 1/.....	14	-	1	-	-	-	1	2	1	3	4	2	-
2005 1/.....	21	-	-	-	2	2	1	2	4	5	4	1	-
2004 1/.....	11	-	-	-	-	-	2	1	1	1	5	1	-
2003 1/.....	20	-	-	-	-	4	1	2	3	3	7	-	-
2002.....	22	1	-	-	-	2	-	3	3	8	4	1	-
2001.....	15	-	-	-	1	1	-	1	2	3	7	-	-
2000.....	13	-	-	-	1	2	-	-	1	3	5	1	-
1999.....	18	-	-	-	1	1	1	2	4	2	3	4	-
Asian or Pacific Islander 3/, male													
2007 1/.....	8	-	-	-	-	1	-	1	2	-	4	-	-
2006 1/.....	7	-	1	-	-	-	1	1	-	2	2	-	-
2005 1/.....	10	-	-	-	2	2	-	1	1	2	2	-	-
2004 1/.....	5	-	-	-	-	-	2	1	1	-	1	-	-
2003 1/.....	13	-	-	-	-	3	1	1	3	2	3	-	-
2002.....	13	-	-	-	-	1	-	2	3	4	2	1	-
2001.....	9	-	-	-	-	-	-	1	1	3	4	-	-
2000.....	7	-	-	-	1	2	-	-	1	-	2	1	-
1999.....	7	-	-	-	-	-	-	1	2	-	2	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other tuberculosis (A17-A19)

Asian or Pacific Islander 3/, female

2007 1/.....	6	-	-	-	-	-	-	1	-	2	2	1	-
2006 1/.....	7	-	-	-	-	-	-	1	1	1	2	2	-
2005 1/.....	11	-	-	-	-	-	1	1	3	3	2	1	-
2004 1/.....	6	-	-	-	-	-	-	-	-	1	4	1	-
2003 1/.....	7	-	-	-	-	1	-	1	-	1	4	-	-
2002.....	9	1	-	-	-	1	-	1	-	4	2	-	-
2001.....	6	-	-	-	1	1	-	-	1	-	3	-	-
2000.....	6	-	-	-	-	-	-	-	-	3	3	-	-
1999.....	11	-	-	-	1	1	1	1	2	2	1	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Scarlet fever and erysipelas (A38,A46)

All races, female

2007.....	2	-	-	-	-	1	-	-	-	-	-	1	-
2006.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2005.....	1	-	-	-	-	-	-	-	-	-	-	1	-
2004.....	2	-	-	-	-	-	-	-	-	-	-	2	-
2003.....	1	-	-	-	-	-	-	-	-	-	-	1	-
2002.....	2	-	-	-	-	-	-	-	-	-	1	1	-
2001.....	1	-	-	-	-	-	-	-	-	1	-	-	-
2000.....	1	-	-	-	-	-	-	1	-	-	-	-	-
1999.....	5	-	-	-	-	-	-	-	-	2	1	2	-

White, both sexes

2007 1/.....	2	-	-	-	-	-	-	-	-	-	-	2	-
2006 1/.....	2	-	-	1	-	-	-	1	-	-	-	-	-
2005 1/.....	3	-	1	-	-	-	-	-	-	-	-	2	-
2004 1/.....	3	-	-	-	-	-	-	-	-	1	-	2	-
2003 1/.....	1	-	-	-	-	-	-	-	-	-	-	1	-
2002.....	2	-	-	-	-	-	-	-	-	-	1	1	-
2001.....	3	-	-	1	-	-	-	-	-	1	-	1	-
2000.....	2	-	-	-	-	-	-	1	-	-	1	-	-
1999.....	4	-	-	-	-	-	-	-	-	1	1	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Scarlet fever and erysipelas (A38,A46)

White, male

2007 1/.....	1	-	-	-	-	-	-	-	-	-	-	-	1	-
2006 1/.....	2	-	-	1	-	-	-	1	-	-	-	-	-	-
2005 1/.....	2	-	1	-	-	-	-	-	-	-	-	-	1	-
2004 1/.....	1	-	-	-	-	-	-	-	-	1	-	-	-	-
2003 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	2	-	-	1	-	-	-	-	-	-	-	-	1	-
2000.....	1	-	-	-	-	-	-	-	-	-	1	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-	-	-

White, female

2007 1/.....	1	-	-	-	-	-	-	-	-	-	-	-	1	-
2006 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2005 1/.....	1	-	-	-	-	-	-	-	-	-	-	-	1	-
2004 1/.....	2	-	-	-	-	-	-	-	-	-	-	-	2	-
2003 1/.....	1	-	-	-	-	-	-	-	-	-	-	-	1	-
2002.....	2	-	-	-	-	-	-	-	-	-	1	1	-	-
2001.....	1	-	-	-	-	-	-	-	-	1	-	-	-	-
2000.....	1	-	-	-	-	-	-	1	-	-	-	-	-	-
1999.....	4	-	-	-	-	-	-	-	-	1	1	2	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Meningococcal infection (A39)

All races, both sexes

2007.....	87	11	5	11	21	6	9	8	8	4	1	3	-
2006.....	105	11	14	7	22	13	6	13	9	4	4	2	-
2005.....	123	17	11	7	29	8	8	18	10	4	6	5	-
2004.....	138	8	18	13	31	14	15	15	6	7	3	8	-
2003.....	161	12	18	11	39	12	19	18	11	9	9	3	-
2002.....	161	17	12	20	33	16	9	19	13	7	10	5	-
2001.....	199	16	19	26	55	10	13	13	15	15	8	9	-
2000.....	211	22	19	18	56	15	13	29	9	8	15	7	-
1999.....	227	32	30	25	39	23	27	17	4	9	13	8	-

All races, male

2007.....	48	6	3	8	12	3	3	5	3	3	1	1	-
2006.....	67	10	10	3	16	9	4	8	4	2	1	-	-
2005.....	68	10	6	5	17	8	4	11	4	2	-	1	-
2004.....	79	2	11	9	19	10	8	12	4	3	1	-	-
2003.....	75	9	10	3	18	7	9	10	5	2	1	1	-
2002.....	83	8	9	13	17	10	5	10	4	2	3	2	-
2001.....	94	6	9	15	27	6	6	8	7	5	3	2	-
2000.....	106	14	12	7	35	6	9	11	2	2	7	1	-
1999.....	101	14	19	13	19	12	9	4	-	5	4	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Meningococcal infection (A39)													
All races, female													
2007.....	39	5	2	3	9	3	6	3	5	1	-	2	-
2006.....	38	1	4	4	6	4	2	5	5	2	3	2	-
2005.....	55	7	5	2	12	-	4	7	6	2	6	4	-
2004.....	59	6	7	4	12	4	7	3	2	4	2	8	-
2003.....	86	3	8	8	21	5	10	8	6	7	8	2	-
2002.....	78	9	3	7	16	6	4	9	9	5	7	3	-
2001.....	105	10	10	11	28	4	7	5	8	10	5	7	-
2000.....	105	8	7	11	21	9	4	18	7	6	8	6	-
1999.....	126	18	11	12	20	11	18	13	4	4	9	6	-
White, both sexes													
2007 1/.....	58	7	4	7	13	4	5	3	7	4	1	3	-
2006 1/.....	79	10	13	4	16	7	4	8	8	4	3	2	-
2005 1/.....	101	13	11	6	23	6	5	15	9	3	5	5	-
2004 1/.....	110	7	15	12	24	10	10	11	5	6	3	7	-
2003 1/.....	121	12	11	7	30	8	14	11	7	9	9	3	-
2002.....	125	14	8	10	27	13	5	16	11	6	10	5	-
2001.....	158	14	14	20	40	7	11	12	11	14	7	8	-
2000.....	161	19	14	10	45	10	7	19	8	7	15	7	-
1999.....	184	30	26	18	29	14	19	15	4	9	12	8	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Meningococcal infection (A39)

White, male

2007 1/.....	32	4	3	5	8	2	1	1	3	3	1	1	-
2006 1/.....	50	9	9	1	11	5	2	6	4	2	1	-	-
2005 1/.....	53	7	6	4	13	6	2	9	4	1	-	1	-
2004 1/.....	62	2	10	8	13	8	6	9	3	2	1	-	-
2003 1/.....	56	9	6	2	14	5	7	7	2	2	1	1	-
2002.....	67	8	6	9	14	9	3	8	4	1	3	2	-
2001.....	72	5	7	12	20	4	4	7	5	5	2	1	-
2000.....	75	11	8	4	28	2	5	5	2	2	7	1	-
1999.....	79	14	17	9	13	6	6	3	-	5	4	2	-

White, female

2007 1/.....	26	3	1	2	5	2	4	2	4	1	-	2	-
2006 1/.....	29	1	4	3	5	2	2	2	4	2	2	2	-
2005 1/.....	48	6	5	2	10	-	3	6	5	2	5	4	-
2004 1/.....	48	5	5	4	11	2	4	2	2	4	2	7	-
2003 1/.....	65	3	5	5	16	3	7	4	5	7	8	2	-
2002.....	58	6	2	1	13	4	2	8	7	5	7	3	-
2001.....	86	9	7	8	20	3	7	5	6	9	5	7	-
2000.....	86	8	6	6	17	8	2	14	6	5	8	6	-
1999.....	105	16	9	9	16	8	13	12	4	4	8	6	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Meningococcal infection (A39)

Black, both sexes

2007 1/.....	26	4	1	3	8	2	2	5	1	-	-	-	-
2006 1/.....	23	1	1	3	6	3	2	5	1	-	1	-	-
2005 1/.....	17	2	-	-	6	2	2	2	1	1	1	-	-
2004 1/.....	22	-	3	1	5	3	4	3	1	1	-	1	-
2003 1/.....	36	-	7	4	7	3	5	6	4	-	-	-	-
2002.....	29	3	3	8	5	2	4	3	1	-	-	-	-
2001.....	36	2	4	6	13	2	1	1	4	1	1	1	-
2000.....	44	3	5	8	7	4	6	9	1	1	-	-	-
1999.....	39	2	4	6	9	8	8	2	-	-	-	-	-

Black, male

2007 1/.....	14	2	-	2	4	1	1	4	-	-	-	-	-
2006 1/.....	15	1	1	2	5	2	2	2	-	-	-	-	-
2005 1/.....	11	1	-	-	4	2	1	2	-	1	-	-	-
2004 1/.....	13	-	1	1	4	2	1	2	1	1	-	-	-
2003 1/.....	17	-	4	1	3	1	2	3	3	-	-	-	-
2002.....	12	-	2	4	2	-	2	2	-	-	-	-	-
2001.....	18	1	1	3	6	1	1	1	2	-	1	1	-
2000.....	27	3	4	3	5	3	4	5	-	-	-	-	-
1999.....	20	-	2	3	6	5	3	1	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Meningococcal infection (A39)

Black, female

2007 1/.....	12	2	1	1	4	1	1	1	1	-	-	-	-
2006 1/.....	8	-	-	1	1	1	-	3	1	-	1	-	-
2005 1/.....	6	1	-	-	2	-	1	-	1	-	1	-	-
2004 1/.....	9	-	2	-	1	1	3	1	-	-	-	1	-
2003 1/.....	19	-	3	3	4	2	3	3	1	-	-	-	-
2002.....	17	3	1	4	3	2	2	1	1	-	-	-	-
2001.....	18	1	3	3	7	1	-	-	2	1	-	-	-
2000.....	17	-	1	5	2	1	2	4	1	1	-	-	-
1999.....	19	2	2	3	3	3	5	1	-	-	-	-	-

American Indian 2/, both sexes

2007 1/.....	2	-	-	-	-	-	2	-	-	-	-	-	-
2006 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2005 1/.....	2	1	-	-	-	-	-	1	-	-	-	-	-
2004 1/.....	3	-	-	-	-	1	1	1	-	-	-	-	-
2003 1/.....	3	-	-	-	2	-	-	1	-	-	-	-	-
2002.....	2	-	1	1	-	-	-	-	-	-	-	-	-
2001.....	1	-	1	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	1	-	-	-	-	-	-	-	-
1999.....	2	-	-	1	-	-	-	-	-	-	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Meningococcal infection (A39)

American Indian 2/, male

2007 1/.....	1	-	-	-	-	-	1	-	-	-	-	-	-
2006 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2005 1/.....	1	1	-	-	-	-	-	-	-	-	-	-	-
2004 1/.....	2	-	-	-	-	-	1	1	-	-	-	-	-
2003 1/.....	1	-	-	-	1	-	-	-	-	-	-	-	-
2002.....	1	-	1	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	1	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	1	-	-	-	-	-	-	-	-	-

American Indian 2/, female

2007 1/.....	1	-	-	-	-	-	1	-	-	-	-	-	-
2006 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2005 1/.....	1	-	-	-	-	-	-	1	-	-	-	-	-
2004 1/.....	1	-	-	-	-	1	-	-	-	-	-	-	-
2003 1/.....	2	-	-	-	1	-	-	1	-	-	-	-	-
2002.....	1	-	-	1	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	1	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	-	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Meningococcal infection (A39)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	1	-	-	1	-	-	-	-	-	-	-	-	-
2006 1/.....	3	-	-	-	-	3	-	-	-	-	-	-	-
2005 1/.....	3	1	-	1	-	-	1	-	-	-	-	-	-
2004 1/.....	3	1	-	-	2	-	-	-	-	-	-	-	-
2003 1/.....	1	-	-	-	-	1	-	-	-	-	-	-	-
2002.....	5	-	-	1	1	1	-	-	1	1	-	-	-
2001.....	4	-	-	-	2	1	1	-	-	-	-	-	-
2000.....	5	-	-	-	3	1	-	1	-	-	-	-	-
1999.....	2	-	-	-	1	1	-	-	-	-	-	-	-

Asian or Pacific Islander 3/, male

2007 1/.....	1	-	-	1	-	-	-	-	-	-	-	-	-
2006 1/.....	2	-	-	-	-	2	-	-	-	-	-	-	-
2005 1/.....	3	1	-	1	-	-	1	-	-	-	-	-	-
2004 1/.....	2	-	-	-	2	-	-	-	-	-	-	-	-
2003 1/.....	1	-	-	-	-	1	-	-	-	-	-	-	-
2002.....	3	-	-	-	1	1	-	-	-	1	-	-	-
2001.....	3	-	-	-	1	1	1	-	-	-	-	-	-
2000.....	4	-	-	-	2	1	-	1	-	-	-	-	-
1999.....	1	-	-	-	-	1	-	-	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Septicemia (A40-A41)													
All races, both sexes													
2007.....	34,828	283	78	74	160	297	910	2,431	4,231	6,345	10,403	9,614	2
2006.....	34,234	269	88	84	139	291	870	2,260	4,032	6,064	10,745	9,392	-
2005.....	34,136	302	85	81	148	311	840	2,211	3,912	6,073	10,626	9,544	3
2004.....	33,373	271	84	66	135	328	846	2,251	3,745	5,983	10,586	9,075	3
2003.....	34,069	278	85	77	154	309	910	2,157	3,651	5,970	10,932	9,543	3
2002.....	33,865	296	79	95	118	314	856	2,074	3,360	6,336	11,010	9,324	3
2001.....	32,238	312	108	69	133	281	829	1,977	3,111	5,998	10,351	9,069	-
2000.....	31,224	274	99	63	100	280	877	1,845	2,899	5,704	9,938	9,144	1
1999.....	30,680	283	87	77	121	275	812	1,685	2,714	5,750	9,710	9,166	-
All races, male													
2007.....	15,839	152	44	45	85	153	463	1,305	2,203	3,256	4,768	3,364	1
2006.....	15,522	143	43	52	68	152	452	1,247	2,144	3,019	4,914	3,288	-
2005.....	15,322	171	47	49	82	152	414	1,175	2,074	3,095	4,823	3,238	2
2004.....	15,011	167	41	31	71	165	433	1,207	1,934	3,017	4,870	3,074	1
2003.....	14,987	151	54	34	82	161	455	1,155	1,887	3,023	4,894	3,089	2
2002.....	14,947	158	41	52	59	166	441	1,124	1,748	3,154	5,028	2,973	3
2001.....	14,307	174	60	39	75	139	451	1,043	1,606	3,044	4,762	2,914	-
2000.....	13,537	139	53	38	60	158	451	1,039	1,513	2,867	4,482	2,736	1
1999.....	13,395	175	48	36	67	147	460	919	1,412	2,919	4,426	2,786	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Septicemia (A40-A41)													
All races, female													
2007.....	18,989	131	34	29	75	144	447	1,126	2,028	3,089	5,635	6,250	1
2006.....	18,712	126	45	32	71	139	418	1,013	1,888	3,045	5,831	6,104	-
2005.....	18,814	131	38	32	66	159	426	1,036	1,838	2,978	5,803	6,306	1
2004.....	18,362	104	43	35	64	163	413	1,044	1,811	2,966	5,716	6,001	2
2003.....	19,082	127	31	43	72	148	455	1,002	1,764	2,947	6,038	6,454	1
2002.....	18,918	138	38	43	59	148	415	950	1,612	3,182	5,982	6,351	-
2001.....	17,931	138	48	30	58	142	378	934	1,505	2,954	5,589	6,155	-
2000.....	17,687	135	46	25	40	122	426	806	1,386	2,837	5,456	6,408	-
1999.....	17,285	108	39	41	54	128	352	766	1,302	2,831	5,284	6,380	-
White, both sexes													
2007 1/.....	27,750	158	60	59	106	194	608	1,628	3,120	4,852	8,684	8,280	1
2006 1/.....	27,373	156	59	63	90	186	572	1,509	2,929	4,668	9,001	8,140	-
2005 1/.....	27,194	160	59	57	95	191	552	1,499	2,877	4,751	8,829	8,123	1
2004 1/.....	26,678	165	60	50	82	206	559	1,515	2,727	4,649	8,859	7,804	2
2003 1/.....	27,180	142	53	54	103	190	598	1,439	2,666	4,587	9,210	8,136	2
2002.....	27,110	191	58	69	86	202	574	1,410	2,410	4,938	9,202	7,968	2
2001.....	25,806	171	80	54	82	169	531	1,312	2,240	4,677	8,708	7,782	-
2000.....	24,804	169	66	49	60	165	578	1,219	2,066	4,418	8,223	7,790	1
1999.....	24,287	164	60	55	79	182	510	1,107	1,928	4,411	8,035	7,756	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Septicemia (A40-A41)													
White, male													
2007 1/.....	12,600	85	33	38	59	103	319	878	1,616	2,494	4,068	2,906	1
2006 1/.....	12,450	84	29	38	43	99	306	856	1,549	2,373	4,174	2,899	-
2005 1/.....	12,172	96	33	33	56	97	276	800	1,544	2,423	4,058	2,756	-
2004 1/.....	11,986	103	29	26	46	101	294	825	1,411	2,370	4,105	2,675	1
2003 1/.....	11,909	74	35	26	56	95	303	791	1,378	2,344	4,150	2,656	1
2002.....	11,919	100	34	42	45	109	294	776	1,290	2,482	4,193	2,552	2
2001.....	11,371	100	46	32	46	85	309	695	1,160	2,378	4,029	2,491	-
2000.....	10,716	88	35	30	39	95	311	693	1,097	2,264	3,737	2,326	1
1999.....	10,489	106	32	28	49	102	294	607	983	2,247	3,695	2,346	-
White, female													
2007 1/.....	15,150	73	27	21	47	91	289	750	1,504	2,358	4,616	5,374	-
2006 1/.....	14,923	72	30	25	47	87	266	653	1,380	2,295	4,827	5,241	-
2005 1/.....	15,022	64	26	24	39	94	276	699	1,333	2,328	4,771	5,367	1
2004 1/.....	14,692	62	31	24	36	105	265	690	1,316	2,279	4,754	5,129	1
2003 1/.....	15,271	68	18	28	47	95	295	648	1,288	2,243	5,060	5,480	1
2002.....	15,191	91	24	27	41	93	280	634	1,120	2,456	5,009	5,416	-
2001.....	14,435	71	34	22	36	84	222	617	1,080	2,299	4,679	5,291	-
2000.....	14,088	81	31	19	21	70	267	526	969	2,154	4,486	5,464	-
1999.....	13,798	58	28	27	30	80	216	500	945	2,164	4,340	5,410	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Septicemia (A40-A41)													
Black, both sexes													
2007 1/.....	6,297	116	16	11	47	90	274	721	1,004	1,320	1,519	1,179	-
2006 1/.....	6,108	102	21	20	41	89	273	680	997	1,245	1,532	1,108	-
2005 1/.....	6,221	131	20	21	44	108	262	658	942	1,183	1,590	1,260	2
2004 1/.....	6,010	96	19	12	46	104	257	664	916	1,211	1,548	1,136	1
2003 1/.....	6,206	123	24	15	41	102	285	656	891	1,247	1,546	1,275	1
2002.....	6,137	91	17	20	24	103	258	600	876	1,260	1,651	1,236	1
2001.....	5,880	126	22	10	41	98	269	600	814	1,198	1,514	1,188	-
2000.....	5,928	101	26	11	33	108	272	575	768	1,189	1,590	1,255	-
1999.....	5,826	107	17	14	33	85	282	525	722	1,223	1,525	1,293	-
Black, male													
2007 1/.....	2,835	63	11	5	22	41	127	380	528	659	609	390	-
2006 1/.....	2,717	54	12	13	21	45	134	354	534	570	648	332	-
2005 1/.....	2,786	69	13	14	24	50	122	346	488	594	661	403	2
2004 1/.....	2,690	55	9	4	22	53	126	345	468	585	679	344	-
2003 1/.....	2,775	71	14	6	19	56	135	340	466	609	670	388	1
2002.....	2,703	51	5	7	10	48	133	315	423	603	740	367	1
2001.....	2,635	67	12	3	25	46	128	309	417	591	663	374	-
2000.....	2,587	50	13	6	18	59	125	319	387	555	684	371	-
1999.....	2,623	61	11	7	14	41	155	283	394	614	656	387	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Septicemia (A40-A41)													
Black, female													
2007 1/.....	3,462	53	5	6	25	49	147	341	476	661	910	789	-
2006 1/.....	3,391	48	9	7	20	44	139	326	463	675	884	776	-
2005 1/.....	3,435	62	7	7	20	58	140	312	454	589	929	857	-
2004 1/.....	3,320	41	10	8	24	51	131	319	448	626	869	792	1
2003 1/.....	3,431	52	10	9	22	46	150	316	425	638	876	887	-
2002.....	3,434	40	12	13	14	55	125	285	453	657	911	869	-
2001.....	3,245	59	10	7	16	52	141	291	397	607	851	814	-
2000.....	3,341	51	13	5	15	49	147	256	381	634	906	884	-
1999.....	3,203	46	6	7	19	44	127	242	328	609	869	906	-
American Indian 2/, both sexes													
2007 1/.....	230	4	-	2	2	6	12	46	44	49	41	24	-
2006 1/.....	207	3	2	-	3	6	13	37	40	37	45	21	-
2005 1/.....	217	2	1	-	3	4	16	24	45	47	45	30	-
2004 1/.....	183	5	2	4	2	9	16	31	33	25	31	25	-
2003 1/.....	227	4	4	4	4	8	22	23	45	46	42	25	-
2002.....	195	4	3	2	2	4	13	29	27	46	40	25	-
2001.....	155	4	3	2	3	4	15	29	21	39	21	14	-
2000.....	162	1	3	1	2	3	13	21	29	33	34	22	-
1999.....	169	3	1	3	4	3	12	26	29	33	32	23	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Septicemia (A40-A41)

American Indian 2/, male

2007 1/.....	102	2	-	1	1	4	7	24	19	22	16	6	-
2006 1/.....	99	1	-	-	2	-	5	19	23	21	19	9	-
2005 1/.....	104	-	-	-	-	1	10	12	22	29	16	14	-
2004 1/.....	85	5	2	1	1	4	8	15	14	12	13	10	-
2003 1/.....	89	2	2	1	4	5	14	7	20	17	11	6	-
2002.....	95	2	2	2	-	4	6	16	11	23	23	6	-
2001.....	83	2	1	2	1	2	7	17	11	20	13	7	-
2000.....	74	1	2	1	1	2	7	12	14	15	14	5	-
1999.....	81	3	-	-	2	2	7	15	13	18	14	7	-

American Indian 2/, female

2007 1/.....	128	2	-	1	1	2	5	22	25	27	25	18	-
2006 1/.....	108	2	2	-	1	6	8	18	17	16	26	12	-
2005 1/.....	113	2	1	-	3	3	6	12	23	18	29	16	-
2004 1/.....	98	-	-	3	1	5	8	16	19	13	18	15	-
2003 1/.....	138	2	2	3	-	3	8	16	25	29	31	19	-
2002.....	100	2	1	-	2	-	7	13	16	23	17	19	-
2001.....	72	2	2	-	2	2	8	12	10	19	8	7	-
2000.....	88	-	1	-	1	1	6	9	15	18	20	17	-
1999.....	88	-	1	3	2	1	5	11	16	15	18	16	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Septicemia (A40-A41)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	551	5	2	2	5	7	16	36	63	124	159	131	1
2006 1/.....	546	8	6	1	5	10	12	34	66	114	167	123	-
2005 1/.....	504	9	5	3	6	8	10	30	48	92	162	131	-
2004 1/.....	502	5	3	-	5	9	14	41	69	98	148	110	-
2003 1/.....	456	9	4	4	6	9	5	39	49	90	134	107	-
2002.....	423	10	1	4	6	5	11	35	47	92	117	95	-
2001.....	397	11	3	3	7	10	14	36	36	84	108	85	-
2000.....	330	3	4	2	5	4	14	30	36	64	91	77	-
1999.....	398	9	9	5	5	5	8	27	35	83	118	94	-

Asian or Pacific Islander 3/, male

2007 1/.....	302	2	-	1	3	5	10	23	40	81	75	62	-
2006 1/.....	256	4	2	1	2	8	7	18	38	55	73	48	-
2005 1/.....	260	6	1	2	2	4	6	17	20	49	88	65	-
2004 1/.....	250	4	1	-	2	7	5	22	41	50	73	45	-
2003 1/.....	214	4	3	1	3	5	3	17	23	53	63	39	-
2002.....	230	5	-	1	4	5	8	17	24	46	72	48	-
2001.....	218	5	1	2	3	6	7	22	18	55	57	42	-
2000.....	160	-	3	1	2	2	8	15	15	33	47	34	-
1999.....	202	5	5	1	2	2	4	14	22	40	61	46	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Septicemia (A40-A41)

Asian or Pacific Islander 3/, female

2007 1/.....	249	3	2	1	2	2	6	13	23	43	84	69	1
2006 1/.....	290	4	4	-	3	2	5	16	28	59	94	75	-
2005 1/.....	244	3	4	1	4	4	4	13	28	43	74	66	-
2004 1/.....	252	1	2	-	3	2	9	19	28	48	75	65	-
2003 1/.....	242	5	1	3	3	4	2	22	26	37	71	68	-
2002.....	193	5	1	3	2	-	3	18	23	46	45	47	-
2001.....	179	6	2	1	4	4	7	14	18	29	51	43	-
2000.....	170	3	1	1	3	2	6	15	21	31	44	43	-
1999.....	196	4	4	4	3	3	4	13	13	43	57	48	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Syphilis (A50-A53)													
All races, both sexes													
2007.....	42	5	-	-	-	-	4	2	1	8	12	10	-
2006.....	38	-	1	-	1	-	4	4	7	4	7	10	-
2005.....	47	-	-	1	-	4	5	6	8	4	13	6	-
2004.....	43	-	-	-	-	3	3	3	4	6	12	12	-
2003.....	34	-	-	-	-	-	3	5	3	6	8	9	-
2002.....	41	1	1	-	-	-	2	5	5	6	12	9	-
2001.....	36	1	-	1	-	1	3	3	4	5	13	5	-
2000.....	41	1	-	-	-	1	-	6	3	7	13	10	-
1999.....	33	2	-	-	-	1	1	5	4	6	10	4	-
All races, male													
2007.....	25	3	-	-	-	-	2	2	1	8	6	3	-
2006.....	20	-	1	-	-	-	2	3	4	3	3	4	-
2005.....	32	-	-	-	-	4	4	5	5	4	8	2	-
2004.....	21	-	-	-	-	2	2	1	2	4	6	4	-
2003.....	22	-	-	-	-	-	3	5	2	5	4	3	-
2002.....	21	-	1	-	-	-	2	4	2	5	4	3	-
2001.....	19	1	-	1	-	-	2	3	3	3	5	1	-
2000.....	24	1	-	-	-	1	-	5	3	2	7	5	-
1999.....	13	-	-	-	-	1	-	3	3	5	-	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Syphilis (A50-A53)

All races, female

2007.....	17	2	-	-	-	-	2	-	-	-	6	7	-
2006.....	18	-	-	-	1	-	2	1	3	1	4	6	-
2005.....	15	-	-	1	-	-	1	1	3	-	5	4	-
2004.....	22	-	-	-	-	1	1	2	2	2	6	8	-
2003.....	12	-	-	-	-	-	-	-	1	1	4	6	-
2002.....	20	1	-	-	-	-	-	1	3	1	8	6	-
2001.....	17	-	-	-	-	1	1	-	1	2	8	4	-
2000.....	17	-	-	-	-	-	-	1	-	5	6	5	-
1999.....	20	2	-	-	-	-	1	2	1	1	10	3	-

White, both sexes

2007 1/.....	20	2	-	-	-	-	3	-	-	3	7	5	-
2006 1/.....	13	-	-	-	-	-	2	-	2	-	4	5	-
2005 1/.....	16	-	-	-	-	-	1	1	3	1	8	2	-
2004 1/.....	17	-	-	-	-	-	3	1	1	3	6	3	-
2003 1/.....	17	-	-	-	-	-	1	1	2	2	5	6	-
2002.....	18	-	-	-	-	-	1	2	1	3	5	6	-
2001.....	21	1	-	-	-	-	1	1	3	2	10	3	-
2000.....	21	-	-	-	-	1	-	2	-	3	7	8	-
1999.....	20	2	-	-	-	1	1	5	3	3	4	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Syphilis (A50-A53)													
White, male													
2007 1/.....	12	2	-	-	-	-	2	-	-	3	3	2	-
2006 1/.....	5	-	-	-	-	-	2	-	-	-	1	2	-
2005 1/.....	10	-	-	-	-	-	1	1	2	1	5	-	-
2004 1/.....	7	-	-	-	-	-	2	-	1	1	2	1	-
2003 1/.....	10	-	-	-	-	-	1	1	1	2	3	2	-
2002.....	9	-	-	-	-	-	1	2	-	3	1	2	-
2001.....	9	1	-	-	-	-	-	1	2	1	3	1	-
2000.....	7	-	-	-	-	1	-	1	-	-	2	3	-
1999.....	8	-	-	-	-	1	-	3	2	2	-	-	-
White, female													
2007 1/.....	8	-	-	-	-	-	1	-	-	-	4	3	-
2006 1/.....	8	-	-	-	-	-	-	-	2	-	3	3	-
2005 1/.....	6	-	-	-	-	-	-	-	1	-	3	2	-
2004 1/.....	10	-	-	-	-	-	1	1	-	2	4	2	-
2003 1/.....	7	-	-	-	-	-	-	-	1	-	2	4	-
2002.....	9	-	-	-	-	-	-	-	1	-	4	4	-
2001.....	12	-	-	-	-	-	1	-	1	1	7	2	-
2000.....	14	-	-	-	-	-	-	1	-	3	5	5	-
1999.....	12	2	-	-	-	-	1	2	1	1	4	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Syphilis (A50-A53)

Black, both sexes

2007 1/.....	21	3	-	-	-	-	1	2	1	4	5	5	-
2006 1/.....	24	-	1	-	-	-	2	4	5	4	3	5	-
2005 1/.....	30	-	-	1	-	4	4	5	5	2	5	4	-
2004 1/.....	23	-	-	-	-	3	-	1	3	3	5	8	-
2003 1/.....	16	-	-	-	-	-	2	4	1	4	3	2	-
2002.....	23	1	1	-	-	-	1	3	4	3	7	3	-
2001.....	14	-	-	1	-	1	2	2	-	3	3	2	-
2000.....	20	1	-	-	-	-	-	4	3	4	6	2	-
1999.....	12	-	-	-	-	-	-	-	1	3	5	3	-

Black, male

2007 1/.....	12	1	-	-	-	-	-	2	1	4	3	1	-
2006 1/.....	15	-	1	-	-	-	-	3	4	3	2	2	-
2005 1/.....	21	-	-	-	-	4	3	4	3	2	3	2	-
2004 1/.....	13	-	-	-	-	2	-	1	1	3	3	3	-
2003 1/.....	11	-	-	-	-	-	2	4	1	3	1	-	-
2002.....	12	-	1	-	-	-	1	2	2	2	3	1	-
2001.....	9	-	-	1	-	-	2	2	-	2	2	-	-
2000.....	17	1	-	-	-	-	-	4	3	2	5	2	-
1999.....	5	-	-	-	-	-	-	-	1	3	-	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Syphilis (A50-A53)

Asian or Pacific Islander 3/, female

2007 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2006 1/.....	1	-	-	-	1	-	-	-	-	-	-	-	-
2005 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	-	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Arthropod-borne viral encephalitis (A83-A84,A85.2)

All races, both sexes

2007.....	3	-	-	1	-	-	-	-	-	1	-	1	-
2006.....	5	-	-	1	1	-	-	2	-	1	-	-	-
2005.....	6	-	1	1	-	1	-	-	2	-	1	-	-
2004.....	5	-	-	-	-	-	1	-	1	3	-	-	-
2003.....	13	1	-	1	-	-	-	1	3	2	5	-	-
2002.....	9	-	-	-	1	1	-	2	1	1	2	1	-
2001.....	9	1	-	-	-	-	-	1	-	5	2	-	-
2000.....	2	-	-	-	-	-	1	-	-	-	1	-	-
1999.....	3	-	-	-	-	-	-	-	-	2	-	1	-

All races, male

2007.....	2	-	-	-	-	-	-	-	-	1	-	1	-
2006.....	4	-	-	1	1	-	-	2	-	-	-	-	-
2005.....	4	-	-	1	-	1	-	-	1	-	1	-	-
2004.....	5	-	-	-	-	-	1	-	1	3	-	-	-
2003.....	9	1	-	1	-	-	-	1	2	2	2	-	-
2002.....	4	-	-	-	-	-	-	1	1	1	1	-	-
2001.....	6	-	-	-	-	-	-	-	-	4	2	-	-
2000.....	2	-	-	-	-	-	1	-	-	-	1	-	-
1999.....	3	-	-	-	-	-	-	-	-	2	-	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Arthropod-borne viral encephalitis (A83-A84,A85.2)

All races, female

2007.....	1	-	-	1	-	-	-	-	-	-	-	-	-
2006.....	1	-	-	-	-	-	-	-	-	1	-	-	-
2005.....	2	-	1	-	-	-	-	-	1	-	-	-	-
2004.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	4	-	-	-	-	-	-	-	1	-	3	-	-
2002.....	5	-	-	-	1	1	-	1	-	-	1	1	-
2001.....	3	1	-	-	-	-	3	1	-	1	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-	-

White, both sexes

2007 1/.....	2	-	-	1	-	-	-	-	-	-	-	1	-
2006 1/.....	2	-	-	1	1	-	-	-	-	-	-	-	-
2005 1/.....	6	-	1	1	-	1	-	-	2	-	1	-	-
2004 1/.....	3	-	-	-	-	-	-	-	1	2	-	-	-
2003 1/.....	10	-	-	-	-	-	-	-	3	2	5	-	-
2002.....	6	-	-	-	1	1	-	1	-	-	2	1	-
2001.....	6	-	-	-	-	-	-	1	-	4	1	-	-
2000.....	2	-	-	-	-	-	1	-	-	-	1	-	-
1999.....	3	-	-	-	-	-	-	-	-	2	-	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Measles (B05)

All races, female

2007	-	-	-	-	-	-	-	-	-	-	-	-	-
2006	-	-	-	-	-	-	-	-	-	-	-	-	-
2005	-	-	-	-	-	-	-	-	-	-	-	-	-
2004	-	-	-	-	-	-	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	1	-	-	-	-	-	-	-	-	-	-	-
1999	2	1	-	-	-	-	-	-	1	-	-	-	-

White, both sexes

2007 1/	-	-	-	-	-	-	-	-	-	-	-	-	-
2006 1/	-	-	-	-	-	-	-	-	-	-	-	-	-
2005 1/	1	-	1	-	-	-	-	-	-	-	-	-	-
2004 1/	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 1/	1	-	1	-	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	1	-	-	-	-	-	-
2000	1	1	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	1	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Measles (B05)

White, male

2007 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2006 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2005 1/.....	1	-	1	-	-	-	-	-	-	-	-	-	-
2004 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 1/.....	1	-	1	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-	-

White, female

2007 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2006 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2005 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	1	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Viral hepatitis (B15-B19)													
All races, both sexes													
2007.....	7,407	1	1	-	14	72	489	2,815	2,413	871	570	161	-
2006.....	7,250	1	1	-	11	61	534	2,911	2,127	843	598	163	-
2005.....	5,529	-	-	1	14	61	509	2,314	1,377	648	482	123	-
2004.....	5,382	1	-	2	11	61	610	2,226	1,244	634	464	129	-
2003.....	5,431	1	-	-	14	64	652	2,259	1,085	698	543	115	-
2002.....	5,793	2	3	2	23	94	740	2,331	1,072	762	587	177	-
2001.....	5,585	1	-	3	20	74	797	2,060	1,014	836	613	166	1
2000.....	5,357	6	1	2	18	79	779	1,894	904	845	660	168	1
1999.....	4,853	2	3	3	14	88	722	1,506	786	881	651	197	-
All races, male													
2007.....	4,910	1	-	-	9	49	326	1,997	1,761	464	249	54	-
2006.....	4,745	-	-	-	6	39	370	2,102	1,502	432	243	51	-
2005.....	3,612	-	-	1	3	34	349	1,697	977	317	193	41	-
2004.....	3,513	-	-	1	7	33	399	1,645	860	334	189	45	-
2003.....	3,546	-	-	-	7	42	461	1,666	729	352	249	40	-
2002.....	3,770	1	3	1	13	63	519	1,755	702	373	270	70	-
2001.....	3,539	1	-	2	11	42	553	1,570	643	396	263	57	1
2000.....	3,381	2	1	1	11	46	540	1,425	577	396	311	70	1
1999.....	3,043	2	2	2	10	53	520	1,138	491	426	317	82	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

Viral hepatitis (B15-B19)

All races, female

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
2007.....	2,497	-	1	-	5	23	163	818	652	407	321	107	-
2006.....	2,505	1	1	-	5	22	164	809	625	411	355	112	-
2005.....	1,917	-	-	-	11	27	160	617	400	331	289	82	-
2004.....	1,869	1	-	1	4	28	211	581	384	300	275	84	-
2003.....	1,885	1	-	-	7	22	191	593	356	346	294	75	-
2002.....	2,023	1	-	1	10	31	221	576	370	389	317	107	-
2001.....	2,046	-	-	1	9	32	244	490	371	440	350	109	-
2000.....	1,976	4	-	1	7	33	239	469	327	449	349	98	-
1999.....	1,810	-	1	1	4	35	202	368	295	455	334	115	-

White, both sexes

2007 1/.....	5,959	1	1	-	9	55	387	2,345	1,887	682	457	135	-
2006 1/.....	5,759	1	1	-	7	47	421	2,415	1,626	638	468	135	-
2005 1/.....	4,395	-	-	-	9	40	421	1,895	1,059	471	397	103	-
2004 1/.....	4,246	1	-	1	5	40	500	1,789	934	491	375	110	-
2003 1/.....	4,295	-	-	-	9	49	531	1,833	795	533	447	98	-
2002.....	4,620	1	-	2	20	65	600	1,881	804	589	503	155	-
2001.....	4,451	1	-	3	13	53	646	1,632	778	655	527	143	-
2000.....	4,216	3	-	2	8	51	625	1,462	665	678	572	149	1
1999.....	3,835	1	3	2	9	68	571	1,164	592	694	561	170	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Viral hepatitis (B15-B19)

White, male

2007 1/.....	4,005	1	-	-	5	36	260	1,686	1,395	367	207	48	-
2006 1/.....	3,822	-	-	-	2	30	291	1,770	1,154	330	198	47	-
2005 1/.....	2,934	-	-	-	2	24	295	1,409	760	240	169	35	-
2004 1/.....	2,799	-	-	1	3	23	338	1,336	644	262	155	37	-
2003 1/.....	2,848	-	-	-	3	30	379	1,371	549	275	205	36	-
2002.....	3,049	1	-	1	11	46	429	1,432	527	302	239	61	-
2001.....	2,900	1	-	2	6	31	456	1,271	508	331	241	53	-
2000.....	2,686	1	-	1	6	30	439	1,123	432	317	273	63	1
1999.....	2,436	1	2	1	7	41	419	872	384	354	285	70	-

White, female

2007 1/.....	1,954	-	1	-	4	19	127	659	492	315	250	87	-
2006 1/.....	1,937	1	1	-	5	17	130	645	472	308	270	88	-
2005 1/.....	1,461	-	-	-	7	16	126	486	299	231	228	68	-
2004 1/.....	1,447	1	-	-	2	17	162	453	290	229	220	73	-
2003 1/.....	1,447	-	-	-	6	19	152	462	246	258	242	62	-
2002.....	1,571	-	-	1	9	19	171	449	277	287	264	94	-
2001.....	1,551	-	-	1	7	22	190	361	270	324	286	90	-
2000.....	1,530	2	-	1	2	21	186	339	233	361	299	86	-
1999.....	1,399	-	1	1	2	27	152	292	208	340	276	100	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Viral hepatitis (B15-B19)

Black, both sexes

2007 1/.....	1,049	-	-	-	4	14	65	382	409	107	56	12	-
2006 1/.....	1,094	-	-	-	2	10	73	396	396	129	71	17	-
2005 1/.....	872	-	-	1	3	18	65	355	257	107	53	13	-
2004 1/.....	863	-	-	1	4	14	82	369	251	80	50	12	-
2003 1/.....	854	-	-	-	4	11	98	357	222	92	59	11	-
2002.....	865	1	1	-	2	22	112	374	187	103	50	13	-
2001.....	831	-	-	-	6	15	121	355	170	93	54	16	1
2000.....	793	3	-	-	10	21	118	336	143	94	57	11	-
1999.....	731	1	-	1	4	17	121	279	120	113	55	20	-

Black, male

2007 1/.....	690	-	-	-	3	11	41	254	296	61	23	1	-
2006 1/.....	694	-	-	-	2	6	43	259	283	69	29	3	-
2005 1/.....	530	-	-	1	-	7	39	240	175	53	13	2	-
2004 1/.....	564	-	-	-	3	6	40	267	179	48	17	4	-
2003 1/.....	544	-	-	-	3	9	64	254	145	43	22	4	-
2002.....	541	-	1	-	1	11	68	266	124	50	17	3	-
2001.....	484	-	-	-	4	7	79	243	101	34	13	2	1
2000.....	501	1	-	-	5	11	78	241	88	48	25	4	-
1999.....	443	1	-	1	2	9	81	217	66	40	17	9	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Viral hepatitis (B15-B19)

Black, female

2007 1/.....	359	-	-	-	1	3	24	128	113	46	33	11	-
2006 1/.....	400	-	-	-	-	4	30	137	113	60	42	14	-
2005 1/.....	342	-	-	-	3	11	26	115	82	54	40	11	-
2004 1/.....	299	-	-	1	1	8	42	102	72	32	33	8	-
2003 1/.....	310	-	-	-	1	2	34	103	77	49	37	7	-
2002.....	324	1	-	-	1	11	44	108	63	53	33	10	-
2001.....	347	-	-	-	2	8	42	112	69	59	41	14	-
2000.....	292	2	-	-	5	10	40	95	55	46	32	7	-
1999.....	288	-	-	-	2	8	40	62	54	73	38	11	-

American Indian 2/, both sexes

2007 1/.....	105	-	-	-	-	1	19	33	39	11	2	-	-
2006 1/.....	102	-	-	-	-	1	18	41	28	7	6	1	-
2005 1/.....	56	-	-	-	-	-	10	23	14	6	2	1	-
2004 1/.....	62	-	-	-	-	-	13	31	11	5	1	1	-
2003 1/.....	62	-	-	-	1	3	9	27	16	5	1	-	-
2002.....	56	-	-	-	-	2	12	25	13	4	-	-	-
2001.....	55	-	-	-	1	4	13	22	5	8	2	-	-
2000.....	63	-	-	-	-	4	15	25	10	5	3	1	-
1999.....	37	-	-	-	1	1	8	14	4	3	5	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Viral hepatitis (B15-B19)

American Indian 2/, male

2007 1/.....	66	-	-	-	-	1	12	20	23	8	2	-	-
2006 1/.....	66	-	-	-	-	1	16	30	15	2	2	-	-
2005 1/.....	38	-	-	-	-	-	7	18	10	2	-	1	-
2004 1/.....	31	-	-	-	-	-	8	14	7	2	-	-	-
2003 1/.....	32	-	-	-	1	2	6	9	10	3	1	-	-
2002.....	36	-	-	-	-	2	8	17	7	2	-	-	-
2001.....	30	-	-	-	1	2	6	16	3	2	-	-	-
2000.....	33	-	-	-	-	2	8	16	5	1	1	-	-
1999.....	22	-	-	-	1	1	3	12	2	1	2	-	-

American Indian 2/, female

2007 1/.....	39	-	-	-	-	-	7	13	16	3	-	-	-
2006 1/.....	36	-	-	-	-	-	2	11	13	5	4	1	-
2005 1/.....	18	-	-	-	-	-	3	5	4	4	2	-	-
2004 1/.....	31	-	-	-	-	-	5	17	4	3	1	1	-
2003 1/.....	30	-	-	-	-	1	3	18	6	2	-	-	-
2002.....	20	-	-	-	-	-	4	8	6	2	-	-	-
2001.....	25	-	-	-	-	2	7	6	2	6	2	-	-
2000.....	30	-	-	-	-	2	7	9	5	4	2	1	-
1999.....	15	-	-	-	-	-	5	2	2	2	3	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Viral hepatitis (B15-B19)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	294	-	-	-	1	2	18	55	78	71	55	14	-
2006 1/.....	295	-	-	-	2	3	22	59	77	69	53	10	-
2005 1/.....	206	-	-	-	2	3	13	41	47	64	30	6	-
2004 1/.....	211	-	-	-	2	7	15	37	48	58	38	6	-
2003 1/.....	220	1	-	-	-	1	14	42	52	68	36	6	-
2002.....	252	-	2	-	1	5	16	51	68	66	34	9	-
2001.....	248	-	-	-	-	2	17	51	61	80	30	7	-
2000.....	285	-	1	-	-	3	21	71	86	68	28	7	-
1999.....	250	-	-	-	-	2	22	49	70	71	30	6	-

Asian or Pacific Islander 3/, male

2007 1/.....	149	-	-	-	1	1	13	37	47	28	17	5	-
2006 1/.....	163	-	-	-	2	2	20	43	50	31	14	1	-
2005 1/.....	110	-	-	-	1	3	8	30	32	22	11	3	-
2004 1/.....	119	-	-	-	1	4	13	28	30	22	17	4	-
2003 1/.....	122	-	-	-	-	1	12	32	25	31	21	-	-
2002.....	144	-	2	-	1	4	14	40	44	19	14	6	-
2001.....	125	-	-	-	-	2	12	40	31	29	9	2	-
2000.....	161	-	1	-	-	3	15	45	52	30	12	3	-
1999.....	142	-	-	-	-	2	17	37	39	31	13	3	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Viral hepatitis (B15-B19)

Asian or Pacific Islander 3/, female

2007 1/.....	145	-	-	-	-	1	5	18	31	43	38	9	-
2006 1/.....	132	-	-	-	-	1	2	16	27	38	39	9	-
2005 1/.....	96	-	-	-	1	-	5	11	15	42	19	3	-
2004 1/.....	92	-	-	-	1	3	2	9	18	36	21	2	-
2003 1/.....	98	1	-	-	-	-	2	10	27	37	15	6	-
2002.....	108	-	-	-	-	1	2	11	24	47	20	3	-
2001.....	123	-	-	-	-	-	5	11	30	51	21	5	-
2000.....	124	-	-	-	-	-	6	26	34	38	16	4	-
1999.....	108	-	-	-	-	-	5	12	31	40	17	3	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Human immunodeficiency virus (HIV) disease (B20-B24)

All races, both sexes

2007.....	11,295	5	4	10	160	1,091	3,572	4,156	1,721	448	109	16	3
2006.....	12,113	7	-	15	206	1,182	4,010	4,377	1,722	475	107	11	1
2005.....	12,543	2	2	13	159	1,318	4,363	4,516	1,612	438	104	15	1
2004.....	13,063	3	4	25	191	1,468	4,826	4,422	1,562	436	106	16	4
2003.....	13,658	5	2	36	178	1,588	5,340	4,442	1,517	438	95	14	3
2002.....	14,095	7	5	25	178	1,839	5,707	4,474	1,347	401	96	12	4
2001.....	14,175	10	13	38	225	2,101	5,867	4,120	1,317	384	88	10	2
2000.....	14,478	10	14	46	179	2,437	5,919	4,142	1,239	397	85	9	1
1999.....	14,802	14	24	66	198	2,729	6,232	3,907	1,149	398	70	14	1

All races, male

2007.....	8,097	5	3	3	88	653	2,506	3,028	1,344	369	86	9	3
2006.....	8,756	2	-	8	121	729	2,829	3,263	1,352	362	84	5	1
2005.....	9,189	-	1	7	89	809	3,146	3,417	1,292	348	73	6	1
2004.....	9,506	1	1	12	107	917	3,466	3,327	1,262	333	71	5	4
2003.....	10,119	2	2	19	95	1,038	3,884	3,438	1,215	336	75	12	3
2002.....	10,468	4	1	14	93	1,187	4,198	3,485	1,083	323	71	5	4
2001.....	10,590	7	8	23	105	1,421	4,360	3,241	1,051	292	73	7	2
2000.....	10,868	8	6	22	99	1,602	4,441	3,296	1,013	313	65	2	1
1999.....	11,234	9	16	36	95	1,914	4,706	3,134	948	317	50	8	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Human immunodeficiency virus (HIV) disease (B20-B24)													
All races, female													
2007.....	3,198	-	1	7	72	438	1,066	1,128	377	79	23	7	-
2006.....	3,357	5	-	7	85	453	1,181	1,114	370	113	23	6	-
2005.....	3,354	2	1	6	70	509	1,217	1,099	320	90	31	9	-
2004.....	3,557	2	3	13	84	551	1,360	1,095	300	103	35	11	-
2003.....	3,539	3	-	17	83	550	1,456	1,004	302	102	20	2	-
2002.....	3,627	3	4	11	85	652	1,509	989	264	78	25	7	-
2001.....	3,585	3	5	15	120	680	1,507	879	266	92	15	3	-
2000.....	3,610	2	8	24	80	835	1,478	846	226	84	20	7	-
1999.....	3,568	5	8	30	103	815	1,526	773	201	81	20	6	-
White, both sexes													
2007 1/.....	4,672	2	3	2	51	367	1,522	1,747	704	214	53	7	-
2006 1/.....	5,103	3	-	4	59	405	1,760	1,874	733	206	55	4	-
2005 1/.....	5,361	2	2	3	50	464	1,958	1,947	683	186	54	12	-
2004 1/.....	5,608	2	2	6	48	503	2,196	1,900	685	191	61	11	3
2003 1/.....	6,019	1	1	5	57	619	2,432	1,970	687	189	48	8	2
2002.....	6,089	4	-	6	52	688	2,570	1,921	643	151	42	9	3
2001.....	6,171	3	3	11	60	814	2,681	1,792	600	163	37	7	-
2000.....	6,498	1	1	21	53	983	2,785	1,856	588	166	40	3	1
1999.....	6,752	4	7	25	63	1,202	2,951	1,765	519	171	37	8	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Human immunodeficiency virus (HIV) disease (B20-B24)

White, male

2007 1/.....	3,797	2	2	1	32	263	1,249	1,422	584	192	46	4	-
2006 1/.....	4,193	1	-	2	36	305	1,427	1,573	631	172	44	2	-
2005 1/.....	4,379	-	1	-	32	337	1,608	1,613	583	159	41	5	-
2004 1/.....	4,593	1	1	2	33	392	1,777	1,578	600	159	44	3	3
2003 1/.....	4,964	1	1	2	35	479	1,986	1,677	580	153	42	6	2
2002.....	5,028	3	-	4	31	532	2,123	1,619	546	134	29	4	3
2001.....	5,135	2	-	7	35	628	2,238	1,540	507	140	33	5	-
2000.....	5,360	1	-	9	38	749	2,300	1,590	499	140	33	-	1
1999.....	5,613	3	5	13	40	956	2,449	1,510	461	145	29	2	-

White, female

2007 1/.....	875	-	1	1	19	104	273	325	120	22	7	3	-
2006 1/.....	910	2	-	2	23	100	333	301	102	34	11	2	-
2005 1/.....	982	2	1	3	18	127	350	334	100	27	13	7	-
2004 1/.....	1,015	1	1	4	15	111	419	322	85	32	17	8	-
2003 1/.....	1,055	-	-	3	22	140	446	293	107	36	6	2	-
2002.....	1,061	1	-	2	21	156	447	302	97	17	13	5	-
2001.....	1,036	1	3	4	25	186	443	252	93	23	4	2	-
2000.....	1,138	-	1	12	15	234	485	266	89	26	7	3	-
1999.....	1,139	1	2	12	23	246	502	255	58	26	8	6	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Human immunodeficiency virus (HIV) disease (B20-B24)													
Black, both sexes													
2007 1/.....	6,470	1	1	8	109	705	2,002	2,351	995	231	55	9	3
2006 1/.....	6,854	4	-	11	144	770	2,189	2,449	976	254	49	7	1
2005 1/.....	7,022	-	-	9	109	836	2,337	2,520	911	246	50	3	1
2004 1/.....	7,271	1	2	18	142	944	2,554	2,464	860	235	45	5	1
2003 1/.....	7,479	3	1	31	119	948	2,832	2,428	817	247	46	6	1
2002.....	7,835	3	5	19	124	1,123	3,063	2,509	688	243	54	3	1
2001.....	7,844	7	10	26	163	1,258	3,119	2,290	703	214	50	2	2
2000.....	7,848	9	12	23	125	1,432	3,080	2,257	636	224	44	6	-
1999.....	7,893	10	17	41	133	1,497	3,206	2,103	625	222	33	5	1
Black, male													
2007 1/.....	4,186	1	1	2	56	376	1,223	1,563	742	175	39	5	3
2006 1/.....	4,443	1	-	6	84	418	1,358	1,646	709	179	38	3	1
2005 1/.....	4,684	-	-	6	57	458	1,489	1,760	696	184	32	1	1
2004 1/.....	4,769	-	-	9	73	508	1,624	1,706	650	169	27	2	1
2003 1/.....	5,033	1	1	17	59	543	1,840	1,725	625	183	32	6	1
2002.....	5,301	1	1	10	60	634	2,017	1,827	523	184	42	1	1
2001.....	5,328	5	8	15	69	767	2,071	1,668	533	150	39	1	2
2000.....	5,400	7	6	12	60	835	2,096	1,682	501	168	31	2	-
1999.....	5,493	6	11	23	54	935	2,194	1,592	483	168	21	5	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Human immunodeficiency virus (HIV) disease (B20-B24)													
Black, female													
2007 1/.....	2,284	-	-	6	53	329	779	788	253	56	16	4	-
2006 1/.....	2,411	3	-	5	60	352	831	803	267	75	11	4	-
2005 1/.....	2,338	-	-	3	52	378	848	760	215	62	18	2	-
2004 1/.....	2,502	1	2	9	69	436	930	758	210	66	18	3	-
2003 1/.....	2,446	2	-	14	60	405	992	703	192	64	14	-	-
2002.....	2,534	2	4	9	64	489	1,046	682	165	59	12	2	-
2001.....	2,516	2	2	11	94	491	1,048	622	170	64	11	1	-
2000.....	2,448	2	6	11	65	597	984	575	135	56	13	4	-
1999.....	2,400	4	6	18	79	562	1,012	511	142	54	12	-	-
American Indian 2/, both sexes													
2007 1/.....	78	-	-	-	-	5	31	32	7	2	1	-	-
2006 1/.....	69	-	-	-	1	5	34	23	3	3	-	-	-
2005 1/.....	80	-	-	1	-	10	35	26	6	2	-	-	-
2004 1/.....	84	-	-	-	-	9	39	26	9	1	-	-	-
2003 1/.....	70	-	-	-	-	8	37	18	5	1	1	-	-
2002.....	64	-	-	-	1	13	36	11	3	-	-	-	-
2001.....	74	-	-	1	-	11	35	22	2	2	1	-	-
2000.....	57	-	1	1	-	8	24	18	1	3	1	-	-
1999.....	68	-	-	-	-	17	34	13	2	2	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Human immunodeficiency virus (HIV) disease (B20-B24)

American Indian 2/, male

2007 1/.....	53	-	-	-	-	4	21	22	4	1	1	-	-
2006 1/.....	48	-	-	-	-	5	21	18	3	1	-	-	-
2005 1/.....	59	-	-	1	-	7	22	24	4	1	-	-	-
2004 1/.....	63	-	-	-	-	8	33	17	5	-	-	-	-
2003 1/.....	49	-	-	-	-	7	23	14	4	-	1	-	-
2002.....	50	-	-	-	1	10	29	8	2	-	-	-	-
2001.....	57	-	-	1	-	10	24	19	2	-	1	-	-
2000.....	43	-	-	-	-	5	20	14	-	3	1	-	-
1999.....	54	-	-	-	-	13	27	11	1	2	-	-	-

American Indian 2/, female

2007 1/.....	25	-	-	-	-	1	10	10	3	1	-	-	-
2006 1/.....	21	-	-	-	1	-	13	5	-	2	-	-	-
2005 1/.....	21	-	-	-	-	3	13	2	2	1	-	-	-
2004 1/.....	21	-	-	-	-	1	6	9	4	1	-	-	-
2003 1/.....	21	-	-	-	-	1	14	4	1	1	-	-	-
2002.....	14	-	-	-	-	3	7	3	1	-	-	-	-
2001.....	17	-	-	-	-	1	11	3	-	2	-	-	-
2000.....	14	-	1	1	-	3	4	4	1	-	-	-	-
1999.....	14	-	-	-	-	4	7	2	1	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Human immunodeficiency virus (HIV) disease (B20-B24)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	75	2	-	-	-	14	17	26	15	1	-	-	-
2006 1/.....	87	-	-	-	2	2	27	31	10	12	3	-	-
2005 1/.....	80	-	-	-	-	8	33	23	12	4	-	-	-
2004 1/.....	100	-	-	1	1	12	37	32	8	9	-	-	-
2003 1/.....	90	1	-	-	2	13	39	26	8	1	-	-	-
2002.....	107	-	-	-	1	15	38	33	13	7	-	-	-
2001.....	86	-	-	-	2	18	32	16	12	5	-	1	-
2000.....	75	-	-	1	1	14	30	11	14	4	-	-	-
1999.....	89	-	-	-	2	13	41	26	3	3	-	1	-

Asian or Pacific Islander 3/, male

2007 1/.....	61	2	-	-	-	10	13	21	14	1	-	-	-
2006 1/.....	72	-	-	-	1	1	23	26	9	10	2	-	-
2005 1/.....	67	-	-	-	-	7	27	20	9	4	-	-	-
2004 1/.....	81	-	-	1	1	9	32	26	7	5	-	-	-
2003 1/.....	73	-	-	-	1	9	35	22	6	-	-	-	-
2002.....	89	-	-	-	1	11	29	31	12	5	-	-	-
2001.....	70	-	-	-	1	16	27	14	9	2	-	1	-
2000.....	65	-	-	1	1	13	25	10	13	2	-	-	-
1999.....	74	-	-	-	1	10	36	21	3	2	-	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Human immunodeficiency virus (HIV) disease (B20-B24)

Asian or Pacific Islander 3/, female

2007 1/.....	14	-	-	-	-	4	4	5	1	-	-	-	-
2006 1/.....	15	-	-	-	1	1	4	5	1	2	1	-	-
2005 1/.....	13	-	-	-	-	1	6	3	3	-	-	-	-
2004 1/.....	19	-	-	-	-	3	5	6	1	4	-	-	-
2003 1/.....	17	1	-	-	1	4	4	4	2	1	-	-	-
2002.....	18	-	-	-	-	4	9	2	1	2	-	-	-
2001.....	16	-	-	-	1	2	5	2	3	3	-	-	-
2000.....	10	-	-	-	-	1	5	1	1	2	-	-	-
1999.....	15	-	-	-	1	3	5	5	-	1	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malaria (B50-B54)

All races, both sexes

2007.....	5	-	-	-	-	-	2	-	1	1	-	1	-
2006.....	9	-	1	-	-	-	1	2	3	-	2	-	-
2005.....	6	-	-	-	2	1	1	1	-	1	-	-	-
2004.....	8	-	-	-	1	-	4	2	-	1	-	-	-
2003.....	4	-	-	-	-	-	1	-	2	1	-	-	-
2002.....	12	-	-	-	1	1	1	5	2	2	-	-	-
2001.....	9	-	-	1	-	-	2	2	2	1	1	-	-
2000.....	3	-	-	-	1	-	-	1	-	-	1	-	-
1999.....	7	1	-	-	-	2	1	-	1	2	-	-	-

All races, male

2007.....	3	-	-	-	-	-	1	-	-	1	-	1	-
2006.....	6	-	-	-	-	-	1	2	1	-	2	-	-
2005.....	4	-	-	-	-	1	1	1	-	1	-	-	-
2004.....	8	-	-	-	1	-	4	2	-	1	-	-	-
2003.....	4	-	-	-	-	-	1	-	2	1	-	-	-
2002.....	8	-	-	-	-	-	1	4	1	2	-	-	-
2001.....	5	-	-	1	-	-	-	1	1	1	1	-	-
2000.....	2	-	-	-	-	-	-	1	-	-	1	-	-
1999.....	4	1	-	-	-	1	-	-	1	1	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Malaria (B50-B54)

All races, female

2007.....	2	-	-	-	-	-	1	-	1	-	-	-	-
2006.....	3	-	1	-	-	-	-	-	2	-	-	-	-
2005.....	2	-	-	-	2	-	-	-	-	-	-	-	-
2004.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	4	-	-	-	1	1	1	1	-	-	-	-	-
2001.....	4	-	-	-	-	-	2	1	1	-	-	-	-
2000.....	1	-	-	-	1	-	-	-	-	-	-	-	-
1999.....	3	-	-	-	-	1	1	-	-	1	-	-	-

White, both sexes

2007 1/.....	3	-	-	-	-	-	1	-	1	-	-	1	-
2006 1/.....	5	-	-	-	-	-	1	-	2	-	2	-	-
2005 1/.....	2	-	-	-	1	-	1	-	-	-	-	-	-
2004 1/.....	3	-	-	-	1	-	2	-	-	-	-	-	-
2003 1/.....	4	-	-	-	-	-	1	-	2	1	-	-	-
2002.....	7	-	-	-	-	-	1	3	1	2	-	-	-
2001.....	5	-	-	-	-	-	2	1	-	1	1	-	-
2000.....	1	-	-	-	1	-	-	-	-	-	-	-	-
1999.....	4	-	-	-	-	1	1	-	1	1	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Malaria (B50-B54)

White, male

2007 1/.....	2	-	-	-	-	-	1	-	-	-	-	1	-
2006 1/.....	3	-	-	-	-	-	1	-	-	-	2	-	-
2005 1/.....	1	-	-	-	-	-	1	-	-	-	-	-	-
2004 1/.....	3	-	-	-	1	-	2	-	-	-	-	-	-
2003 1/.....	4	-	-	-	-	-	1	-	2	1	-	-	-
2002.....	7	-	-	-	-	-	1	3	1	2	-	-	-
2001.....	2	-	-	-	-	-	-	-	-	1	1	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	3	-	-	-	-	1	-	-	1	1	-	-	-

White, female

2007 1/.....	1	-	-	-	-	-	-	-	1	-	-	-	-
2006 1/.....	2	-	-	-	-	-	-	-	2	-	-	-	-
2005 1/.....	1	-	-	-	1	-	-	-	-	-	-	-	-
2004 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	3	-	-	-	-	-	2	1	-	-	-	-	-
2000.....	1	-	-	-	1	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	1	-	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)

All races, both sexes

2007.....	5,825	154	63	51	97	136	229	511	919	1,098	1,463	1,104	-
2006.....	5,897	167	64	67	87	129	269	562	864	1,106	1,502	1,079	1
2005.....	7,727	169	63	64	86	130	403	1,364	1,273	1,309	1,785	1,081	-
2004.....	7,540	167	65	47	95	144	434	1,236	1,237	1,309	1,714	1,092	-
2003.....	7,427	174	85	51	88	140	447	1,185	1,136	1,344	1,715	1,062	-
2002.....	6,707	224	61	84	97	169	453	916	889	1,223	1,594	997	-
2001.....	5,715	191	64	64	86	147	369	663	771	1,143	1,330	887	-
2000.....	5,532	167	71	67	96	141	350	585	707	1,074	1,375	899	-
1999.....	7,279	203	81	79	90	242	545	836	898	1,443	1,739	1,123	-

All races, male

2007.....	2,982	84	30	36	56	80	130	313	502	584	718	449	-
2006.....	3,028	95	41	35	44	68	169	343	492	581	736	423	1
2005.....	4,216	100	40	29	43	80	270	933	768	711	864	378	-
2004.....	4,097	100	35	25	59	81	257	834	760	710	790	446	-
2003.....	3,964	86	43	25	52	83	284	792	689	711	816	383	-
2002.....	3,622	125	39	40	52	96	291	626	532	663	785	373	-
2001.....	2,931	101	37	29	51	83	227	426	449	565	656	307	-
2000.....	2,861	94	41	35	54	75	220	358	383	597	665	339	-
1999.....	3,655	119	51	44	49	131	333	478	482	760	827	381	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)

All races, female

2007.....	2,843	70	33	15	41	56	99	198	417	514	745	655	-
2006.....	2,869	72	23	32	43	61	100	219	372	525	766	656	-
2005.....	3,511	69	23	35	43	50	133	431	505	598	921	703	-
2004.....	3,443	67	30	22	36	63	177	402	477	599	924	646	-
2003.....	3,463	88	42	26	36	57	163	393	447	633	899	679	-
2002.....	3,085	99	22	44	45	73	162	290	357	560	809	624	-
2001.....	2,784	90	27	35	35	64	142	237	322	578	674	580	-
2000.....	2,671	73	30	32	42	66	130	227	324	477	710	560	-
1999.....	3,624	84	30	35	41	111	212	358	416	683	912	742	-

White, both sexes

2007 1/.....	4,850	107	48	40	68	92	145	374	745	928	1,287	1,016	-
2006 1/.....	4,874	89	47	54	66	79	178	413	692	944	1,333	978	1
2005 1/.....	6,427	117	45	51	55	85	299	1,101	1,039	1,108	1,556	971	-
2004 1/.....	6,270	116	45	32	70	90	324	1,021	1,002	1,084	1,501	985	-
2003 1/.....	6,182	110	53	39	63	97	334	939	909	1,161	1,497	980	-
2002.....	5,528	155	45	66	64	117	307	708	692	1,048	1,412	914	-
2001.....	4,688	120	44	49	58	96	258	508	609	965	1,180	801	-
2000.....	4,550	101	50	52	71	84	242	444	560	913	1,226	807	-
1999.....	5,794	135	56	56	64	144	353	580	691	1,208	1,510	997	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)

White, male

2007 1/.....	2,471	57	20	28	38	51	89	236	414	487	639	412	-
2006 1/.....	2,488	54	30	27	39	43	111	253	383	504	652	391	1
2005 1/.....	3,519	68	27	20	26	53	204	776	628	613	761	343	-
2004 1/.....	3,414	69	24	18	44	50	194	695	615	598	697	410	-
2003 1/.....	3,289	51	26	19	40	57	220	640	553	608	720	355	-
2002.....	2,967	85	28	35	33	63	207	492	410	577	695	342	-
2001.....	2,405	62	25	22	33	61	160	332	360	480	590	280	-
2000.....	2,334	59	27	25	41	48	153	274	311	511	587	298	-
1999.....	2,910	85	37	31	34	78	239	345	377	644	710	330	-

White, female

2007 1/.....	2,379	50	28	12	30	41	56	138	331	441	648	604	-
2006 1/.....	2,386	35	17	27	27	36	67	160	309	440	681	587	-
2005 1/.....	2,908	49	18	31	29	32	95	325	411	495	795	628	-
2004 1/.....	2,856	47	21	14	26	40	130	326	387	486	804	575	-
2003 1/.....	2,893	59	27	20	23	40	114	299	356	553	777	625	-
2002.....	2,561	70	17	31	31	54	100	216	282	471	717	572	-
2001.....	2,283	58	19	27	25	35	98	176	249	485	590	521	-
2000.....	2,216	42	23	27	30	36	89	170	249	402	639	509	-
1999.....	2,884	50	19	25	30	66	114	235	314	564	800	667	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)

Black, both sexes

2007 1/.....	767	41	9	7	24	39	67	114	143	124	130	69	-
2006 1/.....	836	67	13	11	18	39	75	133	139	130	126	85	-
2005 1/.....	976	43	13	10	22	40	84	209	182	137	151	85	-
2004 1/.....	967	44	12	10	20	42	93	166	179	151	164	86	-
2003 1/.....	960	53	25	9	18	33	101	208	168	119	159	67	-
2002.....	943	59	14	10	27	47	125	181	152	122	133	73	-
2001.....	825	62	16	13	23	45	94	131	130	139	105	67	-
2000.....	826	61	14	10	22	50	95	127	119	129	122	77	-
1999.....	1,260	60	23	17	20	89	172	221	171	194	186	107	-

Black, male

2007 1/.....	399	24	6	5	15	27	30	66	70	75	53	28	-
2006 1/.....	428	35	9	6	4	20	47	82	85	58	59	23	-
2005 1/.....	524	27	8	6	11	22	55	120	110	69	71	25	-
2004 1/.....	525	25	8	5	12	23	49	111	115	76	71	30	-
2003 1/.....	523	30	13	5	9	20	57	127	98	76	66	22	-
2002.....	526	33	10	3	15	29	71	118	99	62	59	27	-
2001.....	427	36	10	6	15	19	59	81	71	69	42	19	-
2000.....	432	31	10	5	10	22	58	76	62	70	60	28	-
1999.....	624	31	14	10	12	48	83	115	83	95	94	39	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)

Black, female

2007 1/.....	368	17	3	2	9	12	37	48	73	49	77	41	-
2006 1/.....	408	32	4	5	14	19	28	51	54	72	67	62	-
2005 1/.....	452	16	5	4	11	18	29	89	72	68	80	60	-
2004 1/.....	442	19	4	5	8	19	44	55	64	75	93	56	-
2003 1/.....	437	23	12	4	9	13	44	81	70	43	93	45	-
2002.....	417	26	4	7	12	18	54	63	53	60	74	46	-
2001.....	398	26	6	7	8	26	35	50	59	70	63	48	-
2000.....	394	30	4	5	12	28	37	51	57	59	62	49	-
1999.....	636	29	9	7	8	41	89	106	88	99	92	68	-

American Indian 2/, both sexes

2007 1/.....	51	3	2	1	1	1	6	9	7	9	10	2	-
2006 1/.....	47	5	-	-	-	3	8	5	10	8	5	3	-
2005 1/.....	80	4	1	2	3	1	10	24	16	7	10	2	-
2004 1/.....	78	6	4	1	-	4	6	22	14	10	8	3	-
2003 1/.....	72	4	4	2	2	2	5	19	14	8	8	4	-
2002.....	54	4	1	1	-	-	9	8	12	11	8	-	-
2001.....	51	5	3	2	2	-	8	5	7	8	7	4	-
2000.....	34	2	2	-	-	1	5	5	12	3	3	1	-
1999.....	53	4	1	1	1	-	7	7	9	13	7	3	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)

American Indian 2/, male

2007 1/.....	29	2	1	1	-	1	5	3	5	5	5	1	-
2006 1/.....	29	4	-	-	-	2	6	4	8	2	2	1	-
2005 1/.....	47	3	1	2	2	1	6	15	9	4	3	1	-
2004 1/.....	42	5	2	-	-	2	4	11	7	5	5	1	-
2003 1/.....	33	1	2	1	-	2	2	10	8	3	3	1	-
2002.....	28	3	1	1	-	-	5	6	6	3	3	-	-
2001.....	21	2	2	1	2	-	3	2	2	2	3	2	-
2000.....	13	1	-	-	-	-	2	5	3	-	1	1	-
1999.....	20	1	-	-	-	-	3	1	3	7	3	2	-

American Indian 2/, female

2007 1/.....	22	1	1	-	1	-	1	6	2	4	5	1	-
2006 1/.....	18	1	-	-	-	1	2	1	2	6	3	2	-
2005 1/.....	33	1	-	-	1	-	4	9	7	3	7	1	-
2004 1/.....	36	1	2	1	-	2	2	11	7	5	3	2	-
2003 1/.....	39	3	2	1	2	-	3	9	6	5	5	3	-
2002.....	26	1	-	-	-	-	4	2	6	8	5	-	-
2001.....	30	3	1	1	-	-	5	3	5	6	4	2	-
2000.....	21	1	2	-	-	1	3	-	9	3	2	-	-
1999.....	33	3	1	1	1	-	4	6	6	6	4	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	157	3	4	3	4	4	11	14	24	37	36	17	-
2006 1/.....	140	6	4	2	3	8	8	11	23	24	38	13	-
2005 1/.....	244	5	4	1	6	4	10	30	36	57	68	23	-
2004 1/.....	225	1	4	4	5	8	11	27	42	64	41	18	-
2003 1/.....	213	7	3	1	5	8	7	19	45	56	51	11	-
2002.....	182	6	1	7	6	5	12	19	33	42	41	10	-
2001.....	151	4	1	-	3	6	9	19	25	31	38	15	-
2000.....	122	3	5	5	3	6	8	9	16	29	24	14	-
1999.....	172	4	1	5	5	9	13	28	27	28	36	16	-

Asian or Pacific Islander 3/, male

2007 1/.....	83	1	3	2	3	1	6	8	13	17	21	8	-
2006 1/.....	83	2	2	2	1	3	5	4	16	17	23	8	-
2005 1/.....	126	2	4	1	4	4	5	22	21	25	29	9	-
2004 1/.....	116	1	1	2	3	6	10	17	23	31	17	5	-
2003 1/.....	119	4	2	-	3	4	5	15	30	24	27	5	-
2002.....	101	4	-	1	4	4	8	10	17	21	28	4	-
2001.....	78	1	-	-	1	3	5	11	16	14	21	6	-
2000.....	82	3	4	5	3	5	7	3	7	16	17	12	-
1999.....	101	2	-	3	3	5	8	17	19	14	20	10	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)

Asian or Pacific Islander 3/, female

2007 1/.....	74	2	1	1	1	3	5	6	11	20	15	9	-
2006 1/.....	57	4	2	-	2	5	3	7	7	7	15	5	-
2005 1/.....	118	3	-	-	2	-	5	8	15	32	39	14	-
2004 1/.....	109	-	3	2	2	2	1	10	19	33	24	13	-
2003 1/.....	94	3	1	1	2	4	2	4	15	32	24	6	-
2002.....	81	2	1	6	2	1	4	9	16	21	13	6	-
2001.....	73	3	1	-	2	3	4	8	9	17	17	9	-
2000.....	40	-	1	-	-	1	1	6	9	13	7	2	-
1999.....	71	2	1	2	2	4	5	11	8	14	16	6	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms (C00-C97)

All races, both sexes

2007.....	562,875	72	364	959	1,653	3,463	13,288	50,167	103,171	138,466	163,608	87,656	8
2006.....	559,888	76	377	907	1,644	3,656	13,917	50,334	101,454	137,554	164,889	85,072	8
2005.....	559,312	75	377	1,000	1,717	3,601	14,566	50,405	99,240	138,446	166,421	83,455	9
2004.....	553,888	74	399	1,019	1,709	3,633	14,723	49,520	96,956	139,417	166,085	80,345	8
2003.....	556,902	75	392	1,076	1,651	3,741	15,509	49,843	95,692	141,248	167,617	80,046	12
2002.....	557,271	74	402	1,072	1,730	3,872	16,085	49,637	93,391	144,757	167,062	79,182	7
2001.....	553,768	66	420	1,008	1,704	3,994	16,569	49,562	90,223	147,018	165,445	77,751	8
2000.....	553,091	92	420	1,014	1,713	3,916	16,520	48,034	89,005	150,131	165,099	77,136	11
1999.....	549,838	68	418	1,012	1,724	4,005	16,732	46,681	89,067	152,338	162,770	75,014	9

All races, male

2007.....	292,857	39	191	499	987	1,692	5,701	25,370	56,559	75,924	85,927	39,964	4
2006.....	290,069	39	207	507	1,007	1,770	5,979	25,341	55,348	75,468	86,434	37,964	5
2005.....	290,422	45	217	557	1,035	1,805	6,338	25,403	54,039	76,685	87,096	37,197	5
2004.....	286,830	37	212	572	1,027	1,750	6,408	25,432	52,738	76,487	86,734	35,427	6
2003.....	287,990	34	227	591	979	1,804	6,825	25,542	51,927	77,793	87,382	34,878	8
2002.....	288,768	42	216	600	1,016	1,868	7,056	25,192	51,115	80,085	86,952	34,621	5
2001.....	287,075	31	229	533	1,024	1,860	7,297	25,071	49,244	81,699	86,126	33,954	7
2000.....	286,082	50	233	569	1,022	1,854	7,344	24,215	48,427	83,191	85,907	33,260	10
1999.....	285,832	36	227	555	1,023	1,959	7,471	23,634	48,833	84,544	85,060	32,484	6

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms (C00-C97)

All races, female

2007.....	270,018	33	173	460	666	1,771	7,587	24,797	46,612	62,542	77,681	47,692	4
2006.....	269,819	37	170	400	637	1,886	7,938	24,993	46,106	62,086	78,455	47,108	3
2005.....	268,890	30	160	443	682	1,796	8,228	25,002	45,201	61,761	79,325	46,258	4
2004.....	267,058	37	187	447	682	1,883	8,315	24,088	44,218	62,930	79,351	44,918	2
2003.....	268,912	41	165	485	672	1,937	8,684	24,301	43,765	63,455	80,235	45,168	4
2002.....	268,503	32	186	472	714	2,004	9,029	24,445	42,276	64,672	80,110	44,561	2
2001.....	266,693	35	191	475	680	2,134	9,272	24,491	40,979	65,319	79,319	43,797	1
2000.....	267,009	42	187	445	691	2,062	9,176	23,819	40,578	66,940	79,192	43,876	1
1999.....	264,006	32	191	457	701	2,046	9,261	23,047	40,234	67,794	77,710	42,530	3

White, both sexes

2007 1/.....	483,939	60	287	776	1,296	2,660	10,289	39,557	85,373	118,826	145,585	79,226	4
2006 1/.....	482,575	64	286	720	1,294	2,799	10,755	39,795	84,421	118,280	147,150	77,005	6
2005 1/.....	482,132	64	293	787	1,365	2,694	11,352	39,717	82,439	119,280	148,732	75,404	5
2004 1/.....	478,134	63	311	793	1,323	2,783	11,386	39,028	81,073	120,167	148,497	72,704	6
2003 1/.....	481,556	64	317	853	1,324	2,868	12,066	39,259	80,298	122,223	149,848	72,425	11
2002.....	482,481	56	306	846	1,399	2,997	12,582	39,516	78,140	125,458	149,525	71,652	4
2001.....	479,651	54	353	780	1,359	3,094	12,959	39,686	75,603	127,404	148,005	70,350	4
2000.....	480,011	75	330	818	1,333	3,072	12,764	38,478	74,759	130,570	148,126	69,679	7
1999.....	477,238	53	353	780	1,327	3,147	13,060	37,489	74,568	132,703	146,009	67,742	7

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms (C00-C97)

White, male

2007 1/.....	252,049	31	146	404	781	1,329	4,522	20,133	46,817	65,198	76,691	35,995	2
2006 1/.....	250,322	33	162	401	805	1,396	4,726	20,163	46,069	64,913	77,316	34,334	4
2005 1/.....	250,478	38	168	429	826	1,378	5,072	20,194	44,816	66,080	77,915	33,559	3
2004 1/.....	247,450	31	161	453	801	1,401	5,068	20,108	44,020	65,946	77,484	31,973	4
2003 1/.....	249,053	29	182	472	790	1,432	5,422	20,158	43,526	67,430	78,084	31,520	8
2002.....	249,867	35	168	459	830	1,472	5,604	20,157	42,632	69,588	77,649	31,270	3
2001.....	248,146	24	189	398	824	1,493	5,783	20,069	41,155	70,875	76,782	30,550	4
2000.....	247,403	42	190	456	798	1,500	5,739	19,353	40,457	72,284	76,714	29,863	7
1999.....	247,396	31	194	434	784	1,561	5,934	18,839	40,768	73,679	76,066	29,102	4

White, female

2007 1/.....	231,890	29	141	372	515	1,331	5,767	19,424	38,556	53,628	68,894	43,231	2
2006 1/.....	232,253	31	124	319	489	1,403	6,029	19,632	38,352	53,367	69,834	42,671	2
2005 1/.....	231,654	26	125	358	539	1,316	6,280	19,523	37,623	53,200	70,817	41,845	2
2004 1/.....	230,684	32	150	340	522	1,382	6,318	18,920	37,053	54,221	71,013	40,731	2
2003 1/.....	232,503	35	135	381	534	1,436	6,644	19,101	36,772	54,793	71,764	40,905	3
2002.....	232,614	21	138	387	569	1,525	6,978	19,359	35,508	55,870	71,876	40,382	1
2001.....	231,505	30	164	382	535	1,601	7,176	19,617	34,448	56,529	71,223	39,800	-
2000.....	232,608	33	140	362	535	1,572	7,025	19,125	34,302	58,286	71,412	39,816	-
1999.....	229,842	22	159	346	543	1,586	7,126	18,650	33,800	59,024	69,943	38,640	3

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms (C00-C97)

Black, both sexes

2007 1/.....	64,049	9	55	143	261	618	2,365	8,818	14,762	16,055	14,277	6,682	4
2006 1/.....	63,082	9	65	145	264	639	2,515	8,800	14,171	15,767	14,237	6,469	1
2005 1/.....	63,165	10	64	172	264	688	2,560	8,901	13,960	15,719	14,295	6,528	4
2004 1/.....	62,499	7	65	170	311	667	2,682	8,753	13,296	15,871	14,384	6,291	2
2003 1/.....	62,660	7	57	173	247	667	2,817	8,903	13,015	15,816	14,606	6,352	-
2002.....	62,617	12	74	172	250	669	2,903	8,450	12,825	16,245	14,664	6,350	3
2001.....	62,170	6	48	176	262	697	2,981	8,245	12,323	16,547	14,590	6,291	4
2000.....	61,945	11	70	150	301	649	3,108	8,147	12,119	16,636	14,361	6,390	3
1999.....	61,951	11	49	183	318	671	3,080	7,830	12,433	16,764	14,312	6,299	1

Black, male

2007 1/.....	33,069	6	33	72	145	263	907	4,365	8,181	8,747	7,245	3,103	2
2006 1/.....	32,556	3	35	83	153	272	980	4,369	7,805	8,695	7,314	2,846	1
2005 1/.....	32,726	6	37	103	155	321	973	4,365	7,767	8,724	7,386	2,887	2
2004 1/.....	32,443	2	37	84	176	265	1,035	4,489	7,359	8,730	7,497	2,767	2
2003 1/.....	32,442	4	31	96	143	269	1,127	4,590	7,190	8,680	7,602	2,710	-
2002.....	32,627	4	34	106	140	290	1,172	4,240	7,183	8,931	7,749	2,776	2
2001.....	32,679	3	30	102	153	270	1,217	4,235	6,833	9,222	7,782	2,829	3
2000.....	32,817	4	34	87	179	262	1,309	4,192	6,829	9,321	7,767	2,831	2
1999.....	32,850	3	26	96	184	297	1,267	4,144	6,993	9,314	7,674	2,851	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms (C00-C97)

Black, female

2007 1/.....	30,980	3	22	71	116	355	1,458	4,453	6,581	7,308	7,032	3,579	2
2006 1/.....	30,526	6	30	62	111	367	1,535	4,431	6,366	7,072	6,923	3,623	-
2005 1/.....	30,439	4	27	69	109	367	1,587	4,536	6,193	6,995	6,909	3,641	2
2004 1/.....	30,056	5	28	86	135	402	1,647	4,264	5,937	7,141	6,887	3,524	-
2003 1/.....	30,218	3	26	77	104	398	1,690	4,313	5,825	7,136	7,004	3,642	-
2002.....	29,990	8	40	66	110	379	1,731	4,210	5,642	7,314	6,915	3,574	1
2001.....	29,491	3	18	74	109	427	1,764	4,010	5,490	7,325	6,808	3,462	1
2000.....	29,128	7	36	63	122	387	1,799	3,955	5,290	7,315	6,594	3,559	1
1999.....	29,101	8	23	87	134	374	1,813	3,686	5,440	7,450	6,638	3,448	-

American Indian 2/, both sexes

2007 1/.....	2,561	1	1	6	22	25	83	330	623	685	561	224	-
2006 1/.....	2,447	-	7	5	16	32	93	308	543	680	560	203	-
2005 1/.....	2,465	-	4	4	15	29	116	340	580	676	518	183	-
2004 1/.....	2,392	1	3	8	16	25	106	313	564	670	511	175	-
2003 1/.....	2,154	-	7	8	19	24	101	303	465	557	494	176	-
2002.....	2,175	1	3	12	20	35	100	293	485	589	460	177	-
2001.....	2,155	3	4	5	22	34	114	282	470	596	453	172	-
2000.....	1,914	2	3	8	13	26	105	220	417	539	412	169	-
1999.....	1,836	1	4	6	13	26	85	217	406	535	405	138	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms (C00-C97)													
American Indian 2/, male													
2007 1/.....	1,345	1	-	4	16	18	37	163	350	360	285	111	-
2006 1/.....	1,217	-	2	3	10	13	35	151	280	352	280	91	-
2005 1/.....	1,297	-	3	4	10	16	62	161	321	358	267	95	-
2004 1/.....	1,241	1	2	7	9	11	42	166	305	358	265	75	-
2003 1/.....	1,092	-	7	2	13	11	44	153	239	301	234	88	-
2002.....	1,081	-	2	7	10	14	44	138	265	307	236	58	-
2001.....	1,103	3	3	3	15	19	53	135	244	328	225	75	-
2000.....	997	2	2	4	7	13	49	116	226	291	213	74	-
1999.....	949	-	1	5	7	14	35	113	205	292	208	69	-
American Indian 2/, female													
2007 1/.....	1,216	-	1	2	6	7	46	167	273	325	276	113	-
2006 1/.....	1,230	-	5	2	6	19	58	157	263	328	280	112	-
2005 1/.....	1,168	-	1	-	5	13	54	179	259	318	251	88	-
2004 1/.....	1,151	-	1	1	7	14	64	147	259	312	246	100	-
2003 1/.....	1,062	-	-	6	6	13	57	150	226	256	260	88	-
2002.....	1,094	1	1	5	10	21	56	155	220	282	224	119	-
2001.....	1,052	-	1	2	7	15	61	147	226	268	228	97	-
2000.....	917	-	1	4	6	13	56	104	191	248	199	95	-
1999.....	887	1	3	1	6	12	50	104	201	243	197	69	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms (C00-C97)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	12,326	2	21	34	74	160	551	1,462	2,413	2,900	3,185	1,524	-
2006 1/.....	11,784	3	19	37	70	186	554	1,431	2,319	2,827	2,942	1,395	1
2005 1/.....	11,550	1	16	37	73	190	538	1,447	2,261	2,771	2,876	1,340	-
2004 1/.....	10,863	3	20	48	59	158	549	1,426	2,023	2,709	2,693	1,175	-
2003 1/.....	10,532	4	11	42	61	182	525	1,378	1,914	2,652	2,669	1,093	1
2002.....	9,998	5	19	42	61	171	500	1,378	1,941	2,465	2,413	1,003	-
2001.....	9,792	3	15	47	61	169	515	1,349	1,827	2,471	2,397	938	-
2000.....	9,221	4	17	38	66	169	543	1,189	1,710	2,386	2,200	898	1
1999.....	8,813	3	12	43	66	161	507	1,145	1,660	2,336	2,044	835	1

Asian or Pacific Islander 3/, male

2007 1/.....	6,394	1	12	19	45	82	235	709	1,211	1,619	1,706	755	-
2006 1/.....	5,974	3	8	20	39	89	238	658	1,194	1,508	1,524	693	-
2005 1/.....	5,921	1	9	21	44	90	231	683	1,135	1,523	1,528	656	-
2004 1/.....	5,696	3	12	28	41	73	263	669	1,054	1,453	1,488	612	-
2003 1/.....	5,403	1	7	21	33	92	232	641	972	1,382	1,462	560	-
2002.....	5,193	3	12	28	36	92	236	657	1,035	1,259	1,318	517	-
2001.....	5,147	1	7	30	32	78	244	632	1,012	1,274	1,337	500	-
2000.....	4,865	2	7	22	38	79	247	554	915	1,295	1,213	492	1
1999.....	4,637	2	6	20	48	87	235	538	867	1,259	1,112	462	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms (C00-C97)

Asian or Pacific Islander 3/, female

2007 1/.....	5,932	1	9	15	29	78	316	753	1,202	1,281	1,479	769	-
2006 1/.....	5,810	-	11	17	31	97	316	773	1,125	1,319	1,418	702	1
2005 1/.....	5,629	-	7	16	29	100	307	764	1,126	1,248	1,348	684	-
2004 1/.....	5,167	-	8	20	18	85	286	757	969	1,256	1,205	563	-
2003 1/.....	5,129	3	4	21	28	90	293	737	942	1,270	1,207	533	1
2002.....	4,805	2	7	14	25	79	264	721	906	1,206	1,095	486	-
2001.....	4,645	2	8	17	29	91	271	717	815	1,197	1,060	438	-
2000.....	4,356	2	10	16	28	90	296	635	795	1,091	987	406	-
1999.....	4,176	1	6	23	18	74	272	607	793	1,077	932	373	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

All races, both sexes

2007.....	8,067	-	-	1	18	44	251	1,136	2,019	1,915	1,741	942	-
2006.....	7,720	1	-	2	14	41	272	1,128	1,823	1,814	1,760	865	-
2005.....	7,773	-	1	2	24	54	258	1,123	1,936	1,835	1,673	867	-
2004.....	7,826	-	2	5	23	43	256	1,193	1,781	1,874	1,774	875	-
2003.....	7,778	-	4	8	30	59	274	1,131	1,826	1,874	1,692	879	1
2002.....	7,737	-	3	8	23	57	258	1,101	1,811	1,898	1,721	857	-
2001.....	7,702	-	-	4	29	56	283	1,076	1,740	1,912	1,787	814	1
2000.....	7,492	-	-	4	26	50	293	1,068	1,642	1,959	1,626	824	-
1999.....	7,486	-	2	6	27	43	291	1,061	1,616	1,896	1,744	800	-

All races, male

2007.....	5,510	-	-	-	10	23	168	857	1,582	1,375	1,068	427	-
2006.....	5,266	1	-	-	7	29	189	881	1,423	1,286	1,081	369	-
2005.....	5,273	-	-	1	16	32	188	874	1,501	1,309	999	353	-
2004.....	5,312	-	-	2	18	35	176	937	1,400	1,312	1,057	375	-
2003.....	5,298	-	2	4	21	39	195	921	1,423	1,304	999	389	1
2002.....	5,184	-	2	3	11	39	179	870	1,392	1,317	997	374	-
2001.....	5,156	-	-	1	12	35	202	846	1,372	1,327	1,025	335	1
2000.....	4,952	-	-	4	13	28	215	825	1,270	1,326	939	332	-
1999.....	4,944	-	1	3	20	27	204	820	1,231	1,308	1,007	323	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

All races, female

2007.....	2,557	-	-	1	8	21	83	279	437	540	673	515	-
2006.....	2,454	-	-	2	7	12	83	247	400	528	679	496	-
2005.....	2,500	-	1	1	8	22	70	249	435	526	674	514	-
2004.....	2,514	-	2	3	5	8	80	256	381	562	717	500	-
2003.....	2,480	-	2	4	9	20	79	210	403	570	693	490	-
2002.....	2,553	-	1	5	12	18	79	231	419	581	724	483	-
2001.....	2,546	-	-	3	17	21	81	230	368	585	762	479	-
2000.....	2,540	-	-	-	13	22	78	243	372	633	687	492	-
1999.....	2,542	-	1	3	7	16	87	241	385	588	737	477	-

White, both sexes

2007 1/.....	6,720	-	-	1	11	27	172	856	1,627	1,607	1,551	868	-
2006 1/.....	6,419	1	-	2	7	32	197	855	1,448	1,501	1,571	805	-
2005 1/.....	6,409	-	1	2	12	31	187	828	1,538	1,531	1,487	792	-
2004 1/.....	6,443	-	2	3	14	29	162	853	1,401	1,585	1,581	813	-
2003 1/.....	6,409	-	3	5	19	41	196	824	1,467	1,561	1,483	809	1
2002.....	6,432	-	2	5	15	39	180	795	1,428	1,627	1,558	783	-
2001.....	6,369	-	-	2	21	37	178	793	1,361	1,601	1,620	756	-
2000.....	6,135	-	-	2	18	35	198	766	1,255	1,644	1,461	756	-
1999.....	6,116	-	2	4	14	30	194	717	1,248	1,603	1,560	744	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
White, male													
2007 1/.....	4,555	-	-	-	5	10	118	656	1,290	1,137	949	390	-
2006 1/.....	4,321	1	-	-	2	23	137	678	1,128	1,053	958	341	-
2005 1/.....	4,247	-	-	1	9	19	130	642	1,181	1,074	872	319	-
2004 1/.....	4,270	-	-	-	10	25	109	667	1,099	1,088	934	338	-
2003 1/.....	4,282	-	1	2	13	25	153	675	1,126	1,068	862	356	1
2002.....	4,256	-	2	2	8	28	127	630	1,097	1,118	904	340	-
2001.....	4,183	-	-	1	7	23	130	627	1,062	1,103	920	310	-
2000.....	3,942	-	-	2	7	20	138	580	966	1,096	827	306	-
1999.....	3,955	-	1	2	9	17	134	555	948	1,108	886	295	-
White, female													
2007 1/.....	2,165	-	-	1	6	17	54	200	337	470	602	478	-
2006 1/.....	2,098	-	-	2	5	9	60	177	320	448	613	464	-
2005 1/.....	2,162	-	1	1	3	12	57	186	357	457	615	473	-
2004 1/.....	2,173	-	2	3	4	4	53	186	302	497	647	475	-
2003 1/.....	2,127	-	2	3	6	16	43	149	341	493	621	453	-
2002.....	2,176	-	-	3	7	11	53	165	331	509	654	443	-
2001.....	2,186	-	-	1	14	14	48	166	299	498	700	446	-
2000.....	2,193	-	-	-	11	15	60	186	289	548	634	450	-
1999.....	2,161	-	1	2	5	13	60	162	300	495	674	449	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)													
Black, both sexes													
2007 1/.....	1,062	-	-	-	7	9	54	216	328	252	139	57	-
2006 1/.....	1,015	-	-	-	7	6	47	222	306	249	138	40	-
2005 1/.....	1,080	-	-	-	10	14	49	252	325	246	137	47	-
2004 1/.....	1,149	-	-	2	5	11	76	297	323	239	152	44	-
2003 1/.....	1,107	-	1	3	9	11	63	244	309	264	150	53	-
2002.....	1,044	-	-	2	6	11	65	255	319	218	114	54	-
2001.....	1,065	-	-	1	4	6	72	230	319	259	131	42	1
2000.....	1,102	-	-	2	7	7	61	252	330	256	138	49	-
1999.....	1,145	-	-	2	8	8	67	290	322	250	153	45	-
Black, male													
2007 1/.....	769	-	-	-	5	8	37	160	245	198	88	28	-
2006 1/.....	751	-	-	-	5	5	33	166	243	190	93	16	-
2005 1/.....	841	-	-	-	5	9	41	202	266	194	101	23	-
2004 1/.....	874	-	-	2	5	9	51	237	256	192	95	27	-
2003 1/.....	837	-	1	2	6	10	32	199	260	204	99	24	-
2002.....	766	-	-	1	3	7	43	204	258	161	64	25	-
2001.....	782	-	-	-	2	5	46	180	264	187	80	17	1
2000.....	833	-	-	2	5	4	47	206	265	193	95	16	-
1999.....	828	-	-	1	6	6	47	227	246	171	103	21	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

Black, female

2007 1/.....	293	-	-	-	2	1	17	56	83	54	51	29	-
2006 1/.....	264	-	-	-	2	1	14	56	63	59	45	24	-
2005 1/.....	239	-	-	-	5	5	8	50	59	52	36	24	-
2004 1/.....	275	-	-	-	-	2	25	60	67	47	57	17	-
2003 1/.....	270	-	-	1	3	1	31	45	49	60	51	29	-
2002.....	278	-	-	1	3	4	22	51	61	57	50	29	-
2001.....	283	-	-	1	2	1	26	50	55	72	51	25	-
2000.....	269	-	-	-	2	3	14	46	65	63	43	33	-
1999.....	317	-	-	1	2	2	20	63	76	79	50	24	-

American Indian 2/, both sexes

2007 1/.....	43	-	-	-	-	1	2	4	11	12	12	1	-
2006 1/.....	45	-	-	-	-	-	3	10	6	16	7	3	-
2005 1/.....	38	-	-	-	-	-	2	3	16	10	3	4	-
2004 1/.....	39	-	-	-	1	-	5	3	13	12	3	2	-
2003 1/.....	33	-	-	-	-	-	-	6	9	6	10	2	-
2002.....	43	-	-	-	-	1	2	5	15	14	6	-	-
2001.....	40	-	-	-	-	2	5	5	13	8	6	1	-
2000.....	41	-	-	-	-	-	4	4	10	17	3	3	-
1999.....	33	-	-	-	-	1	1	10	7	11	2	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

American Indian 2/, male

2007 1/.....	26	-	-	-	-	1	-	3	7	8	7	-	-
2006 1/.....	29	-	-	-	-	-	2	6	5	11	4	1	-
2005 1/.....	31	-	-	-	-	-	2	3	13	7	3	3	-
2004 1/.....	31	-	-	-	1	-	4	3	10	9	2	2	-
2003 1/.....	17	-	-	-	-	-	-	5	4	4	3	1	-
2002.....	27	-	-	-	-	1	1	4	9	7	5	-	-
2001.....	33	-	-	-	-	1	4	4	12	8	3	1	-
2000.....	27	-	-	-	-	-	4	3	5	12	2	1	-
1999.....	22	-	-	-	-	1	1	7	5	7	-	1	-

American Indian 2/, female

2007 1/.....	17	-	-	-	-	-	2	1	4	4	5	1	-
2006 1/.....	16	-	-	-	-	-	1	4	1	5	3	2	-
2005 1/.....	7	-	-	-	-	-	-	-	3	3	-	1	-
2004 1/.....	8	-	-	-	-	-	1	-	3	3	1	-	-
2003 1/.....	16	-	-	-	-	-	-	1	5	2	7	1	-
2002.....	16	-	-	-	-	-	1	1	6	7	1	-	-
2001.....	7	-	-	-	-	1	1	1	1	-	3	-	-
2000.....	14	-	-	-	-	-	-	1	5	5	1	2	-
1999.....	11	-	-	-	-	-	-	3	2	4	2	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	242	-	-	-	-	7	23	60	53	44	39	16	-
2006 1/.....	241	-	-	-	-	3	25	41	63	48	44	17	-
2005 1/.....	246	-	-	-	2	9	20	40	57	48	46	24	-
2004 1/.....	195	-	-	-	3	3	13	40	44	38	38	16	-
2003 1/.....	229	-	-	-	2	7	15	57	41	43	49	15	-
2002.....	218	-	1	1	2	6	11	46	49	39	43	20	-
2001.....	228	-	-	1	4	11	28	48	47	44	30	15	-
2000.....	214	-	-	-	1	8	30	46	47	42	24	16	-
1999.....	192	-	-	-	5	4	29	44	39	32	29	10	-

Asian or Pacific Islander 3/, male

2007 1/.....	160	-	-	-	-	4	13	38	40	32	24	9	-
2006 1/.....	165	-	-	-	-	1	17	31	47	32	26	11	-
2005 1/.....	154	-	-	-	2	4	15	27	41	34	23	8	-
2004 1/.....	137	-	-	-	2	1	12	30	35	23	26	8	-
2003 1/.....	162	-	-	-	2	4	10	42	33	28	35	8	-
2002.....	135	-	-	-	-	3	8	32	28	31	24	9	-
2001.....	158	-	-	-	3	6	22	35	34	29	22	7	-
2000.....	150	-	-	-	1	4	26	36	34	25	15	9	-
1999.....	139	-	-	-	5	3	22	31	32	22	18	6	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

Asian or Pacific Islander 3/, female

2007 1/.....	82	-	-	-	-	3	10	22	13	12	15	7	-
2006 1/.....	76	-	-	-	-	2	8	10	16	16	18	6	-
2005 1/.....	92	-	-	-	-	5	5	13	16	14	23	16	-
2004 1/.....	58	-	-	-	1	2	1	10	9	15	12	8	-
2003 1/.....	67	-	-	-	-	3	5	15	8	15	14	7	-
2002.....	83	-	1	1	2	3	3	14	21	8	19	11	-
2001.....	70	-	-	1	1	5	6	13	13	15	8	8	-
2000.....	64	-	-	-	-	4	4	10	13	17	9	7	-
1999.....	53	-	-	-	-	1	7	13	7	10	11	4	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasm of esophagus (C15)

All races, both sexes

2007.....	13,592	-	-	-	5	28	246	1,452	3,379	3,726	3,376	1,380	-
2006.....	13,686	-	-	-	3	29	279	1,524	3,289	3,691	3,515	1,355	1
2005.....	13,499	-	-	1	3	35	316	1,510	3,100	3,682	3,581	1,271	-
2004.....	13,023	-	-	-	5	36	291	1,423	2,991	3,554	3,467	1,256	-
2003.....	12,860	-	-	-	5	29	288	1,407	2,940	3,619	3,392	1,180	-
2002.....	12,701	-	-	1	6	19	311	1,390	2,763	3,691	3,389	1,130	1
2001.....	12,530	-	-	-	6	29	309	1,408	2,554	3,779	3,307	1,137	1
2000.....	12,232	1	-	-	3	39	297	1,348	2,610	3,671	3,163	1,100	-
1999.....	11,918	-	-	-	-	37	293	1,262	2,531	3,694	2,986	1,114	1

All races, male

2007.....	10,750	-	-	-	4	25	207	1,244	2,867	3,006	2,537	860	-
2006.....	10,723	-	-	-	3	25	241	1,274	2,801	2,974	2,585	819	1
2005.....	10,611	-	-	-	3	30	263	1,302	2,619	2,977	2,679	738	-
2004.....	10,081	-	-	-	4	28	242	1,211	2,502	2,817	2,546	731	-
2003.....	9,921	-	-	-	4	23	240	1,202	2,483	2,851	2,464	654	-
2002.....	9,789	-	-	-	3	14	257	1,185	2,321	2,952	2,454	603	-
2001.....	9,607	-	-	-	5	26	255	1,204	2,137	2,989	2,365	625	1
2000.....	9,279	1	-	-	2	31	265	1,161	2,169	2,831	2,233	586	-
1999.....	9,094	-	-	-	-	29	255	1,094	2,081	2,925	2,136	573	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of esophagus (C15)													
All races, female													
2007.....	2,842	-	-	-	1	3	39	208	512	720	839	520	-
2006.....	2,963	-	-	-	-	4	38	250	488	717	930	536	-
2005.....	2,888	-	-	1	-	5	53	208	481	705	902	533	-
2004.....	2,942	-	-	-	1	8	49	212	489	737	921	525	-
2003.....	2,939	-	-	-	1	6	48	205	457	768	928	526	-
2002.....	2,912	-	-	1	3	5	54	205	442	739	935	527	1
2001.....	2,923	-	-	-	1	3	54	204	417	790	942	512	-
2000.....	2,953	-	-	-	1	8	32	187	441	840	930	514	-
1999.....	2,824	-	-	-	-	8	38	168	450	769	850	541	-
White, both sexes													
2007 1/.....	11,878	-	-	-	5	26	206	1,200	2,847	3,270	3,049	1,275	-
2006 1/.....	11,959	-	-	-	1	27	244	1,235	2,786	3,216	3,194	1,255	1
2005 1/.....	11,783	-	-	1	3	31	254	1,235	2,611	3,212	3,249	1,187	-
2004 1/.....	11,251	-	-	-	3	27	246	1,137	2,488	3,063	3,138	1,149	-
2003 1/.....	11,058	-	-	-	5	25	239	1,088	2,424	3,115	3,065	1,097	-
2002.....	10,910	-	-	1	6	17	251	1,128	2,262	3,150	3,051	1,044	-
2001.....	10,624	-	-	-	6	24	250	1,107	2,028	3,169	2,992	1,047	1
2000.....	10,406	-	-	-	3	38	249	1,051	2,082	3,146	2,817	1,020	-
1999.....	10,079	-	-	-	-	32	229	979	2,009	3,139	2,666	1,025	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of esophagus (C15)													
White, male													
2007 1/.....	9,493	-	-	-	4	24	180	1,053	2,456	2,670	2,310	796	-
2006 1/.....	9,476	-	-	-	1	23	213	1,056	2,424	2,615	2,384	759	1
2005 1/.....	9,364	-	-	-	3	28	216	1,099	2,239	2,623	2,455	701	-
2004 1/.....	8,831	-	-	-	3	21	213	1,001	2,126	2,466	2,331	670	-
2003 1/.....	8,617	-	-	-	4	21	204	950	2,088	2,476	2,264	610	-
2002.....	8,510	-	-	-	3	13	216	986	1,932	2,563	2,236	561	-
2001.....	8,205	-	-	-	5	21	209	962	1,726	2,550	2,157	574	1
2000.....	7,986	-	-	-	2	31	229	929	1,766	2,466	2,016	547	-
1999.....	7,740	-	-	-	-	24	203	865	1,678	2,530	1,922	518	-
White, female													
2007 1/.....	2,385	-	-	-	1	2	26	147	391	600	739	479	-
2006 1/.....	2,483	-	-	-	-	4	31	179	362	601	810	496	-
2005 1/.....	2,419	-	-	1	-	3	38	136	372	589	794	486	-
2004 1/.....	2,420	-	-	-	-	6	33	136	362	597	807	479	-
2003 1/.....	2,441	-	-	-	1	4	35	138	336	639	801	487	-
2002.....	2,400	-	-	1	3	4	35	142	330	587	815	483	-
2001.....	2,419	-	-	-	1	3	41	145	302	619	835	473	-
2000.....	2,420	-	-	-	1	7	20	122	316	680	801	473	-
1999.....	2,339	-	-	-	-	8	26	114	331	609	744	507	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of esophagus (C15)													
Black, both sexes													
2007 1/.....	1,453	-	-	-	-	1	34	222	467	392	262	75	-
2006 1/.....	1,452	-	-	-	1	2	31	260	440	398	250	70	-
2005 1/.....	1,468	-	-	-	-	4	51	246	431	397	274	65	-
2004 1/.....	1,527	-	-	-	2	6	39	254	451	428	271	76	-
2003 1/.....	1,591	-	-	-	-	3	47	290	467	437	274	73	-
2002.....	1,590	-	-	-	-	2	56	238	457	481	285	70	1
2001.....	1,688	-	-	-	-	3	55	264	470	540	279	77	-
2000.....	1,655	1	-	-	-	1	44	280	483	466	309	71	-
1999.....	1,648	-	-	-	-	3	54	265	477	491	282	75	1
Black, male													
2007 1/.....	1,055	-	-	-	-	-	21	165	357	291	174	47	-
2006 1/.....	1,045	-	-	-	1	2	25	196	326	300	156	39	-
2005 1/.....	1,075	-	-	-	-	2	39	186	334	298	187	29	-
2004 1/.....	1,060	-	-	-	1	5	26	189	330	298	172	39	-
2003 1/.....	1,133	-	-	-	-	2	36	226	354	315	163	37	-
2002.....	1,133	-	-	-	-	1	38	179	354	345	181	35	-
2001.....	1,231	-	-	-	-	3	42	210	367	382	186	41	-
2000.....	1,170	1	-	-	-	-	32	219	371	321	193	33	-
1999.....	1,203	-	-	-	-	3	45	212	369	344	185	44	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Malignant neoplasm of esophagus (C15)

Black, female

2007 1/.....	398	-	-	-	-	1	13	57	110	101	88	28	-
2006 1/.....	407	-	-	-	-	-	6	64	114	98	94	31	-
2005 1/.....	393	-	-	-	-	2	12	60	97	99	87	36	-
2004 1/.....	467	-	-	-	1	1	13	65	121	130	99	37	-
2003 1/.....	458	-	-	-	-	1	11	64	113	122	111	36	-
2002.....	457	-	-	-	-	1	18	59	103	136	104	35	1
2001.....	457	-	-	-	-	-	13	54	103	158	93	36	-
2000.....	485	-	-	-	-	1	12	61	112	145	116	38	-
1999.....	445	-	-	-	-	-	9	53	108	147	97	31	-

American Indian 2/, both sexes

2007 1/.....	52	-	-	-	-	-	-	10	18	9	10	5	-
2006 1/.....	66	-	-	-	-	-	-	13	18	21	12	2	-
2005 1/.....	69	-	-	-	-	-	4	12	17	19	14	3	-
2004 1/.....	58	-	-	-	-	-	3	5	17	24	3	6	-
2003 1/.....	49	-	-	-	-	-	-	11	12	15	11	-	-
2002.....	47	-	-	-	-	-	2	5	9	12	16	3	-
2001.....	52	-	-	-	-	-	3	6	14	16	10	3	-
2000.....	43	-	-	-	-	-	2	6	11	13	9	2	-
1999.....	38	-	-	-	-	-	1	4	14	11	8	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Malignant neoplasm of esophagus (C15)

American Indian 2/, male

2007 1/.....	43	-	-	-	-	-	-	8	17	7	10	1	-
2006 1/.....	45	-	-	-	-	-	-	9	13	18	4	1	-
2005 1/.....	52	-	-	-	-	-	4	7	14	15	10	2	-
2004 1/.....	45	-	-	-	-	-	1	4	15	20	-	5	-
2003 1/.....	38	-	-	-	-	-	-	10	7	13	8	-	-
2002.....	37	-	-	-	-	-	-	1	5	8	9	12	2
2001.....	38	-	-	-	-	-	-	3	5	10	13	5	2
2000.....	36	-	-	-	-	-	-	2	6	9	12	6	1
1999.....	31	-	-	-	-	-	-	1	4	10	11	5	-

American Indian 2/, female

2007 1/.....	9	-	-	-	-	-	-	2	1	2	-	4	-
2006 1/.....	21	-	-	-	-	-	-	4	5	3	8	1	-
2005 1/.....	17	-	-	-	-	-	-	5	3	4	4	1	-
2004 1/.....	13	-	-	-	-	-	2	1	2	4	3	1	-
2003 1/.....	11	-	-	-	-	-	-	1	5	2	3	-	-
2002.....	10	-	-	-	-	-	-	1	-	1	3	4	1
2001.....	14	-	-	-	-	-	-	-	1	4	3	5	1
2000.....	7	-	-	-	-	-	-	-	2	1	3	1	-
1999.....	7	-	-	-	-	-	-	-	4	-	3	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasm of esophagus (C15)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	209	-	-	-	-	1	6	20	47	55	55	25	-
2006 1/.....	209	-	-	-	1	-	4	16	45	56	59	28	-
2005 1/.....	179	-	-	-	-	-	7	17	41	54	44	16	-
2004 1/.....	187	-	-	-	-	3	3	27	35	39	55	25	-
2003 1/.....	162	-	-	-	-	1	2	18	37	52	42	10	-
2002.....	154	-	-	-	-	-	2	19	35	48	37	13	-
2001.....	166	-	-	-	-	2	1	31	42	54	26	10	-
2000.....	128	-	-	-	-	-	2	11	34	46	28	7	-
1999.....	153	-	-	-	-	2	9	14	31	53	30	14	-

Asian or Pacific Islander 3/, male

2007 1/.....	159	-	-	-	-	1	6	18	37	38	43	16	-
2006 1/.....	157	-	-	-	1	-	3	13	38	41	41	20	-
2005 1/.....	120	-	-	-	-	-	4	10	32	41	27	6	-
2004 1/.....	145	-	-	-	-	2	2	17	31	33	43	17	-
2003 1/.....	133	-	-	-	-	-	-	16	34	47	29	7	-
2002.....	109	-	-	-	-	-	2	15	27	35	25	5	-
2001.....	133	-	-	-	-	2	1	27	34	44	17	8	-
2000.....	87	-	-	-	-	-	2	7	23	32	18	5	-
1999.....	120	-	-	-	-	2	6	13	24	40	24	11	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasm of esophagus (C15)

Asian or Pacific Islander 3/, female

2007 1/.....	50	-	-	-	-	-	-	2	10	17	12	9	-
2006 1/.....	52	-	-	-	-	-	1	3	7	15	18	8	-
2005 1/.....	59	-	-	-	-	-	3	7	9	13	17	10	-
2004 1/.....	42	-	-	-	-	1	1	10	4	6	12	8	-
2003 1/.....	29	-	-	-	-	1	2	2	3	5	13	3	-
2002.....	45	-	-	-	-	-	-	4	8	13	12	8	-
2001.....	33	-	-	-	-	-	-	4	8	10	9	2	-
2000.....	41	-	-	-	-	-	-	4	11	14	10	2	-
1999.....	33	-	-	-	-	-	3	1	7	13	6	3	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of stomach (C16)													
All races, both sexes													
2007.....	11,388	1	-	-	14	127	456	1,151	1,879	2,555	3,255	1,950	-
2006.....	11,345	-	-	1	18	124	465	1,165	1,870	2,462	3,343	1,897	-
2005.....	11,514	-	-	-	20	115	448	1,103	1,783	2,615	3,470	1,960	-
2004.....	11,859	-	-	-	12	122	478	1,142	1,887	2,752	3,415	2,051	-
2003.....	12,110	-	-	1	19	127	480	1,210	1,874	2,793	3,549	2,057	-
2002.....	12,198	-	1	-	29	138	436	1,106	1,831	2,816	3,656	2,185	-
2001.....	12,319	-	-	-	20	122	464	1,127	1,771	3,021	3,793	2,001	-
2000.....	12,645	-	-	1	14	124	484	1,097	1,804	3,054	3,833	2,234	-
1999.....	12,711	-	-	1	13	136	505	1,101	1,860	3,152	3,860	2,083	-
All races, male													
2007.....	6,757	-	-	-	7	60	248	704	1,259	1,700	1,871	908	-
2006.....	6,653	-	-	1	10	58	233	729	1,262	1,564	1,949	847	-
2005.....	6,752	-	-	-	9	58	266	693	1,204	1,681	1,972	869	-
2004.....	6,958	-	-	-	7	63	269	722	1,273	1,734	1,941	949	-
2003.....	7,055	-	-	1	7	63	262	767	1,282	1,826	1,980	867	-
2002.....	7,133	-	1	-	8	80	239	725	1,215	1,867	2,064	934	-
2001.....	7,191	-	-	-	9	71	253	731	1,238	1,928	2,148	813	-
2000.....	7,328	-	-	-	6	69	284	703	1,228	1,966	2,144	928	-
1999.....	7,459	-	-	1	6	76	283	746	1,252	2,005	2,239	851	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of stomach (C16)													
All races, female													
2007.....	4,631	1	-	-	7	67	208	447	620	855	1,384	1,042	-
2006.....	4,692	-	-	-	8	66	232	436	608	898	1,394	1,050	-
2005.....	4,762	-	-	-	11	57	182	410	579	934	1,498	1,091	-
2004.....	4,901	-	-	-	5	59	209	420	614	1,018	1,474	1,102	-
2003.....	5,055	-	-	-	12	64	218	443	592	967	1,569	1,190	-
2002.....	5,065	-	-	-	21	58	197	381	616	949	1,592	1,251	-
2001.....	5,128	-	-	-	11	51	211	396	533	1,093	1,645	1,188	-
2000.....	5,317	-	-	1	8	55	200	394	576	1,088	1,689	1,306	-
1999.....	5,252	-	-	-	7	60	222	355	608	1,147	1,621	1,232	-
White, both sexes													
2007 1/.....	8,551	1	-	-	9	98	332	793	1,321	1,901	2,517	1,579	-
2006 1/.....	8,582	-	-	-	12	89	305	833	1,369	1,836	2,595	1,543	-
2005 1/.....	8,711	-	-	-	14	69	296	779	1,302	1,949	2,709	1,593	-
2004 1/.....	9,067	-	-	-	7	82	334	784	1,376	2,050	2,736	1,698	-
2003 1/.....	9,282	-	-	1	15	88	323	827	1,410	2,149	2,823	1,646	-
2002.....	9,319	-	1	-	21	94	301	770	1,354	2,124	2,902	1,752	-
2001.....	9,501	-	-	-	10	84	310	794	1,315	2,314	3,036	1,638	-
2000.....	9,723	-	-	1	9	78	318	777	1,352	2,336	3,036	1,816	-
1999.....	9,779	-	-	-	8	88	321	771	1,348	2,459	3,098	1,686	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of stomach (C16)													
White, male													
2007 1/.....	5,099	-	-	-	4	48	189	483	899	1,291	1,444	741	-
2006 1/.....	5,102	-	-	-	7	39	155	535	939	1,190	1,540	697	-
2005 1/.....	5,107	-	-	-	6	41	173	489	880	1,251	1,559	708	-
2004 1/.....	5,332	-	-	-	5	46	193	506	938	1,291	1,558	795	-
2003 1/.....	5,428	-	-	1	7	46	183	529	967	1,404	1,592	699	-
2002.....	5,542	-	1	-	5	57	168	516	918	1,439	1,677	761	-
2001.....	5,568	-	-	-	4	48	176	519	921	1,477	1,744	679	-
2000.....	5,688	-	-	-	3	45	181	503	936	1,534	1,722	764	-
1999.....	5,786	-	-	-	4	55	182	534	923	1,593	1,811	684	-
White, female													
2007 1/.....	3,452	1	-	-	5	50	143	310	422	610	1,073	838	-
2006 1/.....	3,480	-	-	-	5	50	150	298	430	646	1,055	846	-
2005 1/.....	3,604	-	-	-	8	28	123	290	422	698	1,150	885	-
2004 1/.....	3,735	-	-	-	2	36	141	278	438	759	1,178	903	-
2003 1/.....	3,854	-	-	-	8	42	140	298	443	745	1,231	947	-
2002.....	3,777	-	-	-	16	37	133	254	436	685	1,225	991	-
2001.....	3,933	-	-	-	6	36	134	275	394	837	1,292	959	-
2000.....	4,035	-	-	1	6	33	137	274	416	802	1,314	1,052	-
1999.....	3,993	-	-	-	4	33	139	237	425	866	1,287	1,002	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasm of stomach (C16)

Black, both sexes

2007 1/.....	1,998	-	-	-	3	18	81	262	400	460	512	262	-
2006 1/.....	1,916	-	-	1	3	20	99	226	351	435	523	258	-
2005 1/.....	1,973	-	-	-	3	24	101	231	343	466	527	278	-
2004 1/.....	2,034	-	-	-	4	28	93	260	371	524	501	253	-
2003 1/.....	2,041	-	-	-	1	22	103	275	350	463	533	294	-
2002.....	2,131	-	-	-	8	33	92	250	340	522	548	338	-
2001.....	2,067	-	-	-	5	25	121	240	328	509	552	287	-
2000.....	2,176	-	-	-	4	36	118	240	338	529	586	325	-
1999.....	2,203	-	-	1	3	36	127	240	378	512	585	321	-

Black, male

2007 1/.....	1,193	-	-	-	2	8	42	176	264	295	293	113	-
2006 1/.....	1,092	-	-	1	2	13	51	145	233	255	298	94	-
2005 1/.....	1,162	-	-	-	2	11	68	151	243	300	264	123	-
2004 1/.....	1,191	-	-	-	2	12	50	164	251	341	270	101	-
2003 1/.....	1,187	-	-	-	-	10	55	184	241	313	282	102	-
2002.....	1,194	-	-	-	3	20	49	166	223	327	278	128	-
2001.....	1,186	-	-	-	2	14	63	163	223	333	296	92	-
2000.....	1,218	-	-	-	2	20	78	155	225	319	301	118	-
1999.....	1,263	-	-	1	1	19	72	159	245	313	327	126	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Malignant neoplasm of stomach (C16)

Black, female

2007 1/.....	805	-	-	-	1	10	39	86	136	165	219	149	-
2006 1/.....	824	-	-	-	1	7	48	81	118	180	225	164	-
2005 1/.....	811	-	-	-	1	13	33	80	100	166	263	155	-
2004 1/.....	843	-	-	-	2	16	43	96	120	183	231	152	-
2003 1/.....	854	-	-	-	1	12	48	91	109	150	251	192	-
2002.....	937	-	-	-	5	13	43	84	117	195	270	210	-
2001.....	881	-	-	-	3	11	58	77	105	176	256	195	-
2000.....	958	-	-	-	2	16	40	85	113	210	285	207	-
1999.....	940	-	-	-	2	17	55	81	133	199	258	195	-

American Indian 2/, both sexes

2007 1/.....	90	-	-	-	-	1	6	14	22	25	17	5	-
2006 1/.....	84	-	-	-	-	2	6	13	16	20	19	8	-
2005 1/.....	85	-	-	-	-	1	8	12	21	21	15	7	-
2004 1/.....	78	-	-	-	-	-	6	12	16	20	15	9	-
2003 1/.....	84	-	-	-	1	4	7	16	10	22	16	8	-
2002.....	91	-	-	-	-	1	6	16	16	21	25	6	-
2001.....	79	-	-	-	1	2	1	7	15	17	28	8	-
2000.....	73	-	-	-	-	-	4	5	16	25	15	8	-
1999.....	67	-	-	-	1	1	5	5	11	23	16	5	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Malignant neoplasm of stomach (C16)

American Indian 2/, male

2007 1/.....	53	-	-	-	-	1	3	8	15	16	8	2	-
2006 1/.....	58	-	-	-	-	2	5	6	10	15	12	8	-
2005 1/.....	52	-	-	-	-	1	2	7	12	14	12	4	-
2004 1/.....	46	-	-	-	-	-	2	7	8	13	13	3	-
2003 1/.....	50	-	-	-	-	2	2	10	6	17	8	5	-
2002.....	45	-	-	-	-	1	4	8	5	12	12	3	-
2001.....	51	-	-	-	1	2	1	4	13	10	14	6	-
2000.....	41	-	-	-	-	-	3	4	9	13	8	4	-
1999.....	44	-	-	-	-	1	3	3	7	15	11	4	-

American Indian 2/, female

2007 1/.....	37	-	-	-	-	-	3	6	7	9	9	3	-
2006 1/.....	26	-	-	-	-	-	1	7	6	5	7	-	-
2005 1/.....	33	-	-	-	-	-	6	5	9	7	3	3	-
2004 1/.....	32	-	-	-	-	-	4	5	8	7	2	6	-
2003 1/.....	34	-	-	-	1	2	5	6	4	5	8	3	-
2002.....	46	-	-	-	-	-	2	8	11	9	13	3	-
2001.....	28	-	-	-	-	-	-	3	2	7	14	2	-
2000.....	32	-	-	-	-	-	1	1	7	12	7	4	-
1999.....	23	-	-	-	1	-	2	2	4	8	5	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasm of stomach (C16)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	749	-	-	-	2	10	37	82	136	169	209	104	-
2006 1/.....	763	-	-	-	3	13	55	93	134	171	206	88	-
2005 1/.....	745	-	-	-	3	21	43	81	117	179	219	82	-
2004 1/.....	680	-	-	-	1	12	45	86	124	158	163	91	-
2003 1/.....	703	-	-	-	2	13	47	92	104	159	177	109	-
2002.....	657	-	-	-	-	10	37	70	121	149	181	89	-
2001.....	672	-	-	-	4	11	32	86	113	181	177	68	-
2000.....	673	-	-	-	1	10	44	75	98	164	196	85	-
1999.....	662	-	-	-	1	11	52	85	123	158	161	71	-

Asian or Pacific Islander 3/, male

2007 1/.....	412	-	-	-	1	3	14	37	81	98	126	52	-
2006 1/.....	401	-	-	-	1	4	22	43	80	104	99	48	-
2005 1/.....	431	-	-	-	1	5	23	46	69	116	137	34	-
2004 1/.....	389	-	-	-	-	5	24	45	76	89	100	50	-
2003 1/.....	390	-	-	-	-	5	22	44	68	92	98	61	-
2002.....	352	-	-	-	-	2	18	35	69	89	97	42	-
2001.....	386	-	-	-	2	7	13	45	81	108	94	36	-
2000.....	381	-	-	-	1	4	22	41	58	100	113	42	-
1999.....	366	-	-	-	1	1	26	50	77	84	90	37	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasm of stomach (C16)

Asian or Pacific Islander 3/, female

2007 1/.....	337	-	-	-	1	7	23	45	55	71	83	52	-
2006 1/.....	362	-	-	-	2	9	33	50	54	67	107	40	-
2005 1/.....	314	-	-	-	2	16	20	35	48	63	82	48	-
2004 1/.....	291	-	-	-	1	7	21	41	48	69	63	41	-
2003 1/.....	313	-	-	-	2	8	25	48	36	67	79	48	-
2002.....	305	-	-	-	-	8	19	35	52	60	84	47	-
2001.....	286	-	-	-	2	4	19	41	32	73	83	32	-
2000.....	292	-	-	-	-	6	22	34	40	64	83	43	-
1999.....	296	-	-	-	-	10	26	35	46	74	71	34	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of colon, rectum and anus (C18-C21)													
All races, both sexes													
2007.....	53,586	-	-	1	35	275	1,302	4,793	9,058	11,634	15,417	11,069	2
2006.....	53,549	-	-	1	42	323	1,330	4,499	8,626	11,709	16,024	10,992	3
2005.....	53,252	1	-	4	47	305	1,297	4,343	8,153	11,792	16,254	11,054	2
2004.....	53,772	-	-	1	54	279	1,326	4,273	8,208	11,954	16,714	10,963	-
2003.....	55,958	-	-	4	40	291	1,315	4,442	8,304	12,934	17,331	11,296	1
2002.....	56,741	1	-	1	50	290	1,425	4,573	8,421	13,392	17,498	11,089	1
2001.....	56,887	-	-	-	46	299	1,433	4,548	7,984	13,791	17,527	11,259	-
2000.....	57,477	-	-	3	40	272	1,369	4,237	7,940	14,101	18,053	11,462	-
1999.....	57,155	1	-	1	49	283	1,333	4,057	7,967	14,249	17,929	11,286	-
All races, male													
2007.....	27,125	-	-	1	26	155	705	2,657	5,339	6,617	7,652	3,972	1
2006.....	26,921	-	-	1	20	166	697	2,474	5,018	6,662	7,921	3,960	2
2005.....	26,843	1	-	4	27	167	694	2,401	4,812	6,789	8,003	3,943	2
2004.....	26,930	-	-	1	34	152	695	2,442	4,785	6,792	8,189	3,840	-
2003.....	28,007	-	-	3	25	158	714	2,509	4,831	7,346	8,471	3,949	1
2002.....	28,501	1	-	1	34	161	797	2,564	5,035	7,580	8,504	3,823	1
2001.....	28,225	-	-	-	24	157	781	2,512	4,652	7,773	8,451	3,875	-
2000.....	28,462	-	-	2	24	153	759	2,359	4,608	7,924	8,760	3,873	-
1999.....	28,240	-	-	1	27	160	678	2,250	4,621	7,883	8,754	3,866	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of colon, rectum and anus (C18-C21)													
All races, female													
2007.....	26,461	-	-	-	9	120	597	2,136	3,719	5,017	7,765	7,097	1
2006.....	26,628	-	-	-	22	157	633	2,025	3,608	5,047	8,103	7,032	1
2005.....	26,409	-	-	-	20	138	603	1,942	3,341	5,003	8,251	7,111	-
2004.....	26,842	-	-	-	20	127	631	1,831	3,423	5,162	8,525	7,123	-
2003.....	27,951	-	-	1	15	133	601	1,933	3,473	5,588	8,860	7,347	-
2002.....	28,240	-	-	-	16	129	628	2,009	3,386	5,812	8,994	7,266	-
2001.....	28,662	-	-	-	22	142	652	2,036	3,332	6,018	9,076	7,384	-
2000.....	29,015	-	-	1	16	119	610	1,878	3,332	6,177	9,293	7,589	-
1999.....	28,915	1	-	-	22	123	655	1,807	3,346	6,366	9,175	7,420	-
White, both sexes													
2007 1/.....	45,174	-	-	1	21	205	997	3,674	7,174	9,693	13,474	9,934	1
2006 1/.....	45,245	-	-	-	34	229	984	3,462	6,940	9,789	13,934	9,871	2
2005 1/.....	44,994	1	-	4	35	228	993	3,285	6,503	9,860	14,201	9,883	1
2004 1/.....	45,824	-	-	1	35	209	997	3,294	6,678	10,019	14,731	9,860	-
2003 1/.....	47,711	-	-	4	29	217	965	3,370	6,734	10,910	15,330	10,151	1
2002.....	48,562	1	-	1	38	234	1,071	3,553	6,858	11,393	15,417	9,995	1
2001.....	48,748	-	-	-	34	223	1,091	3,535	6,515	11,726	15,449	10,175	-
2000.....	49,488	-	-	3	31	201	1,015	3,300	6,489	12,085	16,032	10,332	-
1999.....	49,374	-	-	1	32	212	1,020	3,199	6,537	12,228	15,944	10,201	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

White, male

2007 1/.....	22,926	-	-	1	15	114	540	2,058	4,293	5,569	6,772	3,564	-
2006 1/.....	22,775	-	-	-	19	122	520	1,916	4,106	5,620	6,928	3,543	1
2005 1/.....	22,782	1	-	4	20	120	553	1,848	3,885	5,744	7,062	3,544	1
2004 1/.....	23,038	-	-	1	21	114	542	1,919	3,960	5,756	7,248	3,477	-
2003 1/.....	24,013	-	-	3	20	118	539	1,941	3,975	6,298	7,552	3,566	1
2002.....	24,439	1	-	1	24	126	600	2,023	4,156	6,547	7,519	3,441	1
2001.....	24,335	-	-	-	16	119	599	2,000	3,867	6,715	7,522	3,497	-
2000.....	24,517	-	-	2	16	119	567	1,874	3,788	6,861	7,817	3,473	-
1999.....	24,539	-	-	1	16	118	540	1,811	3,828	6,891	7,832	3,502	-

White, female

2007 1/.....	22,248	-	-	-	6	91	457	1,616	2,881	4,124	6,702	6,370	1
2006 1/.....	22,470	-	-	-	15	107	464	1,546	2,834	4,169	7,006	6,328	1
2005 1/.....	22,212	-	-	-	15	108	440	1,437	2,618	4,116	7,139	6,339	-
2004 1/.....	22,786	-	-	-	14	95	455	1,375	2,718	4,263	7,483	6,383	-
2003 1/.....	23,698	-	-	1	9	99	426	1,429	2,759	4,612	7,778	6,585	-
2002.....	24,123	-	-	-	14	108	471	1,530	2,702	4,846	7,898	6,554	-
2001.....	24,413	-	-	-	18	104	492	1,535	2,648	5,011	7,927	6,678	-
2000.....	24,971	-	-	1	15	82	448	1,426	2,701	5,224	8,215	6,859	-
1999.....	24,835	-	-	-	16	94	480	1,388	2,709	5,337	8,112	6,699	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of colon, rectum and anus (C18-C21)													
Black, both sexes													
2007 1/.....	6,922	-	-	-	10	54	238	947	1,578	1,600	1,591	903	1
2006 1/.....	6,895	-	-	-	5	71	286	888	1,392	1,606	1,718	928	1
2005 1/.....	6,884	-	-	-	10	62	244	897	1,382	1,607	1,702	979	1
2004 1/.....	6,640	-	-	-	12	53	252	818	1,299	1,639	1,642	925	-
2003 1/.....	6,943	-	-	-	8	50	288	899	1,329	1,716	1,688	965	-
2002.....	6,874	-	-	-	10	40	284	863	1,308	1,674	1,770	925	-
2001.....	6,905	-	-	-	8	56	286	851	1,262	1,736	1,773	933	-
2000.....	6,867	-	-	-	6	54	293	823	1,225	1,732	1,743	991	-
1999.....	6,777	1	-	-	16	56	273	719	1,256	1,746	1,750	960	-
Black, male													
2007 1/.....	3,443	-	-	-	8	30	131	509	870	860	720	314	1
2006 1/.....	3,438	-	-	-	1	34	147	482	743	874	810	346	1
2005 1/.....	3,344	-	-	-	6	39	116	465	784	844	771	318	1
2004 1/.....	3,205	-	-	-	10	27	122	435	691	869	771	280	-
2003 1/.....	3,335	-	-	-	4	25	141	487	725	894	755	304	-
2002.....	3,351	-	-	-	8	25	157	455	720	852	828	306	-
2001.....	3,251	-	-	-	7	30	155	431	661	885	776	306	-
2000.....	3,326	-	-	-	5	26	161	431	681	899	796	327	-
1999.....	3,196	-	-	-	10	31	125	377	698	861	795	299	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of colon, rectum and anus (C18-C21)													
Black, female													
2007 1/.....	3,479	-	-	-	2	24	107	438	708	740	871	589	-
2006 1/.....	3,457	-	-	-	4	37	139	406	649	732	908	582	-
2005 1/.....	3,540	-	-	-	4	23	128	432	598	763	931	661	-
2004 1/.....	3,435	-	-	-	2	26	130	383	608	770	871	645	-
2003 1/.....	3,608	-	-	-	4	25	147	412	604	822	933	661	-
2002.....	3,523	-	-	-	2	15	127	408	588	822	942	619	-
2001.....	3,654	-	-	-	1	26	131	420	601	851	997	627	-
2000.....	3,541	-	-	-	1	28	132	392	544	833	947	664	-
1999.....	3,581	1	-	-	6	25	148	342	558	885	955	661	-
American Indian 2/, both sexes													
2007 1/.....	252	-	-	-	1	3	11	35	64	61	55	22	-
2006 1/.....	230	-	-	-	1	3	3	32	64	52	58	17	-
2005 1/.....	240	-	-	-	-	4	15	43	50	53	54	21	-
2004 1/.....	226	-	-	-	1	3	9	24	54	61	58	16	-
2003 1/.....	205	-	-	-	-	-	10	33	38	49	51	24	-
2002.....	246	-	-	-	-	3	11	30	64	68	46	24	-
2001.....	192	-	-	-	-	1	11	19	42	61	42	16	-
2000.....	194	-	-	-	-	3	13	18	42	51	43	24	-
1999.....	170	-	-	-	1	2	7	25	30	37	46	22	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

American Indian 2/, male

2007 1/.....	132	-	-	-	1	2	7	14	40	35	22	11	-
2006 1/.....	124	-	-	-	-	-	1	19	37	32	28	7	-
2005 1/.....	136	-	-	-	-	2	8	26	30	35	25	10	-
2004 1/.....	131	-	-	-	1	3	4	15	35	33	32	8	-
2003 1/.....	111	-	-	-	-	-	5	17	20	30	24	15	-
2002.....	131	-	-	-	-	1	6	18	47	33	21	5	-
2001.....	95	-	-	-	-	-	5	13	20	33	16	8	-
2000.....	110	-	-	-	-	1	8	9	30	30	22	10	-
1999.....	87	-	-	-	1	2	2	12	17	20	24	9	-

American Indian 2/, female

2007 1/.....	120	-	-	-	-	1	4	21	24	26	33	11	-
2006 1/.....	106	-	-	-	1	3	2	13	27	20	30	10	-
2005 1/.....	104	-	-	-	-	2	7	17	20	18	29	11	-
2004 1/.....	95	-	-	-	-	-	5	9	19	28	26	8	-
2003 1/.....	94	-	-	-	-	-	5	16	18	19	27	9	-
2002.....	115	-	-	-	-	2	5	12	17	35	25	19	-
2001.....	97	-	-	-	-	1	6	6	22	28	26	8	-
2000.....	84	-	-	-	-	2	5	9	12	21	21	14	-
1999.....	83	-	-	-	-	-	5	13	13	17	22	13	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	1,238	-	-	-	3	13	56	137	242	280	297	210	-
2006 1/.....	1,179	-	-	1	2	20	57	117	230	262	314	176	-
2005 1/.....	1,134	-	-	-	2	11	45	118	218	272	297	171	-
2004 1/.....	1,082	-	-	-	6	14	68	137	177	235	283	162	-
2003 1/.....	1,099	-	-	-	3	24	52	140	203	259	262	156	-
2002.....	1,059	-	-	-	2	13	59	127	191	257	265	145	-
2001.....	1,042	-	-	-	4	19	45	143	165	268	263	135	-
2000.....	928	-	-	-	3	14	48	96	184	233	235	115	-
1999.....	834	-	-	-	-	13	33	114	144	238	189	103	-

Asian or Pacific Islander 3/, male

2007 1/.....	624	-	-	-	2	9	27	76	136	153	138	83	-
2006 1/.....	584	-	-	1	-	10	29	57	132	136	155	64	-
2005 1/.....	581	-	-	-	1	6	17	62	113	166	145	71	-
2004 1/.....	556	-	-	-	2	8	27	73	99	134	138	75	-
2003 1/.....	548	-	-	-	1	15	29	64	111	124	140	64	-
2002.....	580	-	-	-	2	9	34	68	112	148	136	71	-
2001.....	544	-	-	-	1	8	22	68	104	140	137	64	-
2000.....	509	-	-	-	3	7	23	45	109	134	125	63	-
1999.....	418	-	-	-	-	9	11	50	78	111	103	56	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

Asian or Pacific Islander 3/, female

2007 1/.....	614	-	-	-	1	4	29	61	106	127	159	127	-
2006 1/.....	595	-	-	-	2	10	28	60	98	126	159	112	-
2005 1/.....	553	-	-	-	1	5	28	56	105	106	152	100	-
2004 1/.....	526	-	-	-	4	6	41	64	78	101	145	87	-
2003 1/.....	551	-	-	-	2	9	23	76	92	135	122	92	-
2002.....	479	-	-	-	-	4	25	59	79	109	129	74	-
2001.....	498	-	-	-	3	11	23	75	61	128	126	71	-
2000.....	419	-	-	-	-	7	25	51	75	99	110	52	-
1999.....	416	-	-	-	-	4	22	64	66	127	86	47	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

All races, both sexes

2007.....	17,146	6	19	25	38	90	368	2,503	4,181	3,884	4,266	1,766	-
2006.....	16,525	2	15	16	31	87	361	2,406	3,842	3,830	4,161	1,774	-
2005.....	16,076	8	21	16	30	89	380	2,515	3,438	3,747	4,176	1,655	1
2004.....	15,321	1	22	16	43	84	380	2,348	3,123	3,670	4,059	1,575	-
2003.....	14,706	6	23	14	36	89	395	2,223	2,765	3,630	3,949	1,576	-
2002.....	14,047	3	28	18	27	105	414	1,976	2,558	3,581	3,728	1,609	-
2001.....	13,351	5	18	15	29	75	401	1,868	2,351	3,440	3,649	1,500	-
2000.....	12,916	2	23	20	30	86	429	1,686	2,256	3,439	3,558	1,387	-
1999.....	12,382	8	29	11	43	78	444	1,492	2,091	3,491	3,425	1,270	-

All races, male

2007.....	11,343	4	13	16	17	58	258	1,969	3,241	2,552	2,446	769	-
2006.....	10,759	2	10	9	14	54	251	1,890	2,892	2,537	2,361	739	-
2005.....	10,548	6	13	9	18	63	258	2,003	2,615	2,470	2,374	719	-
2004.....	9,832	1	12	8	25	55	289	1,889	2,318	2,382	2,227	626	-
2003.....	9,482	2	18	9	20	53	288	1,779	1,995	2,330	2,301	687	-
2002.....	9,011	2	19	12	18	62	308	1,611	1,858	2,367	2,112	642	-
2001.....	8,483	3	7	7	15	43	302	1,490	1,691	2,224	2,047	654	-
2000.....	8,176	1	14	8	20	59	297	1,324	1,607	2,246	2,052	548	-
1999.....	7,689	5	20	9	24	48	318	1,161	1,465	2,284	1,877	478	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

All races, female

2007.....	5,803	2	6	9	21	32	110	534	940	1,332	1,820	997	-
2006.....	5,766	-	5	7	17	33	110	516	950	1,293	1,800	1,035	-
2005.....	5,528	2	8	7	12	26	122	512	823	1,277	1,802	936	1
2004.....	5,489	-	10	8	18	29	91	459	805	1,288	1,832	949	-
2003.....	5,224	4	5	5	16	36	107	444	770	1,300	1,648	889	-
2002.....	5,036	1	9	6	9	43	106	365	700	1,214	1,616	967	-
2001.....	4,868	2	11	8	14	32	99	378	660	1,216	1,602	846	-
2000.....	4,740	1	9	12	10	27	132	362	649	1,193	1,506	839	-
1999.....	4,693	3	9	2	19	30	126	331	626	1,207	1,548	792	-

White, both sexes

2007 1/.....	13,613	6	14	22	31	52	243	1,841	3,087	3,128	3,650	1,539	-
2006 1/.....	13,103	1	12	14	28	51	224	1,727	2,856	3,075	3,539	1,576	-
2005 1/.....	12,735	7	19	11	19	46	240	1,823	2,564	2,989	3,564	1,453	-
2004 1/.....	12,111	1	19	10	27	47	248	1,669	2,334	2,918	3,472	1,366	-
2003 1/.....	11,660	4	19	8	23	45	264	1,556	2,080	2,903	3,390	1,368	-
2002.....	11,250	2	22	13	19	57	275	1,383	1,928	2,909	3,237	1,405	-
2001.....	10,697	5	14	8	22	48	237	1,313	1,772	2,805	3,144	1,329	-
2000.....	10,304	1	17	13	20	44	261	1,164	1,680	2,784	3,093	1,227	-
1999.....	9,986	6	26	6	29	36	283	1,049	1,554	2,872	2,998	1,127	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of liver and intrahepatic bile ducts (C22)													
White, male													
2007 1/.....	8,884	4	9	14	13	32	159	1,448	2,371	2,049	2,117	668	-
2006 1/.....	8,419	1	8	8	13	24	143	1,361	2,114	2,036	2,046	665	-
2005 1/.....	8,278	6	12	8	9	32	155	1,442	1,941	1,976	2,047	650	-
2004 1/.....	7,694	1	11	5	14	31	183	1,348	1,713	1,903	1,942	543	-
2003 1/.....	7,489	2	15	7	13	23	183	1,242	1,513	1,900	1,995	596	-
2002.....	7,114	2	14	7	10	27	201	1,129	1,397	1,930	1,830	567	-
2001.....	6,774	3	5	4	12	24	175	1,044	1,280	1,860	1,783	584	-
2000.....	6,447	1	11	7	13	27	179	901	1,183	1,833	1,814	478	-
1999.....	6,151	3	18	5	17	21	200	799	1,094	1,898	1,658	438	-
White, female													
2007 1/.....	4,729	2	5	8	18	20	84	393	716	1,079	1,533	871	-
2006 1/.....	4,684	-	4	6	15	27	81	366	742	1,039	1,493	911	-
2005 1/.....	4,457	1	7	3	10	14	85	381	623	1,013	1,517	803	-
2004 1/.....	4,417	-	8	5	13	16	65	321	621	1,015	1,530	823	-
2003 1/.....	4,171	2	4	1	10	22	81	314	567	1,003	1,395	772	-
2002.....	4,136	-	8	6	9	30	74	254	531	979	1,407	838	-
2001.....	3,923	2	9	4	10	24	62	269	492	945	1,361	745	-
2000.....	3,857	-	6	6	7	17	82	263	497	951	1,279	749	-
1999.....	3,835	3	8	1	12	15	83	250	460	974	1,340	689	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

Black, both sexes

2007 1/.....	2,264	-	4	1	5	28	77	447	770	448	347	137	-
2006 1/.....	2,187	-	2	2	2	20	73	489	702	437	340	120	-
2005 1/.....	2,118	1	2	3	7	28	84	488	619	440	336	109	1
2004 1/.....	2,014	-	2	5	14	22	79	475	504	443	342	128	-
2003 1/.....	1,877	2	1	3	7	25	78	470	450	391	307	143	-
2002.....	1,734	1	4	3	4	27	84	405	392	388	298	128	-
2001.....	1,670	-	3	5	4	15	96	389	375	363	306	114	-
2000.....	1,598	1	2	2	6	21	102	365	344	391	263	101	-
1999.....	1,507	1	3	4	7	21	91	293	315	398	266	108	-

Black, male

2007 1/.....	1,592	-	3	-	4	20	60	349	619	297	181	59	-
2006 1/.....	1,515	-	1	1	1	16	56	369	568	296	168	39	-
2005 1/.....	1,483	-	1	1	6	20	58	397	488	292	186	34	-
2004 1/.....	1,349	-	-	2	9	14	60	381	387	291	158	47	-
2003 1/.....	1,234	-	-	2	4	15	59	379	319	238	157	61	-
2002.....	1,165	-	3	3	4	17	61	325	285	264	164	39	-
2001.....	1,048	-	2	2	1	10	71	305	267	207	146	37	-
2000.....	1,061	-	1	-	5	15	66	290	256	255	131	42	-
1999.....	947	1	2	3	2	14	65	240	212	248	134	26	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

Black, female

2007 1/.....	672	-	1	1	1	8	17	98	151	151	166	78	-
2006 1/.....	672	-	1	1	1	4	17	120	134	141	172	81	-
2005 1/.....	635	1	1	2	1	8	26	91	131	148	150	75	1
2004 1/.....	665	-	2	3	5	8	19	94	117	152	184	81	-
2003 1/.....	643	2	1	1	3	10	19	91	131	153	150	82	-
2002.....	569	1	1	-	-	10	23	80	107	124	134	89	-
2001.....	622	-	1	3	3	5	25	84	108	156	160	77	-
2000.....	537	1	1	2	1	6	36	75	88	136	132	59	-
1999.....	560	-	1	1	5	7	26	53	103	150	132	82	-

American Indian 2/, both sexes

2007 1/.....	151	-	-	-	-	-	5	38	40	30	28	10	-
2006 1/.....	134	-	-	-	-	1	2	24	33	38	31	5	-
2005 1/.....	119	-	-	-	-	-	6	24	28	35	20	6	-
2004 1/.....	143	-	-	-	-	2	2	28	39	21	41	10	-
2003 1/.....	105	-	2	1	-	1	4	28	27	22	16	4	-
2002.....	94	-	1	-	-	1	2	17	20	27	17	9	-
2001.....	109	-	-	-	1	1	3	22	20	36	19	7	-
2000.....	98	-	-	1	-	2	4	20	23	21	17	10	-
1999.....	76	-	-	-	1	1	3	11	18	20	17	5	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

American Indian 2/, male

2007 1/.....	100	-	-	-	-	-	4	28	33	18	13	4	-
2006 1/.....	88	-	-	-	-	-	1	21	25	22	18	1	-
2005 1/.....	73	-	-	-	-	-	3	18	23	21	5	3	-
2004 1/.....	77	-	-	-	-	2	2	22	23	10	15	3	-
2003 1/.....	71	-	2	-	-	1	4	22	15	14	12	1	-
2002.....	60	-	1	-	-	1	2	14	15	17	8	2	-
2001.....	71	-	-	-	-	-	1	19	13	27	10	1	-
2000.....	65	-	-	1	-	1	4	17	15	16	8	3	-
1999.....	42	-	-	-	1	-	2	8	13	8	9	1	-

American Indian 2/, female

2007 1/.....	51	-	-	-	-	-	1	10	7	12	15	6	-
2006 1/.....	46	-	-	-	-	1	1	3	8	16	13	4	-
2005 1/.....	46	-	-	-	-	-	3	6	5	14	15	3	-
2004 1/.....	66	-	-	-	-	-	-	6	16	11	26	7	-
2003 1/.....	34	-	-	1	-	-	-	6	12	8	4	3	-
2002.....	34	-	-	-	-	-	-	3	5	10	9	7	-
2001.....	38	-	-	-	1	1	2	3	7	9	9	6	-
2000.....	33	-	-	-	-	1	-	3	8	5	9	7	-
1999.....	34	-	-	-	-	1	1	3	5	12	8	4	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	1,118	-	1	2	2	10	43	177	284	278	241	80	-
2006 1/.....	1,101	1	1	-	1	15	62	166	251	280	251	73	-
2005 1/.....	1,104	-	-	2	4	15	50	180	227	283	256	87	-
2004 1/.....	1,053	-	1	1	2	13	51	176	246	288	204	71	-
2003 1/.....	1,064	-	1	2	6	18	49	169	208	314	236	61	-
2002.....	969	-	1	2	4	20	53	171	218	257	176	67	-
2001.....	875	-	1	2	2	11	65	144	184	236	180	50	-
2000.....	916	-	4	4	4	19	62	137	209	243	185	49	-
1999.....	813	1	-	1	6	20	67	139	204	201	144	30	-

Asian or Pacific Islander 3/, male

2007 1/.....	767	-	1	2	-	6	35	144	218	188	135	38	-
2006 1/.....	737	1	1	-	-	14	51	139	185	183	129	34	-
2005 1/.....	714	-	-	-	3	11	42	146	163	181	136	32	-
2004 1/.....	712	-	1	1	2	8	44	138	195	178	112	33	-
2003 1/.....	688	-	1	-	3	14	42	136	148	178	137	29	-
2002.....	672	-	1	2	4	17	44	143	161	156	110	34	-
2001.....	590	-	-	1	2	9	55	122	131	130	108	32	-
2000.....	603	-	2	-	2	16	48	116	153	142	99	25	-
1999.....	549	1	-	1	4	13	51	114	146	130	76	13	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

Asian or Pacific Islander 3/, female

2007 1/.....	351	-	-	-	2	4	8	33	66	90	106	42	-
2006 1/.....	364	-	-	-	1	1	11	27	66	97	122	39	-
2005 1/.....	390	-	-	2	1	4	8	34	64	102	120	55	-
2004 1/.....	341	-	-	-	-	5	7	38	51	110	92	38	-
2003 1/.....	376	-	-	2	3	4	7	33	60	136	99	32	-
2002.....	297	-	-	-	-	3	9	28	57	101	66	33	-
2001.....	285	-	1	1	-	2	10	22	53	106	72	18	-
2000.....	313	-	2	4	2	3	14	21	56	101	86	24	-
1999.....	264	-	-	-	2	7	16	25	58	71	68	17	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of pancreas (C25)													
All races, both sexes													
2007.....	34,117	-	-	2	5	52	538	2,808	6,507	8,671	10,317	5,217	-
2006.....	33,454	-	2	1	5	49	506	2,730	6,298	8,473	10,260	5,130	-
2005.....	32,760	-	-	2	3	64	565	2,744	5,984	8,288	10,147	4,962	1
2004.....	31,772	-	-	-	11	62	521	2,641	5,756	8,178	9,853	4,749	1
2003.....	30,777	-	-	1	10	60	548	2,540	5,320	8,104	9,708	4,486	-
2002.....	30,264	1	-	-	7	39	596	2,433	5,105	8,187	9,555	4,340	1
2001.....	29,803	-	1	2	2	55	574	2,410	4,868	8,158	9,322	4,410	1
2000.....	29,332	2	1	1	5	55	548	2,362	4,708	8,103	9,459	4,087	1
1999.....	29,082	1	-	-	3	55	594	2,353	4,707	8,293	8,933	4,142	1
All races, male													
2007.....	17,132	-	-	-	3	32	325	1,690	3,872	4,664	4,682	1,864	-
2006.....	16,559	-	1	-	3	21	308	1,661	3,735	4,537	4,569	1,724	-
2005.....	16,147	-	-	2	1	39	332	1,613	3,551	4,421	4,540	1,648	-
2004.....	15,776	-	-	-	6	39	323	1,647	3,377	4,430	4,416	1,538	-
2003.....	15,060	-	-	1	6	35	329	1,548	3,106	4,251	4,402	1,382	-
2002.....	14,877	-	-	-	5	26	364	1,526	3,012	4,270	4,270	1,403	1
2001.....	14,467	-	1	-	2	30	321	1,467	2,874	4,215	4,191	1,365	1
2000.....	14,238	1	1	-	3	33	324	1,486	2,749	4,189	4,189	1,262	1
1999.....	14,176	-	-	-	1	33	369	1,482	2,721	4,347	3,940	1,283	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of pancreas (C25)													
All races, female													
2007.....	16,985	-	-	2	2	20	213	1,118	2,635	4,007	5,635	3,353	-
2006.....	16,895	-	1	1	2	28	198	1,069	2,563	3,936	5,691	3,406	-
2005.....	16,613	-	-	-	2	25	233	1,131	2,433	3,867	5,607	3,314	1
2004.....	15,996	-	-	-	5	23	198	994	2,379	3,748	5,437	3,211	1
2003.....	15,717	-	-	-	4	25	219	992	2,214	3,853	5,306	3,104	-
2002.....	15,387	1	-	-	2	13	232	907	2,093	3,917	5,285	2,937	-
2001.....	15,336	-	-	2	-	25	253	943	1,994	3,943	5,131	3,045	-
2000.....	15,094	1	-	1	2	22	224	876	1,959	3,914	5,270	2,825	-
1999.....	14,906	1	-	-	2	22	225	871	1,986	3,946	4,993	2,859	1
White, both sexes													
2007 1/.....	29,096	-	-	1	4	35	425	2,204	5,411	7,335	9,054	4,627	-
2006 1/.....	28,677	-	-	1	5	38	392	2,170	5,224	7,191	9,081	4,575	-
2005 1/.....	28,092	-	-	2	1	47	451	2,163	4,980	7,069	8,942	4,436	1
2004 1/.....	27,229	-	-	-	6	45	402	2,091	4,748	6,945	8,738	4,253	1
2003 1/.....	26,418	-	-	-	7	48	421	1,986	4,399	6,936	8,601	4,020	-
2002.....	26,031	-	-	-	7	28	473	1,938	4,248	6,944	8,478	3,915	-
2001.....	25,769	-	1	2	2	42	461	1,945	4,107	6,987	8,274	3,947	1
2000.....	25,301	2	1	1	4	45	429	1,927	3,925	6,944	8,389	3,633	1
1999.....	25,086	-	-	-	2	40	468	1,913	3,897	7,140	7,913	3,712	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of pancreas (C25)													
White, male													
2007 1/.....	14,774	-	-	-	2	23	264	1,341	3,238	4,031	4,192	1,683	-
2006 1/.....	14,374	-	-	-	3	18	244	1,342	3,140	3,926	4,129	1,572	-
2005 1/.....	13,958	-	-	2	-	28	269	1,287	2,992	3,843	4,050	1,487	-
2004 1/.....	13,610	-	-	-	2	29	250	1,326	2,820	3,793	3,989	1,401	-
2003 1/.....	13,033	-	-	-	3	26	251	1,218	2,611	3,711	3,959	1,254	-
2002.....	12,921	-	-	-	5	19	291	1,236	2,544	3,693	3,856	1,277	-
2001.....	12,641	-	1	-	2	25	259	1,204	2,466	3,655	3,775	1,253	1
2000.....	12,392	1	1	-	3	27	258	1,224	2,317	3,658	3,767	1,135	1
1999.....	12,273	-	-	-	1	25	297	1,204	2,265	3,795	3,527	1,159	-
White, female													
2007 1/.....	14,322	-	-	1	2	12	161	863	2,173	3,304	4,862	2,944	-
2006 1/.....	14,303	-	-	1	2	20	148	828	2,084	3,265	4,952	3,003	-
2005 1/.....	14,134	-	-	-	1	19	182	876	1,988	3,226	4,892	2,949	1
2004 1/.....	13,619	-	-	-	4	16	152	765	1,928	3,152	4,749	2,852	1
2003 1/.....	13,385	-	-	-	4	22	170	768	1,788	3,225	4,642	2,766	-
2002.....	13,110	-	-	-	2	9	182	702	1,704	3,251	4,622	2,638	-
2001.....	13,128	-	-	2	-	17	202	741	1,641	3,332	4,499	2,694	-
2000.....	12,909	1	-	1	1	18	171	703	1,608	3,286	4,622	2,498	-
1999.....	12,813	-	-	-	1	15	171	709	1,632	3,345	4,386	2,553	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasm of pancreas (C25)

Black, both sexes

2007 1/.....	4,069	-	-	1	1	12	88	508	917	1,079	1,006	457	-
2006 1/.....	3,868	-	1	-	-	8	89	474	898	1,031	930	437	-
2005 1/.....	3,783	-	-	-	1	13	86	487	824	1,002	953	417	-
2004 1/.....	3,674	-	-	-	5	12	92	459	824	1,004	875	403	-
2003 1/.....	3,616	-	-	1	3	10	103	481	754	963	914	387	-
2002.....	3,494	1	-	-	-	6	102	403	730	1,025	867	359	1
2001.....	3,394	-	-	-	-	10	90	401	647	975	876	395	-
2000.....	3,419	-	-	-	1	9	97	382	672	978	897	383	-
1999.....	3,391	1	-	-	-	12	112	390	673	979	852	372	-

Black, male

2007 1/.....	1,893	-	-	-	1	6	50	293	542	504	371	126	-
2006 1/.....	1,768	-	-	-	-	2	52	269	509	496	338	102	-
2005 1/.....	1,756	-	-	-	1	8	44	275	464	475	381	108	-
2004 1/.....	1,737	-	-	-	4	7	55	259	467	522	322	101	-
2003 1/.....	1,675	-	-	1	3	8	64	294	409	449	347	100	-
2002.....	1,617	-	-	-	-	5	58	244	401	479	335	94	1
2001.....	1,523	-	-	-	-	4	47	230	343	470	341	88	-
2000.....	1,561	-	-	-	-	5	54	231	373	450	353	95	-
1999.....	1,611	-	-	-	-	8	63	246	387	462	347	98	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of pancreas (C25)													
Black, female													
2007 1/.....	2,176	-	-	1	-	6	38	215	375	575	635	331	-
2006 1/.....	2,100	-	1	-	-	6	37	205	389	535	592	335	-
2005 1/.....	2,027	-	-	-	-	5	42	212	360	527	572	309	-
2004 1/.....	1,937	-	-	-	1	5	37	200	357	482	553	302	-
2003 1/.....	1,941	-	-	-	-	2	39	187	345	514	567	287	-
2002.....	1,877	1	-	-	-	1	44	159	329	546	532	265	-
2001.....	1,871	-	-	-	-	6	43	171	304	505	535	307	-
2000.....	1,858	-	-	-	1	4	43	151	299	528	544	288	-
1999.....	1,780	1	-	-	-	4	49	144	286	517	505	274	-
American Indian 2/, both sexes													
2007 1/.....	143	-	-	-	-	-	6	20	35	36	32	14	-
2006 1/.....	138	-	-	-	-	-	5	8	34	38	38	15	-
2005 1/.....	133	-	-	-	-	2	2	18	28	34	41	8	-
2004 1/.....	139	-	-	-	-	-	4	23	41	42	22	7	-
2003 1/.....	103	-	-	-	-	1	7	11	29	30	19	6	-
2002.....	112	-	-	-	-	1	4	20	22	34	25	6	-
2001.....	89	-	-	-	-	1	3	11	15	32	18	9	-
2000.....	91	-	-	-	-	-	8	9	25	24	18	7	-
1999.....	84	-	-	-	-	1	2	9	19	21	22	10	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Malignant neoplasm of pancreas (C25)

American Indian 2/, male

2007 1/.....	75	-	-	-	-	-	4	11	19	19	15	7	-
2006 1/.....	55	-	-	-	-	-	3	5	18	13	10	6	-
2005 1/.....	73	-	-	-	-	2	1	8	20	21	17	4	-
2004 1/.....	71	-	-	-	-	-	2	17	19	19	10	4	-
2003 1/.....	51	-	-	-	-	-	3	6	17	15	8	2	-
2002.....	50	-	-	-	-	-	4	11	8	16	8	3	-
2001.....	38	-	-	-	-	-	1	4	8	17	6	2	-
2000.....	46	-	-	-	-	-	5	5	16	12	6	2	-
1999.....	40	-	-	-	-	-	1	5	9	14	9	2	-

American Indian 2/, female

2007 1/.....	68	-	-	-	-	-	2	9	16	17	17	7	-
2006 1/.....	83	-	-	-	-	-	2	3	16	25	28	9	-
2005 1/.....	60	-	-	-	-	-	1	10	8	13	24	4	-
2004 1/.....	68	-	-	-	-	-	2	6	22	23	12	3	-
2003 1/.....	52	-	-	-	-	1	4	5	12	15	11	4	-
2002.....	62	-	-	-	-	1	-	9	14	18	17	3	-
2001.....	51	-	-	-	-	1	2	7	7	15	12	7	-
2000.....	45	-	-	-	-	-	3	4	9	12	12	5	-
1999.....	44	-	-	-	-	1	1	4	10	7	13	8	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Malignant neoplasm of pancreas (C25)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	809	-	-	-	-	5	19	76	144	221	225	119	-
2006 1/.....	771	-	1	-	-	3	20	78	142	213	211	103	-
2005 1/.....	752	-	-	-	1	2	26	76	152	183	211	101	-
2004 1/.....	730	-	-	-	-	5	23	68	143	187	218	86	-
2003 1/.....	640	-	-	-	-	1	17	62	138	175	174	73	-
2002.....	627	-	-	-	-	4	17	72	105	184	185	60	-
2001.....	551	-	-	-	-	2	20	53	99	164	154	59	-
2000.....	521	-	-	-	-	1	14	44	86	157	155	64	-
1999.....	521	-	-	-	1	2	12	41	118	153	146	48	-

Asian or Pacific Islander 3/, male

2007 1/.....	390	-	-	-	-	3	7	45	73	110	104	48	-
2006 1/.....	362	-	1	-	-	1	9	45	68	102	92	44	-
2005 1/.....	360	-	-	-	-	1	18	43	75	82	92	49	-
2004 1/.....	358	-	-	-	-	3	16	45	71	96	95	32	-
2003 1/.....	301	-	-	-	-	1	11	30	69	76	88	26	-
2002.....	289	-	-	-	-	2	11	35	59	82	71	29	-
2001.....	265	-	-	-	-	1	14	29	57	73	69	22	-
2000.....	239	-	-	-	-	1	7	26	43	69	63	30	-
1999.....	252	-	-	-	-	-	8	27	60	76	57	24	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasm of pancreas (C25)

Asian or Pacific Islander 3/, female

2007 1/.....	419	-	-	-	-	2	12	31	71	111	121	71	-
2006 1/.....	409	-	-	-	-	2	11	33	74	111	119	59	-
2005 1/.....	392	-	-	-	1	1	8	33	77	101	119	52	-
2004 1/.....	372	-	-	-	-	2	7	23	72	91	123	54	-
2003 1/.....	339	-	-	-	-	-	6	32	69	99	86	47	-
2002.....	338	-	-	-	-	2	6	37	46	102	114	31	-
2001.....	286	-	-	-	-	1	6	24	42	91	85	37	-
2000.....	282	-	-	-	-	-	7	18	43	88	92	34	-
1999.....	269	-	-	-	1	2	4	14	58	77	89	24	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of larynx (C32)													
All races, both sexes													
2007.....	3,634	-	-	-	2	2	55	450	932	989	871	333	-
2006.....	3,821	-	-	-	-	2	58	451	978	1,071	928	333	-
2005.....	3,797	-	-	-	-	3	64	423	916	1,156	914	320	1
2004.....	3,668	-	-	1	1	3	59	443	936	1,051	858	316	-
2003.....	3,792	-	-	-	-	1	79	420	956	1,136	891	308	1
2002.....	3,723	-	-	-	-	11	76	425	904	1,084	969	253	1
2001.....	3,797	-	-	-	-	3	86	419	913	1,170	910	296	-
2000.....	3,861	-	-	2	1	4	64	426	873	1,219	996	276	-
1999.....	3,815	-	-	-	-	6	78	405	904	1,239	892	291	-
All races, male													
2007.....	2,890	-	-	-	1	1	40	371	779	765	686	247	-
2006.....	2,991	-	-	-	-	1	40	364	781	846	709	250	-
2005.....	2,983	-	-	-	-	2	47	320	752	916	693	252	1
2004.....	2,935	-	-	-	1	3	48	362	756	854	676	235	-
2003.....	3,020	-	-	-	-	1	60	354	777	899	708	221	-
2002.....	2,922	-	-	-	-	6	58	349	738	832	755	183	1
2001.....	3,018	-	-	-	-	2	66	346	737	919	710	238	-
2000.....	3,041	-	-	-	-	2	49	340	726	945	765	214	-
1999.....	2,992	-	-	-	-	5	64	346	735	948	681	213	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of larynx (C32)													
All races, female													
2007.....	744	-	-	-	1	1	15	79	153	224	185	86	-
2006.....	830	-	-	-	-	1	18	87	197	225	219	83	-
2005.....	814	-	-	-	-	1	17	103	164	240	221	68	-
2004.....	733	-	-	1	-	-	11	81	180	197	182	81	-
2003.....	772	-	-	-	-	-	19	66	179	237	183	87	1
2002.....	801	-	-	-	-	5	18	76	166	252	214	70	-
2001.....	779	-	-	-	-	1	20	73	176	251	200	58	-
2000.....	820	-	-	2	1	2	15	86	147	274	231	62	-
1999.....	823	-	-	-	-	1	14	59	169	291	211	78	-
White, both sexes													
2007 1/.....	2,905	-	-	-	2	2	42	329	708	792	739	291	-
2006 1/.....	3,123	-	-	-	-	1	40	329	777	870	811	295	-
2005 1/.....	3,060	-	-	-	-	2	49	308	684	934	807	275	1
2004 1/.....	2,985	-	-	1	1	3	42	324	735	861	740	278	-
2003 1/.....	3,052	-	-	-	-	1	62	292	744	923	764	265	1
2002.....	3,012	-	-	-	-	7	48	296	700	901	827	233	-
2001.....	3,049	-	-	-	-	3	61	288	700	954	780	263	-
2000.....	3,096	-	-	2	1	4	49	299	643	987	859	252	-
1999.....	3,106	-	-	-	-	4	45	284	691	1,035	794	253	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of larynx (C32)													
White, male													
2007 1/.....	2,283	-	-	-	1	1	30	265	588	608	577	213	-
2006 1/.....	2,420	-	-	-	-	-	26	260	620	676	615	223	-
2005 1/.....	2,388	-	-	-	-	1	36	236	553	736	609	216	1
2004 1/.....	2,370	-	-	-	1	3	36	262	585	699	580	204	-
2003 1/.....	2,410	-	-	-	-	1	49	248	605	718	599	190	-
2002.....	2,351	-	-	-	-	3	37	247	561	689	647	167	-
2001.....	2,417	-	-	-	-	2	50	234	567	751	598	215	-
2000.....	2,426	-	-	-	-	2	35	237	530	763	664	195	-
1999.....	2,419	-	-	-	-	3	38	245	562	789	601	181	-
White, female													
2007 1/.....	622	-	-	-	1	1	12	64	120	184	162	78	-
2006 1/.....	703	-	-	-	-	1	14	69	157	194	196	72	-
2005 1/.....	672	-	-	-	-	1	13	72	131	198	198	59	-
2004 1/.....	615	-	-	1	-	-	6	62	150	162	160	74	-
2003 1/.....	642	-	-	-	-	-	13	44	139	205	165	75	1
2002.....	661	-	-	-	-	4	11	49	139	212	180	66	-
2001.....	632	-	-	-	-	1	11	54	133	203	182	48	-
2000.....	670	-	-	2	1	2	14	62	113	224	195	57	-
1999.....	687	-	-	-	-	1	7	39	129	246	193	72	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of larynx (C32)													
Black, both sexes													
2007 1/.....	659	-	-	-	-	-	13	113	204	179	117	33	-
2006 1/.....	635	-	-	-	-	1	16	111	189	186	101	31	-
2005 1/.....	680	-	-	-	-	1	13	111	214	203	100	38	-
2004 1/.....	638	-	-	-	-	-	17	112	190	179	111	29	-
2003 1/.....	693	-	-	-	-	-	16	121	206	197	115	38	-
2002.....	675	-	-	-	-	4	27	123	198	174	132	16	1
2001.....	678	-	-	-	-	-	24	125	200	191	111	27	-
2000.....	719	-	-	-	-	-	15	123	219	211	129	22	-
1999.....	663	-	-	-	-	2	32	117	205	190	91	26	-
Black, male													
2007 1/.....	547	-	-	-	-	-	10	98	177	139	96	27	-
2006 1/.....	515	-	-	-	-	1	12	93	152	155	81	21	-
2005 1/.....	548	-	-	-	-	1	9	80	186	163	78	31	-
2004 1/.....	530	-	-	-	-	-	12	95	163	147	89	24	-
2003 1/.....	571	-	-	-	-	-	10	100	167	169	99	26	-
2002.....	542	-	-	-	-	3	20	96	172	135	102	13	1
2001.....	541	-	-	-	-	-	15	106	161	147	94	18	-
2000.....	578	-	-	-	-	-	14	100	187	166	94	17	-
1999.....	536	-	-	-	-	2	25	97	169	148	74	21	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of larynx (C32)													
Black, female													
2007 1/.....	112	-	-	-	-	-	3	15	27	40	21	6	-
2006 1/.....	120	-	-	-	-	-	4	18	37	31	20	10	-
2005 1/.....	132	-	-	-	-	-	4	31	28	40	22	7	-
2004 1/.....	108	-	-	-	-	-	5	17	27	32	22	5	-
2003 1/.....	122	-	-	-	-	-	6	21	39	28	16	12	-
2002.....	133	-	-	-	-	1	7	27	26	39	30	3	-
2001.....	137	-	-	-	-	-	9	19	39	44	17	9	-
2000.....	141	-	-	-	-	-	1	23	32	45	35	5	-
1999.....	127	-	-	-	-	-	7	20	36	42	17	5	-
American Indian 2/, both sexes													
2007 1/.....	21	-	-	-	-	-	-	-	10	7	2	2	-
2006 1/.....	18	-	-	-	-	-	1	5	3	2	4	3	-
2005 1/.....	21	-	-	-	-	-	-	2	9	7	1	2	-
2004 1/.....	15	-	-	-	-	-	-	2	4	7	2	-	-
2003 1/.....	14	-	-	-	-	-	-	5	2	3	3	1	-
2002.....	14	-	-	-	-	-	-	3	3	4	4	-	-
2001.....	23	-	-	-	-	-	1	2	3	10	5	2	-
2000.....	16	-	-	-	-	-	-	1	6	8	1	-	-
1999.....	12	-	-	-	-	-	-	2	3	4	-	3	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Malignant neoplasm of larynx (C32)

American Indian 2/, male

2007 1/.....	18	-	-	-	-	-	-	-	8	7	1	2	-
2006 1/.....	15	-	-	-	-	-	1	5	3	2	2	2	-
2005 1/.....	17	-	-	-	-	-	-	2	7	6	1	1	-
2004 1/.....	10	-	-	-	-	-	-	1	2	5	2	-	-
2003 1/.....	11	-	-	-	-	-	-	4	2	1	3	1	-
2002.....	11	-	-	-	-	-	-	3	3	4	1	-	-
2001.....	18	-	-	-	-	-	1	2	2	7	4	2	-
2000.....	11	-	-	-	-	-	-	1	4	5	1	-	-
1999.....	10	-	-	-	-	-	-	2	2	3	-	3	-

American Indian 2/, female

2007 1/.....	3	-	-	-	-	-	-	-	2	-	1	-	-
2006 1/.....	3	-	-	-	-	-	-	-	-	-	2	1	-
2005 1/.....	4	-	-	-	-	-	-	-	2	1	-	1	-
2004 1/.....	5	-	-	-	-	-	-	1	2	2	-	-	-
2003 1/.....	3	-	-	-	-	-	-	1	-	2	-	-	-
2002.....	3	-	-	-	-	-	-	-	-	-	3	-	-
2001.....	5	-	-	-	-	-	-	-	1	3	1	-	-
2000.....	5	-	-	-	-	-	-	-	2	3	-	-	-
1999.....	2	-	-	-	-	-	-	-	1	1	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Malignant neoplasm of larynx (C32)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	49	-	-	-	-	-	-	8	10	11	13	7	-
2006 1/.....	45	-	-	-	-	-	1	6	9	13	12	4	-
2005 1/.....	36	-	-	-	-	2	-	2	9	12	6	5	-
2004 1/.....	30	-	-	-	-	-	-	5	7	4	5	9	-
2003 1/.....	33	-	-	-	-	-	1	2	4	13	9	4	-
2002.....	22	-	-	-	-	-	1	3	3	5	6	4	-
2001.....	47	-	-	-	-	-	-	4	10	15	14	4	-
2000.....	30	-	-	-	-	-	-	3	5	13	7	2	-
1999.....	34	-	-	-	-	-	1	2	5	10	7	9	-

Asian or Pacific Islander 3/, male

2007 1/.....	42	-	-	-	-	-	-	8	6	11	12	5	-
2006 1/.....	41	-	-	-	-	-	1	6	6	13	11	4	-
2005 1/.....	30	-	-	-	-	-	2	2	6	11	5	4	-
2004 1/.....	25	-	-	-	-	-	-	4	6	3	5	7	-
2003 1/.....	28	-	-	-	-	-	1	2	3	11	7	4	-
2002.....	18	-	-	-	-	-	1	3	2	4	5	3	-
2001.....	42	-	-	-	-	-	-	4	7	14	14	3	-
2000.....	26	-	-	-	-	-	-	2	5	11	6	2	-
1999.....	27	-	-	-	-	-	1	2	2	8	6	8	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasm of larynx (C32)

Asian or Pacific Islander 3/, female

2007 1/.....	7	-	-	-	-	-	-	-	4	-	1	2	-
2006 1/.....	4	-	-	-	-	-	-	-	3	-	1	-	-
2005 1/.....	6	-	-	-	-	-	-	-	3	1	1	1	-
2004 1/.....	5	-	-	-	-	-	-	1	1	1	-	2	-
2003 1/.....	5	-	-	-	-	-	-	-	1	2	2	-	-
2002.....	4	-	-	-	-	-	-	-	1	1	1	1	-
2001.....	5	-	-	-	-	-	-	-	3	1	-	1	-
2000.....	4	-	-	-	-	-	-	1	-	2	1	-	-
1999.....	7	-	-	-	-	-	-	-	3	2	1	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of trachea, bronchus and lung (C33-C34)													
All races, both sexes													
2007.....	158,760	1	1	4	25	135	1,852	12,480	31,216	48,157	48,358	16,528	3
2006.....	158,664	1	1	2	27	155	2,023	12,608	31,317	47,882	48,732	15,915	1
2005.....	159,292	-	3	1	16	133	2,323	12,624	31,363	48,390	49,032	15,404	3
2004.....	158,091	2	4	4	23	147	2,426	12,448	31,086	49,029	48,464	14,455	3
2003.....	158,086	-	2	2	25	154	2,478	12,374	30,956	49,386	48,619	14,088	2
2002.....	157,713	-	1	4	21	165	2,698	12,150	30,684	50,256	48,082	13,652	-
2001.....	156,058	-	5	2	20	157	2,796	12,041	29,777	51,225	46,700	13,331	4
2000.....	155,521	2	2	1	22	197	2,756	11,915	29,721	52,258	45,836	12,806	5
1999.....	152,156	-	-	5	20	179	2,724	11,434	29,344	51,848	44,261	12,339	2
All races, male													
2007.....	88,372	1	1	4	17	79	911	6,939	18,326	27,568	26,475	8,049	2
2006.....	89,279	1	1	1	10	85	1,034	7,196	18,514	27,695	27,015	7,727	-
2005.....	90,187	-	3	-	9	77	1,197	7,327	18,663	28,207	27,196	7,506	2
2004.....	89,630	1	1	3	14	87	1,280	7,317	18,440	28,453	26,992	7,039	3
2003.....	89,964	-	1	1	14	78	1,344	7,313	18,347	28,935	27,064	6,865	2
2002.....	90,171	-	1	3	16	90	1,386	7,194	18,415	29,543	26,821	6,702	-
2001.....	90,426	-	4	1	12	77	1,504	7,190	17,873	30,699	26,406	6,656	4
2000.....	90,469	2	1	-	14	102	1,542	7,123	17,937	31,382	25,966	6,395	5
1999.....	89,453	-	-	4	13	97	1,474	6,851	18,013	31,287	25,397	6,315	2

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

All races, female

2007.....	70,388	-	-	-	8	56	941	5,541	12,890	20,589	21,883	8,479	1
2006.....	69,385	-	-	1	17	70	989	5,412	12,803	20,187	21,717	8,188	1
2005.....	69,105	-	-	1	7	56	1,126	5,297	12,700	20,183	21,836	7,898	1
2004.....	68,461	1	3	1	9	60	1,146	5,131	12,646	20,576	21,472	7,416	-
2003.....	68,122	-	1	1	11	76	1,134	5,061	12,609	20,451	21,555	7,223	-
2002.....	67,542	-	-	1	5	75	1,312	4,956	12,269	20,713	21,261	6,950	-
2001.....	65,632	-	1	1	8	80	1,292	4,851	11,904	20,526	20,294	6,675	-
2000.....	65,052	-	1	1	8	95	1,214	4,792	11,784	20,876	19,870	6,411	-
1999.....	62,703	-	-	1	7	82	1,250	4,583	11,331	20,561	18,864	6,024	-

White, both sexes

2007 1/.....	138,730	-	1	2	20	111	1,484	9,895	26,366	42,163	43,671	15,016	1
2006 1/.....	138,814	-	1	-	18	112	1,593	9,935	26,499	41,987	44,102	14,566	1
2005 1/.....	139,442	-	3	1	14	98	1,851	10,046	26,469	42,527	44,428	14,003	2
2004 1/.....	138,425	1	3	4	18	109	1,900	9,888	26,364	43,136	43,770	13,230	2
2003 1/.....	138,743	-	2	1	22	127	1,950	9,786	26,435	43,558	43,988	12,872	2
2002.....	138,661	-	1	3	17	127	2,120	9,714	26,055	44,453	43,678	12,493	-
2001.....	137,274	-	4	2	16	130	2,214	9,674	25,506	45,259	42,297	12,170	2
2000.....	136,912	1	2	1	14	155	2,093	9,537	25,487	46,208	41,726	11,685	3
1999.....	133,752	-	-	3	11	149	2,128	9,147	25,090	45,840	40,131	11,252	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

White, male

2007 1/.....	76,502	-	1	2	13	61	726	5,418	15,264	23,937	23,838	7,241	1
2006 1/.....	77,476	-	1	-	6	61	816	5,625	15,443	24,082	24,388	7,054	-
2005 1/.....	78,379	-	3	-	8	59	943	5,814	15,517	24,638	24,606	6,790	1
2004 1/.....	77,730	1	-	3	11	64	988	5,706	15,394	24,873	24,261	6,427	2
2003 1/.....	78,215	-	1	-	12	62	1,041	5,724	15,426	25,364	24,330	6,253	2
2002.....	78,465	-	1	2	13	72	1,063	5,656	15,391	25,974	24,174	6,119	-
2001.....	78,665	-	3	1	10	65	1,165	5,660	15,095	26,909	23,711	6,044	2
2000.....	78,747	1	1	-	10	82	1,136	5,594	15,179	27,534	23,415	5,792	3
1999.....	77,846	-	-	2	7	81	1,138	5,365	15,201	27,450	22,864	5,737	1

White, female

2007 1/.....	62,228	-	-	-	7	50	758	4,477	11,102	18,226	19,833	7,775	-
2006 1/.....	61,338	-	-	-	12	51	777	4,310	11,056	17,905	19,714	7,512	1
2005 1/.....	61,063	-	-	1	6	39	908	4,232	10,952	17,889	19,822	7,213	1
2004 1/.....	60,695	-	3	1	7	45	912	4,182	10,970	18,263	19,509	6,803	-
2003 1/.....	60,528	-	1	1	10	65	909	4,062	11,009	18,194	19,658	6,619	-
2002.....	60,196	-	-	1	4	55	1,057	4,058	10,664	18,479	19,504	6,374	-
2001.....	58,609	-	1	1	6	65	1,049	4,014	10,411	18,350	18,586	6,126	-
2000.....	58,165	-	1	1	4	73	957	3,943	10,308	18,674	18,311	5,893	-
1999.....	55,906	-	-	1	4	68	990	3,782	9,889	18,390	17,267	5,515	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

Black, both sexes

2007 1/.....	16,494	1	-	2	4	18	290	2,252	4,139	4,973	3,679	1,134	2
2006 1/.....	16,493	1	-	1	8	32	347	2,334	4,123	4,910	3,717	1,020	-
2005 1/.....	16,567	-	-	-	1	19	388	2,250	4,211	4,912	3,703	1,082	1
2004 1/.....	16,472	1	1	-	4	26	444	2,236	4,106	4,845	3,824	984	1
2003 1/.....	16,372	-	-	1	3	19	442	2,272	3,979	4,932	3,761	963	-
2002.....	16,316	-	-	1	4	23	505	2,180	4,026	4,983	3,677	917	-
2001.....	16,019	-	1	-	3	16	519	2,079	3,726	5,117	3,642	914	2
2000.....	16,066	1	-	-	5	32	584	2,132	3,726	5,226	3,438	921	1
1999.....	15,963	-	-	2	8	20	533	2,056	3,770	5,148	3,515	911	-

Black, male

2007 1/.....	9,839	1	-	2	3	14	146	1,326	2,672	3,037	2,043	594	1
2006 1/.....	9,823	1	-	1	4	19	176	1,365	2,650	3,028	2,084	495	-
2005 1/.....	9,887	-	-	-	1	11	204	1,334	2,721	3,013	2,069	533	1
2004 1/.....	9,995	-	1	-	3	14	236	1,423	2,675	2,962	2,195	485	1
2003 1/.....	9,988	-	-	1	2	13	252	1,411	2,586	3,035	2,206	482	-
2002.....	10,075	-	-	1	3	14	289	1,387	2,631	3,115	2,183	452	-
2001.....	10,131	-	1	-	1	7	301	1,369	2,434	3,290	2,242	484	2
2000.....	10,153	1	-	-	2	15	362	1,384	2,437	3,341	2,122	488	1
1999.....	10,135	-	-	2	5	11	303	1,359	2,521	3,294	2,150	490	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of trachea, bronchus and lung (C33-C34)													
Black, female													
2007 1/.....	6,655	-	-	-	1	4	144	926	1,467	1,936	1,636	540	1
2006 1/.....	6,670	-	-	-	4	13	171	969	1,473	1,882	1,633	525	-
2005 1/.....	6,680	-	-	-	-	8	184	916	1,490	1,899	1,634	549	-
2004 1/.....	6,477	1	-	-	1	12	208	813	1,431	1,883	1,629	499	-
2003 1/.....	6,384	-	-	-	1	6	190	861	1,393	1,897	1,555	481	-
2002.....	6,241	-	-	-	1	9	216	793	1,395	1,868	1,494	465	-
2001.....	5,888	-	-	-	2	9	218	710	1,292	1,827	1,400	430	-
2000.....	5,913	-	-	-	3	17	222	748	1,289	1,885	1,316	433	-
1999.....	5,828	-	-	-	3	9	230	697	1,249	1,854	1,365	421	-
American Indian 2/, both sexes													
2007 1/.....	695	-	-	-	-	-	11	54	184	246	157	43	-
2006 1/.....	635	-	-	-	1	-	10	63	162	223	145	31	-
2005 1/.....	674	-	-	-	-	2	20	78	172	221	149	32	-
2004 1/.....	677	-	-	-	-	1	15	65	167	253	143	33	-
2003 1/.....	556	-	-	-	-	-	11	62	134	191	130	28	-
2002.....	556	-	-	-	-	1	5	49	151	191	124	35	-
2001.....	547	-	-	-	-	1	8	63	133	181	134	27	-
2000.....	480	-	-	-	-	1	11	45	126	165	106	26	-
1999.....	497	-	-	-	-	-	8	40	142	183	103	21	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

American Indian 2/, male

2007 1/.....	384	-	-	-	-	-	4	30	109	129	87	25	-
2006 1/.....	339	-	-	-	-	-	5	37	81	122	81	13	-
2005 1/.....	357	-	-	-	-	-	14	39	99	110	80	15	-
2004 1/.....	373	-	-	-	-	1	9	38	100	132	80	13	-
2003 1/.....	301	-	-	-	-	-	7	32	85	99	67	11	-
2002.....	304	-	-	-	-	-	2	27	94	94	72	15	-
2001.....	299	-	-	-	-	-	8	35	75	100	73	8	-
2000.....	270	-	-	-	-	1	5	24	76	83	68	13	-
1999.....	285	-	-	-	-	-	3	25	79	109	55	14	-

American Indian 2/, female

2007 1/.....	311	-	-	-	-	-	7	24	75	117	70	18	-
2006 1/.....	296	-	-	-	1	-	5	26	81	101	64	18	-
2005 1/.....	317	-	-	-	-	2	6	39	73	111	69	17	-
2004 1/.....	304	-	-	-	-	-	6	27	67	121	63	20	-
2003 1/.....	255	-	-	-	-	-	4	30	49	92	63	17	-
2002.....	252	-	-	-	-	1	3	22	57	97	52	20	-
2001.....	248	-	-	-	-	1	-	28	58	81	61	19	-
2000.....	210	-	-	-	-	-	6	21	50	82	38	13	-
1999.....	212	-	-	-	-	-	5	15	63	74	48	7	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	2,841	-	-	-	1	6	67	279	527	775	851	335	-
2006 1/.....	2,722	-	-	1	-	11	73	276	533	762	768	298	-
2005 1/.....	2,609	-	-	-	1	14	64	250	511	730	752	287	-
2004 1/.....	2,517	-	-	-	1	11	67	259	449	795	727	208	-
2003 1/.....	2,415	-	-	-	-	8	75	254	408	705	740	225	-
2002.....	2,180	-	-	-	-	14	68	207	452	629	603	207	-
2001.....	2,218	-	-	-	1	10	55	225	412	668	627	220	-
2000.....	2,063	-	-	-	3	9	68	201	382	659	566	174	1
1999.....	1,944	-	-	-	1	10	55	191	342	677	512	155	1

Asian or Pacific Islander 3/, male

2007 1/.....	1,647	-	-	-	1	4	35	165	281	465	507	189	-
2006 1/.....	1,641	-	-	-	-	5	37	169	340	463	462	165	-
2005 1/.....	1,564	-	-	-	-	7	36	140	326	446	441	168	-
2004 1/.....	1,532	-	-	-	-	8	47	150	271	486	456	114	-
2003 1/.....	1,460	-	-	-	-	3	44	146	250	437	461	119	-
2002.....	1,327	-	-	-	-	4	32	124	299	360	392	116	-
2001.....	1,331	-	-	-	1	5	30	126	269	400	380	120	-
2000.....	1,299	-	-	-	2	4	39	121	245	424	361	102	1
1999.....	1,187	-	-	-	1	5	30	102	212	434	328	74	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

Asian or Pacific Islander 3/, female

2007 1/.....	1,194	-	-	-	-	2	32	114	246	310	344	146	-
2006 1/.....	1,081	-	-	1	-	6	36	107	193	299	306	133	-
2005 1/.....	1,045	-	-	-	1	7	28	110	185	284	311	119	-
2004 1/.....	985	-	-	-	1	3	20	109	178	309	271	94	-
2003 1/.....	955	-	-	-	-	5	31	108	158	268	279	106	-
2002.....	853	-	-	-	-	10	36	83	153	269	211	91	-
2001.....	887	-	-	-	-	5	25	99	143	268	247	100	-
2000.....	764	-	-	-	1	5	29	80	137	235	205	72	-
1999.....	757	-	-	-	-	5	25	89	130	243	184	81	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant melanoma of skin (C43)													
All races, both sexes													
2007.....	8,461	-	1	2	31	175	466	1,174	1,742	1,737	2,035	1,098	-
2006.....	8,441	-	2	2	47	187	505	1,168	1,642	1,780	2,041	1,067	-
2005.....	8,345	-	1	3	40	216	524	1,197	1,575	1,748	2,031	1,010	-
2004.....	7,952	-	2	-	39	194	526	1,162	1,547	1,711	1,912	859	-
2003.....	7,818	-	1	2	38	191	561	1,179	1,547	1,593	1,875	831	-
2002.....	7,514	-	3	1	41	194	604	1,135	1,351	1,616	1,748	820	1
2001.....	7,542	-	-	-	41	186	622	1,208	1,333	1,689	1,678	785	-
2000.....	7,420	-	-	1	41	186	626	1,199	1,303	1,637	1,624	803	-
1999.....	7,215	1	-	1	39	194	625	1,120	1,263	1,652	1,639	681	-
All races, male													
2007.....	5,506	-	-	-	20	95	266	759	1,214	1,198	1,318	636	-
2006.....	5,477	-	2	1	28	111	308	745	1,122	1,186	1,355	619	-
2005.....	5,283	-	1	-	23	138	313	748	1,030	1,177	1,294	559	-
2004.....	5,125	-	2	-	17	117	312	749	1,058	1,158	1,239	473	-
2003.....	4,963	-	-	-	24	119	336	753	1,042	1,050	1,203	436	-
2002.....	4,725	-	2	1	20	119	363	746	915	1,075	1,078	406	-
2001.....	4,763	-	-	-	24	103	380	756	893	1,156	1,063	388	-
2000.....	4,592	-	-	1	21	110	355	769	866	1,067	1,013	390	-
1999.....	4,529	1	-	-	23	110	397	717	851	1,096	1,006	328	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant melanoma of skin (C43)													
All races, female													
2007.....	2,955	-	1	2	11	80	200	415	528	539	717	462	-
2006.....	2,964	-	-	1	19	76	197	423	520	594	686	448	-
2005.....	3,062	-	-	3	17	78	211	449	545	571	737	451	-
2004.....	2,827	-	-	-	22	77	214	413	489	553	673	386	-
2003.....	2,855	-	1	2	14	72	225	426	505	543	672	395	-
2002.....	2,789	-	1	-	21	75	241	389	436	541	670	414	1
2001.....	2,779	-	-	-	17	83	242	452	440	533	615	397	-
2000.....	2,828	-	-	-	20	76	271	430	437	570	611	413	-
1999.....	2,686	-	-	1	16	84	228	403	412	556	633	353	-
White, both sexes													
2007 1/.....	8,258	-	1	2	30	171	451	1,145	1,703	1,699	1,987	1,069	-
2006 1/.....	8,250	-	2	-	45	183	486	1,148	1,605	1,751	1,995	1,035	-
2005 1/.....	8,146	-	1	2	35	211	506	1,170	1,531	1,716	1,992	982	-
2004 1/.....	7,801	-	1	-	36	191	516	1,145	1,525	1,676	1,871	840	-
2003 1/.....	7,648	-	1	2	36	186	549	1,151	1,513	1,564	1,834	812	-
2002.....	7,353	-	3	1	40	192	590	1,113	1,319	1,574	1,721	799	1
2001.....	7,403	-	-	-	40	184	615	1,192	1,312	1,644	1,645	771	-
2000.....	7,245	-	-	1	40	180	606	1,179	1,273	1,598	1,588	780	-
1999.....	7,086	1	-	1	36	188	618	1,107	1,244	1,623	1,604	664	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant melanoma of skin (C43)													
White, male													
2007 1/.....	5,408	-	-	-	20	92	261	741	1,191	1,181	1,300	622	-
2006 1/.....	5,381	-	2	-	27	108	300	735	1,099	1,174	1,332	604	-
2005 1/.....	5,184	-	1	-	20	133	299	734	1,005	1,166	1,276	550	-
2004 1/.....	5,053	-	1	-	16	115	306	740	1,045	1,146	1,218	466	-
2003 1/.....	4,887	-	-	-	24	117	332	739	1,024	1,038	1,183	430	-
2002.....	4,639	-	2	1	19	117	357	732	895	1,051	1,066	399	-
2001.....	4,702	-	-	-	23	101	376	749	883	1,135	1,050	385	-
2000.....	4,514	-	-	1	21	108	345	760	851	1,047	996	385	-
1999.....	4,459	1	-	-	20	106	394	708	838	1,083	988	321	-
White, female													
2007 1/.....	2,850	-	1	2	10	79	190	404	512	518	687	447	-
2006 1/.....	2,869	-	-	-	18	75	186	413	506	577	663	431	-
2005 1/.....	2,962	-	-	2	15	78	207	436	526	550	716	432	-
2004 1/.....	2,748	-	-	-	20	76	210	405	480	530	653	374	-
2003 1/.....	2,761	-	1	2	12	69	217	412	489	526	651	382	-
2002.....	2,714	-	1	-	21	75	233	381	424	523	655	400	1
2001.....	2,701	-	-	-	17	83	239	443	429	509	595	386	-
2000.....	2,731	-	-	-	19	72	261	419	422	551	592	395	-
1999.....	2,627	-	-	1	16	82	224	399	406	540	616	343	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant melanoma of skin (C43)													
Black, both sexes													
2007 1/.....	136	-	-	-	1	2	11	18	26	22	33	23	-
2006 1/.....	126	-	-	1	1	3	11	12	26	17	34	21	-
2005 1/.....	124	-	-	1	2	3	10	13	26	19	30	20	-
2004 1/.....	119	-	1	-	3	2	8	12	16	27	34	16	-
2003 1/.....	121	-	-	-	1	3	11	16	23	26	28	13	-
2002.....	102	-	-	-	1	1	9	12	21	25	18	15	-
2001.....	104	-	-	-	1	1	7	9	13	37	25	11	-
2000.....	135	-	-	-	1	3	15	13	20	30	32	21	-
1999.....	96	-	-	-	1	4	4	9	14	25	25	14	-
Black, male													
2007 1/.....	59	-	-	-	-	2	4	10	14	7	12	10	-
2006 1/.....	66	-	-	1	1	2	5	7	17	6	17	10	-
2005 1/.....	57	-	-	-	1	3	8	7	15	5	13	5	-
2004 1/.....	57	-	1	-	1	1	5	6	10	9	18	6	-
2003 1/.....	50	-	-	-	-	2	4	8	12	12	10	2	-
2002.....	52	-	-	-	1	1	4	7	13	12	8	6	-
2001.....	37	-	-	-	1	1	4	3	4	17	6	1	-
2000.....	60	-	-	-	-	2	9	6	11	14	15	3	-
1999.....	48	-	-	-	1	2	1	6	10	10	12	6	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Malignant melanoma of skin (C43)

Black, female

2007 1/.....	77	-	-	-	1	-	7	8	12	15	21	13	-
2006 1/.....	60	-	-	-	-	1	6	5	9	11	17	11	-
2005 1/.....	67	-	-	1	-	-	2	6	11	14	17	15	-
2004 1/.....	62	-	-	-	2	1	3	6	6	18	16	10	-
2003 1/.....	71	-	-	-	1	1	7	8	11	14	18	11	-
2002.....	50	-	-	-	-	-	5	5	8	13	10	9	-
2001.....	67	-	-	-	-	-	3	6	9	20	19	10	-
2000.....	75	-	-	-	1	1	6	7	9	16	17	18	-
1999.....	48	-	-	-	-	2	3	3	4	15	13	8	-

American Indian 2/, both sexes

2007 1/.....	22	-	-	-	-	2	-	4	5	4	5	2	-
2006 1/.....	21	-	-	-	-	-	3	2	4	3	4	5	-
2005 1/.....	20	-	-	-	2	-	5	2	7	2	1	1	-
2004 1/.....	11	-	-	-	-	-	1	3	2	4	1	-	-
2003 1/.....	14	-	-	-	1	-	-	2	4	2	5	-	-
2002.....	17	-	-	-	-	-	-	2	4	6	2	3	-
2001.....	10	-	-	-	-	1	-	1	3	-	3	2	-
2000.....	8	-	-	-	-	-	-	2	3	1	2	-	-
1999.....	10	-	-	-	1	-	-	3	3	2	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant melanoma of skin (C43)

American Indian 2/, male

2007 1/.....	14	-	-	-	-	1	-	4	5	1	1	2	-
2006 1/.....	12	-	-	-	-	-	1	-	3	2	3	3	-
2005 1/.....	15	-	-	-	2	-	4	2	4	2	1	-	-
2004 1/.....	7	-	-	-	-	-	1	1	2	3	-	-	-
2003 1/.....	5	-	-	-	-	-	-	1	2	-	2	-	-
2002.....	9	-	-	-	-	-	-	2	3	4	-	-	-
2001.....	9	-	-	-	-	1	-	1	3	-	2	2	-
2000.....	5	-	-	-	-	-	-	1	2	1	1	-	-
1999.....	7	-	-	-	1	-	-	3	1	2	-	-	-

American Indian 2/, female

2007 1/.....	8	-	-	-	-	1	-	-	-	3	4	-	-
2006 1/.....	9	-	-	-	-	-	2	2	1	1	1	2	-
2005 1/.....	5	-	-	-	-	-	1	-	3	-	-	1	-
2004 1/.....	4	-	-	-	-	-	-	2	-	1	1	-	-
2003 1/.....	9	-	-	-	1	-	-	1	2	2	3	-	-
2002.....	8	-	-	-	-	-	-	-	1	2	2	3	-
2001.....	1	-	-	-	-	-	-	-	-	-	1	-	-
2000.....	3	-	-	-	-	-	-	1	1	-	1	-	-
1999.....	3	-	-	-	-	-	-	-	2	-	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant melanoma of skin (C43)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	45	-	-	-	-	-	4	7	8	12	10	4	-
2006 1/.....	44	-	-	1	1	1	5	6	7	9	8	6	-
2005 1/.....	55	-	-	-	1	2	3	12	11	11	8	7	-
2004 1/.....	21	-	-	-	-	1	1	2	4	4	6	3	-
2003 1/.....	35	-	-	-	-	2	1	10	7	1	8	6	-
2002.....	42	-	-	-	-	1	5	8	7	11	7	3	-
2001.....	25	-	-	-	-	-	-	6	5	8	5	1	-
2000.....	32	-	-	-	-	3	5	5	7	8	2	2	-
1999.....	23	-	-	-	1	2	3	1	2	2	9	3	-

Asian or Pacific Islander 3/, male

2007 1/.....	25	-	-	-	-	-	1	4	4	9	5	2	-
2006 1/.....	18	-	-	-	-	1	2	3	3	4	3	2	-
2005 1/.....	27	-	-	-	-	2	2	5	6	4	4	4	-
2004 1/.....	8	-	-	-	-	1	-	2	1	-	3	1	-
2003 1/.....	21	-	-	-	-	-	-	5	4	-	8	4	-
2002.....	25	-	-	-	-	1	2	5	4	8	4	1	-
2001.....	15	-	-	-	-	-	-	3	3	4	5	-	-
2000.....	13	-	-	-	-	-	1	2	2	5	1	2	-
1999.....	15	-	-	-	1	2	2	-	2	1	6	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant melanoma of skin (C43)

Asian or Pacific Islander 3/, female

2007 1/.....	20	-	-	-	-	-	3	3	4	3	5	2	-
2006 1/.....	26	-	-	1	1	-	3	3	4	5	5	4	-
2005 1/.....	28	-	-	-	1	-	1	7	5	7	4	3	-
2004 1/.....	13	-	-	-	-	-	1	-	3	4	3	2	-
2003 1/.....	14	-	-	-	-	2	1	5	3	1	-	2	-
2002.....	17	-	-	-	-	-	3	3	3	3	3	2	-
2001.....	10	-	-	-	-	-	-	3	2	4	-	1	-
2000.....	19	-	-	-	-	3	4	3	5	3	1	-	-
1999.....	8	-	-	-	-	-	1	1	-	1	3	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of breast (C50)													
All races, both sexes													
2007.....	40,970	-	-	-	15	344	2,184	5,990	8,756	8,179	9,075	6,426	1
2006.....	41,210	-	1	-	13	368	2,352	6,115	8,871	7,961	9,346	6,182	1
2005.....	41,491	1	-	-	12	356	2,497	6,232	8,646	8,113	9,382	6,252	-
2004.....	41,316	-	-	-	11	396	2,511	6,232	8,474	8,285	9,371	6,036	-
2003.....	42,000	-	-	-	16	407	2,716	6,365	8,267	8,338	9,644	6,245	2
2002.....	41,883	-	-	-	10	425	2,722	6,454	7,838	8,505	9,748	6,181	-
2001.....	41,809	-	-	-	9	463	2,823	6,593	7,632	8,715	9,663	5,911	-
2000.....	42,300	-	-	-	11	467	2,841	6,385	7,573	9,024	9,750	6,249	-
1999.....	41,528	-	1	1	13	439	2,741	6,174	7,399	9,109	9,651	6,000	-
All races, male													
2007.....	371	-	-	-	-	-	10	36	70	91	104	60	-
2006.....	389	-	-	-	1	3	6	43	88	86	111	51	-
2005.....	375	1	-	-	-	-	9	33	73	109	111	39	-
2004.....	362	-	-	-	1	1	10	33	67	91	109	50	-
2003.....	380	-	-	-	-	2	9	48	71	89	104	56	1
2002.....	369	-	-	-	-	3	12	44	67	84	112	47	-
2001.....	415	-	-	-	-	2	14	46	72	120	115	46	-
2000.....	428	-	-	-	-	3	19	46	85	115	109	51	-
1999.....	384	-	-	-	-	3	15	41	70	103	98	54	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of breast (C50)													
All races, female													
2007.....	40,599	-	-	-	15	344	2,174	5,954	8,686	8,088	8,971	6,366	1
2006.....	40,821	-	1	-	12	365	2,346	6,072	8,783	7,875	9,235	6,131	1
2005.....	41,116	-	-	-	12	356	2,488	6,199	8,573	8,004	9,271	6,213	-
2004.....	40,954	-	-	-	10	395	2,501	6,199	8,407	8,194	9,262	5,986	-
2003.....	41,620	-	-	-	16	405	2,707	6,317	8,196	8,249	9,540	6,189	1
2002.....	41,514	-	-	-	10	422	2,710	6,410	7,771	8,421	9,636	6,134	-
2001.....	41,394	-	-	-	9	461	2,809	6,547	7,560	8,595	9,548	5,865	-
2000.....	41,872	-	-	-	11	464	2,822	6,339	7,488	8,909	9,641	6,198	-
1999.....	41,144	-	1	1	13	436	2,726	6,133	7,329	9,006	9,553	5,946	-
White, both sexes													
2007 1/.....	34,160	-	-	-	8	218	1,551	4,511	7,064	6,960	8,035	5,813	-
2006 1/.....	34,464	-	1	-	6	245	1,692	4,616	7,220	6,803	8,316	5,565	-
2005 1/.....	34,663	1	-	-	7	235	1,787	4,646	7,002	6,936	8,393	5,656	-
2004 1/.....	34,759	-	-	-	7	263	1,781	4,700	6,966	7,140	8,436	5,466	-
2003 1/.....	35,358	-	-	-	11	269	1,981	4,875	6,786	7,158	8,638	5,639	1
2002.....	35,382	-	-	-	6	285	1,950	4,963	6,494	7,333	8,733	5,618	-
2001.....	35,422	-	-	-	4	306	2,069	5,143	6,307	7,512	8,707	5,374	-
2000.....	36,127	-	-	-	5	325	2,097	5,000	6,282	7,925	8,842	5,651	-
1999.....	35,385	-	-	1	8	304	2,001	4,826	6,111	7,922	8,714	5,498	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of breast (C50)													
White, male													
2007 1/.....	315	-	-	-	-	-	8	28	56	78	91	54	-
2006 1/.....	327	-	-	-	-	3	4	28	72	73	99	48	-
2005 1/.....	311	1	-	-	-	-	5	26	60	88	96	35	-
2004 1/.....	301	-	-	-	-	-	6	25	53	76	100	41	-
2003 1/.....	326	-	-	-	-	1	4	39	59	76	95	51	1
2002.....	300	-	-	-	-	2	9	32	51	68	96	42	-
2001.....	344	-	-	-	-	1	8	34	61	101	100	39	-
2000.....	360	-	-	-	-	1	11	37	67	103	98	43	-
1999.....	312	-	-	-	-	2	8	30	54	85	86	47	-
White, female													
2007 1/.....	33,845	-	-	-	8	218	1,543	4,483	7,008	6,882	7,944	5,759	-
2006 1/.....	34,137	-	1	-	6	242	1,688	4,588	7,148	6,730	8,217	5,517	-
2005 1/.....	34,352	-	-	-	7	235	1,782	4,620	6,942	6,848	8,297	5,621	-
2004 1/.....	34,458	-	-	-	7	263	1,775	4,675	6,913	7,064	8,336	5,425	-
2003 1/.....	35,032	-	-	-	11	268	1,977	4,836	6,727	7,082	8,543	5,588	-
2002.....	35,082	-	-	-	6	283	1,941	4,931	6,443	7,265	8,637	5,576	-
2001.....	35,078	-	-	-	4	305	2,061	5,109	6,246	7,411	8,607	5,335	-
2000.....	35,767	-	-	-	5	324	2,086	4,963	6,215	7,822	8,744	5,608	-
1999.....	35,073	-	-	1	8	302	1,993	4,796	6,057	7,837	8,628	5,451	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of breast (C50)													
Black, both sexes													
2007 1/.....	5,852	-	-	-	6	107	548	1,273	1,428	1,043	898	548	1
2006 1/.....	5,752	-	-	-	6	108	576	1,259	1,375	972	908	548	-
2005 1/.....	5,858	-	-	-	5	103	614	1,370	1,386	1,011	829	540	-
2004 1/.....	5,597	-	-	-	2	116	627	1,280	1,268	976	819	509	-
2003 1/.....	5,746	-	-	-	5	119	640	1,270	1,254	1,013	891	554	-
2002.....	5,636	-	-	-	3	125	679	1,258	1,137	1,014	906	514	-
2001.....	5,571	-	-	-	5	142	645	1,221	1,131	1,060	872	495	-
2000.....	5,424	-	-	-	5	123	645	1,178	1,110	969	830	564	-
1999.....	5,425	-	1	-	5	126	665	1,167	1,103	1,040	856	462	-
Black, male													
2007 1/.....	53	-	-	-	-	-	2	8	14	13	11	5	-
2006 1/.....	58	-	-	-	1	-	2	15	14	13	10	3	-
2005 1/.....	56	-	-	-	-	-	2	6	12	21	13	2	-
2004 1/.....	56	-	-	-	-	1	3	8	12	14	9	9	-
2003 1/.....	51	-	-	-	-	1	5	8	12	13	7	5	-
2002.....	64	-	-	-	-	1	3	11	15	16	13	5	-
2001.....	65	-	-	-	-	1	5	12	10	17	14	6	-
2000.....	63	-	-	-	-	2	8	8	16	11	10	8	-
1999.....	67	-	-	-	-	1	7	9	16	16	11	7	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of breast (C50)													
Black, female													
2007 1/.....	5,799	-	-	-	6	107	546	1,265	1,414	1,030	887	543	1
2006 1/.....	5,694	-	-	-	5	108	574	1,244	1,361	959	898	545	-
2005 1/.....	5,802	-	-	-	5	103	612	1,364	1,374	990	816	538	-
2004 1/.....	5,541	-	-	-	2	115	624	1,272	1,256	962	810	500	-
2003 1/.....	5,695	-	-	-	5	118	635	1,262	1,242	1,000	884	549	-
2002.....	5,572	-	-	-	3	124	676	1,247	1,122	998	893	509	-
2001.....	5,506	-	-	-	5	141	640	1,209	1,121	1,043	858	489	-
2000.....	5,361	-	-	-	5	121	637	1,170	1,094	958	820	556	-
1999.....	5,358	-	1	-	5	125	658	1,158	1,087	1,024	845	455	-
American Indian 2/, both sexes													
2007 1/.....	170	-	-	-	-	3	9	40	50	28	21	19	-
2006 1/.....	160	-	-	-	-	3	17	34	42	30	20	14	-
2005 1/.....	167	-	-	-	-	5	12	27	27	44	42	10	-
2004 1/.....	169	-	-	-	-	3	15	42	39	28	29	13	-
2003 1/.....	148	-	-	-	-	4	9	28	36	24	33	14	-
2002.....	148	-	-	-	-	5	17	36	31	29	18	12	-
2001.....	127	-	-	-	-	2	15	35	32	25	8	10	-
2000.....	131	-	-	-	-	2	16	25	39	23	20	6	-
1999.....	131	-	-	-	-	2	11	26	28	36	17	11	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Malignant neoplasm of breast (C50)

American Indian 2/, male

2007 1/.....	2	-	-	-	-	-	-	-	-	-	1	1	-
2006 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2005 1/.....	1	-	-	-	-	-	-	-	-	-	-	1	-
2004 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 1/.....	1	-	-	-	-	-	-	-	-	-	1	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	1	-	-	1	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	1	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	1	-	-	-

American Indian 2/, female

2007 1/.....	168	-	-	-	-	3	9	40	50	28	20	18	-
2006 1/.....	160	-	-	-	-	3	17	34	42	30	20	14	-
2005 1/.....	166	-	-	-	-	5	12	27	27	44	42	9	-
2004 1/.....	169	-	-	-	-	3	15	42	39	28	29	13	-
2003 1/.....	147	-	-	-	-	4	9	28	36	24	32	14	-
2002.....	148	-	-	-	-	5	17	36	31	29	18	12	-
2001.....	126	-	-	-	-	2	15	35	32	24	8	10	-
2000.....	130	-	-	-	-	2	16	25	38	23	20	6	-
1999.....	130	-	-	-	-	2	11	26	28	35	17	11	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasm of breast (C50)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	788	-	-	-	1	16	76	166	214	148	121	46	-
2006 1/.....	834	-	-	-	1	12	67	206	234	156	102	55	1
2005 1/.....	803	-	-	-	-	13	84	189	231	122	118	46	-
2004 1/.....	791	-	-	-	2	14	88	210	201	141	87	48	-
2003 1/.....	748	-	-	-	-	15	86	192	191	143	82	38	1
2002.....	717	-	-	-	1	10	76	197	176	129	91	37	-
2001.....	689	-	-	-	-	13	94	194	162	118	76	32	-
2000.....	618	-	-	-	1	17	83	182	142	107	58	28	-
1999.....	587	-	-	-	-	7	64	155	157	111	64	29	-

Asian or Pacific Islander 3/, male

2007 1/.....	1	-	-	-	-	-	-	-	-	-	1	-	-
2006 1/.....	4	-	-	-	-	-	-	-	2	-	2	-	-
2005 1/.....	7	-	-	-	-	-	2	1	1	-	2	1	-
2004 1/.....	5	-	-	-	1	-	1	-	2	1	-	-	-
2003 1/.....	2	-	-	-	-	-	-	1	-	-	1	-	-
2002.....	5	-	-	-	-	-	-	1	1	-	3	-	-
2001.....	5	-	-	-	-	-	1	-	1	1	1	1	-
2000.....	4	-	-	-	-	-	-	1	1	1	1	-	-
1999.....	4	-	-	-	-	-	-	2	-	1	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasm of breast (C50)

Asian or Pacific Islander 3/, female

2007 1/.....	787	-	-	-	1	16	76	166	214	148	120	46	-
2006 1/.....	830	-	-	-	1	12	67	206	232	156	100	55	1
2005 1/.....	796	-	-	-	-	13	82	188	230	122	116	45	-
2004 1/.....	786	-	-	-	1	14	87	210	199	140	87	48	-
2003 1/.....	746	-	-	-	-	15	86	191	191	143	81	38	1
2002.....	712	-	-	-	1	10	76	196	175	129	88	37	-
2001.....	684	-	-	-	-	13	93	194	161	117	75	31	-
2000.....	614	-	-	-	1	17	83	181	141	106	57	28	-
1999.....	583	-	-	-	-	7	64	153	157	110	63	29	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of cervix uteri (C53)													
All races, female													
2007.....	4,021	-	-	-	8	183	645	915	892	611	506	261	-
2006.....	3,976	-	-	1	8	181	627	963	821	604	493	278	-
2005.....	3,924	-	-	-	14	168	622	958	796	575	518	273	-
2004.....	3,850	-	-	-	19	213	622	850	815	601	484	246	-
2003.....	3,919	-	-	-	16	201	633	890	793	564	552	270	-
2002.....	3,952	-	-	-	17	208	666	894	725	630	528	284	-
2001.....	4,092	-	-	1	12	209	711	953	723	585	583	315	-
2000.....	4,200	-	-	1	17	230	694	915	748	686	599	310	-
1999.....	4,205	-	-	-	12	217	699	914	769	696	568	329	1
White, both sexes													
2007 1/.....	3,037	-	-	-	7	129	480	691	683	460	390	197	-
2006 1/.....	3,014	-	-	1	5	126	466	736	654	443	383	200	-
2005 1/.....	2,982	-	-	-	10	133	471	713	595	443	399	218	-
2004 1/.....	2,899	-	-	-	14	158	467	635	633	438	376	178	-
2003 1/.....	2,953	-	-	-	13	139	478	675	613	414	411	210	-
2002.....	2,967	-	-	-	12	160	503	667	553	455	407	210	-
2001.....	3,136	-	-	-	11	162	551	734	555	430	446	247	-
2000.....	3,158	-	-	-	13	174	511	686	557	498	478	241	-
1999.....	3,229	-	-	-	9	176	535	731	573	503	446	255	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Malignant neoplasm of cervix uteri (C53)

White, male

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
2007 1/.....
2006 1/.....
2005 1/.....
2004 1/.....
2003 1/.....
2002.....
2001.....
2000.....
1999.....

White, female

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
2007 1/.....	3,037	-	-	-	7	129	480	691	683	460	390	197	-
2006 1/.....	3,014	-	-	1	5	126	466	736	654	443	383	200	-
2005 1/.....	2,982	-	-	-	10	133	471	713	595	443	399	218	-
2004 1/.....	2,899	-	-	-	14	158	467	635	633	438	376	178	-
2003 1/.....	2,953	-	-	-	13	139	478	675	613	414	411	210	-
2002.....	2,967	-	-	-	12	160	503	667	553	455	407	210	-
2001.....	3,136	-	-	-	11	162	551	734	555	430	446	247	-
2000.....	3,158	-	-	-	13	174	511	686	557	498	478	241	-
1999.....	3,229	-	-	-	9	176	535	731	573	503	446	255	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasm of cervix uteri (C53)

Black, female

2007 1/.....	805	-	-	-	1	50	137	183	160	126	96	52	-
2006 1/.....	784	-	-	-	3	42	137	184	128	135	87	68	-
2005 1/.....	783	-	-	-	4	30	133	189	159	112	107	49	-
2004 1/.....	789	-	-	-	4	45	128	182	154	131	88	57	-
2003 1/.....	794	-	-	-	2	50	127	175	153	126	111	50	-
2002.....	829	-	-	-	3	43	146	191	131	152	104	59	-
2001.....	787	-	-	1	1	42	139	169	131	130	112	62	-
2000.....	869	-	-	1	4	51	150	180	153	158	109	63	-
1999.....	838	-	-	-	3	36	144	140	169	176	102	68	-

American Indian 2/, both sexes

2007 1/.....	32	-	-	-	-	-	5	8	9	6	1	3	-
2006 1/.....	26	-	-	-	-	3	3	6	7	6	1	-	-
2005 1/.....	34	-	-	-	-	1	4	17	7	4	-	1	-
2004 1/.....	31	-	-	-	-	4	10	7	4	3	1	2	-
2003 1/.....	31	-	-	-	-	2	7	5	8	3	4	2	-
2002.....	34	-	-	-	2	2	3	12	4	5	3	3	-
2001.....	32	-	-	-	-	1	7	12	6	2	2	2	-
2000.....	30	-	-	-	-	2	7	9	5	3	2	2	-
1999.....	19	-	-	-	-	2	3	6	5	1	2	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Malignant neoplasm of cervix uteri (C53)

American Indian 2/, male

2007 1/.....
2006 1/.....
2005 1/.....
2004 1/.....
2003 1/.....
2002.....
2001.....
2000.....
1999.....

American Indian 2/, female

2007 1/.....	32	-	-	-	-	-	5	8	9	6	1	3	-
2006 1/.....	26	-	-	-	-	3	3	6	7	6	1	-	-
2005 1/.....	34	-	-	-	-	1	4	17	7	4	-	1	-
2004 1/.....	31	-	-	-	-	4	10	7	4	3	1	2	-
2003 1/.....	31	-	-	-	-	2	7	5	8	3	4	2	-
2002.....	34	-	-	-	2	2	3	12	4	5	3	3	-
2001.....	32	-	-	-	-	1	7	12	6	2	2	2	-
2000.....	30	-	-	-	-	2	7	9	5	3	2	2	-
1999.....	19	-	-	-	-	2	3	6	5	1	2	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasm of cervix uteri (C53)

Asian or Pacific Islander 3/, female

2007 1/.....	147	-	-	-	-	4	23	33	40	19	19	9	-
2006 1/.....	152	-	-	-	-	10	21	37	32	20	22	10	-
2005 1/.....	125	-	-	-	-	4	14	39	35	16	12	5	-
2004 1/.....	131	-	-	-	1	6	17	26	24	29	19	9	-
2003 1/.....	141	-	-	-	1	10	21	35	19	21	26	8	-
2002.....	122	-	-	-	-	3	14	24	37	18	14	12	-
2001.....	137	-	-	-	-	4	14	38	31	23	23	4	-
2000.....	143	-	-	-	-	3	26	40	33	27	10	4	-
1999.....	119	-	-	-	-	3	17	37	22	16	18	6	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

All races, female

2007.....	7,456	-	-	-	2	31	168	584	1,583	2,021	1,960	1,107	-
2006.....	7,384	1	-	-	2	27	145	599	1,522	1,897	2,049	1,142	-
2005.....	7,096	-	-	-	4	22	146	568	1,468	1,808	1,955	1,125	-
2004.....	6,990	-	-	-	5	20	168	557	1,290	1,856	2,011	1,083	-
2003.....	6,899	-	-	-	4	27	154	534	1,301	1,851	1,951	1,077	-
2002.....	6,853	-	-	-	-	23	146	565	1,199	1,835	1,957	1,128	-
2001.....	6,784	-	-	-	-	38	139	566	1,105	1,784	2,063	1,088	1
2000.....	6,586	-	-	-	3	27	138	507	1,122	1,781	1,958	1,049	1
1999.....	6,468	-	-	-	2	23	142	468	1,019	1,807	2,008	999	-

White, both sexes

2007 1/.....	5,962	-	-	-	1	24	119	463	1,247	1,517	1,606	985	-
2006 1/.....	5,954	1	-	-	-	20	118	474	1,160	1,428	1,722	1,031	-
2005 1/.....	5,742	-	-	-	2	13	110	449	1,118	1,414	1,639	997	-
2004 1/.....	5,713	-	-	-	3	16	131	433	1,011	1,411	1,725	983	-
2003 1/.....	5,641	-	-	-	3	16	108	428	1,016	1,452	1,650	968	-
2002.....	5,567	-	-	-	-	17	116	443	934	1,419	1,648	990	-
2001.....	5,606	-	-	-	-	26	107	446	872	1,404	1,776	975	-
2000.....	5,397	-	-	-	1	20	106	420	890	1,373	1,651	936	-
1999.....	5,386	-	-	-	2	18	109	386	816	1,435	1,739	881	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)													
White, male													
2007 1/.....
2006 1/.....
2005 1/.....
2004 1/.....
2003 1/.....
2002.....
2001.....
2000.....
1999.....
White, female													
2007 1/.....	5,962	-	-	-	1	24	119	463	1,247	1,517	1,606	985	-
2006 1/.....	5,954	1	-	-	-	20	118	474	1,160	1,428	1,722	1,031	-
2005 1/.....	5,742	-	-	-	2	13	110	449	1,118	1,414	1,639	997	-
2004 1/.....	5,713	-	-	-	3	16	131	433	1,011	1,411	1,725	983	-
2003 1/.....	5,641	-	-	-	3	16	108	428	1,016	1,452	1,650	968	-
2002.....	5,567	-	-	-	-	17	116	443	934	1,419	1,648	990	-
2001.....	5,606	-	-	-	-	26	107	446	872	1,404	1,776	975	-
2000.....	5,397	-	-	-	1	20	106	420	890	1,373	1,651	936	-
1999.....	5,386	-	-	-	2	18	109	386	816	1,435	1,739	881	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

Black, female

2007 1/.....	1,295	-	-	-	1	6	33	97	286	452	312	108	-
2006 1/.....	1,240	-	-	-	1	7	19	89	314	415	295	100	-
2005 1/.....	1,180	-	-	-	2	7	25	95	301	355	278	117	-
2004 1/.....	1,114	-	-	-	2	4	30	95	232	398	260	93	-
2003 1/.....	1,107	-	-	-	1	8	31	84	248	359	273	103	-
2002.....	1,119	-	-	-	-	5	22	89	223	376	273	131	-
2001.....	1,047	-	-	-	-	9	22	93	195	354	268	105	1
2000.....	1,061	-	-	-	1	6	22	68	202	375	280	106	1
1999.....	987	-	-	-	-	5	28	57	189	350	247	111	-

American Indian 2/, both sexes

2007 1/.....	32	-	-	-	-	-	3	5	5	9	7	3	-
2006 1/.....	30	-	-	-	-	-	2	5	5	8	6	4	-
2005 1/.....	22	-	-	-	-	-	-	3	4	7	6	2	-
2004 1/.....	21	-	-	-	-	-	-	5	4	7	4	1	-
2003 1/.....	11	-	-	-	-	1	1	2	1	4	2	-	-
2002.....	32	-	-	-	-	-	4	7	6	4	9	2	-
2001.....	25	-	-	-	-	-	2	3	6	6	5	3	-
2000.....	19	-	-	-	-	-	1	1	2	8	6	1	-
1999.....	13	-	-	-	-	-	2	2	2	4	1	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

American Indian 2/, male

2007 1/.....
2006 1/.....
2005 1/.....
2004 1/.....
2003 1/.....
2002.....
2001.....
2000.....
1999.....

American Indian 2/, female

2007 1/.....	32	-	-	-	-	-	3	5	5	9	7	3	-
2006 1/.....	30	-	-	-	-	-	2	5	5	8	6	4	-
2005 1/.....	22	-	-	-	-	-	-	3	4	7	6	2	-
2004 1/.....	21	-	-	-	-	-	-	5	4	7	4	1	-
2003 1/.....	11	-	-	-	-	1	1	2	1	4	2	-	-
2002.....	32	-	-	-	-	-	4	7	6	4	9	2	-
2001.....	25	-	-	-	-	-	2	3	6	6	5	3	-
2000.....	19	-	-	-	-	-	1	1	2	8	6	1	-
1999.....	13	-	-	-	-	-	2	2	2	4	1	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

Asian or Pacific Islander 3/, female

2007 1/.....	167	-	-	-	-	1	13	19	45	43	35	11	-
2006 1/.....	160	-	-	-	1	-	6	31	43	46	26	7	-
2005 1/.....	152	-	-	-	-	2	11	21	45	32	32	9	-
2004 1/.....	142	-	-	-	-	-	7	24	43	40	22	6	-
2003 1/.....	140	-	-	-	-	2	14	20	36	36	26	6	-
2002.....	135	-	-	-	-	1	4	26	36	36	27	5	-
2001.....	106	-	-	-	-	3	8	24	32	20	14	5	-
2000.....	109	-	-	-	1	1	9	18	28	25	21	6	-
1999.....	82	-	-	-	-	-	3	23	12	18	21	5	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of ovary (C56)													
All races, female													
2007.....	14,621	-	-	-	28	79	352	1,532	2,997	3,616	3,946	2,071	-
2006.....	14,857	1	1	3	23	79	401	1,666	3,048	3,573	4,061	2,001	-
2005.....	14,787	-	-	3	18	83	413	1,682	3,048	3,569	4,031	1,940	-
2004.....	14,716	-	-	2	29	94	454	1,654	2,794	3,693	4,165	1,831	-
2003.....	14,657	-	-	-	28	84	452	1,629	2,891	3,569	4,194	1,810	-
2002.....	14,682	1	1	2	37	97	469	1,717	2,748	3,601	4,259	1,750	-
2001.....	14,414	-	-	2	21	102	472	1,626	2,612	3,666	4,170	1,743	-
2000.....	14,060	-	-	2	23	101	472	1,583	2,444	3,601	4,098	1,736	-
1999.....	13,627	-	1	2	27	77	496	1,549	2,426	3,612	3,875	1,562	-
White, both sexes													
2007 1/.....	13,093	-	-	-	23	61	302	1,304	2,630	3,239	3,610	1,924	-
2006 1/.....	13,289	1	-	3	20	67	335	1,411	2,683	3,212	3,712	1,845	-
2005 1/.....	13,199	-	-	2	18	67	332	1,446	2,696	3,151	3,683	1,804	-
2004 1/.....	13,127	-	-	2	20	70	375	1,393	2,460	3,270	3,846	1,691	-
2003 1/.....	13,123	-	-	-	21	69	373	1,387	2,571	3,181	3,858	1,663	-
2002.....	13,202	1	1	2	30	77	391	1,488	2,441	3,235	3,917	1,619	-
2001.....	12,950	-	-	1	14	81	390	1,426	2,284	3,289	3,856	1,609	-
2000.....	12,731	-	-	1	17	84	397	1,379	2,187	3,241	3,814	1,611	-
1999.....	12,302	-	1	2	22	57	424	1,349	2,164	3,254	3,570	1,459	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Malignant neoplasm of ovary (C56)

White, male

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
2007 1/.....
2006 1/.....
2005 1/.....
2004 1/.....
2003 1/.....
2002.....
2001.....
2000.....
1999.....

White, female

2007 1/.....	13,093	-	-	-	23	61	302	1,304	2,630	3,239	3,610	1,924	-
2006 1/.....	13,289	1	-	3	20	67	335	1,411	2,683	3,212	3,712	1,845	-
2005 1/.....	13,199	-	-	2	18	67	332	1,446	2,696	3,151	3,683	1,804	-
2004 1/.....	13,127	-	-	2	20	70	375	1,393	2,460	3,270	3,846	1,691	-
2003 1/.....	13,123	-	-	-	21	69	373	1,387	2,571	3,181	3,858	1,663	-
2002.....	13,202	1	1	2	30	77	391	1,488	2,441	3,235	3,917	1,619	-
2001.....	12,950	-	-	1	14	81	390	1,426	2,284	3,289	3,856	1,609	-
2000.....	12,731	-	-	1	17	84	397	1,379	2,187	3,241	3,814	1,611	-
1999.....	12,302	-	1	2	22	57	424	1,349	2,164	3,254	3,570	1,459	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of ovary (C56)													
Black, female													
2007 1/.....	1,160	-	-	-	3	17	37	155	278	301	260	109	-
2006 1/.....	1,185	-	1	-	2	8	49	180	274	286	256	129	-
2005 1/.....	1,217	-	-	1	-	14	55	160	263	328	279	117	-
2004 1/.....	1,256	-	-	-	7	17	53	192	250	349	272	116	-
2003 1/.....	1,196	-	-	-	6	12	58	184	238	302	270	126	-
2002.....	1,163	-	-	-	7	13	59	160	240	289	278	117	-
2001.....	1,153	-	-	1	4	15	66	136	253	305	254	119	-
2000.....	1,073	-	-	1	2	12	60	140	205	299	244	110	-
1999.....	1,079	-	-	-	4	13	47	144	217	304	260	90	-
American Indian 2/, both sexes													
2007 1/.....	56	-	-	-	-	-	-	9	11	15	12	9	-
2006 1/.....	62	-	-	-	-	-	1	12	10	15	20	4	-
2005 1/.....	55	-	-	-	-	1	2	14	11	16	9	2	-
2004 1/.....	57	-	-	-	-	-	5	9	18	12	7	6	-
2003 1/.....	62	-	-	-	-	-	2	12	15	12	16	5	-
2002.....	50	-	-	-	-	2	4	13	8	12	9	2	-
2001.....	68	-	-	-	-	-	4	11	16	20	14	3	-
2000.....	28	-	-	-	1	-	1	3	7	5	9	2	-
1999.....	29	-	-	-	-	-	1	3	3	11	9	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Malignant neoplasm of ovary (C56)

American Indian 2/, male

2007 1/.....
2006 1/.....
2005 1/.....
2004 1/.....
2003 1/.....
2002.....
2001.....
2000.....
1999.....

American Indian 2/, female

2007 1/.....	56	-	-	-	-	-	-	9	11	15	12	9	-
2006 1/.....	62	-	-	-	-	-	1	12	10	15	20	4	-
2005 1/.....	55	-	-	-	-	1	2	14	11	16	9	2	-
2004 1/.....	57	-	-	-	-	-	5	9	18	12	7	6	-
2003 1/.....	62	-	-	-	-	-	2	12	15	12	16	5	-
2002.....	50	-	-	-	-	2	4	13	8	12	9	2	-
2001.....	68	-	-	-	-	-	4	11	16	20	14	3	-
2000.....	28	-	-	-	1	-	1	3	7	5	9	2	-
1999.....	29	-	-	-	-	-	1	3	3	11	9	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasm of ovary (C56)

Asian or Pacific Islander 3/, female

2007 1/.....	312	-	-	-	2	1	13	64	78	61	64	29	-
2006 1/.....	321	-	-	-	1	4	16	63	81	60	73	23	-
2005 1/.....	316	-	-	-	-	1	24	62	78	74	60	17	-
2004 1/.....	276	-	-	-	2	7	21	60	66	62	40	18	-
2003 1/.....	276	-	-	-	1	3	19	46	67	74	50	16	-
2002.....	267	-	-	-	-	5	15	56	59	65	55	12	-
2001.....	243	-	-	-	3	6	12	53	59	52	46	12	-
2000.....	228	-	-	-	3	5	14	61	45	56	31	13	-
1999.....	217	-	-	-	1	7	24	53	42	43	36	11	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasm of prostate (C61)

All races, both sexes

2007.....	29,093	-	1	-	1	1	21	428	2,271	5,716	11,257	9,397	-
2006.....	28,372	-	-	1	1	5	40	405	2,273	5,603	11,157	8,887	-
2005.....	28,905	-	1	-	1	3	24	395	2,154	5,764	11,666	8,897	-
2004.....	29,004	-	1	1	1	2	23	434	2,044	5,711	11,993	8,792	2
2003.....	29,554	-	-	1	1	1	25	418	2,074	6,033	12,284	8,717	-
2002.....	30,446	-	-	-	1	1	32	382	2,020	6,300	12,753	8,957	-
2001.....	30,719	-	-	-	1	1	30	412	1,907	6,543	12,976	8,849	-
2000.....	31,078	-	-	-	-	1	25	371	1,929	6,808	13,071	8,873	-
1999.....	31,729	1	-	-	1	2	33	374	1,946	7,316	13,435	8,620	1

All races, male

2007.....	29,093	-	1	-	1	1	21	428	2,271	5,716	11,257	9,397	-
2006.....	28,372	-	-	1	1	5	40	405	2,273	5,603	11,157	8,887	-
2005.....	28,905	-	1	-	1	3	24	395	2,154	5,764	11,666	8,897	-
2004.....	29,004	-	1	1	1	2	23	434	2,044	5,711	11,993	8,792	2
2003.....	29,554	-	-	1	1	1	25	418	2,074	6,033	12,284	8,717	-
2002.....	30,446	-	-	-	1	1	32	382	2,020	6,300	12,753	8,957	-
2001.....	30,719	-	-	-	1	1	30	412	1,907	6,543	12,976	8,849	-
2000.....	31,078	-	-	-	-	1	25	371	1,929	6,808	13,071	8,873	-
1999.....	31,729	1	-	-	1	2	33	374	1,946	7,316	13,435	8,620	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Malignant neoplasm of prostate (C61)

All races, female

2007
2006
2005
2004
2003
2002
2001
2000
1999

White, both sexes

2007 1/	23,666	-	-	-	1	1	16	304	1,656	4,324	9,295	8,069	-
2006 1/	23,202	-	-	1	-	2	27	288	1,683	4,237	9,262	7,702	-
2005 1/	23,597	-	1	-	1	2	13	265	1,566	4,402	9,700	7,647	-
2004 1/	23,727	-	1	1	1	2	17	288	1,518	4,426	9,925	7,547	1
2003 1/	24,232	-	-	1	1	1	17	289	1,503	4,690	10,175	7,555	-
2002	24,918	-	-	-	-	1	25	255	1,463	4,858	10,566	7,750	-
2001	25,066	-	-	-	1	-	19	285	1,396	5,087	10,730	7,548	-
2000	25,340	-	-	-	-	1	16	262	1,404	5,231	10,804	7,622	-
1999	26,005	1	-	-	1	2	25	266	1,404	5,715	11,240	7,350	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of prostate (C61)													
Black, both sexes													
2007 1/.....	4,908	-	1	-	-	-	5	119	575	1,282	1,754	1,172	-
2006 1/.....	4,698	-	-	-	1	3	12	116	541	1,253	1,719	1,053	-
2005 1/.....	4,823	-	-	-	-	-	10	120	546	1,264	1,777	1,106	-
2004 1/.....	4,816	-	-	-	-	-	4	140	498	1,199	1,874	1,100	1
2003 1/.....	4,894	-	-	-	-	-	8	125	541	1,262	1,926	1,032	-
2002.....	5,145	-	-	-	1	-	7	119	512	1,370	2,030	1,106	-
2001.....	5,265	-	-	-	-	1	11	124	487	1,384	2,076	1,182	-
2000.....	5,346	-	-	-	-	-	9	102	497	1,491	2,107	1,140	-
1999.....	5,357	-	-	-	-	-	7	106	515	1,534	2,046	1,149	-
Black, male													
2007 1/.....	4,908	-	1	-	-	-	5	119	575	1,282	1,754	1,172	-
2006 1/.....	4,698	-	-	-	1	3	12	116	541	1,253	1,719	1,053	-
2005 1/.....	4,823	-	-	-	-	-	10	120	546	1,264	1,777	1,106	-
2004 1/.....	4,816	-	-	-	-	-	4	140	498	1,199	1,874	1,100	1
2003 1/.....	4,894	-	-	-	-	-	8	125	541	1,262	1,926	1,032	-
2002.....	5,145	-	-	-	1	-	7	119	512	1,370	2,030	1,106	-
2001.....	5,265	-	-	-	-	1	11	124	487	1,384	2,076	1,182	-
2000.....	5,346	-	-	-	-	-	9	102	497	1,491	2,107	1,140	-
1999.....	5,357	-	-	-	-	-	7	106	515	1,534	2,046	1,149	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Malignant neoplasm of prostate (C61)

Black, female

2007 1/.....
2006 1/.....
2005 1/.....
2004 1/.....
2003 1/.....
2002.....
2001.....
2000.....
1999.....

American Indian 2/, both sexes

2007 1/.....	105	-	-	-	-	-	-	2	17	19	43	24	-
2006 1/.....	104	-	-	-	-	-	-	1	15	31	38	19	-
2005 1/.....	113	-	-	-	-	-	1	3	11	24	47	27	-
2004 1/.....	99	-	-	-	-	-	1	2	10	26	38	22	-
2003 1/.....	100	-	-	-	-	-	-	1	12	23	39	25	-
2002.....	85	-	-	-	-	-	-	2	12	20	36	15	-
2001.....	92	-	-	-	-	-	-	2	9	22	35	24	-
2000.....	92	-	-	-	-	-	-	4	6	30	33	19	-
1999.....	63	-	-	-	-	-	-	1	6	12	30	14	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasm of prostate (C61)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	414	-	-	-	-	-	-	3	23	91	165	132	-
2006 1/.....	368	-	-	-	-	-	1	-	34	82	138	113	-
2005 1/.....	372	-	-	-	-	1	-	7	31	74	142	117	-
2004 1/.....	362	-	-	-	-	-	1	4	18	60	156	123	-
2003 1/.....	328	-	-	-	-	-	-	3	18	58	144	105	-
2002.....	298	-	-	-	-	-	-	6	33	52	121	86	-
2001.....	296	-	-	-	-	-	-	1	15	50	135	95	-
2000.....	300	-	-	-	-	-	-	3	22	56	127	92	-
1999.....	304	-	-	-	-	-	1	1	21	55	119	107	-

Asian or Pacific Islander 3/, male

2007 1/.....	414	-	-	-	-	-	-	3	23	91	165	132	-
2006 1/.....	368	-	-	-	-	-	1	-	34	82	138	113	-
2005 1/.....	372	-	-	-	-	1	-	7	31	74	142	117	-
2004 1/.....	362	-	-	-	-	-	1	4	18	60	156	123	-
2003 1/.....	328	-	-	-	-	-	-	3	18	58	144	105	-
2002.....	298	-	-	-	-	-	-	6	33	52	121	86	-
2001.....	296	-	-	-	-	-	-	1	15	50	135	95	-
2000.....	300	-	-	-	-	-	-	3	22	56	127	92	-
1999.....	304	-	-	-	-	-	1	1	21	55	119	107	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of kidney and renal pelvis (C64-C65)													
All races, both sexes													
2007.....	12,703	2	9	25	34	50	240	1,243	2,584	3,159	3,539	1,818	-
2006.....	12,379	2	11	30	20	59	267	1,216	2,462	3,085	3,412	1,815	-
2005.....	12,517	8	16	30	17	49	294	1,277	2,514	3,143	3,510	1,659	-
2004.....	12,313	4	18	29	16	49	268	1,256	2,534	3,075	3,474	1,590	-
2003.....	12,286	1	14	39	24	56	321	1,201	2,498	3,090	3,475	1,567	-
2002.....	12,165	4	8	30	20	48	306	1,322	2,340	3,239	3,355	1,493	-
2001.....	12,078	1	22	35	20	52	358	1,311	2,328	3,221	3,306	1,424	-
2000.....	11,736	2	15	29	20	42	334	1,239	2,195	3,146	3,282	1,432	-
1999.....	11,116	2	19	33	17	54	342	1,147	2,111	3,076	3,010	1,305	-
All races, male													
2007.....	7,964	1	2	10	20	32	161	885	1,821	2,045	2,083	904	-
2006.....	7,699	2	4	18	11	37	188	858	1,733	2,004	1,964	880	-
2005.....	7,783	4	9	13	11	35	181	890	1,752	2,078	1,999	811	-
2004.....	7,640	2	9	11	9	34	183	889	1,786	2,015	1,993	709	-
2003.....	7,667	1	4	19	16	36	231	859	1,787	1,975	2,036	703	-
2002.....	7,587	1	-	19	7	31	217	947	1,658	2,087	1,924	696	-
2001.....	7,524	1	10	9	16	27	240	941	1,634	2,093	1,866	687	-
2000.....	7,249	1	8	15	10	26	228	880	1,530	2,010	1,865	676	-
1999.....	6,867	1	11	16	11	37	236	816	1,474	1,958	1,717	590	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

Malignant neoplasms of kidney and renal pelvis (C64-C65)

All races, female

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
2007.....	4,739	1	7	15	14	18	79	358	763	1,114	1,456	914	-
2006.....	4,680	-	7	12	9	22	79	358	729	1,081	1,448	935	-
2005.....	4,734	4	7	17	6	14	113	387	762	1,065	1,511	848	-
2004.....	4,673	2	9	18	7	15	85	367	748	1,060	1,481	881	-
2003.....	4,619	-	10	20	8	20	90	342	711	1,115	1,439	864	-
2002.....	4,578	3	8	11	13	17	89	375	682	1,152	1,431	797	-
2001.....	4,554	-	12	26	4	25	118	370	694	1,128	1,440	737	-
2000.....	4,487	1	7	14	10	16	106	359	665	1,136	1,417	756	-
1999.....	4,249	1	8	17	6	17	106	331	637	1,118	1,293	715	-

White, both sexes

2007 1/.....	11,142	1	7	17	17	35	184	1,018	2,231	2,790	3,189	1,653	-
2006 1/.....	10,961	2	10	22	11	31	201	1,004	2,164	2,736	3,119	1,661	-
2005 1/.....	11,091	7	12	21	7	23	234	1,076	2,186	2,806	3,204	1,515	-
2004 1/.....	10,988	4	13	24	11	28	213	1,077	2,252	2,730	3,161	1,475	-
2003 1/.....	10,930	1	14	30	16	35	247	1,018	2,175	2,772	3,183	1,439	-
2002.....	10,856	4	8	24	16	30	248	1,116	2,065	2,910	3,056	1,379	-
2001.....	10,720	-	16	23	8	31	274	1,111	2,030	2,908	3,010	1,309	-
2000.....	10,480	-	10	19	13	31	270	1,066	1,934	2,813	3,001	1,323	-
1999.....	9,878	2	12	25	10	32	271	962	1,868	2,757	2,752	1,187	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of kidney and renal pelvis (C64-C65)													
White, male													
2007 1/.....	6,996	-	2	5	9	22	123	728	1,571	1,802	1,900	834	-
2006 1/.....	6,811	2	4	11	5	14	141	716	1,515	1,781	1,810	812	-
2005 1/.....	6,888	3	8	7	4	16	140	744	1,527	1,857	1,831	751	-
2004 1/.....	6,839	2	6	9	7	17	147	770	1,596	1,797	1,827	661	-
2003 1/.....	6,840	1	4	14	11	20	180	736	1,574	1,773	1,873	654	-
2002.....	6,803	1	-	15	5	19	176	815	1,467	1,886	1,766	653	-
2001.....	6,695	-	6	5	5	16	192	807	1,426	1,901	1,697	640	-
2000.....	6,475	-	5	9	7	17	192	761	1,353	1,795	1,709	627	-
1999.....	6,112	1	8	11	7	20	189	682	1,310	1,759	1,587	538	-
White, female													
2007 1/.....	4,146	1	5	12	8	13	61	290	660	988	1,289	819	-
2006 1/.....	4,150	-	6	11	6	17	60	288	649	955	1,309	849	-
2005 1/.....	4,203	4	4	14	3	7	94	332	659	949	1,373	764	-
2004 1/.....	4,149	2	7	15	4	11	66	307	656	933	1,334	814	-
2003 1/.....	4,090	-	10	16	5	15	67	282	601	999	1,310	785	-
2002.....	4,053	3	8	9	11	11	72	301	598	1,024	1,290	726	-
2001.....	4,025	-	10	18	3	15	82	304	604	1,007	1,313	669	-
2000.....	4,005	-	5	10	6	14	78	305	581	1,018	1,292	696	-
1999.....	3,766	1	4	14	3	12	82	280	558	998	1,165	649	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of kidney and renal pelvis (C64-C65)

Black, both sexes

2007 1/.....	1,222	1	1	8	15	13	46	178	286	276	277	121	-
2006 1/.....	1,138	-	-	8	8	24	55	185	238	266	225	129	-
2005 1/.....	1,155	-	2	9	10	22	49	167	272	269	241	114	-
2004 1/.....	1,091	-	4	4	5	20	44	151	232	274	257	100	-
2003 1/.....	1,116	-	-	9	8	19	64	154	262	250	234	116	-
2002.....	1,069	-	-	6	3	17	46	164	226	260	252	95	-
2001.....	1,113	-	4	12	10	19	72	165	241	251	242	97	-
2000.....	1,044	1	5	8	6	10	52	145	214	273	235	95	-
1999.....	1,035	-	4	6	7	21	61	154	206	265	213	98	-

Black, male

2007 1/.....	744	1	-	5	11	9	30	126	199	176	138	49	-
2006 1/.....	714	-	-	7	5	20	42	124	173	169	119	55	-
2005 1/.....	723	-	-	6	7	17	35	122	190	176	128	42	-
2004 1/.....	655	-	3	1	2	16	29	102	156	173	134	39	-
2003 1/.....	677	-	-	5	5	14	44	108	170	158	131	42	-
2002.....	630	-	-	4	2	11	32	103	164	151	126	37	-
2001.....	678	-	2	4	10	10	40	112	168	154	137	41	-
2000.....	632	-	3	4	2	8	27	96	143	180	126	43	-
1999.....	622	-	2	3	4	16	42	109	140	160	104	42	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of kidney and renal pelvis (C64-C65)

Black, female

2007 1/.....	478	-	1	3	4	4	16	52	87	100	139	72	-
2006 1/.....	424	-	-	1	3	4	13	61	65	97	106	74	-
2005 1/.....	432	-	2	3	3	5	14	45	82	93	113	72	-
2004 1/.....	436	-	1	3	3	4	15	49	76	101	123	61	-
2003 1/.....	439	-	-	4	3	5	20	46	92	92	103	74	-
2002.....	439	-	-	2	1	6	14	61	62	109	126	58	-
2001.....	435	-	2	8	-	9	32	53	73	97	105	56	-
2000.....	412	1	2	4	4	2	25	49	71	93	109	52	-
1999.....	413	-	2	3	3	5	19	45	66	105	109	56	-

American Indian 2/, both sexes

2007 1/.....	102	-	-	-	1	-	2	22	23	29	18	7	-
2006 1/.....	79	-	1	-	-	-	4	8	18	24	18	6	-
2005 1/.....	87	-	-	-	-	2	6	11	24	24	17	3	-
2004 1/.....	91	-	-	1	-	-	2	15	22	30	18	3	-
2003 1/.....	79	-	-	-	-	-	5	12	20	23	17	2	-
2002.....	98	-	-	-	-	-	7	18	19	34	15	5	-
2001.....	87	1	1	-	1	-	4	15	25	22	13	5	-
2000.....	64	1	-	1	1	1	5	12	10	19	10	4	-
1999.....	78	-	2	-	-	-	3	12	16	21	20	4	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Malignant neoplasms of kidney and renal pelvis (C64-C65)

American Indian 2/, male

2007 1/.....	71	-	-	-	-	-	2	16	16	19	12	6	-
2006 1/.....	52	-	-	-	-	-	3	4	13	17	11	4	-
2005 1/.....	58	-	-	-	-	2	3	5	17	14	15	2	-
2004 1/.....	59	-	-	1	-	-	1	11	14	19	12	1	-
2003 1/.....	48	-	-	-	-	-	3	6	16	16	7	-	-
2002.....	63	-	-	-	-	-	4	11	11	26	10	1	-
2001.....	54	1	1	-	1	-	2	10	14	17	6	2	-
2000.....	43	1	-	1	1	1	4	8	8	11	7	1	-
1999.....	50	-	-	-	-	-	1	9	10	15	13	2	-

American Indian 2/, female

2007 1/.....	31	-	-	-	1	-	-	6	7	10	6	1	-
2006 1/.....	27	-	1	-	-	-	1	4	5	7	7	2	-
2005 1/.....	29	-	-	-	-	-	3	6	7	10	2	1	-
2004 1/.....	32	-	-	-	-	-	1	4	8	11	6	2	-
2003 1/.....	31	-	-	-	-	-	2	6	4	7	10	2	-
2002.....	35	-	-	-	-	-	3	7	8	8	5	4	-
2001.....	33	-	-	-	-	-	2	5	11	5	7	3	-
2000.....	21	-	-	-	-	-	1	4	2	8	3	3	-
1999.....	28	-	2	-	-	-	2	3	6	6	7	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of kidney and renal pelvis (C64-C65)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	237	-	1	-	1	2	8	25	44	64	55	37	-
2006 1/.....	201	-	-	-	1	4	7	19	42	59	50	19	-
2005 1/.....	184	1	2	-	-	2	5	23	32	44	48	27	-
2004 1/.....	143	-	1	-	-	1	9	13	28	41	38	12	-
2003 1/.....	161	-	-	-	-	2	5	17	41	45	41	10	-
2002.....	142	-	-	-	1	1	5	24	30	35	32	14	-
2001.....	158	-	1	-	1	2	8	20	32	40	41	13	-
2000.....	148	-	-	1	-	-	7	16	37	41	36	10	-
1999.....	125	-	1	2	-	1	7	19	21	33	25	16	-

Asian or Pacific Islander 3/, male

2007 1/.....	153	-	-	-	-	1	6	15	35	48	33	15	-
2006 1/.....	122	-	-	-	1	3	2	14	32	37	24	9	-
2005 1/.....	114	1	1	-	-	-	3	19	18	31	25	16	-
2004 1/.....	87	-	-	-	-	1	6	6	20	26	20	8	-
2003 1/.....	102	-	-	-	-	2	4	9	27	28	25	7	-
2002.....	91	-	-	-	-	1	5	18	16	24	22	5	-
2001.....	97	-	1	-	-	1	6	12	26	21	26	4	-
2000.....	99	-	-	1	-	-	5	15	26	24	23	5	-
1999.....	83	-	1	2	-	1	4	16	14	24	13	8	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of kidney and renal pelvis (C64-C65)

Asian or Pacific Islander 3/, female

2007 1/.....	84	-	1	-	1	1	2	10	9	16	22	22	-
2006 1/.....	79	-	-	-	-	1	5	5	10	22	26	10	-
2005 1/.....	70	-	1	-	-	2	2	4	14	13	23	11	-
2004 1/.....	56	-	1	-	-	-	3	7	8	15	18	4	-
2003 1/.....	59	-	-	-	-	-	1	8	14	17	16	3	-
2002.....	51	-	-	-	1	-	-	6	14	11	10	9	-
2001.....	61	-	-	-	1	1	2	8	6	19	15	9	-
2000.....	49	-	-	-	-	-	2	1	11	17	13	5	-
1999.....	42	-	-	-	-	-	3	3	7	9	12	8	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of bladder (C67)													
All races, both sexes													
2007.....	13,843	1	-	-	-	7	93	570	1,564	2,817	5,009	3,782	-
2006.....	13,474	-	-	-	1	18	104	528	1,568	2,810	4,874	3,571	-
2005.....	13,253	-	-	-	-	15	122	523	1,482	2,772	4,859	3,480	-
2004.....	13,030	-	-	1	1	12	120	545	1,452	2,753	4,847	3,299	-
2003.....	12,483	-	-	-	1	4	98	522	1,391	2,736	4,679	3,052	-
2002.....	12,628	-	-	3	1	14	123	503	1,310	2,907	4,707	3,059	1
2001.....	12,225	-	-	1	3	15	128	475	1,265	2,814	4,488	3,036	-
2000.....	12,002	-	-	1	3	10	111	440	1,223	2,855	4,407	2,952	-
1999.....	11,911	-	-	1	1	16	100	463	1,244	2,923	4,317	2,845	1
All races, male													
2007.....	9,644	1	-	-	-	4	57	406	1,176	2,103	3,540	2,357	-
2006.....	9,412	-	-	-	1	11	69	387	1,192	2,059	3,472	2,221	-
2005.....	9,190	-	-	-	-	9	81	358	1,115	2,072	3,436	2,119	-
2004.....	8,995	-	-	1	1	7	69	411	1,103	2,021	3,376	2,006	-
2003.....	8,576	-	-	-	1	2	64	401	1,049	1,998	3,254	1,807	-
2002.....	8,504	-	-	1	-	9	78	352	984	2,089	3,199	1,791	1
2001.....	8,343	-	-	-	3	8	70	351	956	2,088	3,116	1,751	-
2000.....	8,163	-	-	-	3	2	62	322	916	2,094	3,054	1,710	-
1999.....	8,105	-	-	1	-	9	63	350	933	2,132	3,008	1,608	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of bladder (C67)													
All races, female													
2007.....	4,199	-	-	-	-	3	36	164	388	714	1,469	1,425	-
2006.....	4,062	-	-	-	-	7	35	141	376	751	1,402	1,350	-
2005.....	4,063	-	-	-	-	6	41	165	367	700	1,423	1,361	-
2004.....	4,035	-	-	-	-	5	51	134	349	732	1,471	1,293	-
2003.....	3,907	-	-	-	-	2	34	121	342	738	1,425	1,245	-
2002.....	4,124	-	-	2	1	5	45	151	326	818	1,508	1,268	-
2001.....	3,882	-	-	1	-	7	58	124	309	726	1,372	1,285	-
2000.....	3,839	-	-	1	-	8	49	118	307	761	1,353	1,242	-
1999.....	3,806	-	-	-	1	7	37	113	311	791	1,309	1,237	-
White, both sexes													
2007 1/.....	12,620	-	-	-	-	5	73	476	1,371	2,562	4,612	3,521	-
2006 1/.....	12,286	-	-	-	1	13	82	458	1,392	2,522	4,495	3,323	-
2005 1/.....	12,100	-	-	-	-	13	110	440	1,320	2,509	4,498	3,210	-
2004 1/.....	11,957	-	-	1	-	9	85	459	1,319	2,514	4,523	3,047	-
2003 1/.....	11,409	-	-	-	1	4	75	433	1,253	2,496	4,310	2,837	-
2002.....	11,512	-	-	3	-	10	105	443	1,170	2,625	4,344	2,811	1
2001.....	11,180	-	-	1	3	14	104	407	1,120	2,573	4,138	2,820	-
2000.....	11,036	-	-	1	2	9	99	391	1,109	2,605	4,056	2,764	-
1999.....	10,863	-	-	1	-	10	80	397	1,093	2,643	4,008	2,630	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of bladder (C67)													
White, male													
2007 1/.....	8,950	-	-	-	-	3	48	354	1,044	1,937	3,327	2,237	-
2006 1/.....	8,735	-	-	-	1	8	56	340	1,067	1,896	3,270	2,097	-
2005 1/.....	8,523	-	-	-	-	8	74	314	1,010	1,907	3,225	1,985	-
2004 1/.....	8,407	-	-	1	-	5	48	353	1,015	1,889	3,200	1,896	-
2003 1/.....	7,965	-	-	-	1	2	49	344	962	1,848	3,047	1,712	-
2002.....	7,908	-	-	1	-	6	68	319	906	1,931	2,990	1,686	1
2001.....	7,780	-	-	-	3	8	57	308	869	1,946	2,929	1,660	-
2000.....	7,620	-	-	-	2	2	57	291	843	1,949	2,861	1,615	-
1999.....	7,520	-	-	1	-	5	49	297	843	1,977	2,828	1,519	1
White, female													
2007 1/.....	3,670	-	-	-	-	2	25	122	327	625	1,285	1,284	-
2006 1/.....	3,551	-	-	-	-	5	26	118	325	626	1,225	1,226	-
2005 1/.....	3,577	-	-	-	-	5	36	126	310	602	1,273	1,225	-
2004 1/.....	3,550	-	-	-	-	4	37	106	304	625	1,323	1,151	-
2003 1/.....	3,444	-	-	-	-	2	26	89	291	648	1,263	1,125	-
2002.....	3,604	-	-	2	-	4	37	124	264	694	1,354	1,125	-
2001.....	3,400	-	-	1	-	6	47	99	251	627	1,209	1,160	-
2000.....	3,416	-	-	1	-	7	42	100	266	656	1,195	1,149	-
1999.....	3,343	-	-	-	-	5	31	100	250	666	1,180	1,111	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasm of bladder (C67)

Black, both sexes

2007 1/.....	1,034	1	-	-	-	1	17	84	165	220	330	216	-
2006 1/.....	1,007	-	-	-	-	4	17	64	154	242	316	210	-
2005 1/.....	961	-	-	-	-	2	12	74	141	218	299	215	-
2004 1/.....	913	-	-	-	1	3	27	77	118	203	271	213	-
2003 1/.....	914	-	-	-	-	-	21	79	109	205	317	183	-
2002.....	940	-	-	-	-	4	13	49	120	242	299	213	-
2001.....	877	-	-	-	-	1	23	56	125	198	295	179	-
2000.....	852	-	-	-	1	-	10	44	100	220	318	159	-
1999.....	918	-	-	-	1	4	18	62	133	247	264	189	-

Black, male

2007 1/.....	565	1	-	-	-	1	8	46	109	139	168	93	-
2006 1/.....	558	-	-	-	-	2	11	44	108	134	155	104	-
2005 1/.....	546	-	-	-	-	1	7	39	90	134	166	109	-
2004 1/.....	478	-	-	-	1	2	18	52	76	105	142	82	-
2003 1/.....	502	-	-	-	-	-	13	49	66	124	173	77	-
2002.....	481	-	-	-	-	3	7	26	66	128	166	85	-
2001.....	447	-	-	-	-	-	13	36	72	111	146	69	-
2000.....	466	-	-	-	1	-	4	27	63	120	174	77	-
1999.....	496	-	-	-	-	2	13	49	78	130	150	74	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Malignant neoplasm of bladder (C67)

Black, female

2007 1/.....	469	-	-	-	-	-	9	38	56	81	162	123	-
2006 1/.....	449	-	-	-	-	2	6	20	46	108	161	106	-
2005 1/.....	415	-	-	-	-	1	5	35	51	84	133	106	-
2004 1/.....	435	-	-	-	-	1	9	25	42	98	129	131	-
2003 1/.....	412	-	-	-	-	-	8	30	43	81	144	106	-
2002.....	459	-	-	-	-	1	6	23	54	114	133	128	-
2001.....	430	-	-	-	-	1	10	20	53	87	149	110	-
2000.....	386	-	-	-	-	-	6	17	37	100	144	82	-
1999.....	422	-	-	-	1	2	5	13	55	117	114	115	-

American Indian 2/, both sexes

2007 1/.....	31	-	-	-	-	-	-	3	5	5	12	6	-
2006 1/.....	37	-	-	-	-	1	3	2	4	7	14	6	-
2005 1/.....	30	-	-	-	-	-	-	2	6	9	10	3	-
2004 1/.....	20	-	-	-	-	-	1	4	3	4	4	4	-
2003 1/.....	30	-	-	-	-	-	-	2	10	4	11	3	-
2002.....	31	-	-	-	1	-	-	2	4	9	8	7	-
2001.....	26	-	-	-	-	-	-	4	4	8	7	3	-
2000.....	16	-	-	-	-	-	1	1	2	3	7	2	-
1999.....	21	-	-	-	-	1	-	-	5	10	2	3	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasm of bladder (C67)

American Indian 2/, male

2007 1/.....	22	-	-	-	-	-	-	1	3	4	9	5	-
2006 1/.....	26	-	-	-	-	1	1	-	4	5	12	3	-
2005 1/.....	18	-	-	-	-	-	-	2	3	7	5	1	-
2004 1/.....	15	-	-	-	-	-	-	3	3	3	4	2	-
2003 1/.....	16	-	-	-	-	-	-	-	6	3	5	2	-
2002.....	17	-	-	-	-	-	-	2	1	7	4	3	-
2001.....	16	-	-	-	-	-	-	2	2	4	6	2	-
2000.....	11	-	-	-	-	-	1	-	2	3	3	2	-
1999.....	16	-	-	-	-	1	-	-	3	9	1	2	-

American Indian 2/, female

2007 1/.....	9	-	-	-	-	-	-	2	2	1	3	1	-
2006 1/.....	11	-	-	-	-	-	2	2	-	2	2	3	-
2005 1/.....	12	-	-	-	-	-	-	-	3	2	5	2	-
2004 1/.....	5	-	-	-	-	-	1	1	-	1	-	2	-
2003 1/.....	14	-	-	-	-	-	-	2	4	1	6	1	-
2002.....	14	-	-	-	1	-	-	-	3	2	4	4	-
2001.....	10	-	-	-	-	-	-	2	2	4	1	1	-
2000.....	5	-	-	-	-	-	-	1	-	-	4	-	-
1999.....	5	-	-	-	-	-	-	-	2	1	1	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasm of bladder (C67)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	158	-	-	-	-	1	3	7	23	30	55	39	-
2006 1/.....	144	-	-	-	-	-	2	4	18	39	49	32	-
2005 1/.....	162	-	-	-	-	-	-	7	15	36	52	52	-
2004 1/.....	140	-	-	-	-	-	7	5	12	32	49	35	-
2003 1/.....	130	-	-	-	-	-	2	8	19	31	41	29	-
2002.....	145	-	-	-	-	-	5	9	16	31	56	28	-
2001.....	142	-	-	-	-	-	1	8	16	35	48	34	-
2000.....	98	-	-	-	-	1	1	4	12	27	26	27	-
1999.....	109	-	-	-	-	1	2	4	13	23	43	23	-

Asian or Pacific Islander 3/, male

2007 1/.....	107	-	-	-	-	-	1	5	20	23	36	22	-
2006 1/.....	93	-	-	-	-	-	1	3	13	24	35	17	-
2005 1/.....	103	-	-	-	-	-	-	3	12	24	40	24	-
2004 1/.....	95	-	-	-	-	-	3	3	9	24	30	26	-
2003 1/.....	93	-	-	-	-	-	2	8	15	23	29	16	-
2002.....	98	-	-	-	-	-	3	5	11	23	39	17	-
2001.....	100	-	-	-	-	-	-	5	13	27	35	20	-
2000.....	66	-	-	-	-	-	-	4	8	22	16	16	-
1999.....	73	-	-	-	-	1	1	4	9	16	29	13	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasm of bladder (C67)

Asian or Pacific Islander 3/, female

2007 1/.....	51	-	-	-	-	1	2	2	3	7	19	17	-
2006 1/.....	51	-	-	-	-	-	1	1	5	15	14	15	-
2005 1/.....	59	-	-	-	-	-	-	4	3	12	12	28	-
2004 1/.....	45	-	-	-	-	-	4	2	3	8	19	9	-
2003 1/.....	37	-	-	-	-	-	-	-	4	8	12	13	-
2002.....	47	-	-	-	-	-	2	4	5	8	17	11	-
2001.....	42	-	-	-	-	-	1	3	3	8	13	14	-
2000.....	32	-	-	-	-	1	1	-	4	5	10	11	-
1999.....	36	-	-	-	-	-	1	-	4	7	14	10	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

All races, both sexes

2007.....	13,234	15	109	302	204	369	845	1,916	2,975	3,002	2,608	889	-
2006.....	12,886	16	101	305	206	377	863	1,951	2,925	2,781	2,476	885	-
2005.....	13,152	18	84	324	234	379	948	1,943	2,839	2,943	2,633	807	-
2004.....	12,829	19	102	325	223	359	904	1,902	2,750	2,895	2,543	807	-
2003.....	12,901	18	104	336	236	405	1,079	1,962	2,698	2,836	2,510	717	-
2002.....	12,830	12	114	337	210	388	1,009	2,058	2,560	2,954	2,435	753	-
2001.....	12,609	16	100	300	224	416	1,028	1,951	2,464	2,823	2,566	721	-
2000.....	12,655	20	104	316	225	406	1,096	1,924	2,337	3,032	2,478	717	-
1999.....	12,765	10	113	308	218	444	1,065	1,880	2,453	3,119	2,492	663	-

All races, male

2007.....	7,315	6	62	144	113	229	527	1,150	1,782	1,645	1,282	375	-
2006.....	7,198	5	56	166	118	215	535	1,161	1,737	1,575	1,272	358	-
2005.....	7,317	12	38	171	144	248	604	1,160	1,653	1,643	1,335	309	-
2004.....	7,074	10	56	173	133	218	560	1,191	1,653	1,551	1,237	292	-
2003.....	7,122	9	56	175	136	232	639	1,233	1,530	1,556	1,267	289	-
2002.....	7,125	8	63	191	136	223	618	1,225	1,494	1,678	1,210	279	-
2001.....	7,015	8	52	145	129	262	638	1,173	1,470	1,589	1,288	261	-
2000.....	7,011	10	58	176	132	264	704	1,148	1,379	1,678	1,187	275	-
1999.....	7,001	8	56	154	131	276	638	1,120	1,417	1,728	1,226	247	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

All races, female

2007.....	5,919	9	47	158	91	140	318	766	1,193	1,357	1,326	514	-
2006.....	5,688	11	45	139	88	162	328	790	1,188	1,206	1,204	527	-
2005.....	5,835	6	46	153	90	131	344	783	1,186	1,300	1,298	498	-
2004.....	5,755	9	46	152	90	141	344	711	1,097	1,344	1,306	515	-
2003.....	5,779	9	48	161	100	173	440	729	1,168	1,280	1,243	428	-
2002.....	5,705	4	51	146	74	165	391	833	1,066	1,276	1,225	474	-
2001.....	5,594	8	48	155	95	154	390	778	994	1,234	1,278	460	-
2000.....	5,644	10	46	140	93	142	392	776	958	1,354	1,291	442	-
1999.....	5,764	2	57	154	87	168	427	760	1,036	1,391	1,266	416	-

White, both sexes

2007 1/.....	12,177	13	93	252	170	316	754	1,720	2,747	2,790	2,486	836	-
2006 1/.....	11,850	13	77	246	164	327	770	1,788	2,705	2,583	2,347	830	-
2005 1/.....	12,079	15	68	257	197	329	850	1,766	2,624	2,744	2,472	757	-
2004 1/.....	11,793	18	82	249	184	306	803	1,710	2,564	2,711	2,407	759	-
2003 1/.....	11,849	15	85	271	197	355	976	1,775	2,496	2,637	2,375	667	-
2002.....	11,794	7	80	265	174	338	922	1,882	2,359	2,764	2,292	711	-
2001.....	11,638	12	84	244	183	374	942	1,783	2,270	2,628	2,440	678	-
2000.....	11,672	16	82	255	183	360	969	1,771	2,161	2,864	2,340	671	-
1999.....	11,819	7	96	242	171	403	966	1,725	2,276	2,945	2,363	625	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

White, male

2007 1/.....	6,759	5	53	122	99	192	470	1,042	1,648	1,535	1,233	360	-
2006 1/.....	6,650	5	47	140	93	192	485	1,068	1,610	1,468	1,206	336	-
2005 1/.....	6,715	10	28	132	121	214	549	1,055	1,526	1,539	1,249	292	-
2004 1/.....	6,508	9	44	134	113	186	492	1,068	1,542	1,474	1,165	281	-
2003 1/.....	6,543	8	48	139	112	206	572	1,120	1,423	1,448	1,196	271	-
2002.....	6,554	5	46	143	117	191	572	1,132	1,369	1,577	1,136	266	-
2001.....	6,498	6	46	113	107	238	589	1,081	1,355	1,500	1,222	241	-
2000.....	6,486	9	46	141	109	227	630	1,058	1,284	1,593	1,128	261	-
1999.....	6,492	6	49	125	101	250	577	1,023	1,316	1,641	1,169	235	-

White, female

2007 1/.....	5,418	8	40	130	71	124	284	678	1,099	1,255	1,253	476	-
2006 1/.....	5,200	8	30	106	71	135	285	720	1,095	1,115	1,141	494	-
2005 1/.....	5,364	5	40	125	76	115	301	711	1,098	1,205	1,223	465	-
2004 1/.....	5,285	9	38	115	71	120	311	642	1,022	1,237	1,242	478	-
2003 1/.....	5,306	7	37	132	85	149	404	655	1,073	1,189	1,179	396	-
2002.....	5,240	2	34	122	57	147	350	750	990	1,187	1,156	445	-
2001.....	5,140	6	38	131	76	136	353	702	915	1,128	1,218	437	-
2000.....	5,186	7	36	114	74	133	339	713	877	1,271	1,212	410	-
1999.....	5,327	1	47	117	70	153	389	702	960	1,304	1,194	390	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Black, both sexes

2007 1/.....	791	2	9	40	21	33	56	153	177	174	90	36	-
2006 1/.....	765	3	18	47	29	25	66	130	170	144	96	37	-
2005 1/.....	793	3	13	52	24	30	65	138	164	147	122	35	-
2004 1/.....	765	-	18	50	33	37	72	138	142	139	99	37	-
2003 1/.....	811	2	17	48	34	36	83	147	153	152	103	36	-
2002.....	781	3	27	54	28	33	68	127	149	146	114	32	-
2001.....	741	1	10	44	29	28	63	133	148	153	94	38	-
2000.....	772	2	18	52	35	36	91	122	139	134	109	34	-
1999.....	764	3	12	55	37	36	77	129	142	135	107	31	-

Black, male

2007 1/.....	404	1	4	15	6	21	36	86	103	87	35	10	-
2006 1/.....	409	-	7	20	20	13	39	75	101	76	46	12	-
2005 1/.....	440	2	8	28	14	23	36	85	95	76	64	9	-
2004 1/.....	409	-	10	24	14	20	50	88	87	57	51	8	-
2003 1/.....	447	1	6	27	21	19	56	89	82	85	54	7	-
2002.....	409	2	12	35	14	22	35	57	95	73	57	7	-
2001.....	382	-	3	25	16	14	33	75	89	70	42	15	-
2000.....	414	1	10	31	21	28	54	73	77	67	43	9	-
1999.....	404	2	5	25	24	23	48	82	76	68	41	10	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Black, female

2007 1/.....	387	1	5	25	15	12	20	67	74	87	55	26	-
2006 1/.....	356	3	11	27	9	12	27	55	69	68	50	25	-
2005 1/.....	353	1	5	24	10	7	29	53	69	71	58	26	-
2004 1/.....	356	-	8	26	19	17	22	50	55	82	48	29	-
2003 1/.....	364	1	11	21	13	17	27	58	71	67	49	29	-
2002.....	372	1	15	19	14	11	33	70	54	73	57	25	-
2001.....	359	1	7	19	13	14	30	58	59	83	52	23	-
2000.....	358	1	8	21	14	8	37	49	62	67	66	25	-
1999.....	360	1	7	30	13	13	29	47	66	67	66	21	-

American Indian 2/, both sexes

2007 1/.....	44	-	-	2	3	3	6	6	8	5	8	3	-
2006 1/.....	40	-	3	2	1	4	4	3	7	10	5	1	-
2005 1/.....	44	-	1	2	3	2	7	5	15	2	6	1	-
2004 1/.....	34	1	1	2	2	1	1	13	5	3	5	-	-
2003 1/.....	45	-	1	4	2	1	3	11	9	8	3	3	-
2002.....	41	1	1	7	2	4	3	8	5	6	3	1	-
2001.....	43	2	1	-	1	2	7	9	8	8	5	-	-
2000.....	41	-	2	4	2	1	11	5	3	4	8	1	-
1999.....	34	-	1	1	2	-	9	4	8	6	2	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

American Indian 2/, male

2007 1/.....	24	-	-	2	2	3	4	3	5	3	1	1	-
2006 1/.....	21	-	1	1	1	1	2	2	2	6	5	-	-
2005 1/.....	28	-	1	2	2	2	4	2	11	-	3	1	-
2004 1/.....	25	1	1	2	2	1	1	10	4	1	2	-	-
2003 1/.....	26	-	1	2	1	-	2	8	4	3	3	2	-
2002.....	23	-	-	4	1	1	2	6	4	4	1	-	-
2001.....	25	2	1	-	-	2	6	1	5	4	4	-	-
2000.....	20	-	1	2	1	-	5	2	1	2	5	1	-
1999.....	19	-	-	-	1	-	7	1	6	2	1	1	-

American Indian 2/, female

2007 1/.....	20	-	-	-	1	-	2	3	3	2	7	2	-
2006 1/.....	19	-	2	1	-	3	2	1	5	4	-	1	-
2005 1/.....	16	-	-	-	1	-	3	3	4	2	3	-	-
2004 1/.....	9	-	-	-	-	-	-	3	1	2	3	-	-
2003 1/.....	19	-	-	2	1	1	1	3	5	5	-	1	-
2002.....	18	1	1	3	1	3	1	2	1	2	2	1	-
2001.....	18	-	-	-	1	-	1	8	3	4	1	-	-
2000.....	21	-	1	2	1	1	6	3	2	2	3	-	-
1999.....	15	-	1	1	1	-	2	3	2	4	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	222	-	7	8	10	17	29	37	43	33	24	14	-
2006 1/.....	231	-	3	10	12	21	23	30	43	44	28	17	-
2005 1/.....	236	-	2	13	10	18	26	34	36	50	33	14	-
2004 1/.....	237	-	1	24	4	15	28	41	39	42	32	11	-
2003 1/.....	196	1	1	13	3	13	17	29	40	39	29	11	-
2002.....	214	1	6	11	6	13	16	41	47	38	26	9	-
2001.....	187	1	5	12	11	12	16	26	38	34	27	5	-
2000.....	170	2	2	5	5	9	25	26	34	30	21	11	-
1999.....	148	-	4	10	8	5	13	22	27	33	20	6	-

Asian or Pacific Islander 3/, male

2007 1/.....	128	-	5	5	6	13	17	19	26	20	13	4	-
2006 1/.....	118	-	1	5	4	9	9	16	24	25	15	10	-
2005 1/.....	134	-	1	9	7	9	15	18	21	28	19	7	-
2004 1/.....	132	-	1	13	4	11	17	25	20	19	19	3	-
2003 1/.....	106	-	1	7	2	7	9	16	21	20	14	9	-
2002.....	139	1	5	9	4	9	9	30	26	24	16	6	-
2001.....	110	-	2	7	6	8	10	16	21	15	20	5	-
2000.....	91	-	1	2	1	9	15	15	17	16	11	4	-
1999.....	86	-	2	4	5	3	6	14	19	17	15	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Asian or Pacific Islander 3/, female

2007 1/.....	94	-	2	3	4	4	12	18	17	13	11	10	-
2006 1/.....	113	-	2	5	8	12	14	14	19	19	13	7	-
2005 1/.....	102	-	1	4	3	9	11	16	15	22	14	7	-
2004 1/.....	105	-	-	11	-	4	11	16	19	23	13	8	-
2003 1/.....	90	1	-	6	1	6	8	13	19	19	15	2	-
2002.....	75	-	1	2	2	4	7	11	21	14	10	3	-
2001.....	77	1	3	5	5	4	6	10	17	19	7	-	-
2000.....	79	2	1	3	4	-	10	11	17	14	10	7	-
1999.....	62	-	2	6	3	2	7	8	8	16	5	5	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)													
All races, both sexes													
2007.....	54,991	26	111	352	630	771	1,464	3,606	7,694	12,223	17,884	10,228	2
2006.....	55,045	32	132	298	662	828	1,536	3,696	7,695	12,335	17,974	9,856	1
2005.....	55,028	24	128	335	655	811	1,498	3,687	7,623	12,487	18,109	9,671	-
2004.....	54,645	25	122	359	645	822	1,514	3,741	7,491	12,588	18,132	9,206	-
2003.....	55,679	31	138	394	606	865	1,728	3,785	7,471	12,885	18,442	9,334	-
2002.....	56,225	28	119	379	703	916	1,751	3,960	7,500	13,508	18,289	9,072	-
2001.....	56,341	27	157	367	705	988	1,857	4,064	7,455	13,659	18,257	8,805	-
2000.....	56,518	39	146	372	718	918	1,853	4,025	7,284	14,041	18,448	8,674	-
1999.....	56,318	30	137	351	720	960	1,913	3,950	7,554	14,332	18,089	8,282	-
All races, male													
2007.....	30,198	18	56	207	377	461	889	2,199	4,631	7,175	9,681	4,503	1
2006.....	30,007	18	71	177	435	500	906	2,234	4,694	7,141	9,699	4,131	1
2005.....	30,031	12	84	206	411	464	880	2,175	4,543	7,385	9,662	4,209	-
2004.....	29,522	10	72	214	399	489	915	2,253	4,497	7,368	9,462	3,843	-
2003.....	30,076	10	80	225	375	535	1,043	2,294	4,557	7,550	9,524	3,883	-
2002.....	30,302	15	63	217	417	545	1,038	2,341	4,470	7,903	9,553	3,740	-
2001.....	30,060	12	82	217	426	574	1,133	2,442	4,412	7,859	9,327	3,576	-
2000.....	29,888	21	76	223	455	539	1,075	2,399	4,205	8,038	9,399	3,458	-
1999.....	29,721	12	75	206	437	557	1,144	2,368	4,426	8,067	9,100	3,329	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)													
All races, female													
2007.....	24,793	8	55	145	253	310	575	1,407	3,063	5,048	8,203	5,725	1
2006.....	25,038	14	61	121	227	328	630	1,462	3,001	5,194	8,275	5,725	-
2005.....	24,997	12	44	129	244	347	618	1,512	3,080	5,102	8,447	5,462	-
2004.....	25,123	15	50	145	246	333	599	1,488	2,994	5,220	8,670	5,363	-
2003.....	25,603	21	58	169	231	330	685	1,491	2,914	5,335	8,918	5,451	-
2002.....	25,923	13	56	162	286	371	713	1,619	3,030	5,605	8,736	5,332	-
2001.....	26,281	15	75	150	279	414	724	1,622	3,043	5,800	8,930	5,229	-
2000.....	26,630	18	70	149	263	379	778	1,626	3,079	6,003	9,049	5,216	-
1999.....	26,597	18	62	145	283	403	769	1,582	3,128	6,265	8,989	4,953	-
White, both sexes													
2007 1/.....	48,377	21	84	283	493	599	1,108	2,855	6,436	10,664	16,302	9,530	2
2006 1/.....	48,549	26	108	236	523	647	1,191	2,953	6,447	10,812	16,460	9,145	1
2005 1/.....	48,703	20	93	274	522	583	1,172	2,912	6,457	10,966	16,663	9,041	-
2004 1/.....	48,240	20	94	289	515	626	1,200	3,011	6,368	10,995	16,582	8,540	-
2003 1/.....	49,420	29	115	314	483	670	1,366	3,055	6,429	11,362	16,892	8,705	-
2002.....	49,943	19	98	303	575	710	1,405	3,185	6,366	11,977	16,836	8,469	-
2001.....	49,989	22	136	281	562	753	1,470	3,344	6,318	12,108	16,791	8,204	-
2000.....	50,394	36	121	306	549	707	1,485	3,297	6,255	12,558	17,073	8,007	-
1999.....	50,189	24	118	271	564	751	1,539	3,281	6,440	12,826	16,687	7,688	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)													
White, male													
2007 1/.....	26,765	14	39	167	296	365	687	1,766	3,951	6,345	8,921	4,213	1
2006 1/.....	26,521	14	57	137	348	397	702	1,799	3,970	6,312	8,953	3,831	1
2005 1/.....	26,707	10	60	164	332	334	703	1,729	3,895	6,566	8,936	3,978	-
2004 1/.....	26,204	8	52	175	319	386	741	1,824	3,880	6,491	8,734	3,594	-
2003 1/.....	26,943	8	67	184	301	419	836	1,869	3,967	6,783	8,857	3,652	-
2002.....	27,087	11	53	170	352	426	832	1,923	3,843	7,094	8,870	3,513	-
2001.....	26,767	9	69	157	346	451	892	2,042	3,770	7,040	8,641	3,350	-
2000.....	26,766	19	64	180	351	420	890	1,977	3,639	7,249	8,760	3,217	-
1999.....	26,558	12	63	160	349	431	926	1,984	3,817	7,287	8,444	3,085	-
White, female													
2007 1/.....	21,612	7	45	116	197	234	421	1,089	2,485	4,319	7,381	5,317	1
2006 1/.....	22,028	12	51	99	175	250	489	1,154	2,477	4,500	7,507	5,314	-
2005 1/.....	21,996	10	33	110	190	249	469	1,183	2,562	4,400	7,727	5,063	-
2004 1/.....	22,036	12	42	114	196	240	459	1,187	2,488	4,504	7,848	4,946	-
2003 1/.....	22,477	21	48	130	182	251	530	1,186	2,462	4,579	8,035	5,053	-
2002.....	22,856	8	45	133	223	284	573	1,262	2,523	4,883	7,966	4,956	-
2001.....	23,222	13	67	124	216	302	578	1,302	2,548	5,068	8,150	4,854	-
2000.....	23,628	17	57	126	198	287	595	1,320	2,616	5,309	8,313	4,790	-
1999.....	23,631	12	55	111	215	320	613	1,297	2,623	5,539	8,243	4,603	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Black, both sexes

2007 1/.....	5,255	3	20	51	98	133	294	619	1,030	1,257	1,195	555	-
2006 1/.....	5,267	4	17	48	107	135	287	627	1,055	1,242	1,192	553	-
2005 1/.....	5,046	4	27	45	104	182	261	623	936	1,210	1,142	512	-
2004 1/.....	5,233	3	19	49	106	164	256	621	940	1,301	1,228	546	-
2003 1/.....	5,103	1	14	57	88	156	301	624	858	1,250	1,238	516	-
2002.....	5,151	5	16	53	90	157	276	631	949	1,282	1,179	513	-
2001.....	5,227	4	14	65	111	183	323	589	930	1,302	1,199	507	-
2000.....	5,071	1	20	43	132	160	306	623	875	1,241	1,109	561	-
1999.....	5,088	4	15	63	134	160	314	561	927	1,253	1,149	508	-

Black, male

2007 1/.....	2,663	2	13	31	56	72	168	358	549	650	550	214	-
2006 1/.....	2,813	2	10	32	65	75	173	368	617	685	568	218	-
2005 1/.....	2,633	2	18	32	64	100	142	364	527	651	556	177	-
2004 1/.....	2,650	-	14	26	63	87	139	361	513	711	541	195	-
2003 1/.....	2,537	1	8	31	52	88	172	369	493	618	528	177	-
2002.....	2,622	2	7	33	45	93	168	345	523	685	532	189	-
2001.....	2,676	3	9	44	61	100	200	330	520	685	542	182	-
2000.....	2,558	-	9	26	80	91	153	351	485	656	512	195	-
1999.....	2,590	-	9	39	74	98	182	324	503	649	524	188	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Black, female

2007 1/.....	2,592	1	7	20	42	61	126	261	481	607	645	341	-
2006 1/.....	2,454	2	7	16	42	60	114	259	438	557	624	335	-
2005 1/.....	2,413	2	9	13	40	82	119	259	409	559	586	335	-
2004 1/.....	2,583	3	5	23	43	77	117	260	427	590	687	351	-
2003 1/.....	2,566	-	6	26	36	68	129	255	365	632	710	339	-
2002.....	2,529	3	9	20	45	64	108	286	426	597	647	324	-
2001.....	2,551	1	5	21	50	83	123	259	410	617	657	325	-
2000.....	2,513	1	11	17	52	69	153	272	390	585	597	366	-
1999.....	2,498	4	6	24	60	62	132	237	424	604	625	320	-

American Indian 2/, both sexes

2007 1/.....	209	1	-	3	12	5	8	23	38	54	49	16	-
2006 1/.....	194	-	1	-	7	6	12	24	34	46	44	20	-
2005 1/.....	222	-	2	1	6	6	10	29	53	62	38	15	-
2004 1/.....	199	-	1	4	5	6	12	18	39	48	52	14	-
2003 1/.....	210	-	3	3	12	5	12	19	41	48	50	17	-
2002.....	171	-	1	2	11	7	14	19	35	34	36	12	-
2001.....	225	-	1	4	10	14	18	24	46	55	36	17	-
2000.....	199	1	1	2	5	9	9	20	34	46	48	24	-
1999.....	161	1	1	2	3	7	9	15	23	50	37	13	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

American Indian 2/, male

2007 1/.....	118	1	-	1	9	5	4	15	19	33	26	5	-
2006 1/.....	105	-	1	-	5	3	5	17	23	22	20	9	-
2005 1/.....	121	-	2	1	3	4	7	15	23	38	23	5	-
2004 1/.....	112	-	1	3	2	3	6	14	22	29	29	3	-
2003 1/.....	99	-	3	-	9	5	6	11	18	25	16	6	-
2002.....	89	-	1	1	8	3	10	11	16	17	19	3	-
2001.....	113	-	1	2	7	9	9	13	26	28	13	5	-
2000.....	101	1	1	-	3	6	4	14	17	26	21	8	-
1999.....	92	-	1	2	1	4	4	10	12	29	21	8	-

American Indian 2/, female

2007 1/.....	91	-	-	2	3	-	4	8	19	21	23	11	-
2006 1/.....	89	-	-	-	2	3	7	7	11	24	24	11	-
2005 1/.....	101	-	-	-	3	2	3	14	30	24	15	10	-
2004 1/.....	87	-	-	1	3	3	6	4	17	19	23	11	-
2003 1/.....	111	-	-	3	3	-	6	8	23	23	34	11	-
2002.....	82	-	-	1	3	4	4	8	19	17	17	9	-
2001.....	112	-	-	2	3	5	9	11	20	27	23	12	-
2000.....	98	-	-	2	2	3	5	6	17	20	27	16	-
1999.....	69	1	-	-	2	3	5	5	11	21	16	5	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	1,150	1	7	15	27	34	54	109	190	248	338	127	-
2006 1/.....	1,035	2	6	14	25	40	46	92	159	235	278	138	-
2005 1/.....	1,057	-	6	15	23	40	55	123	177	249	266	103	-
2004 1/.....	973	2	8	17	19	26	46	91	144	244	270	106	-
2003 1/.....	946	1	6	20	23	34	49	87	143	225	262	96	-
2002.....	960	4	4	21	27	42	56	125	150	215	238	78	-
2001.....	900	1	6	17	22	38	46	107	161	194	231	77	-
2000.....	854	1	4	21	32	42	53	85	120	196	218	82	-
1999.....	880	1	3	15	19	42	51	93	164	203	216	73	-

Asian or Pacific Islander 3/, male

2007 1/.....	652	1	4	8	16	19	30	60	112	147	184	71	-
2006 1/.....	568	2	3	8	17	25	26	50	84	122	158	73	-
2005 1/.....	570	-	4	9	12	26	28	67	98	130	147	49	-
2004 1/.....	556	2	5	10	15	13	29	54	82	137	158	51	-
2003 1/.....	497	1	2	10	13	23	29	45	79	124	123	48	-
2002.....	504	2	2	13	12	23	28	62	88	107	132	35	-
2001.....	504	-	3	14	12	14	32	57	96	106	131	39	-
2000.....	463	1	2	17	21	22	28	57	64	107	106	38	-
1999.....	481	-	2	5	13	24	32	50	94	102	111	48	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Asian or Pacific Islander 3/, female

2007 1/.....	498	-	3	7	11	15	24	49	78	101	154	56	-
2006 1/.....	467	-	3	6	8	15	20	42	75	113	120	65	-
2005 1/.....	487	-	2	6	11	14	27	56	79	119	119	54	-
2004 1/.....	417	-	3	7	4	13	17	37	62	107	112	55	-
2003 1/.....	449	-	4	10	10	11	20	42	64	101	139	48	-
2002.....	456	2	2	8	15	19	28	63	62	108	106	43	-
2001.....	396	1	3	3	10	24	14	50	65	88	100	38	-
2000.....	391	-	2	4	11	20	25	28	56	89	112	44	-
1999.....	399	1	1	10	6	18	19	43	70	101	105	25	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hodgkin's disease (C81)													
All races, both sexes													
2007.....	1,271	-	-	3	65	120	135	145	176	209	286	132	-
2006.....	1,327	-	-	5	65	136	154	148	188	205	288	138	-
2005.....	1,272	-	-	11	61	116	118	168	184	205	304	105	-
2004.....	1,276	-	-	6	65	134	120	168	205	222	249	107	-
2003.....	1,347	-	-	5	64	167	154	154	198	216	277	112	-
2002.....	1,352	-	-	6	86	152	159	177	168	232	247	125	-
2001.....	1,323	-	-	6	78	167	161	182	143	211	278	97	-
2000.....	1,287	-	-	8	73	145	151	161	158	232	255	104	-
1999.....	1,403	-	1	9	94	148	165	165	178	283	262	98	-
All races, male													
2007.....	706	-	-	1	36	75	88	103	99	115	140	49	-
2006.....	770	-	-	3	42	83	103	97	125	119	136	62	-
2005.....	684	-	-	7	27	65	71	94	121	118	131	50	-
2004.....	730	-	-	4	37	70	70	103	132	136	129	49	-
2003.....	725	-	-	4	30	88	95	91	119	128	132	38	-
2002.....	751	-	-	5	41	81	92	105	106	146	125	50	-
2001.....	725	-	-	3	43	90	101	110	85	129	130	34	-
2000.....	686	-	-	4	35	79	82	99	87	134	121	45	-
1999.....	785	-	1	3	55	86	103	98	115	160	119	45	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hodgkin's disease (C81)													
All races, female													
2007.....	565	-	-	2	29	45	47	42	77	94	146	83	-
2006.....	557	-	-	2	23	53	51	51	63	86	152	76	-
2005.....	588	-	-	4	34	51	47	74	63	87	173	55	-
2004.....	546	-	-	2	28	64	50	65	73	86	120	58	-
2003.....	622	-	-	1	34	79	59	63	79	88	145	74	-
2002.....	601	-	-	1	45	71	67	72	62	86	122	75	-
2001.....	598	-	-	3	35	77	60	72	58	82	148	63	-
2000.....	601	-	-	4	38	66	69	62	71	98	134	59	-
1999.....	618	-	-	6	39	62	62	67	63	123	143	53	-
White, both sexes													
2007 1/.....	1,121	-	-	3	51	100	109	116	147	192	276	127	-
2006 1/.....	1,160	-	-	5	52	107	131	123	164	183	269	126	-
2005 1/.....	1,095	-	-	11	48	73	88	137	163	189	289	97	-
2004 1/.....	1,118	-	-	5	51	98	101	146	180	201	236	100	-
2003 1/.....	1,192	-	-	5	51	134	132	139	170	196	258	107	-
2002.....	1,193	-	-	5	72	127	130	151	141	213	234	120	-
2001.....	1,170	-	-	5	62	125	138	161	124	200	260	95	-
2000.....	1,156	-	-	5	57	130	127	141	138	211	248	99	-
1999.....	1,263	-	1	7	75	122	146	146	163	260	254	89	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hodgkin's disease (C81)													
White, male													
2007 1/.....	621	-	-	1	29	66	73	81	82	107	135	47	-
2006 1/.....	666	-	-	3	36	63	87	81	108	105	126	57	-
2005 1/.....	582	-	-	7	22	38	55	76	103	108	124	49	-
2004 1/.....	647	-	-	4	29	56	59	90	116	122	123	48	-
2003 1/.....	638	-	-	4	24	74	84	81	101	112	123	35	-
2002.....	655	-	-	4	37	65	73	85	86	136	120	49	-
2001.....	644	-	-	3	38	69	85	97	74	124	122	32	-
2000.....	616	-	-	2	30	70	70	87	74	122	118	43	-
1999.....	708	-	1	3	45	72	92	83	107	148	116	41	-
White, female													
2007 1/.....	500	-	-	2	22	34	36	35	65	85	141	80	-
2006 1/.....	494	-	-	2	16	44	44	42	56	78	143	69	-
2005 1/.....	513	-	-	4	26	35	33	61	60	81	165	48	-
2004 1/.....	471	-	-	1	22	42	42	56	64	79	113	52	-
2003 1/.....	554	-	-	1	27	60	48	58	69	84	135	72	-
2002.....	538	-	-	1	35	62	57	66	55	77	114	71	-
2001.....	526	-	-	2	24	56	53	64	50	76	138	63	-
2000.....	540	-	-	3	27	60	57	54	64	89	130	56	-
1999.....	555	-	-	4	30	50	54	63	56	112	138	48	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hodgkin's disease (C81)													
Black, both sexes													
2007 1/.....	130	-	-	-	10	19	22	27	27	12	9	4	-
2006 1/.....	137	-	-	-	8	23	21	24	20	18	14	9	-
2005 1/.....	145	-	-	-	9	36	26	28	17	13	11	5	-
2004 1/.....	135	-	-	1	14	32	17	20	19	18	11	3	-
2003 1/.....	120	-	-	-	10	31	18	14	21	17	9	-	-
2002.....	130	-	-	1	11	24	26	20	21	15	9	3	-
2001.....	137	-	-	1	15	40	21	17	18	11	12	2	-
2000.....	110	-	-	2	14	13	24	17	17	15	5	3	-
1999.....	128	-	-	1	19	26	18	19	14	18	6	7	-
Black, male													
2007 1/.....	74	-	-	-	5	8	13	21	17	4	4	2	-
2006 1/.....	82	-	-	-	4	15	14	15	13	11	7	3	-
2005 1/.....	85	-	-	-	4	22	14	15	15	8	6	1	-
2004 1/.....	71	-	-	-	8	13	9	13	12	11	5	-	-
2003 1/.....	68	-	-	-	5	12	10	9	15	13	4	-	-
2002.....	82	-	-	1	4	15	18	15	16	8	4	1	-
2001.....	73	-	-	-	4	21	15	11	10	5	5	2	-
2000.....	57	-	-	1	5	8	12	9	11	7	3	1	-
1999.....	75	-	-	-	10	14	11	15	8	12	3	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hodgkin's disease (C81)													
Black, female													
2007 1/.....	56	-	-	-	5	11	9	6	10	8	5	2	-
2006 1/.....	55	-	-	-	4	8	7	9	7	7	7	6	-
2005 1/.....	60	-	-	-	5	14	12	13	2	5	5	4	-
2004 1/.....	64	-	-	1	6	19	8	7	7	7	6	3	-
2003 1/.....	52	-	-	-	5	19	8	5	6	4	5	-	-
2002.....	48	-	-	-	7	9	8	5	5	7	5	2	-
2001.....	64	-	-	1	11	19	6	6	8	6	7	-	-
2000.....	53	-	-	1	9	5	12	8	6	8	2	2	-
1999.....	53	-	-	1	9	12	7	4	6	6	3	5	-
American Indian 2/, both sexes													
2007 1/.....	3	-	-	-	1	-	-	1	-	-	1	-	-
2006 1/.....	6	-	-	-	1	1	2	-	-	-	2	-	-
2005 1/.....	3	-	-	-	-	1	-	1	-	1	-	-	-
2004 1/.....	4	-	-	-	-	2	1	-	-	1	-	-	-
2003 1/.....	9	-	-	-	-	-	1	-	2	1	4	1	-
2002.....	1	-	-	-	-	-	-	-	1	-	-	-	-
2001.....	3	-	-	-	-	1	1	1	-	-	-	-	-
2000.....	5	-	-	-	-	-	-	-	1	4	-	-	-
1999.....	5	-	-	-	-	-	1	-	-	3	-	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hodgkin's disease (C81)													
American Indian 2/, male													
2007 1/.....	3	-	-	-	1	-	-	1	-	-	1	-	-
2006 1/.....	4	-	-	-	-	1	2	-	-	-	1	-	-
2005 1/.....	3	-	-	-	-	1	-	1	-	-	-	-	-
2004 1/.....	3	-	-	-	-	1	1	-	-	1	-	-	-
2003 1/.....	4	-	-	-	-	-	1	-	1	1	1	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	4	-	-	-	-	-	-	-	-	4	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	-	-	1	-
American Indian 2/, female													
2007 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2006 1/.....	2	-	-	-	1	-	-	-	-	-	1	-	-
2005 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 1/.....	1	-	-	-	-	1	-	-	-	-	-	-	-
2003 1/.....	5	-	-	-	-	-	-	-	1	-	3	1	-
2002.....	1	-	-	-	-	-	-	-	1	-	-	-	-
2001.....	3	-	-	-	-	1	1	1	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	1	-	-	-	-
1999.....	4	-	-	-	-	-	1	-	-	3	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Hodgkin's disease (C81)

Asian or Pacific Islander 3/, female

2007 1/.....	9	-	-	-	2	-	2	1	2	1	-	1	-
2006 1/.....	6	-	-	-	2	1	-	-	-	1	1	1	-
2005 1/.....	15	-	-	-	3	2	2	-	1	1	3	3	-
2004 1/.....	10	-	-	-	-	2	-	2	2	-	1	3	-
2003 1/.....	11	-	-	-	2	-	3	-	3	-	2	1	-
2002.....	14	-	-	-	3	-	2	1	1	2	3	2	-
2001.....	5	-	-	-	-	1	-	1	-	-	3	-	-
2000.....	7	-	-	-	2	1	-	-	-	1	2	1	-
1999.....	6	-	-	1	-	-	-	-	1	2	2	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Non-Hodgkin's lymphoma (C82-C85)													
All races, both sexes													
2007.....	20,528	2	5	33	133	206	516	1,392	2,922	4,476	6,868	3,975	-
2006.....	20,594	1	7	36	131	219	515	1,417	2,886	4,543	7,007	3,831	1
2005.....	20,873	1	6	48	127	222	566	1,410	2,925	4,626	7,111	3,831	-
2004.....	20,938	1	2	41	144	256	543	1,538	2,866	4,820	7,080	3,647	-
2003.....	21,475	-	5	45	140	252	630	1,548	3,007	4,919	7,217	3,712	-
2002.....	21,910	2	15	57	142	281	687	1,581	3,049	5,264	7,283	3,549	-
2001.....	22,305	1	1	38	148	298	756	1,732	3,098	5,358	7,369	3,506	-
2000.....	22,729	1	7	47	180	293	751	1,745	3,170	5,598	7,535	3,402	-
1999.....	22,802	2	16	37	144	289	789	1,741	3,154	5,848	7,554	3,228	-
All races, male													
2007.....	11,004	1	3	23	87	131	340	884	1,788	2,629	3,490	1,628	-
2006.....	10,933	1	6	28	94	134	320	918	1,795	2,577	3,546	1,513	1
2005.....	11,131	-	2	40	90	134	365	909	1,745	2,677	3,557	1,612	-
2004.....	11,056	-	2	28	93	164	367	983	1,725	2,738	3,511	1,445	-
2003.....	11,399	-	5	26	88	165	406	972	1,848	2,829	3,563	1,497	-
2002.....	11,538	1	6	40	93	178	439	978	1,838	2,986	3,594	1,385	-
2001.....	11,710	1	1	29	94	179	500	1,113	1,825	3,025	3,558	1,385	-
2000.....	11,812	-	5	28	126	172	482	1,112	1,849	3,105	3,634	1,299	-
1999.....	11,794	-	9	30	91	178	510	1,111	1,847	3,176	3,596	1,246	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Non-Hodgkin's lymphoma (C82-C85)													
All races, female													
2007.....	9,524	1	2	10	46	75	176	508	1,134	1,847	3,378	2,347	-
2006.....	9,661	-	1	8	37	85	195	499	1,091	1,966	3,461	2,318	-
2005.....	9,742	1	4	8	37	88	201	501	1,180	1,949	3,554	2,219	-
2004.....	9,882	1	-	13	51	92	176	555	1,141	2,082	3,569	2,202	-
2003.....	10,076	-	-	19	52	87	224	576	1,159	2,090	3,654	2,215	-
2002.....	10,372	1	9	17	49	103	248	603	1,211	2,278	3,689	2,164	-
2001.....	10,595	-	-	9	54	119	256	619	1,273	2,333	3,811	2,121	-
2000.....	10,917	1	2	19	54	121	269	633	1,321	2,493	3,901	2,103	-
1999.....	11,008	2	7	7	53	111	279	630	1,307	2,672	3,958	1,982	-
White, both sexes													
2007 1/.....	18,581	2	2	28	101	144	393	1,127	2,523	4,057	6,417	3,787	-
2006 1/.....	18,636	1	5	28	94	162	371	1,148	2,491	4,114	6,585	3,636	1
2005 1/.....	18,970	1	5	38	92	152	431	1,140	2,547	4,206	6,707	3,651	-
2004 1/.....	19,036	1	2	30	110	186	439	1,255	2,525	4,376	6,660	3,452	-
2003 1/.....	19,598	-	4	31	112	177	499	1,278	2,681	4,478	6,793	3,545	-
2002.....	19,996	1	14	45	106	203	540	1,309	2,699	4,799	6,873	3,407	-
2001.....	20,335	1	1	32	109	221	601	1,461	2,732	4,903	6,952	3,322	-
2000.....	20,892	-	4	40	131	218	610	1,482	2,854	5,161	7,157	3,235	-
1999.....	20,941	-	13	26	110	214	627	1,504	2,818	5,415	7,157	3,057	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Non-Hodgkin's lymphoma (C82-C85)													
White, male													
2007 1/.....	9,941	1	1	19	66	95	260	724	1,563	2,380	3,282	1,550	-
2006 1/.....	9,871	1	4	23	70	101	235	741	1,560	2,345	3,360	1,430	1
2005 1/.....	10,118	-	1	30	70	89	291	733	1,530	2,450	3,378	1,546	-
2004 1/.....	10,045	-	2	22	69	123	306	809	1,524	2,493	3,319	1,378	-
2003 1/.....	10,447	-	4	17	72	120	327	805	1,664	2,620	3,384	1,434	-
2002.....	10,535	-	6	32	75	131	343	830	1,637	2,745	3,408	1,328	-
2001.....	10,642	1	1	23	68	140	396	947	1,609	2,781	3,365	1,311	-
2000.....	10,838	-	4	24	96	131	401	944	1,658	2,875	3,465	1,240	-
1999.....	10,768	-	8	23	71	130	407	962	1,643	2,953	3,405	1,166	-
White, female													
2007 1/.....	8,640	1	1	9	35	49	133	403	960	1,677	3,135	2,237	-
2006 1/.....	8,765	-	1	5	24	61	136	407	931	1,769	3,225	2,206	-
2005 1/.....	8,852	1	4	8	22	63	140	407	1,017	1,756	3,329	2,105	-
2004 1/.....	8,991	1	-	8	41	63	133	446	1,001	1,883	3,341	2,074	-
2003 1/.....	9,151	-	-	14	40	57	172	473	1,017	1,858	3,409	2,111	-
2002.....	9,461	1	8	13	31	72	197	479	1,062	2,054	3,465	2,079	-
2001.....	9,693	-	-	9	41	81	205	514	1,123	2,122	3,587	2,011	-
2000.....	10,054	-	-	16	35	87	209	538	1,196	2,286	3,692	1,995	-
1999.....	10,173	-	5	3	39	84	220	542	1,175	2,462	3,752	1,891	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Non-Hodgkin's lymphoma (C82-C85)													
Black, both sexes													
2007 1/.....	1,390	-	2	5	24	50	105	207	307	301	264	125	-
2006 1/.....	1,475	-	2	6	29	46	120	229	322	307	283	131	-
2005 1/.....	1,392	-	1	9	30	57	112	214	280	297	266	126	-
2004 1/.....	1,414	-	-	8	30	59	85	235	262	320	278	137	-
2003 1/.....	1,371	-	-	10	23	58	108	230	246	311	278	107	-
2002.....	1,474	1	1	10	25	63	118	218	286	358	291	103	-
2001.....	1,496	-	-	5	32	63	129	224	276	351	288	128	-
2000.....	1,369	1	2	5	35	56	121	210	257	320	246	116	-
1999.....	1,443	2	2	9	32	58	138	192	264	331	288	127	-
Black, male													
2007 1/.....	749	-	2	4	15	29	68	125	174	177	112	43	-
2006 1/.....	801	-	2	4	18	26	74	153	196	163	118	47	-
2005 1/.....	752	-	1	9	17	34	61	143	166	167	111	43	-
2004 1/.....	739	-	-	3	20	36	50	148	153	169	118	42	-
2003 1/.....	687	-	-	8	12	34	63	144	138	145	110	33	-
2002.....	775	1	-	7	10	40	79	119	161	195	123	40	-
2001.....	804	-	-	5	22	33	88	138	154	191	125	48	-
2000.....	728	-	-	3	19	31	71	135	158	170	103	38	-
1999.....	786	-	1	6	18	39	87	123	155	174	132	51	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Non-Hodgkin's lymphoma (C82-C85)													
Black, female													
2007 1/.....	641	-	-	1	9	21	37	82	133	124	152	82	-
2006 1/.....	674	-	-	2	11	20	46	76	126	144	165	84	-
2005 1/.....	640	-	-	-	13	23	51	71	114	130	155	83	-
2004 1/.....	675	-	-	5	10	23	35	87	109	151	160	95	-
2003 1/.....	684	-	-	2	11	24	45	86	108	166	168	74	-
2002.....	699	-	1	3	15	23	39	99	125	163	168	63	-
2001.....	692	-	-	-	10	30	41	86	122	160	163	80	-
2000.....	641	1	2	2	16	25	50	75	99	150	143	78	-
1999.....	657	2	1	3	14	19	51	69	109	157	156	76	-
American Indian 2/, both sexes													
2007 1/.....	77	-	-	-	1	2	2	7	12	23	22	8	-
2006 1/.....	68	-	-	-	-	1	5	7	8	18	17	12	-
2005 1/.....	83	-	-	-	-	3	5	10	25	19	13	8	-
2004 1/.....	67	-	-	-	-	2	5	4	14	21	16	5	-
2003 1/.....	69	-	1	-	-	2	3	8	17	20	15	3	-
2002.....	53	-	-	1	1	1	8	7	11	11	9	4	-
2001.....	93	-	-	1	1	4	10	9	17	24	19	8	-
2000.....	83	-	-	-	1	4	4	12	10	21	26	5	-
1999.....	57	-	-	1	-	3	6	3	9	18	12	5	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Non-Hodgkin's lymphoma (C82-C85)													
American Indian 2/, male													
2007 1/.....	42	-	-	-	1	2	2	4	5	15	9	4	-
2006 1/.....	31	-	-	-	-	-	-	5	4	8	8	6	-
2005 1/.....	39	-	-	-	-	2	3	4	11	11	6	2	-
2004 1/.....	39	-	-	-	-	-	2	3	9	16	7	2	-
2003 1/.....	34	-	1	-	-	2	2	3	10	10	6	-	-
2002.....	29	-	-	1	1	-	7	3	7	6	4	-	-
2001.....	48	-	-	1	1	2	5	5	11	13	7	3	-
2000.....	41	-	-	-	1	3	1	8	5	8	14	1	-
1999.....	31	-	-	1	-	1	4	2	6	7	8	2	-
American Indian 2/, female													
2007 1/.....	35	-	-	-	-	-	-	3	7	8	13	4	-
2006 1/.....	37	-	-	-	-	1	5	2	4	10	9	6	-
2005 1/.....	44	-	-	-	-	1	2	6	14	8	7	6	-
2004 1/.....	28	-	-	-	-	2	3	1	5	5	9	3	-
2003 1/.....	35	-	-	-	-	-	1	5	7	10	9	3	-
2002.....	24	-	-	-	-	1	1	4	4	5	5	4	-
2001.....	45	-	-	-	-	2	5	4	6	11	12	5	-
2000.....	42	-	-	-	-	1	3	4	5	13	12	4	-
1999.....	26	-	-	-	-	2	2	1	3	11	4	3	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Non-Hodgkin's lymphoma (C82-C85)													
Asian or Pacific Islander 3/, both sexes													
2007 1/.....	480	-	1	-	7	10	16	51	80	95	165	55	-
2006 1/.....	415	-	-	2	8	10	19	33	65	104	122	52	-
2005 1/.....	428	-	-	1	5	10	18	46	73	104	125	46	-
2004 1/.....	421	-	-	3	4	9	14	44	65	103	126	53	-
2003 1/.....	437	-	-	4	5	15	20	32	63	110	131	57	-
2002.....	387	-	-	1	10	14	21	47	53	96	110	35	-
2001.....	381	-	-	-	6	10	16	38	73	80	110	48	-
2000.....	385	-	1	2	13	15	16	41	49	96	106	46	-
1999.....	361	-	1	1	2	14	18	42	63	84	97	39	-
Asian or Pacific Islander 3/, male													
2007 1/.....	272	-	-	-	5	5	10	31	46	57	87	31	-
2006 1/.....	230	-	-	1	6	7	11	19	35	61	60	30	-
2005 1/.....	222	-	-	1	3	9	10	29	38	49	62	21	-
2004 1/.....	233	-	-	3	4	5	9	23	39	60	67	23	-
2003 1/.....	231	-	-	1	4	9	14	20	36	54	63	30	-
2002.....	199	-	-	-	7	7	10	26	33	40	59	17	-
2001.....	216	-	-	-	3	4	11	23	51	40	61	23	-
2000.....	205	-	1	1	10	7	9	25	28	52	52	20	-
1999.....	209	-	-	-	2	8	12	24	43	42	51	27	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Non-Hodgkin's lymphoma (C82-C85)

Asian or Pacific Islander 3/, female

2007 1/.....	208	-	1	-	2	5	6	20	34	38	78	24	-
2006 1/.....	185	-	-	1	2	3	8	14	30	43	62	22	-
2005 1/.....	206	-	-	-	2	1	8	17	35	55	63	25	-
2004 1/.....	188	-	-	-	-	4	5	21	26	43	59	30	-
2003 1/.....	206	-	-	3	1	6	6	12	27	56	68	27	-
2002.....	188	-	-	1	3	7	11	21	20	56	51	18	-
2001.....	165	-	-	-	3	6	5	15	22	40	49	25	-
2000.....	180	-	-	1	3	8	7	16	21	44	54	26	-
1999.....	152	-	1	1	-	6	6	18	20	42	46	12	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Leukemia (C91-C95)													
All races, both sexes													
2007.....	21,825	21	106	314	428	438	657	1,362	2,801	4,611	6,858	4,228	1
2006.....	21,944	31	124	256	465	460	715	1,398	2,823	4,735	6,863	4,074	-
2005.....	21,623	22	120	275	466	461	680	1,419	2,662	4,673	6,832	4,013	-
2004.....	21,395	23	120	312	435	416	713	1,367	2,688	4,694	6,797	3,830	-
2003.....	21,535	30	133	342	402	433	796	1,365	2,605	4,706	6,937	3,786	-
2002.....	21,498	21	104	316	472	471	767	1,459	2,611	4,818	6,763	3,696	-
2001.....	21,451	25	155	323	479	513	779	1,475	2,535	4,886	6,696	3,585	-
2000.....	21,339	38	137	315	464	473	801	1,411	2,414	5,010	6,733	3,543	-
1999.....	21,014	27	119	305	482	500	799	1,370	2,516	5,012	6,460	3,424	-
All races, male													
2007.....	12,388	14	53	181	251	251	364	786	1,715	2,809	4,020	1,943	1
2006.....	12,385	17	64	146	298	274	393	796	1,733	2,869	4,021	1,774	-
2005.....	12,231	11	81	159	293	257	358	762	1,649	2,902	3,957	1,802	-
2004.....	12,007	9	70	182	268	242	398	782	1,663	2,903	3,821	1,669	-
2003.....	12,073	9	75	195	257	274	444	801	1,593	2,930	3,880	1,615	-
2002.....	12,016	11	57	172	282	280	429	821	1,558	3,010	3,804	1,592	-
2001.....	11,850	10	81	185	289	300	432	813	1,552	2,962	3,699	1,527	-
2000.....	11,777	21	69	190	293	283	423	791	1,452	3,090	3,695	1,470	-
1999.....	11,519	11	65	173	291	276	435	777	1,489	3,054	3,525	1,423	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Leukemia (C91-C95)													
All races, female													
2007.....	9,437	7	53	133	177	187	293	576	1,086	1,802	2,838	2,285	-
2006.....	9,559	14	60	110	167	186	322	602	1,090	1,866	2,842	2,300	-
2005.....	9,392	11	39	116	173	204	322	657	1,013	1,771	2,875	2,211	-
2004.....	9,388	14	50	130	167	174	315	585	1,025	1,791	2,976	2,161	-
2003.....	9,462	21	58	147	145	159	352	564	1,012	1,776	3,057	2,171	-
2002.....	9,482	10	47	144	190	191	338	638	1,053	1,808	2,959	2,104	-
2001.....	9,601	15	74	138	190	213	347	662	983	1,924	2,997	2,058	-
2000.....	9,562	17	68	125	171	190	378	620	962	1,920	3,038	2,073	-
1999.....	9,495	16	54	132	191	224	364	593	1,027	1,958	2,935	2,001	-
White, both sexes													
2007 1/.....	19,481	19	82	251	339	349	507	1,125	2,420	4,082	6,338	3,968	1
2006 1/.....	19,654	25	102	203	377	370	580	1,182	2,445	4,239	6,329	3,802	-
2005 1/.....	19,414	18	87	224	381	351	558	1,160	2,332	4,192	6,334	3,777	-
2004 1/.....	19,077	18	92	254	353	330	563	1,134	2,333	4,154	6,270	3,576	-
2003 1/.....	19,353	28	111	276	320	349	631	1,145	2,285	4,224	6,433	3,551	-
2002.....	19,273	16	84	253	394	370	644	1,196	2,246	4,361	6,262	3,447	-
2001.....	19,239	20	134	244	391	402	628	1,227	2,210	4,386	6,223	3,374	-
2000.....	19,191	36	115	259	361	355	648	1,169	2,088	4,560	6,298	3,302	-
1999.....	18,899	23	103	238	379	401	660	1,143	2,196	4,531	6,007	3,218	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Leukemia (C91-C95)													
White, male													
2007 1/.....	11,147	13	38	146	199	201	286	653	1,511	2,514	3,756	1,829	1
2006 1/.....	11,110	13	52	111	242	228	313	676	1,515	2,582	3,726	1,652	-
2005 1/.....	11,034	9	58	127	239	203	291	630	1,462	2,628	3,672	1,715	-
2004 1/.....	10,753	7	50	149	220	197	316	645	1,467	2,576	3,556	1,570	-
2003 1/.....	10,938	7	63	163	205	218	351	682	1,412	2,659	3,650	1,528	-
2002.....	10,835	10	47	134	239	225	362	678	1,359	2,736	3,547	1,498	-
2001.....	10,672	7	68	131	240	239	342	691	1,367	2,681	3,458	1,448	-
2000.....	10,617	19	58	153	225	216	353	658	1,267	2,827	3,471	1,370	-
1999.....	10,395	11	54	134	233	219	362	646	1,314	2,784	3,299	1,339	-
White, female													
2007 1/.....	8,334	6	44	105	140	148	221	472	909	1,568	2,582	2,139	-
2006 1/.....	8,544	12	50	92	135	142	267	506	930	1,657	2,603	2,150	-
2005 1/.....	8,380	9	29	97	142	148	267	530	870	1,564	2,662	2,062	-
2004 1/.....	8,324	11	42	105	133	133	247	489	866	1,578	2,714	2,006	-
2003 1/.....	8,415	21	48	113	115	131	280	463	873	1,565	2,783	2,023	-
2002.....	8,438	6	37	119	155	145	282	518	887	1,625	2,715	1,949	-
2001.....	8,567	13	66	113	151	163	286	536	843	1,705	2,765	1,926	-
2000.....	8,574	17	57	106	136	139	295	511	821	1,733	2,827	1,932	-
1999.....	8,504	12	49	104	146	182	298	497	882	1,747	2,708	1,879	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Leukemia (C91-C95)													
Black, both sexes													
2007 1/.....	1,831	1	18	45	64	63	116	190	306	415	405	208	-
2006 1/.....	1,804	4	15	41	69	62	105	170	306	408	410	214	-
2005 1/.....	1,718	4	25	36	65	84	89	201	257	374	388	195	-
2004 1/.....	1,860	3	19	40	62	71	114	195	298	436	409	213	-
2003 1/.....	1,762	1	14	47	55	65	132	178	264	395	408	203	-
2002.....	1,770	4	15	42	54	68	91	201	295	376	406	218	-
2001.....	1,765	4	14	59	64	75	119	187	249	415	393	186	-
2000.....	1,742	-	18	36	82	88	117	204	261	375	350	211	-
1999.....	1,708	2	13	53	83	68	107	175	252	407	373	175	-
Black, male													
2007 1/.....	951	-	11	26	36	34	62	110	160	227	195	90	-
2006 1/.....	999	2	8	28	42	30	64	90	179	240	221	95	-
2005 1/.....	911	2	17	23	43	40	48	102	145	211	213	67	-
2004 1/.....	980	-	14	23	35	37	60	111	165	267	188	80	-
2003 1/.....	918	1	8	23	35	41	75	100	151	221	189	74	-
2002.....	923	1	7	25	31	37	49	112	161	221	198	81	-
2001.....	928	3	9	39	35	44	68	93	147	235	187	68	-
2000.....	929	-	9	22	55	50	53	107	151	216	181	85	-
1999.....	892	-	8	33	46	38	54	103	142	224	178	66	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Leukemia (C91-C95)													
Black, female													
2007 1/.....	880	1	7	19	28	29	54	80	146	188	210	118	-
2006 1/.....	805	2	7	13	27	32	41	80	127	168	189	119	-
2005 1/.....	807	2	8	13	22	44	41	99	112	163	175	128	-
2004 1/.....	880	3	5	17	27	34	54	84	133	169	221	133	-
2003 1/.....	844	-	6	24	20	24	57	78	113	174	219	129	-
2002.....	847	3	8	17	23	31	42	89	134	155	208	137	-
2001.....	837	1	5	20	29	31	51	94	102	180	206	118	-
2000.....	813	-	9	14	27	38	64	97	110	159	169	126	-
1999.....	816	2	5	20	37	30	53	72	110	183	195	109	-
American Indian 2/, both sexes													
2007 1/.....	78	-	-	3	9	3	5	9	13	15	15	6	-
2006 1/.....	71	-	1	-	6	4	4	10	15	12	13	6	-
2005 1/.....	73	-	2	1	6	2	4	7	10	17	17	7	-
2004 1/.....	76	-	1	4	5	2	6	5	11	16	19	7	-
2003 1/.....	77	-	2	3	12	3	7	8	10	14	12	6	-
2002.....	73	-	1	1	10	6	4	4	13	12	15	7	-
2001.....	77	-	1	3	9	9	5	8	18	15	5	4	-
2000.....	72	1	1	2	4	5	4	4	15	16	10	10	-
1999.....	56	1	1	1	3	4	1	9	8	13	10	5	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Leukemia (C91-C95)													
American Indian 2/, male													
2007 1/.....	42	-	-	1	6	3	2	6	8	9	7	-	-
2006 1/.....	45	-	1	-	5	2	2	9	11	8	6	1	-
2005 1/.....	43	-	2	1	3	1	3	4	4	12	10	3	-
2004 1/.....	42	-	1	3	2	2	3	3	9	7	11	1	-
2003 1/.....	38	-	2	-	9	3	3	6	3	7	4	1	-
2002.....	40	-	1	-	7	3	1	3	7	7	9	2	-
2001.....	42	-	1	1	6	7	3	5	10	7	2	-	-
2000.....	38	1	1	-	2	3	2	3	7	11	4	4	-
1999.....	38	-	1	1	1	3	-	7	4	11	7	3	-
American Indian 2/, female													
2007 1/.....	36	-	-	2	3	-	3	3	5	6	8	6	-
2006 1/.....	26	-	-	-	1	2	2	1	4	4	7	5	-
2005 1/.....	30	-	-	-	3	1	1	3	6	5	7	4	-
2004 1/.....	34	-	-	1	3	-	3	2	2	9	8	6	-
2003 1/.....	39	-	-	3	3	-	4	2	7	7	8	5	-
2002.....	33	-	-	1	3	3	3	1	6	5	6	5	-
2001.....	35	-	-	2	3	2	2	3	8	8	3	4	-
2000.....	34	-	-	2	2	2	2	1	8	5	6	6	-
1999.....	18	1	-	-	2	1	1	2	4	2	3	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Leukemia (C91-C95)													
Asian or Pacific Islander 3/, both sexes													
2007 1/.....	435	1	6	15	16	23	29	38	62	99	100	46	-
2006 1/.....	415	2	6	12	13	24	26	36	57	76	111	52	-
2005 1/.....	418	-	6	14	14	24	29	51	63	90	93	34	-
2004 1/.....	382	2	8	14	15	13	30	33	46	88	99	34	-
2003 1/.....	343	1	6	16	15	16	26	34	46	73	84	26	-
2002.....	382	1	4	20	14	27	28	58	57	69	80	24	-
2001.....	370	1	6	17	15	27	27	53	58	70	75	21	-
2000.....	334	1	3	18	17	25	32	34	50	59	75	20	-
1999.....	351	1	2	13	17	27	31	43	60	61	70	26	-
Asian or Pacific Islander 3/, male													
2007 1/.....	248	1	4	8	10	13	14	17	36	59	62	24	-
2006 1/.....	231	2	3	7	9	14	14	21	28	39	68	26	-
2005 1/.....	243	-	4	8	8	13	16	26	38	51	62	17	-
2004 1/.....	232	2	5	7	11	6	19	23	22	53	66	18	-
2003 1/.....	179	1	2	9	8	12	15	13	27	43	37	12	-
2002.....	218	-	2	13	5	15	17	28	31	46	50	11	-
2001.....	208	-	3	14	8	10	19	24	28	39	52	11	-
2000.....	193	1	1	15	11	14	15	23	27	36	39	11	-
1999.....	194	-	2	5	11	16	19	21	29	35	41	15	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Leukemia (C91-C95)

Asian or Pacific Islander 3/, female

2007 1/.....	187	-	2	7	6	10	15	21	26	40	38	22	-
2006 1/.....	184	-	3	5	4	10	12	15	29	37	43	26	-
2005 1/.....	175	-	2	6	6	11	13	25	25	39	31	17	-
2004 1/.....	150	-	3	7	4	7	11	10	24	35	33	16	-
2003 1/.....	164	-	4	7	7	4	11	21	19	30	47	14	-
2002.....	164	1	2	7	9	12	11	30	26	23	30	13	-
2001.....	162	1	3	3	7	17	8	29	30	31	23	10	-
2000.....	141	-	2	3	6	11	17	11	23	23	36	9	-
1999.....	157	1	-	8	6	11	12	22	31	26	29	11	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Multiple myeloma and immunoproliferative neoplasms (C88,C90)													
All races, both sexes													
2007.....	11,307	-	-	-	2	7	155	705	1,788	2,917	3,853	1,879	1
2006.....	11,111	-	-	-	1	12	150	731	1,789	2,843	3,793	1,792	-
2005.....	11,200	-	-	-	1	11	131	688	1,842	2,969	3,846	1,712	-
2004.....	10,965	1	-	-	1	14	136	663	1,720	2,840	3,980	1,610	-
2003.....	11,267	-	-	1	-	12	147	711	1,659	3,029	3,997	1,711	-
2002.....	11,392	2	-	-	2	12	138	736	1,669	3,176	3,971	1,686	-
2001.....	11,172	-	-	-	-	9	157	671	1,671	3,181	3,886	1,597	-
2000.....	11,063	-	-	1	-	7	146	700	1,530	3,179	3,896	1,604	-
1999.....	11,016	-	-	-	-	22	156	665	1,694	3,172	3,788	1,519	-
All races, male													
2007.....	6,066	-	-	-	1	4	96	425	1,024	1,617	2,021	878	-
2006.....	5,882	-	-	-	1	8	89	422	1,035	1,569	1,985	773	-
2005.....	5,945	-	-	-	1	8	83	409	1,021	1,678	2,006	739	-
2004.....	5,694	1	-	-	1	11	78	381	972	1,587	1,990	673	-
2003.....	5,848	-	-	-	-	7	97	426	995	1,654	1,941	728	-
2002.....	5,953	-	-	-	-	6	78	433	966	1,746	2,018	706	-
2001.....	5,724	-	-	-	-	4	97	403	945	1,725	1,926	624	-
2000.....	5,553	-	-	-	-	5	85	391	808	1,697	1,932	635	-
1999.....	5,584	-	-	-	-	16	95	378	968	1,669	1,846	612	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Multiple myeloma and immunoproliferative neoplasms (C88,C90)													
All races, female													
2007.....	5,241	-	-	-	1	3	59	280	764	1,300	1,832	1,001	1
2006.....	5,229	-	-	-	-	4	61	309	754	1,274	1,808	1,019	-
2005.....	5,255	-	-	-	-	3	48	279	821	1,291	1,840	973	-
2004.....	5,271	-	-	-	-	3	58	282	748	1,253	1,990	937	-
2003.....	5,419	-	-	1	-	5	50	285	664	1,375	2,056	983	-
2002.....	5,439	2	-	-	2	6	60	303	703	1,430	1,953	980	-
2001.....	5,448	-	-	-	-	5	60	268	726	1,456	1,960	973	-
2000.....	5,510	-	-	1	-	2	61	309	722	1,482	1,964	969	-
1999.....	5,432	-	-	-	-	6	61	287	726	1,503	1,942	907	-
White, both sexes													
2007 1/.....	9,148	-	-	-	1	6	99	486	1,340	2,324	3,254	1,637	1
2006 1/.....	9,038	-	-	-	-	8	108	499	1,340	2,268	3,254	1,561	-
2005 1/.....	9,171	-	-	-	1	6	92	473	1,406	2,367	3,319	1,507	-
2004 1/.....	8,948	1	-	-	1	11	95	472	1,320	2,252	3,394	1,402	-
2003 1/.....	9,227	-	-	1	-	9	103	488	1,291	2,450	3,395	1,490	-
2002.....	9,420	1	-	-	2	10	91	523	1,277	2,587	3,447	1,482	-
2001.....	9,168	-	-	-	-	5	100	491	1,247	2,598	3,333	1,394	-
2000.....	9,066	-	-	1	-	4	97	499	1,164	2,605	3,344	1,352	-
1999.....	9,010	-	-	-	-	13	103	481	1,252	2,603	3,246	1,312	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Multiple myeloma and immunoproliferative neoplasms (C88,C90)													
White, male													
2007 1/.....	5,032	-	-	-	1	3	68	307	791	1,340	1,738	784	-
2006 1/.....	4,843	-	-	-	-	5	67	301	782	1,274	1,730	684	-
2005 1/.....	4,937	-	-	-	1	4	63	289	794	1,372	1,752	662	-
2004 1/.....	4,727	1	-	-	1	9	58	276	768	1,296	1,726	592	-
2003 1/.....	4,892	-	-	-	-	6	73	299	788	1,383	1,693	650	-
2002.....	5,024	-	-	-	-	5	54	326	759	1,463	1,785	632	-
2001.....	4,765	-	-	-	-	3	67	304	716	1,437	1,684	554	-
2000.....	4,644	-	-	-	-	3	64	284	632	1,414	1,690	557	-
1999.....	4,651	-	-	-	-	9	64	290	747	1,394	1,611	536	-
White, female													
2007 1/.....	4,116	-	-	-	-	3	31	179	549	984	1,516	853	1
2006 1/.....	4,195	-	-	-	-	3	41	198	558	994	1,524	877	-
2005 1/.....	4,234	-	-	-	-	2	29	184	612	995	1,567	845	-
2004 1/.....	4,221	-	-	-	-	2	37	196	552	956	1,668	810	-
2003 1/.....	4,335	-	-	1	-	3	30	189	503	1,067	1,702	840	-
2002.....	4,396	1	-	-	2	5	37	197	518	1,124	1,662	850	-
2001.....	4,403	-	-	-	-	2	33	187	531	1,161	1,649	840	-
2000.....	4,422	-	-	1	-	1	33	215	532	1,191	1,654	795	-
1999.....	4,359	-	-	-	-	4	39	191	505	1,209	1,635	776	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

Black, both sexes

2007 1/.....	1,893	-	-	-	-	1	51	194	389	528	515	215	-
2006 1/.....	1,844	-	-	-	1	3	40	203	405	508	485	199	-
2005 1/.....	1,785	-	-	-	-	5	34	180	381	524	476	185	-
2004 1/.....	1,816	-	-	-	-	2	40	170	359	527	527	191	-
2003 1/.....	1,847	-	-	-	-	2	43	201	327	526	543	205	-
2002.....	1,771	-	-	-	-	2	41	192	347	532	470	187	-
2001.....	1,817	-	-	-	-	4	53	161	384	523	502	190	-
2000.....	1,841	-	-	-	-	3	44	190	339	531	505	229	-
1999.....	1,803	-	-	-	-	8	50	173	396	497	481	198	-

Black, male

2007 1/.....	882	-	-	-	-	1	25	102	197	241	239	77	-
2006 1/.....	926	-	-	-	1	3	20	109	228	270	222	73	-
2005 1/.....	882	-	-	-	-	4	19	104	200	263	226	66	-
2004 1/.....	859	-	-	-	-	1	20	89	183	264	230	72	-
2003 1/.....	863	-	-	-	-	1	24	115	189	239	225	70	-
2002.....	840	-	-	-	-	1	22	99	185	260	206	67	-
2001.....	864	-	-	-	-	1	28	88	208	253	223	63	-
2000.....	837	-	-	-	-	2	17	98	164	263	224	69	-
1999.....	834	-	-	-	-	7	30	82	197	239	210	69	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Multiple myeloma and immunoproliferative neoplasms (C88,C90)													
Black, female													
2007 1/.....	1,011	-	-	-	-	-	26	92	192	287	276	138	-
2006 1/.....	918	-	-	-	-	-	20	94	177	238	263	126	-
2005 1/.....	903	-	-	-	-	1	15	76	181	261	250	119	-
2004 1/.....	957	-	-	-	-	1	20	81	176	263	297	119	-
2003 1/.....	984	-	-	-	-	1	19	86	138	287	318	135	-
2002.....	931	-	-	-	-	1	19	93	162	272	264	120	-
2001.....	953	-	-	-	-	3	25	73	176	270	279	127	-
2000.....	1,004	-	-	-	-	1	27	92	175	268	281	160	-
1999.....	969	-	-	-	-	1	20	91	199	258	271	129	-
American Indian 2/, both sexes													
2007 1/.....	49	-	-	-	-	-	1	6	13	16	11	2	-
2006 1/.....	49	-	-	-	-	-	1	7	11	16	12	2	-
2005 1/.....	63	-	-	-	-	-	1	11	18	25	8	-	-
2004 1/.....	52	-	-	-	-	-	-	9	14	10	17	2	-
2003 1/.....	55	-	-	-	-	-	1	3	12	13	19	7	-
2002.....	43	-	-	-	-	-	2	8	10	11	11	1	-
2001.....	51	-	-	-	-	-	2	6	11	16	11	5	-
2000.....	38	-	-	-	-	-	1	4	8	4	12	9	-
1999.....	43	-	-	-	-	-	1	3	6	16	15	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Multiple myeloma and immunoproliferative neoplasms (C88,C90)													
American Indian 2/, male													
2007 1/.....	29	-	-	-	-	-	-	4	6	9	9	1	-
2006 1/.....	25	-	-	-	-	-	1	3	8	6	5	2	-
2005 1/.....	36	-	-	-	-	-	1	6	8	14	7	-	-
2004 1/.....	28	-	-	-	-	-	-	8	4	5	11	-	-
2003 1/.....	23	-	-	-	-	-	-	2	4	7	5	5	-
2002.....	20	-	-	-	-	-	2	5	2	4	6	1	-
2001.....	23	-	-	-	-	-	1	3	5	8	4	2	-
2000.....	17	-	-	-	-	-	1	3	5	2	3	3	-
1999.....	22	-	-	-	-	-	-	1	2	11	6	2	-
American Indian 2/, female													
2007 1/.....	20	-	-	-	-	-	1	2	7	7	2	1	-
2006 1/.....	24	-	-	-	-	-	-	4	3	10	7	-	-
2005 1/.....	27	-	-	-	-	-	-	5	10	11	1	-	-
2004 1/.....	24	-	-	-	-	-	-	1	10	5	6	2	-
2003 1/.....	32	-	-	-	-	-	1	1	8	6	14	2	-
2002.....	23	-	-	-	-	-	-	3	8	7	5	-	-
2001.....	28	-	-	-	-	-	1	3	6	8	7	3	-
2000.....	21	-	-	-	-	-	-	1	3	2	9	6	-
1999.....	21	-	-	-	-	-	1	2	4	5	9	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Multiple myeloma and immunoproliferative neoplasms (C88,C90)													
Asian or Pacific Islander 3/, both sexes													
2007 1/.....	217	-	-	-	1	-	4	19	46	49	73	25	-
2006 1/.....	180	-	-	-	-	1	1	22	33	51	42	30	-
2005 1/.....	181	-	-	-	-	-	4	24	37	53	43	20	-
2004 1/.....	149	-	-	-	-	1	1	12	27	51	42	15	-
2003 1/.....	138	-	-	-	-	1	-	19	29	40	40	9	-
2002.....	158	1	-	-	-	-	4	13	35	46	43	16	-
2001.....	136	-	-	-	-	-	2	13	29	44	40	8	-
2000.....	118	-	-	-	-	-	4	7	19	39	35	14	-
1999.....	160	-	-	-	-	1	2	8	40	56	46	7	-
Asian or Pacific Islander 3/, male													
2007 1/.....	123	-	-	-	-	-	3	12	30	27	35	16	-
2006 1/.....	88	-	-	-	-	-	1	9	17	19	28	14	-
2005 1/.....	90	-	-	-	-	-	-	10	19	29	21	11	-
2004 1/.....	80	-	-	-	-	1	-	8	17	22	23	9	-
2003 1/.....	70	-	-	-	-	-	-	10	14	25	18	3	-
2002.....	69	-	-	-	-	-	-	3	20	19	21	6	-
2001.....	72	-	-	-	-	-	1	8	16	27	15	5	-
2000.....	55	-	-	-	-	-	3	6	7	18	15	6	-
1999.....	77	-	-	-	-	-	1	5	22	25	19	5	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

Asian or Pacific Islander 3/, female

2007 1/.....	94	-	-	-	1	-	1	7	16	22	38	9	-
2006 1/.....	92	-	-	-	-	1	-	13	16	32	14	16	-
2005 1/.....	91	-	-	-	-	-	4	14	18	24	22	9	-
2004 1/.....	69	-	-	-	-	-	1	4	10	29	19	6	-
2003 1/.....	68	-	-	-	-	1	-	9	15	15	22	6	-
2002.....	89	1	-	-	-	-	4	10	15	27	22	10	-
2001.....	64	-	-	-	-	-	1	5	13	17	25	3	-
2000.....	63	-	-	-	-	-	1	1	12	21	20	8	-
1999.....	83	-	-	-	-	1	1	3	18	31	27	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)

All races, both sexes

2007.....	60	3	-	2	2	-	1	2	7	10	19	14	-
2006.....	69	-	1	1	-	1	2	2	9	9	23	21	-
2005.....	60	1	2	1	-	1	3	2	10	14	16	10	-
2004.....	71	-	-	-	-	2	2	5	12	12	26	12	-
2003.....	55	1	-	1	-	1	1	7	2	15	14	13	-
2002.....	73	3	-	-	1	-	-	7	3	18	25	16	-
2001.....	90	1	1	-	-	1	4	4	8	23	28	20	-
2000.....	100	-	2	1	1	-	4	8	12	22	29	21	-
1999.....	83	1	1	-	-	1	4	9	12	17	25	13	-

All races, male

2007.....	34	3	-	2	2	-	1	1	5	5	10	5	-
2006.....	37	-	1	-	-	1	1	1	6	7	11	9	-
2005.....	40	1	1	-	-	-	3	1	7	10	11	6	-
2004.....	35	-	-	-	-	2	2	4	5	4	11	7	-
2003.....	31	1	-	-	-	1	1	4	2	9	8	5	-
2002.....	44	3	-	-	1	-	-	4	2	15	12	7	-
2001.....	51	1	-	-	-	1	3	3	5	18	14	6	-
2000.....	60	-	2	1	1	-	3	6	9	12	17	9	-
1999.....	39	1	-	-	-	1	1	4	7	8	14	3	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)

All races, female

2007.....	26	-	-	-	-	-	-	1	2	5	9	9	-
2006.....	32	-	-	1	-	-	1	1	3	2	12	12	-
2005.....	20	-	1	1	-	1	-	1	3	4	5	4	-
2004.....	36	-	-	-	-	-	-	1	7	8	15	5	-
2003.....	24	-	-	1	-	-	-	3	-	6	6	8	-
2002.....	29	-	-	-	-	-	-	3	1	3	13	9	-
2001.....	39	-	1	-	-	-	1	1	3	5	14	14	-
2000.....	40	-	-	-	-	-	1	2	3	10	12	12	-
1999.....	44	-	1	-	-	-	3	5	5	9	11	10	-

White, both sexes

2007 1/.....	46	-	-	1	1	-	-	1	6	9	17	11	-
2006 1/.....	61	-	1	-	-	-	1	1	7	8	23	20	-
2005 1/.....	53	1	1	1	-	1	3	2	9	12	14	9	-
2004 1/.....	61	-	-	-	-	1	2	4	10	12	22	10	-
2003 1/.....	50	1	-	1	-	1	1	5	2	14	13	12	-
2002.....	61	1	-	-	1	-	-	6	3	17	20	13	-
2001.....	77	1	1	-	-	-	3	4	5	21	23	19	-
2000.....	89	-	2	1	-	-	3	6	11	21	26	19	-
1999.....	76	1	1	-	-	1	3	7	11	17	23	12	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)

White, male

2007 1/.....	24	-	-	1	1	-	-	1	4	4	10	3	-
2006 1/.....	31	-	1	-	-	-	-	-	5	6	11	8	-
2005 1/.....	36	1	1	-	-	-	3	1	6	8	10	6	-
2004 1/.....	32	-	-	-	-	1	2	4	5	4	10	6	-
2003 1/.....	28	1	-	-	-	1	1	2	2	9	7	5	-
2002.....	38	1	-	-	1	-	-	4	2	14	10	6	-
2001.....	44	1	-	-	-	-	2	3	4	17	12	5	-
2000.....	51	-	2	1	-	-	2	4	8	11	16	7	-
1999.....	36	1	-	-	-	1	1	3	6	8	13	3	-

White, female

2007 1/.....	22	-	-	-	-	-	-	-	2	5	7	8	-
2006 1/.....	30	-	-	-	-	-	1	1	2	2	12	12	-
2005 1/.....	17	-	-	1	-	1	-	1	3	4	4	3	-
2004 1/.....	29	-	-	-	-	-	-	-	5	8	12	4	-
2003 1/.....	22	-	-	1	-	-	-	3	-	5	6	7	-
2002.....	23	-	-	-	-	-	-	2	1	3	10	7	-
2001.....	33	-	1	-	-	-	1	1	1	4	11	14	-
2000.....	38	-	-	-	-	-	1	2	3	10	10	12	-
1999.....	40	-	1	-	-	-	2	4	5	9	10	9	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)

Black, both sexes

2007 1/.....	11	2	-	1	-	-	-	1	1	1	2	3	-
2006 1/.....	7	-	-	1	-	1	1	1	2	1	-	-	-
2005 1/.....	6	-	1	-	-	-	-	-	1	2	1	1	-
2004 1/.....	8	-	-	-	-	-	-	1	2	-	3	2	-
2003 1/.....	3	-	-	-	-	-	-	1	-	1	-	1	-
2002.....	6	-	-	-	-	-	-	-	-	1	3	2	-
2001.....	12	-	-	-	-	1	1	-	3	2	4	1	-
2000.....	9	-	-	-	1	-	-	2	1	-	3	2	-
1999.....	6	-	-	-	-	-	1	2	1	-	1	1	-

Black, male

2007 1/.....	7	2	-	1	-	-	-	-	1	1	-	2	-
2006 1/.....	5	-	-	-	-	1	1	1	1	1	-	-	-
2005 1/.....	3	-	-	-	-	-	-	-	1	2	-	-	-
2004 1/.....	1	-	-	-	-	-	-	-	-	-	-	1	-
2003 1/.....	1	-	-	-	-	-	-	1	-	-	-	-	-
2002.....	2	-	-	-	-	-	-	-	-	1	1	-	-
2001.....	7	-	-	-	-	1	1	-	1	1	2	1	-
2000.....	7	-	-	-	1	-	-	2	1	-	1	2	-
1999.....	3	-	-	-	-	-	-	1	1	-	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)

Asian or Pacific Islander 3/, female

2007 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2006 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2005 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	-	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

All races, both sexes

2007.....	63,192	20	113	245	558	700	1,742	5,436	10,942	13,854	18,188	11,394	-
2006.....	63,100	20	111	244	521	717	1,783	5,516	10,584	14,193	18,283	11,127	1
2005.....	62,851	15	122	279	579	701	1,827	5,558	10,422	14,019	18,480	10,848	1
2004.....	61,911	23	126	275	548	696	1,876	5,276	9,997	14,187	18,549	10,356	2
2003.....	62,639	19	106	274	516	690	1,885	5,611	9,820	14,277	18,880	10,556	5
2002.....	62,969	24	124	288	527	734	2,043	5,493	9,723	14,757	18,685	10,570	1
2001.....	62,708	17	117	279	516	728	2,055	5,506	9,441	15,023	18,700	10,326	-
2000.....	63,060	24	129	260	511	701	2,090	5,307	9,293	15,716	18,860	10,165	4
1999.....	66,251	14	116	291	519	762	2,314	5,477	9,863	16,834	19,656	10,403	2

All races, male

2007.....	32,887	8	56	117	371	437	908	3,076	6,329	7,704	9,245	4,636	-
2006.....	32,364	10	62	132	345	449	934	3,039	6,083	7,713	9,214	4,382	1
2005.....	32,194	9	68	151	362	440	1,001	3,111	6,002	7,687	9,137	4,226	-
2004.....	31,654	13	59	158	357	420	1,014	2,945	5,679	7,798	9,281	3,929	1
2003.....	31,845	12	66	152	329	427	1,046	3,143	5,573	7,800	9,321	3,973	3
2002.....	32,122	15	65	152	340	459	1,110	3,131	5,521	8,141	9,146	4,041	1
2001.....	31,663	7	73	153	346	442	1,108	3,164	5,326	8,177	9,032	3,835	-
2000.....	31,728	13	75	140	319	432	1,141	2,959	5,223	8,572	9,161	3,689	4
1999.....	33,449	8	64	160	329	490	1,300	3,098	5,597	9,157	9,439	3,806	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

All races, female

2007.....	30,305	12	57	128	187	263	834	2,360	4,613	6,150	8,943	6,758	-
2006.....	30,736	10	49	112	176	268	849	2,477	4,501	6,480	9,069	6,745	-
2005.....	30,657	6	54	128	217	261	826	2,447	4,420	6,332	9,343	6,622	1
2004.....	30,257	10	67	117	191	276	862	2,331	4,318	6,389	9,268	6,427	1
2003.....	30,794	7	40	122	187	263	839	2,468	4,247	6,477	9,559	6,583	2
2002.....	30,847	9	59	136	187	275	933	2,362	4,202	6,616	9,539	6,529	-
2001.....	31,045	10	44	126	170	286	947	2,342	4,115	6,846	9,668	6,491	-
2000.....	31,332	11	54	120	192	269	949	2,348	4,070	7,144	9,699	6,476	-
1999.....	32,802	6	52	131	190	272	1,014	2,379	4,266	7,677	10,217	6,597	1

White, both sexes

2007 1/.....	54,780	18	87	195	443	545	1,350	4,278	9,064	11,932	16,368	10,500	-
2006 1/.....	54,834	19	75	194	414	559	1,408	4,373	8,809	12,288	16,512	10,182	1
2005 1/.....	54,604	13	95	210	468	533	1,446	4,367	8,693	12,122	16,702	9,955	-
2004 1/.....	53,795	19	96	208	421	563	1,467	4,137	8,333	12,279	16,739	9,531	2
2003 1/.....	54,660	15	78	216	422	532	1,476	4,444	8,250	12,442	17,078	9,702	5
2002.....	54,810	22	90	225	423	574	1,608	4,384	8,143	12,807	16,857	9,676	1
2001.....	54,510	15	98	216	422	572	1,616	4,366	7,835	13,006	16,874	9,490	-
2000.....	55,066	19	97	212	410	581	1,596	4,206	7,794	13,730	17,066	9,352	3
1999.....	57,818	12	98	223	408	615	1,804	4,400	8,205	14,764	17,782	9,505	2

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

White, male

2007 1/.....	28,674	8	42	93	299	341	703	2,448	5,301	6,704	8,425	4,310	-
2006 1/.....	28,332	10	43	104	280	362	757	2,416	5,139	6,774	8,396	4,050	1
2005 1/.....	28,050	7	55	111	293	343	814	2,470	5,039	6,670	8,342	3,906	-
2004 1/.....	27,536	10	46	124	278	357	797	2,305	4,736	6,778	8,472	3,632	1
2003 1/.....	27,830	10	46	121	268	344	829	2,495	4,703	6,835	8,505	3,671	3
2002.....	28,060	15	49	117	269	365	862	2,526	4,642	7,170	8,316	3,728	1
2001.....	27,506	6	59	117	283	351	887	2,513	4,411	7,145	8,203	3,531	-
2000.....	27,697	11	62	114	254	371	875	2,365	4,351	7,572	8,316	3,403	3
1999.....	29,229	7	55	127	252	401	1,034	2,471	4,687	8,078	8,623	3,493	1

White, female

2007 1/.....	26,106	10	45	102	144	204	647	1,830	3,763	5,228	7,943	6,190	-
2006 1/.....	26,502	9	32	90	134	197	651	1,957	3,670	5,514	8,116	6,132	-
2005 1/.....	26,554	6	40	99	175	190	632	1,897	3,654	5,452	8,360	6,049	-
2004 1/.....	26,259	9	50	84	143	206	670	1,832	3,597	5,501	8,267	5,899	1
2003 1/.....	26,830	5	32	95	154	188	647	1,949	3,547	5,607	8,573	6,031	2
2002.....	26,750	7	41	108	154	209	746	1,858	3,501	5,637	8,541	5,948	-
2001.....	27,004	9	39	99	139	221	729	1,853	3,424	5,861	8,671	5,959	-
2000.....	27,369	8	35	98	156	210	721	1,841	3,443	6,158	8,750	5,949	-
1999.....	28,589	5	43	96	156	214	770	1,929	3,518	6,686	9,159	6,012	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Black, both sexes

2007 1/.....	6,670	1	20	40	85	116	306	972	1,548	1,519	1,379	684	-
2006 1/.....	6,659	1	26	37	80	120	298	950	1,495	1,543	1,392	717	-
2005 1/.....	6,672	2	20	61	81	130	310	990	1,417	1,513	1,459	688	1
2004 1/.....	6,658	3	20	60	102	101	341	954	1,378	1,573	1,464	662	-
2003 1/.....	6,618	2	24	51	71	124	333	993	1,332	1,508	1,473	707	-
2002.....	6,824	2	27	53	82	125	363	933	1,347	1,624	1,520	748	-
2001.....	6,799	1	16	47	77	125	347	971	1,342	1,683	1,483	707	-
2000.....	6,696	4	25	41	85	88	434	935	1,267	1,657	1,485	674	1
1999.....	7,067	1	14	50	84	108	426	931	1,422	1,721	1,549	761	-

Black, male

2007 1/.....	3,342	-	12	19	49	72	157	536	872	772	611	242	-
2006 1/.....	3,293	-	17	20	46	65	144	535	810	765	652	239	-
2005 1/.....	3,352	2	10	36	48	76	154	532	806	818	632	238	-
2004 1/.....	3,381	2	8	29	61	50	175	549	787	840	656	224	-
2003 1/.....	3,324	2	16	27	46	62	176	554	753	791	665	232	-
2002.....	3,381	-	12	29	56	67	201	516	751	818	682	249	-
2001.....	3,436	-	13	27	52	70	171	549	763	883	658	250	-
2000.....	3,378	1	11	24	56	46	231	513	742	838	695	220	1
1999.....	3,547	-	8	22	57	61	222	542	808	906	671	250	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Black, female

2007 1/.....	3,328	1	8	21	36	44	149	436	676	747	768	442	-
2006 1/.....	3,366	1	9	17	34	55	154	415	685	778	740	478	-
2005 1/.....	3,320	-	10	25	33	54	156	458	611	695	827	450	1
2004 1/.....	3,277	1	12	31	41	51	166	405	591	733	808	438	-
2003 1/.....	3,294	-	8	24	25	62	157	439	579	717	808	475	-
2002.....	3,443	2	15	24	26	58	162	417	596	806	838	499	-
2001.....	3,363	1	3	20	25	55	176	422	579	800	825	457	-
2000.....	3,318	3	14	17	29	42	203	422	525	819	790	454	-
1999.....	3,520	1	6	28	27	47	204	389	614	815	878	511	-

American Indian 2/, both sexes

2007 1/.....	311	-	1	1	5	7	9	33	68	85	72	30	-
2006 1/.....	344	-	2	3	6	9	14	43	61	90	76	40	-
2005 1/.....	292	-	1	1	4	3	12	35	74	82	45	35	-
2004 1/.....	285	-	1	1	7	5	14	33	67	65	65	27	-
2003 1/.....	275	-	1	-	3	5	23	37	48	68	58	32	-
2002.....	265	-	-	3	4	7	16	29	57	59	58	32	-
2001.....	289	-	1	1	8	6	22	31	60	67	63	30	-
2000.....	250	-	-	-	4	5	8	30	47	73	55	28	-
1999.....	300	-	-	3	4	8	20	39	63	72	70	21	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

American Indian 2/, male

2007 1/.....	158	-	-	1	4	5	5	20	37	42	29	15	-
2006 1/.....	144	-	-	2	4	6	5	19	28	34	32	14	-
2005 1/.....	152	-	-	1	3	3	9	22	34	44	20	16	-
2004 1/.....	140	-	-	1	3	1	8	18	38	36	26	9	-
2003 1/.....	147	-	1	-	3	3	12	20	25	38	28	17	-
2002.....	130	-	-	2	1	6	8	14	29	37	27	6	-
2001.....	150	-	-	1	6	4	12	20	32	37	28	10	-
2000.....	118	-	-	-	2	3	4	18	25	35	22	9	-
1999.....	140	-	-	3	2	5	10	23	25	35	29	8	-

American Indian 2/, female

2007 1/.....	153	-	1	-	1	2	4	13	31	43	43	15	-
2006 1/.....	200	-	2	1	2	3	9	24	33	56	44	26	-
2005 1/.....	140	-	1	-	1	-	3	13	40	38	25	19	-
2004 1/.....	145	-	1	-	4	4	6	15	29	29	39	18	-
2003 1/.....	128	-	-	-	-	2	11	17	23	30	30	15	-
2002.....	135	-	-	1	3	1	8	15	28	22	31	26	-
2001.....	139	-	1	-	2	2	10	11	28	30	35	20	-
2000.....	132	-	-	-	2	2	4	12	22	38	33	19	-
1999.....	160	-	-	-	2	3	10	16	38	37	41	13	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	1,431	1	5	9	25	32	77	153	262	318	369	180	-
2006 1/.....	1,263	-	8	10	21	29	63	150	219	272	303	188	-
2005 1/.....	1,283	-	6	7	26	35	59	166	238	302	274	170	-
2004 1/.....	1,173	1	9	6	18	27	54	152	219	270	281	136	-
2003 1/.....	1,086	2	3	7	20	29	53	137	190	259	271	115	-
2002.....	1,070	-	7	7	18	28	56	147	176	267	250	114	-
2001.....	1,110	1	2	15	9	25	70	138	204	267	280	99	-
2000.....	1,048	1	7	7	12	27	52	136	185	256	254	111	-
1999.....	1,066	1	4	15	23	31	64	107	173	277	255	116	-

Asian or Pacific Islander 3/, male

2007 1/.....	713	-	2	4	19	19	43	72	119	186	180	69	-
2006 1/.....	595	-	2	6	15	16	28	69	106	140	134	79	-
2005 1/.....	640	-	3	3	18	18	24	87	123	155	143	66	-
2004 1/.....	597	1	5	4	15	12	34	73	118	144	127	64	-
2003 1/.....	544	-	3	4	12	18	29	74	92	136	123	53	-
2002.....	551	-	4	4	14	21	39	75	99	116	121	58	-
2001.....	571	1	1	8	5	17	38	82	120	112	143	44	-
2000.....	535	1	2	2	7	12	31	63	105	127	128	57	-
1999.....	533	1	1	8	18	23	34	62	77	138	116	55	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Asian or Pacific Islander 3/, female

2007 1/.....	718	1	3	5	6	13	34	81	143	132	189	111	-
2006 1/.....	668	-	6	4	6	13	35	81	113	132	169	109	-
2005 1/.....	643	-	3	4	8	17	35	79	115	147	131	104	-
2004 1/.....	576	-	4	2	3	15	20	79	101	126	154	72	-
2003 1/.....	542	2	-	3	8	11	24	63	98	123	148	62	-
2002.....	519	-	3	3	4	7	17	72	77	151	129	56	-
2001.....	539	-	1	7	4	8	32	56	84	155	137	55	-
2000.....	513	-	5	5	5	15	21	73	80	129	126	54	-
1999.....	533	-	3	7	5	8	30	45	96	139	139	61	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

All races, both sexes

2007.....	14,204	59	59	84	81	138	340	699	1,416	2,402	4,717	4,209	-
2006.....	14,122	65	60	76	100	143	316	794	1,315	2,369	4,750	4,134	-
2005.....	13,710	59	52	76	93	162	368	729	1,318	2,350	4,594	3,909	-
2004.....	13,580	71	53	84	110	173	318	698	1,310	2,343	4,644	3,774	2
2003.....	13,563	62	51	79	96	178	339	682	1,238	2,354	4,689	3,795	-
2002.....	13,299	69	60	89	87	163	366	714	1,193	2,348	4,597	3,613	-
2001.....	13,676	51	58	105	102	151	366	719	1,253	2,574	4,654	3,643	-
2000.....	13,546	60	53	99	95	165	368	690	1,224	2,546	4,645	3,600	1
1999.....	13,227	58	63	101	98	171	342	662	1,133	2,684	4,472	3,443	-

All races, male

2007.....	7,256	24	28	47	43	89	181	382	838	1,354	2,570	1,700	-
2006.....	6,984	33	24	41	51	67	167	432	732	1,360	2,519	1,558	-
2005.....	6,820	22	31	36	54	86	186	404	754	1,342	2,419	1,486	-
2004.....	6,764	35	28	46	63	94	172	382	750	1,322	2,378	1,492	2
2003.....	6,687	32	31	43	54	98	195	366	685	1,334	2,379	1,470	-
2002.....	6,536	30	30	49	50	84	196	365	680	1,315	2,332	1,405	-
2001.....	6,735	26	30	52	53	77	193	391	673	1,439	2,403	1,398	-
2000.....	6,559	32	33	49	62	82	188	385	657	1,415	2,319	1,336	1
1999.....	6,520	30	33	49	54	96	197	362	656	1,522	2,218	1,303	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

All races, female

2007.....	6,948	35	31	37	38	49	159	317	578	1,048	2,147	2,509	-
2006.....	7,138	32	36	35	49	76	149	362	583	1,009	2,231	2,576	-
2005.....	6,890	37	21	40	39	76	182	325	564	1,008	2,175	2,423	-
2004.....	6,816	36	25	38	47	79	146	316	560	1,021	2,266	2,282	-
2003.....	6,876	30	20	36	42	80	144	316	553	1,020	2,310	2,325	-
2002.....	6,763	39	30	40	37	79	170	349	513	1,033	2,265	2,208	-
2001.....	6,941	25	28	53	49	74	173	328	580	1,135	2,251	2,245	-
2000.....	6,987	28	20	50	33	83	180	305	567	1,131	2,326	2,264	-
1999.....	6,707	28	30	52	44	75	145	300	477	1,162	2,254	2,140	-

White, both sexes

2007 1/.....	12,655	47	49	60	59	104	263	569	1,190	2,085	4,301	3,928	-
2006 1/.....	12,553	50	51	59	70	107	233	641	1,109	2,089	4,335	3,809	-
2005 1/.....	12,249	42	42	57	76	129	282	577	1,109	2,067	4,220	3,648	-
2004 1/.....	12,092	59	43	59	86	129	240	557	1,093	2,065	4,244	3,515	2
2003 1/.....	12,181	48	39	55	78	144	262	549	1,061	2,081	4,331	3,533	-
2002.....	11,913	59	43	67	68	121	286	583	1,023	2,091	4,222	3,350	-
2001.....	12,261	37	48	77	74	126	290	610	1,075	2,288	4,292	3,344	-
2000.....	12,171	48	44	76	73	129	292	558	1,058	2,239	4,302	3,352	-
1999.....	11,849	41	48	84	70	124	269	534	961	2,385	4,120	3,213	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)													
White, male													
2007 1/.....	6,470	18	25	31	30	67	140	319	705	1,177	2,370	1,588	-
2006 1/.....	6,254	26	19	33	39	52	130	364	629	1,215	2,322	1,425	-
2005 1/.....	6,161	16	26	29	47	69	146	326	644	1,206	2,251	1,401	-
2004 1/.....	6,071	30	22	33	52	74	132	312	631	1,186	2,192	1,405	2
2003 1/.....	6,036	25	22	30	46	77	158	302	583	1,200	2,215	1,378	-
2002.....	5,902	27	21	37	38	70	165	308	582	1,189	2,155	1,310	-
2001.....	6,089	20	24	37	37	64	159	342	583	1,304	2,227	1,292	-
2000.....	5,900	25	25	39	48	70	147	319	564	1,247	2,168	1,248	-
1999.....	5,891	21	22	40	38	67	164	302	569	1,372	2,062	1,234	-
White, female													
2007 1/.....	6,185	29	24	29	29	37	123	250	485	908	1,931	2,340	-
2006 1/.....	6,299	24	32	26	31	55	103	277	480	874	2,013	2,384	-
2005 1/.....	6,088	26	16	28	29	60	136	251	465	861	1,969	2,247	-
2004 1/.....	6,021	29	21	26	34	55	108	245	462	879	2,052	2,110	-
2003 1/.....	6,145	23	17	25	32	67	104	247	478	881	2,116	2,155	-
2002.....	6,011	32	22	30	30	51	121	275	441	902	2,067	2,040	-
2001.....	6,172	17	24	40	37	62	131	268	492	984	2,065	2,052	-
2000.....	6,271	23	19	37	25	59	145	239	494	992	2,134	2,104	-
1999.....	5,958	20	26	44	32	57	105	232	392	1,013	2,058	1,979	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Black, both sexes

2007 1/.....	1,190	8	9	21	19	29	65	106	174	251	310	198	-
2006 1/.....	1,232	13	9	15	25	31	64	129	174	210	322	240	-
2005 1/.....	1,151	15	5	14	11	27	77	125	165	226	282	204	-
2004 1/.....	1,192	10	8	21	19	38	67	112	180	222	313	202	-
2003 1/.....	1,123	11	9	18	16	29	67	118	144	223	290	198	-
2002.....	1,143	9	14	18	15	31	70	114	144	212	302	214	-
2001.....	1,157	11	5	26	21	18	61	90	149	239	298	239	-
2000.....	1,136	9	9	19	16	33	63	114	135	248	284	205	1
1999.....	1,173	15	13	17	26	30	66	106	149	258	299	194	-

Black, male

2007 1/.....	594	4	3	15	11	19	33	53	105	136	142	73	-
2006 1/.....	567	6	5	7	11	13	25	56	90	111	153	90	-
2005 1/.....	499	6	2	4	5	13	35	63	85	103	124	59	-
2004 1/.....	551	4	5	12	10	16	33	57	100	111	141	62	-
2003 1/.....	525	5	8	9	7	18	31	55	83	115	130	64	-
2002.....	500	2	8	9	9	8	26	47	83	99	135	74	-
2001.....	525	5	3	13	13	11	30	38	74	119	141	78	-
2000.....	520	4	8	8	9	11	32	55	76	131	119	66	1
1999.....	521	8	10	9	15	20	30	46	75	129	125	54	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Black, female

2007 1/.....	596	4	6	6	8	10	32	53	69	115	168	125	-
2006 1/.....	665	7	4	8	14	18	39	73	84	99	169	150	-
2005 1/.....	652	9	3	10	6	14	42	62	80	123	158	145	-
2004 1/.....	641	6	3	9	9	22	34	55	80	111	172	140	-
2003 1/.....	598	6	1	9	9	11	36	63	61	108	160	134	-
2002.....	643	7	6	9	6	23	44	67	61	113	167	140	-
2001.....	632	6	2	13	8	7	31	52	75	120	157	161	-
2000.....	616	5	1	11	7	22	31	59	59	117	165	139	-
1999.....	652	7	3	8	11	10	36	60	74	129	174	140	-

American Indian 2/, both sexes

2007 1/.....	51	-	-	1	-	-	1	2	9	14	15	9	-
2006 1/.....	65	1	-	1	3	-	6	4	6	11	15	18	-
2005 1/.....	51	-	-	-	2	1	2	8	6	5	14	13	-
2004 1/.....	48	-	1	1	1	1	4	4	7	9	9	11	-
2003 1/.....	39	-	-	1	-	2	2	1	8	9	9	7	-
2002.....	34	-	2	1	-	1	3	2	4	9	8	4	-
2001.....	58	1	-	1	4	1	6	3	3	16	14	9	-
2000.....	45	-	-	1	4	2	4	2	6	10	7	9	-
1999.....	42	-	-	-	1	1	-	3	7	13	9	8	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

American Indian 2/, male

2007 1/.....	26	-	-	-	-	-	-	1	8	7	7	3	-
2006 1/.....	31	-	-	-	-	-	4	3	4	5	8	7	-
2005 1/.....	23	-	-	-	1	1	1	5	4	2	6	3	-
2004 1/.....	23	-	-	1	-	1	3	1	2	5	5	5	-
2003 1/.....	20	-	-	1	-	1	-	1	5	3	6	3	-
2002.....	15	-	-	1	-	-	-	-	3	4	5	2	-
2001.....	29	1	-	1	2	-	2	3	2	7	6	5	-
2000.....	20	-	-	1	3	1	1	-	3	6	3	2	-
1999.....	22	-	-	-	1	1	-	2	4	5	6	3	-

American Indian 2/, female

2007 1/.....	25	-	-	1	-	-	1	1	1	7	8	6	-
2006 1/.....	34	1	-	1	3	-	2	1	2	6	7	11	-
2005 1/.....	28	-	-	-	1	-	1	3	2	3	8	10	-
2004 1/.....	25	-	1	-	1	-	1	3	5	4	4	6	-
2003 1/.....	19	-	-	-	-	1	2	-	3	6	3	4	-
2002.....	19	-	2	-	-	1	3	2	1	5	3	2	-
2001.....	29	-	-	-	2	1	4	-	1	9	8	4	-
2000.....	25	-	-	-	1	1	3	2	3	4	4	7	-
1999.....	20	-	-	-	-	-	-	1	3	8	3	5	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	308	4	1	2	3	5	11	22	43	52	91	74	-
2006 1/.....	272	1	-	1	2	5	13	20	26	59	78	67	-
2005 1/.....	259	2	5	5	4	7	19	38	52	78	44	44	-
2004 1/.....	248	2	1	3	4	5	7	25	30	47	78	46	-
2003 1/.....	220	3	3	5	2	3	8	14	25	41	59	57	-
2002.....	209	1	1	3	4	10	7	15	22	36	65	45	-
2001.....	200	2	5	1	3	6	9	16	26	31	50	51	-
2000.....	194	3	-	3	2	1	9	16	25	49	52	34	-
1999.....	163	2	2	-	1	16	7	19	16	28	44	28	-

Asian or Pacific Islander 3/, male

2007 1/.....	166	2	-	1	2	3	8	9	20	34	51	36	-
2006 1/.....	132	1	-	1	1	2	8	9	9	29	36	36	-
2005 1/.....	137	-	3	3	1	3	4	10	21	31	38	23	-
2004 1/.....	119	1	1	-	1	3	4	12	17	20	40	20	-
2003 1/.....	106	2	1	3	1	2	6	8	14	16	28	25	-
2002.....	119	1	1	2	3	6	5	10	12	23	37	19	-
2001.....	92	-	3	1	1	2	2	8	14	9	29	23	-
2000.....	119	3	-	1	2	-	8	11	14	31	29	20	-
1999.....	86	1	1	-	-	8	3	12	8	16	25	12	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Asian or Pacific Islander 3/, female

2007 1/.....	142	2	1	1	1	2	3	13	23	18	40	38	-
2006 1/.....	140	-	-	-	1	3	5	11	17	30	42	31	-
2005 1/.....	122	2	2	2	3	2	3	9	17	21	40	21	-
2004 1/.....	129	1	-	3	3	2	3	13	13	27	38	26	-
2003 1/.....	114	1	2	2	1	1	2	6	11	25	31	32	-
2002.....	90	-	-	1	1	4	2	5	10	13	28	26	-
2001.....	108	2	2	-	2	4	7	8	12	22	21	28	-
2000.....	75	-	-	2	-	1	1	5	11	18	23	14	-
1999.....	77	1	1	-	1	8	4	7	8	12	19	16	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Anemias (D50-D64)

All races, both sexes

2007.....	4,829	17	17	32	90	136	187	255	322	560	1,224	1,989	-
2006.....	3,996	11	26	34	93	141	192	232	304	401	975	1,587	-
2005.....	4,624	19	23	26	103	157	189	253	304	494	1,193	1,863	-
2004.....	4,575	15	33	46	128	164	201	236	309	473	1,119	1,850	1
2003.....	4,594	22	24	47	105	157	183	241	276	484	1,136	1,919	-
2002.....	4,614	18	21	53	111	180	196	250	264	508	1,161	1,852	-
2001.....	4,627	16	22	55	109	160	215	232	249	492	1,226	1,851	-
2000.....	4,494	10	35	52	85	153	190	227	260	515	1,171	1,796	-
1999.....	4,503	12	30	53	121	160	194	232	231	486	1,164	1,820	-

All races, male

2007.....	1,940	6	6	21	49	71	98	125	154	264	536	610	-
2006.....	1,592	5	15	16	55	71	89	99	143	179	421	499	-
2005.....	1,787	13	9	14	57	85	97	108	141	224	502	537	-
2004.....	1,833	9	18	21	69	90	107	112	164	226	466	550	1
2003.....	1,832	11	14	23	52	83	91	125	156	251	477	549	-
2002.....	1,781	12	9	26	67	101	96	109	131	244	454	532	-
2001.....	1,774	11	12	27	59	95	91	115	111	237	496	520	-
2000.....	1,669	6	16	29	39	70	78	108	130	236	465	492	-
1999.....	1,746	8	17	23	67	81	101	116	105	226	484	518	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Anemias (D50-D64)													
All races, female													
2007.....	2,889	11	11	11	41	65	89	130	168	296	688	1,379	-
2006.....	2,404	6	11	18	38	70	103	133	161	222	554	1,088	-
2005.....	2,837	6	14	12	46	72	92	145	163	270	691	1,326	-
2004.....	2,742	6	15	25	59	74	94	124	145	247	653	1,300	-
2003.....	2,762	11	10	24	53	74	92	116	120	233	659	1,370	-
2002.....	2,833	6	12	27	44	79	100	141	133	264	707	1,320	-
2001.....	2,853	5	10	28	50	65	124	117	138	255	730	1,331	-
2000.....	2,825	4	19	23	46	83	112	119	130	279	706	1,304	-
1999.....	2,757	4	13	30	54	79	93	116	126	260	680	1,302	-
White, both sexes													
2007 1/.....	3,734	4	5	18	22	31	42	105	211	437	1,060	1,799	-
2006 1/.....	2,989	3	13	13	25	23	43	95	198	320	849	1,407	-
2005 1/.....	3,499	9	11	11	22	25	37	114	202	372	1,021	1,675	-
2004 1/.....	3,456	8	13	20	35	33	55	96	184	379	981	1,651	1
2003 1/.....	3,476	9	10	21	23	39	45	90	153	374	985	1,727	-
2002.....	3,460	8	12	23	30	43	45	99	161	399	990	1,650	-
2001.....	3,543	6	11	27	33	39	53	92	168	393	1,059	1,662	-
2000.....	3,450	4	15	21	21	29	43	94	169	410	1,031	1,613	-
1999.....	3,393	6	9	20	34	17	48	96	122	389	1,032	1,620	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Anemias (D50-D64)													
White, male													
2007 1/.....	1,447	2	2	13	11	13	19	50	103	212	471	551	-
2006 1/.....	1,163	2	5	6	16	13	22	43	88	145	368	455	-
2005 1/.....	1,288	6	4	5	10	16	15	50	96	171	428	487	-
2004 1/.....	1,322	4	6	8	19	18	31	53	95	186	402	499	1
2003 1/.....	1,319	4	5	11	11	22	22	51	85	189	424	495	-
2002.....	1,267	4	5	11	17	24	26	51	83	189	389	468	-
2001.....	1,303	5	5	16	18	20	21	40	75	187	441	475	-
2000.....	1,235	2	8	12	8	15	21	43	87	199	407	433	-
1999.....	1,244	4	6	7	14	6	24	51	48	182	434	468	-
White, female													
2007 1/.....	2,287	2	3	5	11	18	23	55	108	225	589	1,248	-
2006 1/.....	1,826	1	8	7	9	10	21	52	110	175	481	952	-
2005 1/.....	2,211	3	7	6	12	9	22	64	106	201	593	1,188	-
2004 1/.....	2,134	4	7	12	16	15	24	43	89	193	579	1,152	-
2003 1/.....	2,157	5	5	10	12	17	23	39	68	185	561	1,232	-
2002.....	2,193	4	7	12	13	19	19	48	78	210	601	1,182	-
2001.....	2,240	1	6	11	15	19	32	52	93	206	618	1,187	-
2000.....	2,215	2	7	9	13	14	22	51	82	211	624	1,180	-
1999.....	2,149	2	3	13	20	11	24	45	74	207	598	1,152	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Anemias (D50-D64)													
Black, both sexes													
2007 1/.....	970	7	12	13	67	102	142	140	100	110	134	143	-
2006 1/.....	917	3	12	20	64	112	146	131	100	74	105	150	-
2005 1/.....	1,032	6	11	15	77	131	147	136	92	109	149	159	-
2004 1/.....	1,029	4	19	23	93	129	140	134	116	83	117	171	-
2003 1/.....	1,037	6	12	25	79	114	134	145	116	95	142	169	-
2002.....	1,064	6	9	26	79	130	145	145	94	100	154	176	-
2001.....	1,001	6	11	26	73	119	154	137	75	91	151	158	-
2000.....	967	5	19	31	62	124	141	128	83	89	122	163	-
1999.....	1,019	3	21	33	84	140	142	126	101	82	121	166	-
Black, male													
2007 1/.....	437	2	4	8	37	56	78	69	46	45	53	39	-
2006 1/.....	388	-	9	9	37	55	65	55	51	30	44	33	-
2005 1/.....	456	6	5	9	45	69	79	58	35	47	63	40	-
2004 1/.....	472	3	12	12	50	71	74	57	64	36	50	43	-
2003 1/.....	487	4	7	12	40	58	69	74	67	56	53	47	-
2002.....	479	5	4	14	50	73	68	56	46	51	59	53	-
2001.....	446	4	7	10	41	74	66	75	35	46	51	37	-
2000.....	395	4	8	17	30	55	55	64	37	31	46	48	-
1999.....	457	2	11	16	50	73	73	61	54	37	42	38	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Anemias (D50-D64)

Black, female

2007 1/.....	533	5	8	5	30	46	64	71	54	65	81	104	-
2006 1/.....	529	3	3	11	27	57	81	76	49	44	61	117	-
2005 1/.....	576	-	6	6	32	62	68	78	57	62	86	119	-
2004 1/.....	557	1	7	11	43	58	66	77	52	47	67	128	-
2003 1/.....	550	2	5	13	39	56	65	71	49	39	89	122	-
2002.....	585	1	5	12	29	57	77	89	48	49	95	123	-
2001.....	555	2	4	16	32	45	88	62	40	45	100	121	-
2000.....	572	1	11	14	32	69	86	64	46	58	76	115	-
1999.....	562	1	10	17	34	67	69	65	47	45	79	128	-

American Indian 2/, both sexes

2007 1/.....	21	-	-	-	1	-	2	1	4	5	3	5	-
2006 1/.....	14	-	-	-	-	1	-	2	2	1	3	5	-
2005 1/.....	24	-	-	-	1	-	-	1	4	7	2	9	-
2004 1/.....	13	-	-	1	-	-	1	-	1	3	3	4	-
2003 1/.....	15	-	-	-	-	1	2	2	2	5	2	1	-
2002.....	15	-	-	1	1	2	1	2	2	-	2	4	-
2001.....	27	-	-	1	2	-	4	1	1	1	7	10	-
2000.....	20	-	-	-	-	-	2	1	1	4	8	4	-
1999.....	21	-	-	-	2	1	2	4	4	3	-	5	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Anemias (D50-D64)

American Indian 2/, male

2007 1/.....	9	-	-	-	1	-	-	1	1	2	2	2	-
2006 1/.....	5	-	-	-	-	-	-	-	1	1	1	2	-
2005 1/.....	13	-	-	-	-	-	-	4	4	1	4	4	-
2004 1/.....	6	-	-	-	-	-	1	-	1	1	2	1	-
2003 1/.....	4	-	-	-	-	1	-	-	1	2	-	-	-
2002.....	4	-	-	-	-	1	-	-	1	-	1	1	-
2001.....	7	-	-	-	-	-	2	-	1	1	1	2	-
2000.....	9	-	-	-	-	-	1	-	1	1	6	-	-
1999.....	9	-	-	-	2	-	2	1	2	-	-	2	-

American Indian 2/, female

2007 1/.....	12	-	-	-	-	-	2	-	3	3	1	3	-
2006 1/.....	9	-	-	-	-	1	-	2	1	-	2	3	-
2005 1/.....	11	-	-	-	1	-	-	1	-	3	1	5	-
2004 1/.....	7	-	-	1	-	-	-	-	-	2	1	3	-
2003 1/.....	11	-	-	-	-	-	2	2	1	3	2	1	-
2002.....	11	-	-	1	1	1	1	2	1	-	1	3	-
2001.....	20	-	-	1	2	-	2	1	-	-	6	8	-
2000.....	11	-	-	-	-	-	1	1	-	3	2	4	-
1999.....	12	-	-	-	-	1	-	3	2	3	-	3	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Anemias (D50-D64)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	104	6	-	1	-	3	1	9	7	8	27	42	-
2006 1/.....	76	5	1	1	4	5	3	4	4	6	18	25	-
2005 1/.....	69	4	1	-	3	1	5	2	6	6	21	20	-
2004 1/.....	77	3	1	2	-	2	5	6	8	8	18	24	-
2003 1/.....	66	7	2	1	3	3	2	4	5	10	7	22	-
2002.....	75	4	-	3	1	5	5	4	7	9	15	22	-
2001.....	56	4	-	1	1	2	4	2	5	7	9	21	-
2000.....	57	1	1	-	2	-	4	4	7	12	10	16	-
1999.....	70	3	-	-	1	2	2	6	4	12	11	29	-

Asian or Pacific Islander 3/, male

2007 1/.....	47	2	-	-	-	2	1	5	4	5	10	18	-
2006 1/.....	36	3	1	1	2	3	2	1	3	3	8	9	-
2005 1/.....	30	1	-	-	2	-	3	-	6	2	10	6	-
2004 1/.....	33	2	-	1	-	1	1	2	4	3	12	7	-
2003 1/.....	22	3	2	-	1	2	-	-	3	4	-	7	-
2002.....	31	3	-	1	-	3	2	2	1	4	5	10	-
2001.....	18	2	-	1	-	1	2	-	-	3	3	6	-
2000.....	30	-	-	-	1	-	1	1	5	5	6	11	-
1999.....	36	2	-	-	1	2	2	3	1	7	8	10	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Anemias (D50-D64)

Asian or Pacific Islander 3/, female

2007 1/.....	57	4	-	1	-	1	-	4	3	3	17	24	-
2006 1/.....	40	2	-	-	2	2	1	3	1	3	10	16	-
2005 1/.....	39	3	1	-	1	1	2	2	-	4	11	14	-
2004 1/.....	44	1	1	1	-	1	4	4	4	5	6	17	-
2003 1/.....	44	4	-	1	2	1	2	4	2	6	7	15	-
2002.....	44	1	-	2	1	2	3	2	6	5	10	12	-
2001.....	38	2	-	-	1	1	2	2	5	4	6	15	-
2000.....	27	1	1	-	1	-	3	3	2	7	4	5	-
1999.....	34	1	-	-	-	-	-	3	3	5	3	19	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diabetes mellitus (E10-E14)													
All races, both sexes													
2007.....	71,382	7	5	21	168	610	1,984	5,753	11,304	15,112	21,189	15,227	2
2006.....	72,449	6	7	28	165	673	2,094	5,692	11,432	15,483	21,763	15,105	1
2005.....	75,119	1	2	38	202	617	2,045	5,691	11,301	16,183	23,136	15,903	-
2004.....	73,138	1	8	37	161	599	2,026	5,567	10,780	16,093	22,945	14,918	3
2003.....	74,219	3	8	31	160	657	2,049	5,658	10,731	16,656	23,299	14,964	3
2002.....	73,249	1	3	34	171	642	2,164	5,496	10,022	16,709	23,282	14,724	1
2001.....	71,372	2	6	39	151	595	1,958	5,343	9,570	16,731	22,805	14,171	1
2000.....	69,301	4	6	26	162	623	1,926	4,954	9,186	16,674	22,184	13,556	-
1999.....	68,399	2	8	32	155	582	1,942	4,735	9,097	16,908	21,757	13,178	3
All races, male													
2007.....	35,478	5	3	11	98	359	1,228	3,513	6,644	8,279	9,874	5,462	2
2006.....	36,006	4	3	13	96	376	1,279	3,515	6,670	8,344	10,286	5,419	1
2005.....	36,538	-	-	15	115	372	1,270	3,467	6,570	8,505	10,701	5,523	-
2004.....	35,267	1	5	16	83	340	1,258	3,376	6,153	8,430	10,569	5,035	1
2003.....	35,438	-	6	16	96	379	1,264	3,341	6,044	8,687	10,679	4,923	3
2002.....	34,301	-	1	22	98	385	1,330	3,225	5,683	8,417	10,295	4,844	1
2001.....	32,841	1	2	23	85	341	1,181	3,154	5,241	8,437	9,918	4,457	1
2000.....	31,602	4	2	13	94	376	1,151	2,821	4,943	8,226	9,738	4,234	-
1999.....	31,150	-	4	16	80	319	1,139	2,791	4,856	8,279	9,539	4,125	2

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diabetes mellitus (E10-E14)													
All races, female													
2007.....	35,904	2	2	10	70	251	756	2,240	4,660	6,833	11,315	9,765	-
2006.....	36,443	2	4	15	69	297	815	2,177	4,762	7,139	11,477	9,686	-
2005.....	38,581	1	2	23	87	245	775	2,224	4,731	7,678	12,435	10,380	-
2004.....	37,871	-	3	21	78	259	768	2,191	4,627	7,663	12,376	9,883	2
2003.....	38,781	3	2	15	64	278	785	2,317	4,687	7,969	12,620	10,041	-
2002.....	38,948	1	2	12	73	257	834	2,271	4,339	8,292	12,987	9,880	-
2001.....	38,531	1	4	16	66	254	777	2,189	4,329	8,294	12,887	9,714	-
2000.....	37,699	-	4	13	68	247	775	2,133	4,243	8,448	12,446	9,322	-
1999.....	37,249	2	4	16	75	263	803	1,944	4,241	8,629	12,218	9,053	1
White, both sexes													
2007 1/.....	56,390	5	4	15	109	379	1,344	4,123	8,378	11,652	17,411	12,968	2
2006 1/.....	57,204	5	5	19	106	406	1,441	4,059	8,482	11,915	17,898	12,867	1
2005 1/.....	59,755	1	1	25	124	372	1,428	4,079	8,458	12,476	19,212	13,579	-
2004 1/.....	58,087	-	6	19	99	396	1,421	3,993	8,030	12,456	18,988	12,677	2
2003 1/.....	59,099	2	6	22	111	431	1,439	4,003	7,987	12,934	19,358	12,803	3
2002.....	58,459	1	3	20	109	439	1,554	3,908	7,440	12,942	19,433	12,610	-
2001.....	57,180	1	6	21	82	383	1,399	3,842	7,093	13,085	19,070	12,197	1
2000.....	55,561	3	6	17	105	420	1,419	3,514	6,879	13,075	18,478	11,645	-
1999.....	54,599	2	6	22	107	409	1,384	3,355	6,609	13,230	18,217	11,257	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diabetes mellitus (E10-E14)													
White, male													
2007 1/.....	28,744	3	3	9	73	234	861	2,591	5,091	6,653	8,425	4,799	2
2006 1/.....	29,060	4	3	11	60	224	894	2,583	5,094	6,658	8,785	4,743	1
2005 1/.....	29,628	-	-	11	73	227	912	2,555	5,045	6,778	9,170	4,857	-
2004 1/.....	28,629	-	4	7	52	227	905	2,508	4,728	6,741	9,061	4,396	-
2003 1/.....	28,939	-	5	13	67	246	917	2,465	4,654	7,002	9,210	4,357	3
2002.....	28,110	-	1	17	60	266	978	2,368	4,371	6,800	8,920	4,329	-
2001.....	26,917	-	2	15	50	212	876	2,326	4,031	6,917	8,520	3,967	1
2000.....	26,009	3	2	9	64	267	866	2,077	3,823	6,739	8,419	3,740	-
1999.....	25,545	-	3	10	59	226	829	2,055	3,684	6,787	8,293	3,598	1
White, female													
2007 1/.....	27,646	2	1	6	36	145	483	1,532	3,287	4,999	8,986	8,169	-
2006 1/.....	28,144	1	2	8	46	182	547	1,476	3,388	5,257	9,113	8,124	-
2005 1/.....	30,127	1	1	14	51	145	516	1,524	3,413	5,698	10,042	8,722	-
2004 1/.....	29,458	-	2	12	47	169	516	1,485	3,302	5,715	9,927	8,281	2
2003 1/.....	30,160	2	1	9	44	185	522	1,538	3,333	5,932	10,148	8,446	-
2002.....	30,349	1	2	3	49	173	576	1,540	3,069	6,142	10,513	8,281	-
2001.....	30,263	1	4	6	32	171	523	1,516	3,062	6,168	10,550	8,230	-
2000.....	29,552	-	4	8	41	153	553	1,437	3,056	6,336	10,059	7,905	-
1999.....	29,054	2	3	12	48	183	555	1,300	2,925	6,443	9,924	7,659	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diabetes mellitus (E10-E14)													
Black, both sexes													
2007 1/.....	12,459	2	1	6	56	213	562	1,433	2,486	2,861	3,068	1,771	-
2006 1/.....	12,813	1	2	9	55	240	583	1,418	2,500	3,008	3,181	1,816	-
2005 1/.....	12,970	-	1	12	73	219	541	1,410	2,440	3,128	3,251	1,895	-
2004 1/.....	12,834	1	1	16	61	183	547	1,385	2,334	3,107	3,310	1,888	1
2003 1/.....	12,892	1	2	7	46	208	549	1,445	2,367	3,160	3,299	1,808	-
2002.....	12,687	-	-	14	59	195	560	1,399	2,224	3,175	3,249	1,811	1
2001.....	12,305	1	-	15	60	202	511	1,324	2,154	3,145	3,168	1,725	-
2000.....	12,021	1	-	9	55	189	462	1,279	2,015	3,118	3,215	1,678	-
1999.....	11,927	-	2	10	44	162	507	1,231	2,151	3,137	3,022	1,659	2
Black, male													
2007 1/.....	5,493	2	-	2	22	116	314	795	1,293	1,317	1,169	463	-
2006 1/.....	5,772	-	-	2	32	137	346	804	1,306	1,415	1,211	519	-
2005 1/.....	5,730	-	-	3	39	128	310	780	1,287	1,430	1,253	500	-
2004 1/.....	5,565	1	-	8	31	104	317	754	1,184	1,432	1,224	509	1
2003 1/.....	5,425	-	1	3	27	122	310	748	1,182	1,401	1,196	435	-
2002.....	5,207	-	-	5	36	113	319	748	1,114	1,332	1,150	389	1
2001.....	5,049	1	-	6	30	123	278	713	1,043	1,296	1,162	397	-
2000.....	4,771	1	-	4	28	99	254	654	962	1,264	1,113	392	-
1999.....	4,759	-	1	6	19	86	277	650	995	1,268	1,024	432	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diabetes mellitus (E10-E14)													
Black, female													
2007 1/.....	6,966	-	1	4	34	97	248	638	1,193	1,544	1,899	1,308	-
2006 1/.....	7,041	1	2	7	23	103	237	614	1,194	1,593	1,970	1,297	-
2005 1/.....	7,240	-	1	9	34	91	231	630	1,153	1,698	1,998	1,395	-
2004 1/.....	7,269	-	1	8	30	79	230	631	1,150	1,675	2,086	1,379	-
2003 1/.....	7,467	1	1	4	19	86	239	697	1,185	1,759	2,103	1,373	-
2002.....	7,480	-	-	9	23	82	241	651	1,110	1,843	2,099	1,422	-
2001.....	7,256	-	-	9	30	79	233	611	1,111	1,849	2,006	1,328	-
2000.....	7,250	-	-	5	27	90	208	625	1,053	1,854	2,102	1,286	-
1999.....	7,168	-	1	4	25	76	230	581	1,156	1,869	1,998	1,227	1
American Indian 2/, both sexes													
2007 1/.....	790	-	-	-	-	6	45	92	167	208	173	99	-
2006 1/.....	811	-	-	-	1	15	39	103	190	190	184	89	-
2005 1/.....	818	-	-	-	4	17	41	98	177	234	159	88	-
2004 1/.....	746	-	-	-	1	11	36	85	193	197	173	50	-
2003 1/.....	783	-	-	-	-	12	42	115	162	206	173	73	-
2002.....	744	-	-	-	2	3	33	104	167	223	162	50	-
2001.....	644	-	-	1	6	4	18	90	142	171	156	56	-
2000.....	616	-	-	-	2	7	23	78	134	186	133	53	-
1999.....	725	-	-	-	2	8	29	84	165	219	146	72	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Diabetes mellitus (E10-E14)

American Indian 2/, male

2007 1/.....	381	-	-	-	-	3	29	60	92	104	59	34	-
2006 1/.....	362	-	-	-	1	6	19	56	103	82	66	29	-
2005 1/.....	412	-	-	-	2	11	27	63	94	115	69	31	-
2004 1/.....	365	-	-	-	-	5	25	54	106	81	79	15	-
2003 1/.....	367	-	-	-	-	8	26	67	74	95	72	25	-
2002.....	336	-	-	-	1	3	21	57	85	96	57	16	-
2001.....	276	-	-	1	3	2	7	51	65	68	63	16	-
2000.....	275	-	-	-	2	6	16	41	68	70	50	22	-
1999.....	323	-	-	-	2	5	20	53	78	82	62	21	-

American Indian 2/, female

2007 1/.....	409	-	-	-	-	3	16	32	75	104	114	65	-
2006 1/.....	449	-	-	-	-	9	20	47	87	108	118	60	-
2005 1/.....	406	-	-	-	2	6	14	35	83	119	90	57	-
2004 1/.....	381	-	-	-	1	6	11	31	87	116	94	35	-
2003 1/.....	416	-	-	-	-	4	16	48	88	111	101	48	-
2002.....	408	-	-	-	1	-	12	47	82	127	105	34	-
2001.....	368	-	-	-	3	2	11	39	77	103	93	40	-
2000.....	341	-	-	-	-	1	7	37	66	116	83	31	-
1999.....	402	-	-	-	-	3	9	31	87	137	84	51	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diabetes mellitus (E10-E14)													
Asian or Pacific Islander 3/, both sexes													
2007 1/.....	1,743	-	-	-	3	12	33	105	273	391	537	389	-
2006 1/.....	1,621	-	-	-	3	12	31	112	260	370	500	333	-
2005 1/.....	1,576	-	-	1	1	9	35	104	226	345	514	341	-
2004 1/.....	1,471	-	1	2	-	9	22	104	223	333	474	303	-
2003 1/.....	1,445	-	-	2	3	6	19	95	215	356	469	280	-
2002.....	1,359	-	-	-	1	5	17	85	191	369	438	253	-
2001.....	1,243	-	-	2	3	6	30	87	181	330	411	193	-
2000.....	1,103	-	-	-	-	7	22	83	158	295	358	180	-
1999.....	1,148	-	-	-	2	3	22	65	172	322	372	190	-
Asian or Pacific Islander 3/, male													
2007 1/.....	860	-	-	-	3	6	24	67	168	205	221	166	-
2006 1/.....	812	-	-	-	3	9	20	72	167	189	224	128	-
2005 1/.....	768	-	-	1	1	6	21	69	144	182	209	135	-
2004 1/.....	708	-	1	1	-	4	11	60	135	176	205	115	-
2003 1/.....	707	-	-	-	2	3	11	61	134	189	201	106	-
2002.....	648	-	-	-	1	3	12	52	113	189	168	110	-
2001.....	599	-	-	1	2	4	20	64	102	156	173	77	-
2000.....	547	-	-	-	-	4	15	49	90	153	156	80	-
1999.....	523	-	-	-	-	2	13	33	99	142	160	74	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Diabetes mellitus (E10-E14)

Asian or Pacific Islander 3/, female

2007 1/.....	883	-	-	-	-	6	9	38	105	186	316	223	-
2006 1/.....	809	-	-	-	-	3	11	40	93	181	276	205	-
2005 1/.....	808	-	-	-	-	3	14	35	82	163	305	206	-
2004 1/.....	763	-	-	1	-	5	11	44	88	157	269	188	-
2003 1/.....	738	-	-	2	1	3	8	34	81	167	268	174	-
2002.....	711	-	-	-	-	2	5	33	78	180	270	143	-
2001.....	644	-	-	1	1	2	10	23	79	174	238	116	-
2000.....	556	-	-	-	-	3	7	34	68	142	202	100	-
1999.....	625	-	-	-	2	1	9	32	73	180	212	116	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nutritional deficiencies (E40-E64)													
All races, both sexes													
2007.....	2,852	7	5	3	4	19	36	93	201	333	808	1,343	-
2006.....	2,556	9	2	4	6	11	46	98	185	275	720	1,200	-
2005.....	3,183	5	3	4	3	19	38	118	157	353	915	1,568	-
2004.....	3,019	7	2	3	6	13	39	117	162	298	897	1,474	1
2003.....	3,338	10	2	2	7	10	35	100	168	340	992	1,672	-
2002.....	3,779	4	1	1	7	16	60	102	168	382	1,132	1,906	-
2001.....	3,704	4	4	2	10	14	44	110	153	374	1,100	1,889	-
2000.....	4,242	14	2	5	10	18	56	99	173	460	1,285	2,120	-
1999.....	4,289	9	2	1	8	12	64	106	170	429	1,248	2,240	-
All races, male													
2007.....	1,071	5	3	2	1	12	13	45	97	168	333	392	-
2006.....	913	7	-	2	3	7	20	45	82	133	299	315	-
2005.....	1,116	3	1	1	1	7	14	56	88	157	341	447	-
2004.....	1,087	5	1	2	5	6	15	50	87	145	357	413	1
2003.....	1,176	7	2	2	4	7	19	48	79	166	400	442	-
2002.....	1,321	3	1	-	4	7	31	48	93	186	434	514	-
2001.....	1,233	4	1	2	7	4	21	56	81	174	426	457	-
2000.....	1,412	10	2	2	2	9	16	46	97	221	471	536	-
1999.....	1,486	4	2	1	5	4	28	52	98	221	494	577	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nutritional deficiencies (E40-E64)													
All races, female													
2007.....	1,781	2	2	1	3	7	23	48	104	165	475	951	-
2006.....	1,643	2	2	2	3	4	26	53	103	142	421	885	-
2005.....	2,067	2	2	3	2	12	24	62	69	196	574	1,121	-
2004.....	1,932	2	1	1	1	7	24	67	75	153	540	1,061	-
2003.....	2,162	3	-	-	3	3	16	52	89	174	592	1,230	-
2002.....	2,458	1	-	1	3	9	29	54	75	196	698	1,392	-
2001.....	2,471	-	3	-	3	10	23	54	72	200	674	1,432	-
2000.....	2,830	4	-	3	8	9	40	53	76	239	814	1,584	-
1999.....	2,803	5	-	-	3	8	36	54	72	208	754	1,663	-
White, both sexes													
2007 1/.....	2,375	4	3	3	4	11	26	74	163	252	681	1,154	-
2006 1/.....	2,161	6	2	4	2	7	35	71	148	219	614	1,053	-
2005 1/.....	2,677	2	2	3	1	11	28	88	121	281	763	1,377	-
2004 1/.....	2,529	5	1	2	3	10	25	82	125	234	765	1,277	-
2003 1/.....	2,826	6	2	2	5	8	29	65	127	277	848	1,457	-
2002.....	3,213	3	1	-	6	13	41	81	121	303	976	1,668	-
2001.....	3,152	3	4	1	8	10	29	73	117	303	922	1,682	-
2000.....	3,627	8	1	2	8	9	40	73	130	370	1,112	1,874	-
1999.....	3,628	5	1	-	5	9	49	76	118	325	1,050	1,990	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nutritional deficiencies (E40-E64)													
White, male													
2007 1/.....	875	2	2	2	1	8	10	34	74	124	280	338	-
2006 1/.....	740	5	-	2	1	4	13	32	60	103	244	276	-
2005 1/.....	924	1	-	-	-	4	11	42	64	127	282	393	-
2004 1/.....	884	3	-	1	3	5	7	33	64	110	300	358	-
2003 1/.....	969	5	2	2	3	6	16	31	54	132	331	387	-
2002.....	1,102	2	1	-	3	6	22	34	59	150	369	456	-
2001.....	1,000	3	1	1	7	3	12	28	55	133	359	398	-
2000.....	1,175	6	1	2	2	4	11	33	65	172	404	475	-
1999.....	1,199	2	1	-	3	3	20	39	64	158	407	502	-
White, female													
2007 1/.....	1,500	2	1	1	3	3	16	40	89	128	401	816	-
2006 1/.....	1,421	1	2	2	1	3	22	39	88	116	370	777	-
2005 1/.....	1,753	1	2	3	1	7	17	46	57	154	481	984	-
2004 1/.....	1,645	2	1	1	-	5	18	49	61	124	465	919	-
2003 1/.....	1,857	1	-	-	2	2	13	34	73	145	517	1,070	-
2002.....	2,111	1	-	-	3	7	19	47	62	153	607	1,212	-
2001.....	2,152	-	3	-	1	7	17	45	62	170	563	1,284	-
2000.....	2,452	2	-	-	6	5	29	40	65	198	708	1,399	-
1999.....	2,429	3	-	-	2	6	29	37	54	167	643	1,488	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Nutritional deficiencies (E40-E64)

Black, both sexes

2007 1/.....	393	3	2	-	-	7	9	17	32	70	107	146	-
2006 1/.....	341	2	-	-	4	4	11	27	35	53	87	118	-
2005 1/.....	430	3	1	1	1	7	9	28	34	61	126	159	-
2004 1/.....	416	1	1	1	2	3	12	32	34	58	105	166	1
2003 1/.....	455	3	-	-	1	2	4	33	39	54	130	189	-
2002.....	502	-	-	1	1	3	16	20	46	71	140	204	-
2001.....	475	1	-	1	2	4	15	33	30	61	150	178	-
2000.....	540	5	1	3	2	8	16	24	41	78	148	214	-
1999.....	589	4	1	1	3	3	15	27	47	90	176	222	-

Black, male

2007 1/.....	164	3	1	-	-	4	3	10	21	36	47	39	-
2006 1/.....	153	2	-	-	2	3	7	13	20	30	44	32	-
2005 1/.....	156	2	1	1	1	2	3	14	23	24	45	40	-
2004 1/.....	173	1	1	1	2	1	7	16	21	32	44	46	1
2003 1/.....	185	1	-	-	-	1	2	17	24	31	63	46	-
2002.....	188	-	-	-	1	1	7	14	33	31	57	44	-
2001.....	204	1	-	1	-	1	9	25	22	38	56	51	-
2000.....	215	3	1	-	-	5	5	13	30	44	59	55	-
1999.....	250	2	1	1	2	1	8	12	30	55	74	64	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Nutritional deficiencies (E40-E64)

Black, female

2007 1/.....	229	-	1	-	-	3	6	7	11	34	60	107	-
2006 1/.....	188	-	-	-	2	1	4	14	15	23	43	86	-
2005 1/.....	274	1	-	-	-	5	6	14	11	37	81	119	-
2004 1/.....	243	-	-	-	-	2	5	16	13	26	61	120	-
2003 1/.....	270	2	-	-	1	1	2	16	15	23	67	143	-
2002.....	314	-	-	1	-	2	9	6	13	40	83	160	-
2001.....	271	-	-	-	2	3	6	8	8	23	94	127	-
2000.....	325	2	-	3	2	3	11	11	11	34	89	159	-
1999.....	339	2	-	-	1	2	7	15	17	35	102	158	-

American Indian 2/, both sexes

2007 1/.....	26	-	-	-	-	-	-	2	5	4	6	9	-
2006 1/.....	14	1	-	-	-	-	-	-	1	1	3	8	-
2005 1/.....	20	-	-	-	-	-	-	2	2	6	4	6	-
2004 1/.....	25	-	-	-	-	-	2	1	2	3	10	7	-
2003 1/.....	21	-	-	-	-	-	1	2	-	4	7	7	-
2002.....	25	-	-	-	-	-	1	1	-	4	6	13	-
2001.....	28	-	-	-	-	-	-	2	3	6	9	8	-
2000.....	31	1	-	-	-	-	-	1	-	6	8	15	-
1999.....	28	-	-	-	-	-	-	3	4	6	7	8	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Nutritional deficiencies (E40-E64)

American Indian 2/, male

2007 1/.....	10	-	-	-	-	-	-	1	1	3	2	3	-
2006 1/.....	4	-	-	-	-	-	-	-	1	-	1	2	-
2005 1/.....	11	-	-	-	-	-	-	-	1	4	3	3	-
2004 1/.....	10	-	-	-	-	-	1	-	2	1	5	1	-
2003 1/.....	4	-	-	-	-	-	-	-	-	2	2	-	-
2002.....	8	-	-	-	-	-	-	-	-	3	1	4	-
2001.....	8	-	-	-	-	-	-	1	2	2	3	-	-
2000.....	8	1	-	-	-	-	-	-	-	2	3	2	-
1999.....	12	-	-	-	-	-	-	1	3	2	5	1	-

American Indian 2/, female

2007 1/.....	16	-	-	-	-	-	-	1	4	1	4	6	-
2006 1/.....	10	1	-	-	-	-	-	-	-	1	2	6	-
2005 1/.....	9	-	-	-	-	-	-	2	1	2	1	3	-
2004 1/.....	15	-	-	-	-	-	1	1	-	2	5	6	-
2003 1/.....	17	-	-	-	-	-	1	2	-	2	5	7	-
2002.....	17	-	-	-	-	-	1	1	-	1	5	9	-
2001.....	20	-	-	-	-	-	-	1	1	4	6	8	-
2000.....	23	-	-	-	-	-	-	1	-	4	5	13	-
1999.....	16	-	-	-	-	-	-	2	1	4	2	7	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Nutritional deficiencies (E40-E64)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	58	-	-	-	-	1	1	-	1	7	14	34	-
2006 1/.....	40	-	-	-	-	-	-	-	1	2	16	21	-
2005 1/.....	56	-	-	-	1	1	1	-	-	5	22	26	-
2004 1/.....	49	1	-	-	1	-	-	2	1	3	17	24	-
2003 1/.....	36	1	-	-	1	-	1	-	2	5	7	19	-
2002.....	39	1	-	-	-	-	2	-	1	4	10	21	-
2001.....	49	-	-	-	-	-	-	2	3	4	19	21	-
2000.....	44	-	-	-	-	1	-	1	2	6	17	17	-
1999.....	44	-	-	-	-	-	-	-	1	8	15	20	-

Asian or Pacific Islander 3/, male

2007 1/.....	22	-	-	-	-	-	-	-	1	5	4	12	-
2006 1/.....	16	-	-	-	-	-	-	-	1	-	10	5	-
2005 1/.....	25	-	-	-	-	1	-	-	-	2	11	11	-
2004 1/.....	20	1	-	-	-	-	-	1	-	2	8	8	-
2003 1/.....	18	1	-	-	1	-	1	-	1	1	4	9	-
2002.....	23	1	-	-	-	-	2	-	1	2	7	10	-
2001.....	21	-	-	-	-	-	-	2	2	1	8	8	-
2000.....	14	-	-	-	-	-	-	-	2	3	5	4	-
1999.....	25	-	-	-	-	-	-	-	1	6	8	10	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Nutritional deficiencies (E40-E64)

Asian or Pacific Islander 3/, female

2007 1/.....	36	-	-	-	-	1	1	-	-	2	10	22	-
2006 1/.....	24	-	-	-	-	-	-	-	-	2	6	16	-
2005 1/.....	31	-	-	-	1	-	1	-	-	3	11	15	-
2004 1/.....	29	-	-	-	1	-	-	1	1	1	9	16	-
2003 1/.....	18	-	-	-	-	-	-	-	1	4	3	10	-
2002.....	16	-	-	-	-	-	-	-	-	2	3	11	-
2001.....	28	-	-	-	-	-	-	-	1	3	11	13	-
2000.....	30	-	-	-	-	1	-	1	-	3	12	13	-
1999.....	19	-	-	-	-	-	-	-	-	2	7	10	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malnutrition (E40-E46)													
All races, both sexes													
2007.....	2,644	6	2	3	4	18	35	87	188	314	755	1,232	-
2006.....	2,377	7	2	2	5	11	44	93	173	262	673	1,105	-
2005.....	3,003	5	2	3	3	18	36	113	149	333	866	1,475	-
2004.....	2,847	6	1	2	5	11	35	109	152	288	864	1,373	1
2003.....	3,153	5	1	1	6	8	33	94	160	320	952	1,573	-
2002.....	3,510	2	1	1	6	13	55	91	156	354	1,067	1,764	-
2001.....	3,454	2	-	2	9	13	39	99	146	347	1,025	1,772	-
2000.....	3,926	11	1	3	8	16	53	90	158	423	1,199	1,964	-
1999.....	3,985	7	2	1	7	11	61	99	154	407	1,159	2,077	-
All races, male													
2007.....	991	4	2	2	1	11	12	43	89	154	313	360	-
2006.....	848	5	-	1	3	7	18	43	76	123	276	296	-
2005.....	1,059	3	-	1	1	6	14	53	83	149	327	422	-
2004.....	1,026	5	1	2	5	6	13	48	81	141	342	381	1
2003.....	1,098	4	1	1	3	6	19	47	75	155	379	408	-
2002.....	1,233	1	1	-	4	5	28	42	84	174	406	488	-
2001.....	1,148	2	-	2	6	4	19	47	75	157	400	436	-
2000.....	1,302	8	1	1	2	8	15	41	86	207	434	499	-
1999.....	1,371	4	2	1	4	4	27	48	92	207	452	530	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malnutrition (E40-E46)													
All races, female													
2007.....	1,653	2	-	1	3	7	23	44	99	160	442	872	-
2006.....	1,529	2	2	1	2	4	26	50	97	139	397	809	-
2005.....	1,944	2	2	2	2	12	22	60	66	184	539	1,053	-
2004.....	1,821	1	-	-	-	5	22	61	71	147	522	992	-
2003.....	2,055	1	-	-	3	2	14	47	85	165	573	1,165	-
2002.....	2,277	1	-	1	2	8	27	49	72	180	661	1,276	-
2001.....	2,306	-	-	-	3	9	20	52	71	190	625	1,336	-
2000.....	2,624	3	-	2	6	8	38	49	72	216	765	1,465	-
1999.....	2,614	3	-	-	3	7	34	51	62	200	707	1,547	-
White, both sexes													
2007 1/.....	2,194	3	1	3	4	11	25	70	153	234	633	1,057	-
2006 1/.....	2,000	5	2	2	1	7	33	66	140	206	574	964	-
2005 1/.....	2,518	2	2	2	1	11	27	84	114	262	719	1,294	-
2004 1/.....	2,373	4	-	1	3	8	23	76	118	226	734	1,180	-
2003 1/.....	2,657	2	1	1	5	6	28	60	120	259	810	1,365	-
2002.....	2,979	2	1	-	5	10	38	71	112	285	917	1,538	-
2001.....	2,939	1	-	1	7	9	25	67	113	282	859	1,575	-
2000.....	3,352	7	-	1	7	8	37	66	119	335	1,039	1,733	-
1999.....	3,368	4	1	-	5	8	47	73	107	305	976	1,842	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malnutrition (E40-E46)													
White, male													
2007 1/.....	802	1	1	2	1	8	9	32	68	110	261	309	-
2006 1/.....	682	4	-	1	1	4	11	30	56	93	225	257	-
2005 1/.....	875	1	-	-	-	4	11	39	60	119	270	371	-
2004 1/.....	829	3	-	1	3	5	6	31	59	107	286	328	-
2003 1/.....	899	2	1	1	3	5	16	30	51	122	312	356	-
2002.....	1,029	1	1	-	3	4	20	29	53	142	345	431	-
2001.....	932	1	-	1	6	3	10	24	52	121	335	379	-
2000.....	1,079	5	-	1	2	4	10	29	58	159	371	440	-
1999.....	1,100	2	1	-	3	3	19	36	61	146	373	456	-
White, female													
2007 1/.....	1,392	2	-	1	3	3	16	38	85	124	372	748	-
2006 1/.....	1,318	1	2	1	-	3	22	36	84	113	349	707	-
2005 1/.....	1,643	1	2	2	1	7	16	45	54	143	449	923	-
2004 1/.....	1,544	1	-	-	-	3	17	45	59	119	448	852	-
2003 1/.....	1,758	-	-	-	2	1	12	30	69	137	498	1,009	-
2002.....	1,950	1	-	-	2	6	18	42	59	143	572	1,107	-
2001.....	2,007	-	-	-	1	6	15	43	61	161	524	1,196	-
2000.....	2,273	2	-	-	5	4	27	37	61	176	668	1,293	-
1999.....	2,268	2	-	-	2	5	28	37	46	159	603	1,386	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malnutrition (E40-E46)													
Black, both sexes													
2007 1/.....	374	3	1	-	-	6	9	15	30	69	102	139	-
2006 1/.....	325	1	-	-	4	4	11	27	31	53	81	113	-
2005 1/.....	412	3	-	1	1	7	8	27	33	60	121	151	-
2004 1/.....	402	1	1	1	2	3	10	30	31	56	103	163	1
2003 1/.....	442	2	-	-	1	2	4	32	38	52	128	183	-
2002.....	472	-	-	1	1	3	14	19	44	61	135	194	-
2001.....	441	1	-	1	2	4	14	28	27	56	140	168	-
2000.....	508	3	1	2	1	7	16	23	37	77	137	204	-
1999.....	547	3	1	1	2	3	14	24	42	88	162	207	-
Black, male													
2007 1/.....	160	3	1	-	-	3	3	10	19	36	46	39	-
2006 1/.....	146	1	-	-	2	3	7	13	18	30	40	32	-
2005 1/.....	149	2	-	1	1	2	3	14	22	24	43	37	-
2004 1/.....	167	1	1	1	2	1	6	16	20	31	43	44	1
2003 1/.....	179	1	-	-	-	1	2	17	23	30	61	44	-
2002.....	175	-	-	-	1	1	6	13	31	27	53	43	-
2001.....	188	1	-	1	-	1	9	20	19	34	54	49	-
2000.....	203	2	1	-	-	4	5	12	26	43	57	53	-
1999.....	234	2	1	1	1	1	8	11	27	53	66	63	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malnutrition (E40-E46)

Black, female

2007 1/.....	214	-	-	-	-	3	6	5	11	33	56	100	-
2006 1/.....	179	-	-	-	2	1	4	14	13	23	41	81	-
2005 1/.....	263	1	-	-	-	5	5	13	11	36	78	114	-
2004 1/.....	235	-	-	-	-	2	4	14	11	25	60	119	-
2003 1/.....	263	1	-	-	1	1	2	15	15	22	67	139	-
2002.....	297	-	-	1	-	2	8	6	13	34	82	151	-
2001.....	253	-	-	-	2	3	5	8	8	22	86	119	-
2000.....	305	1	-	2	1	3	11	11	11	34	80	151	-
1999.....	313	1	-	-	1	2	6	13	15	35	96	144	-

American Indian 2/, both sexes

2007 1/.....	24	-	-	-	-	-	-	2	4	4	6	8	-
2006 1/.....	14	1	-	-	-	-	-	-	1	1	3	8	-
2005 1/.....	20	-	-	-	-	-	-	2	2	6	4	6	-
2004 1/.....	25	-	-	-	-	-	2	1	2	3	10	7	-
2003 1/.....	20	-	-	-	-	-	-	2	-	4	7	7	-
2002.....	23	-	-	-	-	-	1	1	-	4	5	12	-
2001.....	28	-	-	-	-	-	-	2	3	6	9	8	-
2000.....	25	1	-	-	-	-	-	-	-	5	7	12	-
1999.....	27	-	-	-	-	-	-	2	4	6	7	8	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malnutrition (E40-E46)													
American Indian 2/, male													
2007 1/.....	9	-	-	-	-	-	-	1	1	3	2	2	-
2006 1/.....	4	-	-	-	-	-	-	-	1	-	1	2	-
2005 1/.....	11	-	-	-	-	-	-	-	1	4	3	3	-
2004 1/.....	10	-	-	-	-	-	1	-	2	1	5	1	-
2003 1/.....	4	-	-	-	-	-	-	-	-	2	2	-	-
2002.....	8	-	-	-	-	-	-	-	-	3	1	4	-
2001.....	8	-	-	-	-	-	-	1	2	2	3	-	-
2000.....	7	1	-	-	-	-	-	-	-	2	2	2	-
1999.....	12	-	-	-	-	-	-	1	3	2	5	1	-
American Indian 2/, female													
2007 1/.....	15	-	-	-	-	-	-	1	3	1	4	6	-
2006 1/.....	10	1	-	-	-	-	-	-	-	1	2	6	-
2005 1/.....	9	-	-	-	-	-	-	2	1	2	1	3	-
2004 1/.....	15	-	-	-	-	-	1	1	-	2	5	6	-
2003 1/.....	16	-	-	-	-	-	-	2	-	2	5	7	-
2002.....	15	-	-	-	-	-	1	1	-	1	4	8	-
2001.....	20	-	-	-	-	-	-	1	1	4	6	8	-
2000.....	18	-	-	-	-	-	-	-	-	3	5	10	-
1999.....	15	-	-	-	-	-	-	1	1	4	2	7	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Malnutrition (E40-E46)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	52	-	-	-	-	1	1	-	1	7	14	28	-
2006 1/.....	38	-	-	-	-	-	-	-	1	2	15	20	-
2005 1/.....	53	-	-	-	1	-	1	-	-	5	22	24	-
2004 1/.....	47	1	-	-	-	-	-	2	1	3	17	23	-
2003 1/.....	34	1	-	-	-	-	1	-	2	5	7	18	-
2002.....	36	-	-	-	-	-	2	-	-	4	10	20	-
2001.....	46	-	-	-	-	-	-	2	3	3	17	21	-
2000.....	41	-	-	-	-	1	-	1	2	6	16	15	-
1999.....	43	-	-	-	-	-	-	-	1	8	14	20	-

Asian or Pacific Islander 3/, male

2007 1/.....	20	-	-	-	-	-	-	-	1	5	4	10	-
2006 1/.....	16	-	-	-	-	-	-	-	1	-	10	5	-
2005 1/.....	24	-	-	-	-	-	-	-	-	2	11	11	-
2004 1/.....	20	1	-	-	-	-	-	1	-	2	8	8	-
2003 1/.....	16	1	-	-	-	-	1	-	1	1	4	8	-
2002.....	21	-	-	-	-	-	2	-	-	2	7	10	-
2001.....	20	-	-	-	-	-	-	2	2	-	8	8	-
2000.....	13	-	-	-	-	-	-	-	2	3	4	4	-
1999.....	25	-	-	-	-	-	-	-	1	6	8	10	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malnutrition (E40-E46)

Asian or Pacific Islander 3/, female

2007 1/.....	32	-	-	-	-	1	1	-	-	2	10	18	-
2006 1/.....	22	-	-	-	-	-	-	-	-	2	5	15	-
2005 1/.....	29	-	-	-	1	-	1	-	-	3	11	13	-
2004 1/.....	27	-	-	-	-	-	-	1	1	1	9	15	-
2003 1/.....	18	-	-	-	-	-	-	-	1	4	3	10	-
2002.....	15	-	-	-	-	-	-	-	-	2	3	10	-
2001.....	26	-	-	-	-	-	-	-	1	3	9	13	-
2000.....	28	-	-	-	-	1	-	1	-	3	12	11	-
1999.....	18	-	-	-	-	-	-	-	-	2	6	10	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other nutritional deficiencies (E50-E64)

All races, both sexes

2007.....	208	1	3	-	-	1	1	6	13	19	53	111	-
2006.....	179	2	-	2	1	-	2	5	12	13	47	95	-
2005.....	180	-	1	1	-	1	2	5	8	20	49	93	-
2004.....	172	1	1	1	1	2	4	8	10	10	33	101	-
2003.....	185	5	1	1	1	2	2	6	8	20	40	99	-
2002.....	269	2	-	-	1	3	5	11	12	28	65	142	-
2001.....	250	2	4	-	1	1	5	11	7	27	75	117	-
2000.....	316	3	1	2	2	2	3	9	15	37	86	156	-
1999.....	304	2	-	-	1	1	3	7	16	22	89	163	-

All races, male

2007.....	80	1	1	-	-	1	1	2	8	14	20	32	-
2006.....	65	2	-	1	-	-	2	2	6	10	23	19	-
2005.....	57	-	1	-	-	1	-	3	5	8	14	25	-
2004.....	61	-	-	-	-	-	2	2	6	4	15	32	-
2003.....	78	3	1	1	1	1	-	1	4	11	21	34	-
2002.....	88	2	-	-	-	2	3	6	9	12	28	26	-
2001.....	85	2	1	-	1	-	2	9	6	17	26	21	-
2000.....	110	2	1	1	-	1	1	5	11	14	37	37	-
1999.....	115	-	-	-	1	-	1	4	6	14	42	47	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other nutritional deficiencies (E50-E64)													
All races, female													
2007.....	128	-	2	-	-	-	-	4	5	5	33	79	-
2006.....	114	-	-	1	1	-	-	3	6	3	24	76	-
2005.....	123	-	-	1	-	-	2	2	3	12	35	68	-
2004.....	111	1	1	1	1	2	2	6	4	6	18	69	-
2003.....	107	2	-	-	-	1	2	5	4	9	19	65	-
2002.....	181	-	-	-	1	1	2	5	3	16	37	116	-
2001.....	165	-	3	-	-	1	3	2	1	10	49	96	-
2000.....	206	1	-	1	2	1	2	4	4	23	49	119	-
1999.....	189	2	-	-	-	1	2	3	10	8	47	116	-
White, both sexes													
2007 1/.....	181	1	2	-	-	-	1	4	10	18	48	97	-
2006 1/.....	161	1	-	2	1	-	2	5	8	13	40	89	-
2005 1/.....	159	-	-	1	-	-	1	4	7	19	44	83	-
2004 1/.....	156	1	1	1	-	2	2	6	7	8	31	97	-
2003 1/.....	169	4	1	1	-	2	1	5	7	18	38	92	-
2002.....	234	1	-	-	1	3	3	10	9	18	59	130	-
2001.....	213	2	4	-	1	1	4	6	4	21	63	107	-
2000.....	275	1	1	1	1	1	3	7	11	35	73	141	-
1999.....	260	1	-	-	-	1	2	3	11	20	74	148	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other nutritional deficiencies (E50-E64)													
White, male													
2007 1/.....	73	1	1	-	-	-	1	2	6	14	19	29	-
2006 1/.....	58	1	-	1	-	-	2	2	4	10	19	19	-
2005 1/.....	49	-	-	-	-	-	-	3	4	8	12	22	-
2004 1/.....	55	-	-	-	-	-	1	2	5	3	14	30	-
2003 1/.....	70	3	1	1	-	1	-	1	3	10	19	31	-
2002.....	73	1	-	-	-	2	2	5	6	8	24	25	-
2001.....	68	2	1	-	1	-	2	4	3	12	24	19	-
2000.....	96	1	1	1	-	-	1	4	7	13	33	35	-
1999.....	99	-	-	-	-	-	1	3	3	12	34	46	-
White, female													
2007 1/.....	108	-	1	-	-	-	-	2	4	4	29	68	-
2006 1/.....	103	-	-	1	1	-	-	3	4	3	21	70	-
2005 1/.....	110	-	-	1	-	-	1	1	3	11	32	61	-
2004 1/.....	101	1	1	1	-	2	1	4	2	5	17	67	-
2003 1/.....	99	1	-	-	-	1	1	4	4	8	19	61	-
2002.....	161	-	-	-	1	1	1	5	3	10	35	105	-
2001.....	145	-	3	-	-	1	2	2	1	9	39	88	-
2000.....	179	-	-	-	1	1	2	3	4	22	40	106	-
1999.....	161	1	-	-	-	1	1	-	8	8	40	102	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other nutritional deficiencies (E50-E64)

Black, both sexes

2007 1/.....	19	-	1	-	-	1	-	2	2	1	5	7	-
2006 1/.....	16	1	-	-	-	-	-	-	4	-	6	5	-
2005 1/.....	18	-	1	-	-	-	1	1	1	1	5	8	-
2004 1/.....	14	-	-	-	-	-	2	2	3	2	2	3	-
2003 1/.....	13	1	-	-	-	-	-	1	1	2	2	6	-
2002.....	30	-	-	-	-	-	2	1	2	10	5	10	-
2001.....	34	-	-	-	-	-	1	5	3	5	10	10	-
2000.....	32	2	-	1	1	1	-	1	4	1	11	10	-
1999.....	42	1	-	-	1	-	1	3	5	2	14	15	-

Black, male

2007 1/.....	4	-	-	-	-	1	-	-	2	-	1	-	-
2006 1/.....	7	1	-	-	-	-	-	-	2	-	4	-	-
2005 1/.....	7	-	1	-	-	-	-	-	1	-	2	3	-
2004 1/.....	6	-	-	-	-	-	1	-	1	1	1	2	-
2003 1/.....	6	-	-	-	-	-	-	-	1	1	2	2	-
2002.....	13	-	-	-	-	-	1	1	2	4	4	1	-
2001.....	16	-	-	-	-	-	-	5	3	4	2	2	-
2000.....	12	1	-	-	-	1	-	1	4	1	2	2	-
1999.....	16	-	-	-	1	-	-	1	3	2	8	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other nutritional deficiencies (E50-E64)

Black, female

2007 1/.....	15	-	1	-	-	-	-	2	-	1	4	7	-
2006 1/.....	9	-	-	-	-	-	-	-	2	-	2	5	-
2005 1/.....	11	-	-	-	-	-	1	1	-	1	3	5	-
2004 1/.....	8	-	-	-	-	-	1	2	2	1	1	1	-
2003 1/.....	7	1	-	-	-	-	-	1	-	1	-	4	-
2002.....	17	-	-	-	-	-	1	-	-	6	1	9	-
2001.....	18	-	-	-	-	-	1	-	-	1	8	8	-
2000.....	20	1	-	1	1	-	-	-	-	-	9	8	-
1999.....	26	1	-	-	-	-	1	2	2	-	6	14	-

American Indian 2/, both sexes

2007 1/.....	2	-	-	-	-	-	-	-	1	-	-	1	-
2006 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2005 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 1/.....	1	-	-	-	-	-	1	-	-	-	-	-	-
2002.....	2	-	-	-	-	-	-	-	-	-	1	1	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	6	-	-	-	-	-	-	1	-	1	1	3	-
1999.....	1	-	-	-	-	-	-	1	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other nutritional deficiencies (E50-E64)

American Indian 2/, male

2007 1/.....	1	-	-	-	-	-	-	-	-	-	-	-	1	-
2006 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2005 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	1	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-	-	-

American Indian 2/, female

2007 1/.....	1	-	-	-	-	-	-	-	1	-	-	-	-	-
2006 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2005 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 1/.....	1	-	-	-	-	-	1	-	-	-	-	-	-	-
2002.....	2	-	-	-	-	-	-	-	-	-	1	1	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	5	-	-	-	-	-	-	1	-	1	-	-	3	-
1999.....	1	-	-	-	-	-	-	1	-	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other nutritional deficiencies (E50-E64)

Asian or Pacific Islander 3/, female

2007 1/.....	4	-	-	-	-	-	-	-	-	-	-	4	-
2006 1/.....	2	-	-	-	-	-	-	-	-	-	1	1	-
2005 1/.....	2	-	-	-	-	-	-	-	-	-	-	2	-
2004 1/.....	2	-	-	-	1	-	-	-	-	-	-	1	-
2003 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	1	-
2001.....	2	-	-	-	-	-	-	-	-	-	2	-	-
2000.....	2	-	-	-	-	-	-	-	-	-	-	2	-
1999.....	1	-	-	-	-	-	-	-	-	-	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Meningitis (G00,G03)													
All races, both sexes													
2007.....	655	82	16	25	35	34	66	111	89	72	88	37	-
2006.....	634	61	20	15	25	35	80	108	77	76	89	48	-
2005.....	669	57	32	26	28	32	74	118	106	74	76	45	1
2004.....	674	66	25	25	26	37	67	117	92	92	85	42	-
2003.....	730	77	24	17	37	44	67	117	108	96	90	52	1
2002.....	700	74	19	31	26	36	61	116	88	109	84	55	1
2001.....	727	78	26	24	40	41	69	105	113	94	96	41	-
2000.....	758	91	36	30	15	43	72	100	90	108	123	50	-
1999.....	850	116	45	27	41	48	76	108	90	115	122	62	-
All races, male													
2007.....	346	35	10	12	18	17	42	68	51	41	40	12	-
2006.....	315	29	6	8	15	20	43	63	40	33	39	19	-
2005.....	353	29	21	15	15	24	44	68	53	37	32	14	1
2004.....	349	32	15	13	10	22	46	73	48	33	38	19	-
2003.....	387	38	12	10	20	28	40	73	61	45	40	19	1
2002.....	389	41	13	16	15	19	43	71	48	62	41	19	1
2001.....	394	39	16	15	23	25	42	60	61	51	47	15	-
2000.....	405	48	20	16	4	20	46	56	54	60	62	19	-
1999.....	442	61	26	18	26	24	45	68	45	60	49	20	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Meningitis (G00,G03)

All races, female

2007.....	309	47	6	13	17	17	24	43	38	31	48	25	-
2006.....	319	32	14	7	10	15	37	45	37	43	50	29	-
2005.....	316	28	11	11	13	8	30	50	53	37	44	31	-
2004.....	325	34	10	12	16	15	21	44	44	59	47	23	-
2003.....	343	39	12	7	17	16	27	44	47	51	50	33	-
2002.....	311	33	6	15	11	17	18	45	40	47	43	36	-
2001.....	333	39	10	9	17	16	27	45	52	43	49	26	-
2000.....	353	43	16	14	11	23	26	44	36	48	61	31	-
1999.....	408	55	19	9	15	24	31	40	45	55	73	42	-

White, both sexes

2007 1/.....	499	51	12	19	23	30	52	77	63	57	80	35	-
2006 1/.....	487	44	15	7	17	28	48	85	56	65	77	45	-
2005 1/.....	496	35	20	18	21	20	49	88	82	60	63	39	1
2004 1/.....	489	46	19	19	19	19	49	76	67	69	69	37	-
2003 1/.....	524	52	18	13	25	26	43	78	72	76	78	42	1
2002.....	498	52	13	20	18	20	39	71	58	88	68	50	1
2001.....	526	47	19	18	26	29	40	72	84	78	82	31	-
2000.....	536	54	23	21	10	25	44	71	65	79	104	40	-
1999.....	601	68	33	19	21	34	49	72	71	89	95	50	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Meningitis (G00,G03)													
White, male													
2007 1/.....	257	19	7	7	12	14	30	49	38	33	36	12	-
2006 1/.....	241	20	4	5	13	16	24	49	26	28	38	18	-
2005 1/.....	262	17	13	9	13	14	30	54	41	30	28	12	1
2004 1/.....	259	25	12	11	7	13	34	51	31	25	32	18	-
2003 1/.....	293	32	9	9	16	17	26	50	43	37	37	16	1
2002.....	270	29	8	11	10	9	26	42	32	51	36	15	1
2001.....	275	22	13	10	13	18	24	41	45	41	39	9	-
2000.....	294	31	14	10	3	10	27	43	42	48	52	14	-
1999.....	308	46	20	12	12	13	27	41	37	46	38	16	-
White, female													
2007 1/.....	242	32	5	12	11	16	22	28	25	24	44	23	-
2006 1/.....	246	24	11	2	4	12	24	36	30	37	39	27	-
2005 1/.....	234	18	7	9	8	6	19	34	41	30	35	27	-
2004 1/.....	230	21	7	8	12	6	15	25	36	44	37	19	-
2003 1/.....	231	20	9	4	9	9	17	28	29	39	41	26	-
2002.....	228	23	5	9	8	11	13	29	26	37	32	35	-
2001.....	251	25	6	8	13	11	16	31	39	37	43	22	-
2000.....	242	23	9	11	7	15	17	28	23	31	52	26	-
1999.....	293	22	13	7	9	21	22	31	34	43	57	34	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Meningitis (G00,G03)													
Black, both sexes													
2007 1/.....	132	23	3	5	9	4	13	29	23	14	7	2	-
2006 1/.....	131	16	4	8	8	7	26	20	21	8	10	3	-
2005 1/.....	143	18	6	7	5	11	22	28	20	13	8	5	-
2004 1/.....	159	17	5	6	5	15	18	37	21	19	13	3	-
2003 1/.....	187	23	5	3	12	18	22	33	35	18	9	9	-
2002.....	178	20	6	10	6	15	20	40	23	21	13	4	-
2001.....	175	29	5	4	11	11	26	31	25	14	11	8	-
2000.....	204	35	13	9	5	18	27	26	22	27	13	9	-
1999.....	222	42	11	8	17	11	22	35	18	21	25	12	-
Black, male													
2007 1/.....	77	12	3	4	5	3	11	17	12	7	3	-	-
2006 1/.....	66	9	1	3	2	4	16	12	14	3	1	1	-
2005 1/.....	76	9	3	6	1	9	13	13	10	7	3	2	-
2004 1/.....	76	7	2	2	2	7	12	20	13	6	5	-	-
2003 1/.....	84	6	2	1	4	11	12	18	18	8	2	2	-
2002.....	107	11	5	5	5	9	15	26	12	11	5	3	-
2001.....	105	16	2	3	8	6	17	18	14	9	6	6	-
2000.....	102	16	6	6	1	10	18	11	11	11	7	5	-
1999.....	121	13	6	6	11	9	15	27	8	12	10	4	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Meningitis (G00,G03)

Black, female

2007 1/.....	55	11	-	1	4	1	2	12	11	7	4	2	-
2006 1/.....	65	7	3	5	6	3	10	8	7	5	9	2	-
2005 1/.....	67	9	3	1	4	2	9	15	10	6	5	3	-
2004 1/.....	83	10	3	4	3	8	6	17	8	13	8	3	-
2003 1/.....	103	17	3	2	8	7	10	15	17	10	7	7	-
2002.....	71	9	1	5	1	6	5	14	11	10	8	1	-
2001.....	70	13	3	1	3	5	9	13	11	5	5	2	-
2000.....	102	19	7	3	4	8	9	15	11	16	6	4	-
1999.....	101	29	5	2	6	2	7	8	10	9	15	8	-

American Indian 2/, both sexes

2007 1/.....	6	4	-	-	-	-	1	-	1	-	-	-	-
2006 1/.....	8	1	-	-	-	-	4	3	-	-	-	-	-
2005 1/.....	11	2	3	1	-	-	2	-	1	-	2	-	-
2004 1/.....	9	1	-	-	-	1	-	3	2	-	2	-	-
2003 1/.....	8	1	1	-	-	-	1	3	1	1	-	-	-
2002.....	8	1	-	1	1	-	1	1	2	-	1	-	-
2001.....	10	1	2	-	1	-	-	2	1	2	1	-	-
2000.....	5	1	-	-	-	-	1	-	-	-	3	-	-
1999.....	10	2	-	-	2	2	1	1	-	1	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Meningitis (G00,G03)

American Indian 2/, male

2007 1/.....	4	3	-	-	-	-	1	-	-	-	-	-	-
2006 1/.....	4	-	-	-	-	-	2	2	-	-	-	-	-
2005 1/.....	6	2	2	-	-	-	1	1	1	-	-	-	-
2004 1/.....	6	-	-	-	-	1	-	2	2	-	1	-	-
2003 1/.....	5	-	1	-	-	-	1	3	-	-	-	-	-
2002.....	5	1	-	-	-	-	1	1	2	-	-	-	-
2001.....	5	-	1	-	1	-	-	1	1	1	-	-	-
2000.....	3	1	-	-	-	-	1	-	-	-	1	-	-
1999.....	7	1	-	-	2	2	-	-	-	1	1	-	-

American Indian 2/, female

2007 1/.....	2	1	-	-	-	-	-	-	1	-	-	-	-
2006 1/.....	4	1	-	-	-	-	2	1	-	-	-	-	-
2005 1/.....	5	-	1	1	-	-	1	-	-	-	2	-	-
2004 1/.....	3	1	-	-	-	-	-	1	-	-	1	-	-
2003 1/.....	3	1	-	-	-	-	-	-	1	1	-	-	-
2002.....	3	-	-	1	1	-	-	-	-	-	1	-	-
2001.....	5	1	1	-	-	-	-	1	-	1	1	-	-
2000.....	2	-	-	-	-	-	-	-	-	-	2	-	-
1999.....	3	1	-	-	-	-	1	1	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Meningitis (G00,G03)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	18	4	1	1	3	-	-	5	2	1	1	-	-
2006 1/.....	8	-	1	-	-	-	2	-	-	3	2	-	-
2005 1/.....	19	2	3	-	2	1	1	2	3	1	3	1	-
2004 1/.....	17	2	1	-	2	2	-	1	2	4	1	2	-
2003 1/.....	11	1	-	1	-	-	1	3	-	1	3	1	-
2002.....	16	1	-	-	1	1	1	4	5	-	2	1	-
2001.....	16	1	-	2	2	1	3	-	3	-	2	2	-
2000.....	13	1	-	-	-	-	-	3	3	2	3	1	-
1999.....	17	4	1	-	1	1	4	-	1	4	1	-	-

Asian or Pacific Islander 3/, male

2007 1/.....	8	1	-	1	1	-	-	2	1	1	1	-	-
2006 1/.....	4	-	1	-	-	-	1	-	-	2	-	-	-
2005 1/.....	9	1	3	-	1	1	-	1	1	-	1	-	-
2004 1/.....	8	-	1	-	1	1	-	-	2	2	-	1	-
2003 1/.....	5	-	-	-	-	-	1	2	-	-	1	1	-
2002.....	7	-	-	-	-	1	1	2	2	-	-	1	-
2001.....	9	1	-	2	1	1	1	-	1	-	2	-	-
2000.....	6	-	-	-	-	-	-	2	1	1	2	-	-
1999.....	6	1	-	-	1	-	3	-	-	1	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Meningitis (G00,G03)

Asian or Pacific Islander 3/, female

2007 1/.....	10	3	1	-	2	-	-	3	1	-	-	-	-
2006 1/.....	4	-	-	-	-	-	1	-	-	1	2	-	-
2005 1/.....	10	1	-	-	1	-	1	1	2	1	2	1	-
2004 1/.....	9	2	-	-	1	1	-	1	-	2	1	1	-
2003 1/.....	6	1	-	1	-	-	-	1	-	1	2	-	-
2002.....	9	1	-	-	1	-	-	2	3	-	2	-	-
2001.....	7	-	-	-	1	-	2	-	2	-	-	2	-
2000.....	7	1	-	-	-	-	-	1	2	1	1	1	-
1999.....	11	3	1	-	-	1	1	-	1	3	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Parkinson's disease (G20-G21)													
All races, both sexes													
2007.....	20,058	-	-	-	2	2	12	60	396	2,310	9,363	7,911	2
2006.....	19,566	1	1	-	2	4	15	66	397	2,310	9,101	7,669	-
2005.....	19,544	-	-	-	3	4	7	66	434	2,414	9,294	7,322	-
2004.....	17,989	-	1	-	1	2	8	64	349	2,214	8,752	6,598	-
2003.....	17,997	1	-	1	-	2	11	64	352	2,330	8,723	6,513	-
2002.....	16,959	-	-	-	-	-	6	58	318	2,226	8,139	6,212	-
2001.....	16,544	-	-	-	-	2	15	45	301	2,147	8,125	5,909	-
2000.....	15,682	-	1	-	-	4	7	49	270	2,108	7,652	5,590	1
1999.....	14,593	1	-	-	2	2	5	43	242	2,020	7,110	5,168	-
All races, male													
2007.....	11,554	-	-	-	1	1	9	38	251	1,507	5,694	4,051	2
2006.....	11,300	1	-	-	1	2	10	45	262	1,522	5,546	3,911	-
2005.....	11,247	-	-	-	3	2	3	44	275	1,596	5,658	3,666	-
2004.....	10,366	-	-	-	1	2	6	49	227	1,457	5,306	3,318	-
2003.....	10,180	1	-	-	-	-	6	33	219	1,517	5,191	3,213	-
2002.....	9,593	-	-	-	-	-	3	40	193	1,487	4,811	3,059	-
2001.....	9,405	-	-	-	-	1	11	33	214	1,394	4,837	2,915	-
2000.....	8,804	-	-	-	-	3	5	29	170	1,338	4,530	2,728	1
1999.....	8,262	-	-	-	2	2	3	30	161	1,303	4,307	2,454	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Parkinson's disease (G20-G21)													
All races, female													
2007.....	8,504	-	-	-	1	1	3	22	145	803	3,669	3,860	-
2006.....	8,266	-	1	-	1	2	5	21	135	788	3,555	3,758	-
2005.....	8,297	-	-	-	-	2	4	22	159	818	3,636	3,656	-
2004.....	7,623	-	1	-	-	-	2	15	122	757	3,446	3,280	-
2003.....	7,817	-	-	1	-	2	5	31	133	813	3,532	3,300	-
2002.....	7,366	-	-	-	-	-	3	18	125	739	3,328	3,153	-
2001.....	7,139	-	-	-	-	1	4	12	87	753	3,288	2,994	-
2000.....	6,878	-	1	-	-	1	2	20	100	770	3,122	2,862	-
1999.....	6,331	1	-	-	-	-	2	13	81	717	2,803	2,714	-
White, both sexes													
2007 1/.....	18,864	-	-	-	2	1	11	54	351	2,115	8,832	7,497	1
2006 1/.....	18,517	1	-	-	2	3	13	51	354	2,120	8,641	7,332	-
2005 1/.....	18,496	-	-	-	3	4	7	60	395	2,224	8,797	7,006	-
2004 1/.....	17,041	-	1	-	-	2	7	55	321	2,075	8,313	6,267	-
2003 1/.....	17,045	1	-	-	-	2	10	55	307	2,158	8,295	6,217	-
2002.....	16,135	-	-	-	-	-	3	53	290	2,061	7,780	5,948	-
2001.....	15,757	-	-	-	-	1	11	40	270	2,011	7,772	5,652	-
2000.....	14,942	-	-	-	-	3	4	41	251	1,982	7,306	5,354	1
1999.....	13,886	1	-	-	2	1	5	39	215	1,895	6,787	4,941	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Parkinson's disease (G20-G21)													
White, male													
2007 1/.....	10,895	-	-	-	1	-	9	35	228	1,374	5,399	3,848	1
2006 1/.....	10,706	1	-	-	1	1	9	38	235	1,398	5,263	3,760	-
2005 1/.....	10,649	-	-	-	3	2	3	39	250	1,472	5,374	3,506	-
2004 1/.....	9,829	-	-	-	-	2	5	42	207	1,367	5,058	3,148	-
2003 1/.....	9,630	1	-	-	-	-	6	29	189	1,411	4,946	3,048	-
2002.....	9,130	-	-	-	-	-	2	35	179	1,384	4,604	2,926	-
2001.....	8,982	-	-	-	-	-	8	30	197	1,311	4,641	2,795	-
2000.....	8,367	-	-	-	-	3	3	24	158	1,262	4,307	2,609	1
1999.....	7,880	-	-	-	2	1	3	29	147	1,236	4,123	2,339	-
White, female													
2007 1/.....	7,969	-	-	-	1	1	2	19	123	741	3,433	3,649	-
2006 1/.....	7,811	-	-	-	1	2	4	13	119	722	3,378	3,572	-
2005 1/.....	7,847	-	-	-	-	2	4	21	145	752	3,423	3,500	-
2004 1/.....	7,212	-	1	-	-	-	2	13	114	708	3,255	3,119	-
2003 1/.....	7,415	-	-	-	-	2	4	26	118	747	3,349	3,169	-
2002.....	7,005	-	-	-	-	-	1	18	111	677	3,176	3,022	-
2001.....	6,775	-	-	-	-	1	3	10	73	700	3,131	2,857	-
2000.....	6,575	-	-	-	-	-	1	17	93	720	2,999	2,745	-
1999.....	6,006	1	-	-	-	-	2	10	68	659	2,664	2,602	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Parkinson's disease (G20-G21)													
Black, both sexes													
2007 1/.....	802	-	-	-	-	1	-	4	36	135	361	264	1
2006 1/.....	682	-	1	-	-	1	2	10	31	133	306	198	-
2005 1/.....	711	-	-	-	-	-	-	5	25	127	342	212	-
2004 1/.....	627	-	-	-	-	-	1	5	21	93	290	217	-
2003 1/.....	628	-	-	1	-	-	1	7	33	120	285	181	-
2002.....	557	-	-	-	-	-	1	4	19	114	245	174	-
2001.....	551	-	-	-	-	1	4	5	20	91	238	192	-
2000.....	534	-	-	-	-	1	3	6	14	94	251	165	-
1999.....	507	-	-	-	-	-	-	4	22	97	221	163	-
Black, male													
2007 1/.....	437	-	-	-	-	1	-	2	20	95	201	117	1
2006 1/.....	381	-	-	-	-	1	1	6	19	88	187	79	-
2005 1/.....	412	-	-	-	-	-	-	4	15	90	198	105	-
2004 1/.....	345	-	-	-	-	-	1	5	15	61	156	107	-
2003 1/.....	351	-	-	-	-	-	-	3	21	73	161	93	-
2002.....	313	-	-	-	-	-	-	4	10	78	138	83	-
2001.....	294	-	-	-	-	1	3	3	13	59	127	88	-
2000.....	308	-	-	-	-	-	2	4	9	60	158	75	-
1999.....	270	-	-	-	-	-	-	1	13	59	123	74	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Parkinson's disease (G20-G21)

Black, female

2007 1/.....	365	-	-	-	-	-	-	2	16	40	160	147	-
2006 1/.....	301	-	1	-	-	-	1	4	12	45	119	119	-
2005 1/.....	299	-	-	-	-	-	-	1	10	37	144	107	-
2004 1/.....	282	-	-	-	-	-	-	-	6	32	134	110	-
2003 1/.....	277	-	-	1	-	-	1	4	12	47	124	88	-
2002.....	244	-	-	-	-	-	1	-	9	36	107	91	-
2001.....	257	-	-	-	-	-	1	2	7	32	111	104	-
2000.....	226	-	-	-	-	1	1	2	5	34	93	90	-
1999.....	237	-	-	-	-	-	-	3	9	38	98	89	-

American Indian 2/, both sexes

2007 1/.....	55	-	-	-	-	-	-	1	1	12	17	24	-
2006 1/.....	57	-	-	-	-	-	-	1	1	11	24	20	-
2005 1/.....	66	-	-	-	-	-	-	1	6	10	31	18	-
2004 1/.....	46	-	-	-	1	-	-	2	1	3	25	14	-
2003 1/.....	60	-	-	-	-	-	-	-	4	13	26	17	-
2002.....	53	-	-	-	-	-	1	-	2	14	22	14	-
2001.....	36	-	-	-	-	-	-	-	3	10	14	9	-
2000.....	45	-	1	-	-	-	-	-	2	9	20	13	-
1999.....	26	-	-	-	-	-	-	-	-	4	16	6	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Parkinson's disease (G20-G21)

American Indian 2/, male

2007 1/.....	32	-	-	-	-	-	-	1	-	9	11	11	-
2006 1/.....	32	-	-	-	-	-	-	-	-	7	17	8	-
2005 1/.....	34	-	-	-	-	-	-	1	3	3	18	9	-
2004 1/.....	28	-	-	-	1	-	-	1	1	2	17	6	-
2003 1/.....	32	-	-	-	-	-	-	-	3	7	12	10	-
2002.....	38	-	-	-	-	-	1	-	2	10	16	9	-
2001.....	18	-	-	-	-	-	-	-	-	6	10	2	-
2000.....	25	-	-	-	-	-	-	-	1	4	12	8	-
1999.....	16	-	-	-	-	-	-	-	-	-	14	2	-

American Indian 2/, female

2007 1/.....	23	-	-	-	-	-	-	-	1	3	6	13	-
2006 1/.....	25	-	-	-	-	-	-	1	1	4	7	12	-
2005 1/.....	32	-	-	-	-	-	-	-	3	7	13	9	-
2004 1/.....	18	-	-	-	-	-	-	1	-	1	8	8	-
2003 1/.....	28	-	-	-	-	-	-	-	1	6	14	7	-
2002.....	15	-	-	-	-	-	-	-	-	4	6	5	-
2001.....	18	-	-	-	-	-	-	-	3	4	4	7	-
2000.....	20	-	1	-	-	-	-	-	1	5	8	5	-
1999.....	10	-	-	-	-	-	-	-	-	4	2	4	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Parkinson's disease (G20-G21)													
Asian or Pacific Islander 3/, both sexes													
2007 1/.....	337	-	-	-	-	-	1	1	8	48	153	126	-
2006 1/.....	310	-	-	-	-	-	-	4	11	46	130	119	-
2005 1/.....	271	-	-	-	-	-	-	-	8	53	124	86	-
2004 1/.....	275	-	-	-	-	-	-	2	6	43	124	100	-
2003 1/.....	264	-	-	-	-	-	-	2	8	39	117	98	-
2002.....	214	-	-	-	-	-	1	1	7	37	92	76	-
2001.....	200	-	-	-	-	-	-	-	8	35	101	56	-
2000.....	161	-	-	-	-	-	-	2	3	23	75	58	-
1999.....	174	-	-	-	-	1	-	-	5	24	86	58	-
Asian or Pacific Islander 3/, male													
2007 1/.....	190	-	-	-	-	-	-	-	3	29	83	75	-
2006 1/.....	181	-	-	-	-	-	-	1	8	29	79	64	-
2005 1/.....	152	-	-	-	-	-	-	-	7	31	68	46	-
2004 1/.....	164	-	-	-	-	-	-	1	4	27	75	57	-
2003 1/.....	167	-	-	-	-	-	-	1	6	26	72	62	-
2002.....	112	-	-	-	-	-	-	1	2	15	53	41	-
2001.....	111	-	-	-	-	-	-	-	4	18	59	30	-
2000.....	104	-	-	-	-	-	-	1	2	12	53	36	-
1999.....	96	-	-	-	-	1	-	-	1	8	47	39	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Parkinson's disease (G20-G21)

Asian or Pacific Islander 3/, female

2007 1/.....	147	-	-	-	-	-	1	1	5	19	70	51	-
2006 1/.....	129	-	-	-	-	-	-	3	3	17	51	55	-
2005 1/.....	119	-	-	-	-	-	-	-	1	22	56	40	-
2004 1/.....	111	-	-	-	-	-	-	1	2	16	49	43	-
2003 1/.....	97	-	-	-	-	-	-	1	2	13	45	36	-
2002.....	102	-	-	-	-	-	1	-	5	22	39	35	-
2001.....	89	-	-	-	-	-	-	-	4	17	42	26	-
2000.....	57	-	-	-	-	-	-	1	1	11	22	22	-
1999.....	78	-	-	-	-	-	-	-	4	16	39	19	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alzheimer's disease (G30)													
All races, both sexes													
2007.....	74,632	-	-	-	-	1	8	95	728	3,984	23,009	46,804	3
2006.....	72,432	-	-	-	-	1	12	104	655	3,812	22,915	44,933	-
2005.....	71,599	-	-	-	-	2	10	80	648	3,813	23,139	43,906	1
2004.....	65,965	-	-	-	1	1	7	101	542	3,641	21,882	39,790	-
2003.....	63,457	-	-	-	-	-	11	78	554	3,836	21,157	37,821	-
2002.....	58,866	-	-	1	-	2	12	52	510	3,602	20,135	34,552	-
2001.....	53,852	-	-	-	-	2	3	73	529	3,422	18,542	31,281	-
2000.....	49,558	-	-	-	-	2	6	64	491	3,431	17,253	28,309	2
1999.....	44,536	-	-	-	1	-	7	60	448	3,204	15,836	24,980	-
All races, male													
2007.....	21,800	-	-	-	-	-	3	43	330	1,694	8,250	11,479	1
2006.....	21,151	-	-	-	-	-	8	53	304	1,709	8,295	10,782	-
2005.....	20,559	-	-	-	-	-	6	30	287	1,648	8,326	10,262	-
2004.....	18,974	-	-	-	-	1	4	38	253	1,558	7,777	9,343	-
2003.....	18,335	-	-	-	-	-	4	34	243	1,684	7,606	8,764	-
2002.....	16,989	-	-	-	-	2	8	27	228	1,547	7,227	7,950	-
2001.....	15,762	-	-	-	-	2	3	38	228	1,472	6,795	7,224	-
2000.....	14,438	-	-	-	-	1	4	34	210	1,554	6,228	6,406	1
1999.....	13,391	-	-	-	1	-	3	28	222	1,437	5,904	5,796	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alzheimer's disease (G30)													
All races, female													
2007.....	52,832	-	-	-	-	1	5	52	398	2,290	14,759	35,325	2
2006.....	51,281	-	-	-	-	1	4	51	351	2,103	14,620	34,151	-
2005.....	51,040	-	-	-	-	2	4	50	361	2,165	14,813	33,644	1
2004.....	46,991	-	-	-	1	-	3	63	289	2,083	14,105	30,447	-
2003.....	45,122	-	-	-	-	-	7	44	311	2,152	13,551	29,057	-
2002.....	41,877	-	-	1	-	-	4	25	282	2,055	12,908	26,602	-
2001.....	38,090	-	-	-	-	-	-	35	301	1,950	11,747	24,057	-
2000.....	35,120	-	-	-	-	1	2	30	281	1,877	11,025	21,903	1
1999.....	31,145	-	-	-	-	-	4	32	226	1,767	9,932	19,184	-
White, both sexes													
2007 1/.....	68,933	-	-	-	-	1	6	86	676	3,563	21,115	43,484	2
2006 1/.....	67,088	-	-	-	-	1	9	94	598	3,433	21,126	41,827	-
2005 1/.....	66,191	-	-	-	-	2	10	72	592	3,383	21,272	40,860	-
2004 1/.....	61,087	-	-	-	1	1	6	97	495	3,263	20,130	37,094	-
2003 1/.....	59,184	-	-	-	-	-	11	74	518	3,452	19,625	35,504	-
2002.....	55,058	-	-	1	-	1	7	49	468	3,282	18,791	32,459	-
2001.....	50,348	-	-	-	-	2	3	67	499	3,136	17,226	29,415	-
2000.....	46,460	-	-	-	-	2	6	55	450	3,139	16,124	26,683	1
1999.....	41,877	-	-	-	1	-	5	53	413	2,926	14,833	23,646	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alzheimer's disease (G30)													
White, male													
2007 1/.....	20,185	-	-	-	-	-	3	40	302	1,522	7,608	10,710	-
2006 1/.....	19,654	-	-	-	-	-	7	50	278	1,546	7,701	10,072	-
2005 1/.....	18,990	-	-	-	-	-	6	26	265	1,465	7,669	9,559	-
2004 1/.....	17,605	-	-	-	-	1	3	37	232	1,400	7,208	8,724	-
2003 1/.....	17,086	-	-	-	-	-	4	32	224	1,522	7,068	8,236	-
2002.....	15,874	-	-	-	-	1	5	24	210	1,409	6,746	7,479	-
2001.....	14,714	-	-	-	-	2	3	34	210	1,349	6,327	6,789	-
2000.....	13,524	-	-	-	-	1	4	29	193	1,432	5,854	6,010	1
1999.....	12,585	-	-	-	1	-	2	26	205	1,316	5,564	5,471	-
White, female													
2007 1/.....	48,748	-	-	-	-	1	3	46	374	2,041	13,507	32,774	2
2006 1/.....	47,434	-	-	-	-	1	2	44	320	1,887	13,425	31,755	-
2005 1/.....	47,201	-	-	-	-	2	4	46	327	1,918	13,603	31,301	-
2004 1/.....	43,482	-	-	-	1	-	3	60	263	1,863	12,922	28,370	-
2003 1/.....	42,098	-	-	-	-	-	7	42	294	1,930	12,557	27,268	-
2002.....	39,184	-	-	1	-	-	2	25	258	1,873	12,045	24,980	-
2001.....	35,634	-	-	-	-	-	-	33	289	1,787	10,899	22,626	-
2000.....	32,936	-	-	-	-	1	2	26	257	1,707	10,270	20,673	-
1999.....	29,292	-	-	-	-	-	3	27	208	1,610	9,269	18,175	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alzheimer's disease (G30)													
Black, both sexes													
2007 1/.....	4,760	-	-	-	-	-	2	8	38	360	1,627	2,724	1
2006 1/.....	4,455	-	-	-	-	-	3	9	53	336	1,495	2,559	-
2005 1/.....	4,620	-	-	-	-	-	-	6	47	390	1,596	2,580	1
2004 1/.....	4,232	-	-	-	-	-	1	4	37	334	1,523	2,333	-
2003 1/.....	3,718	-	-	-	-	-	-	4	29	328	1,356	2,001	-
2002.....	3,341	-	-	-	-	1	4	3	32	283	1,187	1,831	-
2001.....	3,114	-	-	-	-	-	-	5	27	254	1,178	1,650	-
2000.....	2,728	-	-	-	-	-	-	5	30	261	1,008	1,423	1
1999.....	2,342	-	-	-	-	-	1	5	29	249	890	1,168	-
Black, male													
2007 1/.....	1,301	-	-	-	-	-	-	3	17	152	539	589	1
2006 1/.....	1,190	-	-	-	-	-	1	3	25	143	491	527	-
2005 1/.....	1,292	-	-	-	-	-	-	4	18	163	549	558	-
2004 1/.....	1,147	-	-	-	-	-	1	1	15	138	491	501	-
2003 1/.....	1,087	-	-	-	-	-	-	2	17	144	483	441	-
2002.....	959	-	-	-	-	1	3	3	14	126	427	385	-
2001.....	910	-	-	-	-	-	-	3	18	111	413	365	-
2000.....	776	-	-	-	-	-	-	3	16	113	322	322	-
1999.....	692	-	-	-	-	-	-	2	15	112	292	271	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alzheimer's disease (G30)													
Black, female													
2007 1/.....	3,459	-	-	-	-	-	2	5	21	208	1,088	2,135	-
2006 1/.....	3,265	-	-	-	-	-	2	6	28	193	1,004	2,032	-
2005 1/.....	3,328	-	-	-	-	-	-	2	29	227	1,047	2,022	1
2004 1/.....	3,085	-	-	-	-	-	-	3	22	196	1,032	1,832	-
2003 1/.....	2,631	-	-	-	-	-	-	2	12	184	873	1,560	-
2002.....	2,382	-	-	-	-	-	1	-	18	157	760	1,446	-
2001.....	2,204	-	-	-	-	-	-	2	9	143	765	1,285	-
2000.....	1,952	-	-	-	-	-	-	2	14	148	686	1,101	1
1999.....	1,650	-	-	-	-	-	1	3	14	137	598	897	-
American Indian 2/, both sexes													
2007 1/.....	191	-	-	-	-	-	-	1	5	24	65	96	-
2006 1/.....	169	-	-	-	-	-	-	-	1	15	73	80	-
2005 1/.....	173	-	-	-	-	-	-	-	-	10	69	94	-
2004 1/.....	112	-	-	-	-	-	-	-	2	11	43	56	-
2003 1/.....	122	-	-	-	-	-	-	-	1	15	41	65	-
2002.....	115	-	-	-	-	-	-	-	1	14	42	58	-
2001.....	93	-	-	-	-	-	-	1	1	10	40	41	-
2000.....	95	-	-	-	-	-	-	-	3	14	39	39	-
1999.....	86	-	-	-	-	-	-	-	3	13	29	41	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alzheimer's disease (G30)													
American Indian 2/, male													
2007 1/.....	69	-	-	-	-	-	-	-	5	5	25	34	-
2006 1/.....	62	-	-	-	-	-	-	-	-	9	25	28	-
2005 1/.....	48	-	-	-	-	-	-	-	-	6	19	23	-
2004 1/.....	32	-	-	-	-	-	-	-	2	6	14	10	-
2003 1/.....	22	-	-	-	-	-	-	-	-	4	9	9	-
2002.....	35	-	-	-	-	-	-	-	1	5	15	14	-
2001.....	29	-	-	-	-	-	-	1	-	4	17	7	-
2000.....	26	-	-	-	-	-	-	-	-	4	13	9	-
1999.....	23	-	-	-	-	-	-	-	1	5	11	6	-
American Indian 2/, female													
2007 1/.....	122	-	-	-	-	-	-	1	-	19	40	62	-
2006 1/.....	107	-	-	-	-	-	-	-	1	6	48	52	-
2005 1/.....	125	-	-	-	-	-	-	-	-	4	50	71	-
2004 1/.....	80	-	-	-	-	-	-	-	-	5	29	46	-
2003 1/.....	100	-	-	-	-	-	-	-	1	11	32	56	-
2002.....	80	-	-	-	-	-	-	-	-	9	27	44	-
2001.....	64	-	-	-	-	-	-	-	1	6	23	34	-
2000.....	69	-	-	-	-	-	-	-	3	10	26	30	-
1999.....	63	-	-	-	-	-	-	-	2	8	18	35	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Alzheimer's disease (G30)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	748	-	-	-	-	-	-	9	37	202	500	-
2006 1/.....	720	-	-	-	-	-	1	3	28	221	467	-
2005 1/.....	615	-	-	-	-	-	2	9	30	202	372	-
2004 1/.....	534	-	-	-	-	-	-	8	33	186	307	-
2003 1/.....	433	-	-	-	-	-	-	6	41	135	251	-
2002.....	352	-	-	-	-	-	1	9	23	115	204	-
2001.....	297	-	-	-	-	-	-	2	22	98	175	-
2000.....	275	-	-	-	-	-	4	8	17	82	164	-
1999.....	231	-	-	-	-	-	1	2	3	16	125	-

Asian or Pacific Islander 3/, male

2007 1/.....	245	-	-	-	-	-	-	6	15	78	146	-
2006 1/.....	245	-	-	-	-	-	-	1	11	78	155	-
2005 1/.....	229	-	-	-	-	-	-	4	14	89	122	-
2004 1/.....	190	-	-	-	-	-	-	4	14	64	108	-
2003 1/.....	140	-	-	-	-	-	-	2	14	46	78	-
2002.....	121	-	-	-	-	-	-	3	7	39	72	-
2001.....	109	-	-	-	-	-	-	-	8	38	63	-
2000.....	112	-	-	-	-	-	2	1	5	39	65	-
1999.....	91	-	-	-	-	-	1	-	1	4	48	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Alzheimer's disease (G30)

Asian or Pacific Islander 3/, female

2007 1/.....	503	-	-	-	-	-	-	-	3	22	124	354	-
2006 1/.....	475	-	-	-	-	-	-	1	2	17	143	312	-
2005 1/.....	386	-	-	-	-	-	-	2	5	16	113	250	-
2004 1/.....	344	-	-	-	-	-	-	-	4	19	122	199	-
2003 1/.....	293	-	-	-	-	-	-	-	4	27	89	173	-
2002.....	231	-	-	-	-	-	1	-	6	16	76	132	-
2001.....	188	-	-	-	-	-	-	-	2	14	60	112	-
2000.....	163	-	-	-	-	-	-	2	7	12	43	99	-
1999.....	140	-	-	-	-	-	-	2	2	12	47	77	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Major cardiovascular diseases (I00-I78)

All races, both sexes

2007.....	806,156	571	230	338	1,369	3,950	14,867	46,280	80,797	115,623	229,050	313,044	37
2006.....	823,746	508	218	363	1,376	4,099	15,420	47,037	80,714	119,102	240,862	314,006	41
2005.....	856,030	503	217	367	1,391	4,041	15,852	46,928	79,896	124,366	256,362	326,066	41
2004.....	861,190	564	234	336	1,333	4,006	16,138	46,153	77,984	128,816	264,037	321,544	45
2003.....	902,443	555	234	347	1,447	4,089	16,930	46,279	79,497	137,220	279,358	336,449	38
2002.....	918,628	640	223	358	1,283	3,986	16,962	45,800	78,430	144,071	287,415	339,409	51
2001.....	922,334	600	285	365	1,280	4,038	16,686	44,450	76,224	148,409	291,995	337,930	72
2000.....	936,923	636	234	362	1,309	3,792	16,624	43,583	77,482	156,187	296,202	340,450	62
1999.....	950,314	644	234	354	1,315	3,873	17,010	42,512	77,833	163,428	302,552	340,510	49

All races, male

2007.....	388,049	323	104	181	874	2,567	10,107	32,038	54,646	69,432	113,012	104,737	28
2006.....	394,840	283	115	207	882	2,765	10,370	32,633	54,055	71,210	118,431	103,859	30
2005.....	405,780	282	126	200	914	2,657	10,664	32,548	53,484	73,733	125,037	106,106	29
2004.....	406,458	307	120	170	867	2,602	10,806	32,057	52,014	76,179	128,490	102,808	38
2003.....	423,810	310	116	204	893	2,609	11,302	32,260	52,804	81,512	135,868	105,903	29
2002.....	429,682	343	122	194	845	2,568	11,356	31,928	52,037	85,319	138,770	106,164	36
2001.....	428,065	315	149	184	807	2,555	11,180	30,860	50,320	87,756	140,789	103,097	53
2000.....	435,912	341	128	206	783	2,378	11,110	30,535	51,423	92,510	142,797	103,658	43
1999.....	441,827	347	127	199	799	2,442	11,471	29,745	51,525	96,904	145,567	102,663	38

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Major cardiovascular diseases (I00-I78)

All races, female

2007.....	418,107	248	126	157	495	1,383	4,760	14,242	26,151	46,191	116,038	208,307	9
2006.....	428,906	225	103	156	494	1,334	5,050	14,404	26,659	47,892	122,431	210,147	11
2005.....	450,250	221	91	167	477	1,384	5,188	14,380	26,412	50,633	131,325	219,960	12
2004.....	454,732	257	114	166	466	1,404	5,332	14,096	25,970	52,637	135,547	218,736	7
2003.....	478,633	245	118	143	554	1,480	5,628	14,019	26,693	55,708	143,490	230,546	9
2002.....	488,946	297	101	164	438	1,418	5,606	13,872	26,393	58,752	148,645	233,245	15
2001.....	494,269	285	136	181	473	1,483	5,506	13,590	25,904	60,653	151,206	234,833	19
2000.....	501,011	295	106	156	526	1,414	5,514	13,048	26,059	63,677	153,405	236,792	19
1999.....	508,487	297	107	155	516	1,431	5,539	12,767	26,308	66,524	156,985	237,847	11

White, both sexes

2007 1/.....	691,283	359	151	240	905	2,541	10,311	33,801	62,439	93,995	200,863	285,653	25
2006 1/.....	707,419	314	151	249	955	2,642	10,764	34,227	62,546	97,032	211,545	286,961	33
2005 1/.....	737,248	318	141	240	926	2,574	11,085	33,972	61,969	102,044	225,723	298,229	27
2004 1/.....	742,775	377	149	228	894	2,574	11,418	33,462	60,692	106,114	233,121	293,716	30
2003 1/.....	780,020	343	148	234	951	2,663	11,819	33,491	61,975	113,130	247,328	307,907	31
2002.....	796,748	421	160	244	863	2,565	12,029	33,531	61,434	119,894	254,894	310,678	35
2001.....	801,052	383	185	258	838	2,678	11,628	32,580	59,631	123,621	259,517	309,688	45
2000.....	816,219	434	161	242	858	2,505	11,679	32,222	60,871	131,092	263,970	312,140	45
1999.....	829,464	434	161	250	886	2,598	11,884	31,589	61,248	137,588	270,242	312,552	32

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Major cardiovascular diseases (I00-I78)

White, male

2007 1/.....	331,560	201	65	131	561	1,691	7,203	24,251	43,196	57,458	100,338	96,445	20
2006 1/.....	337,670	172	76	156	636	1,840	7,531	24,569	42,742	58,935	105,402	95,587	24
2005 1/.....	348,454	177	86	137	610	1,748	7,753	24,447	42,448	61,578	111,597	97,854	19
2004 1/.....	349,806	207	82	117	588	1,737	7,985	24,177	41,407	64,018	114,985	94,477	26
2003 1/.....	365,390	184	67	140	593	1,754	8,258	24,303	42,278	68,570	121,648	97,573	22
2002.....	372,088	230	92	131	577	1,708	8,399	24,343	41,782	72,502	124,636	97,662	26
2001.....	370,918	212	95	136	535	1,748	8,157	23,571	40,443	74,628	126,557	94,802	34
2000.....	379,095	233	96	146	528	1,645	8,189	23,533	41,460	79,359	128,696	95,176	34
1999.....	384,572	234	90	140	531	1,697	8,383	22,953	41,561	83,294	131,574	94,090	25

White, female

2007 1/.....	359,723	158	86	109	344	850	3,108	9,550	19,243	36,537	100,525	189,208	5
2006 1/.....	369,749	142	75	93	319	802	3,233	9,658	19,804	38,097	106,143	191,374	9
2005 1/.....	388,794	141	55	103	316	826	3,332	9,525	19,521	40,466	114,126	200,375	8
2004 1/.....	392,969	170	67	111	306	837	3,433	9,285	19,285	42,096	118,136	199,239	4
2003 1/.....	414,630	159	81	94	358	909	3,561	9,188	19,697	44,560	125,680	210,334	9
2002.....	424,660	191	68	113	286	857	3,630	9,188	19,652	47,392	130,258	213,016	9
2001.....	430,134	171	90	122	303	930	3,471	9,009	19,188	48,993	132,960	214,886	11
2000.....	437,124	201	65	96	330	860	3,490	8,689	19,411	51,733	135,274	216,964	11
1999.....	444,892	200	71	110	355	901	3,501	8,636	19,687	54,294	138,668	218,462	7

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Major cardiovascular diseases (I00-I78)

Black, both sexes

2007 1/.....	96,051	175	59	78	418	1,220	3,970	11,163	16,086	18,303	22,826	21,742	11
2006 1/.....	97,389	170	57	90	373	1,287	4,109	11,424	15,841	18,620	23,874	21,536	8
2005 1/.....	100,099	153	60	104	405	1,284	4,212	11,565	15,671	18,845	25,274	22,513	13
2004 1/.....	100,477	145	75	100	379	1,255	4,257	11,387	15,241	19,448	25,592	22,584	14
2003 1/.....	104,181	173	73	95	438	1,251	4,535	11,455	15,368	20,631	26,636	23,521	5
2002.....	104,215	186	51	95	363	1,267	4,414	11,044	15,008	20,693	27,232	23,847	15
2001.....	104,316	186	87	88	399	1,223	4,543	10,653	14,653	21,304	27,327	23,828	25
2000.....	104,280	171	63	96	399	1,171	4,456	10,206	14,665	21,779	27,221	24,036	17
1999.....	104,693	177	61	93	394	1,134	4,660	9,820	14,654	22,492	27,411	23,781	16

Black, male

2007 1/.....	46,698	106	32	40	282	743	2,489	6,838	9,955	10,027	10,082	6,097	7
2006 1/.....	47,276	98	35	42	217	802	2,449	7,093	9,773	10,274	10,423	6,064	6
2005 1/.....	47,584	89	30	53	266	779	2,524	7,136	9,589	10,139	10,830	6,140	9
2004 1/.....	47,360	80	37	49	248	734	2,495	6,964	9,270	10,337	10,896	6,239	11
2003 1/.....	48,853	105	43	53	271	732	2,653	7,032	9,143	10,958	11,578	6,280	5
2002.....	48,264	97	25	51	224	749	2,582	6,727	8,974	10,903	11,460	6,463	9
2001.....	48,237	94	46	40	244	718	2,655	6,461	8,607	11,190	11,674	6,490	18
2000.....	47,980	91	27	48	219	657	2,590	6,213	8,687	11,208	11,565	6,666	9
1999.....	48,578	94	32	53	244	660	2,751	6,052	8,730	11,720	11,470	6,760	12

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Major cardiovascular diseases (I00-I78)

Black, female

2007 1/.....	49,353	69	27	38	136	477	1,481	4,325	6,131	8,276	12,744	15,645	4
2006 1/.....	50,113	72	22	48	156	485	1,660	4,331	6,068	8,346	13,451	15,472	2
2005 1/.....	52,515	64	30	51	139	505	1,688	4,429	6,082	8,706	14,444	16,373	4
2004 1/.....	53,117	65	38	51	131	521	1,762	4,423	5,971	9,111	14,696	16,345	3
2003 1/.....	55,328	68	30	42	167	519	1,882	4,423	6,225	9,673	15,058	17,241	-
2002.....	55,951	89	26	44	139	518	1,832	4,317	6,034	9,790	15,772	17,384	6
2001.....	56,079	92	41	48	155	505	1,888	4,192	6,046	10,114	15,653	17,338	7
2000.....	56,300	80	36	48	180	514	1,866	3,993	5,978	10,571	15,656	17,370	8
1999.....	56,115	83	29	40	150	474	1,909	3,768	5,924	10,772	15,941	17,021	4

American Indian 2/, both sexes

2007 1/.....	3,482	13	8	1	16	65	171	384	590	716	851	667	-
2006 1/.....	3,496	9	2	5	19	40	178	404	583	725	875	656	-
2005 1/.....	3,534	8	1	2	15	52	157	414	604	757	839	685	-
2004 1/.....	3,408	12	2	3	15	58	152	351	539	716	858	702	-
2003 1/.....	3,488	9	5	3	14	52	176	385	592	759	839	652	2
2002.....	3,259	6	6	7	20	41	147	319	517	723	845	627	1
2001.....	3,162	11	3	4	10	36	162	330	491	740	783	592	-
2000.....	3,198	7	2	7	12	30	150	309	498	718	824	641	-
1999.....	3,121	8	5	6	10	43	139	278	471	730	824	607	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Major cardiovascular diseases (I00-I78)

American Indian 2/, male

2007 1/.....	1,913	5	5	1	8	47	109	275	385	420	432	226	-
2006 1/.....	1,855	4	2	2	7	26	129	289	379	420	395	202	-
2005 1/.....	1,837	3	1	-	8	37	102	261	387	419	389	230	-
2004 1/.....	1,786	6	-	2	5	41	97	220	349	411	418	237	-
2003 1/.....	1,882	5	2	-	10	38	115	258	369	433	399	251	2
2002.....	1,752	4	2	4	16	29	102	230	331	408	396	229	1
2001.....	1,663	3	2	2	8	20	103	220	325	420	355	205	-
2000.....	1,670	3	1	3	8	16	96	208	307	424	396	208	-
1999.....	1,613	5	2	2	7	19	93	189	280	432	395	189	-

American Indian 2/, female

2007 1/.....	1,569	8	3	-	8	18	62	109	205	296	419	441	-
2006 1/.....	1,641	5	-	3	12	14	49	115	204	305	480	454	-
2005 1/.....	1,697	5	-	2	7	15	55	153	217	338	450	455	-
2004 1/.....	1,622	6	2	1	10	17	55	131	190	305	440	465	-
2003 1/.....	1,606	4	3	3	4	14	61	127	223	326	440	401	-
2002.....	1,507	2	4	3	4	12	45	89	186	315	449	398	-
2001.....	1,499	8	1	2	2	16	59	110	166	320	428	387	-
2000.....	1,528	4	1	4	4	14	54	101	191	294	428	433	-
1999.....	1,508	3	3	4	3	24	46	89	191	298	429	418	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Major cardiovascular diseases (I00-I78)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	15,340	24	12	19	30	124	415	932	1,682	2,609	4,510	4,982	1
2006 1/.....	15,442	15	8	19	29	130	369	982	1,744	2,725	4,568	4,853	-
2005 1/.....	15,149	24	15	21	45	131	398	977	1,652	2,720	4,526	4,639	1
2004 1/.....	14,530	30	8	5	45	119	311	953	1,512	2,538	4,466	4,542	1
2003 1/.....	14,754	30	8	15	44	123	400	948	1,562	2,700	4,555	4,369	-
2002.....	14,406	27	6	12	37	113	372	906	1,471	2,761	4,444	4,257	-
2001.....	13,804	20	10	15	33	101	353	887	1,449	2,744	4,368	3,822	2
2000.....	13,226	24	8	17	40	86	339	846	1,448	2,598	4,187	3,633	-
1999.....	13,036	25	7	5	25	98	327	825	1,460	2,618	4,075	3,570	1

Asian or Pacific Islander 3/, male

2007 1/.....	7,878	11	2	9	23	86	306	674	1,110	1,527	2,160	1,969	1
2006 1/.....	8,039	9	2	7	22	97	261	682	1,161	1,581	2,211	2,006	-
2005 1/.....	7,905	13	9	10	30	93	285	704	1,060	1,597	2,221	1,882	1
2004 1/.....	7,506	14	1	2	26	90	229	696	988	1,413	2,191	1,855	1
2003 1/.....	7,685	16	4	11	19	85	276	667	1,014	1,551	2,243	1,799	-
2002.....	7,578	12	3	8	28	82	273	628	950	1,506	2,278	1,810	-
2001.....	7,247	6	6	6	20	69	265	608	945	1,518	2,203	1,600	1
2000.....	7,167	14	4	9	28	60	235	581	969	1,519	2,140	1,608	-
1999.....	7,064	14	3	4	17	66	244	551	954	1,458	2,128	1,624	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Major cardiovascular diseases (I00-I78)

Asian or Pacific Islander 3/, female

2007 1/.....	7,462	13	10	10	7	38	109	258	572	1,082	2,350	3,013	-
2006 1/.....	7,403	6	6	12	7	33	108	300	583	1,144	2,357	2,847	-
2005 1/.....	7,244	11	6	11	15	38	113	273	592	1,123	2,305	2,757	-
2004 1/.....	7,024	16	7	3	19	29	82	257	524	1,125	2,275	2,687	-
2003 1/.....	7,069	14	4	4	25	38	124	281	548	1,149	2,312	2,570	-
2002.....	6,828	15	3	4	9	31	99	278	521	1,255	2,166	2,447	-
2001.....	6,557	14	4	9	13	32	88	279	504	1,226	2,165	2,222	1
2000.....	6,059	10	4	8	12	26	104	265	479	1,079	2,047	2,025	-
1999.....	5,972	11	4	1	8	32	83	274	506	1,160	1,947	1,946	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of heart (I00-I09,I11,I13,I20-I51)													
All races, both sexes													
2007.....	616,067	424	173	241	1,084	3,223	11,839	37,434	65,527	89,589	171,257	235,249	27
2006.....	631,636	346	161	253	1,076	3,307	12,339	38,095	65,477	92,752	180,451	237,339	40
2005.....	652,091	358	151	252	1,119	3,249	12,688	38,103	65,208	96,729	190,693	243,504	37
2004.....	652,486	421	187	245	1,038	3,163	12,925	37,556	63,613	99,999	195,379	237,924	36
2003.....	685,089	439	186	264	1,133	3,250	13,600	37,732	65,060	107,263	207,331	248,796	35
2002.....	696,947	500	165	255	1,022	3,165	13,688	37,570	64,234	112,547	213,581	250,173	47
2001.....	700,142	479	225	272	999	3,160	13,326	36,399	62,486	116,299	216,992	249,439	66
2000.....	710,760	493	181	271	1,031	2,958	13,181	35,480	63,399	122,405	220,060	251,242	59
1999.....	725,192	524	183	277	1,069	3,066	13,600	34,994	64,167	129,253	226,152	251,860	47
All races, male													
2007.....	309,821	237	82	133	700	2,177	8,345	26,911	45,558	55,532	88,005	82,120	21
2006.....	315,706	187	89	140	711	2,298	8,625	27,433	45,177	57,265	92,378	81,373	30
2005.....	322,841	197	81	131	770	2,206	8,931	27,486	44,895	59,050	96,580	82,489	25
2004.....	321,973	228	94	122	691	2,143	9,011	27,050	43,788	61,000	98,812	79,002	32
2003.....	336,095	247	89	155	718	2,129	9,463	27,304	44,522	65,568	104,646	81,227	27
2002.....	340,933	267	87	141	686	2,131	9,638	27,239	43,900	68,658	107,217	80,935	34
2001.....	339,095	243	121	138	647	2,068	9,346	26,280	42,519	70,637	108,576	78,473	47
2000.....	344,807	260	107	163	638	1,922	9,286	25,927	43,285	74,589	109,693	78,896	41
1999.....	351,617	270	104	157	664	1,988	9,640	25,489	43,662	78,836	112,470	78,300	37

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of heart (I00-I09,I11,I13,I20-I51)													
All races, female													
2007.....	306,246	187	91	108	384	1,046	3,494	10,523	19,969	34,057	83,252	153,129	6
2006.....	315,930	159	72	113	365	1,009	3,714	10,662	20,300	35,487	88,073	155,966	10
2005.....	329,250	161	70	121	349	1,043	3,757	10,617	20,313	37,679	94,113	161,015	12
2004.....	330,513	193	93	123	347	1,020	3,914	10,506	19,825	38,999	96,567	158,922	4
2003.....	348,994	192	97	109	415	1,121	4,137	10,428	20,538	41,695	102,685	167,569	8
2002.....	356,014	233	78	114	336	1,034	4,050	10,331	20,334	43,889	106,364	169,238	13
2001.....	361,047	236	104	134	352	1,092	3,980	10,119	19,967	45,662	108,416	170,966	19
2000.....	365,953	233	74	108	393	1,036	3,895	9,553	20,114	47,816	110,367	172,346	18
1999.....	373,575	254	79	120	405	1,078	3,960	9,505	20,505	50,417	113,682	173,560	10
White, both sexes													
2007 1/.....	531,636	254	112	166	706	2,073	8,406	28,135	51,585	73,600	151,041	215,539	19
2006 1/.....	545,974	210	110	170	730	2,143	8,879	28,462	51,715	76,373	159,346	217,804	32
2005 1/.....	564,796	228	93	159	736	2,084	9,101	28,374	51,407	80,273	168,711	223,603	27
2004 1/.....	565,703	287	116	159	678	2,040	9,351	27,969	50,449	83,145	173,433	218,052	24
2003 1/.....	594,842	273	117	174	736	2,120	9,780	28,057	51,648	89,233	184,357	228,319	28
2002.....	606,876	332	115	176	667	2,040	9,967	28,279	51,089	94,492	190,122	229,565	32
2001.....	610,638	312	141	187	639	2,086	9,510	27,472	49,719	97,764	193,525	229,241	42
2000.....	621,719	343	126	177	656	1,937	9,532	26,991	50,581	103,449	196,833	231,050	44
1999.....	635,118	352	124	193	700	2,048	9,718	26,693	51,274	109,578	202,692	231,714	32

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of heart (I00-I09,I11,I13,I20-I51)													
White, male													
2007 1/.....	266,908	143	47	90	445	1,433	6,058	20,876	36,646	46,500	78,676	75,979	15
2006 1/.....	272,117	114	60	99	505	1,537	6,437	21,158	36,407	47,936	82,625	75,215	24
2005 1/.....	279,324	125	55	90	509	1,469	6,644	21,173	36,235	49,895	86,700	76,410	19
2004 1/.....	278,914	156	61	82	459	1,431	6,796	20,911	35,512	51,740	88,944	72,801	21
2003 1/.....	291,560	150	52	105	464	1,449	7,093	21,062	36,294	55,619	94,151	75,101	20
2002.....	296,904	181	65	99	457	1,420	7,288	21,283	35,789	58,873	96,734	74,691	24
2001.....	295,556	170	74	98	414	1,408	6,981	20,620	34,724	60,615	98,006	72,415	31
2000.....	301,551	182	80	111	413	1,316	7,020	20,480	35,422	64,464	99,295	72,735	33
1999.....	307,585	183	74	108	429	1,376	7,185	20,144	35,695	68,247	102,139	71,980	25
White, female													
2007 1/.....	264,728	111	65	76	261	640	2,348	7,259	14,939	27,100	72,365	139,560	4
2006 1/.....	273,857	96	50	71	225	606	2,442	7,304	15,308	28,437	76,721	142,589	8
2005 1/.....	285,472	103	38	69	227	615	2,457	7,201	15,172	30,378	82,011	147,193	8
2004 1/.....	286,789	131	55	77	219	609	2,555	7,058	14,937	31,405	84,489	145,251	3
2003 1/.....	303,282	123	65	69	272	671	2,687	6,995	15,354	33,614	90,206	153,218	8
2002.....	309,972	151	50	77	210	620	2,679	6,996	15,300	35,619	93,388	154,874	8
2001.....	315,082	142	67	89	225	678	2,529	6,852	14,995	37,149	95,519	156,826	11
2000.....	320,168	161	46	66	243	621	2,512	6,511	15,159	38,985	97,538	158,315	11
1999.....	327,533	169	50	85	271	672	2,533	6,549	15,579	41,331	100,553	159,734	7

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

Black, both sexes

2007 1/.....	71,209	139	43	63	348	997	2,992	8,357	12,289	13,642	16,559	15,773	7
2006 1/.....	72,253	116	44	67	308	1,033	3,078	8,612	12,050	13,932	17,370	15,635	8
2005 1/.....	74,159	107	45	77	334	1,022	3,189	8,716	12,115	14,005	18,351	16,189	9
2004 1/.....	74,225	101	62	80	316	980	3,236	8,680	11,679	14,552	18,344	16,184	11
2003 1/.....	77,372	133	57	76	354	991	3,428	8,704	11,821	15,605	19,261	16,937	5
2002.....	77,621	144	40	65	306	1,004	3,356	8,413	11,696	15,601	19,757	17,225	14
2001.....	77,674	141	74	73	326	970	3,451	8,044	11,348	16,120	19,924	17,181	22
2000.....	77,523	122	49	76	333	937	3,305	7,680	11,385	16,590	19,802	17,229	15
1999.....	78,574	144	50	77	340	914	3,546	7,525	11,475	17,240	20,038	17,210	15

Black, male

2007 1/.....	35,669	79	28	35	235	632	1,951	5,304	7,762	7,595	7,478	4,565	5
2006 1/.....	36,230	62	27	35	183	663	1,892	5,511	7,568	7,854	7,871	4,558	6
2005 1/.....	36,343	62	19	36	229	633	1,989	5,569	7,505	7,676	8,069	4,551	5
2004 1/.....	36,222	55	32	37	207	597	1,961	5,468	7,243	7,914	8,030	4,668	10
2003 1/.....	37,466	81	32	42	232	580	2,087	5,502	7,155	8,495	8,603	4,652	5
2002.....	37,094	76	18	33	192	619	2,060	5,286	7,124	8,357	8,539	4,781	9
2001.....	37,016	65	41	35	209	586	2,092	5,016	6,810	8,604	8,731	4,812	15
2000.....	36,740	62	24	42	193	547	2,009	4,837	6,882	8,672	8,579	4,885	8
1999.....	37,576	72	27	45	215	546	2,193	4,774	7,004	9,155	8,533	5,000	12

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of heart (I00-I09,I11,I13,I20-I51)													
Black, female													
2007 1/.....	35,540	60	15	28	113	365	1,041	3,053	4,527	6,047	9,081	11,208	2
2006 1/.....	36,023	54	17	32	125	370	1,186	3,101	4,482	6,078	9,499	11,077	2
2005 1/.....	37,816	45	26	41	105	389	1,200	3,147	4,610	6,329	10,282	11,638	4
2004 1/.....	38,003	46	30	43	109	383	1,275	3,212	4,436	6,638	10,314	11,516	1
2003 1/.....	39,906	52	25	34	122	411	1,341	3,202	4,666	7,110	10,658	12,285	-
2002.....	40,527	68	22	32	114	385	1,296	3,127	4,572	7,244	11,218	12,444	5
2001.....	40,658	76	33	38	117	384	1,359	3,028	4,538	7,516	11,193	12,369	7
2000.....	40,783	60	25	34	140	390	1,296	2,843	4,503	7,918	11,223	12,344	7
1999.....	40,998	72	23	32	125	368	1,353	2,751	4,471	8,085	11,505	12,210	3
American Indian 2/, both sexes													
2007 1/.....	2,648	10	7	1	12	60	137	315	468	543	629	466	-
2006 1/.....	2,736	7	-	3	14	35	141	332	492	571	671	470	-
2005 1/.....	2,659	5	1	2	13	46	120	330	485	552	597	508	-
2004 1/.....	2,598	11	2	2	13	48	118	282	439	535	624	524	-
2003 1/.....	2,712	9	4	2	11	44	135	308	492	592	623	490	2
2002.....	2,467	5	6	5	17	34	107	246	412	542	633	459	1
2001.....	2,402	9	3	3	10	29	125	270	416	554	568	415	-
2000.....	2,417	7	1	5	11	24	122	249	394	548	597	459	-
1999.....	2,404	6	5	3	9	30	106	231	385	584	616	429	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

American Indian 2/, male

2007 1/.....	1,520	5	5	1	5	43	90	234	312	332	327	166	-
2006 1/.....	1,532	3	-	1	5	23	104	242	322	346	324	162	-
2005 1/.....	1,471	1	1	-	7	33	84	221	328	319	294	183	-
2004 1/.....	1,437	6	-	1	5	37	83	181	296	329	313	186	-
2003 1/.....	1,529	5	1	-	7	32	95	218	315	349	308	197	2
2002.....	1,412	3	2	4	14	25	82	189	277	324	314	177	1
2001.....	1,358	3	2	1	8	18	85	191	287	332	285	146	-
2000.....	1,341	3	-	3	8	16	83	179	252	336	306	155	-
1999.....	1,302	3	2	1	6	14	75	163	241	353	303	141	-

American Indian 2/, female

2007 1/.....	1,128	5	2	-	7	17	47	81	156	211	302	300	-
2006 1/.....	1,204	4	-	2	9	12	37	90	170	225	347	308	-
2005 1/.....	1,188	4	-	2	6	13	36	109	157	233	303	325	-
2004 1/.....	1,161	5	2	1	8	11	35	101	143	206	311	338	-
2003 1/.....	1,183	4	3	2	4	12	40	90	177	243	315	293	-
2002.....	1,055	2	4	1	3	9	25	57	135	218	319	282	-
2001.....	1,044	6	1	2	2	11	40	79	129	222	283	269	-
2000.....	1,076	4	1	2	3	8	39	70	142	212	291	304	-
1999.....	1,102	3	3	2	3	16	31	68	144	231	313	288	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of heart (I00-I09,I11,I13,I20-I51)													
Asian or Pacific Islander 3/, both sexes													
2007 1/.....	10,574	21	11	11	18	93	304	627	1,185	1,804	3,028	3,471	1
2006 1/.....	10,673	13	7	13	24	96	241	689	1,220	1,876	3,064	3,430	-
2005 1/.....	10,477	18	12	14	36	97	278	683	1,201	1,899	3,034	3,204	1
2004 1/.....	9,960	22	7	4	31	95	220	625	1,046	1,767	2,978	3,164	1
2003 1/.....	10,163	24	8	12	32	95	257	663	1,099	1,833	3,090	3,050	-
2002.....	9,983	19	4	9	32	87	258	632	1,037	1,912	3,069	2,924	-
2001.....	9,428	17	7	9	24	75	240	613	1,003	1,861	2,975	2,602	2
2000.....	9,101	21	5	13	31	60	222	560	1,039	1,818	2,828	2,504	-
1999.....	9,096	22	4	4	20	74	230	545	1,033	1,851	2,806	2,507	-
Asian or Pacific Islander 3/, male													
2007 1/.....	5,724	10	2	7	15	69	246	497	838	1,105	1,524	1,410	1
2006 1/.....	5,827	8	2	5	18	75	192	522	880	1,129	1,558	1,438	-
2005 1/.....	5,703	9	6	5	25	71	214	523	827	1,160	1,517	1,345	1
2004 1/.....	5,400	11	1	2	20	78	171	490	737	1,017	1,525	1,347	1
2003 1/.....	5,540	11	4	8	15	68	188	522	758	1,105	1,584	1,277	-
2002.....	5,523	7	2	5	23	67	208	481	710	1,104	1,630	1,286	-
2001.....	5,165	5	4	4	16	56	188	453	698	1,086	1,554	1,100	1
2000.....	5,175	13	3	7	24	43	174	431	729	1,117	1,513	1,121	-
1999.....	5,154	12	1	3	14	52	187	408	722	1,081	1,495	1,179	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

Asian or Pacific Islander 3/, female

2007 1/.....	4,850	11	9	4	3	24	58	130	347	699	1,504	2,061	-
2006 1/.....	4,846	5	5	8	6	21	49	167	340	747	1,506	1,992	-
2005 1/.....	4,774	9	6	9	11	26	64	160	374	739	1,517	1,859	-
2004 1/.....	4,560	11	6	2	11	17	49	135	309	750	1,453	1,817	-
2003 1/.....	4,623	13	4	4	17	27	69	141	341	728	1,506	1,773	-
2002.....	4,460	12	2	4	9	20	50	151	327	808	1,439	1,638	-
2001.....	4,263	12	3	5	8	19	52	160	305	775	1,421	1,502	1
2000.....	3,926	8	2	6	7	17	48	129	310	701	1,315	1,383	-
1999.....	3,942	10	3	1	6	22	43	137	311	770	1,311	1,328	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

All races, both sexes

2007.....	3,201	2	1	1	12	28	70	177	337	529	1,055	989	-
2006.....	3,257	3	-	7	7	31	66	176	352	608	1,123	884	-
2005.....	3,365	1	1	4	9	27	82	191	354	613	1,114	969	-
2004.....	3,254	2	3	4	7	27	96	207	321	612	1,146	829	-
2003.....	3,624	1	1	6	11	25	99	224	367	693	1,233	964	-
2002.....	3,579	4	1	6	18	36	95	237	433	703	1,227	819	-
2001.....	3,489	3	1	5	11	34	130	232	361	744	1,188	780	-
2000.....	3,582	6	2	4	17	33	141	252	420	819	1,176	712	-
1999.....	3,676	6	1	5	13	42	124	245	411	854	1,227	748	-

All races, male

2007.....	1,026	-	1	-	6	14	27	78	133	178	334	255	-
2006.....	1,008	2	-	2	5	17	28	81	135	208	327	203	-
2005.....	1,044	1	-	2	7	14	38	81	141	210	326	224	-
2004.....	1,009	2	3	2	4	9	38	88	107	202	380	174	-
2003.....	1,153	1	-	4	7	14	44	78	146	241	384	234	-
2002.....	1,078	3	1	2	8	18	45	95	156	245	341	164	-
2001.....	1,031	3	1	2	7	15	54	87	135	231	330	166	-
2000.....	1,072	5	2	3	11	12	61	104	166	237	321	150	-
1999.....	1,042	2	1	2	10	17	49	90	133	290	325	123	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)													
All races, female													
2007.....	2,175	2	-	1	6	14	43	99	204	351	721	734	-
2006.....	2,249	1	-	5	2	14	38	95	217	400	796	681	-
2005.....	2,321	-	1	2	2	13	44	110	213	403	788	745	-
2004.....	2,245	-	-	2	3	18	58	119	214	410	766	655	-
2003.....	2,471	-	1	2	4	11	55	146	221	452	849	730	-
2002.....	2,501	1	-	4	10	18	50	142	277	458	886	655	-
2001.....	2,458	-	-	3	4	19	76	145	226	513	858	614	-
2000.....	2,510	1	-	1	6	21	80	148	254	582	855	562	-
1999.....	2,634	4	-	3	3	25	75	155	278	564	902	625	-
White, both sexes													
2007 1/.....	2,853	1	1	-	9	18	51	125	258	458	990	942	-
2006 1/.....	2,903	3	-	2	3	20	42	124	277	528	1,046	858	-
2005 1/.....	3,029	1	1	4	4	17	57	134	284	546	1,044	937	-
2004 1/.....	2,911	2	2	2	4	16	57	153	257	542	1,078	798	-
2003 1/.....	3,226	1	1	4	5	18	61	149	277	611	1,172	927	-
2002.....	3,204	4	1	3	10	24	68	180	348	630	1,152	784	-
2001.....	3,086	1	1	3	10	15	85	157	293	649	1,125	747	-
2000.....	3,229	5	2	4	10	25	96	197	350	734	1,129	677	-
1999.....	3,302	4	1	3	8	25	84	187	341	776	1,161	712	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)													
White, male													
2007 1/.....	907	-	1	-	5	9	20	58	103	158	307	246	-
2006 1/.....	887	2	-	-	3	11	22	59	105	180	309	196	-
2005 1/.....	926	1	-	2	2	11	28	60	116	188	305	213	-
2004 1/.....	898	2	2	1	3	2	23	65	83	184	364	169	-
2003 1/.....	1,023	1	-	3	4	9	26	56	107	216	375	226	-
2002.....	950	3	1	2	4	11	30	72	126	219	323	159	-
2001.....	910	1	1	1	7	9	40	64	115	202	313	157	-
2000.....	965	4	2	3	6	10	46	87	147	210	309	141	-
1999.....	933	1	1	-	5	12	35	72	111	270	306	120	-
White, female													
2007 1/.....	1,946	1	-	-	4	9	31	67	155	300	683	696	-
2006 1/.....	2,016	1	-	2	-	9	20	65	172	348	737	662	-
2005 1/.....	2,103	-	1	2	2	6	29	74	168	358	739	724	-
2004 1/.....	2,013	-	-	1	1	14	34	88	174	358	714	629	-
2003 1/.....	2,203	-	1	1	1	9	35	93	170	395	797	701	-
2002.....	2,254	1	-	1	6	13	38	108	222	411	829	625	-
2001.....	2,176	-	-	2	3	6	45	93	178	447	812	590	-
2000.....	2,264	1	-	1	4	15	50	110	203	524	820	536	-
1999.....	2,369	3	-	3	3	13	49	115	230	506	855	592	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

Black, both sexes

2007 1/.....	245	-	-	1	3	4	14	43	57	43	46	34	-
2006 1/.....	249	-	-	3	2	7	20	39	54	53	54	17	-
2005 1/.....	227	-	-	-	2	8	16	48	44	42	44	23	-
2004 1/.....	247	-	1	2	3	6	33	49	46	46	40	21	-
2003 1/.....	308	-	-	2	4	6	29	63	72	66	40	26	-
2002.....	252	-	-	2	5	9	20	44	63	49	42	18	-
2001.....	302	2	-	1	1	13	41	60	53	69	40	22	-
2000.....	265	1	-	-	7	8	35	43	50	58	36	27	-
1999.....	275	1	-	2	4	13	32	49	52	56	40	26	-

Black, male

2007 1/.....	84	-	-	-	1	2	4	16	24	11	19	7	-
2006 1/.....	87	-	-	1	1	4	6	16	23	19	14	3	-
2005 1/.....	81	-	-	-	2	2	8	18	17	13	15	6	-
2004 1/.....	84	-	1	1	1	4	14	20	19	13	8	3	-
2003 1/.....	110	-	-	1	2	4	15	20	33	21	8	6	-
2002.....	93	-	-	-	3	4	12	17	22	21	11	3	-
2001.....	86	2	-	-	-	3	13	20	17	17	8	6	-
2000.....	79	1	-	-	5	2	12	13	14	16	8	8	-
1999.....	88	1	-	2	4	3	12	18	17	16	14	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)													
Black, female													
2007 1/.....	161	-	-	1	2	2	10	27	33	32	27	27	-
2006 1/.....	162	-	-	2	1	3	14	23	31	34	40	14	-
2005 1/.....	146	-	-	-	-	6	8	30	27	29	29	17	-
2004 1/.....	163	-	-	1	2	2	19	29	27	33	32	18	-
2003 1/.....	198	-	-	1	2	2	14	43	39	45	32	20	-
2002.....	159	-	-	2	2	5	8	27	41	28	31	15	-
2001.....	216	-	-	1	1	10	28	40	36	52	32	16	-
2000.....	186	-	-	-	2	6	23	30	36	42	28	19	-
1999.....	187	-	-	-	-	10	20	31	35	40	26	25	-
American Indian 2/, both sexes													
2007 1/.....	17	1	-	-	-	1	2	2	3	7	1	-	-
2006 1/.....	20	-	-	1	2	-	-	3	6	2	6	-	-
2005 1/.....	20	-	-	-	-	-	4	3	3	3	5	2	-
2004 1/.....	15	-	-	-	-	1	1	2	3	2	5	1	-
2003 1/.....	11	-	-	-	-	-	1	1	3	4	1	1	-
2002.....	23	-	-	-	1	1	1	1	5	5	8	1	-
2001.....	21	-	-	-	-	-	1	4	4	7	4	1	-
2000.....	18	-	-	-	-	-	2	1	6	4	3	2	-
1999.....	19	-	-	-	-	2	1	1	5	6	3	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

American Indian 2/, male

2007 1/.....	7	-	-	-	-	-	2	2	1	2	-	-	-
2006 1/.....	6	-	-	1	1	-	-	2	2	-	-	-	-
2005 1/.....	7	-	-	-	-	-	1	1	1	2	1	1	-
2004 1/.....	6	-	-	-	-	-	1	2	1	-	2	-	-
2003 1/.....	3	-	-	-	-	-	-	-	2	1	-	-	-
2002.....	9	-	-	-	-	1	-	1	2	-	4	1	-
2001.....	11	-	-	-	-	-	1	2	1	3	3	1	-
2000.....	3	-	-	-	-	-	-	-	1	1	1	-	-
1999.....	2	-	-	-	-	1	1	-	-	-	-	-	-

American Indian 2/, female

2007 1/.....	10	1	-	-	-	1	-	-	2	5	1	-	-
2006 1/.....	14	-	-	-	1	-	-	1	4	2	6	-	-
2005 1/.....	13	-	-	-	-	-	3	2	2	1	4	1	-
2004 1/.....	9	-	-	-	-	1	-	-	2	2	3	1	-
2003 1/.....	8	-	-	-	-	-	1	1	1	3	1	1	-
2002.....	14	-	-	-	1	-	1	-	3	5	4	-	-
2001.....	10	-	-	-	-	-	-	2	3	4	1	-	-
2000.....	15	-	-	-	-	-	2	1	5	3	2	2	-
1999.....	17	-	-	-	-	1	-	1	5	6	3	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	86	-	-	-	-	5	3	7	19	21	18	13	-
2006 1/.....	85	-	-	1	-	4	4	10	15	25	17	9	-
2005 1/.....	89	-	-	-	3	2	5	6	23	22	21	7	-
2004 1/.....	81	-	-	-	-	4	5	3	15	22	23	9	-
2003 1/.....	79	-	-	-	2	1	8	11	15	12	20	10	-
2002.....	100	-	-	1	2	2	6	12	17	19	25	16	-
2001.....	80	-	-	1	-	6	3	11	11	19	19	10	-
2000.....	70	-	-	-	-	-	8	11	14	23	8	6	-
1999.....	80	1	-	-	1	2	7	8	13	16	23	9	-

Asian or Pacific Islander 3/, male

2007 1/.....	28	-	-	-	-	3	1	2	5	7	8	2	-
2006 1/.....	28	-	-	-	-	2	-	4	5	9	4	4	-
2005 1/.....	30	-	-	-	3	1	1	2	7	7	5	4	-
2004 1/.....	21	-	-	-	-	3	-	1	4	5	6	2	-
2003 1/.....	17	-	-	-	1	1	3	2	4	3	1	2	-
2002.....	26	-	-	-	1	2	3	5	6	5	3	1	-
2001.....	24	-	-	1	-	3	-	1	2	9	6	2	-
2000.....	25	-	-	-	-	-	3	4	4	10	3	1	-
1999.....	19	-	-	-	1	1	1	-	5	4	5	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

Asian or Pacific Islander 3/, female

2007 1/.....	58	-	-	-	-	2	2	5	14	14	10	11	-
2006 1/.....	57	-	-	1	-	2	4	6	10	16	13	5	-
2005 1/.....	59	-	-	-	-	1	4	4	16	15	16	3	-
2004 1/.....	60	-	-	-	-	1	5	2	11	17	17	7	-
2003 1/.....	62	-	-	-	1	-	5	9	11	9	19	8	-
2002.....	74	-	-	1	1	-	3	7	11	14	22	15	-
2001.....	56	-	-	-	-	3	3	10	9	10	13	8	-
2000.....	45	-	-	-	-	-	5	7	10	13	5	5	-
1999.....	61	1	-	-	-	1	6	8	8	12	18	7	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart disease (I11)													
All races, both sexes													
2007.....	30,780	1	1	-	44	338	1,372	3,604	4,487	4,009	6,324	10,598	2
2006.....	29,788	1	-	2	52	288	1,358	3,586	4,130	3,860	6,232	10,277	2
2005.....	29,282	2	-	3	36	328	1,368	3,293	3,954	3,806	6,370	10,119	3
2004.....	28,585	-	-	2	44	302	1,317	3,265	3,885	4,001	6,398	9,368	3
2003.....	28,345	-	1	-	53	305	1,306	3,230	3,651	4,045	6,565	9,189	-
2002.....	26,551	-	-	2	30	297	1,213	2,882	3,320	3,895	6,321	8,587	4
2001.....	24,689	-	-	1	24	246	1,091	2,464	2,769	3,619	6,119	8,345	11
2000.....	23,761	-	1	2	32	213	981	2,104	2,672	3,605	5,962	8,186	3
1999.....	22,702	1	-	-	33	210	964	1,930	2,537	3,517	5,830	7,679	1
All races, male													
2007.....	14,249	1	1	-	34	237	953	2,473	3,009	2,266	2,673	2,600	2
2006.....	13,677	1	-	1	42	211	925	2,486	2,737	2,226	2,631	2,415	2
2005.....	13,219	1	-	2	24	222	958	2,278	2,649	2,147	2,570	2,365	3
2004.....	12,929	-	-	-	32	223	896	2,192	2,584	2,303	2,504	2,192	3
2003.....	12,536	-	1	-	40	208	877	2,157	2,308	2,264	2,627	2,054	-
2002.....	11,660	-	-	1	26	215	849	1,938	2,159	2,088	2,509	1,871	4
2001.....	10,451	-	-	-	17	169	740	1,646	1,749	1,961	2,414	1,746	9
2000.....	9,876	-	-	1	24	139	643	1,436	1,684	1,971	2,269	1,707	2
1999.....	9,316	-	-	-	25	149	660	1,259	1,560	1,816	2,252	1,594	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart disease (I11)													
All races, female													
2007.....	16,531	-	-	-	10	101	419	1,131	1,478	1,743	3,651	7,998	-
2006.....	16,111	-	-	1	10	77	433	1,100	1,393	1,634	3,601	7,862	-
2005.....	16,063	1	-	1	12	106	410	1,015	1,305	1,659	3,800	7,754	-
2004.....	15,656	-	-	2	12	79	421	1,073	1,301	1,698	3,894	7,176	-
2003.....	15,809	-	-	-	13	97	429	1,073	1,343	1,781	3,938	7,135	-
2002.....	14,891	-	-	1	4	82	364	944	1,161	1,807	3,812	6,716	-
2001.....	14,238	-	-	1	7	77	351	818	1,020	1,658	3,705	6,599	2
2000.....	13,885	-	1	1	8	74	338	668	988	1,634	3,693	6,479	1
1999.....	13,386	1	-	-	8	61	304	671	977	1,701	3,578	6,085	-
White, both sexes													
2007 1/.....	22,938	1	1	-	16	164	771	2,256	2,908	2,773	4,921	9,126	1
2006 1/.....	22,220	-	-	-	34	140	814	2,183	2,697	2,577	4,877	8,896	2
2005 1/.....	21,686	1	-	2	12	171	774	1,936	2,609	2,598	4,922	8,658	3
2004 1/.....	21,156	-	-	1	23	144	757	1,915	2,475	2,801	4,991	8,046	3
2003 1/.....	20,904	-	-	-	27	157	734	1,828	2,335	2,753	5,181	7,889	-
2002.....	19,530	-	-	1	13	141	680	1,704	2,076	2,659	4,894	7,361	1
2001.....	18,232	-	-	-	10	125	586	1,382	1,713	2,434	4,805	7,171	6
2000.....	17,425	-	-	2	15	95	542	1,168	1,588	2,411	4,626	6,977	1
1999.....	16,646	1	-	-	12	109	491	1,059	1,529	2,326	4,517	6,602	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart disease (I11)													
White, male													
2007 1/.....	10,209	1	1	-	15	118	567	1,626	2,013	1,586	2,068	2,213	1
2006 1/.....	9,755	-	-	-	28	108	600	1,618	1,838	1,488	2,028	2,045	2
2005 1/.....	9,408	1	-	2	8	130	577	1,420	1,801	1,486	2,002	1,978	3
2004 1/.....	9,218	-	-	-	18	120	557	1,383	1,677	1,638	1,966	1,856	3
2003 1/.....	8,948	-	-	-	21	121	556	1,334	1,554	1,560	2,078	1,724	-
2002.....	8,234	-	-	1	12	108	511	1,229	1,401	1,452	1,950	1,569	1
2001.....	7,395	-	-	-	7	96	441	986	1,144	1,373	1,888	1,455	5
2000.....	6,927	-	-	1	12	69	385	863	1,049	1,365	1,748	1,434	1
1999.....	6,499	-	-	-	9	91	370	749	987	1,223	1,735	1,335	-
White, female													
2007 1/.....	12,729	-	-	-	1	46	204	630	895	1,187	2,853	6,913	-
2006 1/.....	12,465	-	-	-	6	32	214	565	859	1,089	2,849	6,851	-
2005 1/.....	12,278	-	-	-	4	41	197	516	808	1,112	2,920	6,680	-
2004 1/.....	11,938	-	-	1	5	24	200	532	798	1,163	3,025	6,190	-
2003 1/.....	11,956	-	-	-	6	36	178	494	781	1,193	3,103	6,165	-
2002.....	11,296	-	-	-	1	33	169	475	675	1,207	2,944	5,792	-
2001.....	10,837	-	-	-	3	29	145	396	569	1,061	2,917	5,716	1
2000.....	10,498	-	-	1	3	26	157	305	539	1,046	2,878	5,543	-
1999.....	10,147	1	-	-	3	18	121	310	542	1,103	2,782	5,267	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart disease (I11)													
Black, both sexes													
2007 1/.....	7,108	-	-	-	28	162	564	1,270	1,467	1,132	1,235	1,249	1
2006 1/.....	6,845	1	-	2	17	139	507	1,315	1,346	1,171	1,192	1,155	-
2005 1/.....	6,923	1	-	1	23	140	545	1,274	1,261	1,115	1,298	1,265	-
2004 1/.....	6,796	-	-	1	21	148	530	1,292	1,308	1,119	1,261	1,116	-
2003 1/.....	6,843	-	1	-	25	137	544	1,339	1,234	1,184	1,240	1,139	-
2002.....	6,480	-	-	1	15	150	500	1,130	1,173	1,147	1,283	1,079	2
2001.....	5,970	-	-	1	14	115	479	1,030	993	1,095	1,176	1,062	5
2000.....	5,886	-	1	-	17	114	424	895	1,031	1,106	1,209	1,087	2
1999.....	5,619	-	-	-	20	98	450	848	945	1,105	1,195	957	1
Black, male													
2007 1/.....	3,669	-	-	-	19	109	363	790	921	622	530	314	1
2006 1/.....	3,559	1	-	1	14	96	294	808	842	667	531	305	-
2005 1/.....	3,459	-	-	-	15	78	345	796	794	600	507	324	-
2004 1/.....	3,391	-	-	-	14	93	316	766	835	625	473	269	-
2003 1/.....	3,298	-	1	-	19	79	302	776	703	656	486	276	-
2002.....	3,146	-	-	-	12	101	315	673	716	586	492	249	2
2001.....	2,813	-	-	-	10	68	280	625	563	546	466	251	4
2000.....	2,722	-	-	-	12	66	248	543	599	556	460	237	1
1999.....	2,606	-	-	-	15	55	270	493	537	547	463	225	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart disease (I11)													
Black, female													
2007 1/.....	3,439	-	-	-	9	53	201	480	546	510	705	935	-
2006 1/.....	3,286	-	-	1	3	43	213	507	504	504	661	850	-
2005 1/.....	3,464	1	-	1	8	62	200	478	467	515	791	941	-
2004 1/.....	3,405	-	-	1	7	55	214	526	473	494	788	847	-
2003 1/.....	3,545	-	-	-	6	58	242	563	531	528	754	863	-
2002.....	3,334	-	-	1	3	49	185	457	457	561	791	830	-
2001.....	3,157	-	-	1	4	47	199	405	430	549	710	811	1
2000.....	3,164	-	1	-	5	48	176	352	432	550	749	850	1
1999.....	3,013	-	-	-	5	43	180	355	408	558	732	732	-
American Indian 2/, both sexes													
2007 1/.....	135	-	-	-	-	5	17	30	26	20	20	17	-
2006 1/.....	133	-	-	-	-	4	15	33	20	23	23	15	-
2005 1/.....	127	-	-	-	-	8	13	22	25	19	19	21	-
2004 1/.....	109	-	-	-	-	3	8	14	31	12	20	21	-
2003 1/.....	112	-	-	-	-	6	8	20	25	18	14	21	-
2002.....	68	-	-	-	2	-	11	7	10	13	13	11	1
2001.....	64	-	-	-	-	2	6	8	10	10	16	12	-
2000.....	59	-	-	-	-	-	-	11	11	11	16	10	-
1999.....	56	-	-	-	-	-	6	2	14	10	15	9	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Hypertensive heart disease (I11)

American Indian 2/, male

2007 1/.....	79	-	-	-	-	5	10	21	16	14	8	5	-
2006 1/.....	80	-	-	-	-	2	14	20	13	16	10	5	-
2005 1/.....	68	-	-	-	-	5	8	16	17	9	7	6	-
2004 1/.....	59	-	-	-	-	3	5	8	22	7	9	5	-
2003 1/.....	54	-	-	-	-	4	5	11	17	8	6	3	-
2002.....	32	-	-	-	2	-	6	6	5	4	7	1	1
2001.....	29	-	-	-	-	1	3	5	6	7	6	1	-
2000.....	32	-	-	-	-	-	-	8	5	8	8	3	-
1999.....	26	-	-	-	-	-	5	2	5	7	5	2	-

American Indian 2/, female

2007 1/.....	56	-	-	-	-	-	7	9	10	6	12	12	-
2006 1/.....	53	-	-	-	-	2	1	13	7	7	13	10	-
2005 1/.....	59	-	-	-	-	3	5	6	8	10	12	15	-
2004 1/.....	50	-	-	-	-	-	3	6	9	5	11	16	-
2003 1/.....	58	-	-	-	-	2	3	9	8	10	8	18	-
2002.....	36	-	-	-	-	-	5	1	5	9	6	10	-
2001.....	35	-	-	-	-	1	3	3	4	3	10	11	-
2000.....	27	-	-	-	-	-	-	3	6	3	8	7	-
1999.....	30	-	-	-	-	-	1	-	9	3	10	7	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart disease (I11)													
Asian or Pacific Islander 3/, both sexes													
2007 1/.....	599	-	-	-	-	7	20	48	86	84	148	206	-
2006 1/.....	590	-	-	-	1	5	22	55	67	89	140	211	-
2005 1/.....	546	-	-	-	1	9	36	61	59	74	131	175	-
2004 1/.....	524	-	-	-	-	7	22	44	71	69	126	185	-
2003 1/.....	486	-	-	-	1	5	20	43	57	90	130	140	-
2002.....	473	-	-	-	-	6	22	41	61	76	131	136	-
2001.....	423	-	-	-	-	4	20	44	53	80	122	100	-
2000.....	391	-	-	-	-	4	15	30	42	77	111	112	-
1999.....	381	-	-	-	1	3	17	21	49	76	103	111	-
Asian or Pacific Islander 3/, male													
2007 1/.....	292	-	-	-	-	5	13	36	59	44	67	68	-
2006 1/.....	283	-	-	-	-	5	17	40	44	55	62	60	-
2005 1/.....	284	-	-	-	1	9	28	46	37	52	54	57	-
2004 1/.....	261	-	-	-	-	7	18	35	50	33	56	62	-
2003 1/.....	236	-	-	-	-	4	14	36	34	40	57	51	-
2002.....	248	-	-	-	-	6	17	30	37	46	60	52	-
2001.....	214	-	-	-	-	4	16	30	36	35	54	39	-
2000.....	195	-	-	-	-	4	10	22	31	42	53	33	-
1999.....	185	-	-	-	1	3	15	15	31	39	49	32	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Hypertensive heart disease (I11)

Asian or Pacific Islander 3/, female

2007 1/.....	307	-	-	-	-	2	7	12	27	40	81	138	-
2006 1/.....	307	-	-	-	1	-	5	15	23	34	78	151	-
2005 1/.....	262	-	-	-	-	-	8	15	22	22	77	118	-
2004 1/.....	263	-	-	-	-	-	4	9	21	36	70	123	-
2003 1/.....	250	-	-	-	1	1	6	7	23	50	73	89	-
2002.....	225	-	-	-	-	-	5	11	24	30	71	84	-
2001.....	209	-	-	-	-	-	4	14	17	45	68	61	-
2000.....	196	-	-	-	-	-	5	8	11	35	58	79	-
1999.....	196	-	-	-	-	-	2	6	18	37	54	79	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart and renal disease (I13)													
All races, both sexes													
2007.....	2,987	-	-	-	5	32	77	178	293	379	792	1,231	-
2006.....	2,918	-	-	-	6	31	81	179	259	359	806	1,197	-
2005.....	3,172	-	-	-	5	33	83	189	240	426	908	1,288	-
2004.....	3,046	-	-	1	7	23	96	209	271	379	854	1,206	-
2003.....	3,108	-	1	-	5	28	93	201	267	401	903	1,209	-
2002.....	2,895	-	-	-	5	28	77	165	273	402	866	1,079	-
2001.....	2,826	1	-	-	7	28	99	146	247	417	855	1,026	-
2000.....	2,785	-	-	1	6	28	83	147	208	417	863	1,032	-
1999.....	3,327	-	-	2	6	21	114	156	262	470	1,087	1,209	-
All races, male													
2007.....	1,318	-	-	-	4	17	56	120	178	191	344	408	-
2006.....	1,290	-	-	-	3	24	57	101	162	177	365	401	-
2005.....	1,369	-	-	-	4	20	52	125	142	209	378	439	-
2004.....	1,301	-	-	-	4	14	69	123	158	189	377	367	-
2003.....	1,326	-	1	-	4	13	62	130	159	189	382	386	-
2002.....	1,205	-	-	-	3	19	44	105	158	198	357	321	-
2001.....	1,169	1	-	-	4	17	63	99	126	206	339	314	-
2000.....	1,163	-	-	-	4	18	49	87	108	206	350	341	-
1999.....	1,390	-	-	1	3	16	67	100	146	223	460	374	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart and renal disease (I13)													
All races, female													
2007.....	1,669	-	-	-	1	15	21	58	115	188	448	823	-
2006.....	1,628	-	-	-	3	7	24	78	97	182	441	796	-
2005.....	1,803	-	-	-	1	13	31	64	98	217	530	849	-
2004.....	1,745	-	-	1	3	9	27	86	113	190	477	839	-
2003.....	1,782	-	-	-	1	15	31	71	108	212	521	823	-
2002.....	1,690	-	-	-	2	9	33	60	115	204	509	758	-
2001.....	1,657	-	-	-	3	11	36	47	121	211	516	712	-
2000.....	1,622	-	-	1	2	10	34	60	100	211	513	691	-
1999.....	1,937	-	-	1	3	5	47	56	116	247	627	835	-
White, both sexes													
2007 1/.....	2,064	-	-	-	1	12	20	68	132	213	582	1,036	-
2006 1/.....	1,985	-	-	-	-	13	27	63	111	190	587	994	-
2005 1/.....	2,186	-	-	-	2	7	28	61	94	240	658	1,096	-
2004 1/.....	2,071	-	-	1	2	8	27	60	119	212	635	1,007	-
2003 1/.....	2,103	-	1	-	2	7	21	53	111	229	685	994	-
2002.....	1,925	-	-	-	2	11	21	40	107	218	655	871	-
2001.....	1,922	-	-	-	3	7	25	52	116	239	642	838	-
2000.....	1,904	-	-	-	1	8	37	54	90	245	624	845	-
1999.....	2,235	-	-	1	-	7	28	52	110	249	806	982	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart and renal disease (I13)													
White, male													
2007 1/.....	864	-	-	-	1	6	14	52	75	111	261	344	-
2006 1/.....	836	-	-	-	-	11	22	35	70	88	268	342	-
2005 1/.....	886	-	-	-	2	4	19	45	57	116	275	368	-
2004 1/.....	846	-	-	-	2	7	17	36	74	113	285	312	-
2003 1/.....	865	-	1	-	1	2	14	37	68	118	307	317	-
2002.....	760	-	-	-	1	6	15	24	65	114	276	259	-
2001.....	762	-	-	-	3	3	17	32	56	118	266	267	-
2000.....	775	-	-	-	1	6	22	31	46	126	273	270	-
1999.....	906	-	-	1	-	6	17	30	60	123	358	311	-
White, female													
2007 1/.....	1,200	-	-	-	-	6	6	16	57	102	321	692	-
2006 1/.....	1,149	-	-	-	-	2	5	28	41	102	319	652	-
2005 1/.....	1,300	-	-	-	-	3	9	16	37	124	383	728	-
2004 1/.....	1,225	-	-	1	-	1	10	24	45	99	350	695	-
2003 1/.....	1,238	-	-	-	1	5	7	16	43	111	378	677	-
2002.....	1,165	-	-	-	1	5	6	16	42	104	379	612	-
2001.....	1,160	-	-	-	-	4	8	20	60	121	376	571	-
2000.....	1,129	-	-	-	-	2	15	23	44	119	351	575	-
1999.....	1,329	-	-	-	-	1	11	22	50	126	448	671	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart and renal disease (I13)													
Black, both sexes													
2007 1/.....	825	-	-	-	4	18	56	105	153	152	179	158	-
2006 1/.....	824	-	-	-	4	16	51	110	140	153	189	161	-
2005 1/.....	889	-	-	-	3	25	54	124	138	169	216	160	-
2004 1/.....	883	-	-	-	5	15	67	140	143	147	195	171	-
2003 1/.....	913	-	-	-	3	20	71	142	149	154	192	182	-
2002.....	903	-	-	-	2	17	54	125	160	167	189	189	-
2001.....	832	1	-	-	4	21	68	93	126	166	190	163	-
2000.....	809	-	-	1	5	19	46	91	113	164	209	161	-
1999.....	1,000	-	-	1	6	13	85	97	144	205	244	205	-
Black, male													
2007 1/.....	412	-	-	-	3	10	42	63	99	73	72	50	-
2006 1/.....	404	-	-	-	2	11	33	63	87	79	85	44	-
2005 1/.....	432	-	-	-	2	16	33	78	82	82	85	54	-
2004 1/.....	407	-	-	-	2	7	51	80	79	63	83	42	-
2003 1/.....	413	-	-	-	3	10	47	88	87	63	67	48	-
2002.....	405	-	-	-	1	13	27	81	88	71	67	57	-
2001.....	376	1	-	-	1	14	43	66	67	82	63	39	-
2000.....	356	-	-	-	3	12	27	55	58	77	69	55	-
1999.....	434	-	-	-	3	9	49	65	81	92	84	51	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Hypertensive heart and renal disease (I13)

Black, female

2007 1/.....	413	-	-	-	1	8	14	42	54	79	107	108	-
2006 1/.....	420	-	-	-	2	5	18	47	53	74	104	117	-
2005 1/.....	457	-	-	-	1	9	21	46	56	87	131	106	-
2004 1/.....	476	-	-	-	3	8	16	60	64	84	112	129	-
2003 1/.....	500	-	-	-	-	10	24	54	62	91	125	134	-
2002.....	498	-	-	-	1	4	27	44	72	96	122	132	-
2001.....	456	-	-	-	3	7	25	27	59	84	127	124	-
2000.....	453	-	-	1	2	7	19	36	55	87	140	106	-
1999.....	566	-	-	1	3	4	36	32	63	113	160	154	-

American Indian 2/, both sexes

2007 1/.....	8	-	-	-	-	-	-	1	1	-	3	3	-
2006 1/.....	17	-	-	-	-	1	1	2	2	3	3	5	-
2005 1/.....	12	-	-	-	-	1	-	1	2	2	3	3	-
2004 1/.....	13	-	-	-	-	-	-	1	3	2	5	2	-
2003 1/.....	8	-	-	-	-	1	-	-	2	1	2	2	-
2002.....	5	-	-	-	-	-	-	-	-	1	1	3	-
2001.....	8	-	-	-	-	-	2	-	2	-	2	2	-
2000.....	9	-	-	-	-	-	-	1	1	1	2	4	-
1999.....	14	-	-	-	-	-	-	1	1	6	4	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Hypertensive heart and renal disease (I13)

American Indian 2/, male

2007 1/.....	4	-	-	-	-	-	1	1	-	1	1	-
2006 1/.....	8	-	-	-	-	1	-	1	3	1	2	-
2005 1/.....	7	-	-	-	-	-	-	1	2	2	2	-
2004 1/.....	7	-	-	-	-	-	-	1	2	2	2	-
2003 1/.....	6	-	-	-	-	1	-	1	1	1	2	-
2002.....	2	-	-	-	-	-	-	-	1	-	1	-
2001.....	5	-	-	-	-	-	2	2	-	-	1	-
2000.....	3	-	-	-	-	-	-	1	-	1	1	-
1999.....	8	-	-	-	-	-	1	1	3	2	1	-

American Indian 2/, female

2007 1/.....	4	-	-	-	-	-	-	-	-	2	2	-
2006 1/.....	9	-	-	-	-	-	1	2	1	2	3	-
2005 1/.....	5	-	-	-	-	1	-	1	1	1	1	-
2004 1/.....	6	-	-	-	-	-	-	1	2	3	-	-
2003 1/.....	2	-	-	-	-	-	-	-	1	-	1	-
2002.....	3	-	-	-	-	-	-	-	-	1	2	-
2001.....	3	-	-	-	-	-	-	-	-	2	1	-
2000.....	6	-	-	-	-	-	-	1	-	1	3	-
1999.....	6	-	-	-	-	-	-	-	3	2	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart and renal disease (I13)													
Asian or Pacific Islander 3/, both sexes													
2007 1/.....	90	-	-	-	-	2	1	4	7	14	28	34	-
2006 1/.....	92	-	-	-	2	1	2	4	6	13	27	37	-
2005 1/.....	85	-	-	-	-	-	1	3	6	15	31	29	-
2004 1/.....	79	-	-	-	-	-	2	8	6	18	19	26	-
2003 1/.....	84	-	-	-	-	-	1	6	5	17	24	31	-
2002.....	62	-	-	-	1	-	2	-	6	16	21	16	-
2001.....	64	-	-	-	-	-	4	1	3	12	21	23	-
2000.....	63	-	-	-	-	1	-	1	4	7	28	22	-
1999.....	78	-	-	-	-	1	1	6	7	10	33	20	-
Asian or Pacific Islander 3/, male													
2007 1/.....	38	-	-	-	-	1	-	4	3	7	10	13	-
2006 1/.....	42	-	-	-	1	1	2	3	4	7	11	13	-
2005 1/.....	44	-	-	-	-	-	-	2	2	9	16	15	-
2004 1/.....	41	-	-	-	-	-	1	7	4	11	7	11	-
2003 1/.....	42	-	-	-	-	-	1	5	3	7	7	19	-
2002.....	38	-	-	-	1	-	2	-	5	12	14	4	-
2001.....	26	-	-	-	-	-	1	1	1	6	10	7	-
2000.....	29	-	-	-	-	-	-	1	3	3	7	15	-
1999.....	42	-	-	-	-	1	1	4	4	5	16	11	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Hypertensive heart and renal disease (I13)

Asian or Pacific Islander 3/, female

2007 1/.....	52	-	-	-	-	1	1	-	4	7	18	21	-
2006 1/.....	50	-	-	-	1	-	-	1	2	6	16	24	-
2005 1/.....	41	-	-	-	-	-	1	1	4	6	15	14	-
2004 1/.....	38	-	-	-	-	-	1	1	2	7	12	15	-
2003 1/.....	42	-	-	-	-	-	-	1	2	10	17	12	-
2002.....	24	-	-	-	-	-	-	-	1	4	7	12	-
2001.....	38	-	-	-	-	-	3	-	2	6	11	16	-
2000.....	34	-	-	-	-	1	-	-	1	4	21	7	-
1999.....	36	-	-	-	-	-	-	2	3	5	17	9	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Ischemic heart diseases (I20-I25)													
All races, both sexes													
2007.....	406,351	24	8	21	151	1,048	6,219	24,390	46,164	63,027	116,152	149,126	21
2006.....	425,425	22	7	13	138	1,054	6,699	24,990	46,706	66,685	125,225	153,855	31
2005.....	445,687	13	7	14	151	1,014	6,860	25,310	46,799	70,121	134,435	160,935	28
2004.....	451,326	13	10	17	125	1,092	7,132	24,923	46,002	73,102	139,128	159,756	26
2003.....	480,028	22	12	24	160	1,082	7,547	25,361	47,377	78,880	149,370	170,164	29
2002.....	494,382	23	7	20	133	996	7,683	25,575	47,262	83,867	155,391	173,388	37
2001.....	502,189	26	9	20	163	1,116	7,534	25,073	46,574	87,463	159,854	174,309	48
2000.....	515,204	30	10	16	123	935	7,519	24,890	47,737	92,885	163,692	177,319	48
1999.....	529,659	27	5	22	137	1,014	7,751	24,775	48,377	98,840	169,296	179,376	39
All races, male													
2007.....	216,050	11	3	12	107	809	4,732	18,526	33,476	40,777	62,693	54,889	15
2006.....	224,510	15	5	7	106	829	5,075	18,972	33,593	42,848	67,275	55,763	22
2005.....	232,115	7	4	9	121	774	5,188	19,233	33,548	44,497	71,352	57,364	18
2004.....	233,538	8	1	9	96	822	5,366	18,986	32,868	46,273	73,591	55,495	23
2003.....	246,342	13	6	10	107	813	5,679	19,481	33,767	49,999	78,573	57,872	22
2002.....	252,760	15	4	13	92	741	5,840	19,551	33,656	53,136	81,239	58,446	27
2001.....	254,005	17	3	13	119	830	5,712	19,141	33,006	55,072	82,953	57,106	33
2000.....	260,574	17	7	9	81	704	5,787	19,149	33,795	58,554	84,747	57,690	34
1999.....	267,268	13	3	17	93	739	5,970	19,061	34,123	62,187	87,305	57,726	31

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Ischemic heart diseases (I20-I25)													
All races, female													
2007.....	190,301	13	5	9	44	239	1,487	5,864	12,688	22,250	53,459	94,237	6
2006.....	200,915	7	2	6	32	225	1,624	6,018	13,113	23,837	57,950	98,092	9
2005.....	213,572	6	3	5	30	240	1,672	6,077	13,251	25,624	63,083	103,571	10
2004.....	217,788	5	9	8	29	270	1,766	5,937	13,134	26,829	65,537	104,261	3
2003.....	233,686	9	6	14	53	269	1,868	5,880	13,610	28,881	70,797	112,292	7
2002.....	241,622	8	3	7	41	255	1,843	6,024	13,606	30,731	74,152	114,942	10
2001.....	248,184	9	6	7	44	286	1,822	5,932	13,568	32,391	76,901	117,203	15
2000.....	254,630	13	3	7	42	231	1,732	5,741	13,942	34,331	78,945	119,629	14
1999.....	262,391	14	2	5	44	275	1,781	5,714	14,254	36,653	81,991	121,650	8
White, both sexes													
2007 1/.....	354,481	19	4	18	98	743	4,807	19,266	37,468	52,574	102,896	136,574	14
2006 1/.....	371,445	15	5	10	106	770	5,209	19,610	37,909	55,721	110,990	141,074	26
2005 1/.....	390,421	7	5	12	106	738	5,325	19,853	37,912	59,065	119,496	147,883	19
2004 1/.....	395,705	9	7	11	72	784	5,610	19,564	37,527	61,507	124,112	146,485	17
2003 1/.....	421,482	16	6	15	112	796	5,886	19,896	38,649	66,539	133,245	156,299	23
2002.....	435,170	14	5	15	94	740	6,010	20,227	38,577	71,269	138,903	159,288	28
2001.....	442,536	19	6	13	104	820	5,842	19,867	37,929	74,390	143,195	160,321	30
2000.....	455,400	23	10	13	92	676	5,851	19,835	38,955	79,442	147,075	163,392	36
1999.....	469,149	18	4	14	85	753	5,988	19,813	39,666	84,853	152,630	165,298	27

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Ischemic heart diseases (I20-I25)													
White, male													
2007 1/.....	189,056	10	1	9	70	575	3,723	15,001	27,723	34,689	56,396	50,849	10
2006 1/.....	196,396	12	3	7	86	620	4,065	15,294	27,772	36,434	60,452	51,632	19
2005 1/.....	203,924	5	2	9	89	564	4,136	15,478	27,773	38,208	64,375	53,273	12
2004 1/.....	205,475	6	1	7	57	607	4,345	15,343	27,398	39,722	66,674	51,300	15
2003 1/.....	216,751	9	3	7	76	608	4,542	15,711	28,222	42,966	70,935	53,656	16
2002.....	223,262	10	2	10	66	573	4,696	15,943	28,094	46,128	73,652	54,067	21
2001.....	224,464	14	2	9	78	624	4,560	15,640	27,537	47,795	75,330	52,854	21
2000.....	230,951	14	7	7	61	529	4,662	15,753	28,221	51,202	77,100	53,368	27
1999.....	237,193	8	3	12	63	555	4,742	15,672	28,548	54,493	79,823	53,253	21
White, female													
2007 1/.....	165,425	9	3	9	28	168	1,084	4,265	9,745	17,885	46,500	85,725	4
2006 1/.....	175,049	3	2	3	20	150	1,144	4,316	10,137	19,287	50,538	89,442	7
2005 1/.....	186,497	2	3	3	17	174	1,189	4,375	10,139	20,857	55,121	94,610	7
2004 1/.....	190,230	3	6	4	15	177	1,265	4,221	10,129	21,785	57,438	95,185	2
2003 1/.....	204,731	7	3	8	36	188	1,344	4,185	10,427	23,573	62,310	102,643	7
2002.....	211,908	4	3	5	28	167	1,314	4,284	10,483	25,141	65,251	105,221	7
2001.....	218,072	5	4	4	26	196	1,282	4,227	10,392	26,595	67,865	107,467	9
2000.....	224,449	9	3	6	31	147	1,189	4,082	10,734	28,240	69,975	110,024	9
1999.....	231,956	10	1	2	22	198	1,246	4,141	11,118	30,360	72,807	112,045	6

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Ischemic heart diseases (I20-I25)													
Black, both sexes													
2007 1/.....	42,679	3	2	2	50	254	1,149	4,469	7,504	8,711	10,651	9,878	6
2006 1/.....	44,530	6	2	2	32	234	1,278	4,704	7,526	9,159	11,498	10,084	5
2005 1/.....	46,027	5	1	2	36	220	1,316	4,769	7,680	9,230	12,267	10,493	8
2004 1/.....	46,695	3	1	6	46	259	1,332	4,732	7,396	9,853	12,370	10,689	8
2003 1/.....	49,274	4	5	8	43	234	1,425	4,765	7,545	10,515	13,380	11,346	4
2002.....	50,174	8	2	5	31	213	1,460	4,724	7,607	10,735	13,719	11,661	9
2001.....	50,933	7	3	6	53	250	1,486	4,576	7,552	11,235	13,976	11,773	16
2000.....	51,265	4	-	2	24	232	1,473	4,469	7,685	11,619	13,966	11,779	12
1999.....	51,995	8	1	8	47	213	1,570	4,403	7,632	12,094	14,071	11,936	12
Black, male													
2007 1/.....	21,768	1	1	2	35	191	794	2,994	4,884	4,981	4,942	2,939	4
2006 1/.....	22,713	2	2	-	20	166	833	3,140	4,890	5,300	5,384	2,973	3
2005 1/.....	22,933	2	1	-	26	161	868	3,212	4,902	5,172	5,588	2,996	5
2004 1/.....	23,060	2	-	2	36	171	861	3,153	4,686	5,498	5,523	3,121	7
2003 1/.....	24,273	3	3	3	28	162	948	3,204	4,702	5,879	6,186	3,151	4
2002.....	24,322	4	2	3	18	133	959	3,100	4,793	5,891	6,096	3,317	6
2001.....	24,593	3	1	3	36	171	983	3,012	4,669	6,174	6,203	3,327	11
2000.....	24,625	2	-	2	14	152	963	2,924	4,788	6,205	6,199	3,369	7
1999.....	25,104	4	-	5	26	148	1,057	2,938	4,806	6,543	6,085	3,482	10

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Ischemic heart diseases (I20-I25)													
Black, female													
2007 1/.....	20,911	2	1	-	15	63	355	1,475	2,620	3,730	5,709	6,939	2
2006 1/.....	21,817	4	-	2	12	68	445	1,564	2,636	3,859	6,114	7,111	2
2005 1/.....	23,094	3	-	2	10	59	448	1,557	2,778	4,058	6,679	7,497	3
2004 1/.....	23,635	1	1	4	10	88	471	1,579	2,710	4,355	6,847	7,568	1
2003 1/.....	25,001	1	2	5	15	72	477	1,561	2,843	4,636	7,194	8,195	-
2002.....	25,852	4	-	2	13	80	501	1,624	2,814	4,844	7,623	8,344	3
2001.....	26,340	4	2	3	17	79	503	1,564	2,883	5,061	7,773	8,446	5
2000.....	26,640	2	-	-	10	80	510	1,545	2,897	5,414	7,767	8,410	5
1999.....	26,891	4	1	3	21	65	513	1,465	2,826	5,551	7,986	8,454	2
American Indian 2/, both sexes													
2007 1/.....	1,777	1	2	-	1	19	74	201	330	394	445	310	-
2006 1/.....	1,880	-	-	-	-	14	67	211	357	420	490	321	-
2005 1/.....	1,788	-	-	-	1	16	60	216	355	390	419	331	-
2004 1/.....	1,843	1	1	-	1	21	68	197	312	410	460	372	-
2003 1/.....	1,905	-	-	-	2	16	82	215	340	450	465	333	2
2002.....	1,777	1	-	-	4	16	59	173	323	426	465	310	-
2001.....	1,710	-	-	-	3	11	64	196	324	428	406	278	-
2000.....	1,740	1	-	-	4	11	73	176	302	414	438	321	-
1999.....	1,720	1	-	-	1	16	56	166	286	447	453	294	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Ischemic heart diseases (I20-I25)

American Indian 2/, male

2007 1/.....	1,048	-	1	-	1	13	49	153	232	247	238	114	-
2006 1/.....	1,082	-	-	-	-	11	49	164	241	256	248	113	-
2005 1/.....	1,028	-	-	-	-	13	48	154	250	225	220	118	-
2004 1/.....	1,058	-	-	-	1	18	51	136	217	259	246	130	-
2003 1/.....	1,133	-	-	-	1	11	63	163	229	281	248	135	2
2002.....	1,041	1	-	-	4	13	48	136	224	261	232	122	-
2001.....	995	-	-	-	3	8	46	140	233	259	202	104	-
2000.....	1,008	-	-	-	3	9	56	134	201	257	231	117	-
1999.....	977	1	-	-	1	8	45	125	191	275	234	97	-

American Indian 2/, female

2007 1/.....	729	1	1	-	-	6	25	48	98	147	207	196	-
2006 1/.....	798	-	-	-	-	3	18	47	116	164	242	208	-
2005 1/.....	760	-	-	-	1	3	12	62	105	165	199	213	-
2004 1/.....	785	1	1	-	-	3	17	61	95	151	214	242	-
2003 1/.....	772	-	-	-	1	5	19	52	111	169	217	198	-
2002.....	736	-	-	-	-	3	11	37	99	165	233	188	-
2001.....	715	-	-	-	-	3	18	56	91	169	204	174	-
2000.....	732	1	-	-	1	2	17	42	101	157	207	204	-
1999.....	743	-	-	-	-	8	11	41	95	172	219	197	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Ischemic heart diseases (I20-I25)													
Asian or Pacific Islander 3/, both sexes													
2007 1/.....	7,414	1	-	1	2	32	189	454	862	1,348	2,160	2,364	1
2006 1/.....	7,570	1	-	1	-	36	145	465	914	1,385	2,247	2,376	-
2005 1/.....	7,451	1	1	-	8	40	159	472	852	1,436	2,253	2,228	1
2004 1/.....	7,083	-	1	-	6	28	122	430	767	1,332	2,186	2,210	1
2003 1/.....	7,367	2	1	1	3	36	154	485	843	1,376	2,280	2,186	-
2002.....	7,261	-	-	-	4	27	154	451	755	1,437	2,304	2,129	-
2001.....	7,010	-	-	1	3	35	142	434	769	1,410	2,277	1,937	2
2000.....	6,799	2	-	1	3	16	122	410	795	1,410	2,213	1,827	-
1999.....	6,795	-	-	-	4	32	137	393	793	1,446	2,142	1,848	-
Asian or Pacific Islander 3/, male													
2007 1/.....	4,178	-	-	1	1	30	166	378	637	860	1,117	987	1
2006 1/.....	4,319	1	-	-	-	32	128	374	690	858	1,191	1,045	-
2005 1/.....	4,230	-	1	-	6	36	136	389	623	892	1,169	977	1
2004 1/.....	3,945	-	-	-	2	26	109	354	567	794	1,148	944	1
2003 1/.....	4,185	1	-	-	2	32	126	403	614	873	1,204	930	-
2002.....	4,135	-	-	-	4	22	137	372	545	856	1,259	940	-
2001.....	3,953	-	-	1	2	27	123	349	567	844	1,218	821	1
2000.....	3,990	1	-	-	3	14	106	338	585	890	1,217	836	-
1999.....	3,994	-	-	-	3	28	126	326	578	876	1,163	894	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Ischemic heart diseases (I20-I25)

Asian or Pacific Islander 3/, female

2007 1/.....	3,236	1	-	-	1	2	23	76	225	488	1,043	1,377	-
2006 1/.....	3,251	-	-	1	-	4	17	91	224	527	1,056	1,331	-
2005 1/.....	3,221	1	-	-	2	4	23	83	229	544	1,084	1,251	-
2004 1/.....	3,138	-	1	-	4	2	13	76	200	538	1,038	1,266	-
2003 1/.....	3,182	1	1	1	1	4	28	82	229	503	1,076	1,256	-
2002.....	3,126	-	-	-	-	5	17	79	210	581	1,045	1,189	-
2001.....	3,057	-	-	-	1	8	19	85	202	566	1,059	1,116	1
2000.....	2,809	1	-	1	-	2	16	72	210	520	996	991	-
1999.....	2,801	-	-	-	1	4	11	67	215	570	979	954	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute myocardial infarction (I21-I22)													
All races, both sexes													
2007.....	132,968	10	4	11	54	400	2,402	9,467	17,835	23,441	37,629	41,711	4
2006.....	141,462	11	5	6	60	422	2,676	9,746	18,373	24,795	41,624	43,739	5
2005.....	151,004	11	2	7	68	392	2,734	10,070	18,553	26,674	45,449	47,041	3
2004.....	156,816	9	2	5	51	447	2,838	10,084	18,758	28,327	48,037	48,252	6
2003.....	170,564	9	6	12	73	413	3,094	10,420	19,767	30,923	53,108	52,734	5
2002.....	179,514	11	5	7	62	400	3,161	10,619	19,933	34,003	56,590	54,718	5
2001.....	184,757	15	2	7	64	475	3,176	10,817	19,830	35,380	59,254	55,727	10
2000.....	192,898	20	1	6	52	375	3,132	10,877	21,013	38,459	62,210	56,745	8
1999.....	199,454	13	2	11	60	434	3,249	10,823	21,406	41,441	64,711	57,297	7
All races, male													
2007.....	71,712	5	-	5	33	297	1,768	7,129	12,744	14,824	19,698	15,206	3
2006.....	76,089	8	4	3	42	315	2,011	7,278	13,031	15,767	21,819	15,807	4
2005.....	80,079	6	2	3	54	297	2,025	7,563	13,151	16,676	23,540	16,759	3
2004.....	82,909	7	-	4	39	322	2,113	7,632	13,342	17,658	24,893	16,895	4
2003.....	89,515	4	2	6	46	300	2,301	7,984	13,972	19,290	27,350	18,256	4
2002.....	93,830	7	2	2	41	292	2,347	7,995	13,966	21,264	28,919	18,991	4
2001.....	95,875	10	2	4	48	333	2,384	8,158	13,858	21,963	30,125	18,984	6
2000.....	100,306	10	1	4	30	269	2,387	8,265	14,638	23,823	31,631	19,243	5
1999.....	104,169	6	1	7	40	306	2,470	8,300	15,029	25,708	32,943	19,357	2

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute myocardial infarction (I21-I22)													
All races, female													
2007.....	61,256	5	4	6	21	103	634	2,338	5,091	8,617	17,931	26,505	1
2006.....	65,373	3	1	3	18	107	665	2,468	5,342	9,028	19,805	27,932	1
2005.....	70,925	5	-	4	14	95	709	2,507	5,402	9,998	21,909	30,282	-
2004.....	73,907	2	2	1	12	125	725	2,452	5,416	10,669	23,144	31,357	2
2003.....	81,049	5	4	6	27	113	793	2,436	5,795	11,633	25,758	34,478	1
2002.....	85,684	4	3	5	21	108	814	2,624	5,967	12,739	27,671	35,727	1
2001.....	88,882	5	-	3	16	142	792	2,659	5,972	13,417	29,129	36,743	4
2000.....	92,592	10	-	2	22	106	745	2,612	6,375	14,636	30,579	37,502	3
1999.....	95,285	7	1	4	20	128	779	2,523	6,377	15,733	31,768	37,940	5
White, both sexes													
2007 1/.....	115,900	9	2	10	36	284	1,906	7,694	14,880	19,845	33,310	37,922	2
2006 1/.....	123,549	10	3	4	49	316	2,140	7,879	15,341	21,065	36,950	39,788	4
2005 1/.....	132,364	6	-	5	50	278	2,160	8,172	15,493	22,852	40,417	42,929	2
2004 1/.....	137,643	6	2	3	27	337	2,269	8,125	15,742	24,124	43,019	43,985	4
2003 1/.....	149,915	7	3	9	53	310	2,452	8,359	16,569	26,495	47,440	48,214	4
2002.....	157,922	9	5	5	43	301	2,524	8,621	16,721	29,251	50,565	49,873	4
2001.....	162,943	12	2	2	42	355	2,513	8,805	16,626	30,469	53,186	50,925	6
2000.....	170,584	16	1	4	39	272	2,496	8,900	17,659	33,285	55,966	51,939	7
1999.....	177,075	9	2	6	39	325	2,596	8,953	18,084	36,106	58,482	52,468	5

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute myocardial infarction (I21-I22)													
White, male													
2007 1/.....	63,011	5	-	4	22	213	1,430	5,952	10,875	12,793	17,717	13,998	2
2006 1/.....	66,946	8	2	3	38	240	1,651	6,049	11,078	13,689	19,660	14,525	3
2005 1/.....	70,791	4	-	3	42	214	1,646	6,308	11,219	14,615	21,276	15,462	2
2004 1/.....	73,448	5	-	3	21	255	1,746	6,328	11,472	15,345	22,702	15,568	3
2003 1/.....	79,157	3	1	4	35	229	1,876	6,592	12,004	16,861	24,701	16,848	3
2002.....	83,288	6	2	2	30	229	1,921	6,689	12,025	18,752	26,199	17,430	3
2001.....	85,280	9	2	1	34	254	1,951	6,879	11,937	19,355	27,434	17,419	5
2000.....	89,383	9	1	2	24	205	1,969	7,003	12,586	21,089	28,832	17,659	4
1999.....	93,189	3	1	4	29	229	2,032	7,076	12,967	22,869	30,234	17,744	1
White, female													
2007 1/.....	52,889	4	2	6	14	71	476	1,742	4,005	7,052	15,593	23,924	-
2006 1/.....	56,603	2	1	1	11	76	489	1,830	4,263	7,376	17,290	25,263	1
2005 1/.....	61,573	2	-	2	8	64	514	1,864	4,274	8,237	19,141	27,467	-
2004 1/.....	64,195	1	2	-	6	82	523	1,797	4,270	8,779	20,317	28,417	1
2003 1/.....	70,758	4	2	5	18	81	576	1,767	4,565	9,634	22,739	31,366	1
2002.....	74,634	3	3	3	13	72	603	1,932	4,696	10,499	24,366	32,443	1
2001.....	77,663	3	-	1	8	101	562	1,926	4,689	11,114	25,752	33,506	1
2000.....	81,201	7	-	2	15	67	527	1,897	5,073	12,196	27,134	34,280	3
1999.....	83,886	6	1	2	10	96	564	1,877	5,117	13,237	28,248	34,724	4

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute myocardial infarction (I21-I22)													
Black, both sexes													
2007 1/.....	14,097	1	1	1	18	96	415	1,526	2,554	3,000	3,477	3,006	2
2006 1/.....	14,776	1	2	2	11	92	471	1,639	2,596	3,109	3,753	3,099	1
2005 1/.....	15,536	4	1	2	16	95	498	1,668	2,616	3,173	4,124	3,338	1
2004 1/.....	16,206	2	-	2	21	94	502	1,743	2,641	3,547	4,178	3,475	1
2003 1/.....	17,441	1	2	3	19	88	580	1,797	2,725	3,761	4,711	3,753	1
2002.....	18,322	2	-	2	15	89	561	1,765	2,824	4,030	4,988	4,045	1
2001.....	18,581	3	-	4	21	99	579	1,752	2,783	4,216	5,082	4,038	4
2000.....	19,112	2	-	2	11	94	569	1,737	2,935	4,447	5,236	4,078	1
1999.....	19,276	4	-	5	19	88	587	1,659	2,882	4,596	5,298	4,136	2
Black, male													
2007 1/.....	6,997	-	-	1	11	69	276	985	1,581	1,653	1,533	887	1
2006 1/.....	7,392	-	2	-	4	64	308	1,055	1,648	1,718	1,681	911	1
2005 1/.....	7,527	2	1	-	12	67	315	1,072	1,617	1,683	1,804	953	1
2004 1/.....	7,811	2	-	1	17	54	314	1,146	1,597	1,918	1,746	1,016	-
2003 1/.....	8,472	-	1	2	11	59	374	1,175	1,642	2,011	2,128	1,068	1
2002.....	8,680	1	-	-	7	55	362	1,118	1,666	2,103	2,173	1,194	1
2001.....	8,717	1	-	2	13	64	360	1,074	1,626	2,190	2,168	1,218	1
2000.....	9,045	1	-	2	5	56	362	1,075	1,748	2,295	2,270	1,230	1
1999.....	9,186	3	-	3	10	61	379	1,060	1,756	2,404	2,230	1,279	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute myocardial infarction (I21-I22)													
Black, female													
2007 1/.....	7,100	1	1	-	7	27	139	541	973	1,347	1,944	2,119	1
2006 1/.....	7,384	1	-	2	7	28	163	584	948	1,391	2,072	2,188	-
2005 1/.....	8,009	2	-	2	4	28	183	596	999	1,490	2,320	2,385	-
2004 1/.....	8,395	-	-	1	4	40	188	597	1,044	1,629	2,432	2,459	1
2003 1/.....	8,969	1	1	1	8	29	206	622	1,083	1,750	2,583	2,685	-
2002.....	9,642	1	-	2	8	34	199	647	1,158	1,927	2,815	2,851	-
2001.....	9,864	2	-	2	8	35	219	678	1,157	2,026	2,914	2,820	3
2000.....	10,067	1	-	-	6	38	207	662	1,187	2,152	2,966	2,848	-
1999.....	10,090	1	-	2	9	27	208	599	1,126	2,192	3,068	2,857	1
American Indian 2/, both sexes													
2007 1/.....	591	-	1	-	-	8	24	77	128	131	133	89	-
2006 1/.....	665	-	-	-	-	7	26	81	131	142	175	103	-
2005 1/.....	589	-	-	-	-	6	19	72	125	127	156	84	-
2004 1/.....	634	1	-	-	-	7	26	74	112	166	126	122	-
2003 1/.....	706	-	-	-	-	5	27	85	152	172	172	93	-
2002.....	657	-	-	-	2	3	21	73	132	165	166	95	-
2001.....	660	-	-	-	-	7	30	87	125	162	139	110	-
2000.....	656	1	-	-	1	5	30	72	113	163	166	105	-
1999.....	674	-	-	-	-	8	22	65	116	183	177	103	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute myocardial infarction (I21-I22)													
American Indian 2/, male													
2007 1/.....	359	-	-	-	-	5	14	55	92	79	75	39	-
2006 1/.....	371	-	-	-	-	5	17	65	86	76	87	35	-
2005 1/.....	339	-	-	-	-	4	17	54	84	67	77	36	-
2004 1/.....	359	-	-	-	-	6	18	50	76	109	65	35	-
2003 1/.....	430	-	-	-	-	3	22	70	101	105	90	39	-
2002.....	384	-	-	-	2	3	17	60	99	90	73	40	-
2001.....	404	-	-	-	-	5	26	65	89	96	73	50	-
2000.....	384	-	-	-	-	4	23	55	73	101	87	41	-
1999.....	382	-	-	-	-	4	20	47	77	113	84	37	-
American Indian 2/, female													
2007 1/.....	232	-	1	-	-	3	10	22	36	52	58	50	-
2006 1/.....	294	-	-	-	-	2	9	16	45	66	88	68	-
2005 1/.....	250	-	-	-	-	2	2	18	41	60	79	48	-
2004 1/.....	275	1	-	-	-	1	8	24	36	57	61	87	-
2003 1/.....	276	-	-	-	-	2	5	15	51	67	82	54	-
2002.....	273	-	-	-	-	-	4	13	33	75	93	55	-
2001.....	256	-	-	-	-	2	4	22	36	66	66	60	-
2000.....	272	1	-	-	1	1	7	17	40	62	79	64	-
1999.....	292	-	-	-	-	4	2	18	39	70	93	66	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute myocardial infarction (I21-I22)													
Asian or Pacific Islander 3/, both sexes													
2007 1/.....	2,380	-	-	-	-	12	57	170	273	465	709	694	-
2006 1/.....	2,472	-	-	-	-	7	39	147	305	479	746	749	-
2005 1/.....	2,515	1	1	-	2	13	57	158	319	522	752	690	-
2004 1/.....	2,333	-	-	-	3	9	41	142	263	490	714	670	1
2003 1/.....	2,502	1	1	-	1	10	35	179	321	495	785	674	-
2002.....	2,613	-	-	-	2	7	55	160	256	557	871	705	-
2001.....	2,573	-	-	1	1	14	54	173	296	533	847	654	-
2000.....	2,546	1	-	-	1	4	37	168	306	564	842	623	-
1999.....	2,429	-	-	-	2	13	44	146	324	556	754	590	-
Asian or Pacific Islander 3/, male													
2007 1/.....	1,345	-	-	-	-	10	48	137	196	299	373	282	-
2006 1/.....	1,380	-	-	-	-	6	35	109	219	284	391	336	-
2005 1/.....	1,422	-	1	-	-	12	47	129	231	311	383	308	-
2004 1/.....	1,291	-	-	-	1	7	35	108	197	286	380	276	1
2003 1/.....	1,456	1	-	-	-	9	29	147	225	313	431	301	-
2002.....	1,478	-	-	-	2	5	47	128	176	319	474	327	-
2001.....	1,474	-	-	1	1	10	47	140	206	322	450	297	-
2000.....	1,494	-	-	-	1	4	33	132	231	338	442	313	-
1999.....	1,412	-	-	-	1	12	39	117	229	322	395	297	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Acute myocardial infarction (I21-I22)

Asian or Pacific Islander 3/, female

2007 1/.....	1,035	-	-	-	-	2	9	33	77	166	336	412	-
2006 1/.....	1,092	-	-	-	-	1	4	38	86	195	355	413	-
2005 1/.....	1,093	1	-	-	2	1	10	29	88	211	369	382	-
2004 1/.....	1,042	-	-	-	2	2	6	34	66	204	334	394	-
2003 1/.....	1,046	-	1	-	1	1	6	32	96	182	354	373	-
2002.....	1,135	-	-	-	-	2	8	32	80	238	397	378	-
2001.....	1,099	-	-	-	-	4	7	33	90	211	397	357	-
2000.....	1,052	1	-	-	-	-	4	36	75	226	400	310	-
1999.....	1,017	-	-	-	1	1	5	29	95	234	359	293	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other acute ischemic heart diseases (I24)

All races, both sexes

2007.....	4,092	2	-	-	3	17	109	376	679	740	1,021	1,145	-
2006.....	3,932	2	1	-	7	21	92	354	596	676	1,041	1,139	3
2005.....	3,565	-	1	1	3	11	82	351	515	606	935	1,060	-
2004.....	2,462	1	2	1	5	10	80	217	359	399	670	717	1
2003.....	3,158	4	1	1	5	11	98	277	403	579	837	941	1
2002.....	3,407	2	-	1	4	18	77	345	460	631	898	971	-
2001.....	3,351	3	-	2	7	14	94	307	463	645	871	945	-
2000.....	3,363	1	1	-	8	16	92	282	484	644	906	928	1
1999.....	3,232	3	1	1	3	6	81	252	448	674	855	908	-

All races, male

2007.....	2,159	1	-	-	2	14	76	268	487	479	500	332	-
2006.....	2,016	1	-	-	6	17	54	254	400	402	511	368	3
2005.....	1,823	-	-	1	2	7	59	245	371	364	450	324	-
2004.....	1,301	-	-	-	4	6	59	151	251	254	329	246	1
2003.....	1,621	2	1	-	5	7	73	197	278	361	414	282	1
2002.....	1,767	1	-	1	1	12	50	250	320	390	444	298	-
2001.....	1,720	2	-	1	6	10	65	230	320	393	424	269	-
2000.....	1,749	1	1	-	6	13	71	217	341	398	446	254	1
1999.....	1,665	1	1	1	3	4	63	178	315	410	423	266	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other acute ischemic heart diseases (I24)													
All races, female													
2007.....	1,933	1	-	-	1	3	33	108	192	261	521	813	-
2006.....	1,916	1	1	-	1	4	38	100	196	274	530	771	-
2005.....	1,742	-	1	-	1	4	23	106	144	242	485	736	-
2004.....	1,161	1	2	1	1	4	21	66	108	145	341	471	-
2003.....	1,537	2	-	1	-	4	25	80	125	218	423	659	-
2002.....	1,640	1	-	-	3	6	27	95	140	241	454	673	-
2001.....	1,631	1	-	1	1	4	29	77	143	252	447	676	-
2000.....	1,614	-	-	-	2	3	21	65	143	246	460	674	-
1999.....	1,567	2	-	-	-	2	18	74	133	264	432	642	-
White, both sexes													
2007 1/.....	3,441	2	-	-	1	8	78	286	537	604	891	1,034	-
2006 1/.....	3,289	2	1	-	5	15	60	250	474	548	902	1,030	2
2005 1/.....	2,924	-	1	1	3	7	60	237	383	493	801	938	-
2004 1/.....	2,025	-	2	1	3	3	60	143	268	317	571	656	1
2003 1/.....	2,663	2	-	1	4	6	69	205	311	473	729	862	1
2002.....	2,907	1	-	1	3	12	55	261	367	524	796	887	-
2001.....	2,821	3	-	2	4	11	63	226	355	531	771	855	-
2000.....	2,839	1	1	-	6	12	68	214	367	530	790	850	-
1999.....	2,752	1	1	1	3	5	53	176	357	563	750	842	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other acute ischemic heart diseases (I24)													
White, male													
2007 1/.....	1,792	1	-	-	1	7	59	206	384	396	437	301	-
2006 1/.....	1,664	1	-	-	4	13	36	188	320	331	438	331	2
2005 1/.....	1,480	-	-	1	2	4	45	183	280	296	385	284	-
2004 1/.....	1,042	-	-	-	2	3	46	102	189	194	285	220	1
2003 1/.....	1,338	1	-	-	4	5	56	148	213	302	351	257	1
2002.....	1,494	1	-	1	1	8	42	187	259	323	396	276	-
2001.....	1,431	2	-	1	3	9	46	172	250	322	382	244	-
2000.....	1,453	1	1	-	4	9	54	168	266	325	392	233	-
1999.....	1,389	1	1	1	3	4	41	127	246	343	375	247	-
White, female													
2007 1/.....	1,649	1	-	-	-	1	19	80	153	208	454	733	-
2006 1/.....	1,625	1	1	-	1	2	24	62	154	217	464	699	-
2005 1/.....	1,444	-	1	-	1	3	15	54	103	197	416	654	-
2004 1/.....	983	-	2	1	1	-	14	41	79	123	286	436	-
2003 1/.....	1,325	1	-	1	-	1	13	57	98	171	378	605	-
2002.....	1,413	-	-	-	2	4	13	74	108	201	400	611	-
2001.....	1,390	1	-	1	1	2	17	54	105	209	389	611	-
2000.....	1,386	-	-	-	2	3	14	46	101	205	398	617	-
1999.....	1,363	-	-	-	-	1	12	49	111	220	375	595	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other acute ischemic heart diseases (I24)													
Black, both sexes													
2007 1/.....	539	-	-	-	2	7	24	75	117	113	114	87	-
2006 1/.....	534	-	-	-	2	6	28	86	109	108	114	80	1
2005 1/.....	558	-	-	-	-	4	19	101	115	98	119	102	-
2004 1/.....	395	1	-	-	1	7	18	69	86	71	88	54	-
2003 1/.....	436	1	1	-	1	3	27	62	82	94	97	68	-
2002.....	446	1	-	-	1	6	20	78	79	100	90	71	-
2001.....	454	-	-	-	3	2	29	69	98	95	80	78	-
2000.....	456	-	-	-	1	4	22	63	103	93	99	70	1
1999.....	431	2	-	-	-	1	26	73	85	93	97	54	-
Black, male													
2007 1/.....	303	-	-	-	1	5	15	52	84	69	53	24	-
2006 1/.....	291	-	-	-	2	4	15	54	73	60	56	26	1
2005 1/.....	296	-	-	-	-	3	12	56	79	58	58	30	-
2004 1/.....	235	-	-	-	1	3	12	44	59	51	42	23	-
2003 1/.....	256	1	1	-	1	1	16	43	59	54	57	23	-
2002.....	243	-	-	-	-	4	6	58	50	64	41	20	-
2001.....	249	-	-	-	3	-	19	50	62	59	35	21	-
2000.....	258	-	-	-	1	4	16	46	67	59	46	18	1
1999.....	248	-	-	-	-	-	20	49	65	56	41	17	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other acute ischemic heart diseases (I24)													
Black, female													
2007 1/.....	236	-	-	-	1	2	9	23	33	44	61	63	-
2006 1/.....	243	-	-	-	-	2	13	32	36	48	58	54	-
2005 1/.....	262	-	-	-	-	1	7	45	36	40	61	72	-
2004 1/.....	160	1	-	-	-	4	6	25	27	20	46	31	-
2003 1/.....	180	-	-	-	-	2	11	19	23	40	40	45	-
2002.....	203	1	-	-	1	2	14	20	29	36	49	51	-
2001.....	205	-	-	-	-	2	10	19	36	36	45	57	-
2000.....	198	-	-	-	-	-	6	17	36	34	53	52	-
1999.....	183	2	-	-	-	1	6	24	20	37	56	37	-
American Indian 2/, both sexes													
2007 1/.....	61	-	-	-	-	1	6	9	14	12	7	12	-
2006 1/.....	52	-	-	-	-	-	3	11	9	8	12	9	-
2005 1/.....	44	-	-	-	-	-	3	8	10	11	5	7	-
2004 1/.....	23	-	-	-	1	-	2	2	3	4	5	6	-
2003 1/.....	37	-	-	-	-	1	2	6	8	9	6	5	-
2002.....	36	-	-	-	-	-	2	5	8	5	8	8	-
2001.....	47	-	-	-	-	1	1	10	7	14	8	6	-
2000.....	43	-	-	-	1	-	2	5	9	15	8	3	-
1999.....	37	-	-	-	-	-	1	3	5	14	6	8	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other acute ischemic heart diseases (I24)

American Indian 2/, male

2007 1/.....	34	-	-	-	-	1	1	6	11	6	6	3	-
2006 1/.....	29	-	-	-	-	-	2	6	5	3	9	4	-
2005 1/.....	26	-	-	-	-	-	2	2	7	8	3	4	-
2004 1/.....	11	-	-	-	1	-	1	2	2	3	-	2	-
2003 1/.....	15	-	-	-	-	-	1	4	5	3	1	1	-
2002.....	21	-	-	-	-	-	2	4	7	2	5	1	-
2001.....	28	-	-	-	-	1	-	7	6	10	3	1	-
2000.....	26	-	-	-	1	-	1	3	5	10	4	2	-
1999.....	20	-	-	-	-	-	1	2	3	8	5	1	-

American Indian 2/, female

2007 1/.....	27	-	-	-	-	-	5	3	3	6	1	9	-
2006 1/.....	23	-	-	-	-	-	1	5	4	5	3	5	-
2005 1/.....	18	-	-	-	-	-	1	6	3	3	2	3	-
2004 1/.....	12	-	-	-	-	-	1	-	1	1	5	4	-
2003 1/.....	22	-	-	-	-	1	1	2	3	6	5	4	-
2002.....	15	-	-	-	-	-	-	1	1	3	3	7	-
2001.....	19	-	-	-	-	-	1	3	1	4	5	5	-
2000.....	17	-	-	-	-	-	1	2	4	5	4	1	-
1999.....	17	-	-	-	-	-	-	1	2	6	1	7	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other acute ischemic heart diseases (I24)													
Asian or Pacific Islander 3/, both sexes													
2007 1/.....	51	-	-	-	-	1	1	6	11	11	9	12	-
2006 1/.....	57	-	-	-	-	-	1	7	4	12	13	20	-
2005 1/.....	39	-	-	-	-	-	-	5	7	4	10	13	-
2004 1/.....	19	-	-	-	-	-	-	3	2	7	6	1	-
2003 1/.....	22	1	-	-	-	1	-	4	2	3	5	6	-
2002.....	18	-	-	-	-	-	-	1	6	2	4	5	-
2001.....	29	-	-	-	-	-	1	2	3	5	12	6	-
2000.....	25	-	-	-	-	-	-	-	5	6	9	5	-
1999.....	12	-	-	-	-	-	1	-	1	4	2	4	-
Asian or Pacific Islander 3/, male													
2007 1/.....	30	-	-	-	-	1	1	4	8	8	4	4	-
2006 1/.....	32	-	-	-	-	-	1	6	2	8	8	7	-
2005 1/.....	21	-	-	-	-	-	-	4	5	2	4	6	-
2004 1/.....	13	-	-	-	-	-	-	3	1	6	2	1	-
2003 1/.....	12	-	-	-	-	1	-	2	1	2	5	1	-
2002.....	9	-	-	-	-	-	-	1	4	1	2	1	-
2001.....	12	-	-	-	-	-	-	1	2	2	4	3	-
2000.....	12	-	-	-	-	-	-	-	3	4	4	1	-
1999.....	8	-	-	-	-	-	1	-	1	3	2	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other acute ischemic heart diseases (I24)

Asian or Pacific Islander 3/, female

2007 1/.....	21	-	-	-	-	-	-	2	3	3	5	8	-
2006 1/.....	25	-	-	-	-	-	-	1	2	4	5	13	-
2005 1/.....	18	-	-	-	-	-	-	1	2	2	6	7	-
2004 1/.....	6	-	-	-	-	-	-	-	1	1	4	-	-
2003 1/.....	10	1	-	-	-	-	-	2	1	1	-	5	-
2002.....	9	-	-	-	-	-	-	-	2	1	2	4	-
2001.....	17	-	-	-	-	-	1	1	1	3	8	3	-
2000.....	13	-	-	-	-	-	-	-	2	2	5	4	-
1999.....	4	-	-	-	-	-	-	-	-	1	-	3	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)													
All races, both sexes													
2007.....	269,291	12	4	10	94	631	3,708	14,547	27,650	38,846	77,502	106,270	17
2006.....	280,031	9	1	7	71	611	3,931	14,890	27,737	41,214	82,560	108,977	23
2005.....	291,118	2	4	6	80	611	4,044	14,889	27,731	42,841	88,051	112,834	25
2004.....	292,048	3	6	11	69	635	4,214	14,622	26,885	44,376	90,421	110,787	19
2003.....	306,306	9	5	11	82	658	4,355	14,664	27,207	47,378	95,425	116,489	23
2002.....	311,461	10	2	12	67	578	4,445	14,611	26,869	49,233	97,903	117,699	32
2001.....	314,081	8	7	11	92	627	4,264	13,949	26,281	51,438	99,729	117,637	38
2000.....	318,943	9	8	10	63	544	4,295	13,731	26,240	53,782	100,576	119,646	39
1999.....	326,973	11	2	10	74	574	4,421	13,700	26,523	56,725	103,730	121,171	32
All races, male													
2007.....	142,179	5	3	7	72	498	2,888	11,129	20,245	25,474	42,495	39,351	12
2006.....	146,405	6	1	4	58	497	3,010	11,440	20,162	26,679	44,945	39,588	15
2005.....	150,213	1	2	5	65	470	3,104	11,425	20,026	27,457	47,362	40,281	15
2004.....	149,328	1	1	5	53	494	3,194	11,203	19,275	28,361	48,369	38,354	18
2003.....	155,206	7	3	4	56	506	3,305	11,300	19,517	30,348	50,809	39,334	17
2002.....	157,163	7	2	10	50	437	3,443	11,306	19,370	31,482	51,876	39,157	23
2001.....	156,410	5	1	8	65	487	3,263	10,753	18,828	32,716	52,404	37,853	27
2000.....	158,519	6	5	5	45	422	3,329	10,667	18,816	34,333	52,670	38,193	28
1999.....	161,434	6	1	9	50	429	3,437	10,583	18,779	36,069	53,939	38,103	29

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)													
All races, female													
2007.....	127,112	7	1	3	22	133	820	3,418	7,405	13,372	35,007	66,919	5
2006.....	133,626	3	-	3	13	114	921	3,450	7,575	14,535	37,615	69,389	8
2005.....	140,905	1	2	1	15	141	940	3,464	7,705	15,384	40,689	72,553	10
2004.....	142,720	2	5	6	16	141	1,020	3,419	7,610	16,015	42,052	72,433	1
2003.....	151,100	2	2	7	26	152	1,050	3,364	7,690	17,030	44,616	77,155	6
2002.....	154,298	3	-	2	17	141	1,002	3,305	7,499	17,751	46,027	78,542	9
2001.....	157,671	3	6	3	27	140	1,001	3,196	7,453	18,722	47,325	79,784	11
2000.....	160,424	3	3	5	18	122	966	3,064	7,424	19,449	47,906	81,453	11
1999.....	165,539	5	1	1	24	145	984	3,117	7,744	20,656	49,791	83,068	3
White, both sexes													
2007 1/.....	235,140	8	2	8	61	451	2,823	11,286	22,051	32,125	68,695	97,618	12
2006 1/.....	244,607	3	1	6	52	439	3,009	11,481	22,094	34,108	73,138	100,256	20
2005 1/.....	255,133	1	4	6	53	453	3,105	11,444	22,036	35,720	78,278	104,016	17
2004 1/.....	256,037	3	3	7	42	444	3,281	11,296	21,517	37,066	80,522	101,844	12
2003 1/.....	268,904	7	3	5	55	480	3,365	11,332	21,769	39,571	85,076	107,223	18
2002.....	274,341	4	-	9	48	427	3,431	11,345	21,489	41,494	87,542	108,528	24
2001.....	276,772	4	4	9	58	454	3,266	10,836	20,948	43,390	89,238	108,541	24
2000.....	281,977	6	8	9	47	392	3,287	10,721	20,929	45,627	90,319	110,603	29
1999.....	289,322	8	1	7	43	423	3,339	10,684	21,225	48,184	93,398	111,988	22

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)													
White, male													
2007 1/.....	124,253	4	1	5	47	355	2,234	8,843	16,464	21,500	38,242	36,550	8
2006 1/.....	127,786	3	1	4	44	367	2,378	9,057	16,374	22,414	40,354	36,776	14
2005 1/.....	131,653	1	2	5	45	346	2,445	8,987	16,274	23,297	42,714	37,527	10
2004 1/.....	130,985	1	1	4	34	349	2,553	8,913	15,737	24,183	43,687	35,512	11
2003 1/.....	136,256	5	2	3	37	374	2,610	8,971	16,005	25,803	45,883	36,551	12
2002.....	138,480	3	-	7	35	336	2,733	9,067	15,810	27,053	47,057	36,361	18
2001.....	137,753	3	-	7	41	361	2,563	8,589	15,350	28,118	47,514	35,191	16
2000.....	140,115	4	5	5	33	315	2,639	8,582	15,369	29,788	47,876	35,476	23
1999.....	142,615	4	1	7	31	322	2,669	8,469	15,335	31,281	49,214	35,262	20
White, female													
2007 1/.....	110,887	4	1	3	14	96	589	2,443	5,587	10,625	30,453	61,068	4
2006 1/.....	116,821	-	-	2	8	72	631	2,424	5,720	11,694	32,784	63,480	6
2005 1/.....	123,480	-	2	1	8	107	660	2,457	5,762	12,423	35,564	66,489	7
2004 1/.....	125,052	2	2	3	8	95	728	2,383	5,780	12,883	36,835	66,332	1
2003 1/.....	132,648	2	1	2	18	106	755	2,361	5,764	13,768	39,193	70,672	6
2002.....	135,861	1	-	2	13	91	698	2,278	5,679	14,441	40,485	72,167	6
2001.....	139,019	1	4	2	17	93	703	2,247	5,598	15,272	41,724	73,350	8
2000.....	141,862	2	3	4	14	77	648	2,139	5,560	15,839	42,443	75,127	6
1999.....	146,707	4	-	-	12	101	670	2,215	5,890	16,903	44,184	76,726	2

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)													
Black, both sexes													
2007 1/.....	28,043	2	1	1	30	151	710	2,868	4,833	5,598	7,060	6,785	4
2006 1/.....	29,220	5	-	-	19	136	779	2,979	4,821	5,942	7,631	6,905	3
2005 1/.....	29,933	1	-	-	20	121	799	3,000	4,949	5,959	8,024	7,053	7
2004 1/.....	30,094	-	1	4	24	158	812	2,920	4,669	6,235	8,104	7,160	7
2003 1/.....	31,397	2	2	5	23	143	818	2,906	4,738	6,660	8,572	7,525	3
2002.....	31,406	5	2	3	15	118	879	2,881	4,704	6,605	8,641	7,545	8
2001.....	31,898	4	3	2	29	149	878	2,755	4,671	6,924	8,814	7,657	12
2000.....	31,697	2	-	-	12	134	882	2,669	4,647	7,079	8,631	7,631	10
1999.....	32,288	2	1	3	28	124	957	2,671	4,665	7,405	8,676	7,746	10
Black, male													
2007 1/.....	14,468	1	1	1	23	117	503	1,957	3,219	3,259	3,356	2,028	3
2006 1/.....	15,030	2	-	-	14	98	510	2,031	3,169	3,522	3,647	2,036	1
2005 1/.....	15,110	-	-	-	14	91	541	2,084	3,206	3,431	3,726	2,013	4
2004 1/.....	15,014	-	-	1	18	114	535	1,963	3,030	3,529	3,735	2,082	7
2003 1/.....	15,545	2	1	1	16	102	558	1,986	3,001	3,814	4,001	2,060	3
2002.....	15,399	3	2	3	11	74	591	1,924	3,077	3,724	3,882	2,103	5
2001.....	15,627	2	1	1	20	107	604	1,888	2,981	3,925	4,000	2,088	10
2000.....	15,322	1	-	-	8	92	585	1,803	2,973	3,851	3,883	2,121	5
1999.....	15,670	1	-	2	16	87	658	1,829	2,985	4,083	3,814	2,186	9

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)													
Black, female													
2007 1/.....	13,575	1	-	-	7	34	207	911	1,614	2,339	3,704	4,757	1
2006 1/.....	14,190	3	-	-	5	38	269	948	1,652	2,420	3,984	4,869	2
2005 1/.....	14,823	1	-	-	6	30	258	916	1,743	2,528	4,298	5,040	3
2004 1/.....	15,080	-	1	3	6	44	277	957	1,639	2,706	4,369	5,078	-
2003 1/.....	15,852	-	1	4	7	41	260	920	1,737	2,846	4,571	5,465	-
2002.....	16,007	2	-	-	4	44	288	957	1,627	2,881	4,759	5,442	3
2001.....	16,271	2	2	1	9	42	274	867	1,690	2,999	4,814	5,569	2
2000.....	16,375	1	-	-	4	42	297	866	1,674	3,228	4,748	5,510	5
1999.....	16,618	1	1	1	12	37	299	842	1,680	3,322	4,862	5,560	1
American Indian 2/, both sexes													
2007 1/.....	1,125	1	1	-	1	10	44	115	188	251	305	209	-
2006 1/.....	1,163	-	-	-	-	7	38	119	217	270	303	209	-
2005 1/.....	1,155	-	-	-	1	10	38	136	220	252	258	240	-
2004 1/.....	1,186	-	1	-	-	14	40	121	197	240	329	244	-
2003 1/.....	1,162	-	-	-	2	10	53	124	180	269	287	235	2
2002.....	1,084	1	-	-	2	13	36	95	183	256	291	207	-
2001.....	1,003	-	-	-	3	3	33	99	192	252	259	162	-
2000.....	1,041	-	-	-	2	6	41	99	180	236	264	213	-
1999.....	1,009	1	-	-	1	8	33	98	165	250	270	183	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other forms of chronic ischemic heart disease (I20,I25)

American Indian 2/, male

2007 1/.....	655	-	1	-	1	7	34	92	129	162	157	72	-
2006 1/.....	682	-	-	-	-	6	30	93	150	177	152	74	-
2005 1/.....	663	-	-	-	-	9	29	98	159	150	140	78	-
2004 1/.....	688	-	-	-	-	12	32	84	139	147	181	93	-
2003 1/.....	688	-	-	-	1	8	40	89	123	173	157	95	2
2002.....	636	1	-	-	2	10	29	72	118	169	154	81	-
2001.....	563	-	-	-	3	2	20	68	138	153	126	53	-
2000.....	598	-	-	-	2	5	32	76	123	146	140	74	-
1999.....	575	1	-	-	1	4	24	76	111	154	145	59	-

American Indian 2/, female

2007 1/.....	470	1	-	-	-	3	10	23	59	89	148	137	-
2006 1/.....	481	-	-	-	-	1	8	26	67	93	151	135	-
2005 1/.....	492	-	-	-	1	1	9	38	61	102	118	162	-
2004 1/.....	498	-	1	-	-	2	8	37	58	93	148	151	-
2003 1/.....	474	-	-	-	1	2	13	35	57	96	130	140	-
2002.....	448	-	-	-	-	3	7	23	65	87	137	126	-
2001.....	440	-	-	-	-	1	13	31	54	99	133	109	-
2000.....	443	-	-	-	-	1	9	23	57	90	124	139	-
1999.....	434	-	-	-	-	4	9	22	54	96	125	124	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other forms of chronic ischemic heart disease (I20,I25)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	4,983	1	-	1	2	19	131	278	578	872	1,442	1,658	1
2006 1/.....	5,041	1	-	1	-	29	105	311	605	894	1,488	1,607	-
2005 1/.....	4,897	-	-	-	6	27	102	309	526	910	1,491	1,525	1
2004 1/.....	4,731	-	1	-	3	19	81	285	502	835	1,466	1,539	-
2003 1/.....	4,843	-	-	1	2	25	119	302	520	878	1,490	1,506	-
2002.....	4,630	-	-	-	2	20	99	290	493	878	1,429	1,419	-
2001.....	4,408	-	-	-	2	21	87	259	470	872	1,418	1,277	2
2000.....	4,228	1	-	1	2	12	85	242	484	840	1,362	1,199	-
1999.....	4,354	-	-	-	2	19	92	247	468	886	1,386	1,254	-

Asian or Pacific Islander 3/, male

2007 1/.....	2,803	-	-	1	1	19	117	237	433	553	740	701	1
2006 1/.....	2,907	1	-	-	-	26	92	259	469	566	792	702	-
2005 1/.....	2,787	-	-	-	6	24	89	256	387	579	782	663	1
2004 1/.....	2,641	-	-	-	1	19	74	243	369	502	766	667	-
2003 1/.....	2,717	-	-	-	2	22	97	254	388	558	768	628	-
2002.....	2,648	-	-	-	2	17	90	243	365	536	783	612	-
2001.....	2,467	-	-	-	1	17	76	208	359	520	764	521	1
2000.....	2,484	1	-	-	2	10	73	206	351	548	771	522	-
1999.....	2,574	-	-	-	2	16	86	209	348	551	766	596	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other forms of chronic ischemic heart disease (I20,I25)

Asian or Pacific Islander 3/, female

2007 1/.....	2,180	1	-	-	1	-	14	41	145	319	702	957	-
2006 1/.....	2,134	-	-	1	-	3	13	52	136	328	696	905	-
2005 1/.....	2,110	-	-	-	-	3	13	53	139	331	709	862	-
2004 1/.....	2,090	-	1	-	2	-	7	42	133	333	700	872	-
2003 1/.....	2,126	-	-	1	-	3	22	48	132	320	722	878	-
2002.....	1,982	-	-	-	-	3	9	47	128	342	646	807	-
2001.....	1,941	-	-	-	1	4	11	51	111	352	654	756	1
2000.....	1,744	-	-	1	-	2	12	36	133	292	591	677	-
1999.....	1,780	-	-	-	-	3	6	38	120	335	620	658	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)													
All races, both sexes													
2007.....	59,051	1	-	-	17	211	1,395	5,910	10,055	9,884	14,221	17,347	10
2006.....	61,030	-	-	-	20	210	1,441	5,820	9,900	10,420	14,905	18,298	16
2005.....	62,799	1	-	-	16	182	1,420	5,884	9,565	10,246	16,286	19,182	17
2004.....	63,661	-	1	-	9	193	1,463	5,665	9,297	10,830	16,730	19,459	14
2003.....	67,232	2	-	-	16	211	1,488	5,684	9,110	11,494	18,339	20,874	14
2002.....	68,129	2	-	1	17	181	1,515	5,366	8,793	11,817	18,618	21,796	23
2001.....	68,103	-	-	-	20	179	1,457	5,047	8,438	12,005	19,196	21,736	25
2000.....	69,554	1	-	1	15	144	1,376	4,980	8,242	12,567	19,545	22,664	19
1999.....	71,926	-	1	1	13	160	1,488	4,737	8,383	13,395	20,321	23,403	24
All races, male													
2007.....	33,401	1	-	-	12	176	1,109	4,618	7,583	6,549	7,552	5,794	7
2006.....	34,157	-	-	-	17	173	1,119	4,612	7,427	6,854	7,777	6,168	10
2005.....	34,453	-	-	-	15	146	1,120	4,643	7,160	6,709	8,384	6,266	10
2004.....	34,574	-	-	-	8	157	1,143	4,495	6,864	7,046	8,606	6,242	13
2003.....	36,042	2	-	-	12	181	1,157	4,552	6,694	7,422	9,333	6,677	12
2002.....	36,047	1	-	-	13	144	1,216	4,239	6,549	7,617	9,416	6,835	17
2001.....	35,440	-	-	-	16	133	1,130	4,037	6,245	7,668	9,671	6,521	19
2000.....	35,896	-	-	1	12	115	1,099	3,977	6,125	8,069	9,725	6,758	15
1999.....	36,633	-	-	1	10	119	1,192	3,752	6,123	8,546	9,963	6,905	22

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)													
All races, female													
2007.....	25,650	-	-	-	5	35	286	1,292	2,472	3,335	6,669	11,553	3
2006.....	26,873	-	-	-	3	37	322	1,208	2,473	3,566	7,128	12,130	6
2005.....	28,346	1	-	-	1	36	300	1,241	2,405	3,537	7,902	12,916	7
2004.....	29,087	-	1	-	1	36	320	1,170	2,433	3,784	8,124	13,217	1
2003.....	31,190	-	-	-	4	30	331	1,132	2,416	4,072	9,006	14,197	2
2002.....	32,082	1	-	1	4	37	299	1,127	2,244	4,200	9,202	14,961	6
2001.....	32,663	-	-	-	4	46	327	1,010	2,193	4,337	9,525	15,215	6
2000.....	33,658	1	-	-	3	29	277	1,003	2,117	4,498	9,820	15,906	4
1999.....	35,293	-	1	-	3	41	296	985	2,260	4,849	10,358	16,498	2
White, both sexes													
2007 1/.....	48,660	1	-	-	10	144	1,049	4,492	7,770	7,737	11,997	15,451	9
2006 1/.....	50,146	-	-	-	16	138	1,072	4,352	7,579	8,114	12,529	16,333	13
2005 1/.....	51,717	-	-	-	14	124	1,073	4,420	7,200	8,026	13,720	17,129	11
2004 1/.....	52,585	-	-	-	7	132	1,138	4,242	7,108	8,487	14,132	17,332	7
2003 1/.....	55,760	1	-	-	12	156	1,144	4,262	6,927	8,976	15,572	18,700	10
2002.....	56,569	1	-	1	15	132	1,130	4,048	6,682	9,222	15,783	19,539	16
2001.....	56,414	-	-	-	12	133	1,089	3,783	6,232	9,316	16,316	19,517	16
2000.....	57,794	-	-	1	13	102	1,034	3,732	6,113	9,858	16,609	20,319	13
1999.....	59,711	-	-	1	5	110	1,097	3,488	6,211	10,472	17,317	20,995	15

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)													
White, male													
2007 1/.....	27,365	1	-	-	7	117	851	3,590	5,957	5,220	6,418	5,198	6
2006 1/.....	27,777	-	-	-	13	121	857	3,541	5,764	5,377	6,581	5,514	9
2005 1/.....	28,155	-	-	-	13	97	876	3,583	5,503	5,304	7,102	5,672	5
2004 1/.....	28,370	-	-	-	6	107	913	3,496	5,314	5,638	7,326	5,564	6
2003 1/.....	29,560	1	-	-	9	133	899	3,503	5,194	5,845	7,974	5,994	8
2002.....	29,604	1	-	-	11	109	935	3,292	5,067	6,040	8,000	6,137	12
2001.....	29,039	-	-	-	8	103	864	3,113	4,736	6,044	8,296	5,864	11
2000.....	29,467	-	-	1	10	83	852	3,067	4,660	6,447	8,286	6,049	12
1999.....	29,977	-	-	1	5	85	909	2,839	4,633	6,801	8,526	6,165	13
White, female													
2007 1/.....	21,295	-	-	-	3	27	198	902	1,813	2,517	5,579	10,253	3
2006 1/.....	22,369	-	-	-	3	17	215	811	1,815	2,737	5,948	10,819	4
2005 1/.....	23,562	-	-	-	1	27	197	837	1,697	2,722	6,618	11,457	6
2004 1/.....	24,215	-	-	-	1	25	225	746	1,794	2,849	6,806	11,768	1
2003 1/.....	26,200	-	-	-	3	23	245	759	1,733	3,131	7,598	12,706	2
2002.....	26,965	-	-	1	4	23	195	756	1,615	3,182	7,783	13,402	4
2001.....	27,375	-	-	-	4	30	225	670	1,496	3,272	8,020	13,653	5
2000.....	28,327	-	-	-	3	19	182	665	1,453	3,411	8,323	14,270	1
1999.....	29,734	-	-	-	-	25	188	649	1,578	3,671	8,791	14,830	2

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)													
Black, both sexes													
2007 1/.....	8,921	-	-	-	6	54	279	1,254	2,012	1,877	1,853	1,585	1
2006 1/.....	9,362	-	-	-	4	58	317	1,296	2,017	1,987	2,030	1,650	3
2005 1/.....	9,576	1	-	-	1	42	296	1,288	2,070	1,928	2,188	1,757	5
2004 1/.....	9,666	-	1	-	2	49	277	1,268	1,934	2,080	2,257	1,791	7
2003 1/.....	9,923	1	-	-	3	47	285	1,234	1,926	2,190	2,372	1,862	3
2002.....	10,195	1	-	-	2	41	335	1,188	1,885	2,300	2,498	1,938	7
2001.....	10,321	-	-	-	7	38	325	1,130	1,965	2,394	2,518	1,936	8
2000.....	10,418	1	-	-	1	36	295	1,140	1,911	2,408	2,553	2,067	6
1999.....	10,823	-	1	-	8	41	361	1,117	1,942	2,608	2,639	2,097	9
Black, male													
2007 1/.....	5,155	-	-	-	5	47	203	887	1,413	1,166	952	481	1
2006 1/.....	5,410	-	-	-	4	40	216	917	1,428	1,268	1,015	521	1
2005 1/.....	5,372	-	-	-	1	34	202	911	1,432	1,225	1,078	485	4
2004 1/.....	5,331	-	-	-	2	39	187	867	1,353	1,241	1,100	535	7
2003 1/.....	5,543	1	-	-	2	40	207	897	1,306	1,352	1,175	560	3
2002.....	5,598	-	-	-	2	28	234	832	1,311	1,382	1,245	559	5
2001.....	5,578	-	-	-	7	24	231	811	1,328	1,438	1,195	537	7
2000.....	5,615	-	-	-	1	26	210	821	1,308	1,417	1,221	608	3
1999.....	5,815	-	-	-	5	27	257	802	1,309	1,559	1,236	611	9

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)													
Black, female													
2007 1/.....	3,766	-	-	-	1	7	76	367	599	711	901	1,104	-
2006 1/.....	3,952	-	-	-	-	18	101	379	589	719	1,015	1,129	2
2005 1/.....	4,204	1	-	-	-	8	94	377	638	703	1,110	1,272	1
2004 1/.....	4,335	-	1	-	-	10	90	401	581	839	1,157	1,256	-
2003 1/.....	4,380	-	-	-	1	7	78	337	620	838	1,197	1,302	-
2002.....	4,597	1	-	-	-	13	101	356	574	918	1,253	1,379	2
2001.....	4,743	-	-	-	-	14	94	319	637	956	1,323	1,399	1
2000.....	4,803	1	-	-	-	10	85	319	603	991	1,332	1,459	3
1999.....	5,008	-	1	-	3	14	104	315	633	1,049	1,403	1,486	-
American Indian 2/, both sexes													
2007 1/.....	325	-	-	-	-	4	13	51	61	76	81	39	-
2006 1/.....	317	-	-	-	-	4	13	47	77	70	69	37	-
2005 1/.....	344	-	-	-	-	6	19	57	80	72	66	44	-
2004 1/.....	331	-	-	-	-	5	14	46	70	66	71	59	-
2003 1/.....	358	-	-	-	-	2	18	53	61	94	70	59	1
2002.....	318	-	-	-	-	3	18	35	60	81	66	55	-
2001.....	335	-	-	-	1	2	14	44	77	83	71	43	-
2000.....	266	-	-	-	1	4	12	33	52	55	58	51	-
1999.....	295	-	-	-	-	3	6	37	52	70	79	48	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Atherosclerotic cardiovascular disease, so described (I25.0)

American Indian 2/, male

2007 1/.....	200	-	-	-	-	3	10	40	44	51	41	11	-
2006 1/.....	210	-	-	-	-	3	10	39	55	50	35	18	-
2005 1/.....	216	-	-	-	-	5	14	44	57	40	39	17	-
2004 1/.....	207	-	-	-	-	4	11	32	53	40	43	24	-
2003 1/.....	227	-	-	-	-	2	15	40	42	66	37	24	1
2002.....	196	-	-	-	-	2	16	30	38	55	32	23	-
2001.....	196	-	-	-	1	1	9	32	55	50	34	14	-
2000.....	159	-	-	-	1	4	8	25	39	33	33	16	-
1999.....	179	-	-	-	-	2	4	29	40	44	44	16	-

American Indian 2/, female

2007 1/.....	125	-	-	-	-	1	3	11	17	25	40	28	-
2006 1/.....	107	-	-	-	-	1	3	8	22	20	34	19	-
2005 1/.....	128	-	-	-	-	1	5	13	23	32	27	27	-
2004 1/.....	124	-	-	-	-	1	3	14	17	26	28	35	-
2003 1/.....	131	-	-	-	-	-	3	13	19	28	33	35	-
2002.....	122	-	-	-	-	1	2	5	22	26	34	32	-
2001.....	139	-	-	-	-	1	5	12	22	33	37	29	-
2000.....	107	-	-	-	-	-	4	8	13	22	25	35	-
1999.....	116	-	-	-	-	1	2	8	12	26	35	32	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Atherosclerotic cardiovascular disease, so described (I25.0)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	1,145	-	-	-	1	9	54	113	212	194	290	272	-
2006 1/.....	1,205	-	-	-	-	10	39	125	227	249	277	278	-
2005 1/.....	1,162	-	-	-	1	10	32	119	215	220	312	252	1
2004 1/.....	1,079	-	-	-	-	7	34	109	185	197	270	277	-
2003 1/.....	1,191	-	-	-	1	6	41	135	196	234	325	253	-
2002.....	1,047	-	-	-	-	5	32	95	166	214	271	264	-
2001.....	1,033	-	-	-	-	6	29	90	164	212	291	240	1
2000.....	1,076	-	-	-	-	2	35	75	166	246	325	227	-
1999.....	1,097	-	-	-	-	6	24	95	178	245	286	263	-

Asian or Pacific Islander 3/, male

2007 1/.....	681	-	-	-	-	9	45	101	169	112	141	104	-
2006 1/.....	760	-	-	-	-	9	36	115	180	159	146	115	-
2005 1/.....	710	-	-	-	1	10	28	105	168	140	165	92	1
2004 1/.....	666	-	-	-	-	7	32	100	144	127	137	119	-
2003 1/.....	712	-	-	-	1	6	36	112	152	159	147	99	-
2002.....	649	-	-	-	-	5	31	85	133	140	139	116	-
2001.....	627	-	-	-	-	5	26	81	126	136	146	106	1
2000.....	655	-	-	-	-	2	29	64	118	172	185	85	-
1999.....	662	-	-	-	-	5	22	82	141	142	157	113	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Atherosclerotic cardiovascular disease, so described (I25.0)

Asian or Pacific Islander 3/, female

2007 1/.....	464	-	-	-	1	-	9	12	43	82	149	168	-
2006 1/.....	445	-	-	-	-	1	3	10	47	90	131	163	-
2005 1/.....	452	-	-	-	-	-	4	14	47	80	147	160	-
2004 1/.....	413	-	-	-	-	-	2	9	41	70	133	158	-
2003 1/.....	479	-	-	-	-	-	5	23	44	75	178	154	-
2002.....	398	-	-	-	-	-	1	10	33	74	132	148	-
2001.....	406	-	-	-	-	1	3	9	38	76	145	134	-
2000.....	421	-	-	-	-	-	6	11	48	74	140	142	-
1999.....	435	-	-	-	-	1	2	13	37	103	129	150	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

All races, both sexes

2007.....	210,240	11	4	10	77	420	2,313	8,637	17,595	28,962	63,281	88,923	7
2006.....	219,001	9	1	7	51	401	2,490	9,070	17,837	30,794	67,655	90,679	7
2005.....	228,319	1	4	6	64	429	2,624	9,005	18,166	32,595	71,765	93,652	8
2004.....	228,387	3	5	11	60	442	2,751	8,957	17,588	33,546	73,691	91,328	5
2003.....	239,074	7	5	11	66	447	2,867	8,980	18,097	35,884	77,086	95,615	9
2002.....	243,332	8	2	11	50	397	2,930	9,245	18,076	37,416	79,285	95,903	9
2001.....	245,978	8	7	11	72	448	2,807	8,902	17,843	39,433	80,533	95,901	13
2000.....	249,389	8	8	9	48	400	2,919	8,751	17,998	41,215	81,031	96,982	20
1999.....	255,047	11	1	9	61	414	2,933	8,963	18,140	43,330	83,409	97,768	8

All races, male

2007.....	108,778	4	3	7	60	322	1,779	6,511	12,662	18,925	34,943	33,557	5
2006.....	112,248	6	1	4	41	324	1,891	6,828	12,735	19,825	37,168	33,420	5
2005.....	115,760	1	2	5	50	324	1,984	6,782	12,866	20,748	38,978	34,015	5
2004.....	114,754	1	1	5	45	337	2,051	6,708	12,411	21,315	39,763	32,112	5
2003.....	119,164	5	3	4	44	325	2,148	6,748	12,823	22,926	41,476	32,657	5
2002.....	121,116	6	2	10	37	293	2,227	7,067	12,821	23,865	42,460	32,322	6
2001.....	120,970	5	1	8	49	354	2,133	6,716	12,583	25,048	42,733	31,332	8
2000.....	122,623	6	5	4	33	307	2,230	6,690	12,691	26,264	42,945	31,435	13
1999.....	124,801	6	1	8	40	310	2,245	6,831	12,656	27,523	43,976	31,198	7

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

All races, female

2007.....	101,462	7	1	3	17	98	534	2,126	4,933	10,037	28,338	55,366	2
2006.....	106,753	3	-	3	10	77	599	2,242	5,102	10,969	30,487	57,259	2
2005.....	112,559	-	2	1	14	105	640	2,223	5,300	11,847	32,787	59,637	3
2004.....	113,633	2	4	6	15	105	700	2,249	5,177	12,231	33,928	59,216	-
2003.....	119,910	2	2	7	22	122	719	2,232	5,274	12,958	35,610	62,958	4
2002.....	122,216	2	-	1	13	104	703	2,178	5,255	13,551	36,825	63,581	3
2001.....	125,008	3	6	3	23	94	674	2,186	5,260	14,385	37,800	64,569	5
2000.....	126,766	2	3	5	15	93	689	2,061	5,307	14,951	38,086	65,547	7
1999.....	130,246	5	-	1	21	104	688	2,132	5,484	15,807	39,433	66,570	1

White, both sexes

2007 1/.....	186,480	7	2	8	51	307	1,774	6,794	14,281	24,388	56,698	82,167	3
2006 1/.....	194,461	3	1	6	36	301	1,937	7,129	14,515	25,994	60,609	83,923	7
2005 1/.....	203,416	1	4	6	39	329	2,032	7,024	14,836	27,694	64,558	86,887	6
2004 1/.....	203,452	3	3	7	35	312	2,143	7,054	14,409	28,579	66,390	84,512	5
2003 1/.....	213,144	6	3	5	43	324	2,221	7,070	14,842	30,595	69,504	88,523	8
2002.....	217,772	3	-	8	33	295	2,301	7,297	14,807	32,272	71,759	88,989	8
2001.....	220,358	4	4	9	46	321	2,177	7,053	14,716	34,074	72,922	89,024	8
2000.....	224,183	6	8	8	34	290	2,253	6,989	14,816	35,769	73,710	90,284	16
1999.....	229,611	8	1	6	38	313	2,242	7,196	15,014	37,712	76,081	90,993	7

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

White, male

2007 1/.....	96,888	3	1	5	40	238	1,383	5,253	10,507	16,280	31,824	31,352	2
2006 1/.....	100,009	3	1	4	31	246	1,521	5,516	10,610	17,037	33,773	31,262	5
2005 1/.....	103,498	1	2	5	32	249	1,569	5,404	10,771	17,993	35,612	31,855	5
2004 1/.....	102,615	1	1	4	28	242	1,640	5,417	10,423	18,545	36,361	29,948	5
2003 1/.....	106,696	4	2	3	28	241	1,711	5,468	10,811	19,958	37,909	30,557	4
2002.....	108,876	2	-	7	24	227	1,798	5,775	10,743	21,013	39,057	30,224	6
2001.....	108,714	3	-	7	33	258	1,699	5,476	10,614	22,074	39,218	29,327	5
2000.....	110,648	4	5	4	23	232	1,787	5,515	10,709	23,341	39,590	29,427	11
1999.....	112,638	4	1	6	26	237	1,760	5,630	10,702	24,480	40,688	29,097	7

White, female

2007 1/.....	89,592	4	1	3	11	69	391	1,541	3,774	8,108	24,874	50,815	1
2006 1/.....	94,452	-	-	2	5	55	416	1,613	3,905	8,957	26,836	52,661	2
2005 1/.....	99,918	-	2	1	7	80	463	1,620	4,065	9,701	28,946	55,032	1
2004 1/.....	100,837	2	2	3	7	70	503	1,637	3,986	10,034	30,029	54,564	-
2003 1/.....	106,448	2	1	2	15	83	510	1,602	4,031	10,637	31,595	57,966	4
2002.....	108,896	1	-	1	9	68	503	1,522	4,064	11,259	32,702	58,765	2
2001.....	111,644	1	4	2	13	63	478	1,577	4,102	12,000	33,704	59,697	3
2000.....	113,535	2	3	4	11	58	466	1,474	4,107	12,428	34,120	60,857	5
1999.....	116,973	4	-	-	12	76	482	1,566	4,312	13,232	35,393	61,896	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)													
Black, both sexes													
2007 1/.....	19,122	2	1	1	24	97	431	1,614	2,821	3,721	5,207	5,200	3
2006 1/.....	19,858	5	-	-	15	78	462	1,683	2,804	3,955	5,601	5,255	-
2005 1/.....	20,357	-	-	-	19	79	503	1,712	2,879	4,031	5,836	5,296	2
2004 1/.....	20,428	-	-	4	22	109	535	1,652	2,735	4,155	5,847	5,369	-
2003 1/.....	21,474	1	2	5	20	96	533	1,672	2,812	4,470	6,200	5,663	-
2002.....	21,211	4	2	3	13	77	544	1,693	2,819	4,305	6,143	5,607	1
2001.....	21,577	4	3	2	22	111	553	1,625	2,706	4,530	6,296	5,721	4
2000.....	21,279	1	-	-	11	98	587	1,529	2,736	4,671	6,078	5,564	4
1999.....	21,465	2	-	3	20	83	596	1,554	2,723	4,797	6,037	5,649	1
Black, male													
2007 1/.....	9,313	1	1	1	18	70	300	1,070	1,806	2,093	2,404	1,547	2
2006 1/.....	9,620	2	-	-	10	58	294	1,114	1,741	2,254	2,632	1,515	-
2005 1/.....	9,738	-	-	-	13	57	339	1,173	1,774	2,206	2,648	1,528	-
2004 1/.....	9,683	-	-	1	16	75	348	1,096	1,677	2,288	2,635	1,547	-
2003 1/.....	10,002	1	1	1	14	62	351	1,089	1,695	2,462	2,826	1,500	-
2002.....	9,801	3	2	3	9	46	357	1,092	1,766	2,342	2,637	1,544	-
2001.....	10,049	2	1	1	13	83	373	1,077	1,653	2,487	2,805	1,551	3
2000.....	9,707	1	-	-	7	66	375	982	1,665	2,434	2,662	1,513	2
1999.....	9,855	1	-	2	11	60	401	1,027	1,676	2,524	2,578	1,575	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Black, female

2007 1/.....	9,809	1	-	-	6	27	131	544	1,015	1,628	2,803	3,653	1
2006 1/.....	10,238	3	-	-	5	20	168	569	1,063	1,701	2,969	3,740	-
2005 1/.....	10,619	-	-	-	6	22	164	539	1,105	1,825	3,188	3,768	2
2004 1/.....	10,745	-	-	3	6	34	187	556	1,058	1,867	3,212	3,822	-
2003 1/.....	11,472	-	1	4	6	34	182	583	1,117	2,008	3,374	4,163	-
2002.....	11,410	1	-	-	4	31	187	601	1,053	1,963	3,506	4,063	1
2001.....	11,528	2	2	1	9	28	180	548	1,053	2,043	3,491	4,170	1
2000.....	11,572	-	-	-	4	32	212	547	1,071	2,237	3,416	4,051	2
1999.....	11,610	1	-	1	9	23	195	527	1,047	2,273	3,459	4,074	1

American Indian 2/, both sexes

2007 1/.....	800	1	1	-	1	6	31	64	127	175	224	170	-
2006 1/.....	846	-	-	-	-	3	25	72	140	200	234	172	-
2005 1/.....	811	-	-	-	1	4	19	79	140	180	192	196	-
2004 1/.....	855	-	1	-	-	9	26	75	127	174	258	185	-
2003 1/.....	804	-	-	-	2	8	35	71	119	175	217	176	1
2002.....	766	1	-	-	2	10	18	60	123	175	225	152	-
2001.....	668	-	-	-	2	1	19	55	115	169	188	119	-
2000.....	775	-	-	-	1	2	29	66	128	181	206	162	-
1999.....	714	1	-	-	1	5	27	61	113	180	191	135	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

American Indian 2/, male

2007 1/.....	455	-	1	-	1	4	24	52	85	111	116	61	-
2006 1/.....	472	-	-	-	-	3	20	54	95	127	117	56	-
2005 1/.....	447	-	-	-	-	4	15	54	102	110	101	61	-
2004 1/.....	481	-	-	-	-	8	21	52	86	107	138	69	-
2003 1/.....	461	-	-	-	1	6	25	49	81	107	120	71	1
2002.....	440	1	-	-	2	8	13	42	80	114	122	58	-
2001.....	367	-	-	-	2	1	11	36	83	103	92	39	-
2000.....	439	-	-	-	1	1	24	51	84	113	107	58	-
1999.....	396	1	-	-	1	2	20	47	71	110	101	43	-

American Indian 2/, female

2007 1/.....	345	1	-	-	-	2	7	12	42	64	108	109	-
2006 1/.....	374	-	-	-	-	-	5	18	45	73	117	116	-
2005 1/.....	364	-	-	-	1	-	4	25	38	70	91	135	-
2004 1/.....	374	-	1	-	-	1	5	23	41	67	120	116	-
2003 1/.....	343	-	-	-	1	2	10	22	38	68	97	105	-
2002.....	326	-	-	-	-	2	5	18	43	61	103	94	-
2001.....	301	-	-	-	-	-	8	19	32	66	96	80	-
2000.....	336	-	-	-	-	1	5	15	44	68	99	104	-
1999.....	318	-	-	-	-	3	7	14	42	70	90	92	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)													
Asian or Pacific Islander 3/, both sexes													
2007 1/.....	3,838	1	-	1	1	10	77	165	366	678	1,152	1,386	1
2006 1/.....	3,836	1	-	1	-	19	66	186	378	645	1,211	1,329	-
2005 1/.....	3,735	-	-	-	5	17	70	190	311	690	1,179	1,273	-
2004 1/.....	3,652	-	1	-	3	12	47	176	317	638	1,196	1,262	-
2003 1/.....	3,652	-	-	1	1	19	78	167	324	644	1,165	1,253	-
2002.....	3,583	-	-	-	2	15	67	195	327	664	1,158	1,155	-
2001.....	3,375	-	-	-	2	15	58	169	306	660	1,127	1,037	1
2000.....	3,152	1	-	1	2	10	50	167	318	594	1,037	972	-
1999.....	3,257	-	-	-	2	13	68	152	290	641	1,100	991	-
Asian or Pacific Islander 3/, male													
2007 1/.....	2,122	-	-	1	1	10	72	136	264	441	599	597	1
2006 1/.....	2,147	1	-	-	-	17	56	144	289	407	646	587	-
2005 1/.....	2,077	-	-	-	5	14	61	151	219	439	617	571	-
2004 1/.....	1,975	-	-	-	1	12	42	143	225	375	629	548	-
2003 1/.....	2,005	-	-	-	1	16	61	142	236	399	621	529	-
2002.....	1,999	-	-	-	2	12	59	158	232	396	644	496	-
2001.....	1,840	-	-	-	1	12	50	127	233	384	618	415	-
2000.....	1,829	1	-	-	2	8	44	142	233	376	586	437	-
1999.....	1,912	-	-	-	2	11	64	127	207	409	609	483	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Asian or Pacific Islander 3/, female

2007 1/.....	1,716	1	-	-	-	-	5	29	102	237	553	789	-
2006 1/.....	1,689	-	-	1	-	2	10	42	89	238	565	742	-
2005 1/.....	1,658	-	-	-	-	3	9	39	92	251	562	702	-
2004 1/.....	1,677	-	1	-	2	-	5	33	92	263	567	714	-
2003 1/.....	1,647	-	-	1	-	3	17	25	88	245	544	724	-
2002.....	1,584	-	-	-	-	3	8	37	95	268	514	659	-
2001.....	1,535	-	-	-	1	3	8	42	73	276	509	622	1
2000.....	1,323	-	-	1	-	2	6	25	85	218	451	535	-
1999.....	1,345	-	-	-	-	2	4	25	83	232	491	508	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other heart diseases (I26-I51)													
All races, both sexes													
2007.....	172,748	397	163	219	872	1,777	4,101	9,085	14,246	21,645	46,934	73,305	4
2006.....	170,248	320	154	231	873	1,903	4,135	9,164	14,030	21,240	47,065	71,126	7
2005.....	170,585	342	143	231	918	1,847	4,295	9,120	13,861	21,763	47,866	70,193	6
2004.....	166,275	406	174	221	855	1,719	4,284	8,952	13,134	21,905	47,853	66,765	7
2003.....	169,984	416	171	234	904	1,810	4,555	8,716	13,398	23,244	49,260	67,270	6
2002.....	169,540	473	157	227	836	1,808	4,620	8,711	12,946	23,680	49,776	66,300	6
2001.....	166,949	449	215	246	794	1,736	4,472	8,484	12,535	24,056	48,976	64,979	7
2000.....	165,428	457	168	248	853	1,749	4,457	8,087	12,362	24,679	48,367	63,993	8
1999.....	165,828	490	177	248	880	1,779	4,647	7,888	12,580	25,572	48,712	62,848	7
All races, male													
2007.....	77,178	225	77	121	549	1,100	2,577	5,714	8,762	12,120	21,961	23,968	4
2006.....	75,221	169	84	130	555	1,217	2,540	5,793	8,550	11,806	21,780	22,591	6
2005.....	75,094	188	77	118	614	1,176	2,695	5,769	8,415	11,987	21,954	22,097	4
2004.....	73,196	218	90	111	555	1,075	2,642	5,661	8,071	12,033	21,960	20,774	6
2003.....	74,738	233	81	141	560	1,081	2,801	5,458	8,142	12,875	22,680	20,681	5
2002.....	74,230	249	82	125	557	1,138	2,860	5,550	7,771	12,991	22,771	20,133	3
2001.....	72,439	222	117	123	500	1,037	2,777	5,307	7,503	13,167	22,540	19,141	5
2000.....	72,122	238	98	150	518	1,049	2,746	5,151	7,532	13,621	22,006	19,008	5
1999.....	72,601	255	100	137	533	1,067	2,894	4,979	7,700	14,320	22,128	18,483	5

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other heart diseases (I26-I51)													
All races, female													
2007.....	95,570	172	86	98	323	677	1,524	3,371	5,484	9,525	24,973	49,337	-
2006.....	95,027	151	70	101	318	686	1,595	3,371	5,480	9,434	25,285	48,535	1
2005.....	95,491	154	66	113	304	671	1,600	3,351	5,446	9,776	25,912	48,096	2
2004.....	93,079	188	84	110	300	644	1,642	3,291	5,063	9,872	25,893	45,991	1
2003.....	95,246	183	90	93	344	729	1,754	3,258	5,256	10,369	26,580	46,589	1
2002.....	95,310	224	75	102	279	670	1,760	3,161	5,175	10,689	27,005	46,167	3
2001.....	94,510	227	98	123	294	699	1,695	3,177	5,032	10,889	26,436	45,838	2
2000.....	93,306	219	70	98	335	700	1,711	2,936	4,830	11,058	26,361	44,985	3
1999.....	93,227	235	77	111	347	712	1,753	2,909	4,880	11,252	26,584	44,365	2
White, both sexes													
2007 1/.....	149,300	233	106	148	582	1,136	2,757	6,420	10,819	17,582	41,652	67,861	4
2006 1/.....	147,421	192	105	158	587	1,200	2,787	6,482	10,721	17,357	41,846	65,982	4
2005 1/.....	147,474	219	87	141	612	1,151	2,917	6,390	10,508	17,824	42,591	65,029	5
2004 1/.....	143,860	276	107	144	577	1,088	2,900	6,277	10,071	18,083	42,617	61,716	4
2003 1/.....	147,127	256	109	155	590	1,142	3,078	6,131	10,276	19,101	44,074	62,210	5
2002.....	147,047	314	109	157	548	1,124	3,188	6,128	9,981	19,716	44,518	61,261	3
2001.....	144,862	292	134	171	512	1,119	2,972	6,014	9,668	20,052	43,758	60,164	6
2000.....	143,761	315	114	158	538	1,133	3,006	5,737	9,598	20,617	43,379	59,159	7
1999.....	143,786	329	119	175	595	1,154	3,127	5,582	9,628	21,374	43,578	58,120	5

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other heart diseases (I26-I51)													
White, male													
2007 1/.....	65,872	132	44	81	354	725	1,734	4,139	6,732	9,956	19,644	22,327	4
2006 1/.....	64,243	100	57	92	388	787	1,728	4,152	6,622	9,746	19,568	21,000	3
2005 1/.....	64,180	118	53	77	408	760	1,884	4,170	6,488	9,897	19,743	20,578	4
2004 1/.....	62,477	148	58	74	379	695	1,854	4,084	6,280	10,083	19,655	19,164	3
2003 1/.....	63,973	140	48	95	362	709	1,955	3,924	6,343	10,759	20,456	19,178	4
2002.....	63,698	168	62	86	374	722	2,036	4,015	6,103	10,960	20,533	18,637	2
2001.....	62,025	155	71	88	319	676	1,923	3,898	5,872	11,127	20,209	17,682	5
2000.....	61,933	164	71	100	333	702	1,905	3,746	5,959	11,561	19,865	17,522	5
1999.....	62,054	174	70	95	352	712	2,021	3,621	5,989	12,138	19,917	16,961	4
White, female													
2007 1/.....	83,428	101	62	67	228	411	1,023	2,281	4,087	7,626	22,008	45,534	-
2006 1/.....	83,178	92	48	66	199	413	1,059	2,330	4,099	7,611	22,278	44,982	1
2005 1/.....	83,294	101	34	64	204	391	1,033	2,220	4,020	7,927	22,848	44,451	1
2004 1/.....	81,383	128	49	70	198	393	1,046	2,193	3,791	8,000	22,962	42,552	1
2003 1/.....	83,154	116	61	60	228	433	1,123	2,207	3,933	8,342	23,618	43,032	1
2002.....	83,349	146	47	71	174	402	1,152	2,113	3,878	8,756	23,985	42,624	1
2001.....	82,837	137	63	83	193	443	1,049	2,116	3,796	8,925	23,549	42,482	1
2000.....	81,828	151	43	58	205	431	1,101	1,991	3,639	9,056	23,514	41,637	2
1999.....	81,732	155	49	80	243	442	1,106	1,961	3,639	9,236	23,661	41,159	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other heart diseases (I26-I51)													
Black, both sexes													
2007 1/.....	20,352	136	41	60	263	559	1,209	2,470	3,108	3,604	4,448	4,454	-
2006 1/.....	19,805	109	42	60	253	637	1,222	2,444	2,984	3,396	4,437	4,218	3
2005 1/.....	20,093	101	44	74	270	629	1,258	2,501	2,992	3,449	4,526	4,248	1
2004 1/.....	19,604	98	60	71	241	552	1,274	2,467	2,786	3,387	4,478	4,187	3
2003 1/.....	20,034	129	51	66	279	594	1,359	2,395	2,821	3,686	4,409	4,244	1
2002.....	19,812	136	38	57	253	615	1,322	2,390	2,693	3,503	4,524	4,278	3
2001.....	19,637	131	71	65	254	571	1,377	2,285	2,624	3,555	4,542	4,161	1
2000.....	19,298	117	48	73	280	564	1,327	2,182	2,506	3,643	4,382	4,175	1
1999.....	19,685	135	49	66	263	577	1,409	2,128	2,702	3,780	4,488	4,086	2
Black, male													
2007 1/.....	9,736	78	27	33	177	320	748	1,441	1,834	1,908	1,915	1,255	-
2006 1/.....	9,467	59	25	33	146	386	726	1,484	1,726	1,789	1,857	1,233	3
2005 1/.....	9,438	60	18	36	184	376	735	1,465	1,710	1,809	1,874	1,171	-
2004 1/.....	9,280	53	31	34	154	322	719	1,449	1,624	1,715	1,943	1,233	3
2003 1/.....	9,372	78	28	38	180	325	775	1,414	1,630	1,876	1,856	1,171	1
2002.....	9,128	72	16	30	158	368	747	1,415	1,505	1,788	1,873	1,155	1
2001.....	9,148	59	40	32	162	330	773	1,293	1,494	1,785	1,991	1,189	-
2000.....	8,958	59	24	40	159	315	759	1,302	1,423	1,818	1,843	1,216	-
1999.....	9,344	67	27	38	167	331	805	1,260	1,563	1,957	1,887	1,241	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other heart diseases (I26-I51)													
Black, female													
2007 1/.....	10,616	58	14	27	86	239	461	1,029	1,274	1,696	2,533	3,199	-
2006 1/.....	10,338	50	17	27	107	251	496	960	1,258	1,607	2,580	2,985	-
2005 1/.....	10,655	41	26	38	86	253	523	1,036	1,282	1,640	2,652	3,077	1
2004 1/.....	10,324	45	29	37	87	230	555	1,018	1,162	1,672	2,535	2,954	-
2003 1/.....	10,662	51	23	28	99	269	584	981	1,191	1,810	2,553	3,073	-
2002.....	10,684	64	22	27	95	247	575	975	1,188	1,715	2,651	3,123	2
2001.....	10,489	72	31	33	92	241	604	992	1,130	1,770	2,551	2,972	1
2000.....	10,340	58	24	33	121	249	568	880	1,083	1,825	2,539	2,959	1
1999.....	10,341	68	22	28	96	246	604	868	1,139	1,823	2,601	2,845	1
American Indian 2/, both sexes													
2007 1/.....	711	8	5	1	11	35	44	81	108	122	160	136	-
2006 1/.....	686	7	-	2	12	16	58	83	107	123	149	129	-
2005 1/.....	712	5	1	2	12	21	43	88	100	138	151	151	-
2004 1/.....	618	10	1	2	12	23	41	68	90	109	134	128	-
2003 1/.....	676	9	4	2	9	21	44	72	122	119	141	133	-
2002.....	594	4	6	5	10	17	36	65	74	97	146	134	-
2001.....	599	9	3	3	7	16	52	62	76	109	140	122	-
2000.....	591	6	1	5	7	13	47	60	74	118	138	122	-
1999.....	595	5	5	3	8	12	43	61	79	115	141	123	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other heart diseases (I26-I51)													
American Indian 2/, male													
2007 1/.....	382	5	4	1	4	25	29	57	62	69	80	46	-
2006 1/.....	356	3	-	-	4	9	41	56	65	71	65	42	-
2005 1/.....	361	1	1	-	7	15	27	50	59	81	64	56	-
2004 1/.....	307	6	-	1	4	16	26	35	55	61	54	49	-
2003 1/.....	333	5	1	-	6	16	27	44	66	58	53	57	-
2002.....	328	2	2	4	8	11	28	46	46	58	71	52	-
2001.....	318	3	2	1	5	9	33	44	45	63	74	39	-
2000.....	295	3	-	3	5	7	27	37	44	70	65	34	-
1999.....	289	2	2	1	5	5	24	35	44	68	62	41	-
American Indian 2/, female													
2007 1/.....	329	3	1	-	7	10	15	24	46	53	80	90	-
2006 1/.....	330	4	-	2	8	7	17	27	42	52	84	87	-
2005 1/.....	351	4	-	2	5	6	16	38	41	57	87	95	-
2004 1/.....	311	4	1	1	8	7	15	33	35	48	80	79	-
2003 1/.....	343	4	3	2	3	5	17	28	56	61	88	76	-
2002.....	266	2	4	1	2	6	8	19	28	39	75	82	-
2001.....	281	6	1	2	2	7	19	18	31	46	66	83	-
2000.....	296	3	1	2	2	6	20	23	30	48	73	88	-
1999.....	306	3	3	2	3	7	19	26	35	47	79	82	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other heart diseases (I26-I51)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	2,385	20	11	10	16	47	91	114	211	337	674	854	-
2006 1/.....	2,336	12	7	11	21	50	68	155	218	364	633	797	-
2005 1/.....	2,306	17	11	14	24	46	77	141	261	352	598	765	-
2004 1/.....	2,193	22	6	4	25	56	69	140	187	326	624	734	-
2003 1/.....	2,147	22	7	11	26	53	74	118	179	338	636	683	-
2002.....	2,087	19	4	8	25	52	74	128	198	364	588	627	-
2001.....	1,851	17	7	7	21	30	71	123	167	340	536	532	-
2000.....	1,778	19	5	12	28	39	77	108	184	301	468	537	-
1999.....	1,762	21	4	4	14	36	68	117	171	303	505	519	-

Asian or Pacific Islander 3/, male

2007 1/.....	1,188	10	2	6	14	30	66	77	134	187	322	340	-
2006 1/.....	1,155	7	2	5	17	35	45	101	137	200	290	316	-
2005 1/.....	1,115	9	5	5	15	25	49	84	158	200	273	292	-
2004 1/.....	1,132	11	1	2	18	42	43	93	112	174	308	328	-
2003 1/.....	1,060	10	4	8	12	31	44	76	103	182	315	275	-
2002.....	1,076	7	2	5	17	37	49	74	117	185	294	289	-
2001.....	948	5	4	2	14	22	48	72	92	192	266	231	-
2000.....	936	12	3	7	21	25	55	66	106	172	233	236	-
1999.....	914	12	1	3	9	19	44	63	104	157	262	240	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other heart diseases (I26-I51)

Asian or Pacific Islander 3/, female

2007 1/.....	1,197	10	9	4	2	17	25	37	77	150	352	514	-
2006 1/.....	1,181	5	5	6	4	15	23	54	81	164	343	481	-
2005 1/.....	1,191	8	6	9	9	21	28	57	103	152	325	473	-
2004 1/.....	1,061	11	5	2	7	14	26	47	75	152	316	406	-
2003 1/.....	1,087	12	3	3	14	22	30	42	76	156	321	408	-
2002.....	1,011	12	2	3	8	15	25	54	81	179	294	338	-
2001.....	903	12	3	5	7	8	23	51	75	148	270	301	-
2000.....	842	7	2	5	7	14	22	42	78	129	235	301	-
1999.....	848	9	3	1	5	17	24	54	67	146	243	279	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute and subacute endocarditis (I33)													
All races, both sexes													
2007.....	1,225	3	1	1	9	32	95	198	207	236	308	134	1
2006.....	1,216	-	1	2	10	33	85	203	228	219	319	116	-
2005.....	1,209	4	-	3	8	28	78	173	198	234	328	155	-
2004.....	1,196	1	1	-	12	35	84	190	165	234	318	155	1
2003.....	1,245	1	-	1	16	33	112	183	209	232	318	140	-
2002.....	1,154	2	2	2	7	44	90	174	168	238	306	121	-
2001.....	1,178	1	-	3	8	29	117	186	160	247	316	111	-
2000.....	1,172	3	1	3	13	40	116	186	165	224	302	119	-
1999.....	1,116	2	1	-	7	30	114	143	171	240	289	119	-
All races, male													
2007.....	718	2	1	1	4	28	65	124	132	127	172	61	1
2006.....	660	-	-	1	4	18	51	140	142	120	145	39	-
2005.....	695	3	-	-	5	20	51	113	125	124	173	81	-
2004.....	659	-	-	-	8	22	55	122	98	119	162	72	1
2003.....	682	1	-	1	6	19	63	125	123	119	166	59	-
2002.....	643	1	1	1	6	31	60	117	91	133	153	49	-
2001.....	668	1	-	1	5	17	62	129	95	132	177	49	-
2000.....	665	2	-	1	7	24	73	120	96	122	169	51	-
1999.....	639	2	1	-	3	24	75	103	103	118	154	56	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute and subacute endocarditis (I33)													
All races, female													
2007.....	507	1	-	-	5	4	30	74	75	109	136	73	-
2006.....	556	-	1	1	6	15	34	63	86	99	174	77	-
2005.....	514	1	-	3	3	8	27	60	73	110	155	74	-
2004.....	537	1	1	-	4	13	29	68	67	115	156	83	-
2003.....	563	-	-	-	10	14	49	58	86	113	152	81	-
2002.....	511	1	1	1	1	13	30	57	77	105	153	72	-
2001.....	510	-	-	2	3	12	55	57	65	115	139	62	-
2000.....	507	1	1	2	6	16	43	66	69	102	133	68	-
1999.....	477	-	-	-	4	6	39	40	68	122	135	63	-
White, both sexes													
2007 1/.....	967	2	-	1	9	23	70	132	151	181	277	120	1
2006 1/.....	958	-	1	2	9	22	62	140	159	184	278	101	-
2005 1/.....	949	3	-	1	6	20	46	112	135	197	286	143	-
2004 1/.....	957	1	-	-	10	27	66	114	126	194	279	140	-
2003 1/.....	990	-	-	1	14	25	78	118	149	187	284	134	-
2002.....	930	1	2	-	6	34	66	109	125	192	283	112	-
2001.....	917	1	-	3	6	21	74	125	117	191	274	105	-
2000.....	927	1	1	2	6	27	80	122	122	191	270	105	-
1999.....	879	1	1	-	6	24	69	89	117	196	264	112	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute and subacute endocarditis (I33)													
White, male													
2007 1/.....	554	1	-	1	4	20	50	78	94	96	154	55	1
2006 1/.....	518	-	-	1	4	12	36	96	101	102	128	38	-
2005 1/.....	553	2	-	-	4	14	31	74	84	108	159	77	-
2004 1/.....	536	-	-	-	6	16	47	75	79	101	145	67	-
2003 1/.....	541	-	-	1	6	14	46	77	91	97	151	58	-
2002.....	524	-	1	-	5	26	45	79	68	110	145	45	-
2001.....	517	1	-	1	4	12	41	87	70	98	157	46	-
2000.....	536	1	-	-	3	19	50	83	76	104	153	47	-
1999.....	509	1	1	-	2	18	45	65	73	102	147	55	-
White, female													
2007 1/.....	413	1	-	-	5	3	20	54	57	85	123	65	-
2006 1/.....	440	-	1	1	5	10	26	44	58	82	150	63	-
2005 1/.....	396	1	-	1	2	6	15	38	51	89	127	66	-
2004 1/.....	421	1	-	-	4	11	19	39	47	93	134	73	-
2003 1/.....	449	-	-	-	8	11	32	41	58	90	133	76	-
2002.....	406	1	1	-	1	8	21	30	57	82	138	67	-
2001.....	400	-	-	2	2	9	33	38	47	93	117	59	-
2000.....	391	-	1	2	3	8	30	39	46	87	117	58	-
1999.....	370	-	-	-	4	6	24	24	44	94	117	57	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute and subacute endocarditis (I33)													
Black, both sexes													
2007 1/.....	223	1	1	-	-	5	23	58	50	50	24	11	-
2006 1/.....	228	-	-	-	1	11	20	56	62	30	35	13	-
2005 1/.....	229	1	-	2	2	6	28	55	58	31	36	10	-
2004 1/.....	204	-	1	-	1	6	14	68	35	36	27	15	1
2003 1/.....	225	1	-	-	2	5	32	61	53	38	29	4	-
2002.....	199	-	-	1	1	10	23	57	41	39	19	8	-
2001.....	235	-	-	-	1	8	41	57	41	43	39	5	-
2000.....	214	2	-	-	6	11	31	57	39	31	26	11	-
1999.....	210	-	-	-	1	6	41	49	48	42	17	6	-
Black, male													
2007 1/.....	144	1	1	-	-	5	14	41	35	29	13	5	-
2006 1/.....	122	-	-	-	-	6	12	37	37	15	15	-	-
2005 1/.....	124	1	-	-	1	5	19	34	37	13	11	3	-
2004 1/.....	104	-	-	-	1	4	7	42	18	16	10	5	1
2003 1/.....	122	1	-	-	-	4	15	44	27	18	13	-	-
2002.....	106	-	-	-	1	5	15	33	22	20	6	4	-
2001.....	135	-	-	-	-	5	20	39	24	26	19	2	-
2000.....	111	1	-	-	4	3	21	35	18	16	11	2	-
1999.....	117	-	-	-	1	6	27	35	27	15	5	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Acute and subacute endocarditis (I33)

Black, female

2007 1/.....	79	-	-	-	-	-	9	17	15	21	11	6	-
2006 1/.....	106	-	-	-	1	5	8	19	25	15	20	13	-
2005 1/.....	105	-	-	2	1	1	9	21	21	18	25	7	-
2004 1/.....	100	-	1	-	-	2	7	26	17	20	17	10	-
2003 1/.....	103	-	-	-	2	1	17	17	26	20	16	4	-
2002.....	93	-	-	1	-	5	8	24	19	19	13	4	-
2001.....	100	-	-	-	1	3	21	18	17	17	20	3	-
2000.....	103	1	-	-	2	8	10	22	21	15	15	9	-
1999.....	93	-	-	-	-	-	14	14	21	27	12	5	-

American Indian 2/, both sexes

2007 1/.....	14	-	-	-	-	3	-	5	3	1	2	-	-
2006 1/.....	9	-	-	-	-	-	2	3	1	2	1	-	-
2005 1/.....	12	-	-	-	-	-	2	5	2	3	-	-	-
2004 1/.....	13	-	-	-	-	1	2	5	2	1	2	-	-
2003 1/.....	12	-	-	-	-	2	2	2	2	2	1	1	-
2002.....	6	-	-	-	-	-	-	3	1	-	2	-	-
2001.....	6	-	-	-	1	-	-	3	-	2	-	-	-
2000.....	13	-	-	-	-	-	3	5	2	-	2	1	-
1999.....	7	-	-	-	-	-	3	1	1	-	2	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Acute and subacute endocarditis (I33)

American Indian 2/, male

2007 1/.....	9	-	-	-	-	3	-	3	2	-	1	-	-
2006 1/.....	6	-	-	-	-	-	2	3	-	1	-	-	-
2005 1/.....	6	-	-	-	-	-	-	4	1	1	-	-	-
2004 1/.....	5	-	-	-	-	1	-	3	-	-	1	-	-
2003 1/.....	7	-	-	-	-	1	2	2	-	-	1	1	-
2002.....	2	-	-	-	-	-	-	1	-	-	1	-	-
2001.....	5	-	-	-	1	-	-	2	-	2	-	-	-
2000.....	7	-	-	-	-	-	2	2	1	-	2	-	-
1999.....	4	-	-	-	-	-	2	1	-	-	1	-	-

American Indian 2/, female

2007 1/.....	5	-	-	-	-	-	-	2	1	1	1	-	-
2006 1/.....	3	-	-	-	-	-	-	-	1	1	1	-	-
2005 1/.....	6	-	-	-	-	-	2	1	1	2	-	-	-
2004 1/.....	8	-	-	-	-	-	2	2	2	1	1	-	-
2003 1/.....	5	-	-	-	-	1	-	-	2	2	-	-	-
2002.....	4	-	-	-	-	-	-	2	1	-	1	-	-
2001.....	1	-	-	-	-	-	-	1	-	-	-	-	-
2000.....	6	-	-	-	-	-	1	3	1	-	-	1	-
1999.....	3	-	-	-	-	-	1	-	1	-	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Acute and subacute endocarditis (I33)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	21	-	-	-	-	1	2	3	3	4	5	3	-
2006 1/.....	21	-	-	-	-	-	1	4	6	3	5	2	-
2005 1/.....	19	-	-	-	-	2	2	1	3	3	6	2	-
2004 1/.....	22	-	-	-	1	1	2	3	2	3	10	-	-
2003 1/.....	18	-	-	-	-	1	-	2	5	5	4	1	-
2002.....	19	1	-	1	-	-	1	5	1	7	2	1	-
2001.....	20	-	-	-	-	-	2	1	2	11	3	1	-
2000.....	18	-	-	1	1	2	2	2	2	2	4	2	-
1999.....	20	1	-	-	-	-	1	4	5	2	6	1	-

Asian or Pacific Islander 3/, male

2007 1/.....	11	-	-	-	-	-	1	2	1	2	4	1	-
2006 1/.....	14	-	-	-	-	-	1	4	4	2	2	1	-
2005 1/.....	12	-	-	-	-	1	1	1	3	2	3	1	-
2004 1/.....	14	-	-	-	1	1	1	2	1	2	6	-	-
2003 1/.....	12	-	-	-	-	-	-	2	5	4	1	-	-
2002.....	11	1	-	1	-	-	-	4	1	3	1	-	-
2001.....	11	-	-	-	-	-	1	1	1	6	1	1	-
2000.....	11	-	-	1	-	2	-	-	1	2	3	2	-
1999.....	9	1	-	-	-	-	1	2	3	1	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Acute and subacute endocarditis (I33)

Asian or Pacific Islander 3/, female

2007 1/.....	10	-	-	-	-	1	1	1	2	2	1	2	-
2006 1/.....	7	-	-	-	-	-	-	-	2	1	3	1	-
2005 1/.....	7	-	-	-	-	1	1	-	-	1	3	1	-
2004 1/.....	8	-	-	-	-	-	1	1	1	1	4	-	-
2003 1/.....	6	-	-	-	-	1	-	-	-	1	3	1	-
2002.....	8	-	-	-	-	-	1	1	-	4	1	1	-
2001.....	9	-	-	-	-	-	1	-	1	5	2	-	-
2000.....	7	-	-	-	1	-	2	2	1	-	1	-	-
1999.....	11	-	-	-	-	-	-	2	2	1	5	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)													
All races, both sexes													
2007.....	867	25	19	24	48	53	87	116	123	119	137	116	-
2006.....	816	18	24	21	36	48	78	107	122	116	151	95	-
2005.....	864	21	17	30	34	53	84	132	110	132	155	96	-
2004.....	779	21	25	17	31	39	67	107	111	115	159	87	-
2003.....	868	33	22	28	55	48	86	111	104	151	153	77	-
2002.....	848	26	22	22	36	46	75	107	118	143	157	96	-
2001.....	839	32	26	21	40	50	93	111	106	120	154	85	1
2000.....	767	20	24	20	38	53	86	85	89	131	145	76	-
1999.....	890	27	25	21	32	50	114	105	125	146	160	85	-
All races, male													
2007.....	454	16	13	17	26	32	57	57	71	66	59	40	-
2006.....	431	12	14	8	19	30	45	68	71	63	71	30	-
2005.....	439	9	10	9	21	31	46	77	57	71	78	30	-
2004.....	404	11	15	10	18	18	40	63	65	62	68	34	-
2003.....	449	15	8	12	34	29	52	71	57	84	65	22	-
2002.....	434	9	11	7	24	32	49	69	69	71	60	33	-
2001.....	401	18	15	7	22	33	41	59	61	59	61	24	1
2000.....	404	10	13	9	21	29	58	46	52	67	73	26	-
1999.....	442	13	14	7	20	34	64	56	70	69	65	30	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Diseases of pericardium and acute myocarditis (I30-I31,I40)

All races, female

2007.....	413	9	6	7	22	21	30	59	52	53	78	76	-
2006.....	385	6	10	13	17	18	33	39	51	53	80	65	-
2005.....	425	12	7	21	13	22	38	55	53	61	77	66	-
2004.....	375	10	10	7	13	21	27	44	46	53	91	53	-
2003.....	419	18	14	16	21	19	34	40	47	67	88	55	-
2002.....	414	17	11	15	12	14	26	38	49	72	97	63	-
2001.....	438	14	11	14	18	17	52	52	45	61	93	61	-
2000.....	363	10	11	11	17	24	28	39	37	64	72	50	-
1999.....	448	14	11	14	12	16	50	49	55	77	95	55	-

White, both sexes

2007 1/.....	698	13	14	18	37	41	64	88	95	99	127	102	-
2006 1/.....	675	11	19	12	27	38	58	82	103	99	138	88	-
2005 1/.....	661	8	4	15	26	32	61	102	82	114	131	86	-
2004 1/.....	595	11	12	9	18	23	40	79	94	96	135	78	-
2003 1/.....	677	15	13	17	38	24	55	84	90	133	136	72	-
2002.....	666	13	13	12	26	33	55	79	91	123	131	90	-
2001.....	658	14	14	16	27	38	60	81	85	104	141	77	1
2000.....	590	11	15	10	25	34	58	59	72	108	130	68	-
1999.....	703	12	12	13	24	35	79	82	106	122	142	76	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)													
White, male													
2007 1/.....	366	8	10	12	19	28	43	43	57	56	54	36	-
2006 1/.....	364	7	11	6	15	25	32	53	64	55	68	28	-
2005 1/.....	344	3	4	6	18	18	33	63	42	63	66	28	-
2004 1/.....	305	7	7	5	14	10	26	43	56	48	59	30	-
2003 1/.....	359	9	4	7	25	13	36	56	49	77	61	22	-
2002.....	344	6	7	3	17	24	36	50	55	63	52	31	-
2001.....	310	10	8	6	12	25	29	42	48	51	55	23	1
2000.....	309	6	11	5	13	18	39	32	45	52	65	23	-
1999.....	351	7	5	6	15	24	47	42	61	57	60	27	-
White, female													
2007 1/.....	332	5	4	6	18	13	21	45	38	43	73	66	-
2006 1/.....	311	4	8	6	12	13	26	29	39	44	70	60	-
2005 1/.....	317	5	-	9	8	14	28	39	40	51	65	58	-
2004 1/.....	290	4	5	4	4	13	14	36	38	48	76	48	-
2003 1/.....	318	6	9	10	13	11	19	28	41	56	75	50	-
2002.....	322	7	6	9	9	9	19	29	36	60	79	59	-
2001.....	348	4	6	10	15	13	31	39	37	53	86	54	-
2000.....	281	5	4	5	12	16	19	27	27	56	65	45	-
1999.....	352	5	7	7	9	11	32	40	45	65	82	49	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Diseases of pericardium and acute myocarditis (I30-I31,I40)

Black, both sexes

2007 1/.....	145	11	3	4	10	11	19	26	26	17	8	10	-
2006 1/.....	109	5	4	4	8	9	14	22	15	13	11	4	-
2005 1/.....	164	11	11	9	5	15	17	28	25	14	21	8	-
2004 1/.....	164	8	12	8	10	12	26	27	16	17	22	6	-
2003 1/.....	168	14	9	9	15	22	30	23	12	16	13	5	-
2002.....	154	12	7	8	8	10	17	26	22	19	20	5	-
2001.....	158	14	12	4	13	10	25	30	19	13	11	7	-
2000.....	142	7	8	8	8	16	24	21	15	18	10	7	-
1999.....	160	10	12	8	6	14	31	20	17	20	16	6	-

Black, male

2007 1/.....	74	7	2	4	7	3	12	12	13	8	4	2	-
2006 1/.....	55	3	3	1	3	4	10	14	6	8	2	1	-
2005 1/.....	73	5	5	1	3	10	7	13	12	5	11	1	-
2004 1/.....	89	3	7	5	4	6	14	19	8	13	9	1	-
2003 1/.....	81	4	4	5	9	15	16	13	7	5	3	-	-
2002.....	75	3	3	3	5	6	10	17	13	7	7	1	-
2001.....	79	5	7	1	10	7	8	17	11	8	5	-	-
2000.....	72	4	2	3	3	10	16	10	6	11	5	2	-
1999.....	77	4	9	1	3	10	14	12	9	11	4	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Diseases of pericardium and acute myocarditis (I30-I31,I40)

Black, female

2007 1/.....	71	4	1	-	3	8	7	14	13	9	4	8	-
2006 1/.....	54	2	1	3	5	5	4	8	9	5	9	3	-
2005 1/.....	91	6	6	8	2	5	10	15	13	9	10	7	-
2004 1/.....	75	5	5	3	6	6	12	8	8	4	13	5	-
2003 1/.....	87	10	5	4	6	7	14	10	5	11	10	5	-
2002.....	79	9	4	5	3	4	7	9	9	12	13	4	-
2001.....	79	9	5	3	3	3	17	13	8	5	6	7	-
2000.....	70	3	6	5	5	6	8	11	9	7	5	5	-
1999.....	83	6	3	7	3	4	17	8	8	9	12	6	-

American Indian 2/, both sexes

2007 1/.....	4	-	1	-	1	-	1	1	-	-	-	-	-
2006 1/.....	10	-	-	1	-	1	3	2	1	1	1	-	-
2005 1/.....	3	-	-	-	-	2	-	1	-	-	-	-	-
2004 1/.....	5	1	-	-	1	1	1	-	-	-	-	1	-
2003 1/.....	6	3	-	-	-	-	-	1	1	1	-	-	-
2002.....	8	-	1	1	1	-	1	2	-	-	2	-	-
2001.....	8	3	-	1	-	-	3	-	1	-	-	-	-
2000.....	6	1	-	-	1	-	-	1	-	1	2	-	-
1999.....	9	1	1	-	1	1	1	1	1	2	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Diseases of pericardium and acute myocarditis (I30-I31,I40)

American Indian 2/, male

2007 1/.....	3	-	1	-	-	-	1	1	-	-	-	-	-
2006 1/.....	4	-	-	-	-	1	2	1	-	-	-	-	-
2005 1/.....	3	-	-	-	-	2	-	1	-	-	-	-	-
2004 1/.....	1	-	-	-	-	-	-	-	-	-	-	1	-
2003 1/.....	5	2	-	-	-	-	-	1	1	1	-	-	-
2002.....	5	-	-	1	1	-	1	2	-	-	-	-	-
2001.....	5	2	-	-	-	-	2	-	1	-	-	-	-
2000.....	4	-	-	-	1	-	-	1	-	1	1	-	-
1999.....	2	-	-	-	1	-	-	1	-	-	-	-	-

American Indian 2/, female

2007 1/.....	1	-	-	-	1	-	-	-	-	-	-	-	-
2006 1/.....	6	-	-	1	-	-	1	1	1	1	1	-	-
2005 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 1/.....	4	1	-	-	1	1	1	-	-	-	-	-	-
2003 1/.....	1	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	3	-	1	-	-	-	-	-	-	-	2	-	-
2001.....	3	1	-	1	-	-	1	-	-	-	-	-	-
2000.....	2	1	-	-	-	-	-	-	-	-	1	-	-
1999.....	7	1	1	-	-	1	1	-	1	2	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Diseases of pericardium and acute myocarditis (I30-I31,I40)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	20	1	1	2	-	1	3	1	2	3	2	4	-
2006 1/.....	22	2	1	4	1	-	3	1	3	3	1	3	-
2005 1/.....	36	2	2	6	3	4	6	1	3	4	3	2	-
2004 1/.....	15	1	1	-	2	3	-	1	1	2	2	2	-
2003 1/.....	17	1	-	2	2	2	1	3	1	1	4	-	-
2002.....	20	1	1	1	1	3	2	-	5	1	4	1	-
2001.....	15	1	-	-	-	2	5	-	1	3	2	1	-
2000.....	29	1	1	2	4	3	4	4	2	4	3	1	-
1999.....	18	4	-	-	1	-	3	2	1	2	2	3	-

Asian or Pacific Islander 3/, male

2007 1/.....	11	1	-	1	-	1	1	1	1	2	1	2	-
2006 1/.....	8	2	-	1	1	-	1	-	1	-	1	1	-
2005 1/.....	19	1	1	2	-	1	6	-	3	3	1	1	-
2004 1/.....	9	1	1	-	-	2	-	1	1	1	-	2	-
2003 1/.....	4	-	-	-	-	1	-	1	-	1	1	-	-
2002.....	10	-	1	-	1	2	2	-	1	1	1	1	-
2001.....	7	1	-	-	-	1	2	-	1	-	1	1	-
2000.....	19	-	-	1	4	1	3	3	1	3	2	1	-
1999.....	12	2	-	-	1	-	3	1	-	1	1	3	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Diseases of pericardium and acute myocarditis (I30-I31,I40)

Asian or Pacific Islander 3/, female

2007 1/.....	9	-	1	1	-	-	2	-	1	1	1	2	-
2006 1/.....	14	-	1	3	-	-	2	1	2	3	-	2	-
2005 1/.....	17	1	1	4	3	3	-	1	-	1	2	1	-
2004 1/.....	6	-	-	-	2	1	-	-	-	1	2	-	-
2003 1/.....	13	1	-	2	2	1	1	2	1	-	3	-	-
2002.....	10	1	-	1	-	1	-	-	4	-	3	-	-
2001.....	8	-	-	-	-	1	3	-	-	3	1	-	-
2000.....	10	1	1	1	-	2	1	1	1	1	1	-	-
1999.....	6	2	-	-	-	-	-	1	1	1	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Heart failure (I50)													
All races, both sexes													
2007.....	56,565	21	11	12	43	87	317	1,073	2,758	5,749	15,935	30,558	1
2006.....	60,337	19	9	12	37	86	332	1,225	2,801	5,960	17,309	32,544	3
2005.....	58,933	27	6	11	42	94	316	1,087	2,610	5,888	17,331	31,521	-
2004.....	57,120	30	11	10	33	90	313	1,039	2,444	5,951	17,012	30,186	1
2003.....	57,448	38	5	12	38	97	315	981	2,469	6,138	17,136	30,219	-
2002.....	56,494	23	11	10	27	101	282	993	2,342	6,168	16,910	29,627	-
2001.....	56,934	24	17	15	34	92	307	954	2,374	6,274	17,134	29,708	1
2000.....	55,704	31	9	21	34	63	262	811	2,211	6,403	16,820	29,035	4
1999.....	54,913	11	10	12	31	65	272	782	2,107	6,524	16,771	28,327	1
All races, male													
2007.....	22,914	11	3	6	25	54	207	655	1,643	3,201	7,286	9,822	1
2006.....	23,918	11	5	5	22	56	199	780	1,648	3,202	7,875	10,113	2
2005.....	23,026	14	4	5	27	61	207	653	1,509	3,069	7,804	9,673	-
2004.....	22,292	16	4	7	19	59	186	615	1,420	3,163	7,673	9,130	-
2003.....	22,427	28	3	7	24	60	193	587	1,454	3,313	7,658	9,100	-
2002.....	21,698	14	6	5	15	60	173	617	1,345	3,254	7,573	8,636	-
2001.....	21,632	16	5	8	16	56	182	588	1,347	3,344	7,672	8,398	-
2000.....	21,175	22	7	9	20	36	159	496	1,264	3,414	7,498	8,248	2
1999.....	20,755	5	7	6	23	32	164	486	1,228	3,526	7,318	7,959	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Heart failure (I50)													
All races, female													
2007.....	33,651	10	8	6	18	33	110	418	1,115	2,548	8,649	20,736	-
2006.....	36,419	8	4	7	15	30	133	445	1,153	2,758	9,434	22,431	1
2005.....	35,907	13	2	6	15	33	109	434	1,101	2,819	9,527	21,848	-
2004.....	34,828	14	7	3	14	31	127	424	1,024	2,788	9,339	21,056	1
2003.....	35,021	10	2	5	14	37	122	394	1,015	2,825	9,478	21,119	-
2002.....	34,796	9	5	5	12	41	109	376	997	2,914	9,337	20,991	-
2001.....	35,302	8	12	7	18	36	125	366	1,027	2,930	9,462	21,310	1
2000.....	34,529	9	2	12	14	27	103	315	947	2,989	9,322	20,787	2
1999.....	34,158	6	3	6	8	33	108	296	879	2,998	9,453	20,368	-
White, both sexes													
2007 1/.....	50,367	10	8	9	24	47	176	680	2,030	4,732	14,245	28,405	1
2006 1/.....	53,980	8	8	8	18	38	192	757	2,123	4,936	15,566	30,323	3
2005 1/.....	52,570	15	3	4	25	47	178	686	1,941	4,844	15,471	29,356	-
2004 1/.....	51,205	20	10	5	21	50	193	654	1,826	5,013	15,311	28,101	1
2003 1/.....	51,454	20	4	7	20	58	178	635	1,859	5,118	15,442	28,113	-
2002.....	50,672	15	8	7	16	53	174	639	1,812	5,160	15,199	27,589	-
2001.....	51,252	19	15	11	18	57	188	655	1,842	5,277	15,466	27,703	1
2000.....	50,142	19	5	16	17	38	164	558	1,715	5,384	15,216	27,006	4
1999.....	49,317	8	8	9	18	34	174	509	1,632	5,511	15,081	26,333	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Heart failure (I50)													
White, male													
2007 1/.....	20,262	6	1	4	11	29	113	427	1,215	2,682	6,574	9,199	1
2006 1/.....	21,236	4	4	3	10	26	109	490	1,284	2,674	7,164	9,466	2
2005 1/.....	20,414	8	3	1	18	32	111	421	1,135	2,546	7,047	9,092	-
2004 1/.....	19,806	10	4	4	13	30	120	395	1,063	2,706	6,947	8,514	-
2003 1/.....	19,987	13	2	4	12	38	116	390	1,112	2,820	6,976	8,504	-
2002.....	19,335	10	5	4	7	30	112	410	1,054	2,746	6,886	8,071	-
2001.....	19,332	12	5	5	10	31	118	425	1,076	2,849	6,962	7,839	-
2000.....	18,949	14	5	7	10	22	109	351	1,007	2,941	6,831	7,650	2
1999.....	18,469	3	5	5	14	17	111	322	972	3,038	6,643	7,339	-
White, female													
2007 1/.....	30,105	4	7	5	13	18	63	253	815	2,050	7,671	19,206	-
2006 1/.....	32,744	4	4	5	8	12	83	267	839	2,262	8,402	20,857	1
2005 1/.....	32,156	7	-	3	7	15	67	265	806	2,298	8,424	20,264	-
2004 1/.....	31,399	10	6	1	8	20	73	259	763	2,307	8,364	19,587	1
2003 1/.....	31,467	7	2	3	8	20	62	245	747	2,298	8,466	19,609	-
2002.....	31,337	5	3	3	9	23	62	229	758	2,414	8,313	19,518	-
2001.....	31,920	7	10	6	8	26	70	230	766	2,428	8,504	19,864	1
2000.....	31,193	5	-	9	7	16	55	207	708	2,443	8,385	19,356	2
1999.....	30,848	5	3	4	4	17	63	187	660	2,473	8,438	18,994	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Heart failure (I50)													
Black, both sexes													
2007 1/.....	5,497	8	3	2	18	38	134	378	679	920	1,473	1,844	-
2006 1/.....	5,564	11	1	4	17	44	128	434	627	930	1,499	1,869	-
2005 1/.....	5,609	8	3	6	16	46	131	372	618	946	1,631	1,832	-
2004 1/.....	5,230	8	1	4	11	36	111	368	564	835	1,496	1,796	-
2003 1/.....	5,330	15	-	5	18	37	131	328	569	928	1,476	1,823	-
2002.....	5,186	8	3	3	9	45	104	335	490	898	1,513	1,778	-
2001.....	5,122	5	2	3	16	34	112	285	498	910	1,486	1,771	-
2000.....	4,965	12	4	4	13	24	94	241	455	917	1,416	1,785	-
1999.....	4,998	3	2	3	12	30	94	260	451	927	1,481	1,734	1
Black, male													
2007 1/.....	2,341	4	2	1	14	23	87	220	402	467	611	510	-
2006 1/.....	2,335	7	1	2	12	27	78	267	337	476	609	519	-
2005 1/.....	2,275	4	1	3	9	29	90	216	344	465	659	455	-
2004 1/.....	2,174	5	-	3	6	25	59	211	323	402	633	507	-
2003 1/.....	2,150	13	-	3	12	20	73	183	324	449	581	492	-
2002.....	2,069	4	1	1	6	27	59	194	269	453	608	447	-
2001.....	2,052	4	-	2	6	24	61	153	251	451	623	477	-
2000.....	1,951	8	2	2	8	13	47	138	233	423	573	504	-
1999.....	2,021	2	2	1	8	15	51	157	243	443	583	515	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Heart failure (I50)													
Black, female													
2007 1/.....	3,156	4	1	1	4	15	47	158	277	453	862	1,334	-
2006 1/.....	3,229	4	-	2	5	17	50	167	290	454	890	1,350	-
2005 1/.....	3,334	4	2	3	7	17	41	156	274	481	972	1,377	-
2004 1/.....	3,056	3	1	1	5	11	52	157	241	433	863	1,289	-
2003 1/.....	3,180	2	-	2	6	17	58	145	245	479	895	1,331	-
2002.....	3,117	4	2	2	3	18	45	141	221	445	905	1,331	-
2001.....	3,070	1	2	1	10	10	51	132	247	459	863	1,294	-
2000.....	3,014	4	2	2	5	11	47	103	222	494	843	1,281	-
1999.....	2,977	1	-	2	4	15	43	103	208	484	898	1,219	-
American Indian 2/, both sexes													
2007 1/.....	188	1	-	-	1	1	2	7	27	37	50	62	-
2006 1/.....	218	-	-	-	2	1	9	15	22	40	65	64	-
2005 1/.....	228	2	-	-	1	-	3	15	23	44	72	68	-
2004 1/.....	193	-	-	1	1	2	5	8	21	34	56	65	-
2003 1/.....	180	1	-	-	-	1	2	9	24	30	56	57	-
2002.....	184	-	-	-	1	2	3	11	16	39	52	60	-
2001.....	164	-	-	-	-	-	3	8	12	33	51	57	-
2000.....	175	-	-	-	1	-	1	7	14	42	58	52	-
1999.....	170	-	-	-	1	-	4	7	7	34	58	59	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Heart failure (I50)													
American Indian 2/, male													
2007 1/.....	85	-	-	-	-	1	2	3	13	17	27	22	-
2006 1/.....	101	-	-	-	-	-	9	8	12	24	24	24	-
2005 1/.....	106	1	-	-	-	-	3	8	13	27	27	27	-
2004 1/.....	87	-	-	-	-	2	4	3	13	18	22	25	-
2003 1/.....	75	1	-	-	-	1	1	6	13	14	18	21	-
2002.....	92	-	-	-	1	2	2	9	11	22	22	23	-
2001.....	71	-	-	-	-	-	-	7	8	17	26	13	-
2000.....	79	-	-	-	1	-	1	4	6	23	29	15	-
1999.....	65	-	-	-	1	-	2	3	3	19	21	16	-
American Indian 2/, female													
2007 1/.....	103	1	-	-	1	-	-	4	14	20	23	40	-
2006 1/.....	117	-	-	-	2	1	-	7	10	16	41	40	-
2005 1/.....	122	1	-	-	1	-	-	7	10	17	45	41	-
2004 1/.....	106	-	-	1	1	-	1	5	8	16	34	40	-
2003 1/.....	105	-	-	-	-	-	1	3	11	16	38	36	-
2002.....	92	-	-	-	-	-	1	2	5	17	30	37	-
2001.....	93	-	-	-	-	-	3	1	4	16	25	44	-
2000.....	96	-	-	-	-	-	-	3	8	19	29	37	-
1999.....	105	-	-	-	-	-	2	4	4	15	37	43	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Heart failure (I50)													
Asian or Pacific Islander 3/, both sexes													
2007 1/.....	513	2	-	1	-	1	5	8	22	60	167	247	-
2006 1/.....	575	-	-	-	-	3	3	19	29	54	179	288	-
2005 1/.....	526	2	-	1	-	1	4	14	28	54	157	265	-
2004 1/.....	492	2	-	-	-	2	4	9	33	69	149	224	-
2003 1/.....	484	2	1	-	-	1	4	9	17	62	162	226	-
2002.....	452	-	-	-	1	1	1	8	24	71	146	200	-
2001.....	396	-	-	1	-	1	4	6	22	54	131	177	-
2000.....	422	-	-	1	3	1	3	5	27	60	130	192	-
1999.....	428	-	-	-	-	1	-	6	17	52	151	201	-
Asian or Pacific Islander 3/, male													
2007 1/.....	226	1	-	1	-	1	5	5	13	35	74	91	-
2006 1/.....	246	-	-	-	-	3	3	15	15	28	78	104	-
2005 1/.....	231	1	-	1	-	-	3	8	17	31	71	99	-
2004 1/.....	225	1	-	-	-	2	3	6	21	37	71	84	-
2003 1/.....	215	1	1	-	-	1	3	8	5	30	83	83	-
2002.....	202	-	-	-	1	1	-	4	11	33	57	95	-
2001.....	177	-	-	1	-	1	3	3	12	27	61	69	-
2000.....	196	-	-	-	1	1	2	3	18	27	65	79	-
1999.....	200	-	-	-	-	-	-	4	10	26	71	89	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Heart failure (I50)

Asian or Pacific Islander 3/, female

2007 1/.....	287	1	-	-	-	-	-	3	9	25	93	156	-
2006 1/.....	329	-	-	-	-	-	-	4	14	26	101	184	-
2005 1/.....	295	1	-	-	-	1	1	6	11	23	86	166	-
2004 1/.....	267	1	-	-	-	-	1	3	12	32	78	140	-
2003 1/.....	269	1	-	-	-	-	1	1	12	32	79	143	-
2002.....	250	-	-	-	-	-	1	4	13	38	89	105	-
2001.....	219	-	-	-	-	-	1	3	10	27	70	108	-
2000.....	226	-	-	1	2	-	1	2	9	33	65	113	-
1999.....	228	-	-	-	-	1	-	2	7	26	80	112	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)													
All races, both sexes													
2007.....	114,091	348	132	182	772	1,605	3,602	7,698	11,158	15,541	30,554	42,497	2
2006.....	107,879	283	120	196	790	1,736	3,640	7,629	10,879	14,945	29,286	38,371	4
2005.....	109,579	290	120	187	834	1,672	3,817	7,728	10,943	15,509	30,052	38,421	6
2004.....	107,180	354	137	194	779	1,555	3,820	7,616	10,414	15,605	30,364	36,337	5
2003.....	110,423	344	144	193	795	1,632	4,042	7,441	10,616	16,723	31,653	36,834	6
2002.....	111,044	422	122	193	766	1,617	4,173	7,437	10,318	17,131	32,403	36,456	6
2001.....	107,998	392	172	207	712	1,565	3,955	7,233	9,895	17,415	31,372	35,075	5
2000.....	107,785	403	134	204	768	1,593	3,993	7,005	9,897	17,921	31,100	34,763	4
1999.....	108,909	450	141	215	810	1,634	4,147	6,858	10,177	18,662	31,492	34,317	6
All races, male													
2007.....	53,092	196	60	97	494	986	2,248	4,878	6,916	8,726	14,444	14,045	2
2006.....	50,212	146	65	116	510	1,113	2,245	4,805	6,689	8,421	13,689	12,409	4
2005.....	50,934	162	63	104	561	1,064	2,391	4,926	6,724	8,723	13,899	12,313	4
2004.....	49,841	191	71	94	510	976	2,361	4,861	6,488	8,689	14,057	11,538	5
2003.....	51,180	189	70	121	496	973	2,493	4,675	6,508	9,359	14,791	11,500	5
2002.....	51,455	225	64	112	512	1,015	2,578	4,747	6,266	9,533	14,985	11,415	3
2001.....	49,738	187	97	107	457	931	2,492	4,531	6,000	9,632	14,630	10,670	4
2000.....	49,878	204	78	131	470	960	2,456	4,489	6,120	10,018	14,266	10,683	3
1999.....	50,765	235	78	124	487	977	2,591	4,334	6,299	10,607	14,591	10,438	4

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)													
All races, female													
2007.....	60,999	152	72	85	278	619	1,354	2,820	4,242	6,815	16,110	28,452	-
2006.....	57,667	137	55	80	280	623	1,395	2,824	4,190	6,524	15,597	25,962	-
2005.....	58,645	128	57	83	273	608	1,426	2,802	4,219	6,786	16,153	26,108	2
2004.....	57,339	163	66	100	269	579	1,459	2,755	3,926	6,916	16,307	24,799	-
2003.....	59,243	155	74	72	299	659	1,549	2,766	4,108	7,364	16,862	25,334	1
2002.....	59,589	197	58	81	254	602	1,595	2,690	4,052	7,598	17,418	25,041	3
2001.....	58,260	205	75	100	255	634	1,463	2,702	3,895	7,783	16,742	24,405	1
2000.....	57,907	199	56	73	298	633	1,537	2,516	3,777	7,903	16,834	24,080	1
1999.....	58,144	215	63	91	323	657	1,556	2,524	3,878	8,055	16,901	23,879	2
White, both sexes													
2007 1/.....	97,268	208	84	120	512	1,025	2,447	5,520	8,543	12,570	27,003	39,234	2
2006 1/.....	91,808	173	77	136	533	1,102	2,475	5,503	8,336	12,138	25,864	35,470	1
2005 1/.....	93,294	193	80	121	555	1,052	2,632	5,490	8,350	12,669	26,703	35,444	5
2004 1/.....	91,103	244	85	130	528	988	2,601	5,430	8,025	12,780	26,892	33,397	3
2003 1/.....	94,006	221	92	130	518	1,035	2,767	5,294	8,178	13,663	28,212	33,891	5
2002.....	94,779	285	86	138	500	1,004	2,893	5,301	7,953	14,241	28,905	33,470	3
2001.....	92,035	258	105	141	461	1,003	2,650	5,153	7,624	14,480	27,877	32,279	4
2000.....	92,102	284	93	130	490	1,034	2,704	4,998	7,689	14,934	27,763	31,980	3
1999.....	92,887	308	98	153	547	1,061	2,805	4,902	7,773	15,545	28,091	31,599	5

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

White, male

2007 1/.....	44,690	117	33	64	320	648	1,528	3,591	5,366	7,122	12,862	13,037	2
2006 1/.....	42,125	89	42	82	359	724	1,551	3,513	5,173	6,915	12,208	11,468	1
2005 1/.....	42,869	105	46	70	368	696	1,709	3,612	5,227	7,180	12,471	11,381	4
2004 1/.....	41,830	131	47	65	346	639	1,661	3,571	5,082	7,228	12,504	10,553	3
2003 1/.....	43,086	118	42	83	319	644	1,757	3,401	5,091	7,765	13,268	10,594	4
2002.....	43,495	152	49	79	345	642	1,843	3,476	4,926	8,041	13,450	10,490	2
2001.....	41,866	132	58	76	293	608	1,735	3,344	4,678	8,129	13,035	9,774	4
2000.....	42,139	143	55	88	307	643	1,707	3,280	4,831	8,464	12,816	9,802	3
1999.....	42,725	163	59	84	321	653	1,818	3,192	4,883	8,941	13,067	9,540	4

White, female

2007 1/.....	52,578	91	51	56	192	377	919	1,929	3,177	5,448	14,141	26,197	-
2006 1/.....	49,683	84	35	54	174	378	924	1,990	3,163	5,223	13,656	24,002	-
2005 1/.....	50,425	88	34	51	187	356	923	1,878	3,123	5,489	14,232	24,063	1
2004 1/.....	49,273	113	38	65	182	349	940	1,859	2,943	5,552	14,388	22,844	-
2003 1/.....	50,920	103	50	47	199	391	1,010	1,893	3,087	5,898	14,944	23,297	1
2002.....	51,284	133	37	59	155	362	1,050	1,825	3,027	6,200	15,455	22,980	1
2001.....	50,169	126	47	65	168	395	915	1,809	2,946	6,351	14,842	22,505	-
2000.....	49,963	141	38	42	183	391	997	1,718	2,858	6,470	14,947	22,178	-
1999.....	50,162	145	39	69	226	408	987	1,710	2,890	6,604	15,024	22,059	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)													
Black, both sexes													
2007 1/.....	14,487	116	34	54	235	505	1,033	2,008	2,353	2,617	2,943	2,589	-
2006 1/.....	13,904	93	37	52	227	573	1,060	1,932	2,280	2,423	2,892	2,332	3
2005 1/.....	14,091	81	30	57	247	562	1,082	2,046	2,291	2,458	2,838	2,398	1
2004 1/.....	14,006	82	46	59	219	498	1,123	2,004	2,171	2,499	2,933	2,370	2
2003 1/.....	14,311	99	42	52	244	530	1,166	1,983	2,187	2,704	2,891	2,412	1
2002.....	14,273	116	28	45	235	550	1,178	1,972	2,140	2,547	2,972	2,487	3
2001.....	14,122	112	57	58	224	519	1,199	1,913	2,066	2,589	3,006	2,378	1
2000.....	13,977	96	36	61	253	513	1,178	1,863	1,997	2,677	2,930	2,372	1
1999.....	14,317	122	35	55	244	527	1,243	1,799	2,186	2,791	2,974	2,340	1
Black, male													
2007 1/.....	7,177	66	22	28	156	289	635	1,168	1,384	1,404	1,287	738	-
2006 1/.....	6,955	49	21	30	131	349	626	1,166	1,346	1,290	1,231	713	3
2005 1/.....	6,966	50	12	32	171	332	619	1,202	1,317	1,326	1,193	712	-
2004 1/.....	6,913	45	24	26	143	287	639	1,177	1,275	1,284	1,291	720	2
2003 1/.....	7,019	60	24	30	159	286	671	1,174	1,272	1,404	1,259	679	1
2002.....	6,878	65	12	26	146	330	663	1,171	1,201	1,308	1,252	703	1
2001.....	6,882	50	33	29	146	294	684	1,084	1,208	1,300	1,344	710	-
2000.....	6,824	46	20	35	144	289	675	1,119	1,166	1,368	1,254	708	-
1999.....	7,129	61	16	36	155	300	713	1,056	1,284	1,488	1,295	725	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

Black, female

2007 1/.....	7,310	50	12	26	79	216	398	840	969	1,213	1,656	1,851	-
2006 1/.....	6,949	44	16	22	96	224	434	766	934	1,133	1,661	1,619	-
2005 1/.....	7,125	31	18	25	76	230	463	844	974	1,132	1,645	1,686	1
2004 1/.....	7,093	37	22	33	76	211	484	827	896	1,215	1,642	1,650	-
2003 1/.....	7,292	39	18	22	85	244	495	809	915	1,300	1,632	1,733	-
2002.....	7,395	51	16	19	89	220	515	801	939	1,239	1,720	1,784	2
2001.....	7,240	62	24	29	78	225	515	829	858	1,289	1,662	1,668	1
2000.....	7,153	50	16	26	109	224	503	744	831	1,309	1,676	1,664	1
1999.....	7,188	61	19	19	89	227	530	743	902	1,303	1,679	1,615	1

American Indian 2/, both sexes

2007 1/.....	505	7	4	1	9	31	41	68	78	84	108	74	-
2006 1/.....	449	7	-	1	10	14	44	63	83	80	82	65	-
2005 1/.....	469	3	1	2	11	19	38	67	75	91	79	83	-
2004 1/.....	407	9	1	1	10	19	33	55	67	74	76	62	-
2003 1/.....	478	5	4	2	9	18	40	60	95	86	84	75	-
2002.....	396	4	5	4	8	15	32	49	57	58	90	74	-
2001.....	421	6	3	2	6	16	46	51	63	74	89	65	-
2000.....	397	5	1	5	5	13	43	47	58	75	76	69	-
1999.....	409	4	4	3	6	11	35	52	70	79	81	64	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

American Indian 2/, male

2007 1/.....	285	5	3	1	4	21	26	50	47	52	52	24	-
2006 1/.....	245	3	-	-	4	8	28	44	53	46	41	18	-
2005 1/.....	246	-	1	-	7	13	24	37	45	53	37	29	-
2004 1/.....	214	6	-	1	4	13	22	29	42	43	31	23	-
2003 1/.....	246	2	1	-	6	14	24	35	52	43	34	35	-
2002.....	229	2	2	3	6	9	25	34	35	36	48	29	-
2001.....	237	1	2	1	4	9	31	35	36	44	48	26	-
2000.....	205	3	-	3	3	7	24	30	37	46	33	19	-
1999.....	218	2	2	1	3	5	20	30	41	49	40	25	-

American Indian 2/, female

2007 1/.....	220	2	1	-	5	10	15	18	31	32	56	50	-
2006 1/.....	204	4	-	1	6	6	16	19	30	34	41	47	-
2005 1/.....	223	3	-	2	4	6	14	30	30	38	42	54	-
2004 1/.....	193	3	1	-	6	6	11	26	25	31	45	39	-
2003 1/.....	232	3	3	2	3	4	16	25	43	43	50	40	-
2002.....	167	2	3	1	2	6	7	15	22	22	42	45	-
2001.....	184	5	1	1	2	7	15	16	27	30	41	39	-
2000.....	192	2	1	2	2	6	19	17	21	29	43	50	-
1999.....	191	2	2	2	3	6	15	22	29	30	41	39	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	1,831	17	10	7	16	44	81	102	184	270	500	600	-
2006 1/.....	1,718	10	6	7	20	47	61	131	180	304	448	504	-
2005 1/.....	1,725	13	9	7	21	39	65	125	227	291	432	496	-
2004 1/.....	1,664	19	5	4	22	50	63	127	151	252	463	508	-
2003 1/.....	1,628	19	6	9	24	49	69	104	156	270	466	456	-
2002.....	1,596	17	3	6	23	48	70	115	168	285	436	425	-
2001.....	1,420	16	7	6	21	27	60	116	142	272	400	353	-
2000.....	1,309	18	4	8	20	33	68	97	153	235	331	342	-
1999.....	1,296	16	4	4	13	35	64	105	148	247	346	314	-

Asian or Pacific Islander 3/, male

2007 1/.....	940	8	2	4	14	28	59	69	119	148	243	246	-
2006 1/.....	887	5	2	4	16	32	40	82	117	170	209	210	-
2005 1/.....	853	7	4	2	15	23	39	75	135	164	198	191	-
2004 1/.....	884	9	-	2	17	37	39	84	89	134	231	242	-
2003 1/.....	829	9	3	8	12	29	41	65	93	147	230	192	-
2002.....	853	6	1	4	15	34	47	66	104	148	235	193	-
2001.....	753	4	4	1	14	20	42	68	78	159	203	160	-
2000.....	710	12	3	5	16	21	50	60	86	140	163	154	-
1999.....	693	9	1	3	8	19	40	56	91	129	189	148	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

Asian or Pacific Islander 3/, female

2007 1/.....	891	9	8	3	2	16	22	33	65	122	257	354	-
2006 1/.....	831	5	4	3	4	15	21	49	63	134	239	294	-
2005 1/.....	872	6	5	5	6	16	26	50	92	127	234	305	-
2004 1/.....	780	10	5	2	5	13	24	43	62	118	232	266	-
2003 1/.....	799	10	3	1	12	20	28	39	63	123	236	264	-
2002.....	743	11	2	2	8	14	23	49	64	137	201	232	-
2001.....	667	12	3	5	7	7	18	48	64	113	197	193	-
2000.....	599	6	1	3	4	12	18	37	67	95	168	188	-
1999.....	603	7	3	1	5	16	24	49	57	118	157	166	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Essential hypertension and hypertensive renal disease (I10,I12,I15)

All races, both sexes

2007.....	23,965	1	1	-	23	85	384	1,235	2,124	3,133	6,442	10,536	1
2006.....	23,855	-	1	3	21	110	385	1,299	2,178	3,174	6,654	10,030	-
2005.....	24,902	2	-	3	22	84	406	1,166	1,954	3,306	7,256	10,703	-
2004.....	23,076	2	1	1	21	103	348	1,138	1,839	3,152	6,819	9,648	4
2003.....	21,940	1	2	2	22	98	372	1,031	1,755	3,104	6,649	8,904	-
2002.....	20,261	5	1	1	21	85	354	923	1,526	2,922	6,138	8,285	-
2001.....	19,250	-	1	1	23	104	331	931	1,462	2,832	5,993	7,572	-
2000.....	18,073	1	1	4	17	70	348	878	1,440	2,784	5,623	6,907	-
1999.....	16,968	3	3	1	13	78	313	799	1,318	2,799	5,324	6,317	-

All races, male

2007.....	9,417	1	1	-	13	46	250	760	1,273	1,601	2,667	2,805	-
2006.....	9,415	-	-	2	11	66	242	813	1,273	1,606	2,714	2,688	-
2005.....	9,458	1	-	1	12	45	243	733	1,151	1,639	2,917	2,716	-
2004.....	8,869	1	1	-	13	59	222	714	1,036	1,535	2,782	2,503	3
2003.....	8,213	1	2	1	12	58	215	660	989	1,535	2,522	2,218	-
2002.....	7,647	2	1	1	11	47	219	560	850	1,419	2,405	2,132	-
2001.....	7,254	-	-	-	16	59	206	569	784	1,379	2,282	1,959	-
2000.....	6,895	1	-	3	4	39	216	555	818	1,308	2,204	1,747	-
1999.....	6,488	2	1	1	8	48	195	482	732	1,372	2,067	1,580	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Essential hypertension and hypertensive renal disease (I10,I12,I15)

All races, female

2007.....	14,548	-	-	-	10	39	134	475	851	1,532	3,775	7,731	1
2006.....	14,440	-	1	1	10	44	143	486	905	1,568	3,940	7,342	-
2005.....	15,444	1	-	2	10	39	163	433	803	1,667	4,339	7,987	-
2004.....	14,207	1	-	1	8	44	126	424	803	1,617	4,037	7,145	1
2003.....	13,727	-	-	1	10	40	157	371	766	1,569	4,127	6,686	-
2002.....	12,614	3	-	-	10	38	135	363	676	1,503	3,733	6,153	-
2001.....	11,996	-	1	1	7	45	125	362	678	1,453	3,711	5,613	-
2000.....	11,178	-	1	1	13	31	132	323	622	1,476	3,419	5,160	-
1999.....	10,480	1	2	-	5	30	118	317	586	1,427	3,257	4,737	-

White, both sexes

2007 1/.....	18,583	1	1	-	13	34	201	669	1,313	2,190	5,113	9,048	-
2006 1/.....	18,264	-	1	3	10	44	164	693	1,361	2,180	5,266	8,542	-
2005 1/.....	19,254	-	-	3	10	29	194	602	1,211	2,222	5,805	9,178	-
2004 1/.....	17,693	-	1	-	13	44	161	597	1,079	2,148	5,405	8,242	3
2003 1/.....	16,927	-	1	1	12	43	186	566	1,039	2,065	5,309	7,705	-
2002.....	15,587	4	1	1	14	35	156	487	925	2,027	4,862	7,075	-
2001.....	14,752	-	1	1	14	45	140	463	863	1,939	4,776	6,510	-
2000.....	13,775	1	1	3	7	32	158	438	851	1,906	4,498	5,880	-
1999.....	12,830	2	1	-	7	36	131	407	764	1,949	4,184	5,349	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Essential hypertension and hypertensive renal disease (I10,I12,I15)													
White, male													
2007 1/.....	7,106	1	1	-	7	22	145	432	828	1,131	2,137	2,402	-
2006 1/.....	6,990	-	-	2	4	28	109	458	827	1,098	2,164	2,300	-
2005 1/.....	7,018	-	-	1	7	15	130	397	727	1,101	2,326	2,314	-
2004 1/.....	6,640	-	1	-	10	29	111	394	640	1,073	2,210	2,170	2
2003 1/.....	6,108	-	1	-	7	21	118	370	600	1,062	2,030	1,899	-
2002.....	5,719	2	1	1	8	21	100	311	542	1,006	1,901	1,826	-
2001.....	5,351	-	-	-	9	28	93	291	476	962	1,820	1,672	-
2000.....	5,120	1	-	3	3	22	109	301	503	926	1,783	1,469	-
1999.....	4,744	1	-	-	3	27	92	255	448	982	1,630	1,306	-
White, female													
2007 1/.....	11,477	-	-	-	6	12	56	237	485	1,059	2,976	6,646	-
2006 1/.....	11,274	-	1	1	6	16	55	235	534	1,082	3,102	6,242	-
2005 1/.....	12,236	-	-	2	3	14	64	205	484	1,121	3,479	6,864	-
2004 1/.....	11,053	-	-	-	3	15	50	203	439	1,075	3,195	6,072	1
2003 1/.....	10,819	-	-	1	5	22	68	196	439	1,003	3,279	5,806	-
2002.....	9,868	2	-	-	6	14	56	176	383	1,021	2,961	5,249	-
2001.....	9,401	-	1	1	5	17	47	172	387	977	2,956	4,838	-
2000.....	8,655	-	1	-	4	10	49	137	348	980	2,715	4,411	-
1999.....	8,086	1	1	-	4	9	39	152	316	967	2,554	4,043	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Essential hypertension and hypertensive renal disease (I10,I12,I15)													
Black, both sexes													
2007 1/.....	4,640	-	-	-	10	49	176	535	752	830	1,095	1,192	1
2006 1/.....	4,900	-	-	-	10	64	209	570	740	887	1,193	1,227	-
2005 1/.....	4,953	2	-	-	11	52	197	532	691	958	1,234	1,276	-
2004 1/.....	4,747	2	-	1	7	55	178	501	693	899	1,220	1,190	1
2003 1/.....	4,489	1	1	-	10	53	176	446	667	930	1,173	1,032	-
2002.....	4,196	1	-	-	7	45	191	417	554	803	1,138	1,040	-
2001.....	4,038	-	-	-	9	55	179	443	558	792	1,064	938	-
2000.....	3,887	-	-	1	10	35	179	409	559	806	982	906	-
1999.....	3,732	1	1	1	6	38	177	371	513	767	1,012	845	-
Black, male													
2007 1/.....	1,979	-	-	-	6	23	101	307	406	411	436	289	-
2006 1/.....	2,126	-	-	-	6	36	127	332	404	450	467	304	-
2005 1/.....	2,128	1	-	-	5	28	104	310	400	472	497	311	-
2004 1/.....	1,964	1	-	-	3	29	104	293	366	410	492	265	1
2003 1/.....	1,895	1	1	-	5	35	93	282	361	430	424	263	-
2002.....	1,717	-	-	-	3	24	113	237	276	364	444	256	-
2001.....	1,715	-	-	-	7	29	104	266	288	376	402	243	-
2000.....	1,592	-	-	-	1	16	100	233	303	345	360	234	-
1999.....	1,565	1	1	1	5	18	98	218	261	349	386	227	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Essential hypertension and hypertensive renal disease (I10,I12,I15)

Black, female

2007 1/.....	2,661	-	-	-	4	26	75	228	346	419	659	903	1
2006 1/.....	2,774	-	-	-	4	28	82	238	336	437	726	923	-
2005 1/.....	2,825	1	-	-	6	24	93	222	291	486	737	965	-
2004 1/.....	2,783	1	-	1	4	26	74	208	327	489	728	925	-
2003 1/.....	2,594	-	-	-	5	18	83	164	306	500	749	769	-
2002.....	2,479	1	-	-	4	21	78	180	278	439	694	784	-
2001.....	2,323	-	-	-	2	26	75	177	270	416	662	695	-
2000.....	2,295	-	-	1	9	19	79	176	256	461	622	672	-
1999.....	2,167	-	-	-	1	20	79	153	252	418	626	618	-

American Indian 2/, both sexes

2007 1/.....	108	-	-	-	-	-	1	8	16	20	37	26	-
2006 1/.....	106	-	-	-	1	-	4	10	13	22	23	33	-
2005 1/.....	113	-	-	-	-	1	2	10	7	28	38	27	-
2004 1/.....	107	-	-	-	1	2	6	12	18	23	29	16	-
2003 1/.....	88	-	-	-	-	-	3	7	17	27	15	19	-
2002.....	76	-	-	-	-	1	4	5	11	18	17	20	-
2001.....	63	-	-	-	-	2	5	2	6	16	16	16	-
2000.....	73	-	-	-	-	1	3	5	14	13	16	21	-
1999.....	55	-	-	-	-	2	3	2	9	10	11	18	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Essential hypertension and hypertensive renal disease (I10,I12,I15)

American Indian 2/, male

2007 1/.....	46	-	-	-	-	-	1	5	10	9	13	8	-
2006 1/.....	52	-	-	-	1	-	2	9	7	13	11	9	-
2005 1/.....	56	-	-	-	-	1	-	9	4	16	18	8	-
2004 1/.....	48	-	-	-	-	-	5	8	7	15	9	4	-
2003 1/.....	39	-	-	-	-	-	-	3	12	11	9	4	-
2002.....	30	-	-	-	-	-	4	4	7	7	6	2	-
2001.....	25	-	-	-	-	1	3	1	3	8	4	5	-
2000.....	24	-	-	-	-	-	2	3	5	3	5	6	-
1999.....	25	-	-	-	-	1	3	2	4	5	3	7	-

American Indian 2/, female

2007 1/.....	62	-	-	-	-	-	-	3	6	11	24	18	-
2006 1/.....	54	-	-	-	-	-	2	1	6	9	12	24	-
2005 1/.....	57	-	-	-	-	-	2	1	3	12	20	19	-
2004 1/.....	59	-	-	-	1	2	1	4	11	8	20	12	-
2003 1/.....	49	-	-	-	-	-	3	4	5	16	6	15	-
2002.....	46	-	-	-	-	1	-	1	4	11	11	18	-
2001.....	38	-	-	-	-	1	2	1	3	8	12	11	-
2000.....	49	-	-	-	-	1	1	2	9	10	11	15	-
1999.....	30	-	-	-	-	1	-	-	5	5	8	11	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Essential hypertension and hypertensive renal disease (I10,I12,I15)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	634	-	-	-	-	2	6	23	43	93	197	270	-
2006 1/.....	585	-	-	-	-	2	8	26	64	85	172	228	-
2005 1/.....	582	-	-	-	1	2	13	22	45	98	179	222	-
2004 1/.....	529	-	-	-	-	2	3	28	49	82	165	200	-
2003 1/.....	436	-	-	1	-	2	7	12	32	82	152	148	-
2002.....	402	-	-	-	-	4	3	14	36	74	121	150	-
2001.....	397	-	-	-	-	2	7	23	35	85	137	108	-
2000.....	338	-	-	-	-	2	8	26	16	59	127	100	-
1999.....	351	-	1	-	-	2	2	19	32	73	117	105	-

Asian or Pacific Islander 3/, male

2007 1/.....	286	-	-	-	-	1	3	16	29	50	81	106	-
2006 1/.....	247	-	-	-	-	2	4	14	35	45	72	75	-
2005 1/.....	256	-	-	-	-	1	9	17	20	50	76	83	-
2004 1/.....	217	-	-	-	-	1	2	19	23	37	71	64	-
2003 1/.....	171	-	-	1	-	2	4	5	16	32	59	52	-
2002.....	181	-	-	-	-	2	2	8	25	42	54	48	-
2001.....	163	-	-	-	-	1	6	11	17	33	56	39	-
2000.....	159	-	-	-	-	1	5	18	7	34	56	38	-
1999.....	154	-	-	-	-	2	2	7	19	36	48	40	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Essential hypertension and hypertensive renal disease (I10,I12,I15)

Asian or Pacific Islander 3/, female

2007 1/.....	348	-	-	-	-	1	3	7	14	43	116	164	-
2006 1/.....	338	-	-	-	-	-	4	12	29	40	100	153	-
2005 1/.....	326	-	-	-	1	1	4	5	25	48	103	139	-
2004 1/.....	312	-	-	-	-	1	1	9	26	45	94	136	-
2003 1/.....	265	-	-	-	-	-	3	7	16	50	93	96	-
2002.....	221	-	-	-	-	2	1	6	11	32	67	102	-
2001.....	234	-	-	-	-	1	1	12	18	52	81	69	-
2000.....	179	-	-	-	-	1	3	8	9	25	71	62	-
1999.....	197	-	1	-	-	-	-	12	13	37	69	65	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Cerebrovascular diseases (I60-I69)													
All races, both sexes													
2007.....	135,952	132	52	83	195	505	2,133	6,385	10,500	18,007	41,979	55,975	6
2006.....	137,119	142	54	95	210	527	2,221	6,341	10,518	18,223	43,719	55,068	1
2005.....	143,579	126	62	95	196	546	2,260	6,381	10,028	18,839	46,859	58,183	4
2004.....	150,074	127	41	77	211	567	2,361	6,181	9,966	19,901	50,092	60,545	5
2003.....	157,689	101	46	69	221	583	2,460	6,127	9,946	20,708	52,847	64,579	2
2002.....	162,672	117	53	91	171	567	2,425	6,055	9,897	21,992	54,889	66,412	3
2001.....	163,538	108	54	80	196	601	2,491	5,910	9,608	22,598	55,822	66,066	4
2000.....	167,661	124	47	76	199	602	2,599	6,011	9,956	23,649	57,020	67,376	2
1999.....	167,366	103	46	65	182	580	2,574	5,563	9,652	24,092	57,427	67,080	2
All races, male													
2007.....	54,111	77	20	37	113	252	1,143	3,507	5,992	9,351	17,700	15,915	4
2006.....	54,524	83	26	56	115	290	1,163	3,482	5,897	9,363	18,307	15,742	-
2005.....	56,586	73	41	55	94	301	1,148	3,446	5,636	9,688	19,688	16,412	4
2004.....	58,800	71	23	40	116	284	1,240	3,409	5,523	10,207	21,025	16,859	3
2003.....	61,426	57	25	41	116	314	1,283	3,350	5,478	10,668	22,234	17,859	1
2002.....	62,622	67	30	47	103	284	1,178	3,278	5,456	11,207	22,655	18,316	1
2001.....	63,177	63	25	42	103	311	1,267	3,205	5,277	11,649	23,301	17,930	4
2000.....	64,769	74	19	33	106	308	1,297	3,235	5,496	12,042	23,946	18,212	1
1999.....	64,485	65	22	34	95	299	1,308	2,996	5,305	12,220	24,088	18,052	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Cerebrovascular diseases (I60-I69)													
All races, female													
2007.....	81,841	55	32	46	82	253	990	2,878	4,508	8,656	24,279	40,060	2
2006.....	82,595	59	28	39	95	237	1,058	2,859	4,621	8,860	25,412	39,326	1
2005.....	86,993	53	21	40	102	245	1,112	2,935	4,392	9,151	27,171	41,771	-
2004.....	91,274	56	18	37	95	283	1,121	2,772	4,443	9,694	29,067	43,686	2
2003.....	96,263	44	21	28	105	269	1,177	2,777	4,468	10,040	30,613	46,720	1
2002.....	100,050	50	23	44	68	283	1,247	2,777	4,441	10,785	32,234	48,096	2
2001.....	100,361	45	29	38	93	290	1,224	2,705	4,331	10,949	32,521	48,136	-
2000.....	102,892	50	28	43	93	294	1,302	2,776	4,460	11,607	33,074	49,164	1
1999.....	102,881	38	24	31	87	281	1,266	2,567	4,347	11,872	33,339	49,028	1
White, both sexes													
2007 1/.....	114,695	94	34	63	138	332	1,356	4,139	7,438	14,023	36,340	50,734	4
2006 1/.....	115,864	89	39	67	159	345	1,389	4,135	7,417	14,230	37,984	50,009	1
2005 1/.....	121,868	79	45	64	136	356	1,450	4,083	7,140	14,829	40,832	52,854	-
2004 1/.....	127,868	79	28	58	152	373	1,557	3,943	7,103	15,799	43,672	55,101	3
2003 1/.....	134,705	63	30	51	152	388	1,508	3,859	7,019	16,465	46,329	58,839	2
2002.....	139,719	71	40	63	130	376	1,547	3,874	7,146	17,572	48,298	60,600	2
2001.....	140,465	60	39	59	138	416	1,605	3,736	6,874	18,047	49,265	60,225	1
2000.....	144,580	75	30	55	148	420	1,645	3,914	7,239	19,224	50,367	61,462	1
1999.....	144,827	70	34	49	135	402	1,646	3,609	6,940	19,577	50,917	61,448	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Cerebrovascular diseases (I60-I69)													
White, male													
2007 1/.....	44,714	51	16	32	76	167	746	2,330	4,242	7,272	15,322	14,457	3
2006 1/.....	45,198	48	16	46	91	195	749	2,279	4,110	7,312	16,041	14,311	-
2005 1/.....	47,194	46	28	35	65	197	754	2,239	3,986	7,645	17,264	14,935	-
2004 1/.....	49,296	45	18	29	82	198	842	2,195	3,906	8,149	18,457	15,372	3
2003 1/.....	51,646	32	14	31	87	210	802	2,160	3,855	8,571	19,588	16,295	1
2002.....	52,959	41	22	29	79	195	773	2,132	3,937	9,021	20,037	16,692	1
2001.....	53,428	35	18	34	81	220	834	2,055	3,790	9,393	20,664	16,303	1
2000.....	54,938	46	14	26	84	225	842	2,138	3,998	9,828	21,210	16,526	1
1999.....	54,867	43	16	27	72	213	856	1,949	3,843	10,000	21,413	16,435	-
White, female													
2007 1/.....	69,981	43	18	31	62	165	610	1,809	3,196	6,751	21,018	36,277	1
2006 1/.....	70,666	41	23	21	68	150	640	1,856	3,307	6,918	21,943	35,698	1
2005 1/.....	74,674	33	17	29	71	159	696	1,844	3,154	7,184	23,568	37,919	-
2004 1/.....	78,572	34	10	29	70	175	715	1,748	3,197	7,650	25,215	39,729	-
2003 1/.....	83,059	31	16	20	65	178	706	1,699	3,164	7,894	26,741	42,544	1
2002.....	86,760	30	18	34	51	181	774	1,742	3,209	8,551	28,261	43,908	1
2001.....	87,037	25	21	25	57	196	771	1,681	3,084	8,654	28,601	43,922	-
2000.....	89,642	29	16	29	64	195	803	1,776	3,241	9,396	29,157	44,936	-
1999.....	89,960	27	18	22	63	189	790	1,660	3,097	9,577	29,504	45,013	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Cerebrovascular diseases (I60-I69)													
Black, both sexes													
2007 1/.....	17,085	34	16	13	46	148	655	1,946	2,591	3,257	4,385	3,992	2
2006 1/.....	17,045	49	12	21	44	151	697	1,919	2,613	3,209	4,442	3,888	-
2005 1/.....	17,541	40	14	24	51	163	689	2,004	2,430	3,235	4,746	4,141	4
2004 1/.....	18,118	40	12	17	49	173	703	1,920	2,433	3,384	5,090	4,295	2
2003 1/.....	18,806	32	15	16	59	168	793	1,968	2,481	3,438	5,213	4,623	-
2002.....	18,856	37	11	24	38	168	744	1,893	2,324	3,630	5,326	4,660	1
2001.....	19,002	43	12	14	51	158	765	1,906	2,310	3,714	5,295	4,731	3
2000.....	19,221	46	14	15	43	160	836	1,836	2,291	3,689	5,399	4,891	1
1999.....	18,884	28	10	15	41	150	816	1,683	2,295	3,813	5,351	4,681	1
Black, male													
2007 1/.....	7,549	25	4	3	30	71	335	1,006	1,500	1,714	1,840	1,020	1
2006 1/.....	7,424	33	8	7	20	75	342	1,044	1,523	1,644	1,741	987	-
2005 1/.....	7,519	23	10	15	24	89	332	1,050	1,417	1,653	1,844	1,058	4
2004 1/.....	7,644	23	5	10	31	77	345	1,021	1,388	1,697	1,979	1,068	-
2003 1/.....	7,866	20	10	9	26	87	387	1,039	1,381	1,687	2,088	1,132	-
2002.....	7,828	20	7	16	22	74	332	1,000	1,298	1,842	2,049	1,168	-
2001.....	7,907	27	5	5	20	79	361	1,008	1,265	1,859	2,086	1,189	3
2000.....	8,026	27	3	5	19	71	397	969	1,258	1,835	2,177	1,265	-
1999.....	7,894	18	4	7	19	72	393	910	1,251	1,868	2,116	1,236	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Cerebrovascular diseases (I60-I69)													
Black, female													
2007 1/.....	9,536	9	12	10	16	77	320	940	1,091	1,543	2,545	2,972	1
2006 1/.....	9,621	16	4	14	24	76	355	875	1,090	1,565	2,701	2,901	-
2005 1/.....	10,022	17	4	9	27	74	357	954	1,013	1,582	2,902	3,083	-
2004 1/.....	10,474	17	7	7	18	96	358	899	1,045	1,687	3,111	3,227	2
2003 1/.....	10,940	12	5	7	33	81	406	929	1,100	1,751	3,125	3,491	-
2002.....	11,028	17	4	8	16	94	412	893	1,026	1,788	3,277	3,492	1
2001.....	11,095	16	7	9	31	79	404	898	1,045	1,855	3,209	3,542	-
2000.....	11,195	19	11	10	24	89	439	867	1,033	1,854	3,222	3,626	1
1999.....	10,990	10	6	8	22	78	423	773	1,044	1,945	3,235	3,445	1
American Indian 2/, both sexes													
2007 1/.....	586	2	1	-	2	3	28	51	83	114	154	148	-
2006 1/.....	548	2	2	2	3	5	31	52	66	105	147	133	-
2005 1/.....	627	2	-	-	2	4	32	64	89	142	163	129	-
2004 1/.....	581	1	-	1	1	5	24	49	64	133	168	135	-
2003 1/.....	552	-	1	1	1	5	34	58	63	113	167	109	-
2002.....	567	1	-	2	2	5	33	54	71	116	158	125	-
2001.....	574	2	-	1	-	5	30	50	54	141	160	131	-
2000.....	572	-	1	2	1	5	22	47	79	108	178	129	-
1999.....	546	2	-	1	1	11	30	40	63	105	166	127	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Cerebrovascular diseases (I60-I69)

American Indian 2/, male

2007 1/.....	267	-	-	-	1	2	14	29	49	53	77	42	-
2006 1/.....	231	1	2	1	1	3	21	33	44	50	48	27	-
2005 1/.....	249	1	-	-	1	2	16	27	43	65	60	34	-
2004 1/.....	246	-	-	1	-	2	6	27	34	61	78	37	-
2003 1/.....	242	-	1	-	1	4	18	29	33	54	66	36	-
2002.....	236	1	-	-	1	4	14	28	35	49	61	43	-
2001.....	217	-	-	1	-	1	13	22	23	64	51	42	-
2000.....	250	-	1	-	-	-	11	22	43	65	69	39	-
1999.....	236	2	-	-	1	4	15	21	26	59	75	33	-

American Indian 2/, female

2007 1/.....	319	2	1	-	1	1	14	22	34	61	77	106	-
2006 1/.....	317	1	-	1	2	2	10	19	22	55	99	106	-
2005 1/.....	378	1	-	-	1	2	16	37	46	77	103	95	-
2004 1/.....	335	1	-	-	1	3	18	22	30	72	90	98	-
2003 1/.....	310	-	-	1	-	1	16	29	30	59	101	73	-
2002.....	331	-	-	2	1	1	19	26	36	67	97	82	-
2001.....	357	2	-	-	-	4	17	28	31	77	109	89	-
2000.....	322	-	-	2	1	5	11	25	36	43	109	90	-
1999.....	310	-	-	1	-	7	15	19	37	46	91	94	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Cerebrovascular diseases (I60-I69)													
Asian or Pacific Islander 3/, both sexes													
2007 1/.....	3,586	2	1	7	9	22	94	249	388	613	1,100	1,101	-
2006 1/.....	3,662	2	1	5	4	26	104	235	422	679	1,146	1,038	-
2005 1/.....	3,543	5	3	7	7	23	89	230	369	633	1,118	1,059	-
2004 1/.....	3,507	7	1	1	9	16	77	269	366	585	1,162	1,014	-
2003 1/.....	3,626	6	-	1	9	22	125	242	383	692	1,138	1,008	-
2002.....	3,530	8	2	2	1	18	101	234	356	674	1,107	1,027	-
2001.....	3,497	3	3	6	7	22	91	218	370	696	1,102	979	-
2000.....	3,288	3	2	4	7	17	96	214	347	628	1,076	894	-
1999.....	3,109	3	2	-	5	17	82	231	354	597	993	824	1
Asian or Pacific Islander 3/, male													
2007 1/.....	1,581	1	-	2	6	12	48	142	201	312	461	396	-
2006 1/.....	1,671	1	-	2	3	17	51	126	220	357	477	417	-
2005 1/.....	1,624	3	3	5	4	13	46	130	190	325	520	385	-
2004 1/.....	1,614	3	-	-	3	7	47	166	195	300	511	382	-
2003 1/.....	1,672	5	-	1	2	13	76	122	209	356	492	396	-
2002.....	1,599	5	1	2	1	11	59	118	186	295	508	413	-
2001.....	1,625	1	2	2	2	11	59	120	199	333	500	396	-
2000.....	1,555	1	1	2	3	12	47	106	197	314	490	382	-
1999.....	1,488	2	2	-	3	10	44	116	185	293	484	348	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Cerebrovascular diseases (I60-I69)

Asian or Pacific Islander 3/, female

2007 1/.....	2,005	1	1	5	3	10	46	107	187	301	639	705	-
2006 1/.....	1,991	1	1	3	1	9	53	109	202	322	669	621	-
2005 1/.....	1,919	2	-	2	3	10	43	100	179	308	598	674	-
2004 1/.....	1,893	4	1	1	6	9	30	103	171	285	651	632	-
2003 1/.....	1,954	1	-	-	7	9	49	120	174	336	646	612	-
2002.....	1,931	3	1	-	-	7	42	116	170	379	599	614	-
2001.....	1,872	2	1	4	5	11	32	98	171	363	602	583	-
2000.....	1,733	2	1	2	4	5	49	108	150	314	586	512	-
1999.....	1,621	1	-	-	2	7	38	115	169	304	509	476	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Atherosclerosis (I70)													
All races, both sexes													
2007.....	8,232	1	-	-	2	1	27	134	350	829	2,298	4,590	-
2006.....	8,652	2	1	-	1	7	31	140	388	849	2,469	4,764	-
2005.....	11,841	-	1	-	2	6	31	193	440	1,008	3,387	6,773	-
2004.....	11,861	-	1	-	2	6	43	153	459	1,132	3,406	6,659	-
2003.....	13,053	3	-	-	3	7	47	190	466	1,211	3,837	7,288	1
2002.....	13,821	4	-	-	3	9	50	182	487	1,300	4,062	7,723	1
2001.....	14,086	1	1	1	1	3	40	186	473	1,335	4,156	7,888	1
2000.....	14,393	2	-	-	3	8	48	176	451	1,423	4,277	8,005	-
1999.....	14,979	2	-	-	2	8	45	185	486	1,476	4,385	8,390	-
All races, male													
2007.....	3,220	1	-	-	-	1	19	95	243	474	1,027	1,360	-
2006.....	3,359	1	-	-	1	5	21	99	244	463	1,085	1,440	-
2005.....	4,472	-	1	-	1	3	20	134	269	560	1,532	1,952	-
2004.....	4,510	-	-	-	2	3	25	104	275	619	1,549	1,933	-
2003.....	4,859	2	-	-	2	5	35	131	298	649	1,734	2,002	1
2002.....	5,171	1	-	-	2	5	30	119	305	728	1,803	2,177	1
2001.....	5,143	1	1	1	1	2	26	118	286	743	1,791	2,172	1
2000.....	5,357	-	-	-	2	6	35	119	294	800	1,935	2,166	-
1999.....	5,443	1	-	-	-	5	32	121	307	795	1,927	2,255	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Atherosclerosis (I70)													
All races, female													
2007.....	5,012	-	-	-	2	-	8	39	107	355	1,271	3,230	-
2006.....	5,293	1	1	-	-	2	10	41	144	386	1,384	3,324	-
2005.....	7,369	-	-	-	1	3	11	59	171	448	1,855	4,821	-
2004.....	7,351	-	1	-	-	3	18	49	184	513	1,857	4,726	-
2003.....	8,194	1	-	-	1	2	12	59	168	562	2,103	5,286	-
2002.....	8,650	3	-	-	1	4	20	63	182	572	2,259	5,546	-
2001.....	8,943	-	-	-	-	1	14	68	187	592	2,365	5,716	-
2000.....	9,036	2	-	-	1	2	13	57	157	623	2,342	5,839	-
1999.....	9,536	1	-	-	2	3	13	64	179	681	2,458	6,135	-
White, both sexes													
2007 1/.....	7,444	1	-	-	1	1	20	102	285	706	2,075	4,253	-
2006 1/.....	7,820	2	-	-	1	5	21	104	315	720	2,217	4,435	-
2005 1/.....	10,845	-	-	-	1	5	26	148	366	855	3,088	6,356	-
2004 1/.....	10,877	-	1	-	2	5	30	125	386	975	3,134	6,219	-
2003 1/.....	11,885	1	-	-	2	5	39	148	385	1,031	3,484	6,789	1
2002.....	12,631	3	-	-	3	6	40	129	397	1,132	3,666	7,254	1
2001.....	12,961	1	1	1	-	2	33	151	382	1,166	3,814	7,409	1
2000.....	13,182	2	-	-	3	5	36	123	366	1,246	3,910	7,491	-
1999.....	13,737	1	-	-	2	6	33	151	402	1,300	4,019	7,823	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Atherosclerosis (I70)													
White, male													
2007 1/.....	2,898	1	-	-	-	1	14	72	192	406	946	1,266	-
2006 1/.....	2,992	1	-	-	1	3	13	78	199	389	971	1,337	-
2005 1/.....	4,093	-	-	-	-	2	17	107	225	478	1,410	1,854	-
2004 1/.....	4,146	-	-	-	2	3	19	91	232	545	1,425	1,829	-
2003 1/.....	4,368	-	-	-	1	4	28	102	249	549	1,562	1,872	1
2002.....	4,688	1	-	-	2	4	24	84	250	640	1,630	2,052	1
2001.....	4,696	1	1	1	-	2	21	94	237	655	1,638	2,045	1
2000.....	4,875	-	-	-	2	5	28	86	247	713	1,783	2,011	-
1999.....	4,932	1	-	-	-	4	24	102	254	701	1,755	2,091	-
White, female													
2007 1/.....	4,546	-	-	-	1	-	6	30	93	300	1,129	2,987	-
2006 1/.....	4,828	1	-	-	-	2	8	26	116	331	1,246	3,098	-
2005 1/.....	6,752	-	-	-	1	3	9	41	141	377	1,678	4,502	-
2004 1/.....	6,731	-	1	-	-	2	11	34	154	430	1,709	4,390	-
2003 1/.....	7,517	1	-	-	1	1	11	46	136	482	1,922	4,917	-
2002.....	7,943	2	-	-	1	2	16	45	147	492	2,036	5,202	-
2001.....	8,265	-	-	-	-	-	12	57	145	511	2,176	5,364	-
2000.....	8,307	2	-	-	1	-	8	37	119	533	2,127	5,480	-
1999.....	8,805	-	-	-	2	2	9	49	148	599	2,264	5,732	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Atherosclerosis (I70)													
Black, both sexes													
2007 1/.....	647	-	-	-	1	-	5	30	54	100	177	280	-
2006 1/.....	709	-	1	-	-	2	9	29	68	115	209	276	-
2005 1/.....	844	-	1	-	1	1	5	38	63	131	238	366	-
2004 1/.....	837	-	-	-	-	1	12	26	67	128	238	365	-
2003 1/.....	995	2	-	-	1	2	5	39	72	158	301	415	-
2002.....	1,017	1	-	-	-	3	9	46	77	135	352	394	-
2001.....	970	-	-	-	1	1	6	31	84	148	290	409	-
2000.....	1,054	-	-	-	-	3	12	47	79	148	323	442	-
1999.....	1,069	1	-	-	-	2	9	29	77	151	316	484	-
Black, male													
2007 1/.....	259	-	-	-	-	-	4	22	42	54	63	74	-
2006 1/.....	312	-	-	-	-	2	7	17	43	65	94	84	-
2005 1/.....	312	-	1	-	1	1	3	21	38	71	95	81	-
2004 1/.....	305	-	-	-	-	-	5	12	39	62	108	79	-
2003 1/.....	410	2	-	-	1	1	4	27	44	88	143	100	-
2002.....	414	-	-	-	-	1	6	30	49	71	154	103	-
2001.....	370	-	-	-	1	-	4	21	43	75	126	100	-
2000.....	415	-	-	-	-	1	7	28	43	75	133	128	-
1999.....	432	-	-	-	-	1	6	15	47	80	143	140	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Atherosclerosis (I70)

Black, female

2007 1/.....	388	-	-	-	1	-	1	8	12	46	114	206	-
2006 1/.....	397	-	1	-	-	-	2	12	25	50	115	192	-
2005 1/.....	532	-	-	-	-	-	2	17	25	60	143	285	-
2004 1/.....	532	-	-	-	-	1	7	14	28	66	130	286	-
2003 1/.....	585	-	-	-	-	1	1	12	28	70	158	315	-
2002.....	603	1	-	-	-	2	3	16	28	64	198	291	-
2001.....	600	-	-	-	-	1	2	10	41	73	164	309	-
2000.....	639	-	-	-	-	2	5	19	36	73	190	314	-
1999.....	637	1	-	-	-	1	3	14	30	71	173	344	-

American Indian 2/, both sexes

2007 1/.....	30	-	-	-	-	-	-	1	4	9	7	9	-
2006 1/.....	27	-	-	-	-	-	-	4	1	4	7	11	-
2005 1/.....	51	-	-	-	-	-	-	3	10	12	16	10	-
2004 1/.....	36	-	-	-	-	-	1	-	3	10	10	12	-
2003 1/.....	34	-	-	-	-	-	2	2	5	8	6	11	-
2002.....	54	-	-	-	-	-	1	5	7	17	16	8	-
2001.....	44	-	-	-	-	-	-	2	4	8	14	16	-
2000.....	49	-	-	-	-	-	-	-	3	16	13	17	-
1999.....	44	-	-	-	-	-	-	3	2	11	12	16	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Atherosclerosis (I70)

American Indian 2/, male

2007 1/.....	15	-	-	-	-	-	-	1	3	7	1	3	-
2006 1/.....	6	-	-	-	-	-	-	2	1	2	-	1	-
2005 1/.....	24	-	-	-	-	-	-	3	5	6	6	4	-
2004 1/.....	17	-	-	-	-	-	1	-	3	3	7	3	-
2003 1/.....	17	-	-	-	-	-	2	1	2	5	4	3	-
2002.....	21	-	-	-	-	-	-	3	4	7	5	2	-
2001.....	21	-	-	-	-	-	-	2	3	5	6	5	-
2000.....	18	-	-	-	-	-	-	-	2	6	6	4	-
1999.....	18	-	-	-	-	-	-	2	2	6	6	2	-

American Indian 2/, female

2007 1/.....	15	-	-	-	-	-	-	-	1	2	6	6	-
2006 1/.....	21	-	-	-	-	-	-	2	-	2	7	10	-
2005 1/.....	27	-	-	-	-	-	-	-	5	6	10	6	-
2004 1/.....	19	-	-	-	-	-	-	-	-	7	3	9	-
2003 1/.....	17	-	-	-	-	-	-	1	3	3	2	8	-
2002.....	33	-	-	-	-	-	1	2	3	10	11	6	-
2001.....	23	-	-	-	-	-	-	-	1	3	8	11	-
2000.....	31	-	-	-	-	-	-	-	1	10	7	13	-
1999.....	26	-	-	-	-	-	-	1	-	5	6	14	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Atherosclerosis (I70)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	111	-	-	-	-	-	2	1	7	14	39	48	-
2006 1/.....	96	-	-	-	-	-	1	3	4	10	36	42	-
2005 1/.....	101	-	-	-	-	-	-	4	1	10	45	41	-
2004 1/.....	111	-	-	-	-	-	-	2	3	19	24	63	-
2003 1/.....	139	-	-	-	-	-	1	1	4	14	46	73	-
2002.....	119	-	-	-	-	-	-	2	6	16	28	67	-
2001.....	111	-	-	-	-	-	1	2	3	13	38	54	-
2000.....	108	-	-	-	-	-	-	6	3	13	31	55	-
1999.....	129	-	-	-	-	-	3	2	5	14	38	67	-

Asian or Pacific Islander 3/, male

2007 1/.....	48	-	-	-	-	-	1	-	6	7	17	17	-
2006 1/.....	49	-	-	-	-	-	1	2	1	7	20	18	-
2005 1/.....	43	-	-	-	-	-	-	3	1	5	21	13	-
2004 1/.....	42	-	-	-	-	-	-	1	1	9	9	22	-
2003 1/.....	64	-	-	-	-	-	1	1	3	7	25	27	-
2002.....	48	-	-	-	-	-	-	2	2	10	14	20	-
2001.....	56	-	-	-	-	-	1	1	3	8	21	22	-
2000.....	49	-	-	-	-	-	-	5	2	6	13	23	-
1999.....	61	-	-	-	-	-	2	2	4	8	23	22	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Atherosclerosis (I70)

Asian or Pacific Islander 3/, female

2007 1/.....	63	-	-	-	-	-	1	1	1	7	22	31	-
2006 1/.....	47	-	-	-	-	-	-	1	3	3	16	24	-
2005 1/.....	58	-	-	-	-	-	-	1	-	5	24	28	-
2004 1/.....	69	-	-	-	-	-	-	1	2	10	15	41	-
2003 1/.....	75	-	-	-	-	-	-	-	1	7	21	46	-
2002.....	71	-	-	-	-	-	-	-	4	6	14	47	-
2001.....	55	-	-	-	-	-	-	1	-	5	17	32	-
2000.....	59	-	-	-	-	-	-	1	1	7	18	32	-
1999.....	68	-	-	-	-	-	1	-	1	6	15	45	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other diseases of circulatory system (I71-I78)

All races, both sexes

2007.....	21,940	13	4	14	65	136	484	1,092	2,296	4,065	7,074	6,694	3
2006.....	22,484	18	1	12	68	148	444	1,162	2,153	4,104	7,569	6,805	-
2005.....	23,617	17	3	17	52	156	467	1,085	2,266	4,484	8,167	6,903	-
2004.....	23,693	14	4	13	61	167	461	1,125	2,107	4,632	8,341	6,768	-
2003.....	24,672	11	-	12	68	151	451	1,199	2,270	4,934	8,694	6,882	-
2002.....	24,927	14	4	11	66	160	445	1,070	2,286	5,310	8,745	6,816	-
2001.....	25,318	12	4	11	61	170	498	1,024	2,195	5,345	9,032	6,965	1
2000.....	26,036	16	5	11	59	154	448	1,038	2,236	5,926	9,222	6,920	1
1999.....	25,809	12	2	11	49	141	478	971	2,210	5,808	9,264	6,863	-

All races, male

2007.....	11,480	7	1	11	48	91	350	765	1,580	2,474	3,613	2,537	3
2006.....	11,836	12	-	9	44	106	319	806	1,464	2,513	3,947	2,616	-
2005.....	12,423	11	3	13	37	102	322	749	1,533	2,796	4,320	2,537	-
2004.....	12,306	7	2	8	45	113	308	780	1,392	2,818	4,322	2,511	-
2003.....	13,217	3	-	7	45	103	306	815	1,517	3,092	4,732	2,597	-
2002.....	13,309	6	4	5	43	101	291	732	1,526	3,307	4,690	2,604	-
2001.....	13,396	8	2	3	40	115	335	688	1,454	3,348	4,839	2,563	1
2000.....	14,084	6	2	7	33	103	276	699	1,530	3,771	5,019	2,637	1
1999.....	13,794	9	-	7	32	102	296	657	1,519	3,681	5,015	2,476	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other diseases of circulatory system (I71-I78)

All races, female

2007.....	10,460	6	3	3	17	45	134	327	716	1,591	3,461	4,157	-
2006.....	10,648	6	1	3	24	42	125	356	689	1,591	3,622	4,189	-
2005.....	11,194	6	-	4	15	54	145	336	733	1,688	3,847	4,366	-
2004.....	11,387	7	2	5	16	54	153	345	715	1,814	4,019	4,257	-
2003.....	11,455	8	-	5	23	48	145	384	753	1,842	3,962	4,285	-
2002.....	11,618	8	-	6	23	59	154	338	760	2,003	4,055	4,212	-
2001.....	11,922	4	2	8	21	55	163	336	741	1,997	4,193	4,402	-
2000.....	11,952	10	3	4	26	51	172	339	706	2,155	4,203	4,283	-
1999.....	12,015	3	2	4	17	39	182	314	691	2,127	4,249	4,387	-

White, both sexes

2007 1/.....	18,925	9	4	11	47	101	328	756	1,818	3,476	6,294	6,079	2
2006 1/.....	19,497	13	1	9	55	105	311	833	1,738	3,529	6,732	6,171	-
2005 1/.....	20,485	11	3	14	43	100	314	765	1,845	3,865	7,287	6,238	-
2004 1/.....	20,634	11	3	11	49	112	319	828	1,675	4,047	7,477	6,102	-
2003 1/.....	21,661	6	-	8	49	107	306	861	1,884	4,336	7,849	6,255	-
2002.....	21,935	11	4	4	49	108	319	762	1,877	4,671	7,946	6,184	-
2001.....	22,236	10	3	10	47	129	340	758	1,793	4,705	8,137	6,303	1
2000.....	22,963	13	4	7	44	111	308	756	1,834	5,267	8,362	6,257	-
1999.....	22,952	9	2	8	42	106	356	729	1,868	5,184	8,430	6,218	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of circulatory system (I71-I78)													
White, male													
2007 1/.....	9,934	5	1	9	33	68	240	541	1,288	2,149	3,257	2,341	2
2006 1/.....	10,373	9	-	9	35	77	223	596	1,199	2,200	3,601	2,424	-
2005 1/.....	10,825	6	3	11	29	65	208	531	1,275	2,459	3,897	2,341	-
2004 1/.....	10,810	6	2	6	35	76	217	586	1,117	2,511	3,949	2,305	-
2003 1/.....	11,708	2	-	4	34	70	217	609	1,280	2,769	4,317	2,406	-
2002.....	11,818	5	4	2	31	68	214	533	1,264	2,962	4,334	2,401	-
2001.....	11,887	6	2	3	31	90	228	511	1,216	3,003	4,429	2,367	1
2000.....	12,611	4	2	6	26	77	190	528	1,290	3,428	4,625	2,435	-
1999.....	12,444	6	-	5	27	77	226	503	1,321	3,364	4,637	2,278	-
White, female													
2007 1/.....	8,991	4	3	2	14	33	88	215	530	1,327	3,037	3,738	-
2006 1/.....	9,124	4	1	-	20	28	88	237	539	1,329	3,131	3,747	-
2005 1/.....	9,660	5	-	3	14	35	106	234	570	1,406	3,390	3,897	-
2004 1/.....	9,824	5	1	5	14	36	102	242	558	1,536	3,528	3,797	-
2003 1/.....	9,953	4	-	4	15	37	89	252	604	1,567	3,532	3,849	-
2002.....	10,117	6	-	2	18	40	105	229	613	1,709	3,612	3,783	-
2001.....	10,349	4	1	7	16	39	112	247	577	1,702	3,708	3,936	-
2000.....	10,352	9	2	1	18	34	118	228	544	1,839	3,737	3,822	-
1999.....	10,508	3	2	3	15	29	130	226	547	1,820	3,793	3,940	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other diseases of circulatory system (I71-I78)

Black, both sexes

2007 1/.....	2,470	2	-	2	13	26	142	295	400	474	610	505	1
2006 1/.....	2,482	5	-	2	11	37	116	294	370	477	660	510	-
2005 1/.....	2,602	4	-	3	8	46	132	275	372	516	705	541	-
2004 1/.....	2,550	2	1	2	7	46	128	260	369	485	700	550	-
2003 1/.....	2,519	5	-	3	14	37	133	298	327	500	688	514	-
2002.....	2,525	3	-	6	12	47	114	275	357	524	659	528	-
2001.....	2,632	2	1	1	12	39	142	229	353	530	754	569	-
2000.....	2,595	3	-	4	13	36	124	234	351	546	715	568	1
1999.....	2,434	3	-	-	7	30	112	212	294	521	694	561	-

Black, male

2007 1/.....	1,242	2	-	2	11	17	98	199	245	253	265	149	1
2006 1/.....	1,184	3	-	-	8	26	81	189	235	261	250	131	-
2005 1/.....	1,282	3	-	2	7	28	96	186	229	267	325	139	-
2004 1/.....	1,225	1	-	2	7	31	80	170	234	254	287	159	-
2003 1/.....	1,216	1	-	2	7	29	82	182	202	258	320	133	-
2002.....	1,211	1	-	2	7	31	71	174	227	269	274	155	-
2001.....	1,229	2	-	-	7	24	94	150	201	276	329	146	-
2000.....	1,207	2	-	1	6	22	77	146	201	281	316	154	1
1999.....	1,111	3	-	-	5	23	61	135	167	268	292	157	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other diseases of circulatory system (I71-I78)

Black, female

2007 1/.....	1,228	-	-	-	2	9	44	96	155	221	345	356	-
2006 1/.....	1,298	2	-	2	3	11	35	105	135	216	410	379	-
2005 1/.....	1,320	1	-	1	1	18	36	89	143	249	380	402	-
2004 1/.....	1,325	1	1	-	-	15	48	90	135	231	413	391	-
2003 1/.....	1,303	4	-	1	7	8	51	116	125	242	368	381	-
2002.....	1,314	2	-	4	5	16	43	101	130	255	385	373	-
2001.....	1,403	-	1	1	5	15	48	79	152	254	425	423	-
2000.....	1,388	1	-	3	7	14	47	88	150	265	399	414	-
1999.....	1,323	-	-	-	2	7	51	77	127	253	402	404	-

American Indian 2/, both sexes

2007 1/.....	110	1	-	-	2	2	5	9	19	30	24	18	-
2006 1/.....	79	-	-	-	1	-	2	6	11	23	27	9	-
2005 1/.....	84	1	-	-	-	1	3	7	13	23	25	11	-
2004 1/.....	86	-	-	-	-	3	3	8	15	15	27	15	-
2003 1/.....	102	-	-	-	2	3	2	10	15	19	28	23	-
2002.....	95	-	-	-	1	1	2	9	16	30	21	15	-
2001.....	79	-	-	-	-	-	2	6	11	21	25	14	-
2000.....	87	-	-	-	-	-	3	8	8	33	20	15	-
1999.....	72	-	-	2	-	-	-	2	12	20	19	17	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other diseases of circulatory system (I71-I78)

American Indian 2/, male

2007 1/.....	65	-	-	-	2	2	4	6	11	19	14	7	-
2006 1/.....	34	-	-	-	-	-	2	3	5	9	12	3	-
2005 1/.....	37	1	-	-	-	1	2	1	7	13	11	1	-
2004 1/.....	38	-	-	-	-	2	2	4	9	3	11	7	-
2003 1/.....	55	-	-	-	2	2	-	7	7	14	12	11	-
2002.....	53	-	-	-	1	-	2	6	8	21	10	5	-
2001.....	42	-	-	-	-	-	2	4	9	11	9	7	-
2000.....	37	-	-	-	-	-	-	4	5	14	10	4	-
1999.....	32	-	-	1	-	-	-	1	7	9	8	6	-

American Indian 2/, female

2007 1/.....	45	1	-	-	-	-	1	3	8	11	10	11	-
2006 1/.....	45	-	-	-	1	-	-	3	6	14	15	6	-
2005 1/.....	47	-	-	-	-	-	1	6	6	10	14	10	-
2004 1/.....	48	-	-	-	-	1	1	4	6	12	16	8	-
2003 1/.....	47	-	-	-	-	1	2	3	8	5	16	12	-
2002.....	42	-	-	-	-	1	-	3	8	9	11	10	-
2001.....	37	-	-	-	-	-	-	2	2	10	16	7	-
2000.....	50	-	-	-	-	-	3	4	3	19	10	11	-
1999.....	40	-	-	1	-	-	-	1	5	11	11	11	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other diseases of circulatory system (I71-I78)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	435	1	-	1	3	7	9	32	59	85	146	92	-
2006 1/.....	426	-	-	1	1	6	15	29	34	75	150	115	-
2005 1/.....	446	1	-	-	1	9	18	38	36	80	150	113	-
2004 1/.....	423	1	-	-	5	6	11	29	48	85	137	101	-
2003 1/.....	390	-	-	1	3	4	10	30	44	79	129	90	-
2002.....	372	-	-	1	4	4	10	24	36	85	119	89	-
2001.....	371	-	-	-	2	2	14	31	38	89	116	79	-
2000.....	391	-	1	-	2	7	13	40	43	80	125	80	-
1999.....	351	-	-	1	-	5	10	28	36	83	121	67	-

Asian or Pacific Islander 3/, male

2007 1/.....	239	-	-	-	2	4	8	19	36	53	77	40	-
2006 1/.....	245	-	-	-	1	3	13	18	25	43	84	58	-
2005 1/.....	279	1	-	-	1	8	16	31	22	57	87	56	-
2004 1/.....	233	-	-	-	3	4	9	20	32	50	75	40	-
2003 1/.....	238	-	-	1	2	2	7	17	28	51	83	47	-
2002.....	227	-	-	1	4	2	4	19	27	55	72	43	-
2001.....	238	-	-	-	2	1	11	23	28	58	72	43	-
2000.....	229	-	-	-	1	4	9	21	34	48	68	44	-
1999.....	207	-	-	1	-	2	9	18	24	40	78	35	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other diseases of circulatory system (I71-I78)

Asian or Pacific Islander 3/, female

2007 1/.....	196	1	-	1	1	3	1	13	23	32	69	52	-
2006 1/.....	181	-	-	1	-	3	2	11	9	32	66	57	-
2005 1/.....	167	-	-	-	-	1	2	7	14	23	63	57	-
2004 1/.....	190	1	-	-	2	2	2	9	16	35	62	61	-
2003 1/.....	152	-	-	-	1	2	3	13	16	28	46	43	-
2002.....	145	-	-	-	-	2	6	5	9	30	47	46	-
2001.....	133	-	-	-	-	1	3	8	10	31	44	36	-
2000.....	162	-	1	-	1	3	4	19	9	32	57	36	-
1999.....	144	-	-	-	-	3	1	10	12	43	43	32	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Aortic aneurysm and dissection (I71)													
All races, both sexes													
2007.....	12,986	1	-	7	38	99	364	752	1,483	2,616	4,346	3,279	1
2006.....	13,238	2	-	7	43	107	340	795	1,366	2,635	4,622	3,321	-
2005.....	13,843	1	-	4	34	114	329	758	1,456	2,870	4,971	3,306	-
2004.....	13,753	1	1	5	45	126	318	729	1,318	2,932	5,131	3,147	-
2003.....	14,810	1	-	3	41	112	316	820	1,477	3,271	5,471	3,298	-
2002.....	14,818	2	-	5	39	110	305	710	1,460	3,482	5,518	3,187	-
2001.....	15,234	-	2	4	43	117	332	661	1,394	3,517	5,751	3,412	1
2000.....	15,810	-	-	5	36	107	303	662	1,493	4,012	5,835	3,357	-
1999.....	15,807	1	1	3	31	93	329	638	1,437	4,001	5,936	3,337	-
All races, male													
2007.....	7,543	-	-	6	32	74	278	567	1,093	1,698	2,341	1,453	1
2006.....	7,732	1	-	7	32	87	269	600	1,004	1,687	2,566	1,479	-
2005.....	8,168	1	-	4	26	83	265	577	1,063	1,911	2,778	1,460	-
2004.....	8,140	1	-	3	36	93	246	552	972	1,946	2,859	1,432	-
2003.....	8,976	-	-	2	31	83	246	616	1,107	2,174	3,192	1,525	-
2002.....	9,013	-	-	3	32	82	229	540	1,092	2,369	3,185	1,481	-
2001.....	9,114	-	2	3	31	90	264	506	1,031	2,382	3,313	1,491	1
2000.....	9,661	-	-	3	23	78	219	492	1,111	2,730	3,443	1,562	-
1999.....	9,615	1	-	2	20	78	232	486	1,088	2,712	3,501	1,495	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Aortic aneurysm and dissection (I71)													
All races, female													
2007.....	5,443	1	-	1	6	25	86	185	390	918	2,005	1,826	-
2006.....	5,506	1	-	-	11	20	71	195	362	948	2,056	1,842	-
2005.....	5,675	-	-	-	8	31	64	181	393	959	2,193	1,846	-
2004.....	5,613	-	1	2	9	33	72	177	346	986	2,272	1,715	-
2003.....	5,834	1	-	1	10	29	70	204	370	1,097	2,279	1,773	-
2002.....	5,805	2	-	2	7	28	76	170	368	1,113	2,333	1,706	-
2001.....	6,120	-	-	1	12	27	68	155	363	1,135	2,438	1,921	-
2000.....	6,149	-	-	2	13	29	84	170	382	1,282	2,392	1,795	-
1999.....	6,192	-	1	1	11	15	97	152	349	1,289	2,435	1,842	-
White, both sexes													
2007 1/.....	11,348	1	-	6	26	73	233	515	1,192	2,295	3,959	3,047	1
2006 1/.....	11,613	-	-	7	36	74	240	561	1,122	2,302	4,213	3,058	-
2005 1/.....	12,156	1	-	3	31	76	213	517	1,209	2,539	4,532	3,035	-
2004 1/.....	12,189	1	-	5	35	85	214	528	1,060	2,637	4,698	2,926	-
2003 1/.....	13,246	-	-	1	33	77	214	586	1,243	2,949	5,063	3,080	-
2002.....	13,285	2	-	2	31	73	219	492	1,231	3,152	5,128	2,955	-
2001.....	13,652	-	2	4	32	92	227	478	1,167	3,181	5,293	3,175	1
2000.....	14,210	-	-	2	26	77	204	496	1,257	3,651	5,398	3,099	-
1999.....	14,336	1	1	2	27	70	240	467	1,250	3,652	5,529	3,097	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Aortic aneurysm and dissection (I71)													
White, male													
2007 1/.....	6,610	-	-	5	21	54	179	398	909	1,514	2,158	1,371	1
2006 1/.....	6,836	-	-	7	27	62	194	443	843	1,499	2,386	1,375	-
2005 1/.....	7,203	1	-	3	23	53	168	405	903	1,723	2,563	1,361	-
2004 1/.....	7,226	1	-	3	28	62	169	409	789	1,770	2,652	1,343	-
2003 1/.....	8,074	-	-	-	25	54	174	455	949	1,997	2,982	1,438	-
2002.....	8,132	-	-	1	24	55	164	382	926	2,174	3,000	1,406	-
2001.....	8,224	-	2	3	23	73	183	375	877	2,185	3,087	1,415	1
2000.....	8,797	-	-	2	18	56	149	384	956	2,533	3,227	1,472	-
1999.....	8,817	1	-	1	17	58	172	362	969	2,525	3,311	1,401	-
White, female													
2007 1/.....	4,738	1	-	1	5	19	54	117	283	781	1,801	1,676	-
2006 1/.....	4,777	-	-	-	9	12	46	118	279	803	1,827	1,683	-
2005 1/.....	4,953	-	-	-	8	23	45	112	306	816	1,969	1,674	-
2004 1/.....	4,963	-	-	2	7	23	45	119	271	867	2,046	1,583	-
2003 1/.....	5,172	-	-	1	8	23	40	131	294	952	2,081	1,642	-
2002.....	5,153	2	-	1	7	18	55	110	305	978	2,128	1,549	-
2001.....	5,428	-	-	1	9	19	44	103	290	996	2,206	1,760	-
2000.....	5,413	-	-	-	8	21	55	112	301	1,118	2,171	1,627	-
1999.....	5,519	-	1	1	10	12	68	105	281	1,127	2,218	1,696	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Aortic aneurysm and dissection (I71)													
Black, both sexes													
2007 1/.....	1,257	-	-	1	8	18	118	209	232	238	271	162	-
2006 1/.....	1,267	2	-	-	6	28	85	207	213	259	285	182	-
2005 1/.....	1,304	-	-	1	3	29	98	205	213	255	312	188	-
2004 1/.....	1,228	-	1	-	6	33	93	177	219	226	318	155	-
2003 1/.....	1,230	1	-	1	4	29	95	203	197	247	294	159	-
2002.....	1,218	-	-	3	5	34	80	195	197	246	288	170	-
2001.....	1,251	-	-	-	10	23	92	155	192	244	355	180	-
2000.....	1,277	-	-	3	9	25	90	137	200	281	333	199	-
1999.....	1,181	-	-	-	4	21	80	152	155	274	306	189	-
Black, male													
2007 1/.....	705	-	-	1	8	14	87	150	149	129	118	49	-
2006 1/.....	678	1	-	-	5	22	62	140	138	143	110	57	-
2005 1/.....	719	-	-	1	3	21	81	144	138	133	143	55	-
2004 1/.....	697	-	-	-	6	26	68	125	152	131	132	57	-
2003 1/.....	686	-	-	1	3	26	66	141	133	128	139	49	-
2002.....	677	-	-	2	5	25	61	141	142	133	122	46	-
2001.....	671	-	-	-	7	16	70	109	127	138	161	43	-
2000.....	671	-	-	1	5	19	64	91	125	151	155	60	-
1999.....	619	-	-	-	3	19	52	109	97	149	125	65	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Aortic aneurysm and dissection (I71)													
Black, female													
2007 1/.....	552	-	-	-	-	4	31	59	83	109	153	113	-
2006 1/.....	589	1	-	-	1	6	23	67	75	116	175	125	-
2005 1/.....	585	-	-	-	-	8	17	61	75	122	169	133	-
2004 1/.....	531	-	1	-	-	7	25	52	67	95	186	98	-
2003 1/.....	544	1	-	-	1	3	29	62	64	119	155	110	-
2002.....	541	-	-	1	-	9	19	54	55	113	166	124	-
2001.....	580	-	-	-	3	7	22	46	65	106	194	137	-
2000.....	606	-	-	2	4	6	26	46	75	130	178	139	-
1999.....	562	-	-	-	1	2	28	43	58	125	181	124	-
American Indian 2/, both sexes													
2007 1/.....	60	-	-	-	2	2	5	5	7	15	12	12	-
2006 1/.....	38	-	-	-	1	-	-	3	4	11	13	6	-
2005 1/.....	43	-	-	-	-	1	1	2	4	13	15	7	-
2004 1/.....	38	-	-	-	-	2	2	2	5	6	15	6	-
2003 1/.....	46	-	-	-	1	2	1	5	6	8	16	7	-
2002.....	49	-	-	-	-	1	1	4	6	19	13	5	-
2001.....	47	-	-	-	-	-	2	3	8	12	15	7	-
2000.....	41	-	-	-	-	-	-	1	3	21	12	4	-
1999.....	36	-	-	1	-	-	-	-	5	12	11	7	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Aortic aneurysm and dissection (I71)													
American Indian 2/, male													
2007 1/.....	41	-	-	-	2	2	4	4	4	12	8	5	-
2006 1/.....	20	-	-	-	-	-	-	2	3	5	8	2	-
2005 1/.....	21	-	-	-	-	1	1	-	2	10	7	-	-
2004 1/.....	23	-	-	-	-	1	1	2	5	2	9	3	-
2003 1/.....	26	-	-	-	1	1	-	4	4	5	7	4	-
2002.....	30	-	-	-	-	-	1	3	2	15	7	2	-
2001.....	30	-	-	-	-	-	2	3	6	7	8	4	-
2000.....	21	-	-	-	-	-	-	1	1	11	7	1	-
1999.....	19	-	-	1	-	-	-	-	3	6	6	3	-
American Indian 2/, female													
2007 1/.....	19	-	-	-	-	-	1	1	3	3	4	7	-
2006 1/.....	18	-	-	-	1	-	-	1	1	6	5	4	-
2005 1/.....	22	-	-	-	-	-	-	2	2	3	8	7	-
2004 1/.....	15	-	-	-	-	1	1	-	-	4	6	3	-
2003 1/.....	20	-	-	-	-	1	1	1	2	3	9	3	-
2002.....	19	-	-	-	-	1	-	1	4	4	6	3	-
2001.....	17	-	-	-	-	-	-	-	2	5	7	3	-
2000.....	20	-	-	-	-	-	-	-	2	10	5	3	-
1999.....	17	-	-	-	-	-	-	-	2	6	5	4	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Aortic aneurysm and dissection (I71)													
Asian or Pacific Islander 3/, both sexes													
2007 1/.....	321	-	-	-	2	6	8	23	52	68	104	58	-
2006 1/.....	320	-	-	-	-	5	15	24	27	63	111	75	-
2005 1/.....	340	-	-	-	-	8	17	34	30	63	112	76	-
2004 1/.....	298	-	-	-	4	6	9	22	34	63	100	60	-
2003 1/.....	288	-	-	1	3	4	6	26	31	67	98	52	-
2002.....	266	-	-	-	3	2	5	19	26	65	89	57	-
2001.....	284	-	-	-	1	2	11	25	27	80	88	50	-
2000.....	282	-	-	-	1	5	9	28	33	59	92	55	-
1999.....	254	-	-	-	-	2	9	19	27	63	90	44	-
Asian or Pacific Islander 3/, male													
2007 1/.....	187	-	-	-	1	4	8	15	31	43	57	28	-
2006 1/.....	198	-	-	-	-	3	13	15	20	40	62	45	-
2005 1/.....	225	-	-	-	-	8	15	28	20	45	65	44	-
2004 1/.....	194	-	-	-	2	4	8	16	26	43	66	29	-
2003 1/.....	190	-	-	1	2	2	6	16	21	44	64	34	-
2002.....	174	-	-	-	3	2	3	14	22	47	56	27	-
2001.....	189	-	-	-	1	1	9	19	21	52	57	29	-
2000.....	172	-	-	-	-	3	6	16	29	35	54	29	-
1999.....	160	-	-	-	-	1	8	15	19	32	59	26	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Aortic aneurysm and dissection (I71)

Asian or Pacific Islander 3/, female

2007 1/.....	134	-	-	-	1	2	-	8	21	25	47	30	-
2006 1/.....	122	-	-	-	-	2	2	9	7	23	49	30	-
2005 1/.....	115	-	-	-	-	-	2	6	10	18	47	32	-
2004 1/.....	104	-	-	-	2	2	1	6	8	20	34	31	-
2003 1/.....	98	-	-	-	1	2	-	10	10	23	34	18	-
2002.....	92	-	-	-	-	-	2	5	4	18	33	30	-
2001.....	95	-	-	-	-	1	2	6	6	28	31	21	-
2000.....	110	-	-	-	1	2	3	12	4	24	38	26	-
1999.....	94	-	-	-	-	1	1	4	8	31	31	18	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)													
All races, both sexes													
2007.....	8,954	12	4	7	27	37	120	340	813	1,449	2,728	3,415	2
2006.....	9,246	16	1	5	25	41	104	367	787	1,469	2,947	3,484	-
2005.....	9,774	16	3	13	18	42	138	327	810	1,614	3,196	3,597	-
2004.....	9,940	13	3	8	16	41	143	396	789	1,700	3,210	3,621	-
2003.....	9,862	10	-	9	27	39	135	379	793	1,663	3,223	3,584	-
2002.....	10,109	12	4	6	27	50	140	360	826	1,828	3,227	3,629	-
2001.....	10,084	12	2	7	18	53	166	363	801	1,828	3,281	3,553	-
2000.....	10,226	16	5	6	23	47	145	376	743	1,914	3,387	3,563	1
1999.....	10,002	11	1	8	18	48	149	333	773	1,807	3,328	3,526	-
All races, male													
2007.....	3,937	7	1	5	16	17	72	198	487	776	1,272	1,084	2
2006.....	4,104	11	-	2	12	19	50	206	460	826	1,381	1,137	-
2005.....	4,255	10	3	9	11	19	57	172	470	885	1,542	1,077	-
2004.....	4,166	6	2	5	9	20	62	228	420	872	1,463	1,079	-
2003.....	4,241	3	-	5	14	20	60	199	410	918	1,540	1,072	-
2002.....	4,296	6	4	2	11	19	62	192	434	938	1,505	1,123	-
2001.....	4,282	8	-	-	9	25	71	182	423	966	1,526	1,072	-
2000.....	4,423	6	2	4	10	25	57	207	419	1,041	1,576	1,075	1
1999.....	4,179	8	-	5	12	24	64	171	431	969	1,514	981	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)													
All races, female													
2007.....	5,017	5	3	2	11	20	48	142	326	673	1,456	2,331	-
2006.....	5,142	5	1	3	13	22	54	161	327	643	1,566	2,347	-
2005.....	5,519	6	-	4	7	23	81	155	340	729	1,654	2,520	-
2004.....	5,774	7	1	3	7	21	81	168	369	828	1,747	2,542	-
2003.....	5,621	7	-	4	13	19	75	180	383	745	1,683	2,512	-
2002.....	5,813	6	-	4	16	31	78	168	392	890	1,722	2,506	-
2001.....	5,802	4	2	7	9	28	95	181	378	862	1,755	2,481	-
2000.....	5,803	10	3	2	13	22	88	169	324	873	1,811	2,488	-
1999.....	5,823	3	1	3	6	24	85	162	342	838	1,814	2,545	-
White, both sexes													
2007 1/.....	7,577	8	4	5	21	28	95	241	626	1,181	2,335	3,032	1
2006 1/.....	7,884	13	1	2	19	31	71	272	616	1,227	2,519	3,113	-
2005 1/.....	8,329	10	3	11	12	24	101	248	636	1,326	2,755	3,203	-
2004 1/.....	8,445	10	3	6	14	27	105	300	615	1,410	2,779	3,176	-
2003 1/.....	8,415	6	-	7	16	30	92	275	641	1,387	2,786	3,175	-
2002.....	8,650	9	4	2	18	35	100	270	646	1,519	2,818	3,229	-
2001.....	8,584	10	1	6	15	37	113	280	626	1,524	2,844	3,128	-
2000.....	8,753	13	4	5	18	34	104	260	577	1,616	2,964	3,158	-
1999.....	8,616	8	1	6	15	36	116	262	618	1,532	2,901	3,121	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

White, male

2007 1/.....	3,324	5	1	4	12	14	61	143	379	635	1,099	970	1
2006 1/.....	3,537	9	-	2	8	15	29	153	356	701	1,215	1,049	-
2005 1/.....	3,622	5	3	8	6	12	40	126	372	736	1,334	980	-
2004 1/.....	3,584	5	2	3	7	14	48	177	328	741	1,297	962	-
2003 1/.....	3,634	2	-	4	9	16	43	154	331	772	1,335	968	-
2002.....	3,686	5	4	1	7	13	50	151	338	788	1,334	995	-
2001.....	3,663	6	-	-	8	17	45	136	339	818	1,342	952	-
2000.....	3,814	4	2	4	8	21	41	144	334	895	1,398	963	-
1999.....	3,627	5	-	4	10	19	54	141	352	839	1,326	877	-

White, female

2007 1/.....	4,253	3	3	1	9	14	34	98	247	546	1,236	2,062	-
2006 1/.....	4,347	4	1	-	11	16	42	119	260	526	1,304	2,064	-
2005 1/.....	4,707	5	-	3	6	12	61	122	264	590	1,421	2,223	-
2004 1/.....	4,861	5	1	3	7	13	57	123	287	669	1,482	2,214	-
2003 1/.....	4,781	4	-	3	7	14	49	121	310	615	1,451	2,207	-
2002.....	4,964	4	-	1	11	22	50	119	308	731	1,484	2,234	-
2001.....	4,921	4	1	6	7	20	68	144	287	706	1,502	2,176	-
2000.....	4,939	9	2	1	10	13	63	116	243	721	1,566	2,195	-
1999.....	4,989	3	1	2	5	17	62	121	266	693	1,575	2,244	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

Black, both sexes

2007 1/.....	1,213	2	-	1	5	8	24	86	168	236	339	343	1
2006 1/.....	1,215	3	-	2	5	9	31	87	157	218	375	328	-
2005 1/.....	1,298	4	-	2	5	17	34	70	159	261	393	353	-
2004 1/.....	1,322	2	-	2	1	13	35	83	150	259	382	395	-
2003 1/.....	1,289	4	-	2	10	8	38	95	130	253	394	355	-
2002.....	1,307	3	-	3	7	13	34	80	160	278	371	358	-
2001.....	1,381	2	1	1	2	16	50	74	161	286	399	389	-
2000.....	1,318	3	-	1	4	11	34	97	151	265	382	369	1
1999.....	1,253	3	-	-	3	9	32	60	139	247	388	372	-

Black, male

2007 1/.....	537	2	-	1	3	3	11	49	96	124	147	100	1
2006 1/.....	506	2	-	-	3	4	19	49	97	118	140	74	-
2005 1/.....	563	3	-	1	4	7	15	42	91	134	182	84	-
2004 1/.....	528	1	-	2	1	5	12	45	82	123	155	102	-
2003 1/.....	530	1	-	1	4	3	16	41	69	130	181	84	-
2002.....	534	1	-	-	2	6	10	33	85	136	152	109	-
2001.....	558	2	-	-	-	8	24	41	74	138	168	103	-
2000.....	536	2	-	-	1	3	13	55	76	130	161	94	1
1999.....	492	3	-	-	2	4	9	26	70	119	167	92	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

Black, female

2007 1/.....	676	-	-	-	2	5	13	37	72	112	192	243	-
2006 1/.....	709	1	-	2	2	5	12	38	60	100	235	254	-
2005 1/.....	735	1	-	1	1	10	19	28	68	127	211	269	-
2004 1/.....	794	1	-	-	-	8	23	38	68	136	227	293	-
2003 1/.....	759	3	-	1	6	5	22	54	61	123	213	271	-
2002.....	773	2	-	3	5	7	24	47	75	142	219	249	-
2001.....	823	-	1	1	2	8	26	33	87	148	231	286	-
2000.....	782	1	-	1	3	8	21	42	75	135	221	275	-
1999.....	761	-	-	-	1	5	23	34	69	128	221	280	-

American Indian 2/, both sexes

2007 1/.....	50	1	-	-	-	-	-	4	12	15	12	6	-
2006 1/.....	41	-	-	-	-	-	2	3	7	12	14	3	-
2005 1/.....	41	1	-	-	-	-	2	5	9	10	10	4	-
2004 1/.....	48	-	-	-	-	1	1	6	10	9	12	9	-
2003 1/.....	56	-	-	-	1	1	1	5	9	11	12	16	-
2002.....	46	-	-	-	1	-	1	5	10	11	8	10	-
2001.....	32	-	-	-	-	-	-	3	3	9	10	7	-
2000.....	46	-	-	-	-	-	3	7	5	12	8	11	-
1999.....	36	-	-	1	-	-	-	2	7	8	8	10	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

American Indian 2/, male

2007 1/.....	24	-	-	-	-	-	-	2	7	7	6	2	-
2006 1/.....	14	-	-	-	-	-	2	1	2	4	4	1	-
2005 1/.....	16	1	-	-	-	-	1	1	5	3	4	1	-
2004 1/.....	15	-	-	-	-	1	1	2	4	1	2	4	-
2003 1/.....	29	-	-	-	1	1	-	3	3	9	5	7	-
2002.....	23	-	-	-	1	-	1	3	6	6	3	3	-
2001.....	12	-	-	-	-	-	-	1	3	4	1	3	-
2000.....	16	-	-	-	-	-	-	3	4	3	3	3	-
1999.....	13	-	-	-	-	-	-	1	4	3	2	3	-

American Indian 2/, female

2007 1/.....	26	1	-	-	-	-	-	2	5	8	6	4	-
2006 1/.....	27	-	-	-	-	-	-	2	5	8	10	2	-
2005 1/.....	25	-	-	-	-	-	1	4	4	7	6	3	-
2004 1/.....	33	-	-	-	-	-	-	4	6	8	10	5	-
2003 1/.....	27	-	-	-	-	-	1	2	6	2	7	9	-
2002.....	23	-	-	-	-	-	-	2	4	5	5	7	-
2001.....	20	-	-	-	-	-	-	2	-	5	9	4	-
2000.....	30	-	-	-	-	-	3	4	1	9	5	8	-
1999.....	23	-	-	1	-	-	-	1	3	5	6	7	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	114	1	-	1	1	1	1	9	7	17	42	34	-
2006 1/.....	106	-	-	1	1	1	-	5	7	12	39	40	-
2005 1/.....	106	1	-	-	1	1	1	4	6	17	38	37	-
2004 1/.....	125	1	-	-	1	-	2	7	14	22	37	41	-
2003 1/.....	102	-	-	-	-	-	4	4	13	12	31	38	-
2002.....	106	-	-	1	1	2	5	5	10	20	30	32	-
2001.....	87	-	-	-	1	-	3	6	11	9	28	29	-
2000.....	109	-	1	-	1	2	4	12	10	21	33	25	-
1999.....	97	-	-	1	-	3	1	9	9	20	31	23	-

Asian or Pacific Islander 3/, male

2007 1/.....	52	-	-	-	1	-	-	4	5	10	20	12	-
2006 1/.....	47	-	-	-	1	-	-	3	5	3	22	13	-
2005 1/.....	54	1	-	-	1	-	1	3	2	12	22	12	-
2004 1/.....	39	-	-	-	1	-	1	4	6	7	9	11	-
2003 1/.....	48	-	-	-	-	-	1	1	7	7	19	13	-
2002.....	53	-	-	1	1	-	1	5	5	8	16	16	-
2001.....	49	-	-	-	1	-	2	4	7	6	15	14	-
2000.....	57	-	-	-	1	1	3	5	5	13	14	15	-
1999.....	47	-	-	1	-	1	1	3	5	8	19	9	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

Asian or Pacific Islander 3/, female

2007 1/.....	62	1	-	1	-	1	1	5	2	7	22	22	-
2006 1/.....	59	-	-	1	-	1	-	2	2	9	17	27	-
2005 1/.....	52	-	-	-	-	1	-	1	4	5	16	25	-
2004 1/.....	86	1	-	-	-	-	1	3	8	15	28	30	-
2003 1/.....	54	-	-	-	-	-	3	3	6	5	12	25	-
2002.....	53	-	-	-	-	2	4	-	5	12	14	16	-
2001.....	38	-	-	-	-	-	1	2	4	3	13	15	-
2000.....	52	-	1	-	-	1	1	7	5	8	19	10	-
1999.....	50	-	-	-	-	2	-	6	4	12	12	14	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other disorders of circulatory system (I80-I99)

All races, both sexes

2007.....	4,101	53	3	6	40	120	322	570	586	552	888	960	1
2006.....	3,995	35	3	5	46	114	278	537	540	553	908	976	-
2005.....	4,813	26	6	5	36	132	332	620	641	713	1,128	1,174	-
2004.....	4,673	29	2	5	35	137	313	559	623	672	1,165	1,133	-
2003.....	4,737	36	4	4	50	139	342	555	569	744	1,161	1,133	-
2002.....	4,711	27	4	2	35	139	305	531	595	697	1,253	1,123	-
2001.....	4,665	22	5	-	38	124	285	454	563	841	1,183	1,149	1
2000.....	4,603	27	4	12	37	133	319	515	516	795	1,161	1,084	-
1999.....	4,025	23	1	2	30	93	271	378	445	692	1,117	972	1

All races, male

2007.....	1,902	31	1	4	22	67	188	335	330	279	364	281	-
2006.....	1,772	18	1	3	19	54	149	323	314	248	369	274	-
2005.....	2,156	13	3	3	15	62	164	382	381	379	446	308	-
2004.....	2,083	18	-	3	19	72	162	329	351	328	505	296	-
2003.....	1,966	21	3	2	20	60	167	312	320	348	437	276	-
2002.....	1,944	15	1	-	21	69	154	284	309	323	483	285	-
2001.....	1,981	11	5	-	17	53	149	262	304	405	488	286	1
2000.....	1,981	16	2	6	16	57	190	295	264	390	472	273	-
1999.....	1,714	13	1	1	13	44	159	198	224	350	463	248	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other disorders of circulatory system (I80-I99)

All races, female

2007.....	2,199	22	2	2	18	53	134	235	256	273	524	679	1
2006.....	2,223	17	2	2	27	60	129	214	226	305	539	702	-
2005.....	2,657	13	3	2	21	70	168	238	260	334	682	866	-
2004.....	2,590	11	2	2	16	65	151	230	272	344	660	837	-
2003.....	2,771	15	1	2	30	79	175	243	249	396	724	857	-
2002.....	2,767	12	3	2	14	70	151	247	286	374	770	838	-
2001.....	2,684	11	-	-	21	71	136	192	259	436	695	863	-
2000.....	2,622	11	2	6	21	76	129	220	252	405	689	811	-
1999.....	2,311	10	-	1	17	49	112	180	221	342	654	724	1

White, both sexes

2007 1/.....	3,336	40	3	5	27	79	217	412	449	460	785	858	1
2006 1/.....	3,213	29	3	3	28	72	179	377	419	456	784	863	-
2005 1/.....	3,936	21	3	5	21	87	223	457	492	584	968	1,075	-
2004 1/.....	3,795	22	2	3	23	82	198	401	490	545	1,013	1,016	-
2003 1/.....	3,843	28	4	3	30	87	224	423	442	605	993	1,004	-
2002.....	3,853	20	2	1	21	84	203	399	457	575	1,081	1,010	-
2001.....	3,843	17	3	-	21	80	189	340	432	692	1,031	1,038	-
2000.....	3,787	20	3	9	22	82	230	392	397	637	1,017	978	-
1999.....	3,345	18	-	2	15	64	183	277	367	568	978	872	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other disorders of circulatory system (I80-I99)													
White, male													
2007 1/.....	1,523	22	1	4	10	47	131	244	246	240	323	255	-
2006 1/.....	1,414	14	1	2	9	33	104	233	250	209	315	244	-
2005 1/.....	1,723	11	2	3	8	41	114	286	295	305	382	276	-
2004 1/.....	1,683	15	-	2	10	41	108	245	285	270	439	268	-
2003 1/.....	1,569	16	3	1	11	36	129	230	252	282	367	242	-
2002.....	1,545	13	-	-	12	41	102	212	239	264	412	250	-
2001.....	1,603	8	3	-	10	34	99	204	231	333	426	255	-
2000.....	1,618	13	1	5	9	34	141	228	201	323	419	244	-
1999.....	1,399	9	-	1	8	30	106	142	192	287	401	223	-
White, female													
2007 1/.....	1,813	18	2	1	17	32	86	168	203	220	462	603	1
2006 1/.....	1,799	15	2	1	19	39	75	144	169	247	469	619	-
2005 1/.....	2,213	10	1	2	13	46	109	171	197	279	586	799	-
2004 1/.....	2,112	7	2	1	13	41	90	156	205	275	574	748	-
2003 1/.....	2,274	12	1	2	19	51	95	193	190	323	626	762	-
2002.....	2,308	7	2	1	9	43	101	187	218	311	669	760	-
2001.....	2,240	9	-	-	11	46	90	136	201	359	605	783	-
2000.....	2,169	7	2	4	13	48	89	164	196	314	598	734	-
1999.....	1,946	9	-	1	7	34	77	135	175	281	577	649	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other disorders of circulatory system (I80-I99)

Black, both sexes

2007 1/.....	701	12	-	-	11	40	101	145	125	83	92	92	-
2006 1/.....	713	4	-	2	18	41	92	148	110	88	111	99	-
2005 1/.....	796	4	2	-	15	42	99	152	137	113	141	91	-
2004 1/.....	799	7	-	1	10	51	111	145	122	115	135	102	-
2003 1/.....	824	6	-	-	19	49	112	124	119	129	151	115	-
2002.....	797	7	1	1	13	54	95	121	130	114	160	101	-
2001.....	742	5	2	-	14	43	89	100	123	131	136	98	1
2000.....	761	5	1	3	14	49	87	118	109	147	132	96	-
1999.....	616	4	1	-	14	28	81	96	69	109	125	89	-

Black, male

2007 1/.....	348	9	-	-	10	20	54	82	77	37	36	23	-
2006 1/.....	324	2	-	1	10	20	40	85	60	38	43	25	-
2005 1/.....	388	2	-	-	7	18	44	89	77	65	59	27	-
2004 1/.....	359	3	-	-	7	28	53	78	60	51	54	25	-
2003 1/.....	368	5	-	-	9	23	36	76	64	62	62	31	-
2002.....	366	2	1	-	9	27	49	64	66	54	64	30	-
2001.....	339	3	2	-	5	18	47	49	72	62	54	26	1
2000.....	335	2	1	1	7	22	48	66	56	62	44	26	-
1999.....	282	3	1	-	4	14	49	53	28	56	55	19	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other disorders of circulatory system (I80-I99)

Black, female

2007 1/.....	353	3	-	-	1	20	47	63	48	46	56	69	-
2006 1/.....	389	2	-	1	8	21	52	63	50	50	68	74	-
2005 1/.....	408	2	2	-	8	24	55	63	60	48	82	64	-
2004 1/.....	440	4	-	1	3	23	58	67	62	64	81	77	-
2003 1/.....	456	1	-	-	10	26	76	48	55	67	89	84	-
2002.....	431	5	-	1	4	27	46	57	64	60	96	71	-
2001.....	403	2	-	-	9	25	42	51	51	69	82	72	-
2000.....	426	3	-	2	7	27	39	52	53	85	88	70	-
1999.....	334	1	-	-	10	14	32	43	41	53	70	70	-

American Indian 2/, both sexes

2007 1/.....	17	-	-	-	1	-	2	6	2	2	3	1	-
2006 1/.....	22	1	-	-	-	1	3	6	3	3	1	4	-
2005 1/.....	27	1	1	-	-	-	6	7	2	4	6	-	-
2004 1/.....	20	-	-	-	-	2	2	3	4	3	3	3	-
2003 1/.....	15	-	-	-	-	-	2	3	3	1	4	2	-
2002.....	23	-	-	-	-	-	2	6	5	3	3	4	-
2001.....	25	-	-	-	1	1	2	8	3	6	2	2	-
2000.....	18	-	-	-	1	1	-	2	2	7	4	1	-
1999.....	13	-	-	-	-	1	1	2	2	1	6	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other disorders of circulatory system (I80-I99)

American Indian 2/, male

2007 1/.....	6	-	-	-	1	-	2	3	-	-	-	-	-
2006 1/.....	8	1	-	-	-	1	2	3	-	-	-	1	-
2005 1/.....	16	-	1	-	-	-	4	5	2	2	2	-	-
2004 1/.....	12	-	-	-	-	2	1	2	2	2	2	1	-
2003 1/.....	7	-	-	-	-	-	1	3	1	-	2	-	-
2002.....	13	-	-	-	-	-	1	4	2	3	2	1	-
2001.....	14	-	-	-	-	1	1	6	-	5	-	1	-
2000.....	8	-	-	-	-	1	-	-	1	3	3	-	-
1999.....	4	-	-	-	-	-	-	2	-	1	1	-	-

American Indian 2/, female

2007 1/.....	11	-	-	-	-	-	-	3	2	2	3	1	-
2006 1/.....	14	-	-	-	-	-	1	3	3	3	1	3	-
2005 1/.....	11	1	-	-	-	-	2	2	-	2	4	-	-
2004 1/.....	8	-	-	-	-	-	1	1	2	1	1	2	-
2003 1/.....	8	-	-	-	-	-	1	-	2	1	2	2	-
2002.....	10	-	-	-	-	-	1	2	3	-	1	3	-
2001.....	11	-	-	-	1	-	1	2	3	1	2	1	-
2000.....	10	-	-	-	1	-	-	2	1	4	1	1	-
1999.....	9	-	-	-	-	1	1	-	2	-	5	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other disorders of circulatory system (I80-I99)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	47	1	-	1	1	1	2	7	10	7	8	9	-
2006 1/.....	47	1	-	-	-	-	4	6	8	6	12	10	-
2005 1/.....	54	-	-	-	-	3	4	4	10	12	13	8	-
2004 1/.....	59	-	-	1	2	2	2	10	7	9	14	12	-
2003 1/.....	55	2	-	1	1	3	4	5	5	9	13	12	-
2002.....	38	-	1	-	1	1	5	5	3	5	9	8	-
2001.....	55	-	-	-	2	-	5	6	5	12	14	11	-
2000.....	37	2	-	-	-	1	2	3	8	4	8	9	-
1999.....	51	1	-	-	1	-	6	3	7	14	8	11	-

Asian or Pacific Islander 3/, male

2007 1/.....	25	-	-	-	1	-	1	6	7	2	5	3	-
2006 1/.....	26	1	-	-	-	-	3	2	4	1	11	4	-
2005 1/.....	29	-	-	-	-	3	2	2	7	7	3	5	-
2004 1/.....	29	-	-	1	2	1	-	4	4	5	10	2	-
2003 1/.....	22	-	-	1	-	1	1	3	3	4	6	3	-
2002.....	20	-	-	-	-	1	2	4	2	2	5	4	-
2001.....	25	-	-	-	2	-	2	3	1	5	8	4	-
2000.....	20	1	-	-	-	-	1	1	6	2	6	3	-
1999.....	29	1	-	-	1	-	4	1	4	6	6	6	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other disorders of circulatory system (I80-I99)

Asian or Pacific Islander 3/, female

2007 1/.....	22	1	-	1	-	1	1	1	3	5	3	6	-
2006 1/.....	21	-	-	-	-	-	1	4	4	5	1	6	-
2005 1/.....	25	-	-	-	-	-	2	2	3	5	10	3	-
2004 1/.....	30	-	-	-	-	1	2	6	3	4	4	10	-
2003 1/.....	33	2	-	-	1	2	3	2	2	5	7	9	-
2002.....	18	-	1	-	1	-	3	1	1	3	4	4	-
2001.....	30	-	-	-	-	-	3	3	4	7	6	7	-
2000.....	17	1	-	-	-	1	1	2	2	2	2	6	-
1999.....	22	-	-	-	-	-	2	2	3	8	2	5	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Influenza and pneumonia (J09-J18)													
All races, both sexes													
2007.....	52,717	222	109	103	163	331	784	1,909	3,152	5,547	14,859	25,535	3
2006.....	56,326	263	125	68	184	335	841	2,007	3,154	6,061	16,668	26,617	3
2005.....	63,001	265	110	106	172	354	934	2,183	3,422	6,623	18,563	30,267	2
2004.....	59,664	273	119	82	185	303	891	1,897	3,154	6,382	18,066	28,312	-
2003.....	65,163	322	163	147	224	373	992	2,140	3,130	6,831	19,442	31,397	2
2002.....	65,681	263	110	91	167	345	971	1,918	2,987	6,847	19,984	31,995	3
2001.....	62,034	299	112	92	181	339	983	1,801	2,704	6,650	18,677	30,191	5
2000.....	65,313	289	103	87	189	364	1,068	1,774	2,879	7,189	19,821	31,547	3
1999.....	63,730	320	130	93	179	339	1,063	1,697	2,625	6,861	19,192	31,229	2
All races, male													
2007.....	24,071	135	55	52	91	187	443	1,126	1,881	3,044	7,412	9,642	3
2006.....	25,650	163	63	35	101	175	462	1,186	1,877	3,371	8,315	9,900	2
2005.....	28,052	148	56	53	117	196	540	1,299	1,988	3,648	9,203	10,802	2
2004.....	26,861	166	47	47	118	194	494	1,142	1,823	3,605	9,013	10,212	-
2003.....	28,778	192	91	70	124	229	560	1,279	1,771	3,882	9,629	10,950	1
2002.....	28,918	162	63	38	97	190	546	1,138	1,738	3,850	9,895	11,199	2
2001.....	27,342	167	57	47	106	191	556	1,099	1,587	3,732	9,294	10,502	4
2000.....	28,658	163	54	48	102	204	630	1,066	1,680	4,013	9,751	10,945	2
1999.....	27,718	171	66	53	90	197	592	1,059	1,462	3,893	9,458	10,676	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Influenza and pneumonia (J09-J18)													
All races, female													
2007.....	28,646	87	54	51	72	144	341	783	1,271	2,503	7,447	15,893	-
2006.....	30,676	100	62	33	83	160	379	821	1,277	2,690	8,353	16,717	1
2005.....	34,949	117	54	53	55	158	394	884	1,434	2,975	9,360	19,465	-
2004.....	32,803	107	72	35	67	109	397	755	1,331	2,777	9,053	18,100	-
2003.....	36,385	130	72	77	100	144	432	861	1,359	2,949	9,813	20,447	1
2002.....	36,763	101	47	53	70	155	425	780	1,249	2,997	10,089	20,796	1
2001.....	34,692	132	55	45	75	148	427	702	1,117	2,918	9,383	19,689	1
2000.....	36,655	126	49	39	87	160	438	708	1,199	3,176	10,070	20,602	1
1999.....	36,012	149	64	40	89	142	471	638	1,163	2,968	9,734	20,553	1
White, both sexes													
2007 1/.....	45,947	132	73	77	129	225	580	1,403	2,483	4,551	13,074	23,217	3
2006 1/.....	49,401	161	83	53	138	238	628	1,492	2,557	5,061	14,750	24,237	3
2005 1/.....	55,540	158	76	77	134	262	683	1,623	2,700	5,490	16,547	27,788	2
2004 1/.....	52,430	161	77	57	142	228	635	1,399	2,510	5,330	16,014	25,877	-
2003 1/.....	57,645	209	115	106	173	258	709	1,579	2,452	5,703	17,422	28,918	1
2002.....	58,346	162	63	69	123	248	697	1,456	2,357	5,830	17,898	29,440	3
2001.....	54,774	168	78	67	126	232	678	1,316	2,141	5,644	16,621	27,699	4
2000.....	57,914	172	69	70	138	261	767	1,297	2,258	5,969	17,811	29,101	1
1999.....	56,694	184	75	72	121	222	714	1,223	2,059	5,832	17,405	28,785	2

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Influenza and pneumonia (J09-J18)													
White, male													
2007 1/.....	20,720	85	32	36	74	127	323	810	1,486	2,475	6,517	8,752	3
2006 1/.....	22,310	103	43	31	75	128	349	874	1,522	2,810	7,382	8,991	2
2005 1/.....	24,425	96	36	36	90	142	391	959	1,545	3,014	8,225	9,889	2
2004 1/.....	23,348	100	32	32	98	143	348	846	1,454	2,985	7,979	9,331	-
2003 1/.....	25,009	127	67	50	95	165	396	949	1,347	3,212	8,577	10,024	-
2002.....	25,381	105	35	26	74	137	393	857	1,358	3,291	8,843	10,260	2
2001.....	23,744	100	41	34	79	127	382	791	1,239	3,144	8,225	9,578	4
2000.....	25,002	98	30	39	79	138	455	759	1,302	3,314	8,742	10,045	1
1999.....	24,281	103	33	42	62	134	390	750	1,137	3,288	8,566	9,775	1
White, female													
2007 1/.....	25,227	47	41	41	55	98	257	593	997	2,076	6,557	14,465	-
2006 1/.....	27,091	58	40	22	63	110	279	618	1,035	2,251	7,368	15,246	1
2005 1/.....	31,115	62	40	41	44	120	292	664	1,155	2,476	8,322	17,899	-
2004 1/.....	29,082	61	45	25	44	85	287	553	1,056	2,345	8,035	16,546	-
2003 1/.....	32,636	82	48	56	78	93	313	630	1,105	2,491	8,845	18,894	1
2002.....	32,965	57	28	43	49	111	304	599	999	2,539	9,055	19,180	1
2001.....	31,030	68	37	33	47	105	296	525	902	2,500	8,396	18,121	-
2000.....	32,912	74	39	31	59	123	312	538	956	2,655	9,069	19,056	-
1999.....	32,413	81	42	30	59	88	324	473	922	2,544	8,839	19,010	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Influenza and pneumonia (J09-J18)													
Black, both sexes													
2007 1/.....	5,155	75	29	21	23	94	178	439	573	800	1,312	1,611	-
2006 1/.....	5,311	90	34	9	41	77	186	457	511	806	1,425	1,675	-
2005 1/.....	5,780	98	29	22	31	78	217	499	611	918	1,514	1,763	-
2004 1/.....	5,700	93	35	20	40	66	229	447	552	863	1,564	1,791	-
2003 1/.....	5,872	93	28	30	42	102	252	497	569	927	1,530	1,801	1
2002.....	5,871	87	41	19	38	83	252	403	560	848	1,603	1,937	-
2001.....	5,771	106	26	17	45	89	276	436	480	806	1,597	1,893	-
2000.....	5,990	103	30	11	44	93	257	425	538	997	1,569	1,921	2
1999.....	5,876	115	43	15	49	107	312	431	498	889	1,426	1,991	-
Black, male													
2007 1/.....	2,498	42	16	13	12	51	103	271	332	456	642	560	-
2006 1/.....	2,486	55	16	3	23	37	98	276	306	438	669	565	-
2005 1/.....	2,729	47	17	13	23	45	129	302	377	512	692	572	-
2004 1/.....	2,775	55	12	12	19	44	130	268	322	523	788	602	-
2003 1/.....	2,906	54	14	15	24	59	143	292	358	556	775	615	1
2002.....	2,768	48	22	10	20	45	138	241	341	462	798	643	-
2001.....	2,813	53	14	9	23	53	156	275	298	479	823	630	-
2000.....	2,915	56	21	7	21	59	152	272	327	571	773	655	1
1999.....	2,825	58	25	8	23	58	178	282	289	523	695	686	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Influenza and pneumonia (J09-J18)

Black, female

2007 1/.....	2,657	33	13	8	11	43	75	168	241	344	670	1,051	-
2006 1/.....	2,825	35	18	6	18	40	88	181	205	368	756	1,110	-
2005 1/.....	3,051	51	12	9	8	33	88	197	234	406	822	1,191	-
2004 1/.....	2,925	38	23	8	21	22	99	179	230	340	776	1,189	-
2003 1/.....	2,966	39	14	15	18	43	109	205	211	371	755	1,186	-
2002.....	3,103	39	19	9	18	38	114	162	219	386	805	1,294	-
2001.....	2,958	53	12	8	22	36	120	161	182	327	774	1,263	-
2000.....	3,075	47	9	4	23	34	105	153	211	426	796	1,266	1
1999.....	3,051	57	18	7	26	49	134	149	209	366	731	1,305	-

American Indian 2/, both sexes

2007 1/.....	280	8	2	-	4	4	18	28	35	39	67	75	-
2006 1/.....	267	3	4	2	3	7	16	24	26	40	67	75	-
2005 1/.....	354	5	1	1	1	5	18	27	43	50	98	105	-
2004 1/.....	291	12	4	1	-	7	13	19	32	38	91	74	-
2003 1/.....	390	12	5	3	4	6	21	35	49	56	82	117	-
2002.....	293	6	1	-	3	7	13	23	25	35	81	99	-
2001.....	318	17	3	4	3	9	19	20	25	42	85	91	-
2000.....	289	6	-	3	2	5	28	21	24	45	71	84	-
1999.....	315	9	3	3	5	4	20	26	30	37	72	106	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Influenza and pneumonia (J09-J18)

American Indian 2/, male

2007 1/.....	150	5	2	-	1	3	12	23	27	22	32	23	-
2006 1/.....	137	-	3	-	1	2	10	14	11	27	36	33	-
2005 1/.....	180	2	-	1	-	4	12	16	24	24	52	45	-
2004 1/.....	142	7	1	-	-	5	7	12	13	18	46	33	-
2003 1/.....	184	7	2	2	2	1	14	21	27	31	40	37	-
2002.....	133	3	1	-	1	4	9	14	13	19	33	36	-
2001.....	160	11	-	1	1	6	13	12	13	23	42	38	-
2000.....	149	2	-	-	1	3	18	16	16	29	33	31	-
1999.....	147	4	3	2	4	1	12	17	13	16	39	36	-

American Indian 2/, female

2007 1/.....	130	3	-	-	3	1	6	5	8	17	35	52	-
2006 1/.....	130	3	1	2	2	5	6	10	15	13	31	42	-
2005 1/.....	174	3	1	-	1	1	6	11	19	26	46	60	-
2004 1/.....	149	5	3	1	-	2	6	7	19	20	45	41	-
2003 1/.....	206	5	3	1	2	5	7	14	22	25	42	80	-
2002.....	160	3	-	-	2	3	4	9	12	16	48	63	-
2001.....	158	6	3	3	2	3	6	8	12	19	43	53	-
2000.....	140	4	-	3	1	2	10	5	8	16	38	53	-
1999.....	168	5	-	1	1	3	8	9	17	21	33	70	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Influenza and pneumonia (J09-J18)													
Asian or Pacific Islander 3/, both sexes													
2007 1/.....	1,335	7	5	5	7	8	8	39	61	157	406	632	-
2006 1/.....	1,347	9	4	4	2	13	11	34	60	154	426	630	-
2005 1/.....	1,327	4	4	6	6	9	16	34	68	165	404	611	-
2004 1/.....	1,243	7	3	4	3	2	14	32	60	151	397	570	-
2003 1/.....	1,256	8	15	8	5	7	10	29	60	145	408	561	-
2002.....	1,171	8	5	3	3	7	9	36	45	134	402	519	-
2001.....	1,171	8	5	4	7	9	10	29	58	158	374	508	1
2000.....	1,120	8	4	3	5	5	16	31	59	178	370	441	-
1999.....	845	12	9	3	4	6	17	17	38	103	289	347	-
Asian or Pacific Islander 3/, male													
2007 1/.....	703	3	5	3	4	6	5	22	36	91	221	307	-
2006 1/.....	717	5	1	1	2	8	5	22	38	96	228	311	-
2005 1/.....	718	3	3	3	4	5	8	22	42	98	234	296	-
2004 1/.....	596	4	2	3	1	2	9	16	34	79	200	246	-
2003 1/.....	679	4	8	3	3	4	7	17	39	83	237	274	-
2002.....	636	6	5	2	2	4	6	26	26	78	221	260	-
2001.....	625	3	2	3	3	5	5	21	37	86	204	256	-
2000.....	592	7	3	2	1	4	5	19	35	99	203	214	-
1999.....	465	6	5	1	1	4	12	10	23	66	158	179	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Influenza and pneumonia (J09-J18)

Asian or Pacific Islander 3/, female

2007 1/.....	632	4	-	2	3	2	3	17	25	66	185	325	-
2006 1/.....	630	4	3	3	-	5	6	12	22	58	198	319	-
2005 1/.....	609	1	1	3	2	4	8	12	26	67	170	315	-
2004 1/.....	647	3	1	1	2	-	5	16	26	72	197	324	-
2003 1/.....	577	4	7	5	2	3	3	12	21	62	171	287	-
2002.....	535	2	-	1	1	3	3	10	19	56	181	259	-
2001.....	546	5	3	1	4	4	5	8	21	72	170	252	1
2000.....	528	1	1	1	4	1	11	12	24	79	167	227	-
1999.....	380	6	4	2	3	2	5	7	15	37	131	168	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Influenza (J09-J11)													
All races, both sexes													
2007.....	411	13	19	35	10	9	13	19	37	38	79	139	-
2006.....	849	18	19	25	13	6	10	26	43	81	236	372	-
2005.....	1,812	19	19	20	12	11	19	45	66	137	482	982	-
2004.....	1,100	18	18	13	7	4	16	25	55	99	291	554	-
2003.....	1,792	31	59	46	30	13	24	61	85	183	457	803	-
2002.....	727	7	5	12	7	4	8	17	30	56	177	404	-
2001.....	257	7	6	12	7	10	6	15	21	21	56	96	-
2000.....	1,765	9	10	11	13	13	22	48	93	199	492	855	-
1999.....	1,665	13	12	11	11	9	16	26	71	164	451	881	-
All races, male													
2007.....	187	9	13	17	1	1	6	11	20	19	37	53	-
2006.....	362	11	12	11	5	4	4	13	20	42	104	136	-
2005.....	657	9	10	10	6	3	11	26	37	61	192	292	-
2004.....	421	10	8	8	2	2	7	8	25	58	125	168	-
2003.....	699	18	33	19	12	9	10	33	42	89	189	245	-
2002.....	289	5	3	2	4	2	4	12	20	32	81	124	-
2001.....	101	6	2	4	6	6	2	9	9	8	20	29	-
2000.....	664	5	4	5	6	5	14	25	44	106	196	254	-
1999.....	602	6	9	5	2	5	9	15	37	74	202	238	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Influenza (J09-J11)

All races, female

2007.....	224	4	6	18	9	8	7	8	17	19	42	86	-
2006.....	487	7	7	14	8	2	6	13	23	39	132	236	-
2005.....	1,155	10	9	10	6	8	8	19	29	76	290	690	-
2004.....	679	8	10	5	5	2	9	17	30	41	166	386	-
2003.....	1,093	13	26	27	18	4	14	28	43	94	268	558	-
2002.....	438	2	2	10	3	2	4	5	10	24	96	280	-
2001.....	156	1	4	8	1	4	4	6	12	13	36	67	-
2000.....	1,101	4	6	6	7	8	8	23	49	93	296	601	-
1999.....	1,063	7	3	6	9	4	7	11	34	90	249	643	-

White, both sexes

2007 1/.....	374	9	13	32	10	7	11	16	32	34	74	136	-
2006 1/.....	788	13	11	20	9	4	8	23	41	75	225	359	-
2005 1/.....	1,710	14	15	13	10	10	13	35	59	128	459	954	-
2004 1/.....	1,033	12	14	11	6	3	14	20	50	92	271	540	-
2003 1/.....	1,653	23	42	30	20	10	20	49	76	170	439	774	-
2002.....	686	5	2	10	5	4	8	16	25	47	172	392	-
2001.....	229	7	5	7	5	7	4	15	17	19	52	91	-
2000.....	1,662	7	9	9	12	11	18	40	87	180	469	820	-
1999.....	1,574	7	8	10	10	8	10	22	65	148	429	857	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Influenza (J09-J11)													
White, male													
2007 1/.....	169	6	8	15	1	1	5	11	18	18	34	52	-
2006 1/.....	332	9	8	9	2	3	3	12	19	40	97	130	-
2005 1/.....	609	9	8	7	4	3	8	19	32	58	183	278	-
2004 1/.....	388	5	5	7	2	1	6	6	23	54	114	165	-
2003 1/.....	640	14	25	12	7	8	6	27	36	84	182	239	-
2002.....	270	3	1	2	2	2	4	12	16	28	78	122	-
2001.....	88	6	2	3	4	3	2	9	6	6	20	27	-
2000.....	617	5	4	4	5	3	10	19	41	99	187	240	-
1999.....	569	1	6	5	2	5	6	13	36	67	194	234	-
White, female													
2007 1/.....	205	3	5	17	9	6	6	5	14	16	40	84	-
2006 1/.....	456	4	3	11	7	1	5	11	22	35	128	229	-
2005 1/.....	1,101	5	7	6	6	7	5	16	27	70	276	676	-
2004 1/.....	645	7	9	4	4	2	8	14	27	38	157	375	-
2003 1/.....	1,013	9	17	18	13	2	14	22	40	86	257	535	-
2002.....	416	2	1	8	3	2	4	4	9	19	94	270	-
2001.....	141	1	3	4	1	4	2	6	11	13	32	64	-
2000.....	1,045	2	5	5	7	8	8	21	46	81	282	580	-
1999.....	1,005	6	2	5	8	3	4	9	29	81	235	623	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Influenza (J09-J11)

Black, both sexes

2007 1/.....	26	4	5	2	-	2	2	2	4	2	2	1	-
2006 1/.....	41	5	4	1	3	2	1	3	2	3	10	7	-
2005 1/.....	69	5	3	5	1	1	6	7	7	5	14	15	-
2004 1/.....	50	4	3	1	1	1	1	3	4	7	16	9	-
2003 1/.....	97	7	6	11	7	3	1	12	7	10	13	20	-
2002.....	30	1	3	2	2	-	-	-	4	6	4	8	-
2001.....	20	-	-	2	1	2	1	-	3	2	4	5	-
2000.....	81	2	1	1	1	1	3	7	6	16	13	30	-
1999.....	66	4	1	-	1	1	5	3	5	13	15	18	-

Black, male

2007 1/.....	13	3	4	1	-	-	1	-	2	1	1	-	-
2006 1/.....	21	2	3	1	2	1	-	1	1	1	6	3	-
2005 1/.....	32	-	2	3	1	-	3	5	5	2	3	8	-
2004 1/.....	25	4	2	-	-	1	-	2	1	4	10	1	-
2003 1/.....	37	3	3	4	3	1	1	6	5	4	4	3	-
2002.....	15	1	2	-	2	-	-	-	3	3	2	2	-
2001.....	10	-	-	-	1	2	-	-	3	2	-	2	-
2000.....	35	-	-	1	1	1	3	5	3	5	5	11	-
1999.....	24	3	1	-	-	-	2	2	1	6	6	3	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Influenza (J09-J11)

Black, female

2007 1/.....	13	1	1	1	-	2	1	2	2	1	1	1	-
2006 1/.....	20	3	1	-	1	1	1	2	1	2	4	4	-
2005 1/.....	37	5	1	2	-	1	3	2	2	3	11	7	-
2004 1/.....	25	-	1	1	1	-	1	1	3	3	6	8	-
2003 1/.....	60	4	3	7	4	2	-	6	2	6	9	17	-
2002.....	15	-	1	2	-	-	-	-	1	3	2	6	-
2001.....	10	-	-	2	-	-	1	-	-	-	4	3	-
2000.....	46	2	1	-	-	-	-	2	3	11	8	19	-
1999.....	42	1	-	-	1	1	3	1	4	7	9	15	-

American Indian 2/, both sexes

2007 1/.....	5	-	1	-	-	-	-	-	1	1	1	1	-
2006 1/.....	3	-	-	1	-	-	1	-	-	-	-	1	-
2005 1/.....	11	-	-	-	-	-	-	1	-	3	2	5	-
2004 1/.....	3	-	-	-	-	-	-	1	-	-	1	1	-
2003 1/.....	17	1	3	1	2	-	1	-	-	1	3	5	-
2002.....	5	-	-	-	-	-	-	-	1	2	1	1	-
2001.....	3	-	1	2	-	-	-	-	-	-	-	-	-
2000.....	4	-	-	1	-	-	-	-	-	-	1	2	-
1999.....	5	1	-	1	-	-	-	-	-	1	2	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Influenza (J09-J11)

American Indian 2/, male

2007 1/.....	2	-	1	-	-	-	-	-	-	-	-	1	-
2006 1/.....	1	-	-	-	-	-	1	-	-	-	-	-	-
2005 1/.....	6	-	-	-	-	-	-	1	-	1	2	2	-
2004 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 1/.....	7	1	1	-	1	-	1	-	-	1	1	1	-
2002.....	3	-	-	-	-	-	-	-	1	1	1	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	1	-
1999.....	2	1	-	-	-	-	-	-	-	-	1	-	-

American Indian 2/, female

2007 1/.....	3	-	-	-	-	-	-	-	1	1	1	-	-
2006 1/.....	2	-	-	1	-	-	-	-	-	-	-	1	-
2005 1/.....	5	-	-	-	-	-	-	-	-	2	-	3	-
2004 1/.....	3	-	-	-	-	-	-	1	-	-	1	1	-
2003 1/.....	10	-	2	1	1	-	-	-	-	-	2	4	-
2002.....	2	-	-	-	-	-	-	-	-	1	-	1	-
2001.....	3	-	1	2	-	-	-	-	-	-	-	-	-
2000.....	3	-	-	1	-	-	-	-	-	-	1	1	-
1999.....	3	-	-	1	-	-	-	-	-	1	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Influenza (J09-J11)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	6	-	-	1	-	-	-	1	-	1	2	1	-
2006 1/.....	17	-	4	3	1	-	-	-	-	3	1	5	-
2005 1/.....	22	-	1	2	1	-	-	2	-	1	7	8	-
2004 1/.....	14	2	1	1	-	-	1	1	1	-	3	4	-
2003 1/.....	25	-	8	4	1	-	2	-	2	2	2	4	-
2002.....	6	1	-	-	-	-	-	1	-	1	-	3	-
2001.....	5	-	-	1	1	1	1	-	1	-	-	-	-
2000.....	18	-	-	-	-	1	1	1	-	3	9	3	-
1999.....	20	1	3	-	-	-	1	1	1	2	5	6	-

Asian or Pacific Islander 3/, male

2007 1/.....	3	-	-	1	-	-	-	-	-	-	2	-	-
2006 1/.....	8	-	1	1	1	-	-	-	-	1	1	3	-
2005 1/.....	10	-	-	-	1	-	-	1	-	-	4	4	-
2004 1/.....	8	1	1	1	-	-	1	-	1	-	1	2	-
2003 1/.....	15	-	4	3	1	-	2	-	1	-	2	2	-
2002.....	1	1	-	-	-	-	-	-	-	-	-	-	-
2001.....	3	-	-	1	1	1	-	-	-	-	-	-	-
2000.....	11	-	-	-	-	1	1	1	-	2	4	2	-
1999.....	7	1	2	-	-	-	1	-	-	1	1	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Influenza (J09-J11)

Asian or Pacific Islander 3/, female

2007 1/.....	3	-	-	-	-	-	-	1	-	1	-	1	-
2006 1/.....	9	-	3	2	-	-	-	-	-	2	-	2	-
2005 1/.....	12	-	1	2	-	-	-	1	-	1	3	4	-
2004 1/.....	6	1	-	-	-	-	-	1	-	-	2	2	-
2003 1/.....	10	-	4	1	-	-	-	-	1	2	-	2	-
2002.....	5	-	-	-	-	-	-	1	-	1	-	3	-
2001.....	2	-	-	-	-	-	1	-	1	-	-	-	-
2000.....	7	-	-	-	-	-	-	-	-	1	5	1	-
1999.....	13	-	1	-	-	-	-	1	1	1	4	5	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumonia (J12-J18)													
All races, both sexes													
2007.....	52,306	209	90	68	153	322	771	1,890	3,115	5,509	14,780	25,396	3
2006.....	55,477	245	106	43	171	329	831	1,981	3,111	5,980	16,432	26,245	3
2005.....	61,189	246	91	86	160	343	915	2,138	3,356	6,486	18,081	29,285	2
2004.....	58,564	255	101	69	178	299	875	1,872	3,099	6,283	17,775	27,758	-
2003.....	63,371	291	104	101	194	360	968	2,079	3,045	6,648	18,985	30,594	2
2002.....	64,954	256	105	79	160	341	963	1,901	2,957	6,791	19,807	31,591	3
2001.....	61,777	292	106	80	174	329	977	1,786	2,683	6,629	18,621	30,095	5
2000.....	63,548	280	93	76	176	351	1,046	1,726	2,786	6,990	19,329	30,692	3
1999.....	62,065	307	118	82	168	330	1,047	1,671	2,554	6,697	18,741	30,348	2
All races, male													
2007.....	23,884	126	42	35	90	186	437	1,115	1,861	3,025	7,375	9,589	3
2006.....	25,288	152	51	24	96	171	458	1,173	1,857	3,329	8,211	9,764	2
2005.....	27,395	139	46	43	111	193	529	1,273	1,951	3,587	9,011	10,510	2
2004.....	26,440	156	39	39	116	192	487	1,134	1,798	3,547	8,888	10,044	-
2003.....	28,079	174	58	51	112	220	550	1,246	1,729	3,793	9,440	10,705	1
2002.....	28,629	157	60	36	93	188	542	1,126	1,718	3,818	9,814	11,075	2
2001.....	27,241	161	55	43	100	185	554	1,090	1,578	3,724	9,274	10,473	4
2000.....	27,994	158	50	43	96	199	616	1,041	1,636	3,907	9,555	10,691	2
1999.....	27,116	165	57	48	88	192	583	1,044	1,425	3,819	9,256	10,438	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumonia (J12-J18)													
All races, female													
2007.....	28,422	83	48	33	63	136	334	775	1,254	2,484	7,405	15,807	-
2006.....	30,189	93	55	19	75	158	373	808	1,254	2,651	8,221	16,481	1
2005.....	33,794	107	45	43	49	150	386	865	1,405	2,899	9,070	18,775	-
2004.....	32,124	99	62	30	62	107	388	738	1,301	2,736	8,887	17,714	-
2003.....	35,292	117	46	50	82	140	418	833	1,316	2,855	9,545	19,889	1
2002.....	36,325	99	45	43	67	153	421	775	1,239	2,973	9,993	20,516	1
2001.....	34,536	131	51	37	74	144	423	696	1,105	2,905	9,347	19,622	1
2000.....	35,554	122	43	33	80	152	430	685	1,150	3,083	9,774	20,001	1
1999.....	34,949	142	61	34	80	138	464	627	1,129	2,878	9,485	19,910	1
White, both sexes													
2007 1/.....	45,573	123	60	45	119	218	569	1,387	2,451	4,517	13,000	23,081	3
2006 1/.....	48,613	148	72	33	129	234	620	1,469	2,516	4,986	14,525	23,878	3
2005 1/.....	53,830	144	61	64	124	252	670	1,588	2,641	5,362	16,088	26,834	2
2004 1/.....	51,397	149	63	46	136	225	621	1,379	2,460	5,238	15,743	25,337	-
2003 1/.....	55,992	186	73	76	153	248	689	1,530	2,376	5,533	16,983	28,144	1
2002.....	57,660	157	61	59	118	244	689	1,440	2,332	5,783	17,726	29,048	3
2001.....	54,545	161	73	60	121	225	674	1,301	2,124	5,625	16,569	27,608	4
2000.....	56,252	165	60	61	126	250	749	1,257	2,171	5,789	17,342	28,281	1
1999.....	55,120	177	67	62	111	214	704	1,201	1,994	5,684	16,976	27,928	2

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumonia (J12-J18)													
White, male													
2007 1/.....	20,551	79	24	21	73	126	318	799	1,468	2,457	6,483	8,700	3
2006 1/.....	21,978	94	35	22	73	125	346	862	1,503	2,770	7,285	8,861	2
2005 1/.....	23,816	87	28	29	86	139	383	940	1,513	2,956	8,042	9,611	2
2004 1/.....	22,960	95	27	25	96	142	342	840	1,431	2,931	7,865	9,166	-
2003 1/.....	24,369	113	42	38	88	157	390	922	1,311	3,128	8,395	9,785	-
2002.....	25,111	102	34	24	72	135	389	845	1,342	3,263	8,765	10,138	2
2001.....	23,656	94	39	31	75	124	380	782	1,233	3,138	8,205	9,551	4
2000.....	24,385	93	26	35	74	135	445	740	1,261	3,215	8,555	9,805	1
1999.....	23,712	102	27	37	60	129	384	737	1,101	3,221	8,372	9,541	1
White, female													
2007 1/.....	25,022	44	36	24	46	92	251	588	983	2,060	6,517	14,381	-
2006 1/.....	26,635	54	37	11	56	109	274	607	1,013	2,216	7,240	15,017	1
2005 1/.....	30,014	57	33	35	38	113	287	648	1,128	2,406	8,046	17,223	-
2004 1/.....	28,437	54	36	21	40	83	279	539	1,029	2,307	7,878	16,171	-
2003 1/.....	31,623	73	31	38	65	91	299	608	1,065	2,405	8,588	18,359	1
2002.....	32,549	55	27	35	46	109	300	595	990	2,520	8,961	18,910	1
2001.....	30,889	67	34	29	46	101	294	519	891	2,487	8,364	18,057	-
2000.....	31,867	72	34	26	52	115	304	517	910	2,574	8,787	18,476	-
1999.....	31,408	75	40	25	51	85	320	464	893	2,463	8,604	18,387	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumonia (J12-J18)													
Black, both sexes													
2007 1/.....	5,129	71	24	19	23	92	176	437	569	798	1,310	1,610	-
2006 1/.....	5,270	85	30	8	38	75	185	454	509	803	1,415	1,668	-
2005 1/.....	5,711	93	26	17	30	77	211	492	604	913	1,500	1,748	-
2004 1/.....	5,650	89	32	19	39	65	228	444	548	856	1,548	1,782	-
2003 1/.....	5,775	86	22	19	35	99	251	485	562	917	1,517	1,781	1
2002.....	5,841	86	38	17	36	83	252	403	556	842	1,599	1,929	-
2001.....	5,751	106	26	15	44	87	275	436	477	804	1,593	1,888	-
2000.....	5,909	101	29	10	43	92	254	418	532	981	1,556	1,891	2
1999.....	5,810	111	42	15	48	106	307	428	493	876	1,411	1,973	-
Black, male													
2007 1/.....	2,485	39	12	12	12	51	102	271	330	455	641	560	-
2006 1/.....	2,465	53	13	2	21	36	98	275	305	437	663	562	-
2005 1/.....	2,697	47	15	10	22	45	126	297	372	510	689	564	-
2004 1/.....	2,750	51	10	12	19	43	130	266	321	519	778	601	-
2003 1/.....	2,869	51	11	11	21	58	142	286	353	552	771	612	1
2002.....	2,753	47	20	10	18	45	138	241	338	459	796	641	-
2001.....	2,803	53	14	9	22	51	156	275	295	477	823	628	-
2000.....	2,880	56	21	6	20	58	149	267	324	566	768	644	1
1999.....	2,801	55	24	8	23	58	176	280	288	517	689	683	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumonia (J12-J18)													
Black, female													
2007 1/.....	2,644	32	12	7	11	41	74	166	239	343	669	1,050	-
2006 1/.....	2,805	32	17	6	17	39	87	179	204	366	752	1,106	-
2005 1/.....	3,014	46	11	7	8	32	85	195	232	403	811	1,184	-
2004 1/.....	2,900	38	22	7	20	22	98	178	227	337	770	1,181	-
2003 1/.....	2,906	35	11	8	14	41	109	199	209	365	746	1,169	-
2002.....	3,088	39	18	7	18	38	114	162	218	383	803	1,288	-
2001.....	2,948	53	12	6	22	36	119	161	182	327	770	1,260	-
2000.....	3,029	45	8	4	23	34	105	151	208	415	788	1,247	1
1999.....	3,009	56	18	7	25	48	131	148	205	359	722	1,290	-
American Indian 2/, both sexes													
2007 1/.....	275	8	1	-	4	4	18	28	34	38	66	74	-
2006 1/.....	264	3	4	1	3	7	15	24	26	40	67	74	-
2005 1/.....	343	5	1	1	1	5	18	26	43	47	96	100	-
2004 1/.....	288	12	4	1	-	7	13	18	32	38	90	73	-
2003 1/.....	373	11	2	2	2	6	20	35	49	55	79	112	-
2002.....	288	6	1	-	3	7	13	23	24	33	80	98	-
2001.....	315	17	2	2	3	9	19	20	25	42	85	91	-
2000.....	285	6	-	2	2	5	28	21	24	45	70	82	-
1999.....	310	8	3	2	5	4	20	26	30	36	70	106	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumonia (J12-J18)													
American Indian 2/, male													
2007 1/.....	148	5	1	-	1	3	12	23	27	22	32	22	-
2006 1/.....	136	-	3	-	1	2	9	14	11	27	36	33	-
2005 1/.....	174	2	-	1	-	4	12	15	24	23	50	43	-
2004 1/.....	142	7	1	-	-	5	7	12	13	18	46	33	-
2003 1/.....	177	6	1	2	1	1	13	21	27	30	39	36	-
2002.....	130	3	1	-	1	4	9	14	12	18	32	36	-
2001.....	160	11	-	1	1	6	13	12	13	23	42	38	-
2000.....	148	2	-	-	1	3	18	16	16	29	33	30	-
1999.....	145	3	3	2	4	1	12	17	13	16	38	36	-
American Indian 2/, female													
2007 1/.....	127	3	-	-	3	1	6	5	7	16	34	52	-
2006 1/.....	128	3	1	1	2	5	6	10	15	13	31	41	-
2005 1/.....	169	3	1	-	1	1	6	11	19	24	46	57	-
2004 1/.....	146	5	3	1	-	2	6	6	19	20	44	40	-
2003 1/.....	196	5	1	-	1	5	7	14	22	25	40	76	-
2002.....	158	3	-	-	2	3	4	9	12	15	48	62	-
2001.....	155	6	2	1	2	3	6	8	12	19	43	53	-
2000.....	137	4	-	2	1	2	10	5	8	16	37	52	-
1999.....	165	5	-	-	1	3	8	9	17	20	32	70	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumonia (J12-J18)													
Asian or Pacific Islander 3/, both sexes													
2007 1/.....	1,329	7	5	4	7	8	8	38	61	156	404	631	-
2006 1/.....	1,330	9	-	1	1	13	11	34	60	151	425	625	-
2005 1/.....	1,305	4	3	4	5	9	16	32	68	164	397	603	-
2004 1/.....	1,229	5	2	3	3	2	13	31	59	151	394	566	-
2003 1/.....	1,231	8	7	4	4	7	8	29	58	143	406	557	-
2002.....	1,165	7	5	3	3	7	9	35	45	133	402	516	-
2001.....	1,166	8	5	3	6	8	9	29	57	158	374	508	1
2000.....	1,102	8	4	3	5	4	15	30	59	175	361	438	-
1999.....	825	11	6	3	4	6	16	16	37	101	284	341	-
Asian or Pacific Islander 3/, male													
2007 1/.....	700	3	5	2	4	6	5	22	36	91	219	307	-
2006 1/.....	709	5	-	-	1	8	5	22	38	95	227	308	-
2005 1/.....	708	3	3	3	3	5	8	21	42	98	230	292	-
2004 1/.....	588	3	1	2	1	2	8	16	33	79	199	244	-
2003 1/.....	664	4	4	-	2	4	5	17	38	83	235	272	-
2002.....	635	5	5	2	2	4	6	26	26	78	221	260	-
2001.....	622	3	2	2	2	4	5	21	37	86	204	256	-
2000.....	581	7	3	2	1	3	4	18	35	97	199	212	-
1999.....	458	5	3	1	1	4	11	10	23	65	157	178	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Pneumonia (J12-J18)

Asian or Pacific Islander 3/, female

2007 1/.....	629	4	-	2	3	2	3	16	25	65	185	324	-
2006 1/.....	621	4	-	1	-	5	6	12	22	56	198	317	-
2005 1/.....	597	1	-	1	2	4	8	11	26	66	167	311	-
2004 1/.....	641	2	1	1	2	-	5	15	26	72	195	322	-
2003 1/.....	567	4	3	4	2	3	3	12	20	60	171	285	-
2002.....	530	2	-	1	1	3	3	9	19	55	181	256	-
2001.....	544	5	3	1	4	4	4	8	20	72	170	252	1
2000.....	521	1	1	1	4	1	11	12	24	78	162	226	-
1999.....	367	6	3	2	3	2	5	6	14	36	127	163	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other acute lower respiratory infections (J20-J22,U04)

All races, both sexes

2007.....	255	46	17	3	2	5	5	14	19	27	36	81	-
2006.....	297	52	14	3	4	-	8	10	13	24	61	108	-
2005.....	404	51	16	4	4	6	15	15	19	26	78	170	-
2004.....	319	60	16	1	4	3	9	10	14	21	54	127	-
2003.....	421	54	28	6	6	2	11	19	21	30	80	164	-
2002.....	386	52	23	8	3	5	8	18	18	33	66	152	-
2001.....	347	50	18	3	1	2	12	14	16	28	58	145	-
2000.....	425	43	11	7	5	8	12	12	10	43	82	192	-
1999.....	540	63	25	5	5	4	11	15	21	35	107	249	-

All races, male

2007.....	121	32	8	2	2	3	3	8	12	17	17	17	-
2006.....	119	30	6	-	4	-	4	7	6	8	23	31	-
2005.....	172	30	10	4	3	2	6	12	5	10	41	49	-
2004.....	137	34	9	1	3	2	6	8	6	12	18	38	-
2003.....	170	35	17	2	2	1	6	8	12	17	35	35	-
2002.....	154	39	9	7	-	1	3	10	6	13	27	39	-
2001.....	117	26	8	2	1	-	7	7	6	10	20	30	-
2000.....	160	25	4	3	4	6	7	4	3	22	35	47	-
1999.....	196	36	15	2	1	3	5	8	10	17	51	48	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other acute lower respiratory infections (J20-J22,U04)

All races, female

2007.....	134	14	9	1	-	2	2	6	7	10	19	64	-
2006.....	178	22	8	3	-	-	4	3	7	16	38	77	-
2005.....	232	21	6	-	1	4	9	3	14	16	37	121	-
2004.....	182	26	7	-	1	1	3	2	8	9	36	89	-
2003.....	251	19	11	4	4	1	5	11	9	13	45	129	-
2002.....	232	13	14	1	3	4	5	8	12	20	39	113	-
2001.....	230	24	10	1	-	2	5	7	10	18	38	115	-
2000.....	265	18	7	4	1	2	5	8	7	21	47	145	-
1999.....	344	27	10	3	4	1	6	7	11	18	56	201	-

White, both sexes

2007 1/.....	217	28	10	3	2	5	4	12	17	26	35	75	-
2006 1/.....	251	35	8	2	3	-	6	6	8	22	57	104	-
2005 1/.....	351	30	13	3	4	5	12	11	18	22	72	161	-
2004 1/.....	279	43	9	1	3	3	9	8	13	19	49	122	-
2003 1/.....	375	42	17	5	3	1	11	17	18	26	76	159	-
2002.....	334	34	20	8	2	4	7	14	14	28	59	144	-
2001.....	299	31	13	3	1	1	12	10	14	22	54	138	-
2000.....	372	25	8	3	5	5	8	9	10	41	77	181	-
1999.....	462	36	15	4	4	3	9	11	17	27	99	237	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other acute lower respiratory infections (J20-J22,U04)

White, male

2007 1/.....	102	23	4	2	2	3	3	7	10	16	16	16	-
2006 1/.....	93	21	3	-	3	-	3	4	3	7	20	29	-
2005 1/.....	147	18	9	3	3	2	4	8	5	8	40	47	-
2004 1/.....	115	24	4	1	2	2	6	6	6	11	17	36	-
2003 1/.....	147	27	10	2	-	1	6	8	10	15	33	35	-
2002.....	130	26	7	7	-	-	3	9	4	12	25	37	-
2001.....	94	16	5	2	1	-	7	4	5	8	18	28	-
2000.....	134	12	4	2	4	4	6	4	3	20	33	42	-
1999.....	156	19	8	2	1	2	4	8	9	12	48	43	-

White, female

2007 1/.....	115	5	6	1	-	2	1	5	7	10	19	59	-
2006 1/.....	158	14	5	2	-	-	3	2	5	15	37	75	-
2005 1/.....	204	12	4	-	1	3	8	3	13	14	32	114	-
2004 1/.....	164	19	5	-	1	1	3	2	7	8	32	86	-
2003 1/.....	228	15	7	3	3	-	5	9	8	11	43	124	-
2002.....	204	8	13	1	2	4	4	5	10	16	34	107	-
2001.....	205	15	8	1	-	1	5	6	9	14	36	110	-
2000.....	238	13	4	1	1	1	2	5	7	21	44	139	-
1999.....	306	17	7	2	3	1	5	3	8	15	51	194	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other acute lower respiratory infections (J20-J22,U04)

Black, both sexes

2007 1/.....	33	16	5	-	-	-	1	2	2	1	1	5	-
2006 1/.....	37	14	5	1	1	-	2	3	5	2	2	2	-
2005 1/.....	41	18	2	1	-	1	3	2	1	3	4	6	-
2004 1/.....	31	13	6	-	1	-	-	1	-	2	5	3	-
2003 1/.....	40	12	7	1	3	1	-	2	2	4	3	5	-
2002.....	45	16	2	-	1	1	1	4	3	5	7	5	-
2001.....	34	15	2	-	-	1	-	4	1	4	-	7	-
2000.....	46	14	3	3	-	2	4	3	-	2	5	10	-
1999.....	64	24	6	-	1	1	2	4	2	8	6	10	-

Black, male

2007 1/.....	16	7	3	-	-	-	-	1	2	1	1	1	-
2006 1/.....	22	7	3	-	1	-	1	2	3	1	2	2	-
2005 1/.....	17	9	-	1	-	-	2	2	-	1	-	2	-
2004 1/.....	15	6	4	-	1	-	-	1	-	1	1	1	-
2003 1/.....	21	8	6	-	2	-	-	-	1	2	2	-	-
2002.....	23	13	1	-	-	1	-	1	2	1	2	2	-
2001.....	16	9	1	-	-	-	-	3	-	1	-	2	-
2000.....	22	10	-	1	-	2	1	-	-	2	2	4	-
1999.....	30	15	3	-	-	1	1	-	1	5	1	3	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other acute lower respiratory infections (J20-J22,U04)

Black, female

2007 1/.....	17	9	2	-	-	-	1	1	-	-	-	4	-
2006 1/.....	15	7	2	1	-	-	1	1	2	1	-	-	-
2005 1/.....	24	9	2	-	-	1	1	-	1	2	4	4	-
2004 1/.....	16	7	2	-	-	-	-	-	-	1	4	2	-
2003 1/.....	19	4	1	1	1	1	-	2	1	2	1	5	-
2002.....	22	3	1	-	1	-	1	3	1	4	5	3	-
2001.....	18	6	1	-	-	1	-	1	1	3	-	5	-
2000.....	24	4	3	2	-	-	3	3	-	-	3	6	-
1999.....	34	9	3	-	1	-	1	4	1	3	5	7	-

American Indian 2/, both sexes

2007 1/.....	2	1	1	-	-	-	-	-	-	-	-	-	-
2006 1/.....	2	-	-	-	-	-	-	1	-	-	1	-	-
2005 1/.....	2	-	-	-	-	-	-	1	-	-	-	1	-
2004 1/.....	2	1	-	-	-	-	-	1	-	-	-	-	-
2003 1/.....	2	-	1	-	-	-	-	-	-	-	1	-	-
2002.....	3	1	1	-	-	-	-	-	1	-	-	-	-
2001.....	7	2	3	-	-	-	-	-	1	-	1	-	-
2000.....	2	1	-	-	-	-	-	-	-	-	-	1	-
1999.....	5	1	2	1	-	-	-	-	1	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other acute lower respiratory infections (J20-J22,U04)

American Indian 2/, male

2007 1/.....	1	1	-	-	-	-	-	-	-	-	-	-	-
2006 1/.....	2	-	-	-	-	-	-	1	-	-	1	-	-
2005 1/.....	1	-	-	-	-	-	-	1	-	-	-	-	-
2004 1/.....	2	1	-	-	-	-	-	1	-	-	-	-	-
2003 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	1	-	-	-	-	-	-	-	-	-	-
2001.....	4	1	2	-	-	-	-	-	1	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	1	-
1999.....	3	1	2	-	-	-	-	-	-	-	-	-	-

American Indian 2/, female

2007 1/.....	1	-	1	-	-	-	-	-	-	-	-	-	-
2006 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2005 1/.....	1	-	-	-	-	-	-	-	-	-	-	1	-
2004 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 1/.....	2	-	1	-	-	-	-	-	-	-	1	-	-
2002.....	2	1	-	-	-	-	-	-	1	-	-	-	-
2001.....	3	1	1	-	-	-	-	-	-	-	1	-	-
2000.....	1	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	-	-	1	-	-	-	-	1	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other acute lower respiratory infections (J20-J22,U04)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	3	1	1	-	-	-	-	-	-	-	-	1	-
2006 1/.....	7	3	1	-	-	-	-	-	-	-	1	2	-
2005 1/.....	10	3	1	-	-	-	-	1	-	1	2	2	-
2004 1/.....	7	3	1	-	-	-	-	-	1	-	-	2	-
2003 1/.....	4	-	3	-	-	-	-	-	1	-	-	-	-
2002.....	4	1	-	-	-	-	-	-	-	-	-	3	-
2001.....	7	2	-	-	-	-	-	-	-	2	3	-	-
2000.....	5	3	-	1	-	1	-	-	-	-	-	-	-
1999.....	9	2	2	-	-	-	-	-	1	-	2	2	-

Asian or Pacific Islander 3/, male

2007 1/.....	2	1	1	-	-	-	-	-	-	-	-	-	-
2006 1/.....	2	2	-	-	-	-	-	-	-	-	-	-	-
2005 1/.....	7	3	1	-	-	-	-	1	-	1	1	-	-
2004 1/.....	5	3	1	-	-	-	-	-	-	-	-	1	-
2003 1/.....	2	-	1	-	-	-	-	-	1	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	3	-	-	-	-	-	-	-	-	1	2	-	-
2000.....	3	3	-	-	-	-	-	-	-	-	-	-	-
1999.....	7	1	2	-	-	-	-	-	-	-	2	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other acute lower respiratory infections (J20-J22,U04)

Asian or Pacific Islander 3/, female

2007 1/.....	1	-	-	-	-	-	-	-	-	-	-	-	1	-
2006 1/.....	5	1	1	-	-	-	-	-	-	-	-	1	2	-
2005 1/.....	3	-	-	-	-	-	-	-	-	-	-	1	2	-
2004 1/.....	2	-	-	-	-	-	-	-	1	-	-	-	1	-
2003 1/.....	2	-	2	-	-	-	-	-	-	-	-	-	-	-
2002.....	4	1	-	-	-	-	-	-	-	-	-	-	3	-
2001.....	4	2	-	-	-	-	-	-	-	1	1	-	-	-
2000.....	2	-	-	1	-	1	-	-	-	-	-	-	-	-
1999.....	2	1	-	-	-	-	-	-	1	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)													
All races, both sexes													
2007.....	213	45	16	3	1	5	5	11	18	22	28	59	-
2006.....	214	52	14	3	4	-	7	9	9	18	40	58	-
2005.....	283	50	14	4	3	3	13	15	13	17	53	98	-
2004.....	226	58	16	1	4	3	8	7	10	18	32	69	-
2003.....	304	53	27	6	6	2	9	16	17	22	45	101	-
2002.....	279	49	23	8	2	5	6	17	13	23	39	94	-
2001.....	263	49	18	3	1	2	11	12	14	23	39	91	-
2000.....	290	42	11	6	4	6	10	9	7	31	53	111	-
1999.....	388	62	24	4	5	3	10	14	18	29	69	150	-
All races, male													
2007.....	104	32	8	2	1	3	3	5	11	13	14	12	-
2006.....	96	30	6	-	4	-	4	6	5	7	14	20	-
2005.....	123	29	8	4	2	-	5	12	3	7	26	27	-
2004.....	108	33	9	1	3	2	5	5	6	10	7	27	-
2003.....	126	34	16	2	2	1	5	7	11	12	17	19	-
2002.....	122	37	9	7	-	1	3	9	5	10	18	23	-
2001.....	89	25	8	2	1	-	6	6	6	8	10	17	-
2000.....	119	24	4	2	3	4	6	3	3	14	27	29	-
1999.....	154	36	14	2	1	2	4	7	8	15	34	31	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Acute bronchitis and bronchiolitis (J20-J21)

All races, female

2007.....	109	13	8	1	-	2	2	6	7	9	14	47	-
2006.....	118	22	8	3	-	-	3	3	4	11	26	38	-
2005.....	160	21	6	-	1	3	8	3	10	10	27	71	-
2004.....	118	25	7	-	1	1	3	2	4	8	25	42	-
2003.....	178	19	11	4	4	1	4	9	6	10	28	82	-
2002.....	157	12	14	1	2	4	3	8	8	13	21	71	-
2001.....	174	24	10	1	-	2	5	6	8	15	29	74	-
2000.....	171	18	7	4	1	2	4	6	4	17	26	82	-
1999.....	234	26	10	2	4	1	6	7	10	14	35	119	-

White, both sexes

2007 1/.....	180	28	9	3	1	5	4	9	16	21	27	57	-
2006 1/.....	173	35	8	2	3	-	6	6	5	16	37	55	-
2005 1/.....	240	29	12	3	3	2	11	11	13	14	50	92	-
2004 1/.....	190	42	9	1	3	3	8	6	9	16	28	65	-
2003 1/.....	263	41	16	5	3	1	9	15	14	18	43	98	-
2002.....	237	31	20	8	1	4	6	13	10	21	36	87	-
2001.....	218	30	13	3	1	1	11	8	12	18	36	85	-
2000.....	244	25	8	2	4	3	6	6	7	29	49	105	-
1999.....	320	35	15	3	4	3	8	10	14	21	63	144	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)													
White, male													
2007 1/.....	86	23	4	2	1	3	3	4	9	12	13	12	-
2006 1/.....	73	21	3	-	3	-	3	4	2	6	12	19	-
2005 1/.....	101	17	8	3	2	-	4	8	3	6	25	25	-
2004 1/.....	87	23	4	1	2	2	5	4	6	9	6	25	-
2003 1/.....	104	26	9	2	-	1	5	7	9	10	16	19	-
2002.....	100	24	7	7	-	-	3	8	4	9	17	21	-
2001.....	67	15	5	2	1	-	6	3	5	6	8	16	-
2000.....	98	12	4	1	3	2	5	3	3	12	26	27	-
1999.....	119	19	8	2	1	2	3	7	7	10	32	28	-
White, female													
2007 1/.....	94	5	5	1	-	2	1	5	7	9	14	45	-
2006 1/.....	100	14	5	2	-	-	3	2	3	10	25	36	-
2005 1/.....	139	12	4	-	1	2	7	3	10	8	25	67	-
2004 1/.....	103	19	5	-	1	1	3	2	3	7	22	40	-
2003 1/.....	159	15	7	3	3	-	4	8	5	8	27	79	-
2002.....	137	7	13	1	1	4	3	5	6	12	19	66	-
2001.....	151	15	8	1	-	1	5	5	7	12	28	69	-
2000.....	146	13	4	1	1	1	3	4	4	17	23	78	-
1999.....	201	16	7	1	3	1	5	3	7	11	31	116	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Acute bronchitis and bronchiolitis (J20-J21)

Black, both sexes

2007 1/.....	29	15	5	-	-	-	1	2	2	1	1	2	-
2006 1/.....	34	14	5	1	1	-	1	3	4	2	2	1	-
2005 1/.....	32	18	2	1	-	1	2	2	-	2	1	3	-
2004 1/.....	28	12	6	-	1	-	-	1	-	2	4	2	-
2003 1/.....	35	12	7	1	3	1	-	1	2	4	1	3	-
2002.....	36	16	2	-	1	1	-	4	2	2	3	5	-
2001.....	32	15	2	-	-	1	-	4	1	3	-	6	-
2000.....	40	13	3	3	-	2	4	3	-	2	4	6	-
1999.....	55	24	5	-	1	-	2	4	2	8	4	5	-

Black, male

2007 1/.....	15	7	3	-	-	-	-	1	2	1	1	-	-
2006 1/.....	21	7	3	-	1	-	1	2	3	1	2	1	-
2005 1/.....	15	9	-	1	-	-	1	2	-	-	-	2	-
2004 1/.....	15	6	4	-	1	-	-	1	-	1	1	1	-
2003 1/.....	20	8	6	-	2	-	-	-	1	2	1	-	-
2002.....	21	13	1	-	-	1	-	1	1	1	1	2	-
2001.....	15	9	1	-	-	-	-	3	-	1	-	1	-
2000.....	18	9	-	1	-	2	1	-	-	2	1	2	-
1999.....	26	15	2	-	-	-	1	-	1	5	-	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Acute bronchitis and bronchiolitis (J20-J21)

Black, female

2007 1/.....	14	8	2	-	-	-	1	1	-	-	-	2	-
2006 1/.....	13	7	2	1	-	-	-	1	1	1	-	-	-
2005 1/.....	17	9	2	-	-	1	1	-	-	2	1	1	-
2004 1/.....	13	6	2	-	-	-	-	-	-	1	3	1	-
2003 1/.....	15	4	1	1	1	1	-	1	1	2	-	3	-
2002.....	15	3	1	-	1	-	-	3	1	1	2	3	-
2001.....	17	6	1	-	-	1	-	1	1	2	-	5	-
2000.....	22	4	3	2	-	-	3	3	-	-	3	4	-
1999.....	29	9	3	-	1	-	1	4	1	3	4	3	-

American Indian 2/, both sexes

2007 1/.....	2	1	1	-	-	-	-	-	-	-	-	-	-
2006 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2005 1/.....	2	-	-	-	-	-	-	1	-	-	-	1	-
2004 1/.....	1	1	-	-	-	-	-	-	-	-	-	-	-
2003 1/.....	2	-	1	-	-	-	-	-	-	-	1	-	-
2002.....	3	1	1	-	-	-	-	-	1	-	-	-	-
2001.....	6	2	3	-	-	-	-	-	1	-	-	-	-
2000.....	1	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	5	1	2	1	-	-	-	-	1	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Acute bronchitis and bronchiolitis (J20-J21)

American Indian 2/, male

2007 1/.....	1	1	-	-	-	-	-	-	-	-	-	-	-
2006 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2005 1/.....	1	-	-	-	-	-	1	-	-	-	-	-	-
2004 1/.....	1	1	-	-	-	-	-	-	-	-	-	-	-
2003 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	1	-	-	-	-	-	-	-	-	-	-
2001.....	4	1	2	-	-	-	-	1	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	3	1	2	-	-	-	-	-	-	-	-	-	-

American Indian 2/, female

2007 1/.....	1	-	1	-	-	-	-	-	-	-	-	-	-
2006 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2005 1/.....	1	-	-	-	-	-	-	-	-	-	-	1	-
2004 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 1/.....	2	-	1	-	-	-	-	-	-	1	-	-	-
2002.....	2	1	-	-	-	-	-	1	-	-	-	-	-
2001.....	2	1	1	-	-	-	-	-	-	-	-	-	-
2000.....	1	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	-	-	1	-	-	-	1	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Acute bronchitis and bronchiolitis (J20-J21)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	2	1	1	-	-	-	-	-	-	-	-	-	-
2006 1/.....	7	3	1	-	-	-	-	-	-	-	1	2	-
2005 1/.....	9	3	-	-	-	-	1	-	1	1	2	2	-
2004 1/.....	7	3	1	-	-	-	-	-	1	-	-	2	-
2003 1/.....	4	-	3	-	-	-	-	-	1	-	-	-	-
2002.....	3	1	-	-	-	-	-	-	-	-	-	2	-
2001.....	7	2	-	-	-	-	-	-	-	2	3	-	-
2000.....	5	3	-	1	-	1	-	-	-	-	-	-	-
1999.....	8	2	2	-	-	-	-	-	1	-	2	1	-

Asian or Pacific Islander 3/, male

2007 1/.....	2	1	1	-	-	-	-	-	-	-	-	-	-
2006 1/.....	2	2	-	-	-	-	-	-	-	-	-	-	-
2005 1/.....	6	3	-	-	-	-	1	-	1	1	-	-	-
2004 1/.....	5	3	1	-	-	-	-	-	-	-	-	1	-
2003 1/.....	2	-	1	-	-	-	-	-	1	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	3	-	-	-	-	-	-	-	-	1	2	-	-
2000.....	3	3	-	-	-	-	-	-	-	-	-	-	-
1999.....	6	1	2	-	-	-	-	-	-	-	2	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Acute bronchitis and bronchiolitis (J20-J21)

Asian or Pacific Islander 3/, female

2007 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2006 1/.....	5	1	1	-	-	-	-	-	-	-	1	2	-
2005 1/.....	3	-	-	-	-	-	-	-	-	-	1	2	-
2004 1/.....	2	-	-	-	-	-	-	-	1	-	-	1	-
2003 1/.....	2	-	2	-	-	-	-	-	-	-	-	-	-
2002.....	3	1	-	-	-	-	-	-	-	-	-	2	-
2001.....	4	2	-	-	-	-	-	-	-	1	1	-	-
2000.....	2	-	-	1	-	1	-	-	-	-	-	-	-
1999.....	2	1	-	-	-	-	-	-	1	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other and unspecified acute lower respiratory infections (J22,U04)

All races, both sexes

2007.....	42	1	1	-	1	-	-	3	1	5	8	22	-
2006.....	83	-	-	-	-	-	1	1	4	6	21	50	-
2005.....	121	1	2	-	1	3	2	-	6	9	25	72	-
2004.....	93	2	-	-	-	-	1	3	4	3	22	58	-
2003.....	117	1	1	-	-	-	2	3	4	8	35	63	-
2002.....	107	3	-	-	1	-	2	1	5	10	27	58	-
2001.....	84	1	-	-	-	-	1	2	2	5	19	54	-
2000.....	135	1	-	1	1	2	2	3	3	12	29	81	-
1999.....	152	1	1	1	-	1	1	1	3	6	38	99	-

All races, male

2007.....	17	-	-	-	1	-	-	3	1	4	3	5	-
2006.....	23	-	-	-	-	-	-	1	1	1	9	11	-
2005.....	49	1	2	-	1	2	1	-	2	3	15	22	-
2004.....	29	1	-	-	-	-	1	3	-	2	11	11	-
2003.....	44	1	1	-	-	-	1	1	1	5	18	16	-
2002.....	32	2	-	-	-	-	-	1	1	3	9	16	-
2001.....	28	1	-	-	-	-	1	1	-	2	10	13	-
2000.....	41	1	-	1	1	2	1	1	-	8	8	18	-
1999.....	42	-	1	-	-	1	1	1	2	2	17	17	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other and unspecified acute lower respiratory infections (J22,U04)

All races, female

2007.....	25	1	1	-	-	-	-	-	-	1	5	17	-
2006.....	60	-	-	-	-	-	1	-	3	5	12	39	-
2005.....	72	-	-	-	-	1	1	-	4	6	10	50	-
2004.....	64	1	-	-	-	-	-	-	4	1	11	47	-
2003.....	73	-	-	-	-	-	1	2	3	3	17	47	-
2002.....	75	1	-	-	1	-	2	-	4	7	18	42	-
2001.....	56	-	-	-	-	-	-	1	2	3	9	41	-
2000.....	94	-	-	-	-	-	1	2	3	4	21	63	-
1999.....	110	1	-	1	-	-	-	-	1	4	21	82	-

White, both sexes

2007 1/.....	37	-	1	-	1	-	-	3	1	5	8	18	-
2006 1/.....	78	-	-	-	-	-	-	-	3	6	20	49	-
2005 1/.....	111	1	1	-	1	3	1	-	5	8	22	69	-
2004 1/.....	89	1	-	-	-	-	1	2	4	3	21	57	-
2003 1/.....	112	1	1	-	-	-	2	2	4	8	33	61	-
2002.....	97	3	-	-	1	-	1	1	4	7	23	57	-
2001.....	81	1	-	-	-	-	1	2	2	4	18	53	-
2000.....	128	-	-	1	1	2	2	3	3	12	28	76	-
1999.....	142	1	-	1	-	-	1	1	3	6	36	93	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other and unspecified acute lower respiratory infections (J22,U04)

White, male

2007 1/.....	16	-	-	-	1	-	-	3	1	4	3	4	-
2006 1/.....	20	-	-	-	-	-	-	-	1	1	8	10	-
2005 1/.....	46	1	1	-	1	2	-	-	2	2	15	22	-
2004 1/.....	28	1	-	-	-	-	1	2	-	2	11	11	-
2003 1/.....	43	1	1	-	-	-	1	1	1	5	17	16	-
2002.....	30	2	-	-	-	-	-	1	-	3	8	16	-
2001.....	27	1	-	-	-	-	1	1	-	2	10	12	-
2000.....	36	-	-	1	1	2	1	1	-	8	7	15	-
1999.....	37	-	-	-	-	-	1	1	2	2	16	15	-

White, female

2007 1/.....	21	-	1	-	-	-	-	-	-	1	5	14	-
2006 1/.....	58	-	-	-	-	-	-	-	2	5	12	39	-
2005 1/.....	65	-	-	-	-	1	1	-	3	6	7	47	-
2004 1/.....	61	-	-	-	-	-	-	-	4	1	10	46	-
2003 1/.....	69	-	-	-	-	-	1	1	3	3	16	45	-
2002.....	67	1	-	-	1	-	1	-	4	4	15	41	-
2001.....	54	-	-	-	-	-	-	1	2	2	8	41	-
2000.....	92	-	-	-	-	-	1	2	3	4	21	61	-
1999.....	105	1	-	1	-	-	-	-	1	4	20	78	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other and unspecified acute lower respiratory infections (J22,U04)

Black, both sexes

2007 1/.....	4	1	-	-	-	-	-	-	-	-	-	3	-
2006 1/.....	3	-	-	-	-	-	1	-	1	-	-	1	-
2005 1/.....	9	-	-	-	-	-	1	-	1	1	3	3	-
2004 1/.....	3	1	-	-	-	-	-	-	-	-	1	1	-
2003 1/.....	5	-	-	-	-	-	-	1	-	-	2	2	-
2002.....	9	-	-	-	-	-	1	-	1	3	4	-	-
2001.....	2	-	-	-	-	-	-	-	-	1	-	1	-
2000.....	6	1	-	-	-	-	-	-	-	-	1	4	-
1999.....	9	-	1	-	-	1	-	-	-	-	2	5	-

Black, male

2007 1/.....	1	-	-	-	-	-	-	-	-	-	-	1	-
2006 1/.....	1	-	-	-	-	-	-	-	-	-	-	1	-
2005 1/.....	2	-	-	-	-	-	1	-	-	1	-	-	-
2004 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 1/.....	1	-	-	-	-	-	-	-	-	-	1	-	-
2002.....	2	-	-	-	-	-	-	-	1	-	1	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	1	-
2000.....	4	1	-	-	-	-	-	-	-	-	1	2	-
1999.....	4	-	1	-	-	1	-	-	-	-	1	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Chronic lower respiratory diseases (J40-J47)

All races, both sexes

2007.....	127,924	43	57	118	149	263	796	4,153	12,777	28,664	48,041	32,857	6
2006.....	124,583	30	43	115	151	255	841	3,924	12,375	28,236	47,406	31,203	4
2005.....	130,933	33	56	104	148	258	890	3,977	12,747	29,910	50,333	32,473	4
2004.....	121,987	35	48	120	179	255	887	3,511	11,754	28,390	47,568	29,239	1
2003.....	126,382	31	55	118	191	282	950	3,537	12,077	29,919	49,286	29,934	2
2002.....	124,816	39	65	136	192	301	1,008	3,475	11,280	29,788	49,241	29,284	7
2001.....	123,013	41	43	104	171	290	969	3,324	11,166	30,751	47,762	28,391	1
2000.....	122,009	36	51	139	190	296	930	3,251	10,739	31,157	47,722	27,496	2
1999.....	124,181	36	54	139	209	330	890	3,110	11,297	32,644	48,635	26,833	4

All races, male

2007.....	61,235	27	30	78	96	150	392	2,125	6,733	14,679	23,439	13,480	6
2006.....	59,260	20	29	78	93	137	396	1,958	6,382	14,152	23,167	12,845	3
2005.....	62,435	19	33	63	95	130	426	2,034	6,589	15,414	24,479	13,150	3
2004.....	58,646	24	30	75	110	151	431	1,806	6,029	14,503	23,255	12,231	1
2003.....	60,714	23	38	75	114	156	429	1,826	6,243	15,333	23,963	12,512	2
2002.....	60,713	23	44	74	120	165	497	1,795	5,783	15,339	24,432	12,436	5
2001.....	59,697	22	29	61	99	146	455	1,702	5,702	15,877	23,691	11,912	1
2000.....	60,004	23	28	80	120	165	437	1,661	5,564	16,210	23,838	11,876	2
1999.....	62,415	19	30	93	123	152	410	1,551	5,958	17,477	24,705	11,894	3

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Chronic lower respiratory diseases (J40-J47)

All races, female

2007.....	66,689	16	27	40	53	113	404	2,028	6,044	13,985	24,602	19,377	-
2006.....	65,323	10	14	37	58	118	445	1,966	5,993	14,084	24,239	18,358	1
2005.....	68,498	14	23	41	53	128	464	1,943	6,158	14,496	25,854	19,323	1
2004.....	63,341	11	18	45	69	104	456	1,705	5,725	13,887	24,313	17,008	-
2003.....	65,668	8	17	43	77	126	521	1,711	5,834	14,586	25,323	17,422	-
2002.....	64,103	16	21	62	72	136	511	1,680	5,497	14,449	24,809	16,848	2
2001.....	63,316	19	14	43	72	144	514	1,622	5,464	14,874	24,071	16,479	-
2000.....	62,005	13	23	59	70	131	493	1,590	5,175	14,947	23,884	15,620	-
1999.....	61,766	17	24	46	86	178	480	1,559	5,339	15,167	23,930	14,939	1

White, both sexes

2007 1/.....	118,081	26	22	54	79	159	578	3,395	11,412	26,389	44,986	30,976	5
2006 1/.....	114,993	15	26	42	69	141	591	3,236	11,000	26,055	44,390	29,425	3
2005 1/.....	120,884	20	24	45	71	152	647	3,208	11,372	27,517	47,232	30,593	3
2004 1/.....	112,914	23	22	53	91	150	644	2,830	10,483	26,273	44,742	27,602	1
2003 1/.....	116,917	23	24	54	101	169	675	2,873	10,770	27,606	46,331	28,289	2
2002.....	115,395	25	30	55	109	189	696	2,812	10,053	27,501	46,258	27,662	5
2001.....	113,819	22	20	48	85	156	665	2,703	9,967	28,385	44,996	26,772	-
2000.....	112,840	23	28	63	103	166	624	2,584	9,532	28,765	45,025	25,926	1
1999.....	114,735	16	31	71	125	176	608	2,498	10,056	30,111	45,698	25,342	3

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Chronic lower respiratory diseases (J40-J47)

White, male

2007 1/.....	55,934	13	13	34	49	86	278	1,777	5,956	13,399	21,735	12,589	5
2006 1/.....	54,043	10	17	32	40	70	282	1,604	5,595	12,916	21,486	11,989	2
2005 1/.....	56,911	13	11	26	46	72	311	1,641	5,821	14,050	22,693	12,225	2
2004 1/.....	53,622	13	15	30	55	82	312	1,462	5,284	13,285	21,681	11,402	1
2003 1/.....	55,397	20	14	31	59	79	305	1,495	5,511	14,002	22,192	11,687	2
2002.....	55,409	16	21	31	60	101	336	1,453	5,080	14,015	22,681	11,611	4
2001.....	54,561	13	12	28	45	74	316	1,391	5,043	14,499	22,042	11,098	-
2000.....	54,816	16	13	33	65	84	292	1,321	4,893	14,786	22,254	11,058	1
1999.....	57,000	10	14	45	68	78	271	1,237	5,247	15,973	22,944	11,111	2

White, female

2007 1/.....	62,147	13	9	20	30	73	300	1,618	5,456	12,990	23,251	18,387	-
2006 1/.....	60,950	5	9	10	29	71	309	1,632	5,405	13,139	22,904	17,436	1
2005 1/.....	63,973	7	13	19	25	80	336	1,567	5,551	13,467	24,539	18,368	1
2004 1/.....	59,292	10	7	23	36	68	332	1,368	5,199	12,988	23,061	16,200	-
2003 1/.....	61,520	3	10	23	42	90	370	1,378	5,259	13,604	24,139	16,602	-
2002.....	59,986	9	9	24	49	88	360	1,359	4,973	13,486	23,577	16,051	1
2001.....	59,258	9	8	20	40	82	349	1,312	4,924	13,886	22,954	15,674	-
2000.....	58,024	7	15	30	38	82	332	1,263	4,639	13,979	22,771	14,868	-
1999.....	57,735	6	17	26	57	98	337	1,261	4,809	14,138	22,754	14,231	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Chronic lower respiratory diseases (J40-J47)

Black, both sexes

2007 1/.....	7,901	15	30	60	65	96	193	672	1,178	1,897	2,340	1,355	-
2006 1/.....	7,730	11	15	69	75	108	226	621	1,205	1,818	2,303	1,279	-
2005 1/.....	8,229	12	31	53	75	97	219	703	1,207	2,020	2,439	1,372	1
2004 1/.....	7,400	11	22	64	79	101	225	622	1,106	1,770	2,216	1,184	-
2003 1/.....	7,709	8	26	63	82	104	247	601	1,118	1,934	2,316	1,210	-
2002.....	7,831	14	30	76	79	104	298	599	1,076	1,916	2,398	1,240	1
2001.....	7,589	17	23	54	77	129	279	558	1,031	1,985	2,212	1,224	-
2000.....	7,607	11	21	70	81	116	275	599	1,073	2,014	2,152	1,195	-
1999.....	7,915	18	18	68	78	144	264	561	1,091	2,112	2,380	1,180	1

Black, male

2007 1/.....	4,207	13	13	41	44	58	99	304	672	1,061	1,296	606	-
2006 1/.....	4,136	6	12	44	47	61	103	319	690	1,028	1,253	573	-
2005 1/.....	4,464	6	22	34	48	53	103	354	669	1,159	1,375	640	1
2004 1/.....	4,046	10	14	44	48	66	108	317	648	1,013	1,208	570	-
2003 1/.....	4,247	3	21	43	49	72	108	295	622	1,102	1,355	577	-
2002.....	4,341	7	20	41	59	58	155	309	615	1,107	1,367	602	1
2001.....	4,187	7	17	32	48	70	128	276	563	1,160	1,294	592	-
2000.....	4,238	6	14	43	51	72	130	294	595	1,209	1,223	601	-
1999.....	4,500	9	13	48	52	69	126	286	629	1,255	1,413	599	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Chronic lower respiratory diseases (J40-J47)

Black, female

2007 1/.....	3,694	2	17	19	21	38	94	368	506	836	1,044	749	-
2006 1/.....	3,594	5	3	25	28	47	123	302	515	790	1,050	706	-
2005 1/.....	3,765	6	9	19	27	44	116	349	538	861	1,064	732	-
2004 1/.....	3,354	1	8	20	31	35	117	305	458	757	1,008	614	-
2003 1/.....	3,462	5	5	20	33	32	139	306	496	832	961	633	-
2002.....	3,490	7	10	35	20	46	143	290	461	809	1,031	638	-
2001.....	3,402	10	6	22	29	59	151	282	468	825	918	632	-
2000.....	3,369	5	7	27	30	44	145	305	478	805	929	594	-
1999.....	3,415	9	5	20	26	75	138	275	462	857	967	581	-

American Indian 2/, both sexes

2007 1/.....	611	1	2	2	2	2	10	48	100	173	185	86	-
2006 1/.....	508	2	-	2	1	1	12	33	86	137	166	68	-
2005 1/.....	518	1	1	2	-	3	12	33	91	142	164	69	-
2004 1/.....	486	1	1	1	3	2	8	29	89	136	161	55	-
2003 1/.....	512	-	1	-	2	3	14	35	92	130	163	72	-
2002.....	452	-	1	1	2	1	6	25	72	135	139	70	-
2001.....	427	1	-	1	3	2	10	18	72	122	129	68	1
2000.....	429	1	2	-	3	4	10	17	71	124	126	71	-
1999.....	406	1	-	-	1	2	9	19	63	123	139	49	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Chronic lower respiratory diseases (J40-J47)

American Indian 2/, male

2007 1/.....	299	-	1	1	1	1	7	24	49	92	86	37	-
2006 1/.....	234	2	-	1	1	1	7	15	44	62	73	28	-
2005 1/.....	257	-	-	1	-	1	3	18	45	65	92	32	-
2004 1/.....	235	1	-	-	2	1	5	13	46	68	75	24	-
2003 1/.....	269	-	-	-	2	1	8	19	47	67	91	34	-
2002.....	220	-	1	-	-	1	2	12	35	64	77	28	-
2001.....	200	1	-	-	2	1	3	8	34	56	63	31	1
2000.....	228	1	1	-	3	3	4	12	41	50	77	36	-
1999.....	197	-	-	-	-	1	5	10	28	68	63	22	-

American Indian 2/, female

2007 1/.....	312	1	1	1	1	1	3	24	51	81	99	49	-
2006 1/.....	274	-	-	1	-	-	5	18	42	75	93	40	-
2005 1/.....	261	1	1	1	-	2	9	15	46	77	72	37	-
2004 1/.....	251	-	1	1	1	1	3	16	43	68	86	31	-
2003 1/.....	243	-	1	-	-	2	6	16	45	63	72	38	-
2002.....	232	-	-	1	2	-	4	13	37	71	62	42	-
2001.....	227	-	-	1	1	1	7	10	38	66	66	37	-
2000.....	201	-	1	-	-	1	6	5	30	74	49	35	-
1999.....	209	1	-	-	1	1	4	9	35	55	76	27	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Chronic lower respiratory diseases (J40-J47)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	1,331	1	3	2	3	6	15	38	87	205	530	440	1
2006 1/.....	1,352	2	2	2	6	5	12	34	84	226	547	431	1
2005 1/.....	1,302	-	-	4	2	6	12	33	77	231	498	439	-
2004 1/.....	1,187	-	3	2	6	2	10	30	76	211	449	398	-
2003 1/.....	1,244	-	4	1	6	6	14	28	97	249	476	363	-
2002.....	1,138	-	4	4	2	7	8	39	79	236	446	312	1
2001.....	1,178	1	-	1	6	3	15	45	96	259	425	327	-
2000.....	1,133	1	-	6	3	10	21	51	63	254	419	304	1
1999.....	1,125	1	5	-	5	8	9	32	87	298	418	262	-

Asian or Pacific Islander 3/, male

2007 1/.....	795	1	3	2	2	5	8	20	56	127	322	248	1
2006 1/.....	847	2	-	1	5	5	4	20	53	146	355	255	1
2005 1/.....	803	-	-	2	1	4	9	21	54	140	319	253	-
2004 1/.....	743	-	1	1	5	2	6	14	51	137	291	235	-
2003 1/.....	801	-	3	1	4	4	8	17	63	162	325	214	-
2002.....	743	-	2	2	1	5	4	21	53	153	307	195	-
2001.....	749	1	-	1	4	1	8	27	62	162	292	191	-
2000.....	722	-	-	4	1	6	11	34	35	165	284	181	1
1999.....	718	-	3	-	3	4	8	18	54	181	285	162	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Chronic lower respiratory diseases (J40-J47)

Asian or Pacific Islander 3/, female

2007 1/.....	536	-	-	-	1	1	7	18	31	78	208	192	-
2006 1/.....	505	-	2	1	1	-	8	14	31	80	192	176	-
2005 1/.....	499	-	-	2	1	2	3	12	23	91	179	186	-
2004 1/.....	444	-	2	1	1	-	4	16	25	74	158	163	-
2003 1/.....	443	-	1	-	2	2	6	11	34	87	151	149	-
2002.....	395	-	2	2	1	2	4	18	26	83	139	117	1
2001.....	429	-	-	-	2	2	7	18	34	97	133	136	-
2000.....	411	1	-	2	2	4	10	17	28	89	135	123	-
1999.....	407	1	2	-	2	4	1	14	33	117	133	100	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Bronchitis, chronic and unspecified (J40-J42)													
All races, both sexes													
2007.....	667	24	10	2	1	5	14	24	52	88	181	266	-
2006.....	740	19	12	2	9	5	10	27	49	89	200	318	-
2005.....	866	25	12	3	5	6	15	27	73	119	205	376	-
2004.....	773	17	12	4	7	5	12	35	67	97	199	318	-
2003.....	850	20	13	6	7	10	17	35	47	104	246	345	-
2002.....	955	24	13	5	6	6	18	36	54	135	286	372	-
2001.....	959	21	8	-	5	6	21	42	61	133	261	401	-
2000.....	1,167	18	14	5	5	8	20	38	67	158	333	501	-
1999.....	1,172	20	19	4	7	11	18	38	97	173	339	446	-
All races, male													
2007.....	273	13	6	1	1	4	5	13	22	34	86	88	-
2006.....	299	12	9	1	6	4	9	13	23	40	82	100	-
2005.....	361	16	5	2	3	3	10	18	35	52	100	117	-
2004.....	338	12	9	4	4	4	5	22	35	54	83	106	-
2003.....	351	15	6	3	6	3	9	23	26	46	113	101	-
2002.....	420	15	8	2	3	2	12	27	28	74	127	122	-
2001.....	396	11	5	-	3	2	8	21	36	76	115	119	-
2000.....	503	9	7	3	4	7	14	16	32	79	174	158	-
1999.....	498	9	11	3	4	3	5	19	51	94	157	142	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Bronchitis, chronic and unspecified (J40-J42)													
All races, female													
2007.....	394	11	4	1	-	1	9	11	30	54	95	178	-
2006.....	441	7	3	1	3	1	1	14	26	49	118	218	-
2005.....	505	9	7	1	2	3	5	9	38	67	105	259	-
2004.....	435	5	3	-	3	1	7	13	32	43	116	212	-
2003.....	499	5	7	3	1	7	8	12	21	58	133	244	-
2002.....	535	9	5	3	3	4	6	9	26	61	159	250	-
2001.....	563	10	3	-	2	4	13	21	25	57	146	282	-
2000.....	664	9	7	2	1	1	6	22	35	79	159	343	-
1999.....	674	11	8	1	3	8	13	19	46	79	182	304	-
White, both sexes													
2007 1/.....	592	13	6	1	-	3	11	22	44	76	171	245	-
2006 1/.....	657	9	9	1	8	4	8	21	42	80	178	297	-
2005 1/.....	784	15	10	3	4	6	11	23	58	106	188	360	-
2004 1/.....	691	12	6	3	5	5	8	26	56	91	186	293	-
2003 1/.....	751	15	8	3	6	6	13	24	38	94	225	319	-
2002.....	876	16	7	3	4	5	16	29	46	127	267	356	-
2001.....	878	11	6	-	4	5	18	36	57	116	242	383	-
2000.....	1,073	9	7	4	4	6	14	32	61	140	310	486	-
1999.....	1,056	11	15	4	6	8	16	24	88	155	306	423	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Bronchitis, chronic and unspecified (J40-J42)													
White, male													
2007 1/.....	238	5	4	1	-	3	4	12	19	28	81	81	-
2006 1/.....	259	6	7	-	5	4	7	9	18	36	71	96	-
2005 1/.....	315	10	4	2	2	3	7	16	31	44	89	107	-
2004 1/.....	295	7	5	3	3	4	2	14	30	51	79	97	-
2003 1/.....	309	12	5	1	5	2	7	17	23	40	102	95	-
2002.....	376	11	3	2	2	2	10	20	23	68	118	117	-
2001.....	354	7	5	-	2	1	7	17	34	67	103	111	-
2000.....	458	5	3	2	3	5	8	15	28	71	165	153	-
1999.....	440	6	8	3	3	2	5	11	45	85	138	134	-
White, female													
2007 1/.....	354	8	2	-	-	-	7	10	25	48	90	164	-
2006 1/.....	398	3	2	1	3	-	1	12	24	44	107	201	-
2005 1/.....	469	5	6	1	2	3	4	7	27	62	99	253	-
2004 1/.....	396	5	1	-	2	1	6	12	26	40	107	196	-
2003 1/.....	442	3	3	2	1	4	6	7	15	54	123	224	-
2002.....	500	5	4	1	2	3	6	9	23	59	149	239	-
2001.....	524	4	1	-	2	4	11	19	23	49	139	272	-
2000.....	615	4	4	2	1	1	6	17	33	69	145	333	-
1999.....	616	5	7	1	3	6	11	13	43	70	168	289	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Bronchitis, chronic and unspecified (J40-J42)

Black, both sexes

2007 1/.....	56	9	3	1	1	2	1	2	4	10	9	14	-
2006 1/.....	66	8	3	1	-	1	2	5	7	9	16	14	-
2005 1/.....	54	9	1	-	1	-	3	4	11	9	10	6	-
2004 1/.....	65	5	5	1	2	-	4	9	9	5	8	17	-
2003 1/.....	75	5	3	3	1	3	4	11	8	6	12	19	-
2002.....	66	8	6	2	2	1	2	7	5	6	13	14	-
2001.....	63	10	2	-	1	1	3	5	4	15	11	11	-
2000.....	80	7	6	1	1	1	6	6	4	16	20	12	-
1999.....	89	7	1	-	1	3	2	12	8	15	26	14	-

Black, male

2007 1/.....	24	7	1	-	1	1	-	1	-	5	4	4	-
2006 1/.....	31	4	2	1	-	-	2	3	5	4	8	2	-
2005 1/.....	33	6	1	-	1	-	2	2	4	6	7	4	-
2004 1/.....	35	5	4	1	1	-	3	8	3	3	2	5	-
2003 1/.....	30	3	-	2	1	1	2	6	3	2	7	3	-
2002.....	35	4	5	-	1	-	2	7	2	4	5	5	-
2001.....	32	4	-	-	1	1	1	4	2	8	6	5	-
2000.....	35	3	3	1	1	1	6	1	3	6	7	3	-
1999.....	44	3	1	-	1	1	-	7	5	7	16	3	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Bronchitis, chronic and unspecified (J40-J42)

Black, female

2007 1/.....	32	2	2	1	-	1	1	1	4	5	5	10	-
2006 1/.....	35	4	1	-	-	1	-	2	2	5	8	12	-
2005 1/.....	21	3	-	-	-	-	1	2	7	3	3	2	-
2004 1/.....	30	-	1	-	1	-	1	1	6	2	6	12	-
2003 1/.....	45	2	3	1	-	2	2	5	5	4	5	16	-
2002.....	31	4	1	2	1	1	-	-	3	2	8	9	-
2001.....	31	6	2	-	-	-	2	1	2	7	5	6	-
2000.....	45	4	3	-	-	-	-	5	1	10	13	9	-
1999.....	45	4	-	-	-	2	2	5	3	8	10	11	-

American Indian 2/, both sexes

2007 1/.....	2	1	-	-	-	-	-	-	-	1	-	-	-
2006 1/.....	2	-	-	-	-	-	-	-	-	-	1	1	-
2005 1/.....	5	1	1	-	-	-	-	-	1	-	2	-	-
2004 1/.....	2	-	-	-	-	-	-	-	-	-	1	1	-
2003 1/.....	5	-	1	-	-	-	-	-	1	-	2	1	-
2002.....	1	-	-	-	-	-	-	-	1	-	-	-	-
2001.....	2	-	-	-	-	-	-	1	-	-	1	-	-
2000.....	2	1	1	-	-	-	-	-	-	-	-	-	-
1999.....	7	1	-	-	-	-	-	2	-	1	1	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Bronchitis, chronic and unspecified (J40-J42)

American Indian 2/, male

2007 1/.....	1	-	-	-	-	-	-	-	-	1	-	-	-
2006 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2005 1/.....	1	-	-	-	-	-	-	-	-	-	1	-	-
2004 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 1/.....	1	-	-	-	-	-	-	-	-	-	-	1	-
2002.....	1	-	-	-	-	-	-	-	1	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	1	-	-
2000.....	2	1	1	-	-	-	-	-	-	-	-	-	-
1999.....	2	-	-	-	-	-	-	1	-	1	-	-	-

American Indian 2/, female

2007 1/.....	1	1	-	-	-	-	-	-	-	-	-	-	-
2006 1/.....	2	-	-	-	-	-	-	-	-	-	1	1	-
2005 1/.....	4	1	1	-	-	-	-	-	1	-	1	-	-
2004 1/.....	2	-	-	-	-	-	-	-	-	-	1	1	-
2003 1/.....	4	-	1	-	-	-	-	-	1	-	2	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	1	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	5	1	-	-	-	-	-	1	-	-	1	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Bronchitis, chronic and unspecified (J40-J42)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	17	1	1	-	-	-	2	-	4	1	1	7	-
2006 1/.....	15	2	-	-	1	-	-	1	-	-	5	6	-
2005 1/.....	23	-	-	-	-	-	1	-	3	4	5	10	-
2004 1/.....	15	-	1	-	-	-	-	-	2	1	4	7	-
2003 1/.....	19	-	1	-	-	1	-	-	-	4	7	6	-
2002.....	12	-	-	-	-	-	-	-	2	2	6	2	-
2001.....	16	-	-	-	-	-	-	-	-	2	7	7	-
2000.....	12	1	-	-	-	1	-	-	2	2	3	3	-
1999.....	20	1	3	-	-	-	-	-	1	2	6	7	-

Asian or Pacific Islander 3/, male

2007 1/.....	10	1	1	-	-	-	1	-	3	-	1	3	-
2006 1/.....	9	2	-	-	1	-	-	1	-	-	3	2	-
2005 1/.....	12	-	-	-	-	-	1	-	-	2	3	6	-
2004 1/.....	8	-	-	-	-	-	-	-	2	-	2	4	-
2003 1/.....	11	-	1	-	-	-	-	-	-	4	4	2	-
2002.....	8	-	-	-	-	-	-	-	2	2	4	-	-
2001.....	9	-	-	-	-	-	-	-	-	1	5	3	-
2000.....	8	-	-	-	-	1	-	-	1	2	2	2	-
1999.....	12	-	2	-	-	-	-	-	1	1	3	5	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Bronchitis, chronic and unspecified (J40-J42)

Asian or Pacific Islander 3/, female

2007 1/.....	7	-	-	-	-	-	1	-	1	1	-	4	-
2006 1/.....	6	-	-	-	-	-	-	-	-	-	2	4	-
2005 1/.....	11	-	-	-	-	-	-	-	3	2	2	4	-
2004 1/.....	7	-	1	-	-	-	-	-	-	1	2	3	-
2003 1/.....	8	-	-	-	-	1	-	-	-	-	3	4	-
2002.....	4	-	-	-	-	-	-	-	-	-	2	2	-
2001.....	7	-	-	-	-	-	-	-	-	1	2	4	-
2000.....	4	1	-	-	-	-	-	-	1	-	1	1	-
1999.....	8	1	1	-	-	-	-	-	-	1	3	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Emphysema (J43)													
All races, both sexes													
2007.....	12,790	3	-	1	1	10	60	486	1,590	3,294	4,835	2,509	1
2006.....	12,551	2	-	2	1	9	86	454	1,523	3,273	4,799	2,400	2
2005.....	14,002	2	1	-	-	8	77	444	1,677	3,749	5,400	2,643	1
2004.....	13,639	3	2	1	3	10	80	413	1,583	3,793	5,324	2,427	-
2003.....	14,861	-	1	-	4	5	87	450	1,755	4,063	5,920	2,574	2
2002.....	15,489	1	3	-	2	10	94	454	1,703	4,388	6,109	2,724	1
2001.....	16,242	3	-	1	2	7	75	448	1,829	4,804	6,337	2,736	-
2000.....	16,700	3	2	-	5	11	82	474	1,807	5,120	6,511	2,685	-
1999.....	17,787	1	-	1	1	9	97	468	2,019	5,656	6,955	2,580	-
All races, male													
2007.....	6,598	2	-	-	-	5	41	305	903	1,789	2,457	1,095	1
2006.....	6,318	2	-	1	-	6	49	267	811	1,697	2,408	1,075	2
2005.....	7,264	-	1	-	-	5	54	277	973	2,039	2,767	1,147	1
2004.....	7,075	2	-	1	3	5	49	244	873	2,051	2,725	1,122	-
2003.....	7,549	-	1	-	2	3	49	279	932	2,141	2,995	1,145	2
2002.....	8,084	-	3	-	-	8	66	267	925	2,335	3,228	1,251	1
2001.....	8,415	1	-	1	1	4	52	266	966	2,554	3,255	1,315	-
2000.....	8,671	3	-	-	5	5	46	280	943	2,738	3,373	1,278	-
1999.....	9,480	-	-	-	1	6	49	268	1,085	3,136	3,677	1,258	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Emphysema (J43)													
All races, female													
2007.....	6,192	1	-	1	1	5	19	181	687	1,505	2,378	1,414	-
2006.....	6,233	-	-	1	1	3	37	187	712	1,576	2,391	1,325	-
2005.....	6,738	2	-	-	-	3	23	167	704	1,710	2,633	1,496	-
2004.....	6,564	1	2	-	-	5	31	169	710	1,742	2,599	1,305	-
2003.....	7,312	-	-	-	2	2	38	171	823	1,922	2,925	1,429	-
2002.....	7,405	1	-	-	2	2	28	187	778	2,053	2,881	1,473	-
2001.....	7,827	2	-	-	1	3	23	182	863	2,250	3,082	1,421	-
2000.....	8,029	-	2	-	-	6	36	194	864	2,382	3,138	1,407	-
1999.....	8,307	1	-	1	-	3	48	200	934	2,520	3,278	1,322	-
White, both sexes													
2007 1/.....	11,886	2	-	1	1	5	50	410	1,442	3,063	4,530	2,381	1
2006 1/.....	11,689	1	-	1	1	7	66	396	1,379	3,049	4,511	2,276	2
2005 1/.....	12,987	1	1	-	-	6	58	367	1,522	3,480	5,054	2,497	1
2004 1/.....	12,757	1	1	1	2	7	61	342	1,431	3,559	5,052	2,300	-
2003 1/.....	13,856	-	1	-	4	3	70	387	1,595	3,795	5,574	2,425	2
2002.....	14,476	1	1	-	1	8	67	390	1,560	4,081	5,781	2,585	1
2001.....	15,243	2	-	-	1	4	64	383	1,663	4,497	6,034	2,595	-
2000.....	15,607	3	-	-	2	10	63	383	1,645	4,783	6,193	2,525	-
1999.....	16,674	-	-	1	-	7	70	390	1,851	5,320	6,597	2,438	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Emphysema (J43)													
White, male													
2007 1/.....	6,032	1	-	-	-	2	32	256	802	1,647	2,269	1,022	1
2006 1/.....	5,784	1	-	1	-	4	38	221	709	1,567	2,230	1,011	2
2005 1/.....	6,618	-	1	-	-	4	39	225	863	1,868	2,553	1,064	1
2004 1/.....	6,511	-	-	1	2	3	35	196	772	1,896	2,557	1,049	-
2003 1/.....	6,915	-	1	-	2	1	39	234	831	1,980	2,774	1,051	2
2002.....	7,426	-	1	-	-	6	48	223	828	2,134	3,016	1,169	1
2001.....	7,798	-	-	-	1	2	44	228	859	2,362	3,066	1,236	-
2000.....	7,985	3	-	-	2	4	34	218	847	2,519	3,188	1,170	-
1999.....	8,754	-	-	-	-	4	35	212	975	2,900	3,449	1,179	-
White, female													
2007 1/.....	5,854	1	-	1	1	3	18	154	640	1,416	2,261	1,359	-
2006 1/.....	5,905	-	-	-	1	3	28	175	670	1,482	2,281	1,265	-
2005 1/.....	6,369	1	-	-	-	2	19	142	659	1,612	2,501	1,433	-
2004 1/.....	6,246	1	1	-	-	4	26	146	659	1,663	2,495	1,251	-
2003 1/.....	6,941	-	-	-	2	2	31	153	764	1,815	2,800	1,374	-
2002.....	7,050	1	-	-	1	2	19	167	732	1,947	2,765	1,416	-
2001.....	7,445	2	-	-	-	2	20	155	804	2,135	2,968	1,359	-
2000.....	7,622	-	-	-	-	6	29	165	798	2,264	3,005	1,355	-
1999.....	7,920	-	-	1	-	3	35	178	876	2,420	3,148	1,259	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Emphysema (J43)													
Black, both sexes													
2007 1/.....	733	1	-	-	-	5	8	69	129	192	233	96	-
2006 1/.....	707	1	-	-	-	1	20	51	136	181	225	92	-
2005 1/.....	815	1	-	-	-	2	17	69	138	221	265	102	-
2004 1/.....	719	2	-	-	1	2	15	64	131	186	220	98	-
2003 1/.....	810	-	-	-	-	2	15	56	142	221	268	106	-
2002.....	850	-	1	-	1	2	26	57	117	268	270	108	-
2001.....	811	1	-	1	1	3	10	59	142	256	223	115	-
2000.....	901	-	1	-	3	1	17	85	147	281	247	119	-
1999.....	930	1	-	-	-	-	25	75	146	281	284	118	-
Black, male													
2007 1/.....	457	1	-	-	-	3	7	42	88	123	139	54	-
2006 1/.....	430	1	-	-	-	1	11	40	96	104	132	45	-
2005 1/.....	509	-	-	-	-	1	13	47	100	139	154	55	-
2004 1/.....	460	2	-	-	1	1	11	45	89	123	131	57	-
2003 1/.....	498	-	-	-	-	2	8	38	90	129	163	68	-
2002.....	541	-	1	-	-	2	18	39	77	174	170	60	-
2001.....	487	1	-	1	-	2	8	33	91	156	135	60	-
2000.....	559	-	-	-	3	1	11	56	88	187	137	76	-
1999.....	598	-	-	-	-	-	13	54	94	197	178	62	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Emphysema (J43)													
Black, female													
2007 1/.....	276	-	-	-	-	2	1	27	41	69	94	42	-
2006 1/.....	277	-	-	-	-	-	9	11	40	77	93	47	-
2005 1/.....	306	1	-	-	-	1	4	22	38	82	111	47	-
2004 1/.....	259	-	-	-	-	1	4	19	42	63	89	41	-
2003 1/.....	312	-	-	-	-	-	7	18	52	92	105	38	-
2002.....	309	-	-	-	1	-	8	18	40	94	100	48	-
2001.....	324	-	-	-	1	1	2	26	51	100	88	55	-
2000.....	342	-	1	-	-	-	6	29	59	94	110	43	-
1999.....	332	1	-	-	-	-	12	21	52	84	106	56	-
American Indian 2/, both sexes													
2007 1/.....	49	-	-	-	-	-	2	4	9	11	12	11	-
2006 1/.....	32	-	-	-	-	1	-	2	3	13	10	3	-
2005 1/.....	54	-	-	-	-	-	1	4	11	14	18	6	-
2004 1/.....	46	-	-	-	-	-	2	4	12	13	10	5	-
2003 1/.....	46	-	-	-	-	-	-	4	8	9	18	7	-
2002.....	47	-	-	-	-	-	1	2	10	14	13	7	-
2001.....	52	-	-	-	-	-	-	-	10	19	19	4	-
2000.....	59	-	1	-	-	-	1	2	11	19	18	7	-
1999.....	45	-	-	-	-	1	2	2	7	16	13	4	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Emphysema (J43)													
American Indian 2/, male													
2007 1/.....	25	-	-	-	-	-	2	4	5	5	4	5	-
2006 1/.....	16	-	-	-	-	1	-	2	2	5	6	-	-
2005 1/.....	30	-	-	-	-	-	1	1	6	8	12	2	-
2004 1/.....	24	-	-	-	-	-	2	1	6	7	6	2	-
2003 1/.....	28	-	-	-	-	-	-	4	5	4	13	2	-
2002.....	30	-	-	-	-	-	-	-	7	8	11	4	-
2001.....	29	-	-	-	-	-	-	-	7	12	7	3	-
2000.....	29	-	-	-	-	-	1	2	5	6	9	6	-
1999.....	23	-	-	-	-	1	1	1	4	8	6	2	-
American Indian 2/, female													
2007 1/.....	24	-	-	-	-	-	-	-	4	6	8	6	-
2006 1/.....	16	-	-	-	-	-	-	-	1	8	4	3	-
2005 1/.....	24	-	-	-	-	-	-	3	5	6	6	4	-
2004 1/.....	22	-	-	-	-	-	-	3	6	6	4	3	-
2003 1/.....	18	-	-	-	-	-	-	-	3	5	5	5	-
2002.....	17	-	-	-	-	-	1	2	3	6	2	3	-
2001.....	23	-	-	-	-	-	-	-	3	7	12	1	-
2000.....	30	-	1	-	-	-	-	-	6	13	9	1	-
1999.....	22	-	-	-	-	-	1	1	3	8	7	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Emphysema (J43)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	122	-	-	-	-	-	-	3	10	28	60	21	-
2006 1/.....	123	-	-	1	-	-	-	5	5	30	53	29	-
2005 1/.....	146	-	-	-	-	-	1	4	6	34	63	38	-
2004 1/.....	117	-	1	-	-	1	2	3	9	35	42	24	-
2003 1/.....	149	-	-	-	-	-	2	3	10	38	60	36	-
2002.....	116	-	1	-	-	-	-	5	16	25	45	24	-
2001.....	136	-	-	-	-	-	1	6	14	32	61	22	-
2000.....	133	-	-	-	-	-	1	4	4	37	53	34	-
1999.....	138	-	-	-	1	1	-	1	15	39	61	20	-

Asian or Pacific Islander 3/, male

2007 1/.....	84	-	-	-	-	-	-	3	8	14	45	14	-
2006 1/.....	88	-	-	-	-	-	-	4	4	21	40	19	-
2005 1/.....	107	-	-	-	-	-	1	4	4	24	48	26	-
2004 1/.....	80	-	-	-	-	1	1	2	6	25	31	14	-
2003 1/.....	108	-	-	-	-	-	2	3	6	28	45	24	-
2002.....	87	-	1	-	-	-	-	5	13	19	31	18	-
2001.....	101	-	-	-	-	-	-	5	9	24	47	16	-
2000.....	98	-	-	-	-	-	-	4	3	26	39	26	-
1999.....	105	-	-	-	1	1	-	1	12	31	44	15	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Emphysema (J43)

Asian or Pacific Islander 3/, female

2007 1/.....	38	-	-	-	-	-	-	-	2	14	15	7	-
2006 1/.....	35	-	-	1	-	-	-	1	1	9	13	10	-
2005 1/.....	39	-	-	-	-	-	-	-	2	10	15	12	-
2004 1/.....	37	-	1	-	-	-	1	1	3	10	11	10	-
2003 1/.....	41	-	-	-	-	-	-	-	4	10	15	12	-
2002.....	29	-	-	-	-	-	-	-	3	6	14	6	-
2001.....	35	-	-	-	-	-	1	1	5	8	14	6	-
2000.....	35	-	-	-	-	-	1	-	1	11	14	8	-
1999.....	33	-	-	-	-	-	-	-	3	8	17	5	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Asthma (J45-J46)													
All races, both sexes													
2007.....	3,447	4	41	107	133	201	320	538	461	412	569	659	2
2006.....	3,613	6	26	99	135	194	373	566	492	443	626	653	-
2005.....	3,884	4	37	97	131	207	369	595	520	475	709	740	-
2004.....	3,816	7	29	105	159	197	376	562	520	504	673	684	-
2003.....	4,099	7	37	110	158	227	411	632	562	532	752	671	-
2002.....	4,261	4	43	123	169	235	472	608	536	583	812	675	1
2001.....	4,269	10	31	99	140	233	455	603	553	634	802	709	-
2000.....	4,487	8	32	129	167	249	458	614	540	734	849	707	-
1999.....	4,657	5	27	126	182	262	447	607	583	773	925	720	-
All races, male													
2007.....	1,274	2	21	70	89	119	134	209	194	135	172	127	2
2006.....	1,296	3	17	71	82	105	155	211	168	146	185	153	-
2005.....	1,315	1	25	61	83	101	151	215	170	150	218	140	-
2004.....	1,386	7	19	64	99	113	158	213	187	178	181	167	-
2003.....	1,493	4	29	71	91	127	160	236	214	173	234	154	-
2002.....	1,580	3	30	69	110	129	198	233	202	198	256	152	-
2001.....	1,479	6	22	57	80	116	195	219	182	220	225	157	-
2000.....	1,632	6	20	74	102	142	196	238	200	253	243	158	-
1999.....	1,620	3	16	85	108	128	184	197	207	267	273	152	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Asthma (J45-J46)													
All races, female													
2007.....	2,173	2	20	37	44	82	186	329	267	277	397	532	-
2006.....	2,317	3	9	28	53	89	218	355	324	297	441	500	-
2005.....	2,569	3	12	36	48	106	218	380	350	325	491	600	-
2004.....	2,430	-	10	41	60	84	218	349	333	326	492	517	-
2003.....	2,606	3	8	39	67	100	251	396	348	359	518	517	-
2002.....	2,681	1	13	54	59	106	274	375	334	385	556	523	1
2001.....	2,790	4	9	42	60	117	260	384	371	414	577	552	-
2000.....	2,855	2	12	55	65	107	262	376	340	481	606	549	-
1999.....	3,037	2	11	41	74	134	263	410	376	506	652	568	-
White, both sexes													
2007 1/.....	2,376	2	11	45	68	116	187	311	295	297	462	581	1
2006 1/.....	2,497	4	13	33	55	98	220	341	311	320	525	577	-
2005 1/.....	2,714	2	9	39	58	116	225	362	343	322	574	664	-
2004 1/.....	2,658	5	13	45	76	103	226	320	338	386	546	600	-
2003 1/.....	2,888	4	13	49	78	131	242	377	379	389	619	607	-
2002.....	3,014	2	18	47	90	145	266	361	351	424	701	609	-
2001.....	2,990	5	11	45	67	114	247	375	349	477	673	627	-
2000.....	3,144	5	18	54	89	132	256	360	347	547	726	610	-
1999.....	3,328	1	11	59	106	127	266	374	390	570	798	626	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Asthma (J45-J46)													
White, male													
2007 1/.....	789	-	7	27	45	64	73	127	117	91	130	107	1
2006 1/.....	805	2	7	27	31	48	90	115	95	99	153	138	-
2005 1/.....	843	1	4	24	37	53	91	134	110	99	166	124	-
2004 1/.....	851	5	9	24	47	51	92	115	106	129	133	140	-
2003 1/.....	961	4	8	29	44	60	91	139	136	127	184	139	-
2002.....	1,017	1	15	28	52	74	97	139	128	137	212	134	-
2001.....	937	4	6	26	35	51	104	127	111	160	180	133	-
2000.....	1,057	3	9	28	55	70	105	150	119	185	207	126	-
1999.....	1,046	1	5	38	60	58	99	118	136	183	231	117	-
White, female													
2007 1/.....	1,587	2	4	18	23	52	114	184	178	206	332	474	-
2006 1/.....	1,692	2	6	6	24	50	130	226	216	221	372	439	-
2005 1/.....	1,871	1	5	15	21	63	134	228	233	223	408	540	-
2004 1/.....	1,807	-	4	21	29	52	134	205	232	257	413	460	-
2003 1/.....	1,927	-	5	20	34	71	151	238	243	262	435	468	-
2002.....	1,997	1	3	19	38	71	169	222	223	287	489	475	-
2001.....	2,053	1	5	19	32	63	143	248	238	317	493	494	-
2000.....	2,087	2	9	26	34	62	151	210	228	362	519	484	-
1999.....	2,282	-	6	21	46	69	167	256	254	387	567	509	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Asthma (J45-J46)													
Black, both sexes													
2007 1/.....	902	2	27	58	60	78	121	204	145	94	66	47	-
2006 1/.....	957	2	12	63	74	91	139	204	164	99	61	48	-
2005 1/.....	1,016	2	28	52	71	85	133	211	164	127	99	44	-
2004 1/.....	1,008	2	14	57	74	91	142	221	163	98	91	55	-
2003 1/.....	1,030	3	22	60	75	90	157	232	141	120	89	41	-
2002.....	1,096	2	22	71	75	85	199	223	170	132	77	40	-
2001.....	1,108	4	20	52	67	116	190	206	172	128	100	53	-
2000.....	1,158	3	14	69	74	104	179	228	168	153	95	71	-
1999.....	1,145	4	14	67	71	130	168	213	162	158	94	64	-
Black, male													
2007 1/.....	406	2	12	40	41	50	55	71	66	37	25	7	-
2006 1/.....	423	1	10	42	46	52	62	84	68	35	16	7	-
2005 1/.....	411	-	21	34	45	45	56	69	54	40	39	8	-
2004 1/.....	455	2	9	39	45	60	61	89	67	40	31	12	-
2003 1/.....	445	-	20	41	43	63	63	87	52	38	31	7	-
2002.....	497	2	14	39	57	51	98	84	67	51	24	10	-
2001.....	459	1	16	30	41	63	81	85	52	47	30	13	-
2000.....	481	3	11	42	44	64	79	74	67	54	21	22	-
1999.....	481	2	10	47	46	68	74	72	57	62	27	16	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Asthma (J45-J46)													
Black, female													
2007 1/.....	496	-	15	18	19	28	66	133	79	57	41	40	-
2006 1/.....	534	1	2	21	28	39	77	120	96	64	45	41	-
2005 1/.....	605	2	7	18	26	40	77	142	110	87	60	36	-
2004 1/.....	553	-	5	18	29	31	81	132	96	58	60	43	-
2003 1/.....	585	3	2	19	32	27	94	145	89	82	58	34	-
2002.....	599	-	8	32	18	34	101	139	103	81	53	30	-
2001.....	649	3	4	22	26	53	109	121	120	81	70	40	-
2000.....	677	-	3	27	30	40	100	154	101	99	74	49	-
1999.....	664	2	4	20	25	62	94	141	105	96	67	48	-
American Indian 2/, both sexes													
2007 1/.....	34	-	1	2	2	2	3	7	8	5	2	2	-
2006 1/.....	20	-	-	2	1	-	5	3	2	4	2	1	-
2005 1/.....	31	-	-	2	-	2	7	6	4	6	1	3	-
2004 1/.....	19	-	1	1	3	2	2	4	3	1	2	-	-
2003 1/.....	38	-	-	-	1	3	5	9	8	3	7	2	-
2002.....	25	-	-	1	2	-	4	3	3	7	4	1	-
2001.....	33	1	-	1	1	2	7	4	7	4	2	4	-
2000.....	31	-	-	-	2	4	7	1	7	6	2	2	-
1999.....	29	-	-	-	1	1	4	4	7	6	4	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Asthma (J45-J46)													
American Indian 2/, male													
2007 1/.....	12	-	-	1	1	1	2	3	2	1	-	1	-
2006 1/.....	7	-	-	1	1	-	1	1	-	1	2	-	-
2005 1/.....	11	-	-	1	-	-	-	4	2	3	1	-	-
2004 1/.....	8	-	-	-	2	1	1	2	2	-	-	-	-
2003 1/.....	18	-	-	-	1	1	4	2	5	-	4	1	-
2002.....	9	-	-	-	-	-	2	1	-	3	3	-	-
2001.....	11	1	-	-	1	1	2	-	4	1	-	1	-
2000.....	14	-	-	-	2	3	2	-	3	1	2	1	-
1999.....	12	-	-	-	-	-	3	-	2	3	2	2	-
American Indian 2/, female													
2007 1/.....	22	-	1	1	1	1	1	4	6	4	2	1	-
2006 1/.....	13	-	-	1	-	-	4	2	2	3	-	1	-
2005 1/.....	20	-	-	1	-	2	7	2	2	3	-	3	-
2004 1/.....	11	-	1	1	1	1	1	2	1	1	2	-	-
2003 1/.....	20	-	-	-	-	2	1	7	3	3	3	1	-
2002.....	16	-	-	1	2	-	2	2	3	4	1	1	-
2001.....	22	-	-	1	-	1	5	4	3	3	2	3	-
2000.....	17	-	-	-	-	1	5	1	4	5	-	1	-
1999.....	17	-	-	-	1	1	1	4	5	3	2	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Asthma (J45-J46)													
Asian or Pacific Islander 3/, both sexes													
2007 1/.....	135	-	2	2	3	5	9	16	13	16	39	29	1
2006 1/.....	139	-	1	1	5	5	9	18	15	20	38	27	-
2005 1/.....	123	-	-	4	2	4	4	16	9	20	35	29	-
2004 1/.....	131	-	1	2	6	1	6	17	16	19	34	29	-
2003 1/.....	143	-	2	1	4	3	7	14	34	20	37	21	-
2002.....	126	-	3	4	2	5	3	21	12	20	30	25	1
2001.....	138	-	-	1	5	1	11	18	25	25	27	25	-
2000.....	154	-	-	6	2	9	16	25	18	28	26	24	-
1999.....	155	-	2	-	4	4	9	16	24	39	29	28	-
Asian or Pacific Islander 3/, male													
2007 1/.....	67	-	2	2	2	4	4	8	9	6	17	12	1
2006 1/.....	61	-	-	1	4	5	2	11	5	11	14	8	-
2005 1/.....	50	-	-	2	1	3	4	8	4	8	12	8	-
2004 1/.....	72	-	1	1	5	1	4	7	12	9	17	15	-
2003 1/.....	69	-	1	1	3	3	2	8	21	8	15	7	-
2002.....	57	-	1	2	1	4	1	9	7	7	17	8	-
2001.....	72	-	-	1	3	1	8	7	15	12	15	10	-
2000.....	80	-	-	4	1	5	10	14	11	13	13	9	-
1999.....	81	-	1	-	2	2	8	7	12	19	13	17	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Asthma (J45-J46)

Asian or Pacific Islander 3/, female

2007 1/.....	68	-	-	-	1	1	5	8	4	10	22	17	-
2006 1/.....	78	-	1	-	1	-	7	7	10	9	24	19	-
2005 1/.....	73	-	-	2	1	1	-	8	5	12	23	21	-
2004 1/.....	59	-	-	1	1	-	2	10	4	10	17	14	-
2003 1/.....	74	-	1	-	1	-	5	6	13	12	22	14	-
2002.....	69	-	2	2	1	1	2	12	5	13	13	17	1
2001.....	66	-	-	-	2	-	3	11	10	13	12	15	-
2000.....	74	-	-	2	1	4	6	11	7	15	13	15	-
1999.....	74	-	1	-	2	2	1	9	12	20	16	11	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other chronic lower respiratory diseases (J44,J47)													
All races, both sexes													
2007.....	111,020	12	6	8	14	47	402	3,105	10,674	24,870	42,456	29,423	3
2006.....	107,679	3	5	12	6	47	372	2,877	10,311	24,431	41,781	27,832	2
2005.....	112,181	2	6	4	12	37	429	2,911	10,477	25,567	44,019	28,714	3
2004.....	103,759	8	5	10	10	43	419	2,501	9,584	23,996	41,372	25,810	1
2003.....	106,572	4	4	2	22	40	435	2,420	9,713	25,220	42,368	26,344	-
2002.....	104,111	10	6	8	15	50	424	2,377	8,987	24,682	42,034	25,513	5
2001.....	101,543	7	4	4	24	44	418	2,231	8,723	25,180	40,362	24,545	1
2000.....	99,655	7	3	5	13	28	370	2,125	8,325	25,145	40,029	23,603	2
1999.....	100,565	10	8	8	19	48	328	1,997	8,598	26,042	40,416	23,087	4
All races, male													
2007.....	53,090	10	3	7	6	22	212	1,598	5,614	12,721	20,724	12,170	3
2006.....	51,347	3	3	5	5	22	183	1,467	5,380	12,269	20,492	11,517	1
2005.....	53,495	2	2	-	9	21	211	1,524	5,411	13,173	21,394	11,746	2
2004.....	49,847	3	2	6	4	29	219	1,327	4,934	12,220	20,266	10,836	1
2003.....	51,321	4	2	1	15	23	211	1,288	5,071	12,973	20,621	11,112	-
2002.....	50,629	5	3	3	7	26	221	1,268	4,628	12,732	20,821	10,911	4
2001.....	49,407	4	2	3	15	24	200	1,196	4,518	13,027	20,096	10,321	1
2000.....	49,198	5	1	3	9	11	181	1,127	4,389	13,140	20,048	10,282	2
1999.....	50,817	7	3	5	10	15	172	1,067	4,615	13,980	20,598	10,342	3

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other chronic lower respiratory diseases (J44,J47)													
All races, female													
2007.....	57,930	2	3	1	8	25	190	1,507	5,060	12,149	21,732	17,253	-
2006.....	56,332	-	2	7	1	25	189	1,410	4,931	12,162	21,289	16,315	1
2005.....	58,686	-	4	4	3	16	218	1,387	5,066	12,394	22,625	16,968	1
2004.....	53,912	5	3	4	6	14	200	1,174	4,650	11,776	21,106	14,974	-
2003.....	55,251	-	2	1	7	17	224	1,132	4,642	12,247	21,747	15,232	-
2002.....	53,482	5	3	5	8	24	203	1,109	4,359	11,950	21,213	14,602	1
2001.....	52,136	3	2	1	9	20	218	1,035	4,205	12,153	20,266	14,224	-
2000.....	50,457	2	2	2	4	17	189	998	3,936	12,005	19,981	13,321	-
1999.....	49,748	3	5	3	9	33	156	930	3,983	12,062	19,818	12,745	1
White, both sexes													
2007 1/.....	103,227	9	5	7	10	35	330	2,652	9,631	22,953	39,823	27,769	3
2006 1/.....	100,150	1	4	7	5	32	297	2,478	9,268	22,606	39,176	26,275	1
2005 1/.....	104,399	2	4	3	9	24	353	2,456	9,449	23,609	41,416	27,072	2
2004 1/.....	96,808	5	2	4	8	35	349	2,142	8,658	22,237	38,958	24,409	1
2003 1/.....	99,422	4	2	2	13	29	350	2,085	8,758	23,328	39,913	24,938	-
2002.....	97,029	6	4	5	14	31	347	2,032	8,096	22,869	39,509	24,112	4
2001.....	94,708	4	3	3	13	33	336	1,909	7,898	23,295	38,047	23,167	-
2000.....	93,016	6	3	5	8	18	291	1,809	7,479	23,295	37,796	22,305	1
1999.....	93,677	4	5	7	13	34	256	1,710	7,727	24,066	37,997	21,855	3

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other chronic lower respiratory diseases (J44,J47)													
White, male													
2007 1/.....	48,875	7	2	6	4	17	169	1,382	5,018	11,633	19,255	11,379	3
2006 1/.....	47,195	1	3	4	4	14	147	1,259	4,773	11,214	19,032	10,744	-
2005 1/.....	49,135	2	2	-	7	12	174	1,266	4,817	12,039	19,885	10,930	1
2004 1/.....	45,965	1	1	2	3	24	183	1,137	4,376	11,209	18,912	10,116	1
2003 1/.....	47,212	4	-	1	8	16	168	1,105	4,521	11,855	19,132	10,402	-
2002.....	46,590	4	2	1	6	19	181	1,071	4,101	11,676	19,335	10,191	3
2001.....	45,472	2	1	2	7	20	161	1,019	4,039	11,910	18,693	9,618	-
2000.....	45,316	5	1	3	5	5	145	938	3,899	12,011	18,694	9,609	1
1999.....	46,760	3	1	4	5	14	132	896	4,091	12,805	19,126	9,681	2
White, female													
2007 1/.....	54,352	2	3	1	6	18	161	1,270	4,613	11,320	20,568	16,390	-
2006 1/.....	52,955	-	1	3	1	18	150	1,219	4,495	11,392	20,144	15,531	1
2005 1/.....	55,264	-	2	3	2	12	179	1,190	4,632	11,570	21,531	16,142	1
2004 1/.....	50,843	4	1	2	5	11	166	1,005	4,282	11,028	20,046	14,293	-
2003 1/.....	52,210	-	2	1	5	13	182	980	4,237	11,473	20,781	14,536	-
2002.....	50,439	2	2	4	8	12	166	961	3,995	11,193	20,174	13,921	1
2001.....	49,236	2	2	1	6	13	175	890	3,859	11,385	19,354	13,549	-
2000.....	47,700	1	2	2	3	13	146	871	3,580	11,284	19,102	12,696	-
1999.....	46,917	1	4	3	8	20	124	814	3,636	11,261	18,871	12,174	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other chronic lower respiratory diseases (J44,J47)													
Black, both sexes													
2007 1/.....	6,210	3	-	1	4	11	63	397	900	1,601	2,032	1,198	-
2006 1/.....	6,000	-	-	5	1	15	65	361	898	1,529	2,001	1,125	-
2005 1/.....	6,344	-	2	1	3	10	66	419	894	1,663	2,065	1,220	1
2004 1/.....	5,608	2	3	6	2	8	64	328	803	1,481	1,897	1,014	-
2003 1/.....	5,794	-	1	-	6	9	71	302	827	1,587	1,947	1,044	-
2002.....	5,819	4	1	3	1	16	71	312	784	1,510	2,038	1,078	1
2001.....	5,607	2	1	1	8	9	76	288	713	1,586	1,878	1,045	-
2000.....	5,468	1	-	-	3	10	73	280	754	1,564	1,790	993	-
1999.....	5,751	6	3	1	6	11	69	261	775	1,658	1,976	984	1
Black, male													
2007 1/.....	3,320	3	-	1	2	4	37	190	518	896	1,128	541	-
2006 1/.....	3,252	-	-	1	1	8	28	192	521	885	1,097	519	-
2005 1/.....	3,511	-	-	-	2	7	32	236	511	974	1,175	573	1
2004 1/.....	3,096	1	1	4	1	5	33	175	489	847	1,044	496	-
2003 1/.....	3,274	-	1	-	5	6	35	164	477	933	1,154	499	-
2002.....	3,268	1	-	2	1	5	37	179	469	878	1,168	527	1
2001.....	3,209	1	1	1	6	4	38	154	418	949	1,123	514	-
2000.....	3,163	-	-	-	3	6	34	163	437	962	1,058	500	-
1999.....	3,377	4	2	1	5	-	39	153	473	989	1,192	518	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other chronic lower respiratory diseases (J44,J47)													
Black, female													
2007 1/.....	2,890	-	-	-	2	7	26	207	382	705	904	657	-
2006 1/.....	2,748	-	-	4	-	7	37	169	377	644	904	606	-
2005 1/.....	2,833	-	2	1	1	3	34	183	383	689	890	647	-
2004 1/.....	2,512	1	2	2	1	3	31	153	314	634	853	518	-
2003 1/.....	2,520	-	-	-	1	3	36	138	350	654	793	545	-
2002.....	2,551	3	1	1	-	11	34	133	315	632	870	551	-
2001.....	2,398	1	-	-	2	5	38	134	295	637	755	531	-
2000.....	2,305	1	-	-	-	4	39	117	317	602	732	493	-
1999.....	2,374	2	1	-	1	11	30	108	302	669	784	466	-
American Indian 2/, both sexes													
2007 1/.....	526	-	1	-	-	-	5	37	83	156	171	73	-
2006 1/.....	454	2	-	-	-	-	7	28	81	120	153	63	-
2005 1/.....	428	-	-	-	-	1	4	23	75	122	143	60	-
2004 1/.....	419	1	-	-	-	-	4	21	74	122	148	49	-
2003 1/.....	423	-	-	-	1	-	9	22	75	118	136	62	-
2002.....	379	-	1	-	-	1	1	20	58	114	122	62	-
2001.....	340	-	-	-	2	-	3	13	55	99	107	60	1
2000.....	337	-	-	-	1	-	2	14	53	99	106	62	-
1999.....	325	-	-	-	-	-	3	11	49	100	121	41	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other chronic lower respiratory diseases (J44,J47)

American Indian 2/, male

2007 1/.....	261	-	1	-	-	-	3	17	42	85	82	31	-
2006 1/.....	211	2	-	-	-	-	6	12	42	56	65	28	-
2005 1/.....	215	-	-	-	-	1	2	13	37	54	78	30	-
2004 1/.....	203	1	-	-	-	-	2	10	38	61	69	22	-
2003 1/.....	222	-	-	-	1	-	4	13	37	63	74	30	-
2002.....	180	-	1	-	-	1	-	11	27	53	63	24	-
2001.....	159	-	-	-	1	-	1	8	23	43	55	27	1
2000.....	183	-	-	-	1	-	1	10	33	43	66	29	-
1999.....	160	-	-	-	-	-	1	8	22	56	55	18	-

American Indian 2/, female

2007 1/.....	265	-	-	-	-	-	2	20	41	71	89	42	-
2006 1/.....	243	-	-	-	-	-	1	16	39	64	88	35	-
2005 1/.....	213	-	-	-	-	-	2	10	38	68	65	30	-
2004 1/.....	216	-	-	-	-	-	2	11	36	61	79	27	-
2003 1/.....	201	-	-	-	-	-	5	9	38	55	62	32	-
2002.....	199	-	-	-	-	-	1	9	31	61	59	38	-
2001.....	181	-	-	-	1	-	2	5	32	56	52	33	-
2000.....	154	-	-	-	-	-	1	4	20	56	40	33	-
1999.....	165	-	-	-	-	-	2	3	27	44	66	23	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other chronic lower respiratory diseases (J44,J47)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	1,057	-	-	-	-	1	4	19	60	160	430	383	-
2006 1/.....	1,075	-	1	-	-	-	3	10	64	176	451	369	1
2005 1/.....	1,010	-	-	-	-	2	6	13	59	173	395	362	-
2004 1/.....	924	-	-	-	-	-	2	10	49	156	369	338	-
2003 1/.....	933	-	1	-	2	2	5	11	53	187	372	300	-
2002.....	884	-	-	-	-	2	5	13	49	189	365	261	-
2001.....	888	1	-	-	1	2	3	21	57	200	330	273	-
2000.....	834	-	-	-	1	-	4	22	39	187	337	243	1
1999.....	812	-	-	-	-	3	-	15	47	218	322	207	-

Asian or Pacific Islander 3/, male

2007 1/.....	634	-	-	-	-	1	3	9	36	107	259	219	-
2006 1/.....	689	-	-	-	-	-	2	4	44	114	298	226	1
2005 1/.....	634	-	-	-	-	1	3	9	46	106	256	213	-
2004 1/.....	583	-	-	-	-	-	1	5	31	103	241	202	-
2003 1/.....	613	-	1	-	1	1	4	6	36	122	261	181	-
2002.....	591	-	-	-	-	1	3	7	31	125	255	169	-
2001.....	567	1	-	-	1	-	-	15	38	125	225	162	-
2000.....	536	-	-	-	-	-	1	16	20	124	230	144	1
1999.....	520	-	-	-	-	1	-	10	29	130	225	125	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other chronic lower respiratory diseases (J44,J47)

Asian or Pacific Islander 3/, female

2007 1/.....	423	-	-	-	-	-	1	10	24	53	171	164	-
2006 1/.....	386	-	1	-	-	-	1	6	20	62	153	143	-
2005 1/.....	376	-	-	-	-	1	3	4	13	67	139	149	-
2004 1/.....	341	-	-	-	-	-	1	5	18	53	128	136	-
2003 1/.....	320	-	-	-	1	1	1	5	17	65	111	119	-
2002.....	293	-	-	-	-	1	2	6	18	64	110	92	-
2001.....	321	-	-	-	-	2	3	6	19	75	105	111	-
2000.....	298	-	-	-	1	-	3	6	19	63	107	99	-
1999.....	292	-	-	-	-	2	-	5	18	88	97	82	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Pneumoconioses and chemical effects (J60-J66,J68)

All races, both sexes

2007.....	915	-	-	-	-	3	2	20	59	175	384	272	-
2006.....	924	2	1	-	1	1	3	20	50	183	392	271	-
2005.....	1,007	-	1	1	-	-	3	22	69	175	447	288	1
2004.....	1,033	-	-	-	1	-	4	17	70	209	442	290	-
2003.....	1,114	-	-	-	1	1	7	14	70	233	487	301	-
2002.....	1,114	2	-	-	-	-	5	17	75	223	480	312	-
2001.....	1,136	-	1	-	-	1	5	11	64	266	516	272	-
2000.....	1,161	-	-	-	-	3	7	12	57	281	484	317	-
1999.....	1,113	-	3	-	1	-	7	15	57	274	503	253	-

All races, male

2007.....	875	-	-	-	-	3	2	18	53	169	372	258	-
2006.....	887	2	1	-	-	1	2	16	50	174	383	258	-
2005.....	965	-	-	1	-	-	3	19	65	173	428	275	1
2004.....	993	-	-	-	1	-	4	14	68	204	425	277	-
2003.....	1,065	-	-	-	1	1	6	13	67	222	469	286	-
2002.....	1,057	-	-	-	-	-	4	13	69	214	457	300	-
2001.....	1,095	-	1	-	-	1	5	9	62	261	494	262	-
2000.....	1,113	-	-	-	-	3	6	12	54	265	467	306	-
1999.....	1,079	-	3	-	1	-	5	15	57	269	486	243	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)													
All races, female													
2007.....	40	-	-	-	-	-	-	2	6	6	12	14	-
2006.....	37	-	-	-	1	-	1	4	-	9	9	13	-
2005.....	42	-	1	-	-	-	-	3	4	2	19	13	-
2004.....	40	-	-	-	-	-	-	3	2	5	17	13	-
2003.....	49	-	-	-	-	-	1	1	3	11	18	15	-
2002.....	57	2	-	-	-	-	1	4	6	9	23	12	-
2001.....	41	-	-	-	-	-	-	2	2	5	22	10	-
2000.....	48	-	-	-	-	-	1	-	3	16	17	11	-
1999.....	34	-	-	-	-	-	2	-	-	5	17	10	-
White, both sexes													
2007 1/.....	868	-	-	-	-	2	1	16	55	167	360	267	-
2006 1/.....	868	1	1	-	-	1	3	16	43	173	370	260	-
2005 1/.....	953	-	1	1	-	-	2	20	62	169	424	273	1
2004 1/.....	977	-	-	-	1	-	4	16	65	194	426	271	-
2003 1/.....	1,058	-	-	-	1	-	5	12	67	221	460	292	-
2002.....	1,048	1	-	-	-	-	4	14	67	213	451	298	-
2001.....	1,073	-	1	-	-	1	5	7	55	252	493	259	-
2000.....	1,099	-	-	-	-	3	4	11	55	266	456	304	-
1999.....	1,037	-	-	-	-	-	4	11	47	255	484	236	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Pneumoconioses and chemical effects (J60-J66,J68)

White, male

2007 1/.....	832	-	-	-	-	2	1	14	50	161	351	253	-
2006 1/.....	835	1	1	-	-	1	2	12	43	164	363	248	-
2005 1/.....	915	-	-	1	-	-	2	17	60	167	407	260	1
2004 1/.....	938	-	-	-	1	-	4	13	63	189	409	259	-
2003 1/.....	1,012	-	-	-	1	-	5	11	64	210	443	278	-
2002.....	998	-	-	-	-	-	3	10	64	204	430	287	-
2001.....	1,034	-	1	-	-	1	5	7	53	247	471	249	-
2000.....	1,053	-	-	-	-	3	3	11	52	250	441	293	-
1999.....	1,006	-	-	-	-	-	3	11	47	251	467	227	-

White, female

2007 1/.....	36	-	-	-	-	-	-	2	5	6	9	14	-
2006 1/.....	33	-	-	-	-	-	1	4	-	9	7	12	-
2005 1/.....	38	-	1	-	-	-	-	3	2	2	17	13	-
2004 1/.....	39	-	-	-	-	-	-	3	2	5	17	12	-
2003 1/.....	46	-	-	-	-	-	-	1	3	11	17	14	-
2002.....	50	1	-	-	-	-	1	4	3	9	21	11	-
2001.....	39	-	-	-	-	-	-	-	2	5	22	10	-
2000.....	46	-	-	-	-	-	1	-	3	16	15	11	-
1999.....	31	-	-	-	-	-	1	-	-	4	17	9	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Pneumoconioses and chemical effects (J60-J66,J68)

Black, both sexes

2007 1/.....	45	-	-	-	-	1	1	4	4	8	22	5	-
2006 1/.....	49	1	-	-	1	-	-	4	6	8	19	10	-
2005 1/.....	47	-	-	-	-	-	1	2	6	5	19	14	-
2004 1/.....	54	-	-	-	-	-	-	1	5	14	16	18	-
2003 1/.....	48	-	-	-	-	1	1	2	1	11	23	9	-
2002.....	58	1	-	-	-	-	1	3	6	10	23	14	-
2001.....	57	-	-	-	-	-	-	4	8	12	21	12	-
2000.....	55	-	-	-	-	-	3	1	2	15	23	11	-
1999.....	67	-	3	-	1	-	3	4	8	16	16	16	-

Black, male

2007 1/.....	42	-	-	-	-	1	1	4	3	8	20	5	-
2006 1/.....	46	1	-	-	-	-	-	4	6	8	18	9	-
2005 1/.....	43	-	-	-	-	-	1	2	4	5	17	14	-
2004 1/.....	54	-	-	-	-	-	-	1	5	14	16	18	-
2003 1/.....	45	-	-	-	-	1	-	2	1	11	22	8	-
2002.....	52	-	-	-	-	-	1	3	4	10	21	13	-
2001.....	55	-	-	-	-	-	-	2	8	12	21	12	-
2000.....	53	-	-	-	-	-	3	1	2	15	21	11	-
1999.....	64	-	3	-	1	-	2	4	8	15	16	15	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Pneumoconioses and chemical effects (J60-J66,J68)

Black, female

2007 1/.....	3	-	-	-	-	-	-	-	1	-	2	-	-
2006 1/.....	3	-	-	-	1	-	-	-	-	-	1	1	-
2005 1/.....	4	-	-	-	-	-	-	-	2	-	2	-	-
2004 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 1/.....	3	-	-	-	-	-	1	-	-	-	1	1	-
2002.....	6	1	-	-	-	-	-	-	2	-	2	1	-
2001.....	2	-	-	-	-	-	2	-	-	-	-	-	-
2000.....	2	-	-	-	-	-	-	-	-	-	2	-	-
1999.....	3	-	-	-	-	-	1	-	-	1	-	1	-

American Indian 2/, both sexes

2007 1/.....	1	-	-	-	-	-	-	-	-	-	1	-	-
2006 1/.....	6	-	-	-	-	-	-	-	1	2	2	1	-
2005 1/.....	2	-	-	-	-	-	-	-	-	1	1	-	-
2004 1/.....	1	-	-	-	-	-	-	-	-	-	-	1	-
2003 1/.....	4	-	-	-	-	-	-	-	-	1	3	-	-
2002.....	5	-	-	-	-	-	-	-	1	-	4	-	-
2001.....	4	-	-	-	-	-	-	-	1	1	2	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	1	-
1999.....	7	-	-	-	-	-	-	-	2	1	3	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Pneumoconioses and chemical effects (J60-J66,J68)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	1	-	-	-	-	-	-	-	-	-	1	-	-
2006 1/.....	1	-	-	-	-	-	-	-	-	-	1	-	-
2005 1/.....	5	-	-	-	-	-	-	1	-	-	3	1	-
2004 1/.....	1	-	-	-	-	-	-	-	-	1	-	-	-
2003 1/.....	4	-	-	-	-	-	1	-	2	-	1	-	-
2002.....	3	-	-	-	-	-	-	-	1	-	2	-	-
2001.....	2	-	-	-	-	-	-	-	-	1	-	1	-
2000.....	6	-	-	-	-	-	-	-	-	-	5	1	-
1999.....	2	-	-	-	-	-	-	-	-	2	-	-	-

Asian or Pacific Islander 3/, male

2007 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2006 1/.....	1	-	-	-	-	-	-	-	-	-	1	-	-
2005 1/.....	5	-	-	-	-	-	-	-	1	-	3	1	-
2004 1/.....	1	-	-	-	-	-	-	-	-	1	-	-	-
2003 1/.....	4	-	-	-	-	-	1	-	2	-	1	-	-
2002.....	2	-	-	-	-	-	-	-	-	-	2	-	-
2001.....	2	-	-	-	-	-	-	-	-	1	-	1	-
2000.....	6	-	-	-	-	-	-	-	-	-	5	1	-
1999.....	2	-	-	-	-	-	-	-	-	2	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Pneumonitis due to solids and liquids (J69)

All races, both sexes

2007.....	16,988	10	8	16	47	70	154	436	884	1,724	5,187	8,451	1
2006.....	16,887	11	9	13	30	78	166	437	855	1,750	5,284	8,254	-
2005.....	17,279	17	8	13	43	66	172	476	833	1,741	5,558	8,352	-
2004.....	16,780	12	9	11	39	68	159	394	743	1,767	5,564	8,014	-
2003.....	17,335	13	10	16	46	69	166	392	773	1,754	5,789	8,307	-
2002.....	17,593	2	7	16	44	74	192	351	671	1,793	5,900	8,543	-
2001.....	17,301	12	14	15	34	79	179	378	659	1,830	5,759	8,342	-
2000.....	16,636	10	11	13	37	62	181	366	598	1,899	5,505	7,954	-
1999.....	15,268	16	12	16	32	78	166	284	586	1,754	5,030	7,294	-

All races, male

2007.....	8,934	5	5	12	31	43	98	278	549	1,042	2,926	3,944	1
2006.....	8,916	5	3	11	15	53	103	272	537	1,092	3,029	3,796	-
2005.....	8,760	10	3	8	23	43	88	280	506	1,065	3,057	3,677	-
2004.....	8,593	4	5	6	20	40	97	250	451	1,085	3,059	3,576	-
2003.....	8,702	8	5	9	31	42	89	239	471	1,069	3,181	3,558	-
2002.....	8,876	-	3	9	27	42	119	226	393	1,083	3,243	3,731	-
2001.....	8,576	6	6	7	22	40	117	232	384	1,106	3,145	3,511	-
2000.....	8,390	4	5	9	23	35	115	225	354	1,195	3,038	3,387	-
1999.....	7,755	9	3	6	22	47	97	165	365	1,083	2,875	3,083	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Pneumonitis due to solids and liquids (J69)

All races, female

2007.....	8,054	5	3	4	16	27	56	158	335	682	2,261	4,507	-
2006.....	7,971	6	6	2	15	25	63	165	318	658	2,255	4,458	-
2005.....	8,519	7	5	5	20	23	84	196	327	676	2,501	4,675	-
2004.....	8,187	8	4	5	19	28	62	144	292	682	2,505	4,438	-
2003.....	8,633	5	5	7	15	27	77	153	302	685	2,608	4,749	-
2002.....	8,717	2	4	7	17	32	73	125	278	710	2,657	4,812	-
2001.....	8,725	6	8	8	12	39	62	146	275	724	2,614	4,831	-
2000.....	8,246	6	6	4	14	27	66	141	244	704	2,467	4,567	-
1999.....	7,513	7	9	10	10	31	69	119	221	671	2,155	4,211	-

White, both sexes

2007 1/.....	15,229	7	7	12	35	52	125	355	713	1,483	4,647	7,792	1
2006 1/.....	15,136	7	6	10	19	57	126	359	721	1,464	4,751	7,616	-
2005 1/.....	15,338	13	3	9	32	53	143	382	681	1,460	4,932	7,630	-
2004 1/.....	14,968	5	7	9	28	52	122	314	588	1,497	4,994	7,352	-
2003 1/.....	15,395	7	8	13	29	55	124	326	615	1,478	5,135	7,605	-
2002.....	15,773	2	5	14	34	59	147	276	554	1,506	5,308	7,868	-
2001.....	15,425	9	12	11	25	62	142	298	550	1,541	5,172	7,603	-
2000.....	14,869	8	9	12	27	46	142	281	485	1,580	4,948	7,331	-
1999.....	13,582	13	8	12	22	57	121	219	473	1,461	4,492	6,704	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)													
White, male													
2007 1/.....	8,010	3	5	9	24	30	80	233	440	897	2,624	3,664	1
2006 1/.....	7,984	2	1	8	10	38	79	227	453	916	2,726	3,524	-
2005 1/.....	7,759	8	1	7	17	32	75	221	413	890	2,711	3,384	-
2004 1/.....	7,644	2	4	5	15	32	77	204	348	912	2,755	3,290	-
2003 1/.....	7,731	4	5	7	21	33	71	201	375	905	2,832	3,277	-
2002.....	7,954	-	3	8	20	35	90	176	327	911	2,931	3,453	-
2001.....	7,705	6	4	5	18	35	94	189	321	945	2,849	3,239	-
2000.....	7,488	4	4	8	18	22	93	171	291	982	2,746	3,149	-
1999.....	6,853	7	2	5	14	33	71	128	291	895	2,562	2,845	-
White, female													
2007 1/.....	7,219	4	2	3	11	22	45	122	273	586	2,023	4,128	-
2006 1/.....	7,152	5	5	2	9	19	47	132	268	548	2,025	4,092	-
2005 1/.....	7,579	5	2	2	15	21	68	161	268	570	2,221	4,246	-
2004 1/.....	7,324	3	3	4	13	20	45	110	240	585	2,239	4,062	-
2003 1/.....	7,664	3	3	6	8	22	53	125	240	573	2,303	4,328	-
2002.....	7,819	2	2	6	14	24	57	100	227	595	2,377	4,415	-
2001.....	7,720	3	8	6	7	27	48	109	229	596	2,323	4,364	-
2000.....	7,381	4	5	4	9	24	49	110	194	598	2,202	4,182	-
1999.....	6,729	6	6	7	8	24	50	91	182	566	1,930	3,859	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)													
Black, both sexes													
2007 1/.....	1,429	3	1	3	7	16	25	71	158	205	438	502	-
2006 1/.....	1,439	4	-	3	9	20	37	73	119	250	429	495	-
2005 1/.....	1,620	3	5	3	10	12	26	87	136	235	530	573	-
2004 1/.....	1,495	6	2	1	11	14	35	65	140	241	467	513	-
2003 1/.....	1,607	4	1	3	14	12	41	61	142	235	534	560	-
2002.....	1,537	-	2	2	8	14	38	65	99	250	503	556	-
2001.....	1,599	3	2	4	7	16	34	73	101	252	503	604	-
2000.....	1,521	1	2	1	9	15	38	80	102	290	472	511	-
1999.....	1,505	2	4	3	10	18	41	59	105	266	473	524	-
Black, male													
2007 1/.....	739	2	-	2	4	13	16	40	100	129	244	189	-
2006 1/.....	757	3	-	3	4	14	22	40	75	155	247	194	-
2005 1/.....	825	2	2	1	5	11	12	56	83	146	290	217	-
2004 1/.....	766	1	1	-	5	8	20	39	92	153	243	204	-
2003 1/.....	781	3	-	2	8	9	17	36	87	139	276	204	-
2002.....	760	-	-	1	5	6	23	42	57	152	256	218	-
2001.....	741	-	2	2	2	4	20	39	60	141	256	215	-
2000.....	765	-	1	1	4	12	21	53	55	190	246	182	-
1999.....	775	1	1	1	8	12	23	33	68	170	263	195	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Pneumonitis due to solids and liquids (J69)

Black, female

2007 1/.....	690	1	1	1	3	3	9	31	58	76	194	313	-
2006 1/.....	682	1	-	-	5	6	15	33	44	95	182	301	-
2005 1/.....	795	1	3	2	5	1	14	31	53	89	240	356	-
2004 1/.....	729	5	1	1	6	6	15	26	48	88	224	309	-
2003 1/.....	826	1	1	1	6	3	24	25	55	96	258	356	-
2002.....	777	-	2	1	3	8	15	23	42	98	247	338	-
2001.....	858	3	-	2	5	12	14	34	41	111	247	389	-
2000.....	756	1	1	-	5	3	17	27	47	100	226	329	-
1999.....	730	1	3	2	2	6	18	26	37	96	210	329	-

American Indian 2/, both sexes

2007 1/.....	76	-	-	-	2	2	3	5	5	13	21	25	-
2006 1/.....	68	-	-	-	1	1	2	3	7	15	16	23	-
2005 1/.....	65	1	-	-	-	1	1	4	5	15	17	21	-
2004 1/.....	64	1	-	1	-	-	1	9	5	9	21	17	-
2003 1/.....	70	2	1	-	-	1	-	1	7	13	22	23	-
2002.....	73	-	-	-	1	1	4	6	8	8	23	22	-
2001.....	59	-	-	-	1	1	2	5	1	10	21	18	-
2000.....	49	-	-	-	-	1	-	3	7	9	14	15	-
1999.....	46	-	-	-	-	3	2	2	5	9	8	17	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Pneumonitis due to solids and liquids (J69)

American Indian 2/, male

2007 1/.....	35	-	-	-	1	-	2	2	2	3	10	15	-
2006 1/.....	39	-	-	-	-	1	1	3	4	9	8	13	-
2005 1/.....	35	-	-	-	-	-	1	1	4	8	12	9	-
2004 1/.....	31	1	-	1	-	-	-	3	3	4	12	7	-
2003 1/.....	33	1	-	-	-	-	-	-	3	5	11	13	-
2002.....	44	-	-	-	1	1	3	4	4	5	13	13	-
2001.....	23	-	-	-	1	1	2	2	1	4	8	4	-
2000.....	27	-	-	-	-	1	-	-	4	7	8	7	-
1999.....	28	-	-	-	-	2	1	2	4	4	5	10	-

American Indian 2/, female

2007 1/.....	41	-	-	-	1	2	1	3	3	10	11	10	-
2006 1/.....	29	-	-	-	1	-	1	-	3	6	8	10	-
2005 1/.....	30	1	-	-	-	1	-	3	1	7	5	12	-
2004 1/.....	33	-	-	-	-	-	1	6	2	5	9	10	-
2003 1/.....	37	1	1	-	-	1	-	1	4	8	11	10	-
2002.....	29	-	-	-	-	-	1	2	4	3	10	9	-
2001.....	36	-	-	-	-	-	-	3	-	6	13	14	-
2000.....	22	-	-	-	-	-	-	3	3	2	6	8	-
1999.....	18	-	-	-	-	1	1	-	1	5	3	7	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Pneumonitis due to solids and liquids (J69)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	254	-	-	1	3	-	1	5	8	23	81	132	-
2006 1/.....	244	-	3	-	1	-	1	2	8	21	88	120	-
2005 1/.....	256	-	-	1	1	-	2	3	11	31	79	128	-
2004 1/.....	253	-	-	-	-	2	1	6	10	20	82	132	-
2003 1/.....	263	-	-	-	3	1	1	4	9	28	98	119	-
2002.....	210	-	-	-	1	-	3	4	10	29	66	97	-
2001.....	218	-	-	-	1	-	1	2	7	27	63	117	-
2000.....	197	1	-	-	1	-	1	2	4	20	71	97	-
1999.....	135	1	-	1	-	-	2	4	3	18	57	49	-

Asian or Pacific Islander 3/, male

2007 1/.....	150	-	-	1	2	-	-	3	7	13	48	76	-
2006 1/.....	136	-	2	-	1	-	1	2	5	12	48	65	-
2005 1/.....	141	-	-	-	1	-	-	2	6	21	44	67	-
2004 1/.....	152	-	-	-	-	-	-	4	8	16	49	75	-
2003 1/.....	157	-	-	-	2	-	1	2	6	20	62	64	-
2002.....	118	-	-	-	1	-	3	4	5	15	43	47	-
2001.....	107	-	-	-	1	-	1	2	2	16	32	53	-
2000.....	110	-	-	-	1	-	1	1	4	16	38	49	-
1999.....	99	1	-	-	-	-	2	2	2	14	45	33	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Pneumonitis due to solids and liquids (J69)

Asian or Pacific Islander 3/, female

2007 1/.....	104	-	-	-	1	-	1	2	1	10	33	56	-
2006 1/.....	108	-	1	-	-	-	-	-	3	9	40	55	-
2005 1/.....	115	-	-	1	-	-	2	1	5	10	35	61	-
2004 1/.....	101	-	-	-	-	2	1	2	2	4	33	57	-
2003 1/.....	106	-	-	-	1	1	-	2	3	8	36	55	-
2002.....	92	-	-	-	-	-	-	-	5	14	23	50	-
2001.....	111	-	-	-	-	-	-	-	5	11	31	64	-
2000.....	87	1	-	-	-	-	-	1	-	4	33	48	-
1999.....	36	-	-	1	-	-	-	2	1	4	12	16	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

All races, both sexes

2007.....	28,508	319	100	66	152	216	493	1,376	3,038	5,556	9,535	7,657	-
2006.....	27,644	334	97	70	103	204	523	1,354	2,805	5,344	9,572	7,238	-
2005.....	27,056	303	94	61	110	181	474	1,360	2,676	5,373	9,466	6,955	3
2004.....	27,310	266	105	62	123	174	537	1,349	2,769	5,523	9,629	6,773	-
2003.....	25,520	279	86	60	136	210	495	1,335	2,442	5,156	8,810	6,510	1
2002.....	25,039	278	85	84	126	185	528	1,288	2,440	5,237	8,658	6,130	-
2001.....	26,178	294	93	78	141	216	570	1,356	2,432	5,510	9,202	6,283	3
2000.....	25,535	273	87	74	131	253	615	1,332	2,411	5,448	8,900	6,008	3
1999.....	25,031	252	91	76	115	257	576	1,219	2,260	5,409	8,646	6,130	-

All races, male

2007.....	14,291	190	66	31	90	116	252	789	1,703	3,122	4,925	3,007	-
2006.....	13,728	195	61	38	66	96	269	699	1,573	3,033	4,934	2,764	-
2005.....	13,168	187	57	33	60	91	226	737	1,486	2,974	4,787	2,529	1
2004.....	13,434	138	59	35	61	90	269	723	1,509	3,106	4,898	2,546	-
2003.....	12,391	163	49	38	76	115	251	704	1,367	2,849	4,395	2,384	-
2002.....	12,057	164	56	39	68	88	281	656	1,339	2,929	4,261	2,176	-
2001.....	12,495	186	47	43	76	104	304	737	1,322	3,025	4,453	2,197	1
2000.....	12,391	154	45	38	73	127	335	712	1,324	2,961	4,415	2,205	2
1999.....	12,110	145	55	39	69	127	317	637	1,213	3,005	4,328	2,175	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

All races, female

2007.....	14,217	129	34	35	62	100	241	587	1,335	2,434	4,610	4,650	-
2006.....	13,916	139	36	32	37	108	254	655	1,232	2,311	4,638	4,474	-
2005.....	13,888	116	37	28	50	90	248	623	1,190	2,399	4,679	4,426	2
2004.....	13,876	128	46	27	62	84	268	626	1,260	2,417	4,731	4,227	-
2003.....	13,129	116	37	22	60	95	244	631	1,075	2,307	4,415	4,126	1
2002.....	12,982	114	29	45	58	97	247	632	1,101	2,308	4,397	3,954	-
2001.....	13,683	108	46	35	65	112	266	619	1,110	2,485	4,749	4,086	2
2000.....	13,144	119	42	36	58	126	280	620	1,087	2,487	4,485	3,803	1
1999.....	12,921	107	36	37	46	130	259	582	1,047	2,404	4,318	3,955	-

White, both sexes

2007 1/.....	25,155	189	65	46	109	151	354	1,027	2,499	4,846	8,760	7,109	-
2006 1/.....	24,463	182	66	50	70	158	377	1,009	2,299	4,726	8,800	6,726	-
2005 1/.....	23,860	191	66	49	77	125	343	1,000	2,207	4,706	8,668	6,426	2
2004 1/.....	24,258	156	66	41	93	118	376	999	2,312	4,904	8,879	6,314	-
2003 1/.....	22,706	169	53	46	88	147	334	992	2,040	4,605	8,137	6,094	1
2002.....	22,184	151	56	57	81	132	379	962	2,016	4,647	7,978	5,725	-
2001.....	23,133	156	61	59	100	139	388	991	2,022	4,891	8,507	5,818	1
2000.....	22,664	166	63	53	86	182	443	1,009	2,017	4,836	8,208	5,600	1
1999.....	22,161	147	62	55	90	178	395	922	1,883	4,787	7,968	5,674	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

White, male

2007 1/.....	12,638	116	39	18	68	83	183	593	1,411	2,755	4,564	2,808	-
2006 1/.....	12,183	99	42	26	43	72	195	541	1,327	2,705	4,571	2,562	-
2005 1/.....	11,574	114	38	25	43	67	170	551	1,221	2,649	4,383	2,312	1
2004 1/.....	11,944	80	39	22	48	61	182	548	1,275	2,780	4,536	2,373	-
2003 1/.....	11,014	92	28	29	46	83	181	547	1,151	2,564	4,065	2,228	-
2002.....	10,693	87	39	27	48	63	201	496	1,116	2,634	3,945	2,037	-
2001.....	11,085	95	28	34	55	64	227	556	1,111	2,731	4,143	2,041	-
2000.....	10,979	100	30	28	46	93	241	540	1,118	2,650	4,088	2,044	1
1999.....	10,764	85	34	29	58	98	228	489	1,014	2,706	4,019	2,004	-

White, female

2007 1/.....	12,517	73	26	28	41	68	171	434	1,088	2,091	4,196	4,301	-
2006 1/.....	12,280	83	24	24	27	86	182	468	972	2,021	4,229	4,164	-
2005 1/.....	12,286	77	28	24	34	58	173	449	986	2,057	4,285	4,114	1
2004 1/.....	12,314	76	27	19	45	57	194	451	1,037	2,124	4,343	3,941	-
2003 1/.....	11,692	77	25	17	42	64	153	445	889	2,041	4,072	3,866	1
2002.....	11,491	64	17	30	33	69	178	466	900	2,013	4,033	3,688	-
2001.....	12,048	61	33	25	45	75	161	435	911	2,160	4,364	3,777	1
2000.....	11,685	66	33	25	40	89	202	469	899	2,186	4,120	3,556	-
1999.....	11,397	62	28	26	32	80	167	433	869	2,081	3,949	3,670	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Black, both sexes

2007 1/.....	2,591	112	30	18	38	58	117	308	450	535	544	381	-
2006 1/.....	2,442	132	22	17	30	43	125	304	418	459	525	367	-
2005 1/.....	2,481	102	23	11	28	50	121	312	371	512	566	384	1
2004 1/.....	2,397	98	30	17	23	49	150	308	388	476	529	329	-
2003 1/.....	2,233	99	27	8	41	56	141	306	336	413	499	307	-
2002.....	2,279	112	29	25	41	47	134	284	358	448	497	304	-
2001.....	2,491	117	26	17	38	68	166	327	345	492	536	357	2
2000.....	2,361	103	21	18	40	59	153	288	327	504	532	314	2
1999.....	2,356	93	26	19	21	74	163	259	319	507	525	350	-

Black, male

2007 1/.....	1,261	65	23	12	20	26	60	170	243	262	246	134	-
2006 1/.....	1,168	82	15	10	21	21	62	135	201	246	245	130	-
2005 1/.....	1,209	68	15	7	15	21	53	157	209	248	274	142	-
2004 1/.....	1,162	53	15	11	11	27	82	153	197	249	248	116	-
2003 1/.....	1,097	62	18	6	27	28	60	139	186	219	243	109	-
2002.....	1,084	67	17	10	19	23	73	140	190	222	229	94	-
2001.....	1,131	77	14	8	19	34	70	158	173	227	236	114	1
2000.....	1,135	51	14	10	23	30	85	153	166	246	241	115	1
1999.....	1,092	54	19	9	9	27	77	133	165	239	236	124	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Black, female

2007 1/.....	1,330	47	7	6	18	32	57	138	207	273	298	247	-
2006 1/.....	1,274	50	7	7	9	22	63	169	217	213	280	237	-
2005 1/.....	1,272	34	8	4	13	29	68	155	162	264	292	242	1
2004 1/.....	1,235	45	15	6	12	22	68	155	191	227	281	213	-
2003 1/.....	1,136	37	9	2	14	28	81	167	150	194	256	198	-
2002.....	1,195	45	12	15	22	24	61	144	168	226	268	210	-
2001.....	1,360	40	12	9	19	34	96	169	172	265	300	243	1
2000.....	1,226	52	7	8	17	29	68	135	161	258	291	199	1
1999.....	1,264	39	7	10	12	47	86	126	154	268	289	226	-

American Indian 2/, both sexes

2007 1/.....	214	3	3	1	1	-	10	18	35	61	45	37	-
2006 1/.....	225	9	4	2	1	1	8	20	34	53	60	33	-
2005 1/.....	185	3	3	-	-	3	6	14	37	46	43	30	-
2004 1/.....	182	6	2	1	5	1	5	23	22	46	43	28	-
2003 1/.....	166	2	-	2	2	-	7	18	26	40	43	26	-
2002.....	146	5	-	-	2	3	6	18	22	33	29	28	-
2001.....	177	7	3	-	2	4	9	18	27	37	43	27	-
2000.....	157	1	2	1	1	7	8	17	24	39	37	20	-
1999.....	148	3	1	-	1	3	5	18	14	28	46	29	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

American Indian 2/, male

2007 1/.....	114	2	2	-	1	-	6	12	22	37	18	14	-
2006 1/.....	110	7	2	1	-	1	3	13	13	25	31	14	-
2005 1/.....	100	1	3	-	-	2	3	8	22	26	25	10	-
2004 1/.....	92	2	1	1	2	-	2	15	12	26	20	11	-
2003 1/.....	74	2	-	1	1	-	3	7	13	17	21	9	-
2002.....	73	4	-	-	1	-	2	7	13	15	17	14	-
2001.....	89	5	3	-	1	4	5	9	12	20	22	8	-
2000.....	73	-	1	-	1	2	5	10	14	15	16	9	-
1999.....	71	3	1	-	1	1	2	8	9	15	19	12	-

American Indian 2/, female

2007 1/.....	100	1	1	1	-	-	4	6	13	24	27	23	-
2006 1/.....	115	2	2	1	1	-	5	7	21	28	29	19	-
2005 1/.....	85	2	-	-	-	1	3	6	15	20	18	20	-
2004 1/.....	90	4	1	-	3	1	3	8	10	20	23	17	-
2003 1/.....	92	-	-	1	1	-	4	11	13	23	22	17	-
2002.....	73	1	-	-	1	3	4	11	9	18	12	14	-
2001.....	88	2	-	-	1	-	4	9	15	17	21	19	-
2000.....	84	1	1	1	-	5	3	7	10	24	21	11	-
1999.....	77	-	-	-	-	2	3	10	5	13	27	17	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	548	15	2	1	4	7	12	23	54	114	186	130	-
2006 1/.....	514	11	5	1	2	2	13	21	54	106	187	112	-
2005 1/.....	530	7	2	1	5	3	4	34	61	109	189	115	-
2004 1/.....	473	6	7	3	2	6	6	19	47	97	178	102	-
2003 1/.....	415	9	6	4	5	7	13	19	40	98	131	83	-
2002.....	430	10	-	2	2	3	9	24	44	109	154	73	-
2001.....	377	14	3	2	1	5	7	20	38	90	116	81	-
2000.....	353	3	1	2	4	5	11	18	43	69	123	74	-
1999.....	366	9	2	2	3	2	13	20	44	87	107	77	-

Asian or Pacific Islander 3/, male

2007 1/.....	278	7	2	1	1	7	3	14	27	68	97	51	-
2006 1/.....	267	7	2	1	2	2	9	10	32	57	87	58	-
2005 1/.....	285	4	1	1	2	1	-	21	34	51	105	65	-
2004 1/.....	236	3	4	1	-	2	3	7	25	51	94	46	-
2003 1/.....	206	7	3	2	2	4	7	11	17	49	66	38	-
2002.....	207	6	-	2	-	2	5	13	20	58	70	31	-
2001.....	190	9	2	1	1	2	2	14	26	47	52	34	-
2000.....	204	3	-	-	3	2	4	9	26	50	70	37	-
1999.....	183	3	1	1	1	1	10	7	25	45	54	35	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Asian or Pacific Islander 3/, female

2007 1/.....	270	8	-	-	3	-	9	9	27	46	89	79	-
2006 1/.....	247	4	3	-	-	-	4	11	22	49	100	54	-
2005 1/.....	245	3	1	-	3	2	4	13	27	58	84	50	-
2004 1/.....	237	3	3	2	2	4	3	12	22	46	84	56	-
2003 1/.....	209	2	3	2	3	3	6	8	23	49	65	45	-
2002.....	223	4	-	-	2	1	4	11	24	51	84	42	-
2001.....	187	5	1	1	-	3	5	6	12	43	64	47	-
2000.....	149	-	1	2	1	3	7	9	17	19	53	37	-
1999.....	183	6	1	1	2	1	3	13	19	42	53	42	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Peptic ulcer (K25-K28)													
All races, both sexes													
2007.....	3,045	2	-	2	6	20	78	293	378	487	828	951	-
2006.....	3,323	1	-	1	8	18	73	285	383	520	937	1,097	-
2005.....	3,478	2	1	1	8	25	108	281	376	544	1,040	1,092	-
2004.....	3,682	2	3	2	10	17	90	269	374	582	1,148	1,185	-
2003.....	3,913	2	1	3	7	25	101	281	383	587	1,279	1,244	-
2002.....	4,079	1	1	1	-	22	113	272	411	665	1,291	1,302	-
2001.....	4,491	-	1	5	6	32	118	284	418	718	1,438	1,470	1
2000.....	4,558	-	-	-	9	14	113	307	443	748	1,415	1,509	-
1999.....	4,604	2	2	-	2	21	123	255	422	836	1,500	1,439	2
All races, male													
2007.....	1,469	2	-	1	5	13	49	200	231	268	381	319	-
2006.....	1,591	-	-	-	5	10	43	196	250	280	436	371	-
2005.....	1,722	1	-	1	6	17	73	203	244	305	512	360	-
2004.....	1,690	1	2	1	6	14	57	176	228	304	530	371	-
2003.....	1,880	-	1	1	6	13	60	192	237	346	606	418	-
2002.....	1,951	-	1	1	-	14	72	177	247	378	622	439	-
2001.....	2,074	-	1	2	3	21	87	186	273	397	649	454	1
2000.....	2,201	-	-	-	7	6	79	217	284	442	667	499	-
1999.....	2,189	-	-	-	2	12	76	169	265	504	712	448	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Peptic ulcer (K25-K28)													
All races, female													
2007.....	1,576	-	-	1	1	7	29	93	147	219	447	632	-
2006.....	1,732	1	-	1	3	8	30	89	133	240	501	726	-
2005.....	1,756	1	1	-	2	8	35	78	132	239	528	732	-
2004.....	1,992	1	1	1	4	3	33	93	146	278	618	814	-
2003.....	2,033	2	-	2	1	12	41	89	146	241	673	826	-
2002.....	2,128	1	-	-	-	8	41	95	164	287	669	863	-
2001.....	2,417	-	-	3	3	11	31	98	145	321	789	1,016	-
2000.....	2,357	-	-	-	2	8	34	90	159	306	748	1,010	-
1999.....	2,415	2	2	-	-	9	47	86	157	332	788	991	1
White, both sexes													
2007 1/.....	2,628	2	-	2	6	19	63	216	296	417	732	875	-
2006 1/.....	2,857	1	-	-	5	13	49	207	307	429	837	1,009	-
2005 1/.....	2,995	1	-	1	4	17	77	209	314	459	923	990	-
2004 1/.....	3,215	1	1	1	9	10	62	198	309	493	1,030	1,101	-
2003 1/.....	3,408	1	1	2	4	20	72	206	312	514	1,130	1,146	-
2002.....	3,587	1	1	-	-	12	77	212	325	576	1,176	1,207	-
2001.....	3,942	-	1	4	3	16	78	214	331	619	1,303	1,372	1
2000.....	4,021	-	-	-	9	9	81	227	370	647	1,286	1,392	-
1999.....	4,040	1	2	-	2	15	77	186	350	716	1,360	1,330	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Peptic ulcer (K25-K28)													
White, male													
2007 1/.....	1,232	2	-	1	5	12	42	145	178	224	334	289	-
2006 1/.....	1,321	-	-	-	4	6	25	136	203	225	388	334	-
2005 1/.....	1,453	1	-	1	3	12	50	146	204	253	458	325	-
2004 1/.....	1,450	-	1	1	5	8	40	131	187	262	468	347	-
2003 1/.....	1,592	-	1	1	3	10	42	138	191	299	528	379	-
2002.....	1,685	-	1	-	-	10	50	133	194	324	564	409	-
2001.....	1,751	-	1	2	2	11	56	133	213	332	580	420	1
2000.....	1,900	-	-	-	7	4	56	164	240	380	598	451	-
1999.....	1,887	-	-	-	2	8	44	121	215	432	650	414	1
White, female													
2007 1/.....	1,396	-	-	1	1	7	21	71	118	193	398	586	-
2006 1/.....	1,536	1	-	-	1	7	24	71	104	204	449	675	-
2005 1/.....	1,542	-	-	-	1	5	27	63	110	206	465	665	-
2004 1/.....	1,765	1	-	-	4	2	22	67	122	231	562	754	-
2003 1/.....	1,816	1	-	1	1	10	30	68	121	215	602	767	-
2002.....	1,902	1	-	-	-	2	27	79	131	252	612	798	-
2001.....	2,191	-	-	2	1	5	22	81	118	287	723	952	-
2000.....	2,121	-	-	-	2	5	25	63	130	267	688	941	-
1999.....	2,153	1	2	-	-	7	33	65	135	284	710	916	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Peptic ulcer (K25-K28)

Black, both sexes

2007 1/.....	305	-	-	-	-	1	13	67	69	52	57	46	-
2006 1/.....	361	-	-	1	3	4	22	68	66	67	73	57	-
2005 1/.....	363	1	1	-	4	8	28	62	50	69	75	65	-
2004 1/.....	366	-	1	1	1	6	28	63	59	73	77	57	-
2003 1/.....	383	1	-	1	1	5	23	71	61	54	108	58	-
2002.....	380	-	-	1	-	10	35	56	68	69	77	64	-
2001.....	409	-	-	1	2	15	35	59	69	74	86	68	-
2000.....	425	-	-	-	-	4	29	70	65	80	92	85	-
1999.....	448	1	-	-	-	4	44	60	66	93	99	80	1

Black, male

2007 1/.....	174	-	-	-	-	1	5	48	44	33	28	15	-
2006 1/.....	209	-	-	-	1	3	16	53	40	43	34	19	-
2005 1/.....	211	-	-	-	3	5	20	50	36	42	33	22	-
2004 1/.....	191	-	1	-	1	5	17	41	38	33	40	15	-
2003 1/.....	223	-	-	-	1	3	13	51	42	37	57	19	-
2002.....	212	-	-	1	-	4	21	41	45	41	40	19	-
2001.....	243	-	-	-	1	10	27	46	48	48	42	21	-
2000.....	229	-	-	-	-	2	21	45	40	48	46	27	-
1999.....	243	-	-	-	-	3	31	43	47	55	40	24	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Peptic ulcer (K25-K28)

Black, female

2007 1/.....	131	-	-	-	-	-	8	19	25	19	29	31	-
2006 1/.....	152	-	-	1	2	1	6	15	26	24	39	38	-
2005 1/.....	152	1	1	-	1	3	8	12	14	27	42	43	-
2004 1/.....	175	-	-	1	-	1	11	22	21	40	37	42	-
2003 1/.....	160	1	-	1	-	2	10	20	19	17	51	39	-
2002.....	168	-	-	-	-	6	14	15	23	28	37	45	-
2001.....	166	-	-	1	1	5	8	13	21	26	44	47	-
2000.....	196	-	-	-	-	2	8	25	25	32	46	58	-
1999.....	205	1	-	-	-	1	13	17	19	38	59	56	1

American Indian 2/, both sexes

2007 1/.....	19	-	-	-	-	-	-	5	5	2	6	1	-
2006 1/.....	17	-	-	-	-	1	1	2	2	5	5	1	-
2005 1/.....	12	-	-	-	-	-	1	3	1	1	4	2	-
2004 1/.....	13	-	-	-	-	-	-	3	1	4	3	2	-
2003 1/.....	16	-	-	-	1	-	3	2	1	2	5	2	-
2002.....	17	-	-	-	-	-	-	-	7	4	3	3	-
2001.....	28	-	-	-	-	1	2	5	6	7	6	1	-
2000.....	19	-	-	-	-	-	1	5	4	5	3	1	-
1999.....	10	-	-	-	-	-	2	2	-	2	3	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Peptic ulcer (K25-K28)

American Indian 2/, male

2007 1/.....	14	-	-	-	-	-	4	5	1	3	1	-
2006 1/.....	15	-	-	-	-	1	1	2	2	5	1	-
2005 1/.....	6	-	-	-	-	-	1	2	-	1	2	-
2004 1/.....	2	-	-	-	-	-	-	-	-	1	1	-
2003 1/.....	7	-	-	-	1	-	2	1	-	2	1	-
2002.....	10	-	-	-	-	-	-	-	3	3	3	1
2001.....	17	-	-	-	-	-	1	4	3	5	3	1
2000.....	11	-	-	-	-	-	1	3	2	3	2	-
1999.....	5	-	-	-	-	-	1	1	-	1	2	-

American Indian 2/, female

2007 1/.....	5	-	-	-	-	-	-	1	-	1	3	-
2006 1/.....	2	-	-	-	-	-	-	-	-	-	2	-
2005 1/.....	6	-	-	-	-	-	-	1	1	-	2	2
2004 1/.....	11	-	-	-	-	-	-	3	1	3	2	2
2003 1/.....	9	-	-	-	-	-	1	1	1	-	4	2
2002.....	7	-	-	-	-	-	-	-	4	1	-	2
2001.....	11	-	-	-	-	1	1	1	3	2	3	-
2000.....	8	-	-	-	-	-	-	2	2	2	1	1
1999.....	5	-	-	-	-	-	1	1	-	1	1	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Peptic ulcer (K25-K28)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	93	-	-	-	-	-	2	5	8	16	33	29	-
2006 1/.....	88	-	-	-	-	-	1	8	8	19	22	30	-
2005 1/.....	108	-	-	-	-	-	2	7	11	15	38	35	-
2004 1/.....	88	1	1	-	-	1	-	5	5	12	38	25	-
2003 1/.....	106	-	-	-	1	-	3	2	9	17	36	38	-
2002.....	95	-	-	-	-	-	1	4	11	16	35	28	-
2001.....	112	-	-	-	1	-	3	6	12	18	43	29	-
2000.....	93	-	-	-	-	1	2	5	4	16	34	31	-
1999.....	106	-	-	-	-	2	-	7	6	25	38	28	-

Asian or Pacific Islander 3/, male

2007 1/.....	49	-	-	-	-	-	2	3	4	10	16	14	-
2006 1/.....	46	-	-	-	-	-	1	5	5	7	11	17	-
2005 1/.....	52	-	-	-	-	-	2	5	4	9	19	13	-
2004 1/.....	47	1	-	-	-	1	-	4	3	8	21	9	-
2003 1/.....	58	-	-	-	1	-	3	2	4	8	20	20	-
2002.....	44	-	-	-	-	-	1	3	5	10	15	10	-
2001.....	63	-	-	-	-	-	3	3	9	12	24	12	-
2000.....	61	-	-	-	-	-	1	5	2	11	21	21	-
1999.....	54	-	-	-	-	1	-	4	3	16	20	10	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Peptic ulcer (K25-K28)

Asian or Pacific Islander 3/, female

2007 1/.....	44	-	-	-	-	-	-	2	4	6	17	15	-
2006 1/.....	42	-	-	-	-	-	-	3	3	12	11	13	-
2005 1/.....	56	-	-	-	-	-	-	2	7	6	19	22	-
2004 1/.....	41	-	1	-	-	-	-	1	2	4	17	16	-
2003 1/.....	48	-	-	-	-	-	-	-	5	9	16	18	-
2002.....	51	-	-	-	-	-	-	1	6	6	20	18	-
2001.....	49	-	-	-	1	-	-	3	3	6	19	17	-
2000.....	32	-	-	-	-	1	1	-	2	5	13	10	-
1999.....	52	-	-	-	-	1	-	3	3	9	18	18	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Diseases of appendix (K35-K38)

All races, both sexes

2007.....	426	1	2	10	16	9	19	40	56	82	110	81	-
2006.....	424	3	4	10	8	11	24	33	58	83	99	91	-
2005.....	439	2	4	12	9	9	19	33	57	76	110	108	-
2004.....	452	3	4	13	8	9	14	45	52	81	138	85	-
2003.....	439	4	5	11	11	16	25	50	43	64	132	78	-
2002.....	480	3	-	7	12	10	27	54	64	89	120	94	-
2001.....	428	1	8	6	15	15	26	40	45	89	90	93	-
2000.....	435	3	5	15	11	11	27	43	60	71	121	68	-
1999.....	390	3	2	26	5	6	24	49	48	69	96	62	-

All races, male

2007.....	259	1	1	5	12	6	16	27	42	46	65	38	-
2006.....	253	1	1	6	6	6	19	21	33	57	65	38	-
2005.....	247	-	2	9	7	4	15	18	37	53	57	45	-
2004.....	253	3	4	11	4	7	13	32	29	43	70	37	-
2003.....	258	2	2	5	9	9	16	30	30	46	75	34	-
2002.....	275	1	-	3	7	9	17	39	45	56	67	31	-
2001.....	252	-	7	5	8	11	17	28	30	57	52	37	-
2000.....	237	2	2	4	7	10	16	29	43	40	60	24	-
1999.....	236	1	2	15	5	5	17	37	34	38	47	35	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Diseases of appendix (K35-K38)

All races, female

2007.....	167	-	1	5	4	3	3	13	14	36	45	43	-
2006.....	171	2	3	4	2	5	5	12	25	26	34	53	-
2005.....	192	2	2	3	2	5	4	15	20	23	53	63	-
2004.....	199	-	-	2	4	2	1	13	23	38	68	48	-
2003.....	181	2	3	6	2	7	9	20	13	18	57	44	-
2002.....	205	2	-	4	5	1	10	15	19	33	53	63	-
2001.....	176	1	1	1	7	4	9	12	15	32	38	56	-
2000.....	198	1	3	11	4	1	11	14	17	31	61	44	-
1999.....	154	2	-	11	-	1	7	12	14	31	49	27	-

White, both sexes

2007 1/.....	342	1	1	5	9	4	14	27	40	71	94	76	-
2006 1/.....	350	2	3	5	6	7	20	23	41	72	86	85	-
2005 1/.....	373	2	4	7	4	8	14	23	46	65	98	102	-
2004 1/.....	377	3	3	8	5	5	10	35	44	69	117	78	-
2003 1/.....	355	3	3	6	7	10	14	38	32	53	117	72	-
2002.....	404	2	-	4	7	8	18	40	56	71	112	86	-
2001.....	342	1	5	4	10	8	17	28	29	80	77	83	-
2000.....	348	3	4	13	8	4	18	29	50	54	100	65	-
1999.....	314	3	-	18	3	4	19	38	30	57	84	58	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Diseases of appendix (K35-K38)

White, male

2007 1/.....	210	1	1	2	7	4	13	17	31	42	57	35	-
2006 1/.....	208	-	1	2	6	4	17	15	25	48	55	35	-
2005 1/.....	208	-	2	6	3	4	10	12	31	46	52	42	-
2004 1/.....	210	3	3	8	3	3	9	25	26	36	63	31	-
2003 1/.....	213	2	2	1	6	5	12	24	23	38	67	33	-
2002.....	225	-	-	2	3	7	13	29	40	44	62	25	-
2001.....	198	-	4	3	5	5	12	20	22	52	46	29	-
2000.....	183	2	2	4	5	4	9	19	34	31	51	22	-
1999.....	184	1	-	10	3	3	13	28	24	30	39	33	-

White, female

2007 1/.....	132	-	-	3	2	-	1	10	9	29	37	41	-
2006 1/.....	142	2	2	3	-	3	3	8	16	24	31	50	-
2005 1/.....	165	2	2	1	1	4	4	11	15	19	46	60	-
2004 1/.....	167	-	-	-	2	2	1	10	18	33	54	47	-
2003 1/.....	142	1	1	5	1	5	2	14	9	15	50	39	-
2002.....	179	2	-	2	4	1	5	11	16	27	50	61	-
2001.....	144	1	1	1	5	3	5	8	7	28	31	54	-
2000.....	165	1	2	9	3	-	9	10	16	23	49	43	-
1999.....	130	2	-	8	-	1	6	10	6	27	45	25	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Diseases of appendix (K35-K38)

Black, both sexes

2007 1/.....	65	-	1	5	5	4	5	12	14	7	8	4	-
2006 1/.....	59	-	1	2	2	4	4	8	16	7	11	4	-
2005 1/.....	55	-	-	4	5	1	5	9	9	9	9	4	-
2004 1/.....	59	-	1	3	2	4	3	8	7	10	16	5	-
2003 1/.....	70	-	2	5	4	5	11	8	10	9	11	5	-
2002.....	63	1	-	3	5	2	5	13	7	14	5	8	-
2001.....	73	-	2	2	5	6	8	10	16	7	11	6	-
2000.....	77	-	-	2	2	7	9	12	10	16	16	3	-
1999.....	68	-	2	7	1	2	5	10	17	10	11	3	-

Black, male

2007 1/.....	39	-	-	3	5	1	3	10	9	3	2	3	-
2006 1/.....	33	-	-	2	-	2	2	5	7	5	9	1	-
2005 1/.....	33	-	-	3	4	-	5	5	5	6	4	1	-
2004 1/.....	39	-	1	2	1	4	3	7	3	7	6	5	-
2003 1/.....	37	-	-	4	3	3	4	4	6	7	5	1	-
2002.....	42	1	-	1	4	2	1	10	4	9	4	6	-
2001.....	47	-	2	2	3	6	5	7	8	4	5	5	-
2000.....	48	-	-	-	1	6	7	8	9	9	6	2	-
1999.....	48	-	2	4	1	2	4	9	9	7	8	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Diseases of appendix (K35-K38)

Black, female

2007 1/.....	26	-	1	2	-	3	2	2	5	4	6	1	-
2006 1/.....	26	-	1	-	2	2	2	3	9	2	2	3	-
2005 1/.....	22	-	-	1	1	1	-	4	4	3	5	3	-
2004 1/.....	20	-	-	1	1	-	-	1	4	3	10	-	-
2003 1/.....	33	-	2	1	1	2	7	4	4	2	6	4	-
2002.....	21	-	-	2	1	-	4	3	3	5	1	2	-
2001.....	26	-	-	-	2	-	3	3	8	3	6	1	-
2000.....	29	-	-	2	1	1	2	4	1	7	10	1	-
1999.....	20	-	-	3	-	-	1	1	8	3	3	1	-

American Indian 2/, both sexes

2007 1/.....	3	-	-	-	1	-	-	-	-	1	1	-	-
2006 1/.....	5	-	-	2	-	-	-	2	-	-	1	-	-
2005 1/.....	3	-	-	-	-	-	-	-	1	-	1	1	-
2004 1/.....	5	-	-	1	1	-	-	2	1	-	-	-	-
2003 1/.....	2	-	-	-	-	-	-	1	1	-	-	-	-
2002.....	3	-	-	-	-	-	2	1	-	-	-	-	-
2001.....	2	-	1	-	-	1	-	-	-	-	-	-	-
2000.....	2	-	1	-	-	-	-	-	-	-	1	-	-
1999.....	2	-	-	-	1	-	-	-	1	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Diseases of appendix (K35-K38)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	16	-	-	-	1	1	-	1	2	3	7	1	-
2006 1/.....	10	1	-	1	-	-	-	-	1	4	1	2	-
2005 1/.....	8	-	-	1	-	-	-	1	1	2	2	1	-
2004 1/.....	11	-	-	1	-	-	-	1	-	2	5	2	-
2003 1/.....	12	1	-	-	-	1	-	3	-	2	4	1	-
2002.....	10	-	-	-	-	-	-	2	-	1	4	3	-
2001.....	11	-	-	-	-	-	-	1	2	-	2	2	4
2000.....	8	-	-	-	1	-	-	2	-	1	4	-	-
1999.....	6	-	-	1	-	-	-	1	-	2	1	1	-

Asian or Pacific Islander 3/, male

2007 1/.....	8	-	-	-	-	1	-	-	2	-	5	-	-
2006 1/.....	9	1	-	1	-	-	-	-	1	4	-	2	-
2005 1/.....	3	-	-	-	-	-	-	1	-	1	-	1	-
2004 1/.....	3	-	-	-	-	-	-	1	-	-	1	1	-
2003 1/.....	7	-	-	-	-	1	-	2	-	1	3	-	-
2002.....	7	-	-	-	-	-	-	2	-	1	3	1	-
2001.....	6	-	-	-	-	-	-	1	-	1	1	3	-
2000.....	6	-	-	-	1	-	-	2	-	-	3	-	-
1999.....	2	-	-	1	-	-	-	-	-	1	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Diseases of appendix (K35-K38)

Asian or Pacific Islander 3/, female

2007 1/.....	8	-	-	-	1	-	-	1	-	3	2	1	-
2006 1/.....	1	-	-	-	-	-	-	-	-	-	1	-	-
2005 1/.....	5	-	-	1	-	-	-	-	1	1	2	-	-
2004 1/.....	8	-	-	1	-	-	-	-	-	2	4	1	-
2003 1/.....	5	1	-	-	-	-	-	1	-	1	1	1	-
2002.....	3	-	-	-	-	-	-	-	-	1	2	-	-
2001.....	5	-	-	-	-	-	1	1	-	1	1	1	-
2000.....	2	-	-	-	-	-	-	-	-	1	1	-	-
1999.....	4	-	-	-	-	-	-	1	-	1	1	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hernia (K40-K46)													
All races, both sexes													
2007.....	1,698	35	5	2	2	8	35	98	170	244	474	625	-
2006.....	1,744	34	3	2	7	9	40	87	178	234	510	640	-
2005.....	1,639	43	4	2	2	5	39	83	181	210	454	616	-
2004.....	1,602	40	4	4	9	10	36	88	130	239	509	533	-
2003.....	1,613	37	6	6	3	5	28	105	161	237	462	563	-
2002.....	1,595	29	3	7	5	11	35	77	126	225	489	588	-
2001.....	1,522	43	2	3	3	15	34	82	131	213	443	553	-
2000.....	1,522	46	4	3	5	5	31	84	113	210	466	555	-
1999.....	1,464	44	3	2	3	16	39	55	107	215	451	529	-
All races, male													
2007.....	708	21	3	1	1	2	18	55	91	111	191	214	-
2006.....	739	20	2	1	5	6	22	45	83	112	215	228	-
2005.....	646	28	1	1	-	1	15	45	91	86	175	203	-
2004.....	652	20	1	2	5	7	22	45	62	120	198	170	-
2003.....	668	20	4	4	1	4	16	59	72	111	199	178	-
2002.....	599	18	3	5	4	7	17	35	56	103	189	162	-
2001.....	587	28	1	-	1	10	16	44	64	94	173	156	-
2000.....	586	28	3	3	3	1	13	45	49	92	195	154	-
1999.....	597	26	2	1	3	8	23	32	60	93	192	157	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hernia (K40-K46)													
All races, female													
2007.....	990	14	2	1	1	6	17	43	79	133	283	411	-
2006.....	1,005	14	1	1	2	3	18	42	95	122	295	412	-
2005.....	993	15	3	1	2	4	24	38	90	124	279	413	-
2004.....	950	20	3	2	4	3	14	43	68	119	311	363	-
2003.....	945	17	2	2	2	1	12	46	89	126	263	385	-
2002.....	996	11	-	2	1	4	18	42	70	122	300	426	-
2001.....	935	15	1	3	2	5	18	38	67	119	270	397	-
2000.....	936	18	1	-	2	4	18	39	64	118	271	401	-
1999.....	867	18	1	1	-	8	16	23	47	122	259	372	-
White, both sexes													
2007 1/.....	1,504	28	3	1	2	5	25	73	143	205	423	596	-
2006 1/.....	1,547	27	3	1	6	7	30	69	154	200	452	598	-
2005 1/.....	1,456	35	4	1	1	4	29	65	149	182	414	572	-
2004 1/.....	1,423	32	2	4	7	8	25	69	107	208	469	492	-
2003 1/.....	1,421	29	3	3	2	3	20	85	131	208	415	522	-
2002.....	1,433	24	2	7	5	6	26	58	102	192	455	556	-
2001.....	1,343	31	2	1	2	13	25	67	107	185	399	511	-
2000.....	1,354	36	2	3	3	4	22	67	99	185	421	512	-
1999.....	1,298	34	3	2	2	10	30	41	94	183	408	491	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hernia (K40-K46)													
White, male													
2007 1/.....	608	18	2	1	1	-	14	41	72	95	163	201	-
2006 1/.....	654	16	2	-	5	6	18	38	71	96	189	213	-
2005 1/.....	568	22	1	-	-	1	12	38	76	76	157	185	-
2004 1/.....	572	15	-	2	4	5	15	38	53	103	182	155	-
2003 1/.....	585	16	3	2	1	2	11	49	62	96	180	163	-
2002.....	533	14	2	5	4	3	14	30	47	86	178	150	-
2001.....	502	18	1	-	1	8	13	32	50	85	155	139	-
2000.....	520	23	2	3	1	1	10	34	43	82	180	141	-
1999.....	535	20	2	1	2	5	20	26	57	86	177	139	-
White, female													
2007 1/.....	896	10	1	-	1	5	11	32	71	110	260	395	-
2006 1/.....	893	11	1	1	1	1	12	31	83	104	263	385	-
2005 1/.....	888	13	3	1	1	3	17	27	73	106	257	387	-
2004 1/.....	851	17	2	2	3	3	10	31	54	105	287	337	-
2003 1/.....	836	13	-	1	1	1	9	36	69	112	235	359	-
2002.....	900	10	-	2	1	3	12	28	55	106	277	406	-
2001.....	841	13	1	1	1	5	12	35	57	100	244	372	-
2000.....	834	13	-	-	2	3	12	33	56	103	241	371	-
1999.....	763	14	1	1	-	5	10	15	37	97	231	352	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Hernia (K40-K46)

Black, both sexes

2007 1/.....	167	7	2	-	-	3	8	20	25	37	41	24	-
2006 1/.....	172	4	-	1	-	2	10	18	24	31	46	36	-
2005 1/.....	152	5	-	1	1	1	8	16	28	25	33	34	-
2004 1/.....	150	6	2	-	2	2	9	17	20	28	31	33	-
2003 1/.....	164	8	3	3	-	2	8	18	28	24	39	31	-
2002.....	137	4	1	-	-	3	8	17	21	27	30	26	-
2001.....	157	11	-	1	1	2	9	15	23	25	34	36	-
2000.....	145	7	2	-	2	1	8	13	13	21	38	40	-
1999.....	140	9	-	-	1	4	9	12	12	24	38	31	-

Black, male

2007 1/.....	86	3	1	-	-	2	3	11	18	14	22	12	-
2006 1/.....	72	2	-	1	-	-	4	7	12	14	19	13	-
2005 1/.....	66	4	-	1	-	-	3	7	12	10	16	13	-
2004 1/.....	67	4	1	-	1	2	5	7	9	16	11	11	-
2003 1/.....	72	4	1	2	-	2	5	9	10	13	15	11	-
2002.....	56	3	1	-	-	2	3	5	7	15	11	9	-
2001.....	76	9	-	-	-	2	3	12	13	7	15	15	-
2000.....	54	3	1	-	2	-	2	9	5	9	12	11	-
1999.....	53	6	-	-	1	1	3	6	2	7	13	14	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hernia (K40-K46)													
Black, female													
2007 1/.....	81	4	1	-	-	1	5	9	7	23	19	12	-
2006 1/.....	100	2	-	-	-	2	6	11	12	17	27	23	-
2005 1/.....	86	1	-	-	1	1	5	9	16	15	17	21	-
2004 1/.....	83	2	1	-	1	-	4	10	11	12	20	22	-
2003 1/.....	92	4	2	1	-	-	3	9	18	11	24	20	-
2002.....	81	1	-	-	-	1	5	12	14	12	19	17	-
2001.....	81	2	-	1	1	-	6	3	10	18	19	21	-
2000.....	91	4	1	-	-	1	6	4	8	12	26	29	-
1999.....	87	3	-	-	-	3	6	6	10	17	25	17	-
American Indian 2/, both sexes													
2007 1/.....	12	-	-	-	-	-	-	3	1	1	5	2	-
2006 1/.....	8	-	-	-	-	-	-	-	-	3	5	-	-
2005 1/.....	17	2	-	-	-	-	2	-	3	3	4	3	-
2004 1/.....	14	1	-	-	-	-	-	2	2	3	3	3	-
2003 1/.....	15	-	-	-	1	-	-	2	1	4	1	6	-
2002.....	14	-	-	-	-	2	1	2	1	3	4	1	-
2001.....	5	-	-	-	-	-	-	-	1	1	1	2	-
2000.....	11	2	-	-	-	-	1	2	1	3	2	-	-
1999.....	8	1	-	-	-	1	-	1	1	2	-	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Hernia (K40-K46)

American Indian 2/, male

2007 1/.....	7	-	-	-	-	-	3	-	1	2	1	-
2006 1/.....	5	-	-	-	-	-	-	-	2	3	-	-
2005 1/.....	5	1	-	-	-	-	-	2	-	1	1	-
2004 1/.....	5	1	-	-	-	-	-	-	1	1	2	-
2003 1/.....	6	-	-	-	-	-	1	-	1	-	4	-
2002.....	4	-	-	-	-	2	-	1	1	-	-	-
2001.....	3	-	-	-	-	-	-	1	1	-	1	-
2000.....	5	1	-	-	-	-	1	1	1	1	-	-
1999.....	2	-	-	-	-	1	-	-	1	-	-	-

American Indian 2/, female

2007 1/.....	5	-	-	-	-	-	-	1	-	3	1	-
2006 1/.....	3	-	-	-	-	-	-	-	1	2	-	-
2005 1/.....	12	1	-	-	-	-	2	-	1	3	3	2
2004 1/.....	9	-	-	-	-	-	2	2	2	2	1	-
2003 1/.....	9	-	-	-	1	-	1	1	3	1	2	-
2002.....	10	-	-	-	-	-	1	2	-	2	4	1
2001.....	2	-	-	-	-	-	-	-	-	-	1	-
2000.....	6	1	-	-	-	-	-	1	-	2	2	-
1999.....	6	1	-	-	-	-	-	1	-	2	-	2

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Hernia (K40-K46)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	15	-	-	1	-	-	2	2	1	1	5	3	-
2006 1/.....	17	3	-	-	1	-	-	-	-	-	7	6	-
2005 1/.....	14	1	-	-	-	-	-	2	1	-	3	7	-
2004 1/.....	15	1	-	-	-	-	2	-	1	-	6	5	-
2003 1/.....	13	-	-	-	-	-	-	-	1	1	7	4	-
2002.....	11	1	-	-	-	-	-	-	2	3	-	5	-
2001.....	17	1	-	1	-	-	-	-	-	2	9	4	-
2000.....	12	1	-	-	-	-	-	2	-	1	5	3	-
1999.....	18	-	-	-	-	1	-	1	-	6	5	5	-

Asian or Pacific Islander 3/, male

2007 1/.....	7	-	-	-	-	-	1	-	1	1	4	-	-
2006 1/.....	8	2	-	-	-	-	-	-	-	-	4	2	-
2005 1/.....	7	1	-	-	-	-	-	-	1	-	1	4	-
2004 1/.....	8	-	-	-	-	-	2	-	-	-	4	2	-
2003 1/.....	5	-	-	-	-	-	-	-	-	1	4	-	-
2002.....	6	1	-	-	-	-	-	-	1	1	-	3	-
2001.....	6	1	-	-	-	-	-	-	-	1	3	1	-
2000.....	7	1	-	-	-	-	-	1	-	-	3	2	-
1999.....	7	-	-	-	-	1	-	-	-	-	2	4	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Hernia (K40-K46)

Asian or Pacific Islander 3/, female

2007 1/.....	8	-	-	1	-	-	1	2	-	-	1	3	-
2006 1/.....	9	1	-	-	1	-	-	-	-	-	3	4	-
2005 1/.....	7	-	-	-	-	-	-	2	-	-	2	3	-
2004 1/.....	7	1	-	-	-	-	-	-	1	-	2	3	-
2003 1/.....	8	-	-	-	-	-	-	-	1	-	3	4	-
2002.....	5	-	-	-	-	-	-	-	1	2	-	2	-
2001.....	11	-	-	1	-	-	-	-	-	1	6	3	-
2000.....	5	-	-	-	-	-	-	1	-	1	2	1	-
1999.....	11	-	-	-	-	-	-	1	-	6	3	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Chronic liver disease and cirrhosis (K70,K73-K74)													
All races, both sexes													
2007.....	29,165	4	4	-	30	384	2,570	8,212	8,004	5,167	3,694	1,093	3
2006.....	27,555	5	3	-	26	316	2,551	7,712	7,217	4,917	3,778	1,026	4
2005.....	27,530	10	1	2	23	311	2,688	7,517	7,126	5,066	3,781	1,002	3
2004.....	27,013	9	1	1	19	309	2,799	7,496	6,569	5,119	3,733	957	1
2003.....	27,503	2	-	-	16	358	3,020	7,466	6,428	5,406	3,858	946	3
2002.....	27,257	6	3	6	31	374	3,154	7,216	6,097	5,381	4,000	985	4
2001.....	27,035	3	-	2	26	387	3,336	7,259	5,750	5,486	3,801	979	6
2000.....	26,552	3	3	3	36	415	3,371	6,654	5,774	5,482	3,829	979	3
1999.....	26,259	5	2	2	26	407	3,302	6,368	5,637	5,642	3,901	965	2
All races, male													
2007.....	19,151	3	2	-	14	261	1,722	5,788	5,732	3,219	1,949	458	3
2006.....	17,866	2	3	-	14	215	1,666	5,517	5,095	3,008	1,902	441	3
2005.....	17,937	6	1	1	13	196	1,786	5,423	5,094	3,052	1,920	442	3
2004.....	17,545	6	-	-	9	221	1,916	5,382	4,587	3,122	1,917	385	-
2003.....	17,912	2	-	-	8	252	2,033	5,514	4,537	3,265	1,901	397	3
2002.....	17,401	6	1	3	14	236	2,079	5,285	4,228	3,197	1,953	395	4
2001.....	17,393	1	-	2	12	259	2,206	5,295	4,028	3,335	1,833	416	6
2000.....	17,214	3	1	2	19	280	2,314	4,975	4,005	3,317	1,874	421	3
1999.....	17,115	4	-	1	11	257	2,314	4,829	3,943	3,454	1,905	395	2

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Chronic liver disease and cirrhosis (K70,K73-K74)													
All races, female													
2007.....	10,014	1	2	-	16	123	848	2,424	2,272	1,948	1,745	635	-
2006.....	9,689	3	-	-	12	101	885	2,195	2,122	1,909	1,876	585	1
2005.....	9,593	4	-	1	10	115	902	2,094	2,032	2,014	1,861	560	-
2004.....	9,468	3	1	1	10	88	883	2,114	1,982	1,997	1,816	572	1
2003.....	9,591	-	-	-	8	106	987	1,952	1,891	2,141	1,957	549	-
2002.....	9,856	-	2	3	17	138	1,075	1,931	1,869	2,184	2,047	590	-
2001.....	9,642	2	-	-	14	128	1,130	1,964	1,722	2,151	1,968	563	-
2000.....	9,338	-	2	1	17	135	1,057	1,679	1,769	2,165	1,955	558	-
1999.....	9,144	1	2	1	15	150	988	1,539	1,694	2,188	1,996	570	-
White, both sexes													
2007 1/.....	25,490	3	4	-	20	310	2,184	7,007	6,879	4,624	3,435	1,021	3
2006 1/.....	24,191	3	2	-	17	250	2,136	6,610	6,285	4,424	3,499	961	4
2005 1/.....	24,003	8	1	2	15	231	2,284	6,342	6,138	4,581	3,472	927	2
2004 1/.....	23,562	7	1	1	12	251	2,327	6,348	5,619	4,613	3,494	888	1
2003 1/.....	24,005	1	-	-	9	286	2,540	6,241	5,519	4,883	3,635	888	3
2002.....	23,809	5	3	5	25	285	2,656	6,020	5,259	4,877	3,744	926	4
2001.....	23,408	3	-	1	10	307	2,745	5,989	4,924	4,939	3,561	924	5
2000.....	22,982	-	3	2	21	312	2,745	5,515	4,948	4,925	3,591	918	2
1999.....	22,620	4	-	1	15	323	2,676	5,230	4,760	5,073	3,640	896	2

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

Chronic liver disease and cirrhosis (K70,K73-K74)

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
White, male													
2007 1/.....	16,801	2	2	-	7	219	1,477	5,007	4,963	2,874	1,812	435	3
2006 1/.....	15,739	2	2	-	10	174	1,424	4,772	4,464	2,708	1,766	414	3
2005 1/.....	15,679	5	1	1	7	152	1,539	4,633	4,407	2,761	1,759	412	2
2004 1/.....	15,334	4	-	-	7	182	1,614	4,604	3,950	2,807	1,802	364	-
2003 1/.....	15,662	1	-	-	5	205	1,748	4,667	3,914	2,962	1,786	371	3
2002.....	15,278	5	1	3	12	193	1,783	4,489	3,672	2,901	1,843	372	4
2001.....	15,048	1	-	1	6	211	1,826	4,431	3,442	3,003	1,728	394	5
2000.....	15,002	-	1	1	11	224	1,921	4,205	3,475	3,006	1,762	394	2
1999.....	14,790	4	-	-	8	213	1,906	4,021	3,372	3,114	1,781	369	2
White, female													
2007 1/.....	8,689	1	2	-	13	91	707	2,000	1,916	1,750	1,623	586	-
2006 1/.....	8,452	1	-	-	7	76	712	1,838	1,821	1,716	1,733	547	1
2005 1/.....	8,324	3	-	1	8	79	745	1,709	1,731	1,820	1,713	515	-
2004 1/.....	8,228	3	1	1	5	69	713	1,744	1,669	1,806	1,692	524	1
2003 1/.....	8,343	-	-	-	4	81	792	1,574	1,605	1,921	1,849	517	-
2002.....	8,531	-	2	2	13	92	873	1,531	1,587	1,976	1,901	554	-
2001.....	8,360	2	-	-	4	96	919	1,558	1,482	1,936	1,833	530	-
2000.....	7,980	-	2	1	10	88	824	1,310	1,473	1,919	1,829	524	-
1999.....	7,830	-	-	1	7	110	770	1,209	1,388	1,959	1,859	527	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Chronic liver disease and cirrhosis (K70,K73-K74)

Black, both sexes

2007 1/.....	2,558	-	-	-	7	28	203	853	867	404	157	39	-
2006 1/.....	2,349	2	1	-	6	22	256	804	710	340	167	41	-
2005 1/.....	2,530	2	-	-	4	33	239	892	763	357	193	46	1
2004 1/.....	2,534	2	-	-	4	22	310	903	733	374	143	43	-
2003 1/.....	2,621	1	-	-	3	41	326	948	713	406	141	42	-
2002.....	2,588	1	-	1	5	51	318	968	660	365	178	41	-
2001.....	2,775	-	-	1	10	47	422	1,011	657	417	175	34	1
2000.....	2,737	2	-	1	12	57	451	937	633	434	165	44	1
1999.....	2,833	-	2	1	10	55	457	922	699	438	195	54	-

Black, male

2007 1/.....	1,682	-	-	-	5	14	124	560	606	272	90	11	-
2006 1/.....	1,519	-	1	-	3	9	155	542	481	220	91	17	-
2005 1/.....	1,656	1	-	-	4	14	144	603	536	230	106	17	1
2004 1/.....	1,670	2	-	-	-	14	199	618	505	244	74	14	-
2003 1/.....	1,727	1	-	-	2	25	191	674	503	237	73	21	-
2002.....	1,622	1	-	-	1	27	183	645	445	227	78	15	-
2001.....	1,843	-	-	1	2	26	274	713	469	270	76	11	1
2000.....	1,730	2	-	1	6	27	284	639	414	258	81	17	1
1999.....	1,842	-	-	1	3	28	296	663	460	277	96	18	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Chronic liver disease and cirrhosis (K70,K73-K74)													
Black, female													
2007 1/.....	876	-	-	-	2	14	79	293	261	132	67	28	-
2006 1/.....	830	2	-	-	3	13	101	262	229	120	76	24	-
2005 1/.....	874	1	-	-	-	19	95	289	227	127	87	29	-
2004 1/.....	864	-	-	-	4	8	111	285	228	130	69	29	-
2003 1/.....	894	-	-	-	1	16	135	274	210	169	68	21	-
2002.....	966	-	-	1	4	24	135	323	215	138	100	26	-
2001.....	932	-	-	-	8	21	148	298	188	147	99	23	-
2000.....	1,007	-	-	-	6	30	167	298	219	176	84	27	-
1999.....	991	-	2	-	7	27	161	259	239	161	99	36	-
American Indian 2/, both sexes													
2007 1/.....	709	1	-	-	3	37	145	259	167	66	20	11	-
2006 1/.....	596	-	-	-	3	31	121	201	135	73	25	7	-
2005 1/.....	596	-	-	-	3	36	125	196	143	61	27	5	-
2004 1/.....	577	-	-	-	3	21	128	184	144	64	28	5	-
2003 1/.....	570	-	-	-	4	25	127	203	134	50	23	4	-
2002.....	547	-	-	-	1	35	142	162	112	67	25	3	-
2001.....	533	-	-	-	6	25	135	187	100	58	18	4	-
2000.....	534	-	-	-	2	39	137	147	123	59	25	2	-
1999.....	513	-	-	-	1	21	135	158	123	58	15	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Chronic liver disease and cirrhosis (K70,K73-K74)

American Indian 2/, male

2007 1/.....	415	1	-	-	2	22	90	153	93	38	12	4	-
2006 1/.....	330	-	-	-	1	21	59	119	82	37	10	1	-
2005 1/.....	348	-	-	-	2	22	70	116	95	29	11	3	-
2004 1/.....	323	-	-	-	2	11	75	105	84	30	15	1	-
2003 1/.....	317	-	-	-	1	17	72	118	72	26	9	2	-
2002.....	319	-	-	-	1	15	82	105	65	38	12	1	-
2001.....	309	-	-	-	4	17	82	99	68	29	8	2	-
2000.....	298	-	-	-	1	23	78	91	71	21	12	1	-
1999.....	299	-	-	-	-	10	84	99	70	28	7	1	-

American Indian 2/, female

2007 1/.....	294	-	-	-	1	15	55	106	74	28	8	7	-
2006 1/.....	266	-	-	-	2	10	62	82	53	36	15	6	-
2005 1/.....	248	-	-	-	1	14	55	80	48	32	16	2	-
2004 1/.....	254	-	-	-	1	10	53	79	60	34	13	4	-
2003 1/.....	253	-	-	-	3	8	55	85	62	24	14	2	-
2002.....	228	-	-	-	-	20	60	57	47	29	13	2	-
2001.....	224	-	-	-	2	8	53	88	32	29	10	2	-
2000.....	236	-	-	-	1	16	59	56	52	38	13	1	-
1999.....	214	-	-	-	1	11	51	59	53	30	8	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Chronic liver disease and cirrhosis (K70,K73-K74)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	408	-	-	-	-	9	38	93	91	73	82	22	-
2006 1/.....	419	-	-	-	-	13	38	97	87	80	87	17	-
2005 1/.....	401	-	-	-	1	11	40	87	82	67	89	24	-
2004 1/.....	340	-	-	-	-	15	34	61	73	68	68	21	-
2003 1/.....	307	-	-	-	-	6	27	74	62	67	59	12	-
2002.....	313	-	-	-	-	3	38	66	66	72	53	15	-
2001.....	319	-	-	-	-	8	34	72	69	72	47	17	-
2000.....	299	1	-	-	1	7	38	55	70	64	48	15	-
1999.....	293	1	-	-	-	8	34	58	55	73	51	13	-

Asian or Pacific Islander 3/, male

2007 1/.....	253	-	-	-	-	6	31	68	70	35	35	8	-
2006 1/.....	278	-	-	-	-	11	28	84	68	43	35	9	-
2005 1/.....	254	-	-	-	-	8	33	71	56	32	44	10	-
2004 1/.....	218	-	-	-	-	14	28	55	48	41	26	6	-
2003 1/.....	206	-	-	-	-	5	22	55	48	40	33	3	-
2002.....	182	-	-	-	-	1	31	46	46	31	20	7	-
2001.....	193	-	-	-	-	5	24	52	49	33	21	9	-
2000.....	184	1	-	-	1	6	31	40	45	32	19	9	-
1999.....	184	-	-	-	-	6	28	46	41	35	21	7	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Chronic liver disease and cirrhosis (K70,K73-K74)

Asian or Pacific Islander 3/, female

2007 1/.....	155	-	-	-	-	3	7	25	21	38	47	14	-
2006 1/.....	141	-	-	-	-	2	10	13	19	37	52	8	-
2005 1/.....	147	-	-	-	1	3	7	16	26	35	45	14	-
2004 1/.....	122	-	-	-	-	1	6	6	25	27	42	15	-
2003 1/.....	101	-	-	-	-	1	5	19	14	27	26	9	-
2002.....	131	-	-	-	-	2	7	20	20	41	33	8	-
2001.....	126	-	-	-	-	3	10	20	20	39	26	8	-
2000.....	115	-	-	-	-	1	7	15	25	32	29	6	-
1999.....	109	1	-	-	-	2	6	12	14	38	30	6	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alcoholic liver disease (K70)													
All races, both sexes													
2007.....	14,406	-	-	-	17	293	1,834	5,126	4,309	1,959	747	118	3
2006.....	13,050	-	-	-	14	245	1,732	4,562	3,823	1,783	773	117	1
2005.....	12,928	-	-	-	7	217	1,806	4,490	3,721	1,844	747	94	2
2004.....	12,548	-	-	-	11	228	1,871	4,392	3,393	1,841	709	102	1
2003.....	12,360	-	-	-	6	243	1,971	4,288	3,247	1,788	729	86	2
2002.....	12,121	-	1	-	8	242	1,971	4,137	3,083	1,849	730	98	2
2001.....	12,207	-	-	-	8	264	2,145	4,203	2,913	1,848	717	104	5
2000.....	12,109	-	-	-	8	279	2,226	3,888	2,969	1,891	751	95	2
1999.....	11,958	-	-	-	8	259	2,186	3,802	2,859	1,935	789	118	2
All races, male													
2007.....	10,549	-	-	-	8	201	1,220	3,680	3,294	1,505	554	84	3
2006.....	9,443	-	-	-	6	175	1,122	3,295	2,870	1,357	538	79	1
2005.....	9,425	-	-	-	6	147	1,189	3,274	2,850	1,352	537	68	2
2004.....	9,155	-	-	-	8	167	1,267	3,176	2,549	1,392	527	69	-
2003.....	9,104	-	-	-	3	182	1,317	3,228	2,459	1,343	518	52	2
2002.....	8,814	-	1	-	6	162	1,305	3,074	2,337	1,367	495	65	2
2001.....	8,932	-	-	-	5	179	1,446	3,099	2,242	1,379	505	72	5
2000.....	8,985	-	-	-	5	202	1,533	2,968	2,216	1,433	561	65	2
1999.....	8,922	-	-	-	6	165	1,520	2,927	2,194	1,497	538	73	2

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alcoholic liver disease (K70)													
All races, female													
2007.....	3,857	-	-	-	9	92	614	1,446	1,015	454	193	34	-
2006.....	3,607	-	-	-	8	70	610	1,267	953	426	235	38	-
2005.....	3,503	-	-	-	1	70	617	1,216	871	492	210	26	-
2004.....	3,393	-	-	-	3	61	604	1,216	844	449	182	33	1
2003.....	3,256	-	-	-	3	61	654	1,060	788	445	211	34	-
2002.....	3,307	-	-	-	2	80	666	1,063	746	482	235	33	-
2001.....	3,275	-	-	-	3	85	699	1,104	671	469	212	32	-
2000.....	3,124	-	-	-	3	77	693	920	753	458	190	30	-
1999.....	3,036	-	-	-	2	94	666	875	665	438	251	45	-
White, both sexes													
2007 1/.....	12,541	-	-	-	12	235	1,565	4,437	3,764	1,727	688	110	3
2006 1/.....	11,413	-	-	-	8	191	1,467	3,961	3,381	1,579	713	112	1
2005 1/.....	11,205	-	-	-	4	164	1,523	3,859	3,243	1,641	682	87	2
2004 1/.....	10,859	-	-	-	7	189	1,582	3,779	2,914	1,638	654	95	1
2003 1/.....	10,625	-	-	-	4	198	1,658	3,619	2,797	1,586	682	79	2
2002.....	10,423	-	1	-	4	187	1,660	3,508	2,656	1,652	665	88	2
2001.....	10,387	-	-	-	2	211	1,771	3,521	2,498	1,629	660	90	5
2000.....	10,272	-	-	-	6	218	1,810	3,246	2,533	1,689	681	88	1
1999.....	10,041	-	-	-	5	205	1,764	3,152	2,405	1,698	708	102	2

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alcoholic liver disease (K70)													
White, male													
2007 1/.....	9,267	-	-	-	6	168	1,049	3,222	2,905	1,324	511	79	3
2006 1/.....	8,346	-	-	-	3	140	977	2,887	2,558	1,206	499	75	1
2005 1/.....	8,250	-	-	-	3	116	1,025	2,837	2,507	1,207	490	63	2
2004 1/.....	7,992	-	-	-	6	140	1,082	2,771	2,203	1,238	487	65	-
2003 1/.....	7,905	-	-	-	2	149	1,131	2,750	2,143	1,198	482	48	2
2002.....	7,688	-	1	-	4	135	1,112	2,649	2,047	1,226	455	57	2
2001.....	7,689	-	-	-	2	151	1,202	2,650	1,931	1,216	469	63	5
2000.....	7,770	-	-	-	4	168	1,264	2,538	1,930	1,288	517	60	1
1999.....	7,602	-	-	-	4	136	1,249	2,469	1,876	1,317	486	63	2
White, female													
2007 1/.....	3,274	-	-	-	6	67	516	1,215	859	403	177	31	-
2006 1/.....	3,067	-	-	-	5	51	490	1,074	823	373	214	37	-
2005 1/.....	2,955	-	-	-	1	48	498	1,022	736	434	192	24	-
2004 1/.....	2,867	-	-	-	1	49	500	1,008	711	400	167	30	1
2003 1/.....	2,720	-	-	-	2	49	527	869	654	388	200	31	-
2002.....	2,735	-	-	-	-	52	548	859	609	426	210	31	-
2001.....	2,698	-	-	-	-	60	569	871	567	413	191	27	-
2000.....	2,502	-	-	-	2	50	546	708	603	401	164	28	-
1999.....	2,439	-	-	-	1	69	515	683	529	381	222	39	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alcoholic liver disease (K70)													
Black, both sexes													
2007 1/.....	1,203	-	-	-	4	20	127	434	398	175	39	6	-
2006 1/.....	1,045	-	-	-	3	17	143	393	303	141	42	3	-
2005 1/.....	1,169	-	-	-	2	15	152	447	347	154	46	6	-
2004 1/.....	1,169	-	-	-	2	9	164	443	362	155	31	3	-
2003 1/.....	1,222	-	-	-	-	19	188	473	345	159	33	5	-
2002.....	1,191	-	-	-	3	26	170	464	324	145	51	8	-
2001.....	1,348	-	-	-	1	27	245	516	328	178	44	9	-
2000.....	1,380	-	-	-	-	34	287	508	325	170	50	5	1
1999.....	1,466	-	-	-	2	29	283	519	361	194	64	14	-
Black, male													
2007 1/.....	851	-	-	-	2	10	77	297	293	139	30	3	-
2006 1/.....	712	-	-	-	2	8	83	269	214	105	28	3	-
2005 1/.....	801	-	-	-	2	6	84	309	250	113	33	4	-
2004 1/.....	812	-	-	-	-	7	106	294	262	122	19	2	-
2003 1/.....	865	-	-	-	-	14	111	351	247	114	24	4	-
2002.....	797	-	-	-	1	14	103	316	220	105	32	6	-
2001.....	935	-	-	-	-	12	161	354	242	135	26	5	-
2000.....	918	-	-	-	-	17	187	341	211	124	33	4	1
1999.....	1,018	-	-	-	2	15	179	368	257	148	41	8	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alcoholic liver disease (K70)													
Black, female													
2007 1/.....	352	-	-	-	2	10	50	137	105	36	9	3	-
2006 1/.....	333	-	-	-	1	9	60	124	89	36	14	-	-
2005 1/.....	368	-	-	-	-	9	68	138	97	41	13	2	-
2004 1/.....	357	-	-	-	2	2	58	149	100	33	12	1	-
2003 1/.....	357	-	-	-	-	5	77	122	98	45	9	1	-
2002.....	394	-	-	-	2	12	67	148	104	40	19	2	-
2001.....	413	-	-	-	1	15	84	162	86	43	18	4	-
2000.....	462	-	-	-	-	17	100	167	114	46	17	1	-
1999.....	448	-	-	-	-	14	104	151	104	46	23	6	-
American Indian 2/, both sexes													
2007 1/.....	500	-	-	-	1	31	115	199	105	41	8	-	-
2006 1/.....	426	-	-	-	3	29	95	155	96	39	9	-	-
2005 1/.....	410	-	-	-	1	29	104	137	98	30	10	1	-
2004 1/.....	389	-	-	-	2	20	104	134	87	27	13	2	-
2003 1/.....	403	-	-	-	2	21	108	156	81	27	7	1	-
2002.....	381	-	-	-	1	28	112	126	74	34	6	-	-
2001.....	359	-	-	-	5	20	110	129	61	28	4	2	-
2000.....	339	-	-	-	1	25	99	102	79	23	10	-	-
1999.....	356	-	-	-	1	19	118	107	75	27	8	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Alcoholic liver disease (K70)

American Indian 2/, male

2007 1/.....	304	-	-	-	-	19	72	118	61	29	5	-	-
2006 1/.....	246	-	-	-	1	20	43	91	61	25	5	-	-
2005 1/.....	258	-	-	-	1	19	58	87	68	18	6	1	-
2004 1/.....	240	-	-	-	2	11	62	79	60	14	11	1	-
2003 1/.....	241	-	-	-	1	14	61	92	51	17	5	-	-
2002.....	230	-	-	-	1	13	66	79	45	23	3	-	-
2001.....	218	-	-	-	3	13	68	66	47	17	2	2	-
2000.....	205	-	-	-	-	15	56	63	51	14	6	-	-
1999.....	224	-	-	-	-	9	73	72	45	19	5	1	-

American Indian 2/, female

2007 1/.....	196	-	-	-	1	12	43	81	44	12	3	-	-
2006 1/.....	180	-	-	-	2	9	52	64	35	14	4	-	-
2005 1/.....	152	-	-	-	-	10	46	50	30	12	4	-	-
2004 1/.....	149	-	-	-	-	9	42	55	27	13	2	1	-
2003 1/.....	162	-	-	-	1	7	47	64	30	10	2	1	-
2002.....	151	-	-	-	-	15	46	47	29	11	3	-	-
2001.....	141	-	-	-	2	7	42	63	14	11	2	-	-
2000.....	134	-	-	-	1	10	43	39	28	9	4	-	-
1999.....	132	-	-	-	1	10	45	35	30	8	3	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Alcoholic liver disease (K70)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	162	-	-	-	-	7	27	56	42	16	12	2	-
2006 1/.....	166	-	-	-	-	8	27	53	43	24	9	2	-
2005 1/.....	144	-	-	-	-	9	27	47	33	19	9	-	-
2004 1/.....	131	-	-	-	-	10	21	36	30	21	11	2	-
2003 1/.....	110	-	-	-	-	5	17	40	24	16	7	1	-
2002.....	126	-	-	-	-	1	29	39	29	18	8	2	-
2001.....	113	-	-	-	-	6	19	37	26	13	9	3	-
2000.....	118	-	-	-	1	2	30	32	32	9	10	2	-
1999.....	95	-	-	-	-	6	21	24	18	16	9	1	-

Asian or Pacific Islander 3/, male

2007 1/.....	127	-	-	-	-	4	22	43	35	13	8	2	-
2006 1/.....	139	-	-	-	-	7	19	48	37	21	6	1	-
2005 1/.....	116	-	-	-	-	6	22	41	25	14	8	-	-
2004 1/.....	111	-	-	-	-	9	17	32	24	18	10	1	-
2003 1/.....	93	-	-	-	-	5	14	35	18	14	7	-	-
2002.....	99	-	-	-	-	-	24	30	25	13	5	2	-
2001.....	90	-	-	-	-	3	15	29	22	11	8	2	-
2000.....	92	-	-	-	1	2	26	26	24	7	5	1	-
1999.....	78	-	-	-	-	5	19	18	16	13	6	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Alcoholic liver disease (K70)

Asian or Pacific Islander 3/, female

2007 1/.....	35	-	-	-	-	3	5	13	7	3	4	-	-
2006 1/.....	27	-	-	-	-	1	8	5	6	3	3	1	-
2005 1/.....	28	-	-	-	-	3	5	6	8	5	1	-	-
2004 1/.....	20	-	-	-	-	1	4	4	6	3	1	1	-
2003 1/.....	17	-	-	-	-	-	3	5	6	2	-	1	-
2002.....	27	-	-	-	-	1	5	9	4	5	3	-	-
2001.....	23	-	-	-	-	3	4	8	4	2	1	1	-
2000.....	26	-	-	-	-	-	4	6	8	2	5	1	-
1999.....	17	-	-	-	-	1	2	6	2	3	3	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other chronic liver disease and cirrhosis (K73-K74)													
All races, both sexes													
2007.....	14,759	4	4	-	13	91	736	3,086	3,695	3,208	2,947	975	-
2006.....	14,505	5	3	-	12	71	819	3,150	3,394	3,134	3,005	909	3
2005.....	14,602	10	1	2	16	94	882	3,027	3,405	3,222	3,034	908	1
2004.....	14,465	9	1	1	8	81	928	3,104	3,176	3,278	3,024	855	-
2003.....	15,143	2	-	-	10	115	1,049	3,178	3,181	3,618	3,129	860	1
2002.....	15,136	6	2	6	23	132	1,183	3,079	3,014	3,532	3,270	887	2
2001.....	14,828	3	-	2	18	123	1,191	3,056	2,837	3,638	3,084	875	1
2000.....	14,443	3	3	3	28	136	1,145	2,766	2,805	3,591	3,078	884	1
1999.....	14,301	5	2	2	18	148	1,116	2,566	2,778	3,707	3,112	847	-
All races, male													
2007.....	8,602	3	2	-	6	60	502	2,108	2,438	1,714	1,395	374	-
2006.....	8,423	2	3	-	8	40	544	2,222	2,225	1,651	1,364	362	2
2005.....	8,512	6	1	1	7	49	597	2,149	2,244	1,700	1,383	374	1
2004.....	8,390	6	-	-	1	54	649	2,206	2,038	1,730	1,390	316	-
2003.....	8,808	2	-	-	5	70	716	2,286	2,078	1,922	1,383	345	1
2002.....	8,587	6	-	3	8	74	774	2,211	1,891	1,830	1,458	330	2
2001.....	8,461	1	-	2	7	80	760	2,196	1,786	1,956	1,328	344	1
2000.....	8,229	3	1	2	14	78	781	2,007	1,789	1,884	1,313	356	1
1999.....	8,193	4	-	1	5	92	794	1,902	1,749	1,957	1,367	322	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other chronic liver disease and cirrhosis (K73-K74)													
All races, female													
2007.....	6,157	1	2	-	7	31	234	978	1,257	1,494	1,552	601	-
2006.....	6,082	3	-	-	4	31	275	928	1,169	1,483	1,641	547	1
2005.....	6,090	4	-	1	9	45	285	878	1,161	1,522	1,651	534	-
2004.....	6,075	3	1	1	7	27	279	898	1,138	1,548	1,634	539	-
2003.....	6,335	-	-	-	5	45	333	892	1,103	1,696	1,746	515	-
2002.....	6,549	-	2	3	15	58	409	868	1,123	1,702	1,812	557	-
2001.....	6,367	2	-	-	11	43	431	860	1,051	1,682	1,756	531	-
2000.....	6,214	-	2	1	14	58	364	759	1,016	1,707	1,765	528	-
1999.....	6,108	1	2	1	13	56	322	664	1,029	1,750	1,745	525	-
White, both sexes													
2007 1/.....	12,949	3	4	-	8	75	619	2,570	3,115	2,897	2,747	911	-
2006 1/.....	12,778	3	2	-	9	59	669	2,649	2,904	2,845	2,786	849	3
2005 1/.....	12,798	8	1	2	11	67	761	2,483	2,895	2,940	2,790	840	-
2004 1/.....	12,703	7	1	1	5	62	745	2,569	2,705	2,975	2,840	793	-
2003 1/.....	13,380	1	-	-	5	88	882	2,622	2,722	3,297	2,953	809	1
2002.....	13,386	5	2	5	21	98	996	2,512	2,603	3,225	3,079	838	2
2001.....	13,021	3	-	1	8	96	974	2,468	2,426	3,310	2,901	834	-
2000.....	12,710	-	3	2	15	94	935	2,269	2,415	3,236	2,910	830	1
1999.....	12,579	4	-	1	10	118	912	2,078	2,355	3,375	2,932	794	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other chronic liver disease and cirrhosis (K73-K74)													
White, male													
2007 1/.....	7,534	2	2	-	1	51	428	1,785	2,058	1,550	1,301	356	-
2006 1/.....	7,393	2	2	-	7	34	447	1,885	1,906	1,502	1,267	339	2
2005 1/.....	7,429	5	1	1	4	36	514	1,796	1,900	1,554	1,269	349	-
2004 1/.....	7,342	4	-	-	1	42	532	1,833	1,747	1,569	1,315	299	-
2003 1/.....	7,757	1	-	-	3	56	617	1,917	1,771	1,764	1,304	323	1
2002.....	7,590	5	-	3	8	58	671	1,840	1,625	1,675	1,388	315	2
2001.....	7,359	1	-	1	4	60	624	1,781	1,511	1,787	1,259	331	-
2000.....	7,232	-	1	1	7	56	657	1,667	1,545	1,718	1,245	334	1
1999.....	7,188	4	-	-	4	77	657	1,552	1,496	1,797	1,295	306	-
White, female													
2007 1/.....	5,415	1	2	-	7	24	191	785	1,057	1,347	1,446	555	-
2006 1/.....	5,385	1	-	-	2	25	222	764	998	1,343	1,519	510	1
2005 1/.....	5,369	3	-	1	7	31	247	687	995	1,386	1,521	491	-
2004 1/.....	5,361	3	1	1	4	20	213	736	958	1,406	1,525	494	-
2003 1/.....	5,623	-	-	-	2	32	265	705	951	1,533	1,649	486	-
2002.....	5,796	-	2	2	13	40	325	672	978	1,550	1,691	523	-
2001.....	5,662	2	-	-	4	36	350	687	915	1,523	1,642	503	-
2000.....	5,478	-	2	1	8	38	278	602	870	1,518	1,665	496	-
1999.....	5,391	-	-	1	6	41	255	526	859	1,578	1,637	488	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other chronic liver disease and cirrhosis (K73-K74)													
Black, both sexes													
2007 1/.....	1,355	-	-	-	3	8	76	419	469	229	118	33	-
2006 1/.....	1,304	2	1	-	3	5	113	411	407	199	125	38	-
2005 1/.....	1,361	2	-	-	2	18	87	445	416	203	147	40	1
2004 1/.....	1,365	2	-	-	2	13	146	460	371	219	112	40	-
2003 1/.....	1,399	1	-	-	3	22	138	475	368	247	108	37	-
2002.....	1,397	1	-	1	2	25	148	504	336	220	127	33	-
2001.....	1,427	-	-	1	9	20	177	495	329	239	131	25	1
2000.....	1,357	2	-	1	12	23	164	429	308	264	115	39	-
1999.....	1,367	-	2	1	8	26	174	403	338	244	131	40	-
Black, male													
2007 1/.....	831	-	-	-	3	4	47	263	313	133	60	8	-
2006 1/.....	807	-	1	-	1	1	72	273	267	115	63	14	-
2005 1/.....	855	1	-	-	2	8	60	294	286	117	73	13	1
2004 1/.....	858	2	-	-	-	7	93	324	243	122	55	12	-
2003 1/.....	862	1	-	-	2	11	80	323	256	123	49	17	-
2002.....	825	1	-	-	-	13	80	329	225	122	46	9	-
2001.....	908	-	-	1	2	14	113	359	227	135	50	6	1
2000.....	812	2	-	1	6	10	97	298	203	134	48	13	-
1999.....	824	-	-	1	1	13	117	295	203	129	55	10	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other chronic liver disease and cirrhosis (K73-K74)

Black, female

2007 1/.....	524	-	-	-	-	4	29	156	156	96	58	25	-
2006 1/.....	497	2	-	-	2	4	41	138	140	84	62	24	-
2005 1/.....	506	1	-	-	-	10	27	151	130	86	74	27	-
2004 1/.....	507	-	-	-	2	6	53	136	128	97	57	28	-
2003 1/.....	537	-	-	-	1	11	58	152	112	124	59	20	-
2002.....	572	-	-	1	2	12	68	175	111	98	81	24	-
2001.....	519	-	-	-	7	6	64	136	102	104	81	19	-
2000.....	545	-	-	-	6	13	67	131	105	130	67	26	-
1999.....	543	-	2	-	7	13	57	108	135	115	76	30	-

American Indian 2/, both sexes

2007 1/.....	209	1	-	-	2	6	30	60	62	25	12	11	-
2006 1/.....	170	-	-	-	-	2	26	46	39	34	16	7	-
2005 1/.....	186	-	-	-	2	7	21	59	45	31	17	4	-
2004 1/.....	188	-	-	-	1	1	24	50	57	37	15	3	-
2003 1/.....	167	-	-	-	2	4	19	47	53	23	16	3	-
2002.....	166	-	-	-	-	7	30	36	38	33	19	3	-
2001.....	174	-	-	-	1	5	25	58	39	30	14	2	-
2000.....	195	-	-	-	1	14	38	45	44	36	15	2	-
1999.....	157	-	-	-	-	2	17	51	48	31	7	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other chronic liver disease and cirrhosis (K73-K74)

American Indian 2/, male

2007 1/.....	111	1	-	-	2	3	18	35	32	9	7	4	-
2006 1/.....	84	-	-	-	-	1	16	28	21	12	5	1	-
2005 1/.....	90	-	-	-	1	3	12	29	27	11	5	2	-
2004 1/.....	83	-	-	-	-	-	13	26	24	16	4	-	-
2003 1/.....	76	-	-	-	-	3	11	26	21	9	4	2	-
2002.....	89	-	-	-	-	2	16	26	20	15	9	1	-
2001.....	91	-	-	-	1	4	14	33	21	12	6	-	-
2000.....	93	-	-	-	1	8	22	28	20	7	6	1	-
1999.....	75	-	-	-	-	1	11	27	25	9	2	-	-

American Indian 2/, female

2007 1/.....	98	-	-	-	-	3	12	25	30	16	5	7	-
2006 1/.....	86	-	-	-	-	1	10	18	18	22	11	6	-
2005 1/.....	96	-	-	-	1	4	9	30	18	20	12	2	-
2004 1/.....	105	-	-	-	1	1	11	24	33	21	11	3	-
2003 1/.....	91	-	-	-	2	1	8	21	32	14	12	1	-
2002.....	77	-	-	-	-	5	14	10	18	18	10	2	-
2001.....	83	-	-	-	-	1	11	25	18	18	8	2	-
2000.....	102	-	-	-	-	6	16	17	24	29	9	1	-
1999.....	82	-	-	-	-	1	6	24	23	22	5	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other chronic liver disease and cirrhosis (K73-K74)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	246	-	-	-	-	2	11	37	49	57	70	20	-
2006 1/.....	253	-	-	-	-	5	11	44	44	56	78	15	-
2005 1/.....	257	-	-	-	1	2	13	40	49	48	80	24	-
2004 1/.....	209	-	-	-	-	5	13	25	43	47	57	19	-
2003 1/.....	197	-	-	-	-	1	10	34	38	51	52	11	-
2002.....	187	-	-	-	-	2	9	27	37	54	45	13	-
2001.....	206	-	-	-	-	2	15	35	43	59	38	14	-
2000.....	181	1	-	-	-	5	8	23	38	55	38	13	-
1999.....	198	1	-	-	-	2	13	34	37	57	42	12	-

Asian or Pacific Islander 3/, male

2007 1/.....	126	-	-	-	-	2	9	25	35	22	27	6	-
2006 1/.....	139	-	-	-	-	4	9	36	31	22	29	8	-
2005 1/.....	138	-	-	-	-	2	11	30	31	18	36	10	-
2004 1/.....	107	-	-	-	-	5	11	23	24	23	16	5	-
2003 1/.....	113	-	-	-	-	-	8	20	30	26	26	3	-
2002.....	83	-	-	-	-	1	7	16	21	18	15	5	-
2001.....	103	-	-	-	-	2	9	23	27	22	13	7	-
2000.....	92	1	-	-	-	4	5	14	21	25	14	8	-
1999.....	106	-	-	-	-	1	9	28	25	22	15	6	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other chronic liver disease and cirrhosis (K73-K74)

Asian or Pacific Islander 3/, female

2007 1/.....	120	-	-	-	-	-	2	12	14	35	43	14	-
2006 1/.....	114	-	-	-	-	1	2	8	13	34	49	7	-
2005 1/.....	119	-	-	-	1	-	2	10	18	30	44	14	-
2004 1/.....	102	-	-	-	-	-	2	2	19	24	41	14	-
2003 1/.....	84	-	-	-	-	1	2	14	8	25	26	8	-
2002.....	104	-	-	-	-	1	2	11	16	36	30	8	-
2001.....	103	-	-	-	-	-	6	12	16	37	25	7	-
2000.....	89	-	-	-	-	1	3	9	17	30	24	5	-
1999.....	92	1	-	-	-	1	4	6	12	35	27	6	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

All races, both sexes

2007.....	3,237	1	2	1	7	23	54	132	240	482	1,011	1,284	-
2006.....	3,114	-	-	1	9	20	44	125	270	467	1,022	1,156	-
2005.....	3,072	-	-	4	8	16	42	113	217	448	1,055	1,169	-
2004.....	3,086	-	-	-	8	17	55	138	247	458	1,008	1,155	-
2003.....	2,948	-	-	1	4	25	40	126	182	489	1,054	1,027	-
2002.....	2,979	-	-	-	9	21	54	129	216	504	992	1,054	-
2001.....	2,965	1	-	1	7	19	44	137	242	526	945	1,043	-
2000.....	2,810	-	-	2	9	19	53	109	189	450	926	1,053	-
1999.....	2,830	2	-	1	10	16	59	124	203	475	948	992	-

All races, male

2007.....	1,441	1	2	-	1	8	29	71	131	269	507	422	-
2006.....	1,376	-	-	1	2	8	22	68	133	266	510	366	-
2005.....	1,363	-	-	1	5	8	17	59	118	271	495	389	-
2004.....	1,370	-	-	-	3	10	30	73	138	248	483	385	-
2003.....	1,318	-	-	-	4	11	21	64	106	265	519	328	-
2002.....	1,341	-	-	-	3	11	29	83	122	258	500	335	-
2001.....	1,341	1	-	-	3	9	25	79	142	269	464	349	-
2000.....	1,244	-	-	1	4	7	27	53	110	242	461	339	-
1999.....	1,248	1	-	1	5	4	30	58	120	252	447	330	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

All races, female

2007.....	1,796	-	-	1	6	15	25	61	109	213	504	862	-
2006.....	1,738	-	-	-	7	12	22	57	137	201	512	790	-
2005.....	1,709	-	-	3	3	8	25	54	99	177	560	780	-
2004.....	1,716	-	-	-	5	7	25	65	109	210	525	770	-
2003.....	1,630	-	-	1	-	14	19	62	76	224	535	699	-
2002.....	1,638	-	-	-	6	10	25	46	94	246	492	719	-
2001.....	1,624	-	-	1	4	10	19	58	100	257	481	694	-
2000.....	1,566	-	-	1	5	12	26	56	79	208	465	714	-
1999.....	1,582	1	-	-	5	12	29	66	83	223	501	662	-

White, both sexes

2007 1/.....	2,839	1	2	1	6	18	37	96	190	409	891	1,188	-
2006 1/.....	2,671	-	-	1	7	16	31	87	207	361	900	1,061	-
2005 1/.....	2,655	-	-	4	6	13	28	74	175	359	936	1,060	-
2004 1/.....	2,672	-	-	-	6	10	45	95	192	374	888	1,062	-
2003 1/.....	2,573	-	-	1	3	18	28	85	142	421	942	933	-
2002.....	2,614	-	-	-	7	16	43	93	177	431	871	976	-
2001.....	2,567	1	-	1	5	15	33	99	191	433	835	954	-
2000.....	2,457	-	-	1	7	10	38	75	151	366	827	982	-
1999.....	2,487	-	-	1	6	13	37	97	153	398	864	918	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)													
White, male													
2007 1/.....	1,280	1	2	-	1	7	19	54	112	236	458	390	-
2006 1/.....	1,195	-	-	1	1	6	19	51	100	213	457	347	-
2005 1/.....	1,193	-	-	1	4	8	13	37	96	223	453	358	-
2004 1/.....	1,205	-	-	-	3	7	24	49	113	211	435	363	-
2003 1/.....	1,150	-	-	-	3	6	13	47	82	232	466	301	-
2002.....	1,179	-	-	-	3	8	24	59	101	226	447	311	-
2001.....	1,157	1	-	-	2	6	20	57	112	225	420	314	-
2000.....	1,111	-	-	1	3	5	22	39	92	206	423	320	-
1999.....	1,101	-	-	1	4	2	20	45	91	217	418	303	-
White, female													
2007 1/.....	1,559	-	-	1	5	11	18	42	78	173	433	798	-
2006 1/.....	1,476	-	-	-	6	10	12	36	107	148	443	714	-
2005 1/.....	1,462	-	-	3	2	5	15	37	79	136	483	702	-
2004 1/.....	1,467	-	-	-	3	3	21	46	79	163	453	699	-
2003 1/.....	1,423	-	-	1	-	12	15	38	60	189	476	632	-
2002.....	1,435	-	-	-	4	8	19	34	76	205	424	665	-
2001.....	1,410	-	-	1	3	9	13	42	79	208	415	640	-
2000.....	1,346	-	-	-	4	5	16	36	59	160	404	662	-
1999.....	1,386	-	-	-	2	11	17	52	62	181	446	615	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

Black, both sexes

2007 1/.....	299	-	-	-	1	5	11	27	43	52	88	72	-
2006 1/.....	324	-	-	-	2	2	13	32	46	71	92	66	-
2005 1/.....	325	-	-	-	2	3	12	31	35	72	93	77	-
2004 1/.....	317	-	-	-	-	4	9	37	46	61	91	69	-
2003 1/.....	286	-	-	-	1	7	11	35	33	50	78	71	-
2002.....	273	-	-	-	1	5	9	29	29	56	94	50	-
2001.....	310	-	-	-	2	4	11	31	42	78	78	64	-
2000.....	280	-	-	1	2	8	14	28	29	65	76	57	-
1999.....	257	2	-	-	4	3	20	21	35	59	65	48	-

Black, male

2007 1/.....	118	-	-	-	-	1	7	11	16	22	38	23	-
2006 1/.....	117	-	-	-	1	1	3	14	23	28	35	12	-
2005 1/.....	135	-	-	-	1	-	4	17	17	38	37	21	-
2004 1/.....	122	-	-	-	-	2	5	22	21	26	32	14	-
2003 1/.....	127	-	-	-	1	5	8	15	18	23	37	20	-
2002.....	110	-	-	-	-	3	4	18	17	19	38	11	-
2001.....	130	-	-	-	1	3	5	18	24	34	21	24	-
2000.....	96	-	-	-	1	2	4	11	14	25	27	12	-
1999.....	102	1	-	-	1	2	9	9	19	25	23	13	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Cholelithiasis and other disorders of gallbladder (K80-K82)

Black, female

2007 1/.....	181	-	-	-	1	4	4	16	27	30	50	49	-
2006 1/.....	207	-	-	-	1	1	10	18	23	43	57	54	-
2005 1/.....	190	-	-	-	1	3	8	14	18	34	56	56	-
2004 1/.....	195	-	-	-	-	2	4	15	25	35	59	55	-
2003 1/.....	159	-	-	-	-	2	3	20	15	27	41	51	-
2002.....	163	-	-	-	1	2	5	11	12	37	56	39	-
2001.....	180	-	-	-	1	1	6	13	18	44	57	40	-
2000.....	184	-	-	1	1	6	10	17	15	40	49	45	-
1999.....	155	1	-	-	3	1	11	12	16	34	42	35	-

American Indian 2/, both sexes

2007 1/.....	23	-	-	-	-	-	3	4	2	3	6	5	-
2006 1/.....	24	-	-	-	-	1	-	1	5	8	3	6	-
2005 1/.....	27	-	-	-	-	-	1	3	5	4	7	7	-
2004 1/.....	27	-	-	-	-	1	-	1	5	5	7	8	-
2003 1/.....	29	-	-	-	-	-	1	2	4	6	9	7	-
2002.....	26	-	-	-	1	-	-	4	4	6	5	6	-
2001.....	15	-	-	-	-	-	-	2	2	1	6	4	-
2000.....	20	-	-	-	-	1	-	2	2	2	8	5	-
1999.....	24	-	-	-	-	-	2	3	3	6	4	6	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Cholelithiasis and other disorders of gallbladder (K80-K82)

American Indian 2/, male

2007 1/.....	10	-	-	-	-	-	1	2	1	1	2	3	-
2006 1/.....	14	-	-	-	-	-	-	-	3	7	2	2	-
2005 1/.....	10	-	-	-	-	-	-	1	3	2	2	2	-
2004 1/.....	12	-	-	-	-	-	-	1	3	2	3	3	-
2003 1/.....	12	-	-	-	-	-	-	-	4	4	4	-	-
2002.....	13	-	-	-	-	-	-	4	1	5	2	1	-
2001.....	11	-	-	-	-	-	-	2	2	1	3	3	-
2000.....	7	-	-	-	-	-	-	1	1	1	3	1	-
1999.....	11	-	-	-	-	-	1	3	2	3	1	1	-

American Indian 2/, female

2007 1/.....	13	-	-	-	-	-	2	2	1	2	4	2	-
2006 1/.....	10	-	-	-	-	1	-	1	2	1	1	4	-
2005 1/.....	17	-	-	-	-	-	1	2	2	2	5	5	-
2004 1/.....	15	-	-	-	-	1	-	-	2	3	4	5	-
2003 1/.....	17	-	-	-	-	-	1	2	-	2	5	7	-
2002.....	13	-	-	-	1	-	-	-	3	1	3	5	-
2001.....	4	-	-	-	-	-	-	-	-	-	3	1	-
2000.....	13	-	-	-	-	1	-	1	1	1	5	4	-
1999.....	13	-	-	-	-	-	1	-	1	3	3	5	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	76	-	-	-	-	-	3	5	5	18	26	19	-
2006 1/.....	95	-	-	-	-	1	-	5	12	27	27	23	-
2005 1/.....	65	-	-	-	-	-	1	5	2	13	19	25	-
2004 1/.....	70	-	-	-	2	2	1	5	4	18	22	16	-
2003 1/.....	60	-	-	-	-	-	-	4	3	12	25	16	-
2002.....	66	-	-	-	-	-	2	3	6	11	22	22	-
2001.....	73	-	-	-	-	-	-	5	7	14	26	21	-
2000.....	53	-	-	-	-	-	1	4	7	17	15	9	-
1999.....	62	-	-	-	-	-	-	3	12	12	15	20	-

Asian or Pacific Islander 3/, male

2007 1/.....	33	-	-	-	-	-	2	4	2	10	9	6	-
2006 1/.....	50	-	-	-	-	1	-	3	7	18	16	5	-
2005 1/.....	25	-	-	-	-	-	-	4	2	8	3	8	-
2004 1/.....	31	-	-	-	-	1	1	1	1	9	13	5	-
2003 1/.....	29	-	-	-	-	-	-	2	2	6	12	7	-
2002.....	39	-	-	-	-	-	1	2	3	8	13	12	-
2001.....	43	-	-	-	-	-	-	2	4	9	20	8	-
2000.....	30	-	-	-	-	-	1	2	3	10	8	6	-
1999.....	34	-	-	-	-	-	-	1	8	7	5	13	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

Asian or Pacific Islander 3/, female

2007 1/.....	43	-	-	-	-	-	1	1	3	8	17	13	-
2006 1/.....	45	-	-	-	-	-	-	2	5	9	11	18	-
2005 1/.....	40	-	-	-	-	-	1	1	-	5	16	17	-
2004 1/.....	39	-	-	-	2	1	-	4	3	9	9	11	-
2003 1/.....	31	-	-	-	-	-	-	2	1	6	13	9	-
2002.....	27	-	-	-	-	-	1	1	3	3	9	10	-
2001.....	30	-	-	-	-	-	-	3	3	5	6	13	-
2000.....	23	-	-	-	-	-	-	2	4	7	7	3	-
1999.....	28	-	-	-	-	-	-	2	4	5	10	7	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)													
All races, both sexes													
2007.....	46,448	144	22	24	86	261	754	2,233	4,440	7,752	14,711	16,021	-
2006.....	45,344	162	14	18	99	276	785	2,245	4,368	7,453	14,536	15,388	-
2005.....	43,901	159	16	23	91	285	742	2,028	4,141	7,320	14,403	14,693	-
2004.....	42,480	174	14	21	87	246	801	2,067	3,963	7,119	14,058	13,928	2
2003.....	42,453	181	14	22	95	282	796	2,001	3,806	7,345	14,093	13,816	2
2002.....	40,974	173	14	21	84	269	749	1,893	3,455	7,164	13,896	13,256	-
2001.....	39,480	133	16	18	92	225	768	1,822	3,284	7,356	13,097	12,668	1
2000.....	37,251	162	14	25	88	244	740	1,651	3,100	6,990	12,458	11,777	2
1999.....	35,525	166	15	31	72	252	707	1,480	2,864	6,841	11,927	11,170	-
All races, male													
2007.....	22,616	78	11	13	42	141	428	1,260	2,478	4,044	7,435	6,686	-
2006.....	22,094	101	9	12	56	157	458	1,338	2,442	3,909	7,246	6,366	-
2005.....	21,268	88	12	10	45	153	423	1,237	2,352	3,733	7,177	6,038	-
2004.....	20,370	110	10	11	50	113	451	1,208	2,120	3,677	7,007	5,612	1
2003.....	20,481	106	7	9	59	147	459	1,193	2,065	3,785	7,049	5,600	2
2002.....	19,695	109	7	12	41	129	407	1,086	1,824	3,802	6,920	5,358	-
2001.....	18,852	76	7	6	45	112	414	1,053	1,714	3,826	6,485	5,114	-
2000.....	17,811	103	7	17	43	129	429	982	1,651	3,564	6,213	4,671	2
1999.....	17,016	90	9	18	33	125	402	846	1,530	3,504	6,031	4,428	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

All races, female

2007.....	23,832	66	11	11	44	120	326	973	1,962	3,708	7,276	9,335	-
2006.....	23,250	61	5	6	43	119	327	907	1,926	3,544	7,290	9,022	-
2005.....	22,633	71	4	13	46	132	319	791	1,789	3,587	7,226	8,655	-
2004.....	22,110	64	4	10	37	133	350	859	1,843	3,442	7,051	8,316	1
2003.....	21,972	75	7	13	36	135	337	808	1,741	3,560	7,044	8,216	-
2002.....	21,279	64	7	9	43	140	342	807	1,631	3,362	6,976	7,898	-
2001.....	20,628	57	9	12	47	113	354	769	1,570	3,530	6,612	7,554	1
2000.....	19,440	59	7	8	45	115	311	669	1,449	3,426	6,245	7,106	-
1999.....	18,509	76	6	13	39	127	305	634	1,334	3,337	5,896	6,742	-

White, both sexes

2007 1/.....	36,871	94	14	18	57	141	406	1,346	2,892	5,655	12,282	13,966	-
2006 1/.....	35,793	112	12	13	66	158	411	1,334	2,875	5,357	12,060	13,395	-
2005 1/.....	34,806	96	13	18	60	152	401	1,221	2,663	5,343	11,995	12,844	-
2004 1/.....	33,691	97	10	17	53	128	465	1,186	2,552	5,254	11,739	12,188	2
2003 1/.....	33,707	117	10	15	56	157	417	1,149	2,517	5,378	11,768	12,122	1
2002.....	32,615	112	10	15	42	152	417	1,082	2,248	5,303	11,659	11,575	-
2001.....	31,345	77	13	10	61	125	432	1,043	2,101	5,458	10,936	11,088	1
2000.....	29,598	87	8	20	63	136	412	936	2,009	5,144	10,470	10,311	2
1999.....	28,041	102	13	22	42	137	381	825	1,868	5,087	9,934	9,630	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

White, male

2007 1/.....	18,242	48	6	8	27	76	242	788	1,655	3,025	6,400	5,967	-
2006 1/.....	17,715	72	8	8	43	95	252	820	1,659	2,918	6,157	5,683	-
2005 1/.....	17,137	51	9	7	32	82	238	768	1,545	2,867	6,139	5,399	-
2004 1/.....	16,349	61	7	8	29	55	272	692	1,411	2,792	6,029	4,992	1
2003 1/.....	16,408	71	5	7	36	80	253	683	1,378	2,848	6,033	5,013	1
2002.....	15,850	70	4	9	22	76	229	622	1,186	2,914	5,972	4,746	-
2001.....	15,241	45	4	5	30	55	256	613	1,144	2,950	5,565	4,574	-
2000.....	14,385	58	5	14	33	73	243	558	1,106	2,769	5,353	4,171	2
1999.....	13,632	62	7	15	19	70	218	474	1,043	2,711	5,119	3,894	-

White, female

2007 1/.....	18,629	46	8	10	30	65	164	558	1,237	2,630	5,882	7,999	-
2006 1/.....	18,078	40	4	5	23	63	159	514	1,216	2,439	5,903	7,712	-
2005 1/.....	17,669	45	4	11	28	70	163	453	1,118	2,476	5,856	7,445	-
2004 1/.....	17,342	36	3	9	24	73	193	494	1,141	2,462	5,710	7,196	1
2003 1/.....	17,299	46	5	8	20	77	164	466	1,139	2,530	5,735	7,109	-
2002.....	16,765	42	6	6	20	76	188	460	1,062	2,389	5,687	6,829	-
2001.....	16,104	32	9	5	31	70	176	430	957	2,508	5,371	6,514	1
2000.....	15,213	29	3	6	30	63	169	378	903	2,375	5,117	6,140	-
1999.....	14,409	40	6	7	23	67	163	351	825	2,376	4,815	5,736	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Black, both sexes

2007 1/.....	8,392	47	6	3	28	113	330	823	1,391	1,841	2,059	1,751	-
2006 1/.....	8,397	43	2	2	30	107	336	841	1,346	1,846	2,138	1,706	-
2005 1/.....	8,075	58	3	4	29	121	313	754	1,346	1,756	2,103	1,588	-
2004 1/.....	7,834	67	4	3	31	103	317	820	1,291	1,653	2,017	1,528	-
2003 1/.....	7,855	60	3	7	34	119	356	785	1,183	1,765	2,032	1,510	1
2002.....	7,488	55	4	5	36	109	309	752	1,082	1,690	1,959	1,487	-
2001.....	7,274	53	3	7	27	95	309	706	1,054	1,711	1,910	1,399	-
2000.....	6,911	71	5	4	24	100	306	669	991	1,653	1,789	1,299	-
1999.....	6,711	58	1	8	30	103	310	617	887	1,569	1,769	1,359	-

Black, male

2007 1/.....	3,772	29	4	2	14	60	177	435	730	877	863	581	-
2006 1/.....	3,812	24	1	2	11	57	183	480	717	863	927	547	-
2005 1/.....	3,645	36	3	2	11	61	165	443	750	756	899	519	-
2004 1/.....	3,537	41	3	2	19	51	167	478	644	780	836	516	-
2003 1/.....	3,642	32	1	2	20	64	192	471	634	856	872	497	1
2002.....	3,427	35	3	2	15	49	163	432	578	806	820	524	-
2001.....	3,186	29	3	-	12	56	141	400	514	787	793	451	-
2000.....	3,074	42	2	2	10	50	175	398	489	716	765	425	-
1999.....	3,008	25	1	3	14	47	173	354	429	700	813	449	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Black, female

2007 1/.....	4,620	18	2	1	14	53	153	388	661	964	1,196	1,170	-
2006 1/.....	4,585	19	1	-	19	50	153	361	629	983	1,211	1,159	-
2005 1/.....	4,430	22	-	2	18	60	148	311	596	1,000	1,204	1,069	-
2004 1/.....	4,297	26	1	1	12	52	150	342	647	873	1,181	1,012	-
2003 1/.....	4,213	28	2	5	14	55	164	314	549	909	1,160	1,013	-
2002.....	4,061	20	1	3	21	60	146	320	504	884	1,139	963	-
2001.....	4,088	24	-	7	15	39	168	306	540	924	1,117	948	-
2000.....	3,837	29	3	2	14	50	131	271	502	937	1,024	874	-
1999.....	3,703	33	-	5	16	56	137	263	458	869	956	910	-

American Indian 2/, both sexes

2007 1/.....	292	2	1	1	-	1	11	23	57	74	68	54	-
2006 1/.....	288	1	-	1	3	7	17	33	52	81	52	41	-
2005 1/.....	265	2	-	1	1	6	15	20	47	65	62	46	-
2004 1/.....	247	1	-	1	2	10	9	29	39	60	59	37	-
2003 1/.....	255	1	-	-	3	2	11	32	42	46	76	42	-
2002.....	222	2	-	1	2	6	10	26	43	56	46	30	-
2001.....	236	2	-	1	2	1	13	32	51	54	52	28	-
2000.....	215	1	-	-	-	3	12	22	39	65	46	27	-
1999.....	194	-	-	-	-	4	7	18	36	55	42	32	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

American Indian 2/, male

2007 1/.....	126	1	1	1	-	1	5	11	26	35	23	22	-
2006 1/.....	135	1	-	1	2	4	10	18	21	36	24	18	-
2005 1/.....	105	1	-	1	1	5	9	6	17	28	18	19	-
2004 1/.....	112	1	-	1	1	3	4	17	19	28	22	16	-
2003 1/.....	127	1	-	-	1	1	6	18	21	19	43	17	-
2002.....	98	2	-	1	2	3	8	14	20	19	20	9	-
2001.....	105	2	-	1	2	-	8	19	16	19	24	14	-
2000.....	98	-	-	-	-	2	7	11	21	28	16	13	-
1999.....	79	-	-	-	-	4	6	9	17	23	12	8	-

American Indian 2/, female

2007 1/.....	166	1	-	-	-	-	6	12	31	39	45	32	-
2006 1/.....	153	-	-	-	1	3	7	15	31	45	28	23	-
2005 1/.....	160	1	-	-	-	1	6	14	30	37	44	27	-
2004 1/.....	135	-	-	-	1	7	5	12	20	32	37	21	-
2003 1/.....	128	-	-	-	2	1	5	14	21	27	33	25	-
2002.....	124	-	-	-	-	3	2	12	23	37	26	21	-
2001.....	131	-	-	-	-	1	5	13	35	35	28	14	-
2000.....	117	1	-	-	-	1	5	11	18	37	30	14	-
1999.....	115	-	-	-	-	-	1	9	19	32	30	24	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	893	1	1	2	1	6	7	41	100	182	302	250	-
2006 1/.....	866	6	-	2	-	4	21	37	95	169	286	246	-
2005 1/.....	755	3	-	-	1	6	13	33	85	156	243	215	-
2004 1/.....	708	9	-	-	1	5	10	32	81	152	243	175	-
2003 1/.....	636	3	1	-	2	4	12	35	64	156	217	142	-
2002.....	649	4	-	-	4	2	13	33	82	115	232	164	-
2001.....	625	1	-	-	2	4	14	41	78	133	199	153	-
2000.....	527	3	1	1	1	5	10	24	61	128	153	140	-
1999.....	579	6	1	1	-	8	9	20	73	130	182	149	-

Asian or Pacific Islander 3/, male

2007 1/.....	476	-	-	2	1	4	4	26	67	107	149	116	-
2006 1/.....	432	4	-	1	-	1	13	20	45	92	138	118	-
2005 1/.....	381	-	-	-	1	5	11	20	40	82	121	101	-
2004 1/.....	372	7	-	-	1	4	8	21	46	77	120	88	-
2003 1/.....	304	2	1	-	2	2	8	21	32	62	101	73	-
2002.....	320	2	-	-	2	1	7	18	40	63	108	79	-
2001.....	320	-	-	-	1	1	9	21	40	70	103	75	-
2000.....	254	3	-	1	-	4	4	15	35	51	79	62	-
1999.....	297	3	1	-	-	4	5	9	41	70	87	77	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Asian or Pacific Islander 3/, female

2007 1/.....	417	1	1	-	-	2	3	15	33	75	153	134	-
2006 1/.....	434	2	-	1	-	3	8	17	50	77	148	128	-
2005 1/.....	374	3	-	-	-	1	2	13	45	74	122	114	-
2004 1/.....	336	2	-	-	-	1	2	11	35	75	123	87	-
2003 1/.....	332	1	-	-	-	2	4	14	32	94	116	69	-
2002.....	329	2	-	-	2	1	6	15	42	52	124	85	-
2001.....	305	1	-	-	1	3	5	20	38	63	96	78	-
2000.....	273	-	1	-	1	1	6	9	26	77	74	78	-
1999.....	282	3	-	1	-	4	4	11	32	60	95	72	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

All races, both sexes

2007.....	206	5	3	3	1	7	6	13	22	35	62	49	-
2006.....	138	5	4	4	4	4	4	11	10	17	41	34	-
2005.....	137	6	2	2	-	3	4	6	15	17	43	39	-
2004.....	157	7	5	1	2	1	2	16	9	27	49	38	-
2003.....	148	8	2	1	4	2	4	7	14	30	50	26	-
2002.....	166	6	5	2	3	1	4	9	16	27	51	42	-
2001.....	156	8	4	1	-	3	8	6	13	27	48	38	-
2000.....	177	8	3	2	1	4	3	16	10	29	52	49	-
1999.....	182	3	2	5	2	5	5	11	16	37	62	34	-

All races, male

2007.....	111	4	-	1	-	5	3	10	13	18	36	21	-
2006.....	70	4	2	3	1	3	1	9	6	7	18	16	-
2005.....	67	2	2	-	-	2	4	4	8	11	19	15	-
2004.....	73	2	4	-	2	1	1	10	4	14	24	11	-
2003.....	65	7	1	-	4	1	2	3	7	15	20	5	-
2002.....	70	3	2	2	1	1	-	5	11	14	17	14	-
2001.....	73	4	3	-	-	2	6	5	9	10	19	15	-
2000.....	91	3	2	2	1	3	1	11	8	12	26	22	-
1999.....	90	3	2	2	2	4	2	6	6	17	32	14	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

All races, female

2007.....	95	1	3	2	1	2	3	3	9	17	26	28	-
2006.....	68	1	2	1	3	1	3	2	4	10	23	18	-
2005.....	70	4	-	2	-	1	-	2	7	6	24	24	-
2004.....	84	5	1	1	-	-	1	6	5	13	25	27	-
2003.....	83	1	1	1	-	1	2	4	7	15	30	21	-
2002.....	96	3	3	-	2	-	4	4	5	13	34	28	-
2001.....	83	4	1	1	-	1	2	1	4	17	29	23	-
2000.....	86	5	1	-	-	1	2	5	2	17	26	27	-
1999.....	92	-	-	3	-	1	3	5	10	20	30	20	-

White, both sexes

2007 1/.....	168	4	2	2	1	4	5	9	17	27	53	44	-
2006 1/.....	111	3	4	3	2	3	3	5	8	12	37	31	-
2005 1/.....	114	4	2	1	-	-	3	4	15	15	38	32	-
2004 1/.....	127	3	4	1	1	-	-	13	8	22	43	32	-
2003 1/.....	120	6	1	1	3	1	3	7	9	26	42	21	-
2002.....	130	5	4	2	1	-	3	5	14	21	42	33	-
2001.....	130	7	3	-	-	1	8	5	11	21	43	31	-
2000.....	141	6	1	1	1	3	1	6	10	22	46	44	-
1999.....	144	3	2	4	1	2	3	7	9	32	52	29	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)													
White, male													
2007 1/.....	91	4	-	-	-	3	3	7	10	13	32	19	-
2006 1/.....	55	3	2	2	1	2	1	4	5	5	15	15	-
2005 1/.....	60	1	2	-	-	-	3	3	8	10	19	14	-
2004 1/.....	59	-	4	-	1	-	-	8	3	14	20	9	-
2003 1/.....	47	5	-	-	3	-	2	3	3	12	14	5	-
2002.....	56	2	1	2	1	-	-	4	10	11	14	11	-
2001.....	60	3	2	-	-	-	6	5	7	6	19	12	-
2000.....	72	3	1	1	1	2	1	5	8	9	23	18	-
1999.....	67	3	2	2	1	1	1	4	3	14	26	10	-
White, female													
2007 1/.....	77	-	2	2	1	1	2	2	7	14	21	25	-
2006 1/.....	56	-	2	1	1	1	2	1	3	7	22	16	-
2005 1/.....	54	3	-	1	-	-	-	1	7	5	19	18	-
2004 1/.....	68	3	-	1	-	-	-	5	5	8	23	23	-
2003 1/.....	73	1	1	1	-	1	1	4	6	14	28	16	-
2002.....	74	3	3	-	-	-	3	1	4	10	28	22	-
2001.....	70	4	1	-	-	1	2	-	4	15	24	19	-
2000.....	69	3	-	-	-	1	-	1	2	13	23	26	-
1999.....	77	-	-	2	-	1	2	3	6	18	26	19	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

Black, both sexes

2007 1/.....	35	1	-	1	-	3	1	4	5	8	7	5	-
2006 1/.....	20	1	-	-	2	1	1	6	1	4	3	1	-
2005 1/.....	18	1	-	1	-	3	1	2	-	2	4	4	-
2004 1/.....	24	2	1	-	1	-	2	3	1	5	5	4	-
2003 1/.....	18	1	-	-	-	1	1	-	3	4	6	2	-
2002.....	29	1	1	-	2	1	1	4	1	5	6	7	-
2001.....	21	-	1	1	-	2	-	1	1	6	5	4	-
2000.....	30	2	2	-	-	1	2	10	-	6	5	2	-
1999.....	28	-	-	1	1	2	2	4	5	3	6	4	-

Black, male

2007 1/.....	18	-	-	1	-	2	-	3	3	5	2	2	-
2006 1/.....	11	1	-	-	-	1	-	5	-	1	2	1	-
2005 1/.....	6	-	-	-	-	2	1	1	-	1	-	1	-
2004 1/.....	10	1	-	-	1	-	1	2	1	-	3	1	-
2003 1/.....	12	1	-	-	-	1	-	-	2	3	5	-	-
2002.....	11	1	1	-	-	1	-	1	1	2	2	2	-
2001.....	9	-	1	-	-	2	-	-	1	4	-	1	-
2000.....	13	-	1	-	-	1	-	6	-	2	2	1	-
1999.....	16	-	-	-	1	2	1	2	2	2	3	3	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

Black, female

2007 1/.....	17	1	-	-	-	1	1	1	2	3	5	3	-
2006 1/.....	9	-	-	-	2	-	1	1	1	3	1	-	-
2005 1/.....	12	1	-	1	-	1	-	1	-	1	4	3	-
2004 1/.....	14	1	1	-	-	-	1	1	-	5	2	3	-
2003 1/.....	6	-	-	-	-	-	1	-	1	1	1	2	-
2002.....	18	-	-	-	2	-	1	3	-	3	4	5	-
2001.....	12	-	-	1	-	-	-	1	-	2	5	3	-
2000.....	17	2	1	-	-	-	2	4	-	4	3	1	-
1999.....	12	-	-	1	-	-	1	2	3	1	3	1	-

American Indian 2/, both sexes

2007 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2006 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2005 1/.....	1	1	-	-	-	-	-	-	-	-	-	-	-
2004 1/.....	2	1	-	-	-	1	-	-	-	-	-	-	-
2003 1/.....	1	-	-	-	-	-	-	-	-	-	-	1	-
2002.....	2	-	-	-	-	-	-	-	-	-	1	1	-
2001.....	2	1	-	-	-	-	-	-	1	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	1	-
1999.....	3	-	-	-	-	1	-	-	1	-	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

American Indian 2/, male

2007 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2006 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2005 1/.....	1	1	-	-	-	-	-	-	-	-	-	-	-
2004 1/.....	2	1	-	-	-	1	-	-	-	-	-	-	-
2003 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	1	-
2001.....	2	1	-	-	-	-	-	1	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	1	-
1999.....	2	-	-	-	-	1	-	-	1	-	-	-	-

American Indian 2/, female

2007 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2006 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2005 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 1/.....	1	-	-	-	-	-	-	-	-	-	-	1	-
2002.....	1	-	-	-	-	-	-	-	-	-	1	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	-	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	3	-	1	-	-	-	-	-	-	-	2	-	-
2006 1/.....	7	1	-	1	-	-	-	-	1	1	1	2	-
2005 1/.....	4	-	-	-	-	-	-	-	-	-	1	3	-
2004 1/.....	4	1	-	-	-	-	-	-	-	-	1	2	-
2003 1/.....	9	1	1	-	1	-	-	-	2	-	2	2	-
2002.....	5	-	-	-	-	-	-	-	1	1	2	1	-
2001.....	3	-	-	-	-	-	-	-	-	-	-	3	-
2000.....	5	-	-	1	-	-	-	-	-	1	1	2	-
1999.....	7	-	-	-	-	-	-	-	1	2	3	1	-

Asian or Pacific Islander 3/, male

2007 1/.....	2	-	-	-	-	-	-	-	-	-	2	-	-
2006 1/.....	4	-	-	1	-	-	-	-	1	1	1	-	-
2005 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 1/.....	2	-	-	-	-	-	-	-	-	-	1	1	-
2003 1/.....	6	1	1	-	1	-	-	-	2	-	1	-	-
2002.....	2	-	-	-	-	-	-	-	-	1	1	-	-
2001.....	2	-	-	-	-	-	-	-	-	-	-	2	-
2000.....	5	-	-	1	-	-	-	-	-	1	1	2	-
1999.....	5	-	-	-	-	-	-	-	-	1	3	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

Asian or Pacific Islander 3/, female

2007 1/.....	1	-	1	-	-	-	-	-	-	-	-	-	-
2006 1/.....	3	1	-	-	-	-	-	-	-	-	-	2	-
2005 1/.....	4	-	-	-	-	-	-	-	-	-	1	3	-
2004 1/.....	2	1	-	-	-	-	-	-	-	-	-	1	-
2003 1/.....	3	-	-	-	-	-	-	-	-	-	1	2	-
2002.....	3	-	-	-	-	-	-	-	1	-	1	1	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	1	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	-	-	-	-	-	-	-	1	1	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

All races, both sexes

2007.....	2,958	1	2	3	8	17	51	126	208	384	928	1,230	-
2006.....	1,841	3	-	1	6	21	34	82	149	260	575	710	-
2005.....	867	2	1	2	11	21	30	48	93	132	252	275	-
2004.....	553	1	-	2	5	21	30	39	64	88	168	135	-
2003.....	537	3	-	-	8	14	20	45	50	102	165	130	-
2002.....	553	2	-	3	4	23	26	53	52	85	178	127	-
2001.....	511	-	2	1	8	12	27	41	40	111	147	122	-
2000.....	556	1	-	4	9	17	34	39	59	103	172	118	-
1999.....	589	-	1	1	7	15	28	44	60	101	200	132	-

All races, male

2007.....	1,452	-	-	1	5	11	24	73	126	217	463	532	-
2006.....	897	3	-	1	3	14	19	44	89	132	302	290	-
2005.....	444	-	1	-	4	15	23	32	53	67	129	120	-
2004.....	275	1	-	-	1	6	17	20	40	46	84	60	-
2003.....	269	1	-	-	5	5	14	35	22	48	80	59	-
2002.....	278	1	-	1	3	13	14	25	27	49	92	53	-
2001.....	264	-	1	-	5	1	13	24	25	63	84	48	-
2000.....	284	-	-	1	2	11	20	27	31	54	83	55	-
1999.....	304	-	-	1	6	10	18	25	32	45	107	60	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

All races, female

2007.....	1,506	1	2	2	3	6	27	53	82	167	465	698	-
2006.....	944	-	-	-	3	7	15	38	60	128	273	420	-
2005.....	423	2	-	2	7	6	7	16	40	65	123	155	-
2004.....	278	-	-	2	4	15	13	19	24	42	84	75	-
2003.....	268	2	-	-	3	9	6	10	28	54	85	71	-
2002.....	275	1	-	2	1	10	12	28	25	36	86	74	-
2001.....	247	-	1	1	3	11	14	17	15	48	63	74	-
2000.....	272	1	-	3	7	6	14	12	28	49	89	63	-
1999.....	285	-	1	-	1	5	10	19	28	56	93	72	-

White, both sexes

2007 1/.....	2,350	1	1	2	7	11	30	80	145	279	754	1,040	-
2006 1/.....	1,421	3	-	1	5	9	14	47	98	182	466	596	-
2005 1/.....	686	1	-	2	8	13	18	28	65	98	215	238	-
2004 1/.....	447	1	-	2	3	13	20	26	47	72	144	119	-
2003 1/.....	430	3	-	-	3	8	16	31	42	79	137	111	-
2002.....	450	-	-	2	3	15	22	34	39	79	149	107	-
2001.....	417	-	2	-	7	9	15	21	27	93	129	114	-
2000.....	439	-	-	3	7	11	20	26	43	85	150	94	-
1999.....	467	-	1	1	4	11	15	32	43	78	173	109	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

White, male

2007 1/.....	1,179	-	-	-	4	7	15	48	91	165	390	459	-
2006 1/.....	692	3	-	1	3	6	10	26	55	97	249	242	-
2005 1/.....	350	-	-	-	4	9	13	20	37	54	108	105	-
2004 1/.....	221	1	-	-	-	5	12	15	28	36	73	51	-
2003 1/.....	214	1	-	-	1	3	12	25	19	38	65	50	-
2002.....	233	-	-	-	2	10	11	16	23	45	78	48	-
2001.....	216	-	1	-	4	1	8	10	16	54	75	47	-
2000.....	226	-	-	1	1	7	12	18	23	45	75	44	-
1999.....	244	-	-	1	3	8	10	19	24	35	94	50	-

White, female

2007 1/.....	1,171	1	1	2	3	4	15	32	54	114	364	581	-
2006 1/.....	729	-	-	-	2	3	4	21	43	85	217	354	-
2005 1/.....	336	1	-	2	4	4	5	8	28	44	107	133	-
2004 1/.....	226	-	-	2	3	8	8	11	19	36	71	68	-
2003 1/.....	216	2	-	-	2	5	4	6	23	41	72	61	-
2002.....	217	-	-	2	1	5	11	18	16	34	71	59	-
2001.....	201	-	1	-	3	8	7	11	11	39	54	67	-
2000.....	213	-	-	2	6	4	8	8	20	40	75	50	-
1999.....	223	-	1	-	1	3	5	13	19	43	79	59	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

Black, both sexes

2007 1/.....	520	-	1	-	1	6	20	43	56	93	145	155	-
2006 1/.....	349	-	-	-	1	10	12	32	49	70	84	91	-
2005 1/.....	161	1	1	-	3	6	10	17	27	28	34	34	-
2004 1/.....	90	-	-	-	2	8	10	10	15	15	18	12	-
2003 1/.....	76	-	-	-	4	4	4	12	6	15	19	12	-
2002.....	82	2	-	-	1	7	3	18	8	5	21	17	-
2001.....	69	-	-	1	-	2	8	15	9	14	13	7	-
2000.....	90	1	-	1	2	4	10	12	12	14	19	15	-
1999.....	94	-	-	-	3	2	11	10	11	18	20	19	-

Black, male

2007 1/.....	229	-	-	-	1	4	9	23	32	47	60	53	-
2006 1/.....	169	-	-	-	-	8	4	16	33	32	39	37	-
2005 1/.....	81	-	1	-	-	4	8	9	16	9	21	13	-
2004 1/.....	42	-	-	-	1	1	5	4	10	9	7	5	-
2003 1/.....	39	-	-	-	3	1	2	9	2	7	10	5	-
2002.....	37	1	-	-	1	3	2	8	3	4	11	4	-
2001.....	37	-	-	-	-	-	5	12	7	7	5	1	-
2000.....	44	-	-	-	1	2	5	8	7	8	6	7	-
1999.....	47	-	-	-	3	1	6	6	6	7	10	8	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

Black, female

2007 1/.....	291	-	1	-	-	2	11	20	24	46	85	102	-
2006 1/.....	180	-	-	-	1	2	8	16	16	38	45	54	-
2005 1/.....	80	1	-	-	3	2	2	8	11	19	13	21	-
2004 1/.....	48	-	-	-	1	7	5	6	5	6	11	7	-
2003 1/.....	37	-	-	-	1	3	2	3	4	8	9	7	-
2002.....	45	1	-	-	-	4	1	10	5	1	10	13	-
2001.....	32	-	-	1	-	2	3	3	2	7	8	6	-
2000.....	46	1	-	1	1	2	5	4	5	6	13	8	-
1999.....	47	-	-	-	-	1	5	4	5	11	10	11	-

American Indian 2/, both sexes

2007 1/.....	15	-	-	1	-	-	-	-	2	1	4	7	-
2006 1/.....	13	-	-	-	-	1	2	-	-	5	3	2	-
2005 1/.....	7	-	-	-	-	1	1	-	-	2	1	2	-
2004 1/.....	5	-	-	-	-	-	-	2	1	1	1	-	-
2003 1/.....	6	-	-	-	-	1	-	1	-	2	2	-	-
2002.....	6	-	-	1	-	1	1	-	1	-	2	-	-
2001.....	7	-	-	-	-	-	1	2	1	-	3	-	-
2000.....	8	-	-	-	-	1	2	1	1	1	1	1	-
1999.....	4	-	-	-	-	-	1	-	2	1	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

American Indian 2/, male

2007 1/.....	5	-	-	1	-	-	-	-	-	-	1	3	-
2006 1/.....	6	-	-	-	-	-	1	-	-	2	2	1	-
2005 1/.....	4	-	-	-	-	1	1	-	-	1	-	1	-
2004 1/.....	3	-	-	-	-	-	-	1	1	1	-	-	-
2003 1/.....	5	-	-	-	-	-	-	1	-	2	2	-	-
2002.....	4	-	-	1	-	-	1	-	1	-	1	-	-
2001.....	4	-	-	-	-	-	-	1	-	-	3	-	-
2000.....	6	-	-	-	-	1	2	1	1	-	1	-	-
1999.....	3	-	-	-	-	-	1	-	1	1	-	-	-

American Indian 2/, female

2007 1/.....	10	-	-	-	-	-	-	-	2	1	3	4	-
2006 1/.....	7	-	-	-	-	1	1	-	-	3	1	1	-
2005 1/.....	3	-	-	-	-	-	-	-	-	1	1	1	-
2004 1/.....	2	-	-	-	-	-	-	1	-	-	1	-	-
2003 1/.....	1	-	-	-	-	1	-	-	-	-	-	-	-
2002.....	2	-	-	-	-	1	-	-	-	-	1	-	-
2001.....	3	-	-	-	-	-	1	1	1	-	-	-	-
2000.....	2	-	-	-	-	-	-	-	-	1	-	1	-
1999.....	1	-	-	-	-	-	-	-	1	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	73	-	-	-	-	-	1	3	5	11	25	28	-
2006 1/.....	58	-	-	-	-	1	6	3	2	3	22	21	-
2005 1/.....	13	-	-	-	-	1	1	3	1	4	2	1	-
2004 1/.....	11	-	-	-	-	-	-	1	1	-	5	4	-
2003 1/.....	25	-	-	-	1	1	-	1	2	6	7	7	-
2002.....	15	-	-	-	-	-	-	1	4	1	6	3	-
2001.....	18	-	-	-	1	1	3	3	3	4	2	1	-
2000.....	19	-	-	-	-	1	2	-	3	3	2	8	-
1999.....	24	-	-	-	-	2	1	2	4	4	7	4	-

Asian or Pacific Islander 3/, male

2007 1/.....	39	-	-	-	-	-	-	2	3	5	12	17	-
2006 1/.....	30	-	-	-	-	-	4	2	1	1	12	10	-
2005 1/.....	9	-	-	-	-	1	1	3	-	3	-	1	-
2004 1/.....	9	-	-	-	-	-	-	-	1	-	4	4	-
2003 1/.....	11	-	-	-	1	1	-	-	1	1	3	4	-
2002.....	4	-	-	-	-	-	-	1	-	-	2	1	-
2001.....	7	-	-	-	1	-	-	1	2	2	1	-	-
2000.....	8	-	-	-	-	1	1	-	-	1	1	4	-
1999.....	10	-	-	-	-	1	1	-	1	2	3	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

Asian or Pacific Islander 3/, female

2007 1/.....	34	-	-	-	-	-	1	1	2	6	13	11	-
2006 1/.....	28	-	-	-	-	1	2	1	1	2	10	11	-
2005 1/.....	4	-	-	-	-	-	-	-	1	1	2	-	-
2004 1/.....	2	-	-	-	-	-	-	1	-	-	1	-	-
2003 1/.....	14	-	-	-	-	-	-	1	1	5	4	3	-
2002.....	11	-	-	-	-	-	-	-	4	1	4	2	-
2001.....	11	-	-	-	-	1	3	2	1	2	1	1	-
2000.....	11	-	-	-	-	-	1	-	3	2	1	4	-
1999.....	14	-	-	-	-	1	-	2	3	2	4	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Renal failure (N17-N19)													
All races, both sexes													
2007.....	43,263	138	16	18	77	237	696	2,091	4,205	7,330	13,718	14,737	-
2006.....	43,344	154	10	12	89	250	745	2,151	4,207	7,171	13,915	14,640	-
2005.....	42,868	151	13	19	79	261	705	1,973	4,025	7,165	14,102	14,375	-
2004.....	41,732	165	9	18	79	223	767	2,009	3,883	6,997	13,830	13,750	2
2003.....	41,737	170	11	21	82	266	771	1,944	3,741	7,206	13,870	13,653	2
2002.....	40,222	165	9	15	77	243	718	1,829	3,383	7,046	13,657	13,080	-
2001.....	38,784	125	9	16	82	209	732	1,770	3,226	7,211	12,898	12,505	1
2000.....	36,471	152	11	19	78	221	701	1,590	3,023	6,848	12,223	11,603	2
1999.....	34,719	163	12	25	63	230	673	1,422	2,784	6,697	11,652	10,998	-
All races, male													
2007.....	21,038	74	11	11	37	125	400	1,174	2,336	3,807	6,933	6,130	-
2006.....	21,115	94	7	7	52	139	436	1,285	2,346	3,767	6,924	6,058	-
2005.....	20,745	86	9	10	41	136	395	1,201	2,288	3,650	7,027	5,902	-
2004.....	20,006	106	6	11	46	105	433	1,175	2,072	3,616	6,895	5,540	1
2003.....	20,135	98	5	9	49	141	443	1,151	2,036	3,719	6,947	5,535	2
2002.....	19,341	105	5	8	37	114	393	1,055	1,786	3,739	6,808	5,291	-
2001.....	18,509	72	3	6	40	109	394	1,022	1,680	3,752	6,381	5,050	-
2000.....	17,416	100	5	14	40	115	407	941	1,607	3,492	6,102	4,591	2
1999.....	16,611	87	7	15	25	110	382	814	1,492	3,440	5,888	4,351	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Renal failure (N17-N19)													
All races, female													
2007.....	22,225	64	5	7	40	112	296	917	1,869	3,523	6,785	8,607	-
2006.....	22,229	60	3	5	37	111	309	866	1,861	3,404	6,991	8,582	-
2005.....	22,123	65	4	9	38	125	310	772	1,737	3,515	7,075	8,473	-
2004.....	21,726	59	3	7	33	118	334	834	1,811	3,381	6,935	8,210	1
2003.....	21,602	72	6	12	33	125	328	793	1,705	3,487	6,923	8,118	-
2002.....	20,881	60	4	7	40	129	325	774	1,597	3,307	6,849	7,789	-
2001.....	20,275	53	6	10	42	100	338	748	1,546	3,459	6,517	7,455	1
2000.....	19,055	52	6	5	38	106	294	649	1,416	3,356	6,121	7,012	-
1999.....	18,108	76	5	10	38	120	291	608	1,292	3,257	5,764	6,647	-
White, both sexes													
2007 1/.....	34,337	89	10	14	49	126	370	1,254	2,728	5,346	11,472	12,879	-
2006 1/.....	34,241	106	8	8	59	145	392	1,281	2,768	5,158	11,552	12,764	-
2005 1/.....	33,983	91	11	15	51	139	378	1,189	2,575	5,225	11,737	12,572	-
2004 1/.....	33,084	92	6	14	48	114	444	1,146	2,490	5,153	11,543	12,032	2
2003 1/.....	33,133	108	8	14	50	148	397	1,107	2,465	5,270	11,582	11,983	1
2002.....	32,008	107	6	11	38	135	391	1,042	2,191	5,197	11,461	11,429	-
2001.....	30,777	70	7	10	53	114	408	1,013	2,059	5,337	10,763	10,942	1
2000.....	28,982	80	7	16	55	121	390	901	1,949	5,028	10,265	10,168	2
1999.....	27,407	99	10	17	37	122	362	784	1,815	4,972	9,701	9,488	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Renal failure (N17-N19)													
White, male													
2007 1/.....	16,961	44	6	8	23	66	223	730	1,554	2,845	5,975	5,487	-
2006 1/.....	16,956	66	6	4	39	86	239	790	1,598	2,813	5,891	5,424	-
2005 1/.....	16,715	50	7	7	28	73	221	745	1,497	2,798	6,010	5,279	-
2004 1/.....	16,055	59	3	8	27	49	260	668	1,376	2,741	5,932	4,931	1
2003 1/.....	16,139	65	4	7	32	77	239	652	1,356	2,797	5,952	4,957	1
2002.....	15,558	68	3	7	19	65	218	602	1,153	2,858	5,878	4,687	-
2001.....	14,961	42	1	5	26	54	241	596	1,121	2,889	5,471	4,515	-
2000.....	14,070	55	4	12	31	64	229	534	1,071	2,709	5,253	4,106	2
1999.....	13,312	59	5	12	15	60	207	450	1,016	2,661	4,996	3,831	-
White, female													
2007 1/.....	17,376	45	4	6	26	60	147	524	1,174	2,501	5,497	7,392	-
2006 1/.....	17,285	40	2	4	20	59	153	491	1,170	2,345	5,661	7,340	-
2005 1/.....	17,268	41	4	8	23	66	157	444	1,078	2,427	5,727	7,293	-
2004 1/.....	17,029	33	3	6	21	65	184	478	1,114	2,412	5,611	7,101	1
2003 1/.....	16,994	43	4	7	18	71	158	455	1,109	2,473	5,630	7,026	-
2002.....	16,450	39	3	4	19	70	173	440	1,038	2,339	5,583	6,742	-
2001.....	15,816	28	6	5	27	60	167	417	938	2,448	5,292	6,427	1
2000.....	14,912	25	3	4	24	57	161	367	878	2,319	5,012	6,062	-
1999.....	14,095	40	5	5	22	62	155	334	799	2,311	4,705	5,657	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Renal failure (N17-N19)													
Black, both sexes													
2007 1/.....	7,832	46	5	2	27	104	309	776	1,327	1,740	1,907	1,589	-
2006 1/.....	8,028	42	2	2	27	96	323	803	1,296	1,772	2,051	1,614	-
2005 1/.....	7,891	56	2	3	26	112	302	734	1,319	1,725	2,064	1,548	-
2004 1/.....	7,715	65	3	3	28	95	304	805	1,275	1,633	1,992	1,512	-
2003 1/.....	7,755	59	3	7	29	114	351	772	1,174	1,743	2,006	1,496	1
2002.....	7,373	52	3	4	33	101	305	729	1,073	1,680	1,931	1,462	-
2001.....	7,178	53	2	5	26	91	301	690	1,043	1,691	1,890	1,386	-
2000.....	6,782	68	3	3	22	94	293	644	979	1,632	1,764	1,280	-
1999.....	6,577	58	1	7	26	99	297	602	868	1,547	1,738	1,334	-
Black, male													
2007 1/.....	3,521	29	4	1	13	54	168	409	692	825	801	525	-
2006 1/.....	3,632	23	1	2	11	48	179	459	684	830	886	509	-
2005 1/.....	3,558	36	2	2	11	55	156	433	734	746	878	505	-
2004 1/.....	3,483	40	3	2	17	50	161	470	633	771	826	510	-
2003 1/.....	3,587	31	1	2	16	62	190	461	630	844	857	492	1
2002.....	3,377	33	2	1	14	45	161	422	574	800	807	518	-
2001.....	3,138	29	2	-	12	54	136	388	506	776	787	448	-
2000.....	3,015	42	1	2	9	47	170	382	482	706	757	417	-
1999.....	2,943	25	1	3	10	44	166	346	421	690	799	438	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Renal failure (N17-N19)													
Black, female													
2007 1/.....	4,311	17	1	1	14	50	141	367	635	915	1,106	1,064	-
2006 1/.....	4,396	19	1	-	16	48	144	344	612	942	1,165	1,105	-
2005 1/.....	4,333	20	-	1	15	57	146	301	585	979	1,186	1,043	-
2004 1/.....	4,232	25	-	1	11	45	143	335	642	862	1,166	1,002	-
2003 1/.....	4,168	28	2	5	13	52	161	311	544	899	1,149	1,004	-
2002.....	3,996	19	1	3	19	56	144	307	499	880	1,124	944	-
2001.....	4,040	24	-	5	14	37	165	302	537	915	1,103	938	-
2000.....	3,767	26	2	1	13	47	123	262	497	926	1,007	863	-
1999.....	3,634	33	-	4	16	55	131	256	447	857	939	896	-
American Indian 2/, both sexes													
2007 1/.....	277	2	1	-	-	1	11	23	55	73	64	47	-
2006 1/.....	274	1	-	1	3	6	15	33	51	76	49	39	-
2005 1/.....	256	1	-	1	1	5	13	20	47	63	61	44	-
2004 1/.....	240	-	-	1	2	9	9	27	38	59	58	37	-
2003 1/.....	248	1	-	-	3	1	11	31	42	44	74	41	-
2002.....	214	2	-	-	2	5	9	26	42	56	43	29	-
2001.....	226	1	-	1	2	1	12	29	49	54	49	28	-
2000.....	205	1	-	-	-	2	10	21	37	64	45	25	-
1999.....	187	-	-	-	-	3	6	18	33	54	41	32	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Renal failure (N17-N19)

American Indian 2/, male

2007 1/.....	121	1	1	-	-	1	5	11	26	35	22	19	-
2006 1/.....	129	1	-	1	2	4	9	18	21	34	22	17	-
2005 1/.....	100	-	-	1	1	4	8	6	17	27	18	18	-
2004 1/.....	107	-	-	1	1	2	4	16	18	27	22	16	-
2003 1/.....	122	1	-	-	1	1	6	17	21	17	41	17	-
2002.....	93	2	-	-	2	3	7	14	19	19	19	8	-
2001.....	99	1	-	1	2	-	8	18	15	19	21	14	-
2000.....	90	-	-	-	-	1	5	10	19	28	15	12	-
1999.....	74	-	-	-	-	3	5	9	15	22	12	8	-

American Indian 2/, female

2007 1/.....	156	1	-	-	-	-	6	12	29	38	42	28	-
2006 1/.....	145	-	-	-	1	2	6	15	30	42	27	22	-
2005 1/.....	156	1	-	-	-	1	5	14	30	36	43	26	-
2004 1/.....	133	-	-	-	1	7	5	11	20	32	36	21	-
2003 1/.....	126	-	-	-	2	-	5	14	21	27	33	24	-
2002.....	121	-	-	-	-	2	2	12	23	37	24	21	-
2001.....	127	-	-	-	-	1	4	11	34	35	28	14	-
2000.....	115	1	-	-	-	1	5	11	18	36	30	13	-
1999.....	113	-	-	-	-	-	1	9	18	32	29	24	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Renal failure (N17-N19)													
Asian or Pacific Islander 3/, both sexes													
2007 1/.....	817	1	-	2	1	6	6	38	95	171	275	222	-
2006 1/.....	801	5	-	1	-	3	15	34	92	165	263	223	-
2005 1/.....	738	3	-	-	1	5	12	30	84	152	240	211	-
2004 1/.....	693	8	-	-	1	5	10	31	80	152	237	169	-
2003 1/.....	601	2	-	-	-	3	12	34	60	149	208	133	-
2002.....	627	4	-	-	4	2	13	32	77	113	222	160	-
2001.....	603	1	-	-	1	3	11	38	75	129	196	149	-
2000.....	502	3	1	-	1	4	8	24	58	124	149	130	-
1999.....	548	6	1	1	-	6	8	18	68	124	172	144	-
Asian or Pacific Islander 3/, male													
2007 1/.....	435	-	-	2	1	4	4	24	64	102	135	99	-
2006 1/.....	398	4	-	-	-	1	9	18	43	90	125	108	-
2005 1/.....	372	-	-	-	1	4	10	17	40	79	121	100	-
2004 1/.....	361	7	-	-	1	4	8	21	45	77	115	83	-
2003 1/.....	287	1	-	-	-	1	8	21	29	61	97	69	-
2002.....	313	2	-	-	2	1	7	17	40	62	104	78	-
2001.....	311	-	-	-	-	1	9	20	38	68	102	73	-
2000.....	241	3	-	-	-	3	3	15	35	49	77	56	-
1999.....	282	3	1	-	-	3	4	9	40	67	81	74	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Renal failure (N17-N19)

Asian or Pacific Islander 3/, female

2007 1/.....	382	1	-	-	-	2	2	14	31	69	140	123	-
2006 1/.....	403	1	-	1	-	2	6	16	49	75	138	115	-
2005 1/.....	366	3	-	-	-	1	2	13	44	73	119	111	-
2004 1/.....	332	1	-	-	-	1	2	10	35	75	122	86	-
2003 1/.....	314	1	-	-	-	2	4	13	31	88	111	64	-
2002.....	314	2	-	-	2	1	6	15	37	51	118	82	-
2001.....	292	1	-	-	1	2	2	18	37	61	94	76	-
2000.....	261	-	1	-	1	1	5	9	23	75	72	74	-
1999.....	266	3	-	1	-	3	4	9	28	57	91	70	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other disorders of kidney (N25,N27)													
All races, both sexes													
2007.....	21	-	1	-	-	-	1	3	5	3	3	5	-
2006.....	21	-	-	1	-	1	2	1	2	5	5	4	-
2005.....	29	-	-	-	1	-	3	1	8	6	6	4	-
2004.....	38	1	-	-	1	1	2	3	7	7	11	5	-
2003.....	31	-	1	-	1	-	1	5	1	7	8	7	-
2002.....	33	-	-	1	-	2	1	2	4	6	10	7	-
2001.....	29	-	1	-	2	1	1	5	5	7	4	3	-
2000.....	47	1	-	-	-	2	2	6	8	10	11	7	-
1999.....	35	-	-	-	-	2	1	3	4	6	13	6	-
All races, male													
2007.....	15	-	-	-	-	-	1	3	3	2	3	3	-
2006.....	12	-	-	1	-	1	2	-	1	3	2	2	-
2005.....	12	-	-	-	-	-	1	-	3	5	2	1	-
2004.....	16	1	-	-	1	1	-	3	4	1	4	1	-
2003.....	12	-	1	-	1	-	-	4	-	3	2	1	-
2002.....	6	-	-	1	-	1	-	1	-	-	3	-	-
2001.....	6	-	-	-	-	-	1	2	-	1	1	1	-
2000.....	20	-	-	-	-	-	1	3	5	6	2	3	-
1999.....	11	-	-	-	-	1	-	1	-	2	4	3	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other disorders of kidney (N25,N27)													
All races, female													
2007.....	6	-	1	-	-	-	-	-	2	1	-	2	-
2006.....	9	-	-	-	-	-	-	1	1	2	3	2	-
2005.....	17	-	-	-	1	-	2	1	5	1	4	3	-
2004.....	22	-	-	-	-	-	2	-	3	6	7	4	-
2003.....	19	-	-	-	-	-	1	1	1	4	6	6	-
2002.....	27	-	-	-	-	1	1	1	4	6	7	7	-
2001.....	23	-	1	-	2	1	-	3	5	6	3	2	-
2000.....	27	1	-	-	-	2	1	3	3	4	9	4	-
1999.....	24	-	-	-	-	1	1	2	4	4	9	3	-
White, both sexes													
2007 1/.....	16	-	1	-	-	-	1	3	2	3	3	3	-
2006 1/.....	20	-	-	1	-	1	2	1	1	5	5	4	-
2005 1/.....	23	-	-	-	1	-	2	-	8	5	5	2	-
2004 1/.....	33	1	-	-	1	1	1	1	7	7	9	5	-
2003 1/.....	24	-	1	-	-	-	1	4	1	3	7	7	-
2002.....	27	-	-	-	-	2	1	1	4	6	7	6	-
2001.....	21	-	1	-	1	1	1	4	4	7	1	1	-
2000.....	36	1	-	-	-	1	1	3	7	9	9	5	-
1999.....	23	-	-	-	-	2	1	2	1	5	8	4	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other disorders of kidney (N25,N27)													
White, male													
2007 1/.....	11	-	-	-	-	-	1	3	-	2	3	2	-
2006 1/.....	12	-	-	1	-	1	2	-	1	3	2	2	-
2005 1/.....	12	-	-	-	-	-	1	-	3	5	2	1	-
2004 1/.....	14	1	-	-	1	1	-	1	4	1	4	1	-
2003 1/.....	8	-	1	-	-	-	-	3	-	1	2	1	-
2002.....	3	-	-	-	-	1	-	-	-	-	2	-	-
2001.....	4	-	-	-	-	-	1	2	-	1	-	-	-
2000.....	17	-	-	-	-	-	1	1	4	6	2	3	-
1999.....	9	-	-	-	-	1	-	1	-	1	3	3	-
White, female													
2007 1/.....	5	-	1	-	-	-	-	-	2	1	-	1	-
2006 1/.....	8	-	-	-	-	-	-	1	-	2	3	2	-
2005 1/.....	11	-	-	-	1	-	1	-	5	-	3	1	-
2004 1/.....	19	-	-	-	-	-	1	-	3	6	5	4	-
2003 1/.....	16	-	-	-	-	-	1	1	1	2	5	6	-
2002.....	24	-	-	-	-	1	1	1	4	6	5	6	-
2001.....	17	-	1	-	1	1	-	2	4	6	1	1	-
2000.....	19	1	-	-	-	1	-	2	3	3	7	2	-
1999.....	14	-	-	-	-	1	1	1	1	4	5	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other disorders of kidney (N25,N27)													
Black, both sexes													
2007 1/.....	5	-	-	-	-	-	-	-	3	-	-	2	-
2006 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2005 1/.....	5	-	-	-	-	-	-	1	-	1	1	2	-
2004 1/.....	5	-	-	-	-	-	1	2	-	-	2	-	-
2003 1/.....	6	-	-	-	1	-	-	1	-	3	1	-	-
2002.....	4	-	-	1	-	-	-	1	-	-	1	1	-
2001.....	6	-	-	-	1	-	-	-	1	-	2	2	-
2000.....	9	-	-	-	-	1	1	3	-	1	1	2	-
1999.....	12	-	-	-	-	-	-	1	3	1	5	2	-
Black, male													
2007 1/.....	4	-	-	-	-	-	-	-	3	-	-	1	-
2006 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2005 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 1/.....	2	-	-	-	-	-	-	2	-	-	-	-	-
2003 1/.....	4	-	-	-	1	-	-	1	-	2	-	-	-
2002.....	2	-	-	1	-	-	-	1	-	-	-	-	-
2001.....	2	-	-	-	-	-	-	-	-	-	1	1	-
2000.....	2	-	-	-	-	-	-	2	-	-	-	-	-
1999.....	2	-	-	-	-	-	-	-	-	1	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Infections of kidney (N10-N12,N13.6,N15.1)

All races, both sexes

2007.....	628	5	1	1	8	6	30	59	64	91	170	193	-
2006.....	673	5	1	2	4	14	30	55	57	90	164	251	-
2005.....	767	6	4	2	9	14	26	51	53	100	239	263	-
2004.....	764	3	4	-	4	15	34	47	65	104	237	251	-
2003.....	823	4	2	1	10	16	36	62	62	105	247	278	-
2002.....	788	2	-	3	5	13	28	40	59	110	238	290	-
2001.....	748	2	4	-	5	11	33	46	46	87	249	265	-
2000.....	813	5	3	-	2	9	21	42	59	104	248	320	-
1999.....	846	3	1	1	4	9	28	55	59	133	242	311	-

All races, male

2007.....	214	-	1	-	3	1	14	20	26	34	56	59	-
2006.....	206	5	-	2	-	5	11	16	22	40	42	63	-
2005.....	236	3	1	-	2	5	7	18	22	37	83	58	-
2004.....	237	2	1	-	1	3	7	12	32	34	77	68	-
2003.....	246	3	-	1	2	1	10	25	21	31	74	78	-
2002.....	251	2	-	1	1	6	12	15	21	47	72	74	-
2001.....	213	1	2	-	2	3	10	11	12	40	76	56	-
2000.....	214	3	-	-	1	3	8	15	15	26	74	69	-
1999.....	263	1	-	-	1	-	7	19	27	44	83	81	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Infections of kidney (N10-N12,N13.6,N15.1)													
All races, female													
2007.....	414	5	-	1	5	5	16	39	38	57	114	134	-
2006.....	467	-	1	-	4	9	19	39	35	50	122	188	-
2005.....	531	3	3	2	7	9	19	33	31	63	156	205	-
2004.....	527	1	3	-	3	12	27	35	33	70	160	183	-
2003.....	577	1	2	-	8	15	26	37	41	74	173	200	-
2002.....	537	-	-	2	4	7	16	25	38	63	166	216	-
2001.....	535	1	2	-	3	8	23	35	34	47	173	209	-
2000.....	599	2	3	-	1	6	13	27	44	78	174	251	-
1999.....	583	2	1	1	3	9	21	36	32	89	159	230	-
White, both sexes													
2007 1/.....	552	2	1	1	4	5	26	46	55	75	154	183	-
2006 1/.....	599	3	1	2	4	11	23	49	48	79	146	233	-
2005 1/.....	660	5	4	2	7	9	16	39	42	84	215	237	-
2004 1/.....	657	2	3	-	4	10	29	34	49	92	205	229	-
2003 1/.....	705	1	1	1	8	11	28	47	46	93	215	254	-
2002.....	692	2	-	1	1	10	22	26	46	101	217	266	-
2001.....	647	1	2	-	2	8	26	34	37	73	219	245	-
2000.....	689	5	3	-	1	7	12	30	48	82	210	291	-
1999.....	749	1	1	1	2	8	20	44	48	110	223	291	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Infections of kidney (N10-N12,N13.6,N15.1)													
White, male													
2007 1/.....	185	-	1	-	2	1	14	15	21	25	49	57	-
2006 1/.....	180	3	-	2	-	4	8	13	17	37	39	57	-
2005 1/.....	204	2	1	-	2	4	5	14	17	32	77	50	-
2004 1/.....	204	1	-	-	1	1	6	9	25	27	70	64	-
2003 1/.....	215	-	-	1	2	1	9	21	15	28	62	76	-
2002.....	219	2	-	-	1	6	10	12	17	40	63	68	-
2001.....	176	-	1	-	1	2	9	9	9	30	65	50	-
2000.....	182	3	-	-	-	3	4	12	13	17	65	65	-
1999.....	232	1	-	-	1	-	5	14	22	34	77	78	-
White, female													
2007 1/.....	367	2	-	1	2	4	12	31	34	50	105	126	-
2006 1/.....	419	-	1	-	4	7	15	36	31	42	107	176	-
2005 1/.....	456	3	3	2	5	5	11	25	25	52	138	187	-
2004 1/.....	453	1	3	-	3	9	23	25	24	65	135	165	-
2003 1/.....	490	1	1	-	6	10	19	26	31	65	153	178	-
2002.....	473	-	-	1	-	4	12	14	29	61	154	198	-
2001.....	471	1	1	-	1	6	17	25	28	43	154	195	-
2000.....	507	2	3	-	1	4	8	18	35	65	145	226	-
1999.....	517	-	1	1	1	8	15	30	26	76	146	213	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Infections of kidney (N10-N12,N13.6,N15.1)

Black, both sexes

2007 1/.....	53	3	-	-	4	1	3	9	7	11	8	7	-
2006 1/.....	58	1	-	-	-	3	6	6	8	7	15	12	-
2005 1/.....	89	1	-	-	2	4	10	10	10	14	19	19	-
2004 1/.....	84	1	1	-	-	5	4	10	12	9	25	17	-
2003 1/.....	95	3	1	-	2	2	7	12	14	10	24	20	-
2002.....	73	-	-	-	3	2	6	8	10	8	15	21	-
2001.....	80	1	-	-	1	3	7	10	7	13	22	16	-
2000.....	101	-	-	-	1	2	7	11	8	19	28	25	-
1999.....	85	1	-	-	2	1	6	10	11	20	16	18	-

Black, male

2007 1/.....	20	-	-	-	1	-	-	3	3	8	3	2	-
2006 1/.....	21	1	-	-	-	1	2	3	5	3	1	5	-
2005 1/.....	27	1	-	-	-	1	2	3	5	5	5	5	-
2004 1/.....	27	1	1	-	-	2	1	2	6	6	5	3	-
2003 1/.....	29	3	-	-	-	-	1	4	5	3	11	2	-
2002.....	22	-	-	-	-	-	2	1	3	6	5	5	-
2001.....	28	1	-	-	1	1	1	2	3	9	6	4	-
2000.....	27	-	-	-	1	-	2	3	2	8	7	4	-
1999.....	29	-	-	-	-	-	2	5	5	10	4	3	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Infections of kidney (N10-N12,N13.6,N15.1)

Black, female

2007 1/.....	33	3	-	-	3	1	3	6	4	3	5	5	-
2006 1/.....	37	-	-	-	-	2	4	3	3	4	14	7	-
2005 1/.....	62	-	-	-	2	3	8	7	5	9	14	14	-
2004 1/.....	57	-	-	-	-	3	3	8	6	3	20	14	-
2003 1/.....	66	-	1	-	2	2	6	8	9	7	13	18	-
2002.....	51	-	-	-	3	2	4	7	7	2	10	16	-
2001.....	52	-	-	-	-	2	6	8	4	4	16	12	-
2000.....	74	-	-	-	-	2	5	8	6	11	21	21	-
1999.....	56	1	-	-	2	1	4	5	6	10	12	15	-

American Indian 2/, both sexes

2007 1/.....	10	-	-	-	-	-	1	3	-	1	5	-	-
2006 1/.....	1	-	-	-	-	-	-	-	-	-	-	1	-
2005 1/.....	4	-	-	-	-	1	-	1	-	-	2	-	-
2004 1/.....	6	-	-	-	-	-	1	-	2	1	2	-	-
2003 1/.....	6	-	-	-	-	1	1	1	1	-	1	1	-
2002.....	8	-	-	1	-	-	-	3	1	-	2	1	-
2001.....	6	-	1	-	1	-	-	1	-	-	2	1	-
2000.....	5	-	-	-	-	-	-	1	-	1	1	2	-
1999.....	3	-	-	-	-	-	1	-	-	2	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Infections of kidney (N10-N12,N13.6,N15.1)

American Indian 2/, male

2007 1/.....	3	-	-	-	-	-	-	1	-	-	2	-	-
2006 1/.....	1	-	-	-	-	-	-	-	-	-	-	1	-
2005 1/.....	1	-	-	-	-	-	-	1	-	-	-	-	-
2004 1/.....	1	-	-	-	-	-	-	-	-	-	1	-	-
2003 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	3	-	-	1	-	-	-	-	-	-	2	-	-
2001.....	2	-	-	-	-	-	-	-	-	-	2	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-	-

American Indian 2/, female

2007 1/.....	7	-	-	-	-	-	1	2	-	1	3	-	-
2006 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2005 1/.....	3	-	-	-	-	1	-	-	-	-	2	-	-
2004 1/.....	5	-	-	-	-	-	1	-	2	1	1	-	-
2003 1/.....	6	-	-	-	-	1	1	1	1	-	1	1	-
2002.....	5	-	-	-	-	-	-	3	1	-	-	1	-
2001.....	4	-	1	-	1	-	-	1	-	-	-	1	-
2000.....	5	-	-	-	-	-	-	1	-	1	1	2	-
1999.....	3	-	-	-	-	-	1	-	-	2	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Infections of kidney (N10-N12,N13.6,N15.1)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	13	-	-	-	-	-	-	1	2	4	3	3	-
2006 1/.....	15	1	-	-	-	-	1	-	1	4	3	5	-
2005 1/.....	14	-	-	-	-	-	1	1	2	3	7	-	-
2004 1/.....	17	-	-	-	-	-	3	2	2	5	5	-	-
2003 1/.....	17	-	-	-	-	2	-	2	1	2	7	3	-
2002.....	15	-	-	1	1	1	-	3	2	1	4	2	-
2001.....	15	-	1	-	1	-	-	1	2	1	6	3	-
2000.....	18	-	-	-	-	-	2	-	3	2	9	2	-
1999.....	9	1	-	-	-	-	1	1	-	1	3	2	-

Asian or Pacific Islander 3/, male

2007 1/.....	6	-	-	-	-	-	-	1	2	1	2	-	-
2006 1/.....	4	1	-	-	-	-	1	-	-	-	2	-	-
2005 1/.....	4	-	-	-	-	-	-	-	-	-	1	3	-
2004 1/.....	5	-	-	-	-	-	-	1	1	1	1	1	-
2003 1/.....	2	-	-	-	-	-	-	-	1	-	1	-	-
2002.....	7	-	-	-	-	-	-	2	1	1	2	1	-
2001.....	7	-	1	-	-	-	-	-	-	1	3	2	-
2000.....	5	-	-	-	-	-	2	-	-	1	2	-	-
1999.....	2	-	-	-	-	-	-	-	-	-	2	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Infections of kidney (N10-N12,N13.6,N15.1)

Asian or Pacific Islander 3/, female

2007 1/.....	7	-	-	-	-	-	-	-	-	3	1	3	-
2006 1/.....	11	-	-	-	-	-	-	-	1	4	1	5	-
2005 1/.....	10	-	-	-	-	-	-	1	1	2	2	4	-
2004 1/.....	12	-	-	-	-	-	-	2	1	1	4	4	-
2003 1/.....	15	-	-	-	-	2	-	2	-	2	6	3	-
2002.....	8	-	-	1	1	1	-	1	1	-	2	1	-
2001.....	8	-	-	-	1	-	-	1	2	-	3	1	-
2000.....	13	-	-	-	-	-	-	-	3	1	7	2	-
1999.....	7	1	-	-	-	-	1	1	-	1	1	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Hyperplasia of prostate (N40)

All races, both sexes

2007.....	491	-	-	-	-	-	1	1	12	47	147	283	-
2006.....	514	-	-	-	-	-	-	-	10	46	167	291	-
2005.....	525	-	-	-	-	-	-	-	10	52	171	292	-
2004.....	507	-	-	-	-	-	-	4	13	45	177	268	-
2003.....	477	1	-	-	-	-	-	1	12	50	165	248	-
2002.....	437	-	-	-	-	-	1	-	10	41	164	220	1
2001.....	439	-	-	-	-	-	-	2	9	45	141	242	-
2000.....	433	-	-	-	-	-	1	1	11	38	159	222	1
1999.....	430	-	-	-	-	-	-	1	13	47	160	209	-

All races, male

2007.....	491	-	-	-	-	-	1	1	12	47	147	283	-
2006.....	514	-	-	-	-	-	-	-	10	46	167	291	-
2005.....	525	-	-	-	-	-	-	-	10	52	171	292	-
2004.....	507	-	-	-	-	-	-	4	13	45	177	268	-
2003.....	477	1	-	-	-	-	-	1	12	50	165	248	-
2002.....	437	-	-	-	-	-	1	-	10	41	164	220	1
2001.....	439	-	-	-	-	-	-	2	9	45	141	242	-
2000.....	433	-	-	-	-	-	1	1	11	38	159	222	1
1999.....	430	-	-	-	-	-	-	1	13	47	160	209	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Hyperplasia of prostate (N40)

All races, female

2007.....
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

White, both sexes

2007 1/.....	433	-	-	-	-	-	1	1	7	39	135	250	-
2006 1/.....	463	-	-	-	-	-	-	-	8	36	151	268	-
2005 1/.....	483	-	-	-	-	-	-	-	7	45	159	272	-
2004 1/.....	458	-	-	-	-	-	-	2	12	39	156	249	-
2003 1/.....	438	-	-	-	-	-	-	-	9	44	150	235	-
2002.....	389	-	-	-	-	-	-	-	7	34	145	203	-
2001.....	403	-	-	-	-	-	-	1	5	40	129	228	-
2000.....	389	-	-	-	-	-	-	1	7	35	145	201	-
1999.....	390	-	-	-	-	-	-	1	9	40	147	193	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Hyperplasia of prostate (N40)

Black, both sexes

2007 1/.....	44	-	-	-	-	-	-	4	7	10	23	-
2006 1/.....	38	-	-	-	-	-	-	2	9	13	14	-
2005 1/.....	30	-	-	-	-	-	-	3	5	8	14	-
2004 1/.....	35	-	-	-	-	-	2	1	6	13	13	-
2003 1/.....	32	1	-	-	-	-	1	3	5	11	11	-
2002.....	39	-	-	-	-	1	-	3	7	14	13	1
2001.....	32	-	-	-	-	-	1	4	4	12	11	-
2000.....	37	-	-	-	-	1	-	4	2	14	15	1
1999.....	38	-	-	-	-	-	-	4	6	13	15	-

Black, male

2007 1/.....	44	-	-	-	-	-	-	4	7	10	23	-
2006 1/.....	38	-	-	-	-	-	-	2	9	13	14	-
2005 1/.....	30	-	-	-	-	-	-	3	5	8	14	-
2004 1/.....	35	-	-	-	-	-	2	1	6	13	13	-
2003 1/.....	32	1	-	-	-	-	1	3	5	11	11	-
2002.....	39	-	-	-	-	1	-	3	7	14	13	1
2001.....	32	-	-	-	-	-	1	4	4	12	11	-
2000.....	37	-	-	-	-	1	-	4	2	14	15	1
1999.....	38	-	-	-	-	-	-	4	6	13	15	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Hyperplasia of prostate (N40)

Black, female

2007 1/.....
2006 1/.....
2005 1/.....
2004 1/.....
2003 1/.....
2002.....
2001.....
2000.....
1999.....

American Indian 2/, both sexes

2007 1/.....	4	-	-	-	-	-	-	-	1	-	-	3	-
2006 1/.....	1	-	-	-	-	-	-	-	-	-	-	1	-
2005 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 1/.....	2	-	-	-	-	-	-	-	-	-	2	-	-
2003 1/.....	2	-	-	-	-	-	-	-	-	-	-	2	-
2002.....	1	-	-	-	-	-	-	-	-	-	1	-	-
2001.....	1	-	-	-	-	-	-	-	-	1	-	-	-
2000.....	2	-	-	-	-	-	-	-	-	1	-	1	-
1999.....	1	-	-	-	-	-	-	-	-	1	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Inflammatory diseases of female pelvic organs (N70-N76)

All races, female

2007.....	116	4	-	-	1	5	6	14	19	22	21	24	-
2006.....	112	1	-	-	2	3	7	13	12	12	35	27	-
2005.....	120	-	-	-	1	5	7	13	9	23	33	29	-
2004.....	112	1	-	-	2	4	8	11	15	16	21	34	-
2003.....	125	1	-	1	6	3	14	14	10	16	29	31	-
2002.....	114	1	-	-	2	4	17	11	6	8	29	36	-
2001.....	127	1	-	-	2	3	10	12	12	18	31	38	-
2000.....	120	2	-	1	2	6	15	9	8	21	29	27	-
1999.....	108	2	1	-	3	6	12	6	13	15	32	18	-

White, both sexes

2007 1/.....	90	2	-	-	-	2	4	10	15	15	21	21	-
2006 1/.....	90	-	-	-	2	1	6	7	9	9	31	25	-
2005 1/.....	98	-	-	-	-	3	6	7	7	19	31	25	-
2004 1/.....	92	1	-	-	2	3	6	6	13	14	16	31	-
2003 1/.....	101	-	-	1	3	2	10	9	8	13	25	30	-
2002.....	92	1	-	-	-	3	9	7	5	7	28	32	-
2001.....	103	1	-	-	2	3	6	8	8	14	27	34	-
2000.....	94	1	-	-	2	4	7	7	6	17	26	24	-
1999.....	80	1	1	-	2	3	4	4	12	10	26	17	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Inflammatory diseases of female pelvic organs (N70-N76)

White, male

2007 1/.....
2006 1/.....
2005 1/.....
2004 1/.....
2003 1/.....
2002.....
2001.....
2000.....
1999.....

White, female

2007 1/.....	90	2	-	-	-	2	4	10	15	15	21	21	-
2006 1/.....	90	-	-	-	2	1	6	7	9	9	31	25	-
2005 1/.....	98	-	-	-	-	3	6	7	7	19	31	25	-
2004 1/.....	92	1	-	-	2	3	6	6	13	14	16	31	-
2003 1/.....	101	-	-	1	3	2	10	9	8	13	25	30	-
2002.....	92	1	-	-	-	3	9	7	5	7	28	32	-
2001.....	103	1	-	-	2	3	6	8	8	14	27	34	-
2000.....	94	1	-	-	2	4	7	7	6	17	26	24	-
1999.....	80	1	1	-	2	3	4	4	12	10	26	17	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Inflammatory diseases of female pelvic organs (N70-N76)

Asian or Pacific Islander 3/, female

2007 1/.....	4	-	-	-	1	1	-	-	1	1	-	-	-
2006 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2005 1/.....	2	-	-	-	-	1	1	-	-	-	-	-	-
2004 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 1/.....	2	-	-	-	-	-	-	-	-	-	2	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	1	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	1	-	-	-	-	-	-
1999.....	2	-	-	-	-	1	1	-	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pregnancy, childbirth and the puerperium (000-099)													
All races, female													
2007.....	769	1	160	326	220	59	2	1	-	-	-
2006.....	760	2	179	324	192	59	3	-	-	1	-
2005.....	760	-	183	312	207	54	3	1	-	-	-
2004.....	697	2	168	287	157	79	1	1	2	-	-
2003.....	545	1	116	229	166	30	2	1	-	-	-
2002.....	379	1	89	175	105	8	-	-	-	1	-
2001.....	416	3	110	190	106	6	-	-	-	1	-
2000.....	404	-	95	182	121	6	-	-	-	-	-
1999.....	406	-	97	181	121	7	-	-	-	-	-
White, both sexes													
2007 1/.....	465	1	86	193	137	45	2	1	-	-	-
2006 1/.....	446	1	116	180	113	32	3	-	-	1	-
2005 1/.....	453	-	110	179	126	35	2	1	-	-	-
2004 1/.....	411	-	88	168	99	53	1	1	1	-	-
2003 1/.....	317	-	66	132	94	23	1	1	-	-	-
2002.....	203	-	40	97	60	5	-	-	-	1	-
2001.....	235	2	64	114	51	3	-	-	-	1	-
2000.....	243	-	55	108	77	3	-	-	-	-	-
1999.....	220	-	45	100	71	4	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pregnancy, childbirth and the puerperium (000-099)													
White, male													
2007 1/.....
2006 1/.....
2005 1/.....
2004 1/.....
2003 1/.....
2002.....
2001.....
2000.....
1999.....
White, female													
2007 1/.....	465	1	86	193	137	45	2	1	-	-	-
2006 1/.....	446	1	116	180	113	32	3	-	-	1	-
2005 1/.....	453	-	110	179	126	35	2	1	-	-	-
2004 1/.....	411	-	88	168	99	53	1	1	1	-	-
2003 1/.....	317	-	66	132	94	23	1	1	-	-	-
2002.....	203	-	40	97	60	5	-	-	-	1	-
2001.....	235	2	64	114	51	3	-	-	-	1	-
2000.....	243	-	55	108	77	3	-	-	-	-	-
1999.....	220	-	45	100	71	4	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Pregnancy, childbirth and the puerperium (000-099)

Black, female

2007 1/.....	251	-	69	108	61	13	-	-	-	-	-
2006 1/.....	259	1	58	118	61	21	-	-	-	-	-
2005 1/.....	267	-	69	117	65	15	1	-	-	-	-
2004 1/.....	251	-	74	109	45	23	-	-	-	-	-
2003 1/.....	193	1	44	83	58	7	-	-	-	-	-
2002.....	157	1	46	66	42	2	-	-	-	-	-
2001.....	158	1	45	66	43	3	-	-	-	-	-
2000.....	142	-	36	65	38	3	-	-	-	-	-
1999.....	162	-	47	71	42	2	-	-	-	-	-

American Indian 2/, both sexes

2007 1/.....	18	-	4	7	7	-	-	-	-	-	-
2006 1/.....	12	-	2	4	3	3	-	-	-	-	-
2005 1/.....	7	-	2	1	4	-	-	-	-	-	-
2004 1/.....	6	-	1	1	3	1	-	-	-	-	-
2003 1/.....	8	-	1	2	5	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-
2001.....	5	-	-	5	-	-	-	-	-	-	-
2000.....	6	-	2	4	-	-	-	-	-	-	-
1999.....	5	-	3	1	1	-	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Pregnancy, childbirth and the puerperium (000-099)

American Indian 2/, male

2007 1/.....
2006 1/.....
2005 1/.....
2004 1/.....
2003 1/.....
2002.....
2001.....
2000.....
1999.....

American Indian 2/, female

2007 1/.....	18	-	4	7	7	-	-	-	-	-	-
2006 1/.....	12	-	2	4	3	3	-	-	-	-	-
2005 1/.....	7	-	2	1	4	-	-	-	-	-	-
2004 1/.....	6	-	1	1	3	1	-	-	-	-	-
2003 1/.....	8	-	1	2	5	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-
2001.....	5	-	-	5	-	-	-	-	-	-	-
2000.....	6	-	2	4	-	-	-	-	-	-	-
1999.....	5	-	3	1	1	-	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Pregnancy, childbirth and the puerperium (000-099)

Asian or Pacific Islander 3/, female

2007 1/.....	35	-	1	18	15	1	-	-	-	-	-
2006 1/.....	43	-	3	22	15	3	-	-	-	-	-
2005 1/.....	33	-	2	15	12	4	-	-	-	-	-
2004 1/.....	29	2	5	9	10	2	-	-	1	-	-
2003 1/.....	27	-	5	12	9	-	1	-	-	-	-
2002.....	19	-	3	12	3	1	-	-	-	-	-
2001.....	18	-	1	5	12	-	-	-	-	-	-
2000.....	13	-	2	5	6	-	-	-	-	-	-
1999.....	19	-	2	9	7	1	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Pregnancy with abortive outcome (000-007)

All races, female

2007.....	31	-	7	16	6	2	-	-	-	-	-
2006.....	26	-	6	15	4	1	-	-	-	-	-
2005.....	33	-	6	14	11	2	-	-	-	-	-
2004.....	32	-	14	11	6	1	-	-	-	-	-
2003.....	40	-	11	19	9	1	-	-	-	-	-
2002.....	22	1	4	9	7	1	-	-	-	-	-
2001.....	38	2	13	14	9	-	-	-	-	-	-
2000.....	37	-	6	17	14	-	-	-	-	-	-
1999.....	31	-	8	15	8	-	-	-	-	-	-

White, both sexes

2007 1/.....	12	-	1	7	3	1	-	-	-	-	-
2006 1/.....	10	-	4	5	-	1	-	-	-	-	-
2005 1/.....	15	-	4	4	5	2	-	-	-	-	-
2004 1/.....	11	-	5	5	-	1	-	-	-	-	-
2003 1/.....	18	-	3	11	4	-	-	-	-	-	-
2002.....	7	-	1	5	1	-	-	-	-	-	-
2001.....	20	1	7	8	4	-	-	-	-	-	-
2000.....	18	-	1	8	9	-	-	-	-	-	-
1999.....	15	-	4	6	5	-	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Pregnancy with abortive outcome (000-007)

White, male

2007 1/.....
2006 1/.....
2005 1/.....
2004 1/.....
2003 1/.....
2002.....
2001.....
2000.....
1999.....

White, female

2007 1/.....	12	-	1	7	3	1	-	-	-	-	-
2006 1/.....	10	-	4	5	-	1	-	-	-	-	-
2005 1/.....	15	-	4	4	5	2	-	-	-	-	-
2004 1/.....	11	-	5	5	-	1	-	-	-	-	-
2003 1/.....	18	-	3	11	4	-	-	-	-	-	-
2002.....	7	-	1	5	1	-	-	-	-	-	-
2001.....	20	1	7	8	4	-	-	-	-	-	-
2000.....	18	-	1	8	9	-	-	-	-	-	-
1999.....	15	-	4	6	5	-	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Pregnancy with abortive outcome (000-007)

Black, female

2007 1/.....	18	-	6	8	3	1	-	-	-	-	-
2006 1/.....	12	-	2	7	3	-	-	-	-	-	-
2005 1/.....	13	-	1	8	4	-	-	-	-	-	-
2004 1/.....	19	-	8	6	5	-	-	-	-	-	-
2003 1/.....	18	-	8	5	4	1	-	-	-	-	-
2002.....	12	1	2	4	5	-	-	-	-	-	-
2001.....	16	1	6	5	4	-	-	-	-	-	-
2000.....	19	-	5	9	5	-	-	-	-	-	-
1999.....	15	-	4	8	3	-	-	-	-	-	-

American Indian 2/, both sexes

2007 1/.....	-	-	-	-	-	-	-	-	-	-	-
2006 1/.....	1	-	-	1	-	-	-	-	-	-	-
2005 1/.....	1	-	1	-	-	-	-	-	-	-	-
2004 1/.....	-	-	-	-	-	-	-	-	-	-	-
2003 1/.....	1	-	-	-	1	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	1	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	1	-	-	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Pregnancy with abortive outcome (000-007)

American Indian 2/, male

2007 1/.....
2006 1/.....
2005 1/.....
2004 1/.....
2003 1/.....
2002.....
2001.....
2000.....
1999.....

American Indian 2/, female

2007 1/.....	-	-	-	-	-	-	-	-	-	-	-
2006 1/.....	1	-	-	1	-	-	-	-	-	-	-
2005 1/.....	1	-	1	-	-	-	-	-	-	-	-
2004 1/.....	-	-	-	-	-	-	-	-	-	-	-
2003 1/.....	1	-	-	-	1	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	1	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	1	-	-	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Pregnancy with abortive outcome (000-007)

Asian or Pacific Islander 3/, both sexes

Table with 14 columns (years 2007-1999) and 14 rows (age groups All ages, Under 1 year, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over, Age not stated). Data values are integers.

Asian or Pacific Islander 3/, male

Table with 14 columns (years 2007-1999) and 14 rows (age groups All ages, Under 1 year, 1-4 years, 5-14 years, 15-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years, 85 years and over, Age not stated). Data values are integers.

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other complications of pregnancy, childbirth and the puerperium (010-099)

All races, female

2007.....	738	1	153	310	214	57	2	1	-	-	-
2006.....	734	2	173	309	188	58	3	-	-	1	-
2005.....	727	-	177	298	196	52	3	1	-	-	-
2004.....	665	2	154	276	151	78	1	1	2	-	-
2003.....	505	1	105	210	157	29	2	1	-	-	-
2002.....	357	-	85	166	98	7	-	-	-	1	-
2001.....	378	1	97	176	97	6	-	-	-	1	-
2000.....	367	-	89	165	107	6	-	-	-	-	-
1999.....	375	-	89	166	113	7	-	-	-	-	-

White, both sexes

2007 1/.....	453	1	85	186	134	44	2	1	-	-	-
2006 1/.....	436	1	112	175	113	31	3	-	-	1	-
2005 1/.....	438	-	106	175	121	33	2	1	-	-	-
2004 1/.....	400	-	83	163	99	52	1	1	1	-	-
2003 1/.....	299	-	63	121	90	23	1	1	-	-	-
2002.....	196	-	39	92	59	5	-	-	-	1	-
2001.....	215	1	57	106	47	3	-	-	-	1	-
2000.....	225	-	54	100	68	3	-	-	-	-	-
1999.....	205	-	41	94	66	4	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other complications of pregnancy, childbirth and the puerperium (010-099)

White, male

2007 1/.....
2006 1/.....
2005 1/.....
2004 1/.....
2003 1/.....
2002.....
2001.....
2000.....
1999.....

White, female

2007 1/.....	453	1	85	186	134	44	2	1	-	-	-
2006 1/.....	436	1	112	175	113	31	3	-	-	1	-
2005 1/.....	438	-	106	175	121	33	2	1	-	-	-
2004 1/.....	400	-	83	163	99	52	1	1	1	-	-
2003 1/.....	299	-	63	121	90	23	1	1	-	-	-
2002.....	196	-	39	92	59	5	-	-	-	1	-
2001.....	215	1	57	106	47	3	-	-	-	1	-
2000.....	225	-	54	100	68	3	-	-	-	-	-
1999.....	205	-	41	94	66	4	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other complications of pregnancy, childbirth and the puerperium (010-099)

Black, female

2007 1/.....	233	-	63	100	58	12	-	-	-	-	-
2006 1/.....	247	1	56	111	58	21	-	-	-	-	-
2005 1/.....	254	-	68	109	61	15	1	-	-	-	-
2004 1/.....	232	-	66	103	40	23	-	-	-	-	-
2003 1/.....	175	1	36	78	54	6	-	-	-	-	-
2002.....	145	-	44	62	37	2	-	-	-	-	-
2001.....	142	-	39	61	39	3	-	-	-	-	-
2000.....	123	-	31	56	33	3	-	-	-	-	-
1999.....	147	-	43	63	39	2	-	-	-	-	-

American Indian 2/, both sexes

2007 1/.....	18	-	4	7	7	-	-	-	-	-	-
2006 1/.....	11	-	2	3	3	3	-	-	-	-	-
2005 1/.....	6	-	1	1	4	-	-	-	-	-	-
2004 1/.....	6	-	1	1	3	1	-	-	-	-	-
2003 1/.....	7	-	1	2	4	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-
2001.....	4	-	-	4	-	-	-	-	-	-	-
2000.....	6	-	2	4	-	-	-	-	-	-	-
1999.....	4	-	3	-	1	-	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other complications of pregnancy, childbirth and the puerperium (010-099)

American Indian 2/, male

2007 1/.....
2006 1/.....
2005 1/.....
2004 1/.....
2003 1/.....
2002.....
2001.....
2000.....
1999.....

American Indian 2/, female

2007 1/.....	18	-	4	7	7	-	-	-	-	-	-
2006 1/.....	11	-	2	3	3	3	-	-	-	-	-
2005 1/.....	6	-	1	1	4	-	-	-	-	-	-
2004 1/.....	6	-	1	1	3	1	-	-	-	-	-
2003 1/.....	7	-	1	2	4	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-
2001.....	4	-	-	4	-	-	-	-	-	-	-
2000.....	6	-	2	4	-	-	-	-	-	-	-
1999.....	4	-	3	-	1	-	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other complications of pregnancy, childbirth and the puerperium (010-099)

Asian or Pacific Islander 3/, female

2007 1/.....	34	-	1	17	15	1	-	-	-	-	-
2006 1/.....	40	-	3	20	14	3	-	-	-	-	-
2005 1/.....	29	-	2	13	10	4	-	-	-	-	-
2004 1/.....	27	2	4	9	9	2	-	-	1	-	-
2003 1/.....	24	-	5	9	9	-	1	-	-	-	-
2002.....	16	-	2	12	2	-	-	-	-	-	-
2001.....	17	-	1	5	11	-	-	-	-	-	-
2000.....	13	-	2	5	6	-	-	-	-	-	-
1999.....	19	-	2	9	7	1	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Certain conditions originating in the perinatal period (P00-P96)

All races, both sexes

2007.....	14,599	14,466	70	22	17	3	4	5	3	3	1	-	5
2006.....	14,442	14,321	65	24	14	4	4	4	1	-	2	-	3
2005.....	14,549	14,423	58	26	18	6	6	6	2	-	1	-	3
2004.....	14,213	14,079	61	37	11	10	6	3	3	1	1	1	-
2003.....	14,378	14,254	79	19	16	3	3	2	-	2	-	-	-
2002.....	14,254	14,106	65	31	21	10	10	2	1	3	2	1	2
2001.....	13,887	13,734	72	36	17	6	9	5	2	3	1	1	1
2000.....	14,069	13,918	79	41	15	7	4	2	2	-	-	-	1
1999.....	14,259	14,084	92	43	16	7	3	2	3	2	1	1	5

All races, male

2007.....	8,218	8,140	34	14	10	3	3	4	3	3	1	-	3
2006.....	8,097	8,025	37	14	11	3	2	2	1	-	1	-	1
2005.....	8,187	8,113	36	11	12	3	4	5	1	-	1	-	1
2004.....	8,093	8,013	38	23	5	3	5	1	3	1	1	-	-
2003.....	8,197	8,129	43	11	8	2	1	1	-	2	-	-	-
2002.....	8,047	7,960	40	19	12	4	5	2	1	1	1	-	2
2001.....	7,868	7,775	44	24	7	2	6	5	1	2	1	-	1
2000.....	7,968	7,872	52	23	11	6	2	2	-	-	-	-	-
1999.....	8,044	7,941	56	23	10	2	2	2	1	1	1	1	4

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Certain conditions originating in the perinatal period (P00-P96)

All races, female

2007.....	6,381	6,326	36	8	7	-	1	1	-	-	-	-	2
2006.....	6,345	6,296	28	10	3	1	2	2	-	-	1	-	2
2005.....	6,362	6,310	22	15	6	3	2	1	1	-	-	-	2
2004.....	6,120	6,066	23	14	6	7	1	2	-	-	-	1	-
2003.....	6,181	6,125	36	8	8	1	2	1	-	-	-	-	-
2002.....	6,207	6,146	25	12	9	6	5	-	-	2	1	1	-
2001.....	6,019	5,959	28	12	10	4	3	-	1	1	-	1	-
2000.....	6,101	6,046	27	18	4	1	2	-	2	-	-	-	1
1999.....	6,215	6,143	36	20	6	5	1	-	2	1	-	-	1

White, both sexes

2007 1/.....	8,890	8,801	40	16	15	3	2	5	2	1	1	-	4
2006 1/.....	8,730	8,646	40	18	11	4	4	4	1	-	-	-	2
2005 1/.....	8,906	8,815	35	22	15	5	5	6	2	-	1	-	-
2004 1/.....	8,732	8,643	33	26	10	8	4	3	3	-	1	1	-
2003 1/.....	8,975	8,891	45	16	15	3	3	1	-	1	-	-	-
2002.....	8,777	8,679	36	21	17	7	8	2	1	2	2	1	1
2001.....	8,551	8,444	47	21	13	6	8	5	1	3	1	1	1
2000.....	8,532	8,431	46	30	13	3	4	2	2	-	-	-	1
1999.....	8,648	8,522	56	33	15	7	3	2	3	2	1	1	3

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Certain conditions originating in the perinatal period (P00-P96)

White, male

2007 1/.....	4,999	4,945	19	11	9	3	1	4	2	1	1	-	3
2006 1/.....	4,938	4,882	26	12	9	3	2	2	1	-	-	-	1
2005 1/.....	5,026	4,971	20	11	11	3	3	5	1	-	1	-	-
2004 1/.....	5,025	4,971	22	16	5	3	3	1	3	-	1	-	-
2003 1/.....	5,134	5,087	25	10	8	2	1	-	-	1	-	-	-
2002.....	5,011	4,959	19	12	9	4	3	2	1	-	1	-	1
2001.....	4,846	4,780	28	16	6	2	5	5	-	2	1	-	1
2000.....	4,828	4,762	30	20	10	2	2	2	-	-	-	-	-
1999.....	4,884	4,815	30	17	10	2	2	2	1	1	1	1	2

White, female

2007 1/.....	3,891	3,856	21	5	6	-	1	1	-	-	-	-	1
2006 1/.....	3,792	3,764	14	6	2	1	2	2	-	-	-	-	1
2005 1/.....	3,880	3,844	15	11	4	2	2	1	1	-	-	-	-
2004 1/.....	3,707	3,672	11	10	5	5	1	2	-	-	-	1	-
2003 1/.....	3,841	3,804	20	6	7	1	2	1	-	-	-	-	-
2002.....	3,766	3,720	17	9	8	3	5	-	-	2	1	1	-
2001.....	3,705	3,664	19	5	7	4	3	-	1	1	-	1	-
2000.....	3,704	3,669	16	10	3	1	2	-	2	-	-	-	1
1999.....	3,764	3,707	26	16	5	5	1	-	2	1	-	-	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)													
All races, both sexes													
2007.....	10,421	5,785	546	374	402	417	513	685	686	341	406	265	1
2006.....	10,489	5,819	515	344	460	437	468	668	720	367	392	297	2
2005.....	10,410	5,552	522	396	504	436	517	693	716	338	408	327	1
2004.....	10,498	5,622	569	389	483	420	493	692	634	428	472	296	-
2003.....	10,518	5,621	541	386	451	426	564	667	666	413	464	319	-
2002.....	10,687	5,623	530	417	492	475	572	709	634	440	489	305	1
2001.....	10,442	5,513	557	376	505	458	586	636	581	448	465	317	-
2000.....	10,578	5,743	495	399	441	477	569	612	569	464	488	320	1
1999.....	10,393	5,473	549	428	434	465	552	647	557	469	491	326	2
All races, male													
2007.....	5,455	3,066	286	194	241	243	257	359	376	166	178	88	1
2006.....	5,587	3,098	244	181	279	263	265	376	395	180	191	114	1
2005.....	5,447	2,965	264	195	326	244	272	356	370	153	191	111	-
2004.....	5,423	2,898	318	211	282	235	279	375	320	208	195	102	-
2003.....	5,460	3,020	275	198	272	233	304	349	336	198	178	97	-
2002.....	5,515	2,942	282	215	282	284	296	381	321	213	201	97	1
2001.....	5,448	2,954	282	210	306	249	317	323	276	228	203	100	-
2000.....	5,517	3,028	259	207	268	279	316	347	264	222	197	130	-
1999.....	5,346	2,875	310	215	262	242	296	347	283	209	204	102	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All races, female

2007.....	4,966	2,719	260	180	161	174	256	326	310	175	228	177	-
2006.....	4,902	2,721	271	163	181	174	203	292	325	187	201	183	1
2005.....	4,963	2,587	258	201	178	192	245	337	346	185	217	216	1
2004.....	5,075	2,724	251	178	201	185	214	317	314	220	277	194	-
2003.....	5,058	2,601	266	188	179	193	260	318	330	215	286	222	-
2002.....	5,172	2,681	248	202	210	191	276	328	313	227	288	208	-
2001.....	4,994	2,559	275	166	199	209	269	313	305	220	262	217	-
2000.....	5,061	2,715	236	192	173	198	253	265	305	242	291	190	1
1999.....	5,047	2,598	239	213	172	223	256	300	274	260	287	224	1

White, both sexes

2007 1/.....	8,155	4,351	396	274	299	322	421	573	615	295	365	243	1
2006 1/.....	8,246	4,383	376	259	343	349	381	555	646	324	356	272	2
2005 1/.....	8,254	4,187	385	285	386	339	427	606	645	316	379	298	1
2004 1/.....	8,331	4,271	400	287	352	345	409	599	574	392	424	278	-
2003 1/.....	8,378	4,292	392	288	342	343	477	569	596	360	424	295	-
2002.....	8,556	4,330	395	321	383	384	463	605	552	393	444	285	1
2001.....	8,338	4,247	387	282	374	375	479	542	527	403	429	293	-
2000.....	8,396	4,371	347	292	332	391	459	539	496	410	457	301	1
1999.....	8,297	4,211	385	301	332	386	461	557	500	416	445	301	2

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

White, male

2007 1/.....	4,297	2,330	211	145	175	185	214	302	340	151	161	82	1
2006 1/.....	4,403	2,326	191	148	206	212	218	311	353	159	175	103	1
2005 1/.....	4,321	2,239	201	145	252	193	220	312	333	146	181	99	-
2004 1/.....	4,260	2,182	222	162	196	189	227	330	288	192	174	98	-
2003 1/.....	4,288	2,282	198	146	206	189	260	293	298	169	157	90	-
2002.....	4,416	2,297	211	159	218	231	242	319	278	190	178	92	1
2001.....	4,376	2,301	195	169	222	212	261	279	249	213	187	88	-
2000.....	4,368	2,316	180	147	201	229	259	300	237	188	188	123	-
1999.....	4,269	2,240	218	148	204	202	245	296	257	186	179	93	1

White, female

2007 1/.....	3,858	2,021	185	129	124	137	207	271	275	144	204	161	-
2006 1/.....	3,843	2,057	185	111	137	137	163	244	293	165	181	169	1
2005 1/.....	3,933	1,948	184	140	134	146	207	294	312	170	198	199	1
2004 1/.....	4,071	2,089	178	125	156	156	182	269	286	200	250	180	-
2003 1/.....	4,090	2,010	194	142	136	154	217	276	298	191	267	205	-
2002.....	4,140	2,033	184	162	165	153	221	286	274	203	266	193	-
2001.....	3,962	1,946	192	113	152	163	218	263	278	190	242	205	-
2000.....	4,028	2,055	167	145	131	162	200	239	259	222	269	178	1
1999.....	4,028	1,971	167	153	128	184	216	261	243	230	266	208	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Black, both sexes

2007 1/.....	1,807	1,137	119	77	83	83	79	82	60	35	34	18	-
2006 1/.....	1,808	1,161	107	67	88	73	74	96	63	28	30	21	-
2005 1/.....	1,719	1,080	106	88	100	81	75	71	59	16	21	22	-
2004 1/.....	1,745	1,077	132	85	111	60	69	83	47	28	39	14	-
2003 1/.....	1,700	1,039	120	74	92	76	70	79	60	40	30	20	-
2002.....	1,758	1,047	114	83	100	75	93	87	69	40	32	18	-
2001.....	1,720	1,032	131	75	114	68	91	77	48	36	28	20	-
2000.....	1,790	1,102	126	91	91	77	93	67	59	43	25	16	-
1999.....	1,720	1,025	138	106	83	61	77	76	46	46	38	24	-

Black, male

2007 1/.....	924	582	58	37	53	53	39	41	32	11	14	4	-
2006 1/.....	947	614	42	29	53	44	38	55	33	16	14	9	-
2005 1/.....	891	564	48	41	64	43	44	33	31	5	9	9	-
2004 1/.....	942	573	77	39	73	37	42	40	25	14	18	4	-
2003 1/.....	923	574	62	39	55	38	33	42	33	25	16	6	-
2002.....	909	522	59	48	60	43	46	49	40	21	16	5	-
2001.....	882	537	69	35	72	31	47	35	23	12	12	9	-
2000.....	944	567	70	52	58	46	51	41	18	26	9	6	-
1999.....	899	531	79	54	51	32	42	44	22	16	19	9	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Black, female

2007 1/.....	883	555	61	40	30	30	40	41	28	24	20	14	-
2006 1/.....	861	547	65	38	35	29	36	41	30	12	16	12	-
2005 1/.....	828	516	58	47	36	38	31	38	28	11	12	13	-
2004 1/.....	803	504	55	46	38	23	27	43	22	14	21	10	-
2003 1/.....	777	465	58	35	37	38	37	37	27	15	14	14	-
2002.....	849	525	55	35	40	32	47	38	29	19	16	13	-
2001.....	838	495	62	40	42	37	44	42	25	24	16	11	-
2000.....	846	535	56	39	33	31	42	26	41	17	16	10	-
1999.....	821	494	59	52	32	29	35	32	24	30	19	15	-

American Indian 2/, both sexes

2007 1/.....	124	82	8	6	6	3	4	8	2	4	-	1	-
2006 1/.....	117	73	10	5	11	2	4	5	2	5	-	-	-
2005 1/.....	117	77	11	6	1	3	8	5	5	-	1	-	-
2004 1/.....	125	90	8	7	5	7	4	1	3	-	-	-	-
2003 1/.....	118	72	10	11	6	1	6	4	3	3	-	2	-
2002.....	93	61	5	6	1	6	3	2	4	-	5	-	-
2001.....	98	60	15	3	5	5	2	4	1	1	2	-	-
2000.....	95	68	4	3	3	-	4	1	5	5	2	-	-
1999.....	106	67	5	6	7	7	1	6	5	1	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

American Indian 2/, male

2007 1/.....	62	39	5	3	4	2	1	5	1	1	-	1	-
2006 1/.....	65	41	3	2	9	-	3	2	2	3	-	-	-
2005 1/.....	61	44	4	1	-	1	4	4	3	-	-	-	-
2004 1/.....	70	50	3	5	3	5	3	-	1	-	-	-	-
2003 1/.....	58	35	4	5	4	1	4	3	2	-	-	-	-
2002.....	42	29	2	3	-	3	1	1	1	-	2	-	-
2001.....	46	30	5	1	5	1	1	2	-	1	-	-	-
2000.....	56	42	1	2	2	-	1	1	4	3	-	-	-
1999.....	50	28	3	4	3	3	-	5	2	1	1	-	-

American Indian 2/, female

2007 1/.....	62	43	3	3	2	1	3	3	1	3	-	-	-
2006 1/.....	52	32	7	3	2	2	1	3	-	2	-	-	-
2005 1/.....	56	33	7	5	1	2	4	1	2	-	1	-	-
2004 1/.....	55	40	5	2	2	2	1	1	2	-	-	-	-
2003 1/.....	60	37	6	6	2	-	2	1	1	3	-	2	-
2002.....	51	32	3	3	1	3	2	1	3	-	3	-	-
2001.....	52	30	10	2	-	4	1	2	1	-	2	-	-
2000.....	39	26	3	1	1	-	3	-	1	2	2	-	-
1999.....	56	39	2	2	4	4	1	1	3	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	335	215	23	17	14	9	9	22	9	7	7	3	-
2006 1/.....	318	202	22	13	18	13	9	12	9	10	6	4	-
2005 1/.....	320	208	20	17	17	13	7	11	7	6	7	7	-
2004 1/.....	297	184	29	10	15	8	11	9	10	8	9	4	-
2003 1/.....	322	218	19	13	11	6	11	15	7	10	10	2	-
2002.....	280	185	16	7	8	10	13	15	9	7	8	2	-
2001.....	286	174	24	16	12	10	14	13	5	8	6	4	-
2000.....	297	202	18	13	15	9	13	5	9	6	4	3	-
1999.....	270	170	21	15	12	11	13	8	6	6	7	1	-

Asian or Pacific Islander 3/, male

2007 1/.....	172	115	12	9	9	3	3	11	3	3	3	1	-
2006 1/.....	172	117	8	2	11	7	6	8	7	2	2	2	-
2005 1/.....	174	118	11	8	10	7	4	7	3	2	1	3	-
2004 1/.....	151	93	16	5	10	4	7	5	6	2	3	-	-
2003 1/.....	191	129	11	8	7	5	7	11	3	4	5	1	-
2002.....	148	94	10	5	4	7	7	12	2	2	5	-	-
2001.....	144	86	13	5	7	5	8	7	4	2	4	3	-
2000.....	149	103	8	6	7	4	5	5	5	5	-	1	-
1999.....	128	76	10	9	4	5	9	2	2	6	5	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Asian or Pacific Islander 3/, female

2007 1/.....	163	100	11	8	5	6	6	11	6	4	4	2	-
2006 1/.....	146	85	14	11	7	6	3	4	2	8	4	2	-
2005 1/.....	146	90	9	9	7	6	3	4	4	4	6	4	-
2004 1/.....	146	91	13	5	5	4	4	4	4	6	6	4	-
2003 1/.....	131	89	8	5	4	1	4	4	4	6	5	1	-
2002.....	132	91	6	2	4	3	6	3	7	5	3	2	-
2001.....	142	88	11	11	5	5	6	6	1	6	2	1	-
2000.....	148	99	10	7	8	5	8	-	4	1	4	2	-
1999.....	142	94	11	6	8	6	4	6	4	-	2	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All races, both sexes

2007.....	33,500	3,617	237	110	575	877	1,424	2,195	2,229	2,499	5,834	13,860	43
2006.....	31,725	3,462	239	126	659	1,013	1,627	2,266	2,039	2,307	5,435	12,508	44
2005.....	31,999	3,589	233	155	775	1,178	2,052	2,831	2,199	2,390	5,088	11,436	73
2004.....	30,126	3,361	189	120	658	1,092	1,900	2,402	2,041	2,323	5,159	10,762	119
2003.....	31,444	3,318	256	178	837	1,327	2,382	2,874	2,067	2,522	5,006	10,565	112
2002.....	29,975	3,456	248	139	740	1,319	2,402	2,672	2,002	2,398	4,936	9,554	109
2001.....	32,326	3,491	272	181	949	1,530	3,126	3,504	2,292	2,774	5,003	9,089	115
2000.....	31,876	3,639	248	177	868	1,554	2,987	3,142	2,253	3,040	5,029	8,824	115
1999.....	26,672	3,538	236	137	609	1,095	2,062	2,151	1,860	2,698	4,413	7,798	75

All races, male

2007.....	14,282	2,098	133	63	396	586	890	1,369	1,394	1,377	2,435	3,511	30
2006.....	13,981	2,012	134	75	436	712	1,039	1,431	1,256	1,287	2,343	3,227	29
2005.....	14,647	2,139	132	94	546	807	1,263	1,822	1,367	1,321	2,150	2,949	57
2004.....	13,712	2,021	100	59	452	732	1,235	1,546	1,268	1,264	2,181	2,768	86
2003.....	14,583	1,974	143	92	563	907	1,526	1,840	1,294	1,431	2,160	2,570	83
2002.....	14,269	2,032	145	82	501	905	1,540	1,723	1,273	1,324	2,228	2,439	77
2001.....	15,779	2,041	162	99	654	1,040	2,007	2,286	1,457	1,497	2,184	2,274	78
2000.....	15,863	2,156	137	99	581	1,060	1,994	2,123	1,429	1,708	2,244	2,242	90
1999.....	12,877	2,072	136	77	422	721	1,339	1,446	1,181	1,513	1,930	1,984	56

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All races, female

2007.....	19,218	1,519	104	47	179	291	534	826	835	1,122	3,399	10,349	13
2006.....	17,744	1,450	105	51	223	301	588	835	783	1,020	3,092	9,281	15
2005.....	17,352	1,450	101	61	229	371	789	1,009	832	1,069	2,938	8,487	16
2004.....	16,414	1,340	89	61	206	360	665	856	773	1,059	2,978	7,994	33
2003.....	16,861	1,344	113	86	274	420	856	1,034	773	1,091	2,846	7,995	29
2002.....	15,706	1,424	103	57	239	414	862	949	729	1,074	2,708	7,115	32
2001.....	16,547	1,450	110	82	295	490	1,119	1,218	835	1,277	2,819	6,815	37
2000.....	16,013	1,483	111	78	287	494	993	1,019	824	1,332	2,785	6,582	25
1999.....	13,795	1,466	100	60	187	374	723	705	679	1,185	2,483	5,814	19

White, both sexes

2007 1/.....	28,126	2,347	172	80	439	641	1,082	1,643	1,780	2,065	5,148	12,700	29
2006 1/.....	26,672	2,244	152	86	512	785	1,265	1,763	1,649	1,917	4,788	11,476	35
2005 1/.....	26,595	2,331	153	91	591	907	1,596	2,191	1,731	1,953	4,498	10,490	63
2004 1/.....	25,030	2,091	127	77	515	835	1,484	1,883	1,612	1,926	4,566	9,829	85
2003 1/.....	26,147	2,125	166	118	659	1,014	1,849	2,173	1,645	2,095	4,488	9,719	96
2002.....	24,749	2,224	161	84	561	1,002	1,854	1,978	1,598	2,018	4,400	8,786	83
2001.....	26,877	2,231	185	124	747	1,202	2,512	2,803	1,848	2,304	4,465	8,368	88
2000.....	26,285	2,294	182	120	656	1,185	2,395	2,451	1,783	2,518	4,474	8,144	83
1999.....	21,591	2,227	159	96	433	798	1,507	1,572	1,447	2,254	3,873	7,164	61

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

White, male

2007 1/.....	11,642	1,390	99	46	306	437	676	1,028	1,115	1,146	2,163	3,215	21
2006 1/.....	11,422	1,337	84	54	344	560	816	1,109	1,032	1,054	2,060	2,947	25
2005 1/.....	11,861	1,435	88	59	421	638	1,007	1,435	1,073	1,081	1,891	2,682	51
2004 1/.....	11,082	1,250	68	42	358	576	985	1,208	997	1,060	1,940	2,532	66
2003 1/.....	11,802	1,313	91	61	453	706	1,189	1,416	1,021	1,194	1,935	2,352	71
2002.....	11,421	1,323	89	52	391	700	1,198	1,271	1,016	1,117	1,965	2,241	58
2001.....	12,868	1,320	106	66	516	840	1,635	1,851	1,158	1,256	1,977	2,082	61
2000.....	12,827	1,363	106	68	458	830	1,649	1,673	1,125	1,408	2,010	2,071	66
1999.....	10,094	1,321	96	54	309	529	1,005	1,050	919	1,269	1,689	1,806	47

White, female

2007 1/.....	16,484	957	73	34	133	204	406	615	665	919	2,985	9,485	8
2006 1/.....	15,250	907	68	32	168	225	449	654	617	863	2,728	8,529	10
2005 1/.....	14,734	896	65	32	170	269	589	756	658	872	2,607	7,808	12
2004 1/.....	13,948	841	59	35	157	259	499	675	615	866	2,626	7,297	19
2003 1/.....	14,345	812	75	57	206	308	660	757	624	901	2,553	7,367	25
2002.....	13,328	901	72	32	170	302	656	707	582	901	2,435	6,545	25
2001.....	14,009	911	79	58	231	362	877	952	690	1,048	2,488	6,286	27
2000.....	13,458	931	76	52	198	355	746	778	658	1,110	2,464	6,073	17
1999.....	11,497	906	63	42	124	269	502	522	528	985	2,184	5,358	14

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Black, both sexes

2007 1/.....	4,651	1,106	49	24	117	214	302	493	396	381	597	961	11
2006 1/.....	4,379	1,059	74	36	120	191	319	457	342	345	565	863	8
2005 1/.....	4,798	1,121	65	57	162	235	400	596	429	394	524	805	10
2004 1/.....	4,490	1,135	50	36	116	223	371	479	387	354	527	783	29
2003 1/.....	4,708	1,051	80	51	154	271	487	644	391	400	445	723	11
2002.....	4,638	1,095	67	50	154	279	495	637	362	340	478	660	21
2001.....	4,691	1,104	69	42	166	270	523	620	375	408	468	624	22
2000.....	5,004	1,212	57	46	186	316	528	623	424	479	509	596	28
1999.....	4,591	1,167	68	33	154	266	514	535	378	403	490	572	11

Black, male

2007 1/.....	2,264	606	25	13	78	133	184	302	252	211	238	215	7
2006 1/.....	2,202	576	42	21	75	127	191	296	195	206	253	217	3
2005 1/.....	2,475	632	36	30	113	146	220	359	269	217	232	215	6
2004 1/.....	2,312	689	28	16	76	137	216	314	246	179	212	183	16
2003 1/.....	2,459	574	44	27	94	176	300	391	250	223	186	186	8
2002.....	2,536	630	45	28	93	177	309	416	232	185	238	169	14
2001.....	2,465	626	44	26	113	154	303	373	264	210	176	163	13
2000.....	2,693	710	29	28	108	198	293	403	275	273	215	140	21
1999.....	2,517	669	35	20	101	176	307	364	245	221	215	157	7

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Black, female

2007 1/.....	2,387	500	24	11	39	81	118	191	144	170	359	746	4
2006 1/.....	2,177	483	32	15	45	64	128	161	147	139	312	646	5
2005 1/.....	2,323	489	29	27	49	89	180	237	160	177	292	590	4
2004 1/.....	2,178	446	22	20	40	86	155	165	141	175	315	600	13
2003 1/.....	2,249	477	36	24	60	95	187	253	141	177	259	537	3
2002.....	2,102	465	22	22	61	102	186	221	130	155	240	491	7
2001.....	2,226	478	25	16	53	116	220	247	111	198	292	461	9
2000.....	2,311	502	28	18	78	118	235	220	149	206	294	456	7
1999.....	2,074	498	33	13	53	90	207	171	133	182	275	415	4

American Indian 2/, both sexes

2007 1/.....	261	92	7	-	10	12	23	32	24	14	16	31	-
2006 1/.....	224	90	8	2	11	10	20	17	16	9	16	24	1
2005 1/.....	220	71	3	1	10	10	24	26	15	14	17	29	-
2004 1/.....	214	64	5	3	16	16	22	15	13	14	14	29	3
2003 1/.....	231	76	4	4	10	12	28	30	16	5	14	30	2
2002.....	248	72	7	1	9	20	31	28	21	19	13	24	3
2001.....	263	81	6	4	9	20	28	23	26	22	16	24	4
2000.....	209	61	3	1	9	17	21	26	19	11	13	27	1
1999.....	215	72	3	2	6	12	15	20	14	20	27	23	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

American Indian 2/, male

2007 1/.....	164	60	5	-	7	11	15	22	16	5	6	17	-
2006 1/.....	125	55	4	-	6	8	17	9	10	3	6	6	1
2005 1/.....	109	38	2	1	3	7	12	14	8	11	6	7	-
2004 1/.....	122	41	1	-	13	9	17	11	7	8	5	7	3
2003 1/.....	135	44	3	3	8	9	21	17	11	2	9	6	2
2002.....	130	41	3	-	6	13	18	14	12	11	6	3	3
2001.....	160	52	5	1	7	16	22	19	11	12	9	3	3
2000.....	127	42	1	-	6	7	14	21	13	7	6	9	1
1999.....	117	39	1	1	5	5	9	17	7	10	15	7	1

American Indian 2/, female

2007 1/.....	97	32	2	-	3	1	8	10	8	9	10	14	-
2006 1/.....	99	35	4	2	5	2	3	8	6	6	10	18	-
2005 1/.....	111	33	1	-	7	3	12	12	7	3	11	22	-
2004 1/.....	92	23	4	3	3	7	5	4	6	6	9	22	-
2003 1/.....	96	32	1	1	2	3	7	13	5	3	5	24	-
2002.....	118	31	4	1	3	7	13	14	9	8	7	21	-
2001.....	103	29	1	3	2	4	6	4	15	10	7	21	1
2000.....	82	19	2	1	3	10	7	5	6	4	7	18	-
1999.....	98	33	2	1	1	7	6	3	7	10	12	16	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	462	72	9	6	9	10	17	27	29	39	73	168	3
2006 1/.....	450	69	5	2	16	27	23	29	32	36	66	145	-
2005 1/.....	386	66	12	6	12	26	32	18	24	29	49	112	-
2004 1/.....	392	71	7	4	11	18	23	25	29	29	52	121	2
2003 1/.....	358	66	6	5	14	30	18	27	15	22	59	93	3
2002.....	340	65	13	4	16	18	22	29	21	21	45	84	2
2001.....	495	75	12	11	27	38	63	58	43	40	54	73	1
2000.....	378	72	6	10	17	36	43	42	27	32	33	57	3
1999.....	275	72	6	6	16	19	26	24	21	21	23	39	2

Asian or Pacific Islander 3/, male

2007 1/.....	212	42	4	4	5	5	15	17	11	15	28	64	2
2006 1/.....	232	44	4	-	11	17	15	17	19	24	24	57	-
2005 1/.....	202	34	6	4	9	16	24	14	17	12	21	45	-
2004 1/.....	196	41	3	1	5	10	17	13	18	17	24	46	1
2003 1/.....	187	43	5	1	8	16	16	16	12	12	30	26	2
2002.....	182	38	8	2	11	15	15	22	13	11	19	26	2
2001.....	286	43	7	6	18	30	47	43	24	19	22	26	1
2000.....	216	41	1	3	9	25	38	26	16	20	13	22	2
1999.....	149	43	4	2	7	11	18	15	10	13	11	14	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Asian or Pacific Islander 3/, female

2007 1/.....	250	30	5	2	4	5	2	10	18	24	45	104	1
2006 1/.....	218	25	1	2	5	10	8	12	13	12	42	88	-
2005 1/.....	184	32	6	2	3	10	8	4	7	17	28	67	-
2004 1/.....	196	30	4	3	6	8	6	12	11	12	28	75	1
2003 1/.....	171	23	1	4	6	14	2	11	3	10	29	67	1
2002.....	158	27	5	2	5	3	7	7	8	10	26	58	-
2001.....	209	32	5	5	9	8	16	15	19	21	32	47	-
2000.....	162	31	5	7	8	11	5	16	11	12	20	35	1
1999.....	126	29	2	4	9	8	8	9	11	8	12	25	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

All races, both sexes

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
2007.....	238,192	1,337	599	876	1,957	3,149	7,068	16,499	22,895	28,838	61,153	93,813	8
2006.....	237,421	1,173	548	792	2,120	3,364	8,029	17,354	22,624	28,683	61,387	91,329	18
2005.....	217,632	1,198	612	857	2,060	3,159	8,118	17,122	21,108	27,236	56,700	79,449	13
2004.....	201,914	1,192	611	893	1,956	3,232	8,072	16,205	19,486	26,104	53,123	71,028	12
2003.....	201,676	1,190	635	927	2,023	3,330	8,407	15,883	18,734	26,475	52,955	71,100	17
2002.....	194,591	1,144	661	868	1,853	3,136	8,506	15,151	17,378	25,689	51,618	68,574	13
2001.....	186,406	1,131	646	860	1,799	3,072	8,084	14,203	16,048	25,803	49,711	65,032	17
2000.....	178,378	1,105	609	879	1,764	3,097	8,113	13,326	15,156	25,007	48,019	61,292	11
1999.....	171,683	1,148	647	921	1,771	3,062	7,989	12,383	14,390	25,165	45,598	58,599	10

All races, male

2007.....	96,447	779	326	518	1,154	1,787	4,038	9,787	12,722	14,363	25,221	25,747	5
2006.....	96,201	687	299	433	1,263	1,954	4,682	10,280	12,597	14,291	25,135	24,564	16
2005.....	88,869	732	352	474	1,245	1,821	4,837	10,253	11,828	13,311	23,071	20,934	11
2004.....	82,725	704	328	490	1,158	1,861	4,739	9,767	10,796	12,805	21,422	18,646	9
2003.....	81,907	711	342	541	1,189	1,885	4,932	9,456	10,274	12,993	21,393	18,178	13
2002.....	79,007	690	363	486	1,072	1,837	5,019	9,115	9,528	12,667	20,667	17,552	11
2001.....	75,162	649	358	454	1,064	1,729	4,774	8,397	8,652	12,732	19,817	16,520	16
2000.....	72,227	629	338	474	1,057	1,811	4,801	7,911	8,236	12,321	19,102	15,536	11
1999.....	69,784	680	354	483	1,045	1,790	4,835	7,272	7,870	12,332	18,495	14,623	5

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

All races, female

2007.....	141,745	558	273	358	803	1,362	3,030	6,712	10,173	14,475	35,932	68,066	3
2006.....	141,220	486	249	359	857	1,410	3,347	7,074	10,027	14,392	36,252	66,765	2
2005.....	128,763	466	260	383	815	1,338	3,281	6,869	9,280	13,925	33,629	58,515	2
2004.....	119,189	488	283	403	798	1,371	3,333	6,438	8,690	13,299	31,701	52,382	3
2003.....	119,769	479	293	386	834	1,445	3,475	6,427	8,460	13,482	31,562	52,922	4
2002.....	115,584	454	298	382	781	1,299	3,487	6,036	7,850	13,022	30,951	51,022	2
2001.....	111,244	482	288	406	735	1,343	3,310	5,806	7,396	13,071	29,894	48,512	1
2000.....	106,151	476	271	405	707	1,286	3,312	5,415	6,920	12,686	28,917	45,756	-
1999.....	101,899	468	293	438	726	1,272	3,154	5,111	6,520	12,833	27,103	43,976	5

White, both sexes

2007 1/.....	206,947	836	399	641	1,471	2,241	5,196	12,821	18,658	24,356	54,476	85,846	6
2006 1/.....	206,332	762	394	595	1,591	2,453	5,975	13,440	18,396	24,269	54,794	83,650	13
2005 1/.....	188,445	801	446	594	1,562	2,283	6,078	13,188	17,210	23,027	50,542	72,703	11
2004 1/.....	174,636	807	433	663	1,440	2,251	6,012	12,466	15,833	22,085	47,503	65,136	7
2003 1/.....	174,660	796	461	687	1,530	2,356	6,238	12,140	15,207	22,548	47,374	65,310	13
2002.....	168,631	788	470	654	1,348	2,233	6,315	11,610	14,110	21,873	46,212	63,010	8
2001.....	161,293	745	471	663	1,335	2,163	5,889	10,734	12,929	22,055	44,598	59,700	11
2000.....	154,181	740	422	652	1,325	2,197	5,916	10,098	12,182	21,360	43,077	56,204	8
1999.....	148,614	750	482	675	1,319	2,254	5,752	9,321	11,627	21,436	40,987	54,005	6

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

White, male

2007 1/.....	82,943	486	219	375	890	1,330	3,074	7,780	10,443	12,160	22,547	23,635	4
2006 1/.....	82,811	447	214	326	1,000	1,519	3,619	8,187	10,320	12,175	22,463	22,529	12
2005 1/.....	76,187	491	257	337	978	1,388	3,789	8,129	9,739	11,307	20,573	19,190	9
2004 1/.....	70,636	481	235	377	886	1,363	3,707	7,717	8,847	10,861	19,119	17,039	4
2003 1/.....	69,977	502	241	406	953	1,414	3,808	7,393	8,395	11,064	19,078	16,714	9
2002.....	67,452	493	257	366	820	1,368	3,913	7,143	7,784	10,793	18,478	16,029	8
2001.....	64,020	416	272	352	824	1,264	3,596	6,480	7,014	10,919	17,785	15,088	10
2000.....	61,390	409	238	369	822	1,354	3,618	6,158	6,641	10,533	17,107	14,133	8
1999.....	59,336	446	264	344	809	1,384	3,648	5,603	6,413	10,563	16,566	13,293	3

White, female

2007 1/.....	124,004	350	180	266	581	911	2,122	5,041	8,215	12,196	31,929	62,211	2
2006 1/.....	123,521	315	180	269	591	934	2,356	5,253	8,076	12,094	32,331	61,121	1
2005 1/.....	112,258	310	189	257	584	895	2,289	5,059	7,471	11,720	29,969	53,513	2
2004 1/.....	104,000	326	198	286	554	888	2,305	4,749	6,986	11,224	28,384	48,097	3
2003 1/.....	104,683	294	220	281	577	942	2,430	4,747	6,812	11,484	28,296	48,596	4
2002.....	101,179	295	213	288	528	865	2,402	4,467	6,326	11,080	27,734	46,981	-
2001.....	97,273	329	199	311	511	899	2,293	4,254	5,915	11,136	26,813	44,612	1
2000.....	92,791	331	184	283	503	843	2,298	3,940	5,541	10,827	25,970	42,071	-
1999.....	89,278	304	218	331	510	870	2,104	3,718	5,214	10,873	24,421	40,712	3

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Black, both sexes

2007 1/.....	26,371	421	155	191	406	774	1,631	3,248	3,703	3,768	5,546	6,526	2
2006 1/.....	26,205	352	118	166	439	762	1,712	3,468	3,684	3,709	5,482	6,309	4
2005 1/.....	24,843	335	130	221	423	747	1,758	3,452	3,399	3,555	5,204	5,617	2
2004 1/.....	23,519	326	138	185	434	844	1,773	3,346	3,212	3,473	4,799	4,985	4
2003 1/.....	23,223	342	137	186	410	827	1,900	3,340	3,077	3,365	4,714	4,921	4
2002.....	22,519	305	150	176	420	789	1,954	3,183	2,855	3,277	4,606	4,800	4
2001.....	21,970	334	133	167	392	763	1,965	3,147	2,719	3,276	4,450	4,618	6
2000.....	21,272	305	148	189	383	816	1,969	2,930	2,625	3,176	4,288	4,440	3
1999.....	20,325	343	131	202	388	708	2,018	2,751	2,465	3,285	3,996	4,034	4

Black, male

2007 1/.....	11,280	245	80	118	215	383	818	1,754	1,998	1,859	2,205	1,604	1
2006 1/.....	11,201	205	66	95	226	349	860	1,843	1,982	1,793	2,199	1,580	3
2005 1/.....	10,603	212	77	120	224	362	876	1,842	1,806	1,686	2,061	1,335	2
2004 1/.....	10,328	188	75	87	223	424	869	1,815	1,712	1,691	1,940	1,300	4
2003 1/.....	10,126	186	80	100	188	393	958	1,824	1,629	1,660	1,930	1,174	4
2002.....	9,847	168	82	98	208	398	962	1,752	1,519	1,607	1,820	1,231	2
2001.....	9,671	195	64	83	200	384	1,042	1,741	1,430	1,587	1,781	1,158	6
2000.....	9,455	187	78	87	204	404	1,047	1,586	1,420	1,561	1,729	1,149	3
1999.....	9,071	199	68	119	205	348	1,052	1,504	1,291	1,538	1,643	1,102	2

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Black, female

2007 1/.....	15,091	176	75	73	191	391	813	1,494	1,705	1,909	3,341	4,922	1
2006 1/.....	15,004	147	52	71	213	413	852	1,625	1,702	1,916	3,283	4,729	1
2005 1/.....	14,240	123	53	101	199	385	882	1,610	1,593	1,869	3,143	4,282	-
2004 1/.....	13,191	138	63	98	211	420	904	1,531	1,500	1,782	2,859	3,685	-
2003 1/.....	13,097	156	57	86	222	434	942	1,516	1,448	1,705	2,784	3,747	-
2002.....	12,672	137	68	78	212	391	992	1,431	1,336	1,670	2,786	3,569	2
2001.....	12,299	139	69	84	192	379	923	1,406	1,289	1,689	2,669	3,460	-
2000.....	11,817	118	70	102	179	412	922	1,344	1,205	1,615	2,559	3,291	-
1999.....	11,254	144	63	83	183	360	966	1,247	1,174	1,747	2,353	2,932	2

American Indian 2/, both sexes

2007 1/.....	1,412	14	12	15	27	64	142	210	223	222	245	238	-
2006 1/.....	1,483	15	11	5	27	67	188	226	243	221	240	240	-
2005 1/.....	1,363	19	14	12	33	50	158	260	206	206	206	199	-
2004 1/.....	1,226	12	15	14	30	73	168	222	187	169	182	153	1
2003 1/.....	1,243	10	12	16	36	59	159	221	189	165	196	180	-
2002.....	1,102	11	11	11	34	52	126	192	175	179	158	153	-
2001.....	1,109	15	11	8	24	71	138	176	174	155	168	169	-
2000.....	945	17	17	7	16	46	132	151	134	148	152	125	-
1999.....	951	16	12	12	25	50	140	164	133	134	142	123	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

American Indian 2/, male

2007 1/.....	669	7	6	10	16	39	84	117	110	96	106	78	-
2006 1/.....	695	9	8	2	11	39	108	119	132	101	91	75	-
2005 1/.....	668	8	8	2	15	30	95	147	124	95	89	55	-
2004 1/.....	601	10	7	6	16	35	90	127	107	84	79	39	1
2003 1/.....	606	3	8	11	24	32	91	141	99	83	75	39	-
2002.....	562	4	8	6	20	34	81	114	94	86	59	56	-
2001.....	547	11	5	4	18	42	85	103	93	84	51	51	-
2000.....	471	8	7	5	11	37	84	82	61	71	60	45	-
1999.....	469	11	7	5	12	32	83	97	73	57	58	34	-

American Indian 2/, female

2007 1/.....	743	7	6	5	11	25	58	93	113	126	139	160	-
2006 1/.....	788	6	3	3	16	28	80	107	111	120	149	165	-
2005 1/.....	695	11	6	10	18	20	63	113	82	111	117	144	-
2004 1/.....	625	2	8	8	14	38	78	95	80	85	103	114	-
2003 1/.....	637	7	4	5	12	27	68	80	90	82	121	141	-
2002.....	540	7	3	5	14	18	45	78	81	93	99	97	-
2001.....	562	4	6	4	6	29	53	73	81	71	117	118	-
2000.....	474	9	10	2	5	9	48	69	73	77	92	80	-
1999.....	482	5	5	7	13	18	57	67	60	77	84	89	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	3,462	66	33	29	53	70	99	220	311	492	886	1,203	-
2006 1/.....	3,401	44	25	26	63	82	154	220	301	484	871	1,130	1
2005 1/.....	2,981	43	22	30	42	79	124	222	293	448	748	930	-
2004 1/.....	2,533	47	25	31	52	64	119	171	254	377	639	754	-
2003 1/.....	2,550	42	25	38	47	88	110	182	261	397	671	689	-
2002.....	2,339	40	30	27	51	62	111	166	238	360	642	611	1
2001.....	2,034	37	31	22	48	75	92	146	226	317	495	545	-
2000.....	1,980	43	22	31	40	38	96	147	215	323	502	523	-
1999.....	1,793	39	22	32	39	50	79	147	165	310	473	437	-

Asian or Pacific Islander 3/, male

2007 1/.....	1,555	41	21	15	33	35	62	136	171	248	363	430	-
2006 1/.....	1,494	26	11	10	26	47	95	131	163	222	382	380	1
2005 1/.....	1,411	21	10	15	28	41	77	135	159	223	348	354	-
2004 1/.....	1,160	25	11	20	33	39	73	108	130	169	284	268	-
2003 1/.....	1,198	20	13	24	24	46	75	98	151	186	310	251	-
2002.....	1,146	25	16	16	24	37	63	106	131	181	310	236	1
2001.....	924	27	17	15	22	39	51	73	115	142	200	223	-
2000.....	911	25	15	13	20	16	52	85	114	156	206	209	-
1999.....	908	24	15	15	19	26	52	68	93	174	228	194	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Asian or Pacific Islander 3/, female

2007 1/.....	1,907	25	12	14	20	35	37	84	140	244	523	773	-
2006 1/.....	1,907	18	14	16	37	35	59	89	138	262	489	750	-
2005 1/.....	1,570	22	12	15	14	38	47	87	134	225	400	576	-
2004 1/.....	1,373	22	14	11	19	25	46	63	124	208	355	486	-
2003 1/.....	1,352	22	12	14	23	42	35	84	110	211	361	438	-
2002.....	1,193	15	14	11	27	25	48	60	107	179	332	375	-
2001.....	1,110	10	14	7	26	36	41	73	111	175	295	322	-
2000.....	1,069	18	7	18	20	22	44	62	101	167	296	314	-
1999.....	885	15	7	17	20	24	27	79	72	136	245	243	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

All races, both sexes

2007.....	123,706	1,285	1,588	2,194	15,897	14,977	16,931	20,315	12,193	8,753	13,736	15,803	34
2006.....	121,599	1,147	1,610	2,258	16,229	14,954	17,534	19,675	11,446	8,420	13,708	14,561	57
2005.....	117,809	1,083	1,664	2,415	15,753	13,997	16,919	18,339	10,853	8,632	13,854	14,243	57
2004.....	112,012	1,052	1,641	2,666	15,449	13,032	16,471	16,942	9,651	8,116	13,457	13,447	88
2003.....	109,277	945	1,717	2,618	15,272	12,541	16,766	15,837	9,170	8,081	13,108	13,146	76
2002.....	106,742	946	1,641	2,718	15,412	12,569	16,710	14,675	8,345	8,086	12,904	12,651	85
2001.....	101,537	976	1,714	2,836	14,411	11,839	15,945	13,344	7,658	7,835	12,688	12,171	120
2000.....	97,900	881	1,826	2,979	14,113	11,769	15,413	12,278	7,505	7,698	11,758	11,595	85
1999.....	97,860	845	1,898	3,091	13,656	11,890	15,231	11,639	7,285	8,208	12,282	11,729	106

All races, male

2007.....	79,827	715	987	1,346	11,905	11,384	12,022	14,017	8,357	5,593	7,268	6,201	32
2006.....	78,941	667	927	1,385	12,234	11,466	12,431	13,837	7,848	5,212	7,194	5,692	48
2005.....	76,375	578	1,027	1,493	11,827	10,665	12,026	13,002	7,507	5,343	7,337	5,521	49
2004.....	72,050	604	964	1,661	11,423	9,871	11,671	11,988	6,562	4,992	7,030	5,210	74
2003.....	70,532	539	1,048	1,662	11,291	9,534	12,003	11,196	6,324	4,921	6,851	5,093	70
2002.....	69,257	553	1,024	1,713	11,438	9,635	12,012	10,492	5,781	5,031	6,687	4,817	74
2001.....	66,060	541	1,045	1,803	10,832	9,092	11,612	9,743	5,228	4,751	6,652	4,657	104
2000.....	63,817	501	1,098	1,861	10,460	9,059	11,444	9,063	5,133	4,639	6,063	4,413	83
1999.....	63,535	494	1,127	1,974	10,040	9,097	11,248	8,602	5,015	4,992	6,372	4,480	94

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

All races, female

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
2007.....	43,879	570	601	848	3,992	3,593	4,909	6,298	3,836	3,160	6,468	9,602	2
2006.....	42,658	480	683	873	3,995	3,488	5,103	5,838	3,598	3,208	6,514	8,869	9
2005.....	41,434	505	637	922	3,926	3,332	4,893	5,337	3,346	3,289	6,517	8,722	8
2004.....	39,962	448	677	1,005	4,026	3,161	4,800	4,954	3,089	3,124	6,427	8,237	14
2003.....	38,745	406	669	956	3,981	3,007	4,763	4,641	2,846	3,160	6,257	8,053	6
2002.....	37,485	393	617	1,005	3,974	2,934	4,698	4,183	2,564	3,055	6,217	7,834	11
2001.....	35,477	435	669	1,033	3,579	2,747	4,333	3,601	2,430	3,084	6,036	7,514	16
2000.....	34,083	380	728	1,118	3,653	2,710	3,969	3,215	2,372	3,059	5,695	7,182	2
1999.....	34,325	351	771	1,117	3,616	2,793	3,983	3,037	2,270	3,216	5,910	7,249	12

White, both sexes

2007 1/.....	106,252	803	1,169	1,625	13,481	12,646	14,211	17,018	10,224	7,529	12,584	14,933	29
2006 1/.....	103,853	730	1,170	1,668	13,824	12,473	14,647	16,245	9,514	7,266	12,529	13,737	50
2005 1/.....	100,406	734	1,262	1,785	13,325	11,628	14,172	15,135	9,032	7,424	12,524	13,333	52
2004 1/.....	95,890	706	1,242	1,953	13,174	10,787	13,841	14,030	8,078	7,006	12,294	12,700	79
2003 1/.....	93,381	624	1,304	1,954	12,990	10,400	14,016	13,020	7,701	6,952	11,960	12,390	70
2002.....	90,866	583	1,199	1,965	13,048	10,378	13,901	12,069	6,985	6,967	11,816	11,877	78
2001.....	85,964	647	1,309	2,060	12,038	9,727	13,143	10,825	6,367	6,778	11,588	11,379	103
2000.....	82,592	569	1,335	2,219	11,823	9,519	12,663	9,901	6,275	6,617	10,740	10,852	79
1999.....	82,245	562	1,374	2,270	11,364	9,661	12,500	9,368	5,962	7,027	11,130	10,938	89

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

White, male

2007 1/.....	68,059	455	738	1,000	10,121	9,624	10,074	11,725	6,984	4,798	6,663	5,848	29
2006 1/.....	66,843	421	675	1,017	10,423	9,564	10,393	11,380	6,483	4,502	6,584	5,358	43
2005 1/.....	64,600	388	793	1,101	10,035	8,876	10,096	10,714	6,185	4,586	6,627	5,153	46
2004 1/.....	61,267	414	734	1,223	9,770	8,180	9,865	9,933	5,438	4,327	6,423	4,892	68
2003 1/.....	59,912	359	807	1,233	9,596	7,964	10,088	9,220	5,308	4,259	6,235	4,779	64
2002.....	58,467	358	736	1,240	9,679	7,986	10,040	8,664	4,807	4,316	6,090	4,484	67
2001.....	55,493	356	815	1,285	9,067	7,518	9,647	7,897	4,319	4,119	6,050	4,329	91
2000.....	53,329	331	808	1,378	8,770	7,369	9,462	7,322	4,242	3,977	5,493	4,100	77
1999.....	52,898	333	811	1,434	8,327	7,442	9,288	6,919	4,100	4,267	5,758	4,140	79

White, female

2007 1/.....	38,193	348	431	625	3,360	3,022	4,137	5,293	3,240	2,731	5,921	9,085	-
2006 1/.....	37,010	309	495	651	3,401	2,909	4,254	4,865	3,031	2,764	5,945	8,379	7
2005 1/.....	35,806	346	469	684	3,290	2,752	4,076	4,421	2,847	2,838	5,897	8,180	6
2004 1/.....	34,623	292	508	730	3,404	2,607	3,976	4,097	2,640	2,679	5,871	7,808	11
2003 1/.....	33,469	265	497	721	3,394	2,436	3,928	3,800	2,393	2,693	5,725	7,611	6
2002.....	32,399	225	463	725	3,369	2,392	3,861	3,405	2,178	2,651	5,726	7,393	11
2001.....	30,471	291	494	775	2,971	2,209	3,496	2,928	2,048	2,659	5,538	7,050	12
2000.....	29,263	238	527	841	3,053	2,150	3,201	2,579	2,033	2,640	5,247	6,752	2
1999.....	29,347	229	563	836	3,037	2,219	3,212	2,449	1,862	2,760	5,372	6,798	10

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Black, both sexes

2007 1/.....	13,559	428	325	459	1,749	1,779	2,158	2,766	1,599	911	764	617	4
2006 1/.....	13,917	358	371	471	1,750	1,933	2,347	2,856	1,557	865	824	580	5
2005 1/.....	13,652	314	327	501	1,780	1,796	2,208	2,689	1,471	927	980	654	5
2004 1/.....	12,670	303	324	570	1,661	1,740	2,104	2,455	1,268	835	866	540	4
2003 1/.....	12,351	287	343	523	1,618	1,635	2,224	2,324	1,169	862	812	549	5
2002.....	12,513	330	359	621	1,726	1,711	2,268	2,179	1,093	832	806	584	4
2001.....	12,462	281	339	644	1,742	1,638	2,321	2,133	1,064	852	836	598	14
2000.....	12,277	274	405	631	1,695	1,770	2,287	1,988	997	868	778	579	5
1999.....	12,728	252	441	707	1,719	1,765	2,285	1,939	1,066	951	936	651	16

Black, male

2007 1/.....	9,268	222	189	279	1,309	1,369	1,563	1,949	1,151	604	398	233	2
2006 1/.....	9,605	213	214	295	1,331	1,503	1,669	2,055	1,122	548	428	223	4
2005 1/.....	9,329	172	184	314	1,334	1,369	1,557	1,936	1,094	594	530	242	3
2004 1/.....	8,603	163	188	351	1,228	1,333	1,457	1,747	912	532	474	215	3
2003 1/.....	8,385	160	204	334	1,227	1,221	1,565	1,652	836	528	437	216	5
2002.....	8,612	176	239	400	1,298	1,309	1,592	1,552	808	539	452	243	4
2001.....	8,537	162	194	442	1,309	1,220	1,633	1,597	764	517	462	225	12
2000.....	8,531	149	239	414	1,261	1,340	1,678	1,475	750	543	438	239	5
1999.....	8,773	141	266	472	1,301	1,335	1,660	1,471	755	600	488	270	14

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Black, female

2007 1/.....	4,291	206	136	180	440	410	595	817	448	307	366	384	2
2006 1/.....	4,312	145	157	176	419	430	678	801	435	317	396	357	1
2005 1/.....	4,323	142	143	187	446	427	651	753	377	333	450	412	2
2004 1/.....	4,067	140	136	219	433	407	647	708	356	303	392	325	1
2003 1/.....	3,966	127	139	189	391	414	659	672	333	334	375	333	-
2002.....	3,901	154	120	221	428	402	676	627	285	293	354	341	-
2001.....	3,925	119	145	202	433	418	688	536	300	335	374	373	2
2000.....	3,746	125	166	217	434	430	609	513	247	325	340	340	-
1999.....	3,955	111	175	235	418	430	625	468	311	351	448	381	2

American Indian 2/, both sexes

2007 1/.....	1,701	31	43	46	319	309	320	280	150	93	77	33	-
2006 1/.....	1,704	36	32	43	328	305	300	313	157	85	68	36	1
2005 1/.....	1,626	19	44	55	321	324	305	258	135	75	60	30	-
2004 1/.....	1,520	23	42	61	295	262	290	234	117	89	65	37	5
2003 1/.....	1,573	20	33	68	315	264	313	241	128	84	62	44	1
2002.....	1,488	15	33	60	300	254	333	215	112	79	57	29	1
2001.....	1,361	32	30	73	293	216	260	193	98	61	63	42	-
2000.....	1,353	20	39	70	299	239	256	197	94	63	51	24	1
1999.....	1,327	13	37	51	289	250	252	163	105	82	59	26	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

American Indian 2/, male

2007 1/.....	1,129	23	24	30	224	214	216	177	95	62	50	14	-
2006 1/.....	1,184	21	15	26	237	225	210	229	114	55	39	13	-
2005 1/.....	1,118	10	31	36	223	240	213	181	99	45	28	12	-
2004 1/.....	1,010	15	26	33	203	184	191	169	94	46	27	19	3
2003 1/.....	1,043	14	19	48	228	177	215	161	82	51	29	18	1
2002.....	1,003	8	19	31	212	186	228	149	72	56	30	11	1
2001.....	908	14	15	46	203	163	188	130	68	33	34	14	-
2000.....	900	11	25	37	202	172	166	150	56	36	32	12	1
1999.....	890	8	24	28	214	163	167	121	75	51	29	10	-

American Indian 2/, female

2007 1/.....	572	8	19	16	95	95	104	103	55	31	27	19	-
2006 1/.....	520	15	17	17	91	80	90	84	43	30	29	23	1
2005 1/.....	508	9	13	19	98	84	92	77	36	30	32	18	-
2004 1/.....	510	8	16	28	92	78	99	65	23	43	38	18	2
2003 1/.....	530	6	14	20	87	87	98	80	46	33	33	26	-
2002.....	485	7	14	29	88	68	105	66	40	23	27	18	-
2001.....	453	18	15	27	90	53	72	63	30	28	29	28	-
2000.....	453	9	14	33	97	67	90	47	38	27	19	12	-
1999.....	437	5	13	23	75	87	85	42	30	31	30	16	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	2,194	23	51	64	348	243	242	251	220	220	311	220	1
2006 1/.....	2,125	23	37	76	327	243	240	261	218	204	287	208	1
2005 1/.....	2,125	16	31	74	327	249	234	257	215	206	290	226	-
2004 1/.....	1,932	20	33	82	319	243	236	223	188	186	232	170	-
2003 1/.....	1,972	14	37	73	349	242	213	252	172	183	274	163	-
2002.....	1,875	18	50	72	338	226	208	212	155	208	225	161	2
2001.....	1,750	16	36	59	338	258	221	193	129	144	201	152	3
2000.....	1,678	18	47	59	296	241	207	192	139	150	189	140	-
1999.....	1,560	18	46	63	284	214	194	169	152	148	157	114	1

Asian or Pacific Islander 3/, male

2007 1/.....	1,371	15	36	37	251	177	169	166	127	129	157	106	1
2006 1/.....	1,309	12	23	47	243	174	159	173	129	107	143	98	1
2005 1/.....	1,328	8	19	42	235	180	160	171	129	118	152	114	-
2004 1/.....	1,170	12	16	54	222	174	158	139	118	87	106	84	-
2003 1/.....	1,192	6	18	47	240	172	135	163	98	83	150	80	-
2002.....	1,175	11	30	42	249	154	152	127	94	120	115	79	2
2001.....	1,122	9	21	30	253	191	144	119	77	82	106	89	1
2000.....	1,057	10	26	32	227	178	138	116	85	83	100	62	-
1999.....	974	12	26	40	198	157	133	91	85	74	97	60	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Asian or Pacific Islander 3/, female

2007 1/.....	823	8	15	27	97	66	73	85	93	91	154	114	-
2006 1/.....	816	11	14	29	84	69	81	88	89	97	144	110	-
2005 1/.....	797	8	12	32	92	69	74	86	86	88	138	112	-
2004 1/.....	762	8	17	28	97	69	78	84	70	99	126	86	-
2003 1/.....	780	8	19	26	109	70	78	89	74	100	124	83	-
2002.....	700	7	20	30	89	72	56	85	61	88	110	82	-
2001.....	628	7	15	29	85	67	77	74	52	62	95	63	2
2000.....	621	8	21	27	69	63	69	76	54	67	89	78	-
1999.....	586	6	20	23	86	57	61	78	67	74	60	54	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Transport accidents (V01-V99,Y85)													
All races, both sexes													
2007.....	46,844	127	581	1,374	10,928	7,452	6,829	7,199	4,838	3,194	2,983	1,326	13
2006.....	48,412	142	615	1,453	11,392	7,753	7,249	7,264	4,986	3,170	3,082	1,290	16
2005.....	48,441	147	646	1,547	11,288	7,638	7,295	7,101	4,918	3,353	3,168	1,329	11
2004.....	48,053	145	659	1,755	11,377	7,521	7,247	6,953	4,510	3,229	3,310	1,320	27
2003.....	48,071	148	649	1,765	11,432	7,280	7,594	6,754	4,441	3,210	3,377	1,401	20
2002.....	48,366	127	640	1,738	11,816	7,484	7,688	6,563	4,165	3,351	3,433	1,322	39
2001.....	47,288	151	678	1,823	11,139	7,454	7,827	6,251	3,861	3,282	3,507	1,272	43
2000.....	46,749	174	691	1,922	11,028	7,400	7,661	5,965	3,883	3,315	3,342	1,335	33
1999.....	46,423	189	712	1,975	10,689	7,434	7,553	5,645	3,812	3,579	3,468	1,330	37
All races, male													
2007.....	33,434	58	328	813	7,943	5,710	5,084	5,348	3,460	2,138	1,798	741	13
2006.....	34,065	70	329	849	8,281	5,951	5,233	5,347	3,483	2,026	1,758	724	14
2005.....	34,086	74	369	910	8,208	5,840	5,303	5,208	3,435	2,157	1,835	737	10
2004.....	33,311	81	359	1,039	8,113	5,641	5,297	5,042	3,114	1,998	1,905	701	21
2003.....	33,291	81	366	1,084	8,171	5,511	5,497	4,865	3,087	1,975	1,885	751	18
2002.....	33,533	72	381	1,039	8,471	5,676	5,499	4,727	2,877	2,101	1,917	737	36
2001.....	32,794	68	392	1,115	8,111	5,661	5,613	4,484	2,631	2,015	1,983	684	37
2000.....	32,166	94	391	1,138	7,890	5,570	5,560	4,260	2,617	2,021	1,858	734	33
1999.....	31,642	101	400	1,206	7,558	5,511	5,376	4,016	2,561	2,162	1,959	758	34

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Transport accidents (V01-V99,Y85)													
All races, female													
2007.....	13,410	69	253	561	2,985	1,742	1,745	1,851	1,378	1,056	1,185	585	-
2006.....	14,347	72	286	604	3,111	1,802	2,016	1,917	1,503	1,144	1,324	566	2
2005.....	14,355	73	277	637	3,080	1,798	1,992	1,893	1,483	1,196	1,333	592	1
2004.....	14,742	64	300	716	3,264	1,880	1,950	1,911	1,396	1,231	1,405	619	6
2003.....	14,780	67	283	681	3,261	1,769	2,097	1,889	1,354	1,235	1,492	650	2
2002.....	14,833	55	259	699	3,345	1,808	2,189	1,836	1,288	1,250	1,516	585	3
2001.....	14,494	83	286	708	3,028	1,793	2,214	1,767	1,230	1,267	1,524	588	6
2000.....	14,583	80	300	784	3,138	1,830	2,101	1,705	1,266	1,294	1,484	601	-
1999.....	14,781	88	312	769	3,131	1,923	2,177	1,629	1,251	1,417	1,509	572	3
White, both sexes													
2007 1/.....	39,086	93	426	1,079	9,115	6,002	5,553	6,029	4,121	2,759	2,683	1,214	12
2006 1/.....	40,422	100	445	1,121	9,598	6,193	5,975	6,055	4,193	2,762	2,764	1,203	13
2005 1/.....	40,647	100	492	1,210	9,478	6,168	6,022	5,942	4,214	2,911	2,859	1,241	10
2004 1/.....	40,279	104	499	1,377	9,648	6,022	5,965	5,740	3,854	2,805	2,995	1,245	25
2003 1/.....	40,282	111	491	1,397	9,643	5,885	6,253	5,607	3,753	2,777	3,048	1,298	19
2002.....	40,614	84	478	1,347	9,991	6,037	6,322	5,476	3,558	2,921	3,137	1,229	34
2001.....	39,544	115	511	1,378	9,285	6,053	6,451	5,175	3,277	2,857	3,217	1,190	35
2000.....	39,088	121	513	1,519	9,279	5,890	6,318	4,924	3,319	2,878	3,060	1,236	31
1999.....	38,767	137	497	1,542	8,953	6,026	6,199	4,665	3,214	3,134	3,154	1,218	28

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Transport accidents (V01-V99,Y85)													
White, male													
2007 1/.....	27,864	46	248	639	6,626	4,592	4,125	4,491	2,954	1,854	1,607	670	12
2006 1/.....	28,429	52	233	645	6,979	4,758	4,319	4,472	2,946	1,764	1,580	669	12
2005 1/.....	28,568	54	278	711	6,898	4,707	4,372	4,380	2,950	1,868	1,654	687	9
2004 1/.....	27,918	54	269	829	6,901	4,510	4,388	4,179	2,649	1,737	1,720	662	20
2003 1/.....	27,925	63	285	851	6,867	4,489	4,540	4,057	2,633	1,733	1,700	690	17
2002.....	28,118	50	278	814	7,161	4,590	4,540	3,973	2,458	1,818	1,729	676	31
2001.....	27,407	52	299	835	6,774	4,620	4,632	3,727	2,230	1,757	1,814	636	31
2000.....	26,815	68	290	896	6,632	4,441	4,589	3,543	2,213	1,744	1,685	683	31
1999.....	26,385	80	273	936	6,316	4,477	4,428	3,331	2,161	1,896	1,773	688	26
White, female													
2007 1/.....	11,222	47	178	440	2,489	1,410	1,428	1,538	1,167	905	1,076	544	-
2006 1/.....	11,993	48	212	476	2,619	1,435	1,656	1,583	1,247	998	1,184	534	1
2005 1/.....	12,079	46	214	499	2,580	1,461	1,650	1,562	1,264	1,043	1,205	554	1
2004 1/.....	12,361	50	230	548	2,747	1,512	1,577	1,561	1,205	1,068	1,275	583	5
2003 1/.....	12,357	48	206	546	2,776	1,396	1,713	1,550	1,120	1,044	1,348	608	2
2002.....	12,496	34	200	533	2,830	1,447	1,782	1,503	1,100	1,103	1,408	553	3
2001.....	12,137	63	212	543	2,511	1,433	1,819	1,448	1,047	1,100	1,403	554	4
2000.....	12,273	53	223	623	2,647	1,449	1,729	1,381	1,106	1,134	1,375	553	-
1999.....	12,382	57	224	606	2,637	1,549	1,771	1,334	1,053	1,238	1,381	530	2

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Transport accidents (V01-V99,Y85)													
Black, both sexes													
2007 1/.....	5,859	28	120	238	1,316	1,112	1,004	934	547	297	191	71	1
2006 1/.....	5,983	31	137	257	1,282	1,209	992	929	601	279	204	60	2
2005 1/.....	5,842	38	119	261	1,328	1,099	990	904	523	316	201	62	1
2004 1/.....	5,868	32	132	295	1,260	1,154	982	959	502	285	215	52	-
2003 1/.....	5,771	28	129	278	1,242	1,056	1,054	870	510	311	220	72	1
2002.....	5,796	36	126	312	1,314	1,112	1,054	828	451	289	209	62	3
2001.....	5,906	19	130	363	1,347	1,080	1,087	842	460	313	200	57	8
2000.....	5,839	40	141	317	1,283	1,169	1,069	818	424	325	183	69	1
1999.....	5,901	44	177	364	1,284	1,089	1,085	772	436	318	236	87	9
Black, male													
2007 1/.....	4,336	11	61	141	975	881	781	706	401	204	128	46	1
2006 1/.....	4,333	15	77	158	946	946	735	685	422	190	116	41	2
2005 1/.....	4,237	16	65	155	975	870	745	654	377	217	126	36	1
2004 1/.....	4,204	20	71	165	907	896	722	705	358	193	138	29	-
2003 1/.....	4,119	12	66	175	933	799	768	643	359	189	133	41	1
2002.....	4,156	18	83	183	949	856	749	598	333	200	141	43	3
2001.....	4,214	7	72	240	986	813	788	624	335	189	123	31	6
2000.....	4,199	21	79	197	928	890	804	579	321	220	120	39	1
1999.....	4,162	18	101	232	926	828	778	567	307	198	141	58	8

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Transport accidents (V01-V99,Y85)													
Black, female													
2007 1/.....	1,523	17	59	97	341	231	223	228	146	93	63	25	-
2006 1/.....	1,650	16	60	99	336	263	257	244	179	89	88	19	-
2005 1/.....	1,605	22	54	106	353	229	245	250	146	99	75	26	-
2004 1/.....	1,664	12	61	130	353	258	260	254	144	92	77	23	-
2003 1/.....	1,652	16	63	103	309	257	286	227	151	122	87	31	-
2002.....	1,640	18	43	129	365	256	305	230	118	89	68	19	-
2001.....	1,692	12	58	123	361	267	299	218	125	124	77	26	2
2000.....	1,640	19	62	120	355	279	265	239	103	105	63	30	-
1999.....	1,739	26	76	132	358	261	307	205	129	120	95	29	1
American Indian 2/, both sexes													
2007 1/.....	850	4	19	24	231	196	152	108	60	35	19	2	-
2006 1/.....	930	8	16	28	261	191	152	138	77	30	26	2	1
2005 1/.....	875	5	27	38	235	201	150	111	59	31	15	3	-
2004 1/.....	851	5	20	35	226	174	144	118	51	48	26	2	2
2003 1/.....	915	7	18	46	263	178	154	123	65	36	16	9	-
2002.....	903	4	16	39	250	171	183	119	63	35	18	5	-
2001.....	832	11	17	48	247	147	146	105	50	31	25	5	-
2000.....	829	7	16	46	241	167	146	96	50	34	19	6	1
1999.....	816	4	15	35	233	163	148	91	59	44	23	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Transport accidents (V01-V99,Y85)

American Indian 2/, male

2007 1/.....	576	1	9	14	155	138	98	74	47	25	13	2	-
2006 1/.....	638	2	8	18	178	134	102	103	56	20	16	1	-
2005 1/.....	622	3	21	25	163	143	107	84	46	20	8	2	-
2004 1/.....	569	4	12	16	148	124	93	85	46	26	13	1	1
2003 1/.....	595	5	8	31	185	121	107	75	37	17	5	4	-
2002.....	603	1	9	20	169	122	123	82	37	23	14	3	-
2001.....	558	5	8	26	164	111	112	65	32	18	15	2	-
2000.....	537	2	10	23	162	117	92	70	25	19	13	3	1
1999.....	547	2	12	21	168	100	96	66	42	26	14	-	-

American Indian 2/, female

2007 1/.....	274	3	10	10	76	58	54	34	13	10	6	-	-
2006 1/.....	292	6	8	10	83	57	50	35	21	10	10	1	1
2005 1/.....	253	2	6	13	72	58	43	27	13	11	7	1	-
2004 1/.....	282	1	8	19	78	50	51	33	5	22	13	1	1
2003 1/.....	320	2	10	15	78	57	47	48	28	19	11	5	-
2002.....	300	3	7	19	81	49	60	37	26	12	4	2	-
2001.....	274	6	9	22	83	36	34	40	18	13	10	3	-
2000.....	292	5	6	23	79	50	54	26	25	15	6	3	-
1999.....	269	2	3	14	65	63	52	25	17	18	9	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Transport accidents (V01-V99,Y85)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	1,049	2	16	33	266	142	120	128	110	103	90	39	-
2006 1/.....	1,077	3	17	47	251	160	130	142	115	99	88	25	-
2005 1/.....	1,077	4	8	38	247	170	133	144	122	95	93	23	-
2004 1/.....	1,055	4	8	48	243	171	156	136	103	91	74	21	-
2003 1/.....	1,103	2	11	44	284	161	133	154	113	86	93	22	-
2002.....	1,053	3	20	40	261	164	129	140	93	106	69	26	2
2001.....	1,006	6	20	34	260	174	143	129	74	81	65	20	-
2000.....	993	6	21	40	225	174	128	127	90	78	80	24	-
1999.....	939	4	23	34	219	156	121	117	103	83	55	24	-

Asian or Pacific Islander 3/, male

2007 1/.....	658	-	10	19	187	99	80	77	58	55	50	23	-
2006 1/.....	665	1	11	28	178	113	77	87	59	52	46	13	-
2005 1/.....	659	1	5	19	172	120	79	90	62	52	47	12	-
2004 1/.....	620	3	7	29	157	111	94	73	61	42	34	9	-
2003 1/.....	652	1	7	27	186	102	82	90	58	36	47	16	-
2002.....	656	3	11	22	192	108	87	74	49	60	33	15	2
2001.....	615	4	13	14	187	117	81	68	34	51	31	15	-
2000.....	615	3	12	22	168	122	75	68	58	38	40	9	-
1999.....	548	1	14	17	148	106	74	52	51	42	31	12	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Transport accidents (V01-V99,Y85)

Asian or Pacific Islander 3/, female

2007 1/.....	391	2	6	14	79	43	40	51	52	48	40	16	-
2006 1/.....	412	2	6	19	73	47	53	55	56	47	42	12	-
2005 1/.....	418	3	3	19	75	50	54	54	60	43	46	11	-
2004 1/.....	435	1	1	19	86	60	62	63	42	49	40	12	-
2003 1/.....	451	1	4	17	98	59	51	64	55	50	46	6	-
2002.....	397	-	9	18	69	56	42	66	44	46	36	11	-
2001.....	391	2	7	20	73	57	62	61	40	30	34	5	-
2000.....	378	3	9	18	57	52	53	59	32	40	40	15	-
1999.....	391	3	9	17	71	50	47	65	52	41	24	12	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

All races, both sexes

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
2007.....	43,945	124	551	1,285	10,568	7,087	6,370	6,530	4,359	2,940	2,845	1,277	9
2006.....	45,316	140	588	1,339	11,015	7,348	6,677	6,615	4,518	2,916	2,906	1,240	14
2005.....	45,343	146	617	1,447	10,908	7,239	6,748	6,397	4,454	3,110	2,991	1,280	6
2004.....	44,933	143	635	1,653	10,987	7,036	6,663	6,276	4,093	2,974	3,180	1,267	26
2003.....	44,757	145	621	1,642	10,972	6,833	6,961	6,074	3,958	2,967	3,210	1,358	16
2002.....	45,380	123	610	1,614	11,459	7,092	7,077	5,937	3,764	3,113	3,271	1,285	35
2001.....	43,788	144	626	1,686	10,725	6,937	7,083	5,564	3,465	2,990	3,312	1,223	33
2000.....	43,354	168	651	1,772	10,560	6,884	6,927	5,361	3,506	3,038	3,173	1,288	26
1999.....	42,401	184	650	1,771	10,128	6,778	6,738	4,972	3,370	3,276	3,243	1,261	30

All races, male

2007.....	31,102	57	309	765	7,667	5,413	4,702	4,789	3,065	1,924	1,694	708	9
2006.....	31,633	70	316	765	7,986	5,628	4,765	4,817	3,119	1,832	1,631	692	12
2005.....	31,631	74	351	848	7,904	5,501	4,869	4,631	3,055	1,976	1,709	708	5
2004.....	30,837	80	344	975	7,814	5,245	4,826	4,486	2,770	1,799	1,811	667	20
2003.....	30,655	80	353	1,009	7,806	5,156	4,984	4,303	2,683	1,778	1,767	722	14
2002.....	31,064	68	361	963	8,178	5,349	4,984	4,189	2,540	1,888	1,793	718	33
2001.....	30,064	66	356	1,029	7,778	5,236	5,016	3,938	2,309	1,786	1,859	664	27
2000.....	29,451	90	368	1,043	7,516	5,135	4,940	3,780	2,306	1,802	1,739	706	26
1999.....	28,552	97	366	1,062	7,127	4,993	4,735	3,473	2,200	1,931	1,827	714	27

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

All races, female

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
2007.....	12,843	67	242	520	2,901	1,674	1,668	1,741	1,294	1,016	1,151	569	-
2006.....	13,683	70	272	574	3,029	1,720	1,912	1,798	1,399	1,084	1,275	548	2
2005.....	13,712	72	266	599	3,004	1,738	1,879	1,766	1,399	1,134	1,282	572	1
2004.....	14,096	63	291	678	3,173	1,791	1,837	1,790	1,323	1,175	1,369	600	6
2003.....	14,102	65	268	633	3,166	1,677	1,977	1,771	1,275	1,189	1,443	636	2
2002.....	14,316	55	249	651	3,281	1,743	2,093	1,748	1,224	1,225	1,478	567	2
2001.....	13,724	78	270	657	2,947	1,701	2,067	1,626	1,156	1,204	1,453	559	6
2000.....	13,903	78	283	729	3,044	1,749	1,987	1,581	1,200	1,236	1,434	582	-
1999.....	13,849	87	284	709	3,001	1,785	2,003	1,499	1,170	1,345	1,416	547	3

White, both sexes

2007 1/.....	36,653	91	401	1,010	8,825	5,706	5,174	5,462	3,706	2,539	2,561	1,169	9
2006 1/.....	37,769	99	426	1,023	9,272	5,864	5,495	5,503	3,777	2,531	2,606	1,161	12
2005 1/.....	38,041	99	470	1,126	9,172	5,842	5,557	5,376	3,797	2,697	2,702	1,197	6
2004 1/.....	37,618	103	480	1,292	9,325	5,617	5,466	5,170	3,488	2,578	2,880	1,195	24
2003 1/.....	37,424	108	467	1,294	9,257	5,507	5,705	5,030	3,319	2,564	2,901	1,257	15
2002.....	38,054	81	452	1,254	9,680	5,697	5,807	4,939	3,209	2,714	2,996	1,195	30
2001.....	36,576	109	466	1,275	8,929	5,613	5,828	4,600	2,936	2,599	3,043	1,149	29
2000.....	36,160	115	484	1,389	8,881	5,448	5,690	4,410	2,984	2,636	2,906	1,193	24
1999.....	35,343	133	455	1,377	8,485	5,469	5,497	4,091	2,831	2,868	2,958	1,157	22

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)													
White, male													
2007 1/.....	25,903	46	232	604	6,400	4,349	3,807	4,022	2,612	1,669	1,514	639	9
2006 1/.....	26,348	52	224	574	6,724	4,495	3,926	4,024	2,626	1,589	1,461	642	11
2005 1/.....	26,508	54	263	661	6,660	4,427	4,005	3,913	2,610	1,709	1,542	659	5
2004 1/.....	25,816	54	258	775	6,662	4,184	3,989	3,707	2,347	1,556	1,635	630	19
2003 1/.....	25,660	62	273	794	6,562	4,189	4,096	3,582	2,270	1,560	1,596	663	13
2002.....	26,000	47	260	755	6,904	4,306	4,104	3,512	2,167	1,634	1,624	659	28
2001.....	25,084	50	268	770	6,488	4,255	4,133	3,265	1,952	1,558	1,701	619	25
2000.....	24,470	64	275	815	6,317	4,067	4,057	3,131	1,937	1,552	1,575	656	24
1999.....	23,763	77	253	823	5,959	4,038	3,879	2,867	1,849	1,692	1,655	651	20
White, female													
2007 1/.....	10,750	45	169	406	2,425	1,357	1,367	1,440	1,094	870	1,047	530	-
2006 1/.....	11,421	47	202	449	2,548	1,369	1,569	1,479	1,151	942	1,145	519	1
2005 1/.....	11,533	45	207	465	2,512	1,415	1,552	1,463	1,187	988	1,160	538	1
2004 1/.....	11,802	49	222	517	2,663	1,433	1,477	1,463	1,141	1,022	1,245	565	5
2003 1/.....	11,764	46	194	500	2,695	1,318	1,609	1,448	1,049	1,004	1,305	594	2
2002.....	12,054	34	192	499	2,776	1,391	1,703	1,427	1,042	1,080	1,372	536	2
2001.....	11,492	59	198	505	2,441	1,358	1,695	1,335	984	1,041	1,342	530	4
2000.....	11,690	51	209	574	2,564	1,381	1,633	1,279	1,047	1,084	1,331	537	-
1999.....	11,580	56	202	554	2,526	1,431	1,618	1,224	982	1,176	1,303	506	2

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Black, both sexes

2007 1/.....	5,519	27	117	223	1,262	1,068	949	851	497	274	181	70	-
2006 1/.....	5,660	30	133	246	1,250	1,153	923	852	561	265	192	54	1
2005 1/.....	5,491	38	114	249	1,277	1,049	929	798	491	297	190	59	-
2004 1/.....	5,513	31	128	282	1,205	1,095	918	877	456	268	201	52	-
2003 1/.....	5,431	28	125	263	1,189	1,007	987	792	471	291	206	71	1
2002.....	5,462	35	123	287	1,279	1,072	980	750	411	268	193	61	3
2001.....	5,506	19	123	331	1,308	1,025	997	755	415	289	186	54	4
2000.....	5,499	40	134	302	1,234	1,111	987	757	394	301	173	65	1
1999.....	5,470	43	160	334	1,230	1,018	995	696	393	295	218	80	8

Black, male

2007 1/.....	4,058	10	59	131	935	847	735	633	358	184	120	46	-
2006 1/.....	4,068	15	76	149	920	898	679	617	388	178	110	37	1
2005 1/.....	3,957	16	63	145	930	830	696	571	351	201	119	35	-
2004 1/.....	3,910	19	68	158	858	843	667	639	320	180	129	29	-
2003 1/.....	3,834	12	65	161	887	759	714	575	324	172	124	40	1
2002.....	3,874	17	81	169	921	821	689	530	296	179	126	42	3
2001.....	3,900	7	67	219	953	771	713	556	298	167	116	31	2
2000.....	3,922	21	74	185	886	842	735	530	295	201	114	38	1
1999.....	3,821	17	89	205	885	771	704	506	271	180	134	52	7

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Black, female

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
2007 1/.....	1,461	17	58	92	327	221	214	218	139	90	61	24	-
2006 1/.....	1,592	15	57	97	330	255	244	235	173	87	82	17	-
2005 1/.....	1,534	22	51	104	347	219	233	227	140	96	71	24	-
2004 1/.....	1,603	12	60	124	347	252	251	238	136	88	72	23	-
2003 1/.....	1,597	16	60	102	302	248	273	217	147	119	82	31	-
2002.....	1,588	18	42	118	358	251	291	220	115	89	67	19	-
2001.....	1,606	12	56	112	355	254	284	199	117	122	70	23	2
2000.....	1,577	19	60	117	348	269	252	227	99	100	59	27	-
1999.....	1,649	26	71	129	345	247	291	190	122	115	84	28	1

American Indian 2/, both sexes

2007 1/.....	781	4	18	22	221	179	137	96	52	32	19	1	-
2006 1/.....	867	8	13	25	253	177	137	125	74	27	25	2	1
2005 1/.....	801	5	26	35	223	188	135	95	51	27	13	3	-
2004 1/.....	802	5	19	34	219	163	138	102	49	43	26	2	2
2003 1/.....	865	7	18	44	254	164	145	116	61	32	16	8	-
2002.....	866	4	16	38	243	165	168	116	62	31	18	5	-
2001.....	764	10	17	47	237	134	129	91	44	27	23	5	-
2000.....	779	7	14	44	231	163	135	82	47	31	18	6	1
1999.....	730	4	14	30	210	150	133	77	51	37	23	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

American Indian 2/, male

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
2007 1/.....	523	1	9	12	148	125	86	63	42	23	13	1	-
2006 1/.....	593	2	6	15	172	126	90	95	53	18	15	1	-
2005 1/.....	556	3	20	23	153	131	94	70	38	16	6	2	-
2004 1/.....	529	4	11	15	142	114	88	73	44	23	13	1	1
2003 1/.....	557	5	8	30	177	109	101	71	35	13	5	3	-
2002.....	572	1	9	19	164	118	110	79	36	19	14	3	-
2001.....	506	5	8	26	156	100	98	57	26	15	13	2	-
2000.....	500	2	9	23	154	114	83	58	23	18	12	3	1
1999.....	481	2	11	19	149	90	84	55	36	21	14	-	-

American Indian 2/, female

2007 1/.....	258	3	9	10	73	54	51	33	10	9	6	-	-
2006 1/.....	274	6	7	10	81	51	47	30	21	9	10	1	1
2005 1/.....	245	2	6	12	70	57	41	25	13	11	7	1	-
2004 1/.....	273	1	8	19	77	49	50	29	5	20	13	1	1
2003 1/.....	308	2	10	14	77	55	44	45	26	19	11	5	-
2002.....	294	3	7	19	79	47	58	37	26	12	4	2	-
2001.....	258	5	9	21	81	34	31	34	18	12	10	3	-
2000.....	279	5	5	21	77	49	52	24	24	13	6	3	-
1999.....	249	2	3	11	61	60	49	22	15	16	9	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	992	2	15	30	260	134	110	121	104	95	84	37	-
2006 1/.....	1,020	3	16	45	240	154	122	135	106	93	83	23	-
2005 1/.....	1,010	4	7	37	236	160	127	128	115	89	86	21	-
2004 1/.....	1,000	4	8	45	238	161	141	127	100	85	73	18	-
2003 1/.....	1,037	2	11	41	272	155	124	136	107	80	87	22	-
2002.....	998	3	19	35	257	158	122	132	82	100	64	24	2
2001.....	942	6	20	33	251	165	129	118	70	75	60	15	-
2000.....	916	6	19	37	214	162	115	112	81	70	76	24	-
1999.....	858	4	21	30	203	141	113	108	95	76	44	23	-

Asian or Pacific Islander 3/, male

2007 1/.....	618	-	9	18	184	92	74	71	53	48	47	22	-
2006 1/.....	624	1	10	27	170	109	70	81	52	47	45	12	-
2005 1/.....	610	1	5	19	161	113	74	77	56	50	42	12	-
2004 1/.....	582	3	7	27	152	104	82	67	59	40	34	7	-
2003 1/.....	604	1	7	24	180	99	73	75	54	33	42	16	-
2002.....	618	3	11	20	189	104	81	68	41	56	29	14	2
2001.....	574	4	13	14	181	110	72	60	33	46	29	12	-
2000.....	559	3	10	20	159	112	65	61	51	31	38	9	-
1999.....	487	1	13	15	134	94	68	45	44	38	24	11	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Asian or Pacific Islander 3/, female

2007 1/.....	374	2	6	12	76	42	36	50	51	47	37	15	-
2006 1/.....	396	2	6	18	70	45	52	54	54	46	38	11	-
2005 1/.....	400	3	2	18	75	47	53	51	59	39	44	9	-
2004 1/.....	418	1	1	18	86	57	59	60	41	45	39	11	-
2003 1/.....	433	1	4	17	92	56	51	61	53	47	45	6	-
2002.....	380	-	8	15	68	54	41	64	41	44	35	10	-
2001.....	368	2	7	19	70	55	57	58	37	29	31	3	-
2000.....	357	3	9	17	55	50	50	51	30	39	38	15	-
1999.....	371	3	8	15	69	47	45	63	51	38	20	12	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

All races, both sexes

2007.....	1,083	1	19	53	174	134	181	235	123	81	61	18	3
2006.....	1,181	1	12	50	168	136	217	244	151	95	78	27	2
2005.....	1,241	1	27	50	192	162	210	267	138	81	84	25	4
2004.....	1,199	1	19	57	190	184	211	222	132	92	63	27	1
2003.....	1,357	2	17	60	264	204	239	234	147	81	80	25	4
2002.....	1,086	2	17	70	159	138	223	211	109	70	66	17	4
2001.....	1,405	3	28	80	198	177	285	247	133	117	98	29	10
2000.....	1,492	2	31	98	243	212	317	234	122	100	91	35	7
1999.....	2,118	4	48	128	362	317	408	304	182	145	159	54	7

All races, male

2007.....	856	-	13	27	129	114	150	197	102	68	40	13	3
2006.....	921	-	5	40	134	104	182	196	112	72	55	19	2
2005.....	957	-	17	35	156	136	160	212	110	54	58	15	4
2004.....	926	1	12	38	152	150	168	182	102	58	46	16	1
2003.....	1,044	-	9	38	205	166	186	192	113	63	52	16	4
2002.....	882	2	10	47	134	108	187	185	92	56	50	8	3
2001.....	1,067	-	20	49	158	152	235	189	104	84	56	10	10
2000.....	1,125	2	17	61	194	171	258	166	98	75	57	19	7
1999.....	1,559	4	26	91	279	244	319	234	135	97	88	35	7

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

All races, female

2007.....	227	1	6	26	45	20	31	38	21	13	21	5	-
2006.....	260	1	7	10	34	32	35	48	39	23	23	8	-
2005.....	284	1	10	15	36	26	50	55	28	27	26	10	-
2004.....	273	-	7	19	38	34	43	40	30	34	17	11	-
2003.....	313	2	8	22	59	38	53	42	34	18	28	9	-
2002.....	204	-	7	23	25	30	36	26	17	14	16	9	1
2001.....	338	3	8	31	40	25	50	58	29	33	42	19	-
2000.....	367	-	14	37	49	41	59	68	24	25	34	16	-
1999.....	559	-	22	37	83	73	89	70	47	48	71	19	-

White, both sexes

2007 1/.....	852	1	14	42	137	101	127	189	103	68	52	16	2
2006 1/.....	959	-	7	42	141	104	168	193	127	83	68	25	1
2005 1/.....	970	1	20	41	148	120	170	196	117	59	73	22	3
2004 1/.....	945	-	15	46	141	138	178	162	107	78	54	25	1
2003 1/.....	1,121	2	14	49	227	165	194	180	126	67	70	23	4
2002.....	870	1	13	53	136	115	170	167	83	57	56	15	4
2001.....	1,105	2	23	56	161	141	208	194	107	97	86	24	6
2000.....	1,221	2	21	83	203	171	250	186	101	85	80	32	7
1999.....	1,755	3	32	99	295	259	340	250	161	123	139	48	6

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

White, male

2007 1/.....	676	-	10	18	105	89	106	159	82	58	36	11	2
2006 1/.....	748	-	3	32	114	80	145	152	91	62	51	17	1
2005 1/.....	738	-	14	28	117	101	129	153	90	38	51	14	3
2004 1/.....	723	-	9	31	109	111	140	133	83	51	40	15	1
2003 1/.....	856	-	8	29	177	134	154	144	95	52	45	14	4
2002.....	708	1	8	38	118	89	143	144	71	45	41	7	3
2001.....	850	-	17	34	132	124	173	152	88	67	50	7	6
2000.....	924	2	10	52	161	139	206	132	82	62	52	19	7
1999.....	1,281	3	15	68	223	199	266	190	119	84	78	30	6

White, female

2007 1/.....	176	1	4	24	32	12	21	30	21	10	16	5	-
2006 1/.....	211	-	4	10	27	24	23	41	36	21	17	8	-
2005 1/.....	232	1	6	13	31	19	41	43	27	21	22	8	-
2004 1/.....	222	-	6	15	32	27	38	29	24	27	14	10	-
2003 1/.....	265	2	6	20	50	31	40	36	31	15	25	9	-
2002.....	162	-	5	15	18	26	27	23	12	12	15	8	1
2001.....	255	2	6	22	29	17	35	42	19	30	36	17	-
2000.....	297	-	11	31	42	32	44	54	19	23	28	13	-
1999.....	474	-	17	31	72	60	74	60	42	39	61	18	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

Black, both sexes

2007 1/.....	165	-	3	9	30	20	36	38	17	6	5	-	1
2006 1/.....	152	1	2	5	18	22	33	40	17	7	5	1	1
2005 1/.....	204	-	5	8	32	32	34	56	13	15	6	2	1
2004 1/.....	199	1	3	8	42	34	28	43	23	8	9	-	-
2003 1/.....	190	-	3	8	27	30	39	46	19	10	7	1	-
2002.....	172	1	3	14	18	19	40	42	18	9	8	-	-
2001.....	226	-	5	23	23	25	60	42	22	15	6	1	4
2000.....	206	-	7	11	32	32	54	35	15	10	7	3	-
1999.....	253	1	13	21	34	42	52	42	17	14	11	5	1

Black, male

2007 1/.....	129	-	2	7	19	15	30	31	17	4	3	-	1
2006 1/.....	122	-	-	5	13	17	25	36	15	6	3	1	1
2005 1/.....	163	-	2	6	27	26	26	46	12	12	4	1	1
2004 1/.....	163	1	2	5	37	30	25	34	17	6	6	-	-
2003 1/.....	155	-	1	7	21	25	28	42	17	9	4	1	-
2002.....	145	1	2	7	12	17	34	40	16	9	7	-	-
2001.....	168	-	3	15	17	19	49	31	14	13	3	-	4
2000.....	158	-	5	8	26	25	43	27	12	9	3	-	-
1999.....	196	1	9	20	28	34	41	34	12	9	3	4	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

Black, female

2007 1/.....	36	-	1	2	11	5	6	7	-	2	2	-	-
2006 1/.....	30	1	2	-	5	5	8	4	2	1	2	-	-
2005 1/.....	41	-	3	2	5	6	8	10	1	3	2	1	-
2004 1/.....	36	-	1	3	5	4	3	9	6	2	3	-	-
2003 1/.....	35	-	2	1	6	5	11	4	2	1	3	-	-
2002.....	27	-	1	7	6	2	6	2	2	-	1	-	-
2001.....	58	-	2	8	6	6	11	11	8	2	3	1	-
2000.....	48	-	2	3	6	7	11	8	3	1	4	3	-
1999.....	57	-	4	1	6	8	11	8	5	5	8	1	-

American Indian 2/, both sexes

2007 1/.....	39	-	1	2	6	10	13	5	1	-	-	1	-
2006 1/.....	40	-	3	2	4	8	11	7	2	3	-	-	-
2005 1/.....	36	-	1	1	6	8	6	8	3	2	1	-	-
2004 1/.....	32	-	1	1	6	7	3	12	1	1	-	-	-
2003 1/.....	28	-	-	2	7	9	4	3	1	1	-	1	-
2002.....	21	-	-	1	5	3	9	-	1	2	-	-	-
2001.....	39	1	-	1	8	9	9	6	2	1	2	-	-
2000.....	28	-	1	2	5	3	8	6	2	1	-	-	-
1999.....	60	-	1	4	20	8	12	10	1	4	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

American Indian 2/, male

2007 1/.....	31	-	-	2	4	7	11	5	1	-	-	1	-
2006 1/.....	29	-	2	2	3	5	8	5	2	2	-	-	-
2005 1/.....	34	-	1	1	6	8	5	7	3	2	1	-	-
2004 1/.....	27	-	1	1	5	7	2	10	1	-	-	-	-
2003 1/.....	21	-	-	1	6	7	2	2	1	1	-	1	-
2002.....	17	-	-	1	4	2	7	-	1	2	-	-	-
2001.....	30	-	-	-	6	8	8	3	2	1	2	-	-
2000.....	19	-	-	-	5	3	6	4	1	-	-	-	-
1999.....	45	-	1	1	17	5	9	8	1	3	-	-	-

American Indian 2/, female

2007 1/.....	8	-	1	-	2	3	2	-	-	-	-	-	-
2006 1/.....	11	-	1	-	1	3	3	2	-	1	-	-	-
2005 1/.....	2	-	-	-	-	-	1	1	-	-	-	-	-
2004 1/.....	5	-	-	-	1	-	1	2	-	1	-	-	-
2003 1/.....	7	-	-	1	1	2	2	1	-	-	-	-	-
2002.....	4	-	-	-	1	1	2	-	-	-	-	-	-
2001.....	9	1	-	1	2	1	1	3	-	-	-	-	-
2000.....	9	-	1	2	-	-	2	2	1	1	-	-	-
1999.....	15	-	-	3	3	3	3	2	-	1	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	27	-	1	-	1	3	5	3	2	7	4	1	-
2006 1/.....	30	-	-	1	5	2	5	4	5	2	5	1	-
2005 1/.....	31	-	1	-	6	2	-	7	5	5	4	1	-
2004 1/.....	23	-	-	2	1	5	2	5	1	5	-	2	-
2003 1/.....	18	-	-	1	3	-	2	5	1	3	3	-	-
2002.....	23	-	1	2	-	1	4	2	7	2	2	2	-
2001.....	35	-	-	-	6	2	8	5	2	4	4	4	-
2000.....	37	-	2	2	3	6	5	7	4	4	4	-	-
1999.....	50	-	2	4	13	8	4	2	3	4	9	1	-

Asian or Pacific Islander 3/, male

2007 1/.....	20	-	1	-	1	3	3	2	2	6	1	1	-
2006 1/.....	22	-	-	1	4	2	4	3	4	2	1	1	-
2005 1/.....	22	-	-	-	6	1	-	6	5	2	2	-	-
2004 1/.....	13	-	-	1	1	2	1	5	1	1	-	1	-
2003 1/.....	12	-	-	1	1	-	2	4	-	1	3	-	-
2002.....	12	-	-	1	-	-	3	1	4	-	2	1	-
2001.....	19	-	-	-	3	1	5	3	-	3	1	3	-
2000.....	24	-	2	1	2	4	3	3	3	4	2	-	-
1999.....	37	-	1	2	11	6	3	2	3	1	7	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

Asian or Pacific Islander 3/, female

2007 1/.....	7	-	-	-	-	-	2	1	-	1	3	-	-
2006 1/.....	8	-	-	-	1	-	1	1	1	-	4	-	-
2005 1/.....	9	-	1	-	-	1	-	1	-	3	2	1	-
2004 1/.....	10	-	-	1	-	3	1	-	-	4	-	1	-
2003 1/.....	6	-	-	-	2	-	-	1	1	2	-	-	-
2002.....	11	-	1	1	-	1	1	1	3	2	-	1	-
2001.....	16	-	-	-	3	1	3	2	2	1	3	1	-
2000.....	13	-	-	1	1	2	2	4	1	-	2	-	-
1999.....	13	-	1	2	2	2	1	-	-	3	2	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)													
All races, both sexes													
2007.....	1,816	2	11	36	186	231	278	434	356	173	77	31	1
2006.....	1,915	1	15	64	209	269	355	405	317	159	98	23	-
2005.....	1,857	-	2	50	188	237	337	437	326	162	93	24	1
2004.....	1,921	1	5	45	200	301	373	455	285	163	67	26	-
2003.....	1,957	1	11	63	196	243	394	446	336	162	87	18	-
2002.....	1,900	2	13	54	198	254	388	415	292	168	96	20	-
2001.....	2,095	4	24	57	216	340	459	440	263	175	97	20	-
2000.....	1,903	4	9	52	225	304	417	370	255	177	78	12	-
1999.....	1,904	1	14	76	199	339	407	369	260	158	66	15	-
All races, male													
2007.....	1,476	1	6	21	147	183	232	362	293	146	64	20	1
2006.....	1,511	-	8	44	161	219	286	334	252	122	72	13	-
2005.....	1,498	-	1	27	148	203	274	365	270	127	68	14	1
2004.....	1,548	-	3	26	147	246	303	374	242	141	48	18	-
2003.....	1,592	1	4	37	160	189	327	370	291	134	66	13	-
2002.....	1,587	2	10	29	159	219	328	353	245	157	74	11	-
2001.....	1,663	2	16	37	175	273	362	357	218	145	68	10	-
2000.....	1,590	2	6	34	180	264	362	314	213	144	62	9	-
1999.....	1,531	-	8	53	152	274	322	309	226	134	44	9	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

All races, female

2007.....	340	1	5	15	39	48	46	72	63	27	13	11	-
2006.....	404	1	7	20	48	50	69	71	65	37	26	10	-
2005.....	359	-	1	23	40	34	63	72	56	35	25	10	-
2004.....	373	1	2	19	53	55	70	81	43	22	19	8	-
2003.....	365	-	7	26	36	54	67	76	45	28	21	5	-
2002.....	313	-	3	25	39	35	60	62	47	11	22	9	-
2001.....	432	2	8	20	41	67	97	83	45	30	29	10	-
2000.....	313	2	3	18	45	40	55	56	42	33	16	3	-
1999.....	373	1	6	23	47	65	85	60	34	24	22	6	-

White, both sexes

2007 1/.....	1,581	1	11	27	153	195	252	378	312	152	70	29	1
2006 1/.....	1,694	1	12	56	185	225	312	359	289	148	90	17	-
2005 1/.....	1,636	-	2	43	158	206	295	370	300	155	84	22	1
2004 1/.....	1,716	1	4	39	182	267	321	408	259	149	61	25	-
2003 1/.....	1,737	1	10	54	159	213	354	397	308	146	77	18	-
2002.....	1,690	2	13	40	175	225	345	370	266	150	85	19	-
2001.....	1,863	4	22	47	195	299	415	381	234	161	88	17	-
2000.....	1,707	4	8	47	195	271	378	328	234	157	74	11	-
1999.....	1,669	1	10	66	173	298	362	324	222	143	57	13	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)													
White, male													
2007 1/.....	1,285	-	6	17	121	154	212	310	260	127	57	20	1
2006 1/.....	1,333	-	6	39	141	183	248	296	229	113	68	10	-
2005 1/.....	1,322	-	1	22	121	179	238	314	250	121	61	14	1
2004 1/.....	1,379	-	2	23	130	215	259	339	219	130	45	17	-
2003 1/.....	1,409	1	4	28	128	166	290	331	268	121	59	13	-
2002.....	1,410	2	10	21	139	195	293	317	220	139	64	10	-
2001.....	1,473	2	14	31	154	241	326	310	190	132	63	10	-
2000.....	1,421	2	5	29	154	235	326	280	194	130	58	8	-
1999.....	1,341	-	5	45	134	240	283	274	193	120	40	7	-
White, female													
2007 1/.....	296	1	5	10	32	41	40	68	52	25	13	9	-
2006 1/.....	361	1	6	17	44	42	64	63	60	35	22	7	-
2005 1/.....	314	-	1	21	37	27	57	56	50	34	23	8	-
2004 1/.....	337	1	2	16	52	52	62	69	40	19	16	8	-
2003 1/.....	328	-	6	26	31	47	64	66	40	25	18	5	-
2002.....	280	-	3	19	36	30	52	53	46	11	21	9	-
2001.....	390	2	8	16	41	58	89	71	44	29	25	7	-
2000.....	286	2	3	18	41	36	52	48	40	27	16	3	-
1999.....	328	1	5	21	39	58	79	50	29	23	17	6	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

Black, both sexes

2007 1/.....	175	1	-	6	24	24	19	45	33	17	5	1	-
2006 1/.....	171	-	2	6	14	34	36	37	23	7	7	5	-
2005 1/.....	147	-	-	4	19	18	27	50	19	4	5	1	-
2004 1/.....	156	-	1	5	13	25	36	39	23	9	5	-	-
2003 1/.....	150	-	1	7	26	19	28	32	20	10	7	-	-
2002.....	162	-	-	11	17	21	34	36	22	12	8	1	-
2001.....	174	-	2	9	16	30	30	45	23	9	8	2	-
2000.....	134	-	-	4	17	26	28	26	15	14	3	1	-
1999.....	178	-	4	9	20	29	38	34	26	9	7	2	-

Black, male

2007 1/.....	149	1	-	3	21	19	16	42	26	16	5	-	-
2006 1/.....	143	-	1	4	13	31	31	32	19	6	3	3	-
2005 1/.....	117	-	-	4	18	14	23	37	14	4	3	-	-
2004 1/.....	131	-	1	2	12	23	30	32	21	7	3	-	-
2003 1/.....	130	-	-	7	25	15	26	26	18	8	5	-	-
2002.....	137	-	-	7	16	18	26	28	21	12	8	1	-
2001.....	146	-	2	6	16	23	26	37	23	9	4	-	-
2000.....	119	-	-	4	16	23	26	22	14	10	3	1	-
1999.....	145	-	3	7	13	23	33	27	24	9	4	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

Black, female

2007 1/.....	26	-	-	3	3	5	3	3	7	1	-	1	-
2006 1/.....	28	-	1	2	1	3	5	5	4	1	4	2	-
2005 1/.....	30	-	-	-	1	4	4	13	5	-	2	1	-
2004 1/.....	25	-	-	3	1	2	6	7	2	2	2	-	-
2003 1/.....	20	-	1	-	1	4	2	6	2	2	2	-	-
2002.....	25	-	-	4	1	3	8	8	1	-	-	-	-
2001.....	28	-	-	3	-	7	4	8	-	-	4	2	-
2000.....	15	-	-	-	1	3	2	4	1	4	-	-	-
1999.....	33	-	1	2	7	6	5	7	2	-	3	-	-

American Indian 2/, both sexes

2007 1/.....	30	-	-	-	4	7	2	7	7	3	-	-	-
2006 1/.....	23	-	-	1	4	6	4	6	1	-	1	-	-
2005 1/.....	38	-	-	2	6	5	9	8	5	2	1	-	-
2004 1/.....	17	-	-	-	1	4	3	4	1	4	-	-	-
2003 1/.....	22	-	-	-	2	5	5	4	3	3	-	-	-
2002.....	16	-	-	-	2	3	6	3	-	2	-	-	-
2001.....	29	-	-	-	2	4	8	8	4	3	-	-	-
2000.....	22	-	1	-	5	1	3	8	1	2	1	-	-
1999.....	26	-	-	1	3	5	3	4	7	3	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

American Indian 2/, male

2007 1/.....	22	-	-	-	3	6	1	6	4	2	-	-	-
2006 1/.....	16	-	-	1	3	3	4	3	1	-	1	-	-
2005 1/.....	32	-	-	1	4	4	8	7	5	2	1	-	-
2004 1/.....	13	-	-	-	1	3	3	2	1	3	-	-	-
2003 1/.....	17	-	-	-	2	5	4	2	1	3	-	-	-
2002.....	14	-	-	-	1	2	6	3	-	2	-	-	-
2001.....	22	-	-	-	2	3	6	5	4	2	-	-	-
2000.....	18	-	1	-	3	-	3	8	1	1	1	-	-
1999.....	21	-	-	1	2	5	3	3	5	2	-	-	-

American Indian 2/, female

2007 1/.....	8	-	-	-	1	1	1	1	3	1	-	-	-
2006 1/.....	7	-	-	-	1	3	-	3	-	-	-	-	-
2005 1/.....	6	-	-	1	2	1	1	1	-	-	-	-	-
2004 1/.....	4	-	-	-	-	1	-	2	-	1	-	-	-
2003 1/.....	5	-	-	-	-	-	1	2	2	-	-	-	-
2002.....	2	-	-	-	1	1	-	-	-	-	-	-	-
2001.....	7	-	-	-	-	1	2	3	-	1	-	-	-
2000.....	4	-	-	-	2	1	-	-	-	1	-	-	-
1999.....	5	-	-	-	1	-	-	1	2	1	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	30	-	-	3	5	5	5	4	4	1	2	1	-
2006 1/.....	27	-	1	1	6	4	3	3	4	4	-	1	-
2005 1/.....	36	-	-	1	5	8	6	9	2	1	3	1	-
2004 1/.....	32	-	-	1	4	5	13	4	2	1	1	1	-
2003 1/.....	48	-	-	2	9	6	7	13	5	3	3	-	-
2002.....	32	-	-	3	4	5	3	6	4	4	3	-	-
2001.....	29	-	-	1	3	7	6	6	2	2	1	1	-
2000.....	40	-	-	1	8	6	8	8	5	4	-	-	-
1999.....	31	-	-	-	3	7	4	7	5	3	2	-	-

Asian or Pacific Islander 3/, male

2007 1/.....	20	-	-	1	2	4	3	4	3	1	2	-	-
2006 1/.....	19	-	1	-	4	2	3	3	3	3	-	-	-
2005 1/.....	27	-	-	-	5	6	5	7	1	-	3	-	-
2004 1/.....	25	-	-	1	4	5	11	1	1	1	-	1	-
2003 1/.....	36	-	-	2	5	3	7	11	4	2	2	-	-
2002.....	26	-	-	1	3	4	3	5	4	4	2	-	-
2001.....	22	-	-	-	3	6	4	5	1	2	1	-	-
2000.....	32	-	-	1	7	6	7	4	4	3	-	-	-
1999.....	24	-	-	-	3	6	3	5	4	3	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

Asian or Pacific Islander 3/, female

2007 1/.....	10	-	-	2	3	1	2	-	1	-	-	1	-
2006 1/.....	8	-	-	1	2	2	-	-	1	1	-	1	-
2005 1/.....	9	-	-	1	-	2	1	2	1	1	-	1	-
2004 1/.....	7	-	-	-	-	-	2	3	1	-	1	-	-
2003 1/.....	12	-	-	-	4	3	-	2	1	1	1	-	-
2002.....	6	-	-	2	1	1	-	1	-	-	1	-	-
2001.....	7	-	-	1	-	1	2	1	1	-	-	1	-
2000.....	8	-	-	-	1	-	1	4	1	1	-	-	-
1999.....	7	-	-	-	-	1	1	2	1	-	2	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nontransport accidents (W00-X59,Y86)													
All races, both sexes													
2007.....	76,862	1,158	1,007	820	4,969	7,525	10,102	13,116	7,355	5,559	10,753	14,477	21
2006.....	73,187	1,005	995	805	4,837	7,201	10,285	12,411	6,460	5,250	10,626	13,271	41
2005.....	69,368	936	1,018	868	4,465	6,359	9,624	11,238	5,935	5,279	10,686	12,914	46
2004.....	63,959	907	982	911	4,072	5,511	9,224	9,989	5,141	4,887	10,147	12,127	61
2003.....	61,206	797	1,068	853	3,840	5,261	9,172	9,083	4,729	4,871	9,731	11,745	56
2002.....	58,376	819	1,001	980	3,596	5,085	9,022	8,112	4,180	4,735	9,471	11,329	46
2001.....	54,249	825	1,036	1,013	3,272	4,385	8,118	7,093	3,797	4,553	9,181	10,899	77
2000.....	51,151	707	1,135	1,057	3,085	4,369	7,752	6,313	3,622	4,383	8,416	10,260	52
1999.....	51,437	656	1,186	1,116	2,967	4,456	7,678	5,994	3,473	4,629	8,814	10,399	69
All races, male													
2007.....	46,393	657	659	533	3,962	5,674	6,938	8,669	4,897	3,455	5,470	5,460	19
2006.....	44,876	597	598	536	3,953	5,515	7,198	8,490	4,365	3,186	5,436	4,968	34
2005.....	42,289	504	658	583	3,619	4,825	6,723	7,794	4,072	3,186	5,502	4,784	39
2004.....	38,739	523	605	622	3,310	4,230	6,374	6,946	3,448	2,994	5,125	4,509	53
2003.....	37,241	458	682	578	3,120	4,023	6,506	6,331	3,237	2,946	4,966	4,342	52
2002.....	35,724	481	643	674	2,967	3,959	6,513	5,765	2,904	2,930	4,770	4,080	38
2001.....	33,266	473	653	688	2,721	3,431	5,999	5,259	2,597	2,736	4,669	3,973	67
2000.....	31,651	407	707	723	2,570	3,489	5,884	4,803	2,516	2,618	4,205	3,679	50
1999.....	31,893	393	727	768	2,482	3,586	5,872	4,586	2,454	2,830	4,413	3,722	60

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nontransport accidents (W00-X59,Y86)													
All races, female													
2007.....	30,469	501	348	287	1,007	1,851	3,164	4,447	2,458	2,104	5,283	9,017	2
2006.....	28,311	408	397	269	884	1,686	3,087	3,921	2,095	2,064	5,190	8,303	7
2005.....	27,079	432	360	285	846	1,534	2,901	3,444	1,863	2,093	5,184	8,130	7
2004.....	25,220	384	377	289	762	1,281	2,850	3,043	1,693	1,893	5,022	7,618	8
2003.....	23,965	339	386	275	720	1,238	2,666	2,752	1,492	1,925	4,765	7,403	4
2002.....	22,652	338	358	306	629	1,126	2,509	2,347	1,276	1,805	4,701	7,249	8
2001.....	20,983	352	383	325	551	954	2,119	1,834	1,200	1,817	4,512	6,926	10
2000.....	19,500	300	428	334	515	880	1,868	1,510	1,106	1,765	4,211	6,581	2
1999.....	19,544	263	459	348	485	870	1,806	1,408	1,019	1,799	4,401	6,677	9
White, both sexes													
2007 1/.....	67,166	710	743	546	4,366	6,644	8,658	10,989	6,103	4,770	9,901	13,719	17
2006 1/.....	63,431	630	725	547	4,226	6,280	8,672	10,190	5,321	4,504	9,765	12,534	37
2005 1/.....	59,759	634	770	575	3,847	5,460	8,150	9,193	4,818	4,513	9,665	12,092	42
2004 1/.....	55,611	602	743	576	3,526	4,765	7,876	8,290	4,224	4,201	9,299	11,455	54
2003 1/.....	53,099	513	813	557	3,347	4,515	7,763	7,413	3,948	4,175	8,912	11,092	51
2002.....	50,252	499	721	618	3,057	4,341	7,579	6,593	3,427	4,046	8,679	10,648	44
2001.....	46,420	532	798	682	2,753	3,674	6,692	5,650	3,090	3,921	8,371	10,189	68
2000.....	43,504	448	822	700	2,544	3,629	6,345	4,977	2,956	3,739	7,680	9,616	48
1999.....	43,478	425	877	728	2,411	3,635	6,301	4,703	2,748	3,893	7,976	9,720	61

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nontransport accidents (W00-X59,Y86)													
White, male													
2007 1/.....	40,195	409	490	361	3,495	5,032	5,949	7,234	4,030	2,944	5,056	5,178	17
2006 1/.....	38,414	369	442	372	3,444	4,806	6,074	6,908	3,537	2,738	5,004	4,689	31
2005 1/.....	36,032	334	515	390	3,137	4,169	5,724	6,334	3,235	2,718	4,973	4,466	37
2004 1/.....	33,349	360	465	394	2,869	3,670	5,477	5,754	2,789	2,590	4,703	4,230	48
2003 1/.....	31,987	296	522	382	2,729	3,475	5,548	5,163	2,675	2,526	4,535	4,089	47
2002.....	30,349	308	458	426	2,518	3,396	5,500	4,691	2,349	2,498	4,361	3,808	36
2001.....	28,086	304	516	450	2,293	2,898	5,015	4,170	2,089	2,362	4,236	3,693	60
2000.....	26,514	263	518	482	2,138	2,928	4,873	3,779	2,029	2,233	3,808	3,417	46
1999.....	26,513	253	538	498	2,011	2,965	4,860	3,588	1,939	2,371	3,985	3,452	53
White, female													
2007 1/.....	26,971	301	253	185	871	1,612	2,709	3,755	2,073	1,826	4,845	8,541	-
2006 1/.....	25,017	261	283	175	782	1,474	2,598	3,282	1,784	1,766	4,761	7,845	6
2005 1/.....	23,727	300	255	185	710	1,291	2,426	2,859	1,583	1,795	4,692	7,626	5
2004 1/.....	22,262	242	278	182	657	1,095	2,399	2,536	1,435	1,611	4,596	7,225	6
2003 1/.....	21,112	217	291	175	618	1,040	2,215	2,250	1,273	1,649	4,377	7,003	4
2002.....	19,903	191	263	192	539	945	2,079	1,902	1,078	1,548	4,318	6,840	8
2001.....	18,334	228	282	232	460	776	1,677	1,480	1,001	1,559	4,135	6,496	8
2000.....	16,990	185	304	218	406	701	1,472	1,198	927	1,506	3,872	6,199	2
1999.....	16,965	172	339	230	400	670	1,441	1,115	809	1,522	3,991	6,268	8

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nontransport accidents (W00-X59,Y86)													
Black, both sexes													
2007 1/.....	7,700	400	205	221	433	667	1,154	1,832	1,052	614	573	546	3
2006 1/.....	7,934	327	234	214	468	724	1,355	1,927	956	586	620	520	3
2005 1/.....	7,810	276	208	240	452	697	1,218	1,785	948	611	779	592	4
2004 1/.....	6,802	271	192	275	401	586	1,122	1,496	766	550	651	488	4
2003 1/.....	6,580	259	214	245	376	579	1,170	1,454	659	551	592	477	4
2002.....	6,717	294	233	309	412	599	1,214	1,351	642	543	597	522	1
2001.....	6,556	262	209	281	395	558	1,234	1,291	604	539	636	541	6
2000.....	6,438	234	264	314	412	601	1,218	1,170	573	543	595	510	4
1999.....	6,827	208	264	343	435	676	1,200	1,167	630	633	700	564	7
Black, male													
2007 1/.....	4,932	211	128	138	334	488	782	1,243	750	400	270	187	1
2006 1/.....	5,272	198	137	137	385	557	934	1,370	700	358	312	182	2
2005 1/.....	5,092	156	119	159	359	499	812	1,282	717	377	404	206	2
2004 1/.....	4,399	143	117	186	321	437	735	1,042	554	339	336	186	3
2003 1/.....	4,266	148	138	159	294	422	797	1,009	477	339	304	175	4
2002.....	4,456	158	156	217	349	453	843	954	475	339	311	200	1
2001.....	4,323	155	122	202	323	407	845	973	429	328	339	194	6
2000.....	4,332	128	160	217	333	450	874	896	429	323	318	200	4
1999.....	4,611	123	165	240	375	507	882	904	448	402	347	212	6

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nontransport accidents (W00-X59,Y86)													
Black, female													
2007 1/.....	2,768	189	77	83	99	179	372	589	302	214	303	359	2
2006 1/.....	2,662	129	97	77	83	167	421	557	256	228	308	338	1
2005 1/.....	2,718	120	89	81	93	198	406	503	231	234	375	386	2
2004 1/.....	2,403	128	75	89	80	149	387	454	212	211	315	302	1
2003 1/.....	2,314	111	76	86	82	157	373	445	182	212	288	302	-
2002.....	2,261	136	77	92	63	146	371	397	167	204	286	322	-
2001.....	2,233	107	87	79	72	151	389	318	175	211	297	347	-
2000.....	2,106	106	104	97	79	151	344	274	144	220	277	310	-
1999.....	2,216	85	99	103	60	169	318	263	182	231	353	352	1
American Indian 2/, both sexes													
2007 1/.....	851	27	24	22	88	113	168	172	90	58	58	31	-
2006 1/.....	774	28	16	15	67	114	148	175	80	55	42	34	-
2005 1/.....	751	14	17	17	86	123	155	147	76	44	45	27	-
2004 1/.....	669	18	22	26	69	88	146	116	66	41	39	35	3
2003 1/.....	658	13	15	22	52	86	159	118	63	48	46	35	1
2002.....	585	11	17	21	50	83	150	96	49	44	39	24	1
2001.....	529	21	13	25	46	69	114	88	48	30	38	37	-
2000.....	524	13	23	24	58	72	110	101	44	29	32	18	-
1999.....	511	9	22	16	56	87	104	72	46	38	36	25	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Nontransport accidents (W00-X59,Y86)

American Indian 2/, male

2007 1/.....	553	22	15	16	69	76	118	103	48	37	37	12	-
2006 1/.....	546	19	7	8	59	91	108	126	58	35	23	12	-
2005 1/.....	496	7	10	11	60	97	106	97	53	25	20	10	-
2004 1/.....	441	11	14	17	55	60	98	84	48	20	14	18	2
2003 1/.....	448	9	11	17	43	56	108	86	45	34	24	14	1
2002.....	400	7	10	11	43	64	105	67	35	33	16	8	1
2001.....	350	9	7	20	39	52	76	65	36	15	19	12	-
2000.....	363	9	15	14	40	55	74	80	31	17	19	9	-
1999.....	343	6	12	7	46	63	71	55	33	25	15	10	-

American Indian 2/, female

2007 1/.....	298	5	9	6	19	37	50	69	42	21	21	19	-
2006 1/.....	228	9	9	7	8	23	40	49	22	20	19	22	-
2005 1/.....	255	7	7	6	26	26	49	50	23	19	25	17	-
2004 1/.....	228	7	8	9	14	28	48	32	18	21	25	17	1
2003 1/.....	210	4	4	5	9	30	51	32	18	14	22	21	-
2002.....	185	4	7	10	7	19	45	29	14	11	23	16	-
2001.....	179	12	6	5	7	17	38	23	12	15	19	25	-
2000.....	161	4	8	10	18	17	36	21	13	12	13	9	-
1999.....	168	3	10	9	10	24	33	17	13	13	21	15	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nontransport accidents (W00-X59,Y86)													
Asian or Pacific Islander 3/, both sexes													
2007 1/.....	1,145	21	35	31	82	101	122	123	110	117	221	181	1
2006 1/.....	1,048	20	20	29	76	83	110	119	103	105	199	183	1
2005 1/.....	1,048	12	23	36	80	79	101	113	93	111	197	203	-
2004 1/.....	877	16	25	34	76	72	80	87	85	95	158	149	-
2003 1/.....	869	12	26	29	65	81	80	98	59	97	181	141	-
2002.....	822	15	30	32	77	62	79	72	62	102	156	135	-
2001.....	744	10	16	25	78	84	78	64	55	63	136	132	3
2000.....	685	12	26	19	71	67	79	65	49	72	109	116	-
1999.....	621	14	23	29	65	58	73	52	49	65	102	90	1
Asian or Pacific Islander 3/, male													
2007 1/.....	713	15	26	18	64	78	89	89	69	74	107	83	1
2006 1/.....	644	11	12	19	65	61	82	86	70	55	97	85	1
2005 1/.....	669	7	14	23	63	60	81	81	67	66	105	102	-
2004 1/.....	550	9	9	25	65	63	64	66	57	45	72	75	-
2003 1/.....	540	5	11	20	54	70	53	73	40	47	103	64	-
2002.....	519	8	19	20	57	46	65	53	45	60	82	64	-
2001.....	507	5	8	16	66	74	63	51	43	31	75	74	1
2000.....	442	7	14	10	59	56	63	48	27	45	60	53	-
1999.....	426	11	12	23	50	51	59	39	34	32	66	48	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Nontransport accidents (W00-X59,Y86)

Asian or Pacific Islander 3/, female

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
2007 1/.....	432	6	9	13	18	23	33	34	41	43	114	98	-
2006 1/.....	404	9	8	10	11	22	28	33	33	50	102	98	-
2005 1/.....	379	5	9	13	17	19	20	32	26	45	92	101	-
2004 1/.....	327	7	16	9	11	9	16	21	28	50	86	74	-
2003 1/.....	329	7	15	9	11	11	27	25	19	50	78	77	-
2002.....	303	7	11	12	20	16	14	19	17	42	74	71	-
2001.....	237	5	8	9	12	10	15	13	12	32	61	58	2
2000.....	243	5	12	9	12	11	16	17	22	27	49	63	-
1999.....	195	3	11	6	15	7	14	13	15	33	36	42	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Falls (W00-W19)													
All races, both sexes													
2007.....	22,631	24	36	32	233	334	593	1,304	1,739	2,594	6,552	9,188	2
2006.....	20,823	23	38	38	242	318	628	1,290	1,592	2,392	6,227	8,031	4
2005.....	19,656	16	34	32	236	295	607	1,181	1,451	2,319	5,957	7,526	2
2004.....	18,807	23	47	37	241	320	659	1,184	1,393	2,255	5,682	6,962	4
2003.....	17,229	14	54	45	230	285	636	1,043	1,220	2,048	5,249	6,404	1
2002.....	16,257	16	37	42	247	307	664	1,017	1,089	1,967	4,880	5,990	1
2001.....	15,019	23	32	66	256	340	647	1,024	1,004	1,833	4,440	5,350	4
2000.....	13,322	8	36	37	237	303	608	871	949	1,660	3,841	4,772	-
1999.....	13,162	12	55	53	242	343	645	824	887	1,663	3,856	4,578	4
All races, male													
2007.....	11,597	14	24	23	197	288	476	982	1,183	1,582	3,331	3,495	2
2006.....	10,808	16	26	23	213	272	510	1,001	1,108	1,431	3,147	3,058	3
2005.....	10,154	6	20	24	198	256	489	897	1,023	1,384	3,051	2,804	2
2004.....	9,856	13	31	22	206	274	526	912	987	1,394	2,858	2,630	3
2003.....	8,910	9	39	30	186	237	525	800	827	1,209	2,661	2,386	1
2002.....	8,463	9	28	27	211	269	522	772	771	1,202	2,444	2,208	-
2001.....	8,089	15	25	46	232	295	544	821	718	1,090	2,277	2,023	3
2000.....	7,122	6	23	30	203	272	496	690	680	991	1,947	1,784	-
1999.....	7,109	8	37	40	209	294	528	661	650	1,027	1,950	1,701	4

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Falls (W00-W19)													
All races, female													
2007.....	11,034	10	12	9	36	46	117	322	556	1,012	3,221	5,693	-
2006.....	10,015	7	12	15	29	46	118	289	484	961	3,080	4,973	1
2005.....	9,502	10	14	8	38	39	118	284	428	935	2,906	4,722	-
2004.....	8,951	10	16	15	35	46	133	272	406	861	2,824	4,332	1
2003.....	8,319	5	15	15	44	48	111	243	393	839	2,588	4,018	-
2002.....	7,794	7	9	15	36	38	142	245	318	765	2,436	3,782	1
2001.....	6,930	8	7	20	24	45	103	203	286	743	2,163	3,327	1
2000.....	6,200	2	13	7	34	31	112	181	269	669	1,894	2,988	-
1999.....	6,053	4	18	13	33	49	117	163	237	636	1,906	2,877	-
White, both sexes													
2007 1/.....	21,020	20	23	28	208	296	498	1,109	1,524	2,320	6,164	8,828	2
2006 1/.....	19,195	17	22	27	215	278	536	1,080	1,367	2,140	5,841	7,669	3
2005 1/.....	18,113	13	19	24	218	248	524	993	1,247	2,088	5,556	7,181	2
2004 1/.....	17,383	18	33	30	217	271	556	993	1,190	2,022	5,364	6,685	4
2003 1/.....	15,855	10	38	39	205	249	531	842	1,051	1,826	4,918	6,146	-
2002.....	15,027	12	23	33	219	271	560	858	944	1,765	4,613	5,728	1
2001.....	13,827	15	28	56	237	297	537	860	859	1,653	4,181	5,100	4
2000.....	12,252	5	23	30	217	265	504	729	802	1,480	3,627	4,570	-
1999.....	11,992	8	43	42	212	284	547	663	722	1,469	3,632	4,366	4

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Falls (W00-W19)													
White, male													
2007 1/.....	10,643	12	16	21	174	256	404	831	1,032	1,406	3,135	3,354	2
2006 1/.....	9,829	12	15	17	187	240	438	836	948	1,279	2,947	2,907	3
2005 1/.....	9,246	5	12	18	184	219	423	750	877	1,253	2,848	2,655	2
2004 1/.....	8,988	11	23	17	187	234	458	765	828	1,259	2,702	2,501	3
2003 1/.....	8,099	6	29	25	169	209	444	648	713	1,075	2,493	2,288	-
2002.....	7,699	6	17	22	186	239	440	650	666	1,079	2,301	2,093	-
2001.....	7,331	10	21	40	217	260	456	682	614	985	2,134	1,909	3
2000.....	6,449	3	13	25	187	239	416	579	574	889	1,830	1,694	-
1999.....	6,352	7	29	32	186	246	442	535	530	904	1,828	1,609	4
White, female													
2007 1/.....	10,377	8	7	7	34	40	94	278	492	914	3,029	5,474	-
2006 1/.....	9,366	5	7	10	28	38	98	244	419	861	2,894	4,762	-
2005 1/.....	8,867	8	7	6	34	29	101	243	370	835	2,708	4,526	-
2004 1/.....	8,395	7	10	13	30	37	98	228	362	763	2,662	4,184	1
2003 1/.....	7,756	4	9	14	36	40	87	194	338	751	2,425	3,858	-
2002.....	7,328	6	6	11	33	32	120	208	278	686	2,312	3,635	1
2001.....	6,496	5	7	16	20	37	81	178	245	668	2,047	3,191	1
2000.....	5,803	2	10	5	30	26	88	150	228	591	1,797	2,876	-
1999.....	5,640	1	14	10	26	38	105	128	192	565	1,804	2,757	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Falls (W00-W19)													
Black, both sexes													
2007 1/.....	1,015	4	5	3	14	23	62	151	153	170	209	221	-
2006 1/.....	1,066	5	12	7	20	24	65	162	152	173	237	208	1
2005 1/.....	1,027	2	12	6	11	30	54	156	146	156	257	197	-
2004 1/.....	999	3	9	5	14	36	81	147	162	175	202	165	-
2003 1/.....	920	2	13	3	17	27	76	156	133	152	190	151	-
2002.....	838	3	8	6	17	24	77	133	111	138	155	166	-
2001.....	854	6	3	7	13	28	91	130	111	145	175	145	-
2000.....	760	2	13	7	13	23	74	114	116	137	137	124	-
1999.....	878	2	11	10	21	43	75	137	131	151	155	142	-
Black, male													
2007 1/.....	602	2	3	2	12	19	46	117	112	113	100	76	-
2006 1/.....	648	3	8	4	20	18	49	126	112	108	124	76	-
2005 1/.....	602	-	5	4	9	21	44	124	107	92	128	68	-
2004 1/.....	628	-	7	3	9	30	56	112	130	104	105	72	-
2003 1/.....	537	1	9	3	10	20	59	116	88	89	88	54	-
2002.....	532	2	7	5	15	21	60	103	79	83	85	72	-
2001.....	547	4	3	4	9	21	72	111	79	87	97	60	-
2000.....	485	2	10	5	11	19	57	91	89	78	70	53	-
1999.....	564	1	8	7	16	35	67	108	93	100	79	50	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Falls (W00-W19)

Black, female

2007 1/.....	413	2	2	1	2	4	16	34	41	57	109	145	-
2006 1/.....	418	2	4	3	-	6	16	36	40	65	113	132	1
2005 1/.....	425	2	7	2	2	9	10	32	39	64	129	129	-
2004 1/.....	371	3	2	2	5	6	25	35	32	71	97	93	-
2003 1/.....	383	1	4	-	7	7	17	40	45	63	102	97	-
2002.....	306	1	1	1	2	3	17	30	32	55	70	94	-
2001.....	307	2	-	3	4	7	19	19	32	58	78	85	-
2000.....	275	-	3	2	2	4	17	23	27	59	67	71	-
1999.....	314	1	3	3	5	8	8	29	38	51	76	92	-

American Indian 2/, both sexes

2007 1/.....	132	-	2	-	2	7	15	16	15	28	34	13	-
2006 1/.....	126	1	2	1	2	6	14	21	22	18	19	20	-
2005 1/.....	117	1	-	1	5	8	16	20	17	12	21	16	-
2004 1/.....	110	2	1	-	5	8	10	21	18	12	14	19	-
2003 1/.....	120	-	-	-	4	5	16	21	16	16	21	20	1
2002.....	97	1	-	2	4	6	13	15	11	11	21	13	-
2001.....	86	-	-	1	3	4	9	16	12	8	12	21	-
2000.....	81	1	-	-	1	9	13	16	12	6	12	11	-
1999.....	75	-	1	-	6	6	7	10	12	10	8	15	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Falls (W00-W19)

American Indian 2/, male

2007 1/.....	89	-	1	-	2	5	12	12	11	17	22	7	-
2006 1/.....	84	1	2	-	2	5	12	16	15	11	9	11	-
2005 1/.....	65	1	-	1	4	7	12	12	10	5	6	7	-
2004 1/.....	69	2	-	-	5	7	3	17	14	5	5	11	-
2003 1/.....	83	-	-	-	4	5	13	15	13	14	11	7	1
2002.....	54	1	-	-	4	5	10	9	9	6	7	3	-
2001.....	51	-	-	1	3	3	7	13	10	3	6	5	-
2000.....	57	1	-	-	1	8	10	11	7	2	10	7	-
1999.....	42	-	-	-	5	4	7	7	8	4	1	6	-

American Indian 2/, female

2007 1/.....	43	-	1	-	-	2	3	4	4	11	12	6	-
2006 1/.....	42	-	-	1	-	1	2	5	7	7	10	9	-
2005 1/.....	52	-	-	-	1	1	4	8	7	7	15	9	-
2004 1/.....	41	-	1	-	-	1	7	4	4	7	9	8	-
2003 1/.....	37	-	-	-	-	-	3	6	3	2	10	13	-
2002.....	43	-	-	2	-	1	3	6	2	5	14	10	-
2001.....	35	-	-	-	-	1	2	3	2	5	6	16	-
2000.....	24	-	-	-	-	1	3	5	5	4	2	4	-
1999.....	33	-	1	-	1	2	-	3	4	6	7	9	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Falls (W00-W19)													
Asian or Pacific Islander 3/, both sexes													
2007 1/.....	464	-	6	1	9	8	18	28	47	76	145	126	-
2006 1/.....	436	-	2	3	5	10	13	27	51	61	130	134	-
2005 1/.....	399	-	3	1	2	9	13	12	41	63	123	132	-
2004 1/.....	315	-	4	2	5	5	12	23	23	46	102	93	-
2003 1/.....	334	2	3	3	4	4	13	24	20	54	120	87	-
2002.....	295	-	6	1	7	6	14	11	23	53	91	83	-
2001.....	252	2	1	2	3	11	10	18	22	27	72	84	-
2000.....	229	-	-	-	6	6	17	12	19	37	65	67	-
1999.....	217	2	-	1	3	10	16	14	22	33	61	55	-
Asian or Pacific Islander 3/, male													
2007 1/.....	263	-	4	-	9	8	14	22	28	46	74	58	-
2006 1/.....	247	-	1	2	4	9	11	23	33	33	67	64	-
2005 1/.....	241	-	3	1	1	9	10	11	29	34	69	74	-
2004 1/.....	171	-	1	2	5	3	9	18	15	26	46	46	-
2003 1/.....	191	2	1	2	3	3	9	21	13	31	69	37	-
2002.....	178	-	4	-	6	4	12	10	17	34	51	40	-
2001.....	160	1	1	1	3	11	9	15	15	15	40	49	-
2000.....	131	-	-	-	4	6	13	9	10	22	37	30	-
1999.....	151	-	-	1	2	9	12	11	19	19	42	36	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Falls (W00-W19)

Asian or Pacific Islander 3/, female

2007 1/.....	201	-	2	1	-	-	4	6	19	30	71	68	-
2006 1/.....	189	-	1	1	1	1	2	4	18	28	63	70	-
2005 1/.....	158	-	-	-	1	-	3	1	12	29	54	58	-
2004 1/.....	144	-	3	-	-	2	3	5	8	20	56	47	-
2003 1/.....	143	-	2	1	1	1	4	3	7	23	51	50	-
2002.....	117	-	2	1	1	2	2	1	6	19	40	43	-
2001.....	92	1	-	1	-	-	1	3	7	12	32	35	-
2000.....	98	-	-	-	2	-	4	3	9	15	28	37	-
1999.....	66	2	-	-	1	1	4	3	3	14	19	19	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental discharge of firearms (W32-W34)													
All races, both sexes													
2007.....	613	1	18	46	155	94	91	82	57	31	31	7	-
2006.....	642	-	13	41	193	113	74	84	49	33	34	8	-
2005.....	789	1	22	52	203	130	114	108	67	44	33	15	-
2004.....	649	1	14	48	172	100	97	106	35	40	28	8	-
2003.....	730	-	7	49	200	120	118	92	62	39	37	6	-
2002.....	762	1	11	48	210	143	123	95	56	36	32	7	-
2001.....	802	-	15	57	206	122	146	103	69	33	41	8	2
2000.....	776	1	18	67	202	131	153	78	50	41	24	11	-
1999.....	824	-	12	76	251	143	136	92	48	33	23	10	-
All races, male													
2007.....	537	1	14	41	147	82	74	71	51	23	27	6	-
2006.....	567	-	11	33	186	100	58	66	45	32	29	7	-
2005.....	683	1	18	39	190	118	94	88	58	37	27	13	-
2004.....	573	-	11	40	160	92	87	90	29	33	23	8	-
2003.....	656	-	7	46	182	110	105	78	59	30	33	6	-
2002.....	667	1	7	39	195	129	98	84	50	32	26	6	-
2001.....	690	-	12	49	193	103	121	87	55	26	36	7	1
2000.....	671	-	11	62	183	109	125	69	44	34	23	11	-
1999.....	707	-	7	60	229	122	112	78	41	31	20	7	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

Accidental discharge of firearms (W32-W34)

All races, female

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
2007.....	76	-	4	5	8	12	17	11	6	8	4	1	-
2006.....	75	-	2	8	7	13	16	18	4	1	5	1	-
2005.....	106	-	4	13	13	12	20	20	9	7	6	2	-
2004.....	76	1	3	8	12	8	10	16	6	7	5	-	-
2003.....	74	-	-	3	18	10	13	14	3	9	4	-	-
2002.....	95	-	4	9	15	14	25	11	6	4	6	1	-
2001.....	112	-	3	8	13	19	25	16	14	7	5	1	1
2000.....	105	1	7	5	19	22	28	9	6	7	1	-	-
1999.....	117	-	5	16	22	21	24	14	7	2	3	3	-

White, both sexes

2007 1/.....	468	-	12	32	99	61	72	73	54	27	31	7	-
2006 1/.....	484	-	4	29	116	80	66	73	46	30	32	8	-
2005 1/.....	601	-	10	37	139	82	92	92	65	41	31	12	-
2004 1/.....	504	1	6	37	117	65	81	99	31	36	24	7	-
2003 1/.....	586	-	4	38	147	83	104	80	59	34	33	4	-
2002.....	591	1	9	36	142	100	102	81	50	35	28	7	-
2001.....	675	-	10	45	152	103	127	98	62	30	38	8	2
2000.....	611	1	11	52	141	94	128	67	47	37	22	11	-
1999.....	649	-	5	57	174	115	116	81	41	30	20	10	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental discharge of firearms (W32-W34)													
White, male													
2007 1/.....	406	-	10	29	95	52	57	62	48	20	27	6	-
2006 1/.....	420	-	3	25	110	69	50	58	42	29	27	7	-
2005 1/.....	516	-	9	29	134	71	73	75	56	34	25	10	-
2004 1/.....	443	-	5	31	109	59	71	86	26	30	19	7	-
2003 1/.....	525	-	4	35	132	75	95	67	57	27	29	4	-
2002.....	510	1	5	28	131	88	80	72	46	31	22	6	-
2001.....	580	-	8	41	141	87	107	83	49	23	33	7	1
2000.....	530	-	6	48	129	75	107	61	41	31	21	11	-
1999.....	554	-	3	46	158	96	94	68	37	28	17	7	-
White, female													
2007 1/.....	62	-	2	3	4	9	15	11	6	7	4	1	-
2006 1/.....	64	-	1	4	6	11	16	15	4	1	5	1	-
2005 1/.....	85	-	1	8	5	11	19	17	9	7	6	2	-
2004 1/.....	61	1	1	6	8	6	10	13	5	6	5	-	-
2003 1/.....	61	-	-	3	15	8	9	13	2	7	4	-	-
2002.....	81	-	4	8	11	12	22	9	4	4	6	1	-
2001.....	95	-	2	4	11	16	20	15	13	7	5	1	1
2000.....	81	1	5	4	12	19	21	6	6	6	1	-	-
1999.....	95	-	2	11	16	19	22	13	4	2	3	3	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Accidental discharge of firearms (W32-W34)

Black, both sexes

2007 1/.....	127	1	5	13	49	31	17	5	3	3	-	-	-
2006 1/.....	140	-	8	9	70	30	6	10	3	3	1	-	-
2005 1/.....	154	1	12	14	50	40	17	11	2	3	1	3	-
2004 1/.....	129	-	7	9	47	33	13	7	4	4	4	1	-
2003 1/.....	128	-	3	9	50	34	12	7	3	5	3	2	-
2002.....	155	-	2	9	61	40	20	13	6	-	4	-	-
2001.....	112	-	5	8	49	17	16	5	6	3	3	-	-
2000.....	137	-	5	12	52	32	19	10	3	3	1	-	-
1999.....	156	-	7	17	67	23	18	11	7	3	3	-	-

Black, male

2007 1/.....	114	1	4	11	45	28	15	5	3	2	-	-	-
2006 1/.....	132	-	7	7	69	28	6	8	3	3	1	-	-
2005 1/.....	138	1	9	9	45	39	17	9	2	3	1	3	-
2004 1/.....	117	-	6	7	45	31	13	4	3	3	4	1	-
2003 1/.....	116	-	3	9	47	33	8	6	2	3	3	2	-
2002.....	142	-	2	8	58	38	17	11	4	-	4	-	-
2001.....	99	-	4	6	47	14	13	4	5	3	3	-	-
2000.....	117	-	3	12	46	29	14	7	3	2	1	-	-
1999.....	135	-	4	12	61	22	16	10	4	3	3	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Accidental discharge of firearms (W32-W34)

Black, female

2007 1/.....	13	-	1	2	4	3	2	-	-	1	-	-	-
2006 1/.....	8	-	1	2	1	2	-	2	-	-	-	-	-
2005 1/.....	16	-	3	5	5	1	-	2	-	-	-	-	-
2004 1/.....	12	-	1	2	2	2	-	3	1	1	-	-	-
2003 1/.....	12	-	-	-	3	1	4	1	1	2	-	-	-
2002.....	13	-	-	1	3	2	3	2	2	-	-	-	-
2001.....	13	-	1	2	2	3	3	1	1	-	-	-	-
2000.....	20	-	2	-	6	3	5	3	-	1	-	-	-
1999.....	21	-	3	5	6	1	2	1	3	-	-	-	-

American Indian 2/, both sexes

2007 1/.....	11	-	1	1	6	-	1	2	-	-	-	-	-
2006 1/.....	7	-	-	2	1	1	1	1	-	-	1	-	-
2005 1/.....	11	-	-	1	4	5	1	-	-	-	-	-	-
2004 1/.....	13	-	1	2	7	1	2	-	-	-	-	-	-
2003 1/.....	13	-	-	2	3	2	2	3	-	-	1	-	-
2002.....	9	-	-	2	3	2	-	1	-	1	-	-	-
2001.....	9	-	-	3	3	1	2	-	-	-	-	-	-
2000.....	14	-	2	2	6	1	1	1	-	1	-	-	-
1999.....	14	-	-	1	9	3	1	-	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Accidental discharge of firearms (W32-W34)

American Indian 2/, male

2007 1/.....	10	-	-	1	6	-	1	2	-	-	-	-	-
2006 1/.....	5	-	-	1	1	1	1	-	-	-	1	-	-
2005 1/.....	9	-	-	1	3	5	-	-	-	-	-	-	-
2004 1/.....	10	-	-	2	5	1	2	-	-	-	-	-	-
2003 1/.....	12	-	-	2	3	1	2	3	-	-	1	-	-
2002.....	9	-	-	2	3	2	-	1	-	1	-	-	-
2001.....	6	-	-	2	3	1	-	-	-	-	-	-	-
2000.....	12	-	2	2	5	1	-	1	-	1	-	-	-
1999.....	13	-	-	1	9	2	1	-	-	-	-	-	-

American Indian 2/, female

2007 1/.....	1	-	1	-	-	-	-	-	-	-	-	-	-
2006 1/.....	2	-	-	1	-	-	-	1	-	-	-	-	-
2005 1/.....	2	-	-	-	1	-	1	-	-	-	-	-	-
2004 1/.....	3	-	1	-	2	-	-	-	-	-	-	-	-
2003 1/.....	1	-	-	-	-	1	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	3	-	-	1	-	-	2	-	-	-	-	-	-
2000.....	2	-	-	-	1	-	1	-	-	-	-	-	-
1999.....	1	-	-	-	-	1	-	-	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Accidental discharge of firearms (W32-W34)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	7	-	-	-	1	2	1	2	-	1	-	-	-
2006 1/.....	11	-	1	1	6	2	1	-	-	-	-	-	-
2005 1/.....	23	-	-	-	10	3	4	5	-	-	1	-	-
2004 1/.....	3	-	-	-	1	1	1	-	-	-	-	-	-
2003 1/.....	3	-	-	-	-	1	-	2	-	-	-	-	-
2002.....	7	-	-	1	4	1	1	-	-	-	-	-	-
2001.....	6	-	-	1	2	1	1	-	1	-	-	-	-
2000.....	14	-	-	1	3	4	5	-	-	-	1	-	-
1999.....	5	-	-	1	1	2	1	-	-	-	-	-	-

Asian or Pacific Islander 3/, male

2007 1/.....	7	-	-	-	1	2	1	2	-	1	-	-	-
2006 1/.....	10	-	1	-	6	2	1	-	-	-	-	-	-
2005 1/.....	20	-	-	-	8	3	4	4	-	-	1	-	-
2004 1/.....	3	-	-	-	1	1	1	-	-	-	-	-	-
2003 1/.....	3	-	-	-	-	1	-	2	-	-	-	-	-
2002.....	6	-	-	1	3	1	1	-	-	-	-	-	-
2001.....	5	-	-	-	2	1	1	-	1	-	-	-	-
2000.....	12	-	-	-	3	4	4	-	-	-	1	-	-
1999.....	5	-	-	1	1	2	1	-	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental drowning and submersion (W65-W74)													
All races, both sexes													
2007.....	3,443	57	458	224	630	381	417	481	324	194	184	88	5
2006.....	3,579	51	458	256	616	436	466	491	310	217	194	73	11
2005.....	3,582	64	493	253	649	385	497	492	266	193	180	92	18
2004.....	3,308	62	430	269	574	385	435	444	271	188	158	78	14
2003.....	3,306	58	456	268	567	356	462	397	265	202	162	83	30
2002.....	3,447	63	454	321	629	433	526	374	212	169	167	79	20
2001.....	3,281	68	458	333	596	374	462	359	206	159	157	75	34
2000.....	3,482	75	493	375	646	419	480	354	217	179	156	64	24
1999.....	3,529	68	490	369	647	446	475	370	219	185	158	68	34
All races, male													
2007.....	2,681	31	310	157	555	330	334	385	254	148	118	54	5
2006.....	2,774	27	289	193	550	393	371	387	235	149	129	41	10
2005.....	2,818	27	336	184	588	328	410	393	218	135	121	62	16
2004.....	2,594	38	265	194	522	332	367	358	208	140	113	43	14
2003.....	2,632	34	301	199	525	323	392	320	211	145	104	48	30
2002.....	2,761	40	312	244	578	380	439	301	161	132	106	50	18
2001.....	2,560	30	291	237	544	328	376	299	155	124	102	45	29
2000.....	2,735	43	316	267	583	364	406	299	179	133	88	34	23
1999.....	2,794	39	310	275	592	397	402	311	170	129	111	28	30

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

Accidental drowning and submersion (W65-W74)

All races, female

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
2007.....	762	26	148	67	75	51	83	96	70	46	66	34	-
2006.....	805	24	169	63	66	43	95	104	75	68	65	32	1
2005.....	764	37	157	69	61	57	87	99	48	58	59	30	2
2004.....	714	24	165	75	52	53	68	86	63	48	45	35	-
2003.....	674	24	155	69	42	33	70	77	54	57	58	35	-
2002.....	686	23	142	77	51	53	87	73	51	37	61	29	2
2001.....	721	38	167	96	52	46	86	60	51	35	55	30	5
2000.....	747	32	177	108	63	55	74	55	38	46	68	30	1
1999.....	735	29	180	94	55	49	73	59	49	56	47	40	4

White, both sexes

2007 1/.....	2,714	39	381	135	461	275	331	406	274	162	165	80	5
2006 1/.....	2,837	45	370	159	449	336	381	408	256	184	174	66	9
2005 1/.....	2,819	56	418	147	470	304	391	404	224	160	147	80	18
2004 1/.....	2,565	51	368	142	407	295	350	362	219	157	134	66	14
2003 1/.....	2,622	44	386	160	419	269	372	326	224	178	145	70	29
2002.....	2,624	49	368	164	455	330	418	291	181	139	143	66	20
2001.....	2,515	52	397	175	424	272	361	295	166	139	138	68	28
2000.....	2,683	47	415	218	460	322	385	277	187	156	137	56	23
1999.....	2,715	46	416	186	462	330	393	317	181	154	138	60	32

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

Accidental drowning and submersion (W65-W74)

White, male

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
2007 1/.....	2,090	19	249	97	404	238	260	324	210	126	107	51	5
2006 1/.....	2,174	23	231	118	390	303	300	321	190	133	118	39	8
2005 1/.....	2,191	21	288	98	426	257	316	320	179	112	102	56	16
2004 1/.....	2,004	29	225	102	369	253	295	292	168	120	99	38	14
2003 1/.....	2,067	24	252	117	389	242	314	257	175	129	96	43	29
2002.....	2,093	30	251	120	417	291	345	235	136	110	94	46	18
2001.....	1,936	25	257	115	384	232	292	244	121	109	91	41	25
2000.....	2,105	28	268	151	413	284	323	232	159	114	81	30	22
1999.....	2,120	26	262	132	419	293	327	265	139	106	97	26	28

White, female

2007 1/.....	624	20	132	38	57	37	71	82	64	36	58	29	-
2006 1/.....	663	22	139	41	59	33	81	87	66	51	56	27	1
2005 1/.....	628	35	130	49	44	47	75	84	45	48	45	24	2
2004 1/.....	561	22	143	40	38	42	55	70	51	37	35	28	-
2003 1/.....	555	20	134	43	30	27	58	69	49	49	49	27	-
2002.....	531	19	117	44	38	39	73	56	45	29	49	20	2
2001.....	579	27	140	60	40	40	69	51	45	30	47	27	3
2000.....	578	19	147	67	47	38	62	45	28	42	56	26	1
1999.....	595	20	154	54	43	37	66	52	42	48	41	34	4

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

Accidental drowning and submersion (W65-W74)

Black, both sexes

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
2007 1/.....	511	16	51	71	129	73	51	53	34	20	7	6	-
2006 1/.....	538	5	69	80	129	65	56	56	39	24	7	6	2
2005 1/.....	556	8	58	88	134	60	72	61	31	25	14	5	-
2004 1/.....	533	9	49	104	118	61	62	64	29	20	10	7	-
2003 1/.....	480	13	54	89	101	61	51	47	34	16	8	5	1
2002.....	613	11	63	136	129	75	74	57	22	18	17	11	-
2001.....	579	12	48	134	134	63	72	51	32	13	13	4	3
2000.....	593	26	56	136	134	58	79	52	17	15	12	7	1
1999.....	634	21	60	159	138	88	60	37	29	21	15	5	1

Black, male

2007 1/.....	419	10	41	46	118	64	44	45	30	15	4	2	-
2006 1/.....	451	4	49	62	124	61	50	48	32	12	5	2	2
2005 1/.....	471	6	36	72	125	54	62	56	29	19	9	3	-
2004 1/.....	426	7	37	75	108	51	53	50	22	15	7	1	-
2003 1/.....	407	9	39	67	95	56	48	44	30	10	5	3	1
2002.....	509	7	44	108	125	65	66	47	19	15	9	4	-
2001.....	478	3	28	103	127	62	60	42	26	12	9	3	3
2000.....	478	14	35	102	126	52	69	45	13	13	4	4	1
1999.....	530	12	40	125	129	80	55	35	23	17	11	2	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Accidental drowning and submersion (W65-W74)

Black, female

2007 1/.....	92	6	10	25	11	9	7	8	4	5	3	4	-
2006 1/.....	87	1	20	18	5	4	6	8	7	12	2	4	-
2005 1/.....	85	2	22	16	9	6	10	5	2	6	5	2	-
2004 1/.....	107	2	12	29	10	10	9	14	7	5	3	6	-
2003 1/.....	73	4	15	22	6	5	3	3	4	6	3	2	-
2002.....	104	4	19	28	4	10	8	10	3	3	8	7	-
2001.....	101	9	20	31	7	1	12	9	6	1	4	1	-
2000.....	115	12	21	34	8	6	10	7	4	2	8	3	-
1999.....	104	9	20	34	9	8	5	2	6	4	4	3	-

American Indian 2/, both sexes

2007 1/.....	67	1	13	3	13	10	12	7	6	2	-	-	-
2006 1/.....	50	1	6	4	8	8	8	9	3	3	-	-	-
2005 1/.....	64	-	5	3	18	9	17	7	-	1	4	-	-
2004 1/.....	57	1	4	9	12	6	10	9	4	1	1	-	-
2003 1/.....	62	1	6	8	10	2	21	11	1	1	1	-	-
2002.....	66	-	8	3	12	12	20	8	2	-	1	-	-
2001.....	52	2	6	9	5	13	9	5	3	-	-	-	-
2000.....	79	2	10	11	18	14	7	11	4	2	-	-	-
1999.....	68	-	7	5	11	15	13	9	4	3	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Accidental drowning and submersion (W65-W74)

American Indian 2/, male

2007 1/.....	57	1	10	3	11	9	11	6	5	1	-	-	-
2006 1/.....	38	-	2	3	8	7	6	7	2	3	-	-	-
2005 1/.....	49	-	4	1	12	7	17	5	-	1	2	-	-
2004 1/.....	46	1	1	7	10	6	9	7	4	1	-	-	-
2003 1/.....	51	1	4	7	9	2	16	9	1	1	1	-	-
2002.....	53	-	6	2	11	11	16	6	1	-	-	-	-
2001.....	46	1	3	9	5	13	7	5	3	-	-	-	-
2000.....	61	1	6	8	15	9	7	9	4	2	-	-	-
1999.....	55	-	4	3	11	11	12	6	4	3	1	-	-

American Indian 2/, female

2007 1/.....	10	-	3	-	2	1	1	1	1	1	-	-	-
2006 1/.....	12	1	4	1	-	1	2	2	1	-	-	-	-
2005 1/.....	15	-	1	2	6	2	-	2	-	-	2	-	-
2004 1/.....	11	-	3	2	2	-	1	2	-	-	1	-	-
2003 1/.....	11	-	2	1	1	-	5	2	-	-	-	-	-
2002.....	13	-	2	1	1	1	4	2	1	-	1	-	-
2001.....	6	1	3	-	-	-	2	-	-	-	-	-	-
2000.....	18	1	4	3	3	5	-	2	-	-	-	-	-
1999.....	13	-	3	2	-	4	1	3	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Accidental drowning and submersion (W65-W74)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	151	1	13	15	27	23	23	15	10	10	12	2	-
2006 1/.....	154	-	13	13	30	27	21	18	12	6	13	1	-
2005 1/.....	143	-	12	15	27	12	17	20	11	7	15	7	-
2004 1/.....	153	1	9	14	37	23	13	9	19	10	13	5	-
2003 1/.....	142	-	10	11	37	24	18	13	6	7	8	8	-
2002.....	144	3	15	18	33	16	14	18	7	12	6	2	-
2001.....	135	2	7	15	33	26	20	8	5	7	6	3	3
2000.....	127	-	12	10	34	25	9	14	9	6	7	1	-
1999.....	112	1	7	19	36	13	9	7	5	7	4	3	1

Asian or Pacific Islander 3/, male

2007 1/.....	115	1	10	11	22	19	19	10	9	6	7	1	-
2006 1/.....	111	-	7	10	28	22	15	11	11	1	6	-	-
2005 1/.....	107	-	8	13	25	10	15	12	10	3	8	3	-
2004 1/.....	118	1	2	10	35	22	10	9	14	4	7	4	-
2003 1/.....	107	-	6	8	32	23	14	10	5	5	2	2	-
2002.....	106	3	11	14	25	13	12	13	5	7	3	-	-
2001.....	100	1	3	10	28	21	17	8	5	3	2	1	1
2000.....	91	-	7	6	29	19	7	13	3	4	3	-	-
1999.....	89	1	4	15	33	13	8	5	4	3	2	-	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Accidental drowning and submersion (W65-W74)

Asian or Pacific Islander 3/, female

2007 1/.....	36	-	3	4	5	4	4	5	1	4	5	1	-
2006 1/.....	43	-	6	3	2	5	6	7	1	5	7	1	-
2005 1/.....	36	-	4	2	2	2	2	8	1	4	7	4	-
2004 1/.....	35	-	7	4	2	1	3	-	5	6	6	1	-
2003 1/.....	35	-	4	3	5	1	4	3	1	2	6	6	-
2002.....	38	-	4	4	8	3	2	5	2	5	3	2	-
2001.....	35	1	4	5	5	5	3	-	-	4	4	2	2
2000.....	36	-	5	4	5	6	2	1	6	2	4	1	-
1999.....	23	-	3	4	3	-	1	2	1	4	2	3	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental exposure to smoke, fire and flames (X00-X09)													
All races, both sexes													
2007.....	3,286	38	201	211	194	222	307	488	492	421	436	272	4
2006.....	3,109	27	199	181	185	199	296	476	461	404	448	228	5
2005.....	3,197	34	204	222	168	224	334	497	398	390	468	254	4
2004.....	3,229	26	223	255	186	240	313	490	412	349	459	273	3
2003.....	3,369	30	224	215	199	252	395	526	392	419	460	253	4
2002.....	3,159	36	221	253	193	266	361	448	340	385	432	221	3
2001.....	3,309	50	225	252	209	247	441	419	378	411	423	251	3
2000.....	3,377	37	290	266	192	241	402	439	369	401	472	263	5
1999.....	3,348	41	302	262	201	282	421	368	319	416	487	245	4
All races, male													
2007.....	1,943	20	130	116	124	137	198	306	330	252	212	115	3
2006.....	1,819	14	117	87	117	132	185	300	292	227	238	106	4
2005.....	1,886	16	124	125	103	137	226	325	244	224	245	113	4
2004.....	1,902	14	120	152	107	165	209	329	251	210	234	109	2
2003.....	1,948	8	131	112	121	154	257	329	254	244	228	107	3
2002.....	1,935	21	126	151	132	170	251	292	219	242	231	97	3
2001.....	1,998	30	142	152	137	161	308	288	236	229	204	108	3
2000.....	2,025	20	174	145	121	157	268	303	241	232	244	115	5
1999.....	2,025	21	173	147	133	197	300	254	194	244	247	111	4

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental exposure to smoke, fire and flames (X00-X09)													
All races, female													
2007.....	1,343	18	71	95	70	85	109	182	162	169	224	157	1
2006.....	1,290	13	82	94	68	67	111	176	169	177	210	122	1
2005.....	1,311	18	80	97	65	87	108	172	154	166	223	141	-
2004.....	1,327	12	103	103	79	75	104	161	161	139	225	164	1
2003.....	1,421	22	93	103	78	98	138	197	138	175	232	146	1
2002.....	1,224	15	95	102	61	96	110	156	121	143	201	124	-
2001.....	1,311	20	83	100	72	86	133	131	142	182	219	143	-
2000.....	1,352	17	116	121	71	84	134	136	128	169	228	148	-
1999.....	1,323	20	129	115	68	85	121	114	125	172	240	134	-
White, both sexes													
2007 1/.....	2,420	23	108	114	153	163	220	375	391	320	347	203	3
2006 1/.....	2,349	18	118	110	143	146	227	370	366	322	347	178	4
2005 1/.....	2,351	23	123	125	123	161	258	390	281	303	366	194	4
2004 1/.....	2,334	15	139	136	140	179	234	356	307	254	356	216	2
2003 1/.....	2,526	22	144	115	156	195	317	395	303	320	356	199	4
2002.....	2,287	19	127	139	138	190	285	341	247	289	342	167	3
2001.....	2,387	34	142	159	157	174	326	303	277	301	321	190	3
2000.....	2,376	23	160	158	141	169	291	330	260	297	345	197	5
1999.....	2,380	26	178	159	153	215	310	272	235	293	360	176	3

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental exposure to smoke, fire and flames (X00-X09)													
White, male													
2007 1/.....	1,426	11	76	60	100	103	138	236	258	185	171	85	3
2006 1/.....	1,389	8	77	54	94	96	140	231	230	180	185	91	3
2005 1/.....	1,392	10	81	75	77	104	171	253	166	171	194	86	4
2004 1/.....	1,375	9	76	80	82	124	153	234	190	153	185	88	1
2003 1/.....	1,480	7	85	58	100	120	214	253	196	189	168	87	3
2002.....	1,419	14	70	90	97	126	194	222	161	179	190	73	3
2001.....	1,436	19	95	92	106	120	228	202	172	163	154	82	3
2000.....	1,435	11	98	85	90	113	195	228	169	169	181	91	5
1999.....	1,466	14	103	88	103	156	223	188	149	169	184	86	3
White, female													
2007 1/.....	994	12	32	54	53	60	82	139	133	135	176	118	-
2006 1/.....	960	10	41	56	49	50	87	139	136	142	162	87	1
2005 1/.....	959	13	42	50	46	57	87	137	115	132	172	108	-
2004 1/.....	959	6	63	56	58	55	81	122	117	101	171	128	1
2003 1/.....	1,046	15	59	57	56	75	103	142	107	131	188	112	1
2002.....	868	5	57	49	41	64	91	119	86	110	152	94	-
2001.....	951	15	47	67	51	54	98	101	105	138	167	108	-
2000.....	941	12	62	73	51	56	96	102	91	128	164	106	-
1999.....	914	12	75	71	50	59	87	84	86	124	176	90	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental exposure to smoke, fire and flames (X00-X09)													
Black, both sexes													
2007 1/.....	772	15	81	87	35	53	74	101	91	91	81	62	1
2006 1/.....	699	7	78	67	35	49	60	98	86	76	96	47	-
2005 1/.....	746	10	73	82	36	52	64	98	107	76	94	54	-
2004 1/.....	799	11	73	105	39	53	66	120	98	84	96	53	1
2003 1/.....	775	7	76	94	39	47	68	124	79	90	100	51	-
2002.....	774	15	88	96	48	61	60	98	87	85	83	53	-
2001.....	846	15	82	88	42	66	104	107	90	102	92	58	-
2000.....	930	14	121	101	44	65	99	100	102	98	121	65	-
1999.....	892	15	111	102	40	53	102	87	80	113	120	68	1
Black, male													
2007 1/.....	459	9	46	49	19	30	51	64	65	61	38	27	-
2006 1/.....	389	5	40	30	17	33	38	61	56	43	51	15	-
2005 1/.....	442	6	38	44	20	27	45	70	73	46	49	24	-
2004 1/.....	475	5	38	64	22	35	45	87	57	54	48	19	1
2003 1/.....	433	1	42	50	19	28	39	73	53	53	57	18	-
2002.....	461	7	53	54	31	35	45	64	55	55	38	24	-
2001.....	511	10	46	56	24	35	73	81	56	62	44	24	-
2000.....	551	9	71	57	27	39	65	69	69	60	61	24	-
1999.....	510	7	63	58	24	30	71	59	44	69	59	25	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental exposure to smoke, fire and flames (X00-X09)													
Black, female													
2007 1/.....	313	6	35	38	16	23	23	37	26	30	43	35	1
2006 1/.....	310	2	38	37	18	16	22	37	30	33	45	32	-
2005 1/.....	304	4	35	38	16	25	19	28	34	30	45	30	-
2004 1/.....	324	6	35	41	17	18	21	33	41	30	48	34	-
2003 1/.....	342	6	34	44	20	19	29	51	26	37	43	33	-
2002.....	313	8	35	42	17	26	15	34	32	30	45	29	-
2001.....	335	5	36	32	18	31	31	26	34	40	48	34	-
2000.....	379	5	50	44	17	26	34	31	33	38	60	41	-
1999.....	382	8	48	44	16	23	31	28	36	44	61	43	-
American Indian 2/, both sexes													
2007 1/.....	42	-	4	4	2	3	7	6	4	7	4	1	-
2006 1/.....	28	1	2	1	4	3	2	4	4	4	1	2	-
2005 1/.....	54	-	6	5	6	8	7	5	6	7	2	2	-
2004 1/.....	47	-	7	5	3	3	7	7	5	5	4	1	-
2003 1/.....	33	-	4	3	2	5	7	3	4	5	-	-	-
2002.....	56	1	3	8	3	11	11	6	3	8	1	1	-
2001.....	45	1	1	4	5	4	9	6	7	5	2	1	-
2000.....	37	-	5	4	4	7	7	2	5	2	1	-	-
1999.....	40	-	5	-	4	7	4	5	3	6	6	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Accidental exposure to smoke, fire and flames (X00-X09)

American Indian 2/, male

2007 1/.....	24	-	2	3	2	2	4	2	2	4	3	-	-
2006 1/.....	22	1	-	1	4	3	2	4	4	2	1	-	-
2005 1/.....	33	-	4	3	4	6	6	1	3	4	1	1	-
2004 1/.....	28	-	5	3	2	2	5	5	3	2	1	-	-
2003 1/.....	16	-	4	2	1	2	4	1	1	1	-	-	-
2002.....	35	-	1	3	1	8	8	4	2	7	1	-	-
2001.....	30	1	1	3	3	3	5	3	6	3	1	1	-
2000.....	23	-	3	2	3	5	4	2	2	1	1	-	-
1999.....	26	-	3	-	4	5	2	4	-	5	3	-	-

American Indian 2/, female

2007 1/.....	18	-	2	1	-	1	3	4	2	3	1	1	-
2006 1/.....	6	-	2	-	-	-	-	-	-	2	-	2	-
2005 1/.....	21	-	2	2	2	2	1	4	3	3	1	1	-
2004 1/.....	19	-	2	2	1	1	2	2	2	3	3	1	-
2003 1/.....	17	-	-	1	1	3	3	2	3	4	-	-	-
2002.....	21	1	2	5	2	3	3	2	1	1	-	1	-
2001.....	15	-	-	1	2	1	4	3	1	2	1	-	-
2000.....	14	-	2	2	1	2	3	-	3	1	-	-	-
1999.....	14	-	2	-	-	2	2	1	3	1	3	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Accidental exposure to smoke, fire and flames (X00-X09)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	52	-	8	6	4	3	6	6	6	3	4	6	-
2006 1/.....	33	1	1	3	3	1	7	4	5	2	4	1	1
2005 1/.....	46	1	2	10	3	3	5	4	4	4	6	4	-
2004 1/.....	49	-	4	9	4	5	6	7	2	6	3	3	-
2003 1/.....	35	1	-	3	2	5	3	4	6	4	4	3	-
2002.....	42	1	3	10	4	4	5	3	3	3	6	-	-
2001.....	31	-	-	1	5	3	2	3	4	3	8	2	-
2000.....	34	-	4	3	3	-	5	7	2	4	5	1	-
1999.....	36	-	8	1	4	7	5	4	1	4	1	1	-

Asian or Pacific Islander 3/, male

2007 1/.....	34	-	6	4	3	2	5	4	5	2	-	3	-
2006 1/.....	19	-	-	2	2	-	5	4	2	2	1	-	1
2005 1/.....	19	-	1	3	2	-	4	1	2	3	1	2	-
2004 1/.....	24	-	1	5	1	4	6	3	1	1	-	2	-
2003 1/.....	19	-	-	2	1	4	-	2	4	1	3	2	-
2002.....	20	-	2	4	3	1	4	2	1	1	2	-	-
2001.....	21	-	-	1	4	3	2	2	2	1	5	1	-
2000.....	16	-	2	1	1	-	4	4	1	2	1	-	-
1999.....	23	-	4	1	2	6	4	3	1	1	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Accidental exposure to smoke, fire and flames (X00-X09)

Asian or Pacific Islander 3/, female

2007 1/.....	18	-	2	2	1	1	1	2	1	1	4	3	-
2006 1/.....	14	1	1	1	1	1	2	-	3	-	3	1	-
2005 1/.....	27	1	1	7	1	3	1	3	2	1	5	2	-
2004 1/.....	25	-	3	4	3	1	-	4	1	5	3	1	-
2003 1/.....	16	1	-	1	1	1	3	2	2	3	1	1	-
2002.....	22	1	1	6	1	3	1	1	2	2	4	-	-
2001.....	10	-	-	-	1	-	-	1	2	2	3	1	-
2000.....	18	-	2	2	2	-	1	3	1	2	4	1	-
1999.....	13	-	4	-	2	1	1	1	-	3	-	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental poisoning and exposure to noxious substances (X40-X49)													
All races, both sexes													
2007.....	29,846	19	34	81	3,159	5,700	7,575	9,006	3,120	602	355	192	3
2006.....	27,531	16	27	58	2,936	5,267	7,542	8,234	2,415	483	340	202	11
2005.....	23,618	20	21	51	2,484	4,386	6,729	6,983	2,007	435	317	179	6
2004.....	20,950	13	18	55	2,259	3,641	6,444	6,033	1,577	421	315	165	9
2003.....	19,457	20	49	61	1,999	3,435	6,230	5,434	1,370	403	284	166	6
2002.....	17,550	26	31	43	1,679	3,116	6,007	4,682	1,098	383	295	181	9
2001.....	14,078	15	31	50	1,362	2,507	5,036	3,547	798	291	279	152	10
2000.....	12,757	14	32	45	1,160	2,380	4,663	3,061	688	278	245	184	7
1999.....	12,186	12	34	40	964	2,355	4,549	2,844	668	300	265	143	12
All races, male													
2007.....	19,644	13	17	50	2,430	4,171	4,947	5,535	1,905	354	154	65	3
2006.....	18,581	7	16	35	2,319	3,894	5,020	5,291	1,512	249	152	78	8
2005.....	15,884	11	12	31	1,940	3,193	4,423	4,537	1,269	228	161	74	5
2004.....	13,934	5	13	30	1,765	2,673	4,176	3,910	929	222	143	64	4
2003.....	13,176	9	27	31	1,562	2,514	4,166	3,598	841	223	142	58	5
2002.....	12,059	21	14	28	1,306	2,319	4,122	3,136	690	209	136	71	7
2001.....	9,885	8	21	23	1,080	1,862	3,522	2,503	500	170	137	49	10
2000.....	9,138	11	21	27	920	1,820	3,384	2,223	401	147	102	76	6
1999.....	8,887	11	17	25	754	1,828	3,349	2,106	448	172	116	53	8

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental poisoning and exposure to noxious substances (X40-X49)													
All races, female													
2007.....	10,202	6	17	31	729	1,529	2,628	3,471	1,215	248	201	127	-
2006.....	8,950	9	11	23	617	1,373	2,522	2,943	903	234	188	124	3
2005.....	7,734	9	9	20	544	1,193	2,306	2,446	738	207	156	105	1
2004.....	7,016	8	5	25	494	968	2,268	2,123	648	199	172	101	5
2003.....	6,281	11	22	30	437	921	2,064	1,836	529	180	142	108	1
2002.....	5,491	5	17	15	373	797	1,885	1,546	408	174	159	110	2
2001.....	4,193	7	10	27	282	645	1,514	1,044	298	121	142	103	-
2000.....	3,619	3	11	18	240	560	1,279	838	287	131	143	108	1
1999.....	3,299	1	17	15	210	527	1,200	738	220	128	149	90	4
White, both sexes													
2007 1/.....	25,973	8	17	59	2,928	5,185	6,650	7,609	2,525	499	315	176	2
2006 1/.....	23,476	9	19	42	2,742	4,714	6,405	6,762	1,909	381	303	179	11
2005 1/.....	20,163	12	11	38	2,313	3,878	5,788	5,755	1,579	344	277	163	5
2004 1/.....	18,042	10	12	40	2,095	3,259	5,581	5,015	1,252	348	273	151	6
2003 1/.....	16,690	12	31	35	1,880	3,038	5,336	4,490	1,137	316	263	148	4
2002.....	14,868	13	19	37	1,555	2,745	5,100	3,782	855	327	264	164	7
2001.....	11,662	9	21	40	1,228	2,176	4,197	2,753	616	245	237	131	9
2000.....	10,508	10	19	30	1,050	2,031	3,842	2,358	543	233	229	157	6
1999.....	9,883	10	20	27	843	1,965	3,713	2,178	510	240	245	123	9

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental poisoning and exposure to noxious substances (X40-X49)													
White, male													
2007 1/.....	17,135	6	12	35	2,272	3,819	4,370	4,649	1,491	281	136	62	2
2006 1/.....	15,784	5	12	24	2,177	3,478	4,272	4,289	1,130	181	136	72	8
2005 1/.....	13,561	7	5	25	1,823	2,844	3,841	3,700	943	162	136	70	5
2004 1/.....	12,022	3	7	25	1,645	2,413	3,648	3,235	692	173	119	59	3
2003 1/.....	11,333	8	16	20	1,472	2,248	3,594	2,958	672	163	131	48	3
2002.....	10,213	12	6	25	1,212	2,052	3,527	2,520	500	170	118	66	5
2001.....	8,232	5	15	19	980	1,638	2,984	1,925	359	142	113	43	9
2000.....	7,558	8	11	18	852	1,568	2,828	1,701	292	115	95	65	5
1999.....	7,240	9	9	17	662	1,551	2,777	1,594	321	138	107	49	6
White, female													
2007 1/.....	8,838	2	5	24	656	1,366	2,280	2,960	1,034	218	179	114	-
2006 1/.....	7,692	4	7	18	565	1,236	2,133	2,473	779	200	167	107	3
2005 1/.....	6,602	5	6	13	490	1,034	1,947	2,055	636	182	141	93	-
2004 1/.....	6,020	7	5	15	450	846	1,933	1,780	560	175	154	92	3
2003 1/.....	5,357	4	15	15	408	790	1,742	1,532	465	153	132	100	1
2002.....	4,655	1	13	12	343	693	1,573	1,262	355	157	146	98	2
2001.....	3,430	4	6	21	248	538	1,213	828	257	103	124	88	-
2000.....	2,950	2	8	12	198	463	1,014	657	251	118	134	92	1
1999.....	2,643	1	11	10	181	414	936	584	189	102	138	74	3

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Accidental poisoning and exposure to noxious substances (X40-X49)

Black, both sexes

2007 1/.....	3,273	8	15	15	152	389	772	1,249	531	96	32	13	1
2006 1/.....	3,526	6	8	7	143	452	995	1,322	462	82	28	21	-
2005 1/.....	2,986	7	7	9	118	401	820	1,107	397	78	30	11	1
2004 1/.....	2,498	3	6	11	119	296	736	929	284	64	35	13	2
2003 1/.....	2,396	7	16	18	87	312	777	861	209	70	20	17	2
2002.....	2,365	12	11	6	88	306	801	833	223	45	27	12	1
2001.....	2,169	6	8	6	95	282	763	747	165	41	34	21	1
2000.....	2,028	4	12	9	79	303	746	655	135	43	15	26	1
1999.....	2,062	1	12	8	94	336	750	628	144	50	18	18	3

Black, male

2007 1/.....	2,150	4	5	11	104	272	483	804	380	69	14	3	1
2006 1/.....	2,439	1	4	6	101	342	651	906	353	57	13	5	-
2005 1/.....	2,015	4	6	4	80	275	505	756	303	60	19	3	-
2004 1/.....	1,638	2	6	3	87	202	441	617	209	45	20	5	1
2003 1/.....	1,609	1	9	7	64	208	504	582	158	54	10	10	2
2002.....	1,631	8	8	3	67	220	526	570	176	35	16	1	1
2001.....	1,490	3	5	2	69	193	487	549	128	27	20	6	1
2000.....	1,433	3	9	7	50	216	507	487	104	31	7	11	1
1999.....	1,494	1	7	6	77	237	519	485	117	30	9	4	2

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental poisoning and exposure to noxious substances (X40-X49)													
Black, female													
2007 1/.....	1,123	4	10	4	48	117	289	445	151	27	18	10	-
2006 1/.....	1,087	5	4	1	42	110	344	416	109	25	15	16	-
2005 1/.....	971	3	1	5	38	126	315	351	94	18	11	8	1
2004 1/.....	860	1	-	8	32	94	295	312	75	19	15	8	1
2003 1/.....	787	6	7	11	23	104	273	279	51	16	10	7	-
2002.....	734	4	3	3	21	86	275	263	47	10	11	11	-
2001.....	679	3	3	4	26	89	276	198	37	14	14	15	-
2000.....	595	1	3	2	29	87	239	168	31	12	8	15	-
1999.....	568	-	5	2	17	99	231	143	27	20	9	14	1
American Indian 2/, both sexes													
2007 1/.....	375	2	2	3	51	74	95	102	37	4	3	2	-
2006 1/.....	324	1	-	4	26	68	87	100	26	10	2	-	-
2005 1/.....	289	-	2	2	32	67	79	80	17	7	2	1	-
2004 1/.....	262	-	-	4	28	59	90	56	18	5	1	-	1
2003 1/.....	231	-	-	3	20	49	85	51	14	9	-	-	-
2002.....	188	1	-	-	16	38	71	45	14	1	-	1	1
2001.....	149	-	1	3	18	31	49	32	11	3	1	-	-
2000.....	137	-	-	4	18	30	49	31	3	1	-	1	-
1999.....	149	1	2	4	15	37	56	23	6	4	-	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Accidental poisoning and exposure to noxious substances (X40-X49)

American Indian 2/, male

2007 1/.....	213	2	-	3	34	45	58	50	15	4	2	-	-
2006 1/.....	220	1	-	2	22	53	55	63	17	7	-	-	-
2005 1/.....	182	-	1	2	21	48	45	50	12	2	1	-	-
2004 1/.....	162	-	-	2	20	35	58	34	11	2	-	-	-
2003 1/.....	141	-	-	2	16	29	48	35	6	5	-	-	-
2002.....	124	1	-	-	13	26	41	31	10	1	-	-	1
2001.....	90	-	-	1	13	16	31	19	8	1	1	-	-
2000.....	84	-	-	-	7	24	27	24	2	-	-	-	-
1999.....	91	1	1	1	10	26	29	15	6	2	-	-	-

American Indian 2/, female

2007 1/.....	162	-	2	-	17	29	37	52	22	-	1	2	-
2006 1/.....	104	-	-	2	4	15	32	37	9	3	2	-	-
2005 1/.....	107	-	1	-	11	19	34	30	5	5	1	1	-
2004 1/.....	100	-	-	2	8	24	32	22	7	3	1	-	1
2003 1/.....	90	-	-	1	4	20	37	16	8	4	-	-	-
2002.....	64	-	-	-	3	12	30	14	4	-	-	1	-
2001.....	59	-	1	2	5	15	18	13	3	2	-	-	-
2000.....	53	-	-	4	11	6	22	7	1	1	-	1	-
1999.....	58	-	1	3	5	11	27	8	-	2	-	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Accidental poisoning and exposure to noxious substances (X40-X49)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	225	1	-	4	28	52	58	46	27	3	5	1	-
2006 1/.....	205	-	-	5	25	33	55	50	18	10	7	2	-
2005 1/.....	180	1	1	2	21	40	42	41	14	6	8	4	-
2004 1/.....	148	-	-	-	17	27	37	33	23	4	6	1	-
2003 1/.....	140	1	2	5	12	36	32	32	10	8	1	1	-
2002.....	129	-	1	-	20	27	35	22	6	10	4	4	-
2001.....	98	-	1	1	21	18	27	15	6	2	7	-	-
2000.....	84	-	1	2	13	16	26	17	7	1	1	-	-
1999.....	92	-	-	1	12	17	30	15	8	6	2	1	-

Asian or Pacific Islander 3/, male

2007 1/.....	146	1	-	1	20	35	36	32	19	-	2	-	-
2006 1/.....	138	-	-	3	19	21	42	33	12	4	3	1	-
2005 1/.....	126	-	-	-	16	26	32	31	11	4	5	1	-
2004 1/.....	112	-	-	-	13	23	29	24	17	2	4	-	-
2003 1/.....	93	-	2	2	10	29	20	23	5	1	1	-	-
2002.....	91	-	-	-	14	21	28	15	4	3	2	4	-
2001.....	73	-	1	1	18	15	20	10	5	-	3	-	-
2000.....	63	-	1	2	11	12	22	11	3	1	-	-	-
1999.....	62	-	-	1	5	14	24	12	4	2	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Accidental poisoning and exposure to noxious substances (X40-X49)

Asian or Pacific Islander 3/, female

2007 1/.....	79	-	-	3	8	17	22	14	8	3	3	1	-
2006 1/.....	67	-	-	2	6	12	13	17	6	6	4	1	-
2005 1/.....	54	1	1	2	5	14	10	10	3	2	3	3	-
2004 1/.....	36	-	-	-	4	4	8	9	6	2	2	1	-
2003 1/.....	47	1	-	3	2	7	12	9	5	7	-	1	-
2002.....	38	-	1	-	6	6	7	7	2	7	2	-	-
2001.....	25	-	-	-	3	3	7	5	1	2	4	-	-
2000.....	21	-	-	-	2	4	4	6	4	-	1	-	-
1999.....	30	-	-	-	7	3	6	3	4	4	2	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

All races, both sexes

2007.....	17,043	1,019	260	226	598	794	1,119	1,755	1,623	1,717	3,195	4,730	7
2006.....	17,503	888	260	231	665	868	1,279	1,836	1,633	1,721	3,383	4,729	10
2005.....	18,526	801	244	258	725	939	1,343	1,977	1,746	1,898	3,731	4,848	16
2004.....	17,016	782	250	247	640	825	1,276	1,732	1,453	1,634	3,505	4,641	31
2003.....	17,115	675	278	215	645	813	1,331	1,591	1,420	1,760	3,539	4,833	15
2002.....	17,201	677	247	273	638	820	1,341	1,496	1,385	1,795	3,665	4,851	13
2001.....	17,760	669	275	255	643	795	1,386	1,641	1,342	1,826	3,841	5,063	24
2000.....	17,437	572	266	267	648	895	1,446	1,510	1,349	1,824	3,678	4,966	16
1999.....	18,388	523	293	316	662	887	1,452	1,496	1,332	2,032	4,025	5,355	15

All races, male

2007.....	9,991	578	164	146	509	666	909	1,390	1,174	1,096	1,628	1,725	6
2006.....	10,327	533	139	165	568	724	1,054	1,445	1,173	1,098	1,741	1,678	9
2005.....	10,864	443	148	180	600	793	1,081	1,554	1,260	1,178	1,897	1,718	12
2004.....	9,880	453	165	184	550	694	1,009	1,347	1,044	995	1,754	1,655	30
2003.....	9,919	398	177	160	544	685	1,061	1,206	1,045	1,095	1,798	1,737	13
2002.....	9,839	389	156	185	545	692	1,081	1,180	1,013	1,113	1,827	1,648	10
2001.....	10,044	390	162	181	535	682	1,128	1,261	933	1,097	1,913	1,741	21
2000.....	9,960	327	162	192	560	767	1,205	1,219	971	1,081	1,801	1,659	16
1999.....	10,371	314	183	221	565	748	1,181	1,176	951	1,227	1,969	1,822	14

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)													
All races, female													
2007.....	7,052	441	96	80	89	128	210	365	449	621	1,567	3,005	1
2006.....	7,176	355	121	66	97	144	225	391	460	623	1,642	3,051	1
2005.....	7,662	358	96	78	125	146	262	423	486	720	1,834	3,130	4
2004.....	7,136	329	85	63	90	131	267	385	409	639	1,751	2,986	1
2003.....	7,196	277	101	55	101	128	270	385	375	665	1,741	3,096	2
2002.....	7,362	288	91	88	93	128	260	316	372	682	1,838	3,203	3
2001.....	7,716	279	113	74	108	113	258	380	409	729	1,928	3,322	3
2000.....	7,477	245	104	75	88	128	241	291	378	743	1,877	3,307	-
1999.....	8,017	209	110	95	97	139	271	320	381	805	2,056	3,533	1
White, both sexes													
2007 1/.....	14,571	620	202	178	517	664	887	1,417	1,335	1,442	2,879	4,425	5
2006 1/.....	15,090	541	192	180	561	726	1,057	1,497	1,377	1,447	3,068	4,434	10
2005 1/.....	15,712	530	189	204	584	787	1,097	1,559	1,422	1,577	3,288	4,462	13
2004 1/.....	14,783	507	185	191	550	696	1,074	1,465	1,225	1,384	3,148	4,330	28
2003 1/.....	14,820	425	210	170	540	681	1,103	1,280	1,174	1,501	3,197	4,525	14
2002.....	14,855	405	175	209	548	705	1,114	1,240	1,150	1,491	3,289	4,516	13
2001.....	15,354	422	200	207	555	652	1,144	1,341	1,110	1,553	3,456	4,692	22
2000.....	15,074	362	194	212	535	748	1,195	1,216	1,117	1,536	3,320	4,625	14
1999.....	15,859	335	215	257	567	726	1,222	1,192	1,059	1,707	3,581	4,985	13

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)													
White, male													
2007 1/.....	8,495	361	127	119	450	564	720	1,132	991	926	1,480	1,620	5
2006 1/.....	8,818	321	104	134	486	620	874	1,173	997	936	1,591	1,573	9
2005 1/.....	9,126	291	120	145	493	674	900	1,236	1,014	986	1,668	1,589	10
2004 1/.....	8,517	308	129	139	477	587	852	1,142	885	855	1,579	1,537	27
2003 1/.....	8,483	251	136	127	467	581	887	980	862	943	1,618	1,619	12
2002.....	8,415	245	109	141	475	600	914	992	840	929	1,636	1,524	10
2001.....	8,571	245	120	143	465	561	948	1,034	774	940	1,711	1,611	19
2000.....	8,437	213	122	155	467	649	1,004	978	794	915	1,600	1,526	14
1999.....	8,781	197	132	183	483	623	997	938	763	1,026	1,752	1,675	12
White, female													
2007 1/.....	6,076	259	75	59	67	100	167	285	344	516	1,399	2,805	-
2006 1/.....	6,272	220	88	46	75	106	183	324	380	511	1,477	2,861	1
2005 1/.....	6,586	239	69	59	91	113	197	323	408	591	1,620	2,873	3
2004 1/.....	6,266	199	56	52	73	109	222	323	340	529	1,569	2,793	1
2003 1/.....	6,337	174	74	43	73	100	216	300	312	558	1,579	2,906	2
2002.....	6,440	160	66	68	73	105	200	248	310	562	1,653	2,992	3
2001.....	6,783	177	80	64	90	91	196	307	336	613	1,745	3,081	3
2000.....	6,637	149	72	57	68	99	191	238	323	621	1,720	3,099	-
1999.....	7,078	138	83	74	84	103	225	254	296	681	1,829	3,310	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)													
Black, both sexes													
2007 1/.....	2,002	356	48	32	54	98	178	273	240	234	244	244	1
2006 1/.....	1,965	304	59	44	71	104	173	279	214	228	251	238	-
2005 1/.....	2,341	248	46	41	103	114	191	352	265	273	383	322	3
2004 1/.....	1,844	245	48	41	64	107	164	229	189	203	304	249	1
2003 1/.....	1,881	230	52	32	82	98	186	259	201	218	271	251	1
2002.....	1,972	253	61	56	69	93	182	217	193	257	311	280	-
2001.....	1,996	223	63	38	62	102	188	251	200	235	319	313	2
2000.....	1,990	188	57	49	90	120	201	239	200	247	309	288	2
1999.....	2,205	169	63	47	75	133	195	267	239	295	389	331	2
Black, male													
2007 1/.....	1,188	185	29	19	36	75	143	208	160	140	114	79	-
2006 1/.....	1,213	185	29	28	54	75	140	221	144	135	118	84	-
2005 1/.....	1,424	139	25	26	80	83	139	267	203	157	198	105	2
2004 1/.....	1,115	129	23	34	50	88	127	172	133	118	152	88	1
2003 1/.....	1,164	136	36	23	59	77	139	188	146	130	141	88	1
2002.....	1,181	134	42	39	53	74	129	159	142	151	159	99	-
2001.....	1,198	135	36	31	47	82	140	186	135	137	166	101	2
2000.....	1,268	100	32	34	73	95	162	197	151	139	175	108	2
1999.....	1,378	102	43	32	68	103	154	207	167	183	186	131	2

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Black, female

2007 1/.....	814	171	19	13	18	23	35	65	80	94	130	165	1
2006 1/.....	752	119	30	16	17	29	33	58	70	93	133	154	-
2005 1/.....	917	109	21	15	23	31	52	85	62	116	185	217	1
2004 1/.....	729	116	25	7	14	19	37	57	56	85	152	161	-
2003 1/.....	717	94	16	9	23	21	47	71	55	88	130	163	-
2002.....	791	119	19	17	16	19	53	58	51	106	152	181	-
2001.....	798	88	27	7	15	20	48	65	65	98	153	212	-
2000.....	722	88	25	15	17	25	39	42	49	108	134	180	-
1999.....	827	67	20	15	7	30	41	60	72	112	203	200	-

American Indian 2/, both sexes

2007 1/.....	224	24	2	11	14	19	38	39	28	17	17	15	-
2006 1/.....	239	24	6	3	26	28	36	40	25	20	19	12	-
2005 1/.....	216	13	4	5	21	26	35	35	36	17	16	8	-
2004 1/.....	180	15	9	6	14	11	27	23	21	18	19	15	2
2003 1/.....	199	12	5	6	13	23	28	29	28	17	23	15	-
2002.....	169	8	6	6	12	14	35	21	19	23	16	9	-
2001.....	188	18	5	5	12	16	36	29	15	14	23	15	-
2000.....	176	10	6	3	11	11	33	40	20	17	19	6	-
1999.....	165	8	7	6	11	19	23	25	21	15	21	9	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

American Indian 2/, male

2007 1/.....	160	19	2	6	14	15	32	31	15	11	10	5	-
2006 1/.....	177	16	3	1	22	22	32	36	20	12	12	1	-
2005 1/.....	158	6	1	3	16	24	26	29	28	13	10	2	-
2004 1/.....	126	8	8	3	13	9	21	21	16	10	8	7	2
2003 1/.....	145	8	3	4	10	17	25	23	24	13	11	7	-
2002.....	125	5	3	4	11	12	30	16	13	18	8	5	-
2001.....	127	7	3	4	12	16	26	25	9	8	11	6	-
2000.....	126	7	4	2	9	8	26	33	16	11	8	2	-
1999.....	116	5	4	2	7	15	20	23	15	11	10	4	-

American Indian 2/, female

2007 1/.....	64	5	-	5	-	4	6	8	13	6	7	10	-
2006 1/.....	62	8	3	2	4	6	4	4	5	8	7	11	-
2005 1/.....	58	7	3	2	5	2	9	6	8	4	6	6	-
2004 1/.....	54	7	1	3	1	2	6	2	5	8	11	8	-
2003 1/.....	54	4	2	2	3	6	3	6	4	4	12	8	-
2002.....	44	3	3	2	1	2	5	5	6	5	8	4	-
2001.....	61	11	2	1	-	-	10	4	6	6	12	9	-
2000.....	50	3	2	1	2	3	7	7	4	6	11	4	-
1999.....	49	3	3	4	4	4	3	2	6	4	11	5	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	246	19	8	5	13	13	16	26	20	24	55	46	1
2006 1/.....	209	19	3	4	7	10	13	20	17	26	45	45	-
2005 1/.....	257	10	5	8	17	12	20	31	23	31	44	56	-
2004 1/.....	209	15	8	9	12	11	11	15	18	29	34	47	-
2003 1/.....	215	8	11	7	10	11	14	23	17	24	48	42	-
2002.....	205	11	5	2	9	8	10	18	23	24	49	46	-
2001.....	222	6	7	5	14	25	18	20	17	24	43	43	-
2000.....	197	12	9	3	12	16	17	15	12	24	30	47	-
1999.....	159	11	8	6	9	9	12	12	13	15	34	30	-

Asian or Pacific Islander 3/, male

2007 1/.....	148	13	6	2	9	12	14	19	8	19	24	21	1
2006 1/.....	119	11	3	2	6	7	8	15	12	15	20	20	-
2005 1/.....	156	7	2	6	11	12	16	22	15	22	21	22	-
2004 1/.....	122	8	5	8	10	10	9	12	10	12	15	23	-
2003 1/.....	127	3	2	6	8	10	10	15	13	9	28	23	-
2002.....	118	5	2	1	6	6	8	13	18	15	24	20	-
2001.....	148	3	3	3	11	23	14	16	15	12	25	23	-
2000.....	129	7	4	1	11	15	13	11	10	16	18	23	-
1999.....	96	10	4	4	7	7	10	8	6	7	21	12	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Asian or Pacific Islander 3/, female

2007 1/.....	98	6	2	3	4	1	2	7	12	5	31	25	-
2006 1/.....	90	8	-	2	1	3	5	5	5	11	25	25	-
2005 1/.....	101	3	3	2	6	-	4	9	8	9	23	34	-
2004 1/.....	87	7	3	1	2	1	2	3	8	17	19	24	-
2003 1/.....	88	5	9	1	2	1	4	8	4	15	20	19	-
2002.....	87	6	3	1	3	2	2	5	5	9	25	26	-
2001.....	74	3	4	2	3	2	4	4	2	12	18	20	-
2000.....	68	5	5	2	1	1	4	4	2	8	12	24	-
1999.....	63	1	4	2	2	2	2	4	7	8	13	18	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

All races, both sexes

2007.....	34,598	184	4,140	5,278	6,722	7,778	5,069	2,444	2,119	858	6
2006.....	33,300	219	4,189	4,985	6,591	7,426	4,583	2,384	2,075	840	8
2005.....	32,637	272	4,212	4,990	6,550	6,991	4,210	2,344	2,200	860	8
2004.....	32,439	285	4,316	5,074	6,638	6,906	4,011	2,279	2,120	799	11
2003.....	31,484	250	3,988	5,065	6,602	6,481	3,843	2,335	2,115	798	7
2002.....	31,655	264	4,010	5,046	6,851	6,308	3,618	2,463	2,259	826	10
2001 4/.....	30,622	279	3,971	5,070	6,635	5,942	3,317	2,432	2,192	769	15
2000.....	29,350	307	3,994	4,792	6,562	5,437	2,945	2,292	2,181	833	7
1999.....	29,199	244	3,901	5,106	6,466	5,081	2,896	2,471	2,217	801	16

All races, male

2007.....	27,269	131	3,481	4,281	5,152	5,824	3,826	2,003	1,824	742	5
2006.....	26,308	153	3,528	4,053	5,065	5,576	3,463	1,968	1,765	729	8
2005.....	25,907	204	3,498	4,066	5,063	5,269	3,250	1,940	1,889	721	7
2004.....	25,566	187	3,596	4,142	5,067	5,078	3,088	1,901	1,817	679	11
2003.....	25,203	194	3,381	4,156	5,138	4,887	2,987	1,954	1,809	690	7
2002.....	25,409	199	3,432	4,135	5,305	4,796	2,837	2,053	1,938	704	10
2001 4/.....	24,672	214	3,409	4,199	5,180	4,504	2,563	2,040	1,885	664	14
2000.....	23,618	244	3,424	3,938	5,114	4,149	2,265	1,888	1,884	705	7
1999.....	23,458	194	3,326	4,194	5,019	3,854	2,255	2,051	1,868	681	16

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

All races, female

2007.....	7,329	53	659	997	1,570	1,954	1,243	441	295	116	1
2006.....	6,992	66	661	932	1,526	1,850	1,120	416	310	111	-
2005.....	6,730	68	714	924	1,487	1,722	960	404	311	139	1
2004.....	6,873	98	720	932	1,571	1,828	923	378	303	120	-
2003.....	6,281	56	607	909	1,464	1,594	856	381	306	108	-
2002.....	6,246	65	578	911	1,546	1,512	781	410	321	122	-
2001 4/.....	5,950	65	562	871	1,455	1,438	754	392	307	105	1
2000.....	5,732	63	570	854	1,448	1,288	680	404	297	128	-
1999.....	5,741	50	575	912	1,447	1,227	641	420	349	120	-

White, both sexes

2007 1/.....	31,348	125	3,442	4,524	6,033	7,259	4,792	2,317	2,017	834	5
2006 1/.....	30,138	167	3,477	4,259	5,945	6,924	4,322	2,243	1,991	806	4
2005 1/.....	29,527	209	3,540	4,229	5,917	6,505	3,979	2,215	2,106	819	8
2004 1/.....	29,251	218	3,610	4,288	6,031	6,420	3,751	2,137	2,019	768	9
2003 1/.....	28,485	193	3,318	4,305	5,983	6,064	3,618	2,198	2,029	770	7
2002.....	28,731	214	3,406	4,298	6,219	5,885	3,414	2,329	2,163	793	10
2001 4/.....	27,710	218	3,334	4,356	6,017	5,540	3,106	2,282	2,100	743	14
2000.....	26,475	254	3,315	4,084	5,942	5,035	2,775	2,168	2,090	806	6
1999.....	26,300	206	3,219	4,346	5,860	4,720	2,698	2,330	2,129	779	13

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

White, male

2007 1/.....	24,725	91	2,896	3,676	4,628	5,436	3,618	1,901	1,750	725	4
2006 1/.....	23,767	117	2,939	3,441	4,548	5,192	3,259	1,862	1,704	701	4
2005 1/.....	23,478	160	2,945	3,447	4,572	4,906	3,080	1,848	1,817	696	7
2004 1/.....	23,081	143	3,016	3,526	4,606	4,716	2,885	1,783	1,738	659	9
2003 1/.....	22,830	152	2,828	3,530	4,651	4,588	2,816	1,845	1,741	672	7
2002.....	23,049	159	2,919	3,495	4,813	4,477	2,684	1,953	1,861	678	10
2001 4/.....	22,328	168	2,860	3,604	4,691	4,218	2,396	1,921	1,810	647	13
2000.....	21,293	200	2,853	3,363	4,620	3,834	2,133	1,784	1,817	683	6
1999.....	21,107	164	2,751	3,567	4,535	3,573	2,099	1,940	1,801	664	13

White, female

2007 1/.....	6,623	34	546	848	1,405	1,823	1,174	416	267	109	1
2006 1/.....	6,371	50	538	818	1,397	1,732	1,063	381	287	105	-
2005 1/.....	6,049	49	595	782	1,345	1,599	899	367	289	123	1
2004 1/.....	6,170	75	594	762	1,425	1,704	866	354	281	109	-
2003 1/.....	5,655	41	490	775	1,332	1,476	802	353	288	98	-
2002.....	5,682	55	487	803	1,406	1,408	730	376	302	115	-
2001 4/.....	5,382	50	474	752	1,326	1,322	710	361	290	96	1
2000.....	5,182	54	462	721	1,322	1,201	642	384	273	123	-
1999.....	5,193	42	468	779	1,325	1,147	599	390	328	115	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)													
Black, both sexes													
2007 1/.....	1,958	42	406	481	424	321	158	71	48	6	1
2006 1/.....	1,954	37	416	474	418	310	156	75	45	22	1
2005 1/.....	1,992	41	437	485	419	314	144	79	56	17	-
2004 1/.....	2,019	49	465	521	381	288	166	76	60	12	1
2003 1/.....	1,955	44	447	509	416	269	137	77	44	12	-
2002.....	1,939	35	403	504	442	273	127	82	53	20	-
2001 4/.....	1,957	41	433	492	425	277	131	91	54	13	-
2000.....	1,962	40	480	476	438	275	103	85	53	12	-
1999.....	1,950	29	470	503	435	230	118	96	52	15	2
Black, male													
2007 1/.....	1,606	30	352	409	334	245	126	64	42	3	1
2006 1/.....	1,669	30	358	421	352	251	128	66	43	19	1
2005 1/.....	1,621	27	382	402	337	240	109	62	47	15	-
2004 1/.....	1,655	30	396	429	304	221	140	71	54	9	1
2003 1/.....	1,597	35	385	432	329	202	107	59	40	8	-
2002.....	1,633	29	351	447	354	219	99	68	48	18	-
2001 4/.....	1,627	32	395	425	336	204	106	74	46	9	-
2000.....	1,636	35	415	402	356	219	82	76	40	11	-
1999.....	1,651	22	414	436	356	187	97	80	45	12	2

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

Black, female

2007 1/.....	352	12	54	72	90	76	32	7	6	3	-
2006 1/.....	285	7	58	53	66	59	28	9	2	3	-
2005 1/.....	371	14	55	83	82	74	35	17	9	2	-
2004 1/.....	364	19	69	92	77	67	26	5	6	3	-
2003 1/.....	358	9	62	77	87	67	30	18	4	4	-
2002.....	306	6	52	57	88	54	28	14	5	2	-
2001 4/.....	330	9	38	67	89	73	25	17	8	4	-
2000.....	326	5	65	74	82	56	21	9	13	1	-
1999.....	299	7	56	67	79	43	21	16	7	3	-

American Indian 2/, both sexes

2007 1/.....	392	9	122	98	74	50	23	11	4	1	-
2006 1/.....	395	12	136	77	86	51	22	8	2	1	-
2005 1/.....	392	14	128	91	84	53	12	5	5	-	-
2004 1/.....	404	10	122	102	90	54	12	11	2	-	1
2003 1/.....	322	7	103	86	77	26	12	6	4	1	-
2002.....	324	6	100	84	68	45	13	3	5	-	-
2001 4/.....	321	12	86	76	80	39	17	6	5	-	-
2000.....	297	7	89	78	67	31	16	8	1	-	-
1999.....	291	2	95	91	49	31	12	9	2	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

American Indian 2/, male

2007 1/.....	310	6	99	81	59	37	17	7	4	-	-
2006 1/.....	309	4	110	63	63	39	20	7	2	1	-
2005 1/.....	313	11	99	73	68	42	11	4	5	-	-
2004 1/.....	306	6	92	76	71	40	9	9	2	-	1
2003 1/.....	265	6	80	76	66	19	9	5	3	1	-
2002.....	258	4	80	74	52	30	10	3	5	-	-
2001 4/.....	259	7	69	63	65	33	13	4	5	-	-
2000.....	237	6	71	59	53	24	15	8	1	-	-
1999.....	233	2	79	72	38	24	7	9	2	-	-

American Indian 2/, female

2007 1/.....	82	3	23	17	15	13	6	4	-	1	-
2006 1/.....	86	8	26	14	23	12	2	1	-	-	-
2005 1/.....	79	3	29	18	16	11	1	1	-	-	-
2004 1/.....	98	4	30	26	19	14	3	2	-	-	-
2003 1/.....	57	1	23	10	11	7	3	1	1	-	-
2002.....	66	2	20	10	16	15	3	-	-	-	-
2001 4/.....	62	5	17	13	15	6	4	2	-	-	-
2000.....	60	1	18	19	14	7	1	-	-	-	-
1999.....	58	-	16	19	11	7	5	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	900	8	170	175	191	148	96	45	50	17	-
2006 1/.....	813	3	160	175	142	141	83	58	37	11	3
2005 1/.....	726	8	107	185	130	119	75	45	33	24	-
2004 1/.....	765	8	119	163	136	144	82	55	39	19	-
2003 1/.....	722	6	120	165	126	122	76	54	38	15	-
2002.....	661	9	101	160	122	105	64	49	38	13	-
2001 4/.....	634	8	118	146	113	86	63	53	33	13	1
2000.....	616	6	110	154	115	96	51	31	37	15	1
1999.....	658	7	117	166	122	100	68	36	34	7	1

Asian or Pacific Islander 3/, male

2007 1/.....	628	4	134	115	131	106	65	31	28	14	-
2006 1/.....	563	2	121	128	102	94	56	33	16	8	3
2005 1/.....	495	6	72	144	86	81	50	26	20	10	-
2004 1/.....	524	8	92	111	86	101	54	38	23	11	-
2003 1/.....	511	1	88	118	92	78	55	45	25	9	-
2002.....	469	7	82	119	86	70	44	29	24	8	-
2001 4/.....	458	7	85	107	88	49	48	41	24	8	1
2000.....	452	3	85	114	85	72	35	20	26	11	1
1999.....	467	6	82	119	90	70	52	22	20	5	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

Asian or Pacific Islander 3/, female

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
2007 1/.....	272	4	36	60	60	42	31	14	22	3	-
2006 1/.....	250	1	39	47	40	47	27	25	21	3	-
2005 1/.....	231	2	35	41	44	38	25	19	13	14	-
2004 1/.....	241	-	27	52	50	43	28	17	16	8	-
2003 1/.....	211	5	32	47	34	44	21	9	13	6	-
2002.....	192	2	19	41	36	35	20	20	14	5	-
2001 4/.....	176	1	33	39	25	37	15	12	9	5	-
2000.....	164	3	25	40	30	24	16	11	11	4	-
1999.....	191	1	35	47	32	30	16	14	14	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

All races, both sexes

2007.....	17,352	53	1,900	2,306	2,879	3,531	2,786	1,700	1,589	606	2
2006.....	16,883	62	1,978	2,150	2,821	3,481	2,562	1,689	1,562	577	1
2005.....	17,002	84	1,962	2,269	2,855	3,472	2,470	1,669	1,649	571	1
2004.....	16,750	59	2,104	2,283	2,868	3,349	2,328	1,631	1,588	537	3
2003.....	16,907	74	2,075	2,381	2,927	3,279	2,317	1,700	1,595	559	-
2002.....	17,108	86	2,088	2,399	3,157	3,136	2,234	1,776	1,680	550	2
2001.....	16,869	90	2,130	2,564	3,030	3,023	2,083	1,758	1,648	537	6
2000.....	16,586	110	2,267	2,453	3,191	2,858	1,833	1,700	1,611	558	5
1999.....	16,599	103	2,315	2,621	3,097	2,677	1,860	1,791	1,606	524	5

All races, male

2007.....	15,181	45	1,727	2,033	2,393	2,977	2,397	1,539	1,483	585	2
2006.....	14,734	52	1,796	1,892	2,358	2,907	2,184	1,522	1,470	552	1
2005.....	14,916	72	1,771	1,981	2,416	2,945	2,140	1,501	1,548	542	-
2004.....	14,523	50	1,889	1,978	2,376	2,761	1,994	1,475	1,479	518	3
2003.....	14,827	63	1,879	2,099	2,470	2,768	2,001	1,538	1,480	529	-
2002.....	15,045	68	1,897	2,078	2,672	2,677	1,930	1,620	1,576	525	2
2001.....	14,758	69	1,935	2,246	2,551	2,551	1,771	1,592	1,531	506	6
2000.....	14,454	90	2,043	2,123	2,657	2,409	1,568	1,528	1,505	526	5
1999.....	14,479	80	2,083	2,270	2,586	2,251	1,594	1,615	1,499	496	5

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

All races, female

2007.....	2,171	8	173	273	486	554	389	161	106	21	-
2006.....	2,149	10	182	258	463	574	378	167	92	25	-
2005.....	2,086	12	191	288	439	527	330	168	101	29	1
2004.....	2,227	9	215	305	492	588	334	156	109	19	-
2003.....	2,080	11	196	282	457	511	316	162	115	30	-
2002.....	2,063	18	191	321	485	459	304	156	104	25	-
2001.....	2,111	21	195	318	479	472	312	166	117	31	-
2000.....	2,132	20	224	330	534	449	265	172	106	32	-
1999.....	2,120	23	232	351	511	426	266	176	107	28	-

White, both sexes

2007 1/.....	16,009	43	1,563	1,993	2,644	3,333	2,660	1,634	1,537	600	2
2006 1/.....	15,539	57	1,632	1,817	2,578	3,303	2,454	1,622	1,520	555	1
2005 1/.....	15,681	75	1,637	1,943	2,612	3,277	2,374	1,602	1,599	561	1
2004 1/.....	15,407	49	1,770	1,937	2,636	3,176	2,220	1,555	1,534	528	2
2003 1/.....	15,587	65	1,728	2,018	2,716	3,114	2,221	1,632	1,543	550	-
2002.....	15,733	73	1,747	2,051	2,900	2,955	2,135	1,711	1,625	534	2
2001.....	15,467	73	1,753	2,220	2,770	2,864	1,988	1,680	1,589	525	5
2000.....	15,174	103	1,878	2,098	2,933	2,680	1,732	1,626	1,573	547	4
1999.....	15,135	94	1,899	2,253	2,833	2,514	1,754	1,714	1,557	512	5

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) by discharge of firearms (X72-X74)													
White, male													
2007 1/.....	13,986	36	1,412	1,754	2,190	2,809	2,287	1,479	1,437	580	2
2006 1/.....	13,527	48	1,485	1,589	2,141	2,756	2,085	1,460	1,429	533	1
2005 1/.....	13,730	65	1,474	1,687	2,197	2,772	2,057	1,446	1,500	532	-
2004 1/.....	13,341	41	1,590	1,671	2,178	2,618	1,901	1,403	1,427	510	2
2003 1/.....	13,649	56	1,566	1,765	2,284	2,626	1,918	1,479	1,433	522	-
2002.....	13,809	58	1,586	1,757	2,451	2,521	1,840	1,562	1,523	509	2
2001.....	13,521	58	1,581	1,936	2,331	2,424	1,686	1,525	1,480	495	5
2000.....	13,214	86	1,693	1,815	2,431	2,261	1,479	1,457	1,473	515	4
1999.....	13,167	72	1,707	1,938	2,349	2,113	1,499	1,545	1,454	485	5
White, female													
2007 1/.....	2,023	7	151	239	454	524	373	155	100	20	-
2006 1/.....	2,012	9	147	228	437	547	369	162	91	22	-
2005 1/.....	1,951	10	163	256	415	505	317	156	99	29	1
2004 1/.....	2,066	8	180	266	458	558	319	152	107	18	-
2003 1/.....	1,938	9	162	253	432	488	303	153	110	28	-
2002.....	1,924	15	161	294	449	434	295	149	102	25	-
2001.....	1,946	15	172	284	439	440	302	155	109	30	-
2000.....	1,960	17	185	283	502	419	253	169	100	32	-
1999.....	1,968	22	192	315	484	401	255	169	103	27	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Black, both sexes

2007 1/.....	987	9	237	251	177	141	83	46	39	4	-
2006 1/.....	1,015	2	250	261	184	129	78	53	39	19	-
2005 1/.....	1,009	6	247	249	189	137	73	57	41	10	-
2004 1/.....	1,009	8	260	265	156	116	87	61	48	8	-
2003 1/.....	1,004	7	269	281	151	119	75	56	39	7	-
2002.....	1,059	9	255	274	201	133	73	56	42	16	-
2001.....	1,091	13	296	267	200	124	70	62	49	10	-
2000.....	1,090	4	308	272	194	138	72	62	32	8	-
1999.....	1,125	5	327	272	199	120	82	65	43	12	-

Black, male

2007 1/.....	881	8	222	224	152	118	75	44	35	3	-
2006 1/.....	922	2	228	238	168	112	71	49	38	16	-
2005 1/.....	911	5	230	224	172	122	62	47	39	10	-
2004 1/.....	904	7	231	239	139	100	75	60	46	7	-
2003 1/.....	902	5	248	262	131	105	63	47	35	6	-
2002.....	962	8	234	259	172	118	65	50	40	16	-
2001.....	971	10	280	242	171	100	64	53	42	9	-
2000.....	965	4	281	240	169	114	62	60	27	8	-
1999.....	1,018	4	298	250	179	104	74	59	39	11	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Black, female

2007 1/.....	106	1	15	27	25	23	8	2	4	1	-
2006 1/.....	93	-	22	23	16	17	7	4	1	3	-
2005 1/.....	98	1	17	25	17	15	11	10	2	-	-
2004 1/.....	105	1	29	26	17	16	12	1	2	1	-
2003 1/.....	102	2	21	19	20	14	12	9	4	1	-
2002.....	97	1	21	15	29	15	8	6	2	-	-
2001.....	120	3	16	25	29	24	6	9	7	1	-
2000.....	125	-	27	32	25	24	10	2	5	-	-
1999.....	107	1	29	22	20	16	8	6	4	1	-

American Indian 2/, both sexes

2007 1/.....	135	1	43	21	21	24	13	8	4	-	-
2006 1/.....	154	2	50	29	33	23	11	4	1	1	-
2005 1/.....	163	2	49	39	29	28	8	3	5	-	-
2004 1/.....	152	1	42	37	32	23	7	7	2	-	1
2003 1/.....	127	1	39	35	26	13	6	4	2	1	-
2002.....	146	1	53	33	27	19	7	2	4	-	-
2001.....	132	2	40	29	32	14	7	4	4	-	-
2000.....	133	1	43	27	25	18	11	7	1	-	-
1999.....	136	1	49	37	22	15	4	6	2	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

American Indian 2/, male

2007 1/.....	116	1	39	18	19	19	11	5	4	-	-
2006 1/.....	132	1	45	27	26	18	9	4	1	1	-
2005 1/.....	145	1	43	35	26	26	7	2	5	-	-
2004 1/.....	130	1	39	32	28	16	6	5	2	-	1
2003 1/.....	117	1	35	31	25	12	6	4	2	1	-
2002.....	128	-	46	30	25	15	6	2	4	-	-
2001.....	115	-	37	26	26	13	6	3	4	-	-
2000.....	109	-	37	20	20	13	11	7	1	-	-
1999.....	120	1	43	32	20	13	3	6	2	-	-

American Indian 2/, female

2007 1/.....	19	-	4	3	2	5	2	3	-	-	-
2006 1/.....	22	1	5	2	7	5	2	-	-	-	-
2005 1/.....	18	1	6	4	3	2	1	1	-	-	-
2004 1/.....	22	-	3	5	4	7	1	2	-	-	-
2003 1/.....	10	-	4	4	1	1	-	-	-	-	-
2002.....	18	1	7	3	2	4	1	-	-	-	-
2001.....	17	2	3	3	6	1	1	1	-	-	-
2000.....	24	1	6	7	5	5	-	-	-	-	-
1999.....	16	-	6	5	2	2	1	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	221	-	57	41	37	33	30	12	9	2	-
2006 1/.....	175	1	46	43	26	26	19	10	2	2	-
2005 1/.....	149	1	29	38	25	30	15	7	4	-	-
2004 1/.....	182	1	32	44	44	34	14	8	4	1	-
2003 1/.....	189	1	39	47	34	33	15	8	11	1	-
2002.....	170	3	33	41	29	29	19	7	9	-	-
2001.....	179	2	41	48	28	21	18	12	6	2	1
2000.....	189	2	38	56	39	22	18	5	5	3	1
1999.....	203	3	40	59	43	28	20	6	4	-	-

Asian or Pacific Islander 3/, male

2007 1/.....	198	-	54	37	32	31	24	11	7	2	-
2006 1/.....	153	1	38	38	23	21	19	9	2	2	-
2005 1/.....	130	1	24	35	21	25	14	6	4	-	-
2004 1/.....	148	1	29	36	31	27	12	7	4	1	-
2003 1/.....	159	1	30	41	30	25	14	8	10	-	-
2002.....	146	2	31	32	24	23	19	6	9	-	-
2001.....	151	1	37	42	23	14	15	11	5	2	1
2000.....	166	-	32	48	37	21	16	4	4	3	1
1999.....	174	3	35	50	38	21	18	5	4	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Asian or Pacific Islander 3/, female

2007 1/.....	23	-	3	4	5	2	6	1	2	-	-
2006 1/.....	22	-	8	5	3	5	-	1	-	-	-
2005 1/.....	19	-	5	3	4	5	1	1	-	-	-
2004 1/.....	34	-	3	8	13	7	2	1	-	-	-
2003 1/.....	30	-	9	6	4	8	1	-	1	1	-
2002.....	24	1	2	9	5	6	-	1	-	-	-
2001.....	28	1	4	6	5	7	3	1	1	-	-
2000.....	23	2	6	8	2	1	2	1	1	-	-
1999.....	29	-	5	9	5	7	2	1	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

All races, both sexes

2007.....	17,246	131	2,240	2,972	3,843	4,247	2,283	744	530	252	4
2006.....	16,417	157	2,211	2,835	3,770	3,945	2,021	695	513	263	7
2005.....	15,635	188	2,250	2,721	3,695	3,519	1,740	675	551	289	7
2004.....	15,689	226	2,212	2,791	3,770	3,557	1,683	648	532	262	8
2003.....	14,577	176	1,913	2,684	3,675	3,202	1,526	635	520	239	7
2002.....	14,547	178	1,922	2,647	3,694	3,172	1,384	687	579	276	8
2001 4/.....	13,753	189	1,841	2,506	3,605	2,919	1,234	674	544	232	9
2000.....	12,764	197	1,727	2,339	3,371	2,579	1,112	592	570	275	2
1999.....	12,600	141	1,586	2,485	3,369	2,404	1,036	680	611	277	11

All races, male

2007.....	12,088	86	1,754	2,248	2,759	2,847	1,429	464	341	157	3
2006.....	11,574	101	1,732	2,161	2,707	2,669	1,279	446	295	177	7
2005.....	10,991	132	1,727	2,085	2,647	2,324	1,110	439	341	179	7
2004.....	11,043	137	1,707	2,164	2,691	2,317	1,094	426	338	161	8
2003.....	10,376	131	1,502	2,057	2,668	2,119	986	416	329	161	7
2002.....	10,364	131	1,535	2,057	2,633	2,119	907	433	362	179	8
2001 4/.....	9,914	145	1,474	1,953	2,629	1,953	792	448	354	158	8
2000.....	9,164	154	1,381	1,815	2,457	1,740	697	360	379	179	2
1999.....	8,979	114	1,243	1,924	2,433	1,603	661	436	369	185	11

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

All races, female

2007.....	5,158	45	486	724	1,084	1,400	854	280	189	95	1
2006.....	4,843	56	479	674	1,063	1,276	742	249	218	86	-
2005.....	4,644	56	523	636	1,048	1,195	630	236	210	110	-
2004.....	4,646	89	505	627	1,079	1,240	589	222	194	101	-
2003.....	4,201	45	411	627	1,007	1,083	540	219	191	78	-
2002.....	4,183	47	387	590	1,061	1,053	477	254	217	97	-
2001 4/.....	3,839	44	367	553	976	966	442	226	190	74	1
2000.....	3,600	43	346	524	914	839	415	232	191	96	-
1999.....	3,621	27	343	561	936	801	375	244	242	92	-

White, both sexes

2007 1/.....	15,339	82	1,879	2,531	3,389	3,926	2,132	683	480	234	3
2006 1/.....	14,599	110	1,845	2,442	3,367	3,621	1,868	621	471	251	3
2005 1/.....	13,846	134	1,903	2,286	3,305	3,228	1,605	613	507	258	7
2004 1/.....	13,844	169	1,840	2,351	3,395	3,244	1,531	582	485	240	7
2003 1/.....	12,898	128	1,590	2,287	3,267	2,950	1,397	566	486	220	7
2002.....	12,998	141	1,659	2,247	3,319	2,930	1,279	618	538	259	8
2001 4/.....	12,243	145	1,581	2,136	3,247	2,676	1,118	602	511	218	9
2000.....	11,301	151	1,437	1,986	3,009	2,355	1,043	542	517	259	2
1999.....	11,165	112	1,320	2,093	3,027	2,206	944	616	572	267	8

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
White, male													
2007 1/.....	10,739	55	1,484	1,922	2,438	2,627	1,331	422	313	145	2
2006 1/.....	10,240	69	1,454	1,852	2,407	2,436	1,174	402	275	168	3
2005 1/.....	9,748	95	1,471	1,760	2,375	2,134	1,023	402	317	164	7
2004 1/.....	9,740	102	1,426	1,855	2,428	2,098	984	380	311	149	7
2003 1/.....	9,181	96	1,262	1,765	2,367	1,962	898	366	308	150	7
2002.....	9,240	101	1,333	1,738	2,362	1,956	844	391	338	169	8
2001 4/.....	8,807	110	1,279	1,668	2,360	1,794	710	396	330	152	8
2000.....	8,079	114	1,160	1,548	2,189	1,573	654	327	344	168	2
1999.....	7,940	92	1,044	1,629	2,186	1,460	600	395	347	179	8
White, female													
2007 1/.....	4,600	27	395	609	951	1,299	801	261	167	89	1
2006 1/.....	4,359	41	391	590	960	1,185	694	219	196	83	-
2005 1/.....	4,098	39	432	526	930	1,094	582	211	190	94	-
2004 1/.....	4,104	67	414	496	967	1,146	547	202	174	91	-
2003 1/.....	3,717	32	328	522	900	988	499	200	178	70	-
2002.....	3,758	40	326	509	957	974	435	227	200	90	-
2001 4/.....	3,436	35	302	468	887	882	408	206	181	66	1
2000.....	3,222	37	277	438	820	782	389	215	173	91	-
1999.....	3,225	20	276	464	841	746	344	221	225	88	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

Black, both sexes

2007 1/.....	971	33	169	230	247	180	75	25	9	2	1
2006 1/.....	939	35	166	213	234	181	78	22	6	3	1
2005 1/.....	983	35	190	236	230	177	71	22	15	7	-
2004 1/.....	1,010	41	205	256	225	172	79	15	12	4	1
2003 1/.....	951	37	178	228	265	150	62	21	5	5	-
2002.....	880	26	148	230	241	140	54	26	11	4	-
2001 4/.....	866	28	137	225	225	153	61	29	5	3	-
2000.....	872	36	172	204	244	137	31	23	21	4	-
1999.....	825	24	143	231	236	110	36	31	9	3	2

Black, male

2007 1/.....	725	22	130	185	182	127	51	20	7	-	1
2006 1/.....	747	28	130	183	184	139	57	17	5	3	1
2005 1/.....	710	22	152	178	165	118	47	15	8	5	-
2004 1/.....	751	23	165	190	165	121	65	11	8	2	1
2003 1/.....	695	30	137	170	198	97	44	12	5	2	-
2002.....	671	21	117	188	182	101	34	18	8	2	-
2001 4/.....	656	22	115	183	165	104	42	21	4	-	-
2000.....	671	31	134	162	187	105	20	16	13	3	-
1999.....	633	18	116	186	177	83	23	21	6	1	2

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

Black, female

2007 1/.....	246	11	39	45	65	53	24	5	2	2	-
2006 1/.....	192	7	36	30	50	42	21	5	1	-	-
2005 1/.....	273	13	38	58	65	59	24	7	7	2	-
2004 1/.....	259	18	40	66	60	51	14	4	4	2	-
2003 1/.....	256	7	41	58	67	53	18	9	-	3	-
2002.....	209	5	31	42	59	39	20	8	3	2	-
2001 4/.....	210	6	22	42	60	49	19	8	1	3	-
2000.....	201	5	38	42	57	32	11	7	8	1	-
1999.....	192	6	27	45	59	27	13	10	3	2	-

American Indian 2/, both sexes

2007 1/.....	257	8	79	77	53	26	10	3	-	1	-
2006 1/.....	241	10	86	48	53	28	11	4	1	-	-
2005 1/.....	229	12	79	52	55	25	4	2	-	-	-
2004 1/.....	252	9	80	65	58	31	5	4	-	-	-
2003 1/.....	195	6	64	51	51	13	6	2	2	-	-
2002.....	178	5	47	51	41	26	6	1	1	-	-
2001 4/.....	189	10	46	47	48	25	10	2	1	-	-
2000.....	164	6	46	51	42	13	5	1	-	-	-
1999.....	155	1	46	54	27	16	8	3	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

American Indian 2/, male

2007 1/.....	194	5	60	63	40	18	6	2	-	-	-
2006 1/.....	177	3	65	36	37	21	11	3	1	-	-
2005 1/.....	168	10	56	38	42	16	4	2	-	-	-
2004 1/.....	176	5	53	44	43	24	3	4	-	-	-
2003 1/.....	148	5	45	45	41	7	3	1	1	-	-
2002.....	130	4	34	44	27	15	4	1	1	-	-
2001 4/.....	144	7	32	37	39	20	7	1	1	-	-
2000.....	128	6	34	39	33	11	4	1	-	-	-
1999.....	113	1	36	40	18	11	4	3	-	-	-

American Indian 2/, female

2007 1/.....	63	3	19	14	13	8	4	1	-	1	-
2006 1/.....	64	7	21	12	16	7	-	1	-	-	-
2005 1/.....	61	2	23	14	13	9	-	-	-	-	-
2004 1/.....	76	4	27	21	15	7	2	-	-	-	-
2003 1/.....	47	1	19	6	10	6	3	1	1	-	-
2002.....	48	1	13	7	14	11	2	-	-	-	-
2001 4/.....	45	3	14	10	9	5	3	1	-	-	-
2000.....	36	-	12	12	9	2	1	-	-	-	-
1999.....	42	-	10	14	9	5	4	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	679	8	113	134	154	115	66	33	41	15	-
2006 1/.....	638	2	114	132	116	115	64	48	35	9	3
2005 1/.....	577	7	78	147	105	89	60	38	29	24	-
2004 1/.....	583	7	87	119	92	110	68	47	35	18	-
2003 1/.....	533	5	81	118	92	89	61	46	27	14	-
2002.....	491	6	68	119	93	76	45	42	29	13	-
2001 4/.....	455	6	77	98	85	65	45	41	27	11	-
2000.....	427	4	72	98	76	74	33	26	32	12	-
1999.....	455	4	77	107	79	72	48	30	30	7	1

Asian or Pacific Islander 3/, male

2007 1/.....	430	4	80	78	99	75	41	20	21	12	-
2006 1/.....	410	1	83	90	79	73	37	24	14	6	3
2005 1/.....	365	5	48	109	65	56	36	20	16	10	-
2004 1/.....	376	7	63	75	55	74	42	31	19	10	-
2003 1/.....	352	-	58	77	62	53	41	37	15	9	-
2002.....	323	5	51	87	62	47	25	23	15	8	-
2001 4/.....	307	6	48	65	65	35	33	30	19	6	-
2000.....	286	3	53	66	48	51	19	16	22	8	-
1999.....	293	3	47	69	52	49	34	17	16	5	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

Asian or Pacific Islander 3/, female

2007 1/.....	249	4	33	56	55	40	25	13	20	3	-
2006 1/.....	228	1	31	42	37	42	27	24	21	3	-
2005 1/.....	212	2	30	38	40	33	24	18	13	14	-
2004 1/.....	207	-	24	44	37	36	26	16	16	8	-
2003 1/.....	181	5	23	41	30	36	20	9	12	5	-
2002.....	168	1	17	32	31	29	20	19	14	5	-
2001 4/.....	148	-	29	33	20	30	12	11	8	5	-
2000.....	141	1	19	32	28	23	14	10	10	4	-
1999.....	162	1	30	38	27	23	14	13	14	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

All races, both sexes

2007.....	18,361	352	398	346	5,551	4,758	3,052	2,140	980	411	268	80	25
2006.....	18,573	336	366	390	5,717	4,725	3,020	2,209	1,012	398	279	102	19
2005.....	18,124	306	375	341	5,466	4,752	3,109	2,060	862	440	283	109	21
2004.....	17,357	325	377	329	5,085	4,495	2,984	2,008	879	446	289	104	36
2003.....	17,732	341	376	324	5,368	4,516	3,110	2,017	786	432	317	104	41
2002.....	17,638	303	423	356	5,219	4,489	3,239	1,915	841	421	296	95	41
2001 4/.....	20,308	332	415	326	5,297	5,204	4,268	2,467	1,018	532	312	105	32
2000.....	16,765	349	356	371	4,939	4,164	3,219	1,755	738	443	301	100	30
1999.....	16,889	331	376	432	4,998	4,231	3,206	1,668	722	473	304	99	49

All races, male

2007.....	14,538	206	213	210	4,829	4,019	2,289	1,572	727	279	149	26	19
2006.....	14,717	199	208	246	4,991	3,983	2,269	1,628	738	260	141	38	16
2005.....	14,376	173	214	204	4,765	4,008	2,317	1,579	636	282	138	44	16
2004.....	13,578	167	204	200	4,380	3,764	2,224	1,492	637	302	138	41	29
2003.....	13,882	205	202	215	4,624	3,741	2,308	1,488	568	285	171	46	29
2002.....	13,640	164	234	197	4,474	3,669	2,386	1,394	614	277	158	41	32
2001 4/.....	15,555	197	238	193	4,541	4,157	3,122	1,832	723	332	155	42	23
2000.....	12,820	203	198	227	4,203	3,361	2,319	1,284	537	279	154	40	15
1999.....	12,785	187	194	250	4,191	3,346	2,307	1,229	541	304	162	41	33

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

All races, female

2007.....	3,823	146	185	136	722	739	763	568	253	132	119	54	6
2006.....	3,856	137	158	144	726	742	751	581	274	138	138	64	3
2005.....	3,748	133	161	137	701	744	792	481	226	158	145	65	5
2004.....	3,779	158	173	129	705	731	760	516	242	144	151	63	7
2003.....	3,850	136	174	109	744	775	802	529	218	147	146	58	12
2002.....	3,998	139	189	159	745	820	853	521	227	144	138	54	9
2001 4/.....	4,753	135	177	133	756	1,047	1,146	635	295	200	157	63	9
2000.....	3,945	146	158	144	736	803	900	471	201	164	147	60	15
1999.....	4,104	144	182	182	807	885	899	439	181	169	142	58	16

White, both sexes

2007 1/.....	8,914	228	216	196	2,193	2,050	1,577	1,271	616	289	196	61	21
2006 1/.....	8,860	220	190	227	2,213	1,902	1,569	1,283	656	275	225	83	17
2005 1/.....	8,770	195	217	168	2,172	2,022	1,615	1,182	573	295	223	88	20
2004 1/.....	8,643	208	201	184	2,126	1,920	1,592	1,170	582	323	219	84	34
2003 1/.....	8,709	216	204	159	2,171	1,911	1,653	1,195	530	304	244	86	36
2002.....	8,685	170	239	195	2,137	1,865	1,739	1,117	587	306	218	76	36
2001 4/.....	11,328	196	228	218	2,264	2,623	2,672	1,639	744	392	240	88	24
2000.....	8,339	197	199	240	1,981	1,814	1,725	1,048	497	303	227	83	25
1999.....	8,628	200	208	261	2,064	1,926	1,766	1,016	496	353	225	79	34

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)													
White, male													
2007 1/.....	6,541	130	116	106	1,792	1,625	1,117	903	430	189	99	17	17
2006 1/.....	6,514	121	109	130	1,824	1,520	1,131	910	458	162	105	30	14
2005 1/.....	6,457	109	119	96	1,800	1,587	1,134	881	410	177	100	28	16
2004 1/.....	6,302	112	113	109	1,727	1,501	1,130	839	409	205	99	31	27
2003 1/.....	6,337	129	96	106	1,781	1,487	1,147	841	364	194	128	39	25
2002.....	6,282	100	133	98	1,746	1,435	1,240	757	409	193	108	34	29
2001 4/.....	8,254	118	132	119	1,811	1,988	1,972	1,196	518	233	118	33	16
2000.....	5,925	125	114	148	1,583	1,356	1,202	718	350	176	109	31	13
1999.....	6,162	121	108	148	1,622	1,445	1,240	737	356	216	117	29	23
White, female													
2007 1/.....	2,373	98	100	90	401	425	460	368	186	100	97	44	4
2006 1/.....	2,346	99	81	97	389	382	438	373	198	113	120	53	3
2005 1/.....	2,313	86	98	72	372	435	481	301	163	118	123	60	4
2004 1/.....	2,341	96	88	75	399	419	462	331	173	118	120	53	7
2003 1/.....	2,372	87	108	53	390	424	506	354	166	110	116	47	11
2002.....	2,403	70	106	97	391	430	499	360	178	113	110	42	7
2001 4/.....	3,074	78	96	99	453	635	700	443	226	159	122	55	8
2000.....	2,414	72	85	92	398	458	523	330	147	127	118	52	12
1999.....	2,466	79	100	113	442	481	526	279	140	137	108	50	11

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)													
Black, both sexes													
2007 1/.....	8,870	110	168	140	3,213	2,560	1,366	802	319	106	66	17	3
2006 1/.....	9,032	109	154	149	3,295	2,670	1,347	825	309	109	45	18	2
2005 1/.....	8,669	95	140	150	3,077	2,591	1,369	797	248	125	58	18	1
2004 1/.....	8,135	103	163	131	2,803	2,445	1,297	752	254	104	63	18	2
2003 1/.....	8,392	106	156	149	3,006	2,475	1,345	740	223	106	65	17	4
2002.....	8,287	117	161	135	2,900	2,469	1,377	722	218	97	71	17	3
2001 4/.....	8,226	123	169	94	2,870	2,371	1,450	719	225	119	63	15	8
2000.....	7,867	140	145	124	2,819	2,224	1,376	628	204	125	64	14	4
1999.....	7,648	118	152	147	2,781	2,157	1,325	569	195	104	67	19	14
Black, male													
2007 1/.....	7,584	70	89	96	2,918	2,283	1,100	623	268	79	49	8	1
2006 1/.....	7,677	72	84	105	2,982	2,346	1,068	650	245	86	30	7	2
2005 1/.....	7,412	54	88	97	2,790	2,319	1,091	633	196	93	37	14	-
2004 1/.....	6,839	48	83	82	2,522	2,163	1,022	592	195	83	37	10	2
2003 1/.....	7,083	60	96	99	2,689	2,156	1,087	584	182	79	41	7	3
2002.....	6,896	56	92	86	2,582	2,129	1,058	582	181	74	48	6	2
2001 4/.....	6,780	70	101	68	2,600	2,028	1,051	556	170	87	34	8	7
2000.....	6,482	73	75	72	2,501	1,911	1,040	508	160	95	39	7	1
1999.....	6,214	60	75	91	2,453	1,800	996	439	160	80	39	11	10

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

Black, female

2007 1/.....	1,286	40	79	44	295	277	266	179	51	27	17	9	2
2006 1/.....	1,355	37	70	44	313	324	279	175	64	23	15	11	-
2005 1/.....	1,257	41	52	53	287	272	278	164	52	32	21	4	1
2004 1/.....	1,296	55	80	49	281	282	275	160	59	21	26	8	-
2003 1/.....	1,309	46	60	50	317	319	258	156	41	27	24	10	1
2002.....	1,391	61	69	49	318	340	319	140	37	23	23	11	1
2001 4/.....	1,446	53	68	26	270	343	399	163	55	32	29	7	1
2000.....	1,385	67	70	52	318	313	336	120	44	30	25	7	3
1999.....	1,434	58	77	56	328	357	329	130	35	24	28	8	4

American Indian 2/, both sexes

2007 1/.....	220	9	5	3	57	65	43	23	8	4	2	1	-
2006 1/.....	254	4	11	2	77	67	48	30	10	3	2	-	-
2005 1/.....	257	5	10	7	82	60	45	29	11	8	-	-	-
2004 1/.....	228	9	7	6	67	54	46	22	8	6	3	-	-
2003 1/.....	237	10	7	4	78	51	44	32	4	5	1	1	-
2002.....	267	10	12	11	70	63	52	29	9	7	3	-	1
2001 4/.....	211	6	12	4	59	54	36	26	5	6	2	1	-
2000.....	203	5	8	1	51	59	46	18	6	4	2	2	1
1999.....	253	5	6	9	60	73	44	33	11	7	5	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

American Indian 2/, male

2007 1/.....	163	4	3	3	45	52	32	16	6	2	-	-	-
2006 1/.....	206	4	9	2	69	53	34	21	9	3	2	-	-
2005 1/.....	193	4	4	3	67	45	32	25	9	4	-	-	-
2004 1/.....	181	5	5	4	53	47	38	18	5	5	1	-	-
2003 1/.....	175	8	5	2	58	41	29	26	2	4	-	-	-
2002.....	185	4	5	3	54	46	40	23	5	4	1	-	-
2001 4/.....	146	3	5	1	45	39	24	19	5	4	1	-	-
2000.....	159	3	6	1	46	46	32	14	4	3	1	2	1
1999.....	179	3	4	4	45	50	30	24	9	6	4	-	-

American Indian 2/, female

2007 1/.....	57	5	2	-	12	13	11	7	2	2	2	1	-
2006 1/.....	48	-	2	-	8	14	14	9	1	-	-	-	-
2005 1/.....	64	1	6	4	15	15	13	4	2	4	-	-	-
2004 1/.....	47	4	2	2	14	7	8	4	3	1	2	-	-
2003 1/.....	62	2	2	2	20	10	15	6	2	1	1	1	-
2002.....	82	6	7	8	16	17	12	6	4	3	2	-	1
2001 4/.....	65	3	7	3	14	15	12	7	-	2	1	1	-
2000.....	44	2	2	-	5	13	14	4	2	1	1	-	-
1999.....	74	2	2	5	15	23	14	9	2	1	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	357	5	9	7	88	83	66	44	37	12	4	1	1
2006 1/.....	427	3	11	12	132	86	56	71	37	11	7	1	-
2005 1/.....	428	11	8	16	135	79	80	52	30	12	2	3	-
2004 1/.....	351	5	6	8	89	76	49	64	35	13	4	2	-
2003 1/.....	394	9	9	12	113	79	68	50	29	17	7	-	1
2002.....	399	6	11	15	112	92	71	47	27	11	4	2	1
2001 4/.....	543	7	6	10	104	156	110	83	44	15	7	1	-
2000.....	356	7	4	6	88	67	72	61	31	11	8	1	-
1999.....	360	8	10	15	93	75	71	50	20	9	7	1	1

Asian or Pacific Islander 3/, male

2007 1/.....	250	2	5	5	74	59	40	30	23	9	1	1	1
2006 1/.....	320	2	6	9	116	64	36	47	26	9	4	1	-
2005 1/.....	314	6	3	8	108	57	60	40	21	8	1	2	-
2004 1/.....	256	2	3	5	78	53	34	43	28	9	1	-	-
2003 1/.....	287	8	5	8	96	57	45	37	20	8	2	-	1
2002.....	277	4	4	10	92	59	48	32	19	6	1	1	1
2001 4/.....	375	6	-	5	85	102	75	61	30	8	2	1	-
2000.....	254	2	3	6	73	48	45	44	23	5	5	-	-
1999.....	230	3	7	7	71	51	41	29	16	2	2	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

Asian or Pacific Islander 3/, female

2007 1/.....	107	3	4	2	14	24	26	14	14	3	3	-	-
2006 1/.....	107	1	5	3	16	22	20	24	11	2	3	-	-
2005 1/.....	114	5	5	8	27	22	20	12	9	4	1	1	-
2004 1/.....	95	3	3	3	11	23	15	21	7	4	3	2	-
2003 1/.....	107	1	4	4	17	22	23	13	9	9	5	-	-
2002.....	122	2	7	5	20	33	23	15	8	5	3	1	-
2001 4/.....	168	1	6	5	19	54	35	22	14	7	5	-	-
2000.....	102	5	1	-	15	19	27	17	8	6	3	1	-
1999.....	130	5	3	8	22	24	30	21	4	7	5	-	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

All races, both sexes

2007.....	12,632	15	48	201	4,669	3,751	2,038	1,159	446	185	88	23	9
2006.....	12,791	6	42	237	4,827	3,767	1,987	1,185	468	158	84	22	8
2005.....	12,352	6	37	187	4,499	3,780	2,010	1,097	405	201	97	24	9
2004.....	11,624	6	36	184	4,127	3,503	1,895	1,062	468	193	100	29	21
2003.....	11,920	8	40	187	4,410	3,540	1,941	1,110	394	161	88	23	18
2002.....	11,829	9	49	205	4,317	3,465	2,042	1,025	409	161	115	19	13
2001.....	11,348	11	55	180	4,200	3,308	1,978	934	364	192	99	16	11
2000.....	10,801	12	28	187	3,963	3,074	1,906	959	367	184	81	28	12
1999.....	10,828	8	50	224	4,038	3,048	1,868	893	373	193	95	23	15

All races, male

2007.....	10,767	11	23	141	4,199	3,324	1,645	902	341	121	48	3	9
2006.....	10,886	2	25	167	4,375	3,322	1,586	909	339	104	41	8	8
2005.....	10,561	6	22	131	4,096	3,344	1,599	859	312	131	44	10	7
2004.....	9,921	2	18	134	3,718	3,111	1,556	821	347	138	41	15	20
2003.....	10,126	7	20	134	3,977	3,125	1,535	860	292	115	35	10	16
2002.....	9,899	3	30	127	3,851	3,017	1,623	761	301	101	64	9	12
2001.....	9,532	7	27	125	3,796	2,864	1,528	726	264	123	54	7	11
2000.....	9,006	9	17	138	3,548	2,650	1,487	730	266	108	37	9	7
1999.....	8,944	5	26	155	3,569	2,609	1,414	676	294	124	48	10	14

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)													
All races, female													
2007.....	1,865	4	25	60	470	427	393	257	105	64	40	20	-
2006.....	1,905	4	17	70	452	445	401	276	129	54	43	14	-
2005.....	1,791	-	15	56	403	436	411	238	93	70	53	14	2
2004.....	1,703	4	18	50	409	392	339	241	121	55	59	14	1
2003.....	1,794	1	20	53	433	415	406	250	102	46	53	13	2
2002.....	1,930	6	19	78	466	448	419	264	108	60	51	10	1
2001.....	1,816	4	28	55	404	444	450	208	100	69	45	9	-
2000.....	1,795	3	11	49	415	424	419	229	101	76	44	19	5
1999.....	1,884	3	24	69	469	439	454	217	79	69	47	13	1
White, both sexes													
2007 1/.....	5,380	10	26	103	1,659	1,454	978	653	270	133	67	19	8
2006 1/.....	5,279	2	22	129	1,676	1,338	964	640	314	102	68	17	7
2005 1/.....	5,266	4	23	87	1,577	1,433	988	630	274	140	82	19	9
2004 1/.....	5,119	3	11	92	1,530	1,342	954	604	311	144	84	24	20
2003 1/.....	5,173	2	20	81	1,634	1,331	966	653	268	116	68	20	14
2002.....	5,185	3	33	106	1,600	1,301	1,027	596	284	116	91	17	11
2001.....	5,188	8	33	119	1,578	1,327	1,065	560	255	141	78	15	9
2000.....	4,806	5	17	114	1,434	1,192	985	567	260	134	64	24	10
1999.....	4,918	4	26	127	1,484	1,237	1,011	526	252	148	72	21	10

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

White, male

2007 1/.....	4,240	8	14	66	1,407	1,208	748	477	191	80	31	2	8
2006 1/.....	4,179	-	14	85	1,465	1,130	731	447	213	57	26	4	7
2005 1/.....	4,177	4	13	61	1,390	1,191	721	468	202	83	32	5	7
2004 1/.....	4,078	1	5	64	1,319	1,141	734	441	219	91	33	11	19
2003 1/.....	4,108	1	8	58	1,426	1,117	705	477	188	81	25	9	13
2002.....	4,050	1	20	64	1,380	1,070	773	411	200	65	47	9	10
2001.....	4,042	5	17	78	1,351	1,083	791	415	170	80	37	6	9
2000.....	3,686	2	9	81	1,224	955	724	406	177	70	25	8	5
1999.....	3,820	4	14	86	1,255	1,008	736	392	186	90	31	9	9

White, female

2007 1/.....	1,140	2	12	37	252	246	230	176	79	53	36	17	-
2006 1/.....	1,100	2	8	44	211	208	233	193	101	45	42	13	-
2005 1/.....	1,089	-	10	26	187	242	267	162	72	57	50	14	2
2004 1/.....	1,041	2	6	28	211	201	220	163	92	53	51	13	1
2003 1/.....	1,065	1	12	23	208	214	261	176	80	35	43	11	1
2002.....	1,135	2	13	42	220	231	254	185	84	51	44	8	1
2001.....	1,146	3	16	41	227	244	274	145	85	61	41	9	-
2000.....	1,120	3	8	33	210	237	261	161	83	64	39	16	5
1999.....	1,098	-	12	41	229	229	275	134	66	58	41	12	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

Black, both sexes

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
2007 1/.....	6,960	5	20	92	2,918	2,217	1,007	468	163	46	20	4	-
2006 1/.....	7,113	4	20	100	2,989	2,331	974	490	135	51	14	4	1
2005 1/.....	6,703	2	13	89	2,762	2,260	961	434	108	54	15	5	-
2004 1/.....	6,201	3	22	81	2,500	2,089	893	414	138	41	14	5	1
2003 1/.....	6,397	6	19	95	2,647	2,125	915	416	109	42	17	3	3
2002.....	6,285	6	14	81	2,589	2,082	945	394	109	40	21	2	2
2001.....	5,885	2	18	57	2,537	1,901	873	337	94	45	18	1	2
2000.....	5,699	7	8	69	2,434	1,812	863	354	85	45	17	4	1
1999.....	5,577	2	18	82	2,450	1,724	804	326	104	39	21	2	5

Black, male

2007 1/.....	6,281	3	8	69	2,709	2,045	857	395	141	36	17	1	-
2006 1/.....	6,374	2	11	75	2,760	2,111	823	420	113	42	13	3	1
2005 1/.....	6,067	2	9	65	2,568	2,080	827	364	91	44	12	5	-
2004 1/.....	5,589	1	11	62	2,315	1,912	779	347	111	39	7	4	1
2003 1/.....	5,740	6	11	67	2,443	1,937	786	352	92	33	10	1	2
2002.....	5,575	2	9	54	2,364	1,885	797	324	89	33	16	-	2
2001.....	5,279	1	9	44	2,373	1,723	708	284	80	39	15	1	2
2000.....	5,084	7	5	53	2,243	1,637	723	296	69	37	12	1	1
1999.....	4,883	1	7	61	2,227	1,537	643	260	94	31	16	1	5

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

Black, female

2007 1/.....	679	2	12	23	209	172	150	73	22	10	3	3	-
2006 1/.....	739	2	9	25	229	220	151	70	22	9	1	1	-
2005 1/.....	636	-	4	24	194	180	134	70	17	10	3	-	-
2004 1/.....	612	2	11	19	185	177	114	67	27	2	7	1	-
2003 1/.....	657	-	8	28	204	188	129	64	17	9	7	2	1
2002.....	710	4	5	27	225	197	148	70	20	7	5	2	-
2001.....	606	1	9	13	164	178	165	53	14	6	3	-	-
2000.....	615	-	3	16	191	175	140	58	16	8	5	3	-
1999.....	694	1	11	21	223	187	161	66	10	8	5	1	-

American Indian 2/, both sexes

2007 1/.....	91	-	2	3	29	30	15	10	1	1	-	-	-
2006 1/.....	119	-	-	2	52	33	18	8	3	2	1	-	-
2005 1/.....	117	-	-	4	57	34	12	4	4	2	-	-	-
2004 1/.....	104	-	2	6	38	29	19	7	2	-	1	-	-
2003 1/.....	109	-	-	4	49	25	17	10	3	-	1	-	-
2002.....	117	-	1	7	44	29	22	10	2	2	-	-	-
2001.....	87	1	4	2	29	27	12	6	3	3	-	-	-
2000.....	86	-	2	-	32	27	17	4	2	1	-	-	1
1999.....	104	2	2	6	33	30	11	12	4	3	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

American Indian 2/, male

2007 1/.....	78	-	1	3	26	27	12	7	1	1	-	-	-
2006 1/.....	105	-	-	2	50	29	12	6	3	2	1	-	-
2005 1/.....	96	-	-	1	50	29	9	2	4	1	-	-	-
2004 1/.....	87	-	2	4	31	24	18	5	2	-	1	-	-
2003 1/.....	81	-	-	2	38	22	10	8	1	-	-	-	-
2002.....	85	-	-	2	32	22	18	8	2	1	-	-	-
2001.....	64	1	1	1	24	20	8	3	3	3	-	-	-
2000.....	70	-	2	-	29	22	11	3	2	-	-	-	1
1999.....	82	-	2	4	29	24	7	8	4	3	1	-	-

American Indian 2/, female

2007 1/.....	13	-	1	-	3	3	3	3	-	-	-	-	-
2006 1/.....	14	-	-	-	2	4	6	2	-	-	-	-	-
2005 1/.....	21	-	-	3	7	5	3	2	-	1	-	-	-
2004 1/.....	17	-	-	2	7	5	1	2	-	-	-	-	-
2003 1/.....	28	-	-	2	11	3	7	2	2	-	1	-	-
2002.....	32	-	1	5	12	7	4	2	-	1	-	-	-
2001.....	23	-	3	1	5	7	4	3	-	-	-	-	-
2000.....	16	-	-	-	3	5	6	1	-	1	-	-	-
1999.....	22	2	-	2	4	6	4	4	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	201	-	-	3	63	50	38	28	12	5	1	-	1
2006 1/.....	280	-	-	6	110	65	31	47	16	3	1	1	-
2005 1/.....	266	-	1	7	103	53	49	29	19	5	-	-	-
2004 1/.....	200	-	1	5	59	43	29	37	17	8	1	-	-
2003 1/.....	241	-	1	7	80	59	43	31	14	3	2	-	1
2002.....	242	-	1	11	84	53	48	25	14	3	3	-	-
2001.....	188	-	-	2	56	53	28	31	12	3	3	-	-
2000.....	210	-	1	4	63	43	41	34	20	4	-	-	-
1999.....	229	-	4	9	71	57	42	29	13	3	1	-	-

Asian or Pacific Islander 3/, male

2007 1/.....	168	-	-	3	57	44	28	23	8	4	-	-	1
2006 1/.....	228	-	-	5	100	52	20	36	10	3	1	1	-
2005 1/.....	221	-	-	4	88	44	42	25	15	3	-	-	-
2004 1/.....	167	-	-	4	53	34	25	28	15	8	-	-	-
2003 1/.....	197	-	1	7	70	49	34	23	11	1	-	49	1
2002.....	189	-	1	7	75	40	35	18	10	2	1	-	-
2001.....	147	-	-	2	48	38	21	24	11	1	2	-	-
2000.....	166	-	1	4	52	36	29	25	18	1	-	-	-
1999.....	159	-	3	4	58	40	28	16	10	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

Asian or Pacific Islander 3/, female

2007 1/.....	33	-	-	-	6	6	10	5	4	1	1	-	-
2006 1/.....	52	-	-	1	10	13	11	11	6	-	-	-	-
2005 1/.....	45	-	1	3	15	9	7	4	4	2	-	-	-
2004 1/.....	33	-	1	1	6	9	4	9	2	-	1	-	-
2003 1/.....	44	-	-	-	10	10	9	8	3	2	2	-	-
2002.....	53	-	-	4	9	13	13	7	4	1	2	-	-
2001.....	41	-	-	-	8	15	7	7	1	2	1	-	-
2000.....	44	-	-	-	11	7	12	9	2	3	-	-	-
1999.....	70	-	1	5	13	17	14	13	3	3	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02.X85-X92,X96-Y09,Y87.1)

All races, both sexes

2007.....	5,729	337	350	145	882	1,007	1,014	981	534	226	180	57	16
2006.....	5,782	330	324	153	890	958	1,033	1,024	544	240	195	80	11
2005.....	5,772	300	338	154	967	972	1,099	963	457	239	186	85	12
2004.....	5,733	319	341	145	958	992	1,089	946	411	253	189	75	15
2003.....	5,812	333	336	137	958	976	1,169	907	392	271	229	81	23
2002.....	5,809	294	374	151	902	1,024	1,197	890	432	260	181	76	28
2001 4/.....	8,960	321	360	146	1,097	1,896	2,290	1,533	654	340	213	89	21
2000.....	5,964	337	328	184	976	1,090	1,313	796	371	259	220	72	18
1999.....	6,061	323	326	208	960	1,183	1,338	775	349	280	209	76	34

All races, male

2007.....	3,771	195	190	69	630	695	644	670	386	158	101	23	10
2006.....	3,831	197	183	79	616	661	683	719	399	156	100	30	8
2005.....	3,815	167	192	73	669	664	718	720	324	151	94	34	9
2004.....	3,657	165	186	66	662	653	668	671	290	164	97	26	9
2003.....	3,756	198	182	81	647	616	773	628	276	170	136	36	13
2002.....	3,741	161	204	70	623	652	763	633	313	176	94	32	20
2001 4/.....	6,023	190	211	68	745	1,293	1,594	1,106	459	209	101	35	12
2000.....	3,814	194	181	89	655	711	832	554	271	171	117	31	8
1999.....	3,841	182	168	95	622	737	893	553	247	180	114	31	19

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

All races, female

2007.....	1,958	142	160	76	252	312	370	311	148	68	79	34	6
2006.....	1,951	133	141	74	274	297	350	305	145	84	95	50	3
2005.....	1,957	133	146	81	298	308	381	243	133	88	92	51	3
2004.....	2,076	154	155	79	296	339	421	275	121	89	92	49	6
2003.....	2,056	135	154	56	311	360	396	279	116	101	93	45	10
2002.....	2,068	133	170	81	279	372	434	257	119	84	87	44	8
2001 4/.....	2,937	131	149	78	352	603	696	427	195	131	112	54	9
2000.....	2,150	143	147	95	321	379	481	242	100	88	103	41	10
1999.....	2,220	141	158	113	338	446	445	222	102	100	95	45	15

White, both sexes

2007 1/.....	3,534	218	190	93	534	596	599	618	346	156	129	42	13
2006 1/.....	3,581	218	168	98	537	564	605	643	342	173	157	66	10
2005 1/.....	3,504	191	194	81	595	589	627	552	299	155	141	69	11
2004 1/.....	3,524	205	190	92	596	578	638	566	271	179	135	60	14
2003 1/.....	3,536	214	184	78	537	580	687	542	262	188	176	66	22
2002.....	3,500	167	206	89	537	564	712	521	303	190	127	59	25
2001 4/.....	6,140	188	195	99	686	1,296	1,607	1,079	489	251	162	73	15
2000.....	3,533	192	182	126	547	622	740	481	237	169	163	59	15
1999.....	3,710	196	182	134	580	689	755	490	244	205	153	58	24

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

White, male

2007 1/.....	2,301	122	102	40	385	417	369	426	239	109	68	15	9
2006 1/.....	2,335	121	95	45	359	390	400	463	245	105	79	26	7
2005 1/.....	2,280	105	106	35	410	396	413	413	208	94	68	23	9
2004 1/.....	2,224	111	108	45	408	360	396	398	190	114	66	20	8
2003 1/.....	2,229	128	88	48	355	370	442	364	176	113	103	30	12
2002.....	2,232	99	113	34	366	365	467	346	209	128	61	25	19
2001 4/.....	4,212	113	115	41	460	905	1,181	781	348	153	81	27	7
2000.....	2,239	123	105	67	359	401	478	312	173	106	84	23	8
1999.....	2,342	117	94	62	367	437	504	345	170	126	86	20	14

White, female

2007 1/.....	1,233	96	88	53	149	179	230	192	107	47	61	27	4
2006 1/.....	1,246	97	73	53	178	174	205	180	97	68	78	40	3
2005 1/.....	1,224	86	88	46	185	193	214	139	91	61	73	46	2
2004 1/.....	1,300	94	82	47	188	218	242	168	81	65	69	40	6
2003 1/.....	1,307	86	96	30	182	210	245	178	86	75	73	36	10
2002.....	1,268	68	93	55	171	199	245	175	94	62	66	34	6
2001 4/.....	1,928	75	80	58	226	391	426	298	141	98	81	46	8
2000.....	1,294	69	77	59	188	221	262	169	64	63	79	36	7
1999.....	1,368	79	88	72	213	252	251	145	74	79	67	38	10

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

Black, both sexes

2007 1/.....	1,910	105	148	48	295	343	359	334	156	60	46	13	3
2006 1/.....	1,919	105	134	49	306	339	373	335	174	58	31	14	1
2005 1/.....	1,966	93	127	61	315	331	408	363	140	71	43	13	1
2004 1/.....	1,934	100	141	50	303	356	404	338	116	63	49	13	1
2003 1/.....	1,995	100	137	54	359	350	430	324	114	64	48	14	1
2002.....	2,002	111	147	54	311	387	432	328	109	57	50	15	1
2001 4/.....	2,341	121	151	37	333	470	577	382	131	74	45	14	6
2000.....	2,168	133	137	55	385	412	513	274	119	80	47	10	3
1999.....	2,071	116	134	65	331	433	521	243	91	65	46	17	9

Black, male

2007 1/.....	1,303	67	81	27	209	238	243	228	127	43	32	7	1
2006 1/.....	1,303	70	73	30	222	235	245	230	132	44	17	4	1
2005 1/.....	1,345	52	79	32	222	239	264	269	105	49	25	9	-
2004 1/.....	1,250	47	72	20	207	251	243	245	84	44	30	6	1
2003 1/.....	1,343	54	85	32	246	219	301	232	90	46	31	6	1
2002.....	1,321	54	83	32	218	244	261	258	92	41	32	6	-
2001 4/.....	1,501	69	92	24	227	305	343	272	90	48	19	7	5
2000.....	1,398	66	70	19	258	274	317	212	91	58	27	6	-
1999.....	1,331	59	68	30	226	263	353	179	66	49	23	10	5

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

Black, female

2007 1/.....	607	38	67	21	86	105	116	106	29	17	14	6	2
2006 1/.....	616	35	61	19	84	104	128	105	42	14	14	10	-
2005 1/.....	621	41	48	29	93	92	144	94	35	22	18	4	1
2004 1/.....	684	53	69	30	96	105	161	93	32	19	19	7	-
2003 1/.....	652	46	52	22	113	131	129	92	24	18	17	8	-
2002.....	681	57	64	22	93	143	171	70	17	16	18	9	1
2001 4/.....	840	52	59	13	106	165	234	110	41	26	26	7	1
2000.....	770	67	67	36	127	138	196	62	28	22	20	4	3
1999.....	740	57	66	35	105	170	168	64	25	16	23	7	4

American Indian 2/, both sexes

2007 1/.....	129	9	3	-	28	35	28	13	7	3	2	1	-
2006 1/.....	135	4	11	-	25	34	30	22	7	1	1	-	-
2005 1/.....	140	5	10	3	25	26	33	25	7	6	-	-	-
2004 1/.....	124	9	5	-	29	25	27	15	6	6	2	-	-
2003 1/.....	128	10	7	-	29	26	27	22	1	5	-	1	-
2002.....	150	10	11	4	26	34	30	19	7	5	3	-	1
2001 4/.....	124	5	8	2	30	27	24	20	2	3	2	1	-
2000.....	117	5	6	1	19	32	29	14	4	3	2	2	-
1999.....	149	3	4	3	27	43	33	21	7	4	4	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

American Indian 2/, male

2007 1/.....	85	4	2	-	19	25	20	9	5	1	-	-	-
2006 1/.....	101	4	9	-	19	24	22	15	6	1	1	-	-
2005 1/.....	97	4	4	2	17	16	23	23	5	3	-	-	-
2004 1/.....	94	5	3	-	22	23	20	13	3	5	-	-	-
2003 1/.....	94	8	5	-	20	19	19	18	1	4	-	-	-
2002.....	100	4	5	1	22	24	22	15	3	3	1	-	-
2001 4/.....	82	2	4	-	21	19	16	16	2	1	1	-	-
2000.....	89	3	4	1	17	24	21	11	2	3	1	2	-
1999.....	97	3	2	-	16	26	23	16	5	3	3	-	-

American Indian 2/, female

2007 1/.....	44	5	1	-	9	10	8	4	2	2	2	1	-
2006 1/.....	34	-	2	-	6	10	8	7	1	-	-	-	-
2005 1/.....	43	1	6	1	8	10	10	2	2	3	-	-	-
2004 1/.....	30	4	2	-	7	2	7	2	3	1	2	-	-
2003 1/.....	34	2	2	-	9	7	8	4	-	1	-	1	-
2002.....	50	6	6	3	4	10	8	4	4	2	2	-	1
2001 4/.....	42	3	4	2	9	8	8	4	-	2	1	1	-
2000.....	28	2	2	-	2	8	8	3	2	-	1	-	-
1999.....	52	-	2	3	11	17	10	5	2	1	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	156	5	9	4	25	33	28	16	25	7	3	1	-
2006 1/.....	147	3	11	6	22	21	25	24	21	8	6	-	-
2005 1/.....	162	11	7	9	32	26	31	23	11	7	2	3	-
2004 1/.....	151	5	5	3	30	33	20	27	18	5	3	2	-
2003 1/.....	153	9	8	5	33	20	25	19	15	14	5	-	-
2002.....	157	6	10	4	28	39	23	22	13	8	1	2	1
2001 4/.....	355	7	6	8	48	103	82	52	32	12	4	1	-
2000.....	146	7	3	2	25	24	31	27	11	7	8	1	-
1999.....	131	8	6	6	22	18	29	21	7	6	6	1	1

Asian or Pacific Islander 3/, male

2007 1/.....	82	2	5	2	17	15	12	7	15	5	1	1	-
2006 1/.....	92	2	6	4	16	12	16	11	16	6	3	-	-
2005 1/.....	93	6	3	4	20	13	18	15	6	5	1	2	-
2004 1/.....	89	2	3	1	25	19	9	15	13	1	1	-	-
2003 1/.....	90	8	4	1	26	8	11	14	9	7	2	-	-
2002.....	88	4	3	3	17	19	13	14	9	4	-	1	1
2001 4/.....	228	6	-	3	37	64	54	37	19	7	-	1	-
2000.....	88	2	2	2	21	12	16	19	5	4	5	-	-
1999.....	71	3	4	3	13	11	13	13	6	2	2	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

Asian or Pacific Islander 3/, female

2007 1/.....	74	3	4	2	8	18	16	9	10	2	2	-	-
2006 1/.....	55	1	5	2	6	9	9	13	5	2	3	-	-
2005 1/.....	69	5	4	5	12	13	13	8	5	2	1	1	-
2004 1/.....	62	3	2	2	5	14	11	12	5	4	2	2	-
2003 1/.....	63	1	4	4	7	12	14	5	6	7	3	-	-
2002.....	69	2	7	1	11	20	10	8	4	4	1	1	-
2001 4/.....	127	1	6	5	11	39	28	15	13	5	4	-	-
2000.....	58	5	1	-	4	12	15	8	6	3	3	1	-
1999.....	60	5	2	3	9	7	16	8	1	4	4	-	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Legal intervention (Y35,Y89.0)

All races, both sexes

2007.....	412	-	-	1	97	127	96	64	19	5	2	1	-
2006.....	434	-	-	1	85	141	115	66	17	7	1	1	-
2005.....	414	-	-	1	83	110	124	56	30	7	3	-	-
2004.....	372	-	-	-	80	106	104	58	12	7	5	-	-
2003.....	423	-	-	2	93	135	117	51	19	4	2	-	-
2002.....	384	-	-	3	91	102	109	56	16	6	1	-	-
2001.....	396	-	-	-	88	107	115	51	24	6	4	1	-
2000.....	359	-	-	3	73	110	101	55	10	5	1	1	-
1999.....	398	-	-	1	67	136	121	50	20	2	1	-	-

All races, male

2007.....	400	-	-	1	94	126	91	61	19	5	2	1	-
2006.....	417	-	-	1	82	139	112	61	15	6	1	-	-
2005.....	401	-	-	1	82	109	119	54	28	5	3	-	-
2004.....	363	-	-	-	79	104	103	55	11	6	5	-	-
2003.....	405	-	-	1	89	131	114	46	18	4	2	-	-
2002.....	369	-	-	3	87	99	105	52	16	6	1	-	-
2001.....	380	-	-	-	84	104	108	51	22	6	4	1	-
2000.....	348	-	-	3	71	108	96	54	9	5	1	1	-
1999.....	392	-	-	1	67	135	117	49	20	2	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Legal intervention (Y35,Y89.0)

All races, female

2007.....	12	-	-	-	3	1	5	3	-	-	-	-	-
2006.....	17	-	-	-	3	2	3	5	2	1	-	1	-
2005.....	13	-	-	-	1	1	5	2	2	2	-	-	-
2004.....	9	-	-	-	1	2	1	3	1	1	-	-	-
2003.....	18	-	-	1	4	4	3	5	1	-	-	-	-
2002.....	15	-	-	-	4	3	4	4	-	-	-	-	-
2001.....	16	-	-	-	4	3	7	-	2	-	-	-	-
2000.....	11	-	-	-	2	2	5	1	1	-	-	-	-
1999.....	6	-	-	-	-	1	4	1	-	-	-	-	-

White, both sexes

2007 1/.....	276	-	-	1	47	82	76	49	13	5	2	1	-
2006 1/.....	300	-	-	-	52	91	82	53	14	7	1	-	-
2005 1/.....	289	-	-	-	59	68	85	43	26	6	2	-	-
2004 1/.....	262	-	-	-	55	66	81	42	8	5	5	-	-
2003 1/.....	284	-	-	1	54	76	90	41	17	4	1	-	-
2002.....	257	-	-	1	49	69	78	41	13	5	1	-	-
2001.....	292	-	-	-	60	78	86	40	18	6	4	-	-
2000.....	237	-	-	1	44	72	63	43	9	3	1	1	-
1999.....	255	-	-	-	31	86	80	40	15	2	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Legal intervention (Y35,Y89.0)

White, male

2007 1/.....	267	-	-	1	46	81	71	47	13	5	2	1	-
2006 1/.....	286	-	-	-	50	89	79	49	12	6	1	-	-
2005 1/.....	277	-	-	-	58	67	81	41	24	4	2	-	-
2004 1/.....	256	-	-	-	54	65	80	40	8	4	5	-	-
2003 1/.....	271	-	-	-	52	74	87	37	16	4	1	-	-
2002.....	246	-	-	1	46	67	76	37	13	5	1	-	-
2001.....	283	-	-	-	58	77	81	40	17	6	4	-	-
2000.....	231	-	-	1	42	70	62	43	8	3	1	1	-
1999.....	250	-	-	-	31	86	76	39	15	2	1	-	-

White, female

2007 1/.....	9	-	-	-	1	1	5	2	-	-	-	-	-
2006 1/.....	14	-	-	-	2	2	3	4	2	1	-	-	-
2005 1/.....	12	-	-	-	1	1	4	2	2	2	-	-	-
2004 1/.....	6	-	-	-	1	1	1	2	-	1	-	-	-
2003 1/.....	13	-	-	1	2	2	3	4	1	-	-	-	-
2002.....	11	-	-	-	3	2	2	4	-	-	-	-	-
2001.....	9	-	-	-	2	1	5	-	1	-	-	-	-
2000.....	6	-	-	-	2	2	1	-	1	-	-	-	-
1999.....	5	-	-	-	-	-	4	1	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Legal intervention (Y35,Y89.0)

Black, both sexes

2007 1/.....	127	-	-	-	48	43	17	13	6	-	-	-	-
2006 1/.....	117	-	-	1	27	44	29	12	3	-	-	1	-
2005 1/.....	112	-	-	1	24	39	31	13	2	1	1	-	-
2004 1/.....	92	-	-	-	20	34	19	13	4	2	-	-	-
2003 1/.....	120	-	-	1	35	53	22	7	1	-	1	-	-
2002.....	110	-	-	2	37	29	28	11	2	1	-	-	-
2001.....	84	-	-	-	26	21	23	10	3	-	-	1	-
2000.....	111	-	-	2	28	33	35	11	-	2	-	-	-
1999.....	126	-	-	1	32	42	36	10	5	-	-	-	-

Black, male

2007 1/.....	124	-	-	-	46	43	17	12	6	-	-	-	-
2006 1/.....	114	-	-	1	26	44	29	11	3	-	-	-	-
2005 1/.....	111	-	-	1	24	39	30	13	2	1	1	-	-
2004 1/.....	90	-	-	-	20	34	19	12	3	2	-	-	-
2003 1/.....	117	-	-	1	33	52	22	7	1	-	1	-	-
2002.....	106	-	-	2	36	28	26	11	2	1	-	-	-
2001.....	79	-	-	-	24	20	21	10	3	-	-	1	-
2000.....	108	-	-	2	28	33	33	10	-	2	-	-	-
1999.....	125	-	-	1	32	41	36	10	5	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Legal intervention (Y35,Y89.0)

Black, female

2007 1/.....	3	-	-	-	2	-	-	1	-	-	-	-	-
2006 1/.....	3	-	-	-	1	-	-	1	-	-	-	1	-
2005 1/.....	1	-	-	-	-	-	1	-	-	-	-	-	-
2004 1/.....	2	-	-	-	-	-	-	1	1	-	-	-	-
2003 1/.....	3	-	-	-	2	1	-	-	-	-	-	-	-
2002.....	4	-	-	-	1	1	2	-	-	-	-	-	-
2001.....	5	-	-	-	2	1	2	-	-	-	-	-	-
2000.....	3	-	-	-	-	-	2	1	-	-	-	-	-
1999.....	1	-	-	-	-	1	-	-	-	-	-	-	-

American Indian 2/, both sexes

2007 1/.....	3	-	-	-	1	1	1	-	-	-	-	-	-
2006 1/.....	9	-	-	-	3	3	3	-	-	-	-	-	-
2005 1/.....	7	-	-	-	-	2	4	-	1	-	-	-	-
2004 1/.....	5	-	-	-	1	1	3	-	-	-	-	-	-
2003 1/.....	12	-	-	-	3	4	3	1	1	-	-	-	-
2002.....	9	-	-	-	3	2	1	3	-	-	-	-	-
2001.....	11	-	-	-	2	2	5	-	2	-	-	-	-
2000.....	4	-	-	-	1	1	1	1	-	-	-	-	-
1999.....	10	-	-	-	1	5	4	-	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Legal intervention (Y35,Y89.0)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	6	-	-	-	1	1	2	2	-	-	-	-	-
2006 1/.....	8	-	-	-	3	3	1	1	-	-	-	-	-
2005 1/.....	6	-	-	-	-	1	4	-	1	-	-	-	-
2004 1/.....	13	-	-	-	4	5	1	3	-	-	-	-	-
2003 1/.....	7	-	-	-	1	2	2	2	-	-	-	-	-
2002.....	8	-	-	-	2	2	2	1	1	-	-	-	-
2001.....	9	-	-	-	-	6	1	1	1	-	-	-	-
2000.....	7	-	-	-	-	4	2	-	1	-	-	-	-
1999.....	7	-	-	-	3	3	1	-	-	-	-	-	-

Asian or Pacific Islander 3/, male

2007 1/.....	6	-	-	-	1	1	2	2	-	-	-	-	-
2006 1/.....	8	-	-	-	3	3	1	1	-	-	-	-	-
2005 1/.....	6	-	-	-	-	1	4	-	1	-	-	-	-
2004 1/.....	12	-	-	-	4	4	1	3	-	-	-	-	-
2003 1/.....	5	-	-	-	1	1	2	1	-	-	-	-	-
2002.....	8	-	-	-	2	2	2	1	1	-	-	-	-
2001.....	9	-	-	-	-	6	1	1	1	-	-	-	-
2000.....	6	-	-	-	-	4	1	-	1	-	-	-	-
1999.....	7	-	-	-	3	3	1	-	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

All races, both sexes

2007.....	5,381	88	49	49	579	872	1,251	1,501	599	183	132	70	8
2006.....	5,131	92	52	59	602	817	1,278	1,362	492	163	136	72	6
2005.....	4,742	104	69	64	484	742	1,194	1,314	424	157	106	74	10
2004.....	4,976	104	47	64	501	806	1,389	1,302	399	156	119	80	9
2003.....	5,072	75	31	55	563	835	1,581	1,267	368	131	96	52	18
2002.....	4,830	101	57	68	487	828	1,516	1,140	320	139	95	65	14
2001.....	4,198	83	44	51	412	728	1,396	990	225	99	108	44	18
2000.....	3,819	65	39	60	370	679	1,286	841	228	104	79	57	11
1999.....	3,917	55	43	49	430	743	1,295	793	222	114	96	58	19

All races, male

2007.....	3,351	55	26	31	445	569	787	859	357	109	78	28	7
2006.....	3,249	57	29	43	452	569	767	832	289	92	76	37	6
2005.....	2,861	67	44	43	355	478	695	758	245	91	49	26	10
2004.....	3,164	65	26	43	380	559	872	808	231	91	52	28	9
2003.....	3,295	50	17	41	412	580	1,013	811	222	65	49	21	14
2002.....	3,114	52	28	38	358	585	981	696	203	83	46	31	13
2001.....	2,833	49	19	38	316	515	931	662	146	64	57	20	16
2000.....	2,635	27	26	50	285	489	885	570	152	70	49	21	11
1999.....	2,692	33	21	38	345	523	887	532	148	72	55	25	13

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

All races, female

2007.....	2,030	33	23	18	134	303	464	642	242	74	54	42	1
2006.....	1,882	35	23	16	150	248	511	530	203	71	60	35	-
2005.....	1,881	37	25	21	129	264	499	556	179	66	57	48	-
2004.....	1,812	39	21	21	121	247	517	494	168	65	67	52	-
2003.....	1,777	25	14	14	151	255	568	456	146	66	47	31	4
2002.....	1,716	49	29	30	129	243	535	444	117	56	49	34	1
2001.....	1,365	34	25	13	96	213	465	328	79	35	51	24	2
2000.....	1,184	38	13	10	85	190	401	271	76	34	30	36	-
1999.....	1,225	22	22	11	85	220	408	261	74	42	41	33	6

White, both sexes

2007 1/.....	4,449	54	36	37	477	754	1,042	1,240	482	151	106	65	5
2006 1/.....	4,220	63	28	41	521	696	1,057	1,114	401	133	107	54	5
2005 1/.....	3,917	68	43	47	406	629	975	1,093	360	135	88	67	6
2004 1/.....	4,142	68	30	36	432	677	1,163	1,088	342	123	106	72	5
2003 1/.....	4,255	52	21	37	488	719	1,333	1,057	299	103	85	50	11
2002.....	3,962	56	45	49	418	679	1,252	925	272	115	79	60	12
2001.....	3,453	45	31	39	351	612	1,146	805	189	83	96	41	15
2000.....	3,053	41	27	38	302	553	1,031	672	177	84	72	48	8
1999.....	3,122	36	23	42	342	606	1,023	612	186	97	89	52	14

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

White, male

2007 1/.....	2,721	34	17	20	364	491	649	688	281	86	59	27	5
2006 1/.....	2,616	41	12	29	391	479	618	658	228	70	55	30	5
2005 1/.....	2,327	45	28	32	299	402	566	613	197	76	38	25	6
2004 1/.....	2,607	44	16	24	329	471	724	659	191	72	46	26	5
2003 1/.....	2,709	30	11	29	351	489	848	656	171	51	45	20	8
2002.....	2,524	29	23	26	306	477	805	548	169	62	39	29	11
2001.....	2,283	28	13	29	263	431	745	520	117	52	53	19	13
2000.....	2,078	18	17	32	228	392	710	446	112	53	45	17	8
1999.....	2,105	22	13	33	278	416	698	390	113	61	49	23	9

White, female

2007 1/.....	1,728	20	19	17	113	263	393	552	201	65	47	38	-
2006 1/.....	1,604	22	16	12	130	217	439	456	173	63	52	24	-
2005 1/.....	1,590	23	15	15	107	227	409	480	163	59	50	42	-
2004 1/.....	1,535	24	14	12	103	206	439	429	151	51	60	46	-
2003 1/.....	1,546	22	10	8	137	230	485	401	128	52	40	30	3
2002.....	1,438	27	22	23	112	202	447	377	103	53	40	31	1
2001.....	1,170	17	18	10	88	181	401	285	72	31	43	22	2
2000.....	975	23	10	6	74	161	321	226	65	31	27	31	-
1999.....	1,017	14	10	9	64	190	325	222	73	36	40	29	5

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Black, both sexes

2007 1/.....	760	29	12	11	70	90	168	229	98	28	20	2	3
2006 1/.....	718	23	22	16	57	87	184	205	75	24	15	9	1
2005 1/.....	685	31	23	11	56	92	189	190	52	18	13	7	3
2004 1/.....	702	32	15	26	56	100	191	191	50	22	10	5	4
2003 1/.....	687	21	9	14	51	96	216	190	53	21	7	2	7
2002.....	759	37	8	15	57	125	244	196	39	17	16	4	1
2001.....	646	36	13	11	44	93	227	168	32	12	5	3	2
2000.....	668	15	11	21	52	105	234	156	43	15	6	7	3
1999.....	696	18	16	7	71	121	240	168	28	14	5	3	5

Black, male

2007 1/.....	527	18	9	10	60	60	114	149	70	20	15	-	2
2006 1/.....	508	12	15	12	45	66	126	149	51	18	11	2	1
2005 1/.....	453	19	14	8	41	62	114	131	40	12	8	1	3
2004 1/.....	479	18	9	18	43	73	124	133	38	14	5	-	4
2003 1/.....	498	19	5	8	43	77	144	141	40	12	2	1	6
2002.....	521	18	3	11	43	90	164	137	30	16	7	1	1
2001.....	480	20	6	8	38	67	170	130	27	10	1	1	2
2000.....	488	3	8	17	47	80	162	115	33	13	4	3	3
1999.....	523	10	8	5	56	93	171	134	27	10	4	1	4

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Black, female

2007 1/.....	233	11	3	1	10	30	54	80	28	8	5	2	1
2006 1/.....	210	11	7	4	12	21	58	56	24	6	4	7	-
2005 1/.....	232	12	9	3	15	30	75	59	12	6	5	6	-
2004 1/.....	223	14	6	8	13	27	67	58	12	8	5	5	-
2003 1/.....	189	2	4	6	8	19	72	49	13	9	5	1	1
2002.....	238	19	5	4	14	35	80	59	9	1	9	3	-
2001.....	166	16	7	3	6	26	57	38	5	2	4	2	-
2000.....	180	12	3	4	5	25	72	41	10	2	2	4	-
1999.....	173	8	8	2	15	28	69	34	1	4	1	2	1

American Indian 2/, both sexes

2007 1/.....	92	4	-	-	20	14	22	13	13	2	3	1	-
2006 1/.....	95	2	2	2	14	17	23	22	9	2	2	-	-
2005 1/.....	64	3	2	1	9	9	11	17	7	2	2	-	1
2004 1/.....	68	3	-	2	6	16	23	10	4	3	1	-	-
2003 1/.....	64	2	-	2	8	11	21	11	7	1	1	-	-
2002.....	48	3	2	2	5	12	8	9	4	2	-	-	1
2001.....	43	1	-	-	7	7	14	9	1	1	2	-	1
2000.....	46	4	1	1	5	9	12	7	6	1	-	-	-
1999.....	54	1	2	-	9	11	18	6	3	2	1	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

American Indian 2/, male

2007 1/.....	51	2	-	-	12	8	12	9	3	2	2	1	-
2006 1/.....	62	1	2	2	8	15	15	10	6	1	2	-	-
2005 1/.....	37	1	2	-	6	5	5	9	5	1	2	-	1
2004 1/.....	41	3	-	1	5	9	15	5	1	2	-	-	-
2003 1/.....	47	1	-	2	7	7	16	6	7	1	-	-	-
2002.....	26	1	1	1	5	7	3	4	2	1	-	-	1
2001.....	32	-	-	-	7	7	9	5	1	-	2	-	1
2000.....	36	3	1	1	3	8	9	5	6	-	-	-	-
1999.....	37	1	-	-	7	10	10	4	3	1	1	-	-

American Indian 2/, female

2007 1/.....	41	2	-	-	8	6	10	4	10	-	1	-	-
2006 1/.....	33	1	-	-	6	2	8	12	3	1	-	-	-
2005 1/.....	27	2	-	1	3	4	6	8	2	1	-	-	-
2004 1/.....	27	-	-	1	1	7	8	5	3	1	1	-	-
2003 1/.....	17	1	-	-	1	4	5	5	-	-	1	-	-
2002.....	22	2	1	1	-	5	5	5	2	1	-	-	-
2001.....	11	1	-	-	-	-	5	4	-	1	-	-	-
2000.....	10	1	-	-	2	1	3	2	-	1	-	-	-
1999.....	17	-	2	-	2	1	8	2	-	1	-	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	80	1	1	1	12	14	19	19	6	2	3	2	-
2006 1/.....	98	4	-	-	10	17	14	21	7	4	12	9	-
2005 1/.....	76	2	1	5	13	12	19	14	5	2	3	-	-
2004 1/.....	64	1	2	-	7	13	12	13	3	8	2	3	-
2003 1/.....	66	-	1	2	16	9	11	9	9	6	3	-	-
2002.....	61	5	2	2	7	12	12	10	5	5	-	1	-
2001.....	56	1	-	1	10	16	9	8	3	3	5	-	-
2000.....	52	5	-	-	11	12	9	6	2	4	1	2	-
1999.....	45	-	2	-	8	5	14	7	5	1	1	2	-

Asian or Pacific Islander 3/, male

2007 1/.....	52	1	-	1	9	10	12	13	3	1	2	-	-
2006 1/.....	63	3	-	-	8	9	8	15	4	3	8	5	-
2005 1/.....	44	2	-	3	9	9	10	5	3	2	1	-	-
2004 1/.....	37	-	1	-	3	6	9	11	1	3	1	2	-
2003 1/.....	41	-	1	2	11	7	5	8	4	1	2	-	-
2002.....	43	4	1	-	4	11	9	7	2	4	-	1	-
2001.....	38	1	-	1	8	10	7	7	1	2	1	-	-
2000.....	33	3	-	-	7	9	4	4	1	4	-	1	-
1999.....	27	-	-	-	4	4	8	4	5	-	1	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Asian or Pacific Islander 3/, female

2007 1/.....	28	-	1	-	3	4	7	6	3	1	1	2	-
2006 1/.....	35	1	-	-	2	8	6	6	3	1	4	4	-
2005 1/.....	32	-	1	2	4	3	9	9	2	-	2	-	-
2004 1/.....	27	1	1	-	4	7	3	2	2	5	1	1	-
2003 1/.....	25	-	-	-	5	2	6	1	5	5	1	-	-
2002.....	18	1	1	2	3	1	3	3	3	1	-	-	-
2001.....	18	-	-	-	2	6	2	1	2	1	4	-	-
2000.....	19	2	-	-	4	3	5	2	1	-	1	1	-
1999.....	18	-	2	-	4	1	6	3	-	1	-	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Discharge of firearms, undetermined intent (Y22-Y24)

All races, both sexes

2007.....	276	-	3	12	82	44	36	40	35	17	6	1	-
2006.....	220	-	2	5	76	45	39	22	20	8	3	-	-
2005.....	221	-	3	12	77	35	28	28	23	10	3	2	-
2004.....	235	-	1	9	58	45	44	29	20	16	8	2	3
2003.....	232	-	1	12	69	38	43	31	20	8	9	1	-
2002.....	243	-	1	6	79	46	50	29	12	15	5	-	-
2001.....	231	-	-	6	64	48	50	29	12	10	11	1	-
2000.....	230	-	-	10	71	52	39	23	15	11	6	2	1
1999.....	324	-	3	12	127	51	57	28	14	15	13	4	-

All races, male

2007.....	223	-	2	9	71	36	29	30	27	13	5	1	-
2006.....	181	-	2	5	68	37	31	16	14	5	3	-	-
2005.....	178	-	2	10	68	29	21	22	14	7	3	2	-
2004.....	179	-	-	7	50	33	30	22	14	13	6	1	3
2003.....	186	-	1	12	60	33	26	22	16	7	8	1	-
2002.....	199	-	1	5	71	39	39	23	7	10	4	-	-
2001.....	190	-	-	6	59	42	33	21	11	8	9	1	-
2000.....	190	-	-	9	62	43	30	18	11	9	5	2	1
1999.....	277	-	3	12	111	44	45	21	11	14	12	4	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Discharge of firearms, undetermined intent (Y22-Y24)

All races, female

2007.....	53	-	1	3	11	8	7	10	8	4	1	-	-
2006.....	39	-	-	-	8	8	8	6	6	3	-	-	-
2005.....	43	-	1	2	9	6	7	6	9	3	-	-	-
2004.....	56	-	1	2	8	12	14	7	6	3	2	1	-
2003.....	46	-	-	-	9	5	17	9	4	1	1	-	-
2002.....	44	-	-	1	8	7	11	6	5	5	1	-	-
2001.....	41	-	-	-	5	6	17	8	1	2	2	-	-
2000.....	40	-	-	1	9	9	9	5	4	2	1	-	-
1999.....	47	-	-	-	16	7	12	7	3	1	1	-	-

White, both sexes

2007 1/.....	208	-	2	9	43	34	33	34	32	15	5	1	-
2006 1/.....	169	-	1	4	52	30	32	19	20	8	3	-	-
2005 1/.....	175	-	-	10	57	28	22	24	21	9	2	2	-
2004 1/.....	187	-	-	4	42	31	39	25	19	15	8	2	2
2003 1/.....	183	-	-	7	47	27	38	29	17	8	9	1	-
2002.....	192	-	-	4	59	33	41	27	10	14	4	-	-
2001.....	188	-	-	4	44	40	45	25	9	10	10	1	-
2000.....	167	-	-	5	48	34	31	20	11	9	6	2	1
1999.....	247	-	1	11	85	36	50	23	12	13	12	4	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Discharge of firearms, undetermined intent (Y22-Y24)

White, male

2007 1/.....	166	-	1	6	37	27	26	27	25	12	4	1	-
2006 1/.....	133	-	1	4	44	23	25	14	14	5	3	-	-
2005 1/.....	141	-	-	8	49	22	17	20	14	7	2	2	-
2004 1/.....	142	-	-	4	37	22	26	18	14	12	6	1	2
2003 1/.....	144	-	-	7	40	23	22	21	15	7	8	1	-
2002.....	153	-	-	3	51	27	32	21	7	9	3	-	-
2001.....	150	-	-	4	40	35	29	17	8	8	8	1	-
2000.....	137	-	-	5	42	28	25	15	7	7	5	2	1
1999.....	212	-	1	11	76	31	41	16	9	12	11	4	-

White, female

2007 1/.....	42	-	1	3	6	7	7	7	7	3	1	-	-
2006 1/.....	36	-	-	-	8	7	7	5	6	3	-	-	-
2005 1/.....	34	-	-	2	8	6	5	4	7	2	-	-	-
2004 1/.....	45	-	-	-	5	9	13	7	5	3	2	1	-
2003 1/.....	39	-	-	-	7	4	16	8	2	1	1	-	-
2002.....	39	-	-	1	8	6	9	6	3	5	1	-	-
2001.....	38	-	-	-	4	5	16	8	1	2	2	-	-
2000.....	30	-	-	-	6	6	6	5	4	2	1	-	-
1999.....	35	-	-	-	9	5	9	7	3	1	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Discharge of firearms, undetermined intent (Y22-Y24)

Black, both sexes

2007 1/.....	57	-	1	2	33	9	1	6	2	2	1	-	-
2006 1/.....	48	-	1	1	23	13	7	3	-	-	-	-	-
2005 1/.....	36	-	2	1	16	6	5	3	2	1	-	-	-
2004 1/.....	39	-	1	4	15	9	4	3	1	1	-	-	1
2003 1/.....	35	-	1	4	17	7	3	2	1	-	-	-	-
2002.....	41	-	1	2	17	11	6	1	1	1	1	-	-
2001.....	32	-	-	2	15	5	2	4	3	-	1	-	-
2000.....	54	-	-	5	20	15	7	2	3	2	-	-	-
1999.....	66	-	2	1	35	14	6	4	1	2	1	-	-

Black, male

2007 1/.....	46	-	1	2	28	8	1	3	1	1	1	-	-
2006 1/.....	45	-	1	1	23	12	6	2	-	-	-	-	-
2005 1/.....	28	-	1	1	15	6	4	1	-	-	-	-	-
2004 1/.....	31	-	-	3	12	8	3	3	-	1	-	-	1
2003 1/.....	32	-	1	4	16	7	2	1	1	-	-	-	-
2002.....	39	-	1	2	17	11	5	1	-	1	1	-	-
2001.....	29	-	-	2	14	4	1	4	3	-	1	-	-
2000.....	47	-	-	4	19	12	5	2	3	2	-	-	-
1999.....	56	-	2	1	29	12	4	4	1	2	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Discharge of firearms, undetermined intent (Y22-Y24)

Black, female

2007 1/.....	11	-	-	-	5	1	-	3	1	1	-	-	-
2006 1/.....	3	-	-	-	-	1	1	1	-	-	-	-	-
2005 1/.....	8	-	1	-	1	-	1	2	2	1	-	-	-
2004 1/.....	8	-	1	1	3	1	1	-	1	-	-	-	-
2003 1/.....	3	-	-	-	1	-	1	1	-	-	-	-	-
2002.....	2	-	-	-	-	-	1	-	1	-	-	-	-
2001.....	3	-	-	-	1	1	1	-	-	-	-	-	-
2000.....	7	-	-	1	1	3	2	-	-	-	-	-	-
1999.....	10	-	-	-	6	2	2	-	-	-	-	-	-

American Indian 2/, both sexes

2007 1/.....	6	-	-	-	5	-	1	-	-	-	-	-	-
2006 1/.....	2	-	-	-	1	1	-	-	-	-	-	-	-
2005 1/.....	7	-	1	-	3	1	-	1	-	-	1	-	-
2004 1/.....	7	-	-	1	1	4	1	-	-	-	-	-	-
2003 1/.....	9	-	-	1	2	4	2	-	-	-	-	-	-
2002.....	7	-	-	-	3	2	1	-	1	-	-	-	-
2001.....	4	-	-	-	2	1	1	-	-	-	-	-	-
2000.....	4	-	-	-	2	1	-	-	1	-	-	-	-
1999.....	6	-	-	-	4	1	1	-	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Discharge of firearms, undetermined intent (Y22-Y24)

American Indian 2/, male

2007 1/.....	6	-	-	-	5	-	1	-	-	-	-	-	-
2006 1/.....	2	-	-	-	1	1	-	-	-	-	-	-	-
2005 1/.....	7	-	1	-	3	1	1	-	-	-	1	-	-
2004 1/.....	4	-	-	-	1	2	1	-	-	-	-	-	-
2003 1/.....	8	-	-	1	2	3	2	-	-	-	-	-	-
2002.....	5	-	-	-	3	1	1	-	-	-	-	-	-
2001.....	4	-	-	-	2	1	1	-	-	-	-	-	-
2000.....	3	-	-	-	1	1	-	-	1	-	-	-	-
1999.....	5	-	-	-	4	1	-	-	-	-	-	-	-

American Indian 2/, female

2007 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2006 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2005 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 1/.....	3	-	-	1	-	2	-	-	-	-	-	-	-
2003 1/.....	1	-	-	-	-	1	-	-	-	-	-	-	-
2002.....	2	-	-	-	-	1	-	-	1	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	1	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	1	-	-	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Discharge of firearms, undetermined intent (Y22-Y24)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	5	-	-	1	1	1	1	-	1	-	-	-	-
2006 1/.....	1	-	-	-	-	1	-	-	-	-	-	-	-
2005 1/.....	3	-	-	1	1	-	1	-	-	-	-	-	-
2004 1/.....	2	-	-	-	-	1	-	1	-	-	-	-	-
2003 1/.....	5	-	-	-	3	-	-	-	2	-	-	-	-
2002.....	3	-	-	-	-	-	2	1	-	-	-	-	-
2001.....	7	-	-	-	3	2	2	-	-	-	-	-	-
2000.....	5	-	-	-	1	2	1	1	-	-	-	-	-
1999.....	5	-	-	-	3	-	-	1	1	-	-	-	-

Asian or Pacific Islander 3/, male

2007 1/.....	5	-	-	1	1	1	1	-	1	-	-	-	-
2006 1/.....	1	-	-	-	-	1	-	-	-	-	-	-	-
2005 1/.....	2	-	-	1	1	-	-	-	-	-	-	-	-
2004 1/.....	2	-	-	-	-	1	-	1	-	-	-	-	-
2003 1/.....	2	-	-	-	2	-	-	-	-	-	-	-	-
2002.....	2	-	-	-	-	-	1	1	-	-	-	-	-
2001.....	7	-	-	-	3	2	2	-	-	-	-	-	-
2000.....	3	-	-	-	-	2	-	1	-	-	-	-	-
1999.....	4	-	-	-	2	-	-	1	1	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

All races, both sexes

2007.....	5,105	88	46	37	497	828	1,215	1,461	564	166	126	69	8
2006.....	4,911	92	50	54	526	772	1,239	1,340	472	155	133	72	6
2005.....	4,521	104	66	52	407	707	1,166	1,286	401	147	103	72	10
2004.....	4,741	104	46	55	443	761	1,345	1,273	379	140	111	78	6
2003.....	4,840	75	30	43	494	797	1,538	1,236	348	123	87	51	18
2002.....	4,587	101	56	62	408	782	1,466	1,111	308	124	90	65	14
2001.....	3,967	83	44	45	348	680	1,346	961	213	89	97	43	18
2000.....	3,589	65	39	50	299	627	1,247	818	213	93	73	55	10
1999.....	3,593	55	40	37	303	692	1,238	765	208	99	83	54	19

All races, male

2007.....	3,128	55	24	22	374	533	758	829	330	96	73	27	7
2006.....	3,068	57	27	38	384	532	736	816	275	87	73	37	6
2005.....	2,683	67	42	33	287	449	674	736	231	84	46	24	10
2004.....	2,985	65	26	36	330	526	842	786	217	78	46	27	6
2003.....	3,109	50	16	29	352	547	987	789	206	58	41	20	14
2002.....	2,915	52	27	33	287	546	942	673	196	73	42	31	13
2001.....	2,643	49	19	32	257	473	898	641	135	56	48	19	16
2000.....	2,445	27	26	41	223	446	855	552	141	61	44	19	10
1999.....	2,415	33	18	26	234	479	842	511	137	58	43	21	13

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

All races, female

2007.....	1,977	33	22	15	123	295	457	632	234	70	53	42	1
2006.....	1,843	35	23	16	142	240	503	524	197	68	60	35	-
2005.....	1,838	37	24	19	120	258	492	550	170	63	57	48	-
2004.....	1,756	39	20	19	113	235	503	487	162	62	65	51	-
2003.....	1,731	25	14	14	142	250	551	447	142	65	46	31	4
2002.....	1,672	49	29	29	121	236	524	438	112	51	48	34	1
2001.....	1,324	34	25	13	91	207	448	320	78	33	49	24	2
2000.....	1,144	38	13	9	76	181	392	266	72	32	29	36	-
1999.....	1,178	22	22	11	69	213	396	254	71	41	40	33	6

White, both sexes

2007 1/.....	4,241	54	34	28	434	720	1,009	1,206	450	136	101	64	5
2006 1/.....	4,051	63	27	37	469	666	1,025	1,095	381	125	104	54	5
2005 1/.....	3,742	68	43	37	349	601	953	1,069	339	126	86	65	6
2004 1/.....	3,955	68	30	32	390	646	1,124	1,063	323	108	98	70	3
2003 1/.....	4,072	52	21	30	441	692	1,295	1,028	282	95	76	49	11
2002.....	3,770	56	45	45	359	646	1,211	898	262	101	75	60	12
2001.....	3,265	45	31	35	307	572	1,101	780	180	73	86	40	15
2000.....	2,886	41	27	33	254	519	1,000	652	166	75	66	46	7
1999.....	2,875	36	22	31	257	570	973	589	174	84	77	48	14

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

White, male

2007 1/.....	2,555	34	16	14	327	464	623	661	256	74	55	26	5
2006 1/.....	2,483	41	11	25	347	456	593	644	214	65	52	30	5
2005 1/.....	2,186	45	28	24	250	380	549	593	183	69	36	23	6
2004 1/.....	2,465	44	16	20	292	449	698	641	177	60	40	25	3
2003 1/.....	2,565	30	11	22	311	466	826	635	156	44	37	19	8
2002.....	2,371	29	23	23	255	450	773	527	162	53	36	29	11
2001.....	2,133	28	13	25	223	396	716	503	109	44	45	18	13
2000.....	1,941	18	17	27	186	364	685	431	105	46	40	15	7
1999.....	1,893	22	12	22	202	385	657	374	104	49	38	19	9

White, female

2007 1/.....	1,686	20	18	14	107	256	386	545	194	62	46	38	-
2006 1/.....	1,568	22	16	12	122	210	432	451	167	60	52	24	-
2005 1/.....	1,556	23	15	13	99	221	404	476	156	57	50	42	-
2004 1/.....	1,490	24	14	12	98	197	426	422	146	48	58	45	-
2003 1/.....	1,507	22	10	8	130	226	469	393	126	51	39	30	3
2002.....	1,399	27	22	22	104	196	438	371	100	48	39	31	1
2001.....	1,132	17	18	10	84	176	385	277	71	29	41	22	2
2000.....	945	23	10	6	68	155	315	221	61	29	26	31	-
1999.....	982	14	10	9	55	185	316	215	70	35	39	29	5

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Black, both sexes

2007 1/.....	703	29	11	9	37	81	167	223	96	26	19	2	3
2006 1/.....	670	23	21	15	34	74	177	202	75	24	15	9	1
2005 1/.....	649	31	21	10	40	86	184	187	50	17	13	7	3
2004 1/.....	663	32	14	22	41	91	187	188	49	21	10	5	3
2003 1/.....	652	21	8	10	34	89	213	188	52	21	7	2	7
2002.....	718	37	7	13	40	114	238	195	38	16	15	4	1
2001.....	614	36	13	9	29	88	225	164	29	12	4	3	2
2000.....	614	15	11	16	32	90	227	154	40	13	6	7	3
1999.....	630	18	14	6	36	107	234	164	27	12	4	3	5

Black, male

2007 1/.....	481	18	8	8	32	52	113	146	69	19	14	-	2
2006 1/.....	463	12	14	11	22	54	120	147	51	18	11	2	1
2005 1/.....	425	19	13	7	26	56	110	130	40	12	8	1	3
2004 1/.....	448	18	9	15	31	65	121	130	38	13	5	-	3
2003 1/.....	466	19	4	4	27	70	142	140	39	12	2	1	6
2002.....	482	18	2	9	26	79	159	136	30	15	6	1	1
2001.....	451	20	6	6	24	63	169	126	24	10	-	1	2
2000.....	441	3	8	13	28	68	157	113	30	11	4	3	3
1999.....	467	10	6	4	27	81	167	130	26	8	3	1	4

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Black, female

2007 1/.....	222	11	3	1	5	29	54	77	27	7	5	2	1
2006 1/.....	207	11	7	4	12	20	57	55	24	6	4	7	-
2005 1/.....	224	12	8	3	14	30	74	57	10	5	5	6	-
2004 1/.....	215	14	5	7	10	26	66	58	11	8	5	5	-
2003 1/.....	186	2	4	6	7	19	71	48	13	9	5	1	1
2002.....	236	19	5	4	14	35	79	59	8	1	9	3	-
2001.....	163	16	7	3	5	25	56	38	5	2	4	2	-
2000.....	173	12	3	3	4	22	70	41	10	2	2	4	-
1999.....	163	8	8	2	9	26	67	34	1	4	1	2	1

American Indian 2/, both sexes

2007 1/.....	86	4	-	-	15	14	21	13	13	2	3	1	-
2006 1/.....	93	2	2	2	13	16	23	22	9	2	2	-	-
2005 1/.....	57	3	1	1	6	8	11	16	7	2	1	-	1
2004 1/.....	61	3	-	1	5	12	22	10	4	3	1	-	-
2003 1/.....	55	2	-	1	6	7	19	11	7	1	1	-	-
2002.....	41	3	2	2	2	10	7	9	3	2	-	-	1
2001.....	39	1	-	-	5	6	13	9	1	1	2	-	1
2000.....	42	4	1	1	3	8	12	7	5	1	-	-	-
1999.....	48	1	2	-	5	10	17	6	3	2	1	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

American Indian 2/, male

2007 1/.....	45	2	-	-	7	8	11	9	3	2	2	1	-
2006 1/.....	60	1	2	2	7	14	15	10	6	1	2	-	-
2005 1/.....	30	1	1	-	3	4	5	8	5	1	1	-	1
2004 1/.....	37	3	-	1	4	7	14	5	1	2	-	-	-
2003 1/.....	39	1	-	1	5	4	14	6	7	1	-	-	-
2002.....	21	1	1	1	2	6	2	4	2	1	-	-	1
2001.....	28	-	-	-	5	6	8	5	1	-	2	-	1
2000.....	33	3	1	1	2	7	9	5	5	-	-	-	-
1999.....	32	1	-	-	3	9	10	4	3	1	1	-	-

American Indian 2/, female

2007 1/.....	41	2	-	-	8	6	10	4	10	-	1	-	-
2006 1/.....	33	1	-	-	6	2	8	12	3	1	-	-	-
2005 1/.....	27	2	-	1	3	4	6	8	2	1	-	-	-
2004 1/.....	24	-	-	-	1	5	8	5	3	1	1	-	-
2003 1/.....	16	1	-	-	1	3	5	5	-	-	1	-	-
2002.....	20	2	1	1	-	4	5	5	1	1	-	-	-
2001.....	11	1	-	-	-	-	5	4	-	1	-	-	-
2000.....	9	1	-	-	1	1	3	2	-	1	-	-	-
1999.....	16	-	2	-	2	1	7	2	-	1	-	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	75	1	1	-	11	13	18	19	5	2	3	2	-
2006 1/.....	97	4	-	-	10	16	14	21	7	4	12	9	-
2005 1/.....	73	2	1	4	12	12	18	14	5	2	3	-	-
2004 1/.....	62	1	2	-	7	12	12	12	3	8	2	3	-
2003 1/.....	61	-	1	2	13	9	11	9	7	6	3	-	-
2002.....	58	5	2	2	7	12	10	9	5	5	-	1	-
2001.....	49	1	-	1	7	14	7	8	3	3	5	-	-
2000.....	47	5	-	-	10	10	8	5	2	4	1	2	-
1999.....	40	-	2	-	5	5	14	6	4	1	1	2	-

Asian or Pacific Islander 3/, male

2007 1/.....	47	1	-	-	8	9	11	13	2	1	2	-	-
2006 1/.....	62	3	-	-	8	8	8	15	4	3	8	5	-
2005 1/.....	42	2	-	2	8	9	10	5	3	2	1	-	-
2004 1/.....	35	-	1	-	3	5	9	10	1	3	1	2	-
2003 1/.....	39	-	1	2	9	7	5	8	4	1	2	-	-
2002.....	41	4	1	-	4	11	8	6	2	4	-	1	-
2001.....	31	1	-	1	5	8	5	7	1	2	1	-	-
2000.....	30	3	-	-	7	7	4	3	1	4	-	1	-
1999.....	23	-	-	-	2	4	8	3	4	-	1	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Asian or Pacific Islander 3/, female

2007 1/.....	28	-	1	-	3	4	7	6	3	1	1	2	-
2006 1/.....	35	1	-	-	2	8	6	6	3	1	4	4	-
2005 1/.....	31	-	1	2	4	3	8	9	2	-	2	-	-
2004 1/.....	27	1	1	-	4	7	3	2	2	5	1	1	-
2003 1/.....	22	-	-	-	4	2	6	1	3	5	1	-	-
2002.....	17	1	1	2	3	1	2	3	3	1	-	-	-
2001.....	18	-	-	-	2	6	2	1	2	1	4	-	-
2000.....	17	2	-	-	3	3	4	2	1	-	1	1	-
1999.....	17	-	2	-	3	1	6	3	-	1	-	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Operations of war and their sequelae (Y36,Y89.1)

All races, both sexes

2007.....	21	-	-	-	3	5	-	-	6	1	5	1	-
2006.....	28	-	-	-	13	4	2	1	2	2	4	-	-
2005.....	27	-	-	-	3	3	4	1	3	2	10	1	-
2004.....	28	-	-	-	4	1	3	-	4	1	14	1	-
2003.....	14	-	-	-	1	-	-	1	5	2	5	-	-
2002.....	20	-	-	-	-	1	1	1	3	3	10	1	-
2001.....	17	-	-	-	1	-	1	2	-	4	8	1	-
2000.....	16	-	-	-	-	-	-	2	1	3	9	1	-
1999.....	23	-	-	-	-	-	1	2	1	8	8	3	-

All races, male

2007.....	21	-	-	-	3	5	-	-	6	1	5	1	-
2006.....	28	-	-	-	13	4	2	1	2	2	4	-	-
2005.....	27	-	-	-	3	3	4	1	3	2	10	1	-
2004.....	28	-	-	-	4	1	3	-	4	1	14	1	-
2003.....	14	-	-	-	1	-	-	1	5	2	5	-	-
2002.....	20	-	-	-	-	1	1	1	3	3	10	1	-
2001.....	16	-	-	-	1	-	1	2	-	4	7	1	-
2000.....	16	-	-	-	-	-	-	2	1	3	9	1	-
1999.....	22	-	-	-	-	-	1	2	1	8	7	3	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Operations of war and their sequelae (Y36,Y89.1)

All races, female

2007.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2006.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2005.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2004.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	1	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	-	1	-	-

White, both sexes

2007 1/.....	21	-	-	-	3	5	-	-	6	1	5	1	-
2006 1/.....	27	-	-	-	13	4	1	1	2	2	4	-	-
2005 1/.....	21	-	-	-	3	1	3	-	2	2	9	1	-
2004 1/.....	26	-	-	-	3	1	2	-	4	1	14	1	-
2003 1/.....	13	-	-	-	1	-	-	1	4	2	5	-	-
2002.....	15	-	-	-	-	1	-	-	2	2	9	1	-
2001.....	16	-	-	-	1	-	1	2	-	3	8	1	-
2000.....	15	-	-	-	-	-	-	1	1	3	9	1	-
1999.....	22	-	-	-	-	-	1	2	1	7	8	3	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Operations of war and their sequelae (Y36,Y89.1)

White, male

2007 1/.....	21	-	-	-	3	5	-	-	6	1	5	1	-
2006 1/.....	27	-	-	-	13	4	1	1	2	2	4	-	-
2005 1/.....	21	-	-	-	3	1	3	-	2	2	9	1	-
2004 1/.....	26	-	-	-	3	1	2	-	4	1	14	1	-
2003 1/.....	13	-	-	-	1	-	-	1	4	2	5	-	-
2002.....	15	-	-	-	-	1	-	-	2	2	9	1	-
2001.....	15	-	-	-	1	-	1	2	-	3	7	1	-
2000.....	15	-	-	-	-	-	-	1	1	3	9	1	-
1999.....	21	-	-	-	-	-	1	2	1	7	7	3	-

White, female

2007 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2006 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2005 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2004 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2003 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	1	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	-	1	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Operations of war and their sequelae (Y36,Y89.1)

Black, both sexes

2007 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2006 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2005 1/.....	6	-	-	-	-	2	1	1	1	-	1	-	-
2004 1/.....	2	-	-	-	1	-	1	-	-	-	-	-	-
2003 1/.....	1	-	-	-	-	-	-	-	1	-	-	-	-
2002.....	5	-	-	-	-	-	1	1	1	1	1	-	-
2001.....	1	-	-	-	-	-	-	-	-	1	-	-	-
2000.....	1	-	-	-	-	-	-	1	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	1	-	-	-

Black, male

2007 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2006 1/.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2005 1/.....	6	-	-	-	-	2	1	1	1	-	1	-	-
2004 1/.....	2	-	-	-	1	-	1	-	-	-	-	-	-
2003 1/.....	1	-	-	-	-	-	-	-	1	-	-	-	-
2002.....	5	-	-	-	-	-	1	1	1	1	1	-	-
2001.....	1	-	-	-	-	-	-	-	-	1	-	-	-
2000.....	1	-	-	-	-	-	-	1	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	1	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Operations of war and their sequelae (Y36,Y89.1)

Black, female

Table with 14 columns (years 2007-1999) and 14 rows (age groups) for Black, female. All cells contain dashes (-).

American Indian 2/, both sexes

Table with 14 columns (years 2007-1999) and 14 rows (age groups) for American Indian 2/, both sexes. All cells contain dashes (-).

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Complications of medical and surgical care (Y40-Y84,Y88)

All races, both sexes

2007.....	2,597	22	23	21	30	72	150	250	398	486	712	433	-
2006.....	2,521	23	22	19	45	62	118	276	351	475	688	442	-
2005.....	2,653	19	16	16	29	70	144	287	364	487	759	462	-
2004.....	2,883	16	19	19	48	71	157	293	403	557	787	513	-
2003.....	2,855	15	24	12	44	69	179	273	380	534	794	530	1
2002.....	2,843	15	19	35	35	79	171	265	351	560	792	521	-
2001.....	3,021	26	24	14	43	79	172	283	384	567	864	564	1
2000.....	3,059	19	22	31	32	85	149	295	376	647	825	578	-
1999.....	2,823	54	30	18	36	78	152	214	315	586	811	529	-

All races, male

2007.....	1,224	10	9	10	19	30	68	119	203	244	338	174	-
2006.....	1,121	13	11	11	23	25	51	145	164	210	319	149	-
2005.....	1,220	16	10	9	16	26	62	131	192	250	335	173	-
2004.....	1,364	6	10	12	24	31	68	148	220	280	385	180	-
2003.....	1,333	8	16	4	26	36	96	146	195	264	355	186	1
2002.....	1,236	8	11	26	19	47	82	127	169	256	320	171	-
2001.....	1,357	15	14	8	19	42	83	143	196	265	383	189	-
2000.....	1,430	12	11	15	16	47	76	143	181	331	402	196	-
1999.....	1,308	31	12	9	21	43	78	98	165	279	375	197	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Complications of medical and surgical care (Y40-Y84,Y88)

All races, female

2007.....	1,373	12	14	11	11	42	82	131	195	242	374	259	-
2006.....	1,400	10	11	8	22	37	67	131	187	265	369	293	-
2005.....	1,433	3	6	7	13	44	82	156	172	237	424	289	-
2004.....	1,519	10	9	7	24	40	89	145	183	277	402	333	-
2003.....	1,522	7	8	8	18	33	83	127	185	270	439	344	-
2002.....	1,607	7	8	9	16	32	89	138	182	304	472	350	-
2001.....	1,664	11	10	6	24	37	89	140	188	302	481	375	1
2000.....	1,629	7	11	16	16	38	73	152	195	316	423	382	-
1999.....	1,515	23	18	9	15	35	74	116	150	307	436	332	-

White, both sexes

2007 1/.....	2,107	9	17	13	24	55	112	182	320	394	607	374	-
2006 1/.....	2,046	16	17	14	35	44	84	201	257	396	588	394	-
2005 1/.....	2,170	11	13	14	23	54	112	203	283	379	657	421	-
2004 1/.....	2,392	11	16	16	33	57	113	226	323	460	682	455	-
2003 1/.....	2,356	9	13	8	37	48	141	192	297	445	696	469	1
2002.....	2,331	8	14	21	24	57	119	185	271	470	697	465	-
2001.....	2,446	17	16	10	33	57	117	207	293	445	767	483	1
2000.....	2,520	14	12	23	24	60	115	231	285	535	718	503	-
1999.....	2,314	35	21	14	30	55	114	159	240	492	699	455	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Complications of medical and surgical care (Y40-Y84,Y88)

White, male

2007 1/.....	1,015	3	6	8	16	24	53	94	163	198	299	151	-
2006 1/.....	926	9	8	8	18	15	42	109	123	177	283	134	-
2005 1/.....	1,014	8	9	8	13	23	53	94	145	207	292	162	-
2004 1/.....	1,162	4	8	10	16	26	54	120	179	241	344	160	-
2003 1/.....	1,121	5	9	3	22	25	81	99	157	229	323	167	1
2002.....	1,025	5	9	16	14	35	65	95	135	212	290	149	-
2001.....	1,084	9	10	5	16	29	54	102	145	207	347	160	-
2000.....	1,210	11	8	11	13	33	64	115	144	273	363	175	-
1999.....	1,078	22	11	7	17	32	60	72	126	235	328	168	-

White, female

2007 1/.....	1,092	6	11	5	8	31	59	88	157	196	308	223	-
2006 1/.....	1,120	7	9	6	17	29	42	92	134	219	305	260	-
2005 1/.....	1,156	3	4	6	10	31	59	109	138	172	365	259	-
2004 1/.....	1,230	7	8	6	17	31	59	106	144	219	338	295	-
2003 1/.....	1,235	4	4	5	15	23	60	93	140	216	373	302	-
2002.....	1,306	3	5	5	10	22	54	90	136	258	407	316	-
2001.....	1,362	8	6	5	17	28	63	105	148	238	420	323	1
2000.....	1,310	3	4	12	11	27	51	116	141	262	355	328	-
1999.....	1,236	13	10	7	13	23	54	87	114	257	371	287	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Complications of medical and surgical care (Y40-Y84,Y88)

Black, both sexes

2007 1/.....	432	11	6	6	5	16	35	62	68	84	90	49	-
2006 1/.....	427	7	4	4	7	17	31	70	85	73	89	40	-
2005 1/.....	434	8	2	2	5	15	29	78	76	94	91	34	-
2004 1/.....	437	2	3	3	13	10	43	60	75	86	93	49	-
2003 1/.....	432	5	10	3	6	16	34	68	74	78	86	52	-
2002.....	448	7	5	14	7	17	49	75	70	77	80	47	-
2001.....	522	9	7	3	7	22	50	70	85	106	87	76	-
2000.....	487	5	8	5	7	23	31	58	86	101	96	67	-
1999.....	449	18	9	4	6	18	35	52	62	86	96	63	-

Black, male

2007 1/.....	188	6	3	2	2	5	14	22	37	41	37	19	-
2006 1/.....	179	4	3	2	3	10	9	34	37	30	35	12	-
2005 1/.....	181	8	-	1	2	3	8	35	42	36	36	10	-
2004 1/.....	173	1	2	2	6	3	14	23	38	33	36	15	-
2003 1/.....	177	3	6	1	3	8	12	40	35	28	27	14	-
2002.....	184	3	2	10	4	9	16	29	30	37	25	19	-
2001.....	250	6	4	2	2	13	27	38	47	54	31	26	-
2000.....	193	1	2	2	2	13	12	25	34	52	34	16	-
1999.....	197	8	1	2	4	10	16	26	30	40	40	20	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Complications of medical and surgical care (Y40-Y84,Y88)

Black, female

2007 1/.....	244	5	3	4	3	11	21	40	31	43	53	30	-
2006 1/.....	248	3	1	2	4	7	22	36	48	43	54	28	-
2005 1/.....	253	-	2	1	3	12	21	43	34	58	55	24	-
2004 1/.....	264	1	1	1	7	7	29	37	37	53	57	34	-
2003 1/.....	255	2	4	2	3	8	22	28	39	50	59	38	-
2002.....	264	4	3	4	3	8	33	46	40	40	55	28	-
2001.....	272	3	3	1	5	9	23	32	38	52	56	50	-
2000.....	294	4	6	3	5	10	19	33	52	49	62	51	-
1999.....	252	10	8	2	2	8	19	26	32	46	56	43	-

American Indian 2/, both sexes

2007 1/.....	16	1	-	-	-	-	-	1	5	1	6	2	-
2006 1/.....	12	-	-	-	-	-	2	2	3	3	2	-	-
2005 1/.....	20	-	-	-	-	-	2	5	2	6	3	2	-
2004 1/.....	11	2	-	-	-	-	-	3	1	4	-	1	-
2003 1/.....	24	1	-	-	1	3	1	5	4	3	3	3	-
2002.....	26	-	-	-	1	2	2	3	5	7	4	2	-
2001.....	17	-	1	1	2	-	2	2	1	3	4	1	-
2000.....	14	-	-	1	-	-	1	1	-	5	3	3	-
1999.....	20	-	-	-	-	3	1	1	3	3	5	4	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Complications of medical and surgical care (Y40-Y84,Y88)

American Indian 2/, male

2007 1/.....	4	-	-	-	-	-	-	1	2	-	-	1	-
2006 1/.....	4	-	-	-	-	-	-	-	1	2	1	-	-
2005 1/.....	9	-	-	-	-	-	1	1	2	3	2	-	-
2004 1/.....	6	1	-	-	-	-	-	3	1	1	-	-	-
2003 1/.....	10	-	-	-	1	2	1	3	-	1	-	2	-
2002.....	12	-	-	-	1	2	-	2	1	4	1	1	-
2001.....	6	-	-	1	-	-	1	1	1	-	2	-	-
2000.....	5	-	-	-	-	-	-	1	-	1	2	1	-
1999.....	13	-	-	-	-	1	1	-	3	2	4	2	-

American Indian 2/, female

2007 1/.....	12	1	-	-	-	-	-	-	3	1	6	1	-
2006 1/.....	8	-	-	-	-	-	2	2	2	1	1	-	-
2005 1/.....	11	-	-	-	-	-	1	4	-	3	1	2	-
2004 1/.....	5	1	-	-	-	-	-	-	-	3	-	1	-
2003 1/.....	14	1	-	-	-	1	-	2	4	2	3	1	-
2002.....	14	-	-	-	-	-	2	1	4	3	3	1	-
2001.....	11	-	1	-	2	-	1	1	-	3	2	1	-
2000.....	9	-	-	1	-	-	1	-	-	4	1	2	-
1999.....	7	-	-	-	-	2	-	1	-	1	1	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Complications of medical and surgical care (Y40-Y84,Y88)

Asian or Pacific Islander 3/, both sexes

2007 1/.....	42	1	-	2	1	1	3	5	5	7	9	8	-
2006 1/.....	36	-	1	1	3	1	1	3	6	3	9	8	-
2005 1/.....	29	-	1	-	1	1	1	1	3	8	8	5	-
2004 1/.....	43	1	-	-	2	4	1	4	4	7	12	8	-
2003 1/.....	43	-	1	1	-	2	3	8	5	8	9	6	-
2002.....	38	-	-	-	3	3	1	2	5	6	11	7	-
2001.....	36	-	-	-	1	-	3	4	5	13	6	4	-
2000.....	38	-	2	2	1	2	2	5	5	6	8	5	-
1999.....	40	1	-	-	-	2	2	2	10	5	11	7	-

Asian or Pacific Islander 3/, male

2007 1/.....	17	1	-	-	1	1	1	2	1	5	2	3	-
2006 1/.....	12	-	-	1	2	-	-	2	3	1	-	3	-
2005 1/.....	16	-	1	-	1	-	-	1	3	4	5	1	-
2004 1/.....	23	-	-	-	2	2	-	2	2	5	5	5	-
2003 1/.....	25	-	1	-	-	1	2	4	3	6	5	3	-
2002.....	15	-	-	-	-	1	1	1	3	3	4	2	-
2001.....	17	-	-	-	1	-	1	2	3	4	3	3	-
2000.....	22	-	1	2	1	1	-	2	3	5	3	4	-
1999.....	20	1	-	-	-	-	1	-	6	2	3	7	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Complications of medical and surgical care (Y40-Y84,Y88)

Asian or Pacific Islander 3/, female

2007 1/.....	25	-	-	2	-	-	2	3	4	2	7	5	-
2006 1/.....	24	-	1	-	1	1	1	1	3	2	9	5	-
2005 1/.....	13	-	-	-	-	1	1	-	-	4	3	4	-
2004 1/.....	20	1	-	-	-	2	1	2	2	2	7	3	-
2003 1/.....	18	-	-	1	-	1	1	4	2	2	4	3	-
2002.....	23	-	-	-	3	2	-	1	2	3	7	5	-
2001.....	19	-	-	-	-	-	2	2	2	9	3	1	-
2000.....	16	-	1	-	-	1	2	3	2	1	5	1	-
1999.....	20	-	-	-	-	2	1	2	4	3	8	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 5/

All races, both sexes

2007.....	23,199	2	-	4	194	735	3,138	8,017	6,597	2,993	1,260	253	6
2006.....	22,073	-	-	2	149	637	3,237	7,596	6,127	2,788	1,269	258	10
2005.....	21,634	-	-	2	149	541	3,281	7,463	5,930	2,822	1,201	236	9
2004.....	21,081	-	-	6	139	610	3,359	7,215	5,442	2,892	1,200	211	7
2003.....	20,687	-	-	5	124	596	3,572	7,063	5,168	2,771	1,183	194	11
2002.....	20,218	1	2	3	105	601	3,636	6,748	4,888	2,835	1,190	201	8
2001.....	20,114	2	-	2	106	645	3,746	6,727	4,599	2,846	1,204	221	16
2000.....	19,643	3	1	4	95	650	3,837	6,143	4,542	2,910	1,223	227	8
1999.....	19,469	1	-	2	100	641	3,829	6,002	4,439	2,926	1,291	230	8

All races, male

2007.....	17,428	2	-	3	146	553	2,210	5,911	5,134	2,349	945	170	5
2006.....	16,472	-	-	2	123	496	2,246	5,636	4,736	2,158	905	160	10
2005.....	16,238	-	-	1	127	405	2,336	5,569	4,616	2,133	895	149	7
2004.....	15,906	-	-	5	107	468	2,434	5,401	4,224	2,232	894	135	6
2003.....	15,630	-	-	3	101	472	2,534	5,414	3,997	2,120	866	113	10
2002.....	15,272	-	1	2	86	458	2,571	5,171	3,837	2,158	845	135	8
2001.....	15,149	2	-	-	88	466	2,665	5,116	3,607	2,181	858	151	15
2000.....	14,993	2	-	4	76	504	2,808	4,792	3,489	2,249	915	146	8
1999.....	14,894	1	-	1	80	459	2,816	4,702	3,481	2,270	936	141	7

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 5/

All races, female

2007.....	5,771	-	-	1	48	182	928	2,106	1,463	644	315	83	1
2006.....	5,601	-	-	-	26	141	991	1,960	1,391	630	364	98	-
2005.....	5,396	-	-	1	22	136	945	1,894	1,314	689	306	87	2
2004.....	5,175	-	-	1	32	142	925	1,814	1,218	660	306	76	1
2003.....	5,057	-	-	2	23	124	1,038	1,649	1,171	651	317	81	1
2002.....	4,946	1	1	1	19	143	1,065	1,577	1,051	677	345	66	-
2001.....	4,965	-	-	2	18	179	1,081	1,611	992	665	346	70	1
2000.....	4,650	1	1	-	19	146	1,029	1,351	1,053	661	308	81	-
1999.....	4,575	-	-	1	20	182	1,013	1,300	958	656	355	89	1

White, both sexes

2007 1/.....	19,921	1	-	3	166	591	2,648	6,860	5,683	2,603	1,133	227	6
2006 1/.....	18,917	-	-	-	124	501	2,694	6,515	5,268	2,411	1,159	238	7
2005 1/.....	18,432	-	-	1	119	429	2,743	6,304	5,092	2,452	1,071	213	8
2004 1/.....	17,875	-	-	4	111	482	2,821	6,123	4,586	2,459	1,089	196	4
2003 1/.....	17,437	-	-	4	96	483	2,964	5,878	4,340	2,413	1,074	177	8
2002.....	16,988	-	1	2	86	476	3,026	5,644	4,058	2,448	1,061	180	6
2001.....	16,640	1	-	2	78	507	3,033	5,483	3,810	2,438	1,083	192	13
2000.....	16,223	3	-	3	75	504	3,095	4,994	3,739	2,506	1,094	204	6
1999.....	15,903	1	-	2	77	514	3,035	4,835	3,617	2,481	1,132	201	8

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 5/

White, male

2007 1/.....	15,027	1	-	3	128	458	1,878	5,085	4,441	2,029	843	156	5
2006 1/.....	14,186	-	-	-	103	404	1,906	4,858	4,071	1,859	830	148	7
2005 1/.....	13,917	-	-	1	106	325	1,986	4,728	3,983	1,850	796	136	6
2004 1/.....	13,525	-	-	3	89	375	2,058	4,611	3,562	1,884	813	127	3
2003 1/.....	13,218	-	-	3	82	383	2,133	4,512	3,364	1,853	779	102	7
2002.....	12,926	-	1	1	73	372	2,172	4,373	3,206	1,850	755	117	6
2001.....	12,588	1	-	-	66	377	2,164	4,229	2,979	1,856	773	131	12
2000.....	12,509	2	-	3	62	400	2,293	3,957	2,899	1,937	821	129	6
1999.....	12,277	1	-	1	63	378	2,276	3,839	2,853	1,912	824	123	7

White, female

2007 1/.....	4,894	-	-	-	38	133	770	1,775	1,242	574	290	71	1
2006 1/.....	4,731	-	-	-	21	97	788	1,657	1,197	552	329	90	-
2005 1/.....	4,515	-	-	-	13	104	757	1,576	1,109	602	275	77	2
2004 1/.....	4,350	-	-	1	22	107	763	1,512	1,024	575	276	69	1
2003 1/.....	4,219	-	-	1	14	100	831	1,366	976	560	295	75	1
2002.....	4,062	-	-	1	13	104	854	1,271	852	598	306	63	-
2001.....	4,052	-	-	2	12	130	869	1,254	831	582	310	61	1
2000.....	3,714	1	-	-	13	104	802	1,037	840	569	273	75	-
1999.....	3,626	-	-	1	14	136	759	996	764	569	308	78	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 5/

Black, both sexes

2007 1/.....	2,248	1	-	-	13	58	263	783	705	311	92	22	-
2006 1/.....	2,172	-	-	1	14	54	303	762	657	283	84	12	2
2005 1/.....	2,316	-	-	-	16	48	321	857	661	292	99	21	1
2004 1/.....	2,351	-	-	-	13	53	324	843	681	348	76	10	3
2003 1/.....	2,406	-	-	1	8	60	389	885	670	289	87	14	3
2002.....	2,434	1	1	-	10	71	389	861	668	308	108	16	1
2001.....	2,723	1	-	-	17	72	503	1,004	656	343	105	19	3
2000.....	2,712	-	1	1	9	94	527	960	645	348	105	20	2
1999.....	2,832	-	-	-	10	74	565	950	695	382	132	24	-

Black, male

2007 1/.....	1,703	1	-	-	10	40	175	575	551	262	77	12	-
2006 1/.....	1,599	-	-	1	12	33	189	552	516	227	59	8	2
2005 1/.....	1,698	-	-	-	11	33	206	632	501	227	76	11	1
2004 1/.....	1,784	-	-	-	8	43	236	616	536	285	52	5	3
2003 1/.....	1,824	-	-	-	6	48	259	693	520	218	67	10	3
2002.....	1,798	-	-	-	8	53	255	629	512	251	76	13	1
2001.....	2,048	1	-	-	13	48	361	737	524	276	74	11	3
2000.....	1,993	-	-	1	8	65	367	704	478	273	80	15	2
1999.....	2,100	-	-	-	8	47	385	705	541	308	92	14	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 5/

Black, female

2007 1/.....	545	-	-	-	3	18	88	208	154	49	15	10	-
2006 1/.....	573	-	-	-	2	21	114	210	141	56	25	4	-
2005 1/.....	618	-	-	-	5	15	115	225	160	65	23	10	-
2004 1/.....	567	-	-	-	5	10	88	227	145	63	24	5	-
2003 1/.....	582	-	-	1	2	12	130	192	150	71	20	4	-
2002.....	636	1	1	-	2	18	134	232	156	57	32	3	-
2001.....	675	-	-	-	4	24	142	267	132	67	31	8	-
2000.....	719	-	1	-	1	29	160	256	167	75	25	5	-
1999.....	732	-	-	-	2	27	180	245	154	74	40	10	-

American Indian 2/, both sexes

2007 1/.....	774	-	-	-	14	67	181	288	147	58	18	1	-
2006 1/.....	717	-	-	1	11	63	189	233	142	59	15	4	-
2005 1/.....	655	-	-	1	12	43	173	230	130	48	16	2	-
2004 1/.....	645	-	-	2	15	55	171	196	135	54	15	2	-
2003 1/.....	664	-	-	-	17	36	185	239	122	51	13	1	-
2002.....	597	-	-	1	8	51	173	185	117	52	9	1	-
2001.....	591	-	-	-	10	56	180	190	94	48	7	6	-
2000.....	537	-	-	-	7	48	172	145	110	42	12	1	-
1999.....	589	-	-	-	11	42	195	177	106	40	16	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 5/

American Indian 2/, male

2007 1/.....	495	-	-	-	8	41	119	182	89	42	14	-	-
2006 1/.....	458	-	-	1	8	42	109	148	97	43	9	1	-
2005 1/.....	432	-	-	-	8	30	108	147	94	33	10	2	-
2004 1/.....	417	-	-	2	10	31	103	127	94	36	13	1	-
2003 1/.....	431	-	-	-	11	26	111	155	84	33	11	-	-
2002.....	387	-	-	1	4	31	105	122	80	38	5	1	-
2001.....	387	-	-	-	8	36	116	111	72	35	3	6	-
2000.....	353	-	-	-	2	35	111	93	75	28	8	1	-
1999.....	397	-	-	-	7	26	123	128	69	30	12	2	-

American Indian 2/, female

2007 1/.....	279	-	-	-	6	26	62	106	58	16	4	1	-
2006 1/.....	259	-	-	-	3	21	80	85	45	16	6	3	-
2005 1/.....	223	-	-	1	4	13	65	83	36	15	6	-	-
2004 1/.....	228	-	-	-	5	24	68	69	41	18	2	1	-
2003 1/.....	233	-	-	-	6	10	74	84	38	18	2	1	-
2002.....	210	-	-	-	4	20	68	63	37	14	4	-	-
2001.....	204	-	-	-	2	20	64	79	22	13	4	-	-
2000.....	184	-	-	-	5	13	61	52	35	14	4	-	-
1999.....	192	-	-	-	4	16	72	49	37	10	4	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 5/

Asian or Pacific Islander 3/, both sexes

2007 1/.....	256	-	-	1	1	19	46	86	62	21	17	3	-
2006 1/.....	267	-	-	-	-	19	51	86	60	35	11	4	1
2005 1/.....	231	-	-	-	2	21	44	72	47	30	15	-	-
2004 1/.....	210	-	-	-	-	20	43	53	40	31	20	3	-
2003 1/.....	180	-	-	-	3	17	34	61	36	18	9	2	-
2002.....	199	-	-	-	1	3	48	58	45	27	12	4	1
2001.....	160	-	-	-	1	10	30	50	39	17	9	4	-
2000.....	171	-	-	-	4	4	43	44	48	14	12	2	-
1999.....	145	-	-	-	2	11	34	40	21	23	11	3	-

Asian or Pacific Islander 3/, male

2007 1/.....	203	-	-	-	-	14	38	69	53	16	11	2	-
2006 1/.....	229	-	-	-	-	17	42	78	52	29	7	3	1
2005 1/.....	191	-	-	-	2	17	36	62	38	23	13	-	-
2004 1/.....	180	-	-	-	-	19	37	47	32	27	16	2	-
2003 1/.....	157	-	-	-	2	15	31	54	29	16	9	1	-
2002.....	161	-	-	-	1	2	39	47	39	19	9	4	1
2001.....	126	-	-	-	1	5	24	39	32	14	8	3	-
2000.....	138	-	-	-	4	4	37	38	37	11	6	1	-
1999.....	120	-	-	-	2	8	32	30	18	20	8	2	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 5/

Asian or Pacific Islander 3/, female

2007 1/.....	53	-	-	1	1	5	8	17	9	5	6	1	-
2006 1/.....	38	-	-	-	-	2	9	8	8	6	4	1	-
2005 1/.....	40	-	-	-	-	4	8	10	9	7	2	-	-
2004 1/.....	30	-	-	-	-	1	6	6	8	4	4	1	-
2003 1/.....	23	-	-	-	1	2	3	7	7	2	-	1	-
2002.....	38	-	-	-	-	1	9	11	6	8	3	-	-
2001.....	34	-	-	-	-	5	6	11	7	3	1	1	-
2000.....	33	-	-	-	-	-	6	6	11	3	6	1	-
1999.....	25	-	-	-	-	3	2	10	3	3	3	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 5/

All races, both sexes

2007.....	38,371	33	44	75	3,675	6,950	9,671	11,792	4,453	905	505	264	4
2006.....	38,396	44	36	58	3,652	6,787	10,166	11,545	3,868	1,008	788	428	16
2005.....	33,541	35	30	51	3,095	5,730	9,333	10,041	3,240	887	706	388	5
2004.....	30,711	30	24	63	2,913	5,061	9,228	9,021	2,651	785	619	304	12
2003.....	28,723	25	35	41	2,619	4,845	9,158	8,174	2,239	755	537	283	12
2002.....	26,040	29	30	42	2,215	4,449	8,858	7,182	1,828	653	484	260	10
2001.....	21,705	22	26	45	1,813	3,739	7,669	5,796	1,361	553	439	226	16
2000.....	19,720	18	15	44	1,567	3,527	7,206	4,986	1,187	486	430	240	14
1999.....	19,128	21	25	30	1,351	3,579	7,095	4,624	1,161	556	467	201	18

All races, male

2007.....	23,883	18	21	43	2,760	4,859	6,021	6,847	2,537	473	203	98	3
2006.....	24,507	23	20	33	2,781	4,808	6,433	7,082	2,261	504	368	181	13
2005.....	21,208	18	23	24	2,319	3,970	5,829	6,143	1,912	452	354	159	5
2004.....	19,362	11	16	25	2,187	3,554	5,737	5,528	1,496	393	278	130	7
2003.....	18,426	17	21	20	1,955	3,361	5,873	5,153	1,266	384	253	112	11
2002.....	16,734	20	9	24	1,622	3,139	5,728	4,517	1,040	311	215	102	7
2001.....	14,253	14	21	22	1,350	2,640	5,079	3,793	760	293	191	74	16
2000.....	13,137	12	13	25	1,166	2,515	4,890	3,358	631	235	187	92	13
1999.....	12,885	14	12	16	981	2,579	4,907	3,133	688	286	184	74	11

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 5/

All races, female

2007.....	14,488	15	23	32	915	2,091	3,650	4,945	1,916	432	302	166	1
2006.....	13,889	21	16	25	871	1,979	3,733	4,463	1,607	504	420	247	3
2005.....	12,333	17	7	27	776	1,760	3,504	3,898	1,328	435	352	229	-
2004.....	11,349	19	8	38	726	1,507	3,491	3,493	1,155	392	341	174	5
2003.....	10,297	8	14	21	664	1,484	3,285	3,021	973	371	284	171	1
2002.....	9,306	9	21	18	593	1,310	3,130	2,665	788	342	269	158	3
2001.....	7,452	8	5	23	463	1,099	2,590	2,003	601	260	248	152	-
2000.....	6,583	6	2	19	401	1,012	2,316	1,628	556	251	243	148	1
1999.....	6,243	7	13	14	370	1,000	2,188	1,491	473	270	283	127	7

White, both sexes

2007 1/.....	33,480	18	22	61	3,413	6,338	8,444	10,015	3,679	783	460	245	2
2006 1/.....	32,866	28	22	45	3,414	6,074	8,649	9,531	3,124	851	719	394	15
2005 1/.....	28,804	25	15	38	2,885	5,066	8,021	8,298	2,661	776	647	367	5
2004 1/.....	26,474	18	15	48	2,702	4,504	7,977	7,487	2,177	690	569	280	7
2003 1/.....	24,683	16	22	31	2,451	4,276	7,809	6,767	1,871	653	513	267	7
2002.....	22,146	14	18	36	2,036	3,890	7,522	5,848	1,502	583	450	239	8
2001.....	18,195	10	17	38	1,651	3,245	6,408	4,599	1,115	487	407	203	15
2000.....	16,388	11	7	37	1,417	3,010	5,936	3,917	990	432	407	214	10
1999.....	15,714	13	14	25	1,194	3,008	5,784	3,611	940	489	442	181	13

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 5/

White, male

2007 1/.....	20,795	10	12	33	2,582	4,459	5,283	5,745	2,004	386	187	93	1
2006 1/.....	20,786	16	12	24	2,617	4,289	5,465	5,735	1,713	394	338	171	12
2005 1/.....	18,152	12	11	21	2,182	3,542	5,050	4,994	1,485	376	321	153	5
2004 1/.....	16,634	7	8	23	2,039	3,191	4,980	4,540	1,151	323	249	119	4
2003 1/.....	15,824	14	14	17	1,844	3,005	5,039	4,215	1,011	317	240	102	6
2002.....	14,170	11	3	20	1,499	2,766	4,895	3,624	795	263	193	96	5
2001.....	11,882	4	15	19	1,235	2,317	4,279	2,936	574	251	171	66	15
2000.....	10,857	7	5	21	1,068	2,158	4,061	2,584	490	196	178	80	9
1999.....	10,506	9	8	14	871	2,184	4,033	2,383	513	245	170	69	7

White, female

2007 1/.....	12,685	8	10	28	831	1,879	3,161	4,270	1,675	397	273	152	1
2006 1/.....	12,080	12	10	21	797	1,785	3,184	3,796	1,411	457	381	223	3
2005 1/.....	10,652	13	4	17	703	1,524	2,971	3,304	1,176	400	326	214	-
2004 1/.....	9,840	11	7	25	663	1,313	2,997	2,947	1,026	367	320	161	3
2003 1/.....	8,859	2	8	14	607	1,271	2,770	2,552	860	336	273	165	1
2002.....	7,976	3	15	16	537	1,124	2,627	2,224	707	320	257	143	3
2001.....	6,313	6	2	19	416	928	2,129	1,663	541	236	236	137	-
2000.....	5,531	4	2	16	349	852	1,875	1,333	500	236	229	134	1
1999.....	5,208	4	6	11	323	824	1,751	1,228	427	244	272	112	6

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 5/

Black, both sexes

2007 1/.....	4,194	11	20	11	165	479	1,040	1,617	689	111	37	12	2
2006 1/.....	4,790	15	12	11	169	572	1,319	1,815	681	124	46	25	1
2005 1/.....	4,098	10	14	10	140	529	1,144	1,567	535	91	43	15	-
2004 1/.....	3,633	12	9	12	144	439	1,073	1,385	417	80	39	19	4
2003 1/.....	3,527	9	13	9	114	446	1,196	1,291	323	85	22	14	5
2002.....	3,463	13	10	5	130	461	1,202	1,245	305	52	26	13	1
2001.....	3,165	11	8	5	116	417	1,159	1,125	224	54	25	20	1
2000.....	3,034	6	6	5	118	451	1,159	1,006	183	50	22	24	4
1999.....	3,100	8	11	4	124	497	1,196	962	198	57	20	18	5

Black, male

2007 1/.....	2,708	6	9	8	116	318	636	1,024	490	84	12	3	2
2006 1/.....	3,261	6	7	7	113	422	849	1,231	505	94	21	5	1
2005 1/.....	2,677	6	11	3	95	347	679	1,043	399	66	24	4	-
2004 1/.....	2,352	4	8	2	103	294	641	891	311	61	25	9	3
2003 1/.....	2,303	3	7	3	78	279	749	863	234	61	12	9	5
2002.....	2,307	8	6	3	91	310	754	843	234	38	17	2	1
2001.....	2,163	9	5	2	88	279	736	813	171	37	15	7	1
2000.....	2,094	4	6	2	79	311	762	735	135	36	9	11	4
1999.....	2,191	5	4	2	94	347	806	715	161	37	11	5	4

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 5/

Black, female

2007 1/.....	1,486	5	11	3	49	161	404	593	199	27	25	9	-
2006 1/.....	1,529	9	5	4	56	150	470	584	176	30	25	20	-
2005 1/.....	1,421	4	3	7	45	182	465	524	136	25	19	11	-
2004 1/.....	1,281	8	1	10	41	145	432	494	106	19	14	10	1
2003 1/.....	1,224	6	6	6	36	167	447	428	89	24	10	5	-
2002.....	1,156	5	4	2	39	151	448	402	71	14	9	11	-
2001.....	1,002	2	3	3	28	138	423	312	53	17	10	13	-
2000.....	940	2	-	3	39	140	397	271	48	14	13	13	-
1999.....	909	3	7	2	30	150	390	247	37	20	9	13	1

American Indian 2/, both sexes

2007 1/.....	388	2	1	1	52	71	102	98	50	5	3	3	-
2006 1/.....	407	1	2	-	35	82	112	116	35	13	9	2	-
2005 1/.....	362	-	1	1	38	77	98	107	23	13	3	1	-
2004 1/.....	354	-	-	3	39	73	119	83	28	6	2	-	1
2003 1/.....	295	-	-	1	27	64	109	66	22	6	-	-	-
2002.....	230	2	1	1	24	49	82	55	10	3	-	2	1
2001.....	184	-	-	1	18	41	59	46	11	5	2	1	-
2000.....	160	1	1	2	13	37	62	35	7	1	-	1	-
1999.....	164	-	-	1	12	44	69	27	8	1	1	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 5/

American Indian 2/, male

2007 1/.....	205	1	-	1	37	45	55	43	18	3	2	-	-
2006 1/.....	253	1	1	-	27	63	62	66	24	6	2	1	-
2005 1/.....	205	-	1	-	22	47	52	62	14	6	1	-	-
2004 1/.....	212	-	-	-	28	42	73	50	16	3	-	-	-
2003 1/.....	168	-	-	-	18	38	59	41	10	2	-	-	-
2002.....	136	1	-	1	18	31	43	32	6	3	-	-	1
2001.....	100	-	-	-	10	20	31	28	7	2	2	-	-
2000.....	99	1	1	2	6	29	33	22	4	1	-	-	-
1999.....	96	-	-	-	8	28	37	16	6	-	1	-	-

American Indian 2/, female

2007 1/.....	183	1	1	-	15	26	47	55	32	2	1	3	-
2006 1/.....	154	-	1	-	8	19	50	50	11	7	7	1	-
2005 1/.....	157	-	-	1	16	30	46	45	9	7	2	1	-
2004 1/.....	142	-	-	3	11	31	46	33	12	3	2	-	1
2003 1/.....	127	-	-	1	9	26	50	25	12	4	-	-	-
2002.....	94	1	1	-	6	18	39	23	4	-	-	2	-
2001.....	84	-	-	1	8	21	28	18	4	3	-	1	-
2000.....	61	-	-	-	7	8	29	13	3	-	-	1	-
1999.....	68	-	-	1	4	16	32	11	2	1	-	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 5/

Asian or Pacific Islander 3/, both sexes

2007 1/.....	309	2	1	2	45	62	85	62	35	6	5	4	-
2006 1/.....	333	-	-	2	34	59	86	83	28	20	14	7	-
2005 1/.....	277	-	-	2	32	58	70	69	21	7	13	5	-
2004 1/.....	250	-	-	-	28	45	59	66	29	9	9	5	-
2003 1/.....	218	-	-	-	27	59	44	50	23	11	2	2	-
2002.....	201	-	1	-	25	49	52	34	11	15	8	6	-
2001.....	161	1	1	1	28	36	43	26	11	7	5	2	-
2000.....	138	-	1	-	19	29	49	28	7	3	1	1	-
1999.....	150	-	-	-	21	30	46	24	15	9	4	1	-

Asian or Pacific Islander 3/, male

2007 1/.....	175	1	-	1	25	37	47	35	25	-	2	2	-
2006 1/.....	207	-	-	2	24	34	57	50	19	10	7	4	-
2005 1/.....	174	-	-	-	20	34	48	44	14	4	8	2	-
2004 1/.....	164	-	-	-	17	27	43	47	18	6	4	2	-
2003 1/.....	131	-	-	-	15	39	26	34	11	4	1	1	-
2002.....	121	-	-	-	14	32	36	18	5	7	5	4	-
2001.....	108	1	1	1	17	24	33	16	8	3	3	1	-
2000.....	87	-	1	-	13	17	34	17	2	2	-	1	-
1999.....	92	-	-	-	8	20	31	19	8	4	2	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 5/

Asian or Pacific Islander 3/, female

2007 1/.....	134	1	1	1	20	25	38	27	10	6	3	2	-
2006 1/.....	126	-	-	-	10	25	29	33	9	10	7	3	-
2005 1/.....	103	-	-	2	12	24	22	25	7	3	5	3	-
2004 1/.....	86	-	-	-	11	18	16	19	11	3	5	3	-
2003 1/.....	87	-	-	-	12	20	18	16	12	7	1	1	-
2002.....	80	-	1	-	11	17	16	16	6	8	3	2	-
2001.....	53	-	-	-	11	12	10	10	3	4	2	1	-
2000.....	51	-	-	-	6	12	15	11	5	1	1	-	-
1999.....	58	-	-	-	13	10	15	5	7	5	2	1	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/

All races, both sexes

2007.....	31,224	16	69	313	6,902	6,302	5,120	4,861	3,338	1,938	1,716	638	11
2006.....	30,896	6	57	346	7,155	6,189	5,012	4,828	3,111	1,892	1,683	608	9
2005.....	30,694	7	62	335	6,821	6,302	5,101	4,744	2,987	1,929	1,784	612	10
2004.....	29,569	7	51	300	6,540	6,021	4,986	4,588	2,859	1,885	1,729	576	27
2003.....	30,136	8	48	324	6,846	6,188	5,118	4,550	2,804	1,912	1,731	589	18
2002.....	30,242	10	61	348	6,780	6,140	5,446	4,321	2,719	1,993	1,833	576	15
2001.....	29,573	11	70	333	6,687	6,137	5,288	4,120	2,544	1,998	1,803	563	19
2000.....	28,663	13	46	377	6,575	5,789	5,358	3,951	2,272	1,941	1,723	600	18
1999.....	28,874	8	65	416	6,795	5,975	5,234	3,722	2,306	2,034	1,738	561	20

All races, male

2007.....	27,047	12	39	237	6,237	5,581	4,212	4,026	2,830	1,701	1,565	596	11
2006.....	26,712	2	38	258	6,503	5,463	4,121	3,949	2,592	1,667	1,543	567	9
2005.....	26,657	7	42	252	6,204	5,559	4,221	3,951	2,544	1,679	1,624	567	7
2004.....	25,498	2	29	231	5,895	5,302	4,130	3,733	2,391	1,663	1,554	542	26
2003.....	26,124	7	28	256	6,186	5,472	4,222	3,761	2,378	1,694	1,558	546	16
2002.....	26,098	4	38	242	6,096	5,347	4,503	3,579	2,296	1,768	1,671	540	14
2001.....	25,480	7	39	249	6,066	5,347	4,312	3,416	2,116	1,754	1,634	522	18
2000.....	24,582	9	28	302	5,906	5,003	4,363	3,258	1,896	1,684	1,571	549	13
1999.....	24,700	5	36	308	6,056	5,156	4,229	3,057	1,951	1,786	1,580	517	19

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/

All races, female

2007.....	4,177	4	30	76	665	721	908	835	508	237	151	42	-
2006.....	4,184	4	19	88	652	726	891	879	519	225	140	41	-
2005.....	4,037	-	20	83	617	743	880	793	443	250	160	45	3
2004.....	4,071	5	22	69	645	719	856	855	468	222	175	34	1
2003.....	4,012	1	20	68	660	716	896	789	426	218	173	43	2
2002.....	4,144	6	23	106	684	793	943	742	423	225	162	36	1
2001.....	4,093	4	31	84	621	790	976	704	428	244	169	41	1
2000.....	4,081	4	18	75	669	786	995	693	376	257	152	51	5
1999.....	4,174	3	29	108	739	819	1,005	665	355	248	158	44	1

White, both sexes

2007 1/.....	22,296	10	40	188	3,410	3,607	3,787	4,133	3,027	1,814	1,642	628	10
2006 1/.....	21,721	2	27	219	3,525	3,338	3,708	4,080	2,845	1,766	1,623	580	8
2005 1/.....	21,958	4	33	209	3,467	3,541	3,780	4,056	2,752	1,797	1,715	594	10
2004 1/.....	21,442	4	17	182	3,514	3,433	3,776	3,936	2,586	1,754	1,655	561	24
2003 1/.....	21,763	2	24	192	3,610	3,521	3,894	3,908	2,575	1,794	1,654	575	14
2002.....	21,902	4	42	220	3,597	3,544	4,125	3,685	2,485	1,880	1,749	558	13
2001.....	21,760	8	43	241	3,586	3,759	4,073	3,575	2,325	1,866	1,719	549	16
2000.....	20,945	6	28	275	3,544	3,477	4,123	3,361	2,056	1,809	1,666	585	15
1999.....	21,143	4	32	289	3,670	3,713	4,064	3,172	2,068	1,907	1,662	547	15

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/

White, male

2007 1/.....	19,020	8	25	138	2,996	3,105	3,076	3,413	2,562	1,596	1,501	590	10
2006 1/.....	18,496	-	18	162	3,151	2,882	3,012	3,316	2,363	1,555	1,485	544	8
2005 1/.....	18,788	4	22	163	3,103	3,025	3,071	3,366	2,345	1,573	1,560	549	7
2004 1/.....	18,223	1	10	140	3,109	2,950	3,074	3,193	2,165	1,539	1,490	529	23
2003 1/.....	18,647	1	12	156	3,216	3,040	3,173	3,219	2,187	1,598	1,496	536	13
2002.....	18,714	2	25	154	3,194	2,999	3,389	3,049	2,099	1,671	1,596	524	12
2001.....	18,527	5	25	181	3,170	3,209	3,320	2,967	1,923	1,641	1,562	509	15
2000.....	17,750	2	15	221	3,129	2,931	3,332	2,770	1,710	1,568	1,525	537	10
1999.....	17,942	4	18	215	3,224	3,145	3,270	2,616	1,740	1,677	1,514	505	14

White, female

2007 1/.....	3,276	2	15	50	414	502	711	720	465	218	141	38	-
2006 1/.....	3,225	2	9	57	374	456	696	764	482	211	138	36	-
2005 1/.....	3,170	-	11	46	364	516	709	690	407	224	155	45	3
2004 1/.....	3,219	3	7	42	405	483	702	743	421	215	165	32	1
2003 1/.....	3,116	1	12	36	394	481	721	689	388	196	158	39	1
2002.....	3,188	2	17	66	403	545	736	636	386	209	153	34	1
2001.....	3,233	3	18	60	416	550	753	608	402	225	157	40	1
2000.....	3,195	4	13	54	415	546	791	591	346	241	141	48	5
1999.....	3,201	-	14	74	446	568	794	556	328	230	148	42	1

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/

Black, both sexes

2007 1/.....	8,242	6	26	116	3,285	2,548	1,215	627	254	97	60	8	-
2006 1/.....	8,409	4	29	113	3,358	2,670	1,190	642	217	107	54	24	1
2005 1/.....	7,984	3	27	110	3,098	2,585	1,192	591	187	115	58	18	-
2004 1/.....	7,448	3	30	102	2,841	2,422	1,079	548	233	108	66	14	2
2003 1/.....	7,659	6	23	116	3,017	2,488	1,095	547	189	103	60	12	3
2002.....	7,623	6	17	103	2,955	2,431	1,188	547	190	98	68	18	2
2001.....	7,184	2	23	80	2,923	2,208	1,105	472	176	110	71	12	2
2000.....	7,054	7	13	92	2,842	2,147	1,103	510	163	114	50	12	1
1999.....	7,017	2	27	106	2,911	2,067	1,047	465	196	109	68	14	5

Black, male

2007 1/.....	7,430	4	13	90	3,050	2,345	1,038	527	223	83	53	4	-
2006 1/.....	7,563	2	19	86	3,105	2,424	1,022	551	188	94	52	19	1
2005 1/.....	7,226	3	19	80	2,881	2,379	1,040	502	157	94	53	18	-
2004 1/.....	6,709	1	17	79	2,622	2,216	947	461	191	104	57	12	2
2003 1/.....	6,882	6	15	86	2,786	2,279	941	467	159	83	49	9	2
2002.....	6,798	2	12	74	2,705	2,216	1,006	460	159	85	61	16	2
2001.....	6,438	1	13	62	2,738	2,000	906	394	155	95	61	11	2
2000.....	6,284	7	8	75	2,617	1,934	929	424	137	103	40	9	1
1999.....	6,184	1	13	79	2,647	1,854	862	382	175	95	59	12	5

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/

Black, female

2007 1/.....	812	2	13	26	235	203	177	100	31	14	7	4	-
2006 1/.....	846	2	10	27	253	246	168	91	29	13	2	5	-
2005 1/.....	758	-	8	30	217	206	152	89	30	21	5	-	-
2004 1/.....	739	2	13	23	219	206	132	87	42	4	9	2	-
2003 1/.....	777	-	8	30	231	209	154	80	30	20	11	3	1
2002.....	825	4	5	29	250	215	182	87	31	13	7	2	-
2001.....	746	1	10	18	185	208	199	78	21	15	10	1	-
2000.....	770	-	5	17	225	213	174	86	26	11	10	3	-
1999.....	833	1	14	27	264	213	185	83	21	14	9	2	-

American Indian 2/, both sexes

2007 1/.....	246	-	3	5	84	52	39	36	14	9	4	-	-
2006 1/.....	291	-	-	6	107	67	55	32	14	6	3	1	-
2005 1/.....	305	-	1	7	113	81	46	33	13	5	6	-	-
2004 1/.....	281	-	3	10	89	72	57	30	9	7	3	-	1
2003 1/.....	269	-	-	8	96	70	50	27	9	4	4	1	-
2002.....	287	-	1	10	105	68	51	33	10	5	4	-	-
2001.....	240	1	4	7	76	60	50	20	11	7	4	-	-
2000.....	240	-	4	3	84	57	44	23	14	9	1	-	1
1999.....	268	2	2	8	96	76	37	27	8	9	3	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/

American Indian 2/, male

2007 1/.....	213	-	1	5	77	46	34	28	12	6	4	-	-
2006 1/.....	253	-	-	4	100	61	42	24	12	6	3	1	-
2005 1/.....	264	-	1	3	99	72	39	29	12	3	6	-	-
2004 1/.....	236	-	2	7	77	60	52	21	8	5	3	-	1
2003 1/.....	229	-	-	6	81	61	42	24	7	4	3	1	-
2002.....	235	-	-	4	86	57	45	27	8	4	4	-	-
2001.....	196	1	1	3	68	49	38	16	10	6	4	-	-
2000.....	196	-	4	2	73	45	31	17	14	8	1	-	1
1999.....	228	-	2	6	86	64	30	21	7	9	3	-	-

American Indian 2/, female

2007 1/.....	33	-	2	-	7	6	5	8	2	3	-	-	-
2006 1/.....	38	-	-	2	7	6	13	8	2	-	-	-	-
2005 1/.....	41	-	-	4	14	9	7	4	1	2	-	-	-
2004 1/.....	45	-	1	3	12	12	5	9	1	2	-	-	-
2003 1/.....	40	-	-	2	15	9	8	3	2	-	1	-	-
2002.....	52	-	1	6	19	11	6	6	2	1	-	-	-
2001.....	44	-	3	4	8	11	12	4	1	1	-	-	-
2000.....	44	-	-	1	11	12	13	6	-	1	-	-	-
1999.....	40	2	-	2	10	12	7	6	1	-	-	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/

Asian or Pacific Islander 3/, both sexes

2007 1/.....	440	-	-	4	123	95	79	65	43	18	10	2	1
2006 1/.....	475	-	1	8	165	114	59	74	35	13	3	3	-
2005 1/.....	447	-	1	9	143	95	83	64	35	12	5	-	-
2004 1/.....	398	-	1	6	96	94	74	74	31	16	5	1	-
2003 1/.....	445	-	1	8	123	109	79	68	31	11	13	1	1
2002.....	430	-	1	15	123	97	82	56	34	10	12	-	-
2001.....	389	-	-	5	102	110	60	53	32	15	9	2	1
2000.....	424	-	1	7	105	108	88	57	39	9	6	3	1
1999.....	446	-	4	13	118	119	86	58	34	9	5	-	-

Asian or Pacific Islander 3/, male

2007 1/.....	384	-	-	4	114	85	64	58	33	16	7	2	1
2006 1/.....	400	-	1	6	147	96	45	58	29	12	3	3	-
2005 1/.....	379	-	-	6	121	83	71	54	30	9	5	-	-
2004 1/.....	330	-	-	5	87	76	57	58	27	15	4	1	-
2003 1/.....	366	-	1	8	103	92	66	51	25	9	10	-	1
2002.....	351	-	1	10	111	75	63	43	30	8	10	-	-
2001.....	319	-	-	3	90	89	48	39	28	12	7	2	1
2000.....	352	-	1	4	87	93	71	47	35	5	5	3	1
1999.....	346	-	3	8	99	93	67	38	29	5	4	-	-

Worktable 290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2007

[Race categories are consistent with the 1977 Office of Management and Budget standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/

Asian or Pacific Islander 3/, female

2007 1/.....	56	-	-	-	9	10	15	7	10	2	3	-	-
2006 1/.....	75	-	-	2	18	18	14	16	6	1	-	-	-
2005 1/.....	68	-	1	3	22	12	12	10	5	3	-	-	-
2004 1/.....	68	-	1	1	9	18	17	16	4	1	1	-	-
2003 1/.....	79	-	-	-	20	17	13	17	6	2	3	1	-
2002.....	79	-	-	5	12	22	19	13	4	2	2	-	-
2001.....	70	-	-	2	12	21	12	14	4	3	2	-	-
2000.....	72	-	-	3	18	15	17	10	4	4	1	-	-
1999.....	100	-	1	5	19	26	19	20	5	4	1	-	-

- Quantity zero.

... Category not applicable.

1/ Multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming in 2007, by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming in 2006, by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming in 2005, by California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and by California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data - mortality."

2/ Includes Aleuts and Eskimos.

3/ Includes Chinese, Filipino, Hawaiian, Japanese, and Other Asian or Pacific Islander.

4/ Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."

5/ Included in selected categories.

Source: CDC/NCHS, National Vital Statistics System, Mortality.