

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, all ages				
...	All causes	2,712,630	100.0	844.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	633,842	23.4	197.2
2	Malignant neoplasms (C00-C97)	595,930	22.0	185.4
3	Chronic lower respiratory diseases (J40-J47)	155,041	5.7	48.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	146,571	5.4	45.6
5	Cerebrovascular diseases (I60-I69)	140,323	5.2	43.7
6	Alzheimer's disease (G30)	110,561	4.1	34.4
7	Diabetes mellitus (E10-E14)	79,535	2.9	24.7
8	Influenza and pneumonia (J09-J18)	57,062	2.1	17.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	49,959	1.8	15.5
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	44,193	1.6	13.7
11	Septicemia (A40-A41)	40,773	1.5	12.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	40,326	1.5	12.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	32,200	1.2	10.0
14	Parkinson's disease (G20-G21)	27,972	1.0	8.7
15	Pneumonitis due to solids and liquids (J69)	19,803	0.7	6.2
...	All other causes (residual)	538,539	19.9	167.6
All races and origins, both sexes, 1-4 years				
...	All causes	3,965	100.0	24.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,235	31.1	7.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	435	11.0	2.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	369	9.3	2.3
4	Malignant neoplasms (C00-C97)	354	8.9	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	147	3.7	0.9
6	Influenza and pneumonia (J09-J18)	88	2.2	0.6
7	Septicemia (A40-A41)	54	1.4	0.3
8	Certain conditions originating in the perinatal period (P00-P96)	50	1.3	0.3
9	Cerebrovascular diseases (I60-I69)	42	1.1	0.3
10	Chronic lower respiratory diseases (J40-J47)	40	1.0	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	0.9	0.2
12	Anemias (D50-D64)	25	0.6	0.2
13	Complications of medical and surgical care (Y40-Y84,Y88)	18	0.5	*
14	Acute bronchitis and bronchiolitis (J20-J21)	17	0.4	*
15	Meningitis (G00,G03)	16	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.4	*
...	All other causes (residual)	1,025	25.9	6.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 5-14 years				
...	All causes	5,411	100.0	13.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,518	28.1	3.7
2	Malignant neoplasms (C00-C97)	865	16.0	2.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	413	7.6	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	337	6.2	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	298	5.5	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	210	3.9	0.5
7	Chronic lower respiratory diseases (J40-J47)	173	3.2	0.4
8	Cerebrovascular diseases (I60-I69)	84	1.6	0.2
9	Influenza and pneumonia (J09-J18)	83	1.5	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	72	1.3	0.2
11	Septicemia (A40-A41)	64	1.2	0.2
12	Anemias (D50-D64)	33	0.6	0.1
13	Diabetes mellitus (E10-E14)	23	0.4	0.1
14	Certain conditions originating in the perinatal period (P00-P96)	21	0.4	0.1
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	17	0.3	*
...	All other causes (residual)	1,183	21.9	2.9
All races and origins, both sexes, 15-24 years				
...	All causes	30,494	100.0	69.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,514	41.0	28.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,491	18.0	12.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,733	15.5	10.8
4	Malignant neoplasms (C00-C97)	1,469	4.8	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	997	3.3	2.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	386	1.3	0.9
7	Chronic lower respiratory diseases (J40-J47)	202	0.7	0.5
8	Diabetes mellitus (E10-E14)	196	0.6	0.4
9	Influenza and pneumonia (J09-J18)	184	0.6	0.4
10	Cerebrovascular diseases (I60-I69)	166	0.5	0.4
11	Pregnancy, childbirth and the puerperium (O00-O99)	161	0.5	0.4
12	Septicemia (A40-A41)	130	0.4	0.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	100	0.3	0.2
14	Legal intervention (Y35,Y89.0)	88	0.3	0.2
15	Anemias (D50-D64)	84	0.3	0.2
...	All other causes (residual)	3,593	11.8	8.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 25-44 years				
...	All causes	124,605	100.0	147.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37,613	30.2	44.4
2	Malignant neoplasms (C00-C97)	14,613	11.7	17.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,909	11.2	16.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	13,883	11.1	16.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,758	6.2	9.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,705	3.0	4.4
7	Diabetes mellitus (E10-E14)	2,784	2.2	3.3
8	Cerebrovascular diseases (I60-I69)	2,355	1.9	2.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,584	1.3	1.9
10	Septicemia (A40-A41)	1,208	1.0	1.4
11	Influenza and pneumonia (J09-J18)	1,105	0.9	1.3
12	Chronic lower respiratory diseases (J40-J47)	990	0.8	1.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	948	0.8	1.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	926	0.7	1.1
15	Pregnancy, childbirth and the puerperium (O00-O99)	734	0.6	0.9
...	All other causes (residual)	20,490	16.4	24.2
All races and origins, both sexes, 45-64 years				
...	All causes	532,279	100.0	633.2
1	Malignant neoplasms (C00-C97)	159,176	29.9	189.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	111,120	20.9	132.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	40,987	7.7	48.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	22,152	4.2	26.4
5	Chronic lower respiratory diseases (J40-J47)	21,802	4.1	25.9
6	Diabetes mellitus (E10-E14)	20,378	3.8	24.2
7	Cerebrovascular diseases (I60-I69)	17,423	3.3	20.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	16,490	3.1	19.6
9	Septicemia (A40-A41)	8,316	1.6	9.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,576	1.4	9.0
11	Influenza and pneumonia (J09-J18)	6,651	1.2	7.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,072	1.0	6.0
13	Viral hepatitis (B15-B19)	5,055	0.9	6.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3,851	0.7	4.6
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,415	0.6	4.1
...	All other causes (residual)	82,815	15.6	98.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 65 years and over				
...	All causes	1,992,283	100.0	4,171.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	507,138	25.5	1,061.8
2	Malignant neoplasms (C00-C97)	419,389	21.1	878.1
3	Chronic lower respiratory diseases (J40-J47)	131,804	6.6	276.0
4	Cerebrovascular diseases (I60-I69)	120,156	6.0	251.6
5	Alzheimer's disease (G30)	109,495	5.5	229.3
6	Diabetes mellitus (E10-E14)	56,142	2.8	117.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	51,395	2.6	107.6
8	Influenza and pneumonia (J09-J18)	48,774	2.4	102.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41,258	2.1	86.4
10	Septicemia (A40-A41)	30,817	1.5	64.5
11	Parkinson's disease (G20-G21)	27,269	1.4	57.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	26,515	1.3	55.5
13	Pneumonitis due to solids and liquids (J69)	17,474	0.9	36.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	14,432	0.7	30.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13,339	0.7	27.9
...	All other causes (residual)	376,886	18.9	789.1
All races and origins, male, all ages				
...	All causes	1,373,404	100.0	868.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	335,002	24.4	211.7
2	Malignant neoplasms (C00-C97)	313,818	22.8	198.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	92,919	6.8	58.7
4	Chronic lower respiratory diseases (J40-J47)	72,498	5.3	45.8
5	Cerebrovascular diseases (I60-I69)	58,288	4.2	36.8
6	Diabetes mellitus (E10-E14)	43,123	3.1	27.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33,994	2.5	21.5
8	Alzheimer's disease (G30)	33,690	2.5	21.3
9	Influenza and pneumonia (J09-J18)	26,903	2.0	17.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	25,666	1.9	16.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25,441	1.9	16.1
12	Septicemia (A40-A41)	19,385	1.4	12.3
13	Parkinson's disease (G20-G21)	16,867	1.2	10.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14,274	1.0	9.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13,934	1.0	8.8
...	All other causes (residual)	247,602	18.0	156.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 1-4 years				
...	All causes	2,281	100.0	28.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	768	33.7	9.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	230	10.1	2.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	227	10.0	2.8
4	Malignant neoplasms (C00-C97)	196	8.6	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	78	3.4	1.0
6	Influenza and pneumonia (J09-J18)	49	2.1	0.6
7	Certain conditions originating in the perinatal period (P00-P96)	34	1.5	0.4
8	Chronic lower respiratory diseases (J40-J47)	27	1.2	0.3
9	Cerebrovascular diseases (I60-I69)	26	1.1	0.3
10	Septicemia (A40-A41)	24	1.1	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.7	*
12	Anemias (D50-D64)	12	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	12	0.5	*
14	Acute bronchitis and bronchiolitis (J20-J21)	11	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.4	*
...	All other causes (residual)	561	24.6	6.9
All races and origins, male, 5-14 years				
...	All causes	3,153	100.0	15.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	939	29.8	4.5
2	Malignant neoplasms (C00-C97)	469	14.9	2.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	255	8.1	1.2
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	194	6.2	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	194	6.2	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	108	3.4	0.5
7	Chronic lower respiratory diseases (J40-J47)	104	3.3	0.5
8	Cerebrovascular diseases (I60-I69)	45	1.4	0.2
9	Influenza and pneumonia (J09-J18)	41	1.3	0.2
10	Septicemia (A40-A41)	33	1.0	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	1.0	0.2
12	Anemias (D50-D64)	14	0.4	*
13	Diabetes mellitus (E10-E14)	11	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.3	*
13	Certain conditions originating in the perinatal period (P00-P96)	11	0.3	*
...	All other causes (residual)	691	21.9	3.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 15-24 years				
...	All causes	22,346	100.0	99.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,149	40.9	40.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,359	19.5	19.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,134	18.5	18.4
4	Malignant neoplasms (C00-C97)	880	3.9	3.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	667	3.0	3.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	220	1.0	1.0
7	Chronic lower respiratory diseases (J40-J47)	130	0.6	0.6
8	Diabetes mellitus (E10-E14)	118	0.5	0.5
9	Influenza and pneumonia (J09-J18)	107	0.5	0.5
10	Cerebrovascular diseases (I60-I69)	89	0.4	0.4
11	Legal intervention (Y35,Y89.0)	84	0.4	0.4
12	Septicemia (A40-A41)	71	0.3	0.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	64	0.3	0.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	46	0.2	0.2
15	Anemias (D50-D64)	42	0.2	0.2
...	All other causes (residual)	2,186	9.8	9.7
All races and origins, male, 25-44 years				
...	All causes	81,451	100.0	191.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27,225	33.4	64.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,726	13.2	25.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,391	11.5	22.1
4	Malignant neoplasms (C00-C97)	6,474	7.9	15.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,430	7.9	15.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,331	2.9	5.5
7	Diabetes mellitus (E10-E14)	1,706	2.1	4.0
8	Cerebrovascular diseases (I60-I69)	1,310	1.6	3.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,071	1.3	2.5
10	Influenza and pneumonia (J09-J18)	636	0.8	1.5
11	Septicemia (A40-A41)	591	0.7	1.4
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	527	0.6	1.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	510	0.6	1.2
14	Chronic lower respiratory diseases (J40-J47)	496	0.6	1.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	384	0.5	0.9
...	All other causes (residual)	11,643	14.3	27.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 45-64 years				
...	All causes	323,173	100.0	788.0
1	Malignant neoplasms (C00-C97)	85,351	26.4	208.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	76,979	23.8	187.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27,797	8.6	67.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	14,841	4.6	36.2
5	Diabetes mellitus (E10-E14)	12,570	3.9	30.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12,107	3.7	29.5
7	Chronic lower respiratory diseases (J40-J47)	11,009	3.4	26.8
8	Cerebrovascular diseases (I60-I69)	9,870	3.1	24.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,448	1.4	10.8
10	Septicemia (A40-A41)	4,428	1.4	10.8
11	Influenza and pneumonia (J09-J18)	3,752	1.2	9.1
12	Viral hepatitis (B15-B19)	3,538	1.1	8.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,224	1.0	7.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	2,935	0.9	7.2
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,578	0.8	6.3
...	All other causes (residual)	47,746	14.8	116.4
All races and origins, male, 65 years and over				
...	All causes	927,897	100.0	4,399.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	247,607	26.7	1,174.0
2	Malignant neoplasms (C00-C97)	220,411	23.8	1,045.1
3	Chronic lower respiratory diseases (J40-J47)	60,714	6.5	287.9
4	Cerebrovascular diseases (I60-I69)	46,890	5.1	222.3
5	Alzheimer's disease (G30)	33,225	3.6	157.5
6	Diabetes mellitus (E10-E14)	28,711	3.1	136.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26,272	2.8	124.6
8	Influenza and pneumonia (J09-J18)	22,204	2.4	105.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20,396	2.2	96.7
10	Parkinson's disease (G20-G21)	16,388	1.8	77.7
11	Septicemia (A40-A41)	14,139	1.5	67.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10,315	1.1	48.9
13	Pneumonitis due to solids and liquids (J69)	9,474	1.0	44.9
14	Chronic liver disease and cirrhosis (K70,K73-K74)	8,470	0.9	40.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7,002	0.8	33.2
...	All other causes (residual)	155,679	16.8	738.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, all ages				
...	All causes	1,339,226	100.0	820.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	298,840	22.3	183.1
2	Malignant neoplasms (C00-C97)	282,112	21.1	172.9
3	Chronic lower respiratory diseases (J40-J47)	82,543	6.2	50.6
4	Cerebrovascular diseases (I60-I69)	82,035	6.1	50.3
5	Alzheimer's disease (G30)	76,871	5.7	47.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	53,652	4.0	32.9
7	Diabetes mellitus (E10-E14)	36,412	2.7	22.3
8	Influenza and pneumonia (J09-J18)	30,159	2.3	18.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24,518	1.8	15.0
10	Septicemia (A40-A41)	21,388	1.6	13.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18,266	1.4	11.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	14,660	1.1	9.0
13	Parkinson's disease (G20-G21)	11,105	0.8	6.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,199	0.8	6.2
15	Pneumonitis due to solids and liquids (J69)	8,892	0.7	5.4
...	All other causes (residual)	287,574	21.5	176.2
All races and origins, female, 1-4 years				
...	All causes	1,684	100.0	21.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	467	27.7	6.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	208	12.4	2.7
3	Malignant neoplasms (C00-C97)	158	9.4	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	139	8.3	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	69	4.1	0.9
6	Influenza and pneumonia (J09-J18)	39	2.3	0.5
7	Septicemia (A40-A41)	30	1.8	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	1.0	*
9	Cerebrovascular diseases (I60-I69)	16	1.0	*
9	Certain conditions originating in the perinatal period (P00-P96)	16	1.0	*
11	Anemias (D50-D64)	13	0.8	*
11	Chronic lower respiratory diseases (J40-J47)	13	0.8	*
13	Meningitis (G00,G03)	11	0.7	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	6	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.4	*
...	All other causes (residual)	469	27.9	6.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015



LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 5-14 years				
...	All causes	2,258	100.0	11.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	579	25.6	2.9
2	Malignant neoplasms (C00-C97)	396	17.5	2.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	158	7.0	0.8
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	143	6.3	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	104	4.6	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	102	4.5	0.5
7	Chronic lower respiratory diseases (J40-J47)	69	3.1	0.3
8	Influenza and pneumonia (J09-J18)	42	1.9	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	39	1.7	0.2
9	Cerebrovascular diseases (I60-I69)	39	1.7	0.2
11	Septicemia (A40-A41)	31	1.4	0.2
12	Anemias (D50-D64)	19	0.8	*
13	Diabetes mellitus (E10-E14)	12	0.5	*
14	Certain conditions originating in the perinatal period (P00-P96)	10	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.4	*
...	All other causes (residual)	506	22.4	2.5
All races and origins, female, 15-24 years				
...	All causes	8,148	100.0	38.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,365	41.3	15.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,132	13.9	5.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	599	7.4	2.8
4	Malignant neoplasms (C00-C97)	589	7.2	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	330	4.1	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	166	2.0	0.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	161	2.0	0.8
8	Diabetes mellitus (E10-E14)	78	1.0	0.4
9	Cerebrovascular diseases (I60-I69)	77	0.9	0.4
9	Influenza and pneumonia (J09-J18)	77	0.9	0.4
11	Chronic lower respiratory diseases (J40-J47)	72	0.9	0.3
12	Septicemia (A40-A41)	59	0.7	0.3
13	Anemias (D50-D64)	42	0.5	0.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	0.5	0.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	36	0.4	0.2
...	All other causes (residual)	1,328	16.3	6.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 25-44 years				
...	All causes	43,154	100.0	102.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,388	24.1	24.6
2	Malignant neoplasms (C00-C97)	8,139	18.9	19.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,518	10.5	10.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,157	7.3	7.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,374	3.2	3.3
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,328	3.1	3.1
7	Diabetes mellitus (E10-E14)	1,078	2.5	2.6
8	Cerebrovascular diseases (I60-I69)	1,045	2.4	2.5
9	Pregnancy, childbirth and the puerperium (O00-O99)	734	1.7	1.7
10	Septicemia (A40-A41)	617	1.4	1.5
11	Human immunodeficiency virus (HIV) disease (B20-B24)	513	1.2	1.2
12	Chronic lower respiratory diseases (J40-J47)	494	1.1	1.2
13	Influenza and pneumonia (J09-J18)	469	1.1	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	438	1.0	1.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	399	0.9	0.9
...	All other causes (residual)	8,463	19.6	20.0
All races and origins, female, 45-64 years				
...	All causes	209,106	100.0	485.7
1	Malignant neoplasms (C00-C97)	73,825	35.3	171.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,141	16.3	79.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,190	6.3	30.6
4	Chronic lower respiratory diseases (J40-J47)	10,793	5.2	25.1
5	Diabetes mellitus (E10-E14)	7,808	3.7	18.1
6	Cerebrovascular diseases (I60-I69)	7,553	3.6	17.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	7,311	3.5	17.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,383	2.1	10.2
9	Septicemia (A40-A41)	3,888	1.9	9.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,128	1.5	7.3
11	Influenza and pneumonia (J09-J18)	2,899	1.4	6.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,848	0.9	4.3
13	Viral hepatitis (B15-B19)	1,517	0.7	3.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	991	0.5	2.3
15	Human immunodeficiency virus (HIV) disease (B20-B24)	916	0.4	2.1
...	All other causes (residual)	34,915	16.7	81.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 65 years and over				
...	All causes	1,064,386	100.0	3,990.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	259,531	24.4	973.1
2	Malignant neoplasms (C00-C97)	198,978	18.7	746.1
3	Alzheimer's disease (G30)	76,270	7.2	286.0
4	Cerebrovascular diseases (I60-I69)	73,266	6.9	274.7
5	Chronic lower respiratory diseases (J40-J47)	71,090	6.7	266.5
6	Diabetes mellitus (E10-E14)	27,431	2.6	102.9
7	Influenza and pneumonia (J09-J18)	26,570	2.5	99.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	25,123	2.4	94.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20,862	2.0	78.2
10	Septicemia (A40-A41)	16,678	1.6	62.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16,200	1.5	60.7
12	Parkinson's disease (G20-G21)	10,881	1.0	40.8
13	Pneumonitis due to solids and liquids (J69)	8,000	0.8	30.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6,337	0.6	23.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	5,962	0.6	22.4
...	All other causes (residual)	221,207	20.8	829.4
Hispanic, both sexes, all ages				
...	All causes	179,457	100.0	317.1
1	Malignant neoplasms (C00-C97)	37,804	21.1	66.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,401	20.3	64.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,806	7.7	24.4
4	Cerebrovascular diseases (I60-I69)	9,795	5.5	17.3
5	Diabetes mellitus (E10-E14)	8,278	4.6	14.6
6	Alzheimer's disease (G30)	6,444	3.6	11.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	6,018	3.4	10.6
8	Chronic lower respiratory diseases (J40-J47)	5,159	2.9	9.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,581	2.0	6.3
10	Influenza and pneumonia (J09-J18)	3,497	1.9	6.2
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,303	1.8	5.8
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,886	1.6	5.1
13	Septicemia (A40-A41)	2,692	1.5	4.8
14	Certain conditions originating in the perinatal period (P00-P96)	2,454	1.4	4.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,384	1.3	4.2
...	All other causes (residual)	34,955	19.5	61.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 1-4 years				
...	All causes	815	100.0	19.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	258	31.7	6.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	115	14.1	2.8
3	Malignant neoplasms (C00-C97)	75	9.2	1.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	69	8.5	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	3.6	0.7
6	Influenza and pneumonia (J09-J18)	18	2.2	*
7	Septicemia (A40-A41)	17	2.1	*
8	Cerebrovascular diseases (I60-I69)	7	0.9	*
8	Certain conditions originating in the perinatal period (P00-P96)	7	0.9	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.7	*
11	Anemias (D50-D64)	4	0.5	*
11	Meningitis (G00,G03)	4	0.5	*
11	Acute bronchitis and bronchiolitis (J20-J21)	4	0.5	*
11	Chronic lower respiratory diseases (J40-J47)	4	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.4	*
...	All other causes (residual)	192	23.6	4.7
Hispanic, both sexes, 5-14 years				
...	All causes	1,063	100.0	10.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	308	29.0	3.0
2	Malignant neoplasms (C00-C97)	211	19.8	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	69	6.5	0.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	5.0	0.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	46	4.3	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	3.3	0.3
7	Chronic lower respiratory diseases (J40-J47)	23	2.2	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	1.6	*
9	Septicemia (A40-A41)	16	1.5	*
9	Cerebrovascular diseases (I60-I69)	16	1.5	*
11	Influenza and pneumonia (J09-J18)	14	1.3	*
12	Anemias (D50-D64)	5	0.5	*
13	Meningitis (G00,G03)	4	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	4	0.4	*
15	Diabetes mellitus (E10-E14)	3	0.3	*
15	Acute bronchitis and bronchiolitis (J20-J21)	3	0.3	*
15	Pneumonitis due to solids and liquids (J69)	3	0.3	*
...	All other causes (residual)	233	21.9	2.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 15-24 years				
...	All causes	5,353	100.0	56.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,261	42.2	23.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	921	17.2	9.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	807	15.1	8.5
4	Malignant neoplasms (C00-C97)	332	6.2	3.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	145	2.7	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	69	1.3	0.7
7	Influenza and pneumonia (J09-J18)	36	0.7	0.4
8	Cerebrovascular diseases (I60-I69)	32	0.6	0.3
9	Septicemia (A40-A41)	30	0.6	0.3
10	Chronic lower respiratory diseases (J40-J47)	29	0.5	0.3
11	Pregnancy, childbirth and the puerperium (O00-O99)	27	0.5	0.3
12	Diabetes mellitus (E10-E14)	20	0.4	0.2
13	Legal intervention (Y35,Y89.0)	17	0.3	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.2	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	12	0.2	*
...	All other causes (residual)	602	11.2	6.3
Hispanic, both sexes, 25-44 years				
...	All causes	16,672	100.0	97.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,843	29.0	28.2
2	Malignant neoplasms (C00-C97)	2,432	14.6	14.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,521	9.1	8.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,351	8.1	7.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,297	7.8	7.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	763	4.6	4.4
7	Cerebrovascular diseases (I60-I69)	442	2.7	2.6
8	Diabetes mellitus (E10-E14)	380	2.3	2.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	287	1.7	1.7
10	Influenza and pneumonia (J09-J18)	158	0.9	0.9
11	Septicemia (A40-A41)	151	0.9	0.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	138	0.8	0.8
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	109	0.7	0.6
14	Pregnancy, childbirth and the puerperium (O00-O99)	107	0.6	0.6
15	Chronic lower respiratory diseases (J40-J47)	86	0.5	0.5
...	All other causes (residual)	2,607	15.6	15.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 45-64 years				
...	All causes	43,176	100.0	398.7
1	Malignant neoplasms (C00-C97)	12,194	28.2	112.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,940	18.4	73.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,527	8.2	32.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,244	7.5	30.0
5	Diabetes mellitus (E10-E14)	2,388	5.5	22.0
6	Cerebrovascular diseases (I60-I69)	1,772	4.1	16.4
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	822	1.9	7.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	782	1.8	7.2
9	Viral hepatitis (B15-B19)	773	1.8	7.1
10	Septicemia (A40-A41)	690	1.6	6.4
11	Chronic lower respiratory diseases (J40-J47)	659	1.5	6.1
12	Influenza and pneumonia (J09-J18)	561	1.3	5.2
13	Human immunodeficiency virus (HIV) disease (B20-B24)	504	1.2	4.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	434	1.0	4.0
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	433	1.0	4.0
...	All other causes (residual)	6,453	14.9	59.6
Hispanic, both sexes, 65 years and over				
...	All causes	107,567	100.0	2,851.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,686	24.8	707.3
2	Malignant neoplasms (C00-C97)	22,547	21.0	597.6
3	Cerebrovascular diseases (I60-I69)	7,506	7.0	198.9
4	Alzheimer's disease (G30)	6,357	5.9	168.5
5	Diabetes mellitus (E10-E14)	5,484	5.1	145.4
6	Chronic lower respiratory diseases (J40-J47)	4,353	4.0	115.4
7	Influenza and pneumonia (J09-J18)	2,680	2.5	71.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,628	2.4	69.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,421	2.3	64.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	2,003	1.9	53.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,865	1.7	49.4
12	Septicemia (A40-A41)	1,747	1.6	46.3
13	Parkinson's disease (G20-G21)	1,484	1.4	39.3
14	Pneumonitis due to solids and liquids (J69)	831	0.8	22.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	600	0.6	15.9
...	All other causes (residual)	18,375	17.1	487.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, all ages				
...	All causes	98,170	100.0	343.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,225	20.6	70.7
2	Malignant neoplasms (C00-C97)	19,847	20.2	69.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,067	10.3	35.2
4	Cerebrovascular diseases (I60-I69)	4,544	4.6	15.9
5	Diabetes mellitus (E10-E14)	4,426	4.5	15.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,109	4.2	14.4
7	Chronic lower respiratory diseases (J40-J47)	2,606	2.7	9.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,587	2.6	9.0
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,391	2.4	8.4
10	Alzheimer's disease (G30)	2,035	2.1	7.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,837	1.9	6.4
12	Influenza and pneumonia (J09-J18)	1,735	1.8	6.1
13	Septicemia (A40-A41)	1,370	1.4	4.8
14	Certain conditions originating in the perinatal period (P00-P96)	1,368	1.4	4.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,139	1.2	4.0
...	All other causes (residual)	17,884	18.2	62.5
Hispanic, male, 1-4 years				
...	All causes	461	100.0	22.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	154	33.4	7.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	12.6	2.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	45	9.8	2.1
4	Malignant neoplasms (C00-C97)	40	8.7	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	4.6	1.0
6	Influenza and pneumonia (J09-J18)	12	2.6	*
7	Septicemia (A40-A41)	8	1.7	*
8	Certain conditions originating in the perinatal period (P00-P96)	5	1.1	*
9	Cerebrovascular diseases (I60-I69)	3	0.7	*
9	Chronic lower respiratory diseases (J40-J47)	3	0.7	*
11	Meningitis (G00,G03)	2	0.4	*
11	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
14	Salmonella infections (A01-A02)	1	0.2	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
14	Anemias (D50-D64)	1	0.2	*
14	Diabetes mellitus (E10-E14)	1	0.2	*
14	Nutritional deficiencies (E40-E64)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (residual)	99	21.5	4.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 5-14 years				
...	All causes	633	100.0	12.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	206	32.5	4.0
2	Malignant neoplasms (C00-C97)	121	19.1	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	43	6.8	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	42	6.6	0.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21	3.3	0.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	2.2	*
7	Chronic lower respiratory diseases (J40-J47)	13	2.1	*
8	Septicemia (A40-A41)	12	1.9	*
9	Cerebrovascular diseases (I60-I69)	8	1.3	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.9	*
11	Influenza and pneumonia (J09-J18)	5	0.8	*
12	Anemias (D50-D64)	2	0.3	*
12	Meningitis (G00,G03)	2	0.3	*
12	Aortic aneurysm and dissection (I71)	2	0.3	*
12	Peptic ulcer (K25-K28)	2	0.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
12	Certain conditions originating in the perinatal period (P00-P96)	2	0.3	*
...	All other causes (residual)	130	20.5	2.5
Hispanic, male, 15-24 years				
...	All causes	4,039	100.0	82.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,748	43.3	35.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	809	20.0	16.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	629	15.6	12.8
4	Malignant neoplasms (C00-C97)	208	5.1	4.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	97	2.4	2.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	0.9	0.7
7	Influenza and pneumonia (J09-J18)	21	0.5	0.4
7	Chronic lower respiratory diseases (J40-J47)	21	0.5	0.4
9	Septicemia (A40-A41)	19	0.5	*
9	Cerebrovascular diseases (I60-I69)	19	0.5	*
11	Legal intervention (Y35,Y89.0)	17	0.4	*
12	Diabetes mellitus (E10-E14)	15	0.4	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	9	0.2	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.1	*
15	Aortic aneurysm and dissection (I71)	5	0.1	*
15	Pneumonitis due to solids and liquids (J69)	5	0.1	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	5	0.1	*
...	All other causes (residual)	370	9.2	7.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015



LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 25-44 years				
...	All causes	11,653	100.0	131.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,858	33.1	43.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,128	9.7	12.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,087	9.3	12.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,085	9.3	12.2
5	Malignant neoplasms (C00-C97)	1,073	9.2	12.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	560	4.8	6.3
7	Cerebrovascular diseases (I60-I69)	287	2.5	3.2
8	Diabetes mellitus (E10-E14)	258	2.2	2.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	242	2.1	2.7
10	Influenza and pneumonia (J09-J18)	101	0.9	1.1
11	Septicemia (A40-A41)	93	0.8	1.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	76	0.7	0.9
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	62	0.5	0.7
14	Legal intervention (Y35,Y89.0)	60	0.5	0.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	58	0.5	0.7
...	All other causes (residual)	1,625	13.9	18.3
Hispanic, male, 45-64 years				
...	All causes	27,596	100.0	513.2
1	Malignant neoplasms (C00-C97)	6,424	23.3	119.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,802	21.0	107.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,684	9.7	49.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,420	8.8	45.0
5	Diabetes mellitus (E10-E14)	1,502	5.4	27.9
6	Cerebrovascular diseases (I60-I69)	1,059	3.8	19.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	615	2.2	11.4
8	Viral hepatitis (B15-B19)	568	2.1	10.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	479	1.7	8.9
10	Human immunodeficiency virus (HIV) disease (B20-B24)	413	1.5	7.7
11	Septicemia (A40-A41)	394	1.4	7.3
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	341	1.2	6.3
13	Chronic lower respiratory diseases (J40-J47)	335	1.2	6.2
14	Influenza and pneumonia (J09-J18)	316	1.1	5.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	282	1.0	5.2
...	All other causes (residual)	3,962	14.4	73.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 65 years and over				
...	All causes	51,164	100.0	3,138.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,148	25.7	806.5
2	Malignant neoplasms (C00-C97)	11,973	23.4	734.4
3	Cerebrovascular diseases (I60-I69)	3,156	6.2	193.6
4	Diabetes mellitus (E10-E14)	2,649	5.2	162.5
5	Chronic lower respiratory diseases (J40-J47)	2,188	4.3	134.2
6	Alzheimer's disease (G30)	1,999	3.9	122.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,312	2.6	80.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,263	2.5	77.5
9	Influenza and pneumonia (J09-J18)	1,260	2.5	77.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,124	2.2	68.9
11	Septicemia (A40-A41)	824	1.6	50.5
12	Parkinson's disease (G20-G21)	821	1.6	50.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	795	1.6	48.8
14	Pneumonitis due to solids and liquids (J69)	429	0.8	26.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	296	0.6	18.2
...	All other causes (residual)	7,927	15.5	486.2
Hispanic, female, all ages				
...	All causes	81,287	100.0	290.4
1	Malignant neoplasms (C00-C97)	17,957	22.1	64.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,176	19.9	57.8
3	Cerebrovascular diseases (I60-I69)	5,251	6.5	18.8
4	Alzheimer's disease (G30)	4,409	5.4	15.8
5	Diabetes mellitus (E10-E14)	3,852	4.7	13.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,739	4.6	13.4
7	Chronic lower respiratory diseases (J40-J47)	2,553	3.1	9.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,909	2.3	6.8
9	Influenza and pneumonia (J09-J18)	1,762	2.2	6.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,744	2.1	6.2
11	Septicemia (A40-A41)	1,322	1.6	4.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,245	1.5	4.4
13	Certain conditions originating in the perinatal period (P00-P96)	1,086	1.3	3.9
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	837	1.0	3.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	716	0.9	2.6
...	All other causes (residual)	16,729	20.6	59.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 1-4 years				
...	All causes	354	100.0	17.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	104	29.4	5.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	16.1	2.8
3	Malignant neoplasms (C00-C97)	35	9.9	1.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	24	6.8	1.2
5	Septicemia (A40-A41)	9	2.5	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	2.3	*
7	Influenza and pneumonia (J09-J18)	6	1.7	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.4	*
9	Cerebrovascular diseases (I60-I69)	4	1.1	*
10	Anemias (D50-D64)	3	0.8	*
11	Meningitis (G00,G03)	2	0.6	*
11	Acute bronchitis and bronchiolitis (J20-J21)	2	0.6	*
11	Certain conditions originating in the perinatal period (P00-P96)	2	0.6	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.6	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.3	*
15	Chronic lower respiratory diseases (J40-J47)	1	0.3	*
15	Pneumonitis due to solids and liquids (J69)	1	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.3	*
...	All other causes (residual)	86	24.3	4.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-14 years				
...	All causes	430	100.0	8.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	102	23.7	2.0
2	Malignant neoplasms (C00-C97)	90	20.9	1.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	26	6.0	0.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25	5.8	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	4.9	0.4
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	2.6	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	2.6	*
8	Chronic lower respiratory diseases (J40-J47)	10	2.3	*
9	Influenza and pneumonia (J09-J18)	9	2.1	*
10	Cerebrovascular diseases (I60-I69)	8	1.9	*
11	Septicemia (A40-A41)	4	0.9	*
12	Anemias (D50-D64)	3	0.7	*
12	Diabetes mellitus (E10-E14)	3	0.7	*
12	Pneumonitis due to solids and liquids (J69)	3	0.7	*
15	Meningitis (G00,G03)	2	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	2	0.5	*
15	Certain conditions originating in the perinatal period (P00-P96)	2	0.5	*
...	All other causes (residual)	98	22.8	2.0
Hispanic, female, 15-24 years				
...	All causes	1,314	100.0	28.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	513	39.0	11.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	178	13.5	3.9
3	Malignant neoplasms (C00-C97)	124	9.4	2.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	112	8.5	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	48	3.7	1.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	2.5	0.7
7	Pregnancy, childbirth and the puerperium (O00-O99)	27	2.1	0.6
8	Influenza and pneumonia (J09-J18)	15	1.1	*
9	Cerebrovascular diseases (I60-I69)	13	1.0	*
10	Septicemia (A40-A41)	11	0.8	*
11	Chronic lower respiratory diseases (J40-J47)	8	0.6	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.5	*
13	Diabetes mellitus (E10-E14)	5	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
15	Meningitis (G00,G03)	4	0.3	*
...	All other causes (residual)	211	16.1	4.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-44 years				
...	All causes	5,019	100.0	60.7
1	Malignant neoplasms (C00-C97)	1,359	27.1	16.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	985	19.6	11.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	393	7.8	4.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	266	5.3	3.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	210	4.2	2.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	203	4.0	2.5
7	Cerebrovascular diseases (I60-I69)	155	3.1	1.9
8	Diabetes mellitus (E10-E14)	122	2.4	1.5
9	Pregnancy, childbirth and the puerperium (O00-O99)	107	2.1	1.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	62	1.2	0.7
11	Septicemia (A40-A41)	58	1.2	0.7
12	Influenza and pneumonia (J09-J18)	57	1.1	0.7
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	47	0.9	0.6
14	Human immunodeficiency virus (HIV) disease (B20-B24)	45	0.9	0.5
15	Chronic lower respiratory diseases (J40-J47)	44	0.9	0.5
...	All other causes (residual)	906	18.1	11.0
Hispanic, female, 45-64 years				
...	All causes	15,580	100.0	285.7
1	Malignant neoplasms (C00-C97)	5,770	37.0	105.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,138	13.7	39.2
3	Diabetes mellitus (E10-E14)	886	5.7	16.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	843	5.4	15.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	824	5.3	15.1
6	Cerebrovascular diseases (I60-I69)	713	4.6	13.1
7	Chronic lower respiratory diseases (J40-J47)	324	2.1	5.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	303	1.9	5.6
9	Septicemia (A40-A41)	296	1.9	5.4
10	Influenza and pneumonia (J09-J18)	245	1.6	4.5
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	207	1.3	3.8
12	Viral hepatitis (B15-B19)	205	1.3	3.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	152	1.0	2.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	92	0.6	1.7
15	Human immunodeficiency virus (HIV) disease (B20-B24)	91	0.6	1.7
...	All other causes (residual)	2,491	16.0	45.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65 years and over				
...	All causes	56,403	100.0	2,632.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,538	24.0	631.9
2	Malignant neoplasms (C00-C97)	10,574	18.7	493.5
3	Alzheimer's disease (G30)	4,358	7.7	203.4
4	Cerebrovascular diseases (I60-I69)	4,350	7.7	203.0
5	Diabetes mellitus (E10-E14)	2,835	5.0	132.3
6	Chronic lower respiratory diseases (J40-J47)	2,165	3.8	101.0
7	Influenza and pneumonia (J09-J18)	1,420	2.5	66.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,365	2.4	63.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,109	2.0	51.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,070	1.9	49.9
11	Septicemia (A40-A41)	923	1.6	43.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	879	1.6	41.0
13	Parkinson's disease (G20-G21)	663	1.2	30.9
14	Pneumonitis due to solids and liquids (J69)	402	0.7	18.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	304	0.5	14.2
...	All other causes (residual)	10,448	18.5	487.6
Non-Hispanic, white, both sexes, all ages				
...	All causes	2,123,631	100.0	1,055.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	503,172	23.7	250.0
2	Malignant neoplasms (C00-C97)	467,208	22.0	232.2
3	Chronic lower respiratory diseases (J40-J47)	136,228	6.4	67.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	111,827	5.3	55.6
5	Cerebrovascular diseases (I60-I69)	106,830	5.0	53.1
6	Alzheimer's disease (G30)	93,325	4.4	46.4
7	Diabetes mellitus (E10-E14)	53,590	2.5	26.6
8	Influenza and pneumonia (J09-J18)	45,242	2.1	22.5
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36,465	1.7	18.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35,481	1.7	17.6
11	Septicemia (A40-A41)	30,240	1.4	15.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	29,343	1.4	14.6
13	Parkinson's disease (G20-G21)	24,339	1.1	12.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	22,628	1.1	11.2
15	Pneumonitis due to solids and liquids (J69)	16,430	0.8	8.2
...	All other causes (residual)	411,283	19.4	204.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 1-4 years				
...	All causes	1,895	100.0	22.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	631	33.3	7.6
2	Malignant neoplasms (C00-C97)	206	10.9	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	196	10.3	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	145	7.7	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	2.9	0.7
6	Influenza and pneumonia (J09-J18)	37	2.0	0.4
7	Septicemia (A40-A41)	20	1.1	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.0	*
9	Cerebrovascular diseases (I60-I69)	17	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	14	0.7	*
11	Chronic lower respiratory diseases (J40-J47)	11	0.6	*
12	Acute bronchitis and bronchiolitis (J20-J21)	10	0.5	*
13	Meningitis (G00,G03)	8	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.4	*
15	Anemias (D50-D64)	7	0.4	*
...	All other causes (residual)	511	27.0	6.2
Non-Hispanic, white, both sexes, 5-14 years				
...	All causes	2,835	100.0	12.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	798	28.1	3.6
2	Malignant neoplasms (C00-C97)	476	16.8	2.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	291	10.3	1.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	186	6.6	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	109	3.8	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	101	3.6	0.5
7	Influenza and pneumonia (J09-J18)	47	1.7	0.2
8	Chronic lower respiratory diseases (J40-J47)	45	1.6	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	1.4	0.2
9	Cerebrovascular diseases (I60-I69)	41	1.4	0.2
11	Septicemia (A40-A41)	28	1.0	0.1
12	Diabetes mellitus (E10-E14)	11	0.4	*
13	Anemias (D50-D64)	10	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	10	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.3	*
...	All other causes (residual)	633	22.3	2.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 15-24 years				
...	All causes	16,536	100.0	67.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,079	48.9	33.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,677	22.2	15.0
3	Malignant neoplasms (C00-C97)	781	4.7	3.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	683	4.1	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	495	3.0	2.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	224	1.4	0.9
7	Influenza and pneumonia (J09-J18)	107	0.6	0.4
8	Diabetes mellitus (E10-E14)	104	0.6	0.4
9	Cerebrovascular diseases (I60-I69)	83	0.5	0.3
10	Chronic lower respiratory diseases (J40-J47)	72	0.4	0.3
11	Pregnancy, childbirth and the puerperium (O00-O99)	67	0.4	0.3
12	Septicemia (A40-A41)	61	0.4	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	49	0.3	0.2
14	Legal intervention (Y35,Y89.0)	32	0.2	0.1
15	Pneumonitis due to solids and liquids (J69)	29	0.2	0.1
...	All other causes (residual)	1,993	12.1	8.1
Non-Hispanic, white, both sexes, 25-44 years				
...	All causes	77,129	100.0	156.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27,034	35.1	55.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,800	14.0	22.0
3	Malignant neoplasms (C00-C97)	8,667	11.2	17.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,911	10.3	16.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,355	3.1	4.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,899	2.5	3.9
7	Diabetes mellitus (E10-E14)	1,476	1.9	3.0
8	Cerebrovascular diseases (I60-I69)	1,052	1.4	2.1
9	Septicemia (A40-A41)	661	0.9	1.3
10	Influenza and pneumonia (J09-J18)	650	0.8	1.3
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	618	0.8	1.3
12	Chronic lower respiratory diseases (J40-J47)	571	0.7	1.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	381	0.5	0.8
14	Pregnancy, childbirth and the puerperium (O00-O99)	332	0.4	0.7
15	Human immunodeficiency virus (HIV) disease (B20-B24)	318	0.4	0.6
...	All other causes (residual)	12,404	16.1	25.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015



LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 45-64 years				
...	All causes	376,090	100.0	649.9
1	Malignant neoplasms (C00-C97)	115,260	30.6	199.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	76,553	20.4	132.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30,452	8.1	52.6
4	Chronic lower respiratory diseases (J40-J47)	18,120	4.8	31.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	15,942	4.2	27.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14,475	3.8	25.0
7	Diabetes mellitus (E10-E14)	12,462	3.3	21.5
8	Cerebrovascular diseases (I60-I69)	10,223	2.7	17.7
9	Septicemia (A40-A41)	5,422	1.4	9.4
10	Influenza and pneumonia (J09-J18)	4,531	1.2	7.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,125	1.1	7.1
12	Viral hepatitis (B15-B19)	3,339	0.9	5.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,856	0.8	4.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,708	0.5	3.0
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,612	0.4	2.8
...	All other causes (residual)	59,010	15.7	102.0
Non-Hispanic, white, both sexes, 65 years and over				
...	All causes	1,638,812	100.0	4,390.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	417,905	25.5	1,119.6
2	Malignant neoplasms (C00-C97)	341,783	20.9	915.6
3	Chronic lower respiratory diseases (J40-J47)	117,397	7.2	314.5
4	Cerebrovascular diseases (I60-I69)	95,376	5.8	255.5
5	Alzheimer's disease (G30)	92,493	5.6	247.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	44,192	2.7	118.4
7	Influenza and pneumonia (J09-J18)	39,792	2.4	106.6
8	Diabetes mellitus (E10-E14)	39,534	2.4	105.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30,897	1.9	82.8
10	Septicemia (A40-A41)	23,976	1.5	64.2
11	Parkinson's disease (G20-G21)	23,778	1.5	63.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	19,533	1.2	52.3
13	Pneumonitis due to solids and liquids (J69)	14,723	0.9	39.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11,313	0.7	30.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	11,028	0.7	29.5
...	All other causes (residual)	315,092	19.2	844.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, all ages				
...	All causes	1,063,705	100.0	1,072.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	264,784	24.9	267.0
2	Malignant neoplasms (C00-C97)	247,645	23.3	249.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	68,715	6.5	69.3
4	Chronic lower respiratory diseases (J40-J47)	62,857	5.9	63.4
5	Cerebrovascular diseases (I60-I69)	43,100	4.1	43.5
6	Diabetes mellitus (E10-E14)	29,813	2.8	30.1
7	Alzheimer's disease (G30)	28,631	2.7	28.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28,045	2.6	28.3
9	Influenza and pneumonia (J09-J18)	20,995	2.0	21.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	18,546	1.7	18.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,351	1.7	18.5
12	Parkinson's disease (G20-G21)	14,778	1.4	14.9
13	Septicemia (A40-A41)	14,309	1.3	14.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9,526	0.9	9.6
15	Pneumonitis due to solids and liquids (J69)	9,085	0.9	9.2
...	All other causes (residual)	184,525	17.3	186.0
Non-Hispanic, white, male, 1-4 years				
...	All causes	1,083	100.0	25.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	401	37.0	9.4
2	Malignant neoplasms (C00-C97)	116	10.7	2.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	104	9.6	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	88	8.1	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	2.1	0.5
6	Influenza and pneumonia (J09-J18)	17	1.6	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.1	*
8	Cerebrovascular diseases (I60-I69)	10	0.9	*
9	Septicemia (A40-A41)	9	0.8	*
9	Certain conditions originating in the perinatal period (P00-P96)	9	0.8	*
11	Acute bronchitis and bronchiolitis (J20-J21)	7	0.6	*
11	Chronic lower respiratory diseases (J40-J47)	7	0.6	*
13	Anemias (D50-D64)	4	0.4	*
14	Pneumonitis due to solids and liquids (J69)	3	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
...	All other causes (residual)	267	24.7	6.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 5-14 years				
...	All causes	1,634	100.0	14.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	493	30.2	4.4
2	Malignant neoplasms (C00-C97)	247	15.1	2.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	187	11.4	1.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	106	6.5	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	3.7	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	3.4	0.5
7	Chronic lower respiratory diseases (J40-J47)	27	1.7	0.2
8	Influenza and pneumonia (J09-J18)	25	1.5	0.2
9	Cerebrovascular diseases (I60-I69)	21	1.3	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.2	0.2
11	Septicemia (A40-A41)	12	0.7	*
12	Certain conditions originating in the perinatal period (P00-P96)	6	0.4	*
13	Anemias (D50-D64)	5	0.3	*
13	Diabetes mellitus (E10-E14)	5	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.3	*
...	All other causes (residual)	354	21.7	3.1
Non-Hispanic, white, male, 15-24 years				
...	All causes	11,753	100.0	93.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,805	49.4	46.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,948	25.1	23.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	481	4.1	3.8
4	Malignant neoplasms (C00-C97)	468	4.0	3.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	337	2.9	2.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	123	1.0	1.0
7	Influenza and pneumonia (J09-J18)	62	0.5	0.5
8	Diabetes mellitus (E10-E14)	58	0.5	0.5
9	Chronic lower respiratory diseases (J40-J47)	46	0.4	0.4
10	Cerebrovascular diseases (I60-I69)	43	0.4	0.3
11	Septicemia (A40-A41)	32	0.3	0.3
12	Legal intervention (Y35,Y89.0)	31	0.3	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.2	0.2
14	Pneumonitis due to solids and liquids (J69)	19	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	19	0.2	*
...	All other causes (residual)	1,255	10.7	10.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 25-44 years				
...	All causes	50,033	100.0	201.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19,179	38.3	77.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,295	16.6	33.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,367	10.7	21.6
4	Malignant neoplasms (C00-C97)	4,002	8.0	16.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,458	2.9	5.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,306	2.6	5.3
7	Diabetes mellitus (E10-E14)	912	1.8	3.7
8	Cerebrovascular diseases (I60-I69)	552	1.1	2.2
9	Influenza and pneumonia (J09-J18)	347	0.7	1.4
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	347	0.7	1.4
11	Septicemia (A40-A41)	313	0.6	1.3
12	Chronic lower respiratory diseases (J40-J47)	265	0.5	1.1
13	Human immunodeficiency virus (HIV) disease (B20-B24)	230	0.5	0.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	213	0.4	0.9
15	Aortic aneurysm and dissection (I71)	186	0.4	0.8
...	All other causes (residual)	7,061	14.1	28.5
Non-Hispanic, white, male, 45-64 years				
...	All causes	229,534	100.0	804.9
1	Malignant neoplasms (C00-C97)	62,633	27.3	219.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	54,020	23.5	189.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20,123	8.8	70.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,622	4.6	37.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	10,543	4.6	37.0
6	Chronic lower respiratory diseases (J40-J47)	9,164	4.0	32.1
7	Diabetes mellitus (E10-E14)	7,884	3.4	27.6
8	Cerebrovascular diseases (I60-I69)	5,708	2.5	20.0
9	Septicemia (A40-A41)	2,916	1.3	10.2
10	Influenza and pneumonia (J09-J18)	2,503	1.1	8.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,441	1.1	8.6
12	Viral hepatitis (B15-B19)	2,320	1.0	8.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,891	0.8	6.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,115	0.5	3.9
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1,107	0.5	3.9
...	All other causes (residual)	34,544	15.0	121.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 65 years and over				
...	All causes	763,873	100.0	4,573.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	204,890	26.8	1,226.6
2	Malignant neoplasms (C00-C97)	180,159	23.6	1,078.6
3	Chronic lower respiratory diseases (J40-J47)	53,339	7.0	319.3
4	Cerebrovascular diseases (I60-I69)	36,742	4.8	220.0
5	Alzheimer's disease (G30)	28,269	3.7	169.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22,312	2.9	133.6
7	Diabetes mellitus (E10-E14)	20,953	2.7	125.4
8	Influenza and pneumonia (J09-J18)	17,993	2.4	107.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,660	2.1	93.8
10	Parkinson's disease (G20-G21)	14,393	1.9	86.2
11	Septicemia (A40-A41)	10,988	1.4	65.8
12	Pneumonitis due to solids and liquids (J69)	8,046	1.1	48.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,483	1.0	44.8
14	Chronic liver disease and cirrhosis (K70,K73-K74)	6,535	0.9	39.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6,046	0.8	36.2
...	All other causes (residual)	130,065	17.0	778.7
Non-Hispanic, white, female, all ages				
...	All causes	1,059,926	100.0	1,038.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	238,388	22.5	233.6
2	Malignant neoplasms (C00-C97)	219,563	20.7	215.1
3	Chronic lower respiratory diseases (J40-J47)	73,371	6.9	71.9
4	Alzheimer's disease (G30)	64,694	6.1	63.4
5	Cerebrovascular diseases (I60-I69)	63,730	6.0	62.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	43,112	4.1	42.2
7	Influenza and pneumonia (J09-J18)	24,247	2.3	23.8
8	Diabetes mellitus (E10-E14)	23,777	2.2	23.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,130	1.6	16.8
10	Septicemia (A40-A41)	15,931	1.5	15.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13,102	1.2	12.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	10,797	1.0	10.6
13	Parkinson's disease (G20-G21)	9,561	0.9	9.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,420	0.8	8.3
15	Pneumonitis due to solids and liquids (J69)	7,345	0.7	7.2
...	All other causes (residual)	226,758	21.4	222.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 1-4 years				
...	All causes	812	100.0	20.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	230	28.3	5.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	92	11.3	2.3
3	Malignant neoplasms (C00-C97)	90	11.1	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	57	7.0	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	3.9	0.8
6	Influenza and pneumonia (J09-J18)	20	2.5	0.5
7	Septicemia (A40-A41)	11	1.4	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.9	*
8	Cerebrovascular diseases (I60-I69)	7	0.9	*
10	Meningitis (G00,G03)	6	0.7	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.6	*
11	Certain conditions originating in the perinatal period (P00-P96)	5	0.6	*
13	Chronic lower respiratory diseases (J40-J47)	4	0.5	*
14	Anemias (D50-D64)	3	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	3	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.4	*
...	All other causes (residual)	237	29.2	5.9
Non-Hispanic, white, female, 5-14 years				
...	All causes	1,201	100.0	11.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	305	25.4	2.8
2	Malignant neoplasms (C00-C97)	229	19.1	2.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	104	8.7	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	80	6.7	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	48	4.0	0.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	46	3.8	0.4
7	Influenza and pneumonia (J09-J18)	22	1.8	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	1.7	0.2
9	Cerebrovascular diseases (I60-I69)	20	1.7	0.2
10	Chronic lower respiratory diseases (J40-J47)	18	1.5	*
11	Septicemia (A40-A41)	16	1.3	*
12	Diabetes mellitus (E10-E14)	6	0.5	*
13	Anemias (D50-D64)	5	0.4	*
14	Certain conditions originating in the perinatal period (P00-P96)	4	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.2	*
...	All other causes (residual)	274	22.8	2.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 15-24 years				
...	All causes	4,783	100.0	40.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,274	47.5	19.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	729	15.2	6.1
3	Malignant neoplasms (C00-C97)	313	6.5	2.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	202	4.2	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	158	3.3	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	101	2.1	0.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	67	1.4	0.6
8	Diabetes mellitus (E10-E14)	46	1.0	0.4
9	Influenza and pneumonia (J09-J18)	45	0.9	0.4
10	Cerebrovascular diseases (I60-I69)	40	0.8	0.3
11	Septicemia (A40-A41)	29	0.6	0.2
12	Chronic lower respiratory diseases (J40-J47)	26	0.5	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.5	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.3	*
15	Pneumonitis due to solids and liquids (J69)	10	0.2	*
...	All other causes (residual)	707	14.8	5.9
Non-Hispanic, white, female, 25-44 years				
...	All causes	27,096	100.0	111.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,855	29.0	32.2
2	Malignant neoplasms (C00-C97)	4,665	17.2	19.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,544	9.4	10.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,505	9.2	10.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	897	3.3	3.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	593	2.2	2.4
7	Diabetes mellitus (E10-E14)	564	2.1	2.3
8	Cerebrovascular diseases (I60-I69)	500	1.8	2.1
9	Septicemia (A40-A41)	348	1.3	1.4
10	Pregnancy, childbirth and the puerperium (O00-O99)	332	1.2	1.4
11	Chronic lower respiratory diseases (J40-J47)	306	1.1	1.3
12	Influenza and pneumonia (J09-J18)	303	1.1	1.2
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	271	1.0	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	168	0.6	0.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	109	0.4	0.4
...	All other causes (residual)	5,136	19.0	21.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 45-64 years				
...	All causes	146,556	100.0	499.3
1	Malignant neoplasms (C00-C97)	52,627	35.9	179.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,533	15.4	76.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,329	7.0	35.2
4	Chronic lower respiratory diseases (J40-J47)	8,956	6.1	30.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,399	3.7	18.4
6	Diabetes mellitus (E10-E14)	4,578	3.1	15.6
7	Cerebrovascular diseases (I60-I69)	4,515	3.1	15.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,853	2.6	13.1
9	Septicemia (A40-A41)	2,506	1.7	8.5
10	Influenza and pneumonia (J09-J18)	2,028	1.4	6.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,684	1.1	5.7
12	Viral hepatitis (B15-B19)	1,019	0.7	3.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	965	0.7	3.3
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	716	0.5	2.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	709	0.5	2.4
...	All other causes (residual)	24,139	16.5	82.2
Non-Hispanic, white, female, 65 years and over				
...	All causes	874,939	100.0	4,242.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	213,015	24.3	1,032.9
2	Malignant neoplasms (C00-C97)	161,624	18.5	783.7
3	Alzheimer's disease (G30)	64,224	7.3	311.4
4	Chronic lower respiratory diseases (J40-J47)	64,058	7.3	310.6
5	Cerebrovascular diseases (I60-I69)	58,634	6.7	284.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21,880	2.5	106.1
7	Influenza and pneumonia (J09-J18)	21,799	2.5	105.7
8	Diabetes mellitus (E10-E14)	18,581	2.1	90.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,237	1.7	73.9
10	Septicemia (A40-A41)	12,988	1.5	63.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12,050	1.4	58.4
12	Parkinson's disease (G20-G21)	9,385	1.1	45.5
13	Pneumonitis due to solids and liquids (J69)	6,677	0.8	32.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,267	0.6	25.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4,493	0.5	21.8
...	All other causes (residual)	185,027	21.1	897.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015



LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, all ages				
...	All causes	315,254	100.0	754.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	74,093	23.5	177.4
2	Malignant neoplasms (C00-C97)	68,523	21.7	164.0
3	Cerebrovascular diseases (I60-I69)	17,760	5.6	42.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,366	4.9	36.8
5	Diabetes mellitus (E10-E14)	13,693	4.3	32.8
6	Chronic lower respiratory diseases (J40-J47)	10,327	3.3	24.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9,083	2.9	21.7
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9,038	2.9	21.6
9	Alzheimer's disease (G30)	8,072	2.6	19.3
10	Septicemia (A40-A41)	6,576	2.1	15.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,740	1.8	13.7
12	Influenza and pneumonia (J09-J18)	5,554	1.8	13.3
13	Certain conditions originating in the perinatal period (P00-P96)	3,827	1.2	9.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3,379	1.1	8.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3,185	1.0	7.6
...	All other causes (residual)	61,038	19.4	146.1
Non-Hispanic, black, both sexes, 1-4 years				
...	All causes	1,016	100.0	41.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	286	28.1	11.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	139	13.7	5.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	93	9.2	3.8
4	Malignant neoplasms (C00-C97)	53	5.2	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	52	5.1	2.1
6	Influenza and pneumonia (J09-J18)	27	2.7	1.1
6	Certain conditions originating in the perinatal period (P00-P96)	27	2.7	1.1
8	Chronic lower respiratory diseases (J40-J47)	23	2.3	0.9
9	Cerebrovascular diseases (I60-I69)	17	1.7	*
10	Septicemia (A40-A41)	14	1.4	*
11	Anemias (D50-D64)	13	1.3	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
15	Meningitis (G00,G03)	3	0.3	*
...	All other causes (residual)	250	24.6	10.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 5-14 years				
...	All causes	1,187	100.0	19.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	327	27.5	5.3
2	Malignant neoplasms (C00-C97)	126	10.6	2.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	121	10.2	2.0
4	Chronic lower respiratory diseases (J40-J47)	102	8.6	1.7
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	66	5.6	1.1
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	58	4.9	0.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	47	4.0	0.8
8	Cerebrovascular diseases (I60-I69)	22	1.9	0.4
9	Influenza and pneumonia (J09-J18)	19	1.6	*
10	Septicemia (A40-A41)	16	1.3	*
10	Anemias (D50-D64)	16	1.3	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.8	*
13	Diabetes mellitus (E10-E14)	9	0.8	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.6	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.5	*
...	All other causes (residual)	235	19.8	3.8
Non-Hispanic, black, both sexes, 15-24 years				
...	All causes	7,096	100.0	104.5
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,013	42.5	44.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,649	23.2	24.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	574	8.1	8.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	302	4.3	4.4
5	Malignant neoplasms (C00-C97)	252	3.6	3.7
6	Chronic lower respiratory diseases (J40-J47)	86	1.2	1.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	73	1.0	1.1
8	Anemias (D50-D64)	70	1.0	1.0
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	67	0.9	1.0
10	Diabetes mellitus (E10-E14)	64	0.9	0.9
11	Pregnancy, childbirth and the puerperium (O00-O99)	63	0.9	0.9
12	Cerebrovascular diseases (I60-I69)	39	0.5	0.6
13	Influenza and pneumonia (J09-J18)	35	0.5	0.5
14	Septicemia (A40-A41)	34	0.5	0.5
14	Legal intervention (Y35,Y89.0)	34	0.5	0.5
...	All other causes (residual)	741	10.4	10.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 25-44 years				
...	All causes	24,618	100.0	216.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,290	17.4	37.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,237	17.2	37.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,799	15.4	33.3
4	Malignant neoplasms (C00-C97)	2,539	10.3	22.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,017	4.1	8.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	919	3.7	8.1
7	Diabetes mellitus (E10-E14)	792	3.2	6.9
8	Cerebrovascular diseases (I60-I69)	675	2.7	5.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	389	1.6	3.4
10	Septicemia (A40-A41)	333	1.4	2.9
11	Chronic lower respiratory diseases (J40-J47)	288	1.2	2.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	258	1.0	2.3
13	Chronic liver disease and cirrhosis (K70,K73-K74)	242	1.0	2.1
14	Influenza and pneumonia (J09-J18)	240	1.0	2.1
15	Pregnancy, childbirth and the puerperium (O00-O99)	238	1.0	2.1
...	All other causes (residual)	4,362	17.7	38.3
Non-Hispanic, black, both sexes, 45-64 years				
...	All causes	92,058	100.0	912.9
1	Malignant neoplasms (C00-C97)	25,007	27.2	248.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,485	24.4	223.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,425	5.9	53.8
4	Diabetes mellitus (E10-E14)	4,544	4.9	45.1
5	Cerebrovascular diseases (I60-I69)	4,432	4.8	43.9
6	Chronic lower respiratory diseases (J40-J47)	2,526	2.7	25.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,352	2.6	23.3
8	Chronic liver disease and cirrhosis (K70,K73-K74)	2,002	2.2	19.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,932	2.1	19.2
10	Septicemia (A40-A41)	1,925	2.1	19.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,570	1.7	15.6
12	Influenza and pneumonia (J09-J18)	1,281	1.4	12.7
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,203	1.3	11.9
14	Viral hepatitis (B15-B19)	737	0.8	7.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	604	0.7	6.0
...	All other causes (residual)	14,033	15.2	139.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 65 years and over				
...	All causes	182,355	100.0	4,271.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	47,304	25.9	1,108.2
2	Malignant neoplasms (C00-C97)	40,537	22.2	949.6
3	Cerebrovascular diseases (I60-I69)	12,548	6.9	294.0
4	Diabetes mellitus (E10-E14)	8,279	4.5	193.9
5	Alzheimer's disease (G30)	7,955	4.4	186.4
6	Chronic lower respiratory diseases (J40-J47)	7,290	4.0	170.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,286	3.4	147.3
8	Septicemia (A40-A41)	4,195	2.3	98.3
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,906	2.1	91.5
10	Influenza and pneumonia (J09-J18)	3,895	2.1	91.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,030	1.7	71.0
12	Pneumonitis due to solids and liquids (J69)	1,328	0.7	31.1
13	Parkinson's disease (G20-G21)	1,126	0.6	26.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,015	0.6	23.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	939	0.5	22.0
...	All other causes (residual)	32,722	17.9	766.6
Non-Hispanic, black, male, all ages				
...	All causes	161,850	100.0	809.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	38,638	23.9	193.2
2	Malignant neoplasms (C00-C97)	34,787	21.5	174.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,516	6.5	52.6
4	Cerebrovascular diseases (I60-I69)	7,962	4.9	39.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,909	4.9	39.6
6	Diabetes mellitus (E10-E14)	6,806	4.2	34.0
7	Chronic lower respiratory diseases (J40-J47)	5,236	3.2	26.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,351	2.7	21.8
9	Septicemia (A40-A41)	3,061	1.9	15.3
10	Influenza and pneumonia (J09-J18)	2,729	1.7	13.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,599	1.6	13.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,230	1.4	11.2
13	Alzheimer's disease (G30)	2,154	1.3	10.8
14	Certain conditions originating in the perinatal period (P00-P96)	2,130	1.3	10.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,951	1.2	9.8
...	All other causes (residual)	28,791	17.8	144.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 1-4 years				
...	All causes	608	100.0	48.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	180	29.6	14.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	90	14.8	7.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	8.6	4.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	4.6	2.3
5	Malignant neoplasms (C00-C97)	26	4.3	2.1
6	Certain conditions originating in the perinatal period (P00-P96)	19	3.1	*
7	Influenza and pneumonia (J09-J18)	17	2.8	*
8	Chronic lower respiratory diseases (J40-J47)	15	2.5	*
9	Cerebrovascular diseases (I60-I69)	12	2.0	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	8	1.3	*
11	Septicemia (A40-A41)	7	1.2	*
11	Anemias (D50-D64)	7	1.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
14	Diabetes mellitus (E10-E14)	1	0.2	*
14	Nutritional deficiencies (E40-E64)	1	0.2	*
14	Meningitis (G00,G03)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Diseases of appendix (K35-K38)	1	0.2	*
...	All other causes (residual)	137	22.5	11.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 5-14 years				
...	All causes	703	100.0	22.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	187	26.6	6.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	83	11.8	2.6
3	Malignant neoplasms (C00-C97)	68	9.7	2.2
4	Chronic lower respiratory diseases (J40-J47)	63	9.0	2.0
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	5.4	1.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	5.0	1.1
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	4.6	1.0
8	Cerebrovascular diseases (I60-I69)	13	1.8	*
9	Influenza and pneumonia (J09-J18)	9	1.3	*
10	Septicemia (A40-A41)	7	1.0	*
10	Anemias (D50-D64)	7	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.9	*
12	Diabetes mellitus (E10-E14)	6	0.9	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
14	Certain conditions originating in the perinatal period (P00-P96)	3	0.4	*
...	All other causes (residual)	143	20.3	4.6
Non-Hispanic, black, male, 15-24 years				
...	All causes	5,486	100.0	159.2
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,744	50.0	79.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,226	22.3	35.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	452	8.2	13.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	194	3.5	5.6
5	Malignant neoplasms (C00-C97)	138	2.5	4.0
6	Chronic lower respiratory diseases (J40-J47)	55	1.0	1.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	45	0.8	1.3
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	45	0.8	1.3
9	Diabetes mellitus (E10-E14)	39	0.7	1.1
10	Anemias (D50-D64)	35	0.6	1.0
11	Legal intervention (Y35,Y89.0)	32	0.6	0.9
12	Influenza and pneumonia (J09-J18)	21	0.4	0.6
13	Cerebrovascular diseases (I60-I69)	20	0.4	0.6
14	Septicemia (A40-A41)	15	0.3	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.2	*
...	All other causes (residual)	414	7.5	12.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 25-44 years				
...	All causes	15,839	100.0	289.5
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,829	24.2	70.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,100	19.6	56.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,398	15.1	43.8
4	Malignant neoplasms (C00-C97)	965	6.1	17.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	848	5.4	15.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	553	3.5	10.1
7	Diabetes mellitus (E10-E14)	448	2.8	8.2
8	Cerebrovascular diseases (I60-I69)	375	2.4	6.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	202	1.3	3.7
10	Chronic lower respiratory diseases (J40-J47)	168	1.1	3.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	161	1.0	2.9
12	Septicemia (A40-A41)	155	1.0	2.8
13	Influenza and pneumonia (J09-J18)	147	0.9	2.7
14	Anemias (D50-D64)	116	0.7	2.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	104	0.7	1.9
...	All other causes (residual)	2,270	14.3	41.5
Non-Hispanic, black, male, 45-64 years				
...	All causes	53,255	100.0	1,140.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,111	26.5	302.1
2	Malignant neoplasms (C00-C97)	12,912	24.2	276.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,887	7.3	83.2
4	Diabetes mellitus (E10-E14)	2,582	4.8	55.3
5	Cerebrovascular diseases (I60-I69)	2,525	4.7	54.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,338	2.5	28.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	1,315	2.5	28.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,245	2.3	26.7
9	Chronic lower respiratory diseases (J40-J47)	1,242	2.3	26.6
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	992	1.9	21.2
11	Septicemia (A40-A41)	976	1.8	20.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	901	1.7	19.3
13	Influenza and pneumonia (J09-J18)	760	1.4	16.3
14	Viral hepatitis (B15-B19)	506	1.0	10.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	465	0.9	10.0
...	All other causes (residual)	7,498	14.1	160.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 65 years and over				
...	All causes	82,132	100.0	4,789.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,827	26.6	1,272.9
2	Malignant neoplasms (C00-C97)	20,673	25.2	1,205.6
3	Cerebrovascular diseases (I60-I69)	5,003	6.1	291.8
4	Diabetes mellitus (E10-E14)	3,728	4.5	217.4
5	Chronic lower respiratory diseases (J40-J47)	3,688	4.5	215.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,788	3.4	162.6
7	Alzheimer's disease (G30)	2,104	2.6	122.7
8	Septicemia (A40-A41)	1,867	2.3	108.9
9	Influenza and pneumonia (J09-J18)	1,738	2.1	101.4
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,713	2.1	99.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,535	1.9	89.5
12	Pneumonitis due to solids and liquids (J69)	660	0.8	38.5
13	Parkinson's disease (G20-G21)	645	0.8	37.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	583	0.7	34.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	463	0.6	27.0
...	All other causes (residual)	13,117	16.0	764.9
Non-Hispanic, black, female, all ages				
...	All causes	153,404	100.0	704.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,455	23.1	162.8
2	Malignant neoplasms (C00-C97)	33,736	22.0	154.9
3	Cerebrovascular diseases (I60-I69)	9,798	6.4	45.0
4	Diabetes mellitus (E10-E14)	6,887	4.5	31.6
5	Alzheimer's disease (G30)	5,918	3.9	27.2
6	Chronic lower respiratory diseases (J40-J47)	5,091	3.3	23.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,850	3.2	22.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,732	3.1	21.7
9	Septicemia (A40-A41)	3,515	2.3	16.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,141	2.0	14.4
11	Influenza and pneumonia (J09-J18)	2,825	1.8	13.0
12	Certain conditions originating in the perinatal period (P00-P96)	1,697	1.1	7.8
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,252	0.8	5.7
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,149	0.7	5.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,129	0.7	5.2
...	All other causes (residual)	32,229	21.0	148.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015



LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 1-4 years				
...	All causes	408	100.0	34.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	106	26.0	8.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	49	12.0	4.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	10.0	3.4
4	Malignant neoplasms (C00-C97)	27	6.6	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	5.9	2.0
6	Influenza and pneumonia (J09-J18)	10	2.5	*
7	Chronic lower respiratory diseases (J40-J47)	8	2.0	*
7	Certain conditions originating in the perinatal period (P00-P96)	8	2.0	*
9	Septicemia (A40-A41)	7	1.7	*
10	Anemias (D50-D64)	6	1.5	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.2	*
11	Cerebrovascular diseases (I60-I69)	5	1.2	*
13	Meningitis (G00,G03)	2	0.5	*
14	Meningococcal infection (A39)	1	0.2	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
14	Nutritional deficiencies (E40-E64)	1	0.2	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
14	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
14	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (residual)	101	24.8	8.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 5-14 years				
...	All causes	484	100.0	15.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	140	28.9	4.6
2	Malignant neoplasms (C00-C97)	58	12.0	1.9
3	Chronic lower respiratory diseases (J40-J47)	39	8.1	1.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	38	7.9	1.2
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	5.8	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	5.4	0.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	2.5	*
8	Influenza and pneumonia (J09-J18)	10	2.1	*
9	Septicemia (A40-A41)	9	1.9	*
9	Anemias (D50-D64)	9	1.9	*
9	Cerebrovascular diseases (I60-I69)	9	1.9	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	5	1.0	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.8	*
14	Diabetes mellitus (E10-E14)	3	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
...	All other causes (residual)	91	18.8	3.0
Non-Hispanic, black, female, 15-24 years				
...	All causes	1,610	100.0	48.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	423	26.3	12.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	269	16.7	8.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	122	7.6	3.7
4	Malignant neoplasms (C00-C97)	114	7.1	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	108	6.7	3.2
6	Pregnancy, childbirth and the puerperium (O00-O99)	63	3.9	1.9
7	Anemias (D50-D64)	35	2.2	1.0
8	Chronic lower respiratory diseases (J40-J47)	31	1.9	0.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	28	1.7	0.8
10	Diabetes mellitus (E10-E14)	25	1.6	0.7
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	1.4	0.7
12	Septicemia (A40-A41)	19	1.2	*
12	Cerebrovascular diseases (I60-I69)	19	1.2	*
14	Influenza and pneumonia (J09-J18)	14	0.9	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.8	*
...	All other causes (residual)	305	18.9	9.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 25-44 years				
...	All causes	8,779	100.0	148.1
1	Malignant neoplasms (C00-C97)	1,574	17.9	26.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,401	16.0	23.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,137	13.0	19.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	461	5.3	7.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	366	4.2	6.2
6	Diabetes mellitus (E10-E14)	344	3.9	5.8
7	Cerebrovascular diseases (I60-I69)	300	3.4	5.1
8	Pregnancy, childbirth and the puerperium (O00-O99)	238	2.7	4.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	187	2.1	3.2
10	Septicemia (A40-A41)	178	2.0	3.0
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	169	1.9	2.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	138	1.6	2.3
13	Chronic lower respiratory diseases (J40-J47)	120	1.4	2.0
14	Anemias (D50-D64)	115	1.3	1.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	97	1.1	1.6
...	All other causes (residual)	1,954	22.3	33.0
Non-Hispanic, black, female, 45-64 years				
...	All causes	38,803	100.0	716.8
1	Malignant neoplasms (C00-C97)	12,095	31.2	223.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,374	21.6	154.7
3	Diabetes mellitus (E10-E14)	1,962	5.1	36.2
4	Cerebrovascular diseases (I60-I69)	1,907	4.9	35.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,538	4.0	28.4
6	Chronic lower respiratory diseases (J40-J47)	1,284	3.3	23.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,014	2.6	18.7
8	Septicemia (A40-A41)	949	2.4	17.5
9	Chronic liver disease and cirrhosis (K70,K73-K74)	757	2.0	14.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	669	1.7	12.4
11	Human immunodeficiency virus (HIV) disease (B20-B24)	617	1.6	11.4
12	Influenza and pneumonia (J09-J18)	521	1.3	9.6
13	Viral hepatitis (B15-B19)	231	0.6	4.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	211	0.5	3.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	155	0.4	2.9
...	All other causes (residual)	6,519	16.8	120.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 65 years and over				
...	All causes	100,223	100.0	3,924.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,477	25.4	997.6
2	Malignant neoplasms (C00-C97)	19,864	19.8	777.8
3	Cerebrovascular diseases (I60-I69)	7,545	7.5	295.4
4	Alzheimer's disease (G30)	5,851	5.8	229.1
5	Diabetes mellitus (E10-E14)	4,551	4.5	178.2
6	Chronic lower respiratory diseases (J40-J47)	3,602	3.6	141.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,498	3.5	137.0
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,371	2.4	92.8
9	Septicemia (A40-A41)	2,328	2.3	91.2
10	Influenza and pneumonia (J09-J18)	2,157	2.2	84.5
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,317	1.3	51.6
12	Pneumonitis due to solids and liquids (J69)	668	0.7	26.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	552	0.6	21.6
14	Parkinson's disease (G20-G21)	481	0.5	18.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	356	0.4	13.9
...	All other causes (residual)	19,605	19.6	767.6
Non-Hispanic, American Indian or Alaska Native, both sexes, all ages				
...	All causes	18,039	100.0	670.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,303	18.3	122.8
2	Malignant neoplasms (C00-C97)	3,192	17.7	118.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,953	10.8	72.6
4	Diabetes mellitus (E10-E14)	1,034	5.7	38.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,006	5.6	37.4
6	Chronic lower respiratory diseases (J40-J47)	859	4.8	31.9
7	Cerebrovascular diseases (I60-I69)	646	3.6	24.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	535	3.0	19.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	340	1.9	12.6
10	Influenza and pneumonia (J09-J18)	330	1.8	12.3
11	Alzheimer's disease (G30)	307	1.7	11.4
12	Septicemia (A40-A41)	296	1.6	11.0
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	262	1.5	9.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	192	1.1	7.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	109	0.6	4.1
...	All other causes (residual)	3,675	20.4	136.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 1-4 years				
...	All causes	82	100.0	51.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30	36.6	18.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	12.2	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	9.8	*
4	Malignant neoplasms (C00-C97)	3	3.7	*
5	Meningitis (G00,G03)	1	1.2	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.2	*
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	1.2	*
5	Cerebrovascular diseases (I60-I69)	1	1.2	*
5	Chronic lower respiratory diseases (J40-J47)	1	1.2	*
5	Pneumonitis due to solids and liquids (J69)	1	1.2	*
...	All other causes (residual)	25	30.5	15.6
Non-Hispanic, American Indian or Alaska Native, both sexes, 5-14 years				
...	All causes	99	100.0	24.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	31.3	7.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	14.1	*
3	Malignant neoplasms (C00-C97)	11	11.1	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	7.1	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	7.1	*
6	Chronic lower respiratory diseases (J40-J47)	3	3.0	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.0	*
7	Certain conditions originating in the perinatal period (P00-P96)	2	2.0	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.0	*
9	Influenza and pneumonia (J09-J18)	1	1.0	*
...	All other causes (residual)	20	20.2	4.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 15-24 years				
...	All causes	525	100.0	119.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	205	39.0	46.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	150	28.6	34.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	52	9.9	11.8
4	Malignant neoplasms (C00-C97)	16	3.0	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	2.7	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	1.7	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	6	1.1	*
8	Chronic lower respiratory diseases (J40-J47)	5	1.0	*
9	Cerebrovascular diseases (I60-I69)	4	0.8	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.6	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.6	*
12	Diabetes mellitus (E10-E14)	2	0.4	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
13	Meningitis (G00,G03)	1	0.2	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
13	Influenza and pneumonia (J09-J18)	1	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
13	Legal intervention (Y35,Y89.0)	1	0.2	*
...	All other causes (residual)	50	9.5	11.4
Non-Hispanic, American Indian or Alaska Native, both sexes, 25-44 years				
...	All causes	2,274	100.0	317.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	744	32.7	103.8
2	Chronic liver disease and cirrhosis (K70,K73-K74)	263	11.6	36.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	215	9.5	30.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	194	8.5	27.1
5	Malignant neoplasms (C00-C97)	131	5.8	18.3
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	121	5.3	16.9
7	Diabetes mellitus (E10-E14)	57	2.5	8.0
8	Cerebrovascular diseases (I60-I69)	32	1.4	4.5
9	Septicemia (A40-A41)	31	1.4	4.3
10	Influenza and pneumonia (J09-J18)	18	0.8	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	17	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.7	*
13	Chronic lower respiratory diseases (J40-J47)	13	0.6	*
13	Pregnancy, childbirth and the puerperium (O00-O99)	13	0.6	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	0.5	*
...	All other causes (residual)	397	17.5	55.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 45-64 years				
...	All causes	5,693	100.0	874.1
1	Malignant neoplasms (C00-C97)	1,094	19.2	168.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,054	18.5	161.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	637	11.2	97.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	574	10.1	88.1
5	Diabetes mellitus (E10-E14)	384	6.7	59.0
6	Chronic lower respiratory diseases (J40-J47)	212	3.7	32.5
7	Cerebrovascular diseases (I60-I69)	148	2.6	22.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	130	2.3	20.0
9	Septicemia (A40-A41)	104	1.8	16.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	94	1.7	14.4
11	Influenza and pneumonia (J09-J18)	75	1.3	11.5
12	Viral hepatitis (B15-B19)	69	1.2	10.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	59	1.0	9.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	56	1.0	8.6
15	Human immunodeficiency virus (HIV) disease (B20-B24)	30	0.5	4.6
...	All other causes (residual)	973	17.1	149.4
Non-Hispanic, American Indian or Alaska Native, both sexes, 65 years and over				
...	All causes	9,066	100.0	3,359.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,035	22.4	754.0
2	Malignant neoplasms (C00-C97)	1,936	21.4	717.3
3	Chronic lower respiratory diseases (J40-J47)	625	6.9	231.6
4	Diabetes mellitus (E10-E14)	591	6.5	219.0
5	Cerebrovascular diseases (I60-I69)	460	5.1	170.4
6	Alzheimer's disease (G30)	304	3.4	112.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	274	3.0	101.5
8	Influenza and pneumonia (J09-J18)	231	2.5	85.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	228	2.5	84.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	163	1.8	60.4
11	Septicemia (A40-A41)	159	1.8	58.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	128	1.4	47.4
13	Parkinson's disease (G20-G21)	86	0.9	31.9
14	Pneumonitis due to solids and liquids (J69)	64	0.7	23.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	50	0.6	18.5
...	All other causes (residual)	1,732	19.1	641.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, all ages				
...	All causes	9,869	100.0	747.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,916	19.4	145.1
2	Malignant neoplasms (C00-C97)	1,699	17.2	128.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,282	13.0	97.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	563	5.7	42.6
5	Diabetes mellitus (E10-E14)	537	5.4	40.7
6	Chronic lower respiratory diseases (J40-J47)	398	4.0	30.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	395	4.0	29.9
8	Cerebrovascular diseases (I60-I69)	291	2.9	22.0
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	209	2.1	15.8
10	Influenza and pneumonia (J09-J18)	181	1.8	13.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	159	1.6	12.0
12	Septicemia (A40-A41)	130	1.3	9.8
13	Alzheimer's disease (G30)	100	1.0	7.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	87	0.9	6.6
15	Viral hepatitis (B15-B19)	70	0.7	5.3
...	All other causes (residual)	1,852	18.8	140.3
Non-Hispanic, American Indian or Alaska Native, male, 1-4 years				
...	All causes	45	100.0	55.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	31.1	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	8.9	*
3	Malignant neoplasms (C00-C97)	3	6.7	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	6.7	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.2	*
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	2.2	*
5	Cerebrovascular diseases (I60-I69)	1	2.2	*
5	Chronic lower respiratory diseases (J40-J47)	1	2.2	*
...	All other causes (residual)	17	37.8	*
Non-Hispanic, American Indian or Alaska Native, male, 5-14 years				
...	All causes	51	100.0	24.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	35.3	*
2	Malignant neoplasms (C00-C97)	8	15.7	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	7.8	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	7.8	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	3.9	*
6	Influenza and pneumonia (J09-J18)	1	2.0	*
6	Chronic lower respiratory diseases (J40-J47)	1	2.0	*
...	All other causes (residual)	13	25.5	*

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015



LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 15-24 years				
...	All causes	362	100.0	161.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	138	38.1	61.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	110	30.4	49.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	49	13.5	21.9
4	Malignant neoplasms (C00-C97)	10	2.8	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	1.9	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	5	1.4	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	1.4	*
8	Diabetes mellitus (E10-E14)	2	0.6	*
8	Cerebrovascular diseases (I60-I69)	2	0.6	*
8	Chronic lower respiratory diseases (J40-J47)	2	0.6	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.3	*
11	Influenza and pneumonia (J09-J18)	1	0.3	*
11	Legal intervention (Y35,Y89.0)	1	0.3	*
...	All other causes (residual)	28	7.7	12.5
Non-Hispanic, American Indian or Alaska Native, male, 25-44 years				
...	All causes	1,471	100.0	413.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	521	35.4	146.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	164	11.1	46.1
3	Chronic liver disease and cirrhosis (K70,K73-K74)	151	10.3	42.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	121	8.2	34.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	96	6.5	27.0
6	Malignant neoplasms (C00-C97)	64	4.4	18.0
7	Diabetes mellitus (E10-E14)	34	2.3	9.6
8	Cerebrovascular diseases (I60-I69)	18	1.2	*
9	Septicemia (A40-A41)	17	1.2	*
10	Influenza and pneumonia (J09-J18)	12	0.8	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.7	*
12	Viral hepatitis (B15-B19)	7	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.5	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	0.5	*
15	Legal intervention (Y35,Y89.0)	6	0.4	*
...	All other causes (residual)	236	16.0	66.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 45-64 years				
...	All causes	3,403	100.0	1,096.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	737	21.7	237.5
2	Malignant neoplasms (C00-C97)	587	17.2	189.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	421	12.4	135.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	327	9.6	105.4
5	Diabetes mellitus (E10-E14)	217	6.4	69.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	97	2.9	31.3
7	Chronic lower respiratory diseases (J40-J47)	90	2.6	29.0
8	Cerebrovascular diseases (I60-I69)	75	2.2	24.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	1.5	16.1
10	Viral hepatitis (B15-B19)	48	1.4	15.5
11	Septicemia (A40-A41)	47	1.4	15.1
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	47	1.4	15.1
13	Influenza and pneumonia (J09-J18)	42	1.2	13.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	37	1.1	11.9
15	Human immunodeficiency virus (HIV) disease (B20-B24)	19	0.6	*
15	Pneumonitis due to solids and liquids (J69)	19	0.6	*
...	All other causes (residual)	543	16.0	175.0
Non-Hispanic, American Indian or Alaska Native, male, 65 years and over				
...	All causes	4,376	100.0	3,622.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,049	24.0	868.3
2	Malignant neoplasms (C00-C97)	1,026	23.4	849.2
3	Chronic lower respiratory diseases (J40-J47)	299	6.8	247.5
4	Diabetes mellitus (E10-E14)	284	6.5	235.1
5	Cerebrovascular diseases (I60-I69)	194	4.4	160.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	153	3.5	126.6
7	Influenza and pneumonia (J09-J18)	123	2.8	101.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	101	2.3	83.6
9	Alzheimer's disease (G30)	99	2.3	81.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	80	1.8	66.2
11	Septicemia (A40-A41)	65	1.5	53.8
12	Parkinson's disease (G20-G21)	51	1.2	42.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	44	1.0	36.4
14	Pneumonitis due to solids and liquids (J69)	38	0.9	31.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.6	21.5
...	All other causes (residual)	744	17.0	615.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, all ages				
...	All causes	8,170	100.0	596.7
1	Malignant neoplasms (C00-C97)	1,493	18.3	109.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,387	17.0	101.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	671	8.2	49.0
4	Diabetes mellitus (E10-E14)	497	6.1	36.3
5	Chronic lower respiratory diseases (J40-J47)	461	5.6	33.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	443	5.4	32.4
7	Cerebrovascular diseases (I60-I69)	355	4.3	25.9
8	Alzheimer's disease (G30)	207	2.5	15.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	181	2.2	13.2
10	Septicemia (A40-A41)	166	2.0	12.1
11	Influenza and pneumonia (J09-J18)	149	1.8	10.9
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	140	1.7	10.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	105	1.3	7.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	0.6	3.9
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	0.6	3.8
...	All other causes (residual)	1,810	22.2	132.2
Non-Hispanic, American Indian or Alaska Native, female, 1-4 years				
...	All causes	37	100.0	47.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16	43.2	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	16.2	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	13.5	*
4	Meningitis (G00,G03)	1	2.7	*
4	Pneumonitis due to solids and liquids (J69)	1	2.7	*
...	All other causes (residual)	8	21.6	*
Non-Hispanic, American Indian or Alaska Native, female, 5-14 years				
...	All causes	48	100.0	23.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13	27.1	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10	20.8	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	10.4	*
4	Malignant neoplasms (C00-C97)	3	6.3	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	6.3	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	4.2	*
6	Chronic lower respiratory diseases (J40-J47)	2	4.2	*
6	Certain conditions originating in the perinatal period (P00-P96)	2	4.2	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	2.1	*
...	All other causes (residual)	7	14.6	*

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 15-24 years				
...	All causes	163	100.0	75.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	67	41.1	31.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	24.5	18.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	4.3	*
4	Malignant neoplasms (C00-C97)	6	3.7	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	2.5	*
6	Chronic lower respiratory diseases (J40-J47)	3	1.8	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	3	1.8	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	1.8	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown	2	1.2	*
9	Cerebrovascular diseases (I60-I69)	2	1.2	*
11	Meningitis (G00,G03)	1	0.6	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.6	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.6	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.6	*
...	All other causes (residual)	22	13.5	10.2
Non-Hispanic, American Indian or Alaska Native, female, 25-44 years				
...	All causes	803	100.0	222.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	223	27.8	61.8
2	Chronic liver disease and cirrhosis (K70,K73-K74)	112	13.9	31.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	73	9.1	20.2
4	Malignant neoplasms (C00-C97)	67	8.3	18.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	51	6.4	14.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	3.1	6.9
7	Diabetes mellitus (E10-E14)	23	2.9	6.4
8	Septicemia (A40-A41)	14	1.7	*
8	Cerebrovascular diseases (I60-I69)	14	1.7	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	13	1.6	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	1.1	*
12	Chronic lower respiratory diseases (J40-J47)	8	1.0	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	7	0.9	*
14	Influenza and pneumonia (J09-J18)	6	0.7	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.6	*
...	All other causes (residual)	153	19.1	42.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 45-64 years				
...	All causes	2,290	100.0	671.5
1	Malignant neoplasms (C00-C97)	507	22.1	148.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	317	13.8	93.0
3	Chronic liver disease and cirrhosis (K70,K73-K74)	247	10.8	72.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	216	9.4	63.3
5	Diabetes mellitus (E10-E14)	167	7.3	49.0
6	Chronic lower respiratory diseases (J40-J47)	122	5.3	35.8
7	Cerebrovascular diseases (I60-I69)	73	3.2	21.4
8	Septicemia (A40-A41)	57	2.5	16.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	44	1.9	12.9
10	Influenza and pneumonia (J09-J18)	33	1.4	9.7
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	1.4	9.7
12	Viral hepatitis (B15-B19)	21	0.9	6.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	19	0.8	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	0.5	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	11	0.5	*
...	All other causes (residual)	411	17.9	120.5
Non-Hispanic, American Indian or Alaska Native, female, 65 years and over				
...	All causes	4,690	100.0	3,145.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	986	21.0	661.3
2	Malignant neoplasms (C00-C97)	910	19.4	610.4
3	Chronic lower respiratory diseases (J40-J47)	326	7.0	218.7
4	Diabetes mellitus (E10-E14)	307	6.5	205.9
5	Cerebrovascular diseases (I60-I69)	266	5.7	178.4
6	Alzheimer's disease (G30)	205	4.4	137.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	127	2.7	85.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	121	2.6	81.2
9	Influenza and pneumonia (J09-J18)	108	2.3	72.4
10	Septicemia (A40-A41)	94	2.0	63.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	84	1.8	56.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	83	1.8	55.7
13	Parkinson's disease (G20-G21)	35	0.7	23.5
14	Pneumonitis due to solids and liquids (J69)	26	0.6	17.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.5	16.1
...	All other causes (residual)	988	21.1	662.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, all ages				
...	All causes	65,277	100.0	341.5
1	Malignant neoplasms (C00-C97)	17,306	26.5	90.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,974	21.4	73.1
3	Cerebrovascular diseases (I60-I69)	4,798	7.4	25.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,839	4.3	14.9
5	Diabetes mellitus (E10-E14)	2,599	4.0	13.6
6	Alzheimer's disease (G30)	2,180	3.3	11.4
7	Influenza and pneumonia (J09-J18)	2,161	3.3	11.3
8	Chronic lower respiratory diseases (J40-J47)	1,874	2.9	9.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,318	2.0	6.9
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,273	2.0	6.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,136	1.7	5.9
12	Septicemia (A40-A41)	798	1.2	4.2
13	Parkinson's disease (G20-G21)	747	1.1	3.9
14	Chronic liver disease and cirrhosis (K70,K73-K74)	598	0.9	3.1
15	Pneumonitis due to solids and liquids (J69)	500	0.8	2.6
...	All other causes (residual)	11,176	17.1	58.5
Non-Hispanic, Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	141	100.0	15.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26	18.4	2.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	14.2	2.2
3	Malignant neoplasms (C00-C97)	13	9.2	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	7.1	*
5	Influenza and pneumonia (J09-J18)	6	4.3	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	4.3	*
7	Septicemia (A40-A41)	3	2.1	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	2.1	*
9	Certain conditions originating in the perinatal period (P00-P96)	2	1.4	*
10	Tuberculosis (A16-A19)	1	0.7	*
10	Anemias (D50-D64)	1	0.7	*
10	Acute bronchitis and bronchiolitis (J20-J21)	1	0.7	*
10	Chronic lower respiratory diseases (J40-J47)	1	0.7	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.7	*
...	All other causes (residual)	47	33.3	5.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 5-14 years				
...	All causes	212	100.0	9.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	49	23.1	2.1
2	Malignant neoplasms (C00-C97)	40	18.9	1.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15	7.1	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	6.6	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	3.3	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	3.3	*
7	Cerebrovascular diseases (I60-I69)	5	2.4	*
8	Septicemia (A40-A41)	4	1.9	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.4	*
10	Anemias (D50-D64)	2	0.9	*
10	Influenza and pneumonia (J09-J18)	2	0.9	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.9	*
13	Meningitis (G00,G03)	1	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.5	*
...	All other causes (residual)	60	28.3	2.6
Non-Hispanic, Asian or Pacific Islander, both sexes, 15-24 years				
...	All causes	873	100.0	33.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	277	31.7	10.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	265	30.4	10.3
3	Malignant neoplasms (C00-C97)	84	9.6	3.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	46	5.3	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	40	4.6	1.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	1.8	*
7	Cerebrovascular diseases (I60-I69)	7	0.8	*
8	Influenza and pneumonia (J09-J18)	5	0.6	*
8	Chronic lower respiratory diseases (J40-J47)	5	0.6	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.5	*
11	Septicemia (A40-A41)	3	0.3	*
11	Diabetes mellitus (E10-E14)	3	0.3	*
11	Aortic aneurysm and dissection (I71)	3	0.3	*
11	Pneumonitis due to solids and liquids (J69)	3	0.3	*
11	Legal intervention (Y35,Y89.0)	3	0.3	*
...	All other causes (residual)	109	12.5	4.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 25-44 years				
...	All causes	3,379	100.0	53.6
1	Malignant neoplasms (C00-C97)	806	23.9	12.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	587	17.4	9.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	446	13.2	7.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	410	12.1	6.5
5	Cerebrovascular diseases (I60-I69)	143	4.2	2.3
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	124	3.7	2.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	67	2.0	1.1
8	Diabetes mellitus (E10-E14)	63	1.9	1.0
9	Pregnancy, childbirth and the puerperium (O00-O99)	42	1.2	0.7
10	Influenza and pneumonia (J09-J18)	35	1.0	0.6
11	Septicemia (A40-A41)	28	0.8	0.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	27	0.8	0.4
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	25	0.7	0.4
14	Chronic lower respiratory diseases (J40-J47)	23	0.7	0.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.7	0.4
...	All other causes (residual)	530	15.7	8.4
Non-Hispanic, Asian or Pacific Islander, both sexes, 45-64 years				
...	All causes	12,221	100.0	264.0
1	Malignant neoplasms (C00-C97)	5,062	41.4	109.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,341	19.2	50.6
3	Cerebrovascular diseases (I60-I69)	758	6.2	16.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	577	4.7	12.5
5	Diabetes mellitus (E10-E14)	485	4.0	10.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	365	3.0	7.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	264	2.2	5.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	187	1.5	4.0
9	Influenza and pneumonia (J09-J18)	149	1.2	3.2
10	Septicemia (A40-A41)	139	1.1	3.0
10	Chronic lower respiratory diseases (J40-J47)	139	1.1	3.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	130	1.1	2.8
13	Viral hepatitis (B15-B19)	91	0.7	2.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	81	0.7	1.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	76	0.6	1.6
...	All other causes (residual)	1,377	11.3	29.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015



LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 65 years and over				
...	All causes	47,558	100.0	2,241.2
1	Malignant neoplasms (C00-C97)	11,296	23.8	532.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,145	23.4	525.2
3	Cerebrovascular diseases (I60-I69)	3,880	8.2	182.8
4	Alzheimer's disease (G30)	2,159	4.5	101.7
5	Diabetes mellitus (E10-E14)	2,048	4.3	96.5
6	Influenza and pneumonia (J09-J18)	1,959	4.1	92.3
7	Chronic lower respiratory diseases (J40-J47)	1,706	3.6	80.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,300	2.7	61.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,101	2.3	51.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	993	2.1	46.8
11	Parkinson's disease (G20-G21)	732	1.5	34.5
12	Septicemia (A40-A41)	614	1.3	28.9
13	Pneumonitis due to solids and liquids (J69)	462	1.0	21.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	322	0.7	15.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	266	0.6	12.5
...	All other causes (residual)	7,575	15.9	357.0
Non-Hispanic, Asian or Pacific Islander, male, all ages				
...	All causes	33,306	100.0	364.9
1	Malignant neoplasms (C00-C97)	8,774	26.3	96.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,612	22.9	83.4
3	Cerebrovascular diseases (I60-I69)	2,153	6.5	23.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,788	5.4	19.6
5	Diabetes mellitus (E10-E14)	1,322	4.0	14.5
6	Influenza and pneumonia (J09-J18)	1,109	3.3	12.2
7	Chronic lower respiratory diseases (J40-J47)	1,061	3.2	11.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	855	2.6	9.4
9	Alzheimer's disease (G30)	675	2.0	7.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	651	2.0	7.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	515	1.5	5.6
12	Parkinson's disease (G20-G21)	448	1.3	4.9
13	Septicemia (A40-A41)	418	1.3	4.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	384	1.2	4.2
15	Pneumonitis due to solids and liquids (J69)	285	0.9	3.1
...	All other causes (residual)	5,256	15.8	57.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 1-4 years				
...	All causes	72	100.0	15.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16	22.2	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	12.5	*
3	Malignant neoplasms (C00-C97)	7	9.7	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	6.9	*
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	4.2	*
5	Influenza and pneumonia (J09-J18)	3	4.2	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	4.2	*
8	Acute bronchitis and bronchiolitis (J20-J21)	1	1.4	*
8	Chronic lower respiratory diseases (J40-J47)	1	1.4	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.4	*
8	Certain conditions originating in the perinatal period (P00-P96)	1	1.4	*
...	All other causes (residual)	22	30.6	4.7
Non-Hispanic, Asian or Pacific Islander, male, 5-14 years				
...	All causes	124	100.0	10.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	25.0	2.6
2	Malignant neoplasms (C00-C97)	25	20.2	2.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	6.5	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	5.6	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	3.2	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	3.2	*
7	Cerebrovascular diseases (I60-I69)	3	2.4	*
8	Septicemia (A40-A41)	2	1.6	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.8	*
9	Meningitis (G00,G03)	1	0.8	*
9	Influenza and pneumonia (J09-J18)	1	0.8	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.8	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.8	*
...	All other causes (residual)	35	28.2	3.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 15-24 years				
...	All causes	630	100.0	48.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	208	33.0	15.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	203	32.2	15.5
3	Malignant neoplasms (C00-C97)	54	8.6	4.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	5.6	2.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	4.9	2.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	1.6	*
7	Cerebrovascular diseases (I60-I69)	5	0.8	*
8	Septicemia (A40-A41)	3	0.5	*
8	Aortic aneurysm and dissection (I71)	3	0.5	*
8	Chronic lower respiratory diseases (J40-J47)	3	0.5	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.3	*
12	Diabetes mellitus (E10-E14)	2	0.3	*
12	Influenza and pneumonia (J09-J18)	2	0.3	*
12	Legal intervention (Y35,Y89.0)	2	0.3	*
...	All other causes (residual)	64	10.2	4.9
Non-Hispanic, Asian or Pacific Islander, male, 25-44 years				
...	All causes	2,099	100.0	70.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	445	21.2	14.9
2	Malignant neoplasms (C00-C97)	350	16.7	11.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	325	15.5	10.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	294	14.0	9.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	93	4.4	3.1
6	Cerebrovascular diseases (I60-I69)	72	3.4	2.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	48	2.3	1.6
8	Diabetes mellitus (E10-E14)	46	2.2	1.5
9	Influenza and pneumonia (J09-J18)	27	1.3	0.9
10	Human immunodeficiency virus (HIV) disease (B20-B24)	24	1.1	0.8
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	0.8	*
12	Aortic aneurysm and dissection (I71)	14	0.7	*
13	Septicemia (A40-A41)	12	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.6	*
15	Chronic lower respiratory diseases (J40-J47)	11	0.5	*
...	All other causes (residual)	309	14.7	10.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 45-64 years				
...	All causes	7,207	100.0	337.1
1	Malignant neoplasms (C00-C97)	2,457	34.1	114.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,723	23.9	80.6
3	Cerebrovascular diseases (I60-I69)	443	6.1	20.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	400	5.6	18.7
5	Diabetes mellitus (E10-E14)	302	4.2	14.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	232	3.2	10.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	213	3.0	10.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	114	1.6	5.3
9	Influenza and pneumonia (J09-J18)	98	1.4	4.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	94	1.3	4.4
11	Chronic lower respiratory diseases (J40-J47)	89	1.2	4.2
12	Septicemia (A40-A41)	78	1.1	3.6
13	Viral hepatitis (B15-B19)	61	0.8	2.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	59	0.8	2.8
15	Aortic aneurysm and dissection (I71)	43	0.6	2.0
...	All other causes (residual)	801	11.1	37.5
Non-Hispanic, Asian or Pacific Islander, male, 65 years and over				
...	All causes	22,673	100.0	2,462.1
1	Malignant neoplasms (C00-C97)	5,879	25.9	638.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,512	24.3	598.6
3	Cerebrovascular diseases (I60-I69)	1,627	7.2	176.7
4	Influenza and pneumonia (J09-J18)	974	4.3	105.8
5	Diabetes mellitus (E10-E14)	972	4.3	105.6
6	Chronic lower respiratory diseases (J40-J47)	957	4.2	103.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	673	3.0	73.1
8	Alzheimer's disease (G30)	664	2.9	72.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	519	2.3	56.4
10	Parkinson's disease (G20-G21)	439	1.9	47.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	411	1.8	44.6
12	Septicemia (A40-A41)	319	1.4	34.6
13	Pneumonitis due to solids and liquids (J69)	261	1.2	28.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	148	0.7	16.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	122	0.5	13.2
...	All other causes (residual)	3,196	14.1	347.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, all ages				
...	All causes	31,971	100.0	320.1
1	Malignant neoplasms (C00-C97)	8,532	26.7	85.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,362	19.9	63.7
3	Cerebrovascular diseases (I60-I69)	2,645	8.3	26.5
4	Alzheimer's disease (G30)	1,505	4.7	15.1
5	Diabetes mellitus (E10-E14)	1,277	4.0	12.8
6	Influenza and pneumonia (J09-J18)	1,052	3.3	10.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,051	3.3	10.5
8	Chronic lower respiratory diseases (J40-J47)	813	2.5	8.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	667	2.1	6.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	621	1.9	6.2
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	418	1.3	4.2
12	Septicemia (A40-A41)	380	1.2	3.8
13	Parkinson's disease (G20-G21)	299	0.9	3.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	222	0.7	2.2
15	Pneumonitis due to solids and liquids (J69)	215	0.7	2.2
...	All other causes (residual)	5,912	18.5	59.2
Non-Hispanic, Asian or Pacific Islander, female, 1-4 years				
...	All causes	69	100.0	15.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	15.9	*
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	14.5	*
3	Malignant neoplasms (C00-C97)	6	8.7	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	7.2	*
5	Septicemia (A40-A41)	3	4.3	*
5	Influenza and pneumonia (J09-J18)	3	4.3	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	4.3	*
8	Tuberculosis (A16-A19)	1	1.4	*
8	Anemias (D50-D64)	1	1.4	*
8	Certain conditions originating in the perinatal period (P00-P96)	1	1.4	*
...	All other causes (residual)	25	36.2	5.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 5-14 years				
...	All causes	88	100.0	7.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	20.5	*
2	Malignant neoplasms (C00-C97)	15	17.0	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	8.0	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	8.0	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	3.4	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	3.4	*
7	Septicemia (A40-A41)	2	2.3	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	2.3	*
7	Anemias (D50-D64)	2	2.3	*
7	Cerebrovascular diseases (I60-I69)	2	2.3	*
11	Influenza and pneumonia (J09-J18)	1	1.1	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.1	*
...	All other causes (residual)	25	28.4	2.2
Non-Hispanic, Asian or Pacific Islander, female, 15-24 years				
...	All causes	243	100.0	19.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	69	28.4	5.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	62	25.5	4.9
3	Malignant neoplasms (C00-C97)	30	12.3	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	4.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	3.7	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	2.5	*
7	Influenza and pneumonia (J09-J18)	3	1.2	*
8	Cerebrovascular diseases (I60-I69)	2	0.8	*
8	Chronic lower respiratory diseases (J40-J47)	2	0.8	*
8	Pneumonitis due to solids and liquids (J69)	2	0.8	*
11	Tuberculosis (A16-A19)	1	0.4	*
11	Anemias (D50-D64)	1	0.4	*
11	Diabetes mellitus (E10-E14)	1	0.4	*
11	Meningitis (G00,G03)	1	0.4	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
11	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.4	*
11	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
11	Legal intervention (Y35,Y89.0)	1	0.4	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
11	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.4	*
...	All other causes (residual)	37	15.2	2.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 25-44 years				
...	All causes	1,280	100.0	38.7
1	Malignant neoplasms (C00-C97)	456	35.6	13.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	152	11.9	4.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	142	11.1	4.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	85	6.6	2.6
5	Cerebrovascular diseases (I60-I69)	71	5.5	2.1
6	Pregnancy, childbirth and the puerperium (O00-O99)	42	3.3	1.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	31	2.4	0.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	19	1.5	*
9	Diabetes mellitus (E10-E14)	17	1.3	*
10	Septicemia (A40-A41)	16	1.3	*
11	Chronic lower respiratory diseases (J40-J47)	12	0.9	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.9	*
13	Influenza and pneumonia (J09-J18)	8	0.6	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.5	*
...	All other causes (residual)	203	15.9	6.1
Non-Hispanic, Asian or Pacific Islander, female, 45-64 years				
...	All causes	5,014	100.0	201.3
1	Malignant neoplasms (C00-C97)	2,605	52.0	104.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	618	12.3	24.8
3	Cerebrovascular diseases (I60-I69)	315	6.3	12.6
4	Diabetes mellitus (E10-E14)	183	3.6	7.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	177	3.5	7.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	133	2.7	5.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	73	1.5	2.9
8	Septicemia (A40-A41)	61	1.2	2.4
9	Influenza and pneumonia (J09-J18)	51	1.0	2.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	51	1.0	2.0
11	Chronic lower respiratory diseases (J40-J47)	50	1.0	2.0
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	0.7	1.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	36	0.7	1.4
14	Viral hepatitis (B15-B19)	30	0.6	1.2
15	Aortic aneurysm and dissection (I71)	24	0.5	1.0
...	All other causes (residual)	571	11.4	22.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 65 years and over				
...	All causes	24,885	100.0	2,071.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,633	22.6	469.0
2	Malignant neoplasms (C00-C97)	5,417	21.8	451.0
3	Cerebrovascular diseases (I60-I69)	2,253	9.1	187.6
4	Alzheimer's disease (G30)	1,495	6.0	124.5
5	Diabetes mellitus (E10-E14)	1,076	4.3	89.6
6	Influenza and pneumonia (J09-J18)	985	4.0	82.0
7	Chronic lower respiratory diseases (J40-J47)	749	3.0	62.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	627	2.5	52.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	582	2.3	48.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	582	2.3	48.5
11	Septicemia (A40-A41)	295	1.2	24.6
12	Parkinson's disease (G20-G21)	293	1.2	24.4
13	Pneumonitis due to solids and liquids (J69)	201	0.8	16.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	174	0.7	14.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	144	0.6	12.0
...	All other causes (residual)	4,379	17.6	364.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015