

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|-------------------------------------|----------------|--------------------|--------------|--------------|----------------|----------------|----------------|----------------|----------------|----------------|
| All causes | | | | | | | | | | |
| All origins 3/, both sexes..... | 847.3 | 695.0 | 31.2 | 15.2 | 19.5 | 67.8 | 95.2 | 94.7 | 111.6 | 161.3 |
| Male..... | 846.6 | 761.5 | 35.2 | 16.7 | 23.1 | 94.0 | 140.8 | 134.4 | 149.4 | 206.6 |
| Female..... | 848.0 | 625.3 | 27.0 | 13.5 | 15.9 | 40.1 | 47.3 | 53.6 | 73.3 | 116.1 |
| Hispanic, both sexes..... | 302.2 | 592.7 | 29.8 | 14.4 | 16.6 | 65.2 | 87.9 | 74.7 | 84.2 | 115.5 |
| Male..... | 328.7 | 644.0 | 34.2 | 15.9 | 19.0 | 97.0 | 128.9 | 108.7 | 116.5 | 151.4 |
| Female..... | 274.0 | 539.1 | 25.3 | 12.9 | 14.0 | 30.8 | 37.2 | 33.3 | 46.9 | 75.6 |
| Mexican, both sexes..... | 254.4 | 565.1 | 30.2 | 14.7 | 16.2 | 70.9 | 88.4 | 73.5 | 80.4 | 105.1 |
| Male..... | 280.8 | 617.3 | 34.3 | 16.5 | 18.9 | 106.4 | 131.0 | 108.3 | 108.4 | 138.6 |
| Female..... | 225.3 | 510.4 | 26.0 | 12.8 | 13.4 | 32.0 | 35.0 | 30.9 | 45.5 | 66.0 |
| Puerto Rican, both sexes..... | 431.6 | 841.8 | 31.5 | 11.0 | 18.6 | 46.8 | 88.1 | 85.5 | 118.4 | 183.7 |
| Male..... | 500.9 | * | 45.3 | * | 18.0 | 69.9 | 132.9 | 128.5 | 170.1 | 239.6 |
| Female..... | 368.1 | * | * | * | 19.3 | 25.8 | 41.6 | 44.5 | 71.9 | 129.3 |
| Cuban, both sexes..... | 828.0 | * | * | * | * | * | * | 57.0 | 88.1 | 100.0 |
| Male..... | 838.0 | * | * | * | * | * | * | * | * | * |
| Female..... | 817.8 | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 307.3 | 635.8 | 29.5 | 16.1 | 18.5 | 57.5 | 88.2 | 76.3 | 83.1 | 121.5 |
| Male..... | 328.1 | 681.2 | 31.2 | 16.2 | 21.8 | 80.5 | 122.1 | 106.8 | 125.0 | 159.2 |
| Female..... | 286.1 | 588.8 | 27.7 | 15.9 | 14.9 | 31.2 | 44.6 | 37.0 | 41.8 | 83.5 |
| Non-Hispanic 5/, both sexes..... | 928.8 | 713.0 | 31.4 | 15.3 | 20.1 | 67.9 | 96.3 | 99.1 | 116.6 | 168.0 |
| Male..... | 928.0 | 782.6 | 35.4 | 16.8 | 23.7 | 93.0 | 142.8 | 140.6 | 155.7 | 215.0 |
| Female..... | 929.5 | 640.1 | 27.2 | 13.7 | 16.2 | 41.6 | 49.1 | 57.8 | 77.8 | 121.9 |
| Non-Hispanic white, both sexes..... | 997.5 | 575.9 | 27.1 | 13.7 | 18.1 | 65.5 | 86.6 | 86.5 | 103.9 | 151.1 |
| Male..... | 983.9 | 643.5 | 30.3 | 14.9 | 21.3 | 87.6 | 126.7 | 121.2 | 139.3 | 194.0 |
| Female..... | 1,010.6 | 504.8 | 23.8 | 12.3 | 14.8 | 42.0 | 45.6 | 51.3 | 68.1 | 108.0 |
| Non-Hispanic black, both sexes..... | 792.8 | 1,280.3 | 48.4 | 22.0 | 28.3 | 83.8 | 151.7 | 179.6 | 207.4 | 292.2 |
| Male..... | 842.3 | 1,369.3 | 55.8 | 24.8 | 34.4 | 124.1 | 235.3 | 266.6 | 281.5 | 376.4 |
| Female..... | 748.0 | 1,187.5 | 40.8 | 19.2 | 22.0 | 42.1 | 70.5 | 100.1 | 141.0 | 217.4 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|-------------------------------------|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
| Septicemia (A40-A41) | | | | | | | | | | |
| All origins 3/, both sexes..... | 11.7 | 7.3 | 0.5 | 0.2 | 0.3 | 0.2 | 0.3 | 0.6 | 0.9 | 1.5 |
| Male..... | 10.6 | 7.7 | 0.5 | 0.3 | 0.2 | 0.2 | 0.4 | 0.7 | 1.0 | 1.5 |
| Female..... | 12.9 | 7.0 | 0.5 | * | 0.3 | 0.3 | 0.3 | 0.6 | 0.9 | 1.5 |
| Hispanic, both sexes..... | 3.9 | 7.8 | * | * | * | * | * | * | 0.8 | 0.9 |
| Male..... | 3.7 | 8.0 | * | * | * | * | * | * | * | * |
| Female..... | 4.1 | 7.6 | * | * | * | * | * | * | * | * |
| Mexican, both sexes..... | 3.1 | 8.5 | * | * | * | * | * | * | * | * |
| Male..... | 3.0 | 8.7 | * | * | * | * | * | * | * | * |
| Female..... | 3.3 | 8.2 | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | 8.5 | * | * | * | * | * | * | * | * | * |
| Male..... | 8.4 | * | * | * | * | * | * | * | * | * |
| Female..... | 8.6 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 11.6 | * | * | * | * | * | * | * | * | * |
| Male..... | 9.5 | * | * | * | * | * | * | * | * | * |
| Female..... | 13.8 | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 3.2 | * | * | * | * | * | * | * | * | * |
| Male..... | 3.3 | * | * | * | * | * | * | * | * | * |
| Female..... | 3.1 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 12.9 | 7.1 | 0.5 | 0.2 | 0.3 | 0.2 | 0.4 | 0.7 | 1.0 | 1.6 |
| Male..... | 11.6 | 7.3 | 0.5 | 0.2 | 0.3 | * | 0.4 | 0.7 | 1.0 | 1.5 |
| Female..... | 14.1 | 6.8 | 0.5 | * | 0.3 | 0.3 | 0.3 | 0.7 | 0.9 | 1.6 |
| Non-Hispanic white, both sexes..... | 12.9 | 5.5 | 0.5 | 0.2 | 0.2 | 0.2 | 0.3 | 0.6 | 0.8 | 1.3 |
| Male..... | 11.5 | 5.6 | 0.5 | * | * | * | 0.4 | 0.6 | 0.7 | 1.2 |
| Female..... | 14.2 | 5.4 | * | * | * | 0.3 | * | 0.5 | 0.8 | 1.3 |
| Non-Hispanic black, both sexes..... | 16.8 | 13.1 | * | * | * | * | * | 1.6 | 2.3 | 3.5 |
| Male..... | 15.5 | 13.9 | * | * | * | * | * | * | 2.5 | 3.8 |
| Female..... | 18.0 | 12.3 | * | * | * | * | * | 2.0 | 2.1 | 3.1 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Measles (B05)

Table with 11 columns representing age groups and 11 rows representing population subgroups (All origins, Hispanic, Mexican, Puerto Rican, Cuban, Other Hispanic, Non-Hispanic white, Non-Hispanic black). All cells contain an asterisk (*).

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
| Human immunodeficiency virus (HIV) disease (B20-B24) | | | | | | | | | | |
| All origins 3/, both sexes..... | 4.9 | * | * | * | 0.1 | 0.2 | 0.7 | 2.6 | 6.4 | 11.9 |
| Male..... | 7.4 | * | * | * | * | * | 0.7 | 3.0 | 8.5 | 17.7 |
| Female..... | 2.5 | * | * | * | * | 0.2 | 0.6 | 2.3 | 4.2 | 6.2 |
| Hispanic, both sexes..... | 4.8 | * | * | * | * | * | 0.6 | 2.1 | 5.1 | 11.2 |
| Male..... | 7.2 | * | * | * | * | * | * | 2.9 | 7.4 | 16.9 |
| Female..... | 2.2 | * | * | * | * | * | * | 1.2 | 2.4 | 4.8 |
| Mexican, both sexes..... | 2.5 | * | * | * | * | * | * | 1.4 | 4.0 | 6.8 |
| Male..... | 4.1 | * | * | * | * | * | * | 2.1 | 6.4 | 12.0 |
| Female..... | 0.7 | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | 21.7 | * | * | * | * | * | * | * | 15.5 | 46.4 |
| Male..... | 30.5 | * | * | * | * | * | * | * | * | 58.1 |
| Female..... | 13.5 | * | * | * | * | * | * | * | * | 35.0 |
| Cuban, both sexes..... | 5.6 | * | * | * | * | * | * | * | * | * |
| Male..... | 9.5 | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 4.7 | * | * | * | * | * | * | 3.3 | 4.9 | 10.8 |
| Male..... | 7.6 | * | * | * | * | * | * | * | 8.1 | 17.5 |
| Female..... | 1.8 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 4.8 | * | * | * | * | 0.2 | 0.7 | 2.7 | 6.6 | 11.9 |
| Male..... | 7.3 | * | * | * | * | * | 0.7 | 3.0 | 8.7 | 17.6 |
| Female..... | 2.5 | * | * | * | * | * | 0.7 | 2.5 | 4.5 | 6.3 |
| Non-Hispanic white, both sexes..... | 2.1 | * | * | * | * | * | * | 0.9 | 2.4 | 5.7 |
| Male..... | 3.7 | * | * | * | * | * | * | 1.3 | 3.8 | 9.6 |
| Female..... | 0.6 | * | * | * | * | * | * | 0.5 | 1.0 | 1.8 |
| Non-Hispanic black, both sexes..... | 21.3 | * | * | * | * | 0.8 | 3.4 | 12.2 | 30.2 | 49.3 |
| Male..... | 30.3 | * | * | * | * | * | 3.5 | 12.2 | 38.0 | 69.9 |
| Female..... | 13.2 | * | * | * | * | * | 3.4 | 12.3 | 23.2 | 31.1 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|--------------------------------------|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
| Malignant neoplasms (C00-C97) | | | | | | | | | | |
| All origins 3/, both sexes..... | 193.2 | 1.8 | 2.6 | 2.7 | 2.5 | 3.5 | 5.0 | 6.6 | 12.5 | 23.5 |
| Male..... | 203.8 | 2.0 | 2.7 | 2.9 | 2.8 | 4.0 | 5.7 | 7.0 | 11.3 | 19.8 |
| Female..... | 183.0 | 1.6 | 2.4 | 2.5 | 2.2 | 3.0 | 4.2 | 6.2 | 13.7 | 27.2 |
| Hispanic, both sexes..... | 59.7 | * | 2.3 | 2.8 | 3.0 | 3.8 | 5.1 | 4.7 | 9.1 | 15.8 |
| Male..... | 61.2 | * | 2.2 | 2.7 | 3.1 | 4.9 | 5.5 | 5.0 | 7.8 | 11.9 |
| Female..... | 58.1 | * | 2.4 | 2.9 | 2.9 | 2.6 | 4.5 | 4.4 | 10.6 | 20.2 |
| Mexican, both sexes..... | 47.5 | * | 2.5 | 2.4 | 3.1 | 4.0 | 5.1 | 4.6 | 8.9 | 14.1 |
| Male..... | 48.4 | * | 2.4 | 2.3 | 3.3 | 5.5 | 5.5 | 5.0 | 7.4 | 10.4 |
| Female..... | 46.6 | * | 2.6 | 2.5 | 3.0 | 2.4 | 4.6 | 4.2 | 10.7 | 18.4 |
| Puerto Rican, both sexes..... | 82.4 | * | * | * | * | * | * | * | 10.0 | 19.2 |
| Male..... | 94.6 | * | * | * | * | * | * | * | * | * |
| Female..... | 71.2 | * | * | * | * | * | * | * | * | 25.0 |
| Cuban, both sexes..... | 183.1 | * | * | * | * | * | * | * | * | * |
| Male..... | 205.8 | * | * | * | * | * | * | * | * | * |
| Female..... | 160.0 | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 67.4 | * | * | 4.2 | * | 4.6 | 5.3 | 5.6 | 9.4 | 19.6 |
| Male..... | 64.9 | * | * | * | * | * | 6.1 | 5.7 | 8.3 | 15.2 |
| Female..... | 70.0 | * | * | * | * | * | * | * | 10.5 | 23.9 |
| Non-Hispanic 5/, both sexes..... | 213.4 | 1.7 | 2.6 | 2.7 | 2.4 | 3.5 | 4.9 | 7.1 | 13.1 | 24.7 |
| Male..... | 226.7 | 1.9 | 2.8 | 3.0 | 2.7 | 3.9 | 5.7 | 7.6 | 12.0 | 21.1 |
| Female..... | 200.8 | 1.5 | 2.4 | 2.4 | 2.1 | 3.1 | 4.1 | 6.6 | 14.2 | 28.2 |
| Non-Hispanic white, both sexes..... | 230.9 | 1.6 | 2.6 | 2.8 | 2.4 | 3.5 | 5.1 | 6.9 | 12.7 | 23.9 |
| Male..... | 243.8 | 2.0 | 2.8 | 2.9 | 2.6 | 3.9 | 5.9 | 7.4 | 11.8 | 20.5 |
| Female..... | 218.5 | * | 2.3 | 2.5 | 2.2 | 3.0 | 4.3 | 6.5 | 13.5 | 27.2 |
| Non-Hispanic black, both sexes..... | 171.5 | * | 3.0 | 2.6 | 2.6 | 3.6 | 4.6 | 8.4 | 17.0 | 32.7 |
| Male..... | 187.8 | * | 2.7 | 3.3 | 3.0 | 4.2 | 5.2 | 8.6 | 14.7 | 26.5 |
| Female..... | 156.7 | * | 3.2 | 1.9 | 2.2 | 3.1 | 4.0 | 8.3 | 19.0 | 38.2 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|-------------------------------------|----------------|--------------------|--------------|--------------|----------------|----------------|----------------|----------------|----------------|----------------|
| Malignant neoplasm of breast (C50) | | | | | | | | | | |
| All origins 3/, both sexes..... | 14.5 | * | * | * | * | * | * | 0.3 | 1.7 | 4.1 |
| Male..... | 0.3 | * | * | * | * | * | * | * | * | * |
| Female..... | 28.3 | * | * | * | * | * | * | 0.7 | 3.5 | 8.2 |
| Hispanic, both sexes..... | 4.5 | * | * | * | * | * | * | * | 1.3 | 2.4 |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | 9.2 | * | * | * | * | * | * | * | 2.7 | 5.1 |
| Mexican, both sexes..... | 3.5 | * | * | * | * | * | * | * | 1.3 | 2.3 |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | 7.3 | * | * | * | * | * | * | * | 2.9 | 5.1 |
| Puerto Rican, both sexes..... | 6.1 | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | 11.7 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 13.4 | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | 27.1 | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 5.3 | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | 10.7 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 16.0 | * | * | * | * | * | * | 0.4 | 1.8 | 4.4 |
| Male..... | 0.3 | * | * | * | * | * | * | * | * | * |
| Female..... | 31.0 | * | * | * | * | * | * | 0.7 | 3.6 | 8.7 |
| Non-Hispanic white, both sexes..... | 16.9 | * | * | * | * | * | * | 0.3 | 1.5 | 3.8 |
| Male..... | 0.3 | * | * | * | * | * | * | * | * | * |
| Female..... | 32.9 | * | * | * | * | * | * | 0.5 | 3.0 | 7.6 |
| Non-Hispanic black, both sexes..... | 15.4 | * | * | * | * | * | * | 0.9 | 3.8 | 8.4 |
| Male..... | 0.4 | * | * | * | * | * | * | * | * | * |
| Female..... | 29.1 | * | * | * | * | * | * | 1.7 | 7.1 | 15.6 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
| Malignant neoplasm of cervix uteri (C53) | | | | | | | | | | |
| All origins 3/, both sexes..... | 1.4 | * | * | * | * | * | * | 0.2 | 0.8 | 1.3 |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 2.7 | * | * | * | * | * | * | 0.5 | 1.5 | 2.5 |
| Hispanic, both sexes..... | 1.0 | * | * | * | * | * | * | * | 0.6 | 1.3 |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 2.1 | * | * | * | * | * | * | * | 1.3 | 2.8 |
| Mexican, both sexes..... | 0.9 | * | * | * | * | * | * | * | * | 1.2 |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 2.0 | * | * | * | * | * | * | * | * | 2.6 |
| Puerto Rican, both sexes..... | 1.2 | * | * | * | * | * | * | * | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 2.3 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 1.1 | * | * | * | * | * | * | * | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 2.3 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 1.4 | * | * | * | * | * | * | 0.3 | 0.8 | 1.2 |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 2.8 | * | * | * | * | * | * | 0.5 | 1.6 | 2.4 |
| Non-Hispanic white, both sexes..... | 1.3 | * | * | * | * | * | * | 0.2 | 0.8 | 1.1 |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 2.5 | * | * | * | * | * | * | 0.5 | 1.6 | 2.3 |
| Non-Hispanic black, both sexes..... | 2.3 | * | * | * | * | * | * | * | 1.1 | 2.1 |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 4.3 | * | * | * | * | * | * | * | 2.1 | 4.0 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|---|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
| Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96) | | | | | | | | | | |
| All origins 3/, both sexes..... | 19.5 | 0.7 | 0.8 | 0.8 | 1.0 | 1.5 | 1.9 | 1.9 | 2.6 | 3.2 |
| Male..... | 21.4 | * | 0.8 | 0.9 | 1.2 | 1.8 | 2.2 | 2.3 | 3.0 | 3.7 |
| Female..... | 17.7 | * | 0.7 | 0.8 | 0.8 | 1.2 | 1.7 | 1.5 | 2.2 | 2.6 |
| Hispanic, both sexes..... | 7.3 | * | 0.9 | 1.2 | 1.3 | 1.9 | 2.3 | 1.7 | 2.2 | 3.3 |
| Male..... | 7.6 | * | * | 1.1 | 1.4 | 2.6 | 2.7 | 2.1 | 2.6 | 3.7 |
| Female..... | 7.0 | * | * | 1.4 | 1.2 | * | 1.9 | * | 1.7 | 2.9 |
| Mexican, both sexes..... | 6.0 | * | 1.0 | 0.9 | 1.4 | 2.0 | 2.3 | 1.6 | 1.8 | 2.5 |
| Male..... | 6.2 | * | * | * | * | 2.8 | 2.6 | 2.1 | 2.4 | 2.5 |
| Female..... | 5.7 | * | * | * | * | * | 2.0 | * | * | 2.5 |
| Puerto Rican, both sexes..... | 9.4 | * | * | * | * | * | * | * | * | * |
| Male..... | 9.2 | * | * | * | * | * | * | * | * | * |
| Female..... | 9.6 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 20.0 | * | * | * | * | * | * | * | * | * |
| Male..... | 21.2 | * | * | * | * | * | * | * | * | * |
| Female..... | 18.8 | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 8.5 | * | * | * | * | * | * | * | 2.7 | 5.2 |
| Male..... | 9.3 | * | * | * | * | * | * | * | * | 6.5 |
| Female..... | 7.6 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 21.3 | 0.8 | 0.7 | 0.7 | 0.9 | 1.4 | 1.8 | 2.0 | 2.7 | 3.1 |
| Male..... | 23.6 | * | 0.8 | 0.8 | 1.2 | 1.6 | 2.1 | 2.4 | 3.1 | 3.7 |
| Female..... | 19.2 | * | 0.7 | 0.7 | 0.7 | 1.2 | 1.6 | 1.6 | 2.3 | 2.6 |
| Non-Hispanic white, both sexes..... | 23.7 | * | 0.8 | 0.8 | 0.9 | 1.4 | 2.0 | 2.0 | 2.6 | 3.0 |
| Male..... | 26.2 | * | 0.9 | 0.8 | 1.1 | 1.7 | 2.2 | 2.4 | 2.9 | 3.5 |
| Female..... | 21.2 | * | 0.7 | 0.8 | 0.7 | 1.1 | 1.7 | 1.6 | 2.3 | 2.5 |
| Non-Hispanic black, both sexes..... | 14.1 | * | * | * | 1.0 | 1.4 | 1.5 | 2.5 | 3.5 | 3.8 |
| Male..... | 15.1 | * | * | * | 1.2 | 1.6 | 1.4 | 2.9 | 4.6 | 5.1 |
| Female..... | 13.1 | * | * | * | * | * | 1.6 | 2.1 | 2.6 | 2.6 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

| | | | | | | | | | | |
|-------------------------------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| All origins 3/, both sexes..... | 21.8 | 0.6 | 0.8 | 0.7 | 0.7 | 1.1 | 1.5 | 1.5 | 2.1 | 3.3 |
| Male..... | 22.7 | * | 0.8 | 0.8 | 0.7 | 1.3 | 1.9 | 2.0 | 2.5 | 3.7 |
| Female..... | 21.0 | * | 0.8 | 0.6 | 0.8 | 0.9 | 1.0 | 1.0 | 1.7 | 2.8 |
| Hispanic, both sexes..... | 7.8 | * | * | 0.6 | 0.9 | 1.2 | 1.9 | 1.1 | 1.8 | 2.2 |
| Male..... | 7.8 | * | * | * | * | 1.5 | 2.2 | 1.5 | 2.0 | 2.3 |
| Female..... | 7.7 | * | * | * | * | * | 1.5 | * | 1.7 | 2.1 |
| Mexican, both sexes..... | 6.6 | * | * | * | 0.9 | 1.2 | 2.0 | 1.1 | 1.9 | 2.4 |
| Male..... | 6.7 | * | * | * | * | 1.9 | 2.3 | 1.6 | 2.1 | 2.7 |
| Female..... | 6.4 | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | 9.5 | * | * | * | * | * | * | * | * | * |
| Male..... | 11.0 | * | * | * | * | * | * | * | * | * |
| Female..... | 8.2 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 18.6 | * | * | * | * | * | * | * | * | * |
| Male..... | 20.4 | * | * | * | * | * | * | * | * | * |
| Female..... | 16.8 | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 8.9 | * | * | * | * | * | * | * | * | * |
| Male..... | 8.1 | * | * | * | * | * | * | * | * | * |
| Female..... | 9.8 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 24.0 | * | 0.9 | 0.7 | 0.7 | 1.1 | 1.4 | 1.6 | 2.2 | 3.4 |
| Male..... | 25.0 | * | 0.9 | 0.8 | 0.7 | 1.3 | 1.9 | 2.2 | 2.6 | 4.0 |
| Female..... | 22.9 | * | 0.8 | 0.6 | 0.7 | 1.0 | 0.8 | 1.1 | 1.7 | 2.9 |
| Non-Hispanic white, both sexes..... | 26.0 | * | 0.8 | 0.7 | 0.7 | 1.2 | 1.3 | 1.6 | 2.1 | 3.4 |
| Male..... | 27.2 | * | 0.8 | 0.7 | 0.6 | 1.3 | 1.8 | 2.1 | 2.6 | 3.9 |
| Female..... | 24.9 | * | 0.7 | 0.6 | 0.8 | 1.0 | 0.9 | 1.1 | 1.6 | 2.9 |
| Non-Hispanic black, both sexes..... | 18.7 | * | 1.1 | 0.8 | 0.8 | 1.1 | 1.6 | 2.0 | 2.8 | 4.7 |
| Male..... | 19.4 | * | * | * | * | 1.4 | 2.3 | 2.6 | 2.9 | 5.4 |
| Female..... | 18.0 | * | * | * | * | * | * | 1.5 | 2.7 | 4.2 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|-------------------------------------|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
| Anemias (D50-D64) | | | | | | | | | | |
| All origins 3/, both sexes..... | 1.6 | * | 0.1 | 0.1 | 0.1 | 0.2 | 0.3 | 0.4 | 0.5 | 0.5 |
| Male..... | 1.3 | * | * | * | * | 0.2 | 0.4 | 0.5 | 0.5 | 0.5 |
| Female..... | 1.9 | * | * | * | * | * | 0.3 | 0.3 | 0.5 | 0.4 |
| Hispanic, both sexes..... | 0.5 | * | * | * | * | * | * | * | * | * |
| Male..... | 0.5 | * | * | * | * | * | * | * | * | * |
| Female..... | 0.5 | * | * | * | * | * | * | * | * | * |
| Mexican, both sexes..... | 0.4 | * | * | * | * | * | * | * | * | * |
| Male..... | 0.4 | * | * | * | * | * | * | * | * | * |
| Female..... | 0.4 | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | 0.7 | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 1.7 | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 0.6 | * | * | * | * | * | * | * | * | * |
| Male..... | 0.6 | * | * | * | * | * | * | * | * | * |
| Female..... | 0.5 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 1.8 | * | * | 0.2 | 0.1 | 0.2 | 0.4 | 0.5 | 0.5 | 0.5 |
| Male..... | 1.4 | * | * | * | * | 0.2 | 0.5 | 0.7 | 0.5 | 0.5 |
| Female..... | 2.1 | * | * | * | * | * | 0.3 | 0.4 | 0.5 | 0.5 |
| Non-Hispanic white, both sexes..... | 1.6 | * | * | * | * | * | * | * | 0.2 | * |
| Male..... | 1.2 | * | * | * | * | * | * | * | * | * |
| Female..... | 2.1 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic black, both sexes..... | 2.9 | * | * | * | * | 0.9 | 1.8 | 2.6 | 2.4 | 2.6 |
| Male..... | 2.7 | * | * | * | * | * | 2.3 | 3.3 | 2.6 | 2.7 |
| Female..... | 3.0 | * | * | * | * | * | * | 1.9 | 2.1 | 2.5 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|-------------------------------------|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
| Diabetes mellitus (E10-E14) | | | | | | | | | | |
| All origins 3/, both sexes..... | 25.4 | * | * | * | 0.1 | 0.3 | 0.6 | 1.2 | 2.0 | 3.5 |
| Male..... | 24.2 | * | * | * | * | 0.3 | 0.7 | 1.3 | 2.4 | 4.1 |
| Female..... | 26.5 | * | * | * | * | 0.3 | 0.5 | 1.1 | 1.5 | 3.0 |
| Hispanic, both sexes..... | 15.3 | * | * | * | * | * | * | 0.9 | 0.7 | 2.3 |
| Male..... | 13.9 | * | * | * | * | * | * | * | * | 2.5 |
| Female..... | 16.7 | * | * | * | * | * | * | * | * | 2.1 |
| Mexican, both sexes..... | 14.9 | * | * | * | * | * | * | 0.8 | * | 2.3 |
| Male..... | 13.6 | * | * | * | * | * | * | * | * | 2.2 |
| Female..... | 16.3 | * | * | * | * | * | * | * | * | 2.4 |
| Puerto Rican, both sexes..... | 21.1 | * | * | * | * | * | * | * | * | * |
| Male..... | 19.8 | * | * | * | * | * | * | * | * | * |
| Female..... | 22.4 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 28.0 | * | * | * | * | * | * | * | * | * |
| Male..... | 26.7 | * | * | * | * | * | * | * | * | * |
| Female..... | 29.3 | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 11.6 | * | * | * | * | * | * | * | * | * |
| Male..... | 10.2 | * | * | * | * | * | * | * | * | * |
| Female..... | 13.1 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 26.9 | * | * | * | 0.2 | 0.3 | 0.6 | 1.3 | 2.2 | 3.7 |
| Male..... | 25.8 | * | * | * | * | 0.3 | 0.7 | 1.4 | 2.8 | 4.4 |
| Female..... | 27.9 | * | * | * | * | 0.3 | 0.5 | 1.2 | 1.7 | 3.1 |
| Non-Hispanic white, both sexes..... | 26.4 | * | * | * | * | 0.2 | 0.5 | 1.0 | 2.0 | 3.2 |
| Male..... | 26.0 | * | * | * | * | * | 0.5 | 1.2 | 2.4 | 3.9 |
| Female..... | 26.8 | * | * | * | * | * | 0.5 | 0.9 | 1.5 | 2.6 |
| Non-Hispanic black, both sexes..... | 34.8 | * | * | * | * | * | 1.4 | 2.9 | 4.5 | 7.5 |
| Male..... | 30.0 | * | * | * | * | * | 1.8 | 3.1 | 6.0 | 8.7 |
| Female..... | 39.2 | * | * | * | * | * | * | 2.8 | 3.3 | 6.4 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|---|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
| Major cardiovascular diseases (I00-I78) | | | | | | | | | | |
| All origins 3/, both sexes..... | 318.6 | 15.9 | 1.4 | 0.6 | 1.1 | 2.4 | 3.9 | 7.1 | 12.6 | 25.3 |
| Male..... | 303.3 | 16.6 | 1.5 | 0.6 | 1.3 | 3.1 | 5.0 | 9.0 | 16.1 | 32.9 |
| Female..... | 333.3 | 15.1 | 1.3 | 0.7 | 0.9 | 1.7 | 2.8 | 5.1 | 9.0 | 17.7 |
| Hispanic, both sexes..... | 94.3 | 13.1 | 1.2 | * | 1.0 | 1.8 | 3.3 | 4.7 | 7.5 | 13.8 |
| Male..... | 94.2 | 12.2 | * | * | * | 2.2 | 4.3 | 6.5 | 9.8 | 17.4 |
| Female..... | 94.4 | 14.0 | 1.3 | * | 1.2 | 1.4 | 2.0 | 2.5 | 4.9 | 9.8 |
| Mexican, both sexes..... | 75.1 | 12.9 | 1.2 | * | * | 1.8 | 3.4 | 4.5 | 7.5 | 13.3 |
| Male..... | 76.2 | 11.5 | * | * | * | 2.2 | 4.5 | 6.4 | 9.5 | 16.4 |
| Female..... | 73.8 | 14.4 | * | * | * | * | 2.0 | 2.1 | 5.0 | 9.7 |
| Puerto Rican, both sexes..... | 138.0 | * | * | * | * | * | * | * | 10.7 | 19.2 |
| Male..... | 148.6 | * | * | * | * | * | * | * | * | 24.3 |
| Female..... | 128.2 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 352.6 | * | * | * | * | * | * | * | * | * |
| Male..... | 334.7 | * | * | * | * | * | * | * | * | * |
| Female..... | 371.0 | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 91.7 | * | * | * | * | * | * | 4.4 | 6.6 | 12.6 |
| Male..... | 89.4 | * | * | * | * | * | * | 5.4 | 9.1 | 16.4 |
| Female..... | 94.1 | * | * | * | * | * | * | * | * | 8.8 |
| Non-Hispanic 5/, both sexes..... | 352.2 | 16.5 | 1.5 | 0.7 | 1.1 | 2.5 | 4.0 | 7.7 | 13.6 | 27.1 |
| Male..... | 336.4 | 17.6 | 1.7 | 0.6 | 1.4 | 3.3 | 5.1 | 9.7 | 17.4 | 35.5 |
| Female..... | 367.2 | 15.3 | 1.3 | 0.8 | 0.9 | 1.7 | 2.9 | 5.7 | 9.8 | 18.9 |
| Non-Hispanic white, both sexes..... | 381.9 | 13.4 | 1.4 | 0.6 | 0.9 | 2.2 | 3.2 | 6.0 | 10.8 | 22.9 |
| Male..... | 362.2 | 14.8 | 1.6 | 0.6 | 1.1 | 2.9 | 4.2 | 7.7 | 14.1 | 30.6 |
| Female..... | 400.8 | 12.0 | 1.1 | 0.7 | 0.7 | 1.5 | 2.2 | 4.2 | 7.5 | 15.2 |
| Non-Hispanic black, both sexes..... | 285.1 | 28.3 | 2.1 | 0.8 | 2.0 | 4.0 | 8.1 | 17.4 | 30.7 | 54.7 |
| Male..... | 277.2 | 29.1 | 2.0 | * | 2.4 | 5.1 | 9.7 | 21.6 | 38.2 | 68.8 |
| Female..... | 292.3 | 27.4 | 2.2 | * | 1.6 | 2.9 | 6.6 | 13.5 | 24.0 | 42.2 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|---|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
| Diseases of heart (I00-I09,I11,I13,I20-I51) | | | | | | | | | | |
| All origins 3/, both sexes..... | 241.7 | 12.4 | 1.1 | 0.5 | 0.8 | 2.0 | 3.1 | 5.7 | 10.0 | 20.3 |
| Male..... | 240.7 | 12.9 | 1.1 | 0.4 | 0.9 | 2.6 | 4.0 | 7.4 | 13.4 | 27.8 |
| Female..... | 242.7 | 11.8 | 1.0 | 0.5 | 0.6 | 1.3 | 2.1 | 3.9 | 6.5 | 12.8 |
| Hispanic, both sexes..... | 71.9 | 10.2 | 0.8 | * | * | 1.2 | 2.5 | 3.4 | 5.5 | 9.9 |
| Male..... | 74.0 | 9.6 | * | * | * | 1.6 | 3.4 | 4.8 | 7.5 | 12.9 |
| Female..... | 69.7 | 10.8 | * | * | * | * | 1.3 | 1.7 | 3.1 | 6.6 |
| Mexican, both sexes..... | 56.0 | 9.9 | * | * | * | 1.2 | 2.4 | 3.3 | 5.3 | 9.6 |
| Male..... | 58.8 | 9.1 | * | * | * | * | 3.5 | 4.9 | 6.8 | 11.8 |
| Female..... | 52.8 | 10.8 | * | * | * | * | * | * | 3.3 | 7.0 |
| Puerto Rican, both sexes..... | 109.2 | * | * | * | * | * | * | * | 7.9 | 13.8 |
| Male..... | 118.7 | * | * | * | * | * | * | * | * | * |
| Female..... | 100.5 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 290.4 | * | * | * | * | * | * | * | * | * |
| Male..... | 281.0 | * | * | * | * | * | * | * | * | * |
| Female..... | 300.0 | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 68.7 | * | * | * | * | * | * | 2.8 | 5.2 | 8.7 |
| Male..... | 69.9 | * | * | * | * | * | * | * | 8.1 | 12.1 |
| Female..... | 67.4 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 267.1 | 12.9 | 1.1 | 0.5 | 0.8 | 2.1 | 3.2 | 6.2 | 10.8 | 21.9 |
| Male..... | 267.0 | 13.7 | 1.2 | 0.4 | 1.0 | 2.8 | 4.1 | 8.1 | 14.6 | 30.3 |
| Female..... | 267.2 | 12.0 | 1.1 | 0.6 | 0.6 | 1.4 | 2.2 | 4.3 | 7.1 | 13.7 |
| Non-Hispanic white, both sexes..... | 290.8 | 10.7 | 1.0 | 0.4 | 0.7 | 1.8 | 2.4 | 4.8 | 8.7 | 19.2 |
| Male..... | 289.2 | 11.6 | 1.1 | 0.4 | 0.9 | 2.4 | 3.2 | 6.4 | 12.0 | 26.9 |
| Female..... | 292.3 | 9.6 | 0.8 | 0.5 | 0.5 | 1.2 | 1.6 | 3.3 | 5.4 | 11.4 |
| Non-Hispanic black, both sexes..... | 212.2 | 21.9 | 1.7 | * | 1.4 | 3.5 | 6.7 | 14.1 | 24.1 | 42.1 |
| Male..... | 212.9 | 23.0 | * | * | 1.6 | 4.4 | 8.3 | 18.5 | 31.1 | 56.0 |
| Female..... | 211.6 | 20.8 | 1.9 | * | * | 2.5 | 5.3 | 10.1 | 17.7 | 29.7 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|-------------------------------------|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
| Hypertensive heart disease (I11) | | | | | | | | | | |
| All origins 3/, both sexes..... | 9.2 | * | * | * | * | * | 0.1 | 0.4 | 1.0 | 1.8 |
| Male..... | 8.2 | * | * | * | * | * | 0.2 | 0.7 | 1.4 | 2.6 |
| Female..... | 10.2 | * | * | * | * | * | * | * | 0.6 | 1.0 |
| Hispanic, both sexes..... | 3.7 | * | * | * | * | * | * | * | 0.6 | 1.2 |
| Male..... | 3.9 | * | * | * | * | * | * | * | * | 1.7 |
| Female..... | 3.4 | * | * | * | * | * | * | * | * | * |
| Mexican, both sexes..... | 2.8 | * | * | * | * | * | * | * | * | 1.3 |
| Male..... | 3.0 | * | * | * | * | * | * | * | * | 1.8 |
| Female..... | 2.6 | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | 6.4 | * | * | * | * | * | * | * | * | * |
| Male..... | 7.5 | * | * | * | * | * | * | * | * | * |
| Female..... | 5.3 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 10.9 | * | * | * | * | * | * | * | * | * |
| Male..... | 10.5 | * | * | * | * | * | * | * | * | * |
| Female..... | 11.3 | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 3.9 | * | * | * | * | * | * | * | * | * |
| Male..... | 4.2 | * | * | * | * | * | * | * | * | * |
| Female..... | 3.6 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 10.0 | * | * | * | * | * | 0.1 | 0.5 | 1.1 | 1.9 |
| Male..... | 8.9 | * | * | * | * | * | * | 0.8 | 1.5 | 2.8 |
| Female..... | 11.1 | * | * | * | * | * | * | * | 0.7 | 1.1 |
| Non-Hispanic white, both sexes..... | 9.1 | * | * | * | * | * | * | 0.3 | 0.6 | 1.3 |
| Male..... | 7.6 | * | * | * | * | * | * | 0.4 | 0.9 | 1.9 |
| Female..... | 10.5 | * | * | * | * | * | * | * | 0.3 | 0.7 |
| Non-Hispanic black, both sexes..... | 17.7 | * | * | * | * | * | * | 1.6 | 4.1 | 6.0 |
| Male..... | 18.0 | * | * | * | * | * | * | 2.5 | 5.6 | 8.7 |
| Female..... | 17.4 | * | * | * | * | * | * | * | 2.8 | 3.5 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|-------------------------------------|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
| Ischemic heart diseases (I20-I25) | | | | | | | | | | |
| All origins 3/, both sexes..... | 171.4 | 0.6 | * | * | * | 0.1 | 0.5 | 1.3 | 3.6 | 10.1 |
| Male..... | 178.4 | * | * | * | * | * | 0.7 | 2.0 | 5.2 | 15.2 |
| Female..... | 164.7 | * | * | * | * | * | 0.3 | 0.6 | 1.9 | 4.9 |
| Hispanic, both sexes..... | 54.0 | * | * | * | * | * | * | 0.8 | 1.6 | 4.6 |
| Male..... | 56.5 | * | * | * | * | * | * | 1.3 | 2.6 | 6.4 |
| Female..... | 51.4 | * | * | * | * | * | * | * | * | 2.5 |
| Mexican, both sexes..... | 41.8 | * | * | * | * | * | * | * | 1.5 | 4.4 |
| Male..... | 44.8 | * | * | * | * | * | * | * | 2.4 | 5.8 |
| Female..... | 38.4 | * | * | * | * | * | * | * | * | 2.6 |
| Puerto Rican, both sexes..... | 82.2 | * | * | * | * | * | * | * | * | * |
| Male..... | 89.5 | * | * | * | * | * | * | * | * | * |
| Female..... | 75.4 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 232.3 | * | * | * | * | * | * | * | * | * |
| Male..... | 224.2 | * | * | * | * | * | * | * | * | * |
| Female..... | 240.6 | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 49.9 | * | * | * | * | * | * | * | * | 3.8 |
| Male..... | 52.4 | * | * | * | * | * | * | * | * | * |
| Female..... | 47.3 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 189.0 | * | * | * | * | 0.2 | 0.5 | 1.4 | 3.9 | 11.0 |
| Male..... | 197.6 | * | * | * | * | * | 0.7 | 2.2 | 5.7 | 16.8 |
| Female..... | 180.7 | * | * | * | * | * | 0.3 | 0.7 | 2.1 | 5.3 |
| Non-Hispanic white, both sexes..... | 208.0 | * | * | * | * | 0.2 | 0.4 | 1.3 | 3.7 | 10.6 |
| Male..... | 217.2 | * | * | * | * | * | 0.6 | 2.1 | 5.5 | 16.3 |
| Female..... | 199.1 | * | * | * | * | * | * | 0.6 | 1.9 | 4.8 |
| Non-Hispanic black, both sexes..... | 137.0 | * | * | * | * | * | 0.9 | 2.2 | 5.8 | 15.5 |
| Male..... | 139.5 | * | * | * | * | * | * | 2.9 | 7.7 | 22.5 |
| Female..... | 134.8 | * | * | * | * | * | * | 1.6 | 4.1 | 9.3 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|---------------------------------------|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
| Acute myocardial infarction (I21-I22) | | | | | | | | | | |
| All origins 3/, both sexes..... | 62.3 | * | * | * | * | * | 0.2 | 0.5 | 1.5 | 4.0 |
| Male..... | 66.2 | * | * | * | * | * | 0.3 | 0.7 | 2.1 | 6.0 |
| Female..... | 58.4 | * | * | * | * | * | * | 0.3 | 0.8 | 2.1 |
| Hispanic, both sexes..... | 19.7 | * | * | * | * | * | * | * | 0.7 | 1.9 |
| Male..... | 20.6 | * | * | * | * | * | * | * | 1.1 | 2.3 |
| Female..... | 18.7 | * | * | * | * | * | * | * | * | 1.3 |
| Mexican, both sexes..... | 16.5 | * | * | * | * | * | * | * | * | 1.7 |
| Male..... | 17.8 | * | * | * | * | * | * | * | * | 1.9 |
| Female..... | 15.0 | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | 26.2 | * | * | * | * | * | * | * | * | * |
| Male..... | 28.7 | * | * | * | * | * | * | * | * | * |
| Female..... | 23.8 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 81.7 | * | * | * | * | * | * | * | * | * |
| Male..... | 77.2 | * | * | * | * | * | * | * | * | * |
| Female..... | 86.2 | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 16.3 | * | * | * | * | * | * | * | * | * |
| Male..... | 16.9 | * | * | * | * | * | * | * | * | * |
| Female..... | 15.7 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 68.7 | * | * | * | * | * | 0.3 | 0.5 | 1.6 | 4.4 |
| Male..... | 73.5 | * | * | * | * | * | 0.3 | 0.8 | 2.3 | 6.6 |
| Female..... | 64.1 | * | * | * | * | * | * | 0.3 | 0.9 | 2.2 |
| Non-Hispanic white, both sexes..... | 75.5 | * | * | * | * | * | 0.2 | 0.5 | 1.6 | 4.3 |
| Male..... | 81.2 | * | * | * | * | * | * | 0.7 | 2.3 | 6.7 |
| Female..... | 70.1 | * | * | * | * | * | * | * | 0.8 | 2.0 |
| Non-Hispanic black, both sexes..... | 50.2 | * | * | * | * | * | * | 1.0 | 2.4 | 5.7 |
| Male..... | 50.1 | * | * | * | * | * | * | * | 3.1 | 7.8 |
| Female..... | 50.4 | * | * | * | * | * | * | * | 1.7 | 3.7 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|---|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
| Other forms of chronic ischemic heart disease (I20,I25) | | | | | | | | | | |
| All origins 3/, both sexes..... | 108.0 | * | * | * | * | * | 0.3 | 0.8 | 2.0 | 6.0 |
| Male..... | 110.9 | * | * | * | * | * | 0.4 | 1.3 | 3.0 | 9.2 |
| Female..... | 105.2 | * | * | * | * | * | * | 0.4 | 1.0 | 2.8 |
| Hispanic, both sexes..... | 34.2 | * | * | * | * | * | * | * | 1.0 | 2.7 |
| Male..... | 35.7 | * | * | * | * | * | * | * | 1.5 | 4.1 |
| Female..... | 32.6 | * | * | * | * | * | * | * | * | * |
| Mexican, both sexes..... | 25.2 | * | * | * | * | * | * | * | 0.9 | 2.7 |
| Male..... | 26.9 | * | * | * | * | * | * | * | * | 4.0 |
| Female..... | 23.3 | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | 55.8 | * | * | * | * | * | * | * | * | * |
| Male..... | 60.5 | * | * | * | * | * | * | * | * | * |
| Female..... | 51.4 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 150.5 | * | * | * | * | * | * | * | * | * |
| Male..... | 146.7 | * | * | * | * | * | * | * | * | * |
| Female..... | 154.3 | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 33.3 | * | * | * | * | * | * | * | * | * |
| Male..... | 35.2 | * | * | * | * | * | * | * | * | * |
| Female..... | 31.4 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 118.9 | * | * | * | * | * | 0.3 | 0.9 | 2.2 | 6.5 |
| Male..... | 122.7 | * | * | * | * | * | 0.4 | 1.3 | 3.3 | 10.0 |
| Female..... | 115.4 | * | * | * | * | * | * | 0.4 | 1.1 | 3.1 |
| Non-Hispanic white, both sexes..... | 131.0 | * | * | * | * | * | 0.2 | 0.9 | 2.1 | 6.1 |
| Male..... | 134.5 | * | * | * | * | * | 0.4 | 1.3 | 3.1 | 9.5 |
| Female..... | 127.7 | * | * | * | * | * | * | 0.4 | 1.0 | 2.8 |
| Non-Hispanic black, both sexes..... | 85.5 | * | * | * | * | * | * | 1.2 | 3.3 | 9.8 |
| Male..... | 88.0 | * | * | * | * | * | * | * | 4.3 | 14.6 |
| Female..... | 83.3 | * | * | * | * | * | * | * | 2.4 | 5.5 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

| | | | | | | | | | | |
|-------------------------------------|-------|---|---|---|---|---|-----|-----|-----|-----|
| All origins 3/, both sexes..... | 84.4 | * | * | * | * | * | 0.2 | 0.6 | 1.3 | 4.1 |
| Male..... | 85.5 | * | * | * | * | * | 0.3 | 0.9 | 1.9 | 6.2 |
| Female..... | 83.3 | * | * | * | * | * | * | 0.3 | 0.8 | 2.0 |
| Hispanic, both sexes..... | 26.0 | * | * | * | * | * | * | * | * | 1.7 |
| Male..... | 25.9 | * | * | * | * | * | * | * | * | 2.8 |
| Female..... | 26.1 | * | * | * | * | * | * | * | * | * |
| Mexican, both sexes..... | 18.8 | * | * | * | * | * | * | * | * | 1.8 |
| Male..... | 19.3 | * | * | * | * | * | * | * | * | 2.8 |
| Female..... | 18.2 | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | 44.1 | * | * | * | * | * | * | * | * | * |
| Male..... | 44.2 | * | * | * | * | * | * | * | * | * |
| Female..... | 44.0 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 128.3 | * | * | * | * | * | * | * | * | * |
| Male..... | 123.4 | * | * | * | * | * | * | * | * | * |
| Female..... | 133.2 | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 23.5 | * | * | * | * | * | * | * | * | * |
| Male..... | 23.5 | * | * | * | * | * | * | * | * | * |
| Female..... | 23.4 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 93.1 | * | * | * | * | * | 0.2 | 0.7 | 1.5 | 4.5 |
| Male..... | 94.9 | * | * | * | * | * | 0.3 | 1.1 | 2.2 | 6.8 |
| Female..... | 91.4 | * | * | * | * | * | * | 0.3 | 0.8 | 2.3 |
| Non-Hispanic white, both sexes..... | 104.3 | * | * | * | * | * | 0.2 | 0.7 | 1.4 | 4.3 |
| Male..... | 106.2 | * | * | * | * | * | * | 1.1 | 2.1 | 6.6 |
| Female..... | 102.4 | * | * | * | * | * | * | * | 0.7 | 2.1 |
| Non-Hispanic black, both sexes..... | 57.8 | * | * | * | * | * | * | 1.0 | 2.0 | 6.2 |
| Male..... | 56.1 | * | * | * | * | * | * | * | 2.4 | 9.0 |
| Female..... | 59.3 | * | * | * | * | * | * | * | 1.6 | 3.7 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|-------------------------------------|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
| Other heart diseases (I26-I51) | | | | | | | | | | |
| All origins 3/, both sexes..... | 58.8 | 11.7 | 1.0 | 0.4 | 0.7 | 1.8 | 2.3 | 3.8 | 5.2 | 8.1 |
| Male..... | 52.4 | 12.1 | 1.0 | 0.4 | 0.8 | 2.4 | 3.0 | 4.6 | 6.6 | 9.6 |
| Female..... | 65.0 | 11.4 | 1.0 | 0.5 | 0.5 | 1.1 | 1.7 | 3.0 | 3.8 | 6.6 |
| Hispanic, both sexes..... | 13.3 | 9.7 | 0.7 | * | * | 1.1 | 1.9 | 2.3 | 2.9 | 3.9 |
| Male..... | 12.9 | 9.1 | * | * | * | 1.5 | 2.6 | 3.1 | 3.7 | 4.6 |
| Female..... | 13.7 | 10.3 | * | * | * | * | * | 1.2 | 1.9 | 3.1 |
| Mexican, both sexes..... | 10.6 | 9.7 | * | * | * | 1.1 | 1.8 | 2.3 | 3.0 | 3.8 |
| Male..... | 10.5 | 8.7 | * | * | * | * | 2.6 | 3.3 | 3.5 | 4.1 |
| Female..... | 10.7 | 10.8 | * | * | * | * | * | * | 2.3 | 3.5 |
| Puerto Rican, both sexes..... | 19.3 | * | * | * | * | * | * | * | * | * |
| Male..... | 20.5 | * | * | * | * | * | * | * | * | * |
| Female..... | 18.2 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 44.4 | * | * | * | * | * | * | * | * | * |
| Male..... | 43.7 | * | * | * | * | * | * | * | * | * |
| Female..... | 45.2 | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 14.0 | * | * | * | * | * | * | * | * | 3.5 |
| Male..... | 12.7 | * | * | * | * | * | * | * | * | * |
| Female..... | 15.3 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 65.7 | 12.2 | 1.1 | 0.5 | 0.7 | 1.9 | 2.4 | 4.2 | 5.6 | 8.7 |
| Male..... | 58.7 | 12.8 | 1.1 | 0.4 | 0.9 | 2.6 | 3.1 | 5.0 | 7.1 | 10.4 |
| Female..... | 72.3 | 11.6 | 1.0 | 0.5 | 0.5 | 1.2 | 1.8 | 3.3 | 4.1 | 7.1 |
| Non-Hispanic white, both sexes..... | 71.3 | 10.0 | 0.9 | 0.4 | 0.6 | 1.7 | 1.9 | 3.2 | 4.3 | 7.1 |
| Male..... | 62.7 | 10.7 | 1.1 | 0.4 | 0.8 | 2.2 | 2.4 | 3.8 | 5.5 | 8.5 |
| Female..... | 79.6 | 9.3 | 0.8 | 0.5 | 0.5 | 1.1 | 1.3 | 2.5 | 3.1 | 5.7 |
| Non-Hispanic black, both sexes..... | 54.3 | 20.9 | 1.6 | * | 1.2 | 3.1 | 5.3 | 10.0 | 13.4 | 19.7 |
| Male..... | 52.6 | 21.8 | * | * | 1.4 | 4.0 | 6.4 | 12.5 | 17.0 | 23.6 |
| Female..... | 55.9 | 19.9 | 1.9 | * | * | 2.3 | 4.2 | 7.7 | 10.3 | 16.1 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
| All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51) | | | | | | | | | | |
| All origins 3/, both sexes..... | 38.5 | 10.5 | 0.8 | 0.3 | 0.6 | 1.6 | 2.1 | 3.5 | 4.6 | 7.3 |
| Male..... | 36.3 | 10.9 | 0.8 | 0.3 | 0.7 | 2.2 | 2.7 | 4.2 | 5.8 | 8.8 |
| Female..... | 40.6 | 10.0 | 0.8 | 0.4 | 0.4 | 1.0 | 1.5 | 2.7 | 3.4 | 5.9 |
| Hispanic, both sexes..... | 9.3 | 8.6 | 0.6 | * | * | 1.0 | 1.8 | 2.2 | 2.6 | 3.6 |
| Male..... | 9.6 | 8.0 | * | * | * | 1.4 | 2.5 | 2.9 | 3.3 | 4.4 |
| Female..... | 9.0 | 9.3 | * | * | * | * | * | 1.2 | 1.8 | 2.7 |
| Mexican, both sexes..... | 7.2 | 8.6 | * | * | * | 1.1 | 1.8 | 2.2 | 2.7 | 3.5 |
| Male..... | 7.3 | 7.5 | * | * | * | * | 2.5 | 3.1 | 3.1 | 3.9 |
| Female..... | 7.1 | 9.8 | * | * | * | * | * | * | 2.1 | 3.2 |
| Puerto Rican, both sexes..... | 13.0 | * | * | * | * | * | * | * | * | * |
| Male..... | 15.5 | * | * | * | * | * | * | * | * | * |
| Female..... | 10.7 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 35.0 | * | * | * | * | * | * | * | * | * |
| Male..... | 35.7 | * | * | * | * | * | * | * | * | * |
| Female..... | 34.3 | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 9.9 | * | * | * | * | * | * | * | * | 3.1 |
| Male..... | 10.1 | * | * | * | * | * | * | * | * | * |
| Female..... | 9.7 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 42.9 | 10.9 | 0.8 | 0.4 | 0.6 | 1.7 | 2.2 | 3.8 | 4.9 | 7.9 |
| Male..... | 40.6 | 11.6 | 0.8 | 0.3 | 0.8 | 2.3 | 2.8 | 4.6 | 6.3 | 9.5 |
| Female..... | 45.1 | 10.1 | 0.8 | 0.4 | 0.4 | 1.1 | 1.6 | 3.0 | 3.7 | 6.4 |
| Non-Hispanic white, both sexes..... | 45.8 | 9.2 | 0.7 | 0.3 | 0.6 | 1.5 | 1.7 | 2.8 | 3.8 | 6.5 |
| Male..... | 42.7 | 9.8 | 0.8 | 0.3 | 0.7 | 2.1 | 2.2 | 3.4 | 4.8 | 7.8 |
| Female..... | 48.9 | 8.5 | 0.6 | 0.4 | 0.4 | 1.0 | 1.1 | 2.3 | 2.7 | 5.1 |
| Non-Hispanic black, both sexes..... | 39.1 | 17.8 | 1.2 | * | 1.0 | 2.9 | 4.9 | 9.1 | 11.8 | 17.7 |
| Male..... | 39.6 | 19.7 | * | * | 1.2 | 3.6 | 5.9 | 11.5 | 14.9 | 21.3 |
| Female..... | 38.7 | 15.8 | * | * | * | 2.1 | 4.0 | 7.0 | 9.1 | 14.5 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|-------------------------------------|----------------|--------------------|--------------|--------------|----------------|----------------|----------------|----------------|----------------|----------------|
| Cerebrovascular diseases (I60-I69) | | | | | | | | | | |
| All origins 3/, both sexes..... | 56.4 | 2.9 | 0.3 | 0.2 | 0.3 | 0.3 | 0.6 | 1.0 | 1.8 | 3.6 |
| Male..... | 44.2 | 3.2 | 0.4 | * | 0.3 | 0.3 | 0.7 | 1.0 | 1.7 | 3.6 |
| Female..... | 68.2 | 2.5 | 0.3 | * | 0.3 | * | 0.5 | 0.9 | 1.9 | 3.7 |
| Hispanic, both sexes..... | 16.6 | 2.4 | * | * | * | * | * | 1.1 | 1.5 | 3.0 |
| Male..... | 15.0 | * | * | * | * | * | * | 1.4 | 1.7 | 3.4 |
| Female..... | 18.4 | * | * | * | * | * | * | * | 1.3 | 2.5 |
| Mexican, both sexes..... | 14.5 | * | * | * | * | * | * | 1.0 | 1.7 | 3.1 |
| Male..... | 13.3 | * | * | * | * | * | * | * | 2.1 | 3.6 |
| Female..... | 15.8 | * | * | * | * | * | * | * | * | 2.4 |
| Puerto Rican, both sexes..... | 20.8 | * | * | * | * | * | * | * | * | * |
| Male..... | 22.2 | * | * | * | * | * | * | * | * | * |
| Female..... | 19.5 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 45.5 | * | * | * | * | * | * | * | * | * |
| Male..... | 37.8 | * | * | * | * | * | * | * | * | * |
| Female..... | 53.3 | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 16.7 | * | * | * | * | * | * | * | * | * |
| Male..... | 13.8 | * | * | * | * | * | * | * | * | * |
| Female..... | 19.6 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 62.4 | 3.0 | 0.3 | 0.2 | 0.2 | 0.3 | 0.6 | 0.9 | 1.9 | 3.7 |
| Male..... | 48.9 | 3.5 | 0.4 | * | 0.3 | 0.3 | 0.7 | 0.9 | 1.8 | 3.6 |
| Female..... | 75.3 | 2.5 | * | * | * | * | 0.5 | 0.9 | 2.0 | 3.9 |
| Non-Hispanic white, both sexes..... | 67.0 | 2.2 | 0.3 | 0.2 | 0.2 | 0.2 | 0.6 | 0.7 | 1.5 | 2.8 |
| Male..... | 51.3 | 2.7 | * | * | * | * | 0.7 | 0.8 | 1.3 | 2.6 |
| Female..... | 82.1 | * | * | * | * | * | 0.5 | 0.7 | 1.7 | 3.0 |
| Non-Hispanic black, both sexes..... | 51.7 | 5.6 | * | * | * | * | 0.9 | 2.0 | 4.4 | 8.5 |
| Male..... | 45.1 | * | * | * | * | * | * | 1.7 | 4.1 | 7.8 |
| Female..... | 57.7 | * | * | * | * | * | * | 2.3 | 4.6 | 9.2 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|-------------------------------------|----------------|--------------------|--------------|--------------|----------------|----------------|----------------|----------------|----------------|----------------|
| Influenza and pneumonia (J10-J18) | | | | | | | | | | |
| All origins 3/, both sexes..... | 22.8 | 6.5 | 0.7 | 0.2 | 0.3 | 0.4 | 0.5 | 0.7 | 1.0 | 1.7 |
| Male..... | 20.4 | 7.8 | 0.8 | * | 0.2 | 0.4 | 0.5 | 0.7 | 1.2 | 1.8 |
| Female..... | 25.1 | 5.1 | 0.6 | 0.2 | 0.3 | 0.3 | 0.4 | 0.7 | 0.8 | 1.6 |
| Hispanic, both sexes..... | 7.3 | 5.8 | * | * | * | * | * | * | 0.7 | 1.2 |
| Male..... | 7.0 | 7.7 | * | * | * | * | * | * | * | 1.4 |
| Female..... | 7.6 | * | * | * | * | * | * | * | * | * |
| Mexican, both sexes..... | 5.7 | 5.7 | * | * | * | * | * | * | * | * |
| Male..... | 5.6 | 7.5 | * | * | * | * | * | * | * | * |
| Female..... | 5.7 | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | 11.8 | * | * | * | * | * | * | * | * | * |
| Male..... | 11.5 | * | * | * | * | * | * | * | * | * |
| Female..... | 12.1 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 23.2 | * | * | * | * | * | * | * | * | * |
| Male..... | 20.5 | * | * | * | * | * | * | * | * | * |
| Female..... | 25.9 | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 7.7 | * | * | * | * | * | * | * | * | * |
| Male..... | 7.4 | * | * | * | * | * | * | * | * | * |
| Female..... | 8.1 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 25.1 | 6.7 | 0.7 | 0.2 | 0.2 | 0.4 | 0.5 | 0.8 | 1.1 | 1.7 |
| Male..... | 22.5 | 7.8 | 0.8 | * | * | 0.4 | 0.6 | 0.7 | 1.2 | 1.9 |
| Female..... | 27.5 | 5.4 | 0.7 | * | 0.3 | 0.4 | 0.4 | 0.8 | 0.9 | 1.6 |
| Non-Hispanic white, both sexes..... | 27.9 | 4.9 | 0.5 | 0.2 | 0.2 | 0.3 | 0.5 | 0.7 | 0.9 | 1.5 |
| Male..... | 24.6 | 6.1 | 0.5 | * | * | 0.4 | 0.6 | 0.7 | 1.1 | 1.5 |
| Female..... | 31.1 | 3.6 | 0.5 | * | 0.3 | * | 0.4 | 0.8 | 0.8 | 1.4 |
| Non-Hispanic black, both sexes..... | 16.1 | 13.3 | 1.7 | * | * | * | 0.7 | 1.1 | 2.0 | 3.6 |
| Male..... | 15.9 | 14.2 | 1.8 | * | * | * | * | * | 2.4 | 4.1 |
| Female..... | 16.2 | 12.3 | * | * | * | * | * | * | 1.6 | 3.2 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|-------------------------------------|----------------|--------------------|--------------|--------------|----------------|----------------|----------------|----------------|----------------|----------------|
| Pneumonia (J12-J18) | | | | | | | | | | |
| All origins 3/, both sexes..... | 22.5 | 6.3 | 0.7 | 0.2 | 0.2 | 0.3 | 0.4 | 0.7 | 1.0 | 1.7 |
| Male..... | 20.2 | 7.6 | 0.8 | * | 0.2 | 0.4 | 0.5 | 0.7 | 1.2 | 1.8 |
| Female..... | 24.8 | 5.0 | 0.6 | * | 0.2 | 0.3 | 0.4 | 0.7 | 0.8 | 1.5 |
| Hispanic, both sexes..... | 7.2 | 5.5 | * | * | * | * | * | * | 0.7 | 1.1 |
| Male..... | 6.9 | 7.3 | * | * | * | * | * | * | * | 1.3 |
| Female..... | 7.5 | * | * | * | * | * | * | * | * | * |
| Mexican, both sexes..... | 5.6 | 5.7 | * | * | * | * | * | * | * | * |
| Male..... | 5.6 | 7.5 | * | * | * | * | * | * | * | * |
| Female..... | 5.7 | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | 11.7 | * | * | * | * | * | * | * | * | * |
| Male..... | 11.4 | * | * | * | * | * | * | * | * | * |
| Female..... | 12.1 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 23.1 | * | * | * | * | * | * | * | * | * |
| Male..... | 20.5 | * | * | * | * | * | * | * | * | * |
| Female..... | 25.8 | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 7.6 | * | * | * | * | * | * | * | * | * |
| Male..... | 7.3 | * | * | * | * | * | * | * | * | * |
| Female..... | 8.0 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 24.8 | 6.5 | 0.7 | 0.2 | 0.2 | 0.3 | 0.5 | 0.7 | 1.1 | 1.7 |
| Male..... | 22.3 | 7.6 | 0.8 | * | * | 0.4 | 0.6 | 0.7 | 1.2 | 1.9 |
| Female..... | 27.2 | 5.3 | 0.6 | * | * | 0.3 | 0.4 | 0.8 | 0.9 | 1.6 |
| Non-Hispanic white, both sexes..... | 27.6 | 4.7 | 0.5 | 0.2 | 0.2 | 0.3 | 0.5 | 0.7 | 0.9 | 1.5 |
| Male..... | 24.3 | 6.0 | 0.5 | * | * | 0.4 | 0.6 | 0.6 | 1.1 | 1.5 |
| Female..... | 30.7 | 3.4 | 0.5 | * | * | * | 0.4 | 0.8 | 0.8 | 1.4 |
| Non-Hispanic black, both sexes..... | 16.0 | 13.1 | 1.5 | * | * | * | 0.7 | 1.1 | 2.0 | 3.6 |
| Male..... | 15.8 | 13.9 | 1.6 | * | * | * | * | * | 2.4 | 4.1 |
| Female..... | 16.1 | 12.3 | * | * | * | * | * | * | 1.6 | 3.2 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|

Chronic lower respiratory diseases (J40-J47)

| | | | | | | | | | | |
|-------------------------------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| All origins 3/, both sexes..... | 43.3 | 1.0 | 0.4 | 0.2 | 0.4 | 0.5 | 0.5 | 0.6 | 0.9 | 1.6 |
| Male..... | 42.9 | 1.1 | 0.6 | 0.2 | 0.5 | 0.6 | 0.5 | 0.7 | 0.9 | 1.5 |
| Female..... | 43.7 | * | 0.3 | * | 0.4 | 0.3 | 0.5 | 0.5 | 0.9 | 1.7 |
| Hispanic, both sexes..... | 7.9 | * | * | * | * | * | * | * | 0.6 | 0.8 |
| Male..... | 8.1 | * | * | * | * | * | * | * | * | * |
| Female..... | 7.6 | * | * | * | * | * | * | * | * | * |
| Mexican, both sexes..... | 5.5 | * | * | * | * | * | * | * | * | * |
| Male..... | 6.0 | * | * | * | * | * | * | * | * | * |
| Female..... | 5.0 | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | 13.4 | * | * | * | * | * | * | * | * | * |
| Male..... | 12.7 | * | * | * | * | * | * | * | * | * |
| Female..... | 14.1 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 33.6 | * | * | * | * | * | * | * | * | * |
| Male..... | 34.9 | * | * | * | * | * | * | * | * | * |
| Female..... | 32.2 | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 8.6 | * | * | * | * | * | * | * | * | * |
| Male..... | 8.6 | * | * | * | * | * | * | * | * | * |
| Female..... | 8.5 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 48.6 | 1.1 | 0.4 | 0.2 | 0.5 | 0.5 | 0.5 | 0.7 | 0.9 | 1.7 |
| Male..... | 48.4 | 1.2 | 0.6 | 0.2 | 0.5 | 0.7 | 0.5 | 0.8 | 0.9 | 1.6 |
| Female..... | 48.9 | * | * | * | 0.5 | 0.3 | 0.5 | 0.6 | 0.9 | 1.8 |
| Non-Hispanic white, both sexes..... | 56.4 | 0.9 | 0.2 | * | 0.2 | 0.3 | 0.4 | 0.5 | 0.7 | 1.4 |
| Male..... | 55.1 | * | * | * | * | 0.4 | 0.4 | 0.5 | 0.8 | 1.1 |
| Female..... | 57.7 | * | * | * | * | * | 0.5 | 0.5 | 0.7 | 1.6 |
| Non-Hispanic black, both sexes..... | 21.4 | * | 1.2 | 0.7 | 1.6 | 1.4 | 1.2 | 1.7 | 2.2 | 3.9 |
| Male..... | 24.9 | * | 1.6 | * | 1.6 | 2.3 | 1.6 | 2.3 | 2.2 | 4.6 |
| Female..... | 18.2 | * | * | * | 1.5 | * | * | * | 2.2 | 3.3 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|-------------------------------------|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
| Asthma (J45-J46) | | | | | | | | | | |
| All origins 3/, both sexes..... | 1.5 | * | 0.3 | 0.2 | 0.4 | 0.4 | 0.4 | 0.5 | 0.6 | 0.9 |
| Male..... | 1.1 | * | 0.4 | 0.2 | 0.5 | 0.6 | 0.5 | 0.7 | 0.6 | 0.8 |
| Female..... | 1.8 | * | * | * | 0.4 | 0.2 | 0.4 | 0.4 | 0.6 | 1.0 |
| Hispanic, both sexes..... | 0.7 | * | * | * | * | * | * | * | * | * |
| Male..... | 0.6 | * | * | * | * | * | * | * | * | * |
| Female..... | 0.9 | * | * | * | * | * | * | * | * | * |
| Mexican, both sexes..... | 0.4 | * | * | * | * | * | * | * | * | * |
| Male..... | 0.3 | * | * | * | * | * | * | * | * | * |
| Female..... | 0.5 | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | 2.8 | * | * | * | * | * | * | * | * | * |
| Male..... | 2.5 | * | * | * | * | * | * | * | * | * |
| Female..... | 3.0 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 1.6 | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 0.8 | * | * | * | * | * | * | * | * | * |
| Male..... | 0.6 | * | * | * | * | * | * | * | * | * |
| Female..... | 1.0 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 1.6 | * | 0.3 | 0.2 | 0.4 | 0.5 | 0.5 | 0.6 | 0.7 | 1.0 |
| Male..... | 1.2 | * | 0.4 | * | 0.5 | 0.7 | 0.5 | 0.7 | 0.7 | 0.9 |
| Female..... | 2.0 | * | * | * | 0.4 | 0.2 | 0.4 | 0.5 | 0.7 | 1.1 |
| Non-Hispanic white, both sexes..... | 1.4 | * | * | * | 0.2 | 0.3 | 0.3 | 0.4 | 0.5 | 0.7 |
| Male..... | 0.9 | * | * | * | * | 0.3 | 0.3 | 0.4 | 0.5 | 0.5 |
| Female..... | 1.8 | * | * | * | * | * | 0.4 | 0.4 | 0.6 | 0.9 |
| Non-Hispanic black, both sexes..... | 3.0 | * | 0.9 | * | 1.6 | 1.3 | 1.2 | 1.6 | 1.6 | 3.0 |
| Male..... | 2.8 | * | * | * | 1.6 | 2.2 | 1.6 | 2.3 | 1.7 | 3.5 |
| Female..... | 3.1 | * | * | * | 1.5 | * | * | * | 1.5 | 2.4 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

| | | | | | | | | | | |
|-------------------------------------|------|------|---|---|---|-----|-----|-----|-----|-----|
| All origins 3/, both sexes..... | 14.2 | 4.3 | * | * | * | 0.1 | 0.3 | 0.6 | 0.8 | 1.3 |
| Male..... | 13.9 | 5.3 | * | * | * | * | 0.3 | 0.5 | 0.8 | 1.4 |
| Female..... | 14.5 | 3.2 | * | * | * | * | 0.3 | 0.7 | 0.7 | 1.1 |
| Hispanic, both sexes..... | 4.9 | 4.2 | * | * | * | * | * | * | 0.6 | 0.7 |
| Male..... | 4.7 | 4.9 | * | * | * | * | * | * | * | * |
| Female..... | 5.1 | * | * | * | * | * | * | * | * | * |
| Mexican, both sexes..... | 4.2 | 4.0 | * | * | * | * | * | * | * | * |
| Male..... | 3.8 | * | * | * | * | * | * | * | * | * |
| Female..... | 4.8 | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | 6.9 | * | * | * | * | * | * | * | * | * |
| Male..... | 8.0 | * | * | * | * | * | * | * | * | * |
| Female..... | 5.8 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 12.0 | * | * | * | * | * | * | * | * | * |
| Male..... | 12.8 | * | * | * | * | * | * | * | * | * |
| Female..... | 11.1 | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 4.9 | * | * | * | * | * | * | * | * | * |
| Male..... | 4.9 | * | * | * | * | * | * | * | * | * |
| Female..... | 4.9 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 15.6 | 4.3 | * | * | * | 0.1 | 0.3 | 0.6 | 0.8 | 1.3 |
| Male..... | 15.4 | 5.4 | * | * | * | * | 0.3 | 0.5 | 0.9 | 1.5 |
| Female..... | 15.8 | 3.2 | * | * | * | * | 0.3 | 0.7 | 0.7 | 1.2 |
| Non-Hispanic white, both sexes..... | 15.4 | 3.3 | * | * | * | * | 0.2 | 0.4 | 0.6 | 0.9 |
| Male..... | 15.3 | 4.1 | * | * | * | * | * | 0.3 | 0.7 | 0.9 |
| Female..... | 15.6 | 2.5 | * | * | * | * | * | 0.4 | 0.5 | 0.8 |
| Non-Hispanic black, both sexes..... | 20.5 | 8.5 | * | * | * | * | 0.9 | 1.9 | 2.3 | 4.1 |
| Male..... | 19.7 | 10.6 | * | * | * | * | * | 1.7 | 2.3 | 4.5 |
| Female..... | 21.2 | 6.3 | * | * | * | * | * | 2.1 | 2.2 | 3.7 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|-------------------------------------|----------------|--------------------|--------------|--------------|----------------|----------------|----------------|----------------|----------------|----------------|
| Renal failure (N17-N19) | | | | | | | | | | |
| All origins 3/, both sexes..... | 13.9 | 4.1 | * | * | * | * | 0.3 | 0.5 | 0.7 | 1.2 |
| Male..... | 13.7 | 5.1 | * | * | * | * | 0.3 | 0.4 | 0.7 | 1.3 |
| Female..... | 14.2 | 3.0 | * | * | * | * | 0.3 | 0.6 | 0.7 | 1.1 |
| Hispanic, both sexes..... | 4.8 | 4.2 | * | * | * | * | * | * | * | 0.7 |
| Male..... | 4.6 | 4.9 | * | * | * | * | * | * | * | * |
| Female..... | 5.0 | * | * | * | * | * | * | * | * | * |
| Mexican, both sexes..... | 4.1 | 4.0 | * | * | * | * | * | * | * | * |
| Male..... | 3.7 | * | * | * | * | * | * | * | * | * |
| Female..... | 4.6 | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | 6.8 | * | * | * | * | * | * | * | * | * |
| Male..... | 8.0 | * | * | * | * | * | * | * | * | * |
| Female..... | 5.8 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 11.9 | * | * | * | * | * | * | * | * | * |
| Male..... | 12.8 | * | * | * | * | * | * | * | * | * |
| Female..... | 11.0 | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 4.7 | * | * | * | * | * | * | * | * | * |
| Male..... | 4.7 | * | * | * | * | * | * | * | * | * |
| Female..... | 4.8 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 15.3 | 4.1 | * | * | * | * | 0.3 | 0.6 | 0.7 | 1.3 |
| Male..... | 15.1 | 5.1 | * | * | * | * | 0.3 | 0.5 | 0.8 | 1.4 |
| Female..... | 15.5 | 2.9 | * | * | * | * | 0.3 | 0.6 | 0.7 | 1.1 |
| Non-Hispanic white, both sexes..... | 15.2 | 3.1 | * | * | * | * | 0.2 | 0.3 | 0.5 | 0.8 |
| Male..... | 15.0 | 3.9 | * | * | * | * | * | * | 0.6 | 0.9 |
| Female..... | 15.3 | 2.2 | * | * | * | * | * | 0.4 | 0.4 | 0.7 |
| Non-Hispanic black, both sexes..... | 20.2 | 8.0 | * | * | * | * | 0.8 | 1.8 | 2.1 | 4.0 |
| Male..... | 19.4 | 10.0 | * | * | * | * | * | * | 2.1 | 4.5 |
| Female..... | 20.9 | * | * | * | * | * | * | 2.0 | 2.1 | 3.6 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|

Pregnancy, childbirth and the puerperium (000-099)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| All origins 3/, both sexes..... | 0.1 | ... | ... | ... | * | 0.1 | 0.3 | 0.4 | 0.4 | 0.4 |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 0.3 | ... | ... | ... | * | 0.3 | 0.6 | 0.9 | 0.9 | 0.7 |
| Hispanic, both sexes..... | 0.2 | ... | ... | ... | * | * | * | * | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 0.4 | ... | ... | ... | * | * | * | * | * | * |
| Mexican, both sexes..... | 0.2 | ... | ... | ... | * | * | * | * | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 0.4 | ... | ... | ... | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | ... | ... | ... | * | * | * | * | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | * | ... | ... | ... | * | * | * | * | * | * |
| Cuban, both sexes..... | * | ... | ... | ... | * | * | * | * | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | * | ... | ... | ... | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | ... | ... | ... | * | * | * | * | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | * | ... | ... | ... | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 0.1 | ... | ... | ... | * | 0.1 | 0.3 | 0.5 | 0.4 | 0.3 |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 0.2 | ... | ... | ... | * | 0.3 | 0.6 | 0.9 | 0.8 | 0.7 |
| Non-Hispanic white, both sexes..... | 0.1 | ... | ... | ... | * | * | * | 0.3 | 0.3 | 0.2 |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 0.1 | ... | ... | ... | * | * | * | 0.5 | 0.6 | 0.3 |
| Non-Hispanic black, both sexes..... | 0.4 | ... | ... | ... | * | * | 1.0 | 1.5 | 1.0 | 1.2 |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 0.8 | ... | ... | ... | * | * | 2.0 | 2.8 | 1.9 | 2.3 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|---|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
| Pregnancy with abortive outcome (000-007) | | | | | | | | | | |
| All origins 3/, both sexes..... | 0.0 | ... | ... | ... | * | * | * | * | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 0.0 | ... | ... | ... | * | * | * | * | * | * |
| Hispanic, both sexes..... | * | ... | ... | ... | * | * | * | * | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | * | ... | ... | ... | * | * | * | * | * | * |
| Mexican, both sexes..... | * | ... | ... | ... | * | * | * | * | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | * | ... | ... | ... | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | ... | ... | ... | * | * | * | * | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | * | ... | ... | ... | * | * | * | * | * | * |
| Cuban, both sexes..... | * | ... | ... | ... | * | * | * | * | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | * | ... | ... | ... | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | ... | ... | ... | * | * | * | * | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | * | ... | ... | ... | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 0.0 | ... | ... | ... | * | * | * | * | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 0.0 | ... | ... | ... | * | * | * | * | * | * |
| Non-Hispanic white, both sexes..... | * | ... | ... | ... | * | * | * | * | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | * | ... | ... | ... | * | * | * | * | * | * |
| Non-Hispanic black, both sexes..... | * | ... | ... | ... | * | * | * | * | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | * | ... | ... | ... | * | * | * | * | * | * |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|

Other complications of pregnancy, childbirth and the puerperium (O10-O99)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| All origins 3/, both sexes..... | 0.1 | ... | ... | ... | * | 0.1 | 0.3 | 0.4 | 0.4 | 0.3 |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 0.2 | ... | ... | ... | * | 0.3 | 0.6 | 0.8 | 0.8 | 0.7 |
| Hispanic, both sexes..... | 0.2 | ... | ... | ... | * | * | * | * | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 0.4 | ... | ... | ... | * | * | * | * | * | * |
| Mexican, both sexes..... | 0.2 | ... | ... | ... | * | * | * | * | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 0.4 | ... | ... | ... | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | ... | ... | ... | * | * | * | * | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | * | ... | ... | ... | * | * | * | * | * | * |
| Cuban, both sexes..... | * | ... | ... | ... | * | * | * | * | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | * | ... | ... | ... | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | ... | ... | ... | * | * | * | * | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | * | ... | ... | ... | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 0.1 | ... | ... | ... | * | 0.1 | 0.3 | 0.4 | 0.4 | 0.3 |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 0.2 | ... | ... | ... | * | 0.3 | 0.5 | 0.9 | 0.8 | 0.6 |
| Non-Hispanic white, both sexes..... | 0.1 | ... | ... | ... | * | * | * | 0.3 | 0.3 | 0.2 |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 0.1 | ... | ... | ... | * | * | * | 0.5 | 0.5 | 0.3 |
| Non-Hispanic black, both sexes..... | 0.4 | ... | ... | ... | * | * | 0.9 | 1.4 | 0.9 | 1.1 |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 0.7 | ... | ... | ... | * | * | 1.9 | 2.7 | 1.8 | 2.0 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

| | | | | | | | | | | |
|-------------------------------------|-----|-------|-----|-----|-----|-----|-----|-----|-----|-----|
| All origins 3/, both sexes..... | 3.7 | 139.4 | 3.4 | 1.0 | 1.0 | 1.2 | 1.2 | 1.1 | 1.3 | 1.2 |
| Male..... | 3.9 | 142.6 | 3.5 | 1.0 | 1.1 | 1.4 | 1.3 | 1.4 | 1.4 | 1.2 |
| Female..... | 3.5 | 136.1 | 3.3 | 1.0 | 1.0 | 1.0 | 1.1 | 0.9 | 1.1 | 1.2 |
| Hispanic, both sexes..... | 4.4 | 150.7 | 3.5 | 1.1 | 0.8 | 1.2 | 0.8 | 0.6 | 0.8 | 0.7 |
| Male..... | 4.5 | 153.1 | 3.8 | * | * | 1.3 | * | * | 1.1 | * |
| Female..... | 4.3 | 148.2 | 3.3 | 1.2 | * | * | * | * | * | * |
| Mexican, both sexes..... | 4.8 | 148.8 | 3.6 | 1.0 | 1.1 | 1.2 | 0.9 | * | * | * |
| Male..... | 4.8 | 152.6 | 3.8 | * | * | * | * | * | * | * |
| Female..... | 4.8 | 144.7 | 3.5 | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | 3.7 | * | * | * | * | * | * | * | * | * |
| Male..... | 4.1 | * | * | * | * | * | * | * | * | * |
| Female..... | 3.4 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 2.0 | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 4.0 | 163.0 | * | * | * | * | * | * | * | * |
| Male..... | 4.1 | 164.6 | * | * | * | * | * | * | * | * |
| Female..... | 3.8 | 161.3 | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 3.6 | 134.9 | 3.3 | 1.0 | 1.1 | 1.2 | 1.3 | 1.2 | 1.3 | 1.3 |
| Male..... | 3.8 | 138.1 | 3.5 | 1.0 | 1.1 | 1.4 | 1.4 | 1.6 | 1.5 | 1.3 |
| Female..... | 3.4 | 131.5 | 3.2 | 1.0 | 1.0 | 1.0 | 1.2 | 0.9 | 1.2 | 1.3 |
| Non-Hispanic white, both sexes..... | 3.4 | 131.2 | 3.0 | 1.0 | 1.0 | 1.2 | 1.3 | 1.3 | 1.4 | 1.3 |
| Male..... | 3.6 | 137.2 | 3.2 | 0.9 | 1.0 | 1.3 | 1.4 | 1.6 | 1.6 | 1.3 |
| Female..... | 3.3 | 125.0 | 2.9 | 1.0 | 1.1 | 1.0 | 1.2 | 1.0 | 1.2 | 1.3 |
| Non-Hispanic black, both sexes..... | 4.7 | 156.9 | 4.7 | 1.2 | 1.3 | 1.7 | 1.6 | 1.4 | 1.5 | 1.4 |
| Male..... | 5.2 | 153.3 | 4.8 | 1.4 | 1.5 | 2.1 | 1.8 | 1.8 | 1.7 | * |
| Female..... | 4.4 | 160.6 | 4.7 | * | * | * | 1.4 | * | * | 1.5 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

| | | | | | | | | | | |
|-------------------------------------|------|-------|-----|-----|-----|-----|-----|-----|-----|------|
| All origins 3/, both sexes..... | 10.4 | 85.7 | 1.6 | 0.3 | 0.4 | 1.3 | 2.4 | 3.0 | 3.6 | 4.6 |
| Male..... | 10.1 | 98.5 | 1.8 | 0.4 | 0.4 | 1.6 | 3.2 | 4.1 | 4.9 | 6.1 |
| Female..... | 10.7 | 72.3 | 1.4 | 0.2 | 0.3 | 0.8 | 1.6 | 1.8 | 2.3 | 3.1 |
| Hispanic, both sexes..... | 4.0 | 56.6 | 1.3 | * | * | 1.1 | 1.6 | 1.8 | 2.3 | 2.6 |
| Male..... | 4.6 | 66.4 | 1.6 | * | * | 1.2 | 2.1 | 2.5 | 3.4 | 3.6 |
| Female..... | 3.3 | 46.4 | * | * | * | * | * | * | * | 1.4 |
| Mexican, both sexes..... | 3.3 | 53.6 | 0.9 | * | * | 1.2 | 1.2 | 1.2 | 1.6 | 1.7 |
| Male..... | 3.8 | 63.7 | * | * | * | * | 1.8 | 1.5 | 2.3 | 2.6 |
| Female..... | 2.8 | 43.1 | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | 6.8 | * | * | * | * | * | * | * | * | * |
| Male..... | 9.8 | * | * | * | * | * | * | * | * | * |
| Female..... | 4.0 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 4.9 | * | * | * | * | * | * | * | * | * |
| Male..... | 5.4 | * | * | * | * | * | * | * | * | * |
| Female..... | 4.4 | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 4.7 | 56.0 | * | * | * | * | * | 2.9 | * | 3.7 |
| Male..... | 5.2 | * | * | * | * | * | * | * | * | * |
| Female..... | 4.3 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 11.3 | 92.8 | 1.7 | 0.3 | 0.4 | 1.3 | 2.5 | 3.2 | 3.9 | 4.9 |
| Male..... | 10.9 | 106.2 | 1.9 | 0.4 | 0.4 | 1.7 | 3.4 | 4.5 | 5.1 | 6.5 |
| Female..... | 11.7 | 78.7 | 1.4 | 0.3 | 0.3 | 0.8 | 1.7 | 2.0 | 2.6 | 3.4 |
| Non-Hispanic white, both sexes..... | 11.6 | 76.0 | 1.3 | 0.2 | 0.3 | 1.2 | 2.5 | 3.1 | 3.7 | 4.7 |
| Male..... | 10.7 | 88.2 | 1.4 | * | 0.4 | 1.6 | 3.4 | 4.4 | 4.9 | 6.2 |
| Female..... | 12.5 | 63.2 | 1.3 | * | * | 0.8 | 1.5 | 1.8 | 2.5 | 3.2 |
| Non-Hispanic black, both sexes..... | 12.6 | 165.4 | 2.7 | 0.8 | 0.8 | 1.9 | 3.3 | 4.8 | 5.9 | 7.8 |
| Male..... | 14.4 | 186.3 | 3.6 | * | * | 2.5 | 3.7 | 6.0 | 8.2 | 10.6 |
| Female..... | 10.9 | 143.6 | 1.9 | * | * | * | 2.9 | 3.7 | 3.8 | 5.3 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

| | | | | | | | | | | |
|-------------------------------------|------|------|-----|-----|-----|-----|-----|------|------|------|
| All origins 3/, both sexes..... | 67.5 | 28.4 | 4.2 | 1.9 | 2.3 | 3.5 | 5.6 | 6.6 | 9.0 | 14.1 |
| Male..... | 55.8 | 33.4 | 4.6 | 2.2 | 2.5 | 4.0 | 6.3 | 7.4 | 10.6 | 16.4 |
| Female..... | 78.8 | 23.0 | 3.9 | 1.7 | 2.1 | 3.0 | 4.9 | 5.8 | 7.3 | 11.9 |
| Hispanic, both sexes..... | 21.6 | 25.1 | 3.8 | 2.0 | 2.2 | 3.1 | 4.2 | 4.6 | 6.6 | 9.7 |
| Male..... | 21.1 | 27.2 | 4.1 | 2.5 | 2.3 | 3.2 | 4.7 | 5.3 | 7.9 | 12.1 |
| Female..... | 22.1 | 22.8 | 3.5 | 1.5 | 2.0 | 3.0 | 3.5 | 3.8 | 5.1 | 7.1 |
| Mexican, both sexes..... | 18.1 | 25.2 | 4.0 | 2.2 | 2.1 | 3.5 | 4.0 | 4.4 | 6.3 | 8.9 |
| Male..... | 17.9 | 28.1 | 4.0 | 2.6 | 2.2 | 3.5 | 4.8 | 4.9 | 7.4 | 11.4 |
| Female..... | 18.2 | 22.2 | 4.0 | 1.8 | 2.0 | 3.4 | 3.1 | 3.7 | 5.0 | 5.9 |
| Puerto Rican, both sexes..... | 31.6 | * | * | * | * | * | * | * | 10.0 | 17.4 |
| Male..... | 33.3 | * | * | * | * | * | * | * | * | 23.5 |
| Female..... | 30.0 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 52.5 | * | * | * | * | * | * | * | * | * |
| Male..... | 45.7 | * | * | * | * | * | * | * | * | * |
| Female..... | 59.5 | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 23.3 | 24.0 | * | * | * | * | 4.0 | 5.3 | 6.3 | 9.4 |
| Male..... | 22.5 | * | * | * | * | * | * | 6.2 | 9.1 | 9.6 |
| Female..... | 24.2 | * | * | * | * | * | * | * | * | 9.1 |
| Non-Hispanic 5/, both sexes..... | 74.4 | 29.0 | 4.3 | 1.9 | 2.3 | 3.6 | 5.9 | 7.1 | 9.4 | 14.8 |
| Male..... | 61.2 | 34.9 | 4.6 | 2.1 | 2.5 | 4.2 | 6.6 | 7.9 | 11.2 | 17.1 |
| Female..... | 86.8 | 22.7 | 4.0 | 1.8 | 2.1 | 3.0 | 5.2 | 6.2 | 7.7 | 12.5 |
| Non-Hispanic white, both sexes..... | 80.5 | 24.7 | 3.8 | 1.8 | 2.2 | 3.5 | 5.2 | 6.3 | 8.3 | 13.2 |
| Male..... | 64.8 | 31.4 | 4.0 | 2.0 | 2.3 | 4.2 | 6.2 | 7.3 | 10.3 | 16.0 |
| Female..... | 95.5 | 17.7 | 3.6 | 1.7 | 2.0 | 2.6 | 4.2 | 5.3 | 6.3 | 10.3 |
| Non-Hispanic black, both sexes..... | 61.7 | 46.1 | 6.1 | 2.3 | 3.0 | 4.4 | 9.8 | 12.7 | 17.2 | 26.5 |
| Male..... | 56.6 | 50.6 | 6.6 | 2.3 | 3.4 | 4.3 | 9.8 | 13.0 | 18.5 | 27.2 |
| Female..... | 66.3 | 41.3 | 5.6 | 2.3 | 2.5 | 4.5 | 9.9 | 12.4 | 16.0 | 25.9 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
| Accidents (unintentional injuries) (V01-X59,Y85-Y86) | | | | | | | | | | |
| All origins 3/, both sexes..... | 37.0 | 23.5 | 10.5 | 5.9 | 7.3 | 35.0 | 40.9 | 32.3 | 30.8 | 34.9 |
| Male..... | 48.9 | 26.8 | 12.9 | 7.1 | 9.2 | 47.6 | 62.4 | 49.5 | 46.1 | 50.5 |
| Female..... | 25.6 | 20.0 | 8.1 | 4.7 | 5.3 | 21.8 | 18.4 | 14.5 | 15.2 | 19.2 |
| Hispanic, both sexes..... | 26.1 | 15.2 | 10.9 | 5.3 | 5.4 | 30.8 | 39.6 | 28.8 | 26.4 | 28.5 |
| Male..... | 38.5 | 19.5 | 13.5 | 6.3 | 6.8 | 45.8 | 60.0 | 44.5 | 41.3 | 42.9 |
| Female..... | 12.8 | 10.8 | 8.2 | 4.3 | 3.8 | 14.4 | 14.5 | 9.7 | 9.2 | 12.4 |
| Mexican, both sexes..... | 25.5 | 14.4 | 11.3 | 5.7 | 5.2 | 33.4 | 40.1 | 29.3 | 25.7 | 28.6 |
| Male..... | 37.6 | 18.7 | 14.1 | 6.7 | 6.8 | 49.4 | 60.8 | 44.9 | 38.5 | 43.2 |
| Female..... | 12.1 | 9.8 | 8.4 | 4.6 | 3.6 | 15.9 | 14.3 | 10.1 | 9.8 | 11.5 |
| Puerto Rican, both sexes..... | 25.1 | * | * | * | * | 18.9 | 29.7 | 25.8 | 27.5 | 29.3 |
| Male..... | 37.8 | * | * | * | * | 30.9 | 47.1 | 43.3 | 49.4 | 44.1 |
| Female..... | 13.5 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 29.0 | * | * | * | * | * | * | * | * | * |
| Male..... | 40.5 | * | * | * | * | * | * | * | * | * |
| Female..... | 17.2 | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 28.0 | * | 11.0 | 6.2 | 6.7 | 28.1 | 41.9 | 29.1 | 28.4 | 29.2 |
| Male..... | 41.5 | * | 12.3 | 7.6 | 8.8 | 40.4 | 61.1 | 45.2 | 48.6 | 43.8 |
| Female..... | 14.2 | * | * | * | * | 13.9 | 17.3 | 8.3 | 8.5 | 14.5 |
| Non-Hispanic 5/, both sexes..... | 38.5 | 25.5 | 10.4 | 6.0 | 7.6 | 35.7 | 41.1 | 33.0 | 31.5 | 35.7 |
| Male..... | 50.3 | 28.6 | 12.7 | 7.2 | 9.6 | 47.8 | 62.7 | 50.6 | 46.8 | 51.5 |
| Female..... | 27.3 | 22.3 | 8.1 | 4.8 | 5.6 | 23.0 | 19.2 | 15.5 | 16.3 | 20.2 |
| Non-Hispanic white, both sexes..... | 40.6 | 19.8 | 9.4 | 5.1 | 7.3 | 39.0 | 44.2 | 34.3 | 33.2 | 36.4 |
| Male..... | 52.0 | 23.3 | 11.1 | 6.1 | 9.2 | 51.6 | 67.2 | 51.9 | 48.9 | 51.9 |
| Female..... | 29.6 | 16.0 | 7.5 | 4.1 | 5.4 | 25.7 | 20.7 | 16.4 | 17.2 | 20.9 |
| Non-Hispanic black, both sexes..... | 34.0 | 50.1 | 14.6 | 9.6 | 9.2 | 24.4 | 32.7 | 34.1 | 30.9 | 39.1 |
| Male..... | 49.0 | 52.1 | 18.9 | 11.4 | 12.4 | 35.1 | 50.7 | 55.8 | 48.8 | 60.3 |
| Female..... | 20.3 | 48.0 | 10.1 | 7.7 | 5.9 | 13.3 | 15.2 | 14.4 | 14.9 | 20.3 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|-------------------------------------|----------------|--------------------|--------------|--------------|----------------|----------------|----------------|----------------|----------------|----------------|
| Transport accidents (V01-V99,Y85) | | | | | | | | | | |
| All origins 3/, both sexes..... | 16.8 | 3.1 | 4.1 | 3.5 | 4.9 | 28.4 | 29.9 | 20.5 | 17.1 | 17.1 |
| Male..... | 23.7 | 3.5 | 4.8 | 4.0 | 5.9 | 37.0 | 44.4 | 31.0 | 25.4 | 24.7 |
| Female..... | 10.1 | 2.8 | 3.4 | 3.1 | 3.9 | 19.3 | 14.6 | 9.7 | 8.7 | 9.6 |
| Hispanic, both sexes..... | 15.3 | 3.1 | 5.8 | 3.7 | 3.9 | 24.6 | 30.2 | 20.2 | 16.2 | 15.2 |
| Male..... | 22.4 | * | 7.1 | 4.2 | 5.1 | 35.4 | 44.7 | 30.3 | 24.5 | 22.4 |
| Female..... | 7.7 | * | 4.5 | 3.2 | 2.7 | 12.8 | 12.4 | 8.0 | 6.6 | 7.2 |
| Mexican, both sexes..... | 15.9 | * | 6.5 | 4.0 | 3.9 | 27.2 | 31.3 | 21.0 | 16.9 | 16.2 |
| Male..... | 23.1 | * | 8.1 | 4.5 | 5.4 | 39.1 | 46.3 | 31.0 | 24.5 | 23.9 |
| Female..... | 7.9 | * | 4.9 | 3.5 | 2.4 | 14.1 | 12.4 | 8.8 | 7.5 | 7.3 |
| Puerto Rican, both sexes..... | 9.9 | * | * | * | * | 12.8 | 21.5 | 14.4 | 9.3 | * |
| Male..... | 14.4 | * | * | * | * | 19.5 | 34.5 | 22.7 | * | * |
| Female..... | 5.7 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 14.7 | * | * | * | * | * | * | * | * | * |
| Male..... | 21.9 | * | * | * | * | * | * | * | * | * |
| Female..... | 7.4 | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 15.8 | * | 4.9 | 4.0 | 4.6 | 21.7 | 30.8 | 20.1 | 16.5 | 15.2 |
| Male..... | 23.4 | * | * | * | * | 29.7 | 43.1 | 31.5 | 27.0 | 21.7 |
| Female..... | 8.0 | * | * | * | * | 12.6 | 14.9 | * | 6.1 | 8.6 |
| Non-Hispanic 5/, both sexes..... | 16.9 | 3.2 | 3.7 | 3.5 | 5.1 | 29.0 | 29.7 | 20.5 | 17.3 | 17.3 |
| Male..... | 23.8 | 3.4 | 4.2 | 3.9 | 6.0 | 37.2 | 44.1 | 31.1 | 25.5 | 24.9 |
| Female..... | 10.4 | 2.9 | 3.1 | 3.1 | 4.1 | 20.4 | 15.0 | 10.0 | 9.1 | 9.9 |
| Non-Hispanic white, both sexes..... | 17.5 | 2.5 | 3.3 | 3.3 | 5.2 | 32.1 | 31.4 | 20.5 | 17.7 | 17.3 |
| Male..... | 24.3 | 2.9 | 3.6 | 3.7 | 6.1 | 40.8 | 46.6 | 30.7 | 25.8 | 24.6 |
| Female..... | 10.9 | 2.1 | 2.9 | 2.8 | 4.2 | 22.9 | 15.8 | 10.2 | 9.5 | 10.0 |
| Non-Hispanic black, both sexes..... | 15.7 | 5.6 | 5.0 | 4.7 | 4.7 | 18.3 | 25.1 | 23.9 | 18.4 | 19.7 |
| Male..... | 23.6 | * | 6.5 | 5.0 | 5.8 | 24.7 | 37.8 | 39.0 | 29.5 | 30.5 |
| Female..... | 8.5 | * | 3.6 | 4.3 | 3.6 | 11.6 | 12.7 | 10.2 | 8.5 | 10.2 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

| | | | | | | | | | | |
|-------------------------------------|------|-----|-----|-----|-----|------|------|------|------|------|
| All origins 3/, both sexes..... | 15.7 | 3.0 | 3.9 | 3.3 | 4.5 | 27.6 | 28.8 | 19.7 | 16.0 | 15.8 |
| Male..... | 21.9 | 3.3 | 4.5 | 3.8 | 5.3 | 36.0 | 42.6 | 29.6 | 23.6 | 22.4 |
| Female..... | 9.8 | 2.8 | 3.3 | 2.9 | 3.6 | 18.9 | 14.3 | 9.4 | 8.3 | 9.2 |
| Hispanic, both sexes..... | 14.6 | 3.1 | 5.6 | 3.5 | 3.7 | 23.9 | 29.0 | 19.6 | 15.6 | 14.5 |
| Male..... | 21.3 | * | 6.8 | 3.9 | 4.9 | 34.3 | 42.7 | 29.3 | 23.4 | 21.3 |
| Female..... | 7.4 | * | 4.4 | 3.0 | 2.4 | 12.5 | 12.2 | 7.8 | 6.6 | 7.0 |
| Mexican, both sexes..... | 15.3 | * | 6.3 | 3.8 | 3.8 | 26.6 | 30.2 | 20.5 | 16.3 | 15.3 |
| Male..... | 22.2 | * | 7.7 | 4.1 | 5.3 | 38.2 | 44.5 | 30.3 | 23.6 | 22.5 |
| Female..... | 7.6 | * | 4.8 | 3.4 | 2.3 | 13.9 | 12.3 | 8.5 | 7.4 | 6.9 |
| Puerto Rican, both sexes..... | 9.3 | * | * | * | * | 12.5 | 20.4 | 13.7 | 8.9 | * |
| Male..... | 13.7 | * | * | * | * | * | 33.1 | 21.3 | * | * |
| Female..... | 5.3 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 13.7 | * | * | * | * | * | * | * | * | * |
| Male..... | 20.1 | * | * | * | * | * | * | * | * | * |
| Female..... | 7.3 | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 14.9 | * | 4.7 | 3.7 | 4.0 | 20.5 | 29.2 | 19.1 | 15.9 | 14.9 |
| Male..... | 21.9 | * | * | * | * | 28.0 | 40.5 | 29.6 | 25.7 | 21.2 |
| Female..... | 7.8 | * | * | * | * | 11.9 | 14.6 | * | 6.1 | 8.6 |
| Non-Hispanic 5/, both sexes..... | 15.9 | 3.0 | 3.5 | 3.3 | 4.6 | 28.3 | 28.7 | 19.6 | 16.1 | 15.9 |
| Male..... | 21.9 | 3.2 | 4.0 | 3.7 | 5.4 | 36.2 | 42.4 | 29.6 | 23.5 | 22.5 |
| Female..... | 10.1 | 2.9 | 3.0 | 2.8 | 3.8 | 20.0 | 14.7 | 9.7 | 8.7 | 9.5 |
| Non-Hispanic white, both sexes..... | 16.3 | 2.4 | 3.0 | 3.1 | 4.8 | 31.3 | 30.2 | 19.5 | 16.4 | 15.8 |
| Male..... | 22.3 | 2.7 | 3.3 | 3.5 | 5.6 | 39.7 | 44.6 | 29.0 | 23.6 | 22.1 |
| Female..... | 10.5 | 2.1 | 2.8 | 2.6 | 3.9 | 22.5 | 15.5 | 9.8 | 9.0 | 9.5 |
| Non-Hispanic black, both sexes..... | 14.8 | 5.4 | 4.9 | 4.3 | 4.3 | 17.7 | 24.5 | 23.3 | 17.6 | 18.4 |
| Male..... | 22.0 | * | 6.3 | 4.8 | 5.1 | 23.9 | 36.9 | 37.7 | 28.0 | 28.2 |
| Female..... | 8.3 | * | 3.5 | 3.9 | 3.4 | 11.4 | 12.5 | 10.1 | 8.2 | 9.7 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|--------------------------------------|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
| Nontransport accidents (W00-X59,Y86) | | | | | | | | | | |
| All origins 3/, both sexes..... | 20.2 | 20.3 | 6.4 | 2.4 | 2.4 | 6.6 | 11.1 | 11.8 | 13.6 | 17.8 |
| Male..... | 25.2 | 23.3 | 8.1 | 3.1 | 3.3 | 10.6 | 18.0 | 18.4 | 20.6 | 25.9 |
| Female..... | 15.4 | 17.2 | 4.7 | 1.6 | 1.5 | 2.5 | 3.9 | 4.9 | 6.5 | 9.7 |
| Hispanic, both sexes..... | 10.8 | 12.1 | 5.1 | 1.6 | 1.4 | 6.2 | 9.4 | 8.6 | 10.2 | 13.2 |
| Male..... | 16.1 | 15.7 | 6.4 | 2.1 | 1.7 | 10.5 | 15.3 | 14.2 | 16.8 | 20.5 |
| Female..... | 5.2 | 8.3 | 3.7 | 1.2 | 1.2 | 1.6 | 2.1 | 1.7 | 2.6 | 5.2 |
| Mexican, both sexes..... | 9.6 | 11.3 | 4.8 | 1.7 | 1.3 | 6.2 | 8.9 | 8.2 | 8.8 | 12.3 |
| Male..... | 14.5 | 15.0 | 6.0 | 2.1 | * | 10.3 | 14.4 | 13.9 | 14.0 | 19.3 |
| Female..... | 4.2 | 7.5 | 3.5 | * | * | * | 1.9 | * | 2.3 | 4.2 |
| Puerto Rican, both sexes..... | 15.2 | * | * | * | * | * | 8.2 | 11.4 | 18.2 | 21.4 |
| Male..... | 23.4 | * | * | * | * | * | * | 20.6 | 33.4 | 31.6 |
| Female..... | 7.7 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 14.2 | * | * | * | * | * | * | * | * | * |
| Male..... | 18.6 | * | * | * | * | * | * | * | * | * |
| Female..... | 9.8 | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 12.2 | * | 6.1 | * | * | 6.3 | 11.1 | 8.9 | 11.9 | 14.0 |
| Male..... | 18.2 | * | * | * | * | 10.7 | 17.9 | 13.7 | 21.6 | 22.0 |
| Female..... | 6.1 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 21.6 | 22.3 | 6.7 | 2.5 | 2.6 | 6.7 | 11.4 | 12.5 | 14.2 | 18.4 |
| Male..... | 26.5 | 25.2 | 8.5 | 3.3 | 3.6 | 10.5 | 18.5 | 19.5 | 21.3 | 26.6 |
| Female..... | 16.9 | 19.4 | 4.9 | 1.7 | 1.5 | 2.7 | 4.2 | 5.5 | 7.2 | 10.3 |
| Non-Hispanic white, both sexes..... | 23.1 | 17.2 | 6.1 | 1.9 | 2.1 | 6.9 | 12.8 | 13.8 | 15.5 | 19.1 |
| Male..... | 27.8 | 20.4 | 7.6 | 2.4 | 3.0 | 10.8 | 20.6 | 21.2 | 23.1 | 27.2 |
| Female..... | 18.6 | 13.9 | 4.6 | 1.2 | 1.2 | 2.8 | 4.8 | 6.2 | 7.8 | 10.9 |
| Non-Hispanic black, both sexes..... | 18.3 | 44.5 | 9.5 | 4.9 | 4.5 | 6.1 | 7.6 | 10.2 | 12.5 | 19.4 |
| Male..... | 25.5 | 46.7 | 12.5 | 6.4 | 6.7 | 10.4 | 12.9 | 16.8 | 19.2 | 29.8 |
| Female..... | 11.8 | 42.3 | 6.5 | 3.4 | 2.4 | 1.7 | 2.5 | 4.2 | 6.4 | 10.1 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|---|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
| Accidental exposure to smoke, fire and flames (X00-X09) | | | | | | | | | | |
| All origins 3/, both sexes..... | 1.1 | 0.9 | 1.4 | 0.8 | 0.5 | 0.4 | 0.5 | 0.6 | 0.7 | 0.7 |
| Male..... | 1.4 | 1.0 | 1.6 | 0.9 | 0.6 | 0.6 | 0.7 | 0.8 | 0.9 | 0.9 |
| Female..... | 0.8 | * | 1.2 | 0.6 | 0.4 | 0.3 | 0.4 | 0.4 | 0.6 | 0.4 |
| Hispanic, both sexes..... | 0.6 | * | 1.2 | * | * | * | * | * | * | * |
| Male..... | 0.7 | * | * | * | * | * | * | * | * | * |
| Female..... | 0.4 | * | * | * | * | * | * | * | * | * |
| Mexican, both sexes..... | 0.5 | * | 1.1 | * | * | * | * | * | * | * |
| Male..... | 0.7 | * | * | * | * | * | * | * | * | * |
| Female..... | 0.4 | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | 0.9 | * | * | * | * | * | * | * | * | * |
| Male..... | 1.2 | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 0.6 | * | * | * | * | * | * | * | * | * |
| Male..... | 0.7 | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 1.2 | 1.0 | 1.5 | 0.8 | 0.5 | 0.4 | 0.6 | 0.7 | 0.8 | 0.7 |
| Male..... | 1.5 | * | 1.7 | 1.0 | 0.6 | 0.6 | 0.7 | 0.9 | 1.0 | 0.9 |
| Female..... | 0.9 | * | 1.2 | 0.7 | 0.4 | 0.3 | 0.4 | 0.4 | 0.6 | 0.5 |
| Non-Hispanic white, both sexes..... | 1.0 | * | 1.0 | 0.5 | 0.4 | 0.4 | 0.5 | 0.6 | 0.8 | 0.7 |
| Male..... | 1.3 | * | 1.1 | 0.6 | 0.5 | 0.5 | 0.7 | 0.8 | 1.0 | 0.8 |
| Female..... | 0.8 | * | 0.8 | 0.4 | * | * | 0.3 | * | 0.6 | 0.5 |
| Non-Hispanic black, both sexes..... | 2.1 | * | 3.7 | 2.1 | 0.9 | 0.7 | 0.9 | 1.2 | 1.1 | 1.0 |
| Male..... | 2.6 | * | 4.3 | 2.3 | * | * | * | * | * | 1.6 |
| Female..... | 1.6 | * | 3.0 | 1.8 | * | * | * | * | * | * |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|

Accidental poisoning and exposure to noxious substances (X40-X49)

| | | | | | | | | | | |
|-------------------------------------|------|-----|-----|---|-----|-----|------|------|------|------|
| All origins 3/, both sexes..... | 6.1 | 0.6 | 0.2 | * | 0.1 | 2.4 | 5.9 | 6.9 | 8.6 | 11.9 |
| Male..... | 8.5 | 1.0 | * | * | * | 3.5 | 9.1 | 10.4 | 12.5 | 16.5 |
| Female..... | 3.7 | * | * | * | * | 1.2 | 2.6 | 3.3 | 4.7 | 7.2 |
| Hispanic, both sexes..... | 4.3 | * | * | * | * | 1.8 | 3.8 | 4.3 | 5.7 | 8.3 |
| Male..... | 6.6 | * | * | * | * | 2.6 | 5.9 | 6.8 | 8.9 | 12.4 |
| Female..... | 1.7 | * | * | * | * | * | 1.3 | * | 2.0 | 3.7 |
| Mexican, both sexes..... | 3.5 | * | * | * | * | 1.7 | 3.5 | 3.7 | 4.1 | 6.9 |
| Male..... | 5.4 | * | * | * | * | 2.6 | 5.4 | 6.2 | 6.1 | 10.3 |
| Female..... | 1.3 | * | * | * | * | * | * | * | * | 2.8 |
| Puerto Rican, both sexes..... | 9.1 | * | * | * | * | * | * | 9.1 | 15.8 | 18.5 |
| Male..... | 14.8 | * | * | * | * | * | * | * | 28.4 | 27.9 |
| Female..... | 3.9 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 3.6 | * | * | * | * | * | * | * | * | * |
| Male..... | 5.2 | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 4.8 | * | * | * | * | * | 4.0 | 4.3 | 7.0 | 9.1 |
| Male..... | 7.5 | * | * | * | * | * | 5.9 | 5.9 | 12.5 | 13.8 |
| Female..... | 2.0 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 6.3 | 0.6 | 0.2 | * | 0.1 | 2.5 | 6.3 | 7.5 | 9.2 | 12.4 |
| Male..... | 8.8 | * | * | * | * | 3.7 | 9.8 | 11.3 | 13.1 | 17.1 |
| Female..... | 4.0 | * | * | * | * | 1.3 | 2.8 | 3.8 | 5.2 | 7.7 |
| Non-Hispanic white, both sexes..... | 6.6 | * | * | * | 0.2 | 3.0 | 7.6 | 8.8 | 10.3 | 13.1 |
| Male..... | 9.1 | * | * | * | * | 4.5 | 11.8 | 13.0 | 14.7 | 18.1 |
| Female..... | 4.3 | * | * | * | * | 1.5 | 3.4 | 4.5 | 5.8 | 8.2 |
| Non-Hispanic black, both sexes..... | 6.4 | * | * | * | * | * | 2.3 | 4.5 | 7.0 | 12.1 |
| Male..... | 9.3 | * | * | * | * | * | 3.7 | 7.2 | 10.1 | 17.3 |
| Female..... | 3.8 | * | * | * | * | * | * | 2.1 | 4.2 | 7.5 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
| Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | | | | | | | | | | |
| All origins 3/, both sexes..... | 11.0 | ... | ... | * | 1.2 | 7.4 | 12.4 | 12.8 | 12.5 | 14.3 |
| Male..... | 17.9 | ... | ... | * | 1.8 | 12.2 | 20.8 | 20.8 | 20.2 | 22.7 |
| Female..... | 4.3 | ... | ... | * | 0.6 | 2.4 | 3.5 | 4.5 | 4.7 | 6.0 |
| Hispanic, both sexes..... | 5.0 | ... | ... | * | 1.0 | 5.7 | 7.6 | 6.6 | 6.6 | 7.0 |
| Male..... | 8.3 | ... | ... | * | 1.1 | 9.1 | 11.9 | 10.8 | 10.6 | 11.3 |
| Female..... | 1.6 | ... | ... | * | * | 2.0 | 2.3 | 1.5 | 1.9 | 2.3 |
| Mexican, both sexes..... | 4.4 | ... | ... | * | 1.0 | 6.0 | 7.4 | 6.3 | 6.5 | 6.6 |
| Male..... | 7.2 | ... | ... | * | * | 9.6 | 11.6 | 10.4 | 10.2 | 10.3 |
| Female..... | 1.3 | ... | ... | * | * | 2.0 | 2.2 | * | 1.9 | 2.3 |
| Puerto Rican, both sexes..... | 5.7 | ... | ... | * | * | * | * | * | 9.6 | * |
| Male..... | 10.4 | ... | ... | * | * | * | * | * | * | * |
| Female..... | 1.4 | ... | ... | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 10.6 | ... | ... | * | * | * | * | * | * | * |
| Male..... | 17.6 | ... | ... | * | * | * | * | * | * | * |
| Female..... | 3.6 | ... | ... | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 5.8 | ... | ... | * | * | 5.2 | 8.9 | 7.5 | 4.9 | 8.2 |
| Male..... | 9.2 | ... | ... | * | * | 8.4 | 13.0 | 11.6 | 7.8 | 14.1 |
| Female..... | 2.2 | ... | ... | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 11.8 | ... | ... | * | 1.3 | 7.7 | 13.3 | 14.2 | 13.7 | 15.5 |
| Male..... | 19.4 | ... | ... | * | 1.9 | 12.7 | 22.8 | 23.4 | 22.2 | 24.6 |
| Female..... | 4.6 | ... | ... | * | 0.6 | 2.4 | 3.7 | 5.1 | 5.2 | 6.5 |
| Non-Hispanic white, both sexes..... | 13.4 | ... | ... | * | 1.3 | 8.6 | 14.5 | 15.5 | 15.3 | 17.5 |
| Male..... | 21.9 | ... | ... | * | 2.0 | 14.2 | 24.7 | 24.7 | 24.4 | 27.3 |
| Female..... | 5.3 | ... | ... | * | 0.6 | 2.7 | 4.1 | 6.1 | 6.0 | 7.6 |
| Non-Hispanic black, both sexes..... | 5.2 | ... | ... | * | 1.0 | 4.0 | 9.4 | 10.8 | 8.5 | 8.3 |
| Male..... | 9.3 | ... | ... | * | 1.6 | 6.9 | 16.6 | 20.5 | 15.6 | 14.7 |
| Female..... | 1.6 | ... | ... | * | * | * | 2.4 | 2.1 | 2.1 | 2.6 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

| | | | | | | | | | | |
|-------------------------------------|------|-----|-----|---|-----|-----|------|------|------|------|
| All origins 3/, both sexes..... | 5.9 | ... | ... | * | 0.4 | 3.6 | 6.7 | 6.4 | 5.6 | 6.6 |
| Male..... | 10.6 | ... | ... | * | 0.6 | 6.4 | 11.9 | 11.1 | 9.6 | 11.3 |
| Female..... | 1.4 | ... | ... | * | * | 0.7 | 1.2 | 1.6 | 1.7 | 2.0 |
| Hispanic, both sexes..... | 2.2 | ... | ... | * | * | 2.8 | 3.8 | 3.0 | 2.4 | 2.5 |
| Male..... | 3.8 | ... | ... | * | * | 4.9 | 6.4 | 4.9 | 4.3 | 4.3 |
| Female..... | 0.4 | ... | ... | * | * | * | * | * | * | * |
| Mexican, both sexes..... | 2.0 | ... | ... | * | * | 3.1 | 3.9 | 2.9 | 2.3 | 2.6 |
| Male..... | 3.5 | ... | ... | * | * | 5.4 | 6.7 | 4.8 | 4.1 | 4.2 |
| Female..... | 0.3 | ... | ... | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | 2.1 | ... | ... | * | * | * | * | * | * | * |
| Male..... | 4.1 | ... | ... | * | * | * | * | * | * | * |
| Female..... | * | ... | ... | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 3.6 | ... | ... | * | * | * | * | * | * | * |
| Male..... | 6.6 | ... | ... | * | * | * | * | * | * | * |
| Female..... | * | ... | ... | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 2.4 | ... | ... | * | * | * | 3.7 | 3.2 | * | 3.1 |
| Male..... | 4.2 | ... | ... | * | * | * | 5.4 | * | * | * |
| Female..... | 0.5 | ... | ... | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 6.5 | ... | ... | * | 0.4 | 3.8 | 7.3 | 7.2 | 6.3 | 7.3 |
| Male..... | 11.7 | ... | ... | * | 0.7 | 6.6 | 13.1 | 12.7 | 10.7 | 12.4 |
| Female..... | 1.6 | ... | ... | * | * | 0.8 | 1.3 | 1.8 | 1.9 | 2.2 |
| Non-Hispanic white, both sexes..... | 7.5 | ... | ... | * | 0.5 | 4.1 | 7.8 | 7.8 | 7.2 | 8.3 |
| Male..... | 13.4 | ... | ... | * | 0.8 | 7.2 | 14.0 | 13.3 | 12.0 | 14.0 |
| Female..... | 1.8 | ... | ... | * | * | 0.9 | 1.5 | 2.3 | 2.3 | 2.6 |
| Non-Hispanic black, both sexes..... | 2.9 | ... | ... | * | * | 2.5 | 6.0 | 6.5 | 4.1 | 4.1 |
| Male..... | 5.5 | ... | ... | * | * | 4.4 | 11.4 | 13.2 | 8.0 | 7.7 |
| Female..... | 0.5 | ... | ... | * | * | * | * | * | * | * |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39
 ages 1/ 1 year 2/ years years years years years years years years

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

| | | | | | | | | | | |
|-------------------------------------|------|-----|-----|---|-----|-----|------|------|------|------|
| All origins 3/, both sexes..... | 5.0 | ... | ... | * | 0.8 | 3.8 | 5.7 | 6.4 | 6.9 | 7.7 |
| Male..... | 7.3 | ... | ... | * | 1.2 | 5.8 | 8.9 | 9.7 | 10.6 | 11.4 |
| Female..... | 2.9 | ... | ... | * | 0.4 | 1.6 | 2.3 | 2.9 | 3.1 | 4.0 |
| Hispanic, both sexes..... | 2.9 | ... | ... | * | 0.7 | 2.9 | 3.8 | 3.6 | 4.2 | 4.5 |
| Male..... | 4.4 | ... | ... | * | * | 4.2 | 5.5 | 5.9 | 6.4 | 7.0 |
| Female..... | 1.2 | ... | ... | * | * | 1.5 | 1.7 | * | 1.7 | 1.7 |
| Mexican, both sexes..... | 2.4 | ... | ... | * | * | 2.9 | 3.5 | 3.4 | 4.1 | 4.0 |
| Male..... | 3.7 | ... | ... | * | * | 4.2 | 4.9 | 5.5 | 6.1 | 6.0 |
| Female..... | 1.0 | ... | ... | * | * | * | 1.8 | * | * | * |
| Puerto Rican, both sexes..... | 3.6 | ... | ... | * | * | * | * | * | * | * |
| Male..... | 6.3 | ... | ... | * | * | * | * | * | * | * |
| Female..... | 1.2 | ... | ... | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 7.1 | ... | ... | * | * | * | * | * | * | * |
| Male..... | 11.0 | ... | ... | * | * | * | * | * | * | * |
| Female..... | 3.0 | ... | ... | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 3.4 | ... | ... | * | * | * | 5.2 | 4.3 | 3.1 | 5.1 |
| Male..... | 5.0 | ... | ... | * | * | * | 7.5 | 6.9 | * | 8.2 |
| Female..... | 1.7 | ... | ... | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 5.3 | ... | ... | * | 0.8 | 3.9 | 6.1 | 7.0 | 7.4 | 8.2 |
| Male..... | 7.7 | ... | ... | * | 1.2 | 6.1 | 9.7 | 10.7 | 11.5 | 12.1 |
| Female..... | 3.1 | ... | ... | * | 0.4 | 1.6 | 2.4 | 3.3 | 3.3 | 4.4 |
| Non-Hispanic white, both sexes..... | 6.0 | ... | ... | * | 0.9 | 4.5 | 6.7 | 7.6 | 8.1 | 9.2 |
| Male..... | 8.5 | ... | ... | * | 1.2 | 7.0 | 10.7 | 11.4 | 12.4 | 13.4 |
| Female..... | 3.5 | ... | ... | * | 0.5 | 1.8 | 2.6 | 3.8 | 3.8 | 5.0 |
| Non-Hispanic black, both sexes..... | 2.4 | ... | ... | * | 0.7 | 1.5 | 3.4 | 4.4 | 4.4 | 4.1 |
| Male..... | 3.8 | ... | ... | * | 1.2 | 2.5 | 5.2 | 7.3 | 7.6 | 7.0 |
| Female..... | 1.1 | ... | ... | * | * | * | 1.6 | 1.7 | 1.4 | 1.6 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
| Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | | | | | | | | | | |
| All origins 3/, both sexes..... | 6.1 | 7.5 | 2.7 | 0.7 | 1.0 | 9.3 | 16.5 | 13.3 | 9.4 | 7.9 |
| Male..... | 9.6 | 7.9 | 2.9 | 0.6 | 1.2 | 15.3 | 27.7 | 22.0 | 14.7 | 11.8 |
| Female..... | 2.7 | 7.1 | 2.5 | 0.8 | 0.8 | 2.9 | 4.6 | 4.2 | 4.1 | 4.1 |
| Hispanic, both sexes..... | 8.1 | 6.2 | 2.6 | 0.8 | 1.0 | 14.8 | 20.2 | 14.8 | 9.8 | 8.7 |
| Male..... | 13.2 | 6.6 | 3.1 | * | 1.2 | 25.5 | 32.9 | 23.9 | 15.4 | 13.0 |
| Female..... | 2.6 | 5.9 | 2.1 | * | * | 3.1 | 4.5 | 3.6 | 3.2 | 3.8 |
| Mexican, both sexes..... | 8.5 | 6.1 | 2.4 | 0.8 | 1.1 | 16.7 | 21.5 | 15.8 | 9.9 | 8.4 |
| Male..... | 14.0 | * | 3.0 | * | * | 29.2 | 35.6 | 26.0 | 15.5 | 12.2 |
| Female..... | 2.5 | * | 1.7 | * | * | 3.1 | 3.9 | 3.3 | 2.9 | 3.9 |
| Puerto Rican, both sexes..... | 7.8 | * | * | * | * | 10.9 | 21.8 | 16.1 | 8.3 | 9.8 |
| Male..... | 13.0 | * | * | * | * | 20.2 | 37.3 | 28.2 | * | * |
| Female..... | 3.1 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 3.8 | * | * | * | * | * | * | * | * | * |
| Male..... | 5.7 | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 7.5 | * | * | * | * | 10.8 | 16.6 | 11.7 | 9.9 | 8.9 |
| Male..... | 11.9 | * | * | * | * | 17.6 | 24.8 | 17.3 | 16.4 | 13.8 |
| Female..... | 3.1 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 5.7 | 7.7 | 2.7 | 0.7 | 1.0 | 8.2 | 15.5 | 12.7 | 9.3 | 7.7 |
| Male..... | 8.9 | 8.0 | 2.9 | 0.6 | 1.2 | 13.3 | 26.2 | 21.3 | 14.3 | 11.5 |
| Female..... | 2.7 | 7.4 | 2.6 | 0.8 | 0.8 | 2.8 | 4.6 | 4.3 | 4.2 | 4.1 |
| Non-Hispanic white, both sexes..... | 2.8 | 5.0 | 1.8 | 0.5 | 0.6 | 2.8 | 4.7 | 4.1 | 3.8 | 4.2 |
| Male..... | 3.8 | 5.8 | 1.9 | 0.3 | 0.6 | 3.9 | 6.6 | 5.6 | 5.0 | 5.6 |
| Female..... | 1.9 | 4.1 | 1.8 | 0.6 | 0.6 | 1.6 | 2.7 | 2.5 | 2.7 | 2.8 |
| Non-Hispanic black, both sexes..... | 22.5 | 17.6 | 6.5 | 1.4 | 2.6 | 31.9 | 65.6 | 56.5 | 38.1 | 27.8 |
| Male..... | 39.4 | 16.4 | 7.3 | 1.5 | 3.5 | 54.8 | 119.6 | 104.5 | 67.2 | 46.6 |
| Female..... | 7.2 | 18.9 | 5.7 | 1.4 | 1.6 | 8.2 | 13.2 | 12.7 | 11.9 | 11.1 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
| Assault (homicide) by discharge of firearms (*U01.4,X93-X95) | | | | | | | | | | |
| All origins 3/, both sexes..... | 4.1 | * | 0.3 | 0.3 | 0.7 | 7.7 | 13.6 | 10.6 | 6.9 | 5.2 |
| Male..... | 7.0 | * | 0.4 | 0.3 | 0.9 | 13.2 | 23.8 | 18.6 | 11.6 | 8.4 |
| Female..... | 1.3 | * | * | 0.3 | 0.5 | 1.8 | 2.9 | 2.4 | 2.1 | 2.0 |
| Hispanic, both sexes..... | 5.6 | * | * | * | 0.7 | 12.6 | 15.9 | 11.6 | 6.8 | 5.4 |
| Male..... | 9.7 | * | * | * | * | 22.3 | 27.0 | 19.3 | 11.6 | 8.7 |
| Female..... | 1.2 | * | * | * | * | 2.0 | 2.2 | 2.2 | 1.4 | 1.7 |
| Mexican, both sexes..... | 6.1 | * | * | * | 0.9 | 14.7 | 17.2 | 12.8 | 7.2 | 5.5 |
| Male..... | 10.5 | * | * | * | * | 25.8 | 29.1 | 21.2 | 11.7 | 8.5 |
| Female..... | 1.3 | * | * | * | * | 2.4 | 2.3 | 2.5 | * | * |
| Puerto Rican, both sexes..... | 4.7 | * | * | * | * | 7.7 | 17.9 | 11.7 | * | * |
| Male..... | 9.3 | * | * | * | * | * | 33.8 | 23.4 | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 2.4 | * | * | * | * | * | * | * | * | * |
| Male..... | 3.6 | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 4.8 | * | * | * | * | 9.1 | 12.1 | 8.1 | 6.3 | 5.1 |
| Male..... | 8.2 | * | * | * | * | 15.6 | 19.6 | 12.8 | 11.7 | 8.8 |
| Female..... | 1.3 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 3.8 | * | 0.3 | 0.3 | 0.7 | 6.7 | 13.0 | 10.3 | 6.9 | 5.1 |
| Male..... | 6.5 | * | 0.3 | * | 0.9 | 11.4 | 22.9 | 18.2 | 11.5 | 8.3 |
| Female..... | 1.3 | * | * | 0.3 | 0.5 | 1.8 | 3.0 | 2.4 | 2.2 | 2.0 |
| Non-Hispanic white, both sexes..... | 1.5 | * | 0.2 | 0.2 | 0.3 | 1.9 | 3.1 | 2.7 | 2.5 | 2.5 |
| Male..... | 2.2 | * | * | * | 0.4 | 2.8 | 4.7 | 4.0 | 3.5 | 3.5 |
| Female..... | 0.9 | * | * | * | * | 1.0 | 1.5 | 1.4 | 1.4 | 1.5 |
| Non-Hispanic black, both sexes..... | 17.1 | * | * | * | 2.0 | 28.1 | 59.0 | 48.9 | 31.0 | 20.1 |
| Male..... | 31.9 | * | * | * | 2.8 | 49.8 | 109.9 | 94.2 | 58.1 | 36.9 |
| Female..... | 3.7 | * | * | * | * | 5.6 | 9.5 | 7.6 | 6.8 | 5.2 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

| | | | | | | | | | | |
|-------------------------------------|-----|------|-----|-----|-----|-----|-----|------|-----|-----|
| All origins 3/, both sexes..... | 2.0 | 7.3 | 2.4 | 0.4 | 0.3 | 1.6 | 2.9 | 2.6 | 2.5 | 2.7 |
| Male..... | 2.6 | 7.8 | 2.6 | 0.4 | 0.3 | 2.1 | 3.9 | 3.4 | 3.0 | 3.4 |
| Female..... | 1.4 | 6.8 | 2.2 | 0.5 | 0.3 | 1.1 | 1.8 | 1.8 | 2.0 | 2.1 |
| Hispanic, both sexes..... | 2.5 | 6.0 | 2.2 | * | * | 2.2 | 4.3 | 3.2 | 2.9 | 3.2 |
| Male..... | 3.5 | 6.6 | 2.6 | * | * | 3.2 | 5.9 | 4.7 | 3.9 | 4.3 |
| Female..... | 1.4 | 5.4 | 1.8 | * | * | * | 2.2 | 1.4 | 1.8 | 2.1 |
| Mexican, both sexes..... | 2.4 | 5.7 | 2.0 | * | * | 2.1 | 4.3 | 3.1 | 2.7 | 2.9 |
| Male..... | 3.4 | * | 2.5 | * | * | 3.4 | 6.5 | 4.8 | 3.8 | 3.8 |
| Female..... | 1.2 | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | 3.2 | * | * | * | * | * | * | * | * | * |
| Male..... | 3.7 | * | * | * | * | * | * | * | * | * |
| Female..... | 2.7 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 1.4 | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 2.8 | * | * | * | * | * | 4.5 | 3.6 | 3.6 | 3.8 |
| Male..... | 3.7 | * | * | * | * | * | 5.2 | * | * | * |
| Female..... | 1.8 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 1.9 | 7.5 | 2.4 | 0.4 | 0.3 | 1.5 | 2.5 | 2.4 | 2.4 | 2.6 |
| Male..... | 2.5 | 7.8 | 2.5 | 0.4 | 0.3 | 1.9 | 3.4 | 3.1 | 2.8 | 3.2 |
| Female..... | 1.4 | 7.1 | 2.3 | 0.5 | 0.3 | 1.1 | 1.7 | 1.8 | 2.0 | 2.1 |
| Non-Hispanic white, both sexes..... | 1.3 | 5.0 | 1.6 | 0.3 | 0.2 | 0.9 | 1.6 | 1.4 | 1.4 | 1.7 |
| Male..... | 1.6 | 5.8 | 1.6 | * | * | 1.1 | 2.0 | 1.6 | 1.5 | 2.1 |
| Female..... | 1.0 | 4.1 | 1.6 | 0.4 | * | 0.6 | 1.3 | 1.1 | 1.3 | 1.4 |
| Non-Hispanic black, both sexes..... | 5.4 | 16.7 | 5.9 | 1.0 | 0.6 | 3.8 | 6.6 | 7.6 | 7.0 | 7.6 |
| Male..... | 7.6 | 15.8 | 6.6 | * | * | 5.0 | 9.7 | 10.3 | 9.1 | 9.7 |
| Female..... | 3.5 | 17.7 | 5.2 | * | * | 2.6 | 3.6 | 5.1 | 5.1 | 5.9 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| All origins 3/, both sexes..... | 1.7 | 2.5 | 0.4 | 0.1 | 0.2 | 0.7 | 1.7 | 1.9 | 2.3 | 3.3 |
| Male..... | 2.2 | 2.5 | 0.4 | * | 0.2 | 1.0 | 2.4 | 2.7 | 3.1 | 4.5 |
| Female..... | 1.2 | 2.5 | 0.4 | * | * | 0.4 | 1.0 | 1.0 | 1.4 | 2.1 |
| Hispanic, both sexes..... | 0.8 | * | * | * | * | * | 1.1 | 0.8 | 1.2 | 1.7 |
| Male..... | 1.2 | * | * | * | * | * | 1.7 | 1.1 | 1.8 | 2.4 |
| Female..... | 0.4 | * | * | * | * | * | * | * | * | * |
| Mexican, both sexes..... | 0.7 | * | * | * | * | * | 0.9 | * | 0.9 | 1.2 |
| Male..... | 1.0 | * | * | * | * | * | 1.5 | * | * | * |
| Female..... | 0.3 | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | 1.8 | * | * | * | * | * | * | * | * | * |
| Male..... | 3.1 | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 1.0 | * | * | * | * | * | * | * | * | * |
| Male..... | 1.4 | * | * | * | * | * | * | * | * | * |
| Female..... | 0.6 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 1.8 | 2.7 | 0.4 | 0.1 | 0.2 | 0.7 | 1.8 | 2.1 | 2.5 | 3.6 |
| Male..... | 2.3 | 2.8 | 0.3 | * | 0.3 | 1.1 | 2.6 | 3.0 | 3.4 | 4.9 |
| Female..... | 1.3 | 2.6 | 0.4 | * | * | 0.4 | 1.1 | 1.2 | 1.6 | 2.3 |
| Non-Hispanic white, both sexes..... | 1.8 | 1.8 | 0.4 | * | 0.2 | 0.8 | 2.1 | 2.3 | 2.6 | 3.7 |
| Male..... | 2.3 | 1.9 | * | * | 0.3 | 1.1 | 2.9 | 3.2 | 3.5 | 4.9 |
| Female..... | 1.3 | 1.8 | * | * | * | 0.5 | 1.2 | 1.3 | 1.7 | 2.5 |
| Non-Hispanic black, both sexes..... | 2.0 | 5.7 | * | * | * | * | 1.3 | 2.0 | 2.5 | 4.1 |
| Male..... | 2.9 | * | * | * | * | * | 1.9 | 3.2 | 3.7 | 6.7 |
| Female..... | 1.2 | * | * | * | * | * | * | * | 1.5 | 1.9 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
| Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9) | | | | | | | | | | |
| All origins 3/, both sexes..... | 1.6 | 2.5 | 0.4 | 0.1 | 0.2 | 0.5 | 1.5 | 1.7 | 2.2 | 3.2 |
| Male..... | 2.1 | 2.5 | 0.3 | * | 0.2 | 0.7 | 2.1 | 2.4 | 3.0 | 4.3 |
| Female..... | 1.1 | 2.5 | 0.4 | * | * | 0.3 | 0.9 | 1.0 | 1.4 | 2.0 |
| Hispanic, both sexes..... | 0.8 | * | * | * | * | * | 0.8 | 0.7 | 1.2 | 1.6 |
| Male..... | 1.1 | * | * | * | * | * | 1.3 | * | 1.7 | 2.3 |
| Female..... | 0.4 | * | * | * | * | * | * | * | * | * |
| Mexican, both sexes..... | 0.6 | * | * | * | * | * | * | * | * | 1.1 |
| Male..... | 0.8 | * | * | * | * | * | * | * | * | * |
| Female..... | 0.3 | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | 1.8 | * | * | * | * | * | * | * | * | * |
| Male..... | 3.1 | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 0.9 | * | * | * | * | * | * | * | * | * |
| Male..... | 1.3 | * | * | * | * | * | * | * | * | * |
| Female..... | 0.6 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 1.7 | 2.7 | 0.3 | 0.1 | 0.2 | 0.6 | 1.6 | 2.0 | 2.4 | 3.4 |
| Male..... | 2.2 | 2.8 | 0.3 | * | 0.3 | 0.8 | 2.2 | 2.8 | 3.2 | 4.6 |
| Female..... | 1.2 | 2.6 | 0.4 | * | * | 0.4 | 1.0 | 1.1 | 1.5 | 2.2 |
| Non-Hispanic white, both sexes..... | 1.7 | 1.8 | 0.4 | * | 0.2 | 0.7 | 1.9 | 2.2 | 2.5 | 3.6 |
| Male..... | 2.2 | 1.9 | * | * | * | 0.9 | 2.6 | 3.0 | 3.3 | 4.7 |
| Female..... | 1.3 | 1.8 | * | * | * | 0.5 | 1.1 | 1.3 | 1.7 | 2.5 |
| Non-Hispanic black, both sexes..... | 1.9 | 5.7 | * | * | * | * | 1.1 | 1.8 | 2.4 | 4.0 |
| Male..... | 2.7 | * | * | * | * | * | 1.4 | 2.7 | 3.3 | 6.4 |
| Female..... | 1.2 | * | * | * | * | * | * | * | 1.5 | 1.8 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|

Drug-induced causes (F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,X40-X44,X60-X64,X85,Y10-Y14) 6/

| | | | | | | | | | | |
|-------------------------------------|------|-----|-----|---|-----|-----|------|------|------|------|
| All origins 3/, both sexes..... | 9.0 | 0.7 | 0.2 | * | 0.2 | 3.1 | 7.9 | 9.8 | 12.4 | 17.5 |
| Male..... | 11.8 | 1.0 | * | * | * | 4.3 | 11.4 | 14.0 | 17.0 | 23.1 |
| Female..... | 6.3 | * | 0.3 | * | * | 1.8 | 4.2 | 5.4 | 7.8 | 11.9 |
| Hispanic, both sexes..... | 5.5 | * | * | * | * | 1.8 | 4.4 | 5.0 | 7.5 | 11.3 |
| Male..... | 8.2 | * | * | * | * | 2.6 | 6.4 | 7.8 | 11.0 | 16.4 |
| Female..... | 2.6 | * | * | * | * | 1.9 | 1.7 | 3.4 | 5.8 | |
| Mexican, both sexes..... | 4.4 | * | * | * | * | 1.8 | 3.8 | 4.1 | 5.8 | 9.3 |
| Male..... | 6.6 | * | * | * | * | 2.6 | 5.6 | 6.6 | 8.0 | 13.5 |
| Female..... | 1.9 | * | * | * | * | * | * | * | 3.1 | 4.4 |
| Puerto Rican, both sexes..... | 12.8 | * | * | * | * | * | * | 11.4 | 21.3 | 29.0 |
| Male..... | 20.8 | * | * | * | * | * | * | 20.6 | 36.3 | 44.8 |
| Female..... | 5.4 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 5.4 | * | * | * | * | * | * | * | * | * |
| Male..... | 7.5 | * | * | * | * | * | * | * | * | * |
| Female..... | 3.3 | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 6.0 | * | * | * | * | * | 5.0 | 5.3 | 7.4 | 11.3 |
| Male..... | 8.6 | * | * | * | * | * | 6.6 | 6.9 | 12.2 | 15.8 |
| Female..... | 3.3 | * | * | * | * | * | * | * | * | 6.8 |
| Non-Hispanic 5/, both sexes..... | 9.5 | 0.8 | 0.2 | * | 0.2 | 3.3 | 8.6 | 10.9 | 13.3 | 18.4 |
| Male..... | 12.3 | * | * | * | * | 4.6 | 12.5 | 15.6 | 18.2 | 24.2 |
| Female..... | 6.9 | * | * | * | * | 1.9 | 4.6 | 6.2 | 8.6 | 12.8 |
| Non-Hispanic white, both sexes..... | 10.0 | * | * | * | 0.2 | 3.9 | 10.4 | 12.6 | 14.9 | 19.7 |
| Male..... | 12.8 | * | * | * | * | 5.5 | 15.0 | 17.9 | 20.3 | 25.6 |
| Female..... | 7.4 | * | * | * | * | 2.3 | 5.6 | 7.3 | 9.4 | 13.8 |
| Non-Hispanic black, both sexes..... | 9.4 | * | * | * | * | 1.0 | 3.3 | 6.8 | 10.6 | 17.7 |
| Male..... | 13.2 | * | * | * | * | 1.4 | 4.8 | 10.1 | 14.3 | 24.7 |
| Female..... | 6.0 | * | * | * | * | * | 1.9 | 3.8 | 7.2 | 11.4 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex

| | All ages 1/ | Under 1 year 2/ | 1-4 years | 5-9 years | 10-14 years | 15-19 years | 20-24 years | 25-29 years | 30-34 years | 35-39 years |
|--|-------------|-----------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
| Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 6/ | | | | | | | | | | |
| All origins 3/, both sexes..... | 10.5 | * | 0.4 | 0.4 | 1.3 | 12.1 | 21.3 | 17.8 | 13.2 | 12.4 |
| Male..... | 18.4 | * | 0.5 | 0.4 | 1.9 | 21.1 | 37.6 | 31.1 | 22.3 | 20.6 |
| Female..... | 2.8 | * | 0.3 | 0.3 | 0.7 | 2.7 | 4.2 | 4.1 | 3.9 | 4.2 |
| Hispanic, both sexes..... | 8.1 | * | * | * | 1.1 | 16.1 | 20.7 | 15.2 | 9.7 | 8.5 |
| Male..... | 14.2 | * | * | * | 1.4 | 28.5 | 35.1 | 25.4 | 16.7 | 13.9 |
| Female..... | 1.6 | * | * | * | * | 2.6 | 2.9 | 2.8 | 1.7 | 2.4 |
| Mexican, both sexes..... | 8.6 | * | * | * | 1.2 | 18.7 | 22.3 | 16.3 | 10.1 | 8.5 |
| Male..... | 14.8 | * | * | * | 1.6 | 32.8 | 37.7 | 27.3 | 16.7 | 13.5 |
| Female..... | 1.7 | * | * | * | * | 3.2 | 2.9 | 2.9 | 1.8 | 2.7 |
| Puerto Rican, both sexes..... | 7.0 | * | * | * | * | 9.0 | 21.5 | 15.4 | 10.7 | * |
| Male..... | 13.9 | * | * | * | * | * | 40.8 | 29.6 | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 6.0 | * | * | * | * | * | * | * | * | * |
| Male..... | 10.2 | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 7.5 | * | * | * | * | 12.3 | 16.3 | 12.1 | 8.4 | 8.9 |
| Male..... | 13.0 | * | * | * | * | 21.3 | 25.9 | 18.9 | 15.6 | 15.8 |
| Female..... | 1.8 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 10.8 | * | 0.4 | 0.3 | 1.3 | 11.3 | 21.3 | 18.3 | 13.8 | 13.0 |
| Male..... | 19.0 | * | 0.5 | 0.3 | 2.0 | 19.5 | 37.9 | 32.4 | 23.4 | 21.7 |
| Female..... | 3.0 | * | * | 0.4 | 0.7 | 2.7 | 4.5 | 4.4 | 4.3 | 4.4 |
| Non-Hispanic white, both sexes..... | 9.4 | * | 0.3 | 0.2 | 1.0 | 6.7 | 11.8 | 11.2 | 10.2 | 11.3 |
| Male..... | 16.3 | * | * | * | 1.5 | 11.2 | 20.2 | 18.3 | 16.6 | 18.3 |
| Female..... | 2.8 | * | * | * | 0.5 | 1.9 | 3.2 | 3.8 | 3.8 | 4.2 |
| Non-Hispanic black, both sexes..... | 20.7 | * | * | * | 2.5 | 32.3 | 67.0 | 57.1 | 36.3 | 25.2 |
| Male..... | 38.9 | * | * | * | 3.7 | 57.5 | 125.0 | 111.0 | 68.1 | 46.3 |
| Female..... | 4.3 | * | * | * | 1.3 | 6.2 | 10.6 | 8.0 | 7.6 | 6.5 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

All causes

| | | | | | | | | | | |
|-------------------------------------|-------|-------|---------|---------|---------|---------|---------|---------|----------|----------|
| All origins 3/, both sexes..... | 242.6 | 357.1 | 512.8 | 770.1 | 1,187.7 | 1,832.7 | 2,845.9 | 4,449.1 | 7,103.8 | 14,828.3 |
| Male..... | 306.4 | 454.7 | 653.5 | 961.8 | 1,476.0 | 2,265.9 | 3,533.0 | 5,522.7 | 8,652.3 | 16,254.5 |
| Female..... | 179.7 | 262.4 | 378.2 | 590.0 | 926.0 | 1,458.7 | 2,296.7 | 3,690.9 | 6,162.2 | 14,209.6 |
| Hispanic, both sexes..... | 181.8 | 266.6 | 393.2 | 583.0 | 905.8 | 1,378.9 | 2,179.7 | 3,411.7 | 5,365.5 | 10,679.0 |
| Male..... | 242.6 | 353.7 | 516.8 | 779.4 | 1,158.2 | 1,765.5 | 2,747.9 | 4,258.0 | 6,447.7 | 11,674.1 |
| Female..... | 117.3 | 178.1 | 273.2 | 402.6 | 685.7 | 1,063.3 | 1,742.1 | 2,807.4 | 4,674.8 | 10,186.0 |
| Mexican, both sexes..... | 168.7 | 252.3 | 379.2 | 543.5 | 890.4 | 1,439.7 | 2,411.7 | 3,216.1 | 5,386.7 | 10,048.7 |
| Male..... | 218.2 | 330.0 | 475.4 | 697.8 | 1,133.0 | 1,709.5 | 3,192.2 | 4,294.5 | 6,010.3 | * |
| Female..... | 110.9 | 171.6 | 275.7 | 393.5 | 682.7 | 1,191.4 | 1,865.8 | 2,507.2 | 4,912.7 | 9,528.4 |
| Puerto Rican, both sexes..... | 313.6 | 430.7 | 547.7 | 880.8 | 1,107.8 | 1,772.4 | 2,367.2 | * | * | * |
| Male..... | 445.8 | 565.5 | 769.0 | 1,307.5 | * | * | * | * | * | * |
| Female..... | 195.6 | 289.7 | 348.4 | 537.8 | 792.4 | * | * | * | * | * |
| Cuban, both sexes..... | 178.2 | 288.9 | 437.4 | 535.9 | 654.4 | 891.3 | 1,552.8 | 3,165.3 | * | * |
| Male..... | 224.2 | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | 551.4 | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 165.0 | 231.9 | 347.4 | 546.2 | 961.6 | 1,366.0 | 1,996.0 | 3,407.1 | * | * |
| Male..... | 237.2 | 316.5 | 495.6 | 703.3 | 1,136.9 | 1,980.2 | 2,249.8 | * | * | * |
| Female..... | 101.0 | 152.4 | 231.9 | 398.3 | 790.8 | 980.8 | 1,774.8 | * | * | * |
| Non-Hispanic 5/, both sexes..... | 249.2 | 365.0 | 521.3 | 782.3 | 1,204.9 | 1,859.4 | 2,879.6 | 4,490.9 | 7,161.0 | 14,958.7 |
| Male..... | 313.3 | 463.0 | 662.5 | 972.1 | 1,493.6 | 2,292.2 | 3,569.8 | 5,572.1 | 8,727.3 | 16,418.0 |
| Female..... | 186.7 | 270.3 | 386.3 | 603.7 | 942.0 | 1,484.2 | 2,326.6 | 3,727.8 | 6,210.5 | 14,329.1 |
| Non-Hispanic white, both sexes..... | 228.1 | 331.7 | 475.5 | 731.3 | 1,145.4 | 1,800.0 | 2,825.5 | 4,446.2 | 7,166.0 | 15,149.1 |
| Male..... | 290.9 | 424.3 | 604.5 | 906.2 | 1,413.9 | 2,216.9 | 3,500.6 | 5,514.6 | 8,733.5 | 16,641.9 |
| Female..... | 165.4 | 240.1 | 349.1 | 563.3 | 896.1 | 1,430.8 | 2,273.2 | 3,681.6 | 6,207.1 | 14,504.3 |
| Non-Hispanic black, both sexes..... | 422.8 | 648.6 | 943.7 | 1,346.7 | 1,911.2 | 2,670.7 | 3,877.8 | 5,630.1 | 8,036.6 | 14,386.7 |
| Male..... | 514.5 | 818.5 | 1,229.3 | 1,751.3 | 2,503.4 | 3,402.3 | 4,983.1 | 7,157.7 | 10,054.7 | 15,776.9 |
| Female..... | 342.0 | 501.4 | 702.1 | 1,014.9 | 1,448.4 | 2,128.3 | 3,136.6 | 4,718.0 | 6,986.6 | 13,855.4 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

40-44 45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years and over years years years years years years years years years and over

Shigellosis and amebiasis (A03,A06)

Table with 11 columns representing age groups (40-44 to 85 years and over) and rows for various demographic categories including All origins, Hispanic, Mexican, Puerto Rican, Cuban, Other Hispanic, Non-Hispanic white, and Non-Hispanic black, each split by sex. All cells contain an asterisk (*).

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Septicemia (A40-A41)

| | | | | | | | | | | |
|-------------------------------------|-----|------|------|------|------|------|-------|-------|-------|-------|
| All origins 3/, both sexes..... | 2.3 | 4.2 | 6.2 | 10.0 | 16.0 | 27.0 | 43.1 | 70.0 | 109.4 | 203.0 |
| Male..... | 2.5 | 4.7 | 6.8 | 10.6 | 17.8 | 29.1 | 48.2 | 80.6 | 127.0 | 213.9 |
| Female..... | 2.2 | 3.8 | 5.6 | 9.5 | 14.5 | 25.2 | 39.0 | 62.5 | 98.7 | 198.3 |
| Hispanic, both sexes..... | 1.5 | 2.7 | 5.1 | 6.2 | 11.2 | 20.1 | 39.9 | 54.9 | 78.4 | 156.3 |
| Male..... | 2.2 | 3.8 | 6.9 | 6.4 | 12.6 | 20.7 | 43.9 | 61.4 | 91.5 | 164.0 |
| Female..... | * | * | 3.2 | 6.1 | 10.1 | 19.6 | 36.8 | 50.3 | 70.1 | 152.5 |
| Mexican, both sexes..... | 1.4 | 2.5 | 4.9 | 6.2 | 9.5 | 19.5 | 40.6 | 43.4 | 70.5 | 152.3 |
| Male..... | * | 3.6 | 6.1 | * | * | 18.8 | 46.1 | 55.3 | * | * |
| Female..... | * | * | * | * | * | 20.1 | 36.7 | 35.6 | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | 31.9 | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 2.4 | 4.4 | 6.3 | 10.3 | 16.4 | 27.4 | 43.2 | 70.6 | 110.5 | 204.2 |
| Male..... | 2.5 | 4.8 | 6.8 | 10.8 | 18.1 | 29.5 | 48.3 | 81.2 | 128.1 | 215.6 |
| Female..... | 2.4 | 4.0 | 5.8 | 9.7 | 14.8 | 25.5 | 39.1 | 63.0 | 99.7 | 199.3 |
| Non-Hispanic white, both sexes..... | 2.0 | 3.6 | 4.9 | 8.5 | 13.6 | 23.7 | 38.9 | 64.6 | 104.0 | 192.6 |
| Male..... | 2.0 | 4.0 | 5.2 | 9.1 | 15.4 | 25.2 | 43.8 | 73.6 | 119.9 | 203.9 |
| Female..... | 2.0 | 3.3 | 4.6 | 7.9 | 11.9 | 22.3 | 34.9 | 58.2 | 94.2 | 187.7 |
| Non-Hispanic black, both sexes..... | 5.6 | 9.6 | 17.4 | 27.6 | 41.8 | 63.3 | 92.0 | 148.3 | 210.3 | 370.8 |
| Male..... | 6.1 | 10.5 | 20.6 | 29.3 | 46.2 | 73.2 | 106.7 | 185.4 | 260.5 | 399.3 |
| Female..... | 5.2 | 8.9 | 14.8 | 26.3 | 38.3 | 56.0 | 82.1 | 126.1 | 184.2 | 359.9 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

40-44 45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years and over years years years years years years years years years and over

Arthropod-borne viral encephalitis (A83-A84,A85.2)

Table with 11 columns representing age groups (40-44 to 85 years and over) and 18 rows representing different demographic categories (All origins, Hispanic, Mexican, Puerto Rican, Cuban, Other Hispanic, Non-Hispanic white, Non-Hispanic black) split by sex. All cells contain an asterisk (*).

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Viral hepatitis (B15-B19)

| | | | | | | | | | | |
|-------------------------------------|-----|------|------|------|------|------|-----|------|-----|-----|
| All origins 3/, both sexes..... | 2.3 | 5.6 | 6.1 | 4.3 | 3.7 | 3.9 | 4.5 | 4.7 | 4.4 | 3.9 |
| Male..... | 3.2 | 8.5 | 9.5 | 6.2 | 4.6 | 4.3 | 4.7 | 5.7 | 4.8 | 5.0 |
| Female..... | 1.4 | 2.8 | 2.8 | 2.5 | 2.9 | 3.4 | 4.4 | 4.1 | 4.2 | 3.3 |
| Hispanic, both sexes..... | 2.7 | 6.8 | 9.4 | 8.1 | 8.6 | 8.8 | 8.4 | 11.5 | 8.9 | * |
| Male..... | 3.9 | 10.8 | 13.5 | 11.9 | 11.0 | 7.4 | * | 12.8 | * | * |
| Female..... | * | 2.8 | 5.5 | 4.6 | 6.5 | 10.0 | 8.8 | 10.5 | * | * |
| Mexican, both sexes..... | 2.1 | 5.1 | 8.2 | 5.6 | 6.5 | 8.5 | * | * | * | * |
| Male..... | 2.9 | 7.9 | 11.8 | 7.5 | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | 17.9 | 23.3 | 27.8 | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | 6.4 | 7.1 | * | * | * | * | * | * | * |
| Male..... | * | 11.3 | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 2.2 | 5.4 | 5.7 | 3.9 | 3.3 | 3.5 | 4.2 | 4.4 | 4.2 | 3.7 |
| Male..... | 3.1 | 8.1 | 9.1 | 5.6 | 4.1 | 4.1 | 4.5 | 5.3 | 4.7 | 4.8 |
| Female..... | 1.4 | 2.8 | 2.6 | 2.3 | 2.6 | 2.9 | 4.1 | 3.8 | 3.9 | 3.2 |
| Non-Hispanic white, both sexes..... | 2.1 | 5.2 | 5.4 | 3.3 | 2.7 | 2.9 | 3.8 | 4.1 | 4.0 | 3.5 |
| Male..... | 3.0 | 7.9 | 8.5 | 4.7 | 3.3 | 3.7 | 4.1 | 5.1 | 4.7 | 4.6 |
| Female..... | 1.3 | 2.6 | 2.4 | 2.0 | 2.2 | 2.2 | 3.5 | 3.4 | 3.6 | 3.1 |
| Non-Hispanic black, both sexes..... | 2.9 | 7.3 | 9.2 | 7.8 | 6.3 | 6.1 | 5.8 | 6.0 | * | * |
| Male..... | 3.9 | 10.6 | 15.1 | 12.6 | 8.1 | 7.9 | * | * | * | * |
| Female..... | 1.9 | 4.5 | 4.3 | 3.9 | 4.9 | 4.7 | 5.7 | 5.6 | * | * |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Human immunodeficiency virus (HIV) disease (B20-B24)

| | | | | | | | | | | |
|-------------------------------------|-------|-------|------|------|------|------|------|------|-----|---|
| All origins 3/, both sexes..... | 13.4 | 12.6 | 9.6 | 5.9 | 4.0 | 2.9 | 1.4 | 1.0 | 0.5 | * |
| Male..... | 19.8 | 19.4 | 15.7 | 9.8 | 6.8 | 5.0 | 2.6 | 1.7 | * | * |
| Female..... | 7.2 | 5.9 | 3.6 | 2.3 | 1.5 | 1.0 | 0.5 | * | * | * |
| Hispanic, both sexes..... | 16.1 | 15.4 | 13.7 | 10.8 | 7.8 | 4.3 | * | * | * | * |
| Male..... | 23.9 | 24.0 | 21.8 | 17.4 | 11.5 | 8.5 | * | * | * | * |
| Female..... | 7.8 | 6.7 | 5.9 | 4.7 | 4.5 | * | * | * | * | * |
| Mexican, both sexes..... | 8.4 | 7.6 | 6.5 | 5.5 | * | * | * | * | * | * |
| Male..... | 13.3 | 12.6 | 10.3 | 9.3 | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | 73.8 | 73.5 | 61.1 | 44.3 | * | * | * | * | * | * |
| Male..... | 105.8 | 101.2 | 96.1 | * | * | * | * | * | * | * |
| Female..... | 45.3 | 44.5 | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 13.2 | 10.8 | 8.4 | * | * | * | * | * | * | * |
| Male..... | 22.5 | 19.1 | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 12.9 | 12.1 | 9.0 | 5.5 | 3.6 | 2.7 | 1.4 | 0.9 | 0.4 | * |
| Male..... | 19.0 | 18.6 | 14.9 | 9.1 | 6.3 | 4.8 | 2.4 | 1.7 | * | * |
| Female..... | 7.0 | 5.8 | 3.4 | 2.0 | 1.2 | 1.0 | 0.6 | * | * | * |
| Non-Hispanic white, both sexes..... | 5.8 | 5.5 | 3.5 | 2.6 | 1.7 | 1.0 | 0.5 | 0.4 | * | * |
| Male..... | 9.9 | 9.5 | 6.3 | 4.7 | 3.2 | 1.9 | 1.0 | * | * | * |
| Female..... | 1.7 | 1.6 | 0.8 | 0.6 | * | * | * | * | * | * |
| Non-Hispanic black, both sexes..... | 58.6 | 58.4 | 50.5 | 30.8 | 20.5 | 17.8 | 10.6 | 7.2 | * | * |
| Male..... | 81.3 | 87.2 | 84.9 | 50.9 | 36.7 | 32.0 | 19.5 | 15.0 | * | * |
| Female..... | 38.6 | 33.4 | 21.4 | 14.2 | 8.0 | 7.2 | 4.6 | * | * | * |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|-----|-----|-----|------|------|------|------|
| All origins 3/, both sexes..... | 1.3 | 1.9 | 2.7 | 3.0 | 3.8 | 5.4 | 8.1 | 11.6 | 13.8 | 21.7 |
| Male..... | 1.7 | 2.6 | 3.8 | 3.8 | 4.7 | 6.5 | 9.7 | 14.0 | 17.7 | 26.8 |
| Female..... | 0.8 | 1.2 | 1.7 | 2.2 | 3.0 | 4.5 | 6.8 | 9.9 | 11.5 | 19.5 |
| Hispanic, both sexes..... | 1.7 | 2.4 | 4.9 | 4.6 | 3.9 | 6.2 | 6.5 | 8.8 | 15.6 | 17.7 |
| Male..... | 2.6 | 3.4 | 7.6 | 6.2 | * | 8.1 | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | 14.6 | * |
| Mexican, both sexes..... | 1.6 | 2.8 | 6.3 | 5.5 | * | * | * | * | * | * |
| Male..... | 2.5 | 3.9 | 8.9 | 7.5 | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 1.2 | 1.8 | 2.5 | 2.8 | 3.8 | 5.4 | 8.2 | 11.7 | 13.7 | 21.8 |
| Male..... | 1.6 | 2.6 | 3.4 | 3.6 | 4.7 | 6.4 | 9.8 | 14.1 | 17.7 | 26.8 |
| Female..... | 0.8 | 1.1 | 1.6 | 2.1 | 3.0 | 4.5 | 6.9 | 10.0 | 11.3 | 19.7 |
| Non-Hispanic white, both sexes..... | 1.0 | 1.7 | 2.2 | 2.4 | 3.5 | 5.3 | 8.0 | 11.4 | 13.7 | 22.2 |
| Male..... | 1.3 | 2.4 | 3.0 | 2.9 | 4.3 | 6.4 | 9.6 | 13.8 | 17.5 | 27.2 |
| Female..... | 0.6 | 1.0 | 1.4 | 1.9 | 2.8 | 4.3 | 6.7 | 9.8 | 11.4 | 20.0 |
| Non-Hispanic black, both sexes..... | 2.5 | 3.2 | 4.9 | 6.2 | 5.5 | 5.4 | 9.9 | 15.3 | 11.5 | 21.8 |
| Male..... | 3.1 | 4.5 | 7.0 | 9.5 | 7.5 | 6.7 | 12.3 | 18.3 | * | 28.4 |
| Female..... | 1.9 | 2.0 | 3.2 | 3.4 | 4.0 | 4.6 | 8.3 | 13.5 | 9.7 | 19.2 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Malignant neoplasms (C00-C97)

| | | | | | | | | | | |
|-------------------------------------|------|-------|-------|-------|-------|---------|---------|---------|---------|---------|
| All origins 3/, both sexes..... | 47.5 | 90.6 | 161.5 | 276.8 | 447.0 | 663.8 | 933.6 | 1,207.0 | 1,458.3 | 1,723.9 |
| Male..... | 42.8 | 89.8 | 171.7 | 308.6 | 519.7 | 796.8 | 1,157.8 | 1,554.1 | 1,951.5 | 2,491.1 |
| Female..... | 52.2 | 91.4 | 151.8 | 246.8 | 381.1 | 548.9 | 754.4 | 961.9 | 1,158.4 | 1,391.1 |
| Hispanic, both sexes..... | 31.9 | 57.4 | 105.5 | 173.2 | 271.9 | 412.8 | 600.5 | 831.8 | 1,000.4 | 1,308.8 |
| Male..... | 26.3 | 54.2 | 111.0 | 204.8 | 323.5 | 518.7 | 756.5 | 1,111.5 | 1,332.6 | 1,869.0 |
| Female..... | 37.9 | 60.7 | 100.3 | 144.2 | 227.0 | 326.2 | 480.3 | 632.0 | 788.4 | 1,031.2 |
| Mexican, both sexes..... | 30.4 | 54.5 | 99.9 | 150.4 | 260.8 | 404.4 | 628.2 | 740.6 | 977.8 | 1,229.0 |
| Male..... | 23.2 | 50.5 | 99.4 | 172.1 | 309.0 | 473.0 | 835.2 | 1,090.3 | 1,200.9 | * |
| Female..... | 38.9 | 58.7 | 100.3 | 129.2 | 219.5 | 341.4 | 483.5 | 510.7 | 808.2 | 955.0 |
| Puerto Rican, both sexes..... | 33.9 | 73.0 | 130.0 | 238.8 | 291.8 | 490.0 | 625.9 | * | * | * |
| Male..... | 39.8 | 74.7 | 157.1 | 343.7 | * | * | * | * | * | * |
| Female..... | 28.7 | 71.2 | 105.6 | 154.4 | 230.3 | * | * | * | * | * |
| Cuban, both sexes..... | 38.0 | 74.4 | 170.8 | 216.9 | 235.4 | 324.5 | 482.9 | 826.6 | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | 208.1 | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 33.6 | 55.1 | 96.5 | 186.7 | 311.3 | 455.8 | 608.4 | 928.7 | * | * |
| Male..... | 27.5 | 49.2 | 105.1 | 190.1 | 327.6 | 610.5 | 677.0 | * | * | * |
| Female..... | 39.0 | 60.6 | 89.9 | 183.6 | 295.4 | 358.8 | 548.5 | * | * | * |
| Non-Hispanic 5/, both sexes..... | 49.4 | 93.8 | 166.2 | 284.5 | 459.2 | 680.4 | 952.6 | 1,224.0 | 1,474.9 | 1,736.8 |
| Male..... | 44.9 | 93.3 | 176.8 | 316.2 | 532.8 | 814.2 | 1,180.0 | 1,573.7 | 1,974.9 | 2,513.2 |
| Female..... | 53.8 | 94.3 | 156.0 | 254.6 | 392.3 | 564.4 | 770.5 | 977.1 | 1,171.5 | 1,401.8 |
| Non-Hispanic white, both sexes..... | 47.2 | 89.0 | 158.8 | 276.0 | 451.8 | 676.5 | 950.9 | 1,222.3 | 1,473.3 | 1,737.9 |
| Male..... | 43.2 | 88.6 | 167.2 | 303.2 | 515.9 | 803.1 | 1,167.1 | 1,556.2 | 1,957.1 | 2,507.7 |
| Female..... | 51.2 | 89.5 | 150.7 | 249.8 | 392.4 | 564.4 | 774.0 | 983.3 | 1,177.4 | 1,405.4 |
| Non-Hispanic black, both sexes..... | 70.0 | 138.6 | 243.5 | 407.7 | 606.9 | 845.5 | 1,143.2 | 1,443.0 | 1,680.9 | 1,904.3 |
| Male..... | 61.9 | 140.8 | 277.8 | 494.5 | 790.8 | 1,094.5 | 1,559.5 | 2,071.8 | 2,539.6 | 3,008.4 |
| Female..... | 77.1 | 136.7 | 214.5 | 336.6 | 463.2 | 660.9 | 863.9 | 1,067.5 | 1,234.1 | 1,482.4 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
|--|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|------|------|------|------|------|------|------|------|
| All origins 3/, both sexes..... | 0.8 | 2.0 | 3.6 | 5.8 | 8.1 | 8.8 | 12.1 | 13.1 | 14.1 | 18.7 |
| Male..... | 1.2 | 3.2 | 5.8 | 9.4 | 12.9 | 13.9 | 18.1 | 19.2 | 20.3 | 26.9 |
| Female..... | 0.4 | 0.9 | 1.4 | 2.5 | 3.7 | 4.4 | 7.3 | 8.8 | 10.3 | 15.1 |
| Hispanic, both sexes..... | * | * | 2.0 | 3.3 | 3.3 | 4.6 | 6.1 | 9.1 | 8.9 | 16.1 |
| Male..... | * | * | 3.2 | 5.9 | 5.4 | 7.8 | 12.3 | 16.0 | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Mexican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 0.8 | 2.1 | 3.7 | 6.0 | 8.4 | 9.1 | 12.4 | 13.3 | 14.3 | 18.7 |
| Male..... | 1.2 | 3.4 | 6.0 | 9.7 | 13.4 | 14.4 | 18.3 | 19.3 | 20.6 | 26.7 |
| Female..... | 0.4 | 1.0 | 1.5 | 2.6 | 3.8 | 4.6 | 7.7 | 9.1 | 10.5 | 15.3 |
| Non-Hispanic white, both sexes..... | 0.7 | 1.8 | 3.2 | 5.5 | 7.8 | 8.6 | 12.7 | 13.5 | 14.4 | 18.9 |
| Male..... | 1.0 | 2.9 | 5.1 | 8.8 | 12.2 | 13.3 | 18.4 | 19.5 | 20.5 | 26.8 |
| Female..... | 0.4 | 0.8 | 1.3 | 2.3 | 3.7 | 4.5 | 8.0 | 9.1 | 10.7 | 15.5 |
| Non-Hispanic black, both sexes..... | 1.5 | 4.3 | 7.3 | 10.4 | 15.0 | 13.5 | 12.2 | 11.6 | 12.8 | 16.0 |
| Male..... | 2.4 | 7.0 | 13.2 | 18.7 | 27.5 | 24.6 | 20.9 | 16.9 | 21.8 | 26.3 |
| Female..... | * | 2.0 | 2.3 | 3.5 | 5.3 | 5.3 | 6.4 | 8.4 | 8.1 | 12.1 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|---------------------------------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Malignant neoplasm of esophagus (C15) | | | | | | | | | | |
| All origins 3/, both sexes..... | 1.0 | 2.4 | 4.7 | 8.6 | 12.7 | 18.1 | 22.5 | 25.7 | 27.8 | 24.6 |
| Male..... | 1.6 | 4.1 | 8.3 | 15.1 | 22.1 | 31.4 | 40.3 | 46.8 | 50.6 | 43.4 |
| Female..... | 0.4 | 0.7 | 1.3 | 2.4 | 4.2 | 6.7 | 8.2 | 10.9 | 14.0 | 16.5 |
| Hispanic, both sexes..... | * | 1.1 | 2.0 | 3.2 | 5.3 | 8.8 | 11.8 | 14.4 | 16.5 | 23.3 |
| Male..... | * | 2.1 | 3.5 | 6.4 | 10.3 | 15.9 | 21.5 | 26.2 | 28.6 | 36.8 |
| Female..... | * | * | * | * | * | * | * | * | * | 16.6 |
| Mexican, both sexes..... | * | * | 2.2 | * | * | 7.9 | 15.2 | 14.6 | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 1.0 | 2.5 | 5.0 | 9.0 | 13.3 | 18.8 | 23.1 | 26.3 | 28.4 | 24.5 |
| Male..... | 1.7 | 4.3 | 8.7 | 15.8 | 22.9 | 32.4 | 41.4 | 47.8 | 51.6 | 43.5 |
| Female..... | 0.4 | 0.8 | 1.4 | 2.6 | 4.5 | 6.9 | 8.4 | 11.1 | 14.2 | 16.3 |
| Non-Hispanic white, both sexes..... | 1.0 | 2.4 | 4.9 | 8.5 | 12.8 | 18.2 | 23.0 | 26.3 | 28.7 | 25.1 |
| Male..... | 1.7 | 4.3 | 8.6 | 15.0 | 22.3 | 31.8 | 41.3 | 48.3 | 52.3 | 44.7 |
| Female..... | 0.3 | 0.6 | 1.2 | 2.3 | 4.0 | 6.1 | 8.0 | 10.5 | 14.3 | 16.6 |
| Non-Hispanic black, both sexes..... | 1.7 | 3.7 | 7.0 | 15.3 | 20.8 | 28.1 | 29.8 | 31.9 | 26.7 | 20.6 |
| Male..... | 2.6 | 5.8 | 11.7 | 27.5 | 35.0 | 46.3 | 54.4 | 54.0 | 49.1 | 38.3 |
| Female..... | * | 1.9 | 3.1 | 5.3 | 9.7 | 14.6 | 13.3 | 18.8 | 15.0 | 13.8 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Malignant neoplasm of stomach (C16)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|------|------|------|------|------|------|------|-------|
| All origins 3/, both sexes..... | 1.2 | 2.1 | 3.5 | 5.6 | 8.5 | 12.3 | 18.8 | 25.6 | 33.0 | 47.6 |
| Male..... | 1.4 | 2.7 | 4.8 | 7.7 | 11.9 | 18.1 | 27.5 | 36.4 | 47.1 | 67.2 |
| Female..... | 1.0 | 1.5 | 2.3 | 3.7 | 5.4 | 7.3 | 11.9 | 18.1 | 24.4 | 39.1 |
| Hispanic, both sexes..... | 2.0 | 3.4 | 5.2 | 8.5 | 14.5 | 16.9 | 28.3 | 43.2 | 53.9 | 72.1 |
| Male..... | 2.1 | 4.0 | 5.5 | 10.8 | 17.7 | 24.0 | 44.3 | 55.7 | 72.1 | 107.1 |
| Female..... | 1.8 | 2.8 | 5.0 | 6.4 | 11.6 | 11.1 | 15.9 | 34.3 | 42.3 | 54.7 |
| Mexican, both sexes..... | 2.0 | 3.6 | 5.4 | 7.1 | 15.8 | 17.5 | 30.7 | 43.0 | 64.5 | 74.0 |
| Male..... | * | 4.3 | 5.1 | 8.7 | 20.2 | 20.7 | 49.8 | 62.2 | * | * |
| Female..... | * | * | 5.7 | * | 12.1 | * | * | 30.3 | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | 4.2 | * | 10.4 | 19.3 | * | 33.6 | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 1.1 | 1.9 | 3.4 | 5.4 | 8.0 | 11.9 | 18.2 | 24.6 | 32.0 | 46.3 |
| Male..... | 1.3 | 2.6 | 4.7 | 7.4 | 11.5 | 17.6 | 26.4 | 35.3 | 45.8 | 65.0 |
| Female..... | 0.9 | 1.3 | 2.1 | 3.5 | 4.9 | 7.0 | 11.6 | 17.1 | 23.6 | 38.3 |
| Non-Hispanic white, both sexes..... | 0.8 | 1.6 | 2.6 | 4.5 | 6.7 | 9.9 | 15.5 | 21.3 | 28.0 | 40.7 |
| Male..... | 1.0 | 2.1 | 3.7 | 6.4 | 9.6 | 14.9 | 22.7 | 31.2 | 40.7 | 58.0 |
| Female..... | 0.6 | 1.0 | 1.5 | 2.7 | 3.9 | 5.4 | 9.6 | 14.2 | 20.2 | 33.3 |
| Non-Hispanic black, both sexes..... | 2.3 | 3.7 | 7.7 | 10.7 | 16.4 | 25.0 | 39.7 | 53.7 | 63.8 | 100.1 |
| Male..... | 2.5 | 5.2 | 11.3 | 15.7 | 24.6 | 37.1 | 60.9 | 74.6 | 91.3 | 136.7 |
| Female..... | 2.1 | 2.4 | 4.6 | 6.7 | 10.0 | 15.9 | 25.5 | 41.2 | 49.5 | 86.1 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Malignant neoplasms of colon, rectum and anus (C18-C21)

| | | | | | | | | | | |
|-------------------------------------|-----|------|------|------|------|-------|-------|-------|-------|-------|
| All origins 3/, both sexes..... | 4.3 | 8.2 | 15.0 | 24.6 | 40.7 | 61.6 | 86.2 | 118.9 | 163.2 | 241.4 |
| Male..... | 4.9 | 9.1 | 17.5 | 30.1 | 51.6 | 77.8 | 106.8 | 147.7 | 197.4 | 275.1 |
| Female..... | 3.6 | 7.4 | 12.6 | 19.5 | 30.8 | 47.6 | 69.7 | 98.6 | 142.3 | 226.8 |
| Hispanic, both sexes..... | 2.9 | 5.0 | 11.3 | 17.3 | 30.7 | 41.9 | 63.0 | 90.1 | 104.7 | 165.7 |
| Male..... | 2.9 | 6.3 | 13.4 | 24.8 | 40.0 | 56.9 | 80.3 | 118.4 | 125.8 | 174.0 |
| Female..... | 3.0 | 3.7 | 9.2 | 10.5 | 22.6 | 29.6 | 49.7 | 69.9 | 91.2 | 161.6 |
| Mexican, both sexes..... | 2.8 | 4.1 | 10.9 | 15.2 | 27.1 | 37.3 | 57.7 | 67.2 | 94.6 | 126.2 |
| Male..... | * | 5.2 | 13.2 | 23.0 | 37.6 | 48.0 | 76.5 | 109.5 | * | * |
| Female..... | 3.5 | * | 8.5 | 7.6 | 18.1 | 27.4 | 44.5 | 39.4 | * | * |
| Puerto Rican, both sexes..... | * | * | 16.7 | 24.8 | 40.4 | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | 45.5 | 56.3 | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | 5.1 | 8.4 | 18.4 | 36.8 | 49.3 | 67.3 | 103.0 | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 4.4 | 8.6 | 15.3 | 25.2 | 41.4 | 62.9 | 87.4 | 120.2 | 165.3 | 244.0 |
| Male..... | 5.2 | 9.4 | 17.8 | 30.5 | 52.3 | 79.1 | 108.1 | 148.8 | 200.3 | 279.2 |
| Female..... | 3.7 | 7.8 | 12.9 | 20.2 | 31.5 | 48.9 | 70.9 | 99.9 | 144.1 | 228.8 |
| Non-Hispanic white, both sexes..... | 4.0 | 7.7 | 14.5 | 23.7 | 39.4 | 61.1 | 85.8 | 118.0 | 162.3 | 243.2 |
| Male..... | 4.7 | 8.6 | 16.8 | 28.9 | 49.8 | 77.2 | 106.6 | 145.8 | 195.9 | 277.6 |
| Female..... | 3.3 | 6.9 | 12.2 | 18.8 | 29.8 | 46.9 | 68.7 | 98.2 | 141.8 | 228.3 |
| Non-Hispanic black, both sexes..... | 7.4 | 14.9 | 24.1 | 40.9 | 63.1 | 88.2 | 116.3 | 162.7 | 221.3 | 278.3 |
| Male..... | 8.6 | 16.1 | 28.7 | 48.6 | 81.5 | 110.8 | 139.2 | 212.2 | 288.6 | 332.6 |
| Female..... | 6.4 | 13.8 | 20.1 | 34.5 | 48.7 | 71.3 | 100.9 | 133.2 | 186.3 | 257.5 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|------|------|------|------|------|------|------|------|
| All origins 3/, both sexes..... | 1.3 | 3.4 | 6.6 | 8.1 | 11.6 | 16.9 | 22.6 | 27.6 | 31.6 | 35.0 |
| Male..... | 2.0 | 5.7 | 11.1 | 12.4 | 17.4 | 24.8 | 32.8 | 39.5 | 44.8 | 46.2 |
| Female..... | 0.6 | 1.2 | 2.4 | 4.0 | 6.4 | 10.1 | 14.4 | 19.2 | 23.6 | 30.2 |
| Hispanic, both sexes..... | 1.4 | 4.6 | 7.7 | 12.8 | 19.6 | 30.1 | 39.1 | 44.0 | 44.6 | 59.9 |
| Male..... | 1.9 | 7.1 | 13.5 | 20.9 | 28.5 | 44.0 | 55.3 | 62.1 | 56.0 | 75.3 |
| Female..... | * | 2.0 | * | 5.4 | 11.9 | 18.8 | 26.7 | 31.1 | 37.2 | 52.2 |
| Mexican, both sexes..... | 1.4 | 4.3 | 7.6 | 12.0 | 20.6 | 37.3 | 49.3 | 45.3 | 52.5 | 67.4 |
| Male..... | * | 6.6 | 12.4 | 18.6 | 28.6 | 46.2 | 76.5 | 76.1 | * | * |
| Female..... | * | * | * | * | 13.7 | 29.1 | 30.3 | 25.0 | * | * |
| Puerto Rican, both sexes..... | * | * | 14.4 | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | 4.0 | * | 14.0 | 18.1 | * | 34.5 | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 1.2 | 3.3 | 6.5 | 7.6 | 10.9 | 15.9 | 21.5 | 26.7 | 30.9 | 33.9 |
| Male..... | 2.0 | 5.5 | 10.8 | 11.6 | 16.3 | 23.3 | 31.3 | 38.1 | 44.0 | 44.8 |
| Female..... | 0.6 | 1.2 | 2.4 | 3.9 | 6.0 | 9.4 | 13.6 | 18.6 | 23.0 | 29.2 |
| Non-Hispanic white, both sexes..... | 1.0 | 2.7 | 5.2 | 6.2 | 9.7 | 14.4 | 19.7 | 25.6 | 29.8 | 32.6 |
| Male..... | 1.6 | 4.5 | 8.5 | 9.3 | 14.3 | 21.2 | 28.4 | 36.5 | 41.9 | 43.6 |
| Female..... | 0.5 | 0.9 | 2.0 | 3.2 | 5.3 | 8.4 | 12.5 | 17.8 | 22.5 | 27.8 |
| Non-Hispanic black, both sexes..... | 2.1 | 5.8 | 12.6 | 15.7 | 14.1 | 19.6 | 28.0 | 30.0 | 32.3 | 38.1 |
| Male..... | 3.4 | 9.9 | 22.3 | 25.6 | 22.8 | 32.3 | 46.2 | 43.2 | 51.5 | 41.6 |
| Female..... | * | 2.3 | 4.4 | 7.5 | 7.3 | 10.2 | 15.8 | 22.1 | 22.3 | 36.8 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--------------------------------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Malignant neoplasm of pancreas (C25) | | | | | | | | | | |
| All origins 3/, both sexes..... | 1.9 | 4.2 | 8.2 | 15.5 | 24.0 | 37.2 | 53.2 | 70.2 | 81.8 | 94.5 |
| Male..... | 2.4 | 5.3 | 10.6 | 19.2 | 29.3 | 43.3 | 60.8 | 78.9 | 91.9 | 100.9 |
| Female..... | 1.4 | 3.1 | 6.0 | 12.0 | 19.2 | 31.9 | 47.1 | 64.1 | 75.6 | 91.7 |
| Hispanic, both sexes..... | 1.3 | 2.8 | 5.6 | 11.2 | 17.4 | 26.8 | 44.7 | 53.9 | 58.4 | 79.3 |
| Male..... | 1.5 | 3.7 | 7.3 | 13.0 | 19.5 | 32.8 | 51.3 | 54.4 | 58.3 | 60.2 |
| Female..... | * | 1.9 | 4.0 | 9.4 | 15.7 | 21.8 | 39.5 | 53.5 | 58.4 | 88.7 |
| Mexican, both sexes..... | * | 2.6 | 4.9 | 12.7 | 17.5 | 24.8 | 49.3 | 48.9 | 54.2 | 84.8 |
| Male..... | * | * | 6.1 | 14.4 | 21.6 | 29.2 | 57.2 | 51.9 | * | * |
| Female..... | * | * | * | 11.1 | 14.1 | 20.7 | 43.8 | 47.0 | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | 6.0 | 9.6 | 17.5 | 33.8 | 56.7 | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 2.0 | 4.3 | 8.4 | 15.8 | 24.4 | 37.9 | 53.7 | 70.9 | 82.6 | 94.9 |
| Male..... | 2.5 | 5.5 | 10.8 | 19.6 | 30.0 | 44.0 | 61.3 | 80.1 | 93.2 | 102.4 |
| Female..... | 1.5 | 3.2 | 6.1 | 12.2 | 19.4 | 32.5 | 47.6 | 64.5 | 76.1 | 91.7 |
| Non-Hispanic white, both sexes..... | 2.0 | 4.2 | 7.9 | 15.3 | 23.8 | 36.7 | 52.3 | 69.8 | 82.2 | 94.6 |
| Male..... | 2.5 | 5.3 | 10.3 | 19.0 | 29.5 | 43.3 | 60.1 | 80.0 | 93.6 | 102.9 |
| Female..... | 1.4 | 3.1 | 5.6 | 11.8 | 18.4 | 30.9 | 45.9 | 62.6 | 75.2 | 91.0 |
| Non-Hispanic black, both sexes..... | 2.5 | 5.8 | 12.6 | 22.9 | 35.2 | 51.3 | 74.8 | 89.3 | 93.2 | 108.3 |
| Male..... | 3.0 | 7.4 | 16.8 | 29.6 | 42.0 | 57.6 | 85.5 | 92.0 | 106.1 | 102.8 |
| Female..... | 2.1 | 4.3 | 9.1 | 17.5 | 29.9 | 46.7 | 67.7 | 87.7 | 86.4 | 110.4 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Malignant neoplasm of larynx (C32)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|------|------|------|------|------|------|------|
| All origins 3/, both sexes..... | 0.2 | 0.7 | 1.4 | 2.7 | 4.2 | 5.3 | 6.6 | 7.9 | 7.2 | 5.5 |
| Male..... | 0.4 | 1.2 | 2.4 | 4.7 | 7.2 | 8.9 | 11.3 | 14.8 | 14.9 | 13.2 |
| Female..... | * | 0.2 | 0.5 | 0.9 | 1.5 | 2.2 | 2.9 | 3.1 | 2.5 | 2.2 |
| Hispanic, both sexes..... | * | * | * | 1.8 | 2.6 | 3.2 | 5.2 | 6.4 | * | * |
| Male..... | * | * | * | 3.9 | * | * | 10.1 | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Mexican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 0.3 | 0.8 | 1.5 | 2.8 | 4.4 | 5.4 | 6.7 | 8.0 | 7.3 | 5.4 |
| Male..... | 0.4 | 1.3 | 2.5 | 4.7 | 7.4 | 9.1 | 11.3 | 14.9 | 15.1 | 13.2 |
| Female..... | * | 0.3 | 0.5 | 1.0 | 1.6 | 2.3 | 3.0 | 3.1 | 2.5 | 2.1 |
| Non-Hispanic white, both sexes..... | 0.2 | 0.6 | 1.3 | 2.5 | 3.9 | 5.2 | 6.4 | 7.5 | 7.1 | 5.5 |
| Male..... | 0.3 | 1.1 | 2.1 | 4.1 | 6.4 | 8.5 | 10.6 | 13.8 | 14.8 | 13.3 |
| Female..... | * | * | 0.4 | 1.0 | 1.6 | 2.3 | 2.9 | 3.0 | 2.3 | 2.2 |
| Non-Hispanic black, both sexes..... | * | 1.9 | 3.7 | 6.3 | 9.6 | 9.4 | 12.1 | 15.6 | 11.2 | * |
| Male..... | * | 3.3 | 6.2 | 11.8 | 19.3 | 17.4 | 22.9 | 33.8 | 22.6 | * |
| Female..... | * | * | * | * | * | * | 4.8 | * | * | * |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

| | | | | | | | | | | |
|-------------------------------------|------|------|------|-------|-------|-------|-------|-------|-------|-------|
| All origins 3/, both sexes..... | 9.2 | 20.4 | 41.6 | 86.3 | 152.9 | 233.4 | 320.9 | 376.9 | 378.5 | 297.2 |
| Male..... | 9.6 | 23.6 | 51.4 | 107.0 | 192.7 | 297.5 | 423.0 | 515.5 | 546.8 | 482.2 |
| Female..... | 8.8 | 17.3 | 32.2 | 66.8 | 116.8 | 177.9 | 239.3 | 279.1 | 276.1 | 217.0 |
| Hispanic, both sexes..... | 2.5 | 6.0 | 15.0 | 32.0 | 53.1 | 87.4 | 139.2 | 181.6 | 187.2 | 189.0 |
| Male..... | 2.7 | 6.8 | 20.3 | 47.0 | 79.8 | 133.1 | 198.8 | 296.3 | 304.3 | 307.9 |
| Female..... | 2.3 | 5.2 | 10.0 | 18.2 | 29.7 | 50.0 | 93.3 | 99.6 | 112.4 | 130.1 |
| Mexican, both sexes..... | 2.2 | 5.1 | 11.6 | 21.2 | 47.0 | 81.0 | 134.7 | 165.0 | 179.7 | 184.9 |
| Male..... | * | 6.2 | 15.0 | 28.7 | 69.0 | 116.1 | 204.6 | 291.6 | * | * |
| Female..... | * | 3.9 | 7.9 | 14.0 | 28.1 | 48.7 | 85.7 | 81.8 | 109.0 | * |
| Puerto Rican, both sexes..... | * | * | 20.0 | 54.1 | 53.0 | 122.5 | 167.9 | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | 91.7 | 70.3 | 77.2 | 135.4 | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | 4.7 | 17.0 | 30.9 | 59.7 | 91.2 | 136.4 | 184.7 | * | * |
| Male..... | * | * | 22.7 | 38.0 | 79.5 | * | * | * | * | * |
| Female..... | * | * | 12.6 | 24.1 | * | 64.6 | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 10.0 | 21.9 | 43.9 | 90.5 | 160.2 | 243.4 | 331.9 | 386.3 | 386.0 | 301.0 |
| Male..... | 10.5 | 25.3 | 54.1 | 111.7 | 200.8 | 308.3 | 436.2 | 525.7 | 556.7 | 488.9 |
| Female..... | 9.6 | 18.5 | 34.1 | 70.7 | 123.3 | 187.2 | 248.2 | 287.9 | 282.4 | 220.0 |
| Non-Hispanic white, both sexes..... | 9.7 | 21.0 | 42.4 | 89.4 | 160.6 | 247.9 | 339.0 | 392.3 | 392.1 | 304.8 |
| Male..... | 9.8 | 23.5 | 50.8 | 107.5 | 195.6 | 307.6 | 438.5 | 525.3 | 557.9 | 493.3 |
| Female..... | 9.5 | 18.5 | 34.2 | 72.1 | 128.1 | 194.9 | 257.7 | 297.2 | 290.6 | 223.4 |
| Non-Hispanic black, both sexes..... | 14.5 | 33.3 | 66.0 | 123.3 | 197.4 | 270.6 | 336.0 | 395.6 | 370.2 | 274.9 |
| Male..... | 17.5 | 44.0 | 93.7 | 175.7 | 297.6 | 403.1 | 515.9 | 638.5 | 626.3 | 490.1 |
| Female..... | 11.9 | 24.1 | 42.5 | 80.4 | 119.2 | 172.3 | 215.4 | 250.6 | 237.0 | 192.7 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Malignant neoplasm of breast (C50)

| | | | | | | | | | | |
|-------------------------------------|------|------|------|------|------|------|-------|-------|-------|-------|
| All origins 3/, both sexes..... | 7.9 | 12.9 | 19.7 | 26.2 | 33.7 | 41.3 | 52.4 | 67.1 | 89.8 | 134.6 |
| Male..... | * | * | 0.3 | 0.4 | 0.7 | 0.8 | 1.3 | 1.9 | 2.7 | 3.4 |
| Female..... | 15.6 | 25.3 | 38.3 | 50.4 | 63.7 | 76.2 | 93.2 | 113.2 | 142.7 | 191.5 |
| Hispanic, both sexes..... | 5.4 | 8.9 | 13.7 | 15.9 | 20.0 | 26.3 | 28.6 | 36.8 | 55.3 | 71.0 |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | 11.0 | 17.9 | 26.4 | 30.5 | 37.3 | 47.6 | 50.0 | 63.1 | 89.1 | 105.3 |
| Mexican, both sexes..... | 4.5 | 8.8 | 13.3 | 14.3 | 19.7 | 24.2 | 28.5 | 27.0 | 47.3 | 58.7 |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | 9.7 | 17.7 | 27.0 | 28.2 | 36.6 | 46.4 | 47.7 | 44.7 | * | * |
| Puerto Rican, both sexes..... | * | * | 14.4 | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 7.6 | 9.5 | 13.1 | 20.0 | 21.1 | 26.5 | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | 14.4 | 18.4 | 22.8 | 38.9 | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 8.2 | 13.3 | 20.2 | 26.9 | 34.7 | 42.2 | 53.7 | 68.6 | 91.0 | 136.8 |
| Male..... | * | * | 0.3 | 0.4 | 0.7 | 0.8 | 1.3 | 2.0 | 2.8 | 3.5 |
| Female..... | 16.1 | 26.0 | 39.3 | 51.9 | 65.7 | 78.2 | 95.8 | 115.6 | 144.6 | 194.4 |
| Non-Hispanic white, both sexes..... | 7.2 | 12.0 | 18.8 | 25.7 | 34.1 | 41.8 | 53.4 | 69.1 | 90.3 | 137.7 |
| Male..... | * | * | * | 0.4 | 0.6 | 0.6 | 1.3 | 1.8 | 2.7 | 3.4 |
| Female..... | 14.3 | 23.7 | 37.0 | 50.0 | 65.2 | 78.3 | 96.0 | 117.2 | 143.9 | 195.8 |
| Non-Hispanic black, both sexes..... | 15.6 | 23.4 | 32.9 | 41.2 | 47.0 | 55.4 | 68.2 | 79.9 | 118.8 | 154.0 |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | 29.3 | 43.4 | 60.1 | 74.3 | 82.3 | 94.3 | 112.9 | 125.3 | 178.6 | 210.7 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Malignant neoplasm of cervix uteri (C53)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|-----|------|------|------|------|------|------|
| All origins 3/, both sexes..... | 1.7 | 2.1 | 2.4 | 2.7 | 2.8 | 3.3 | 3.6 | 4.1 | 4.2 | 6.2 |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 3.4 | 4.1 | 4.7 | 5.2 | 5.3 | 6.1 | 6.5 | 7.0 | 6.7 | 8.9 |
| Hispanic, both sexes..... | 1.7 | 2.0 | 2.8 | 3.5 | 3.7 | 3.7 | 5.9 | 6.4 | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 3.4 | 4.1 | 5.6 | 6.7 | 6.9 | 6.6 | 10.5 | 11.0 | * | * |
| Mexican, both sexes..... | 1.6 | 2.7 | 3.2 | 3.5 | * | * | * | * | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 3.5 | 5.4 | 6.5 | 7.0 | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 1.7 | 2.1 | 2.4 | 2.6 | 2.7 | 3.3 | 3.5 | 4.0 | 4.0 | 6.1 |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 3.4 | 4.0 | 4.7 | 5.0 | 5.2 | 6.1 | 6.2 | 6.8 | 6.5 | 8.7 |
| Non-Hispanic white, both sexes..... | 1.5 | 1.8 | 2.1 | 2.4 | 2.2 | 2.7 | 2.7 | 3.5 | 3.4 | 4.9 |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 3.0 | 3.6 | 4.2 | 4.6 | 4.3 | 5.0 | 5.0 | 5.9 | 5.4 | 7.0 |
| Non-Hispanic black, both sexes..... | 3.0 | 3.9 | 4.6 | 4.3 | 6.2 | 8.4 | 10.3 | 9.5 | 12.5 | 17.5 |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 5.7 | 7.2 | 8.4 | 7.8 | 11.0 | 14.6 | 17.2 | 15.1 | 19.1 | 24.2 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|------|------|------|------|------|------|------|
| All origins 3/, both sexes..... | 0.4 | 0.9 | 2.0 | 3.5 | 5.8 | 8.4 | 11.8 | 14.2 | 17.0 | 24.6 |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 0.9 | 1.7 | 3.9 | 6.9 | 11.0 | 15.7 | 21.2 | 24.2 | 27.4 | 35.2 |
| Hispanic, both sexes..... | * | * | 1.3 | 3.3 | 3.1 | 8.2 | 9.9 | 10.7 | 11.6 | 20.5 |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | * | * | 2.6 | 6.4 | 5.8 | 14.9 | 17.6 | 18.3 | 19.0 | 30.7 |
| Mexican, both sexes..... | * | * | * | 3.4 | * | 6.7 | 9.5 | * | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | * | * | * | 6.7 | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 0.4 | 0.9 | 2.1 | 3.5 | 5.9 | 8.4 | 11.9 | 14.3 | 17.2 | 24.6 |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 0.8 | 1.7 | 4.0 | 6.8 | 11.3 | 15.8 | 21.4 | 24.4 | 27.7 | 35.3 |
| Non-Hispanic white, both sexes..... | 0.4 | 0.8 | 1.9 | 3.1 | 5.4 | 7.1 | 10.8 | 13.2 | 16.5 | 23.8 |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 0.9 | 1.6 | 3.8 | 6.2 | 10.3 | 13.4 | 19.7 | 22.6 | 26.6 | 34.1 |
| Non-Hispanic black, both sexes..... | * | 1.1 | 3.0 | 6.8 | 11.0 | 20.4 | 25.5 | 28.1 | 28.8 | 39.6 |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | * | 2.0 | 5.5 | 12.3 | 19.6 | 35.5 | 42.7 | 44.9 | 43.8 | 54.8 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Malignant neoplasm of ovary (C56)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|------|------|------|------|------|------|------|------|
| All origins 3/, both sexes..... | 1.4 | 3.2 | 5.5 | 8.3 | 12.9 | 17.3 | 22.3 | 30.3 | 37.8 | 38.1 |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 2.8 | 6.4 | 10.7 | 16.2 | 24.6 | 32.3 | 40.1 | 51.7 | 60.8 | 54.6 |
| Hispanic, both sexes..... | 1.4 | 2.2 | 4.5 | 4.1 | 7.6 | 10.8 | 14.1 | 24.0 | 24.5 | 23.8 |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 2.9 | 4.4 | 8.8 | 7.9 | 14.3 | 19.6 | 25.0 | 41.1 | 40.1 | 35.6 |
| Mexican, both sexes..... | 1.4 | 2.4 | 4.4 | 3.4 | 6.7 | 10.2 | 15.2 | 22.4 | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 3.1 | 4.9 | 9.2 | 6.7 | 12.5 | 19.6 | 25.8 | 37.1 | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 1.4 | 3.3 | 5.6 | 8.7 | 13.3 | 17.7 | 22.8 | 30.6 | 38.4 | 38.7 |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 2.7 | 6.5 | 10.9 | 16.8 | 25.4 | 33.1 | 41.1 | 52.2 | 61.6 | 55.3 |
| Non-Hispanic white, both sexes..... | 1.4 | 3.4 | 5.9 | 9.0 | 13.9 | 18.5 | 23.5 | 31.4 | 39.3 | 39.6 |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 2.8 | 6.8 | 11.6 | 17.6 | 26.9 | 34.8 | 42.6 | 53.9 | 63.4 | 56.7 |
| Non-Hispanic black, both sexes..... | 1.2 | 2.9 | 4.2 | 7.8 | 11.1 | 15.1 | 20.3 | 26.9 | 33.1 | 35.4 |
| Male..... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... |
| Female..... | 2.3 | 5.4 | 7.8 | 14.2 | 19.7 | 26.4 | 34.0 | 42.9 | 50.3 | 48.9 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Malignant neoplasms of kidney and renal pelvis (C64-C65)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|------|------|------|------|------|------|------|
| All origins 3/, both sexes..... | 0.9 | 2.4 | 4.3 | 7.2 | 10.9 | 16.0 | 19.6 | 24.6 | 28.7 | 32.5 |
| Male..... | 1.3 | 3.4 | 6.4 | 10.6 | 16.1 | 23.0 | 27.6 | 36.6 | 39.9 | 50.1 |
| Female..... | 0.5 | 1.5 | 2.3 | 3.9 | 6.2 | 10.0 | 13.2 | 16.2 | 21.9 | 24.9 |
| Hispanic, both sexes..... | 1.1 | 2.1 | 3.8 | 6.6 | 10.2 | 11.9 | 14.1 | 23.5 | 25.8 | 26.1 |
| Male..... | 1.5 | 2.6 | 5.8 | 9.9 | 13.3 | 16.9 | 20.6 | 41.0 | 35.5 | 40.2 |
| Female..... | * | * | * | 3.5 | 7.4 | 7.7 | 9.1 | 11.0 | 19.7 | 19.1 |
| Mexican, both sexes..... | * | 2.5 | 4.7 | 7.7 | 12.1 | 13.7 | 17.1 | 21.0 | 31.8 | * |
| Male..... | * | * | 6.9 | 11.4 | 16.4 | 20.1 | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 0.9 | 2.4 | 4.3 | 7.2 | 10.9 | 16.3 | 19.9 | 24.6 | 28.8 | 32.7 |
| Male..... | 1.3 | 3.5 | 6.5 | 10.6 | 16.2 | 23.3 | 28.0 | 36.3 | 40.0 | 50.4 |
| Female..... | 0.5 | 1.5 | 2.3 | 4.0 | 6.1 | 10.1 | 13.5 | 16.4 | 22.0 | 25.1 |
| Non-Hispanic white, both sexes..... | 0.9 | 2.5 | 4.4 | 7.4 | 11.2 | 16.9 | 20.5 | 25.0 | 29.3 | 33.5 |
| Male..... | 1.3 | 3.6 | 6.6 | 10.8 | 16.6 | 24.2 | 28.6 | 37.1 | 40.3 | 52.4 |
| Female..... | 0.5 | 1.4 | 2.2 | 4.1 | 6.2 | 10.4 | 13.9 | 16.3 | 22.6 | 25.3 |
| Non-Hispanic black, both sexes..... | 0.9 | 2.9 | 4.5 | 7.2 | 10.9 | 14.7 | 17.2 | 26.2 | 27.0 | 28.1 |
| Male..... | * | 3.6 | 6.6 | 12.1 | 17.3 | 20.5 | 24.3 | 35.2 | 39.0 | 39.4 |
| Female..... | * | 2.3 | 2.6 | 3.1 | 5.9 | 10.4 | 12.4 | 20.7 | 20.7 | 23.8 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Malignant neoplasm of bladder (C67)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|-----|------|------|------|------|------|-------|
| All origins 3/, both sexes..... | 0.4 | 0.9 | 1.7 | 3.8 | 6.4 | 12.0 | 20.2 | 31.9 | 44.1 | 66.6 |
| Male..... | 0.5 | 1.2 | 2.4 | 5.9 | 10.1 | 19.0 | 32.2 | 54.0 | 76.6 | 128.9 |
| Female..... | 0.3 | 0.5 | 1.0 | 1.9 | 2.9 | 6.0 | 10.6 | 16.3 | 24.2 | 39.6 |
| Hispanic, both sexes..... | * | * | * | 2.4 | 3.6 | 6.1 | 9.7 | 14.1 | 25.4 | 54.9 |
| Male..... | * | * | * | * | 6.2 | 10.5 | 15.8 | 21.8 | 38.9 | 100.4 |
| Female..... | * | * | * | * | * | * | * | * | 16.8 | 32.3 |
| Mexican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 0.4 | 0.9 | 1.8 | 3.9 | 6.5 | 12.4 | 20.8 | 32.8 | 44.8 | 66.9 |
| Male..... | 0.5 | 1.3 | 2.5 | 6.1 | 10.4 | 19.6 | 33.2 | 55.7 | 78.2 | 129.8 |
| Female..... | 0.3 | 0.5 | 1.1 | 1.9 | 3.1 | 6.2 | 10.9 | 16.6 | 24.5 | 39.8 |
| Non-Hispanic white, both sexes..... | 0.4 | 1.0 | 1.9 | 4.0 | 7.0 | 13.0 | 21.6 | 34.0 | 46.0 | 68.0 |
| Male..... | 0.6 | 1.4 | 2.7 | 6.5 | 11.1 | 20.7 | 34.9 | 57.9 | 81.1 | 135.2 |
| Female..... | 0.3 | 0.5 | 1.0 | 1.7 | 3.1 | 6.1 | 10.7 | 16.9 | 24.5 | 39.0 |
| Non-Hispanic black, both sexes..... | * | * | 1.6 | 4.1 | 5.1 | 11.2 | 19.1 | 27.0 | 37.6 | 64.1 |
| Male..... | * | * | * | 4.4 | 6.9 | 14.6 | 24.3 | 42.3 | 57.7 | 91.9 |
| Female..... | * | * | * | 3.8 | 3.7 | 8.7 | 15.6 | 17.9 | 27.2 | 53.5 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|------|------|------|------|------|------|------|
| All origins 3/, both sexes..... | 2.8 | 4.2 | 6.2 | 8.5 | 11.1 | 14.7 | 17.8 | 19.6 | 18.5 | 16.4 |
| Male..... | 3.5 | 5.3 | 7.2 | 10.6 | 13.1 | 18.4 | 22.2 | 24.4 | 22.9 | 20.1 |
| Female..... | 2.1 | 3.1 | 5.1 | 6.5 | 9.2 | 11.4 | 14.2 | 16.2 | 15.8 | 14.8 |
| Hispanic, both sexes..... | 1.3 | 2.4 | 4.6 | 4.7 | 7.4 | 9.1 | 11.1 | 14.4 | 12.5 | 13.3 |
| Male..... | 1.5 | 2.9 | 6.4 | 5.0 | 9.0 | 12.2 | 12.7 | 17.3 | * | * |
| Female..... | * | 1.9 | 2.9 | 4.4 | 6.0 | 6.6 | 9.8 | 12.3 | * | * |
| Mexican, both sexes..... | * | 2.4 | 4.5 | 3.7 | 7.1 | 8.5 | 11.4 | 13.3 | * | * |
| Male..... | * | 3.3 | 6.3 | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 2.9 | 4.4 | 6.3 | 8.8 | 11.3 | 15.1 | 18.2 | 19.8 | 18.7 | 16.5 |
| Male..... | 3.7 | 5.6 | 7.3 | 11.1 | 13.3 | 18.9 | 22.8 | 24.7 | 23.3 | 20.2 |
| Female..... | 2.2 | 3.3 | 5.3 | 6.7 | 9.5 | 11.8 | 14.5 | 16.4 | 15.9 | 14.9 |
| Non-Hispanic white, both sexes..... | 3.3 | 4.9 | 6.9 | 9.5 | 12.3 | 16.4 | 19.5 | 20.9 | 19.6 | 17.3 |
| Male..... | 4.2 | 6.2 | 8.0 | 11.7 | 14.4 | 20.5 | 24.3 | 25.9 | 24.2 | 21.4 |
| Female..... | 2.4 | 3.7 | 5.8 | 7.3 | 10.5 | 12.8 | 15.6 | 17.4 | 16.9 | 15.5 |
| Non-Hispanic black, both sexes..... | 1.4 | 2.0 | 3.7 | 5.7 | 5.5 | 7.8 | 9.6 | 12.1 | 11.5 | 9.7 |
| Male..... | * | 2.2 | 3.3 | 8.3 | 7.7 | 9.0 | 12.0 | 15.0 | 17.9 | * |
| Female..... | * | 1.8 | 4.1 | 3.5 | 3.8 | 7.0 | 8.0 | 10.4 | 8.1 | 10.5 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

| | | | | | | | | | | |
|-------------------------------------|-----|------|------|------|------|------|-------|-------|-------|-------|
| All origins 3/, both sexes..... | 4.6 | 7.5 | 12.6 | 21.5 | 36.9 | 59.2 | 90.1 | 125.6 | 168.8 | 197.5 |
| Male..... | 5.5 | 9.1 | 15.1 | 26.8 | 45.7 | 76.4 | 116.8 | 164.3 | 224.2 | 269.1 |
| Female..... | 3.7 | 6.0 | 10.1 | 16.4 | 28.9 | 44.4 | 68.8 | 98.2 | 135.1 | 166.5 |
| Hispanic, both sexes..... | 4.5 | 6.5 | 9.3 | 18.1 | 27.6 | 47.3 | 61.7 | 89.8 | 119.0 | 132.5 |
| Male..... | 5.1 | 7.3 | 11.2 | 21.3 | 33.1 | 56.5 | 72.0 | 108.1 | 138.4 | 180.7 |
| Female..... | 3.9 | 5.7 | 7.5 | 15.2 | 22.8 | 39.8 | 53.7 | 76.8 | 106.6 | 108.6 |
| Mexican, both sexes..... | 4.0 | 5.2 | 8.2 | 16.1 | 27.9 | 46.3 | 68.7 | 86.4 | 117.0 | 121.8 |
| Male..... | 4.0 | 5.5 | 9.1 | 21.0 | 33.8 | 49.9 | 88.5 | 108.3 | * | * |
| Female..... | 3.9 | 4.9 | 7.2 | 11.3 | 22.9 | 43.1 | 54.8 | 72.0 | 113.5 | * |
| Puerto Rican, both sexes..... | * | * | * | 26.3 | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 4.0 | 7.2 | 9.4 | 18.4 | 32.6 | 61.8 | 57.6 | 80.3 | * | * |
| Male..... | * | 9.4 | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 4.6 | 7.6 | 12.8 | 21.7 | 37.5 | 59.9 | 91.7 | 127.2 | 170.5 | 199.8 |
| Male..... | 5.5 | 9.2 | 15.4 | 27.2 | 46.6 | 77.5 | 119.2 | 166.8 | 227.5 | 272.8 |
| Female..... | 3.6 | 6.0 | 10.3 | 16.5 | 29.3 | 44.6 | 69.7 | 99.2 | 135.9 | 168.3 |
| Non-Hispanic white, both sexes..... | 4.5 | 7.3 | 12.3 | 21.2 | 37.6 | 61.0 | 93.5 | 130.4 | 175.5 | 206.5 |
| Male..... | 5.5 | 8.9 | 15.0 | 26.8 | 46.6 | 79.1 | 121.8 | 171.7 | 234.5 | 283.5 |
| Female..... | 3.6 | 5.6 | 9.7 | 15.8 | 29.2 | 45.0 | 70.3 | 100.9 | 139.5 | 173.2 |
| Non-Hispanic black, both sexes..... | 5.8 | 10.8 | 17.5 | 30.3 | 44.3 | 63.1 | 94.6 | 113.6 | 138.5 | 153.7 |
| Male..... | 7.3 | 13.5 | 20.0 | 36.8 | 56.4 | 80.1 | 125.2 | 140.4 | 180.2 | 203.5 |
| Female..... | 4.4 | 8.5 | 15.4 | 25.0 | 34.8 | 50.5 | 74.1 | 97.6 | 116.9 | 134.6 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Non-Hodgkin's lymphoma (C82-C85)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|------|------|------|------|------|------|-------|
| All origins 3/, both sexes..... | 1.9 | 3.0 | 5.0 | 8.8 | 14.9 | 23.2 | 35.0 | 50.2 | 67.0 | 77.3 |
| Male..... | 2.4 | 3.9 | 6.2 | 11.3 | 18.4 | 30.0 | 42.8 | 61.8 | 84.4 | 99.7 |
| Female..... | 1.3 | 2.2 | 3.8 | 6.4 | 11.7 | 17.3 | 28.7 | 41.9 | 56.4 | 67.6 |
| Hispanic, both sexes..... | 1.7 | 2.5 | 3.5 | 6.7 | 13.1 | 18.6 | 27.9 | 41.3 | 47.7 | 50.4 |
| Male..... | 1.9 | 3.3 | 4.0 | 8.4 | 16.2 | 23.4 | 29.4 | 42.2 | 51.5 | 60.2 |
| Female..... | 1.6 | * | 3.1 | 5.1 | 10.5 | 14.7 | 26.7 | 40.7 | 45.3 | 45.6 |
| Mexican, both sexes..... | 1.6 | 1.9 | 3.6 | 5.8 | 12.8 | 20.4 | 31.1 | 37.9 | 50.7 | 52.2 |
| Male..... | * | * | * | 7.8 | 16.4 | 22.5 | 36.9 | * | * | * |
| Female..... | * | * | * | * | 9.7 | 18.5 | 27.1 | 36.4 | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | 18.1 | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 1.9 | 3.0 | 5.1 | 9.0 | 15.0 | 23.5 | 35.4 | 50.5 | 67.5 | 78.2 |
| Male..... | 2.5 | 3.9 | 6.4 | 11.5 | 18.5 | 30.4 | 43.5 | 62.7 | 85.5 | 101.4 |
| Female..... | 1.3 | 2.2 | 3.9 | 6.5 | 11.8 | 17.5 | 28.8 | 42.0 | 56.6 | 68.2 |
| Non-Hispanic white, both sexes..... | 1.8 | 3.1 | 5.0 | 9.1 | 15.8 | 24.7 | 36.9 | 53.4 | 71.1 | 83.1 |
| Male..... | 2.4 | 4.0 | 6.4 | 11.8 | 19.2 | 31.9 | 45.3 | 66.1 | 89.8 | 107.6 |
| Female..... | 1.3 | 2.1 | 3.7 | 6.6 | 12.5 | 18.4 | 30.0 | 44.3 | 59.7 | 72.6 |
| Non-Hispanic black, both sexes..... | 2.7 | 3.6 | 6.2 | 9.4 | 12.6 | 18.0 | 26.2 | 26.9 | 35.0 | 31.2 |
| Male..... | 3.8 | 4.2 | 7.6 | 11.7 | 16.9 | 23.3 | 34.9 | 31.9 | 42.1 | 43.8 |
| Female..... | 1.7 | 3.1 | 5.0 | 7.5 | 9.2 | 14.0 | 20.4 | 23.8 | 31.2 | 26.3 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Leukemia (C91-C95)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|------|------|------|------|------|------|-------|
| All origins 3/, both sexes..... | 1.9 | 2.8 | 4.5 | 7.4 | 12.9 | 20.9 | 32.4 | 45.7 | 63.4 | 80.5 |
| Male..... | 2.2 | 3.2 | 5.3 | 9.2 | 16.1 | 28.0 | 45.8 | 65.0 | 90.0 | 114.5 |
| Female..... | 1.6 | 2.5 | 3.8 | 5.7 | 10.0 | 14.7 | 21.8 | 32.1 | 47.3 | 65.7 |
| Hispanic, both sexes..... | 2.2 | 2.5 | 2.7 | 6.5 | 8.2 | 13.4 | 18.1 | 21.6 | 37.0 | 47.1 |
| Male..... | 2.5 | 2.1 | 3.7 | 6.6 | 8.0 | 15.2 | 22.8 | 32.0 | 50.3 | 65.3 |
| Female..... | 1.9 | 2.8 | * | 6.4 | 8.5 | 11.9 | 14.5 | 14.2 | 28.5 | 38.1 |
| Mexican, both sexes..... | 1.9 | 1.9 | * | 6.8 | 7.8 | 11.9 | 19.3 | 17.8 | 30.1 | * |
| Male..... | * | * | * | 7.5 | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 1.9 | 2.9 | 4.7 | 7.5 | 13.3 | 21.4 | 33.2 | 46.9 | 64.5 | 81.7 |
| Male..... | 2.2 | 3.3 | 5.4 | 9.4 | 16.7 | 28.8 | 47.0 | 66.5 | 91.7 | 116.5 |
| Female..... | 1.6 | 2.5 | 4.0 | 5.7 | 10.1 | 14.9 | 22.2 | 33.0 | 48.0 | 66.6 |
| Non-Hispanic white, both sexes..... | 2.0 | 2.7 | 4.7 | 7.4 | 13.4 | 22.4 | 34.5 | 48.3 | 66.9 | 84.4 |
| Male..... | 2.3 | 3.1 | 5.5 | 9.4 | 16.9 | 29.9 | 48.5 | 68.7 | 95.1 | 121.3 |
| Female..... | 1.7 | 2.4 | 4.0 | 5.4 | 10.2 | 15.6 | 23.1 | 33.8 | 49.6 | 68.4 |
| Non-Hispanic black, both sexes..... | 1.7 | 3.8 | 5.1 | 9.4 | 14.2 | 17.4 | 29.1 | 40.0 | 47.2 | 65.3 |
| Male..... | 1.8 | 5.0 | 5.6 | 11.4 | 17.7 | 23.8 | 43.4 | 53.5 | 65.5 | 87.5 |
| Female..... | 1.6 | 2.7 | 4.7 | 7.8 | 11.5 | 12.7 | 19.5 | 32.0 | 37.7 | 56.9 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|------|------|------|------|------|------|------|
| All origins 3/, both sexes..... | 0.4 | 1.3 | 2.5 | 4.7 | 8.3 | 14.1 | 21.0 | 27.9 | 35.8 | 36.7 |
| Male..... | 0.5 | 1.5 | 3.0 | 5.6 | 10.2 | 16.8 | 25.8 | 35.5 | 46.2 | 50.8 |
| Female..... | 0.4 | 1.0 | 2.0 | 4.0 | 6.5 | 11.7 | 17.2 | 22.5 | 29.5 | 30.6 |
| Hispanic, both sexes..... | * | 1.2 | 2.7 | 4.3 | 5.9 | 14.3 | 14.3 | 25.1 | 29.4 | 33.3 |
| Male..... | * | * | 2.8 | 5.5 | 8.2 | 17.3 | 18.0 | 32.6 | 29.7 | 53.5 |
| Female..... | * | * | 2.5 | * | * | 11.9 | 11.5 | 19.7 | 29.2 | 23.2 |
| Mexican, both sexes..... | * | * | 2.4 | 3.1 | 6.9 | 12.2 | 16.7 | 28.8 | 31.0 | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | 22.8 | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 0.5 | 1.3 | 2.5 | 4.8 | 8.4 | 14.0 | 21.4 | 27.9 | 36.1 | 36.7 |
| Male..... | 0.5 | 1.5 | 3.0 | 5.5 | 10.3 | 16.8 | 26.3 | 35.5 | 46.8 | 50.6 |
| Female..... | 0.4 | 1.1 | 2.0 | 4.0 | 6.7 | 11.6 | 17.5 | 22.5 | 29.5 | 30.8 |
| Non-Hispanic white, both sexes..... | 0.4 | 1.0 | 2.0 | 4.1 | 7.6 | 12.8 | 20.2 | 26.8 | 35.1 | 35.7 |
| Male..... | 0.4 | 1.3 | 2.7 | 4.9 | 9.5 | 15.6 | 25.3 | 34.6 | 46.0 | 50.1 |
| Female..... | 0.3 | 0.8 | 1.4 | 3.4 | 5.8 | 10.3 | 16.0 | 21.2 | 28.4 | 29.5 |
| Non-Hispanic black, both sexes..... | 1.0 | 3.0 | 5.7 | 10.9 | 16.6 | 26.9 | 38.0 | 45.5 | 55.0 | 55.7 |
| Male..... | * | 3.7 | 6.1 | 12.6 | 20.4 | 32.0 | 45.2 | 53.5 | 71.0 | 71.1 |
| Female..... | * | 2.5 | 5.4 | 9.4 | 13.7 | 23.2 | 33.3 | 40.6 | 46.7 | 49.7 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)

Table with 11 columns representing age groups (40-44 to 85 years and over) and rows for various demographic groups including All origins, Hispanic, Mexican, Puerto Rican, Cuban, Other Hispanic, Non-Hispanic white, and Non-Hispanic black, each split by sex. All data points are marked with an asterisk (*).

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

| | | | | | | | | | | |
|-------------------------------------|-----|------|------|------|------|------|-------|-------|-------|-------|
| All origins 3/, both sexes..... | 5.8 | 10.4 | 17.5 | 28.8 | 46.5 | 66.0 | 97.0 | 131.0 | 168.7 | 230.1 |
| Male..... | 6.2 | 11.7 | 20.8 | 33.0 | 56.5 | 79.9 | 118.9 | 161.8 | 207.8 | 290.8 |
| Female..... | 5.4 | 9.1 | 14.3 | 24.9 | 37.5 | 53.9 | 79.5 | 109.2 | 144.9 | 203.8 |
| Hispanic, both sexes..... | 4.0 | 6.9 | 12.8 | 20.4 | 32.3 | 48.4 | 75.2 | 102.6 | 129.7 | 164.1 |
| Male..... | 4.0 | 7.3 | 14.5 | 25.1 | 35.9 | 55.5 | 86.9 | 127.3 | 159.0 | 204.1 |
| Female..... | 3.9 | 6.5 | 11.2 | 16.0 | 29.1 | 42.6 | 66.2 | 85.0 | 111.0 | 144.2 |
| Mexican, both sexes..... | 4.7 | 7.2 | 12.8 | 19.2 | 31.8 | 47.8 | 85.0 | 92.8 | 125.6 | 147.9 |
| Male..... | 4.4 | 8.4 | 13.8 | 22.8 | 33.3 | 51.1 | 108.8 | 117.6 | * | * |
| Female..... | 5.0 | 6.1 | 11.8 | 15.7 | 30.6 | 44.8 | 68.3 | 76.5 | * | * |
| Puerto Rican, both sexes..... | * | * | 14.4 | 26.3 | 37.7 | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 3.1 | 5.9 | 10.7 | 21.2 | 37.4 | 61.0 | 74.4 | 124.5 | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | 56.2 | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 6.0 | 10.7 | 17.9 | 29.4 | 47.5 | 67.0 | 98.2 | 132.1 | 170.1 | 232.1 |
| Male..... | 6.4 | 12.1 | 21.3 | 33.6 | 57.8 | 81.3 | 120.7 | 163.2 | 209.4 | 294.0 |
| Female..... | 5.6 | 9.4 | 14.6 | 25.5 | 38.1 | 54.6 | 80.2 | 110.2 | 146.2 | 205.4 |
| Non-Hispanic white, both sexes..... | 5.7 | 10.1 | 17.1 | 28.7 | 46.6 | 66.5 | 98.0 | 132.8 | 171.4 | 235.2 |
| Male..... | 5.9 | 11.4 | 20.4 | 32.5 | 56.5 | 81.6 | 120.4 | 163.5 | 212.7 | 300.2 |
| Female..... | 5.5 | 8.8 | 13.9 | 25.0 | 37.4 | 53.2 | 79.7 | 110.7 | 146.2 | 207.0 |
| Non-Hispanic black, both sexes..... | 8.2 | 16.1 | 25.8 | 42.4 | 64.0 | 83.8 | 115.8 | 149.2 | 175.4 | 223.2 |
| Male..... | 9.9 | 18.7 | 31.6 | 51.4 | 82.3 | 99.1 | 145.7 | 190.1 | 210.6 | 265.8 |
| Female..... | 6.8 | 13.9 | 20.9 | 35.0 | 49.6 | 72.5 | 95.7 | 124.7 | 157.1 | 206.9 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|-----|-----|------|------|------|------|-------|
| All origins 3/, both sexes..... | 1.0 | 1.5 | 2.1 | 3.7 | 5.4 | 9.7 | 16.3 | 28.8 | 46.3 | 78.7 |
| Male..... | 1.0 | 1.6 | 2.1 | 4.5 | 6.3 | 12.2 | 20.0 | 37.3 | 59.1 | 101.1 |
| Female..... | 0.9 | 1.5 | 2.0 | 3.0 | 4.6 | 7.5 | 13.4 | 22.8 | 38.5 | 68.9 |
| Hispanic, both sexes..... | 0.8 | 1.4 | 1.5 | 1.8 | 4.8 | 6.2 | 12.2 | 17.3 | 37.0 | 48.8 |
| Male..... | * | * | * | * | 5.4 | 7.8 | 13.6 | 19.8 | 52.6 | 56.9 |
| Female..... | * | * | * | * | * | * | 11.2 | 15.5 | 27.0 | 44.8 |
| Mexican, both sexes..... | * | * | * | * | * | 8.5 | 16.3 | 18.3 | 35.3 | 48.9 |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 1.0 | 1.5 | 2.1 | 3.9 | 5.5 | 9.9 | 16.6 | 29.3 | 46.6 | 79.8 |
| Male..... | 1.1 | 1.6 | 2.2 | 4.7 | 6.4 | 12.5 | 20.4 | 38.0 | 59.2 | 102.9 |
| Female..... | 0.9 | 1.5 | 2.0 | 3.1 | 4.7 | 7.7 | 13.5 | 23.2 | 38.9 | 69.8 |
| Non-Hispanic white, both sexes..... | 0.9 | 1.6 | 2.0 | 3.8 | 5.6 | 10.1 | 17.0 | 30.2 | 47.4 | 81.8 |
| Male..... | 1.1 | 1.7 | 2.2 | 4.5 | 6.5 | 12.8 | 21.1 | 39.2 | 60.2 | 106.0 |
| Female..... | 0.8 | 1.4 | 1.8 | 3.1 | 4.8 | 7.8 | 13.6 | 23.8 | 39.6 | 71.4 |
| Non-Hispanic black, both sexes..... | 1.7 | 2.0 | 3.1 | 5.3 | 6.0 | 10.8 | 15.4 | 24.7 | 42.4 | 64.4 |
| Male..... | * | * | 2.9 | 7.7 | 6.7 | 12.8 | 16.4 | 30.0 | 54.6 | 81.0 |
| Female..... | 2.0 | 2.3 | 3.3 | 3.4 | 5.4 | 9.3 | 14.7 | 21.6 | 36.1 | 58.1 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Anemias (D50-D64)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|-----|-----|-----|------|------|------|------|
| All origins 3/, both sexes..... | 0.4 | 0.6 | 0.6 | 0.8 | 1.2 | 1.9 | 3.7 | 6.4 | 12.9 | 40.3 |
| Male..... | 0.4 | 0.5 | 0.6 | 0.8 | 1.3 | 1.7 | 4.3 | 6.6 | 12.4 | 38.3 |
| Female..... | 0.5 | 0.7 | 0.7 | 0.8 | 1.1 | 2.1 | 3.2 | 6.2 | 13.2 | 41.2 |
| Hispanic, both sexes..... | * | * | * | * | * | * | 4.4 | * | 11.6 | 34.4 |
| Male..... | * | * | * | * | * | * | * | * | * | 53.5 |
| Female..... | * | * | * | * | * | * | * | * | * | 24.9 |
| Mexican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 0.4 | 0.6 | 0.7 | 0.9 | 1.2 | 1.9 | 3.7 | 6.5 | 12.9 | 40.5 |
| Male..... | 0.4 | 0.6 | 0.6 | 0.8 | 1.3 | 1.7 | 4.3 | 6.7 | 12.5 | 37.6 |
| Female..... | 0.5 | 0.7 | 0.7 | 0.9 | 1.1 | 2.1 | 3.2 | 6.3 | 13.2 | 41.8 |
| Non-Hispanic white, both sexes..... | * | 0.3 | 0.3 | 0.6 | 0.9 | 1.7 | 3.3 | 5.9 | 12.6 | 39.9 |
| Male..... | * | 0.3 | 0.4 | 0.6 | 1.0 | 1.5 | 3.6 | 6.3 | 11.9 | 36.3 |
| Female..... | * | 0.3 | 0.3 | 0.6 | 0.8 | 1.8 | 3.0 | 5.6 | 13.0 | 41.4 |
| Non-Hispanic black, both sexes..... | 2.5 | 3.2 | 3.1 | 3.2 | 4.2 | 4.5 | 8.1 | 14.0 | 19.5 | 52.9 |
| Male..... | 2.4 | 2.7 | 2.6 | 3.1 | 5.3 | * | 12.0 | 13.1 | 23.4 | 58.0 |
| Female..... | 2.6 | 3.6 | 3.5 | 3.4 | 3.3 | 4.7 | 5.5 | 14.6 | 17.5 | 51.0 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Diabetes mellitus (E10-E14)

| | | | | | | | | | | |
|-------------------------------------|------|------|------|------|-------|-------|-------|-------|-------|-------|
| All origins 3/, both sexes..... | 6.0 | 9.9 | 18.0 | 30.1 | 47.5 | 75.7 | 108.8 | 157.7 | 217.9 | 320.6 |
| Male..... | 7.7 | 11.9 | 21.5 | 36.0 | 55.5 | 84.3 | 121.1 | 175.9 | 243.5 | 348.5 |
| Female..... | 4.4 | 7.9 | 14.7 | 24.5 | 40.2 | 68.2 | 99.0 | 144.9 | 202.3 | 308.4 |
| Hispanic, both sexes..... | 5.0 | 9.2 | 20.4 | 38.0 | 71.3 | 108.2 | 168.0 | 257.0 | 333.3 | 437.9 |
| Male..... | 6.1 | 10.2 | 25.4 | 43.3 | 80.8 | 110.7 | 186.5 | 282.8 | 350.0 | 425.0 |
| Female..... | 3.8 | 8.2 | 15.4 | 33.1 | 63.1 | 106.2 | 153.8 | 238.6 | 322.7 | 444.3 |
| Mexican, both sexes..... | 6.1 | 9.0 | 23.9 | 43.8 | 88.8 | 141.6 | 220.4 | 296.7 | 409.3 | 462.2 |
| Male..... | 7.6 | 9.3 | 29.8 | 48.8 | 101.4 | 134.4 | 258.1 | 341.2 | * | * |
| Female..... | 4.3 | 8.7 | 17.4 | 39.0 | 78.0 | 148.3 | 194.1 | 267.5 | 416.2 | 474.1 |
| Puerto Rican, both sexes..... | * | 17.4 | 23.3 | 51.1 | 68.2 | 121.2 | 183.5 | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | 38.7 | * | 58.7 | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | 6.1 | 11.5 | 22.8 | 44.0 | 76.5 | 117.8 | 203.4 | * | * |
| Male..... | * | * | * | 28.9 | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | 67.0 | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 6.1 | 9.9 | 17.8 | 29.3 | 45.4 | 73.0 | 104.7 | 151.8 | 212.2 | 315.0 |
| Male..... | 7.9 | 12.0 | 21.1 | 35.4 | 53.3 | 82.0 | 116.7 | 169.5 | 237.7 | 344.0 |
| Female..... | 4.5 | 7.9 | 14.6 | 23.7 | 38.2 | 65.1 | 95.2 | 139.4 | 196.7 | 302.5 |
| Non-Hispanic white, both sexes..... | 5.3 | 8.2 | 14.7 | 24.7 | 38.3 | 61.9 | 92.7 | 138.0 | 197.8 | 296.4 |
| Male..... | 6.9 | 10.2 | 17.9 | 30.7 | 46.4 | 72.5 | 107.2 | 160.1 | 228.4 | 338.4 |
| Female..... | 3.7 | 6.3 | 11.6 | 18.8 | 30.7 | 52.5 | 80.8 | 122.1 | 179.0 | 278.3 |
| Non-Hispanic black, both sexes..... | 12.3 | 22.0 | 41.8 | 69.3 | 107.3 | 164.9 | 225.1 | 303.3 | 396.4 | 544.7 |
| Male..... | 15.4 | 26.1 | 47.7 | 78.6 | 120.6 | 168.4 | 226.5 | 297.7 | 390.8 | 423.4 |
| Female..... | 9.7 | 18.5 | 36.7 | 61.8 | 96.9 | 162.2 | 224.1 | 306.7 | 399.3 | 591.1 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Nutritional deficiencies (E40-E64)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|-----|-----|-----|-----|------|------|------|
| All origins 3/, both sexes..... | 0.1 | 0.2 | 0.3 | 0.5 | 0.8 | 1.6 | 2.7 | 6.1 | 12.8 | 41.5 |
| Male..... | * | 0.2 | 0.3 | 0.6 | 0.9 | 1.6 | 3.0 | 6.3 | 12.0 | 37.0 |
| Female..... | * | * | 0.4 | 0.4 | 0.7 | 1.6 | 2.4 | 6.0 | 13.3 | 43.5 |
| Hispanic, both sexes..... | * | * | * | * | * | * | * | 6.7 | 10.7 | 29.4 |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | 29.0 |
| Mexican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 0.2 | 0.2 | 0.3 | 0.5 | 0.8 | 1.6 | 2.6 | 6.0 | 12.9 | 41.8 |
| Male..... | * | * | 0.3 | 0.6 | 1.0 | 1.6 | 3.0 | 6.2 | 12.0 | 37.1 |
| Female..... | * | * | 0.4 | 0.5 | 0.7 | 1.6 | 2.4 | 5.8 | 13.4 | 43.9 |
| Non-Hispanic white, both sexes..... | 0.1 | 0.2 | 0.3 | 0.5 | 0.7 | 1.4 | 2.4 | 5.7 | 12.3 | 40.6 |
| Male..... | * | * | * | 0.5 | 0.7 | 1.4 | 2.7 | 6.0 | 11.0 | 36.5 |
| Female..... | * | * | 0.4 | 0.5 | 0.7 | 1.4 | 2.2 | 5.5 | 13.2 | 42.3 |
| Non-Hispanic black, both sexes..... | * | * | * | 1.6 | 2.1 | 3.6 | 5.1 | 10.7 | 20.8 | 60.2 |
| Male..... | * | * | * | * | * | * | * | 9.9 | 27.3 | 45.9 |
| Female..... | * | * | * | * | * | * | 4.6 | 11.2 | 17.5 | 65.6 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Malnutrition (E40-E46)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|-----|-----|-----|-----|------|------|------|
| All origins 3/, both sexes..... | 0.1 | 0.2 | 0.3 | 0.5 | 0.7 | 1.4 | 2.5 | 5.6 | 12.2 | 38.4 |
| Male..... | * | * | 0.3 | 0.5 | 0.8 | 1.4 | 2.8 | 5.7 | 11.4 | 35.1 |
| Female..... | * | * | 0.3 | 0.4 | 0.6 | 1.4 | 2.2 | 5.6 | 12.6 | 39.8 |
| Hispanic, both sexes..... | * | * | * | * | * | * | * | 6.7 | 10.7 | 28.3 |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | 28.2 |
| Mexican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 0.1 | 0.2 | 0.3 | 0.5 | 0.7 | 1.4 | 2.5 | 5.5 | 12.2 | 38.7 |
| Male..... | * | * | 0.3 | 0.6 | 0.9 | 1.4 | 2.8 | 5.7 | 11.4 | 35.3 |
| Female..... | * | * | 0.3 | 0.5 | 0.6 | 1.5 | 2.2 | 5.4 | 12.7 | 40.1 |
| Non-Hispanic white, both sexes..... | 0.1 | 0.2 | 0.3 | 0.4 | 0.6 | 1.3 | 2.3 | 5.2 | 11.7 | 37.4 |
| Male..... | * | * | * | 0.4 | 0.6 | 1.3 | 2.6 | 5.4 | 10.5 | 34.5 |
| Female..... | * | * | 0.3 | 0.5 | 0.6 | 1.3 | 2.1 | 5.1 | 12.5 | 38.6 |
| Non-Hispanic black, both sexes..... | * | * | * | 1.5 | 2.0 | 3.1 | 4.4 | 10.4 | 20.0 | 57.2 |
| Male..... | * | * | * | * | * | * | * | 9.4 | 25.0 | 44.9 |
| Female..... | * | * | * | * | * | * | * | 10.9 | 17.5 | 61.9 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Parkinson's disease (G20-G21)

| | | | | | | | | | | |
|-------------------------------------|---|---|-----|-----|-----|------|------|------|-------|-------|
| All origins 3/, both sexes..... | * | * | 0.2 | 0.6 | 1.9 | 6.5 | 18.4 | 45.4 | 89.8 | 135.2 |
| Male..... | * | * | 0.3 | 0.8 | 2.5 | 9.8 | 27.3 | 67.2 | 136.6 | 220.1 |
| Female..... | * | * | * | 0.5 | 1.4 | 3.8 | 11.3 | 30.0 | 61.3 | 98.4 |
| Hispanic, both sexes..... | * | * | * | * | * | 5.9 | 12.0 | 27.7 | 53.0 | 76.5 |
| Male..... | * | * | * | * | * | 8.8 | 19.7 | 40.3 | 67.5 | 113.8 |
| Female..... | * | * | * | * | * | * | * | 18.7 | 43.8 | 58.0 |
| Mexican, both sexes..... | * | * | * | * | * | * | 10.6 | 25.6 | 47.3 | 77.2 |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | * | * | 0.2 | 0.6 | 1.9 | 6.6 | 18.8 | 46.2 | 91.2 | 137.4 |
| Male..... | * | * | 0.3 | 0.8 | 2.4 | 9.8 | 27.7 | 68.5 | 139.5 | 224.4 |
| Female..... | * | * | * | 0.5 | 1.5 | 3.8 | 11.6 | 30.5 | 61.9 | 99.8 |
| Non-Hispanic white, both sexes..... | * | * | 0.3 | 0.6 | 2.1 | 7.0 | 20.0 | 49.3 | 97.4 | 145.8 |
| Male..... | * | * | 0.4 | 0.8 | 2.7 | 10.3 | 29.3 | 72.7 | 148.2 | 237.8 |
| Female..... | * | * | * | 0.5 | 1.6 | 4.0 | 12.4 | 32.6 | 66.3 | 106.0 |
| Non-Hispanic black, both sexes..... | * | * | * | * | * | 4.5 | 9.6 | 21.1 | 32.8 | 51.7 |
| Male..... | * | * | * | * | * | 7.4 | 16.1 | 31.0 | 56.2 | 87.5 |
| Female..... | * | * | * | * | * | * | 5.3 | 15.1 | 20.7 | 38.0 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Alzheimer's disease (G30)

| | | | | | | | | | | |
|-------------------------------------|---|---|-----|-----|-----|------|------|-------|-------|-------|
| All origins 3/, both sexes..... | * | * | 0.2 | 0.9 | 3.2 | 9.8 | 30.6 | 95.4 | 245.7 | 752.3 |
| Male..... | * | * | 0.2 | 0.8 | 3.1 | 9.2 | 29.5 | 90.9 | 220.6 | 572.0 |
| Female..... | * | * | * | 1.0 | 3.3 | 10.3 | 31.5 | 98.5 | 260.9 | 830.5 |
| Hispanic, both sexes..... | * | * | * | * | * | 5.9 | 16.8 | 56.3 | 144.4 | 438.5 |
| Male..... | * | * | * | * | * | * | 16.2 | 54.4 | 127.0 | 341.3 |
| Female..... | * | * | * | * | * | 5.8 | 17.2 | 57.6 | 155.5 | 486.6 |
| Mexican, both sexes..... | * | * | * | * | * | * | 13.7 | 50.3 | 128.1 | 364.4 |
| Male..... | * | * | * | * | * | * | * | 51.9 | * | * |
| Female..... | * | * | * | * | * | * | 49.2 | 139.2 | 397.9 | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | * | * | 0.2 | 1.0 | 3.3 | 10.0 | 31.4 | 97.2 | 249.6 | 763.2 |
| Male..... | * | * | 0.3 | 0.8 | 3.2 | 9.4 | 30.2 | 92.6 | 224.5 | 581.4 |
| Female..... | * | * | * | 1.1 | 3.4 | 10.6 | 32.4 | 100.5 | 264.8 | 841.6 |
| Non-Hispanic white, both sexes..... | * | * | 0.3 | 1.1 | 3.5 | 10.5 | 32.8 | 101.2 | 259.6 | 793.9 |
| Male..... | * | * | * | 0.9 | 3.4 | 9.6 | 31.4 | 96.2 | 232.0 | 605.4 |
| Female..... | * | * | * | 1.2 | 3.6 | 11.4 | 34.0 | 104.8 | 276.6 | 875.3 |
| Non-Hispanic black, both sexes..... | * | * | * | * | 2.5 | 8.4 | 27.5 | 82.1 | 188.2 | 548.7 |
| Male..... | * | * | * | * | * | 9.2 | 29.4 | 82.2 | 189.5 | 414.6 |
| Female..... | * | * | * | * | * | 7.8 | 26.2 | 82.1 | 187.5 | 599.9 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Major cardiovascular diseases (I00-I78)

| | | | | | | | | | | |
|-------------------------------------|-------|-------|-------|-------|-------|---------|---------|---------|---------|---------|
| All origins 3/, both sexes..... | 49.7 | 87.9 | 144.2 | 233.5 | 374.0 | 602.6 | 993.2 | 1,700.2 | 3,034.4 | 7,389.6 |
| Male..... | 67.9 | 123.9 | 206.1 | 326.7 | 512.6 | 802.0 | 1,287.4 | 2,140.7 | 3,633.4 | 7,638.8 |
| Female..... | 31.6 | 52.9 | 85.0 | 146.0 | 248.2 | 430.4 | 758.0 | 1,389.0 | 2,670.1 | 7,281.5 |
| Hispanic, both sexes..... | 28.2 | 56.4 | 96.2 | 171.9 | 299.2 | 471.4 | 810.5 | 1,353.8 | 2,354.6 | 5,369.7 |
| Male..... | 39.5 | 79.0 | 135.0 | 247.0 | 413.5 | 626.1 | 1,077.3 | 1,686.2 | 2,749.8 | 5,494.9 |
| Female..... | 16.2 | 33.4 | 58.5 | 102.9 | 199.5 | 345.0 | 605.0 | 1,116.5 | 2,102.3 | 5,307.7 |
| Mexican, both sexes..... | 26.5 | 54.5 | 94.6 | 161.1 | 296.3 | 485.7 | 904.4 | 1,258.1 | 2,355.4 | 5,023.8 |
| Male..... | 35.9 | 73.3 | 125.0 | 220.0 | 407.1 | 604.3 | 1,274.9 | 1,662.0 | 2,571.0 | * |
| Female..... | 15.5 | 34.9 | 61.9 | 103.9 | 201.4 | 376.6 | 645.3 | 992.6 | 2,191.5 | 4,934.4 |
| Puerto Rican, both sexes..... | 49.5 | 80.2 | 114.4 | 266.6 | 389.6 | 658.1 | 928.7 | * | * | * |
| Male..... | 68.5 | 111.6 | 158.2 | 399.3 | * | * | * | * | * | * |
| Female..... | 32.5 | 47.5 | 75.0 | 159.9 | 254.5 | * | * | * | * | * |
| Cuban, both sexes..... | 38.9 | 71.2 | 122.2 | 170.5 | 212.0 | 319.5 | 581.1 | 1,403.9 | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 22.9 | 48.8 | 86.9 | 151.1 | 298.0 | 442.6 | 694.2 | 1,267.3 | * | * |
| Male..... | 36.3 | 75.0 | 142.1 | 225.6 | 388.7 | 681.1 | 791.1 | * | * | * |
| Female..... | 11.1 | 24.2 | 43.8 | 80.9 | 209.6 | 293.0 | 609.8 | * | * | * |
| Non-Hispanic 5/, both sexes..... | 52.2 | 90.9 | 147.9 | 237.6 | 378.1 | 609.6 | 1,001.0 | 1,713.0 | 3,056.0 | 7,451.4 |
| Male..... | 71.5 | 128.4 | 211.5 | 331.6 | 517.8 | 810.8 | 1,295.2 | 2,157.3 | 3,663.2 | 7,713.8 |
| Female..... | 33.4 | 54.7 | 87.1 | 149.1 | 251.0 | 435.1 | 765.3 | 1,399.4 | 2,687.5 | 7,338.2 |
| Non-Hispanic white, both sexes..... | 45.6 | 78.6 | 129.6 | 214.1 | 346.2 | 572.1 | 961.8 | 1,673.1 | 3,043.0 | 7,546.1 |
| Male..... | 64.9 | 115.3 | 190.6 | 304.1 | 481.2 | 771.3 | 1,257.4 | 2,127.0 | 3,670.7 | 7,845.8 |
| Female..... | 26.4 | 42.2 | 69.9 | 127.6 | 220.8 | 395.7 | 719.9 | 1,348.3 | 2,659.1 | 7,416.6 |
| Non-Hispanic black, both sexes..... | 101.5 | 188.5 | 308.2 | 477.1 | 708.5 | 1,015.7 | 1,529.7 | 2,367.5 | 3,588.7 | 7,136.9 |
| Male..... | 126.0 | 242.6 | 413.5 | 644.4 | 949.5 | 1,310.3 | 1,933.1 | 2,831.4 | 4,127.6 | 6,982.7 |
| Female..... | 80.0 | 141.6 | 219.1 | 339.9 | 520.3 | 797.4 | 1,259.2 | 2,090.4 | 3,308.3 | 7,195.9 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Diseases of heart (I00-I09,I11,I13,I20-I51)

| | | | | | | | | | | |
|-------------------------------------|------|-------|-------|-------|-------|---------|---------|---------|---------|---------|
| All origins 3/, both sexes..... | 40.2 | 71.3 | 119.2 | 192.8 | 304.4 | 479.6 | 766.1 | 1,278.2 | 2,234.3 | 5,446.8 |
| Male..... | 57.7 | 104.7 | 177.0 | 278.2 | 429.1 | 654.3 | 1,025.8 | 1,667.3 | 2,786.9 | 5,823.5 |
| Female..... | 22.9 | 38.8 | 63.9 | 112.5 | 191.2 | 328.8 | 558.5 | 1,003.3 | 1,898.3 | 5,283.3 |
| Hispanic, both sexes..... | 20.3 | 41.0 | 73.9 | 134.3 | 234.7 | 376.2 | 620.0 | 1,028.8 | 1,746.8 | 4,121.9 |
| Male..... | 30.0 | 59.4 | 108.7 | 199.5 | 335.0 | 511.6 | 846.9 | 1,314.4 | 2,109.2 | 4,301.8 |
| Female..... | 10.0 | 22.4 | 40.2 | 74.4 | 147.2 | 265.7 | 445.1 | 824.9 | 1,515.4 | 4,032.8 |
| Mexican, both sexes..... | 19.1 | 38.8 | 71.8 | 127.1 | 227.2 | 383.5 | 674.5 | 931.7 | 1,677.7 | 3,748.0 |
| Male..... | 27.9 | 53.5 | 99.2 | 179.3 | 322.6 | 492.4 | 990.0 | 1,260.9 | 1,893.9 | * |
| Female..... | 8.9 | 23.6 | 42.3 | 76.2 | 145.6 | 283.2 | 453.9 | 715.3 | 1,513.5 | 3,620.4 |
| Puerto Rican, both sexes..... | 37.9 | 61.9 | 93.3 | 204.3 | 319.6 | 527.8 | 749.9 | * | * | * |
| Male..... | 49.9 | 89.8 | 133.6 | 310.0 | * | * | * | * | * | * |
| Female..... | 27.2 | * | 57.0 | 119.2 | 214.2 | * | * | * | * | * |
| Cuban, both sexes..... | * | 58.4 | 100.2 | 138.2 | 172.3 | 272.0 | 472.1 | 1,176.7 | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 15.3 | 35.0 | 65.4 | 115.4 | 235.3 | 349.9 | 528.7 | 928.7 | * | * |
| Male..... | 25.6 | 57.0 | 111.7 | 182.7 | 330.1 | 549.4 | 616.2 | * | * | * |
| Female..... | 6.1 | 14.3 | 29.3 | 52.1 | 142.9 | 224.9 | 452.4 | * | * | * |
| Non-Hispanic 5/, both sexes..... | 42.6 | 74.2 | 122.7 | 196.7 | 308.3 | 484.9 | 772.1 | 1,287.0 | 2,249.3 | 5,484.9 |
| Male..... | 61.3 | 109.3 | 182.2 | 283.1 | 434.2 | 661.3 | 1,032.2 | 1,679.9 | 2,809.3 | 5,874.4 |
| Female..... | 24.4 | 40.4 | 65.8 | 115.3 | 193.8 | 332.0 | 563.7 | 1,009.7 | 1,909.4 | 5,316.8 |
| Non-Hispanic white, both sexes..... | 38.5 | 66.3 | 110.8 | 180.7 | 286.8 | 460.5 | 748.2 | 1,262.7 | 2,248.0 | 5,568.4 |
| Male..... | 57.2 | 100.8 | 168.9 | 264.7 | 409.8 | 636.5 | 1,010.8 | 1,665.4 | 2,827.0 | 5,994.1 |
| Female..... | 19.8 | 32.2 | 54.0 | 99.9 | 172.5 | 304.7 | 533.4 | 974.5 | 1,893.8 | 5,384.5 |
| Non-Hispanic black, both sexes..... | 76.5 | 142.1 | 236.4 | 372.5 | 550.7 | 774.7 | 1,140.0 | 1,731.2 | 2,577.8 | 5,151.0 |
| Male..... | 99.1 | 190.7 | 324.8 | 512.2 | 751.5 | 1,014.9 | 1,464.8 | 2,114.1 | 3,061.4 | 5,161.2 |
| Female..... | 56.6 | 100.0 | 161.6 | 258.0 | 393.8 | 596.6 | 922.2 | 1,502.6 | 2,326.2 | 5,147.0 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|-----|-----|-----|-----|------|------|------|
| All origins 3/, both sexes..... | 0.2 | 0.5 | 0.7 | 1.3 | 2.1 | 2.8 | 5.0 | 8.4 | 11.4 | 17.8 |
| Male..... | 0.2 | 0.4 | 0.6 | 1.0 | 1.5 | 1.9 | 4.2 | 5.8 | 8.2 | 11.8 |
| Female..... | 0.3 | 0.5 | 0.9 | 1.5 | 2.6 | 3.7 | 5.6 | 10.2 | 13.4 | 20.4 |
| Hispanic, both sexes..... | * | * | * | * | 2.4 | 3.2 | 6.3 | 5.9 | 8.9 | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | 8.1 | 9.1 | * | * |
| Mexican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 0.3 | 0.4 | 0.7 | 1.3 | 2.1 | 2.8 | 4.9 | 8.5 | 11.5 | 18.1 |
| Male..... | 0.2 | 0.4 | 0.6 | 1.0 | 1.4 | 1.9 | 4.2 | 6.0 | 8.4 | 12.0 |
| Female..... | 0.3 | 0.5 | 0.9 | 1.5 | 2.6 | 3.6 | 5.4 | 10.2 | 13.4 | 20.8 |
| Non-Hispanic white, both sexes..... | 0.2 | 0.4 | 0.7 | 1.1 | 2.0 | 2.7 | 5.1 | 8.7 | 12.3 | 19.3 |
| Male..... | * | 0.3 | 0.5 | 0.9 | 1.4 | 1.8 | 4.4 | 6.2 | 9.1 | 12.9 |
| Female..... | * | 0.4 | 0.9 | 1.4 | 2.5 | 3.6 | 5.7 | 10.6 | 14.2 | 22.0 |
| Non-Hispanic black, both sexes..... | * | 0.9 | 1.1 | 2.1 | 2.9 | 3.2 | * | 4.9 | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | 3.7 | * | * | 5.6 | * | * |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Hypertensive heart disease (I11)

| | | | | | | | | | | |
|-------------------------------------|------|------|------|------|------|------|------|-------|-------|-------|
| All origins 3/, both sexes..... | 3.5 | 5.9 | 8.6 | 11.1 | 14.3 | 18.9 | 24.0 | 37.9 | 66.0 | 187.0 |
| Male..... | 4.9 | 8.0 | 11.9 | 15.2 | 19.0 | 23.6 | 27.0 | 41.0 | 62.2 | 134.6 |
| Female..... | 2.2 | 3.9 | 5.4 | 7.2 | 10.0 | 14.8 | 21.6 | 35.8 | 68.2 | 209.7 |
| Hispanic, both sexes..... | 2.2 | 4.0 | 6.1 | 10.7 | 14.2 | 19.8 | 25.6 | 43.2 | 65.1 | 165.7 |
| Male..... | 3.7 | 6.2 | 9.0 | 15.4 | 19.2 | 24.4 | 33.8 | 51.8 | 77.8 | 137.2 |
| Female..... | * | * | 3.4 | 6.4 | 9.8 | 16.0 | 19.3 | 37.0 | 56.9 | 179.9 |
| Mexican, both sexes..... | 2.2 | 3.8 | 4.0 | 10.0 | 12.3 | 20.1 | 25.4 | 38.4 | 57.6 | 167.5 |
| Male..... | 3.6 | 5.5 | 6.1 | 14.1 | 14.6 | 24.3 | 36.0 | * | * | * |
| Female..... | * | * | * | * | 10.5 | 16.2 | 18.1 | 33.3 | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | 8.1 | * | 16.9 | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 3.7 | 6.1 | 8.7 | 11.0 | 14.2 | 18.5 | 23.8 | 37.5 | 65.7 | 187.1 |
| Male..... | 5.0 | 8.2 | 12.1 | 15.1 | 18.9 | 23.1 | 26.4 | 40.1 | 61.1 | 133.7 |
| Female..... | 2.3 | 4.1 | 5.6 | 7.2 | 9.9 | 14.5 | 21.6 | 35.6 | 68.5 | 210.2 |
| Non-Hispanic white, both sexes..... | 2.5 | 4.2 | 6.0 | 7.8 | 10.0 | 13.7 | 18.7 | 31.0 | 57.5 | 177.0 |
| Male..... | 3.7 | 6.0 | 8.8 | 11.2 | 13.5 | 17.3 | 21.0 | 33.2 | 53.3 | 123.5 |
| Female..... | 1.2 | 2.5 | 3.3 | 4.5 | 6.8 | 10.5 | 16.8 | 29.5 | 60.0 | 200.1 |
| Non-Hispanic black, both sexes..... | 11.6 | 19.8 | 30.8 | 40.4 | 51.2 | 61.1 | 77.3 | 111.6 | 168.4 | 322.1 |
| Male..... | 14.7 | 24.9 | 40.4 | 54.2 | 72.1 | 78.8 | 91.0 | 126.8 | 166.9 | 264.7 |
| Female..... | 8.9 | 15.4 | 22.6 | 29.2 | 34.8 | 48.0 | 68.1 | 102.6 | 169.2 | 344.1 |

Worktable Orig210R. Death rates for I13 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Hypertensive heart and renal disease (I13)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|-----|-----|-----|------|------|------|------|
| All origins 3/, both sexes..... | 0.2 | 0.4 | 0.5 | 0.9 | 1.2 | 1.6 | 2.8 | 4.9 | 9.5 | 23.5 |
| Male..... | 0.2 | 0.5 | 0.6 | 1.1 | 1.4 | 1.5 | 3.4 | 5.1 | 9.9 | 23.1 |
| Female..... | 0.2 | 0.2 | 0.4 | 0.7 | 1.0 | 1.7 | 2.4 | 4.7 | 9.2 | 23.7 |
| Hispanic, both sexes..... | * | * | * | * | * | * | * | 6.7 | 11.6 | 22.7 |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | 21.6 |
| Mexican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 0.2 | 0.4 | 0.5 | 0.9 | 1.1 | 1.6 | 2.8 | 4.8 | 9.4 | 23.5 |
| Male..... | 0.2 | 0.5 | 0.6 | 1.1 | 1.3 | 1.5 | 3.2 | 5.1 | 9.7 | 22.9 |
| Female..... | 0.2 | 0.2 | 0.4 | 0.7 | 1.0 | 1.6 | 2.4 | 4.5 | 9.1 | 23.7 |
| Non-Hispanic white, both sexes..... | * | * | * | 0.3 | 0.5 | 0.7 | 1.8 | 3.6 | 8.2 | 20.8 |
| Male..... | * | * | * | 0.4 | 0.5 | 0.7 | 2.2 | 4.1 | 8.6 | 20.2 |
| Female..... | * | * | * | * | 0.5 | 0.8 | 1.5 | 3.3 | 8.0 | 21.0 |
| Non-Hispanic black, both sexes..... | 1.1 | 2.3 | 3.3 | 6.1 | 6.3 | 8.1 | 12.5 | 17.6 | 23.2 | 57.2 |
| Male..... | * | 3.1 | 4.7 | 7.8 | 7.5 | 7.4 | 14.0 | 17.4 | 22.6 | 62.4 |
| Female..... | * | 1.5 | 2.2 | 4.7 | 5.4 | 8.5 | 11.5 | 17.7 | 23.5 | 55.2 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|-------------------------------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Ischemic heart diseases (I20-I25) | | | | | | | | | | |
| All origins 3/, both sexes..... | 23.8 | 46.4 | 83.5 | 140.1 | 226.2 | 358.4 | 569.8 | 940.2 | 1,611.2 | 3,775.0 |
| Male..... | 36.5 | 72.3 | 130.3 | 211.4 | 331.4 | 508.5 | 791.5 | 1,274.8 | 2,094.2 | 4,205.3 |
| Female..... | 11.2 | 21.4 | 38.7 | 73.1 | 130.7 | 228.8 | 392.6 | 704.0 | 1,317.5 | 3,588.3 |
| Hispanic, both sexes..... | 11.2 | 26.9 | 52.6 | 99.9 | 184.7 | 293.4 | 493.0 | 805.7 | 1,353.7 | 3,182.9 |
| Male..... | 17.5 | 41.4 | 80.3 | 155.3 | 272.2 | 413.4 | 681.9 | 1,053.3 | 1,662.0 | 3,374.9 |
| Female..... | 4.6 | 12.1 | 25.8 | 49.1 | 108.5 | 195.5 | 347.5 | 628.8 | 1,157.0 | 3,087.8 |
| Mexican, both sexes..... | 11.2 | 24.7 | 53.2 | 95.3 | 180.9 | 298.1 | 541.0 | 729.6 | 1,312.3 | 2,849.6 |
| Male..... | 16.8 | 37.1 | 77.1 | 138.9 | 264.3 | 397.0 | 799.2 | 1,018.9 | 1,519.5 | * |
| Female..... | 4.6 | 11.8 | 27.5 | 53.0 | 109.4 | 207.1 | 360.4 | 539.5 | 1,154.8 | 2,722.9 |
| Puerto Rican, both sexes..... | 19.2 | 40.6 | 62.8 | 144.9 | 245.1 | 423.6 | 593.0 | * | * | * |
| Male..... | * | 61.5 | 87.9 | 234.2 | * | * | * | * | * | * |
| Female..... | * | * | * | * | 151.4 | * | * | * | * | * |
| Cuban, both sexes..... | * | 42.5 | * | 98.1 | 135.6 | 213.7 | 376.2 | 934.7 | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 7.5 | 23.9 | 42.9 | 89.0 | 184.0 | 267.6 | 410.9 | 701.2 | * | * |
| Male..... | 14.1 | 39.9 | 75.2 | 149.6 | 266.5 | * | 496.4 | * | * | * |
| Female..... | * | 8.8 | 17.7 | 31.9 | 103.6 | 156.7 | 336.4 | * | * | * |
| Non-Hispanic 5/, both sexes..... | 25.3 | 48.4 | 85.9 | 142.7 | 228.2 | 361.5 | 572.0 | 943.8 | 1,617.7 | 3,787.0 |
| Male..... | 39.0 | 75.4 | 134.1 | 214.8 | 334.2 | 512.8 | 794.4 | 1,281.4 | 2,107.5 | 4,228.4 |
| Female..... | 12.0 | 22.2 | 39.7 | 74.9 | 131.8 | 230.2 | 393.9 | 705.5 | 1,320.5 | 3,596.5 |
| Non-Hispanic white, both sexes..... | 24.5 | 46.0 | 81.2 | 135.4 | 217.1 | 348.4 | 560.1 | 931.0 | 1,622.0 | 3,848.2 |
| Male..... | 38.5 | 73.5 | 129.4 | 206.7 | 322.1 | 501.2 | 786.3 | 1,278.6 | 2,128.1 | 4,324.5 |
| Female..... | 10.5 | 18.8 | 34.2 | 66.9 | 119.6 | 213.1 | 375.2 | 682.3 | 1,312.3 | 3,642.5 |
| Non-Hispanic black, both sexes..... | 36.1 | 74.4 | 139.0 | 234.9 | 366.7 | 530.0 | 787.4 | 1,206.0 | 1,780.5 | 3,481.1 |
| Male..... | 49.8 | 103.8 | 199.8 | 335.4 | 516.5 | 713.6 | 1,034.1 | 1,508.4 | 2,183.9 | 3,572.8 |
| Female..... | 23.9 | 48.9 | 87.6 | 152.6 | 249.7 | 393.9 | 621.9 | 1,025.5 | 1,570.5 | 3,446.0 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Acute myocardial infarction (I21-I22)

| | | | | | | | | | | |
|-------------------------------------|------|------|------|-------|-------|-------|-------|-------|-------|---------|
| All origins 3/, both sexes..... | 9.9 | 19.0 | 34.9 | 58.4 | 96.2 | 148.5 | 227.5 | 352.5 | 572.7 | 1,191.3 |
| Male..... | 14.8 | 29.2 | 53.7 | 86.6 | 139.0 | 207.6 | 312.0 | 466.1 | 726.6 | 1,366.4 |
| Female..... | 5.1 | 9.1 | 17.0 | 32.0 | 57.4 | 97.6 | 159.8 | 272.2 | 479.1 | 1,115.3 |
| Hispanic, both sexes..... | 4.4 | 9.5 | 20.2 | 40.1 | 75.6 | 112.2 | 188.8 | 297.8 | 475.5 | 1,048.2 |
| Male..... | 6.4 | 14.5 | 29.7 | 60.9 | 108.5 | 155.8 | 255.8 | 383.9 | 595.9 | 1,107.7 |
| Female..... | 2.4 | 4.6 | 11.1 | 20.9 | 47.0 | 76.6 | 137.2 | 236.3 | 398.6 | 1,018.8 |
| Mexican, both sexes..... | 4.4 | 9.6 | 22.6 | 41.5 | 81.9 | 124.1 | 234.5 | 292.1 | 475.5 | 978.9 |
| Male..... | 6.3 | 14.2 | 32.1 | 60.8 | 117.4 | 167.2 | 330.9 | 413.8 | * | * |
| Female..... | * | 4.9 | 12.4 | 22.7 | 51.5 | 84.5 | 167.0 | 212.2 | 422.3 | 938.1 |
| Puerto Rican, both sexes..... | * | 13.1 | 15.6 | 57.8 | 89.8 | 132.9 | 160.0 | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | 41.8 | 73.2 | 149.8 | 348.2 | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 3.1 | 7.4 | 15.7 | 30.1 | 68.8 | 99.2 | 127.5 | 231.5 | * | * |
| Male..... | * | 12.1 | 22.7 | 46.3 | 101.5 | * | * | * | * | * |
| Female..... | * | * | * | * | * | 68.2 | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 10.6 | 20.0 | 36.2 | 59.8 | 97.6 | 150.8 | 229.1 | 354.6 | 575.7 | 1,194.0 |
| Male..... | 15.9 | 30.8 | 55.8 | 88.5 | 141.0 | 210.7 | 314.5 | 469.4 | 731.0 | 1,374.2 |
| Female..... | 5.4 | 9.6 | 17.5 | 32.8 | 58.1 | 98.8 | 160.7 | 273.6 | 481.4 | 1,116.3 |
| Non-Hispanic white, both sexes..... | 10.4 | 19.4 | 35.4 | 58.6 | 95.4 | 148.4 | 226.2 | 351.1 | 575.3 | 1,202.8 |
| Male..... | 15.9 | 30.7 | 55.5 | 88.0 | 140.6 | 210.8 | 315.1 | 469.8 | 734.8 | 1,393.0 |
| Female..... | 5.0 | 8.3 | 15.8 | 30.3 | 53.5 | 93.2 | 153.4 | 266.1 | 477.8 | 1,120.7 |
| Non-Hispanic black, both sexes..... | 14.2 | 28.5 | 51.8 | 86.3 | 138.6 | 197.2 | 299.8 | 440.2 | 648.9 | 1,212.9 |
| Male..... | 19.6 | 38.0 | 72.3 | 117.2 | 180.6 | 252.6 | 375.9 | 539.9 | 783.1 | 1,296.3 |
| Female..... | 9.5 | 20.2 | 34.5 | 60.9 | 105.8 | 156.2 | 248.7 | 380.7 | 579.1 | 1,181.0 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Other acute ischemic heart diseases (I24)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|-----|-----|-----|------|-----|------|------|
| All origins 3/, both sexes..... | 0.3 | 0.6 | 1.2 | 1.4 | 2.2 | 3.0 | 3.9 | 5.7 | 9.0 | 21.1 |
| Male..... | 0.4 | 0.8 | 1.8 | 2.1 | 3.0 | 4.3 | 5.1 | 7.0 | 11.3 | 21.4 |
| Female..... | * | 0.3 | 0.7 | 0.7 | 1.4 | 1.9 | 3.0 | 4.7 | 7.6 | 21.0 |
| Hispanic, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Mexican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 0.3 | 0.6 | 1.3 | 1.5 | 2.2 | 3.2 | 4.1 | 5.8 | 9.3 | 21.8 |
| Male..... | 0.4 | 0.9 | 1.9 | 2.2 | 3.1 | 4.6 | 5.2 | 7.3 | 11.7 | 22.1 |
| Female..... | * | 0.3 | 0.7 | 0.8 | 1.4 | 1.9 | 3.2 | 4.8 | 7.8 | 21.6 |
| Non-Hispanic white, both sexes..... | 0.2 | 0.5 | 1.2 | 1.4 | 2.0 | 3.1 | 3.9 | 5.8 | 9.2 | 22.0 |
| Male..... | 0.4 | 0.8 | 1.7 | 2.2 | 2.9 | 4.4 | 4.8 | 7.3 | 11.5 | 22.7 |
| Female..... | * | 0.3 | 0.7 | 0.8 | 1.2 | 1.9 | 3.1 | 4.7 | 7.8 | 21.8 |
| Non-Hispanic black, both sexes..... | * | 1.1 | 2.5 | 2.1 | 4.2 | 5.1 | 7.3 | 7.7 | 12.0 | 21.5 |
| Male..... | * | 2.0 | 3.8 | 3.2 | 5.9 | 8.2 | 10.9 | * | 16.4 | 21.9 |
| Female..... | * | * | * | * | * | * | 4.8 | 7.0 | 9.7 | 21.3 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Other forms of chronic ischemic heart disease (I20,I25)

| | | | | | | | | | | |
|-------------------------------------|------|------|-------|-------|-------|-------|-------|-------|---------|---------|
| All origins 3/, both sexes..... | 13.6 | 26.8 | 47.3 | 80.2 | 127.8 | 206.8 | 338.4 | 582.1 | 1,029.5 | 2,562.5 |
| Male..... | 21.4 | 42.2 | 74.9 | 122.7 | 189.4 | 296.6 | 474.3 | 801.6 | 1,356.2 | 2,817.4 |
| Female..... | 6.0 | 11.9 | 21.0 | 40.3 | 72.0 | 129.4 | 229.7 | 427.1 | 830.8 | 2,451.9 |
| Hispanic, both sexes..... | 6.7 | 17.1 | 32.1 | 59.5 | 108.1 | 180.7 | 302.3 | 505.5 | 875.6 | 2,129.7 |
| Male..... | 11.1 | 26.6 | 50.1 | 93.8 | 162.1 | 257.3 | 422.2 | 666.8 | 1,062.6 | 2,262.2 |
| Female..... | 2.2 | 7.4 | 14.6 | 28.0 | 61.1 | 118.1 | 209.9 | 390.3 | 756.3 | 2,064.1 |
| Mexican, both sexes..... | 6.7 | 14.9 | 30.3 | 53.7 | 97.5 | 173.4 | 304.3 | 436.1 | 833.3 | 1,864.2 |
| Male..... | 10.4 | 22.9 | 44.6 | 77.8 | 144.6 | 229.8 | 463.7 | 604.0 | * | * |
| Female..... | * | 6.7 | 14.8 | 30.3 | 57.1 | 121.4 | 192.8 | 325.8 | 731.0 | 1,778.0 |
| Puerto Rican, both sexes..... | 10.4 | 26.6 | 46.7 | 87.1 | 154.4 | 290.6 | 433.0 | * | * | * |
| Male..... | * | 39.7 | 64.5 | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | 68.4 | 93.8 | 140.5 | 225.3 | 584.7 | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 4.4 | 16.1 | 26.9 | 57.7 | 115.2 | 166.9 | 280.7 | 465.7 | * | * |
| Male..... | 8.8 | 27.0 | 52.0 | 101.7 | 165.0 | * | * | * | * | * |
| Female..... | * | * | * | * | 66.7 | 87.3 | 237.0 | * | * | * |
| Non-Hispanic 5/, both sexes..... | 14.5 | 27.8 | 48.4 | 81.4 | 128.4 | 207.5 | 338.9 | 583.4 | 1,032.8 | 2,571.2 |
| Male..... | 22.7 | 43.7 | 76.5 | 124.1 | 190.0 | 297.4 | 474.7 | 804.7 | 1,364.8 | 2,832.2 |
| Female..... | 6.4 | 12.3 | 21.5 | 41.2 | 72.3 | 129.5 | 230.0 | 427.1 | 831.3 | 2,458.6 |
| Non-Hispanic white, both sexes..... | 13.8 | 26.0 | 44.7 | 75.4 | 119.7 | 196.9 | 330.1 | 574.2 | 1,037.5 | 2,623.4 |
| Male..... | 22.2 | 42.1 | 72.2 | 116.6 | 178.6 | 285.9 | 466.3 | 801.5 | 1,381.9 | 2,908.8 |
| Female..... | 5.5 | 10.2 | 17.7 | 35.8 | 64.9 | 118.0 | 218.6 | 411.5 | 826.8 | 2,500.1 |
| Non-Hispanic black, both sexes..... | 21.2 | 44.8 | 84.7 | 146.5 | 223.9 | 327.7 | 480.3 | 758.1 | 1,119.5 | 2,246.7 |
| Male..... | 29.9 | 63.8 | 123.8 | 215.0 | 329.9 | 452.8 | 647.2 | 959.6 | 1,384.5 | 2,254.6 |
| Female..... | 13.6 | 28.4 | 51.7 | 90.4 | 141.0 | 234.9 | 368.4 | 637.8 | 981.7 | 2,243.7 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Atherosclerotic cardiovascular disease, so described (I25.0)

| | | | | | | | | | | |
|-------------------------------------|------|------|------|------|-------|-------|-------|-------|-------|-------|
| All origins 3/, both sexes..... | 4.8 | 9.8 | 17.5 | 27.9 | 39.7 | 53.9 | 76.5 | 116.6 | 187.6 | 474.5 |
| Male..... | 7.8 | 15.6 | 28.4 | 44.8 | 59.7 | 78.8 | 106.7 | 154.5 | 232.4 | 491.8 |
| Female..... | 1.8 | 4.1 | 7.1 | 12.1 | 21.4 | 32.4 | 52.5 | 89.7 | 160.4 | 467.1 |
| Hispanic, both sexes..... | 2.9 | 6.8 | 11.8 | 22.7 | 39.2 | 49.8 | 75.4 | 105.3 | 155.5 | 358.6 |
| Male..... | 4.9 | 10.7 | 20.3 | 40.4 | 61.8 | 74.8 | 111.5 | 140.1 | 178.4 | 415.0 |
| Female..... | * | 2.8 | 3.5 | 6.4 | 19.5 | 29.3 | 47.7 | 80.4 | 140.9 | 330.7 |
| Mexican, both sexes..... | 2.9 | 5.3 | 11.9 | 21.2 | 35.1 | 42.0 | 77.0 | 97.8 | 172.0 | 364.4 |
| Male..... | 4.4 | 8.5 | 19.7 | 36.2 | 55.9 | 57.8 | 123.5 | 130.2 | * | * |
| Female..... | * | * | * | 6.7 | 17.3 | 27.4 | 44.5 | 76.5 | 160.4 | 321.7 |
| Puerto Rican, both sexes..... | * | * | * | 27.8 | 45.8 | 82.1 | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | 41.5 | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | 8.1 | 11.8 | 24.8 | 56.7 | 57.3 | 85.9 | 103.0 | * | * |
| Male..... | * | 13.3 | 23.9 | 45.5 | 79.5 | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 5.0 | 10.0 | 17.8 | 28.1 | 39.2 | 53.7 | 75.9 | 116.2 | 188.3 | 477.8 |
| Male..... | 8.2 | 16.0 | 28.7 | 44.5 | 58.7 | 78.4 | 105.3 | 153.9 | 233.6 | 493.3 |
| Female..... | 2.0 | 4.2 | 7.3 | 12.6 | 21.4 | 32.3 | 52.3 | 89.7 | 160.7 | 471.1 |
| Non-Hispanic white, both sexes..... | 4.6 | 9.3 | 15.8 | 24.6 | 34.4 | 46.6 | 68.5 | 107.0 | 180.7 | 473.5 |
| Male..... | 7.7 | 15.1 | 26.0 | 39.3 | 52.0 | 68.2 | 96.0 | 142.1 | 222.9 | 489.2 |
| Female..... | 1.6 | 3.5 | 5.8 | 10.5 | 18.0 | 27.5 | 46.0 | 81.9 | 154.9 | 466.8 |
| Non-Hispanic black, both sexes..... | 8.3 | 17.3 | 35.9 | 60.8 | 86.1 | 120.6 | 158.1 | 233.1 | 303.8 | 580.4 |
| Male..... | 12.2 | 26.0 | 54.5 | 97.1 | 131.4 | 182.2 | 220.6 | 321.1 | 420.4 | 603.9 |
| Female..... | 5.0 | 9.7 | 20.1 | 31.1 | 50.8 | 75.0 | 116.1 | 180.5 | 243.1 | 571.5 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

| | | | | | | | | | | |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

| | | | | | | | | | | |
|-------------------------------------|------|------|------|-------|-------|-------|-------|-------|---------|---------|
| All origins 3/, both sexes..... | 8.8 | 17.1 | 29.9 | 52.3 | 88.2 | 152.9 | 261.8 | 465.6 | 841.9 | 2,088.0 |
| Male..... | 13.6 | 26.6 | 46.5 | 78.0 | 129.6 | 217.8 | 367.6 | 647.1 | 1,123.8 | 2,325.6 |
| Female..... | 4.2 | 7.8 | 13.9 | 28.2 | 50.5 | 97.0 | 177.3 | 337.3 | 670.4 | 1,984.9 |
| Hispanic, both sexes..... | 3.8 | 10.3 | 20.3 | 36.8 | 68.9 | 130.9 | 226.8 | 400.2 | 720.1 | 1,771.1 |
| Male..... | 6.1 | 15.9 | 29.8 | 53.4 | 100.3 | 182.5 | 310.7 | 526.6 | 884.2 | 1,847.2 |
| Female..... | * | 4.6 | 11.1 | 21.6 | 41.6 | 88.7 | 162.2 | 309.9 | 615.4 | 1,733.3 |
| Mexican, both sexes..... | 3.8 | 9.6 | 18.4 | 32.5 | 62.4 | 131.4 | 227.2 | 338.3 | 661.3 | 1,499.9 |
| Male..... | 6.0 | 14.4 | 25.0 | 41.6 | 88.7 | 172.0 | 340.1 | 473.7 | * | * |
| Female..... | * | 4.8 | 11.3 | 23.6 | 39.8 | 94.0 | 148.3 | 249.3 | 570.6 | 1,456.3 |
| Puerto Rican, both sexes..... | * | 16.4 | 33.9 | 59.3 | 108.6 | 208.5 | 334.1 | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | 72.4 | 98.9 | 186.9 | 494.9 | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | 8.0 | 15.2 | 32.9 | 58.5 | 109.5 | 194.8 | 362.7 | * | * |
| Male..... | * | 13.7 | 28.1 | 56.2 | 85.6 | * | * | * | * | * |
| Female..... | * | * | * | * | * | 61.0 | 169.0 | * | * | * |
| Non-Hispanic 5/, both sexes..... | 9.4 | 17.7 | 30.6 | 53.3 | 89.2 | 153.8 | 263.0 | 467.1 | 844.5 | 2,093.4 |
| Male..... | 14.5 | 27.7 | 47.8 | 79.6 | 131.3 | 219.0 | 369.4 | 650.8 | 1,131.1 | 2,338.9 |
| Female..... | 4.5 | 8.1 | 14.1 | 28.7 | 51.0 | 97.2 | 177.8 | 337.4 | 670.6 | 1,987.5 |
| Non-Hispanic white, both sexes..... | 9.2 | 16.7 | 28.9 | 50.8 | 85.3 | 150.3 | 261.5 | 467.2 | 856.7 | 2,149.8 |
| Male..... | 14.5 | 27.0 | 46.2 | 77.2 | 126.6 | 217.7 | 370.3 | 659.5 | 1,158.9 | 2,419.7 |
| Female..... | 4.0 | 6.7 | 12.0 | 25.3 | 46.9 | 90.6 | 172.6 | 329.6 | 671.9 | 2,033.3 |
| Non-Hispanic black, both sexes..... | 12.9 | 27.6 | 48.9 | 85.7 | 137.7 | 207.0 | 322.3 | 525.0 | 815.8 | 1,666.3 |
| Male..... | 17.7 | 37.8 | 69.4 | 117.8 | 198.6 | 270.6 | 426.6 | 638.5 | 964.1 | 1,650.8 |
| Female..... | 8.7 | 18.7 | 31.6 | 59.4 | 90.2 | 160.0 | 252.3 | 457.2 | 738.6 | 1,672.2 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|-------------------------------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Other heart diseases (I26-I51) | | | | | | | | | | |
| All origins 3/, both sexes..... | 12.4 | 18.1 | 25.9 | 39.4 | 60.6 | 97.9 | 164.5 | 286.7 | 536.3 | 1,443.5 |
| Male..... | 15.9 | 23.5 | 33.6 | 49.4 | 75.7 | 118.8 | 199.8 | 340.7 | 612.4 | 1,448.6 |
| Female..... | 9.0 | 12.8 | 18.5 | 30.1 | 46.8 | 79.8 | 136.3 | 248.6 | 490.0 | 1,441.3 |
| Hispanic, both sexes..... | 6.6 | 9.2 | 14.2 | 21.4 | 31.4 | 57.4 | 91.6 | 167.4 | 307.5 | 740.0 |
| Male..... | 8.6 | 11.2 | 18.4 | 26.6 | 38.7 | 69.8 | 122.0 | 202.2 | 354.6 | 758.0 |
| Female..... | 4.4 | 7.2 | 10.2 | 16.7 | 25.0 | 47.3 | 68.3 | 142.6 | 277.4 | 731.1 |
| Mexican, both sexes..... | 5.7 | 9.5 | 13.6 | 19.0 | 29.5 | 59.2 | 98.3 | 150.4 | 282.9 | 697.2 |
| Male..... | 7.4 | 10.4 | 14.8 | 23.0 | 37.1 | 67.5 | 148.4 | 189.0 | * | * |
| Female..... | 3.6 | 8.5 | 12.2 | 15.1 | 22.9 | 51.5 | 63.2 | 125.0 | 266.4 | 684.1 |
| Puerto Rican, both sexes..... | 14.0 | * | 17.2 | 36.0 | 49.4 | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | 37.6 | 69.5 | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 5.6 | 7.6 | 13.6 | 18.0 | 30.2 | 62.5 | 82.4 | 172.6 | * | * |
| Male..... | 7.8 | 10.9 | 22.1 | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 13.1 | 19.0 | 26.8 | 40.8 | 62.7 | 100.6 | 168.7 | 292.4 | 545.0 | 1,469.2 |
| Male..... | 16.7 | 24.8 | 34.8 | 51.1 | 78.4 | 122.0 | 204.1 | 347.2 | 622.6 | 1,477.3 |
| Female..... | 9.5 | 13.3 | 19.2 | 31.1 | 48.4 | 82.1 | 140.3 | 253.8 | 497.9 | 1,465.7 |
| Non-Hispanic white, both sexes..... | 11.3 | 15.6 | 22.8 | 36.0 | 57.1 | 94.9 | 162.5 | 288.2 | 548.1 | 1,503.1 |
| Male..... | 14.8 | 20.8 | 30.1 | 45.5 | 72.3 | 115.5 | 197.0 | 343.3 | 627.9 | 1,512.9 |
| Female..... | 7.8 | 10.5 | 15.6 | 27.0 | 43.1 | 76.7 | 134.2 | 248.8 | 499.2 | 1,498.8 |
| Non-Hispanic black, both sexes..... | 27.1 | 44.8 | 62.1 | 89.0 | 123.6 | 172.3 | 260.2 | 391.1 | 601.9 | 1,285.5 |
| Male..... | 32.8 | 58.3 | 78.8 | 112.9 | 153.6 | 211.9 | 322.9 | 457.7 | 685.6 | 1,258.0 |
| Female..... | 22.2 | 33.1 | 48.1 | 69.3 | 100.2 | 142.9 | 218.2 | 351.3 | 558.4 | 1,296.0 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|-------------------------------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Heart failure (I50) | | | | | | | | | | |
| All origins 3/, both sexes..... | 0.8 | 1.7 | 3.4 | 6.0 | 12.4 | 22.9 | 45.7 | 89.0 | 194.0 | 645.0 |
| Male..... | 1.0 | 2.0 | 4.4 | 7.2 | 14.8 | 27.0 | 53.2 | 103.9 | 218.0 | 621.4 |
| Female..... | 0.5 | 1.3 | 2.4 | 4.9 | 10.1 | 19.3 | 39.8 | 78.4 | 179.3 | 655.3 |
| Hispanic, both sexes..... | * | * | 1.8 | 3.1 | 6.7 | 14.2 | 27.1 | 53.3 | 111.8 | 316.0 |
| Male..... | * | * | * | 4.0 | 8.2 | 17.3 | 30.7 | 64.0 | 124.7 | 296.2 |
| Female..... | * | * | * | * | 5.4 | 11.6 | 24.3 | 45.7 | 103.7 | 325.8 |
| Mexican, both sexes..... | * | * | * | * | 6.3 | 17.2 | 35.7 | 50.7 | 122.1 | 328.5 |
| Male..... | * | * | * | * | * | 20.1 | 47.9 | 70.3 | * | * |
| Female..... | * | * | * | * | * | 27.1 | 37.9 | 110.5 | 325.1 | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 0.8 | 1.8 | 3.5 | 6.3 | 12.8 | 23.5 | 46.7 | 90.7 | 197.0 | 656.8 |
| Male..... | 1.1 | 2.2 | 4.6 | 7.5 | 15.3 | 27.7 | 54.4 | 105.9 | 221.4 | 634.7 |
| Female..... | 0.6 | 1.4 | 2.5 | 5.1 | 10.5 | 19.8 | 40.6 | 79.9 | 182.2 | 666.4 |
| Non-Hispanic white, both sexes..... | 0.6 | 1.2 | 2.8 | 5.5 | 11.7 | 22.2 | 45.1 | 89.8 | 198.8 | 677.3 |
| Male..... | 0.9 | 1.6 | 3.7 | 6.6 | 14.0 | 26.1 | 52.5 | 105.5 | 224.9 | 656.2 |
| Female..... | 0.4 | 0.9 | 1.9 | 4.4 | 9.6 | 18.9 | 39.0 | 78.6 | 182.8 | 686.4 |
| Non-Hispanic black, both sexes..... | 2.3 | 5.6 | 9.4 | 14.1 | 25.3 | 39.3 | 72.8 | 119.4 | 217.6 | 534.5 |
| Male..... | 2.8 | 6.7 | 12.3 | 16.6 | 32.2 | 48.6 | 88.9 | 138.0 | 240.2 | 489.0 |
| Female..... | 1.8 | 4.7 | 7.0 | 12.1 | 19.9 | 32.4 | 61.9 | 108.2 | 205.8 | 551.8 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

| | | | | | | | | | | |
|-------------------------------------|------|------|------|------|-------|-------|-------|-------|-------|-------|
| All origins 3/, both sexes..... | 11.2 | 15.7 | 21.7 | 32.5 | 46.9 | 73.3 | 116.3 | 194.4 | 338.2 | 793.7 |
| Male..... | 14.2 | 20.6 | 28.1 | 41.1 | 59.4 | 89.7 | 143.8 | 233.1 | 389.4 | 821.3 |
| Female..... | 8.2 | 11.0 | 15.6 | 24.4 | 35.6 | 59.2 | 94.3 | 167.1 | 307.1 | 781.7 |
| Hispanic, both sexes..... | 6.0 | 7.8 | 11.9 | 17.2 | 23.2 | 41.7 | 62.4 | 111.4 | 192.1 | 419.6 |
| Male..... | 7.9 | 9.8 | 15.4 | 20.9 | 28.7 | 50.5 | 88.6 | 136.3 | 225.3 | 455.1 |
| Female..... | 4.0 | 5.8 | 8.5 | 13.8 | 18.3 | 34.6 | 42.3 | 93.7 | 170.8 | 402.0 |
| Mexican, both sexes..... | 5.0 | 8.2 | 11.1 | 15.5 | 21.9 | 40.5 | 60.7 | 97.8 | 155.6 | 366.5 |
| Male..... | 6.7 | 9.1 | 12.0 | 17.1 | 25.8 | 46.2 | 97.7 | 116.4 | * | * |
| Female..... | 3.1 | 7.2 | 10.3 | 14.0 | 18.5 | 35.3 | 34.8 | 85.6 | 152.9 | 357.3 |
| Puerto Rican, both sexes..... | 13.6 | * | * | 25.5 | 31.4 | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | 57.5 | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 5.6 | 6.2 | 11.5 | 15.2 | 23.5 | 52.2 | 59.3 | 112.4 | * | * |
| Male..... | 7.8 | 9.8 | 19.7 | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 11.8 | 16.6 | 22.6 | 33.6 | 48.7 | 75.4 | 119.4 | 198.4 | 343.8 | 807.6 |
| Male..... | 14.9 | 21.8 | 29.2 | 42.6 | 61.6 | 92.2 | 146.8 | 237.6 | 396.1 | 836.7 |
| Female..... | 8.7 | 11.5 | 16.2 | 25.1 | 36.9 | 60.9 | 97.4 | 170.8 | 312.1 | 795.0 |
| Non-Hispanic white, both sexes..... | 10.2 | 13.9 | 19.4 | 29.9 | 44.2 | 71.1 | 114.9 | 195.2 | 345.0 | 820.9 |
| Male..... | 13.2 | 18.5 | 25.6 | 38.1 | 56.8 | 87.4 | 141.7 | 234.0 | 397.9 | 850.7 |
| Female..... | 7.1 | 9.3 | 13.3 | 21.9 | 32.6 | 56.6 | 93.1 | 167.4 | 312.7 | 808.1 |
| Non-Hispanic black, both sexes..... | 24.0 | 37.5 | 50.6 | 72.2 | 96.2 | 129.8 | 183.6 | 266.8 | 381.5 | 747.1 |
| Male..... | 28.8 | 49.1 | 64.1 | 92.7 | 119.1 | 159.5 | 230.2 | 316.4 | 440.7 | 763.6 |
| Female..... | 19.8 | 27.5 | 39.3 | 55.3 | 78.3 | 107.8 | 152.3 | 237.2 | 350.6 | 740.8 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|------|------|------|------|------|-------|-------|-------|
| All origins 3/, both sexes..... | 1.0 | 1.8 | 2.9 | 4.5 | 7.3 | 11.8 | 20.7 | 34.9 | 66.7 | 180.4 |
| Male..... | 1.3 | 2.3 | 3.5 | 5.5 | 8.2 | 12.7 | 22.2 | 35.9 | 64.8 | 153.4 |
| Female..... | 0.7 | 1.3 | 2.3 | 3.7 | 6.5 | 11.0 | 19.5 | 34.2 | 67.9 | 192.1 |
| Hispanic, both sexes..... | * | * | 1.4 | 4.4 | 8.7 | 10.0 | 22.3 | 34.7 | 68.2 | 148.6 |
| Male..... | * | * | * | 5.5 | 9.2 | 8.8 | 23.7 | 39.0 | 59.5 | 132.2 |
| Female..... | * | * | * | 3.4 | 8.3 | 11.1 | 21.3 | 31.5 | 73.7 | 156.7 |
| Mexican, both sexes..... | * | * | * | 3.7 | 10.4 | 9.9 | 26.9 | 37.0 | 87.7 | 154.4 |
| Male..... | * | * | * | * | 11.7 | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | 27.7 | 29.5 | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 1.0 | 1.9 | 3.0 | 4.5 | 7.2 | 11.8 | 20.5 | 34.9 | 66.5 | 181.1 |
| Male..... | 1.3 | 2.4 | 3.7 | 5.4 | 8.1 | 12.8 | 22.0 | 35.6 | 64.8 | 153.9 |
| Female..... | 0.7 | 1.4 | 2.3 | 3.7 | 6.3 | 10.9 | 19.3 | 34.3 | 67.5 | 192.8 |
| Non-Hispanic white, both sexes..... | 0.6 | 1.1 | 2.0 | 2.9 | 5.1 | 8.9 | 16.3 | 29.3 | 59.7 | 170.6 |
| Male..... | 0.8 | 1.4 | 2.6 | 3.6 | 6.1 | 9.8 | 17.7 | 30.2 | 57.3 | 145.4 |
| Female..... | 0.3 | 0.7 | 1.4 | 2.2 | 4.2 | 8.0 | 15.1 | 28.7 | 61.1 | 181.4 |
| Non-Hispanic black, both sexes..... | 3.9 | 7.6 | 11.1 | 18.8 | 24.8 | 36.9 | 62.6 | 98.1 | 153.2 | 312.4 |
| Male..... | 4.9 | 9.5 | 13.4 | 22.2 | 26.3 | 40.7 | 68.4 | 107.0 | 166.9 | 276.8 |
| Female..... | 3.1 | 6.0 | 9.1 | 16.0 | 23.7 | 34.2 | 58.7 | 92.8 | 146.1 | 326.1 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Cerebrovascular diseases (I60-I69)

| | | | | | | | | | | |
|-------------------------------------|------|------|------|------|-------|-------|-------|-------|-------|---------|
| All origins 3/, both sexes..... | 7.1 | 12.3 | 18.3 | 28.6 | 48.3 | 83.8 | 160.7 | 308.8 | 601.7 | 1,445.9 |
| Male..... | 6.9 | 13.4 | 20.4 | 32.7 | 55.8 | 97.0 | 178.7 | 334.8 | 615.6 | 1,317.9 |
| Female..... | 7.2 | 11.2 | 16.3 | 24.8 | 41.4 | 72.4 | 146.2 | 290.4 | 593.3 | 1,501.5 |
| Hispanic, both sexes..... | 6.1 | 13.0 | 18.4 | 28.2 | 46.2 | 70.3 | 143.8 | 240.2 | 453.2 | 898.0 |
| Male..... | 7.1 | 16.2 | 22.0 | 36.0 | 57.7 | 86.7 | 173.8 | 269.4 | 462.1 | 856.7 |
| Female..... | 5.1 | 9.7 | 14.9 | 21.1 | 36.2 | 57.0 | 120.7 | 219.4 | 447.5 | 918.5 |
| Mexican, both sexes..... | 5.6 | 13.3 | 19.8 | 26.7 | 50.5 | 76.9 | 178.3 | 247.3 | 507.4 | 916.9 |
| Male..... | 5.8 | 16.7 | 22.7 | 33.8 | 62.9 | 87.5 | 228.6 | 293.9 | * | * |
| Female..... | 5.4 | 9.8 | 16.6 | 19.8 | 39.8 | 67.2 | 143.1 | 216.7 | 522.1 | 946.6 |
| Puerto Rican, both sexes..... | * | 13.1 | 15.6 | 45.1 | 50.3 | 96.4 | 131.8 | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | 69.5 | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 5.9 | 12.5 | 17.3 | 24.8 | 42.2 | 67.6 | 124.9 | 236.9 | * | * |
| Male..... | 8.4 | 15.6 | 22.7 | 28.9 | * | * | * | * | * | * |
| Female..... | * | 9.6 | 13.0 | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 7.2 | 12.2 | 18.3 | 28.6 | 48.3 | 84.6 | 161.4 | 311.7 | 606.9 | 1,465.0 |
| Male..... | 6.9 | 13.0 | 20.2 | 32.3 | 55.4 | 97.5 | 178.7 | 337.3 | 621.1 | 1,336.0 |
| Female..... | 7.5 | 11.3 | 16.4 | 25.1 | 41.7 | 73.4 | 147.5 | 293.5 | 598.3 | 1,520.7 |
| Non-Hispanic white, both sexes..... | 5.4 | 9.0 | 13.4 | 23.3 | 40.4 | 74.2 | 149.9 | 300.9 | 600.4 | 1,479.2 |
| Male..... | 5.2 | 9.8 | 14.6 | 25.9 | 45.8 | 84.7 | 166.1 | 325.1 | 615.2 | 1,345.9 |
| Female..... | 5.5 | 8.1 | 12.2 | 20.7 | 35.3 | 64.9 | 136.6 | 283.5 | 591.4 | 1,536.7 |
| Non-Hispanic black, both sexes..... | 18.0 | 32.7 | 52.4 | 72.4 | 112.1 | 172.6 | 276.8 | 452.2 | 721.0 | 1,398.0 |
| Male..... | 17.4 | 34.8 | 63.0 | 90.8 | 141.7 | 217.1 | 334.6 | 508.9 | 738.6 | 1,266.8 |
| Female..... | 18.4 | 30.9 | 43.4 | 57.4 | 88.9 | 139.7 | 238.1 | 418.3 | 711.8 | 1,448.1 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Atherosclerosis (I70)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|-----|-----|-----|------|------|------|-------|
| All origins 3/, both sexes..... | 0.1 | 0.3 | 0.6 | 1.4 | 2.4 | 5.2 | 9.3 | 20.5 | 47.8 | 168.1 |
| Male..... | 0.2 | 0.4 | 0.8 | 1.7 | 3.2 | 6.4 | 11.5 | 24.4 | 52.5 | 156.6 |
| Female..... | * | * | 0.5 | 1.0 | 1.7 | 4.1 | 7.5 | 17.8 | 45.0 | 173.1 |
| Hispanic, both sexes..... | * | * | * | * | * | 3.7 | 5.0 | 15.5 | 34.8 | 108.6 |
| Male..... | * | * | * | * | * | * | * | 18.6 | 41.2 | 102.1 |
| Female..... | * | * | * | * | * | * | * | 13.3 | 30.7 | 111.9 |
| Mexican, both sexes..... | * | * | * | * | * | * | * | 16.9 | 30.1 | 112.0 |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 0.2 | 0.3 | 0.7 | 1.4 | 2.5 | 5.3 | 9.5 | 20.7 | 48.3 | 170.2 |
| Male..... | 0.2 | 0.4 | 0.9 | 1.8 | 3.4 | 6.5 | 11.8 | 24.6 | 52.9 | 158.7 |
| Female..... | * | * | 0.5 | 1.1 | 1.7 | 4.2 | 7.7 | 17.9 | 45.4 | 175.1 |
| Non-Hispanic white, both sexes..... | 0.2 | 0.3 | 0.6 | 1.4 | 2.3 | 5.4 | 9.4 | 20.5 | 48.8 | 177.0 |
| Male..... | * | 0.4 | 0.7 | 1.7 | 3.1 | 6.7 | 11.6 | 24.2 | 53.5 | 165.7 |
| Female..... | * | * | 0.4 | 1.0 | 1.6 | 4.3 | 7.6 | 17.9 | 45.9 | 181.9 |
| Non-Hispanic black, both sexes..... | * | * | 1.5 | 2.0 | 4.3 | 5.3 | 11.7 | 27.0 | 52.1 | 118.0 |
| Male..... | * | * | 2.4 | * | 6.5 | 6.4 | 15.4 | 35.7 | 60.1 | 109.4 |
| Female..... | * | * | * | * | * | 4.6 | 9.2 | 21.9 | 47.9 | 121.2 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Other diseases of circulatory system (I71-I78)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|------|------|------|------|------|-------|-------|
| All origins 3/, both sexes..... | 1.3 | 2.2 | 3.1 | 6.2 | 11.7 | 22.3 | 36.5 | 57.8 | 83.8 | 148.4 |
| Male..... | 1.8 | 3.2 | 4.4 | 8.6 | 16.3 | 31.7 | 49.2 | 78.3 | 113.7 | 187.4 |
| Female..... | 0.7 | 1.4 | 2.0 | 3.9 | 7.5 | 14.2 | 26.4 | 43.4 | 65.6 | 131.5 |
| Hispanic, both sexes..... | 1.1 | 1.4 | 2.1 | 4.2 | 8.2 | 11.1 | 19.5 | 34.7 | 51.7 | 92.6 |
| Male..... | 1.6 | * | 2.8 | 4.8 | 10.3 | 14.9 | 26.3 | 44.8 | 77.8 | 102.1 |
| Female..... | * | * | * | 3.7 | 6.5 | 8.0 | 14.2 | 27.4 | 35.0 | 87.9 |
| Mexican, both sexes..... | * | * | * | * | 6.7 | 11.7 | 20.1 | 25.1 | 52.5 | 92.4 |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 1.3 | 2.3 | 3.2 | 6.3 | 11.9 | 23.0 | 37.5 | 58.8 | 85.1 | 150.3 |
| Male..... | 1.9 | 3.3 | 4.5 | 8.9 | 16.8 | 32.8 | 50.5 | 79.9 | 115.1 | 190.7 |
| Female..... | 0.7 | 1.4 | 2.0 | 4.0 | 7.5 | 14.6 | 27.1 | 44.0 | 66.9 | 132.8 |
| Non-Hispanic white, both sexes..... | 1.1 | 1.9 | 2.8 | 5.9 | 11.6 | 23.2 | 38.1 | 59.8 | 86.1 | 150.9 |
| Male..... | 1.6 | 2.8 | 3.9 | 8.2 | 16.5 | 33.7 | 51.2 | 82.2 | 117.6 | 194.6 |
| Female..... | 0.6 | 1.0 | 1.7 | 3.7 | 7.2 | 13.9 | 27.3 | 43.7 | 66.9 | 132.1 |
| Non-Hispanic black, both sexes..... | 2.9 | 5.4 | 6.8 | 11.3 | 16.6 | 26.2 | 38.6 | 59.0 | 84.6 | 157.6 |
| Male..... | 4.3 | 6.9 | 9.9 | 16.6 | 23.4 | 31.2 | 49.9 | 65.7 | 100.6 | 168.5 |
| Female..... | 1.6 | 4.2 | 4.2 | 7.0 | 11.3 | 22.4 | 31.0 | 54.9 | 76.3 | 153.4 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|---|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Aortic aneurysm and dissection (I71) | | | | | | | | | | |
| All origins 3/, both sexes..... | 0.9 | 1.5 | 2.0 | 4.0 | 7.4 | 14.7 | 23.9 | 37.5 | 51.5 | 69.4 |
| Male..... | 1.4 | 2.4 | 3.2 | 6.3 | 11.5 | 23.0 | 34.9 | 55.0 | 74.5 | 106.6 |
| Female..... | 0.3 | 0.7 | 1.0 | 1.8 | 3.7 | 7.5 | 15.1 | 25.1 | 37.6 | 53.3 |
| Hispanic, both sexes..... | * | * | * | 2.7 | 5.3 | 6.4 | 9.0 | 19.2 | 24.5 | 31.0 |
| Male..... | * | * | * | 3.9 | 7.7 | 9.5 | 15.4 | 25.6 | 43.5 | 43.5 |
| Female..... | * | * | * | * | * | * | * | 14.6 | * | 24.9 |
| Mexican, both sexes..... | * | * | * | * | * | * | * | 11.9 | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 0.9 | 1.6 | 2.1 | 4.1 | 7.6 | 15.2 | 24.8 | 38.3 | 52.6 | 70.8 |
| Male..... | 1.5 | 2.5 | 3.3 | 6.5 | 11.7 | 23.8 | 36.1 | 56.4 | 75.7 | 109.1 |
| Female..... | 0.3 | 0.8 | 1.0 | 1.8 | 3.8 | 7.7 | 15.8 | 25.5 | 38.6 | 54.3 |
| Non-Hispanic white, both sexes..... | 0.7 | 1.3 | 1.8 | 3.8 | 7.7 | 16.0 | 25.8 | 39.9 | 54.3 | 72.8 |
| Male..... | 1.2 | 2.0 | 2.8 | 6.1 | 11.9 | 25.3 | 37.4 | 59.1 | 79.1 | 114.7 |
| Female..... | 0.3 | 0.6 | 0.8 | 1.7 | 3.8 | 7.7 | 16.3 | 26.2 | 39.2 | 54.7 |
| Non-Hispanic black, both sexes..... | 2.1 | 3.9 | 4.8 | 6.8 | 8.3 | 11.7 | 18.8 | 25.1 | 37.9 | 50.5 |
| Male..... | 3.7 | 5.8 | 7.7 | 11.2 | 13.2 | 14.8 | 25.3 | 31.0 | 41.3 | 49.2 |
| Female..... | * | 2.2 | 2.3 | 3.3 | 4.5 | 9.3 | 14.5 | 21.6 | 36.1 | 51.0 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Other diseases of arteries, arterioles and capillaries (I72-I78)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|-----|------|------|------|------|------|-------|
| All origins 3/, both sexes..... | 0.4 | 0.7 | 1.1 | 2.2 | 4.3 | 7.7 | 12.6 | 20.4 | 32.3 | 79.0 |
| Male..... | 0.4 | 0.8 | 1.2 | 2.3 | 4.9 | 8.7 | 14.2 | 23.4 | 39.2 | 80.8 |
| Female..... | 0.4 | 0.6 | 1.0 | 2.1 | 3.7 | 6.7 | 11.3 | 18.3 | 28.1 | 78.2 |
| Hispanic, both sexes..... | * | * | * | * | 3.0 | 4.7 | 10.5 | 15.5 | 27.2 | 61.5 |
| Male..... | * | * | * | * | * | * | 11.0 | 19.2 | 34.3 | 58.6 |
| Female..... | * | * | * | * | * | * | 10.1 | 12.8 | 22.6 | 63.0 |
| Mexican, both sexes..... | * | * | * | * | * | * | 11.8 | 13.3 | 31.8 | 69.6 |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 0.4 | 0.7 | 1.1 | 2.3 | 4.3 | 7.8 | 12.7 | 20.6 | 32.5 | 79.5 |
| Male..... | 0.4 | 0.8 | 1.2 | 2.4 | 5.0 | 9.0 | 14.4 | 23.5 | 39.3 | 81.7 |
| Female..... | 0.4 | 0.6 | 1.0 | 2.1 | 3.7 | 6.9 | 11.3 | 18.5 | 28.3 | 78.5 |
| Non-Hispanic white, both sexes..... | 0.3 | 0.6 | 1.0 | 2.1 | 3.9 | 7.2 | 12.3 | 19.8 | 31.8 | 78.2 |
| Male..... | 0.3 | 0.8 | 1.1 | 2.1 | 4.5 | 8.4 | 13.8 | 23.1 | 38.5 | 79.9 |
| Female..... | 0.3 | 0.5 | 1.0 | 2.0 | 3.4 | 6.2 | 11.0 | 17.5 | 27.7 | 77.4 |
| Non-Hispanic black, both sexes..... | 0.7 | 1.6 | 2.0 | 4.5 | 8.3 | 14.5 | 19.8 | 33.9 | 46.7 | 107.1 |
| Male..... | * | * | 2.1 | 5.4 | 10.2 | 16.4 | 24.6 | 34.7 | 59.3 | 119.2 |
| Female..... | * | 2.0 | 1.9 | 3.8 | 6.8 | 13.1 | 16.5 | 33.4 | 40.2 | 102.4 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Other disorders of circulatory system (I80-I99)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|-----|-----|-----|-----|------|------|------|
| All origins 3/, both sexes..... | 0.8 | 1.2 | 1.5 | 2.0 | 2.5 | 3.4 | 4.3 | 7.4 | 13.2 | 24.4 |
| Male..... | 0.8 | 1.3 | 1.6 | 2.3 | 2.6 | 3.6 | 4.2 | 7.0 | 13.3 | 20.5 |
| Female..... | 0.8 | 1.1 | 1.4 | 1.7 | 2.5 | 3.2 | 4.4 | 7.7 | 13.1 | 26.2 |
| Hispanic, both sexes..... | * | * | * | * | * | * | 4.6 | * | 8.9 | 18.3 |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | 23.2 |
| Mexican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 0.8 | 1.2 | 1.5 | 2.1 | 2.6 | 3.5 | 4.3 | 7.6 | 13.3 | 24.7 |
| Male..... | 0.8 | 1.4 | 1.6 | 2.4 | 2.6 | 3.6 | 4.3 | 7.1 | 13.5 | 21.1 |
| Female..... | 0.8 | 1.1 | 1.4 | 1.8 | 2.6 | 3.3 | 4.3 | 7.9 | 13.2 | 26.2 |
| Non-Hispanic white, both sexes..... | 0.7 | 1.1 | 1.4 | 1.9 | 2.2 | 3.2 | 4.1 | 7.3 | 12.8 | 24.5 |
| Male..... | 0.6 | 1.2 | 1.5 | 2.1 | 2.3 | 3.3 | 4.0 | 6.8 | 12.5 | 20.5 |
| Female..... | 0.7 | 1.0 | 1.3 | 1.7 | 2.2 | 3.2 | 4.1 | 7.6 | 12.9 | 26.3 |
| Non-Hispanic black, both sexes..... | 1.9 | 2.4 | 2.9 | 4.3 | 5.9 | 6.4 | 7.4 | 13.0 | 22.7 | 29.9 |
| Male..... | 2.1 | 3.1 | 2.9 | 5.5 | 5.7 | 7.7 | 7.9 | 12.2 | 28.9 | 31.7 |
| Female..... | 1.6 | 1.8 | 3.0 | 3.3 | 6.0 | 5.5 | 7.1 | 13.5 | 19.5 | 29.3 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Influenza and pneumonia (J10-J18)

| | | | | | | | | | | |
|-------------------------------------|-----|------|------|------|------|------|------|-------|-------|-------|
| All origins 3/, both sexes..... | 2.6 | 4.2 | 5.5 | 8.5 | 14.8 | 25.3 | 50.9 | 106.5 | 227.3 | 696.6 |
| Male..... | 3.1 | 4.9 | 6.7 | 10.2 | 18.0 | 31.2 | 63.9 | 133.9 | 287.7 | 805.8 |
| Female..... | 2.2 | 3.4 | 4.3 | 6.8 | 11.8 | 20.1 | 40.6 | 87.2 | 190.6 | 649.2 |
| Hispanic, both sexes..... | 1.5 | 2.9 | 4.0 | 8.3 | 13.4 | 22.8 | 47.5 | 102.6 | 209.4 | 566.0 |
| Male..... | 2.1 | 3.9 | 5.8 | 11.0 | 18.0 | 29.8 | 61.9 | 120.9 | 266.5 | 655.9 |
| Female..... | * | 1.9 | * | 5.9 | 9.4 | 17.1 | 36.5 | 89.6 | 173.0 | 521.4 |
| Mexican, both sexes..... | 1.3 | 2.6 | 3.6 | 6.9 | 12.3 | 21.9 | 55.8 | 92.3 | 209.0 | 549.3 |
| Male..... | * | 3.3 | 5.5 | 8.4 | 15.0 | 27.4 | 76.5 | 129.1 | * | * |
| Female..... | * | * | * | * | 10.1 | 16.8 | 41.3 | 68.2 | 192.2 | 501.2 |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | 17.5 | 27.2 | 33.6 | 104.4 | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 2.8 | 4.3 | 5.6 | 8.5 | 14.9 | 25.3 | 51.0 | 106.3 | 227.3 | 700.2 |
| Male..... | 3.2 | 5.0 | 6.8 | 10.1 | 18.0 | 31.1 | 63.7 | 134.2 | 287.4 | 810.7 |
| Female..... | 2.4 | 3.5 | 4.5 | 6.9 | 12.0 | 20.4 | 40.8 | 86.7 | 190.7 | 652.6 |
| Non-Hispanic white, both sexes..... | 2.4 | 3.9 | 5.0 | 7.6 | 13.8 | 24.5 | 49.8 | 105.0 | 227.7 | 712.7 |
| Male..... | 2.8 | 4.4 | 6.0 | 8.8 | 16.3 | 29.6 | 62.0 | 131.3 | 286.1 | 820.6 |
| Female..... | 2.1 | 3.4 | 4.1 | 6.3 | 11.5 | 20.0 | 39.8 | 86.1 | 192.0 | 666.1 |
| Non-Hispanic black, both sexes..... | 5.3 | 7.7 | 10.3 | 17.6 | 26.8 | 35.6 | 70.0 | 124.3 | 233.8 | 579.8 |
| Male..... | 6.2 | 10.1 | 13.2 | 24.2 | 36.7 | 48.1 | 91.0 | 175.1 | 324.5 | 695.8 |
| Female..... | 4.5 | 5.5 | 7.9 | 12.1 | 19.1 | 26.4 | 56.0 | 93.9 | 186.7 | 535.5 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|-------------------------------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
| Pneumonia (J12-J18) | | | | | | | | | | |
| All origins 3/, both sexes..... | 2.6 | 4.1 | 5.4 | 8.4 | 14.6 | 25.1 | 50.5 | 105.7 | 225.1 | 687.8 |
| Male..... | 3.0 | 4.9 | 6.7 | 10.1 | 17.8 | 30.9 | 63.4 | 133.0 | 285.0 | 796.9 |
| Female..... | 2.2 | 3.4 | 4.3 | 6.8 | 11.8 | 20.0 | 40.2 | 86.5 | 188.6 | 640.5 |
| Hispanic, both sexes..... | 1.5 | 2.9 | 4.0 | 8.3 | 13.4 | 22.7 | 47.4 | 102.1 | 209.0 | 563.2 |
| Male..... | 2.1 | 3.8 | 5.8 | 11.0 | 18.0 | 29.5 | 61.9 | 120.9 | 265.4 | 654.2 |
| Female..... | * | 1.9 | * | 5.9 | 9.4 | 17.1 | 36.2 | 88.7 | 173.0 | 518.1 |
| Mexican, both sexes..... | 1.3 | 2.6 | 3.6 | 6.9 | 12.3 | 21.9 | 55.8 | 91.4 | 209.0 | 548.2 |
| Male..... | * | 3.3 | 5.5 | 8.4 | 15.0 | 27.4 | 76.5 | 129.1 | * | * |
| Female..... | * | * | * | * | 10.1 | 16.8 | 41.3 | 66.7 | 192.2 | 499.5 |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | 17.5 | 26.5 | 33.6 | 104.4 | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 2.7 | 4.2 | 5.5 | 8.4 | 14.7 | 25.1 | 50.5 | 105.5 | 225.0 | 691.2 |
| Male..... | 3.1 | 4.9 | 6.7 | 10.0 | 17.7 | 30.8 | 63.1 | 133.3 | 284.7 | 801.5 |
| Female..... | 2.3 | 3.5 | 4.4 | 6.8 | 11.9 | 20.2 | 40.4 | 86.0 | 188.7 | 643.7 |
| Non-Hispanic white, both sexes..... | 2.4 | 3.8 | 5.0 | 7.5 | 13.7 | 24.3 | 49.3 | 104.1 | 225.2 | 703.1 |
| Male..... | 2.8 | 4.3 | 5.9 | 8.7 | 16.1 | 29.4 | 61.4 | 130.3 | 283.2 | 810.6 |
| Female..... | 2.1 | 3.4 | 4.0 | 6.2 | 11.4 | 19.9 | 39.4 | 85.4 | 189.8 | 656.6 |
| Non-Hispanic black, both sexes..... | 5.3 | 7.7 | 10.3 | 17.5 | 26.5 | 35.2 | 69.8 | 124.1 | 233.0 | 577.4 |
| Male..... | 6.2 | 10.1 | 13.2 | 24.1 | 36.3 | 47.4 | 91.0 | 175.1 | 322.9 | 693.6 |
| Female..... | 4.5 | 5.5 | 7.9 | 12.1 | 18.9 | 26.2 | 55.5 | 93.6 | 186.3 | 533.0 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Chronic lower respiratory diseases (J40-J47)

| | | | | | | | | | | |
|-------------------------------------|-----|------|------|------|------|-------|-------|-------|-------|-------|
| All origins 3/, both sexes..... | 2.9 | 5.8 | 12.0 | 28.2 | 60.8 | 119.0 | 211.5 | 330.1 | 465.7 | 637.6 |
| Male..... | 3.0 | 5.9 | 12.8 | 29.5 | 65.9 | 131.5 | 246.0 | 405.6 | 595.9 | 894.8 |
| Female..... | 2.8 | 5.6 | 11.2 | 26.9 | 56.1 | 108.3 | 183.8 | 276.8 | 386.5 | 526.0 |
| Hispanic, both sexes..... | 1.7 | 2.4 | 4.9 | 9.0 | 20.6 | 38.4 | 73.1 | 136.0 | 242.9 | 457.3 |
| Male..... | * | 2.7 | 5.4 | 11.7 | 25.7 | 51.5 | 92.2 | 199.7 | 321.4 | 602.4 |
| Female..... | 2.0 | 2.1 | 4.4 | 6.4 | 16.1 | 27.6 | 58.5 | 90.5 | 192.7 | 385.5 |
| Mexican, both sexes..... | * | * | 4.2 | 7.7 | 14.7 | 33.8 | 69.4 | 117.5 | 218.4 | 418.7 |
| Male..... | * | * | * | 9.6 | 19.3 | 44.4 | 94.9 | 187.9 | * | * |
| Female..... | * | * | * | * | 10.9 | 24.1 | 51.6 | 71.2 | 165.0 | 320.0 |
| Puerto Rican, both sexes..... | * | * | * | * | 34.1 | 73.0 | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | 71.9 | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | 10.8 | 28.4 | 38.2 | 84.1 | 155.2 | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 3.0 | 6.1 | 12.6 | 29.6 | 63.8 | 124.6 | 219.7 | 339.4 | 474.4 | 643.5 |
| Male..... | 3.1 | 6.2 | 13.5 | 30.9 | 68.8 | 136.7 | 254.8 | 415.0 | 606.7 | 906.0 |
| Female..... | 2.9 | 6.0 | 11.7 | 28.5 | 59.2 | 114.1 | 191.6 | 286.1 | 394.1 | 530.3 |
| Non-Hispanic white, both sexes..... | 2.6 | 5.8 | 12.4 | 30.6 | 67.2 | 133.4 | 233.0 | 357.7 | 496.4 | 673.1 |
| Male..... | 2.7 | 5.9 | 13.1 | 31.1 | 70.6 | 143.1 | 264.3 | 428.0 | 624.9 | 935.4 |
| Female..... | 2.5 | 5.8 | 11.7 | 30.2 | 64.0 | 124.8 | 207.4 | 307.4 | 417.7 | 559.8 |
| Non-Hispanic black, both sexes..... | 6.5 | 9.3 | 17.6 | 31.4 | 54.8 | 87.4 | 149.7 | 217.7 | 300.6 | 370.8 |
| Male..... | 6.9 | 10.0 | 20.3 | 38.4 | 72.7 | 118.0 | 216.2 | 345.1 | 479.7 | 653.1 |
| Female..... | 6.2 | 8.7 | 15.4 | 25.7 | 40.7 | 64.7 | 105.1 | 141.6 | 207.4 | 263.0 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Emphysema (J43)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|-----|------|------|------|------|------|------|
| All origins 3/, both sexes..... | 0.3 | 0.7 | 1.6 | 4.3 | 9.1 | 17.4 | 31.3 | 43.2 | 54.7 | 59.3 |
| Male..... | 0.5 | 0.9 | 1.9 | 4.6 | 10.7 | 19.7 | 37.8 | 56.1 | 74.9 | 90.0 |
| Female..... | 0.2 | 0.6 | 1.3 | 4.0 | 7.7 | 15.4 | 26.1 | 34.0 | 42.4 | 46.0 |
| Hispanic, both sexes..... | * | * | * | * | 2.4 | 3.8 | 10.3 | 10.9 | 32.1 | 36.6 |
| Male..... | * | * | * | * | * | 6.8 | 12.3 | 17.3 | 43.5 | 53.5 |
| Female..... | * | * | * | * | * | * | 8.8 | * | 24.8 | 28.2 |
| Mexican, both sexes..... | * | * | * | * | * | * | * | * | 32.7 | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 0.4 | 0.8 | 1.7 | 4.6 | 9.6 | 18.3 | 32.6 | 44.8 | 55.6 | 60.1 |
| Male..... | 0.5 | 1.0 | 2.0 | 4.9 | 11.3 | 20.6 | 39.3 | 58.0 | 76.3 | 91.3 |
| Female..... | 0.2 | 0.6 | 1.4 | 4.3 | 8.2 | 16.4 | 27.2 | 35.5 | 43.1 | 46.6 |
| Non-Hispanic white, both sexes..... | 0.3 | 0.8 | 1.8 | 4.9 | 10.4 | 19.8 | 34.8 | 47.7 | 58.4 | 63.1 |
| Male..... | 0.4 | 1.0 | 2.0 | 5.1 | 11.7 | 21.2 | 41.1 | 60.3 | 78.9 | 94.3 |
| Female..... | * | 0.6 | 1.6 | 4.7 | 9.2 | 18.6 | 29.6 | 38.8 | 45.9 | 49.7 |
| Non-Hispanic black, both sexes..... | 0.8 | 1.1 | 1.5 | 3.4 | 6.1 | 12.1 | 21.4 | 23.7 | 35.0 | 32.1 |
| Male..... | * | * | 2.4 | 4.6 | 9.6 | 20.7 | 31.5 | 42.3 | 61.6 | 65.6 |
| Female..... | * | * | * | * | 3.3 | 5.7 | 14.7 | 12.6 | 21.1 | 19.2 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Asthma (J45-J46)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|-----|-----|-----|------|-----|-----|------|
| All origins 3/, both sexes..... | 1.2 | 1.4 | 1.6 | 1.9 | 2.1 | 2.7 | 3.8 | 5.4 | 7.7 | 14.7 |
| Male..... | 0.9 | 1.1 | 1.3 | 1.5 | 1.7 | 2.0 | 2.8 | 4.3 | 6.2 | 10.9 |
| Female..... | 1.4 | 1.8 | 1.9 | 2.3 | 2.5 | 3.3 | 4.5 | 6.2 | 8.7 | 16.3 |
| Hispanic, both sexes..... | 0.9 | 1.2 | * | * | 2.9 | 3.8 | 3.8 | 5.9 | 8.9 | 12.2 |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | 5.8 | * | * | * | * |
| Mexican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 1.2 | 1.4 | 1.7 | 2.0 | 2.1 | 2.6 | 3.7 | 5.4 | 7.7 | 14.7 |
| Male..... | 1.0 | 1.0 | 1.3 | 1.5 | 1.6 | 2.0 | 2.8 | 4.1 | 6.2 | 11.1 |
| Female..... | 1.4 | 1.8 | 2.0 | 2.4 | 2.5 | 3.1 | 4.5 | 6.2 | 8.5 | 16.3 |
| Non-Hispanic white, both sexes..... | 0.7 | 1.0 | 1.1 | 1.4 | 1.7 | 2.0 | 3.1 | 5.1 | 7.4 | 14.7 |
| Male..... | 0.6 | 0.7 | 0.9 | 1.0 | 1.2 | 1.5 | 2.3 | 3.6 | 5.8 | 10.8 |
| Female..... | 0.9 | 1.4 | 1.3 | 1.7 | 2.0 | 2.5 | 3.8 | 6.1 | 8.4 | 16.4 |
| Non-Hispanic black, both sexes..... | 4.1 | 4.2 | 5.8 | 7.3 | 5.7 | 6.2 | 10.3 | 7.2 | 9.1 | 11.8 |
| Male..... | 3.8 | 3.4 | 4.7 | 6.1 | 5.5 | 6.4 | 8.9 | * | * | * |
| Female..... | 4.3 | 4.8 | 6.8 | 8.2 | 5.9 | 6.1 | 11.2 | 7.8 | 9.7 | 12.5 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Other chronic lower respiratory diseases (J44,J47)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|------|------|------|-------|-------|-------|-------|-------|
| All origins 3/, both sexes..... | 1.4 | 3.5 | 8.7 | 21.8 | 49.3 | 98.4 | 175.4 | 279.8 | 400.3 | 555.5 |
| Male..... | 1.6 | 3.8 | 9.5 | 23.3 | 53.2 | 109.1 | 204.3 | 343.0 | 511.9 | 785.1 |
| Female..... | 1.3 | 3.3 | 7.9 | 20.4 | 45.7 | 89.3 | 152.3 | 235.2 | 332.4 | 455.8 |
| Hispanic, both sexes..... | * | 1.1 | 3.3 | 6.9 | 15.1 | 30.4 | 59.0 | 117.3 | 198.7 | 397.4 |
| Male..... | * | * | 3.7 | 9.9 | 19.8 | 43.0 | 77.2 | 175.3 | 268.8 | 535.4 |
| Female..... | * | * | 2.9 | 4.2 | 11.0 | 20.2 | 45.0 | 75.9 | 154.0 | 329.1 |
| Mexican, both sexes..... | * | * | 2.6 | 5.9 | 11.3 | 28.0 | 57.7 | 101.5 | 177.1 | 356.7 |
| Male..... | * | * | * | 8.4 | 15.5 | 37.7 | 82.0 | 163.7 | * | * |
| Female..... | * | * | * | * | * | 19.0 | 40.6 | 60.6 | 131.7 | 262.5 |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | 57.5 | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | 22.3 | 31.6 | 66.4 | 133.8 | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 1.5 | 3.8 | 9.1 | 22.9 | 51.8 | 103.1 | 182.4 | 287.5 | 408.2 | 560.8 |
| Male..... | 1.6 | 4.0 | 10.0 | 24.3 | 55.6 | 113.4 | 211.5 | 350.7 | 521.4 | 794.9 |
| Female..... | 1.3 | 3.5 | 8.3 | 21.7 | 48.3 | 94.3 | 159.0 | 243.0 | 339.5 | 459.8 |
| Non-Hispanic white, both sexes..... | 1.5 | 3.9 | 9.4 | 24.2 | 54.9 | 111.0 | 194.0 | 303.0 | 427.6 | 586.9 |
| Male..... | 1.7 | 4.1 | 10.1 | 24.9 | 57.4 | 119.7 | 219.6 | 361.6 | 537.3 | 821.0 |
| Female..... | 1.4 | 3.8 | 8.8 | 23.6 | 52.6 | 103.3 | 173.0 | 261.1 | 360.4 | 485.7 |
| Non-Hispanic black, both sexes..... | 1.6 | 3.9 | 10.2 | 20.5 | 42.8 | 68.7 | 117.7 | 186.1 | 254.1 | 322.7 |
| Male..... | 1.8 | 4.9 | 12.9 | 27.5 | 57.4 | 90.1 | 175.5 | 295.3 | 408.7 | 572.1 |
| Female..... | 1.5 | 3.1 | 7.9 | 14.8 | 31.3 | 52.8 | 78.9 | 120.8 | 173.7 | 227.4 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Pneumonitis due to solids and liquids (J69)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|-----|-----|------|------|------|------|-------|
| All origins 3/, both sexes..... | 0.5 | 0.7 | 1.1 | 1.8 | 3.4 | 6.5 | 13.4 | 31.3 | 67.3 | 186.0 |
| Male..... | 0.7 | 0.9 | 1.5 | 2.2 | 4.2 | 8.6 | 18.2 | 43.4 | 95.1 | 268.5 |
| Female..... | 0.4 | 0.5 | 0.8 | 1.4 | 2.8 | 4.8 | 9.6 | 22.8 | 50.4 | 150.2 |
| Hispanic, both sexes..... | * | * | * | * | 2.7 | 3.7 | 7.3 | 19.5 | 45.0 | 131.4 |
| Male..... | * | * | * | * | * | * | * | 26.9 | 72.1 | 167.3 |
| Female..... | * | * | * | * | * | * | 6.8 | 14.2 | 27.7 | 113.6 |
| Mexican, both sexes..... | * | * | * | * | * | * | * | 20.1 | 44.7 | 117.5 |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 0.6 | 0.7 | 1.2 | 1.9 | 3.5 | 6.7 | 13.7 | 31.9 | 68.1 | 187.8 |
| Male..... | 0.7 | 0.9 | 1.6 | 2.2 | 4.2 | 8.8 | 18.7 | 44.2 | 95.8 | 272.1 |
| Female..... | 0.4 | 0.5 | 0.8 | 1.5 | 2.9 | 4.9 | 9.7 | 23.2 | 51.4 | 151.4 |
| Non-Hispanic white, both sexes..... | 0.5 | 0.7 | 1.1 | 1.8 | 3.3 | 6.3 | 13.4 | 31.5 | 69.0 | 191.3 |
| Male..... | 0.7 | 0.9 | 1.4 | 2.1 | 4.1 | 8.2 | 18.0 | 43.3 | 97.3 | 278.4 |
| Female..... | 0.4 | 0.5 | 0.7 | 1.5 | 2.6 | 4.6 | 9.6 | 23.0 | 51.8 | 153.7 |
| Non-Hispanic black, both sexes..... | 1.0 | 0.8 | 2.2 | 2.4 | 5.6 | 12.0 | 19.2 | 42.7 | 68.3 | 167.3 |
| Male..... | * | * | 3.2 | 3.7 | 6.5 | 17.1 | 29.1 | 63.4 | 92.0 | 237.4 |
| Female..... | * | * | * | * | 4.9 | 8.2 | 12.6 | 30.3 | 56.0 | 140.5 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|------|------|------|------|------|-------|-------|
| All origins 3/, both sexes..... | 1.5 | 2.5 | 4.1 | 7.0 | 11.9 | 20.6 | 37.6 | 56.8 | 83.7 | 133.5 |
| Male..... | 1.6 | 2.5 | 4.3 | 7.9 | 13.9 | 24.4 | 47.8 | 70.5 | 104.2 | 156.6 |
| Female..... | 1.4 | 2.4 | 3.9 | 6.2 | 10.2 | 17.3 | 29.4 | 47.0 | 71.1 | 123.4 |
| Hispanic, both sexes..... | 1.0 | 2.2 | 3.4 | 6.2 | 11.1 | 18.7 | 35.3 | 49.3 | 72.6 | 112.0 |
| Male..... | * | 2.2 | 3.2 | 7.9 | 12.6 | 24.4 | 47.8 | 49.9 | 89.2 | 108.8 |
| Female..... | * | 2.1 | 3.6 | 4.7 | 9.8 | 14.1 | 25.7 | 48.9 | 62.0 | 113.6 |
| Mexican, both sexes..... | * | 2.0 | 2.5 | 5.8 | 9.7 | 22.1 | 37.9 | 52.6 | 71.4 | 99.0 |
| Male..... | * | * | * | * | 11.7 | 26.7 | 47.0 | 53.0 | * | * |
| Female..... | * | * | * | * | * | 17.9 | 31.6 | 52.3 | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | 45.2 | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 1.5 | 2.5 | 4.1 | 7.1 | 11.9 | 20.6 | 37.7 | 57.1 | 84.0 | 134.1 |
| Male..... | 1.7 | 2.5 | 4.3 | 7.8 | 13.9 | 24.3 | 47.7 | 71.6 | 104.7 | 158.5 |
| Female..... | 1.4 | 2.4 | 3.9 | 6.4 | 10.1 | 17.4 | 29.6 | 46.9 | 71.4 | 123.6 |
| Non-Hispanic white, both sexes..... | 1.4 | 2.0 | 3.8 | 6.7 | 11.6 | 21.0 | 38.2 | 59.0 | 85.8 | 138.6 |
| Male..... | 1.4 | 2.1 | 4.0 | 7.5 | 13.4 | 25.2 | 48.1 | 73.6 | 107.1 | 164.1 |
| Female..... | 1.3 | 2.0 | 3.6 | 5.9 | 9.9 | 17.2 | 30.2 | 48.5 | 72.8 | 127.6 |
| Non-Hispanic black, both sexes..... | 2.9 | 5.6 | 6.9 | 11.1 | 17.2 | 21.2 | 34.5 | 41.4 | 68.9 | 90.7 |
| Male..... | 3.6 | 5.8 | 7.5 | 12.3 | 21.8 | 22.8 | 45.5 | 54.5 | 88.1 | 102.8 |
| Female..... | 2.3 | 5.4 | 6.3 | 10.2 | 13.5 | 20.1 | 27.1 | 33.6 | 58.8 | 86.1 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Peptic ulcer (K25-K28)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|-----|-----|-----|-----|------|------|------|
| All origins 3/, both sexes..... | 0.3 | 0.6 | 0.8 | 1.4 | 1.7 | 2.7 | 4.7 | 8.0 | 13.2 | 28.3 |
| Male..... | 0.4 | 0.7 | 1.2 | 1.8 | 2.1 | 3.2 | 6.1 | 9.3 | 16.7 | 31.6 |
| Female..... | 0.3 | 0.4 | 0.5 | 1.0 | 1.4 | 2.2 | 3.6 | 7.0 | 11.0 | 26.9 |
| Hispanic, both sexes..... | * | * | * | * | * | 3.2 | * | 7.2 | 11.6 | 20.5 |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | 17.4 |
| Mexican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 0.3 | 0.6 | 0.8 | 1.4 | 1.7 | 2.6 | 4.8 | 8.0 | 13.2 | 28.6 |
| Male..... | 0.4 | 0.6 | 1.2 | 1.7 | 2.0 | 3.0 | 6.1 | 9.3 | 16.8 | 31.7 |
| Female..... | 0.3 | 0.5 | 0.5 | 1.0 | 1.4 | 2.2 | 3.7 | 7.1 | 11.0 | 27.3 |
| Non-Hispanic white, both sexes..... | 0.3 | 0.5 | 0.8 | 1.2 | 1.6 | 2.5 | 4.8 | 7.9 | 13.7 | 29.4 |
| Male..... | 0.3 | 0.6 | 1.0 | 1.5 | 1.9 | 2.9 | 6.0 | 9.2 | 17.3 | 32.6 |
| Female..... | * | 0.5 | 0.5 | 1.0 | 1.3 | 2.2 | 3.9 | 7.1 | 11.5 | 27.9 |
| Non-Hispanic black, both sexes..... | 0.9 | 0.9 | 1.6 | 2.4 | 2.9 | 3.3 | 5.4 | 8.8 | 6.7 | 18.8 |
| Male..... | * | * | 2.9 | 3.7 | 4.3 | * | 8.2 | 11.7 | * | * |
| Female..... | * | * | * | * | * | * | * | 7.0 | * | 18.4 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Chronic liver disease and cirrhosis (K70,K73-K74)

| | | | | | | | | | | |
|-------------------------------------|------|------|------|------|------|------|-------|------|------|------|
| All origins 3/, both sexes..... | 9.7 | 16.0 | 20.3 | 21.7 | 24.5 | 27.9 | 31.2 | 32.5 | 29.9 | 21.4 |
| Male..... | 12.9 | 23.3 | 31.0 | 32.0 | 34.5 | 37.2 | 40.0 | 39.4 | 36.9 | 28.4 |
| Female..... | 6.5 | 8.9 | 10.1 | 12.1 | 15.3 | 19.8 | 24.1 | 27.6 | 25.6 | 18.4 |
| Hispanic, both sexes..... | 13.0 | 23.1 | 31.9 | 37.1 | 40.1 | 50.5 | 63.2 | 52.8 | 54.8 | 45.5 |
| Male..... | 20.7 | 38.0 | 52.4 | 61.1 | 58.5 | 76.5 | 79.4 | 60.8 | 70.9 | 45.2 |
| Female..... | 4.8 | 7.9 | 12.1 | 15.0 | 24.2 | 29.3 | 50.7 | 47.1 | 44.5 | 45.6 |
| Mexican, both sexes..... | 15.2 | 25.3 | 36.4 | 41.5 | 45.7 | 62.9 | 83.1 | 58.1 | 55.9 | 52.2 |
| Male..... | 23.1 | 41.0 | 57.4 | 67.4 | 63.4 | 89.4 | 111.5 | 72.6 | * | * |
| Female..... | 5.9 | 9.0 | 13.7 | 16.3 | 30.6 | 38.6 | 63.2 | 48.5 | * | * |
| Puerto Rican, both sexes..... | 13.6 | 29.5 | 38.9 | 47.3 | 52.1 | * | * | * | * | * |
| Male..... | * | 47.3 | 68.0 | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 9.2 | 17.0 | 22.2 | 27.2 | 32.0 | 39.0 | 57.6 | * | * | * |
| Male..... | 16.6 | 30.1 | 38.2 | 42.2 | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 9.2 | 15.1 | 19.1 | 20.3 | 23.1 | 26.1 | 29.0 | 31.3 | 28.7 | 20.4 |
| Male..... | 11.8 | 21.5 | 28.8 | 29.4 | 32.4 | 34.3 | 37.4 | 38.1 | 35.3 | 27.7 |
| Female..... | 6.7 | 9.0 | 9.9 | 11.8 | 14.6 | 19.0 | 22.3 | 26.6 | 24.7 | 17.3 |
| Non-Hispanic white, both sexes..... | 9.4 | 15.0 | 18.7 | 20.1 | 23.1 | 26.9 | 30.2 | 32.9 | 29.6 | 21.2 |
| Male..... | 12.2 | 21.6 | 28.2 | 29.1 | 32.3 | 34.8 | 38.7 | 40.2 | 36.3 | 28.7 |
| Female..... | 6.7 | 8.5 | 9.5 | 11.5 | 14.7 | 19.8 | 23.3 | 27.7 | 25.5 | 18.0 |
| Non-Hispanic black, both sexes..... | 8.2 | 17.8 | 25.3 | 24.9 | 25.8 | 23.2 | 19.9 | 19.7 | 16.3 | 12.1 |
| Male..... | 10.0 | 23.9 | 38.9 | 37.3 | 39.3 | 35.1 | 28.7 | 22.1 | 21.8 | * |
| Female..... | 6.6 | 12.6 | 13.9 | 14.7 | 15.3 | 14.4 | 14.0 | 18.2 | 13.4 | 10.5 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Alcoholic liver disease (K70)

| | | | | | | | | | | |
|-------------------------------------|------|------|------|------|------|------|------|------|-----|-----|
| All origins 3/, both sexes..... | 5.9 | 9.4 | 11.4 | 11.4 | 11.8 | 10.7 | 9.5 | 6.7 | 4.4 | 2.1 |
| Male..... | 8.0 | 13.8 | 17.7 | 17.6 | 19.2 | 17.4 | 15.4 | 10.9 | 8.0 | 4.7 |
| Female..... | 3.9 | 5.1 | 5.4 | 5.6 | 5.1 | 4.9 | 4.8 | 3.7 | 2.2 | 1.0 |
| Hispanic, both sexes..... | 7.9 | 13.6 | 19.3 | 20.1 | 19.5 | 21.2 | 20.4 | 11.7 | * | * |
| Male..... | 13.0 | 22.6 | 33.0 | 36.7 | 35.7 | 41.3 | 39.1 | 19.2 | * | * |
| Female..... | 2.5 | 4.5 | 6.0 | 4.9 | 5.4 | * | * | * | * | * |
| Mexican, both sexes..... | 9.3 | 15.4 | 22.4 | 22.9 | 21.9 | 27.4 | 28.1 | 11.4 | * | * |
| Male..... | 14.3 | 25.2 | 36.7 | 41.3 | 39.9 | 49.2 | 57.2 | * | * | * |
| Female..... | 3.4 | 5.2 | 7.0 | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | 12.1 | 18.9 | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 6.3 | 11.5 | 14.7 | 16.4 | 17.5 | * | * | * | * | * |
| Male..... | 11.6 | 20.7 | 27.5 | 28.1 | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 5.6 | 8.9 | 10.6 | 10.6 | 11.2 | 9.8 | 8.8 | 6.4 | 4.2 | 1.9 |
| Male..... | 7.3 | 12.7 | 16.2 | 15.9 | 17.8 | 15.5 | 13.9 | 10.3 | 7.6 | 4.3 |
| Female..... | 4.1 | 5.1 | 5.3 | 5.6 | 5.1 | 4.9 | 4.7 | 3.6 | 2.2 | 0.9 |
| Non-Hispanic white, both sexes..... | 5.8 | 9.0 | 10.5 | 10.5 | 11.2 | 10.0 | 9.0 | 6.5 | 4.3 | 1.9 |
| Male..... | 7.4 | 13.0 | 16.0 | 15.9 | 17.8 | 15.6 | 14.1 | 10.5 | 7.7 | 4.1 |
| Female..... | 4.2 | 5.0 | 5.1 | 5.3 | 5.0 | 5.0 | 4.8 | 3.6 | 2.1 | 1.0 |
| Non-Hispanic black, both sexes..... | 4.4 | 8.5 | 12.4 | 12.4 | 12.1 | 9.6 | 7.6 | 5.8 | * | * |
| Male..... | 5.8 | 11.9 | 19.1 | 17.2 | 20.4 | 16.4 | 13.3 | 9.9 | * | * |
| Female..... | 3.2 | 5.5 | 6.7 | 8.6 | 5.6 | 4.6 | * | * | * | * |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Other chronic liver disease and cirrhosis (K73-K74)

| | | | | | | | | | | |
|-------------------------------------|-----|------|------|------|------|------|------|------|------|------|
| All origins 3/, both sexes..... | 3.7 | 6.6 | 8.9 | 10.3 | 12.6 | 17.2 | 21.7 | 25.8 | 25.5 | 19.3 |
| Male..... | 4.9 | 9.5 | 13.3 | 14.4 | 15.3 | 19.8 | 24.6 | 28.6 | 28.9 | 23.7 |
| Female..... | 2.6 | 3.8 | 4.7 | 6.5 | 10.2 | 14.9 | 19.4 | 23.9 | 23.4 | 17.4 |
| Hispanic, both sexes..... | 5.1 | 9.5 | 12.6 | 17.0 | 20.7 | 29.4 | 42.8 | 41.1 | 46.3 | 38.8 |
| Male..... | 7.8 | 15.4 | 19.4 | 24.4 | 22.8 | 35.2 | 40.4 | 41.6 | 54.9 | * |
| Female..... | 2.3 | 3.4 | 6.1 | 10.1 | 18.8 | 24.6 | 44.6 | 40.7 | 40.9 | 42.3 |
| Mexican, both sexes..... | 5.9 | 9.9 | 14.0 | 18.6 | 23.8 | 35.5 | 55.0 | 46.6 | 45.6 | 44.6 |
| Male..... | 8.8 | 15.8 | 20.7 | 26.0 | 23.5 | 40.1 | 54.4 | 50.7 | * | * |
| Female..... | * | 3.8 | 6.8 | 11.3 | 24.1 | 31.3 | 55.4 | 43.9 | * | * |
| Puerto Rican, both sexes..... | * | 17.4 | 20.0 | 25.5 | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | 5.5 | 7.6 | 10.8 | * | * | 39.8 | * | * | * |
| Male..... | * | 9.4 | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 3.6 | 6.3 | 8.5 | 9.7 | 11.9 | 16.2 | 20.2 | 25.0 | 24.5 | 18.5 |
| Male..... | 4.5 | 8.7 | 12.7 | 13.6 | 14.6 | 18.7 | 23.4 | 27.8 | 27.7 | 23.4 |
| Female..... | 2.6 | 3.9 | 4.6 | 6.2 | 9.5 | 14.1 | 17.7 | 23.0 | 22.5 | 16.4 |
| Non-Hispanic white, both sexes..... | 3.6 | 6.0 | 8.2 | 9.7 | 11.9 | 16.9 | 21.2 | 26.5 | 25.3 | 19.3 |
| Male..... | 4.8 | 8.6 | 12.2 | 13.2 | 14.4 | 19.2 | 24.6 | 29.7 | 28.6 | 24.7 |
| Female..... | 2.5 | 3.5 | 4.4 | 6.3 | 9.6 | 14.8 | 18.5 | 24.1 | 23.4 | 17.0 |
| Non-Hispanic black, both sexes..... | 3.8 | 9.4 | 13.0 | 12.4 | 13.8 | 13.6 | 12.4 | 13.9 | 12.3 | 9.7 |
| Male..... | 4.2 | 12.1 | 19.8 | 20.1 | 18.9 | 18.7 | 15.4 | 12.2 | * | * |
| Female..... | 3.4 | 7.0 | 7.2 | 6.2 | 9.7 | 9.9 | 10.3 | 14.9 | 11.0 | 9.6 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Cholelithiasis and other disorders of gallbladder (K80-K82)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|-----|-----|-----|-----|-----|------|------|
| All origins 3/, both sexes..... | 0.2 | 0.2 | 0.4 | 0.6 | 1.1 | 2.0 | 3.6 | 5.7 | 10.8 | 22.9 |
| Male..... | 0.2 | 0.3 | 0.5 | 0.8 | 1.1 | 2.3 | 4.0 | 7.5 | 13.5 | 24.1 |
| Female..... | * | * | 0.3 | 0.4 | 1.0 | 1.8 | 3.2 | 4.4 | 9.1 | 22.4 |
| Hispanic, both sexes..... | * | * | * | * | * | * | 4.8 | 7.7 | 11.6 | 24.4 |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | 23.2 |
| Mexican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 0.2 | 0.2 | 0.4 | 0.6 | 1.0 | 2.0 | 3.5 | 5.5 | 10.7 | 22.8 |
| Male..... | * | 0.3 | 0.5 | 0.8 | 1.1 | 2.3 | 3.9 | 7.3 | 13.6 | 23.9 |
| Female..... | * | * | 0.3 | 0.5 | 0.9 | 1.7 | 3.2 | 4.3 | 8.9 | 22.4 |
| Non-Hispanic white, both sexes..... | 0.2 | 0.2 | 0.3 | 0.6 | 0.9 | 2.0 | 3.4 | 5.3 | 10.6 | 23.4 |
| Male..... | * | 0.3 | 0.4 | 0.7 | 1.0 | 2.4 | 3.8 | 7.3 | 13.5 | 24.5 |
| Female..... | * | * | 0.3 | 0.4 | 0.8 | 1.6 | 3.0 | 3.9 | 8.8 | 22.9 |
| Non-Hispanic black, both sexes..... | * | * | * | * | * | 2.2 | 4.7 | 7.7 | 12.5 | 15.1 |
| Male..... | * | * | * | * | * | * | * | * | 16.4 | * |
| Female..... | * | * | * | * | * | * | 5.5 | 7.8 | 10.6 | 16.3 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

| | | | | | | | | | | |
|-------------------------------------|-----|------|------|------|------|------|-------|-------|-------|-------|
| All origins 3/, both sexes..... | 2.1 | 3.7 | 5.9 | 10.0 | 16.8 | 29.4 | 50.0 | 85.4 | 142.2 | 288.6 |
| Male..... | 2.3 | 4.3 | 6.9 | 11.0 | 18.6 | 33.6 | 59.8 | 105.7 | 182.7 | 385.5 |
| Female..... | 1.9 | 3.1 | 4.9 | 9.2 | 15.2 | 25.8 | 42.1 | 71.1 | 117.6 | 246.6 |
| Hispanic, both sexes..... | 1.6 | 2.7 | 6.3 | 10.5 | 17.7 | 32.9 | 40.3 | 74.9 | 111.8 | 201.2 |
| Male..... | 1.9 | 2.8 | 7.6 | 11.7 | 19.8 | 38.9 | 43.4 | 92.1 | 147.6 | 235.9 |
| Female..... | * | 2.7 | 5.1 | 9.3 | 15.9 | 27.9 | 37.9 | 62.6 | 89.1 | 184.0 |
| Mexican, both sexes..... | 1.2 | 3.3 | 6.4 | 11.4 | 17.5 | 37.6 | 50.1 | 80.5 | 120.4 | 183.8 |
| Male..... | * | 3.8 | 6.5 | 11.7 | 18.8 | 36.5 | 47.9 | 98.0 | * | * |
| Female..... | * | * | 6.3 | 11.1 | 16.5 | 38.6 | 51.6 | 68.9 | * | 186.3 |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | 10.4 | 23.5 | 30.1 | 31.9 | 73.6 | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 2.1 | 3.8 | 5.8 | 9.9 | 16.7 | 29.1 | 50.3 | 85.7 | 143.2 | 291.2 |
| Male..... | 2.3 | 4.5 | 6.8 | 10.9 | 18.4 | 33.1 | 60.5 | 106.0 | 183.7 | 391.0 |
| Female..... | 1.9 | 3.1 | 4.9 | 9.1 | 15.0 | 25.6 | 42.2 | 71.3 | 118.6 | 248.2 |
| Non-Hispanic white, both sexes..... | 1.4 | 2.6 | 3.6 | 7.0 | 12.6 | 23.5 | 43.1 | 78.0 | 136.0 | 281.1 |
| Male..... | 1.5 | 3.0 | 4.2 | 7.6 | 13.8 | 27.5 | 53.2 | 99.0 | 178.5 | 382.6 |
| Female..... | 1.3 | 2.1 | 3.1 | 6.4 | 11.6 | 20.1 | 34.8 | 63.0 | 110.0 | 237.3 |
| Non-Hispanic black, both sexes..... | 6.8 | 12.0 | 22.1 | 34.4 | 51.4 | 80.3 | 128.1 | 179.7 | 244.8 | 444.0 |
| Male..... | 7.8 | 15.0 | 27.4 | 41.0 | 62.3 | 90.4 | 150.9 | 208.4 | 284.7 | 568.9 |
| Female..... | 6.0 | 9.4 | 17.5 | 28.9 | 42.8 | 72.9 | 112.9 | 162.6 | 224.0 | 396.3 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Renal failure (N17-N19)

| | | | | | | | | | | |
|-------------------------------------|-----|------|------|------|------|------|-------|-------|-------|-------|
| All origins 3/, both sexes..... | 2.0 | 3.6 | 5.7 | 9.8 | 16.5 | 29.0 | 49.1 | 83.9 | 139.8 | 284.8 |
| Male..... | 2.2 | 4.2 | 6.7 | 10.6 | 18.4 | 33.1 | 58.8 | 103.9 | 180.0 | 380.7 |
| Female..... | 1.8 | 2.9 | 4.8 | 9.0 | 14.9 | 25.4 | 41.4 | 69.9 | 115.3 | 243.2 |
| Hispanic, both sexes..... | 1.5 | 2.7 | 6.3 | 10.1 | 17.3 | 32.3 | 40.1 | 73.0 | 110.1 | 198.4 |
| Male..... | 1.9 | 2.8 | 7.6 | 11.0 | 19.2 | 38.6 | 43.4 | 88.9 | 145.3 | 232.6 |
| Female..... | * | 2.5 | 5.0 | 9.3 | 15.7 | 27.1 | 37.5 | 61.7 | 87.6 | 181.5 |
| Mexican, both sexes..... | 1.2 | 3.1 | 6.3 | 11.2 | 17.3 | 36.4 | 49.7 | 78.2 | 118.7 | 181.6 |
| Male..... | * | 3.8 | 6.5 | 11.4 | 18.8 | 35.9 | 47.9 | 93.4 | * | * |
| Female..... | * | * | 6.1 | 11.1 | 16.1 | 36.9 | 50.9 | 68.2 | * | 182.9 |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | 22.3 | 30.1 | 31.9 | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 2.1 | 3.6 | 5.6 | 9.7 | 16.4 | 28.6 | 49.5 | 84.2 | 140.7 | 287.4 |
| Male..... | 2.2 | 4.3 | 6.6 | 10.6 | 18.2 | 32.6 | 59.4 | 104.1 | 181.0 | 386.2 |
| Female..... | 1.9 | 3.0 | 4.7 | 8.8 | 14.7 | 25.2 | 41.6 | 70.1 | 116.3 | 244.8 |
| Non-Hispanic white, both sexes..... | 1.3 | 2.4 | 3.5 | 6.8 | 12.4 | 23.1 | 42.2 | 76.7 | 133.7 | 277.6 |
| Male..... | 1.4 | 2.9 | 4.1 | 7.3 | 13.5 | 26.9 | 52.1 | 97.4 | 175.9 | 377.8 |
| Female..... | 1.2 | 2.0 | 2.9 | 6.2 | 11.3 | 19.6 | 34.0 | 61.8 | 107.9 | 234.3 |
| Non-Hispanic black, both sexes..... | 6.8 | 11.6 | 21.5 | 33.9 | 51.1 | 79.9 | 127.3 | 177.1 | 241.6 | 436.5 |
| Male..... | 7.6 | 14.8 | 26.7 | 40.4 | 62.3 | 89.8 | 149.5 | 204.2 | 281.6 | 562.3 |
| Female..... | 6.0 | 8.8 | 17.0 | 28.7 | 42.3 | 72.5 | 112.4 | 160.9 | 220.8 | 388.4 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

| | | | | | | | | | | |
|-------------------------------------|------|------|------|------|------|------|------|------|------|-------|
| All origins 3/, both sexes..... | 6.0 | 6.4 | 6.9 | 7.0 | 8.3 | 10.4 | 16.1 | 28.3 | 53.3 | 208.0 |
| Male..... | 7.6 | 8.5 | 9.0 | 9.0 | 11.2 | 12.7 | 19.7 | 32.9 | 60.6 | 175.5 |
| Female..... | 4.5 | 4.4 | 5.0 | 5.0 | 5.6 | 8.4 | 13.3 | 25.1 | 48.9 | 222.1 |
| Hispanic, both sexes..... | 3.6 | 3.1 | 5.0 | 3.9 | 6.1 | 5.5 | 9.0 | 11.7 | 30.3 | 91.5 |
| Male..... | 4.4 | 3.8 | 5.8 | 4.8 | 8.5 | 7.8 | 11.8 | 15.4 | 37.7 | 88.7 |
| Female..... | 2.8 | 2.4 | 4.2 | * | * | * | 6.8 | 9.1 | 25.5 | 92.8 |
| Mexican, both sexes..... | 2.9 | 2.4 | 3.2 | 3.4 | * | * | 9.9 | 11.4 | 31.0 | 92.4 |
| Male..... | 2.9 | * | * | * | * | * | * | * | * | * |
| Female..... | 3.0 | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 3.8 | * | 6.0 | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 6.3 | 6.7 | 7.0 | 7.2 | 8.3 | 10.7 | 16.6 | 29.1 | 54.1 | 211.9 |
| Male..... | 8.0 | 9.0 | 9.2 | 9.2 | 11.2 | 12.9 | 20.1 | 33.7 | 61.1 | 178.6 |
| Female..... | 4.6 | 4.5 | 5.0 | 5.2 | 5.7 | 8.8 | 13.7 | 25.9 | 49.8 | 226.3 |
| Non-Hispanic white, both sexes..... | 6.0 | 6.1 | 6.1 | 6.6 | 7.9 | 10.1 | 16.3 | 28.7 | 54.1 | 215.8 |
| Male..... | 7.6 | 8.1 | 7.8 | 8.3 | 10.6 | 12.1 | 19.7 | 32.3 | 60.6 | 181.5 |
| Female..... | 4.4 | 4.1 | 4.4 | 4.9 | 5.4 | 8.3 | 13.6 | 26.1 | 50.2 | 230.6 |
| Non-Hispanic black, both sexes..... | 9.8 | 12.2 | 15.9 | 13.9 | 13.8 | 18.0 | 23.3 | 39.7 | 65.4 | 198.1 |
| Male..... | 12.7 | 17.3 | 22.4 | 19.8 | 20.4 | 22.8 | 31.1 | 57.3 | 87.4 | 182.7 |
| Female..... | 7.2 | 7.9 | 10.4 | 9.1 | 8.8 | 14.4 | 18.1 | 29.2 | 54.0 | 204.0 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

| | | | | | | | | | | |
|-------------------------------------|------|------|------|-------|-------|-------|-------|-------|-------|---------|
| All origins 3/, both sexes..... | 23.5 | 33.6 | 42.6 | 56.2 | 77.1 | 110.4 | 173.8 | 297.8 | 555.5 | 1,493.0 |
| Male..... | 28.2 | 41.0 | 52.4 | 64.4 | 87.9 | 122.8 | 186.8 | 309.6 | 555.2 | 1,262.9 |
| Female..... | 18.9 | 26.4 | 33.2 | 48.6 | 67.3 | 99.6 | 163.5 | 289.5 | 555.6 | 1,592.8 |
| Hispanic, both sexes..... | 17.0 | 25.6 | 30.1 | 43.3 | 55.6 | 90.9 | 136.7 | 212.2 | 372.5 | 870.8 |
| Male..... | 23.1 | 34.0 | 37.5 | 51.9 | 62.6 | 109.0 | 147.9 | 233.6 | 391.2 | 808.2 |
| Female..... | 10.4 | 17.0 | 22.9 | 35.4 | 49.4 | 76.0 | 128.1 | 197.0 | 360.6 | 901.9 |
| Mexican, both sexes..... | 15.7 | 25.7 | 29.3 | 39.3 | 50.7 | 101.1 | 140.7 | 205.7 | 369.8 | 803.8 |
| Male..... | 20.0 | 34.4 | 34.7 | 44.6 | 55.9 | 110.0 | 160.4 | 243.2 | * | * |
| Female..... | 10.8 | 16.7 | 23.6 | 34.0 | 46.2 | 92.9 | 127.0 | 181.1 | 372.3 | 833.1 |
| Puerto Rican, both sexes..... | 31.1 | 36.7 | 34.4 | 78.8 | 75.4 | 97.7 | 174.1 | * | * | * |
| Male..... | 45.7 | 42.6 | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | 46.5 | 97.1 | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 16.1 | 23.1 | 30.9 | 40.9 | 70.0 | 94.1 | 135.5 | 231.5 | * | * |
| Male..... | 25.9 | 31.6 | 40.0 | 47.9 | 79.5 | * | * | * | * | * |
| Female..... | 7.5 | 15.1 | 23.7 | 34.2 | 60.7 | 74.2 | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 24.3 | 34.3 | 43.4 | 57.0 | 78.4 | 111.5 | 175.8 | 301.5 | 561.8 | 1,514.9 |
| Male..... | 28.7 | 41.5 | 53.4 | 65.0 | 89.3 | 123.3 | 188.7 | 312.9 | 561.0 | 1,280.6 |
| Female..... | 19.9 | 27.3 | 33.9 | 49.5 | 68.5 | 101.3 | 165.4 | 293.4 | 562.3 | 1,615.9 |
| Non-Hispanic white, both sexes..... | 22.1 | 31.2 | 39.8 | 53.6 | 74.5 | 108.5 | 171.7 | 299.2 | 562.5 | 1,540.5 |
| Male..... | 27.0 | 38.3 | 49.3 | 60.8 | 84.6 | 118.5 | 182.6 | 307.9 | 558.6 | 1,293.4 |
| Female..... | 17.1 | 24.2 | 30.4 | 46.7 | 65.1 | 99.6 | 162.8 | 293.0 | 564.9 | 1,647.2 |
| Non-Hispanic black, both sexes..... | 42.8 | 61.9 | 79.5 | 97.9 | 126.0 | 158.0 | 246.8 | 373.5 | 648.9 | 1,443.4 |
| Male..... | 45.2 | 72.8 | 96.4 | 118.3 | 148.7 | 188.1 | 292.8 | 421.6 | 699.6 | 1,331.3 |
| Female..... | 40.7 | 52.5 | 65.3 | 81.1 | 108.4 | 135.7 | 215.9 | 344.8 | 622.5 | 1,486.2 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

| | | | | | | | | | | |
|-------------------------------------|------|------|------|------|------|------|------|-------|-------|-------|
| All origins 3/, both sexes..... | 39.4 | 38.7 | 34.3 | 30.4 | 32.6 | 38.3 | 50.8 | 80.0 | 131.1 | 275.4 |
| Male..... | 56.8 | 56.6 | 49.6 | 44.8 | 45.7 | 53.4 | 68.9 | 106.4 | 170.1 | 346.6 |
| Female..... | 22.4 | 21.3 | 19.6 | 16.8 | 20.7 | 25.3 | 36.3 | 61.4 | 107.3 | 244.6 |
| Hispanic, both sexes..... | 35.6 | 33.0 | 30.5 | 27.3 | 30.5 | 34.7 | 45.4 | 69.9 | 90.0 | 153.0 |
| Male..... | 55.3 | 50.0 | 47.0 | 43.1 | 44.1 | 56.2 | 61.9 | 97.3 | 121.2 | 224.2 |
| Female..... | 14.5 | 15.8 | 14.6 | 12.8 | 18.6 | 17.1 | 32.8 | 50.3 | 70.1 | 117.7 |
| Mexican, both sexes..... | 34.7 | 33.0 | 30.8 | 27.3 | 32.1 | 34.7 | 49.7 | 64.9 | 95.5 | 149.0 |
| Male..... | 53.7 | 49.1 | 45.0 | 42.5 | 50.2 | 57.1 | 73.7 | 99.1 | * | * |
| Female..... | 12.5 | 16.2 | 15.5 | 12.5 | 16.5 | * | 32.9 | 42.4 | * | * |
| Puerto Rican, both sexes..... | 47.1 | 46.9 | 33.9 | 25.5 | * | * | * | * | * | * |
| Male..... | 74.4 | 68.1 | 51.6 | * | * | * | * | * | * | * |
| Female..... | 22.7 | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | 30.8 | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 34.1 | 27.3 | 29.3 | 32.1 | 32.0 | 41.9 | 47.8 | 79.0 | * | * |
| Male..... | 55.3 | 44.9 | 50.8 | 46.3 | * | * | * | * | * | * |
| Female..... | 15.2 | 10.7 | 12.6 | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 39.7 | 39.1 | 34.4 | 30.4 | 32.6 | 38.4 | 50.9 | 80.3 | 132.4 | 279.8 |
| Male..... | 56.7 | 56.9 | 49.5 | 44.5 | 45.5 | 52.8 | 68.9 | 106.5 | 171.3 | 351.2 |
| Female..... | 23.2 | 21.8 | 20.0 | 17.2 | 20.9 | 25.9 | 36.4 | 61.7 | 108.8 | 249.0 |
| Non-Hispanic white, both sexes..... | 40.8 | 38.8 | 33.5 | 29.4 | 32.0 | 37.3 | 51.0 | 81.2 | 136.1 | 290.9 |
| Male..... | 57.9 | 56.6 | 47.3 | 42.2 | 44.1 | 50.1 | 68.3 | 106.5 | 174.1 | 361.6 |
| Female..... | 23.8 | 21.1 | 20.1 | 17.2 | 20.8 | 25.9 | 36.8 | 63.2 | 112.8 | 260.3 |
| Non-Hispanic black, both sexes..... | 40.8 | 49.1 | 45.9 | 42.7 | 40.8 | 47.4 | 52.6 | 76.0 | 97.7 | 175.4 |
| Male..... | 59.1 | 73.0 | 73.6 | 72.1 | 65.4 | 75.5 | 79.4 | 119.2 | 151.3 | 262.5 |
| Female..... | 24.6 | 28.4 | 22.4 | 18.6 | 21.6 | 26.6 | 34.6 | 50.2 | 69.8 | 142.1 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Transport accidents (V01-V99,Y85)

| | | | | | | | | | | |
|-------------------------------------|------|------|------|------|------|------|------|------|------|------|
| All origins 3/, both sexes..... | 17.1 | 16.9 | 15.8 | 14.9 | 16.7 | 16.7 | 20.2 | 25.0 | 29.7 | 28.8 |
| Male..... | 24.5 | 25.1 | 22.7 | 22.2 | 22.8 | 23.4 | 27.5 | 34.5 | 42.6 | 53.0 |
| Female..... | 9.9 | 8.8 | 9.2 | 7.9 | 11.1 | 10.9 | 14.3 | 18.3 | 21.8 | 18.3 |
| Hispanic, both sexes..... | 17.0 | 15.3 | 15.0 | 13.6 | 17.3 | 16.6 | 20.6 | 27.5 | 24.1 | 25.5 |
| Male..... | 25.6 | 22.1 | 20.9 | 20.9 | 23.9 | 27.8 | 29.4 | 40.3 | 32.0 | 45.2 |
| Female..... | 7.8 | 8.4 | 9.2 | 6.9 | 11.6 | 7.5 | 13.9 | 18.3 | 19.0 | * |
| Mexican, both sexes..... | 18.0 | 16.8 | 16.2 | 13.6 | 18.4 | 17.5 | 22.0 | 27.0 | * | * |
| Male..... | 27.3 | 23.0 | 21.1 | 21.3 | 28.2 | 26.7 | 32.3 | * | * | * |
| Female..... | 7.1 | 10.3 | 10.9 | * | 10.1 | * | * | * | * | * |
| Puerto Rican, both sexes..... | 12.4 | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 15.9 | 12.9 | 15.2 | 18.0 | 19.3 | * | * | * | * | * |
| Male..... | 24.1 | 20.7 | 25.1 | * | * | * | * | * | * | * |
| Female..... | 8.6 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 17.1 | 16.9 | 15.8 | 14.9 | 16.5 | 16.6 | 20.1 | 24.8 | 29.7 | 28.9 |
| Male..... | 24.3 | 25.3 | 22.8 | 22.2 | 22.6 | 23.0 | 27.2 | 34.1 | 42.7 | 53.2 |
| Female..... | 10.1 | 8.9 | 9.1 | 8.0 | 11.0 | 11.1 | 14.3 | 18.3 | 21.9 | 18.4 |
| Non-Hispanic white, both sexes..... | 17.4 | 16.9 | 15.8 | 14.8 | 16.5 | 16.3 | 20.3 | 25.3 | 30.4 | 29.7 |
| Male..... | 24.5 | 25.4 | 22.5 | 21.7 | 22.4 | 22.1 | 27.0 | 33.9 | 42.9 | 54.0 |
| Female..... | 10.3 | 8.6 | 9.2 | 8.1 | 10.9 | 11.2 | 14.8 | 19.1 | 22.8 | 19.2 |
| Non-Hispanic black, both sexes..... | 17.3 | 19.0 | 16.8 | 17.2 | 17.4 | 17.2 | 17.3 | 20.5 | 23.8 | 18.8 |
| Male..... | 25.6 | 29.2 | 26.6 | 29.6 | 27.3 | 28.4 | 29.1 | 39.0 | 43.7 | 47.0 |
| Female..... | 10.1 | 10.2 | 8.4 | 7.0 | 9.7 | 8.9 | 9.4 | 9.5 | 13.4 | * |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

| | | | | | | | | | | |
|-------------------------------------|------|------|------|------|------|------|------|------|------|------|
| All origins 3/, both sexes..... | 15.7 | 15.2 | 14.4 | 13.4 | 15.2 | 15.3 | 18.9 | 23.8 | 28.4 | 28.0 |
| Male..... | 22.1 | 22.2 | 20.3 | 19.6 | 20.2 | 20.8 | 25.0 | 32.3 | 39.9 | 51.7 |
| Female..... | 9.4 | 8.4 | 8.7 | 7.5 | 10.6 | 10.6 | 14.1 | 17.7 | 21.4 | 17.7 |
| Hispanic, both sexes..... | 16.1 | 14.2 | 13.6 | 12.7 | 16.6 | 16.1 | 20.0 | 25.6 | 22.3 | 24.9 |
| Male..... | 24.2 | 20.4 | 18.6 | 19.3 | 22.3 | 26.7 | 28.5 | 37.8 | 28.6 | 43.5 |
| Female..... | 7.5 | 7.9 | 8.7 | 6.6 | 11.6 | 7.5 | 13.5 | 16.9 | 18.2 | * |
| Mexican, both sexes..... | 17.1 | 15.7 | 14.6 | 12.8 | 17.5 | 17.2 | 21.2 | 25.6 | * | * |
| Male..... | 25.9 | 21.3 | 18.7 | 20.1 | 26.3 | 26.1 | * | * | * | * |
| Female..... | 6.9 | 9.8 | 10.3 | * | 10.1 | * | * | * | * | * |
| Puerto Rican, both sexes..... | 11.2 | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 15.1 | 12.3 | 13.9 | 16.0 | 18.1 | * | * | * | * | * |
| Male..... | 22.5 | 19.5 | 22.1 | * | * | * | * | * | * | * |
| Female..... | 8.6 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 15.6 | 15.2 | 14.4 | 13.4 | 15.0 | 15.2 | 18.8 | 23.6 | 28.5 | 28.1 |
| Male..... | 21.8 | 22.2 | 20.3 | 19.5 | 19.9 | 20.3 | 24.6 | 31.9 | 40.0 | 51.9 |
| Female..... | 9.6 | 8.5 | 8.7 | 7.6 | 10.4 | 10.8 | 14.1 | 17.8 | 21.5 | 17.8 |
| Non-Hispanic white, both sexes..... | 15.9 | 15.2 | 14.3 | 13.3 | 14.8 | 15.0 | 19.0 | 24.1 | 29.2 | 28.9 |
| Male..... | 22.0 | 22.3 | 20.0 | 19.1 | 19.7 | 19.5 | 24.5 | 31.9 | 40.2 | 52.6 |
| Female..... | 9.8 | 8.2 | 8.7 | 7.7 | 10.3 | 11.0 | 14.5 | 18.5 | 22.4 | 18.6 |
| Non-Hispanic black, both sexes..... | 16.1 | 17.0 | 15.5 | 15.3 | 16.3 | 15.8 | 16.2 | 18.8 | 22.4 | 18.5 |
| Male..... | 23.3 | 25.5 | 24.0 | 25.8 | 24.8 | 25.1 | 26.3 | 34.3 | 40.6 | 45.9 |
| Female..... | 9.7 | 9.6 | 8.3 | 6.8 | 9.5 | 8.9 | 9.4 | 9.5 | 13.0 | * |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Nontransport accidents (W00-X59,Y86)

| | | | | | | | | | | |
|-------------------------------------|------|------|------|------|------|------|------|------|-------|-------|
| All origins 3/, both sexes..... | 22.3 | 21.8 | 18.4 | 15.5 | 16.0 | 21.7 | 30.6 | 55.0 | 101.4 | 246.7 |
| Male..... | 32.3 | 31.5 | 26.8 | 22.6 | 22.9 | 30.0 | 41.4 | 71.9 | 127.5 | 293.6 |
| Female..... | 12.5 | 12.5 | 10.4 | 8.9 | 9.7 | 14.5 | 21.9 | 43.1 | 85.5 | 226.3 |
| Hispanic, both sexes..... | 18.6 | 17.7 | 15.6 | 13.7 | 13.1 | 18.1 | 24.8 | 42.4 | 65.9 | 127.5 |
| Male..... | 29.7 | 27.9 | 26.1 | 22.2 | 20.3 | 28.4 | 32.5 | 57.0 | 89.2 | 179.0 |
| Female..... | 6.8 | 7.5 | 5.4 | 5.9 | 6.9 | 9.7 | 18.9 | 32.0 | 51.1 | 102.0 |
| Mexican, both sexes..... | 16.7 | 16.2 | 14.6 | 13.7 | 13.6 | 17.2 | 27.7 | 37.9 | 73.1 | 121.8 |
| Male..... | 26.4 | 26.0 | 23.9 | 21.3 | 22.1 | 30.4 | 41.5 | 57.6 | * | * |
| Female..... | 5.4 | 5.9 | * | * | * | * | 18.1 | 25.0 | * | * |
| Puerto Rican, both sexes..... | 34.7 | 35.3 | 23.9 | * | * | * | * | * | * | * |
| Male..... | 56.7 | 51.1 | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 18.2 | 14.4 | 14.1 | 14.0 | * | 22.8 | 30.1 | * | * | * |
| Male..... | 31.3 | 24.2 | 25.7 | * | * | * | * | * | * | * |
| Female..... | 6.6 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 22.7 | 22.1 | 18.6 | 15.5 | 16.1 | 21.8 | 30.8 | 55.4 | 102.7 | 250.9 |
| Male..... | 32.4 | 31.7 | 26.6 | 22.3 | 22.9 | 29.8 | 41.7 | 72.4 | 128.6 | 298.0 |
| Female..... | 13.2 | 12.9 | 10.8 | 9.1 | 9.9 | 14.8 | 22.1 | 43.5 | 86.9 | 230.6 |
| Non-Hispanic white, both sexes..... | 23.4 | 21.8 | 17.7 | 14.6 | 15.6 | 20.9 | 30.7 | 56.0 | 105.6 | 261.2 |
| Male..... | 33.4 | 31.1 | 24.8 | 20.4 | 21.7 | 28.0 | 41.3 | 72.6 | 131.2 | 307.6 |
| Female..... | 13.5 | 12.6 | 10.9 | 9.1 | 9.9 | 14.6 | 22.0 | 44.1 | 90.0 | 241.1 |
| Non-Hispanic black, both sexes..... | 23.4 | 30.1 | 29.1 | 25.5 | 23.4 | 30.2 | 35.3 | 55.5 | 73.9 | 156.7 |
| Male..... | 33.5 | 43.8 | 47.1 | 42.5 | 38.1 | 47.1 | 50.3 | 80.3 | 107.6 | 215.5 |
| Female..... | 14.6 | 18.2 | 14.0 | 11.6 | 11.9 | 17.6 | 25.2 | 40.6 | 56.4 | 134.2 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Falls (W00-W19)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|-----|-----|------|------|------|------|-------|
| All origins 3/, both sexes..... | 1.8 | 2.3 | 2.8 | 3.4 | 4.9 | 7.9 | 13.9 | 27.1 | 54.0 | 130.4 |
| Male..... | 2.9 | 3.6 | 4.3 | 5.2 | 7.1 | 10.7 | 18.9 | 35.1 | 67.9 | 158.9 |
| Female..... | 0.7 | 1.0 | 1.4 | 1.8 | 3.0 | 5.5 | 9.9 | 21.4 | 45.5 | 118.1 |
| Hispanic, both sexes..... | 2.0 | 2.1 | 2.9 | 4.0 | 5.0 | 7.0 | 13.7 | 21.3 | 42.3 | 71.5 |
| Male..... | 3.7 | 3.4 | 4.5 | 6.6 | 7.4 | 11.2 | 17.1 | 25.6 | 57.2 | 97.0 |
| Female..... | * | * | * | * | * | * | 11.2 | 18.3 | 32.8 | 58.9 |
| Mexican, both sexes..... | 2.2 | 2.1 | 2.5 | 4.6 | 5.4 | 6.7 | 15.6 | 19.2 | 47.3 | 60.9 |
| Male..... | 3.9 | * | * | 7.2 | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 1.7 | 2.3 | 2.8 | 3.4 | 4.9 | 8.0 | 13.9 | 27.3 | 54.3 | 132.6 |
| Male..... | 2.8 | 3.6 | 4.2 | 5.1 | 7.0 | 10.6 | 18.9 | 35.5 | 68.0 | 161.3 |
| Female..... | 0.7 | 1.0 | 1.4 | 1.8 | 3.0 | 5.7 | 9.9 | 21.5 | 46.0 | 120.2 |
| Non-Hispanic white, both sexes..... | 1.8 | 2.3 | 2.8 | 3.3 | 5.0 | 8.1 | 14.3 | 28.5 | 57.5 | 140.4 |
| Male..... | 2.8 | 3.6 | 4.1 | 4.9 | 7.1 | 10.6 | 19.3 | 36.7 | 71.3 | 169.2 |
| Female..... | 0.7 | 1.0 | 1.5 | 1.8 | 3.2 | 5.9 | 10.2 | 22.6 | 49.1 | 128.0 |
| Non-Hispanic black, both sexes..... | 1.6 | 2.6 | 3.2 | 3.9 | 4.8 | 6.1 | 10.8 | 14.9 | 18.4 | 49.3 |
| Male..... | 2.5 | 4.1 | 5.5 | 6.3 | 7.5 | 9.2 | 15.4 | 22.5 | 28.9 | 76.6 |
| Female..... | * | * | * | * | * | 3.8 | 7.8 | 10.4 | 13.0 | 38.9 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Accidental exposure to smoke, fire and flames (X00-X09)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|-----|-----|-----|-----|-----|------|------|
| All origins 3/, both sexes..... | 0.9 | 1.0 | 1.3 | 1.2 | 1.4 | 1.8 | 2.4 | 2.8 | 4.2 | 4.8 |
| Male..... | 1.3 | 1.4 | 1.6 | 1.5 | 2.0 | 2.6 | 3.2 | 3.5 | 6.1 | 7.0 |
| Female..... | 0.6 | 0.6 | 0.9 | 0.9 | 0.8 | 1.1 | 1.8 | 2.4 | 3.0 | 3.9 |
| Hispanic, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Mexican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 1.0 | 1.0 | 1.3 | 1.2 | 1.4 | 1.8 | 2.5 | 2.9 | 4.2 | 4.8 |
| Male..... | 1.4 | 1.5 | 1.6 | 1.4 | 2.0 | 2.6 | 3.3 | 3.6 | 6.2 | 7.0 |
| Female..... | 0.6 | 0.6 | 1.0 | 0.9 | 0.9 | 1.1 | 1.8 | 2.4 | 2.9 | 3.8 |
| Non-Hispanic white, both sexes..... | 1.0 | 1.0 | 1.1 | 1.0 | 1.1 | 1.6 | 2.1 | 2.5 | 3.7 | 3.9 |
| Male..... | 1.4 | 1.4 | 1.4 | 1.3 | 1.7 | 2.2 | 2.7 | 3.1 | 5.8 | 5.7 |
| Female..... | 0.6 | 0.6 | 0.9 | 0.8 | 0.7 | 1.0 | 1.6 | 2.0 | 2.4 | 3.2 |
| Non-Hispanic black, both sexes..... | 1.1 | 1.6 | 2.8 | 2.6 | 4.1 | 3.9 | 6.5 | 7.4 | 10.7 | 16.0 |
| Male..... | 1.8 | 2.2 | 4.2 | 3.4 | 6.1 | 6.4 | 9.6 | 9.4 | * | 26.3 |
| Female..... | * | * | * | * | * | * | * | 6.2 | 8.9 | 12.1 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Accidental poisoning and exposure to noxious substances (X40-X49)

| | | | | | | | | | | |
|-------------------------------------|------|------|------|------|------|-----|-----|-----|-----|-----|
| All origins 3/, both sexes..... | 14.8 | 13.8 | 9.2 | 5.0 | 3.0 | 2.3 | 1.9 | 2.1 | 2.7 | 3.9 |
| Male..... | 20.3 | 18.7 | 12.8 | 6.8 | 3.6 | 2.8 | 2.1 | 2.3 | 3.2 | 5.1 |
| Female..... | 9.4 | 9.1 | 5.8 | 3.4 | 2.4 | 1.8 | 1.7 | 1.9 | 2.3 | 3.4 |
| Hispanic, both sexes..... | 12.2 | 11.7 | 7.9 | 4.7 | 3.1 | * | * | * | * | * |
| Male..... | 19.0 | 17.6 | 13.4 | 7.2 | 5.4 | * | * | * | * | * |
| Female..... | 5.0 | 5.7 | 2.6 | * | * | * | * | * | * | * |
| Mexican, both sexes..... | 10.0 | 9.7 | 7.9 | 4.3 | * | * | * | * | * | * |
| Male..... | 15.6 | 15.5 | 12.6 | * | * | * | * | * | * | * |
| Female..... | 3.5 | 3.8 | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | 27.1 | 29.5 | * | * | * | * | * | * | * | * |
| Male..... | 43.1 | 41.6 | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 12.9 | 8.7 | 6.5 | * | * | * | * | * | * | * |
| Male..... | 21.9 | 13.7 | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 15.1 | 14.0 | 9.3 | 5.0 | 2.9 | 2.2 | 1.9 | 2.0 | 2.7 | 4.1 |
| Male..... | 20.3 | 18.7 | 12.7 | 6.6 | 3.4 | 2.7 | 2.0 | 2.2 | 3.2 | 5.3 |
| Female..... | 9.9 | 9.4 | 6.1 | 3.5 | 2.5 | 1.8 | 1.7 | 2.0 | 2.3 | 3.5 |
| Non-Hispanic white, both sexes..... | 15.7 | 13.7 | 8.7 | 4.5 | 2.7 | 2.2 | 1.9 | 2.0 | 2.7 | 4.1 |
| Male..... | 21.0 | 18.2 | 11.4 | 5.5 | 2.7 | 2.5 | 1.9 | 2.2 | 2.9 | 5.4 |
| Female..... | 10.4 | 9.4 | 6.1 | 3.5 | 2.7 | 1.9 | 1.8 | 1.9 | 2.5 | 3.5 |
| Non-Hispanic black, both sexes..... | 16.1 | 19.8 | 16.5 | 10.6 | 6.0 | 2.8 | * | * | * | * |
| Male..... | 22.2 | 27.9 | 26.2 | 18.3 | 11.2 | 5.1 | * | * | * | * |
| Female..... | 10.7 | 12.8 | 8.3 | 4.4 | * | * | * | * | * | * |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|------|------|------|------|------|------|-------|
| All origins 3/, both sexes..... | 3.3 | 3.5 | 4.0 | 4.8 | 5.7 | 8.6 | 11.1 | 21.6 | 38.8 | 105.6 |
| Male..... | 5.4 | 5.8 | 6.2 | 7.4 | 8.7 | 12.0 | 15.0 | 28.6 | 47.2 | 118.6 |
| Female..... | 1.3 | 1.3 | 1.9 | 2.4 | 3.0 | 5.7 | 8.1 | 16.6 | 33.8 | 100.0 |
| Hispanic, both sexes..... | 2.8 | 2.4 | 3.3 | 3.0 | 3.9 | 6.4 | 5.9 | 14.9 | 16.0 | 48.2 |
| Male..... | 4.8 | 4.3 | 5.4 | 5.3 | 5.9 | 9.1 | * | 21.1 | * | 70.3 |
| Female..... | * | * | * | * | * | * | * | 10.5 | * | 37.3 |
| Mexican, both sexes..... | 3.2 | 2.6 | 2.9 | * | 5.0 | 6.7 | * | 13.7 | * | 50.0 |
| Male..... | 5.2 | 4.4 | 4.9 | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 3.4 | 3.6 | 4.0 | 4.9 | 5.8 | 8.8 | 11.4 | 21.9 | 39.8 | 107.6 |
| Male..... | 5.4 | 5.9 | 6.2 | 7.4 | 8.9 | 12.2 | 15.4 | 29.0 | 48.2 | 120.4 |
| Female..... | 1.4 | 1.3 | 1.9 | 2.6 | 3.1 | 5.8 | 8.3 | 16.9 | 34.7 | 102.2 |
| Non-Hispanic white, both sexes..... | 3.5 | 3.6 | 4.0 | 4.7 | 5.7 | 8.0 | 11.3 | 21.6 | 40.1 | 110.9 |
| Male..... | 5.7 | 6.1 | 6.0 | 7.0 | 8.7 | 10.9 | 15.2 | 28.4 | 48.2 | 123.1 |
| Female..... | 1.2 | 1.2 | 2.0 | 2.6 | 2.9 | 5.3 | 8.0 | 16.8 | 35.1 | 105.7 |
| Non-Hispanic black, both sexes..... | 3.4 | 4.4 | 5.2 | 7.3 | 7.4 | 16.3 | 14.7 | 28.4 | 39.8 | 84.4 |
| Male..... | 4.3 | 6.6 | 8.8 | 12.6 | 11.4 | 24.3 | 19.2 | 41.8 | 54.6 | 107.2 |
| Female..... | 2.5 | 2.6 | 2.1 | 3.0 | 4.3 | 10.4 | 11.7 | 20.5 | 32.1 | 75.7 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

| | | | | | | | | | | |
|-------------------------------------|------|------|------|------|------|------|------|------|------|------|
| All origins 3/, both sexes..... | 16.1 | 16.3 | 15.1 | 14.6 | 12.3 | 12.5 | 14.6 | 16.6 | 19.3 | 18.0 |
| Male..... | 24.7 | 24.9 | 23.8 | 23.5 | 20.4 | 22.0 | 27.9 | 34.3 | 44.0 | 50.7 |
| Female..... | 7.7 | 8.0 | 6.7 | 6.2 | 5.0 | 4.3 | 3.9 | 4.1 | 4.4 | 3.8 |
| Hispanic, both sexes..... | 7.0 | 7.2 | 7.0 | 7.9 | 6.2 | 6.8 | 8.4 | 8.0 | 11.1 | 12.7 |
| Male..... | 11.1 | 11.0 | 11.7 | 14.5 | 11.3 | 13.2 | 16.7 | 16.0 | 25.2 | 35.1 |
| Female..... | 2.6 | 3.3 | 2.5 | * | * | * | * | * | * | * |
| Mexican, both sexes..... | 6.3 | 6.4 | 5.5 | 5.6 | 4.8 | * | * | * | * | * |
| Male..... | 10.0 | 10.3 | 8.9 | 9.6 | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 7.2 | 7.6 | 8.1 | 9.6 | * | * | * | * | * | * |
| Male..... | 12.5 | 10.2 | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 17.3 | 17.2 | 15.7 | 15.1 | 12.7 | 12.8 | 14.9 | 17.0 | 19.7 | 18.1 |
| Male..... | 26.5 | 26.3 | 24.8 | 24.1 | 20.9 | 22.3 | 28.5 | 35.2 | 44.8 | 50.9 |
| Female..... | 8.3 | 8.5 | 7.1 | 6.5 | 5.2 | 4.5 | 4.0 | 4.2 | 4.4 | 3.9 |
| Non-Hispanic white, both sexes..... | 19.7 | 19.7 | 17.7 | 16.7 | 14.1 | 14.1 | 16.2 | 18.1 | 21.3 | 19.2 |
| Male..... | 30.0 | 29.4 | 27.7 | 26.5 | 23.2 | 24.3 | 31.0 | 37.3 | 48.2 | 54.3 |
| Female..... | 9.5 | 10.0 | 7.8 | 7.4 | 5.6 | 5.0 | 4.0 | 4.3 | 4.8 | 4.1 |
| Non-Hispanic black, both sexes..... | 7.2 | 6.0 | 5.7 | 4.7 | 5.1 | 3.6 | 6.5 | 6.7 | * | * |
| Male..... | 11.7 | 11.3 | 8.9 | 8.7 | 8.1 | 7.9 | 12.0 | 15.5 | * | * |
| Female..... | 3.3 | 1.5 | 3.0 | * | * | * | * | * | * | * |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

| | | | | | | | | | | |
|-------------------------------------|------|------|------|-----|-----|-----|-----|-----|-----|------|
| All origins 3/, both sexes..... | 8.7 | 8.5 | 7.2 | 5.9 | 4.2 | 3.7 | 3.9 | 4.2 | 5.0 | 6.0 |
| Male..... | 12.1 | 11.6 | 9.9 | 8.2 | 5.7 | 5.0 | 5.5 | 6.5 | 8.1 | 12.9 |
| Female..... | 5.4 | 5.6 | 4.6 | 3.9 | 2.9 | 2.5 | 2.6 | 2.6 | 3.1 | 3.0 |
| Hispanic, both sexes..... | 4.5 | 4.3 | 4.6 | 4.4 | 3.0 | * | * | * | * | * |
| Male..... | 7.0 | 6.1 | 6.9 | 7.7 | * | * | * | * | * | * |
| Female..... | 1.9 | 2.5 | * | * | * | * | * | * | * | * |
| Mexican, both sexes..... | 3.7 | 3.9 | 3.5 | * | * | * | * | * | * | * |
| Male..... | 5.8 | 6.0 | 5.5 | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 5.0 | 4.7 | * | * | * | * | * | * | * | * |
| Male..... | 8.4 | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 9.2 | 9.0 | 7.4 | 6.0 | 4.3 | 3.7 | 3.9 | 4.2 | 5.0 | 5.9 |
| Male..... | 12.7 | 12.1 | 10.0 | 8.1 | 5.6 | 4.9 | 5.4 | 6.4 | 8.0 | 12.5 |
| Female..... | 5.8 | 5.9 | 4.8 | 4.0 | 3.0 | 2.6 | 2.6 | 2.6 | 3.2 | 3.1 |
| Non-Hispanic white, both sexes..... | 10.4 | 10.1 | 8.2 | 6.6 | 4.6 | 3.8 | 4.0 | 4.3 | 5.3 | 6.1 |
| Male..... | 14.2 | 13.3 | 11.2 | 8.7 | 6.1 | 4.9 | 5.7 | 6.7 | 8.3 | 13.0 |
| Female..... | 6.6 | 7.0 | 5.2 | 4.5 | 3.2 | 2.8 | 2.6 | 2.6 | 3.4 | 3.2 |
| Non-Hispanic black, both sexes..... | 4.2 | 3.1 | 2.9 | 2.0 | 2.2 | * | * | * | * | * |
| Male..... | 6.4 | 5.4 | 3.8 | 3.2 | * | * | * | * | * | * |
| Female..... | 2.3 | * | 2.2 | * | * | * | * | * | * | * |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
|--|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

| | | | | | | | | | | |
|-------------------------------------|------|------|------|------|------|------|------|------|-----|-----|
| All origins 3/, both sexes..... | 6.5 | 5.4 | 4.0 | 3.5 | 2.7 | 2.4 | 2.2 | 2.3 | 2.4 | 2.1 |
| Male..... | 9.6 | 8.0 | 6.0 | 5.3 | 4.2 | 3.4 | 3.2 | 3.3 | 2.9 | 3.0 |
| Female..... | 3.5 | 2.9 | 2.2 | 1.8 | 1.4 | 1.5 | 1.4 | 1.6 | 2.1 | 1.7 |
| Hispanic, both sexes..... | 7.2 | 6.2 | 5.5 | 4.9 | 3.7 | * | * | * | * | * |
| Male..... | 11.1 | 10.0 | 8.0 | 8.6 | 6.4 | * | * | * | * | * |
| Female..... | 3.1 | 2.4 | 3.1 | * | * | * | * | * | * | * |
| Mexican, both sexes..... | 7.8 | 7.0 | 4.3 | 5.2 | * | * | * | * | * | * |
| Male..... | 11.5 | 11.4 | 6.3 | 8.7 | * | * | * | * | * | * |
| Female..... | 3.4 | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 6.6 | 5.5 | 7.1 | * | * | * | * | * | * | * |
| Male..... | 11.6 | 9.8 | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 6.4 | 5.3 | 3.9 | 3.3 | 2.6 | 2.4 | 2.2 | 2.3 | 2.2 | 2.1 |
| Male..... | 9.3 | 7.7 | 5.8 | 5.0 | 4.0 | 3.3 | 3.1 | 3.3 | 2.8 | 2.9 |
| Female..... | 3.5 | 2.9 | 2.1 | 1.8 | 1.4 | 1.5 | 1.5 | 1.5 | 1.9 | 1.7 |
| Non-Hispanic white, both sexes..... | 4.0 | 3.4 | 2.6 | 2.6 | 2.0 | 1.9 | 1.9 | 1.8 | 1.9 | 1.8 |
| Male..... | 5.7 | 4.3 | 3.5 | 3.6 | 2.9 | 2.5 | 2.4 | 2.4 | 2.1 | 2.7 |
| Female..... | 2.3 | 2.5 | 1.6 | 1.7 | 1.3 | 1.3 | 1.5 | 1.3 | 1.8 | 1.5 |
| Non-Hispanic black, both sexes..... | 20.9 | 17.9 | 13.2 | 9.0 | 7.7 | 6.0 | 5.5 | 8.2 | 5.9 | * |
| Male..... | 33.1 | 31.4 | 22.8 | 17.2 | 13.8 | 10.2 | 10.9 | 15.0 | * | * |
| Female..... | 10.2 | 6.3 | 5.0 | * | * | * | * | * | * | * |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

| | | | | | | | | | | |
|-------------------------------------|------|------|------|-----|-----|-----|-----|-----|-----|---|
| All origins 3/, both sexes..... | 3.9 | 2.8 | 2.2 | 1.8 | 1.3 | 1.0 | 0.8 | 1.0 | 0.8 | * |
| Male..... | 6.2 | 4.2 | 3.4 | 2.7 | 1.9 | 1.4 | 1.0 | 1.4 | 1.0 | * |
| Female..... | 1.7 | 1.5 | 1.1 | 0.9 | 0.7 | 0.6 | 0.6 | 0.6 | 0.7 | * |
| Hispanic, both sexes..... | 4.6 | 3.1 | 3.2 | 1.8 | * | * | * | * | * | * |
| Male..... | 7.3 | 5.1 | 4.9 | * | * | * | * | * | * | * |
| Female..... | 1.8 | * | * | * | * | * | * | * | * | * |
| Mexican, both sexes..... | 5.1 | 3.8 | 2.7 | * | * | * | * | * | * | * |
| Male..... | 7.7 | 6.2 | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 4.3 | * | * | * | * | * | * | * | * | * |
| Male..... | 7.5 | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 3.8 | 2.8 | 2.1 | 1.7 | 1.3 | 1.0 | 0.8 | 1.0 | 0.8 | * |
| Male..... | 6.0 | 4.1 | 3.3 | 2.7 | 1.9 | 1.4 | 1.0 | 1.4 | * | * |
| Female..... | 1.7 | 1.5 | 1.0 | 0.9 | 0.7 | 0.6 | 0.6 | 0.7 | 0.7 | * |
| Non-Hispanic white, both sexes..... | 2.3 | 1.8 | 1.4 | 1.4 | 0.9 | 0.7 | 0.7 | 0.9 | 0.7 | * |
| Male..... | 3.4 | 2.2 | 2.1 | 2.0 | 1.3 | 0.9 | 0.8 | 1.2 | * | * |
| Female..... | 1.2 | 1.3 | 0.8 | 0.9 | 0.5 | 0.5 | 0.7 | 0.6 | 0.7 | * |
| Non-Hispanic black, both sexes..... | 13.3 | 9.7 | 7.2 | 4.2 | 4.1 | 2.8 | * | * | * | * |
| Male..... | 23.1 | 17.6 | 12.3 | 8.4 | 6.7 | 5.4 | * | * | * | * |
| Female..... | 4.7 | 2.8 | 2.8 | * | * | * | * | * | * | * |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

| | | | | | | | | | | |
|-------------------------------------|------|------|------|-----|-----|-----|-----|------|-----|-----|
| All origins 3/, both sexes..... | 2.6 | 2.6 | 1.8 | 1.7 | 1.5 | 1.4 | 1.4 | 1.3 | 1.6 | 1.7 |
| Male..... | 3.4 | 3.8 | 2.6 | 2.6 | 2.3 | 2.0 | 2.2 | 1.8 | 1.9 | 2.3 |
| Female..... | 1.8 | 1.4 | 1.1 | 1.0 | 0.7 | 0.9 | 0.8 | 0.9 | 1.4 | 1.4 |
| Hispanic, both sexes..... | 2.6 | 3.1 | 2.3 | 3.1 | 2.5 | * | * | * | * | * |
| Male..... | 3.7 | 4.9 | 3.1 | 5.3 | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Mexican, both sexes..... | 2.7 | 3.2 | * | 3.4 | * | * | * | * | * | * |
| Male..... | 3.8 | 5.2 | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 2.5 | 2.5 | 1.7 | 1.6 | 1.4 | 1.4 | 1.4 | 1.3 | 1.5 | 1.7 |
| Male..... | 3.4 | 3.6 | 2.5 | 2.3 | 2.1 | 1.9 | 2.1 | 1.8 | 1.8 | 2.3 |
| Female..... | 1.7 | 1.4 | 1.0 | 1.0 | 0.7 | 0.9 | 0.8 | 0.9 | 1.2 | 1.4 |
| Non-Hispanic white, both sexes..... | 1.7 | 1.6 | 1.1 | 1.2 | 1.1 | 1.1 | 1.1 | 0.9 | 1.2 | 1.4 |
| Male..... | 2.3 | 2.1 | 1.4 | 1.5 | 1.6 | 1.6 | 1.6 | 1.2 | 1.4 | 2.0 |
| Female..... | 1.1 | 1.1 | 0.8 | 0.9 | 0.7 | 0.7 | 0.8 | 0.7 | 1.1 | 1.2 |
| Non-Hispanic black, both sexes..... | 7.6 | 8.2 | 6.0 | 4.8 | 3.6 | 3.2 | 3.7 | 5.8 | * | * |
| Male..... | 10.1 | 13.7 | 10.5 | 8.7 | 7.1 | * | 7.2 | 10.3 | * | * |
| Female..... | 5.5 | 3.5 | 2.2 | * | * | * | * | * | * | * |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

40-44 45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years and over years years years years years years years years years

Discharge of firearms, undetermined intent (Y22-Y24)

Table with 11 columns representing age groups (40-44 to 85 years and over) and rows for various demographic categories including All origins, Hispanic, Mexican, Puerto Rican, Cuban, Other Hispanic, Non-Hispanic white, and Non-Hispanic black, each split by sex. All data points are marked with an asterisk (*).

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

40-44 45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years and over years years years years years years years years years

Operations of war and their sequelae (Y36,Y89.1)

Table with 11 columns representing age groups (40-44 to 85 years and over) and rows for various categories: All origins 3/, both sexes; Male; Female; Hispanic, both sexes; Male; Female; Mexican, both sexes; Male; Female; Puerto Rican, both sexes; Male; Female; Cuban, both sexes; Male; Female; Other Hispanic 4/, both sexes; Male; Female; Non-Hispanic 5/, both sexes; Male; Female; Non-Hispanic white, both sexes; Male; Female; Non-Hispanic black, both sexes; Male; Female. All cells contain an asterisk (*).

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Complications of medical and surgical care (Y40-Y84,Y88)

| | | | | | | | | | | |
|-------------------------------------|-----|-----|-----|-----|-----|-----|-----|-----|------|------|
| All origins 3/, both sexes..... | 0.4 | 0.5 | 0.8 | 1.1 | 1.6 | 2.3 | 3.9 | 5.1 | 7.7 | 11.3 |
| Male..... | 0.4 | 0.6 | 0.7 | 1.2 | 1.5 | 2.5 | 3.8 | 5.4 | 7.7 | 12.3 |
| Female..... | 0.4 | 0.5 | 0.9 | 1.0 | 1.7 | 2.2 | 3.9 | 5.0 | 7.7 | 10.9 |
| Hispanic, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Mexican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 0.5 | 0.6 | 0.8 | 1.1 | 1.6 | 2.3 | 3.9 | 5.2 | 7.7 | 11.6 |
| Male..... | 0.4 | 0.6 | 0.7 | 1.2 | 1.6 | 2.5 | 3.9 | 5.4 | 7.8 | 12.9 |
| Female..... | 0.5 | 0.5 | 0.9 | 1.0 | 1.7 | 2.2 | 4.0 | 5.0 | 7.7 | 11.1 |
| Non-Hispanic white, both sexes..... | 0.4 | 0.5 | 0.7 | 1.0 | 1.5 | 2.1 | 3.9 | 5.0 | 7.6 | 11.5 |
| Male..... | 0.4 | 0.5 | 0.6 | 1.1 | 1.5 | 2.2 | 3.8 | 5.3 | 7.9 | 12.4 |
| Female..... | 0.3 | 0.4 | 0.7 | 0.9 | 1.5 | 2.1 | 4.0 | 4.9 | 7.4 | 11.1 |
| Non-Hispanic black, both sexes..... | 1.0 | 1.3 | 2.2 | 2.6 | 2.9 | 4.1 | 5.4 | 7.7 | 9.6 | 14.2 |
| Male..... | * | * | * | * | * | 5.1 | * | * | * | * |
| Female..... | * | * | 2.5 | 2.5 | 3.2 | * | 5.0 | 8.4 | 10.1 | 11.7 |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

| Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
|--|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|

Alcohol-induced causes (F10,G31.2,G62.1,I42.6,K29.2,K70,R78.0,X45,X65,Y15) 6/

| | | | | | | | | | | |
|-------------------------------------|------|------|------|------|------|------|------|------|------|-----|
| All origins 3/, both sexes..... | 10.5 | 15.3 | 18.0 | 18.0 | 18.3 | 16.4 | 14.3 | 10.8 | 7.2 | 4.3 |
| Male..... | 15.0 | 23.2 | 28.9 | 28.8 | 30.5 | 27.7 | 23.5 | 18.6 | 13.4 | 9.6 |
| Female..... | 6.0 | 7.6 | 7.6 | 7.8 | 7.2 | 6.6 | 6.9 | 5.3 | 3.4 | 2.0 |
| Hispanic, both sexes..... | 11.8 | 18.5 | 24.3 | 26.6 | 25.7 | 27.5 | 25.6 | 16.3 | 11.6 | * |
| Male..... | 19.7 | 31.4 | 42.1 | 48.8 | 47.5 | 53.5 | 50.0 | 28.8 | 22.9 | * |
| Female..... | 3.5 | 5.3 | 7.0 | 6.2 | 6.7 | 6.4 | 6.8 | * | * | * |
| Mexican, both sexes..... | 12.5 | 19.9 | 26.8 | 27.7 | 26.2 | 34.7 | 34.5 | 16.5 | * | * |
| Male..... | 19.7 | 33.8 | 44.8 | 50.3 | 47.9 | 62.0 | 72.8 | * | * | * |
| Female..... | 4.0 | 5.6 | 7.4 | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | 17.4 | 24.4 | 33.8 | 36.8 | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 12.8 | 17.8 | 21.2 | 25.2 | 27.8 | 25.0 | * | * | * | * |
| Male..... | 24.1 | 32.4 | 38.8 | 44.6 | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 10.2 | 14.8 | 17.3 | 17.1 | 17.6 | 15.4 | 13.5 | 10.4 | 7.0 | 4.1 |
| Male..... | 14.3 | 22.0 | 27.4 | 26.9 | 29.0 | 25.6 | 21.7 | 17.9 | 12.9 | 9.0 |
| Female..... | 6.3 | 7.8 | 7.7 | 7.9 | 7.2 | 6.7 | 6.9 | 5.2 | 3.3 | 1.9 |
| Non-Hispanic white, both sexes..... | 10.4 | 14.9 | 17.0 | 16.4 | 16.8 | 15.2 | 13.2 | 10.5 | 6.8 | 4.0 |
| Male..... | 14.6 | 22.1 | 26.9 | 25.7 | 27.6 | 24.7 | 20.7 | 17.9 | 12.5 | 8.6 |
| Female..... | 6.3 | 7.8 | 7.3 | 7.5 | 6.9 | 6.8 | 7.0 | 5.2 | 3.3 | 2.0 |
| Non-Hispanic black, both sexes..... | 9.1 | 15.6 | 21.7 | 25.2 | 24.6 | 18.5 | 17.4 | 12.1 | 9.1 | * |
| Male..... | 12.7 | 24.0 | 35.4 | 41.6 | 44.2 | 35.8 | 34.2 | 22.5 | 19.5 | * |
| Female..... | 5.9 | 8.4 | 10.2 | 11.7 | 9.2 | 5.7 | 6.2 | 5.9 | * | * |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Drug-induced causes (F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,X40-X44,X60-X64,X85,Y10-Y14) 6/

| | | | | | | | | | | |
|-------------------------------------|------|------|------|------|------|-----|-----|-----|-----|-----|
| All origins 3/, both sexes..... | 21.8 | 20.9 | 14.5 | 8.2 | 5.1 | 3.8 | 3.3 | 3.5 | 4.1 | 5.6 |
| Male..... | 28.0 | 26.3 | 19.1 | 10.0 | 5.7 | 4.1 | 3.3 | 3.8 | 4.6 | 7.3 |
| Female..... | 15.8 | 15.6 | 10.2 | 6.6 | 4.5 | 3.5 | 3.2 | 3.3 | 3.7 | 4.8 |
| Hispanic, both sexes..... | 15.8 | 15.9 | 11.1 | 6.5 | 4.4 | * | * | * | * | * |
| Male..... | 24.1 | 23.6 | 17.7 | 9.9 | 6.9 | * | * | * | * | * |
| Female..... | 7.0 | 8.1 | 4.6 | 3.4 | * | * | * | * | * | * |
| Mexican, both sexes..... | 13.1 | 13.2 | 10.3 | 5.8 | * | * | * | * | * | * |
| Male..... | 19.7 | 20.7 | 15.6 | 9.0 | * | * | * | * | * | * |
| Female..... | 5.4 | 5.4 | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | 37.5 | 38.2 | 20.6 | * | * | * | * | * | * | * |
| Male..... | 60.1 | 54.8 | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 15.4 | 12.7 | 9.2 | * | * | * | * | * | * | * |
| Male..... | 25.6 | 18.4 | 16.1 | * | * | * | * | * | * | * |
| Female..... | 6.4 | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 22.5 | 21.3 | 14.8 | 8.3 | 5.1 | 3.8 | 3.2 | 3.5 | 4.1 | 5.7 |
| Male..... | 28.3 | 26.5 | 19.1 | 9.9 | 5.6 | 4.0 | 3.2 | 3.7 | 4.6 | 7.4 |
| Female..... | 16.8 | 16.3 | 10.6 | 6.9 | 4.7 | 3.6 | 3.3 | 3.4 | 3.8 | 5.0 |
| Non-Hispanic white, both sexes..... | 23.4 | 21.2 | 14.0 | 8.0 | 4.9 | 3.8 | 3.4 | 3.6 | 4.3 | 5.8 |
| Male..... | 29.3 | 25.9 | 17.2 | 8.7 | 4.6 | 3.7 | 3.3 | 3.7 | 4.5 | 7.7 |
| Female..... | 17.5 | 16.6 | 10.9 | 7.2 | 5.1 | 4.0 | 3.5 | 3.5 | 4.2 | 5.0 |
| Non-Hispanic black, both sexes..... | 24.6 | 28.9 | 25.3 | 13.9 | 8.8 | 3.6 | * | * | * | * |
| Male..... | 31.8 | 39.7 | 40.5 | 23.3 | 16.1 | 6.7 | * | * | * | * |
| Female..... | 18.3 | 19.5 | 12.5 | 6.2 | 3.2 | * | * | * | * | * |

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2002

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2002. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), of Diseases, Tenth Revision, 1992), race and sex

| | 40-44 years | 45-49 years | 50-54 years | 55-59 years | 60-64 years | 65-69 years | 70-74 years | 75-79 years | 80-84 years | 85 years and over |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------------|

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 6/

| | | | | | | | | | | |
|-------------------------------------|------|------|------|------|------|------|------|------|------|------|
| All origins 3/, both sexes..... | 11.9 | 11.1 | 10.5 | 10.7 | 9.6 | 10.1 | 11.8 | 13.6 | 15.5 | 12.5 |
| Male..... | 19.7 | 18.4 | 18.0 | 18.6 | 17.1 | 18.8 | 24.1 | 29.8 | 37.6 | 38.9 |
| Female..... | 4.2 | 4.0 | 3.2 | 3.3 | 2.8 | 2.5 | 2.0 | 2.2 | 2.0 | 1.1 |
| Hispanic, both sexes..... | 7.2 | 6.1 | 5.9 | 5.4 | 4.4 | 5.2 | 5.3 | * | * | * |
| Male..... | 11.6 | 10.2 | 10.0 | 10.3 | 8.5 | 10.2 | 11.8 | * | * | * |
| Female..... | 2.6 | 1.9 | * | * | * | * | * | * | * | * |
| Mexican, both sexes..... | 8.0 | 6.4 | 4.8 | 5.2 | * | * | * | * | * | * |
| Male..... | 12.3 | 10.6 | 7.7 | 9.6 | * | * | * | * | * | * |
| Female..... | 2.8 | * | * | * | * | * | * | * | * | * |
| Puerto Rican, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Cuban, both sexes..... | * | * | * | * | * | * | * | * | * | * |
| Male..... | * | * | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Other Hispanic 4/, both sexes..... | 6.6 | 5.7 | 8.1 | * | * | * | * | * | * | * |
| Male..... | 11.6 | 9.8 | * | * | * | * | * | * | * | * |
| Female..... | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic 5/, both sexes..... | 12.4 | 11.5 | 10.8 | 11.1 | 9.9 | 10.4 | 12.2 | 14.1 | 15.8 | 12.7 |
| Male..... | 20.7 | 19.1 | 18.6 | 19.3 | 17.6 | 19.3 | 24.9 | 30.8 | 38.3 | 39.5 |
| Female..... | 4.4 | 4.2 | 3.4 | 3.4 | 2.9 | 2.6 | 2.1 | 2.3 | 2.1 | 1.2 |
| Non-Hispanic white, both sexes..... | 12.2 | 11.8 | 11.3 | 12.0 | 10.6 | 11.3 | 13.3 | 14.9 | 17.1 | 13.7 |
| Male..... | 20.2 | 19.2 | 19.3 | 20.4 | 18.8 | 20.9 | 26.8 | 32.3 | 41.3 | 42.5 |
| Female..... | 4.3 | 4.5 | 3.4 | 3.9 | 3.0 | 2.9 | 2.2 | 2.4 | 2.3 | 1.2 |
| Non-Hispanic black, both sexes..... | 16.9 | 13.2 | 10.3 | 7.3 | 7.1 | 5.2 | 6.5 | 8.6 | * | * |
| Male..... | 29.6 | 24.6 | 18.0 | 14.6 | 12.4 | 10.5 | 14.0 | 20.2 | * | * |
| Female..... | 5.7 | 3.3 | 3.7 | * | * | * | * | * | * | * |

* Figure does not meet standards of reliability or precision.

0.0 Quantity more than zero but less than 0.05.

... Category not applicable.

1/ Figures for age not stated included in "All ages" but not distributed among age groups.

2/ Death rates for "Under 1 year" (based on population estimates) differ from infant mortality rates (based on live births).

3/ Figures for origin not stated are included in "All origins" but not distributed among specified origins.

4/ Includes Central and South American and Other and unknown Hispanic.

5/ Includes races other than white and black.

6/ Included in selected categories.

Source: CDC/NCHS, National Vital Statistics System, Mortality.