

Worktable Orig250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 2003

[Rates per 100,000 population, in specified group. Rates based on populations estimated as of July 1, 2003. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)												
All origins 3/, both sexes.....	2.6	4.3	0.5	0.1	0.2	0.4	1.0	2.9	4.1	7.3	13.3	22.5
Male.....	2.8	4.2	0.5	0.1	0.2	0.4	1.3	4.0	5.1	8.5	15.8	26.5
Female.....	2.3	4.5	0.5	0.1	0.2	0.3	0.7	1.9	3.1	6.3	11.7	20.8
Hispanic, both sexes.....	1.7	2.7	*	*	*	0.3	1.3	4.4	5.6	8.4	15.3	13.0
Male.....	2.0	*	*	*	*	*	1.8	6.8	7.7	9.8	16.5	*
Female.....	1.3	*	*	*	*	*	0.7	2.0	3.6	7.4	14.5	*
Non-Hispanic 4/, both sexes.....	2.7	4.8	0.5	0.1	0.2	0.4	1.0	2.7	3.9	7.2	13.2	22.9
Male.....	2.9	4.8	0.5	*	0.2	0.4	1.2	3.6	4.9	8.4	15.8	27.0
Female.....	2.5	4.8	0.5	*	0.2	0.3	0.7	1.9	3.0	6.2	11.5	21.2
Non-Hispanic white, both sexes.....	2.8	3.8	0.4	0.1	0.2	0.3	0.8	2.6	3.6	7.2	12.8	23.4
Male.....	2.9	3.6	*	*	0.2	0.3	1.1	3.4	4.5	8.2	15.4	27.6
Female.....	2.6	4.0	*	*	0.2	0.3	0.6	1.7	2.8	6.3	11.1	21.6
Non-Hispanic black, both sexes.....	2.6	8.4	0.9	*	*	0.6	1.8	4.4	6.1	7.0	16.4	19.9
Male.....	3.0	9.3	*	*	*	0.8	2.2	5.8	8.1	10.6	19.0	23.4
Female.....	2.2	7.4	*	*	*	*	1.4	3.1	4.6	4.4	14.8	18.6
Malignant neoplasms (C00-C97)												
All origins 3/, both sexes.....	191.5	1.9	2.5	2.6	4.0	9.4	35.0	122.2	343.0	770.3	1,302.5	1,698.2
Male.....	201.3	1.7	2.8	2.8	4.6	8.9	30.8	127.4	386.8	931.7	1,695.4	2,413.8
Female.....	182.0	2.1	2.1	2.4	3.4	9.9	39.1	117.1	302.3	635.3	1,040.1	1,381.9
Hispanic, both sexes.....	60.3	*	3.0	2.9	4.1	7.1	22.9	80.7	214.7	485.2	890.1	1,234.9
Male.....	61.5	*	3.6	3.2	4.9	6.8	18.2	81.1	246.5	617.6	1,163.9	1,668.6
Female.....	59.1	*	2.4	2.6	3.2	7.4	28.0	80.2	185.9	379.7	702.1	1,014.8
Non-Hispanic 4/, both sexes.....	211.9	2.0	2.3	2.6	4.0	9.9	36.7	126.0	352.4	788.9	1,321.6	1,714.7
Male.....	224.3	1.9	2.6	2.7	4.5	9.4	32.8	131.9	396.9	951.1	1,720.2	2,444.0
Female.....	200.0	2.1	2.1	2.4	3.4	10.4	40.6	120.5	311.1	652.7	1,055.6	1,394.3
Non-Hispanic white, both sexes.....	229.5	2.2	2.4	2.6	4.0	9.6	35.2	119.4	345.9	788.8	1,322.6	1,717.8
Male.....	241.8	2.1	2.6	2.7	4.6	9.4	31.9	123.8	384.8	942.0	1,707.8	2,441.7
Female.....	217.6	2.3	2.2	2.4	3.4	9.8	38.6	115.1	308.9	657.6	1,062.4	1,399.1
Non-Hispanic black, both sexes.....	170.0	*	2.3	2.7	4.1	12.6	50.5	189.7	480.9	941.1	1,508.2	1,867.5
Male.....	184.8	*	2.4	2.9	4.7	10.6	42.9	211.5	597.1	1,242.1	2,175.8	2,859.6
Female.....	156.5	*	2.1	2.4	3.5	14.5	57.2	170.9	387.6	727.0	1,131.2	1,483.2
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)												
All origins 3/, both sexes.....	2.7	*	*	*	0.1	0.1	0.6	2.8	6.5	10.2	13.1	18.6
Male.....	3.7	*	*	*	0.1	0.2	0.9	4.6	10.6	15.6	19.4	26.9
Female.....	1.7	*	*	*	*	0.1	0.4	1.0	2.8	5.7	9.0	15.0
Hispanic, both sexes.....	0.8	*	*	*	*	*	*	1.3	3.6	6.5	9.8	15.0
Male.....	1.1	*	*	*	*	*	*	2.1	6.5	10.3	16.5	*
Female.....	0.5	*	*	*	*	*	*	*	*	3.5	5.3	*
Non-Hispanic 4/, both sexes.....	3.0	*	*	*	0.1	0.2	0.7	2.9	6.8	10.5	13.3	18.7
Male.....	4.1	*	*	*	*	0.2	1.0	4.8	10.9	16.0	19.5	27.1
Female.....	1.9	*	*	*	*	*	0.4	1.1	2.9	5.9	9.2	15.0
Non-Hispanic white, both sexes.....	3.1	*	*	*	*	0.1	0.6	2.6	6.3	10.1	13.0	19.1
Male.....	4.1	*	*	*	*	0.2	1.0	4.2	9.9	14.9	18.7	27.4
Female.....	2.0	*	*	*	*	*	0.2	0.9	2.9	5.9	9.2	15.4
Non-Hispanic black, both sexes.....	3.0	*	*	*	*	*	1.1	5.2	11.5	15.7	15.6	15.5
Male.....	4.8	*	*	*	*	*	1.2	9.2	21.8	29.1	28.6	25.6
Female.....	1.4	*	*	*	*	*	1.1	1.8	3.3	6.2	8.3	11.6

Worktable Orig250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 2003

[Rates per 100,000 population, in specified group. Rates based on populations estimated as of July 1, 2003. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of esophagus (C15)												
All origins 3/, both sexes.....	4.4	*	*	*	*	0.1	0.6	3.4	10.5	19.7	26.4	25.0
Male.....	6.9	*	*	*	*	0.1	1.1	6.0	18.5	34.1	47.8	45.3
Female.....	2.0	*	*	*	*	*	0.2	1.0	3.2	7.7	12.0	16.1
Hispanic, both sexes.....	1.1	*	*	*	*	*	*	1.3	5.0	10.0	14.4	16.1
Male.....	1.6	*	*	*	*	*	*	2.4	8.0	17.2	28.4	*
Female.....	0.5	*	*	*	*	*	*	*	2.4	4.3	*	*
Non-Hispanic 4/, both sexes.....	4.9	*	*	*	*	0.1	0.7	3.7	10.9	20.4	27.0	25.4
Male.....	7.8	*	*	*	*	0.1	1.2	6.4	19.3	35.3	48.8	46.3
Female.....	2.2	*	*	*	*	*	0.2	1.1	3.2	7.9	12.4	16.1
Non-Hispanic white, both sexes.....	5.3	*	*	*	*	0.1	0.7	3.4	10.6	20.3	27.3	26.1
Male.....	8.5	*	*	*	*	*	1.2	6.0	18.8	35.1	49.9	47.9
Female.....	2.3	*	*	*	*	*	0.2	0.9	2.8	7.7	12.0	16.5
Non-Hispanic black, both sexes.....	4.3	*	*	*	*	*	0.9	6.1	17.3	26.1	28.4	21.4
Male.....	6.5	*	*	*	*	*	1.4	10.4	29.4	45.2	46.7	38.4
Female.....	2.4	*	*	*	*	*	*	2.5	7.5	12.5	18.1	14.9
Malignant neoplasm of stomach (C16)												
All origins 3/, both sexes.....	4.2	*	*	*	*	0.3	1.1	3.0	6.7	15.2	27.6	43.6
Male.....	4.9	*	*	*	*	0.3	1.2	3.8	9.5	21.9	38.4	60.0
Female.....	3.4	*	*	*	*	0.3	1.0	2.1	4.1	9.7	20.3	36.4
Hispanic, both sexes.....	3.1	*	*	*	*	0.5	1.6	5.1	11.4	23.8	43.6	62.7
Male.....	3.4	*	*	*	*	*	1.5	5.7	14.1	33.2	63.7	73.9
Female.....	2.8	*	*	*	*	*	1.6	4.5	9.0	16.3	29.8	57.0
Non-Hispanic 4/, both sexes.....	4.3	*	*	*	*	0.3	1.0	2.7	6.3	14.6	26.7	42.7
Male.....	5.2	*	*	*	*	0.3	1.1	3.6	9.2	21.0	36.9	59.2
Female.....	3.5	*	*	*	*	0.3	0.9	1.9	3.7	9.2	19.8	35.5
Non-Hispanic white, both sexes.....	4.0	*	*	*	*	0.2	0.8	2.1	5.4	12.6	23.3	37.5
Male.....	4.8	*	*	*	*	0.2	0.9	2.8	7.7	18.0	32.4	52.3
Female.....	3.3	*	*	*	*	0.2	0.6	1.4	3.1	8.0	17.2	31.0
Non-Hispanic black, both sexes.....	5.5	*	*	*	*	0.4	1.8	5.8	12.9	27.5	55.0	86.0
Male.....	6.8	*	*	*	*	*	2.0	8.5	20.0	44.8	80.5	107.6
Female.....	4.4	*	*	*	*	*	1.6	3.5	7.3	15.2	40.6	77.6
Malignant neoplasms of colon, rectum and anus (C18-C21)												
All origins 3/, both sexes.....	19.2	*	*	*	0.1	0.7	3.0	10.9	29.8	70.5	134.7	239.7
Male.....	19.6	*	*	*	0.1	0.8	3.2	12.5	36.0	88.0	164.4	273.3
Female.....	18.9	*	*	*	*	0.7	2.7	9.3	24.0	55.9	114.8	224.8
Hispanic, both sexes.....	6.2	*	*	*	*	0.5	2.1	8.2	23.6	49.7	95.5	144.6
Male.....	6.6	*	*	*	*	0.5	2.3	10.0	29.8	64.8	120.4	172.4
Female.....	5.7	*	*	*	*	1.8	6.4	18.0	37.6	78.5	130.5	243.2
Non-Hispanic 4/, both sexes.....	21.3	*	*	*	0.1	0.8	3.1	11.1	30.2	71.9	136.5	243.2
Male.....	21.7	*	*	*	*	0.9	3.4	12.7	36.4	89.4	166.3	277.3
Female.....	20.9	*	*	*	*	0.7	2.8	9.6	24.4	57.2	116.6	228.1
Non-Hispanic white, both sexes.....	22.7	*	*	*	*	0.7	2.8	10.1	28.5	69.9	135.0	242.1
Male.....	23.2	*	*	*	*	0.8	3.1	11.6	34.5	87.4	164.5	276.8
Female.....	22.2	*	*	*	*	0.7	2.5	8.5	22.8	54.9	115.0	226.8
Non-Hispanic black, both sexes.....	18.8	*	*	*	*	1.0	5.2	19.2	48.9	102.2	175.0	282.9
Male.....	19.0	*	*	*	*	1.0	5.4	22.4	59.9	128.2	217.0	321.8
Female.....	18.7	*	*	*	*	0.9	5.0	16.4	40.1	83.8	151.2	267.8

Worktable Orig250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 2003

[Rates per 100,000 population, in specified group. Rates based on populations estimated as of July 1, 2003. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of liver and intrahepatic bile ducts (C22)												
All origins 3/, both sexes.....	5.1	*	0.1	*	0.1	0.2	0.9	5.4	9.9	19.8	30.7	33.4
Male.....	6.6	*	*	*	0.1	0.3	1.3	8.9	14.9	27.9	44.6	47.5
Female.....	3.5	*	*	*	*	0.2	0.5	2.1	5.3	13.0	21.4	27.2
Hispanic, both sexes.....	3.8	*	*	*	*	*	0.8	7.5	16.1	32.5	49.8	51.8
Male.....	4.8	*	*	*	*	1.2	12.5	24.3	45.2	66.0	55.4	
Female.....	2.7	*	*	*	*	*	*	2.5	8.8	22.4	38.7	50.0
Non-Hispanic 4/, both sexes.....	5.2	*	*	*	0.1	0.2	0.9	5.2	9.4	18.8	29.7	32.6
Male.....	6.9	*	*	*	*	0.3	1.3	8.4	14.1	26.6	43.5	47.1
Female.....	3.6	*	*	*	*	0.2	0.5	2.1	5.0	12.3	20.4	26.2
Non-Hispanic white, both sexes.....	5.1	*	*	*	0.1	0.1	0.7	4.2	8.0	17.0	28.2	31.1
Male.....	6.7	*	*	*	*	*	1.0	6.7	11.9	24.3	41.6	45.0
Female.....	3.6	*	*	*	*	*	0.5	1.7	4.2	10.7	19.2	25.0
Non-Hispanic black, both sexes.....	5.1	*	*	*	*	0.5	1.4	10.0	16.5	23.1	31.7	41.9
Male.....	7.0	*	*	*	*	*	2.2	17.3	26.4	33.8	44.7	63.9
Female.....	3.3	*	*	*	*	*	*	3.6	8.6	15.5	24.3	33.4
Malignant neoplasm of pancreas (C25)												
All origins 3/, both sexes.....	10.6	*	*	*	*	0.2	1.2	6.2	19.1	44.2	75.4	95.2
Male.....	10.5	*	*	*	*	0.2	1.5	7.7	23.1	50.9	85.4	95.6
Female.....	10.6	*	*	*	*	0.1	1.0	4.8	15.3	38.6	68.8	95.0
Hispanic, both sexes.....	3.7	*	*	*	*	*	0.8	4.1	14.4	35.1	60.3	79.8
Male.....	3.6	*	*	*	*	*	1.0	5.0	17.8	40.0	67.1	70.8
Female.....	3.8	*	*	*	*	*	*	3.2	11.3	31.2	55.6	84.4
Non-Hispanic 4/, both sexes.....	11.7	*	*	*	*	0.2	1.3	6.4	19.4	44.8	76.1	95.7
Male.....	11.7	*	*	*	*	0.2	1.6	8.0	23.5	51.6	86.2	96.5
Female.....	11.6	*	*	*	*	0.1	1.0	4.9	15.6	39.1	69.3	95.3
Non-Hispanic white, both sexes.....	12.5	*	*	*	*	0.2	1.2	6.0	18.7	44.1	75.4	94.9
Male.....	12.6	*	*	*	*	0.2	1.5	7.5	22.8	51.4	86.0	96.8
Female.....	12.4	*	*	*	*	*	1.0	4.6	14.8	38.0	68.1	94.0
Non-Hispanic black, both sexes.....	9.8	*	*	*	*	*	1.9	10.3	27.7	57.5	94.0	113.3
Male.....	9.6	*	*	*	*	*	2.5	13.6	33.7	64.8	99.0	106.5
Female.....	10.0	*	*	*	*	*	1.3	7.4	22.8	52.4	91.3	116.0
Malignant neoplasm of larynx (C32)												
All origins 3/, both sexes.....	1.3	*	*	*	*	*	0.2	1.0	3.4	6.2	6.9	6.5
Male.....	2.1	*	*	*	*	*	0.3	1.8	5.8	10.8	13.7	15.3
Female.....	0.5	*	*	*	*	*	*	0.3	1.2	2.4	2.4	2.7
Hispanic, both sexes.....	0.4	*	*	*	*	*	*	0.5	2.1	3.6	5.9	*
Male.....	0.7	*	*	*	*	*	*	*	4.3	7.6	11.9	*
Female.....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes.....	1.4	*	*	*	*	*	0.2	1.1	3.5	6.3	6.9	6.5
Male.....	2.3	*	*	*	*	*	0.3	1.8	5.9	10.9	13.7	15.0
Female.....	0.6	*	*	*	*	*	*	0.3	1.3	2.5	2.4	2.7
Non-Hispanic white, both sexes.....	1.4	*	*	*	*	*	0.2	0.9	3.2	6.0	6.6	6.1
Male.....	2.3	*	*	*	*	*	0.3	1.5	5.2	9.9	12.8	14.1
Female.....	0.6	*	*	*	*	*	*	0.3	1.2	2.5	2.4	2.6
Non-Hispanic black, both sexes.....	1.9	*	*	*	*	*	*	2.6	7.7	11.6	11.9	11.3
Male.....	3.3	*	*	*	*	*	*	4.6	14.0	24.0	28.3	27.7
Female.....	0.6	*	*	*	*	*	*	0.8	2.6	2.8	*	*

Worktable Orig250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 2003

[Rates per 100,000 population, in specified group. Rates based on populations estimated as of July 1, 2003. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of trachea, bronchus and lung (C33-C34)												
All origins 3/, both sexes.....	54.4	*	*	*	0.1	0.4	5.6	30.3	111.0	269.3	377.8	298.9
Male.....	62.9	*	*	*	*	0.4	6.1	36.5	136.7	346.6	525.1	475.1
Female.....	46.1	*	*	*	*	0.4	5.1	24.4	87.1	204.8	279.4	221.0
Hispanic, both sexes.....	10.4	*	*	*	*	0.3	1.6	10.2	38.1	105.8	183.8	189.2
Male.....	12.8	*	*	*	*	*	1.7	12.6	55.4	161.8	277.3	290.9
Female.....	7.8	*	*	*	*	*	1.4	7.9	22.4	61.2	119.6	137.5
Non-Hispanic 4/, both sexes.....	61.2	*	*	*	0.1	0.4	6.2	32.3	116.6	280.3	387.2	303.0
Male.....	71.1	*	*	*	*	0.4	6.8	38.9	142.9	358.3	537.2	483.1
Female.....	51.8	*	*	*	*	0.4	5.6	26.0	92.2	214.9	287.2	223.9
Non-Hispanic white, both sexes.....	67.5	*	*	*	*	0.4	6.1	31.0	117.0	286.5	392.6	306.7
Male.....	77.3	*	*	*	*	0.4	6.5	36.5	139.4	359.6	537.0	486.8
Female.....	58.1	*	*	*	*	0.4	5.7	25.6	95.7	223.9	295.1	227.4
Non-Hispanic black, both sexes.....	44.5	*	*	*	*	*	8.0	48.5	147.4	293.9	389.4	283.5
Male.....	57.0	*	*	*	*	*	9.7	65.3	215.1	434.9	632.5	509.3
Female.....	33.1	*	*	*	*	*	6.5	34.1	93.1	193.5	252.1	196.0
Malignant melanoma of skin (C43)												
All origins 3/, both sexes.....	2.7	*	*	*	0.1	0.5	1.3	2.9	5.5	8.7	14.6	17.6
Male.....	3.5	*	*	*	0.1	0.6	1.5	3.8	7.8	12.6	23.3	30.2
Female.....	1.9	*	*	*	*	0.4	1.0	2.1	3.5	5.4	8.7	12.1
Hispanic, both sexes.....	0.4	*	*	*	*	*	0.9	1.4	3.0	3.9	3.9	*
Male.....	0.4	*	*	*	*	*	*	*	3.9	*	*	*
Female.....	0.4	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes.....	3.0	*	*	*	0.1	0.6	1.4	3.1	5.9	9.1	15.1	18.1
Male.....	4.0	*	*	*	0.1	0.7	1.7	4.0	8.2	13.2	24.2	30.9
Female.....	2.2	*	*	*	*	0.4	1.1	2.2	3.6	5.7	9.0	12.4
Non-Hispanic white, both sexes.....	3.8	*	*	*	0.1	0.7	1.8	3.7	6.8	10.4	16.6	19.6
Male.....	4.9	*	*	*	0.2	0.9	2.1	4.8	9.4	14.9	26.5	33.8
Female.....	2.6	*	*	*	*	0.5	1.4	2.6	4.2	6.4	9.8	13.3
Non-Hispanic black, both sexes.....	0.3	*	*	*	*	*	*	*	0.9	1.6	2.9	*
Male.....	0.3	*	*	*	*	*	*	*	*	*	*	*
Female.....	0.4	*	*	*	*	*	*	*	*	*	*	*
Malignant neoplasm of breast (C50)												
All origins 3/, both sexes.....	14.4	*	*	*	*	1.0	6.1	15.6	29.6	45.5	74.9	132.5
Male.....	0.3	*	*	*	*	*	*	0.2	0.5	1.1	2.0	3.9
Female.....	28.2	*	*	*	*	2.1	12.2	30.4	56.6	82.6	123.7	189.4
Hispanic, both sexes.....	4.7	*	*	*	*	0.7	4.3	10.5	20.5	27.1	42.0	77.7
Male.....	*	*	*	*	*	*	*	*	*	*	*	*
Female.....	9.8	*	*	*	*	1.6	9.1	21.0	38.8	48.4	70.8	114.8
Non-Hispanic 4/, both sexes.....	15.9	*	*	*	*	1.1	6.4	16.1	30.3	46.7	76.4	134.7
Male.....	0.3	*	*	*	*	*	*	0.3	0.6	1.1	2.1	3.8
Female.....	30.8	*	*	*	*	2.1	12.6	31.3	57.9	84.9	126.0	192.2
Non-Hispanic white, both sexes.....	16.8	*	*	*	*	0.9	5.7	14.8	29.0	46.3	76.7	134.7
Male.....	0.3	*	*	*	*	*	*	0.2	0.5	1.1	2.2	3.9
Female.....	32.6	*	*	*	*	1.8	11.3	29.1	56.0	85.1	127.1	192.3
Non-Hispanic black, both sexes.....	15.5	*	*	*	*	2.3	11.4	27.1	46.1	60.1	91.3	163.3
Male.....	0.3	*	*	*	*	*	*	*	*	*	*	*
Female.....	29.4	*	*	*	*	4.3	21.4	50.1	82.4	101.6	141.8	224.5

Worktable Orig250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 2003

[Rates per 100,000 population, in specified group. Rates based on populations estimated as of July 1, 2003. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of cervix uteri (C53)												
All origins 3/, both sexes.....	1.3	*	*	*	*	0.5	1.4	2.2	2.8	3.1	4.3	5.7
Male.....
Female.....	2.7	*	*	*	*	1.0	2.8	4.3	5.5	5.6	7.2	8.3
Hispanic, both sexes.....	1.0	*	*	*	*	0.4	1.4	2.9	3.1	3.5	6.6	*
Male.....
Female.....	2.0	*	*	*	*	0.9	3.0	5.7	5.9	6.3	11.1	*
Non-Hispanic 4/, both sexes.....	1.4	*	*	*	*	0.5	1.4	2.1	2.8	3.0	4.2	5.7
Male.....
Female.....	2.7	*	*	*	*	1.0	2.8	4.1	5.4	5.6	7.0	8.2
Non-Hispanic white, both sexes.....	1.3	*	*	*	*	0.4	1.3	1.9	2.5	2.5	3.4	4.8
Male.....
Female.....	2.5	*	*	*	*	0.9	2.6	3.7	4.9	4.7	5.7	7.0
Non-Hispanic black, both sexes.....	2.2	*	*	*	*	0.9	2.3	3.7	5.7	7.5	11.6	14.9
Male.....
Female.....	4.1	*	*	*	*	1.8	4.3	6.9	10.2	12.8	18.1	20.6
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)												
All origins 3/, both sexes.....	2.4	*	*	*	*	0.1	0.3	1.3	4.7	10.1	15.2	22.8
Male.....
Female.....	4.7	*	*	*	*	0.1	0.7	2.6	9.0	18.5	25.3	33.0
Hispanic, both sexes.....	0.8	*	*	*	*	*	0.4	1.2	2.9	7.0	9.8	20.7
Male.....
Female.....	1.7	*	*	*	*	*	0.9	2.3	5.6	12.5	16.6	31.3
Non-Hispanic 4/, both sexes.....	2.6	*	*	*	*	0.1	0.3	1.3	4.8	10.3	15.4	22.9
Male.....
Female.....	5.1	*	*	*	*	0.1	0.7	2.6	9.2	18.9	25.6	33.0
Non-Hispanic white, both sexes.....	2.7	*	*	*	*	*	0.3	1.3	4.4	9.3	14.5	22.8
Male.....
Female.....	5.2	*	*	*	*	*	0.6	2.5	8.5	17.2	24.4	32.8
Non-Hispanic black, both sexes.....	3.0	*	*	*	*	*	0.5	1.7	9.1	21.4	28.0	30.3
Male.....
Female.....	5.7	*	*	*	*	*	1.0	3.2	16.5	36.7	43.8	42.1
Malignant neoplasm of ovary (C56)												
All origins 3/, both sexes.....	5.0	*	*	*	0.1	0.2	1.0	4.0	10.4	19.5	32.6	38.4
Male.....
Female.....	9.9	*	*	*	0.1	0.4	2.0	7.8	20.0	35.7	54.4	55.4
Hispanic, both sexes.....	1.6	*	*	*	*	*	0.8	2.7	6.4	12.7	22.0	28.5
Male.....
Female.....	3.3	*	*	*	*	*	1.7	5.5	12.2	22.8	37.1	43.0
Non-Hispanic 4/, both sexes.....	5.6	*	*	*	0.1	0.2	1.0	4.1	10.7	19.9	33.1	38.8
Male.....
Female.....	10.9	*	*	*	0.1	0.5	2.1	8.1	20.5	36.6	55.2	55.9
Non-Hispanic white, both sexes.....	6.3	*	*	*	*	0.3	1.1	4.2	11.1	20.5	34.2	39.5
Male.....
Female.....	12.3	*	*	*	*	0.5	2.2	8.4	21.6	38.1	57.2	56.8
Non-Hispanic black, both sexes.....	3.2	*	*	*	*	*	1.0	3.9	8.8	17.8	27.8	37.2
Male.....
Female.....	6.2	*	*	*	*	*	1.9	7.2	15.9	30.5	43.5	51.6

Worktable Orig250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 2003

[Rates per 100,000 population, in specified group. Rates based on populations estimated as of July 1, 2003. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of prostate (C61)												
All origins 3/, both sexes.....	10.2	*	*	*	*	*	0.1	1.0	7.4	32.9	95.5	184.9
Male.....	20.7	*	*	*	*	*	0.1	2.1	15.4	72.3	238.3	603.3
Female.....
Hispanic, both sexes.....	3.1	*	*	*	*	*	*	0.7	5.3	25.5	74.5	151.8
Male.....	5.9	*	*	*	*	*	*	1.3	11.2	57.6	182.9	451.0
Female.....
Non-Hispanic 4/, both sexes.....	11.3	*	*	*	*	*	0.1	1.1	7.6	33.3	96.4	186.0
Male.....	23.1	*	*	*	*	*	0.1	2.2	15.8	73.1	240.9	609.4
Female.....
Non-Hispanic white, both sexes.....	11.6	*	*	*	*	*	*	0.9	6.3	29.8	89.1	178.2
Male.....	23.6	*	*	*	*	*	*	1.8	13.0	64.5	221.0	583.0
Female.....
Non-Hispanic black, both sexes.....	13.3	*	*	*	*	*	*	2.7	20.1	74.6	198.8	304.6
Male.....	27.9	*	*	*	*	*	*	5.7	45.1	179.5	550.9	1,091.0
Female.....
Malignant neoplasms of kidney and renal pelvis (C64-C65)												
All origins 3/, both sexes.....	4.2	*	*	0.1	0.1	0.1	0.7	2.9	9.0	16.9	27.0	33.2
Male.....	5.4	*	*	*	*	0.2	1.0	4.3	13.3	23.7	39.5	48.7
Female.....	3.1	*	*	0.1	*	0.1	0.4	1.6	4.9	11.2	18.7	26.4
Hispanic, both sexes.....	1.7	*	*	*	*	*	0.5	2.1	8.1	16.2	23.6	26.9
Male.....	2.1	*	*	*	*	*	0.7	2.8	11.0	22.3	32.2	38.5
Female.....	1.4	*	*	*	*	*	*	1.5	5.6	11.3	17.6	21.1
Non-Hispanic 4/, both sexes.....	4.6	*	*	0.1	*	0.2	0.8	3.0	9.0	16.9	27.2	33.4
Male.....	5.9	*	*	*	*	0.2	1.1	4.4	13.5	23.7	39.8	49.0
Female.....	3.4	*	*	*	*	0.1	0.4	1.7	4.9	11.1	18.7	26.6
Non-Hispanic white, both sexes.....	5.1	*	*	0.1	*	0.1	0.7	3.1	9.2	17.4	27.8	34.0
Male.....	6.6	*	*	*	*	*	1.0	4.5	13.7	24.3	40.7	50.4
Female.....	3.8	*	*	*	*	*	0.4	1.7	4.8	11.6	19.1	26.8
Non-Hispanic black, both sexes.....	3.0	*	*	*	*	*	1.1	3.3	9.7	14.8	24.2	34.2
Male.....	3.8	*	*	*	*	*	1.7	5.0	14.1	22.3	37.2	44.7
Female.....	2.3	*	*	*	*	*	0.7	1.8	6.2	9.5	16.8	30.1
Malignant neoplasm of bladder (C67)												
All origins 3/, both sexes.....	4.3	*	*	*	*	*	0.2	1.3	5.0	14.9	36.4	64.8
Male.....	6.0	*	*	*	*	*	0.3	2.0	7.8	23.9	63.1	125.1
Female.....	2.6	*	*	*	*	*	0.2	0.6	2.4	7.4	18.5	38.1
Hispanic, both sexes.....	1.0	*	*	*	*	*	*	0.7	2.6	7.9	20.8	42.0
Male.....	1.3	*	*	*	*	*	*	*	4.0	13.1	33.4	77.0
Female.....	0.7	*	*	*	*	*	*	*	*	3.7	12.1	24.2
Non-Hispanic 4/, both sexes.....	4.8	*	*	*	*	*	0.2	1.3	5.2	15.4	37.1	65.6
Male.....	6.8	*	*	*	*	*	0.3	2.1	8.1	24.7	64.5	127.2
Female.....	2.9	*	*	*	*	*	0.2	0.6	2.5	7.6	18.8	38.6
Non-Hispanic white, both sexes.....	5.5	*	*	*	*	*	0.2	1.3	5.5	16.3	38.3	67.6
Male.....	7.9	*	*	*	*	*	0.3	2.2	8.6	26.1	67.2	133.6
Female.....	3.3	*	*	*	*	*	0.2	0.5	2.5	7.8	18.7	38.6
Non-Hispanic black, both sexes.....	2.5	*	*	*	*	*	0.4	1.7	4.1	12.3	32.8	53.8
Male.....	2.9	*	*	*	*	*	*	2.2	5.5	17.8	49.6	79.9
Female.....	2.1	*	*	*	*	*	*	1.2	2.9	8.3	23.3	43.7

Worktable Orig250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 2003

[Rates per 100,000 population, in specified group. Rates based on populations estimated as of July 1, 2003. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)												
All origins 3/, both sexes.....	7.4	*	*	0.1	0.3	0.6	1.4	3.8	10.8	26.8	56.1	78.8
Male.....	8.0	*	*	0.1	0.4	0.8	1.8	4.8	13.8	33.9	69.1	103.6
Female.....	6.8	*	*	*	0.3	0.4	1.0	2.8	8.0	20.9	47.4	67.8
Hispanic, both sexes.....	2.6	*	*	*	0.4	0.6	1.4	3.0	8.4	19.1	40.4	54.4
Male.....	2.8	*	*	*	*	0.8	1.8	3.4	10.5	23.8	45.3	57.0
Female.....	2.5	*	*	*	*	*	0.9	2.7	6.6	15.3	37.1	53.1
Non-Hispanic 4/, both sexes.....	8.1	*	*	0.1	0.3	0.6	1.4	3.9	10.9	27.3	56.8	79.6
Male.....	8.8	*	*	0.1	0.4	0.8	1.8	5.0	14.0	34.5	70.3	105.6
Female.....	7.4	*	*	*	0.3	0.5	1.0	2.8	8.1	21.3	47.8	68.2
Non-Hispanic white, both sexes.....	9.3	*	*	0.1	0.3	0.5	1.4	3.8	11.4	28.8	60.0	84.3
Male.....	10.1	*	*	*	0.4	0.7	1.8	4.9	14.6	36.6	74.4	112.0
Female.....	8.5	*	*	*	0.3	0.4	1.0	2.8	8.4	22.1	50.3	72.2
Non-Hispanic black, both sexes.....	3.7	*	*	*	0.4	1.1	2.0	4.9	9.1	18.4	28.6	31.2
Male.....	3.9	*	*	*	*	1.4	2.4	6.7	11.5	20.7	31.2	35.2
Female.....	3.5	*	*	*	*	0.9	1.5	3.4	7.1	16.9	27.2	29.7
Leukemia (C91-C95)												
All origins 3/, both sexes.....	7.4	0.7	0.8	0.8	1.0	1.1	1.8	3.3	9.3	25.7	53.9	80.3
Male.....	8.4	*	0.9	0.9	1.2	1.4	2.0	4.0	11.9	35.1	75.3	111.8
Female.....	6.4	1.1	0.8	0.7	0.7	0.8	1.6	2.7	7.0	17.8	39.6	66.4
Hispanic, both sexes.....	2.9	*	1.2	1.1	1.4	1.1	1.7	3.6	6.3	13.9	32.9	47.2
Male.....	3.2	*	1.4	1.3	1.7	1.3	1.9	4.1	7.6	18.3	40.6	61.6
Female.....	2.6	*	*	1.0	1.0	0.9	1.5	3.2	5.2	10.4	27.7	39.8
Non-Hispanic 4/, both sexes.....	8.1	0.8	0.7	0.8	0.9	1.1	1.8	3.3	9.6	26.4	54.9	81.6
Male.....	9.3	*	0.8	0.8	1.1	1.4	2.0	4.0	12.2	36.2	77.0	113.7
Female.....	7.0	*	0.7	0.7	0.7	0.8	1.6	2.7	7.1	18.3	40.2	67.4
Non-Hispanic white, both sexes.....	9.1	1.0	0.8	0.8	0.9	1.1	1.8	3.3	9.8	27.5	57.1	84.7
Male.....	10.5	*	0.8	0.9	1.1	1.3	1.9	4.0	12.5	37.7	80.6	119.0
Female.....	7.8	*	0.7	0.6	0.7	0.8	1.6	2.6	7.3	18.8	41.3	69.6
Non-Hispanic black, both sexes.....	4.8	*	*	0.7	0.9	1.2	2.4	3.8	9.8	23.4	42.1	60.4
Male.....	5.2	*	*	0.7	1.2	1.6	2.8	4.6	12.6	31.5	54.0	78.8
Female.....	4.4	*	*	0.8	0.7	0.9	1.9	3.1	7.5	17.6	35.4	53.2
Multiple myeloma and immunoproliferative neoplasms (C88,C90)												
All origins 3/, both sexes.....	3.9	*	*	*	*	*	0.3	1.7	5.9	16.5	31.1	36.3
Male.....	4.1	*	*	*	*	*	0.4	2.1	7.4	19.8	37.7	50.4
Female.....	3.7	*	*	*	*	*	0.2	1.4	4.6	13.8	26.7	30.1
Hispanic, both sexes.....	1.3	*	*	*	*	*	*	1.4	5.2	12.6	23.9	24.9
Male.....	1.2	*	*	*	*	*	*	1.6	5.8	13.3	25.7	*
Female.....	1.4	*	*	*	*	*	*	1.2	4.7	12.1	22.6	22.7
Non-Hispanic 4/, both sexes.....	4.3	*	*	*	*	*	0.3	1.8	6.0	16.8	31.4	36.7
Male.....	4.6	*	*	*	*	*	0.5	2.2	7.5	20.2	38.2	51.2
Female.....	4.0	*	*	*	*	*	0.2	1.4	4.6	13.8	26.8	30.3
Non-Hispanic white, both sexes.....	4.4	*	*	*	*	*	0.3	1.4	5.4	15.6	29.8	35.3
Male.....	4.7	*	*	*	*	*	0.4	1.8	6.8	19.3	37.0	50.5
Female.....	4.0	*	*	*	*	*	0.2	1.1	4.0	12.4	24.9	28.7
Non-Hispanic black, both sexes.....	5.0	*	*	*	*	*	0.8	4.3	12.2	31.4	55.8	60.4
Male.....	4.9	*	*	*	*	*	0.9	5.3	15.7	34.5	64.1	74.6
Female.....	5.1	*	*	*	*	*	*	3.4	9.3	29.2	51.2	54.9

Worktable Orig250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 2003

[Rates per 100,000 population, in specified group. Rates based on populations estimated as of July 1, 2003. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)												
All origins 3/, both sexes.....	0.0	*	*	*	*	*	*	*	*	*	*	*
Male.....	0.0	*	*	*	*	*	*	*	*	*	*	*
Female.....	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes.....	*	*	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes.....	0.0	*	*	*	*	*	*	*	*	*	*	*
Male.....	0.0	*	*	*	*	*	*	*	*	*	*	*
Female.....	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes.....	0.0	*	*	*	*	*	*	*	*	*	*	*
Male.....	0.0	*	*	*	*	*	*	*	*	*	*	*
Female.....	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes.....	*	*	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*	*	*
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)												
All origins 3/, both sexes.....	21.5	*	0.7	0.7	1.3	1.7	4.2	13.8	35.2	77.9	146.7	224.0
Male.....	22.3	*	0.8	0.7	1.6	2.1	4.7	15.7	41.5	93.4	180.8	275.0
Female.....	20.8	*	0.5	0.6	0.9	1.3	3.8	11.9	29.3	64.8	123.9	201.4
Hispanic, both sexes.....	7.7	*	0.6	0.7	1.3	1.5	3.3	9.8	23.7	58.7	109.4	168.9
Male.....	7.9	*	*	0.7	1.6	2.1	3.3	11.2	27.7	71.0	127.3	194.0
Female.....	7.6	*	*	0.7	1.1	0.8	3.2	8.4	20.1	48.8	97.2	156.3
Non-Hispanic 4/, both sexes.....	23.7	*	0.7	0.7	1.2	1.8	4.4	14.1	36.0	79.0	148.4	225.8
Male.....	24.6	*	0.8	0.7	1.5	2.1	4.9	16.1	42.4	94.7	183.4	278.2
Female.....	22.8	*	0.6	0.6	0.9	1.5	3.8	12.2	30.0	65.9	125.1	202.7
Non-Hispanic white, both sexes.....	25.9	*	0.6	0.7	1.3	1.7	4.2	13.4	35.4	79.5	150.3	229.9
Male.....	26.8	*	0.7	0.7	1.6	2.1	4.8	15.2	41.5	94.8	186.1	284.6
Female.....	25.0	*	0.6	0.6	0.9	1.3	3.7	11.7	29.6	66.4	126.2	205.8
Non-Hispanic black, both sexes.....	17.9	*	0.9	0.8	1.2	2.4	5.9	21.1	49.0	90.0	151.8	206.4
Male.....	18.8	*	*	0.8	1.5	2.4	6.6	25.3	62.2	113.6	189.9	239.7
Female.....	17.1	*	*	0.8	0.8	2.3	5.3	17.5	38.5	73.2	130.4	193.6
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)												
All origins 3/, both sexes.....	4.7	1.5	0.3	0.2	0.2	0.4	0.8	1.7	4.4	12.8	36.4	80.5
Male.....	4.7	1.6	0.4	0.2	0.3	0.5	0.9	1.8	5.1	16.0	46.2	101.7
Female.....	4.7	1.5	0.3	0.2	0.2	0.4	0.6	1.5	3.8	10.2	29.9	71.1
Hispanic, both sexes.....	1.5	*	*	*	0.3	0.4	0.4	1.1	3.1	8.7	28.9	48.2
Male.....	1.5	*	*	*	*	0.5	*	1.2	3.9	10.0	33.7	52.3
Female.....	1.6	*	*	*	*	*	*	1.0	2.3	7.6	25.5	46.1
Non-Hispanic 4/, both sexes.....	5.2	1.5	0.4	0.2	0.2	0.4	0.8	1.7	4.5	13.1	36.8	81.8
Male.....	5.2	1.6	0.4	0.2	0.2	0.5	0.9	1.9	5.2	16.3	46.7	103.9
Female.....	5.1	1.3	*	0.2	0.2	0.4	0.7	1.6	3.9	10.4	30.2	72.1
Non-Hispanic white, both sexes.....	5.8	1.4	0.3	0.2	0.2	0.5	0.8	1.7	4.5	13.4	38.1	84.4
Male.....	5.9	1.7	*	0.2	0.3	0.5	0.9	1.9	5.1	16.8	48.4	107.9
Female.....	5.8	*	*	*	0.2	0.5	0.6	1.5	4.0	10.4	31.1	74.0
Non-Hispanic black, both sexes.....	3.0	*	*	*	*	0.5	1.2	2.5	5.3	13.3	29.6	58.9
Male.....	3.0	*	*	*	*	*	1.2	2.5	6.9	16.5	36.6	68.2
Female.....	3.1	*	*	*	*	*	1.2	2.5	4.1	11.1	25.6	55.3

Worktable Orig250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 2003

[Rates per 100,000 population, in specified group. Rates based on populations estimated as of July 1, 2003. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Anemias (D50-D64)												
All origins 3/, both sexes.....	1.6	0.5	0.2	0.1	0.3	0.4	0.4	0.6	1.0	2.6	8.8	40.7
Male.....	1.3	*	*	0.1	0.2	0.4	0.4	0.6	1.2	3.0	9.3	38.0
Female.....	1.9	*	*	0.1	0.3	0.4	0.4	0.6	0.8	2.3	8.5	41.9
Hispanic, both sexes.....	0.5	*	*	*	*	*	*	*	*	2.0	7.5	29.0
Male.....	0.4	*	*	*	*	*	*	*	*	*	*	30.8
Female.....	0.6	*	*	*	*	*	*	*	*	*	7.6	28.1
Non-Hispanic 4/, both sexes.....	1.8	0.6	*	0.1	0.3	0.5	0.4	0.6	1.0	2.7	8.9	41.2
Male.....	1.4	*	*	0.1	0.3	0.5	0.5	0.7	1.2	3.1	9.4	38.3
Female.....	2.1	*	*	0.1	0.3	0.4	0.4	0.6	0.8	2.3	8.6	42.4
Non-Hispanic white, both sexes.....	1.6	*	*	*	*	0.1	0.1	0.3	0.6	2.4	8.6	41.0
Male.....	1.3	*	*	*	*	*	*	0.3	0.7	2.6	9.2	38.1
Female.....	2.0	*	*	*	*	*	*	0.2	0.5	2.1	8.2	42.2
Non-Hispanic black, both sexes.....	2.8	*	*	0.4	1.3	2.2	2.4	3.1	4.3	5.6	14.5	50.3
Male.....	2.8	*	*	*	1.3	2.3	2.6	3.4	5.6	8.0	15.3	50.1
Female.....	2.9	*	*	*	1.3	2.1	2.2	2.8	3.3	4.0	14.0	50.3
Diabetes mellitus (E10-E14)												
All origins 3/, both sexes.....	25.5	*	*	0.1	0.4	1.6	4.6	13.9	38.5	90.8	181.1	317.5
Male.....	24.8	*	*	*	0.5	1.9	5.7	16.7	45.0	104.0	207.2	340.7
Female.....	26.2	*	*	*	0.3	1.4	3.5	11.2	32.4	79.8	163.6	407.2
Hispanic, both sexes.....	15.5	*	*	*	*	0.9	3.3	15.5	53.7	131.8	272.1	428.1
Male.....	14.4	*	*	*	*	0.8	3.9	18.1	62.4	146.3	297.2	426.4
Female.....	16.6	*	*	*	*	0.9	2.7	13.0	45.8	120.3	254.9	428.9
Non-Hispanic 4/, both sexes.....	27.0	*	*	0.1	0.4	1.8	4.8	13.6	37.1	87.7	175.8	312.0
Male.....	26.4	*	*	*	0.5	2.1	6.0	16.5	43.5	100.9	201.8	335.9
Female.....	27.6	*	*	*	0.3	1.5	3.6	10.9	31.2	76.7	158.5	301.5
Non-Hispanic white, both sexes.....	26.6	*	*	*	0.3	1.5	4.1	11.3	31.4	76.8	161.6	293.8
Male.....	26.6	*	*	*	0.4	1.7	5.2	14.1	37.9	91.3	191.7	327.3
Female.....	26.5	*	*	*	0.3	1.3	2.9	8.5	25.2	64.4	141.3	279.0
Non-Hispanic black, both sexes.....	35.0	*	*	*	0.7	4.0	9.9	30.8	87.5	188.9	340.0	532.2
Male.....	30.9	*	*	*	0.9	4.9	11.9	34.6	97.8	201.2	342.8	457.1
Female.....	38.7	*	*	*	*	3.2	8.1	27.6	79.2	180.2	338.5	561.3
Nutritional deficiencies (E40-E64)												
All origins 3/, both sexes.....	1.1	*	*	*	*	*	0.1	0.2	0.6	1.9	7.7	35.5
Male.....	0.8	*	*	*	*	*	*	0.2	0.6	2.0	7.8	30.6
Female.....	1.5	*	*	*	*	*	*	0.3	0.6	1.7	7.7	37.6
Hispanic, both sexes.....	0.3	*	*	*	*	*	*	*	*	*	5.6	28.5
Male.....	0.3	*	*	*	*	*	*	*	*	*	*	*
Female.....	0.4	*	*	*	*	*	*	*	*	*	6.1	28.9
Non-Hispanic 4/, both sexes.....	1.3	*	*	*	*	*	0.1	0.3	0.6	1.9	7.8	35.7
Male.....	0.9	*	*	*	*	*	*	0.2	0.6	2.0	7.9	30.7
Female.....	1.6	*	*	*	*	*	*	0.3	0.7	1.8	7.7	37.9
Non-Hispanic white, both sexes.....	1.4	*	*	*	*	*	0.1	0.2	0.6	1.7	7.4	34.4
Male.....	0.9	*	*	*	*	*	*	0.2	0.5	1.8	7.2	29.6
Female.....	1.8	*	*	*	*	*	*	0.2	0.6	1.7	7.6	36.5
Non-Hispanic black, both sexes.....	1.2	*	*	*	*	*	*	0.7	1.4	3.2	13.2	55.9
Male.....	1.0	*	*	*	*	*	*	*	1.8	4.5	17.9	49.0
Female.....	1.4	*	*	*	*	*	*	*	*	2.4	10.6	58.6

Worktable Orig250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 2003

[Rates per 100,000 population, in specified group. Rates based on populations estimated as of July 1, 2003. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Parkinson's disease (G20-G21)												
All origins 3/, both sexes.....	6.2	*	*	*	*	*	*	0.2	1.3	12.7	67.8	138.2
Male.....	7.1	*	*	*	*	*	*	0.2	1.6	18.2	100.7	222.4
Female.....	5.3	*	*	*	*	*	*	0.1	0.9	8.1	45.8	101.0
Hispanic, both sexes.....	1.3	*	*	*	*	*	*	*	*	7.9	40.3	77.7
Male.....	1.5	*	*	*	*	*	*	*	*	12.4	58.3	129.3
Female.....	1.1	*	*	*	*	*	*	*	*	4.3	27.9	51.6
Non-Hispanic 4/, both sexes.....	7.0	*	*	*	*	*	*	0.2	1.3	13.0	69.1	140.6
Male.....	8.0	*	*	*	*	*	*	0.2	1.7	18.5	102.7	226.5
Female.....	5.9	*	*	*	*	*	*	0.2	0.9	8.4	46.6	102.8
Non-Hispanic white, both sexes.....	8.3	*	*	*	*	*	*	0.2	1.3	14.0	73.7	148.8
Male.....	9.5	*	*	*	*	*	*	0.2	1.7	19.8	109.0	237.9
Female.....	7.1	*	*	*	*	*	*	0.2	1.0	9.0	49.8	109.6
Non-Hispanic black, both sexes.....	1.7	*	*	*	*	*	*	*	1.2	7.2	29.4	53.5
Male.....	2.0	*	*	*	*	*	*	*	1.8	10.6	45.6	99.1
Female.....	1.4	*	*	*	*	*	*	*	*	4.8	20.2	35.9
Alzheimer's disease (G30)												
All origins 3/, both sexes.....	21.8	*	*	*	*	*	*	0.2	2.0	20.9	164.4	802.4
Male.....	12.8	*	*	*	*	*	*	0.2	1.8	20.2	147.6	606.5
Female.....	30.5	*	*	*	*	*	*	0.2	2.1	21.5	175.7	889.0
Hispanic, both sexes.....	4.6	*	*	*	*	*	*	*	1.3	14.6	103.7	493.4
Male.....	3.1	*	*	*	*	*	*	*	*	13.8	101.2	426.4
Female.....	6.2	*	*	*	*	*	*	*	*	15.3	105.3	527.3
Non-Hispanic 4/, both sexes.....	24.5	*	*	*	*	*	*	0.2	2.0	21.3	167.3	814.0
Male.....	14.4	*	*	*	*	*	*	0.2	1.9	20.5	149.8	613.6
Female.....	34.1	*	*	*	*	*	*	0.2	2.2	22.0	179.0	902.1
Non-Hispanic white, both sexes.....	28.7	*	*	*	*	*	*	0.2	2.2	22.2	173.9	846.9
Male.....	16.8	*	*	*	*	*	*	0.2	2.0	21.2	154.8	638.1
Female.....	40.2	*	*	*	*	*	*	0.2	2.5	23.0	186.8	938.8
Non-Hispanic black, both sexes.....	10.1	*	*	*	*	*	*	*	1.1	19.6	140.3	591.1
Male.....	6.2	*	*	*	*	*	*	*	*	20.5	138.8	465.6
Female.....	13.7	*	*	*	*	*	*	*	*	18.9	141.1	639.7
Major cardiovascular diseases (I00-I78)												
All origins 3/, both sexes.....	310.3	13.9	1.5	0.8	3.5	10.3	38.2	113.4	284.9	748.3	2,170.8	7,138.0
Male.....	296.3	15.2	1.4	1.0	4.2	12.9	51.1	160.9	393.3	976.3	2,636.1	7,329.3
Female.....	323.9	12.5	1.5	0.7	2.8	7.5	25.3	67.5	184.4	557.8	1,860.0	7,053.5
Hispanic, both sexes.....	93.4	11.0	1.3	0.6	2.5	6.4	20.5	71.9	212.2	594.9	1,683.5	5,176.6
Male.....	92.6	12.2	1.5	0.7	3.3	7.8	27.9	102.8	295.6	779.5	2,007.9	5,079.7
Female.....	94.3	9.7	*	*	1.6	4.6	12.3	41.0	137.0	447.8	1,460.7	5,225.8
Non-Hispanic 4/, both sexes.....	343.8	14.5	1.5	0.9	3.7	11.1	40.7	117.2	289.5	756.6	2,190.9	7,206.9
Male.....	329.5	15.9	1.4	1.0	4.4	14.1	54.6	166.3	399.0	986.2	2,662.3	7,419.6
Female.....	357.5	13.1	1.6	0.8	3.0	8.1	27.1	70.0	187.7	564.0	1,876.4	7,113.4
Non-Hispanic white, both sexes.....	372.4	10.9	1.2	0.8	3.0	9.0	34.8	101.4	262.4	717.8	2,169.7	7,304.7
Male.....	354.1	11.3	0.9	0.9	3.6	11.7	48.5	148.4	368.3	945.5	2,648.3	7,561.6
Female.....	390.0	10.4	1.4	0.6	2.4	6.2	21.2	55.1	161.9	522.8	1,846.4	7,191.6
Non-Hispanic black, both sexes.....	282.2	26.6	2.9	1.5	7.1	23.7	81.3	244.5	565.5	1,224.4	2,745.6	6,923.9
Male.....	277.5	31.7	3.3	1.6	8.7	29.2	101.4	324.7	753.7	1,562.4	3,299.8	6,625.9
Female.....	286.5	21.3	2.4	1.3	5.5	18.8	63.5	175.8	414.5	984.0	2,432.6	7,039.3

Worktable Orig250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 2003

[Rates per 100,000 population, in specified group. Rates based on populations estimated as of July 1, 2003. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of heart (I00-I09,I11,I13,I20-I51)												
All origins 3/, both sexes.....	235.6	11.0	1.2	0.6	2.7	8.2	30.7	92.5	233.2	585.0	1,611.1	5,278.4
Male.....	235.0	12.1	1.1	0.7	3.4	10.5	42.8	136.2	331.7	785.3	2,030.3	5,621.5
Female.....	236.2	9.8	1.3	0.5	2.1	5.7	18.6	50.2	141.9	417.5	1,331.1	5,126.7
Hispanic, both sexes.....	70.9	8.7	1.1	0.4	1.9	4.7	14.9	53.3	165.4	461.2	1,261.0	3,932.9
Male.....	72.2	9.5	*	*	2.7	6.1	21.1	79.6	235.6	625.0	1,543.5	3,874.5
Female.....	69.6	7.8	*	*	1.1	3.1	8.0	27.0	102.1	330.6	1,067.0	3,962.5
Non-Hispanic 4/, both sexes.....	261.0	11.5	1.2	0.7	2.9	8.9	32.9	96.1	237.5	591.6	1,625.0	5,324.5
Male.....	261.4	12.8	1.1	0.8	3.5	11.6	46.1	141.5	337.4	793.4	2,050.1	5,690.6
Female.....	260.5	10.2	1.3	0.6	2.3	6.2	20.1	52.4	144.6	422.4	1,341.5	5,163.6
Non-Hispanic white, both sexes.....	284.0	8.7	0.9	0.6	2.3	7.3	29.2	85.8	219.6	566.7	1,616.7	5,412.2
Male.....	282.9	9.4	0.7	0.7	2.8	9.8	42.2	129.8	317.7	767.3	2,049.9	5,821.0
Female.....	285.1	8.0	1.1	0.5	1.9	4.7	16.2	42.4	126.6	394.8	1,324.0	5,232.2
Non-Hispanic black, both sexes.....	209.4	20.4	2.3	1.2	5.7	18.8	61.4	185.8	434.7	925.6	1,983.5	4,977.8
Male.....	212.6	24.5	2.5	1.3	7.4	23.1	79.7	253.9	589.2	1,210.9	2,449.4	4,897.8
Female.....	206.5	16.1	2.0	1.1	4.0	14.8	45.3	127.4	310.7	722.8	1,720.5	5,008.9
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
All origins 3/, both sexes.....	1.2	*	*	*	*	0.1	0.2	0.5	1.3	3.8	9.6	20.5
Male.....	0.8	*	*	*	*	*	0.2	0.4	1.1	2.9	7.5	16.2
Female.....	1.7	*	*	*	*	*	0.2	0.7	1.5	4.5	11.0	22.3
Hispanic, both sexes.....	0.4	*	*	*	*	*	*	0.5	1.0	3.3	6.7	*
Male.....	0.3	*	*	*	*	*	*	*	*	*	*	*
Female.....	0.5	*	*	*	*	*	*	*	*	4.0	8.4	*
Non-Hispanic 4/, both sexes.....	1.4	*	*	*	*	*	0.2	0.5	1.3	3.8	9.7	20.9
Male.....	0.9	*	*	*	*	*	0.2	0.4	1.1	2.9	7.6	16.6
Female.....	1.8	*	*	*	*	*	0.3	0.7	1.5	4.6	11.1	22.8
Non-Hispanic white, both sexes.....	1.5	*	*	*	*	*	0.2	0.4	1.2	3.9	10.4	22.3
Male.....	1.0	*	*	*	*	*	0.2	0.3	0.9	3.0	8.3	17.7
Female.....	2.1	*	*	*	*	*	0.2	0.5	1.4	4.6	11.8	24.4
Non-Hispanic black, both sexes.....	0.8	*	*	*	*	*	0.5	1.4	2.7	4.0	4.2	7.1
Male.....	0.6	*	*	*	*	*	*	0.9	2.8	3.0	*	*
Female.....	1.0	*	*	*	*	*	*	1.7	2.6	4.6	5.2	*
Hypertensive heart disease (I11)												
All origins 3/, both sexes.....	9.7	*	*	*	0.1	0.8	2.9	7.9	13.1	22.1	51.0	195.0
Male.....	8.8	*	*	*	0.2	1.0	4.0	10.8	17.2	27.1	51.0	142.2
Female.....	10.7	*	*	*	*	0.5	1.9	5.2	9.3	17.8	51.0	218.3
Hispanic, both sexes.....	4.0	*	*	*	*	0.4	1.8	5.7	11.8	22.9	54.2	179.3
Male.....	4.3	*	*	*	*	0.5	2.9	8.3	17.0	30.8	63.7	144.7
Female.....	3.7	*	*	*	*	*	*	3.1	7.2	16.6	47.7	196.9
Non-Hispanic 4/, both sexes.....	10.6	*	*	*	0.1	0.9	3.1	8.1	13.1	21.9	50.7	195.0
Male.....	9.5	*	*	*	0.2	1.1	4.1	10.9	17.1	26.7	50.1	141.7
Female.....	11.7	*	*	*	*	0.6	2.1	5.4	9.4	17.8	51.1	218.5
Non-Hispanic white, both sexes.....	9.7	*	*	*	*	0.5	2.0	5.3	9.5	16.7	44.3	184.9
Male.....	8.2	*	*	*	*	0.8	3.0	7.8	13.0	20.4	43.5	130.8
Female.....	11.1	*	*	*	*	0.3	1.1	2.8	6.3	13.6	44.9	208.7
Non-Hispanic black, both sexes.....	18.5	*	*	*	0.4	2.6	9.8	28.5	45.1	70.3	128.4	334.9
Male.....	18.7	*	*	*	*	3.2	11.5	35.6	57.5	93.4	139.1	291.9
Female.....	18.4	*	*	*	*	2.1	8.2	22.4	35.1	53.9	122.4	351.6

Worktable Orig250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 2003

[Rates per 100,000 population, in specified group. Rates based on populations estimated as of July 1, 2003. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart and renal disease (I13)												
All origins 3/, both sexes.....	1.1	*	*	*	*	0.1	0.2	0.5	1.0	2.2	7.0	25.6
Male.....	0.9	*	*	*	*	*	0.3	0.6	1.2	2.3	7.4	26.7
Female.....	1.2	*	*	*	*	*	0.1	0.3	0.7	2.1	6.8	25.2
Hispanic, both sexes.....	0.4	*	*	*	*	*	*	*	*	2.7	5.9	25.4
Male.....	0.4	*	*	*	*	*	*	*	*	*	*	35.4
Female.....	0.4	*	*	*	*	*	*	*	*	3.1	*	20.3
Non-Hispanic 4/, both sexes.....	1.2	*	*	*	*	0.1	0.2	0.5	1.0	2.2	7.1	25.6
Male.....	1.0	*	*	*	*	*	0.3	0.7	1.2	2.3	7.4	26.3
Female.....	1.3	*	*	*	*	*	0.1	0.4	0.7	2.1	6.8	25.3
Non-Hispanic white, both sexes.....	1.0	*	*	*	*	*	*	0.1	0.4	1.3	5.9	23.2
Male.....	0.8	*	*	*	*	*	*	0.2	0.5	1.6	6.6	23.6
Female.....	1.1	*	*	*	*	*	*	*	0.3	1.1	5.5	23.0
Non-Hispanic black, both sexes.....	2.5	*	*	*	*	0.4	1.3	3.1	5.5	9.3	19.9	53.8
Male.....	2.4	*	*	*	*	*	1.8	4.1	7.2	9.1	19.0	51.1
Female.....	2.6	*	*	*	*	*	0.8	2.2	4.2	9.4	20.4	54.9
Ischemic heart diseases (I20-I25)												
All origins 3/, both sexes.....	165.1	0.5	*	0.1	0.4	2.7	17.0	62.2	169.8	430.2	1,160.7	3,610.2
Male.....	172.2	*	*	*	0.5	4.0	25.7	97.2	251.5	598.8	1,524.4	4,005.2
Female.....	158.1	*	*	*	0.3	1.4	8.4	28.3	94.0	289.2	917.7	3,435.5
Hispanic, both sexes.....	52.1	*	*	*	0.3	1.4	7.5	35.4	122.0	355.0	973.0	2,970.0
Male.....	53.7	*	*	*	0.5	2.1	11.0	55.8	179.7	494.6	1,214.9	2,933.9
Female.....	50.4	*	*	*	*	0.6	3.7	15.0	69.8	243.7	806.9	2,988.3
Non-Hispanic 4/, both sexes.....	182.5	0.6	*	0.1	0.4	3.0	18.4	64.7	172.8	433.9	1,167.1	3,629.3
Male.....	191.5	*	*	*	0.5	4.5	27.9	101.1	255.9	603.6	1,536.0	4,045.5
Female.....	173.9	*	*	*	0.3	1.5	9.0	29.6	95.6	291.5	920.9	3,446.4
Non-Hispanic white, both sexes.....	200.8	*	*	*	0.4	2.8	17.9	61.1	164.5	421.6	1,164.6	3,691.6
Male.....	210.2	*	*	*	0.4	4.3	27.5	97.2	247.3	591.8	1,541.1	4,149.5
Female.....	191.8	*	*	*	0.3	1.4	8.2	25.5	85.9	275.9	910.4	3,490.0
Non-Hispanic black, both sexes.....	133.2	*	*	*	0.7	4.4	25.5	101.8	277.3	622.3	1,375.4	3,328.7
Male.....	137.6	*	*	*	0.9	6.4	36.2	148.3	387.0	836.1	1,758.6	3,309.2
Female.....	129.1	*	*	*	*	2.6	16.0	62.0	189.2	470.4	1,159.1	3,336.2
Acute myocardial infarction (I21-I22)												
All origins 3/, both sexes.....	58.7	*	*	*	0.2	1.0	7.0	25.5	70.9	168.6	412.7	1,118.8
Male.....	62.6	*	*	*	0.2	1.5	10.4	39.8	104.1	231.0	530.6	1,263.5
Female.....	54.8	*	*	*	0.1	0.6	3.6	11.7	40.0	116.5	333.9	1,054.8
Hispanic, both sexes.....	18.5	*	*	*	*	0.5	3.0	13.3	44.5	137.7	341.6	952.0
Male.....	18.9	*	*	*	*	0.7	4.1	20.6	64.1	187.5	409.2	963.6
Female.....	18.0	*	*	*	*	*	1.7	6.0	26.8	98.1	295.2	946.1
Non-Hispanic 4/, both sexes.....	64.9	*	*	*	0.2	1.2	7.6	26.7	72.8	170.4	415.5	1,123.8
Male.....	69.7	*	*	*	0.2	1.7	11.4	41.8	106.9	233.4	535.7	1,274.5
Female.....	60.3	*	*	*	0.2	0.6	3.8	12.3	41.0	117.5	335.3	1,057.5
Non-Hispanic white, both sexes.....	71.5	*	*	*	0.2	1.1	7.5	25.9	71.4	168.3	415.1	1,137.4
Male.....	77.0	*	*	*	0.2	1.6	11.5	41.2	106.5	232.9	537.8	1,300.9
Female.....	66.2	*	*	*	*	0.6	3.5	10.8	38.0	113.0	332.2	1,065.3
Non-Hispanic black, both sexes.....	47.3	*	*	*	*	1.7	10.4	38.6	100.6	223.1	486.5	1,103.6
Male.....	48.2	*	*	*	*	2.4	14.3	54.7	135.9	287.1	606.9	1,121.9
Female.....	46.5	*	*	*	*	1.0	6.9	24.7	72.2	177.6	418.5	1,096.5

Worktable Orig250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 2003

[Rates per 100,000 population, in specified group. Rates based on populations estimated as of July 1, 2003. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute ischemic heart diseases (I24)												
All origins 3/, both sexes.....	1.1	*	*	*	*	*	0.2	0.7	1.4	3.2	6.5	20.0
Male.....	1.1	*	*	*	*	*	0.3	1.0	2.1	4.3	8.0	19.5
Female.....	1.0	*	*	*	*	*	0.1	0.4	0.9	2.2	5.5	20.2
Hispanic, both sexes.....	0.1	*	*	*	*	*	*	*	*	*	*	*
Male.....	0.1	*	*	*	*	*	*	*	*	*	*	*
Female.....	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes.....	1.2	*	*	*	*	*	0.3	0.7	1.5	3.3	6.7	20.6
Male.....	1.3	*	*	*	*	*	0.4	1.1	2.2	4.5	8.4	20.2
Female.....	1.2	*	*	*	*	*	0.1	0.4	0.9	2.3	5.7	20.7
Non-Hispanic white, both sexes.....	1.3	*	*	*	*	*	0.2	0.7	1.4	3.1	6.6	20.9
Male.....	1.3	*	*	*	*	*	0.4	1.0	2.0	4.4	7.9	20.4
Female.....	1.3	*	*	*	*	*	*	0.4	0.8	2.1	5.7	21.1
Non-Hispanic black, both sexes.....	1.2	*	*	*	*	*	0.5	1.3	3.0	5.5	10.1	20.2
Male.....	1.5	*	*	*	*	*	*	2.0	4.9	7.7	16.4	24.5
Female.....	0.9	*	*	*	*	*	*	*	1.5	3.9	6.5	18.6
Other forms of chronic ischemic heart disease (I20,I25)												
All origins 3/, both sexes.....	105.3	*	*	*	0.2	1.7	9.8	35.9	97.5	258.4	741.5	2,471.4
Male.....	108.5	*	*	*	0.3	2.5	14.9	56.4	145.4	363.5	985.8	2,722.2
Female.....	102.3	*	*	*	0.1	0.8	4.7	16.2	53.1	170.5	578.3	2,360.5
Hispanic, both sexes.....	33.5	*	*	*	*	0.9	4.6	22.0	77.2	216.4	629.5	2,012.8
Male.....	34.6	*	*	*	*	1.4	6.9	35.1	115.3	306.1	804.2	1,965.7
Female.....	32.3	*	*	*	*	*	2.0	8.9	42.8	144.9	509.6	2,036.7
Non-Hispanic 4/, both sexes.....	116.3	*	*	*	0.2	1.8	10.6	37.2	98.5	260.2	744.8	2,484.9
Male.....	120.4	*	*	*	0.3	2.8	16.2	58.3	146.7	365.6	991.9	2,750.7
Female.....	112.4	*	*	*	0.1	0.9	5.1	16.9	53.7	171.7	580.0	2,368.1
Non-Hispanic white, both sexes.....	128.0	*	*	*	0.2	1.7	10.2	34.6	91.7	250.2	743.0	2,533.4
Male.....	131.9	*	*	*	0.2	2.6	15.7	55.1	138.9	354.5	995.5	2,828.2
Female.....	124.3	*	*	*	*	0.8	4.6	14.4	47.0	160.9	572.4	2,403.6
Non-Hispanic black, both sexes.....	84.7	*	*	*	0.4	2.7	14.6	61.9	173.6	393.8	878.9	2,204.8
Male.....	87.9	*	*	*	*	4.0	21.3	91.6	246.2	541.3	1,135.2	2,162.8
Female.....	81.7	*	*	*	*	1.5	8.7	36.5	115.4	288.8	734.1	2,221.1
Atherosclerotic cardiovascular disease, so described (I25.0)												
All origins 3/, both sexes.....	23.1	*	*	*	*	0.5	3.4	13.9	32.7	62.7	142.5	442.9
Male.....	25.2	*	*	*	*	0.9	5.2	22.7	49.9	88.9	181.1	462.1
Female.....	21.1	*	*	*	*	0.2	1.5	5.5	16.7	40.8	116.7	434.4
Hispanic, both sexes.....	8.3	*	*	*	*	0.3	1.6	9.8	29.4	57.3	124.6	357.1
Male.....	10.2	*	*	*	*	0.5	2.6	17.1	48.7	88.6	169.5	397.1
Female.....	6.2	*	*	*	*	*	*	2.5	12.0	32.4	93.7	336.7
Non-Hispanic 4/, both sexes.....	25.3	*	*	*	*	0.6	3.6	14.2	32.6	62.6	142.6	445.3
Male.....	27.5	*	*	*	*	1.0	5.6	23.1	49.3	88.2	180.4	463.3
Female.....	23.3	*	*	*	*	0.2	1.6	5.7	16.9	41.2	117.4	437.4
Non-Hispanic white, both sexes.....	26.2	*	*	*	*	0.6	3.4	12.8	28.6	55.9	135.1	441.5
Male.....	28.0	*	*	*	*	0.9	5.3	21.0	43.9	78.5	170.6	459.5
Female.....	24.6	*	*	*	*	0.2	1.5	4.6	14.2	36.5	111.0	433.5
Non-Hispanic black, both sexes.....	26.7	*	*	*	*	0.9	5.1	26.1	70.1	129.6	244.1	546.7
Male.....	31.2	*	*	*	*	1.6	7.8	41.1	106.0	191.8	334.2	588.1
Female.....	22.7	*	*	*	*	*	2.7	13.3	41.2	85.3	193.3	530.7

Worktable Orig250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 2003

[Rates per 100,000 population, in specified group. Rates based on populations estimated as of July 1, 2003. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
All origins 3/, both sexes.....	82.2	*	*	*	0.2	1.1	6.5	22.0	64.9	195.7	599.0	2,028.5
Male.....	83.3	*	*	*	0.2	1.6	9.7	33.7	95.5	274.6	804.7	2,260.1
Female.....	81.1	*	*	*	0.1	0.6	3.2	10.8	36.4	129.7	461.6	1,926.2
Hispanic, both sexes.....	25.2	*	*	*	*	0.6	3.0	12.2	47.8	159.1	504.9	1,655.7
Male.....	24.4	*	*	*	*	0.9	4.3	18.0	66.6	217.6	634.7	1,568.6
Female.....	26.1	*	*	*	*	*	1.6	6.4	30.7	112.5	415.8	1,700.0
Non-Hispanic 4/, both sexes.....	91.0	*	*	*	0.2	1.2	7.0	22.9	65.9	197.6	602.3	2,039.6
Male.....	92.9	*	*	*	0.2	1.8	10.6	35.2	97.4	277.5	811.5	2,287.4
Female.....	89.2	*	*	*	0.1	0.7	3.4	11.2	36.7	130.6	462.6	1,930.7
Non-Hispanic white, both sexes.....	101.8	*	*	*	0.1	1.1	6.7	21.8	63.1	194.3	607.9	2,091.9
Male.....	103.9	*	*	*	0.2	1.7	10.4	34.1	94.9	276.0	824.8	2,368.6
Female.....	99.8	*	*	*	*	0.6	3.1	9.7	32.8	124.3	461.4	1,970.1
Non-Hispanic black, both sexes.....	57.9	*	*	*	*	1.8	9.5	35.8	103.6	264.2	634.8	1,658.1
Male.....	56.7	*	*	*	*	2.4	13.4	50.5	140.2	349.5	801.1	1,574.7
Female.....	59.1	*	*	*	*	1.2	6.1	23.2	74.2	203.5	540.8	1,690.4
Other heart diseases (I26-I51)												
All origins 3/, both sexes.....	58.5	10.4	1.1	0.6	2.2	4.5	10.3	21.4	48.0	126.8	382.8	1,427.2
Male.....	52.3	11.4	1.0	0.7	2.6	5.3	12.7	27.2	60.7	154.2	440.0	1,431.3
Female.....	64.5	9.3	1.2	0.5	1.7	3.7	7.9	15.7	36.3	103.8	344.5	1,425.4
Hispanic, both sexes.....	14.1	8.3	1.0	0.3	1.4	2.8	5.3	11.3	29.7	77.3	221.2	749.9
Male.....	13.6	9.1	*	*	1.8	3.4	7.0	14.7	37.0	95.0	253.5	752.7
Female.....	14.5	7.6	*	*	0.9	2.2	3.3	8.0	23.2	63.2	199.1	748.4
Non-Hispanic 4/, both sexes.....	65.4	10.9	1.1	0.6	2.3	4.9	11.0	22.3	49.3	130.0	390.5	1,453.7
Male.....	58.6	12.0	1.0	0.7	2.8	5.8	13.5	28.4	62.2	158.0	448.9	1,460.5
Female.....	71.8	9.7	1.2	0.5	1.9	4.0	8.5	16.4	37.3	106.5	351.5	1,450.7
Non-Hispanic white, both sexes.....	71.0	8.1	0.8	0.5	1.9	3.8	9.1	18.8	44.1	123.1	391.4	1,490.1
Male.....	62.6	8.7	0.7	0.6	2.3	4.6	11.4	24.3	56.0	150.6	450.4	1,499.3
Female.....	79.1	7.5	1.0	0.4	1.6	3.0	6.8	13.5	32.7	99.6	351.5	1,486.1
Non-Hispanic black, both sexes.....	54.4	19.8	2.0	1.0	4.5	11.3	24.4	51.1	104.2	219.8	455.6	1,253.2
Male.....	53.4	23.6	2.2	1.2	5.8	13.0	29.6	65.1	134.7	269.3	530.4	1,239.1
Female.....	55.4	15.8	1.8	0.9	3.3	9.7	19.8	39.2	79.6	184.6	413.4	1,258.7
Acute and subacute endocarditis (I33)												
All origins 3/, both sexes.....	0.4	*	*	*	*	0.1	0.3	0.4	0.7	1.3	2.5	3.0
Male.....	0.5	*	*	*	*	*	0.3	0.6	0.9	1.4	3.2	4.1
Female.....	0.4	*	*	*	*	*	0.2	0.3	0.6	1.1	2.0	2.5
Hispanic, both sexes.....	0.2	*	*	*	*	*	*	*	*	*	*	*
Male.....	0.2	*	*	*	*	*	*	*	*	*	*	*
Female.....	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes.....	0.5	*	*	*	*	0.1	0.3	0.4	0.8	1.3	2.5	3.0
Male.....	0.5	*	*	*	*	*	0.3	0.6	0.9	1.4	3.3	4.1
Female.....	0.4	*	*	*	*	*	0.2	0.3	0.6	1.1	1.9	2.5
Non-Hispanic white, both sexes.....	0.5	*	*	*	*	0.1	0.2	0.3	0.6	1.2	2.5	3.2
Male.....	0.5	*	*	*	*	*	0.3	0.4	0.8	1.3	3.4	4.5
Female.....	0.4	*	*	*	*	*	0.2	0.2	0.5	1.0	1.9	2.7
Non-Hispanic black, both sexes.....	0.6	*	*	*	*	*	0.6	1.3	2.0	2.2	3.0	*
Male.....	0.7	*	*	*	*	*	*	2.0	2.3	*	*	*
Female.....	0.5	*	*	*	*	*	*	1.7	2.1	2.1	*	*

Worktable Orig250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 2003

[Rates per 100,000 population, in specified group. Rates based on populations estimated as of July 1, 2003. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
All origins 3/, both sexes.....	0.3	0.8	0.1	0.1	0.1	0.1	0.2	0.3	0.4	0.8	1.2	1.6
Male.....	0.3	*	*	*	0.2	0.1	0.2	0.4	0.4	1.0	1.3	1.5
Female.....	0.3	*	*	*	0.1	*	0.2	0.2	0.3	0.7	1.1	1.7
Hispanic, both sexes.....	0.1	*	*	*	*	*	*	*	*	*	*	*
Male.....	0.2	*	*	*	*	*	*	*	*	*	*	*
Female.....	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes.....	0.3	0.9	*	0.1	0.1	0.1	0.2	0.3	0.4	0.8	1.2	1.7
Male.....	0.3	*	*	*	0.2	0.2	0.3	0.4	0.4	1.0	1.3	1.6
Female.....	0.3	*	*	*	*	*	0.2	0.2	0.3	0.7	1.1	1.7
Non-Hispanic white, both sexes.....	0.3	*	*	*	0.1	*	0.2	0.2	0.3	0.9	1.2	1.7
Male.....	0.3	*	*	*	0.2	*	0.2	0.3	0.4	1.0	1.4	1.8
Female.....	0.3	*	*	*	*	*	0.2	0.3	0.7	1.1	1.7	
Non-Hispanic black, both sexes.....	0.5	*	*	*	*	0.4	0.5	0.5	*	*	*	*
Male.....	0.5	*	*	*	*	*	*	*	*	*	*	*
Female.....	0.5	*	*	*	*	*	*	*	*	*	*	*
Heart failure (I50)												
All origins 3/, both sexes.....	19.8	0.9	*	*	0.1	0.2	0.7	2.4	8.8	33.5	133.2	641.1
Male.....	15.7	1.4	*	*	0.1	0.3	0.9	2.9	10.8	39.7	148.6	629.8
Female.....	23.7	*	*	*	*	0.2	0.5	1.9	7.0	28.3	122.9	646.1
Hispanic, both sexes.....	4.0	*	*	*	*	*	*	1.3	5.7	22.5	77.1	330.1
Male.....	3.5	*	*	*	*	*	*	1.3	7.8	26.0	92.8	312.5
Female.....	4.6	*	*	*	*	*	*	1.2	3.7	19.7	66.4	339.1
Non-Hispanic 4/, both sexes.....	22.2	1.0	*	*	0.1	0.3	0.8	2.5	9.1	34.1	135.9	653.2
Male.....	17.7	1.4	*	*	0.1	0.3	1.0	3.1	11.0	40.5	151.2	643.2
Female.....	26.5	*	*	*	*	0.2	0.6	2.0	7.3	28.8	125.6	657.6
Non-Hispanic white, both sexes.....	25.0	*	*	*	*	0.2	0.5	1.9	8.0	32.8	137.1	673.5
Male.....	19.7	*	*	*	*	0.3	0.7	2.4	9.7	39.3	153.2	665.3
Female.....	30.1	*	*	*	*	*	0.4	1.5	6.3	27.2	126.3	677.2
Non-Hispanic black, both sexes.....	14.5	*	*	*	*	0.7	2.4	7.1	21.0	55.3	152.7	538.7
Male.....	12.3	*	*	*	*	0.8	2.8	8.5	26.8	64.3	165.9	521.0
Female.....	16.5	*	*	*	*	*	2.0	5.8	16.3	48.9	145.2	545.6
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)												
All origins 3/, both sexes.....	38.0	8.6	0.9	0.5	1.9	4.1	9.1	18.2	38.1	91.2	246.0	781.5
Male.....	35.8	9.2	0.9	0.6	2.3	4.8	11.3	23.3	48.5	112.1	287.0	795.9
Female.....	40.1	7.9	1.0	0.4	1.5	3.4	7.0	13.3	28.4	73.7	218.6	775.1
Hispanic, both sexes.....	9.7	6.8	0.9	0.3	1.2	2.6	4.8	9.5	23.2	53.0	141.3	418.2
Male.....	9.8	7.0	*	*	1.6	3.0	6.4	12.7	28.3	66.2	159.1	437.2
Female.....	9.6	6.6	*	*	0.8	2.0	3.1	6.3	18.6	42.4	129.0	408.6
Non-Hispanic 4/, both sexes.....	42.4	9.0	0.9	0.5	2.1	4.4	9.7	19.1	39.1	93.7	250.9	795.8
Male.....	40.0	9.9	0.8	0.6	2.5	5.2	12.0	24.3	49.8	115.1	293.1	811.6
Female.....	44.6	8.1	1.0	0.4	1.6	3.6	7.5	14.0	29.1	75.8	222.8	788.8
Non-Hispanic white, both sexes.....	45.2	7.1	0.7	0.5	1.7	3.5	8.2	16.3	35.1	88.3	250.6	811.6
Male.....	42.0	7.5	0.6	0.6	2.0	4.2	10.2	21.0	45.1	108.9	292.5	827.7
Female.....	48.3	6.7	0.8	0.3	1.4	2.7	6.1	11.6	25.6	70.6	222.3	804.5
Non-Hispanic black, both sexes.....	38.8	15.2	1.6	0.8	4.0	10.1	20.9	42.3	80.8	161.3	298.6	711.8
Male.....	39.9	18.3	1.9	0.9	5.1	11.4	25.6	53.9	105.1	201.8	359.8	718.1
Female.....	37.8	11.9	*	0.7	2.8	8.8	16.7	32.3	61.2	132.5	264.0	709.4

Worktable Orig250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 2003

[Rates per 100,000 population, in specified group. Rates based on populations estimated as of July 1, 2003. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Essential (primary) hypertension and hypertensive renal disease (I10,I12)												
All origins 3/, both sexes.....	7.5	*	*	*	0.1	0.2	0.8	2.5	6.3	16.9	51.7	188.9
Male.....	5.7	*	*	*	*	0.3	1.0	3.3	7.4	18.4	48.9	153.5
Female.....	9.3	*	*	*	*	0.2	0.7	1.8	5.3	15.7	53.5	204.6
Hispanic, both sexes.....	2.6	*	*	*	*	*	0.4	2.0	5.8	16.7	44.5	171.0
Male.....	2.1	*	*	*	*	*	*	2.8	7.4	15.5	44.5	133.9
Female.....	3.2	*	*	*	*	*	*	1.2	4.4	17.6	44.5	189.8
Non-Hispanic 4/, both sexes.....	8.3	*	*	*	*	0.3	0.9	2.6	6.3	16.8	51.9	189.4
Male.....	6.3	*	*	*	*	0.3	1.0	3.3	7.3	18.5	49.1	154.4
Female.....	10.2	*	*	*	*	0.2	0.8	1.8	5.4	15.5	53.8	204.7
Non-Hispanic white, both sexes.....	8.0	*	*	*	*	0.1	0.5	1.6	4.2	12.6	46.2	181.0
Male.....	5.8	*	*	*	*	*	0.7	2.1	4.9	14.4	43.6	145.4
Female.....	10.0	*	*	*	*	*	0.4	1.1	3.5	11.2	47.9	196.6
Non-Hispanic black, both sexes.....	12.2	*	*	*	*	1.0	3.2	9.5	24.7	55.1	121.0	304.3
Male.....	10.8	*	*	*	*	1.3	3.5	13.0	30.0	61.3	120.9	280.2
Female.....	13.4	*	*	*	*	*	2.8	6.5	20.5	50.8	121.1	313.6
Cerebrovascular diseases (I60-I69)												
All origins 3/, both sexes.....	54.2	2.5	0.3	0.2	0.5	1.5	5.5	15.0	35.6	112.9	410.7	1,370.1
Male.....	42.9	2.8	0.3	0.2	0.5	1.6	5.8	16.7	40.8	127.8	431.4	1,236.0
Female.....	65.1	2.2	0.3	0.1	0.5	1.4	5.3	13.4	30.9	100.5	396.8	1,429.4
Hispanic, both sexes.....	16.7	*	*	*	0.4	1.2	4.5	14.9	35.3	98.4	316.8	880.5
Male.....	14.9	*	*	*	*	1.3	5.2	18.1	43.5	113.9	337.1	837.4
Female.....	18.6	*	*	*	*	1.1	3.7	11.7	27.8	86.0	302.8	902.3
Non-Hispanic 4/, both sexes.....	60.1	2.6	0.3	0.2	0.6	1.5	5.7	15.0	35.6	113.7	415.0	1,388.4
Male.....	47.6	2.8	*	0.2	0.6	1.6	5.9	16.5	40.4	128.4	435.7	1,252.9
Female.....	72.0	2.4	0.3	0.1	0.5	1.4	5.5	13.5	31.0	101.4	401.1	1,448.0
Non-Hispanic white, both sexes.....	64.2	2.0	0.2	0.1	0.5	1.2	4.1	10.9	28.7	103.6	406.6	1,401.1
Male.....	49.7	1.9	*	0.2	0.5	1.3	4.2	12.1	32.1	116.9	426.0	1,264.2
Female.....	78.2	2.1	*	*	0.4	1.1	4.0	9.7	25.5	92.1	393.6	1,461.3
Non-Hispanic black, both sexes.....	51.1	4.9	*	*	1.0	3.2	14.2	42.0	91.3	204.9	539.1	1,366.0
Male.....	44.9	*	*	*	0.8	3.5	14.9	48.1	114.1	241.8	597.6	1,200.7
Female.....	56.7	*	*	*	1.1	3.0	13.7	36.8	73.0	178.6	506.0	1,430.0
Atherosclerosis (I70)												
All origins 3/, both sexes.....	4.5	*	*	*	*	*	0.1	0.5	1.7	6.6	29.8	154.6
Male.....	3.4	*	*	*	*	*	0.2	0.7	2.2	7.8	33.6	138.6
Female.....	5.5	*	*	*	*	*	*	0.3	1.2	5.6	27.3	161.7
Hispanic, both sexes.....	0.9	*	*	*	*	*	*	*	1.1	4.6	18.9	86.0
Male.....	0.9	*	*	*	*	*	*	*	6.1	24.2	95.4	95.4
Female.....	1.0	*	*	*	*	*	*	*	3.4	15.3	81.3	81.3
Non-Hispanic 4/, both sexes.....	5.0	*	*	*	*	*	0.1	0.5	1.7	6.7	30.2	157.1
Male.....	3.8	*	*	*	*	*	0.2	0.7	2.2	7.9	34.0	140.3
Female.....	6.2	*	*	*	*	*	*	0.3	1.2	5.8	27.7	164.5
Non-Hispanic white, both sexes.....	5.8	*	*	*	*	*	0.1	0.5	1.6	6.6	30.8	162.2
Male.....	4.3	*	*	*	*	*	0.2	0.7	2.2	7.7	34.1	145.2
Female.....	7.2	*	*	*	*	*	*	0.3	1.1	5.8	28.5	169.6
Non-Hispanic black, both sexes.....	2.7	*	*	*	*	*	*	0.8	2.7	9.1	30.8	122.9
Male.....	2.3	*	*	*	*	*	*	1.2	3.7	12.0	40.4	105.5
Female.....	3.0	*	*	*	*	*	*	*	1.9	7.1	25.4	129.6

Worktable Orig250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 2003

[Rates per 100,000 population, in specified group. Rates based on populations estimated as of July 1, 2003. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of circulatory system (I71-I78)												
All origins 3/, both sexes.....	8.5	*	*	*	0.2	0.4	1.0	2.9	8.1	26.9	67.6	146.0
Male.....	9.2	*	*	*	0.2	0.5	1.4	4.1	11.3	37.0	91.8	179.7
Female.....	7.8	*	*	*	0.1	0.2	0.7	1.8	5.2	18.4	51.4	131.1
Hispanic, both sexes.....	2.2	*	*	*	*	0.3	0.7	1.6	4.6	14.1	42.3	106.2
Male.....	2.5	*	*	*	*	*	0.9	2.2	7.4	19.0	58.7	138.5
Female.....	1.9	*	*	*	*	*	*	*	2.1	10.1	31.1	89.8
Non-Hispanic 4/, both sexes.....	9.5	*	*	*	0.2	0.4	1.1	3.1	8.4	27.7	68.7	147.5
Male.....	10.4	*	*	*	0.2	0.5	1.5	4.3	11.6	38.2	93.4	181.5
Female.....	8.6	*	*	*	*	0.3	0.7	1.9	5.5	19.0	52.3	132.6
Non-Hispanic white, both sexes.....	10.4	*	*	*	0.2	0.3	0.9	2.6	8.2	28.2	69.5	148.3
Male.....	11.5	*	*	*	0.2	0.4	1.2	3.8	11.3	39.2	94.7	185.8
Female.....	9.4	*	*	*	*	0.2	0.5	1.5	5.2	18.8	52.4	131.8
Non-Hispanic black, both sexes.....	6.8	*	*	*	*	0.7	2.4	6.4	12.1	29.6	71.1	152.9
Male.....	6.9	*	*	*	*	1.2	3.1	8.4	16.7	36.4	91.5	141.7
Female.....	6.8	*	*	*	*	*	1.7	4.6	8.4	24.7	59.6	157.2
Aortic aneurysm and dissection (I71)												
All origins 3/, both sexes.....	5.1	*	*	*	0.1	0.3	0.7	2.0	5.3	17.8	42.5	70.0
Male.....	6.3	*	*	*	0.1	0.4	1.1	3.1	8.2	26.0	61.9	105.5
Female.....	3.9	*	*	*	*	0.1	0.3	1.0	2.6	11.0	29.5	54.2
Hispanic, both sexes.....	1.3	*	*	*	*	*	0.4	0.9	3.1	8.3	26.7	49.2
Male.....	1.6	*	*	*	*	*	*	1.4	5.1	12.5	38.3	80.0
Female.....	0.9	*	*	*	*	*	*	*	*	5.0	18.7	33.6
Non-Hispanic 4/, both sexes.....	5.7	*	*	*	0.1	0.3	0.8	2.1	5.5	18.5	43.2	70.7
Male.....	7.0	*	*	*	0.2	0.4	1.2	3.2	8.5	26.9	63.0	106.6
Female.....	4.4	*	*	*	*	0.2	0.3	1.0	2.7	11.4	30.0	55.0
Non-Hispanic white, both sexes.....	6.4	*	*	*	0.1	0.2	0.6	1.8	5.4	19.3	44.8	73.2
Male.....	7.9	*	*	*	0.2	0.3	1.0	2.8	8.4	28.3	65.5	111.2
Female.....	4.9	*	*	*	*	*	0.2	0.8	2.5	11.6	30.9	56.5
Non-Hispanic black, both sexes.....	3.3	*	*	*	*	0.6	1.7	4.4	7.3	14.5	30.4	47.3
Male.....	3.9	*	*	*	*	1.1	2.5	6.5	11.0	17.9	39.8	52.2
Female.....	2.8	*	*	*	*	*	1.0	2.5	4.3	12.1	25.1	45.4
Other diseases of arteries, arterioles and capillaries (I72-I78)												
All origins 3/, both sexes.....	3.4	*	*	*	0.1	0.1	0.3	0.9	2.8	9.1	25.0	76.0
Male.....	3.0	*	*	*	*	0.1	0.3	1.0	3.1	11.0	29.9	74.2
Female.....	3.8	*	*	*	*	*	0.3	0.9	2.6	7.5	21.8	76.9
Hispanic, both sexes.....	0.9	*	*	*	*	*	*	0.7	1.5	5.7	15.6	57.0
Male.....	0.9	*	*	*	*	*	*	*	2.2	6.5	20.3	58.5
Female.....	1.0	*	*	*	*	*	*	*	*	5.1	12.4	56.3
Non-Hispanic 4/, both sexes.....	3.8	*	*	*	0.1	0.1	0.3	1.0	2.9	9.3	25.5	76.7
Male.....	3.3	*	*	*	*	*	0.3	1.0	3.1	11.3	30.3	74.9
Female.....	4.2	*	*	*	*	*	0.4	0.9	2.8	7.6	22.3	77.6
Non-Hispanic white, both sexes.....	4.0	*	*	*	*	0.1	0.3	0.8	2.8	8.9	24.6	75.1
Male.....	3.5	*	*	*	*	*	0.2	0.9	2.9	10.8	29.2	74.6
Female.....	4.5	*	*	*	*	*	0.3	0.7	2.7	7.3	21.6	75.4
Non-Hispanic black, both sexes.....	3.5	*	*	*	*	*	0.7	2.0	4.8	15.0	40.7	105.6
Male.....	3.0	*	*	*	*	*	*	1.9	5.7	18.5	51.7	89.5
Female.....	4.0	*	*	*	*	*	0.8	2.1	4.1	12.5	34.5	111.8

Worktable Orig250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 2003

[Rates per 100,000 population, in specified group. Rates based on populations estimated as of July 1, 2003. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Emphysema (J43)												
All origins 3/, both sexes.....	5.1	*	*	*	*	*	0.2	1.1	6.3	22.2	46.0	54.6
Male.....	5.3	*	*	*	*	*	0.2	1.4	6.9	25.6	58.1	79.2
Female.....	4.9	*	*	*	*	*	0.2	0.8	5.7	19.2	37.9	43.7
Hispanic, both sexes.....	0.8	*	*	*	*	*	*	*	1.4	6.0	20.0	34.7
Male.....	0.9	*	*	*	*	*	*	*	2.1	7.6	29.1	53.9
Female.....	0.7	*	*	*	*	*	*	*	*	4.7	13.7	25.0
Non-Hispanic 4/, both sexes.....	5.8	*	*	*	*	*	0.2	1.2	6.6	23.3	47.3	55.4
Male.....	6.0	*	*	*	*	*	0.2	1.5	7.3	26.8	59.5	80.4
Female.....	5.6	*	*	*	*	*	0.2	0.9	6.0	20.3	39.1	44.5
Non-Hispanic white, both sexes.....	6.8	*	*	*	*	*	0.2	1.2	7.1	25.2	49.9	57.9
Male.....	6.9	*	*	*	*	*	0.2	1.5	7.6	28.5	61.3	81.6
Female.....	6.7	*	*	*	*	*	0.2	1.0	6.7	22.5	42.3	47.4
Non-Hispanic black, both sexes.....	2.2	*	*	*	*	*	*	1.2	5.3	13.2	27.7	31.2
Male.....	2.8	*	*	*	*	*	*	1.7	7.5	18.6	46.7	71.4
Female.....	1.6	*	*	*	*	*	*	*	3.5	9.4	16.9	15.7
Asthma (J45-J46)												
All origins 3/, both sexes.....	1.4	*	0.2	0.3	0.4	0.6	0.9	1.5	2.0	2.9	5.8	14.2
Male.....	1.0	*	0.4	0.3	0.4	0.6	0.7	1.2	1.6	2.1	4.5	10.7
Female.....	1.8	*	*	0.2	0.3	0.5	1.1	1.9	2.4	3.6	6.7	15.8
Hispanic, both sexes.....	0.7	*	*	*	*	0.3	0.5	1.6	2.2	3.4	6.1	14.5
Male.....	0.6	*	*	*	*	*	*	1.4	2.2	*	*	*
Female.....	0.9	*	*	*	*	*	*	1.8	2.1	4.1	6.3	17.2
Non-Hispanic 4/, both sexes.....	1.5	*	0.2	0.3	0.4	0.6	1.0	1.5	2.0	2.9	5.8	14.1
Male.....	1.1	*	0.4	0.4	0.5	0.7	0.8	1.1	1.5	2.0	4.5	10.7
Female.....	1.9	*	*	0.2	0.4	0.5	1.2	1.9	2.4	3.6	6.7	15.6
Non-Hispanic white, both sexes.....	1.3	*	*	0.1	0.2	0.4	0.7	1.0	1.5	2.4	5.3	14.1
Male.....	0.9	*	*	*	0.3	0.4	0.5	0.8	1.1	1.7	3.8	10.6
Female.....	1.7	*	*	*	0.2	0.5	0.9	1.3	2.0	2.9	6.3	15.7
Non-Hispanic black, both sexes.....	2.8	*	0.8	0.9	1.2	1.7	2.8	4.9	5.2	7.2	9.3	11.9
Male.....	2.5	*	*	1.3	1.4	2.5	2.4	4.0	4.4	5.5	8.9	*
Female.....	3.0	*	*	*	1.1	1.0	3.2	5.7	5.9	8.4	9.5	13.6
Other chronic lower respiratory diseases (J44,J47)												
All origins 3/, both sexes.....	36.6	*	*	*	0.1	0.1	1.0	5.9	34.8	137.5	329.2	558.9
Male.....	35.9	*	*	*	*	0.1	1.0	6.4	37.8	155.4	400.1	769.0
Female.....	37.4	*	*	*	*	*	1.0	5.5	32.1	122.6	281.9	466.0
Hispanic, both sexes.....	6.3	*	*	*	*	*	0.4	1.7	9.1	43.7	143.5	403.7
Male.....	6.6	*	*	*	*	*	*	1.5	12.2	57.8	194.4	574.2
Female.....	6.0	*	*	*	*	*	*	1.9	6.3	32.5	108.5	317.2
Non-Hispanic 4/, both sexes.....	41.4	*	*	*	*	0.1	1.1	6.3	36.7	143.8	338.3	564.3
Male.....	40.7	*	*	*	*	0.1	1.0	6.9	39.6	161.7	410.1	776.7
Female.....	42.0	*	*	*	*	*	1.1	5.8	34.1	128.9	290.3	471.0
Non-Hispanic white, both sexes.....	48.6	*	*	*	*	0.1	1.1	6.6	39.0	154.4	357.6	592.2
Male.....	46.9	*	*	*	*	*	1.0	7.1	41.1	169.5	423.7	804.2
Female.....	50.2	*	*	*	*	*	1.1	6.1	37.0	141.5	313.1	498.8
Non-Hispanic black, both sexes.....	15.7	*	*	*	*	*	1.3	6.5	30.5	94.6	200.3	306.1
Male.....	18.6	*	*	*	*	*	1.4	7.7	39.3	133.6	329.0	526.3
Female.....	13.0	*	*	*	*	*	1.2	5.5	23.5	66.8	127.6	220.8

Worktable Orig250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 2003

[Rates per 100,000 population, in specified group. Rates based on populations estimated as of July 1, 2003. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumoconioses and chemical effects (J60-J66,J68)												
All origins 3/, both sexes.....	0.4	*	*	*	*	*	*	*	0.3	1.3	3.8	6.4
Male.....	0.7	*	*	*	*	*	*	*	0.5	2.7	9.1	19.8
Female.....	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes.....	0.1	*	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes.....	0.4	*	*	*	*	*	*	*	0.3	1.3	3.9	6.6
Male.....	0.9	*	*	*	*	*	*	*	0.5	2.8	9.3	20.7
Female.....	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes.....	0.5	*	*	*	*	*	*	*	0.3	1.5	4.1	7.1
Male.....	1.0	*	*	*	*	*	*	*	0.6	3.0	9.8	22.3
Female.....	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes.....	0.1	*	*	*	*	*	*	*	*	*	2.4	*
Male.....	0.3	*	*	*	*	*	*	*	*	*	6.3	*
Female.....	*	*	*	*	*	*	*	*	*	*	*	*
Pneumonitis due to solids and liquids (J69)												
All origins 3/, both sexes.....	6.0	*	*	*	0.1	0.2	0.4	1.0	2.8	9.6	45.0	176.2
Male.....	6.1	*	*	*	0.1	0.2	0.4	1.2	3.5	12.8	61.7	246.2
Female.....	5.8	*	*	*	*	0.1	0.3	0.7	2.1	6.9	33.8	145.3
Hispanic, both sexes.....	1.3	*	*	*	*	*	*	*	1.7	6.5	22.8	113.0
Male.....	1.3	*	*	*	*	*	*	*	2.0	8.7	32.2	123.1
Female.....	1.4	*	*	*	*	*	*	*	*	4.9	16.3	107.8
Non-Hispanic 4/, both sexes.....	6.7	*	*	*	0.1	0.2	0.4	1.0	2.9	9.7	46.0	178.5
Male.....	6.9	*	*	*	0.1	0.2	0.4	1.2	3.6	13.0	63.1	251.5
Female.....	6.5	*	*	*	*	0.2	0.4	0.8	2.1	7.0	34.6	146.5
Non-Hispanic white, both sexes.....	7.5	*	*	*	0.1	0.2	0.4	1.0	2.6	9.5	45.7	181.0
Male.....	7.6	*	*	*	*	0.2	0.4	1.2	3.3	12.6	62.4	256.5
Female.....	7.3	*	*	*	*	0.2	0.3	0.8	2.0	6.8	34.4	147.8
Non-Hispanic black, both sexes.....	4.4	*	*	*	*	*	0.7	1.3	5.3	13.9	55.5	165.7
Male.....	4.5	*	*	*	*	*	*	1.7	7.2	19.8	79.6	215.2
Female.....	4.3	*	*	*	*	*	0.8	1.0	3.7	9.7	41.9	146.5
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)												
All origins 3/, both sexes.....	8.8	7.0	0.5	0.1	0.3	0.5	1.1	3.3	8.8	28.1	68.5	138.1
Male.....	8.7	8.0	0.6	0.2	0.4	0.6	1.1	3.5	10.2	34.1	85.3	165.0
Female.....	8.9	5.9	0.5	0.1	0.3	0.5	1.1	3.0	7.4	23.1	57.2	126.2
Hispanic, both sexes.....	3.4	6.6	*	*	*	0.4	0.9	2.5	8.9	25.5	58.2	118.7
Male.....	3.3	6.6	*	*	*	*	1.1	3.1	11.3	30.8	63.3	127.8
Female.....	3.6	6.6	*	*	*	*	1.8	6.7	21.2	54.8	114.1	
Non-Hispanic 4/, both sexes.....	9.6	7.0	0.6	0.1	0.4	0.6	1.2	3.3	8.7	28.2	68.9	138.7
Male.....	9.5	8.3	0.6	0.2	0.4	0.6	1.1	3.5	10.1	34.2	86.3	166.5
Female.....	9.7	5.7	0.5	*	0.3	0.5	1.2	3.2	7.5	23.2	57.3	126.4
Non-Hispanic white, both sexes.....	10.7	4.9	0.4	0.1	0.3	0.5	0.9	3.0	8.5	29.1	71.3	143.7
Male.....	10.6	5.5	0.4	0.2	0.3	0.6	1.0	3.2	9.7	35.2	88.8	172.0
Female.....	10.8	4.4	*	*	0.3	0.4	0.9	2.7	7.3	23.9	59.4	131.2
Non-Hispanic black, both sexes.....	6.0	15.3	1.1	*	0.7	1.0	2.5	6.6	12.3	24.6	51.3	91.0
Male.....	6.2	18.6	*	*	0.9	1.1	2.3	6.4	15.3	31.4	69.0	116.1
Female.....	5.9	11.9	*	*	*	1.0	2.7	6.6	9.8	19.8	41.4	81.3

Worktable Orig250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 2003

[Rates per 100,000 population, in specified group. Rates based on populations estimated as of July 1, 2003. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Peptic ulcer (K25-K28)												
All origins 3/, both sexes.....	1.3	*	*	*	*	0.1	0.2	0.7	1.4	3.2	9.9	26.4
Male.....	1.3	*	*	*	*	*	0.3	1.0	1.8	4.1	11.8	28.9
Female.....	1.4	*	*	*	*	*	0.2	0.4	1.0	2.4	8.7	25.3
Hispanic, both sexes.....	0.5	*	*	*	*	*	*	0.8	1.2	3.4	8.7	17.1
Male.....	0.6	*	*	*	*	*	*	1.1	*	5.5	11.5	*
Female.....	0.4	*	*	*	*	*	*	*	*	*	6.8	20.3
Non-Hispanic 4/, both sexes.....	1.5	*	*	*	*	0.1	0.2	0.7	1.4	3.2	9.9	26.7
Male.....	1.4	*	*	*	*	*	0.3	0.9	1.8	4.0	11.7	29.6
Female.....	1.5	*	*	*	*	*	0.2	0.4	1.0	2.5	8.7	25.4
Non-Hispanic white, both sexes.....	1.6	*	*	*	*	*	0.2	0.6	1.3	3.2	9.8	27.3
Male.....	1.5	*	*	*	*	*	0.2	0.8	1.6	4.0	11.3	29.8
Female.....	1.7	*	*	*	*	*	0.2	0.4	1.0	2.6	8.8	26.2
Non-Hispanic black, both sexes.....	1.0	*	*	*	*	*	0.4	1.5	2.2	3.2	11.1	17.3
Male.....	1.3	*	*	*	*	*	*	2.3	3.4	5.2	16.4	*
Female.....	0.8	*	*	*	*	*	*	*	*	*	8.1	16.1
Diseases of appendix (K35-K38)												
All origins 3/, both sexes.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.3	1.0	1.7
Male.....	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.5	2.4
Female.....	0.1	*	*	*	*	*	*	0.1	*	*	0.7	1.3
Hispanic, both sexes.....	0.1	*	*	*	*	*	*	*	*	*	*	*
Male.....	0.1	*	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.0	1.6
Male.....	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.5	2.2
Female.....	0.1	*	*	*	*	*	*	*	*	*	0.8	1.4
Non-Hispanic white, both sexes.....	0.2	*	*	*	*	*	*	0.1	0.1	0.3	1.0	1.7
Male.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.5	2.4
Female.....	0.1	*	*	*	*	*	*	*	*	*	0.8	1.3
Non-Hispanic black, both sexes.....	0.2	*	*	*	*	*	*	*	*	*	*	*
Male.....	0.2	*	*	*	*	*	*	*	*	*	*	*
Female.....	0.2	*	*	*	*	*	*	*	*	*	*	*
Hernia (K40-K46)												
All origins 3/, both sexes.....	0.6	0.9	*	*	*	*	0.1	0.3	0.6	1.3	3.6	11.9
Male.....	0.5	1.0	*	*	*	*	*	0.3	0.5	1.3	3.9	12.3
Female.....	0.6	*	*	*	*	*	*	0.2	0.6	1.3	3.4	11.8
Hispanic, both sexes.....	0.2	*	*	*	*	*	*	*	*	*	3.6	*
Male.....	0.2	*	*	*	*	*	*	*	*	*	*	*
Female.....	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes.....	0.6	1.1	*	*	*	*	0.1	0.2	0.6	1.3	3.6	12.1
Male.....	0.5	*	*	*	*	*	*	0.3	0.5	1.3	3.9	12.5
Female.....	0.7	*	*	*	*	*	*	0.2	0.6	1.3	3.3	12.0
Non-Hispanic white, both sexes.....	0.7	1.1	*	*	*	*	*	0.2	0.5	1.3	3.6	12.4
Male.....	0.6	*	*	*	*	*	*	0.3	0.5	1.3	3.9	12.6
Female.....	0.8	*	*	*	*	*	*	0.2	0.5	1.3	3.3	12.4
Non-Hispanic black, both sexes.....	0.4	*	*	*	*	*	*	*	1.0	1.4	4.0	9.2
Male.....	0.4	*	*	*	*	*	*	*	*	*	*	*
Female.....	0.5	*	*	*	*	*	*	*	*	*	3.9	8.3

Worktable Orig250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 2003

[Rates per 100,000 population, in specified group. Rates based on populations estimated as of July 1, 2003. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic liver disease and cirrhosis (K70,K73-K74)												
All origins 3/, both sexes.....	9.5	*	*	*	*	0.9	6.8	18.3	23.0	29.5	30.0	20.1
Male.....	12.5	*	*	*	*	1.2	9.2	27.5	33.8	39.1	36.9	27.5
Female.....	6.5	*	*	*	*	0.5	4.4	9.4	13.1	21.4	25.4	16.8
Hispanic, both sexes.....	8.5	*	*	*	*	1.4	7.9	24.3	38.4	51.1	62.0	31.1
Male.....	11.6	*	*	*	*	2.0	12.3	40.2	62.2	61.3	71.3	35.4
Female.....	5.2	*	*	*	*	0.6	3.2	8.3	17.0	42.9	55.6	28.9
Non-Hispanic 4/, both sexes.....	9.6	*	*	*	*	0.8	6.6	17.6	21.7	27.9	28.2	19.6
Male.....	12.6	*	*	*	*	1.1	8.6	26.0	31.3	37.5	35.0	27.1
Female.....	6.7	*	*	*	*	0.5	4.6	9.5	12.7	19.8	23.7	16.3
Non-Hispanic white, both sexes.....	10.3	*	*	*	*	0.8	6.8	17.5	21.5	28.9	29.7	20.3
Male.....	13.6	*	*	*	*	1.0	9.0	25.9	30.8	38.7	36.5	27.9
Female.....	7.2	*	*	*	*	0.5	4.6	9.3	12.6	20.5	25.2	17.0
Non-Hispanic black, both sexes.....	7.1	*	*	*	*	0.8	5.8	20.2	26.1	24.1	14.1	12.5
Male.....	9.8	*	*	*	*	1.0	7.3	31.0	41.2	33.7	20.2	22.4
Female.....	4.6	*	*	*	*	*	4.6	10.9	14.0	17.3	10.6	8.7
Alcoholic liver disease (K70)												
All origins 3/, both sexes.....	4.3	*	*	*	*	0.6	4.4	10.5	11.6	9.8	5.7	1.8
Male.....	6.4	*	*	*	*	0.9	6.0	16.1	18.3	16.1	10.1	3.6
Female.....	2.2	*	*	*	*	0.3	2.9	5.1	5.4	4.5	2.7	1.0
Hispanic, both sexes.....	4.3	*	*	*	*	0.9	5.6	14.4	21.4	17.7	14.1	*
Male.....	6.9	*	*	*	*	1.3	8.8	25.5	38.4	29.0	26.5	*
Female.....	1.5	*	*	*	*	*	2.1	3.4	6.1	8.7	5.5	*
Non-Hispanic 4/, both sexes.....	4.2	*	*	*	*	0.5	4.2	10.1	10.8	9.1	5.2	1.7
Male.....	6.2	*	*	*	*	0.8	5.5	15.0	16.6	15.1	9.2	3.4
Female.....	2.3	*	*	*	*	0.3	3.0	5.3	5.4	4.1	2.6	1.0
Non-Hispanic white, both sexes.....	4.5	*	*	*	*	0.5	4.4	10.1	10.7	9.3	5.4	1.7
Male.....	6.6	*	*	*	*	0.8	5.6	15.1	16.4	15.3	9.4	3.5
Female.....	2.4	*	*	*	*	0.3	3.1	5.3	5.2	4.1	2.8	1.0
Non-Hispanic black, both sexes.....	3.3	*	*	*	*	*	3.4	10.0	12.7	9.4	3.4	*
Male.....	4.9	*	*	*	*	*	4.2	16.0	20.3	16.2	6.9	*
Female.....	1.9	*	*	*	*	*	2.6	4.8	6.6	4.6	*	*
Other chronic liver disease and cirrhosis (K73-K74)												
All origins 3/, both sexes.....	5.2	*	*	*	*	0.3	2.4	7.8	11.4	19.7	24.3	18.2
Male.....	6.2	*	*	*	*	0.3	3.2	11.4	15.5	23.0	26.8	23.9
Female.....	4.3	*	*	*	*	0.2	1.5	4.3	7.6	17.0	22.6	15.8
Hispanic, both sexes.....	4.2	*	*	*	*	0.4	2.3	9.8	17.0	33.4	47.9	26.9
Male.....	4.7	*	*	*	*	0.6	3.5	14.7	23.8	32.3	44.9	*
Female.....	3.7	*	*	*	*	*	1.1	5.0	10.8	34.3	50.0	26.6
Non-Hispanic 4/, both sexes.....	5.4	*	*	*	*	0.2	2.4	7.5	10.9	18.7	23.0	17.9
Male.....	6.4	*	*	*	*	0.3	3.2	11.0	14.7	22.3	25.9	23.7
Female.....	4.4	*	*	*	*	0.2	1.6	4.2	7.3	15.7	21.1	15.3
Non-Hispanic white, both sexes.....	5.9	*	*	*	*	0.2	2.4	7.4	10.8	19.6	24.3	18.6
Male.....	6.9	*	*	*	*	0.2	3.3	10.9	14.4	23.4	27.1	24.5
Female.....	4.8	*	*	*	*	0.2	1.5	4.0	7.4	16.4	22.4	16.0
Non-Hispanic black, both sexes.....	3.8	*	*	*	*	0.4	2.5	10.1	13.4	14.7	10.6	11.0
Male.....	4.9	*	*	*	*	*	3.0	14.9	20.9	17.5	13.3	*
Female.....	2.8	*	*	*	*	*	2.0	6.0	7.4	12.6	9.1	8.3

Worktable Orig250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 2003

[Rates per 100,000 population, in specified group. Rates based on populations estimated as of July 1, 2003. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pregnancy, childbirth and the puerperium (000-099)												
All origins 3/, both sexes.....	0.2	*	0.3	0.6	0.4	0.1	*	*	*	*
Male.....
Female.....	0.4	*	0.6	1.2	0.7	0.1	*	*	*	*
Hispanic, both sexes.....	0.3	*	0.3	0.6	0.5	*	*	*	*	*
Male.....
Female.....	0.5	*	0.7	1.4	1.0	*	*	*	*	*
Non-Hispanic 4/, both sexes.....	0.2	*	0.3	0.6	0.4	0.1	*	*	*	*
Male.....
Female.....	0.3	*	0.5	1.1	0.7	0.1	*	*	*	*
Non-Hispanic white, both sexes.....	0.1	*	0.2	0.4	0.2	*	*	*	*	*
Male.....
Female.....	0.2	*	0.3	0.7	0.4	*	*	*	*	*
Non-Hispanic black, both sexes.....	0.5	*	0.7	1.6	1.1	*	*	*	*	*
Male.....
Female.....	1.0	*	1.4	3.0	2.0	*	*	*	*	*
Pregnancy with abortive outcome (000-007)												
All origins 3/, both sexes.....	0.0	*	*	*	*	*	*	*	*	*
Male.....
Female.....	0.0	*	*	*	*	*	*	*	*	*
Hispanic, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes.....	0.0	*	*	*	*	*	*	*	*	*
Male.....
Female.....	0.0	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*	*	*	*
Other complications of pregnancy, childbirth and the puerperium (010-099)												
All origins 3/, both sexes.....	0.2	*	0.3	0.5	0.4	0.1	*	*	*	*
Male.....
Female.....	0.3	*	0.5	1.1	0.7	0.1	*	*	*	*
Hispanic, both sexes.....	0.2	*	0.3	0.5	0.5	*	*	*	*	*
Male.....
Female.....	0.5	*	0.7	1.2	1.0	*	*	*	*	*
Non-Hispanic 4/, both sexes.....	0.2	*	0.2	0.5	0.3	0.1	*	*	*	*
Male.....
Female.....	0.3	*	0.5	1.0	0.7	0.1	*	*	*	*
Non-Hispanic white, both sexes.....	0.1	*	0.2	0.3	0.2	*	*	*	*	*
Male.....
Female.....	0.2	*	0.3	0.7	0.4	*	*	*	*	*
Non-Hispanic black, both sexes.....	0.5	*	0.6	1.5	1.0	*	*	*	*	*
Male.....
Female.....	0.9	*	1.1	2.8	1.8	*	*	*	*	*

Worktable Orig250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 2003

[Rates per 100,000 population, in specified group. Rates based on populations estimated as of July 1, 2003. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Certain conditions originating in the perinatal period (P00-P96)												
All origins 3/, both sexes.....	4.9	356.0	0.5	*	*	*	*	*	*	*	*	*
Male.....	5.7	397.4	0.5	*	*	*	*	*	*	*	*	*
Female.....	4.2	312.8	0.5	*	*	*	*	*	*	*	*	*
Hispanic, both sexes.....	6.6	300.3	*	*	*	*	*	*	*	*	*	*
Male.....	7.1	326.4	*	*	*	*	*	*	*	*	*	*
Female.....	6.1	273.0	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes.....	4.6	365.3	0.5	*	*	*	*	*	*	*	*	*
Male.....	5.4	409.7	0.5	*	*	*	*	*	*	*	*	*
Female.....	3.8	318.9	0.5	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes.....	3.2	274.8	0.3	*	*	*	*	*	*	*	*	*
Male.....	3.8	310.6	*	*	*	*	*	*	*	*	*	*
Female.....	2.6	237.3	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes.....	12.9	740.3	1.2	*	*	*	*	*	*	*	*	*
Male.....	15.5	827.9	*	*	*	*	*	*	*	*	*	*
Female.....	10.6	649.2	*	*	*	*	*	*	*	*	*	*
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)												
All origins 3/, both sexes.....	3.6	140.4	3.4	0.9	1.1	1.1	1.3	1.6	2.4	2.3	3.6	6.8
Male.....	3.8	147.6	3.4	0.9	1.3	1.2	1.4	1.7	2.5	2.4	3.5	6.7
Female.....	3.4	132.8	3.5	0.9	0.9	1.0	1.2	1.5	2.3	2.2	3.7	6.8
Hispanic, both sexes.....	4.5	153.0	3.7	0.9	0.8	0.8	0.7	1.0	1.7	*	*	*
Male.....	4.5	156.9	3.7	0.7	0.8	0.9	0.8	1.2	*	*	*	*
Female.....	4.4	148.9	3.7	1.0	0.8	0.6	*	*	*	*	*	*
Non-Hispanic 4/, both sexes.....	3.5	136.0	3.4	0.9	1.1	1.1	1.3	1.7	2.4	2.3	3.7	6.7
Male.....	3.7	144.2	3.3	1.0	1.4	1.2	1.5	1.8	2.5	2.4	3.5	6.7
Female.....	3.3	127.6	3.4	0.9	0.9	1.0	1.2	1.6	2.3	2.2	3.8	6.8
Non-Hispanic white, both sexes.....	3.3	129.8	3.0	0.9	1.1	1.2	1.4	1.7	2.6	2.3	3.8	6.9
Male.....	3.4	136.1	2.9	0.9	1.3	1.2	1.5	1.8	2.6	2.3	3.4	6.8
Female.....	3.2	123.2	3.0	0.9	0.9	1.1	1.3	1.7	2.5	2.3	4.0	6.9
Non-Hispanic black, both sexes.....	4.6	160.5	4.8	1.1	1.5	1.4	1.2	1.7	2.2	2.4	3.1	5.9
Male.....	5.2	173.1	5.0	1.2	1.8	1.5	1.2	2.0	2.8	3.6	*	*
Female.....	4.0	147.3	4.7	1.1	1.2	1.4	1.3	1.5	1.8	*	*	*
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)												
All origins 3/, both sexes.....	10.8	82.9	1.6	0.4	2.0	3.3	5.4	7.0	7.4	13.8	38.9	224.1
Male.....	10.2	96.5	1.8	0.4	2.7	4.5	6.9	9.2	9.6	17.1	41.9	177.9
Female.....	11.4	68.6	1.5	0.4	1.4	2.1	3.8	5.0	5.3	10.9	36.9	244.6
Hispanic, both sexes.....	4.1	56.8	0.9	0.3	1.4	2.3	3.1	3.7	3.5	6.1	18.4	110.9
Male.....	4.8	69.9	*	*	1.7	3.4	4.4	5.7	5.6	8.1	18.8	95.4
Female.....	3.2	43.2	*	*	1.0	1.0	1.6	1.7	*	4.6	18.2	118.8
Non-Hispanic 4/, both sexes.....	11.8	89.4	1.8	0.5	2.1	3.5	5.7	7.3	7.7	14.3	39.9	228.4
Male.....	11.0	102.9	2.0	0.5	2.8	4.7	7.2	9.4	9.9	17.7	43.0	181.1
Female.....	12.6	75.3	1.6	0.5	1.4	2.4	4.1	5.3	5.6	11.4	37.8	249.2
Non-Hispanic white, both sexes.....	12.3	72.6	1.5	0.4	2.2	3.4	5.5	6.7	7.1	13.7	40.1	232.9
Male.....	11.0	87.4	1.6	0.4	2.9	4.6	6.9	8.6	9.0	16.8	42.8	183.3
Female.....	13.5	57.1	1.4	0.4	1.4	2.2	4.0	4.8	5.4	11.0	38.3	254.7
Non-Hispanic black, both sexes.....	12.7	161.6	3.3	0.8	2.5	5.1	8.6	13.6	14.3	24.0	45.7	214.5
Male.....	13.8	171.9	3.6	0.8	3.0	7.0	11.2	17.9	20.7	32.2	52.8	197.1
Female.....	11.6	150.9	3.0	0.8	1.9	3.5	6.3	10.0	9.3	18.2	41.7	221.2

Worktable Orig250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 2003

[Rates per 100,000 population, in specified group. Rates based on populations estimated as of July 1, 2003. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12, G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99, N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39, N41-N64,N80-N98)												
All origins 3/, both sexes.....	69.3	29.7	4.0	2.3	4.9	8.4	18.9	38.9	67.1	144.4	411.5	1,508.4
Male.....	57.3	34.8	4.2	2.6	5.6	9.3	22.3	47.2	76.5	155.6	415.1	1,258.1
Female.....	81.0	24.5	3.8	1.9	4.2	7.4	15.6	31.0	58.4	135.0	409.1	1,619.1
Hispanic, both sexes.....	23.1	27.1	4.3	2.1	4.1	6.1	14.0	30.3	53.5	106.9	284.5	887.2
Male.....	22.2	32.6	4.5	2.3	4.4	7.3	16.7	38.4	64.6	116.4	301.8	757.3
Female.....	24.1	21.5	4.0	2.0	3.8	4.7	11.0	22.1	43.5	99.3	272.6	953.1
Non-Hispanic 4/, both sexes.....	76.5	30.1	4.0	2.3	5.0	8.8	19.6	39.6	68.0	146.7	417.4	1,531.7
Male.....	63.0	35.0	4.2	2.6	5.8	9.7	23.1	47.9	77.2	157.9	420.3	1,278.3
Female.....	89.4	25.1	3.7	1.9	4.2	7.9	16.2	31.7	59.5	137.3	415.5	1,643.1
Non-Hispanic white, both sexes.....	83.0	24.6	3.5	2.1	4.8	7.8	17.8	36.2	64.3	144.1	417.9	1,558.9
Male.....	66.9	30.4	3.6	2.5	6.0	9.0	21.5	44.3	72.6	153.3	416.3	1,300.3
Female.....	98.4	18.6	3.5	1.7	3.6	6.5	14.0	28.3	56.3	136.2	419.1	1,672.7
Non-Hispanic black, both sexes.....	63.0	52.6	5.5	2.8	6.7	15.7	34.0	71.3	113.6	200.3	487.6	1,451.6
Male.....	57.6	56.8	6.3	3.0	6.0	15.6	36.7	84.1	134.7	237.5	552.6	1,243.4
Female.....	67.9	48.4	4.6	2.6	7.3	15.7	31.7	60.3	96.6	173.9	450.9	1,532.3
Accidents (unintentional injuries) (V01-X59,Y85-Y86)												
All origins 3/, both sexes.....	37.6	23.6	10.9	6.4	37.1	31.5	37.8	38.8	32.9	44.1	101.9	278.9
Male.....	49.3	26.4	13.0	7.9	53.3	47.1	54.2	55.9	47.1	58.9	132.9	352.5
Female.....	26.2	20.7	8.7	4.8	19.9	15.3	21.4	22.4	19.7	31.6	81.1	246.4
Hispanic, both sexes.....	26.1	15.1	9.9	5.8	33.8	29.4	31.7	32.8	30.6	38.1	70.9	151.8
Male.....	38.3	17.4	11.9	7.5	50.7	45.5	48.0	50.6	45.3	53.3	102.4	195.5
Female.....	13.1	12.7	7.8	3.9	14.3	10.1	13.8	15.1	17.3	26.0	49.2	129.7
Non-Hispanic 4/, both sexes.....	39.2	25.9	11.1	6.5	37.6	31.8	38.5	39.2	32.9	44.3	103.2	283.7
Male.....	50.9	28.7	13.2	8.0	53.7	47.3	54.9	56.0	47.0	59.0	134.1	358.9
Female.....	28.1	22.9	8.9	5.0	20.9	16.3	22.4	23.0	19.8	32.0	82.6	250.6
Non-Hispanic white, both sexes.....	41.6	21.6	10.7	6.2	41.1	33.5	39.8	38.7	32.2	43.9	105.5	296.5
Male.....	53.1	24.1	12.9	7.5	58.2	49.5	56.3	54.7	45.4	58.2	135.7	372.9
Female.....	30.5	19.0	8.4	4.8	23.2	17.2	23.2	22.9	19.6	31.7	85.1	262.8
Non-Hispanic black, both sexes.....	33.3	44.7	13.7	8.0	26.4	30.9	39.7	49.3	42.9	51.1	83.1	161.8
Male.....	47.4	49.0	15.9	10.1	39.8	48.4	59.4	75.8	69.0	74.9	123.8	228.0
Female.....	20.5	40.3	11.4	5.9	12.9	14.9	22.4	26.6	22.0	34.1	60.1	136.2
Transport accidents (V01-V99,Y85)												
All origins 3/, both sexes.....	16.5	3.7	4.1	4.3	27.7	18.3	17.1	16.6	15.9	17.5	26.2	29.7
Male.....	23.3	4.0	4.5	5.2	38.6	27.3	24.8	24.3	23.0	23.7	36.6	52.0
Female.....	10.0	3.4	3.7	3.4	16.3	9.0	9.4	9.1	9.4	12.4	19.3	19.9
Hispanic, both sexes.....	15.2	4.5	4.2	4.1	26.2	19.1	16.4	15.1	15.8	18.6	23.6	24.9
Male.....	22.2	5.7	4.6	5.5	38.2	29.3	23.9	22.4	21.4	26.2	35.3	46.2
Female.....	7.6	*	3.8	2.7	12.3	7.0	8.3	7.9	10.7	12.5	15.5	*
Non-Hispanic 4/, both sexes.....	16.7	3.5	4.0	4.3	28.0	18.0	17.2	16.6	15.8	17.3	26.3	29.8
Male.....	23.4	3.5	4.5	5.1	38.5	26.7	24.9	24.4	23.0	23.3	36.6	52.0
Female.....	10.3	3.5	3.6	3.6	17.0	9.4	9.6	9.2	9.2	12.3	19.5	20.1
Non-Hispanic white, both sexes.....	17.2	3.2	3.8	4.4	30.2	18.2	17.3	16.6	15.6	17.3	26.6	30.6
Male.....	23.9	3.3	4.4	5.1	41.1	26.8	24.9	24.1	22.6	23.3	36.6	52.8
Female.....	10.7	3.0	3.2	3.7	18.8	9.5	9.7	9.2	8.9	12.1	19.8	20.8
Non-Hispanic black, both sexes.....	15.6	4.4	5.1	4.3	20.3	20.0	18.8	18.6	18.7	18.2	22.6	21.4
Male.....	23.3	*	5.1	5.3	30.3	31.8	29.1	29.6	29.6	26.5	38.1	43.7
Female.....	8.5	*	5.2	3.3	10.1	9.3	9.7	9.0	9.9	12.3	13.9	12.8

Worktable Orig250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 2003

[Rates per 100,000 population, in specified group. Rates based on populations estimated as of July 1, 2003. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental drowning and submersion (W65-W74)												
All origins 3/, both sexes.....	1.1	1.4	2.9	0.7	1.4	0.9	1.0	1.0	0.9	1.1	1.3	1.8
Male.....	1.8	1.7	3.7	0.9	2.5	1.6	1.8	1.6	1.6	1.7	2.0	3.3
Female.....	0.5	1.2	2.0	0.3	0.2	0.2	0.3	0.4	0.4	0.6	0.8	1.1
Hispanic, both sexes.....	1.1	*	2.5	0.6	1.6	1.2	0.9	0.7	*	*	*	*
Male.....	1.9	*	4.0	0.9	2.9	2.0	1.5	1.3	*	*	*	*
Female.....	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes.....	1.1	1.5	3.0	0.7	1.3	0.8	1.1	1.0	1.0	1.1	1.3	1.8
Male.....	1.8	1.7	3.7	1.0	2.4	1.5	1.8	1.6	1.6	1.8	2.1	3.4
Female.....	0.5	*	2.3	0.3	0.2	0.2	0.3	0.4	0.4	0.5	0.7	1.1
Non-Hispanic white, both sexes.....	1.1	1.4	3.3	0.5	1.2	0.8	1.1	1.0	0.9	1.1	1.3	1.7
Male.....	1.7	*	4.0	0.7	2.1	1.3	1.8	1.5	1.5	1.8	2.1	3.4
Female.....	0.5	*	2.7	0.3	0.2	0.2	0.3	0.4	0.4	0.5	0.7	0.9
Non-Hispanic black, both sexes.....	1.3	*	2.1	1.4	1.7	1.1	0.9	1.0	1.2	*	*	*
Male.....	2.3	*	3.0	2.0	3.1	2.2	1.7	2.0	2.4	*	*	*
Female.....	0.4	*	*	0.7	*	*	*	*	*	*	*	*
Accidental exposure to smoke, fire and flames (X00-X09)												
All origins 3/, both sexes.....	1.2	0.7	1.4	0.5	0.5	0.6	0.9	1.3	1.4	2.3	3.6	5.4
Male.....	1.4	*	1.6	0.5	0.6	0.8	1.2	1.6	1.9	2.9	4.4	7.4
Female.....	1.0	1.1	1.2	0.5	0.4	0.5	0.6	0.9	1.0	1.8	3.0	4.5
Hispanic, both sexes.....	0.5	*	0.8	0.3	*	0.3	0.4	0.8	*	*	*	*
Male.....	0.6	*	*	*	*	*	*	1.4	*	*	*	*
Female.....	0.4	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes.....	1.3	0.8	1.6	0.6	0.5	0.7	1.0	1.3	1.5	2.4	3.6	5.4
Male.....	1.5	*	1.9	0.6	0.6	0.8	1.3	1.7	1.9	3.0	4.5	7.3
Female.....	1.0	1.3	1.3	0.5	0.4	0.6	0.7	1.0	1.0	1.8	3.1	4.5
Non-Hispanic white, both sexes.....	1.2	*	1.3	0.4	0.5	0.7	1.0	1.2	1.3	2.1	3.1	4.7
Male.....	1.4	*	1.6	0.4	0.7	0.8	1.3	1.5	1.7	2.6	3.6	6.5
Female.....	1.0	*	1.0	0.4	0.4	0.6	0.6	0.9	0.9	1.6	2.8	3.9
Non-Hispanic black, both sexes.....	2.1	*	3.1	1.5	0.6	0.9	1.2	2.6	2.9	5.4	10.3	14.9
Male.....	2.5	*	3.4	1.5	*	1.1	1.5	3.4	4.4	7.7	16.2	*
Female.....	1.8	*	2.8	1.4	0.7	*	0.9	2.0	1.7	3.8	7.0	13.2
Accidental poisoning and exposure to noxious substances (X40-X49)												
All origins 3/, both sexes.....	6.7	0.5	0.3	0.1	4.9	8.6	14.0	13.3	4.9	2.2	2.2	3.5
Male.....	9.2	*	0.3	0.1	7.4	12.4	18.8	18.0	6.3	2.7	2.8	4.0
Female.....	4.3	*	0.3	0.2	2.2	4.7	9.3	8.8	3.7	1.8	1.8	3.3
Hispanic, both sexes.....	4.5	*	*	*	3.1	5.7	10.3	10.5	4.4	1.8	*	*
Male.....	6.9	*	*	*	4.8	8.6	15.6	16.4	7.4	*	*	*
Female.....	1.9	*	*	*	1.2	2.2	4.5	4.6	*	*	*	*
Non-Hispanic 4/, both sexes.....	7.0	*	0.3	0.2	5.2	9.2	14.5	13.5	4.9	2.2	2.2	3.5
Male.....	9.5	*	0.3	0.2	7.9	13.3	19.2	17.9	6.1	2.6	2.8	4.0
Female.....	4.6	*	*	0.2	2.4	5.2	9.9	9.2	3.8	1.9	1.8	3.3
Non-Hispanic white, both sexes.....	7.4	*	*	0.1	6.4	10.6	15.5	13.4	4.8	2.0	2.3	3.5
Male.....	10.1	*	*	*	9.8	15.3	20.3	17.5	5.6	2.1	2.9	3.6
Female.....	4.9	*	*	*	2.9	5.9	10.6	9.4	4.0	1.9	1.9	3.5
Non-Hispanic black, both sexes.....	6.4	*	*	*	1.4	5.8	13.9	18.3	7.6	4.1	*	*
Male.....	9.1	*	*	*	2.1	8.3	19.2	26.6	12.9	7.5	*	*
Female.....	4.1	*	*	*	0.7	3.6	9.3	11.1	3.4	*	*	*

Worktable Orig250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 2003

[Rates per 100,000 population, in specified group. Rates based on populations estimated as of July 1, 2003. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)												
All origins 3/, both sexes.....	5.9	16.9	1.8	0.5	1.6	2.0	3.0	3.9	5.1	9.6	27.5	102.5
Male.....	6.9	19.5	2.2	0.8	2.6	3.4	4.8	6.0	7.8	13.1	34.9	120.2
Female.....	4.9	14.1	1.3	0.3	0.5	0.7	1.2	1.9	2.6	6.7	22.6	94.7
Hispanic, both sexes.....	2.5	8.2	1.6	0.4	1.6	2.0	2.2	3.0	3.9	6.3	16.9	37.3
Male.....	3.6	9.1	1.9	0.5	2.9	3.4	3.7	4.7	6.4	9.0	24.9	44.6
Female.....	1.3	7.3	1.3	*	*	*	*	1.3	*	4.1	11.3	33.6
Non-Hispanic 4/, both sexes.....	6.4	19.2	1.8	0.6	1.5	2.0	3.1	4.0	5.2	9.8	28.0	105.1
Male.....	7.5	22.2	2.3	0.8	2.5	3.4	4.9	6.1	7.9	13.4	35.3	123.5
Female.....	5.4	16.0	1.3	0.3	0.6	0.7	1.3	1.9	2.7	6.8	23.1	97.0
Non-Hispanic white, both sexes.....	6.9	15.5	1.7	0.6	1.6	2.2	3.2	3.8	5.0	9.6	28.3	109.1
Male.....	7.9	18.0	2.2	0.8	2.7	3.6	5.0	5.9	7.5	13.1	35.3	127.5
Female.....	6.0	12.9	1.2	0.3	0.5	0.7	1.3	1.8	2.6	6.7	23.6	101.0
Non-Hispanic black, both sexes.....	5.1	35.7	2.1	0.5	1.3	1.8	3.4	5.5	7.5	13.1	27.8	74.1
Male.....	6.6	41.6	2.9	0.7	1.8	3.0	5.3	8.7	12.2	18.8	39.8	92.7
Female.....	3.7	29.7	*	*	0.8	0.8	1.6	2.8	3.7	9.0	21.0	66.9
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
All origins 3/, both sexes.....	10.8	0.6	9.7	12.7	14.9	15.9	13.8	12.7	16.4	16.9
Male.....	17.6	0.9	16.0	20.6	23.2	24.4	22.3	23.4	35.1	47.8
Female.....	4.3	0.3	3.0	4.6	6.6	7.7	5.9	3.8	4.0	3.3
Hispanic, both sexes.....	5.0	0.6	7.0	6.8	6.8	7.2	6.9	6.5	8.0	10.9
Male.....	8.3	0.8	11.2	10.8	11.0	11.5	13.0	13.1	18.0	*
Female.....	1.5	*	2.2	1.9	2.2	2.9	*	*	*	*
Non-Hispanic 4/, both sexes.....	11.7	0.6	10.2	14.0	16.0	16.7	14.3	13.1	16.9	17.1
Male.....	19.1	0.9	16.9	22.9	25.1	25.6	22.9	24.1	36.0	48.6
Female.....	4.6	0.3	3.2	5.2	7.2	8.1	6.3	4.0	4.1	3.3
Non-Hispanic white, both sexes.....	13.3	0.6	10.9	15.5	18.3	19.1	15.8	14.3	18.2	18.4
Male.....	21.6	1.0	18.2	25.0	28.2	29.0	25.1	26.0	38.6	52.5
Female.....	5.3	0.2	3.3	5.8	8.3	9.3	7.0	4.3	4.4	3.4
Non-Hispanic black, both sexes.....	5.3	0.7	7.3	9.6	7.4	5.7	4.9	4.6	4.5	*
Male.....	9.0	1.0	12.5	17.1	12.5	9.3	8.5	8.5	11.3	*
Female.....	1.8	*	2.0	2.7	2.9	2.7	2.0	*	*	*
Intentional self-harm (suicide) by discharge of firearms (X72-X74)												
All origins 3/, both sexes.....	5.8	0.2	5.0	6.0	6.6	8.0	8.3	9.3	12.4	11.9
Male.....	10.4	0.3	8.9	10.4	11.2	13.8	14.9	18.4	28.7	36.6
Female.....	1.4	*	1.0	1.4	2.1	2.5	2.2	1.6	1.5	0.9
Hispanic, both sexes.....	2.1	*	3.3	2.8	2.5	3.0	2.7	3.1	4.1	*
Male.....	3.8	*	5.6	4.8	4.3	5.3	5.6	7.0	9.6	*
Female.....	0.3	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes.....	6.4	0.2	5.4	6.7	7.2	8.5	8.7	9.7	12.8	12.1
Male.....	11.5	0.3	9.6	11.7	12.2	14.7	15.6	19.2	29.7	37.5
Female.....	1.6	*	1.0	1.6	2.3	2.6	2.3	1.7	1.6	0.9
Non-Hispanic white, both sexes.....	7.4	0.2	5.8	7.4	8.4	9.9	9.9	10.8	14.0	13.2
Male.....	13.2	0.4	10.3	12.7	14.1	16.8	17.4	21.2	32.1	41.0
Female.....	1.8	*	1.1	2.0	2.8	3.1	2.7	1.9	1.7	1.0
Non-Hispanic black, both sexes.....	2.7	*	4.4	5.4	2.7	2.5	2.8	3.4	4.0	*
Male.....	5.1	*	8.0	10.5	5.1	4.8	5.3	6.8	9.8	*
Female.....	0.5	*	0.7	*	*	*	*	*	*	*

Worktable Orig250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 2003

[Rates per 100,000 population, in specified group. Rates based on populations estimated as of July 1, 2003. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)												
All origins 3/, both sexes.....	5.0	0.4	4.6	6.7	8.3	7.8	5.5	3.5	4.0	5.1
Male.....	7.3	0.6	7.1	10.2	12.1	10.6	7.3	5.0	6.4	11.1
Female.....	2.8	0.2	2.1	3.2	4.5	5.2	3.7	2.2	2.5	2.4
Hispanic, both sexes.....	2.9	0.5	3.8	3.9	4.3	4.2	4.2	3.4	3.9	*
Male.....	4.5	0.6	5.6	6.0	6.7	6.2	7.5	6.1	8.4	*
Female.....	1.2	*	1.6	1.5	1.7	2.3	*	*	*	*
Non-Hispanic 4/, both sexes.....	5.3	0.4	4.8	7.3	8.8	8.2	5.5	3.4	4.0	5.1
Male.....	7.7	0.6	7.4	11.2	12.8	11.0	7.3	4.9	6.3	11.2
Female.....	3.1	0.2	2.1	3.5	4.9	5.5	3.9	2.2	2.6	2.4
Non-Hispanic white, both sexes.....	5.9	0.4	5.1	8.1	9.9	9.2	6.0	3.5	4.2	5.2
Male.....	8.4	0.6	7.9	12.3	14.1	12.2	7.7	4.9	6.5	11.6
Female.....	3.4	0.2	2.2	3.9	5.6	6.2	4.3	2.4	2.7	2.4
Non-Hispanic black, both sexes.....	2.5	0.6	2.9	4.2	4.7	3.2	2.1	1.3	*	*
Male.....	3.9	0.9	4.4	6.6	7.4	4.5	3.3	*	*	*
Female.....	1.3	*	1.3	2.0	2.3	2.1	*	*	*	*
Assault (homicide) (* -*U02,X85-Y09,Y87.1)												
All origins 3/, both sexes.....	6.1	8.5	2.4	0.8	13.0	11.3	7.0	4.9	2.8	2.4	2.5	2.2
Male.....	9.7	10.0	2.5	1.0	21.8	18.5	10.4	7.4	4.2	3.4	3.3	3.2
Female.....	2.6	6.9	2.3	0.5	3.7	3.9	3.6	2.5	1.5	1.5	1.9	1.8
Hispanic, both sexes.....	8.4	6.9	2.5	0.7	18.4	13.2	8.5	6.4	3.6	3.8	3.6	*
Male.....	13.6	7.9	2.1	0.9	30.3	21.2	12.9	10.2	6.0	6.8	*	*
Female.....	2.8	5.9	2.9	*	4.5	3.5	3.6	2.7	*	*	*	*
Non-Hispanic 4/, both sexes.....	5.7	8.9	2.3	0.8	11.8	10.8	6.7	4.8	2.7	2.2	2.4	2.1
Male.....	9.0	10.5	2.6	1.0	19.8	17.7	10.0	7.1	4.0	3.1	3.2	3.0
Female.....	2.5	7.2	2.1	0.6	3.5	4.0	3.6	2.5	1.5	1.5	1.8	1.8
Non-Hispanic white, both sexes.....	2.7	6.8	1.3	0.4	3.5	3.9	3.8	3.1	2.1	1.7	2.0	2.0
Male.....	3.6	8.0	1.3	0.6	5.0	5.2	4.9	4.3	2.8	2.3	2.6	2.9
Female.....	1.8	5.5	1.4	0.3	1.9	2.5	2.6	2.0	1.4	1.3	1.6	1.6
Non-Hispanic black, both sexes.....	22.7	16.8	6.4	2.3	49.1	47.1	24.2	15.8	8.1	6.3	6.6	*
Male.....	40.1	18.6	7.8	3.0	87.3	86.1	41.6	26.9	14.9	11.3	11.3	*
Female.....	6.8	14.8	4.9	1.5	10.4	11.6	8.8	6.2	2.7	2.8	3.9	*
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
All origins 3/, both sexes.....	4.1	*	0.3	0.5	10.7	8.9	4.4	2.7	1.4	0.9	0.7	0.5
Male.....	7.1	*	0.2	0.6	18.8	15.5	6.9	4.3	2.2	1.4	0.7	*
Female.....	1.2	*	0.3	0.3	2.2	2.1	1.8	1.2	0.7	0.5	0.7	*
Hispanic, both sexes.....	5.8	*	*	0.4	14.9	9.9	5.3	3.6	1.2	*	*	*
Male.....	10.0	*	*	0.6	25.5	16.8	8.4	5.9	2.2	*	*	*
Female.....	1.3	*	*	*	2.7	1.7	1.9	1.4	*	*	*	*
Non-Hispanic 4/, both sexes.....	3.8	*	0.2	0.5	9.8	8.6	4.2	2.6	1.4	0.8	0.7	0.5
Male.....	6.5	*	*	0.7	17.2	15.0	6.7	4.1	2.1	1.3	0.7	*
Female.....	1.2	*	*	0.3	2.0	2.2	1.8	1.2	0.7	0.5	0.7	*
Non-Hispanic white, both sexes.....	1.4	*	*	0.2	2.4	2.5	2.1	1.7	1.1	0.7	0.6	0.5
Male.....	2.1	*	*	0.3	3.7	3.7	2.9	2.4	1.6	1.0	0.5	*
Female.....	0.8	*	*	*	1.0	1.3	1.3	1.0	0.7	0.4	0.6	*
Non-Hispanic black, both sexes.....	17.3	*	*	1.5	43.3	40.5	16.5	8.9	4.0	2.5	*	*
Male.....	32.6	*	*	2.1	79.4	77.4	30.1	16.4	7.4	4.8	*	*
Female.....	3.4	*	*	0.9	6.7	6.9	4.4	2.6	*	*	*	*

Worktable Orig250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 2003

[Rates per 100,000 population, in specified group. Rates based on populations estimated as of July 1, 2003. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02.X85-X92,X96-Y09,Y87.1)												
All origins 3/, both sexes.....	2.0	8.3	2.1	0.3	2.3	2.4	2.6	2.2	1.4	1.5	1.8	1.7
Male.....	2.6	9.7	2.3	0.4	3.1	3.0	3.5	3.1	2.1	2.0	2.6	2.5
Female.....	1.4	6.9	2.0	0.3	1.6	1.8	1.8	1.3	0.8	1.0	1.2	1.4
Hispanic, both sexes.....	2.6	6.8	2.2	0.3	3.5	3.3	3.2	2.8	2.3	2.4	3.1	*
Male.....	3.6	7.7	1.8	*	4.8	4.5	4.6	4.3	3.7	4.4	*	*
Female.....	1.5	5.9	2.6	*	1.9	1.8	1.7	1.4	*	*	*	*
Non-Hispanic 4/, both sexes.....	1.9	8.6	2.1	0.3	2.1	2.2	2.5	2.1	1.3	1.4	1.7	1.6
Male.....	2.4	10.1	2.4	0.4	2.6	2.7	3.3	3.0	1.9	1.8	2.5	2.3
Female.....	1.4	7.1	1.8	0.3	1.5	1.8	1.8	1.3	0.8	1.0	1.1	1.3
Non-Hispanic white, both sexes.....	1.2	6.8	1.2	0.2	1.1	1.4	1.6	1.4	1.0	1.1	1.4	1.5
Male.....	1.5	8.0	1.2	0.3	1.3	1.5	2.0	1.9	1.3	1.3	2.1	2.2
Female.....	1.0	5.4	1.2	0.2	1.0	1.2	1.3	1.0	0.7	0.9	1.0	1.2
Non-Hispanic black, both sexes.....	5.4	15.8	5.6	0.8	5.8	6.6	7.7	6.8	4.2	3.8	4.9	*
Male.....	7.6	16.8	6.9	0.9	7.9	8.7	11.5	10.5	7.4	6.5	8.7	*
Female.....	3.4	14.8	4.3	0.7	3.7	4.8	4.4	3.6	1.5	*	*	*
Legal intervention (Y35,Y89.0)												
All origins 3/, both sexes.....	0.1	*	*	*	0.2	0.3	0.3	0.1	*	*	*	*
Male.....	0.3	*	*	*	0.4	0.6	0.5	0.2	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes.....	0.2	*	*	*	0.4	0.4	*	*	*	*	*	*
Male.....	0.4	*	*	*	0.7	0.8	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes.....	0.1	*	*	*	0.2	0.3	0.3	0.1	*	*	*	*
Male.....	0.3	*	*	*	0.3	0.6	0.5	0.2	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes.....	0.1	*	*	*	0.1	0.2	0.3	0.1	*	*	*	*
Male.....	0.2	*	*	*	0.2	0.3	0.5	0.2	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes.....	0.3	*	*	*	0.5	1.0	0.4	*	*	*	*	*
Male.....	0.7	*	*	*	1.0	2.1	0.8	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*	*	*
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)												
All origins 3/, both sexes.....	1.7	1.9	0.2	0.1	1.4	2.1	3.6	3.1	1.3	0.7	0.7	1.1
Male.....	2.3	2.4	*	0.2	1.9	2.9	4.6	4.0	1.7	0.8	1.0	1.5
Female.....	1.2	1.3	*	*	0.8	1.3	2.6	2.2	1.0	0.7	0.6	0.9
Hispanic, both sexes.....	0.8	*	*	*	0.7	1.2	1.4	1.7	*	*	*	*
Male.....	1.3	*	*	*	1.0	1.7	2.1	2.6	*	*	*	*
Female.....	0.4	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes.....	1.9	2.2	0.2	0.1	1.5	2.3	3.9	3.2	1.3	0.7	0.7	1.1
Male.....	2.5	3.0	*	0.2	2.2	3.1	4.9	4.2	1.7	0.8	0.9	*
Female.....	1.3	1.3	*	*	0.8	1.5	2.8	2.3	1.0	0.7	0.6	1.0
Non-Hispanic white, both sexes.....	2.0	2.0	*	0.1	1.7	2.6	4.1	3.3	1.3	0.7	0.7	1.2
Male.....	2.5	2.4	*	0.2	2.4	3.4	5.1	4.0	1.5	0.7	0.9	*
Female.....	1.4	*	*	*	1.0	1.7	3.1	2.5	1.1	0.6	0.6	1.1
Non-Hispanic black, both sexes.....	1.8	3.3	*	*	0.8	1.8	3.9	4.1	2.0	1.3	*	*
Male.....	2.8	*	*	*	1.4	3.0	5.4	6.6	3.3	*	*	*
Female.....	1.0	*	*	*	*	*	2.5	2.0	*	*	*	*

Worktable Orig250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 2003

[Rates per 100,000 population, in specified group. Rates based on populations estimated as of July 1, 2003. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Complications of medical and surgical care (Y40-Y84,Y88)												
All origins 3/, both sexes.....	1.0	*	0.2	*	0.1	0.2	0.4	0.7	1.4	2.9	6.2	11.2
Male.....	0.9	*	*	*	0.1	0.2	0.4	0.7	1.5	3.2	6.9	12.9
Female.....	1.0	*	*	*	*	0.2	0.4	0.6	1.3	2.7	5.7	10.5
Hispanic, both sexes.....	0.4	*	*	*	*	*	0.4	*	1.0	3.1	3.7	*
Male.....	0.4	*	*	*	*	*	*	*	*	*	*	*
Female.....	0.4	*	*	*	*	*	*	*	3.1	*	*	*
Non-Hispanic 4/, both sexes.....	1.1	*	*	*	0.1	0.2	0.4	0.7	1.4	2.9	6.3	11.4
Male.....	1.0	*	*	*	0.1	0.2	0.4	0.7	1.5	3.2	7.0	13.2
Female.....	1.1	*	*	*	*	0.2	0.4	0.6	1.3	2.7	5.8	10.6
Non-Hispanic white, both sexes.....	1.1	*	*	*	0.1	0.2	0.4	0.6	1.3	2.8	6.2	11.2
Male.....	1.1	*	*	*	*	0.2	0.4	0.6	1.4	3.1	7.1	13.1
Female.....	1.1	*	*	*	*	*	0.3	0.6	1.1	2.5	5.5	10.4
Non-Hispanic black, both sexes.....	1.2	*	*	*	*	*	0.6	1.4	2.8	4.6	9.0	15.2
Male.....	1.0	*	*	*	*	*	*	1.9	2.9	3.9	7.8	*
Female.....	1.3	*	*	*	*	*	0.8	1.0	2.6	5.0	9.6	15.3
Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) /5												
All origins 3/, both sexes.....	7.1	*	*	*	0.3	1.5	8.1	17.3	18.5	15.1	9.2	4.1
Male.....	10.9	*	*	*	0.5	2.3	11.4	27.0	29.8	25.4	16.8	7.8
Female.....	3.4	*	*	*	0.1	0.6	4.7	7.9	8.1	6.5	4.1	2.5
Hispanic, both sexes.....	6.1	*	*	*	0.6	1.8	8.3	20.2	27.6	23.0	17.8	*
Male.....	9.9	*	*	*	0.9	2.8	13.2	35.2	50.5	40.4	34.9	*
Female.....	1.9	*	*	*	*	0.6	2.9	5.2	6.9	9.1	6.1	*
Non-Hispanic 4/, both sexes.....	7.2	*	*	*	0.2	1.4	8.0	16.9	17.7	14.5	8.7	4.1
Male.....	11.0	*	*	*	0.4	2.2	11.1	26.0	27.9	24.2	15.8	7.8
Female.....	3.6	*	*	*	*	0.6	4.9	8.2	8.2	6.3	4.0	2.5
Non-Hispanic white, both sexes.....	7.5	*	*	*	0.2	1.4	8.1	16.9	17.1	14.4	8.8	4.1
Male.....	11.4	*	*	*	0.4	2.2	11.3	25.5	26.7	24.0	15.7	7.7
Female.....	3.8	*	*	*	*	0.7	4.9	8.3	8.0	6.3	4.2	2.5
Non-Hispanic black, both sexes.....	6.5	*	*	*	*	1.2	7.0	18.7	24.5	17.1	9.1	*
Male.....	10.3	*	*	*	*	1.9	10.0	31.6	42.4	30.9	19.3	*
Female.....	3.0	*	*	*	*	*	4.4	7.6	10.1	7.3	3.3	*
Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) /5												
All origins 3/, both sexes.....	9.9	0.6	0.2	0.1	6.4	12.2	20.6	20.0	8.0	4.1	4.2	6.0
Male.....	12.9	*	0.3	0.1	9.2	16.6	26.5	25.7	9.4	4.6	4.9	7.8
Female.....	7.0	*	*	0.1	3.3	7.6	14.8	14.6	6.7	3.7	3.7	5.2
Hispanic, both sexes.....	5.9	*	*	*	3.5	7.2	13.3	14.9	6.5	3.1	3.7	*
Male.....	8.7	*	*	*	5.2	10.5	19.5	22.1	10.5	5.2	*	*
Female.....	2.9	*	*	*	1.5	3.3	6.5	7.8	2.8	*	*	*
Non-Hispanic 4/, both sexes.....	10.4	0.7	0.2	0.1	6.9	13.2	21.6	20.4	8.1	4.2	4.2	6.0
Male.....	13.5	*	*	*	10.1	18.0	27.5	25.9	9.2	4.5	4.9	7.8
Female.....	7.6	*	*	*	3.7	8.4	15.9	15.2	7.0	3.9	3.7	5.3
Non-Hispanic white, both sexes.....	11.2	*	*	0.1	8.5	15.2	23.0	20.3	7.9	4.2	4.5	6.3
Male.....	14.3	*	*	*	12.5	20.8	29.0	25.1	8.4	4.2	5.1	7.8
Female.....	8.2	*	*	*	4.4	9.5	17.0	15.7	7.4	4.1	4.0	5.7
Non-Hispanic black, both sexes.....	9.5	*	*	*	1.9	8.3	21.4	27.5	11.7	4.9	2.2	*
Male.....	13.0	*	*	*	2.5	10.9	28.4	39.7	19.0	8.4	*	*
Female.....	6.3	*	*	*	1.2	5.9	15.2	17.1	5.9	2.5	*	*

Worktable Orig250R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 2003

[Rates per 100,000 population, in specified group. Rates based on populations estimated as of July 1, 2003. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/												
All origins 3/, both sexes.....	10.4	*	0.3	0.8	16.6	15.5	11.5	11.2	10.1	10.4	13.5	12.5
Male.....	18.3	*	0.3	1.2	29.2	27.1	19.1	18.8	17.7	20.3	30.2	37.8
Female.....	2.7	*	0.3	0.3	3.3	3.6	4.0	3.8	2.9	2.2	2.2	1.3
Hispanic, both sexes.....	8.3	*	*	0.7	19.2	13.3	8.1	6.8	4.1	4.7	4.7	*
Male.....	14.6	*	*	1.0	32.8	22.8	13.3	11.5	8.1	10.0	10.7	*
Female.....	1.7	*	*	*	3.5	2.1	2.4	2.0	*	*	*	*
Non-Hispanic 4/, both sexes.....	10.6	*	0.3	0.8	16.0	16.0	12.0	11.6	10.5	10.8	13.9	12.7
Male.....	18.8	*	0.4	1.3	28.3	28.0	19.9	19.5	18.4	21.0	31.2	38.7
Female.....	2.9	*	*	0.3	3.2	3.9	4.2	4.0	3.1	2.3	2.3	1.3
Non-Hispanic white, both sexes.....	9.3	*	*	0.6	8.8	10.4	11.2	12.0	11.3	11.8	14.9	13.8
Male.....	16.0	*	*	0.9	15.2	17.3	18.1	19.9	19.7	22.7	33.5	42.1
Female.....	2.7	*	*	0.2	2.2	3.4	4.3	4.2	3.4	2.4	2.4	1.4
Non-Hispanic black, both sexes.....	20.7	*	0.9	1.8	49.3	47.4	19.7	11.7	6.9	6.2	6.0	*
Male.....	39.1	*	*	2.6	90.5	91.1	36.1	21.6	13.1	12.0	13.6	*
Female.....	4.0	*	*	0.9	7.6	7.6	5.2	3.2	2.0	2.1	*	*

* Figure does not meet standards of reliability or precision.

0.0 Quantity more than zero but less than 0.05.

... Category not applicable.

1/ Figures for age not stated included in "All ages" but not distributed among age groups.

2/ Death rates for "Under 1 year"(based on population estimates) differ from infant mortality rates (based on live births).

3/ Figures for origin not stated are included in "All origins" but not distributed among specified origins.

4/ Includes races other than white and black.

5/ Included in selected categories.

Source: CDC/NCHS, National Vital Statistics System, Mortality.