

Health, United States, 2015

With Special Feature on Racial and Ethnic Health Disparities

This report was updated on June 22, 2017 to reflect corrections to the 2014 mortality data. Changes appear in the highlighted areas of the individual PDF and spreadsheet versions of Tables 17, 18, 19, 20, 29, 30, and 31, and Figure 3 (also updated in the PPT file). For more information about the corrections to the 2014 mortality data, please refer to "Deaths: Final Data for 2014," available from: https://www.cdc.gov/nchs/data/nvsr/nvsr65/nvsr65_04.pdf.



U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Centers for Disease Control and Prevention
National Center for Health Statistics

Copyright information

Permission has been obtained from the copyright holders to reproduce certain quoted material in this report. Further reproduction of this material is prohibited without specific permission of the copyright holder. All other material contained in this report is in the public domain and may be used and reprinted without special permission; citation as to source, however, is appreciated.


Suggested citation

National Center for Health Statistics.
Health, United States, 2015: With Special Feature on Racial
and Ethnic Health Disparities. Hyattsville, MD. 2016.

Library of Congress Catalog Number 76-641496
For sale by Superintendent of Documents
U.S. Government Printing Office
Washington, DC 20402

Health, United States, 2015

With Special Feature on Racial and Ethnic Health Disparities



U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Centers for Disease Control and Prevention
National Center for Health Statistics

May 2016
DHHS Publication No. 2016-1232

U.S. Department of Health and Human Services

Sylvia M. Burwell
Secretary

Centers for Disease Control and Prevention

Thomas R. Frieden, M.D., M.P.H.
Director

National Center for Health Statistics

Charles J. Rothwell, M.S., M.B.A.
Director

Preface

Health, United States, 2015 is the 39th report on the health status of the nation and is submitted by the Secretary of the Department of Health and Human Services to the President and the Congress of the United States in compliance with Section 308 of the Public Health Service Act. This report was compiled by the Centers for Disease Control and Prevention's (CDC) National Center for Health Statistics (NCHS).

The *Health, United States* series presents an annual overview of national trends in health statistics. The report contains a Chartbook that assesses the nation's health by presenting trends and current information on selected measures of morbidity, mortality, health care utilization and access, health risk factors, prevention, health insurance, and personal health care expenditures. This year's Chartbook includes a Special Feature on racial and ethnic health disparities. The report also contains 114 Trend Tables organized around four major subject areas: health status and determinants, health care utilization, health care resources, and health care expenditures. A companion report—*Health, United States: In Brief*—features information extracted from the full report. The complete report, *In Brief*, and related data products are available on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

The 2015 Edition

Health, United States, 2015 contains a summary At a Glance table that displays selected indicators of health and their determinants, cross-referenced to charts and tables in the report. This is followed by a Highlights section, a Chartbook, detailed Trend Tables, two Appendixes, and an Index. The major sections of the 2015 report are described below.

Chartbook

The 2015 Chartbook contains 27 figures, including 10 figures in this year's Special Feature on racial and ethnic health disparities (Figures 18–27). The special feature topic was chosen to commemorate the 30th anniversary of the Report of the Secretary's Task Force on Black and Minority Health (also known as the Heckler Report), which documented significant disparities in the burden of illness and mortality experienced by blacks and other minority groups compared with white persons (1). This year's Special Feature provides a current overview of racial and ethnic differences in health, by race, race and ethnicity, or detailed Hispanic origin, depending on data availability. Data are presented on selected measures of mortality (life expectancy and infant mortality), natality (preterm births and low-risk cesarean delivery), health status and risk factors (obesity, hypertension, and cigarette smoking), and health care

access and utilization (influenza vaccination, lack of health insurance coverage, and difficulty accessing needed dental care) and summarizes whether racial and ethnic differences have been increasing, decreasing, or remaining stable over time.

Trend Tables

The Chartbook is followed by 114 detailed Trend Tables that highlight major trends in health statistics. Comparability across editions of *Health, United States* is fostered by including similar Trend Tables in each volume, and timeliness is maintained by improving the content of tables to reflect key topics in public health. An important criterion used in selecting these tables is the availability of comparable national data over a period of several years.

Appendixes

Appendix I. Data Sources describes each data source used in *Health, United States, 2015* and provides references for further information about the sources. Data sources are listed alphabetically within two broad categories: Government Sources, and Private and Global Sources.

Appendix II. Definitions and Methods is an alphabetical listing of selected terms used in *Health, United States, 2015*. It also contains information on the statistical methodologies used in the report.

Index

The Index to the Trend Tables and Chartbook figures is a useful tool for locating data by topic. Tables and figures are cross-referenced by such topics as child and adolescent health; older population aged 65 and over; women's health; men's health; state data; American Indian or Alaska Native, Asian, black or African American, Hispanic-origin, and white populations; education; injury; disability; and metropolitan and nonmetropolitan data. Many of the Index topics are also available as conveniently grouped data packages on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

Data Considerations

Racial and Ethnic Data

Many tables in *Health, United States* present data according to race and Hispanic origin, consistent with a department-wide emphasis on expanding racial and ethnic detail when presenting health data. Trend data on race and ethnicity are

presented in the greatest detail possible after taking into account the quality of the data, the amount of missing data, and the number of observations. These issues significantly affect the availability of reportable data for certain populations, such as the Native Hawaiian or Other Pacific Islander population and the American Indian or Alaska Native population. Standards for the classification of federal data on race and ethnicity are described in an appendix. (See Appendix II, Race.)

Education and Income Data

Many Trend Tables in *Health, United States* present data according to socioeconomic status, using education and family income as proxy measures. Education and income data are generally obtained directly from survey respondents and are not usually available from records-based data collection systems. (See Appendix II, Education; Family income; Poverty.)

Disability Data

Disability can include the presence of physical or mental impairments that limit a person's ability to perform an important activity and affect the use of or need for support, accommodation, or intervention to improve functioning. Information on disability in the U.S. population is critical to health planning and policy. Disability may be measured based on a specific disability or a composite measure designed to identify persons with any of a variety of disabilities. *Health, United States* includes data from the National Health Interview Survey to examine specific types of disability and to create composite disability measures consistent with two of the conceptual components that have been identified in disability models and legislation: basic actions difficulty and complex activity limitation.

Basic actions difficulty captures limitations or difficulties in movement and sensory, emotional, or mental functioning that are associated with a health problem. Complex activity limitation describes limitations or restrictions in a person's ability to participate fully in social role activities such as working or maintaining a household. *Health, United States, 2015* includes the following disability-related information: difficulty remembering and difficulty doing errands alone (Figure 6), basic actions difficulty and complex activity limitation (Table 42), vision and hearing limitations for adults (Tables 43 and 44), and disability-related information for Medicare enrollees (Table 108), Medicaid recipients (Table 109), and veterans with service-connected disabilities (Table 111). For more information on disability statistics, see Altman and Bernstein (2) and Brault (3).

Statistical Significance

All statements in the text describing differences, or lack thereof, in estimates indicate that statistical testing was performed. Differences between two point estimates were determined to be statistically significant at the 0.05 level using two-sided significance tests (z-tests) without correction for multiple comparisons. Data tables include point estimates and standard errors for users who would like to perform additional statistical tests. In the text, the standard terminology used when a difference between two point estimates was tested is, "Between (estimate 1) and (estimate 2)." For example, the statement "Between 2013 and 2014" indicates that the difference between the point estimate for 2013 and that for 2014 was tested for statistical significance.

The statistical significance of a time trend was assessed using weighted least squares regression applied to data for all years in the time period. (For a description of the trend testing technique, see the Technical Notes that follow the Chartbook.) The terminology used in the text to indicate testing of a trend is "During (time period 1)-(time period 2)." For example, the statement "During 2004–2014" indicates that a statistical test of trend was conducted that included estimates for all 11 years in the time period. Because statistically significant differences or trends are partly a function of sample size (i.e., the larger the sample, the smaller the change that can be detected), statistically significant differences or trends do not necessarily have public health significance (4).

Terms such as "similar," "stable," and "no difference" indicate that the statistics being compared were not significantly different. Lack of comment regarding the difference between statistics does not necessarily suggest that the difference was tested and found to not be significant.

Overall estimates generally have relatively small standard errors, but estimates for certain population subgroups may be based on small numbers and have relatively large standard errors. Although numbers of births and deaths from the Vital Statistics System represent complete counts (except for births in those states where data are based on a 50% sample for selected years) and are not subject to sampling error, the counts are subject to random variation, which means that the number of events that actually occur in a given year may be considered as one of a large series of possible results that could have arisen under the same circumstances. When the number of events is small and the probability of such an event is small, considerable caution must be observed in interpreting the conditions described by the estimates. Estimates that are unreliable because of large standard errors or small numbers of events are noted with an asterisk. The criteria used to designate or suppress unreliable estimates are indicated in the table footnotes.

For NCHS surveys, point estimates and their corresponding variances were calculated using the SUDAAN software package (5), which takes into consideration the complex survey design. Standard errors for other surveys or data sets were computed using the methodology recommended by the programs providing the data, or were provided directly by those programs. Standard errors are available for selected tables in the spreadsheet version on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

Accessing *Health, United States*

Health, United States can be accessed in its entirety at: <http://www.cdc.gov/nchs/hus.htm>. The website is a user-friendly resource for *Health, United States* and related products. In addition to the full report, the website contains the *In Brief* companion report in PDF format. Also found on the website are data conveniently organized and grouped by topic. The Chartbook figures are provided as PowerPoint slides, and the Trend Tables and Chartbook data tables are provided as spreadsheet and PDF files. Many spreadsheet files include additional years of data not shown in the printed report, along with standard errors where available. Visitors to the website can join the *Health, United States* e-mail list (http://www.cdc.gov/nchs/hus/hus_electronic_mailing.htm) to receive announcements about release dates and notices of updates. Previous editions of *Health, United States*, and their Chartbooks, can also be accessed from the website.

Printed copies of *Health, United States* can be purchased from the U.S. Government Printing Office at: <http://bookstore.gpo.gov>.

Questions?

If you have questions about *Health, United States* or related data products, please contact:

Office of Information Services
Information Dissemination Staff
National Center for Health Statistics
Centers for Disease Control and Prevention
3311 Toledo Road, Room 5419
Hyattsville, MD 20782–2064
Phone: 1–800–CDC–INFO (1–800–232–4636)
TTY: 1–888–232–6348
Internet: <http://www.cdc.gov/nchs>
Online request form: <http://www.cdc.gov/cdc-info/requestform.html>
For e-mail updates on NCHS publication releases, subscribe online at: <http://www.cdc.gov/nchs/govdelivery.htm>.

References

1. Heckler MM. Report of the Secretary's Task Force on Black and Minority Health volume I, Executive Summary. Washington, DC: HHS; 1985. Washington, DC.
2. Altman B, Bernstein A. Disability and health in the United States, 2001–2005. Hyattsville, MD: NCHS; 2008.
3. Brault MW. Americans with disabilities: 2010. Current population reports, P70–131. Washington, DC: U.S. Census Bureau; 2012.
4. CDC. Youth Risk Behavior Surveillance System (YRBSS). Interpretation of YRBS trend data. Atlanta, GA; 2014.
5. SUDAAN, release 11.0.0 [computer software]. Research Triangle Park, NC: RTI International; 2012.

Acknowledgments

Overall responsibility for planning and coordinating the content of this volume rested with the National Center for Health Statistics' (NCHS) Office of Analysis and Epidemiology, under the direction of Julia S. Holmes and Irma E. Arispe.

Production of *Health, United States, 2015* was managed by Sheila J. Franco, Virginia M. Freid, and Julia S. Holmes. Preparation of the volume, including highlights, trend tables, appendixes, and index, was completed by Mary Ann Bush, La-Tonya D. Curl, Anne K. Driscoll, Catherine R. Duran, Sheila J. Franco, Virginia M. Freid, Nancy Han, Hashini S. Khajuria, Ji-Eun Kim, Florence Lee, Xianfen Li, Anita L. Powell, Ilene B. Rosen, and Ashley M. Woodall. Administrative and word processing assistance was provided by Lillie C. Featherstone.

Production of the *Chartbook* was managed by Sheila J. Franco and Virginia M. Freid. Data, analysis, and text for specific charts were provided by Mary Ann Bush, Juanita J. Chinn, La-Tonya D. Curl, Anne K. Driscoll, Catherine R. Duran, Sheila J. Franco, Virginia M. Freid, Nancy Han, Hashini S. Khajuria, Ji-Eun Kim, Florence Lee, Xianfen Li, Diane M. Makuc, Makram Talih, and Ashley M. Woodall.

Publication production was performed by CDC/OSELS/NCHS/OD/Office of Information Services, Information Design and Publishing Staff (IDPS). Project management was provided by Jen Hurlburt, Elom L. Lawson, and Barbara J. Wassell. Editorial review was provided by Jen Hurlburt and Barbara J. Wassell. Graphic design was provided by Odell D. Eldridge (contractor) and Kyung M. Park. Layout and production were done by Jacqueline M. Davis and Zung T. Le. Overview for IDPS publications and electronic products was provided by Kimberly N. Ross and Tommy C. Seibert, Jr. Printing was managed by Nathanael Brown, CDC/OD/OADC.

Electronic access through the NCHS website was provided by Christine J. Brown, La-Tonya D. Curl, Jacqueline M. Davis, Virginia M. Freid, Jen Hurlburt, Elom L. Lawson, Zung T. Le, Florence Lee, Anthony Lipphardt, Anita L. Powell, Anthony R. Quintana, Ilene B. Rosen, Barbara J. Wassell, and Ashley M. Woodall.

Data and technical assistance were provided by staff of the following NCHS organizations: *Division of Health Care Statistics*: Esther S. Hing, Eric Jamoom, Susan M. Schappert, and Victor V. Shigaev; *Division of Health and Nutrition Examination Surveys*: Margaret D. Carroll, Mark S. Eberhardt, Eleanor B. Fleming, Cheryl Fryar, Qiuping Gu, Brian K. Kit, Cynthia L. Ogden, and Ryne Paulose-Ram; *Division of Health Interview Statistics*: Patricia F. Adams, Veronica Benson, Lindsey Black, Debra Blackwell, Barbara Bloom, Tainya Clarke, Robin A. Cohen, Jacqueline Lucas, Michael Martinez, Colleen Nugent, Jeannine Schiller, Charlotte Schoenborn,

Jessica Simpson, Maria Villarroel, Brian Ward, and Emily Zammitti; *Division of Vital Statistics*: Robert N. Anderson, Joyce A. Arbertha, Elizabeth Arias, Amy M. Branum, Sharon E. Kirmeyer, Kenneth D. Kochanek, Annie Liu, Joyce A. Martin, T.J. Mathews, Arialdi M. Miniño, Jaleh Mousavi, Sherry L. Murphy, Steven J. Steimel, Betzaida Tejada-Vera, and Elizabeth C. Wilson; *Office of Analysis and Epidemiology*: Li-Hui Chen, Catherine R. Duran, Deborah D. Ingram, Mitchell Loeb, Laura A. Pratt, and Cheryl V. Rose; *Office of the Center Director*: Juan Albertorio and Francis C. Notzon; and *Office of Research and Methodology*: Meena Khare.

Additional data and technical assistance were provided by the following organizations of the Centers for Disease Control and Prevention (CDC): *National Center for Chronic Disease Prevention and Health Promotion*: Karen Pazol; *National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention*: Lori Elmore, Anna Satcher Johnson, Laura Kann, Steve Kinchen, Jennifer A. Ludovic, and the Surveillance and Data Management Branch; *Office of Public Health Scientific Services, Center for Surveillance, Epidemiology, and Laboratory Services*: Ruth Ann Jajosky; by the following organizations within the Department of Health and Human Services: *Agency for Healthcare Research and Quality*: Kellyn V. Carper and Anne Elixhauser; *Centers for Medicare & Medicaid Services*: Catherine A. Curtis, Tony Dean, Maria Diacogiannis, Nathan Espinosa, Deborah W. Kidd, Barbara S. Klees, Lisa M. Mirel, Maggie S. Murgolo, Debra Reed-Gillette, Joseph F. Regan, Jeffrey S. Silverman, Christopher J. Truffer, and Lekha S. Whittle; *National Institutes of Health*: Nadia Howlader and Marsha Lopez; *Substance Abuse and Mental Health Services Administration*: Jonaki Bose, Kathy Piscopo, Neil Russell, and Albert M. Woodward; and by the following governmental and nongovernmental organizations: *U.S. Census Bureau*: Bernadette D. Proctor, Danielle Taylor, and Monica Wiedemann; *Bureau of Labor Statistics*: Elizabeth Ashack, Christen Byler, and Audrey Watson; *Department of Veterans Affairs*: Tom Garin and Dat Tran; *American Association of Colleges of Pharmacy*: Nancy T. Nguyen, Danielle A. Taylor, and Jamie N. Taylor; *American Association of Colleges of Osteopathic Medicine*: Linsey Jurd; *American Association of Colleges of Podiatric Medicine*: Moraith G. North; *American Dental Education Association*: Adriana R. Menezes and Bradley Munson; *American Medical Association Physician Masterfile*: Cheryl Ashe; *Association of American Medical Colleges*: Briana Gunter and Lindsay Roskovensky; *Association of Schools and Colleges of Optometry*: Paige Pence and Joanne Zuckerman; *Association of Schools & Programs of Public Health*: Emily M. Burke and Christine M. Plepys; *Cowles Research Group*: C. McKeen Cowles; and *NOVA Research Company*: Shilpa Bengeri.

Contents

Preface	iii	Functional Limitations	13
Acknowledgments	vi	Disability	13
List of Chartbook Figures	xi	Health Risk Factors	13
List of Trend Tables	xii	Current Cigarette Smoking	13
		Children and Adolescents with Obesity	14
		Adults with Overweight and Obesity	14
		Utilization	15
		Pap Test Use	15
		Emergency Department Use	15
		Difficulty Accessing Needed Medical Care or Prescription Drugs Due to Cost	16
		Health Care Resources	16
		Electronic Health Record Systems	16
		Physicians Accepting New Patients	17
		Personal Health Care Expenditures	17
		Major Source of Funds	17
		Health Insurance	18
		Coverage Among Adults Aged 18–64	18
		Coverage by Medicaid Expansion State	18
		Special Feature on Racial and Ethnic Health Disparities	
		Introduction	20
		Life Expectancy at Birth	22
		Infant Mortality	23
		Preterm Births	24
		Low-risk Births Delivered by Cesarean Section	25
		Children and Adolescents With Obesity	26
		Hypertension	27
		Current Cigarette Smoking	28
		Influenza Vaccination	29
		Health Insurance Coverage	30
		Difficulty Accessing Needed Dental Care Due to Cost	31
		Chartbook Data Tables	32
		Technical Notes	57
		References	59
At a Glance Table and Highlights			
At a Glance Table	2		
Highlights	4		
Health Status and Determinants	4		
Life Expectancy and Mortality	4		
Fertility and Natality	4		
Health Risk Factors for the Noninstitutionalized Population	5		
Measures of Health and Disease Prevalence for the Noninstitutionalized Population	5		
Utilization of Health Resources for the Noninstitutionalized Population	5		
Use of Health Care Services	5		
Use of Preventive Medical Care Services for the Noninstitutionalized Population	5		
Difficulty Accessing Needed Medical Care, Prescription Drugs, and Dental Care Due to Cost for the Noninstitutionalized Population	6		
Health Care Resources	6		
Health Care Expenditures and Payers	6		
Health Care Expenditures	6		
Health Care Payers	6		
Health Insurance Coverage for the Noninstitutionalized Population	7		
Special Feature on Racial and Ethnic Health Disparities	7		
Chartbook With Special Feature on Racial and Ethnic Health Disparities			
Mortality	10		
Life Expectancy at Birth, by Country	10		
Selected Causes of Death	11		
Suicide and Homicide	11		
Natality	12		
Teenage Childbearing	12		
Morbidity	12		
Notifiable Disease Rates	12		

Trend Tables

Health Status and Determinants	65
Population	65
Fertility and Natality	70
Mortality	86
Determinants and Measures of Health	152
Utilization of Health Resources	226
Ambulatory Care	226
Inpatient Care	277
Health Care Resources	282
Personnel	282
Facilities	288
Health Care Expenditures and Payers	293
National Health Expenditures	293
Health Care Coverage and Major Federal Programs	313
State Health Expenditures and Health Insurance	336

Appendixes

Appendix Contents	345
Appendix I. Sources of Data	349
Appendix II. Definitions and Methods	384

Index

Index	439
-----------------	-----

List of Chartbook Figures

Mortality

- Figure 1. Life expectancy at birth, by sex and country: Organisation for Economic Co-operation and Development (OECD) countries, 2013 10
- Figure 2. Age-adjusted death rates for selected causes of death for all ages, by sex: United States, 2004–2014 11
- Figure 3. Suicide and homicide death rates among persons aged 15 and over, by age and sex: United States, 2014 11

Natality

- Figure 4. Teenage childbearing, by maternal age and race and Hispanic origin: United States, 2004–2014 12

Morbidity

- Figure 5. Selected notifiable disease rates: United States, 2003 and 2013 12

Functional Limitations

- Figure 6. Selected disability indicators among adults aged 18 and over, by sex and age: United States, 2014 13

Health Risk Factors

- Figure 7. Current cigarette smoking among adults aged 18 and over, by sex and age: United States, 2004–2014 13
- Figure 8. Obesity among children and adolescents aged 2–19 years, by age: United States, 1999–2002 through 2011–2014 14
- Figure 9. Overweight and obesity among adults aged 20 and over, by sex and grade of obesity: United States, 1999–2002 through 2011–2014 14

Utilization

- Figure 10. Pap test utilization within the past 3 years, by age: United States, 2003–2013 15
- Figure 11. Emergency department utilization within the past 12 months among adults aged 18–64, by type of coverage: United States, 2004–2014 15
- Figure 12. Delay or nonreceipt of needed medical care and nonreceipt of needed prescription drugs in the past 12 months due to cost among adults aged 18–64, by health insurance coverage: United States, 2004–2014 16

Health Care Resources

- Figure 13. Electronic health record system components in physician offices, by selected component type: United States, 2010 and 2013 16
- Figure 14. Office-based physicians accepting new patients, by patient source of payment and urban-rural status: United States, 2013. 17

Personal Health Care Expenditures

- Figure 15. Personal health care expenditures, by source of funds: United States, 2004–2014. 17

Health Insurance

- Figure 16. Health insurance coverage among adults aged 18–64, by type of coverage: United States, 2004–June 2015 (preliminary data). 18
- Figure 17. Health insurance coverage among adults aged 18–64, by state Medicaid expansion status: United States, 2013 and 2014 18

Special Feature on Racial and Ethnic Health Disparities

- Figure 18. Life expectancy at birth, by sex, race and Hispanic origin: United States, 1980–2014 22
- Figure 19. Infant mortality rates, by race and Hispanic origin and detailed Hispanic origin of mother: United States, 1999–2013 23
- Figure 20. Preterm births, by gestational age and race and Hispanic origin and detailed Hispanic origin of mother: United States, 2014 24
- Figure 21. Low-risk births delivered by cesarean section, by race and Hispanic origin and detailed Hispanic origin of mother: United States, 1999–2014 25
- Figure 22. Obesity among children and adolescents aged 2–19 years, by age and race and Hispanic origin: United States, 2011–2014 26
- Figure 23. Hypertension among adults aged 20 and over, by sex and race and Hispanic origin: United States, 1999–2000 through 2013–2014 27
- Figure 24. Current cigarette smoking among adults aged 18 and over, by sex and race and Hispanic origin: United States, 1999–2014 28
- Figure 25. Influenza vaccination among adults aged 18 and over, by age and race and Hispanic origin: United States, 1999–2014 29
- Figure 26. No health insurance coverage among adults aged 18–64, by race and Hispanic origin: United States, 1999–June 2015 (preliminary data). 30
- Figure 27. Nonreceipt of needed dental care in the past 12 months due to cost among adults aged 18–64, by race and Hispanic origin: United States, 1999–2014 31

Summary List of Trend Tables by Topic

Tables 1–114

Population (Tables 1 and 2)

Resident population
Persons in poverty

Fertility and Natality (Tables 3–9)

Births
Low birthweight
Breastfeeding
and more . . .

Mortality (Tables 10–32)

Infant mortality
Life expectancy
Death rates, by cause
and more . . .

Determinants and Measures of Health (Tables 33–60)

Health status
Cigarette smoking
Alcohol consumption
High blood pressure
Overweight and obesity
and more . . .

Ambulatory Care (Tables 61–80)

Visits: health care, dentists, emergency departments
and more . . .
Prevention: mammograms, pap smears, vaccinations

Inpatient Care (Tables 81–82)

Hospital stays and procedures
Nursing homes
and more . . .

Personnel (Tables 83–88)

Physicians
Dentists
Nurses
Health professions school enrollment
and more . . .

Facilities (Tables 89–92)

Hospitals
Nursing homes
and more . . .

National Health Expenditures (Tables 93–101)

Personal health expenditures
Out-of-pocket costs
Prescription drug expenditures
Nursing home costs
and more . . .

Health Care Coverage and Major Federal Programs (Tables 102–111)

Insurance coverage
 Medicare
 Medicaid
 Private coverage
 Uninsured
and more . . .

State Health Expenditures and Health Insurance (Tables 112–114)

Medicare, Medicaid, managed care expenditures and enrollees
Uninsured persons

List of Trend Tables

Health Status and Determinants

Population

Table 1. Resident population, by age, sex, race, and Hispanic origin: United States, selected years 1950–2014 65

Table 2. Persons below poverty level, by selected characteristics, race, and Hispanic origin: United States, selected years 1973–2014 68

Fertility and Natality

Table 3. Crude birth rates, fertility rates, and birth rates, by age, race, and Hispanic origin of mother: United States, selected years 1950–2014 70

Table 4. Nonmarital childbearing, by detailed race and Hispanic origin of mother, and maternal age: United States, selected years 1970–2014 73

Table 5. Low birthweight live births, by detailed race and Hispanic origin of mother: United States, selected years 1970–2014 74

Table 6. Low birthweight live births, by race and Hispanic origin of mother, state, and territory: United States and U.S. dependent areas, 2000–2002, 2003–2005, and 2012–2014 75

Table 7. Legal abortions, legal abortion rates, and legal abortion ratios: United States and 47 continuous reporting areas, 2003–2012. 78

Table 8. Contraceptive use in the past month among women aged 15–44, by age, race and Hispanic origin, and method of contraception: United States, selected years 1982–2013 79

Table 9. Breastfeeding among mothers aged 15–44, by year of baby's birth and selected characteristics of mother: United States, average annual 1986–1988 through 2008–2010 85

Mortality

Table 10. Infant, neonatal, postneonatal, fetal, and perinatal mortality rates, by detailed race and Hispanic origin of mother: United States, selected years 1983–2013 86

Table 11. Infant mortality rates, by race: United States, selected years 1950–2014 88

Table 12. Infant mortality rates, by race and Hispanic origin of mother, state, and territory: United States and U.S. dependent areas, average annual 1989–1991, 2003–2005, and 2011–2013 89

Table 13. Infant mortality rates and international rankings: Organisation for Economic Co-operation and Development (OECD) countries, selected years 1960–2012 92

Table 14. Life expectancy at birth and at age 65, by sex: Organisation for Economic Co-operation and Development (OECD) countries, selected years 1980–2013 93

Table 15. Life expectancy at birth, at age 65, and at age 75, by sex, race, and Hispanic origin: United States, selected years 1900–2014 95

Table 16. Age-adjusted death rates, by race, Hispanic origin, state, and territory: United States and U.S. dependent areas, average annual 1979–1981, 1989–1991, and 2012–2014 97

Table 17. Age-adjusted death rates for selected causes of death, by sex, race, and Hispanic origin: United States, selected years 1950–2014 99

Table 18. Years of potential life lost before age 75 for selected causes of death, by sex, race, and Hispanic origin: United States, selected years 1980–2014 103

Table 19. Leading causes of death and numbers of deaths, by sex, race, and Hispanic origin: United States, 1980 and 2014 107

Table 20. Leading causes of death and numbers of deaths, by age: United States, 1980 and 2014 111

Table 21. Death rates for all causes, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014 113

Table 22. Death rates for diseases of heart, by sex, race, Hispanic origin, and age: United States selected years 1950–2014 117

Table 23. Death rates for cerebrovascular diseases, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014 120

Table 24. Death rates for malignant neoplasms, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014 123

Table 25. Death rates for malignant neoplasms of trachea, bronchus, and lung, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014 127

Table 26. Death rates for malignant neoplasm of breast among females, by race, Hispanic origin, and age: United States, selected years 1950–2014 130

Table 27. Death rates for drug poisoning and drug poisoning involving opioid analgesics and heroin, by sex, age, race, and Hispanic origin: United States, selected years 1999–2014 132

Table 28. Death rates for motor vehicle-related injuries, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014 136

Table 29. Death rates for homicide, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014 140

Table 30. Death rates for suicide, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014 144

Table 31. Death rates for firearm-related injuries, by sex, race, Hispanic origin, and age: United States, selected years 1970–2014 147

Table 32. Occupational fatal injuries, by industry, sex, age, race, and Hispanic origin: United States, selected years 2003–2013 150

Determinants and Measures of Health

Table 33. Selected nationally notifiable disease rates and number of new cases: United States, selected years 1950–2013	152	Table 47. Current cigarette smoking among adults aged 18 and over, by sex, race, and age: United States: selected years 1965–2014	186
Table 34. Human immunodeficiency virus (HIV) diagnoses, by year of diagnosis and selected characteristics: United States, 2010–2014	154	Table 48. Age-adjusted prevalence of current cigarette smoking among adults aged 25 and over, by sex, race, and education level: United States, selected years 1974–2014	188
Table 35. Health conditions among children under age 18, by selected characteristics: United States, average annual, selected years 1997–1999 through 2012–2014	156	Table 49. Current cigarette smoking among adults aged 18 and over, by sex, race, Hispanic origin, age, and education level: United States, average annual, selected years 1990–1992 through 2012–2014	189
Table 36. Age-adjusted cancer incidence rates for selected cancer sites, by sex, race, and Hispanic origin: United States, selected geographic areas, selected years 1990–2012	161	Table 50. Use of selected substances in the past month among persons aged 12 and over, by age, sex, race, and Hispanic origin: United States, selected years 2002–2014	192
Table 37. Five-year relative cancer survival rates for selected cancer sites, by race and sex: United States, selected geographic areas, selected years 1975–1977 through 2005–2011	165	Table 51. Use of selected substances in the past 30 days among 12th graders, 10th graders, and 8th graders, by sex and race: United States, selected years 1980–2014	194
Table 38. Respondent-reported prevalence of heart disease, cancer, and stroke among adults aged 18 and over, by selected characteristics: United States, average annual, selected years 1997–1998 through 2013–2014	166	Table 52. Health risk behaviors among students in grades 9–12, by sex, grade level, race, and Hispanic origin: United States, selected years 1991–2013	197
Table 39. Number of respondent-reported chronic conditions from 10 selected conditions among adults aged 18 and over, by selected characteristics: United States, selected years 2002–2014	168	Table 53. Selected health conditions and risk factors, by age: United States, selected years 1988–1994 through 2013–2014	200
Table 40. Diabetes prevalence and glycemic control among adults aged 20 and over, by sex, age, and race and Hispanic origin: United States, selected years 1988–1994 through 2011–2014	170	Table 54. Hypertension among adults aged 20 and over, by selected characteristics: United States, selected years 1988–1994 through 2011–2014	202
Table 41. Severe headache or migraine, low back pain, and neck pain among adults aged 18 and over, by selected characteristics: United States, selected years 1997–2014	173	Table 55. Cholesterol among adults aged 20 and over, by selected characteristics: United States, selected years 1988–1994 through 2011–2014	204
Table 42. Disability measures among adults aged 18 and over, by selected characteristics: United States, selected years 1997–2014	176	Table 56. Mean macronutrient intake among adults aged 20 and over, by sex and age: United States, selected years 1988–1994 through 2009–2012	208
Table 43. Vision limitations among adults aged 18 and over, by selected characteristics: United States, selected years 1997–2014	178	Table 57. Participation in leisure-time aerobic and muscle-strengthening activities that meet the federal 2008 <i>Physical Activity Guidelines for Americans</i> among adults aged 18 and over, by selected characteristics: United States, selected years 1998–2014	210
Table 44. Hearing limitations among adults aged 18 and over, by selected characteristics: United States, selected years 2007–2014	180	Table 58. Normal weight, overweight, and obesity among adults aged 20 and over, by selected characteristics: United States, selected years 1988–1994 through 2011–2014	215
Table 45. Respondent-assessed fair-poor health status, by selected characteristics: United States, selected years 1991–2014	182	Table 59. Obesity among children and adolescents aged 2–19 years, by selected characteristics: United States, selected years 1988–1994 through 2011–2014	222
Table 46. Serious psychological distress in the past 30 days among adults aged 18 and over, by selected characteristics: United States, average annual, selected years 1997–1998 through 2013–2014	184	Table 60. Untreated dental caries, by selected characteristics: United States, selected years 1988–1994 through 2011–2012	224

Utilization of Health Resources

Ambulatory Care

Table 61. No usual source of health care among children under age 18, by selected characteristics: United States, average annual, selected years 1993–1994 through 2013–2014	226
Table 62. No usual source of health care among adults aged 18–64, by selected characteristics: United States, average annual, selected years 1993–1994 through 2013–2014	228
Table 63. Delay or nonreceipt of needed medical care, nonreceipt of needed prescription drugs, or nonreceipt of needed dental care during the past 12 months due to cost, by selected characteristics: United States, selected years 1997–2014	230
Table 64. No health care visits to an office or clinic within the past 12 months among children under age 18, by selected characteristics: United States, average annual, selected years 1997–1998 through 2013–2014	233
Table 65. Health care visits to doctor offices, emergency departments, and home visits within the past 12 months, by selected characteristics: United States, selected years 1997–2014	235
Table 66. Vaccination coverage for selected diseases among children aged 19–35 months, by race, Hispanic origin, poverty level, and location of residence in metropolitan statistical area: United States, selected years 1998–2014	238
Table 67. Vaccination coverage for selected diseases among adolescents aged 13–17, by selected characteristics: United States, 2008–2014	241
Table 68. Influenza vaccination among adults aged 18 and over, by selected characteristics: United States, selected years 1989–2014	242
Table 69. Pneumococcal vaccination among adults aged 18 and over, by selected characteristics: United States, selected years 1989–2014	244
Table 70. Use of mammography among women aged 40 and over, by selected characteristics: United States, selected years 1987–2013	246
Table 71. Use of Pap smears among women aged 18 and over, by selected characteristics: United States, selected years 1987–2013	249
Table 72. Use of colorectal tests or procedures among adults aged 50–75, by selected characteristics: United States, selected years 2000–2013	254
Table 73. Emergency department visits within the past 12 months among children under age 18, by selected characteristics: United States, selected years 1997–2014	256
Table 74. Emergency department visits within the past 12 months among adults aged 18 and over, by selected characteristics: United States, selected years 1997–2014	260

Table 75. Initial injury-related visits to hospital emergency departments, by sex, age, and intent and mechanism of injury: United States, average annual, selected years 2005–2006 through 2010–2011	263
Table 76. Visits to physician offices, hospital outpatient departments, and hospital emergency departments, by age, sex, and race: United States, selected years 2000–2012	265
Table 77. Visits to primary care generalist and specialty care physicians, by selected characteristics and type of physician: United States, selected years 1980–2012	268
Table 78. Dental visits in the past year, by selected characteristics: United States, selected years 1997–2014	270
Table 79. Prescription drug use in the past 30 days, by sex, race and Hispanic origin, and age: United States, selected years 1988–1994 through 2009–2012	272
Table 80. Selected prescription drug classes used in the past 30 days, by sex and age: United States, selected years 1988–1994 through 2009–2012	274

Inpatient Care

Table 81. Persons with hospital stays in the past year, by selected characteristics: United States, selected years 1997–2014	277
Table 82. Hospital admissions, average length of stay, outpatient visits, and outpatient surgery, by type of ownership and size of hospital: United States, selected years 1975–2013	281

Health Care Resources

Personnel

Table 83. Active physicians and physicians in patient care, by state: United States, selected years 1975–2013	282
Table 84. Doctors of medicine, by place of medical education and activity: United States and outlying U.S. areas, selected years 1975–2013	283
Table 85. Doctors of medicine in primary care, by specialty: United States and outlying U.S. areas, selected years 1949–2013	284
Table 86. Active dentists, by state: United States, selected years 2001–2013	285
Table 87. Healthcare employment and wages, by selected occupations: United States, selected years 2000–2014	286
Table 88. First-year enrollment and graduates of health professions schools, and number of schools, by selected profession: United States, selected academic years 1980–1981 through 2013–2014	287

Facilities

Table 89. Hospitals, beds, and occupancy rates, by type of ownership and size of hospital: United States, selected years 1975–2013	288
Table 90. Community hospital beds and average annual percent change, by state: United States, selected years 1980–2013	289
Table 91. Occupancy rates in community hospitals and average annual percent change, by state: United States, selected years 1980–2013	290
Table 92. Nursing homes, beds, residents, and occupancy rates, by state: United States, selected years 1995–2014	291

Health Care Expenditures and Payers

National Health Expenditures

Table 93. Gross domestic product, national health expenditures, per capita amounts, percent distribution, and average annual percent change: United States, selected years 1960–2014	293
Table 94. National health expenditures, average annual percent change, and percent distribution, by type of expenditure: United States, selected years 1960–2014	295
Table 95. Personal health care expenditures, by source of funds and type of expenditure: United States, selected years 1960–2014	297
Table 96. Cost of hospital discharges with common hospital operating room procedures in nonfederal community hospitals, by age and selected principal procedure: United States, selected years 2000–2013	300
Table 97. Expenses for health care and prescribed medicine, by selected population characteristics: United States, selected years 1987–2012	303
Table 98. Sources of payment for health care, by selected population characteristics: United States, selected years 1987–2012	306
Table 99. Out-of-pocket health care expenses among persons with medical expenses, by age: United States, selected years 1987–2012	309
Table 100. National health expenditures and percent distribution, by sponsor: United States, selected years 1987–2014	310
Table 101. Employers' costs per employee-hour worked for total compensation, wages and salaries, and health insurance, by selected characteristics: United States, selected years 1999–2015	312

Health Care Coverage and Major Federal Programs

Table 102. Private health insurance coverage among persons under age 65, by selected characteristics: United States, selected years 1984–2014	313
Table 103. Private health insurance coverage obtained through the workplace among persons under age 65, by selected characteristics: United States, selected years 1984–2014	316
Table 104. Medicaid coverage among persons under age 65, by selected characteristics: United States, selected years 1984–2014	319
Table 105. No health insurance coverage among persons under age 65, by selected characteristics: United States, selected years 1984–2014	322
Table 106. Health insurance coverage of noninstitutionalized Medicare beneficiaries aged 65 and over, by type of coverage and selected characteristics: United States, selected years 1992–2013	325
Table 107. Medicare enrollees and expenditures and percent distribution, by Medicare program and type of service: United States and other areas, selected years 1970–2014	327
Table 108. Medicare beneficiaries, by race, Hispanic origin, and selected characteristics: United States, selected years 1992–2012	329
Table 109. Medicaid and Children's Health Insurance Program beneficiaries and payments, by basis of eligibility, and race and Hispanic origin: United States, selected fiscal years 1999–2012	331
Table 110. Medicaid and Children's Health Insurance Program beneficiaries and payments, by type of service: United States, selected fiscal years 1999–2012	333
Table 111. Department of Veterans Affairs health care expenditures and use, and persons treated, by selected characteristics: United States, selected fiscal years 2005–2014	335

State Health Expenditures and Health Insurance

Table 112. Medicare enrollees, enrollees in managed care, payment per fee-for-service enrollee, and short-stay hospital utilization, by state: United States, 1994 and 2014	336
Table 113. Medicaid and Children's Health Insurance Program beneficiaries and payments per beneficiary, by state: United States, selected fiscal years 2000–2012	338
Table 114. Persons under age 65 without health insurance coverage, by age, state, and territory: United States and Puerto Rico, 2009–2014	339

Health, United States, 2015: At a Glance

	Value (year)			Health, United States, 2015 Table No.
Life Expectancy and Mortality				
Life expectancy, in years				Table 15
At birth	76.8 (2000)	78.8 (2013)	78.8 (2014)	
Infant deaths per 1,000 live births				Table 11
All infants	6.91 (2000)	5.96 (2013)	5.82 (2014)	
Deaths per 100,000 population, age-adjusted				Table 17
All causes	869.0 (2000)	731.9 (2013)	724.6 (2014)	
Heart disease	257.6 (2000)	169.8 (2013)	167.0 (2014)	
Cancer	199.6 (2000)	163.2 (2013)	161.2 (2014)	
Chronic lower respiratory diseases	44.2 (2000)	42.1 (2013)	40.5 (2014)	
Unintentional injuries	34.9 (2000)	39.4 (2013)	40.5 (2014)	
Stroke	60.9 (2000)	36.2 (2013)	36.5 (2014)	
Alzheimer's disease	18.1 (2000)	23.5 (2013)	25.4 (2014)	
Diabetes	25.0 (2000)	21.2 (2013)	20.9 (2014)	
Influenza and pneumonia	23.7 (2000)	15.9 (2013)	15.1 (2014)	
Nephritis, nephrotic syndrome and nephrosis	13.5 (2000)	13.2 (2013)	13.2 (2014)	
Suicide	10.4 (2000)	12.6 (2013)	13.0 (2014)	
Morbidity and Risk Factors				
Fair or poor health, percent				Table 45
All ages	8.9 (2000)	10.2 (2013)	9.8 (2014)	
65 years and over	26.9 (2000)	23.1 (2013)	21.7 (2014)	
Heart disease (ever told), percent				Table 38
18 years and over	11.3 (2000–2001)	11.4 (2011–2012)	11.5 (2013–2014)	
65 years and over	30.9 (2000–2001)	30.3 (2011–2012)	29.4 (2013–2014)	
Cancer (ever told), percent				Table 38
18 years and over	5.0 (2000–2001)	6.2 (2011–2012)	6.4 (2013–2014)	
65 years and over	15.2 (2000–2001)	18.5 (2011–2012)	18.2 (2013–2014)	
Hypertension, ¹ percent				Table 54
20 years and over	30.2 (1999–2002)	32.2 (2007–2010)	33.0 (2011–2014)	
Diabetes, ² percent				Table 40
20 years and over	9.8 (1999–2002)	12.0 (2007–2010)	12.6 (2011–2014)	
Hypercholesterolemia, ³ percent				Table 55
20 years and over	25.0 (1999–2002)	28.7 (2007–2010)	29.8 (2011–2014)	
Obese, percent				Tables 58 and 59
Obese, ⁴ 20 years and over	30.5 (1999–2002)	34.9 (2007–2010)	36.5 (2011–2014)	
Obese (BMI at or above sex- and age-specific 95th percentile):				
2–5 years	10.3 (1999–2002)	11.1 (2007–2010)	8.9 (2011–2014)	
6–11 years	15.9 (1999–2002)	18.8 (2007–2010)	17.5 (2011–2014)	
12–19 years	16.0 (1999–2002)	18.2 (2007–2010)	20.5 (2011–2014)	
Cigarette smoking, percent				Table 47
18 years and over	23.2 (2000)	17.8 (2013)	16.8 (2014)	
Aerobic activity and muscle strengthening, ⁵ percent meeting both guidelines				Table 57
18 years and over	15.1 (2000)	20.4 (2013)	20.9 (2014)	

¹Having measured high blood pressure (systolic pressure of at least 140 mm Hg or diastolic pressure of at least 90 mm Hg) and/or respondent report of taking antihypertensive medication.

²Includes physician-diagnosed and undiagnosed diabetes (fasting plasma glucose of at least 126 mg/dL or a hemoglobin A1c of at least 6.5%).

³Having high serum total cholesterol of 240 mg/dL or greater and/or respondent report of taking cholesterol-lowering medication.

⁴Obesity is a body mass index (BMI) greater than or equal to 30 for adults. Height and weight are measured rather than self-reported.

⁵Federal guidelines recommend at least 150 minutes of moderate-intensity or 75 minutes of vigorous-intensity aerobic physical activity a week and muscle-strengthening activities at least twice a week.

Health, United States, 2015: At a Glance

	Value (year)			Health, United States, 2015 Table No.
Health Care Utilization				
No health care visit in past 12 months, percent				Table 65
Under 18 years	12.3 (2000)	8.2 (2013)	7.9 (2014)	
18–44 years	23.4 (2000)	24.8 (2013)	23.2 (2014)	
45–64 years	14.9 (2000)	15.2 (2013)	15.0 (2014)	
65 years and over	7.4 (2000)	6.4 (2013)	5.6 (2014)	
Emergency room visit in past 12 months, percent				Tables 73 and 74
Under 18 years	20.3 (2000)	17.6 (2013)	16.7 (2014)	
18–44 years	20.5 (2000)	18.5 (2013)	18.4 (2014)	
45–64 years	17.6 (2000)	17.6 (2013)	17.5 (2014)	
65 years and over	23.7 (2000)	21.3 (2013)	21.2 (2014)	
Dental visit in past year, percent				Table 78
2–17 years	74.1 (2000)	83.0 (2013)	83.0 (2014)	
18–64 years	65.1 (2000)	61.7 (2013)	62.0 (2014)	
65 years and over	56.6 (2000)	60.6 (2013)	62.4 (2014)	
Prescription drug in past 30 days, percent				Table 79
Under 18 years	23.8 (1999–2002)	24.7 (2003–2006)	23.5 (2009–2012)	
18–44 years	35.9 (1999–2002)	37.4 (2003–2006)	38.1 (2009–2012)	
45–64 years	64.1 (1999–2002)	65.2 (2003–2006)	67.2 (2009–2012)	
65 years and over	84.7 (1999–2002)	89.4 (2003–2006)	89.8 (2009–2012)	
Hospitalization in past year, percent				Table 81
18–44 years	7.0 (2000)	6.1 (2013)	5.8 (2014)	
45–64 years	8.4 (2000)	7.8 (2013)	7.4 (2014)	
65 years and over	18.2 (2000)	15.3 (2013)	15.3 (2014)	
Health Insurance and Access to Care				
Uninsured, percent				Table 105
Under 65 years	17.0 (2000)	16.7 (2013)	13.3 (2014)	
Under 18 years	12.6 (2000)	6.6 (2013)	5.4 (2014)	
18–44 years	22.4 (2000)	24.2 (2013)	19.7 (2014)	
45–64 years	12.6 (2000)	15.4 (2013)	11.8 (2014)	
Delay or nonreceipt of needed medical care in past 12 months due to cost, percent				Table 63
Under 18 years	4.6 (2000)	3.1 (2013)	2.8 (2014)	
18–44 years	9.5 (2000)	11.9 (2013)	10.7 (2014)	
45–64 years	8.8 (2000)	13.2 (2013)	11.7 (2014)	
65 years and over	4.5 (2000)	4.2 (2013)	4.3 (2014)	
Health Care Resources				
Patient care physicians per 10,000 population ⁶				Table 83
United States	22.7 (2000)	26.9 (2012)	27.6 (2013)	
Highest state	54.5 (DC) (2000)	65.9 (DC) (2012)	66.1 (DC) (2013)	
Lowest state	14.4 (ID) (2000)	18.0 (ID,MS) (2012)	18.6 (ID) (2013)	
Community hospital beds per 1,000 population ⁷				Table 90
United States	2.9 (2000)	2.6 (2012)	2.5 (2013)	
Highest state	6.0 (ND) (2000)	5.7 (DC) (2012)	5.6 (DC) (2013)	
Lowest state	1.9 (NM,NV,OR,UT,WA) (2000)	1.7 (OR) (2012)	1.7 (OR,WA) (2013)	
Health Care Expenditures				
Personal health care expenditures, in dollars				Table 95
Total, in trillions	\$1.2 (2000)	\$2.4 (2013)	\$2.6 (2014)	
Per capita	\$4,121 (2000)	\$7,727 (2013)	\$8,054 (2014)	

⁶Copyright 2015. Used with permission of the American Medical Association.

⁷Copyright 2015. Used with permission of Health Forum LLC, an affiliate of the American Hospital Association.

NOTES: Estimates in this table are taken from the PDF, printed, or spreadsheet version of the cited tables. For more information and the spreadsheet version of the tables, see the *Health, United States* website: <http://www.cdc.gov/nchs/hus.htm>.

Highlights

This Highlights section presents selected data from the four major areas included in the report: health status and determinants, utilization of health resources, health care resources, and health care expenditures and payers, and from this year's Special Feature on racial and ethnic health disparities. The Highlights focus on topics of public health importance and illustrate the breadth of material included in *Health, United States, 2015*. The Highlights section generally presents trends for the recent 10-year period or examines information for the most recent data year available. Highlights from the 2015 Special Feature generally present data from 1999 to the most recent year available, or only data from the latest year (see [Technical Notes](#) for additional information). Each highlight includes a reference to the detailed trend table or figure where definitions of terms and additional data can be obtained.

Health Status and Determinants

Life Expectancy and Mortality

In 2014, life expectancy at birth in the United States for the total population was 78.8 years—76.4 years for males and 81.2 years for females ([Table 15](#)).

Between 2004 and 2014, life expectancy at birth increased 1.4 years for males and 1.1 years for females. The gap in life expectancy between males and females narrowed from 5.1 years in 2004 to 4.8 years in 2014 ([Table 15](#)).

Between 2004 and 2014, life expectancy at birth increased more for the black than for the white population, thereby narrowing the gap in life expectancy between these two racial groups. In 2004, life expectancy at birth for the white population was 5.2 years longer than for the black population; by 2014, the difference had narrowed to 3.4 years ([Table 15](#)).

Between 2013 and 2014, life expectancy at birth increased 0.2 years to 81.8 years for Hispanic persons, decreased 0.1 years to 78.8 years for non-Hispanic white persons, and increased 0.1 years to 75.2 years for non-Hispanic black persons. For males between 2013 and 2014, life expectancy at birth increased 0.1 years to 79.2 years for Hispanic males, remained stable at 76.5 years for non-Hispanic white males, and increased 0.2 years to 72.0 years for non-Hispanic black males. For females between 2013 and 2014, life expectancy at birth increased 0.2 years to 84.0 years for Hispanic females, decreased 0.1 years to 81.1 years for non-Hispanic white females, and remained stable at 78.1 years for non-Hispanic black females ([Table 15](#)).

Between 2004 and 2014, the death rate for black men aged 45–54 decreased 28%, from 933.3 to 671.8 deaths per

100,000 resident population, while the death rate for non-Hispanic white men remained stable (511.2 in 2014). Between 2004 and 2014, the death rate for black women aged 45–54 decreased 18%, from 558.9 to 455.8, while the death rate for non-Hispanic white women increased 11%, from 293.4 to 325.5 ([Table 21](#)).

Between 2004 and 2014, the infant mortality rate decreased 14%, from 6.79 to 5.82 deaths per 1,000 live births and the neonatal mortality rate (among infants under age 28 days) decreased 13%, from 4.52 to 3.94. Between 2004 and 2014, the postneonatal mortality rate (among infants aged 28 days through 11 months) decreased 17%, from 2.27 to 1.88 ([Table 11](#)).

In 2014, the 10 leading causes of death were heart disease, cancer, chronic lower respiratory diseases, unintentional injuries, stroke, Alzheimer's disease, diabetes, influenza and pneumonia, kidney disease, and suicide. These 10 causes of death accounted for 74% of the 2.6 million deaths in 2014 ([Table 19](#)).

Between 2004 and 2014, the age-adjusted heart disease death rate decreased 25%, from 221.6 to 167.0 deaths per 100,000 resident population. In 2014, 23% of all deaths in the United States were from heart disease ([Tables 19 and 22](#)).

Between 2004 and 2014, the age-adjusted cancer death rate decreased 14%, from 186.8 to 161.2 deaths per 100,000 resident population. In 2014, 23% of all deaths in the United States were from cancer ([Tables 19 and 24](#)).

Between 2004 and 2014, the suicide death rate increased 21%, from 11.1 to 13.4 deaths per 100,000 resident population. Among adults aged 45–64, suicide death rates increased 27% between 2004 and 2014 ([Table 30](#)).

Between 2004 and 2014, the drug poisoning death rate involving heroin increased more than five times, from 0.6 to 3.3 deaths per 100,000 resident population. In 2014, the drug poisoning death rate involving heroin was highest among those aged 25–34 (8.0), followed by those aged 35–44 (5.9), and those aged 45–54 (4.7) ([Table 27](#)).

Fertility and Natality

Between 2004 and 2014, the birth rate among teenagers aged 15–19 fell 40%, from 40.5 to 24.2 live births per 1,000 females—a record low for the United States ([Table 3](#)).

In 2014, 8.00% of infants were low-birthweight (weighing less than 2,500 grams [5.5 pounds] at birth); low-birthweight was more common among non-Hispanic black infants (13.17%) and Puerto Rican infants (9.54%) than among infants in other racial and ethnic groups ([Table 5](#)).

Health Risk Factors for the Noninstitutionalized Population

Children

In 2011–2014, the prevalence of children with obesity among those aged 2–5 years was 8.9%, 17.5% among children aged 6–11, and 20.5% among adolescents aged 12–19 (Table 59 and Figure 8).

In 2014, 4.9% of adolescents aged 12–17 reported smoking cigarettes in the past month. Smoking prevalence has declined since 2004, when 11.9% of adolescents reported smoking cigarettes in the past month (Table 50).

Adults

In 2014, 20.9% of adults aged 18 and over met the 2008 federal physical activity guidelines for both aerobic activity and muscle strengthening (Table 57).

Between 1999–2002 and 2011–2014, the percentage of adults aged 20 and over with Grade 1 obesity (a body mass index [BMI] of 30.0–34.9) increased from 17.9% to 20.6%. Those with Grade 2 obesity (BMI of 35.0–39.9) rose from 7.6% to 8.8%, and those with Grade 3 obesity (BMI of 40 or higher) increased from 4.9% to 6.9% (percentages are age-adjusted) (Table 58).

In 2014, 16.8% of adults aged 18 and over were current cigarette smokers, a decline from 2004 (20.9%). Men (18.8%) were more likely than women (14.8%) to be current cigarette smokers in 2014 (Table 47).

Measures of Health and Disease Prevalence for the Noninstitutionalized Population

In 2012–2014, 4.9% of children under age 18 had an asthma attack in the past year, and 5.6% had a food allergy (Table 35).

Among children aged 5–17, 10.2% were diagnosed with attention deficit/hyperactivity disorder and 5.4% had serious emotional or behavioral difficulties in 2012–2014 (Table 35).

Between 2003 and 2013, the incidence rates of four selected notifiable diseases—tuberculosis, hepatitis A, hepatitis B, and meningococcal disease—decreased, while the incidence rates of Lyme disease increased 57%, to 11.62 new cases per 100,000 population in 2013, and pertussis (a vaccine-preventable disease also known as whooping cough) more than doubled to 9.12 new cases per 100,000 population in 2013. Despite the long-term decline in acute hepatitis B cases, there was a 5% increase in the number of reported cases from 2012 to 2013 (Table 33 and Figure 5).

In 2014, the percentage of adults who reported their health as fair or poor ranged from 6.1% of those aged 18–44 to 24.9% of those aged 75 and over (Table 45).

In 2013–2014, 12.0% of adults aged 45–64 and 29.4% of adults aged 65 and over had ever been told by a physician or other health professional that they had heart disease (Table 38).

In 2013–2014, 6.7% of adults aged 45–64 and 18.2% of adults aged 65 and over had ever been told by a physician or other health professional that they had cancer (excluding squamous and basal cell skin cancers) (Table 38).

In 2011–2014, one-third of adults aged 20 and over had hypertension (having measured high blood pressure or reporting taking antihypertensive medication). Of these adults aged 20 and over with hypertension, nearly one-half (47.0%) had uncontrolled high blood pressure (measured systolic pressure of at least 140 mm Hg or diastolic pressure of at least 90 mm Hg) (Table 54).

In 2014, the prevalence of self-reported serious difficulty concentrating, remembering, or making decisions was highest among men and women aged 75–84 and 85 and over and was similar among men and women in each age group. Difficulty doing errands alone increased with age, and was higher among women than men in all age groups (Figure 6).

Utilization of Health Resources for the Noninstitutionalized Population

Use of Health Care Services

In 2014, 14.9% of persons had no health care visits in the past 12 months, 49.8% had 1–3 health care visits, 23.3% had 4–9 visits, and 11.9% had 10 or more visits. Health care visits for illness, preventive care, or injury include visits to physician offices, emergency departments, clinics, or other locations, in addition to home visits made by health care professionals (Table 65).

In 2014, 83.0% of children aged 2–17 years, 62.0% of adults aged 18–64, and 62.4% of adults aged 65 and over had visited a dentist in the past year (Table 78).

Use of Preventive Medical Care Services for the Noninstitutionalized Population

In 2014, 71.6% of children aged 19–35 months had completed the combined 7-vaccine series of childhood vaccinations (4 or more doses of diphtheria and tetanus toxoids and pertussis vaccine [DTP], diphtheria and tetanus toxoids vaccine [DT], or diphtheria and tetanus toxoids and acellular pertussis vaccine [DTaP]; 3 or more doses of any poliovirus vaccine; 1 or more doses of a measles-containing vaccine [MCV]; 3 or more doses or 4 or more doses of *Haemophilus influenzae* type b vaccine [Hib] depending on Hib vaccine product type [full series Hib]; 3 or more doses of hepatitis B vaccine; 1 or more doses of varicella vaccine; and

4 or more doses of pneumococcal conjugate vaccine [PCV]) (Table 66).

Between 2013 and 2014, receipt of the recommended three doses of human papillomavirus (HPV) vaccine for adolescents aged 13–17 increased among females from 36.8% to 39.7% and among males from 13.4% to 21.6% (Table 67).

In 2013, Pap test utilization was highest among women currently recommended for routine cervical cancer screening; 81.6% of women aged 21–44 and 73.9% of women aged 45–64 received a Pap test in the past 3 years (Table 71 and Figure 10).

In 2014, 42.2% of adults aged 18 and over had received an influenza vaccination in the past 12 months. Influenza vaccination increased with age, with 30.2% of those aged 18–44, 43.3% of those aged 45–64, and 70.1% of those aged 65 and over reporting an influenza vaccination in the past 12 months (Table 68).

In 2014, 61.3% of adults aged 65 and over had ever received a pneumococcal vaccination (Table 69).

Difficulty Accessing Needed Medical Care, Prescription Drugs, and Dental Care Due to Cost for the Noninstitutionalized Population

In 2014, 8.2% of persons reported delaying or not receiving needed medical care due to cost, 5.6% reported not receiving needed prescription drugs due to cost, and 10.0% reported not receiving needed dental care due to cost in the past 12 months (Table 63).

Among adults aged 18–64, the percentage who reported delaying or not receiving needed medical care, not receiving needed prescription drugs, and not receiving needed dental care due to cost in the past 12 months increased 22%–31% during 2004–2010, and then declined 24%–32% during 2010–2014 (Table 63).

Health Care Resources

In 2013, there were 27.6 physicians in patient care per 10,000 civilian population in the United States. The number of patient care physicians per 10,000 population ranged from 18.6 in Idaho to 43.0 in Massachusetts and 66.1 in the District of Columbia (Table 83).

In 2013, the United States had 4,974 community hospitals and 795,603 community hospital beds. Community hospital occupancy averaged 62.9% in 2013, down from 67.3% in 2005 (Table 89).

In 2013, there were 60.46 professionally active dentists per 100,000 civilian population in the United States. The number of dentists per 100,000 population ranged from 40.90 in

Arkansas to 81.22 in New Jersey and 89.20 in the District of Columbia (Table 86).

In 2013, about 8 of every 10 office-based physicians had computerized electronic health record components that recorded patient history and demographic information, ordered prescriptions, and submitted prescriptions to the pharmacy. About 7 of 10 had components to provide electronic warnings of drug interactions and contraindications and to order lab tests electronically (Figure 13).

In 2014, there were 15,643 certified nursing homes with 1,693,943 nursing home beds. U.S. nursing home occupancy averaged 80.8% in 2014. Nursing home occupancy rates were highest in North Dakota (92.4%), Rhode Island (91.9%), South Dakota (91.9%), and the District of Columbia (91.8%) in 2014. The lowest occupancy rates were in Oregon (60.1%), Utah (64.3%), and Idaho (64.5%) (Table 92).

Health Care Expenditures and Payers

Health Care Expenditures

In 2014, personal health care expenditures in the United States totaled \$2.6 trillion—a 5.0% increase from 2013. The per capita personal health care expenditure for the total U.S. population was \$8,054 in 2014—up from \$7,727 in 2013 (Table 93).

Expenditures for hospital care accounted for 37.9% of all personal health care expenditures in 2014. Physician and clinical services accounted for 23.5% of total personal health care expenditures, prescription drugs for 11.6%, and nursing care facilities and continuing care retirement communities for 6.1%; the remaining spending was for other types of personal health care expenditures (Table 94).

In 2014, prescription drug expenditures totaled \$297.7 billion—up 12.2% from \$265.3 billion in 2013 (Table 94).

In 2013, the average cost for the entire hospitalization involving a heart valve procedure was \$51,415; a coronary artery bypass graft procedure was \$41,274; cardiac pacemaker or defibrillator insertion, revision, replacement, or removal was \$35,074; and spinal fusion was \$28,696 (Table 96).

Health Care Payers

In 2014, 33.9% of all personal health care expenditures were paid by private health insurance, 22.7% were paid by Medicare, and 17.4% by Medicaid; consumers paid 12.9% out-of-pocket; and the remaining expenditures were paid by other types of insurance, payers, and programs (Table 95).

In 2014, the Medicare program had 53.8 million enrollees and expenditures of \$613.3 billion—up from 52.5 million enrollees and \$582.9 billion in expenditures the previous year. Expenditures for the Medicare drug program (Part D)

were \$78.1 billion in 2014—up from \$69.7 billion in 2013 (Table 107).

Health Insurance Coverage for the Noninstitutionalized Population

The Affordable Care Act (ACA) of 2010's major provisions were in effect by January 2014. Many of these provisions are intended to expand health insurance and health benefits coverage. Between 2013 and 2014, the percentage of adults aged 18–64 who were uninsured decreased 20%, from 20.5% to 16.3% (Table 105).

From 2014 to June 2015 (preliminary data), the percentage of adults aged 18–64 who were uninsured declined 22%, to 12.7% (Martinez ME, Cohen RA. Health insurance coverage: Early release of estimates from the National Health Interview Survey, January–June 2015. NCHS; 2015. Available from: (<http://www.cdc.gov/nchs/data/nhis/earlyrelease/insur201511.pdf>) (Table 105).

A provision of ACA requires insurers to extend dependent coverage on a family plan until age 26, effective in 2010. This provision, along with other ACA provisions and changes in insurance coverage, has contributed to the 42% decrease in the percentage of adults aged 19–25 who were uninsured, from 33.8% in 2010 to 19.7% in 2014 (Table 105).

From 2014 to June 2015 (preliminary data), the percentage of adults aged 19–25 who were uninsured declined 19%, to 15.9% (Martinez ME, Cohen RA. Health insurance coverage: Early release of estimates from the National Health Interview Survey, January–June 2015. NCHS; 2015. Available from: (<http://www.cdc.gov/nchs/data/nhis/earlyrelease/insur201511.pdf>) (Table 105).

Between 2004 and 2014, the percentage of the population under age 65 with private health insurance obtained through the workplace declined from 64.0% to 56.8% (Table 103).

Between 2004 and 2014, among children in families with income of 100%–199% of the poverty level, the percentage of uninsured children under age 18 decreased from 15.1% to 8.7%, while Medicaid or Children's Health Insurance Program (CHIP) coverage among children in families with income of 100%–199% of poverty increased from 40.2% to 60.0% (Tables 104 and 105).

In 2014, Massachusetts (3.9%), Vermont (5.4%), Hawaii (5.7%), and the District of Columbia (6.1%) had the lowest percentages of persons uninsured (i.e., without public or private coverage) among those under age 65, while Alaska (19.2%), Florida (20.1%), and Texas (21.2%) had the highest percentages uninsured (Table 114).

Special Feature on Racial and Ethnic Health Disparities

The difference between the highest (non-Hispanic black) and lowest (non-Hispanic Asian or Pacific Islander) infant mortality rates among the five racial and ethnic groups narrowed from 9.41 deaths per 1,000 live births in 1999 to 7.21 in 2013 (Figure 19).

In 2014, non-Hispanic black mothers had the highest percentage of preterm births before 37 weeks gestation (11.1%) and non-Hispanic Asian or Pacific Islander mothers had the lowest percentage (6.8%) (Figure 20).

In 2011–2014, among children and adolescents aged 2–19, Hispanic children and adolescents had the highest prevalence of obesity (21.9%) and non-Hispanic Asian children and adolescents had the lowest prevalence (8.6%) (Figure 22).

In 2011–2014 among men aged 20 and over, non-Hispanic black men had the highest prevalence of hypertension (42.4%) and Hispanic men had the lowest (27.7%); among women aged 20 and over, non-Hispanic black women had the highest prevalence of hypertension (44.0%) and non-Hispanic Asian women had the lowest (25.0%) (percentages are age-adjusted) (Figure 23).

The difference for women between the highest (non-Hispanic white) and lowest (non-Hispanic Asian) percentages of current cigarette smokers among the four racial and ethnic groups narrowed from 17.5 percentage points in 1999 to 13.2 in 2014 (percentages are age-adjusted) (Figure 24).


In 2014 among adults aged 18–64, Hispanic adults had the highest percentage of nonreceipt of dental care in the past 12 months due to cost (15.7%) and non-Hispanic Asian adults had the lowest percentage (6.3%) (Figure 27).

Chartbook: Figures 1–17

Mortality

Life Expectancy at Birth, by Country

Figure 1. Life expectancy at birth, by sex and country: Organisation for Economic Co-operation and Development (OECD) countries, 2013



Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig01>

In 2013, U.S. males and females ranked 25th and 27th, respectively, in life expectancy compared with males and females in other OECD countries.

Life expectancy is often used to evaluate the overall health of a population (1). Life expectancy at birth for males and females in the United States was compared with those for males and females in 30 other countries. In 2013, life expectancy at birth for males ranged from a low of 71.7 years for Mexico to a high of 80.7 years for Switzerland, with the United States (76.4 years) ranking 25th out of 31 countries. Life expectancy at birth for females ranged from a low of 77.4 years for Mexico to a high of 86.6 years for Japan, with the United States (81.2 years) tied with Poland and ranking 27th out of 31 countries.


NOTES: Countries with estimated life expectancies or series breaks for 2013 are not presented. Differences in life expectancy may reflect differences in reporting methods, which can vary by country, in addition to actual differences in mortality rates.

SOURCE: CDC/NCHS, *Health, United States, 2015*, Tables 14 and 15. Data for the United States from the National Vital Statistics System (NVSS); all other data from the Organisation for Economic Co-operation and Development (OECD).

Mortality

Selected Causes of Death

Figure 2. Age-adjusted death rates for selected causes of death for all ages, by sex: United States, 2004–2014



Between 2004 and 2014, the all-cause, age-adjusted death rate decreased 12% among males and 11% among females.

During 2004–2014, age-adjusted death rates among males declined 29% for stroke, 23% for heart disease, 16% for cancer, and 10% for both diabetes and CLRD, and increased 11% for Alzheimer's disease and 4% for unintentional injuries. Among females, age-adjusted death rates declined 29% for stroke, 27% for heart disease, 21% for diabetes, and 13% for cancer, and increased 15% for Alzheimer's disease and 11% for unintentional injuries. In 2014, age-adjusted death rates among males were higher than among females for heart disease, cancer, CLRD, diabetes, stroke, and unintentional injuries and were lower among males than females for Alzheimer's disease.

NOTES: CLRD is chronic lower respiratory diseases. A change in the coding rules for nephritis, nephrotic syndrome and nephrosis caused an increase in the number of deaths attributed to diabetes beginning with 2011 data. Thus, the trend for diabetes death rates should be interpreted with caution.

SOURCE: CDC/NCHS, *Health, United States, 2015*, Table 17. Data from the National Vital Statistics System (NVSS).

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig02>

Mortality

Suicide and Homicide

In 2014, suicide rates were higher than homicide rates for males and females of all age groups.


In 2014, suicide was the 10th and homicide the 17th leading cause of death in the U.S.

(Table 19) (2). Suicide and homicide deaths impose emotional and financial costs on both families and society, and death rates for these causes differ by age and other factors (3–7). Suicide rates were higher among males than among females overall (21.1 deaths per 100,000 population compared with 6.0) (Table 30) and within each age group. Among males in 2014, suicide rates were higher among those aged 45–64 and 65 and over than among younger age groups. Among females, suicide rates were highest among those aged 45–64.

Homicide rates were higher among males than among females overall (8.0 deaths per 100,000 population compared with 2.1) (Table 29) and within each age group. Among both males and females, homicide rates were higher among those aged 15–24 and 25–44 than among older age groups in 2014.

SOURCE: CDC/NCHS, *Health, United States, 2015*, Tables 29 and 30. Data from the National Vital Statistics System (NVSS).

Figure 3. Suicide and homicide death rates among persons aged 15 and over, by age and sex: United States, 2014




Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig03>

Natality

Teenage Childbearing

Figure 4. Teenage childbearing, by maternal age and race and Hispanic origin: United States, 2004–2014



Between 2004 and 2014, teenage birth rates declined among all racial and ethnic groups.

Teen childbearing often limits the mother's educational and occupational opportunities, and female babies born to teen mothers are more likely to become teen mothers themselves (8,9). In 2014, teen childbearing fell to a historic low of 24.2 per 1,000 females overall and for each race and Hispanic-origin group (8). Between 2004 and 2014, birth rates declined 50% for teenagers aged 15–17 and 36% for those aged 18–19 (Table 3). Among teenagers aged 15–17, birth rates decreased 44% for non-Hispanic white, 51% for American Indian or Alaska Native, 54% for non-Hispanic black, 59% for Hispanic, and 61% for Asian or Pacific Islander females. Among teenagers aged 18–19, birth rates decreased 32% for non-Hispanic white, 39% for American Indian or Alaska Native, 39% for non-Hispanic black, 47% for Hispanic, and 48% for Asian or Pacific Islander females.

SOURCE: CDC/NCHS, *Health, United States, 2015*, Table 3. Data from the National Vital Statistics System (NVSS).

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig04>

Morbidity

Notifiable Disease Rates


Between 2003 and 2013, the rates for pertussis—a vaccine-preventable disease—and Lyme disease increased, while rates for tuberculosis, hepatitis A, hepatitis B, and meningococcal disease decreased.

Public health officials rely on regular, frequent, timely reporting of notifiable diseases to identify at-risk groups, monitor trends, and control the spread of infectious diseases (10,11). Between 2003 and 2013, the incidence rates of four selected diseases decreased—hepatitis A (79% decrease), meningococcal disease (70%), hepatitis B (63%), and tuberculosis (41%)—while the rates of Lyme disease (57%) and pertussis (whooping cough) (126%) increased. The hepatitis B rate declined in the past decade, but reported cases increased 5% from 2012 to 2013.

NOTES: Diseases with consistent definitions and the greatest changes between 2003 and 2013 were selected for display. Food-borne illnesses were not selected due to year-to-year variation. Rates used the postcensal total resident population and may differ from those elsewhere if different population estimates were used to calculate rates.

SOURCE: CDC/NCHS, *Health, United States, 2015*, Table 33. Data from the National Notifiable Diseases Surveillance System (NNDSS).


Figure 5. Selected notifiable disease rates: United States, 2003 and 2013



Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig05>

Functional Limitations Disability

Figure 6. Selected disability indicators among adults aged 18 and over, by sex and age: United States, 2014



In 2014, disabilities related to cognition and independent living were highest in older age groups; more women than men in each age group reported difficulty doing errands alone.

In 2014, among noninstitutionalized men and women, the prevalence of self-reported serious difficulty concentrating, remembering, or making decisions was higher among older age groups (75–84 and 85 and older) than among younger age groups (18–64 and 65–74) and was similar among men and women in each age group. Difficulty doing errands alone—another disability measure—increased with age. Women in all age groups were more likely than men to report difficulty doing errands alone, ranging from 26% more likely among women aged 18–64 to 72% more likely among women aged 85 and over, compared with men in the same age groups.

NOTE: See [data table for Figure 6](#).
SOURCE: CDC/NCHS, National Health Interview Survey (NHIS).

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig06>


Health Risk Factors Current Cigarette Smoking

During 2004–2014, cigarette smoking prevalence declined among women aged 18–44 and adults aged 45–64.

Smoking is associated with an increased risk of heart disease, stroke, lung and other types of cancers, and chronic lung diseases (12). During 2004–2014, the percentage of adults who smoked cigarettes declined for women aged 18–44 and for both men and women aged 45–64, and remained stable for men and women aged 65 and over. For men aged 18–44, smoking prevalence was stable from 2004–2009 and then declined through 2014. The prevalence of smoking generally was higher for men aged 18–44 and 45–64 than for women in the same age groups (except for 2012). Among adults aged 65 and over, the prevalence for men and women was similar for most years; from 2011–2014, prevalence was higher among men than women. In 2014, 18.8% of men and 14.8% of women aged 18 and over were current cigarette smokers (Table 47).

SOURCE: CDC/NCHS, *Health, United States, 2015*, Table 47. Data from the National Health Interview Survey (NHIS).

Figure 7. Current cigarette smoking among adults aged 18 and over, by sex and age: United States, 2004–2014




Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig07>

Health Risk Factors

Children and Adolescents With Obesity

Figure 8. Obesity among children and adolescents aged 2–19 years, by age: United States, 1999–2002 through 2011–2014



Between 1999–2002 and 2011–2014, the prevalence of obesity was stable among children aged 6–11; increased among adolescents aged 12–19; and increased from 1999–2002 to 2003–2006 among those aged 2–5, then declined through 2011–2014.

Excess body weight in children is associated with excess morbidity during childhood and excess body weight in adulthood (13–16). Obesity among children is defined as a body mass index at or above the sex- and age-specific 95th percentile of the CDC growth charts (15,16). From 1988–1994 to 1999–2002, obesity increased among children aged 2–19. Among children aged 2–5, the prevalence of obesity increased from 1999–2002 to 2003–2006 and then declined through 2011–2014. Among children aged 6–11, the prevalence of obesity was stable from 1999–2002 to 2011–2014. Between 1999–2002 and 2011–2014, the prevalence of obesity among adolescents aged 12–19 increased 28%.

SOURCE: CDC/NCHS, *Health, United States, 2015*, Table 59. Data from the National Health and Nutrition Examination Survey (NHANES).

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig08>

Health Risk Factors


Adults With Overweight and Obesity

Between 1999–2002 and 2011–2014, the prevalence of obesity among men (Grades 1, 2, and 3) and women (Grade 3 only) increased, while the prevalence of overweight but not obese declined among men and remained stable among women aged 20 and over.

Reducing the prevalence of obesity is a public health priority because obesity is correlated with excess morbidity and mortality (17–19). In particular, Grade 2 or higher obesity significantly increases the risk of death (20). Between 1999–2002 and 2011–2014, the percentage of adults aged 20 and over with Grades 1, 2, and 3 obesity increased among men. For women, the percentage of Grade 1 obesity and Grade 2 obesity remained stable while Grade 3 obesity increased. Meanwhile, the percentage of men aged 20 and over who were overweight but not obese declined and was stable among women. In 2011–2014, women were almost twice as likely to have Grade 3 obesity as men (8.9% compared with 4.9%).

NOTES: BMI is body mass index. Overweight but not obese (25 ≤ BMI < 30), Grade 1 obesity (30 ≤ BMI < 35), Grade 2 obesity (35 ≤ BMI < 40), and Grade 3 obesity (BMI ≥ 40). SOURCE: CDC/NCHS, *Health, United States, 2015*, Table 58. Data from the National Health and Nutrition Examination Survey (NHANES).

Figure 9. Overweight and obesity among adults aged 20 and over, by sex and grade of obesity: United States, 1999–2002 through 2011–2014




Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig09>

Utilization

Pap Test Use

Figure 10. Pap test utilization within the past 3 years, by age: United States, 2003–2013



From 2003 to 2013, Pap test utilization decreased for all age groups; the largest decreases were for women aged 18–20 and 65 and over (age groups no longer recommended for routine testing).

Pap tests have reduced cervical cancer deaths by detecting cases at earlier and more treatable stages (21). Current Pap test recommendations suggest limiting routine testing to women aged 21–65 and vary based on individual risk factors including cervical cancer risk, human papillomavirus (HPV) testing, and screening history (22). From 2003 to 2013, recent Pap testing declined for all age groups. The refined recommendations may help explain the decrease for women aged 21–44 (5%) and 45–64 (9%). The greatest decreases were for age groups for which routine testing is no longer recommended: 18–20 (39%), 65–74 (22%), and 75 and over (45%).

NOTES: Pap tests (Pap smears) may be used for screening or diagnostic purposes; the purpose cannot be determined from NHIS. See Appendix II, Pap smear. The 65–74 group includes women aged 65 who are still recommended to have routine testing.

SOURCE: CDC/NCHS, *Health, United States, 2015*, Table 71. Data from the National Health Interview Survey (NHIS).

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig10>

Utilization

Emergency Department Use


During 2004–2014, adults aged 18–64 with Medicaid coverage were more likely to have visited an emergency department within the past year than those with private coverage or the uninsured.

Emergency departments (EDs) are critical in the U.S. health care system, providing emergency and after hours care (23–25). During 2004–2014, adults aged 18–64 with Medicaid coverage were about twice as likely as those with private coverage or the uninsured to have had an ED visit in the past year. During 2004–2014, the percentage with a recent ED visit was stable for adults with Medicaid; for those with private coverage, the percentage was stable through 2010, then declined through 2014; and for the uninsured, the percentage increased during 2004–2011, then declined through 2014. Although adults with Medicaid were more likely to have an ED visit, only 23.4% of all 2014 ED visits were by those with Medicaid; 15.1% were by the uninsured, and 53.6% were by those with private coverage, reflecting the larger percentage of adults with private coverage.

NOTE: See data table for Figure 11.

SOURCE: CDC/NCHS, *Health, United States, 2015*, Table 74. Data from the National Health Interview Survey (NHIS).

Figure 11. Emergency department utilization within the past 12 months among adults aged 18–64, by type of coverage: United States, 2004–2014




Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig11>

Utilization

Difficulty Accessing Needed Medical Care or Prescription Drugs Due to Cost

Figure 12. Delay or nonreceipt of needed medical care and nonreceipt of needed prescription drugs in the past 12 months due to cost among adults aged 18–64, by health insurance coverage: United States, 2004–2014



Uninsured adults aged 18–64 are more likely than those with Medicaid or private coverage to report difficulties affording needed medical care and prescription drugs.

Uninsured adults are more likely than the insured to delay or forego needed medical care and prescription drugs due to cost (26,27). During 2004–2014, uninsured adults were 4–5 times more likely than those with private coverage and 1½–3 times more likely than those with Medicaid to report medical care and prescription access problems. For adults with Medicaid, medical care access problems were stable until 2008 and then decreased through 2014. For those with private insurance, medical care access problems increased until 2009 and then declined through 2014. For the uninsured, medical care and prescription access problems increased (until 2010 and 2009, respectively) and then were stable for medical care and decreased through 2014 for access to drugs. Drug access problems were stable in 2004–2014 for those with private insurance but decreased for adults with Medicaid.

SOURCE: CDC/NCHS, *Health, United States, 2015*, Table 63. Data from the National Health Interview Survey (NHIS).

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig12>


Health Care Resources

Electronic Health Record Systems

In 2013, most physician offices had electronic health record (EHR) systems that record patient history and demographic information (83.0%), order prescriptions (82.6%), send prescriptions to the pharmacy (78.7%), warn of drug interactions and contraindications (73.8%), and order lab tests (68.9%).

EHR systems are thought to make health care delivery more efficient by improving clinician decision-making, care coordination, health care safety, and patient outcomes (28–30). In 2013, about 8 of every 10 office-based physicians had computerized components that recorded patient history and demographic information, ordered prescriptions, and sent prescriptions to the pharmacy. About 7 of every 10 had a component that warned of drug interactions and contraindications and ordered lab tests. From 2010 to 2013, the percent increase in the use of these EHR components ranged from 12% for components to record patient history and demographic information to 80% for components to send prescriptions to the pharmacy.

Figure 13. Electronic health record system components in physician offices, by selected component type: United States, 2010 and 2013



NOTE: See data table for Figure 13.


SOURCE: CDC/NCHS, National Ambulatory Medical Care Survey (NAMCS)—National Electronic Health Records Survey.

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig13>

Health Care Resources

Physicians Accepting New Patients

Figure 14. Office-based physicians accepting new patients, by patient source of payment and urban–rural status: United States, 2013



In 2013, physicians in urban large fringe areas (suburbs) were less likely to accept new Medicaid patients than physicians in any other urban–rural category.

Under the ACA, more Americans have health care coverage. In some areas, finding a physician who is accepting new patients may be difficult (31–33). Physician acceptance of new patients was examined by urban–rural status, which classifies physicians by the location of their practice (34). In 2013, Medicaid acceptance rates varied across urban–rural categories, with the lowest acceptance rates for physicians in urban large fringe counties (suburbs). Physicians in rural areas (micropolitan and noncore) were more likely to accept new Medicaid patients than those in urban areas. Comparing physicians' acceptance of new private to new Medicaid patients, physicians in urban areas were less likely to accept new Medicaid than new private patients, while acceptance rates for new Medicaid and private patients were similar for physicians in rural areas.

NOTE: See data table for Figure 14.
SOURCE: CDC/NCHS, National Ambulatory Medical Care Survey (NAMCS)—National Electronic Health Records Survey.

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig14>

Personal Health Care Expenditures


Major Source of Funds

Between 2004 and 2014, Medicare expenditures for personal health care grew more rapidly than out-of-pocket, private insurance spending, and total Medicaid.

Between 2004 and 2014, total personal health care expenditures grew from \$1.6 trillion to \$2.6 trillion (Table 95). During 2004–2014, the average annual growth in expenditures was 6.8% for Medicare, 5.4% for Medicaid (federal), 4.6% for Medicaid (state), 5.1% for Medicaid (total), 4.4% for private health insurance, and 2.9% for out-of-pocket spending. In 2014, private health insurance accounted for the highest spending on personal health care at \$868.8 billion, followed by Medicare at \$580.7 billion. Out-of-pocket spending by individuals reached \$329.8 billion in 2014, and spending on Medicaid reached \$273.6 billion in federal dollars and \$171.3 billion in state dollars for a total of \$444.9 billion in Medicaid spending. The remainder was paid for by other types of insurance, payers, and programs (Table 95) (35).

NOTES: Personal health care expenditures are outlays relating directly to patient care. See Appendix II, Health expenditures, national.
SOURCE: CDC/NCHS, Health, United States, 2015, Table 95. Data from the Centers for Medicare & Medicaid Services, National Health Expenditure Accounts (NHEA).

Figure 15. Personal health care expenditures, by source of funds: United States, 2004–2014




Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig15>

Health Insurance

Coverage Among Adults Aged 18–64

Figure 16. Health insurance coverage among adults aged 18–64, by type of coverage: United States, 2004–June 2015 (preliminary data)



From 2004 to June 2015, the percentage of adults aged 18–64 with Medicaid coverage increased, the percentage with private coverage decreased through 2012 and then increased through June 2015, and the percentage uninsured increased through 2013 and then declined through June 2015.

Health insurance is a major determinant of access to health care (26). Among adults aged 18–64, the percentage with private coverage declined from 2004 (71.1%) to 2012 (65.1%) and then increased through June 2015 (70.6%) (Table 102) (36). As of June 2015, 8.9 million adults aged 18–64 were covered by private plans obtained through the Health Insurance Marketplace or state-based exchanges (36). The percentage with Medicaid coverage increased from 2004 (6.8%) to June 2015 (12.2%) (Table 104) (37). The percentage of adults aged 18–64 who were uninsured increased from 2004 (19.3%) to 2013 (20.5%) and then declined through June 2015 (12.7%) (Table 105) (36).

NOTE: Preliminary estimates for the first 6 months of 2015 are shown with a dashed line (36).

SOURCE: CDC/NCHS, *Health, United States, 2015*, Tables 102, 104, 105. Data from the National Health Interview Survey (NHIS).

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig16>


Health Insurance

Coverage by Medicaid Expansion State

Between 2013 and 2014, the percentage of adults aged 18–64 who were uninsured declined in both Medicaid expansion states (by 28%) and nonexpansion states (by 14%), and the percentage covered by Medicaid increased by 25% in Medicaid expansion states.

Under the ACA (38), states are authorized to expand Medicaid coverage to adults with low incomes, up to and including 138% of the poverty level (39). Between 2013 and 2014, the percentage of adults aged 18–64 who were uninsured declined in both Medicaid expansion states and nonexpansion states; however, the decline in the uninsured percentage was greater for states that expanded their Medicaid programs (28% compared with 14%). The percentage covered by private insurance increased by about 4% in both Medicaid expansion and nonexpansion states. Medicaid coverage increased 25% in states that expanded their programs and was stable in states that did not expand their programs.

Figure 17. Health insurance coverage among adults aged 18–64, by state Medicaid expansion status: United States, 2013 and 2014



NOTES: States were classified based on their decision to expand Medicaid as of January 1, 2014 (40). See data table for Figure 17. SOURCE: CDC/NCHS, National Health Interview Survey (NHIS).

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig17>

Chartbook: Figures 18–27

Special Feature on Racial and Ethnic Health Disparities

Special Feature on Racial and Ethnic Health Disparities: 30 Years After the Heckler Report

Introduction

The 1985 Report of the Secretary's Task Force on Black and Minority Health, released by then Secretary of Health and Human Services Margaret Heckler, documented significant disparities in the burden of illness and mortality experienced by blacks and other minority groups in the U.S. population compared with whites (41). The report laid out an ambitious agenda, including improving minority access to high-quality health care, expanding health promotion and health education outreach activities, increasing the number of minority health care providers, and enhancing federal and state data collection activities to better report on minority health issues. In the 30 years since the Heckler Report, national efforts to improve minority health through outreach, programming, and monitoring have included the formation of the Department of Health and Human Services (HHS) Office of Minority Health in 1986 (42); the annual National Healthcare Quality and Disparities Reports first issued in 2003 (43); the adoption of disparities elimination as an overarching goal of Healthy People 2010 (44); and most recently, an HHS Action Plan to Reduce Racial and Ethnic Health Disparities—a comprehensive federal commitment to reduce and eventually eliminate disparities in health and health care (45).

Race is a social construct influenced by a complex set of factors (46,47). Because of the complexity and difficulty in conceptualizing and defining race, as well as the increasing representation of racial and ethnic subgroups in the United States, racial classification and data collection systems continue to evolve and expand. In 1977, the Office of Management and Budget (OMB) required that all federal data collection efforts collect data on a minimum of four race groups (American Indian or Alaskan Native, black, Asian or Pacific Islander, and white) and did not allow the reporting of more than one race (48). In 1997, in response to growing interest in more detailed reporting on race and ethnicity, OMB mandated data collection for a minimum of five race groups, splitting Asian or Pacific Islander into two categories (Asian, and Native Hawaiian or Other Pacific Islander) (49). In addition, the 1997 standards allowed respondents to report more than one race. A minimum of two categories for data collection on ethnicity, “Hispanic or Latino” and “Not Hispanic or Latino,” were also required under the 1997 OMB standards. Consequently, whereas the Heckler Report primarily documented black–white differences in health and mortality due to data limitations, this Special Feature is able to report on more detailed racial and ethnic groups. For example, Figures 19–21 display trends in infant mortality and low-risk cesarean section deliveries, and the current data on preterm births for five Hispanic-origin groups.

At the time of the Heckler Report, 22.3% of the population were considered racial or ethnic minorities (Table 1). Current Census (2014) estimates identify 37.9% of the population as racial or ethnic minorities (50). In 2014, Hispanic persons, who may be of any race, comprised 17.4% of the U.S. population. Non-Hispanic multiple race persons were 2.0% of the population. For the single race groups, non-Hispanic American Indian or Alaska Native persons were 0.7%, non-Hispanic Asian persons were 5.3%, non-Hispanic black persons were 12.4%, non-Hispanic Native Hawaiian or Other Pacific Islander persons were 0.2%, and non-Hispanic white persons were 62.1% of the U.S. population in 2014 (50).

Understanding the demographic and socioeconomic composition of U.S. racial and ethnic groups is important because these characteristics are associated with health risk factors, disease prevalence, and access to care, which in turn drive health care utilization and expenditures. Non-Hispanic white persons are, on average, older than those in other racial and ethnic groups, with a median age of 43.1 years, and Hispanic individuals are the youngest, with a median age of 28.5 years in 2014 (50). About one-quarter of black only persons (26.2%) and Hispanic persons (23.6%) lived in poverty compared with 10.1% of non-Hispanic white only persons and 12.0% of Asian only persons in 2014 (51). Non-Hispanic black only children and Hispanic children were particularly likely to live in poverty (37.3% and 31.9%, respectively, in 2014) (52). However, Hispanic individuals are often found to have quite favorable health and mortality patterns in comparison with non-Hispanic white persons and particularly with non-Hispanic black persons, despite having a disadvantaged socioeconomic profile—a pattern termed the epidemiologic paradox (53).

HHS defines a racial or ethnic health disparity as “a particular type of health difference that is closely linked with social, economic, and/or environmental disadvantage. Health disparities adversely affect groups of people who have systematically experienced greater obstacles to health based on their racial or ethnic group” (54). There are many different ways to measure racial and ethnic differences in health and mortality, which can lead to different conclusions (55–58). This Special Feature on Racial and Ethnic Health Disparities (Special Feature) uses the maximal rate difference, one of three overall measures used in Healthy People 2020 to measure differences among groups of people (see [Technical Notes](#)). The maximal rate difference is an overall measure of health disparities calculated as the absolute difference between the highest and lowest group rates in the population for a given characteristic (59). The identification of groups that experience the highest and lowest rates in this Special Feature was based on observed rates and was not tested for a statistically significant difference against other rates. Ties in highest or lowest rates

were resolved by examining decimal places. With respect to changes in health disparities over time, tracking the maximal rate difference over time enables one to determine whether the absolute difference between the highest and lowest group rates is increasing, decreasing, or stable.

The Special Feature charts that follow provide detailed comparisons of key measures of mortality, natality, health conditions, health behaviors, and health care access and utilization, by race, race and ethnicity, or by detailed Hispanic origin, depending on data availability. A majority of the 10 graphs in this year's Special Feature present trends in health from 1999–2014. Results indicate that trends in health were generally positive for the overall population and several graphs illustrate success in narrowing gaps in health by racial and ethnic group. Differences in life expectancy, infant mortality, cigarette smoking among women, influenza vaccinations among those aged 65 and over, and health insurance coverage narrowed among the racial and ethnic groups. For example, the absolute difference in infant mortality rates between infants born to non-Hispanic black mothers (highest rate) and infants born to non-Hispanic Asian or Pacific Islander mothers (lowest rate) narrowed between 1999–2014. Differences by racial and ethnic group in the prevalence of high blood pressure and smoking among adult men remained stable throughout the study period, with non-Hispanic black adults more likely to have high blood pressure than adults in other racial and ethnic groups throughout the period, and non-Hispanic black and non-Hispanic white males more likely to be current smokers than Hispanic and non-Hispanic Asian men. For low-risk cesarean sections, influenza vaccinations among adults aged 18–64, and unmet dental care needs, the gap widened among the racial and ethnic groups between 1999–2014.

Despite improvements over time in many of the health measures presented in this Special Feature, disparities by race and ethnicity were found in the most recent year for all 10 measures, indicating that although progress has been made in the 30 years since the Heckler Report, elimination of disparities in health and access to health care has yet to be achieved.

Life Expectancy at Birth

In 2014, life expectancy was longer for Hispanic men and women than for non-Hispanic white or non-Hispanic black men and women.


Life expectancy is a measure often used to gauge the overall health of a population. Life expectancy at birth represents the average number of years that a group of infants would live if the group were to experience the age-specific death rates present in the year of birth. Differences in life expectancy among various demographic subpopulations, including racial and ethnic groups, may reflect subpopulation differences in a range of factors such as socioeconomic status, access to medical care, and the prevalence of specific risk factors in a particular subpopulation (60,61).

During 1980–2014, life expectancy at birth in the United States increased from 70.0 to 76.4 years for males and from 77.4 to 81.2 years for females (Table 15, and data table for

Figure 18). During this period, life expectancy at birth for males and females was longest for white persons and shortest for black persons. For both males and females, racial differences in life expectancy at birth narrowed, but persisted during 1980–2014. Life expectancy at birth was 6.9 years longer for white males than for black males in 1980, and this difference narrowed to 4.2 years in 2014. In 1980, life expectancy at birth was 5.6 years longer for white females than for black females, and this difference narrowed to 3.0 years in 2014.

In 2014, Hispanic males and females had the longest life expectancy at birth, and non-Hispanic black males and females had the shortest. In 2014, life expectancy at birth was 7.2 years longer for Hispanic males than for non-Hispanic black males and 5.9 years longer for Hispanic females than for non-Hispanic black females.

Figure 18. Life expectancy at birth, by sex, race and Hispanic origin: United States, 1980–2014



NOTES: Life expectancy data by Hispanic origin were available starting in 2006 and were corrected to address racial and ethnic misclassification. See Technical Notes and data table for Figure 18.

SOURCE: CDC/NCHS, National Vital Statistics System (NVSS).

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig18>

Infant Mortality

During 1999–2013, infant mortality rates were highest among infants born to non-Hispanic black women (11.11 infant deaths per 1,000 live births in 2013).


Infant mortality, the death of a baby before his or her first birthday, is an important indicator of the health and wellbeing of a country. It not only measures the risk of infant death but it is used as an indicator of maternal health, community health status, and availability of quality health services and medical technology (62,63).

The infant mortality rate in the United States decreased from 7.04 infant deaths per 1,000 live births in 1999 to 6.75 in 2007, and then decreased at a faster rate to 5.96 in 2013. Trends in infant mortality rates during 1999–2013 varied among the five racial and ethnic groups. During 1999–2013, infants born to non-Hispanic black mothers experienced the highest rates of infant mortality (11.11 in 2013) and infants born to non-Hispanic Asian or Pacific Islander mothers experienced the lowest rates (3.90 in 2013). The difference between the highest and lowest infant mortality rates among the five racial and ethnic groups was stable from 1999 to 2006 and then narrowed from 2006 to 2013. The difference between the highest

(non-Hispanic black) and lowest (non-Hispanic Asian or Pacific Islander) infant mortality rates was 9.41 deaths per 1,000 live births in 1999, compared with 7.21 in 2013.

For infants born to Hispanic mothers, the infant mortality rate remained stable during 1999–2008 (5.71 infant deaths per 1,000 live births in 1999) and then decreased to 5.00 in 2013. During 1999–2013, the infant mortality rate for Hispanic infants varied by the mother's Hispanic-origin group. Throughout this period, infants born to Puerto Rican mothers experienced the highest mortality rates. In all years except 2009, infants born to Cuban mothers and those born to Central and South American mothers experienced the lowest mortality rates at alternate times throughout 1999–2013. The difference between the highest (Puerto Rican) and lowest (Cuban) infant mortality rates among Hispanic-origin groups narrowed from 3.71 deaths per 1,000 live births in 1999 to 2.88 in 2013. During 1999–2013, the difference in infant mortality rates was narrower for mothers in the Hispanic-origin groups than for mothers in the five racial and ethnic groups.

Figure 19. Infant mortality rates, by race and Hispanic origin and detailed Hispanic origin of mother: United States, 1999–2013



NOTES: Highest and lowest rates are based on observed rates and were not tested for statistically significant differences against other rates. Ties in highest and lowest rates were resolved by looking at additional decimal places. See [Technical Notes](#) and [data table](#) for Figure 19.

SOURCE: CDC/NCHS, National Vital Statistics System (NVSS), Linked Birth/Infant Death Data Set.

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig19>

Preterm Births

In 2014, non-Hispanic black mothers had the highest percentage of preterm births of the five racial and ethnic groups, and Puerto Rican mothers had the highest percentage of preterm births of the five Hispanic-origin groups.


An infant's gestational age is an important predictor of his or her survival and subsequent health (64–70). Preterm birth prior to 37 weeks gestation affects infant mortality rates and racial and ethnic disparities in infant mortality (Figure 19) (71). The degree of prematurity matters—infants born prior to 32 weeks gestation are at greatest risk of death during infancy, with the risk of infant death decreasing as gestational age increases (72).

In 2014, 7.7% of singleton births occurred before 37 weeks of gestation; 5.7% at 34–36 weeks; 0.8% at 32–33 weeks gestation; and 1.2% before 32 weeks (data table for Figure 20). In 2014, among the five racial and ethnic groups, non-Hispanic black women had the highest percentage of singleton births before 37 weeks (11.1%) and non-Hispanic Asian or Pacific Islander women had the lowest percentage (6.8%). Non-Hispanic black women also had the highest

percentage of singleton preterm births at each preterm gestational age. The difference between the highest (non-Hispanic black) and lowest (non-Hispanic Asian or Pacific Islander) percentages of singleton preterm births among the five racial and ethnic groups was 4.3 percentage points (before 37 weeks), 2.0 percentage points (34–36 weeks), 0.6 percentage points (32–33 weeks), and 1.7 percentage points (before 32 weeks).

Among Hispanic-origin groups in 2014, Puerto Rican mothers had the highest percentage of singleton births before 37 weeks (9.1%) and Cuban mothers had the lowest percentage (7.2%). The difference between the highest (Puerto Rican) and lowest (Cuban) percentages of singleton preterm births among the Hispanic-origin groups was 1.9 percentage points (before 37 weeks) and 1.3 percentage points (34–36 weeks). Central and South American mothers had the lowest percentage of singleton births before 34 weeks. For preterm births before 34 weeks, the difference between the highest (Puerto Rican) and lowest (Central and South American) percentages was 0.2 percentage points (32–33 weeks) and 0.6 percentage points (before 32 weeks).

Figure 20. Preterm births, by gestational age and race and Hispanic origin and detailed Hispanic origin of mother: United States, 2014



NOTES: Preterm births are based on the obstetric estimate of gestational age and are for all singleton births (73). Highest and lowest percentages are based on observed percentages and were not tested for statistically significant differences against other percentages. Ties in highest and lowest percentages were

resolved by looking at additional decimal places. See [Technical Notes](#) and [data table for Figure 20](#).

SOURCE: CDC/NCHS, National Vital Statistics Survey (NVSS).

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig20>

Low-risk Births Delivered by Cesarean Section

During 1999–2014 non-Hispanic black mothers experienced the highest percentage of low-risk cesarean deliveries among the five racial and ethnic groups (29.9% in 2014); Cuban mothers experienced the highest percentage of low-risk cesarean deliveries among the five Hispanic-origin groups (41.4% in 2014).


Cesarean deliveries comprise approximately one-third of all births in the United States (32.2% in 2014) and can place mothers and infants at increased risk for poor health outcomes (74). Over the past decade, professional medical groups have attempted to reduce low-risk cesarean deliveries defined as cesarean deliveries among full term (37 or more completed weeks of gestation), singleton, vertex (head first) births to women giving birth for the first time (75,76).

The percentage of low-risk births that were delivered by cesarean section increased from 19.5% to 26.6% during 1999–2005, stabilized during 2005–2009, and then decreased to 26.0% in 2014 (data table for Figure 21). Throughout the period 1999–2014, non-Hispanic black mothers experienced the highest percentage of low-risk cesarean deliveries (29.9% in 2014) among the five racial and ethnic groups, while non-Hispanic American Indian or Alaska Native mothers

experienced the lowest percentage (21.5% in 2014). The difference between the highest (non-Hispanic black) and lowest (non-Hispanic American Indian or Alaska Native) percentages widened from 4.8 percentage points in 1999 to 8.4 percentage points in 2014.

Among Hispanic mothers, the percentage of low-risk births that were delivered by cesarean section increased from 18.7% to 24.6% during 1999–2004, increased at a slower rate from 2004–2009, and then remained stable during 2009–2014 (data table for Figure 21). Throughout the period 1999–2014 Cuban mothers experienced the highest percentage of low-risk cesarean deliveries (41.4% in 2014), while Mexican mothers experienced the lowest percentage (24.1% in 2014). Among Hispanic-origin groups, the difference between the highest and lowest percentages of low-risk cesarean deliveries was stable during 1999–2002, widened sharply during 2002–2006, and then narrowed during 2006–2014. The difference between the highest (Cuban) and lowest (Mexican) percentages was 11.7 percentage points in 1999, 21.5 percentage points in 2006, and 17.3 percentage points in 2014.

Figure 21. Low-risk births delivered by cesarean section, by race and Hispanic origin and detailed Hispanic origin of mother: United States, 1999–2014



NOTES: The term low-risk cesarean delivery is not meant to imply that a cesarean delivery may not be medically necessary for low-risk women (75). Highest and lowest percentages are based on observed percentages and were not tested for statistically

significant differences against other percentages. Ties in highest and lowest percentages were resolved by looking at additional decimal places. See Technical Notes and data table for Figure 21.

SOURCE: CDC/NCHS, National Vital Statistics System (NVSS).

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig21>

Children and Adolescents With Obesity

In 2011–2014 for children and adolescents aged 2–19 years, Hispanic children and adolescents had the highest prevalence of obesity and non-Hispanic Asian children had the lowest prevalence.

Childhood obesity is a serious public health challenge in the United States and many other industrialized nations in the world (Figure 8) (19,77,78). Excess body weight in children is associated with excess morbidity in childhood and excess body weight in adulthood (13,14). Obesity among children and adolescents is defined as a body mass index at or above the sex- and age-specific 95th percentile of the CDC growth charts (15). Between 1999–2000 and 2013–2014, the percentage of children and adolescents aged 2–19 with obesity increased from 13.9% to 17.2% (79). However, among youth aged 2–19, the prevalence of obesity did not change from 2003–2004 through 2013–2014 (79).


In 2011–2014 for children and adolescents aged 2–19, the percentage with obesity was highest for Hispanic children and adolescents and lowest for non-Hispanic Asian children and adolescents. For those aged 2–19, the difference

between the highest (Hispanic) and lowest (non-Hispanic Asian) percentages was 13.3 percentage points.

For children aged 2–5, the percentage with obesity was highest for Hispanic children and lowest for non-Hispanic white children. (The estimate for non-Hispanic Asian children aged 2–5 was not stable and is not shown.) The difference between the highest (Hispanic) and lowest (non-Hispanic white) percentages was 10.4 percentage points for children aged 2–5. For children aged 6–11, the percentage with obesity was highest for Hispanic children and lowest for non-Hispanic Asian children. For children aged 6–11, the difference between the highest (Hispanic) and lowest (non-Hispanic Asian) percentages was 15.2 percentage points.

In 2011–2014 for adolescents aged 12–19, the percentage with obesity was highest for Hispanic adolescents and lowest for non-Hispanic Asian adolescents. The difference between the highest (Hispanic) and lowest (non-Hispanic Asian) percentages was 13.4 percentage points for adolescents aged 12–19 years.

Figure 22. Obesity among children and adolescents aged 2–19 years, by age and race and Hispanic origin: United States, 2011–2014



* Relative standard error (RSE) of 20%–30%. The estimate for non-Hispanic Asian children aged 2–5 is not shown because the RSE is greater than 30%.
 NOTES: Obesity is defined as a body mass index at or above the sex- and age-specific 95th percentile of the CDC growth charts. Highest and lowest percentages

are based on observed percentages and were not tested for statistically significant differences against other percentages. See [Technical Notes](#) and [data table](#) for Figure 22.

SOURCE: CDC/NCHS, National Health and Nutrition Examination Survey (NHANES).

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig22>

Hypertension

In 2011–2014, non-Hispanic black men and women were the most likely to have hypertension compared with adults in the other racial and ethnic groups.


Hypertension is an important risk factor for cardiovascular disease, stroke, kidney failure, and other health conditions (80,81). In 2011–2014, 84.1% of adults with hypertension were aware of their status, and 76.1% were taking medication to lower their blood pressure (82). Despite improvement in increasing the awareness, treatment, and control of hypertension, diagnosis and treatment of hypertension among minority groups remains a challenge (83).

Hypertension is defined as reporting taking antihypertensive medication and/or having a measured systolic blood pressure of at least 140 mm Hg or a measured diastolic blood pressure of at least 90 mm Hg. The age-adjusted percentage of adults aged 20 and over with hypertension was stable during 1999–2014 (30.8% in 2013–2014) (data table for Figure 23). During 1999–2014, non-Hispanic black adults had the highest percentage with

hypertension among the three racial and ethnic groups (42.7%, age-adjusted in 2013–2014), while with the exception of 1999–2000, adults of Mexican origin had the lowest percentage with hypertension (28.8%, age-adjusted in 2013–2014). The difference between the highest and lowest age-adjusted percentages of adults with hypertension among the three racial and ethnic groups was stable during 1999–2014; in 2013–2014, the difference between the highest (non-Hispanic black) and lowest (Mexican-origin) percentages was 13.9 percentage points.

In 2011–2014, the age-adjusted percentage of adult men and women with hypertension was similar (31.0% and 29.7%, respectively, data table for Figure 23). The difference between the highest (non-Hispanic black) and lowest (Hispanic) age-adjusted percentages of men with hypertension among the four racial and ethnic groups was 14.7 percentage points; for women, the difference between the highest (non-Hispanic black) and lowest (non-Hispanic Asian) was 19.0 percentage points in 2011–2014.

Figure 23: Hypertension among adults aged 20 and over, by sex and race and Hispanic origin: United States, 1999–2000 through 2013–2014



NOTES: Estimates are age-adjusted. Hypertension is having measured high blood pressure (systolic pressure of at least 140 mm Hg or diastolic pressure of at least 90 mm Hg) and/or respondent report of taking antihypertensive medication. Data for Hispanic adults were available starting in 2007–2008 and for non-Hispanic Asian adults in 2011–2012. Highest and lowest percentages are based on

observed percentages and were not tested for statistically significant differences against other percentages. See Technical Notes and data table for Figure 23.

SOURCE: CDC/NCHS, National Health and Nutrition Examination Survey (NHANES).

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig23>

Current Cigarette Smoking

During 1999–2014, differences in cigarette smoking between racial and ethnic groups were larger for women than for men.

Smoking causes more than 480,000 deaths each year, accounting for about one in five deaths in the United States (84). Smokers are more likely to develop heart disease, stroke, and cancer. Smoking also increases the risk for diabetes, cataracts, rheumatoid arthritis, and stillbirth (85).


During 1999–2014, the age-adjusted percentage of adults aged 18 and over who were current cigarette smokers decreased from 25.2% to 19.0% for men and from 21.6% to 15.1% for women (data table for Figure 24). Within each of the four racial and ethnic groups, men were more likely to be current cigarette smokers than women.

In 2014 for men, the age-adjusted percentage of current cigarette smokers was highest for non-Hispanic black men (22.0%) and lowest for Hispanic men (13.8%). The difference between the highest and lowest age-adjusted percentages

of current cigarette smokers among the four racial and ethnic groups remained stable during 1999–2014 because levels for men in all racial and ethnic groups declined similarly during this period. The difference between the highest (non-Hispanic black) and lowest (Hispanic) percentages for men was 8.2 percentage points in 2014.

For women, non-Hispanic white women consistently had the highest age-adjusted percentage of current cigarette smokers among the four racial and ethnic groups throughout 1999–2014 (18.3% in 2014), while non-Hispanic Asian women had the lowest age-adjusted percentage (5.1% in 2014). For women, the difference between the highest (non-Hispanic white) and lowest (non-Hispanic Asian) percentages narrowed from 17.5 percentage points in 1999 to 13.2 in 2014. During 1999–2014, racial and ethnic differences in cigarette smoking prevalence were larger for women than for men.

Figure 24. Current cigarette smoking among adults aged 18 and over, by sex and race and Hispanic origin: United States, 1999–2014



NOTES: Estimates are age-adjusted. Three-year average annual estimates for the American Indian or Alaska Native population are available in the data table for Figure 24. Estimates for non-Hispanic Asian women in 2001 and 2005 have a relative standard error of 20%–30%. Highest and lowest percentages are based on

observed percentages and were not tested for statistically significant differences against other percentages. Ties in highest and lowest percentages were resolved by looking at additional decimal places. See Technical Notes and data table for Figure 24.

SOURCE: CDC/NCHS, National Health Interview Survey (NHIS).

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig24>

Influenza Vaccination

During 1999–2014, influenza vaccination was highest for those aged 65 and over and lowest for those aged 18–64, for all racial and ethnic groups.


Influenza is a serious illness that can lead to hospitalization and sometimes death. Influenza vaccination is especially important for people who are at risk of getting seriously ill from influenza, including those with chronic conditions, older adults, and young children.

The percentage of adults aged 18–64 who received an influenza vaccination in the past 12 months remained stable during 1999–2006 and then increased to 35.8% in 2014 (data table for Figure 25). This pattern was present for all racial and ethnic groups. Decreases in influenza vaccination coverage in 2005 were related to a vaccine shortage (86). For those aged 18–64, no racial and ethnic group was consistently the most likely to receive influenza vaccination during 1999–2014. In 2014, non-Hispanic Asian adults had the highest percentage for influenza vaccination receipt (41.3%) and Hispanic adults had the lowest percentage (27.9%). For adults aged 18–64, the difference between the

highest and lowest percentages of adults receiving an influenza vaccination among the four racial and ethnic groups widened from 6.9 percentage points in 1999 (non-Hispanic white compared with Hispanic) to 13.4 in 2014 (non-Hispanic Asian compared with Hispanic).

For adults aged 65 and over, the percentage who received an influenza vaccination in the past 12 months increased from 65.7% to 70.1% during 1999–2014. During this period, trends in influenza vaccination coverage varied by racial and ethnic group, and no racial and ethnic group was consistently the most or least likely to receive influenza vaccination. In 2014, non-Hispanic Asian adults had the highest percentage for receipt of influenza vaccination (72.7%) and non-Hispanic black adults had the lowest (57.4%). For adults age 65 and over, the difference between the highest (non-Hispanic Asian) and lowest (non-Hispanic black) percentages of older adults receiving an influenza vaccination among the four racial and ethnic groups was stable during 1999–2003 and then narrowed to 15.3 percentage points in 2014.

Figure 25. Influenza vaccination among adults aged 18 and over, by age and race and Hispanic origin: United States, 1999–2014



NOTES: Three-year average annual estimates for the American Indian or Alaska Native population are available in the data table for Figure 25. Highest and lowest percentages are based on observed percentages and were not tested for statistically significant differences against other percentages. Ties in highest

and lowest percentages were resolved by looking at additional decimal places. See Technical Notes and data table for Figure 25.

SOURCE: CDC/NCHS, National Health Interview Survey (NHIS).

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig25>

Health Insurance Coverage

During 1999 through the first 6 months of 2015 among adults aged 18–64, lack of health insurance coverage was highest among Hispanic adults.


Health insurance is a major determinant of access to health care. Children are less likely to be uninsured than adults aged 18–64 because they are more likely to qualify for public coverage, primarily Medicaid and the Children’s Health Insurance Program (CHIP) (see [data table for Figure 26](#) for estimates for children) (26,87). Passage of the Affordable Care Act (ACA) in 2010 (38) authorized states to expand Medicaid eligibility (88) and to establish the health insurance marketplace in 2014.

For adults aged 18–64, the percentage without coverage increased from 17.9% to 20.5% during 1999–2013, and then decreased to 12.7% in the first 6 months of 2015 (36). During this period, the trend for lack of coverage varied by racial and ethnic group.

During 1999–June 2015, Hispanic adults aged 18–64 had the highest percentage without coverage (27.2% in the first 6 months of 2015), and non-Hispanic white adults aged 18–64 had the lowest, except in the first 6 months of 2015, when non-Hispanic Asian adults had the lowest percentage without coverage.

The difference between the highest and lowest percentages of adults aged 18–64 without health insurance among the four racial and ethnic groups narrowed from 1999–June 2015. This difference was 24.9 percentage points in 1999 (Hispanic adults compared with non-Hispanic white adults) and 19.9 percentage points in the first 6 months of 2015 (Hispanic adults compared with non-Hispanic Asian adults).

Figure 26. No health insurance coverage among adults aged 18–64, by race and Hispanic origin: United States, 1999–June 2015 (preliminary data)



NOTES: Preliminary estimates for the first 6 months of 2015 are shown with a dashed line (36). Three-year average annual estimates for the American Indian or Alaska Native population are available in the [data table for Figure 26](#). Highest and lowest percentages are based on observed percentages and were not tested for

statistically significant differences against other percentages. Ties in highest and lowest percentages were resolved by looking at additional decimal places. See [Technical Notes](#) and [data table for Figure 26](#).

SOURCE: CDC/NCHS, National Health Interview Survey (NHIS).

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig26>

Difficulty Accessing Needed Dental Care Due to Cost


During 1999–2014 among adults aged 18–64, nonreceipt of needed dental care due to cost was lowest among non-Hispanic Asian adults.

Oral health is integral to general health and wellbeing, and forgoing needed dental health care can have serious health effects (89). In general, fewer adults have dental coverage than medical coverage, and dental coverage tends to be less comprehensive (90–92). In 2012, 44% of dental expenditures among adults aged 18–64 were paid out of pocket, a higher out-of-pocket percentage than for any other type of personal health care expenditure (93).

The percentage of adults aged 18–64 who did not receive needed dental care in the past 12 months due to cost increased from 9.3% to 17.3% during 1999–2010, and then decreased to 12.6% in 2014 (data table for Figure 27).

During 1999–2014, non-Hispanic Asian adults aged 18–64 had the lowest percentage of not receiving needed dental care due to cost (6.3% in 2014) among the four racial and ethnic groups. No racial and ethnic group consistently had the highest percentage of not receiving needed dental care due to cost during 1999–2014. The difference between the highest and lowest percentages of adults not receiving needed dental care due to cost among the four racial and ethnic groups widened during 1999–2010, and then remained stable from 2010–2014 for those aged 18–64. This difference was 5.9 percentage points in 1999 (non-Hispanic black compared with non-Hispanic Asian) and 9.4 percentage points in 2014 (Hispanic compared with non-Hispanic Asian).

Figure 27. Nonreceipt of needed dental care in the past 12 months due to cost among adults aged 18–64, by race and Hispanic origin: United States, 1999–2014



NOTES: Three-year average annual estimates for the American Indian or Alaska Native population are available in the data table for Figure 27. Highest and lowest percentages are based on observed percentages and were not tested for statistically significant differences against other percentages. See Technical Notes and data table for Figure 27.

SOURCE: CDC/NCHS, National Health Interview Survey (NHIS).

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig27>

Chartbook Data Tables

All 27 chartbook figures have an accompanying data table either in this section or the Trend Table section.

Data table for Figure 6. Selected disability indicators among adults aged 18 and over, by sex and age: United States, 2014

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig06>

Sex and age	Serious difficulty concentrating, remembering, or making decisions		Difficulty doing errands alone	
	Percent	SE	Percent	SE
Male				
18–64 years	4.4	0.2	3.1	0.2
65–74 years	6.1	0.6	6.4	0.7
75–84 years	10.8	1.3	12.9	1.3
85 years and over	18.8	2.9	26.2	3.3
Female				
18–64 years	4.5	0.2	3.9	0.2
65–74 years	5.3	0.5	9.4	0.7
75–84 years	11.2	1.1	20.7	1.4
85 years and over	21.5	2.1	45.0	2.6

SE is standard error.

NOTES: Respondents were asked, “Because of a physical, mental, or emotional condition, do you have serious difficulty concentrating, remembering, or making decisions?” See Appendix II, Instrumental activities of daily living (IADL). Proxy reporting was 3.3% for those aged 18–64, 4.8% for those aged 65–74, 6.9% for those aged 75–84, and 14.4% for those aged 85 and over. Respondents were asked, “Because of a physical, mental, or emotional condition, do you have difficulty doing errands alone such as visiting a doctor’s office or shopping?” Proxy reporting was 3.7% for those aged 18–64, 6.5% for those aged 65–74 and 75–84, and 8.8% for those aged 85 and over.

SOURCE: CDC/NCHS, National Health Interview Survey. Sample family disability questionnaire. See Appendix I, National Health Interview Survey (NHIS).

Data table for Figure 11. Distribution of emergency department visits within the past 12 months for adults aged 18–64, by type of coverage: United States, 2014

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig11>

<i>Insurance coverage</i>	<i>Percent</i>	<i>SE</i>
Private	53.6	1.0
Medicaid	23.4	0.8
Uninsured	15.1	0.7
Other	7.9	0.5

SE is standard error.

NOTES: Insurance categories are based on coverage at the time of interview and are mutually exclusive. Persons who reported both Medicaid and private coverage are classified as having private coverage. Medicaid coverage includes persons covered by state-sponsored health plans or the Children’s Health Insurance Program (CHIP). The other insured category includes military, other government, and Medicare coverage. Persons not covered by private insurance, Medicaid, or other plans are classified as uninsured. Persons with only Indian Health Service coverage are considered uninsured. The count of emergency department visits in 2014 was determined by using the midpoint of the response categories and then summing the count. Response category None was recoded to 0 visits. Response category 1 was recoded to 1 visit. Response category 2–3 was recoded to 2.5 visits. Response category 4–5 was recoded to 4.5 visits. Response category 6–7 was recoded to 6.5 visits. Response category 8–9 was recoded to 8.5 visits. Response category 10–12 was recoded to 11 visits. Response category 13–15 was recoded to 14 visits. Response category 16 or more was recoded to 16 visits. See Appendix II, Emergency department or emergency room visit; Health insurance coverage.

SOURCE: CDC/NCHS, National Health Interview Survey (NHIS). Family core and sample adult questionnaires. See Appendix I, National Health Interview Survey (NHIS).

Data table for Figure 13. Electronic health record system components in physician offices, by selected component type: United States, 2010 and 2013

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig13>

<i>Type of component</i>	<i>2010</i>		<i>2013</i>	
	<i>Percent</i>	<i>SE</i>	<i>Percent</i>	<i>SE</i>
Record patient history and demographic information	74.3	0.9	83.0	0.9
Order prescriptions	57.2	1.0	82.6	0.9
Send prescriptions to pharmacy	43.8	1.1	78.7	1.0
Issue warnings of drug interactions and contraindications	43.6	1.0	73.8	1.1
Order lab tests	48.5	1.1	68.9	1.1

SE is standard error.

NOTES: Missing values are included in the denominator. Estimates for 2010 are based on the combined in-person and mail survey file. Estimates for 2013 are based on the mail survey file. The Health Information Technology for Economic and Clinical Health (HITECH) Act authorizes Medicare and Medicaid incentive payments to providers for the “meaningful use” of EHR—that is, using EHR components to improve care. The selected components in Figure 13 are among those designated “meaningful use.” For more information see: Hsiao CJ, Hing E. Use and characteristics of electronic health record systems among office-based physician practices: United States, 2001–2013. NCHS data brief, no 143. Hyattsville, MD: NCHS; 2014. Available from: <http://www.cdc.gov/nchs/data/databriefs/db143.htm>.

SOURCE: CDC/NCHS, National Ambulatory Medical Care Survey—National Electronic Health Records Survey. See Appendix I, National Ambulatory Medical Care Survey (NAMCS).

Data table for Figure 14. Office-based physicians accepting new patients, by patient source of payment and urban–rural status: United States, 2013

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig14>

<i>Insurance coverage and urban–rural category</i>	<i>Percent</i>	<i>SE</i>
Accepting new patients	95.9	0.5
Urban:		
Large central metropolitan	97.1	0.9
Large fringe metropolitan (suburbs)	95.9	1.1
Medium or small metropolitan	94.7	0.8
Rural:		
Micropolitan (city/town)	93.6	1.6
Noncore	97.8	0.8
Accepting new Medicaid patients	69.5	1.2
Urban:		
Large central metropolitan	69.2	2.3
Large fringe metropolitan (suburbs)	58.5	2.4
Medium or small metropolitan	73.0	1.8
Rural:		
Micropolitan (city/town)	83.8	3.2
Noncore	89.6	3.3
Accepting new privately insured patients	85.2	0.9
Urban:		
Large central metropolitan	85.1	1.8
Large fringe metropolitan (suburbs)	84.4	1.8
Medium or small metropolitan	85.8	1.3
Rural:		
Micropolitan (city/town)	86.3	2.2
Noncore	84.7	3.4

SE is standard error.

NOTES: The target universe consists of physicians classified as providing direct patient care in office-based practices. Radiologists, anesthesiologists, and pathologists are excluded. Physician offices were classified by the 2013 NCHS urban–rural classification scheme for counties. The medium and small metropolitan categories were combined for this analysis. For more information, see: Ingram DD, Franco SJ. 2013 NCHS urban–rural classification scheme for counties. Vital and health statistics reports; series 2 no 166. Hyattsville, MD: NCHS. 2014. Available from: http://www.cdc.gov/nchs/data_access/urban_rural.htm. Estimates presented here may differ from estimates based on the same data presented elsewhere if different rules were used for including observations in the analysis.

SOURCE: CDC/NCHS, National Ambulatory Medical Care Survey—National Electronic Health Records Survey. See Appendix I, National Ambulatory Medical Care Survey (NAMCS).

Data table for Figure 17. Health insurance coverage among adults aged 18–64, by state Medicaid expansion status: United States, 2013 and 2014

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig17>

<i>Medicaid expansion status and insurance coverage</i>	<i>2013</i>		<i>2014</i>	
	<i>Percent</i>	<i>SE</i>	<i>Percent</i>	<i>SE</i>
States that expanded Medicaid program				
Private	65.8	0.5	68.2	0.5
Medicaid	11.9	0.3	14.9	0.4
Uninsured	18.5	0.3	13.4	0.3
States that did not expand Medicaid program				
Private	64.3	0.6	66.6	0.6
Medicaid	7.7	0.2	8.3	0.3
Uninsured	22.7	0.5	19.6	0.4

SE is standard error.

NOTES: Insurance categories are mutually exclusive. Insurance is at the time of interview. See Appendix II, Health insurance coverage. Under provisions of the Affordable Care Act (ACA) of 2010 (P.L. 111–148, P.L. 111–152), states are authorized to expand Medicaid to a new adult population. There is no deadline for states to implement the Medicaid expansion, and they may do so at any time. States were classified based on their decision to expand Medicaid as of January 1, 2014. As of January 1, 2014, 25 states and the District of Columbia have expanded their Medicaid program (40). They were: Arizona, Arkansas, California, Colorado, Connecticut, Delaware, District of Columbia, Hawaii, Illinois, Iowa, Kentucky, Maryland, Massachusetts, Michigan, Minnesota, Nevada, New Jersey, New Mexico, New York, North Dakota, Ohio, Oregon, Rhode Island, Vermont, Washington, and West Virginia. States that had not expanded their Medicaid programs as of January 1, 2014, were: Alabama, Alaska, Florida, Georgia, Idaho, Indiana, Kansas, Louisiana, Maine, Mississippi, Missouri, Montana, Nebraska, New Hampshire, North Carolina, Oklahoma, Pennsylvania, South Carolina, South Dakota, Tennessee, Texas, Utah, Virginia, Wisconsin, and Wyoming.

SOURCE: CDC/NCHS, National Health Interview Survey. See Appendix I, National Health Interview Survey (NHIS).

Data table for Figure 18 (page 1 of 2). Life Expectancy at birth, by sex, race and Hispanic origin: United States, 1980–2014

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig18>

Year	All races			White			Black or African American		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
	Life expectancy (years)								
1980.....	73.7	70.0	77.4	74.4	70.7	78.1	68.1	63.8	72.5
1981.....	74.1	70.4	77.8	74.8	71.1	78.4	68.9	64.5	73.2
1982.....	74.5	70.8	78.1	75.1	71.5	78.7	69.4	65.1	73.6
1983.....	74.6	71.0	78.1	75.2	71.6	78.7	69.4	65.2	73.5
1984.....	74.7	71.1	78.2	75.3	71.8	78.7	69.5	65.3	73.6
1985.....	74.7	71.1	78.2	75.3	71.8	78.7	69.3	65.0	73.4
1986.....	74.7	71.2	78.2	75.4	71.9	78.8	69.1	64.8	73.4
1987.....	74.9	71.4	78.3	75.6	72.1	78.9	69.1	64.7	73.4
1988.....	74.9	71.4	78.3	75.6	72.2	78.9	68.9	64.4	73.2
1989.....	75.1	71.7	78.5	75.9	72.5	79.2	68.8	64.3	73.3
1990.....	75.4	71.8	78.8	76.1	72.7	79.4	69.1	64.5	73.6
1991.....	75.5	72.0	78.9	76.3	72.9	79.6	69.3	64.6	73.8
1992.....	75.8	72.3	79.1	76.5	73.2	79.8	69.6	65.0	73.9
1993.....	75.5	72.2	78.8	76.3	73.1	79.5	69.2	64.6	73.7
1994.....	75.7	72.4	79.0	76.5	73.3	79.6	69.5	64.9	73.9
1995.....	75.8	72.5	78.9	76.5	73.4	79.6	69.6	65.2	73.9
1996.....	76.1	73.1	79.1	76.8	73.9	79.7	70.2	66.1	74.2
1997.....	76.5	73.6	79.4	77.1	74.3	79.9	71.1	67.2	74.7
1998.....	76.7	73.8	79.5	77.3	74.5	80.0	71.3	67.6	74.8
1999.....	76.7	73.9	79.4	77.3	74.6	79.9	71.4	67.8	74.7
2000.....	76.8	74.1	79.3	77.3	74.7	79.9	71.8	68.2	75.1
2001.....	77.0	74.3	79.5	77.5	74.9	80.0	72.0	68.5	75.3
2002.....	77.0	74.4	79.6	77.5	74.9	80.1	72.2	68.7	75.4
2003.....	77.2	74.5	79.7	77.7	75.1	80.2	72.4	68.9	75.7
2004.....	77.6	75.0	80.1	78.1	75.5	80.5	72.9	69.4	76.1
2005.....	77.6	75.0	80.1	78.0	75.5	80.5	73.0	69.5	76.2
2006.....	77.8	75.2	80.3	78.3	75.8	80.7	73.4	69.9	76.7
2007.....	78.1	75.5	80.6	78.5	76.0	80.9	73.8	70.3	77.0
2008.....	78.2	75.6	80.6	78.5	76.1	80.9	74.3	70.9	77.3
2009.....	78.5	76.0	80.9	78.8	76.4	81.2	74.7	71.4	77.7
2010.....	78.7	76.2	81.0	78.9	76.5	81.3	75.1	71.8	78.0
2011.....	78.7	76.3	81.1	79.0	76.6	81.3	75.3	72.2	78.2
2012.....	78.8	76.4	81.2	79.1	76.7	81.4	75.5	72.3	78.4
2013.....	78.8	76.4	81.2	79.1	76.7	81.4	75.5	72.3	78.4
2014.....	78.8	76.4	81.2	79.0	76.7	81.4	75.6	72.5	78.4

See footnotes at end of table.

Data table for Figure 18 (page 2 of 2). Life Expectancy at birth, by sex, race and Hispanic origin: United States, 1980–2014

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig18>

Year	Not Hispanic or Latino								
	Hispanic ¹			White			Black or African American		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
2014.....	81.8	79.2	84.0	78.8	76.5	81.1	75.2	72.0	78.1

¹Persons of Hispanic origin may be of any race. See Appendix II, Hispanic origin. Life expectancies for the Hispanic population are adjusted for underreporting of Hispanic ethnicity on the death certificate, but are not adjusted to account for the potential effects of return migration. To address the effects of age misstatement at the oldest ages, the probability of death for Hispanic persons ages 80 and over is estimated as a function of non-Hispanic white mortality with the use of the Brass relational logit model. See Appendix II, Race, for a discussion of sources of bias in death rates by race and Hispanic origin.

NOTES: Populations for computing life expectancy for 1991–1999 are 1990-based postcensal estimates of the U.S. resident population. Populations for computing life expectancy for 2001–2009 were based on intercensal population estimates of the U.S. resident population. Populations for computing life expectancy for 2010 were based on 2010 census counts. Life expectancy for 2011 and beyond was computed using 2010-based postcensal estimates. See Appendix I, Population Census and Population Estimates. In 1997, life table methodology was revised to construct complete life tables by single years of age that extend to age 100. (Anderson RN. Method for constructing complete annual U.S. life tables. NCHS. Vital Health Stat 2(129). 1999.) Previously, abridged life tables were constructed for 5-year age groups ending with 85 and over. In 2000, the life table methodology was revised. The revised methodology is similar to that developed for the 1999–2001 decennial life tables. In 2008, the life table methodology was further refined. See Appendix II, Life expectancy. Starting with 2003 data, some states allowed the reporting of more than one race on the death certificate. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards, for comparability with other states. The race groups, white and black include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. See Appendix II, Race. Life expectancy is not currently available for persons of other racial and ethnic groups. Also see Table 15 and Figure 1.

SOURCE: CDC/NCHS, National Vital Statistics System, public-use Mortality Files; Arias E. United States life tables by Hispanic origin. Vital health statistics; vol 2 no 152. Hyattsville, MD: NCHS. 2010. NCHS. Deaths: Final data for 2014. National vital statistics reports (forthcoming). Available from: <http://www.cdc.gov/nchs/products/nvsr.htm>. See Appendix I, National Vital Statistics System (NVSS).

Data table for Figure 19. Infant mortality rates, by race and Hispanic origin of mother: United States, 1999–2013

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig19>

Year	Race and Hispanic origin of mother ¹					
	Total ²	Hispanic or Latina	Not Hispanic or Latina			
			White	Black or African American	Asian or Pacific Islander	American Indian or Alaska Native
Infant deaths per 1,000 live births ³						
1999	7.04	5.71	5.76	14.14	4.73	9.35
2000	6.89	5.59	5.70	13.59	4.79	8.19
2001	6.84	5.44	5.72	13.46	4.65	9.67
2002	6.95	5.62	5.80	13.89	4.66	8.67
2003	6.84	5.65	5.70	13.60	4.68	8.72
2004	6.78	5.55	5.66	13.60	4.55	8.62
2005	6.86	5.62	5.76	13.63	4.77	8.31
2006	6.68	5.41	5.58	13.35	4.40	8.64
2007	6.75	5.51	5.63	13.32	4.60	9.38
2008	6.61	5.59	5.53	12.67	4.39	8.66
2009	6.39	5.29	5.33	12.40	4.28	9.17
2010	6.14	5.25	5.18	11.46	4.17	8.65
2011	6.07	5.15	5.07	11.45	4.18	8.52
2012	5.98	5.11	5.04	11.19	3.97	8.74
2013	5.96	5.00	5.06	11.11	3.90	7.72

Year	Detailed Hispanic origin of mother ¹				
	Mexican	Puerto Rican	Cuban	Central and South American	Other and unknown Hispanic or Latina
Infant deaths per 1,000 live births ³					
1999	5.51	8.35	4.64	4.67	7.24
2000	5.43	8.20	4.57	4.64	6.88
2001	5.22	8.53	4.25	4.97	6.02
2002	5.42	8.19	3.74	5.06	7.15
2003	5.49	8.18	4.59	5.04	6.66
2004	5.47	7.82	4.57	4.65	6.72
2005	5.53	8.31	4.45	4.69	6.44
2006	5.34	8.02	5.06	4.52	5.78
2007	5.42	7.72	5.21	4.57	6.41
2008	5.58	7.29	4.88	4.76	5.86
2009	5.12	7.19	5.75	4.47	6.06
2010	5.12	7.09	3.81	4.43	6.09
2011	4.99	7.84	4.34	4.35	5.41
2012	5.02	6.86	4.99	4.14	5.59
2013	4.90	5.92	3.04	4.30	5.88

¹Persons of Hispanic origin may be of any race. Starting with 2003 data, some states reported multiple-race data. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards, for comparability with other states. See Appendix II, Hispanic origin; Race.

²Includes all infant deaths not shown separately.

³Infant is under age 1 year.

NOTES: Rates based on a period file using weighted data. Also see Table 10.

SOURCE: CDC/NCHS, National Vital Statistics System, public-use Linked Birth/Infant Death Data Set. See Appendix I, National Vital Statistics System (NVSS).

Data table for Figure 20. Preterm births, by gestational age and race and Hispanic origin and detailed Hispanic origin of mother: United States, 2014

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig20>

Preterm births by gestational age, in weeks ²	Total ³	Race and Hispanic origin of mother ¹				
		Hispanic or Latina	White	Not Hispanic or Latina		
				Black or African American	Asian or Pacific Islander	American Indian or Alaska Native
Percent of live singleton births that were preterm						
Less than 37	7.7	7.7	6.9	11.1	6.8	9.0
34–36	5.7	5.7	5.3	7.2	5.2	6.8
32–33	0.8	0.8	0.7	1.3	0.7	1.0
Less than 32	1.2	1.2	0.9	2.6	0.9	1.2
Standard error						
Less than 37	0.01	0.03	0.02	0.04	0.05	0.15
34–36	0.01	0.02	0.02	0.03	0.04	0.13
32–33	0.00	0.01	0.01	0.02	0.02	0.05
Less than 32	0.01	0.01	0.01	0.02	0.02	0.06

Preterm births by gestational age, in weeks ²	Mexican	Detailed Hispanic origin of mother ¹			
		Puerto Rican	Cuban	Central and South American	Other and unknown Hispanic or Latina
Percent of live singleton births that were preterm					
Less than 37	7.6	9.1	7.2	7.2	8.3
34–36	5.6	6.4	5.1	5.3	6.1
32–33	0.8	1.0	0.9	0.8	0.9
Less than 32	1.1	1.7	1.2	1.1	1.3
Standard error					
Less than 37	0.04	0.11	0.18	0.07	0.07
34–36	0.03	0.09	0.16	0.06	0.06
32–33	0.01	0.04	0.07	0.02	0.03
Less than 32	0.01	0.05	0.08	0.03	0.03

0.00 Quantity more than zero but less than 0.005.

¹Persons of Hispanic origin may be of any race. Starting with 2003 data, some states reported multiple-race data. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards, for comparability with other states. See Appendix II, Hispanic origin; Race.

²Preterm births are based on the obstetric estimate of gestational age and are for all singleton births. For more information on the obstetric estimates, see: Martin JA, Osterman MJK, Kirmeyer SE, Gregory ECW. Measuring gestational age in vital statistics data: Transitioning to the obstetric estimate. National vital statistics reports; vol 64 no 5. Hyattsville, MD: NCHS. 2015. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr64/nvsr64_05.pdf.

³Includes all preterm births not shown separately.

NOTES: Ties in highest and lowest rates were resolved by looking at additional decimal places. See Technical Notes.

SOURCE: CDC/NCHS, National Vital Statistics System, public-use Birth File. See Appendix I, National Vital Statistics System (NVSS).

Data table for Figure 21 (page 1 of 2). Low-risk births delivered by cesarean section, by race and Hispanic origin and detailed Hispanic origin of mother: United States, 1999–2014

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig21>

Year	Race and Hispanic origin of mother ¹					
	Total ²	Not Hispanic or Latina				
		Hispanic or Latina	White	Black or African American	Asian or Pacific Islander	American Indian or Alaska Native
Percent of low-risk births delivered by cesarean section ³						
1999.....	19.5	18.7	19.2	21.9	19.6	17.1
2000.....	20.3	19.4	20.1	23.0	20.0	18.0
2001.....	21.6	20.6	21.4	24.3	21.9	18.8
2002.....	23.1	21.9	22.8	25.9	23.2	20.4
2003.....	24.3	23.2	24.0	27.4	24.9	21.1
2004.....	25.7	24.6	25.4	28.8	26.6	22.2
2005.....	26.6	25.1	26.4	29.8	27.8	23.2
2006.....	27.1	25.3	27.0	30.1	27.7	24.5
2007.....	27.5	26.0	27.4	30.5	28.3	24.5
2008.....	27.8	26.2	27.5	30.7	29.1	23.1
2009.....	28.1	27.0	27.7	31.1	29.6	24.1
2010.....	27.6	26.7	27.0	31.0	29.0	23.8
2011.....	27.3	26.6	26.6	30.9	28.8	22.1
2012.....	27.3	26.8	26.5	31.0	28.5	23.5
2013.....	26.9	26.6	25.9	30.8	28.6	23.0
2014 ⁴	26.0	25.8	25.0	29.9	27.6	21.5
Standard error						
1999.....	0.03	0.08	0.04	0.10	0.15	0.37
2000.....	0.03	0.08	0.04	0.10	0.14	0.37
2001.....	0.04	0.08	0.05	0.10	0.15	0.37
2002.....	0.04	0.08	0.05	0.10	0.15	0.38
2003.....	0.04	0.08	0.05	0.11	0.15	0.38
2004.....	0.04	0.08	0.05	0.11	0.15	0.39
2005.....	0.04	0.08	0.05	0.11	0.15	0.39
2006.....	0.04	0.08	0.05	0.11	0.15	0.39
2007.....	0.04	0.08	0.05	0.11	0.15	0.39
2008.....	0.04	0.08	0.05	0.11	0.15	0.38
2009.....	0.04	0.08	0.05	0.11	0.15	0.39
2010.....	0.04	0.09	0.05	0.11	0.15	0.39
2011.....	0.04	0.09	0.05	0.11	0.15	0.39
2012.....	0.04	0.09	0.05	0.11	0.15	0.40
2013.....	0.04	0.09	0.05	0.11	0.15	0.40
2014 ⁴	0.04	0.09	0.05	0.11	0.14	0.40

See footnotes at end of table.

Data table for Figure 21 (page 2 of 2). Low-risk births delivered by cesarean section, by race and Hispanic origin and detailed Hispanic origin of mother: United States, 1999–2014

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig21>

Year	Detailed Hispanic origin of mother ¹				
	Mexican	Puerto Rican	Cuban	Central and South American	Other and unknown Hispanic or Latina
Percent of low-risk births delivered by cesarean section ³					
1999	18.0	19.2	29.7	20.4	18.4
2000	18.6	20.5	30.7	21.2	19.1
2001	19.8	21.3	31.5	22.1	20.7
2002	21.1	23.2	32.8	23.5	21.7
2003	22.3	24.3	36.6	25.0	22.6
2004	23.5	26.8	38.5	26.8	23.7
2005	23.5	27.3	42.1	27.6	26.4
2006	23.3	28.8	44.8	27.7	27.5
2007	24.0	28.8	45.3	28.6	28.1
2008	24.5	29.0	45.4	28.6	26.8
2009	24.9	29.4	44.2	30.0	28.2
2010	24.6	29.4	42.0	29.5	28.0
2011	24.8	28.9	42.0	29.1	26.8
2012	25.0	28.9	43.3	29.4	27.1
2013	24.7	28.3	42.3	29.1	27.2
2014 ⁴	24.1	27.3	41.4	27.9	26.0
Standard error					
1999	0.09	0.29	0.66	0.22	0.30
2000	0.09	0.30	0.65	0.21	0.31
2001	0.09	0.30	0.63	0.21	0.32
2002	0.09	0.31	0.63	0.21	0.32
2003	0.10	0.31	0.63	0.20	0.33
2004	0.10	0.32	0.64	0.20	0.34
2005	0.10	0.31	0.64	0.20	0.32
2006	0.10	0.31	0.63	0.20	0.30
2007	0.10	0.30	0.62	0.20	0.28
2008	0.10	0.30	0.62	0.21	0.23
2009	0.10	0.30	0.62	0.22	0.23
2010	0.11	0.31	0.62	0.22	0.23
2011	0.11	0.30	0.61	0.23	0.21
2012	0.11	0.31	0.62	0.24	0.21
2013	0.11	0.30	0.58	0.24	0.22
2014 ⁴	0.11	0.30	0.55	0.22	0.21

¹Persons of Hispanic origin may be of any race. Starting with 2003 data, some states reported multiple-race data. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards, for comparability with other states. See Appendix II, Hispanic origin; Race.

²Includes all low-risk cesarean section births not shown separately.

³Low-risk cesarean delivery is defined as singleton, term (37 or more weeks of gestation by last menstrual period estimate for data years 1999–2013 and the obstetric estimate for 2014), vertex (not breech) cesarean delivery to women having a first birth per 100 women delivering singleton, term, vertex, first births.

⁴For 2014, the definition of term birth was based on the obstetric estimate of gestational age. For more information on the obstetric estimate, see: Martin JA, Osterman MJK, Kirmeyer SE, Gregory ECW. Measuring gestational age in vital statistics data: Transitioning to the obstetric estimate. National vital statistics reports; vol 64 no 5. Hyattsville, MD: NCHS. 2015. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr64/nvsr64_05.pdf. Use of the obstetric estimate instead of the last menstrual period had a statistically significant but small impact on the percentage of women with low-risk cesarean section births.

NOTES: Ties in highest and lowest rates were resolved by looking at additional decimal places. See Technical Notes.

SOURCE: CDC/NCHS, National Vital Statistics System, public-use Birth File. See Appendix I, National Vital Statistics System (NVSS).

Data table for Figure 22. Obesity among children and adolescents aged 2–19, by age and race and Hispanic origin: United States, 2011–2014

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig22>

Age	Total ²	Race and Hispanic origin ¹			
		Hispanic or Latino	Not Hispanic or Latino		
			White only	Black or African American only	Asian only
Percent with obesity ³					
2–19.	17.0	21.9	14.7	19.5	8.6
2–5 years.	8.9	15.6	*5.2	10.4	*
6–11 years.	17.5	25.0	13.6	21.4	*9.8
12–19 years.	20.5	22.8	19.6	22.6	9.4
Standard error					
2–19.	0.7	0.9	1.2	1.2	1.1
2–5 years.	0.9	1.6	1.2	1.5	*
6–11 years.	1.2	1.4	2.0	2.0	2.0
12–19 years.	1.4	1.6	2.5	2.3	1.6

¹Persons of Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity*. The three non-Hispanic race categories shown in the table conform to the 1997 Standards. Race-specific estimates are for persons who reported only one racial group. See Appendix II, Hispanic origin; Race.

²Includes all persons not shown separately.

³Obesity is defined as a body mass index at or above the sex- and age-specific 95th percentile of the CDC growth charts. Pregnant women are excluded.

NOTE: Also see Table 59 and Figure 8.

SOURCE: CDC/NCHS, National Health and Nutrition Examination Survey. See Appendix I, National Health and Nutrition Examination Survey (NHANES).

Data table for Figure 23 (page 1 of 2). Hypertension among adults aged 20 and over, by sex and race and Hispanic origin: United States, 1999–2000 through 2013–2014

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig23>

Characteristic	1999–2000	2001–2002	2003–2004	2005–2006	2007–2008	2009–2010	2011–2012	2013–2014
	Percent with hypertension ¹							
Both sexes								
20 years and over, crude ²	28.9	28.9	32.5	31.7	32.6	31.9	32.5	33.5
20 years and over, age-adjusted ^{2,3}	30.0	29.7	32.1	30.5	31.2	30.0	30.0	30.8
Race and Hispanic origin ^{3,4}								
Mexican origin	29.4	25.2	29.9	24.4	28.9	28.2	28.0	28.8
Not Hispanic or Latino:								
White only	28.4	27.9	30.8	29.1	30.7	28.6	28.6	29.6
Black only	40.9	43.4	42.6	44.1	42.4	42.9	43.9	42.7
	Standard error							
Both sexes								
20 years and over, crude ²	1.5	1.3	1.3	1.2	0.9	1.3	1.5	1.0
20 years and over, age-adjusted ^{2,3}	1.4	1.0	1.0	1.2	0.7	0.8	0.7	0.8
Race and Hispanic origin ^{3,4}								
Mexican origin	1.5	1.1	2.0	1.6	1.4	1.1	2.5	1.5
Not Hispanic or Latino:								
White only	1.7	1.1	1.2	1.3	1.0	1.1	0.8	0.9
Black only	1.1	1.9	1.8	1.7	1.8	1.6	1.0	1.5

See footnotes at end of table.

Data table for Figure 23 (page 2 of 2). Hypertension among adults aged 20 and over, by sex and race and Hispanic origin: United States, 1999–2000 through 2013–2014

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig23>

Characteristic	2011–2014	
	Percent	Standard error
Both sexes		
20 years and over, crude ²	33.0	0.9
20 years and over, age-adjusted ^{2,3}	30.4	0.5
Race and Hispanic origin ^{3,4}		
Hispanic or Latino	28.2	1.1
Not Hispanic or Latino:		
White only	29.1	0.6
Black only	43.3	0.9
Asian only	26.5	1.1
Men		
20 years and over, crude ²	32.6	1.1
20 years and over, age-adjusted ^{2,3}	31.0	0.7
Race and Hispanic origin ^{3,4}		
Hispanic or Latino	27.7	1.5
Not Hispanic or Latino:		
White only	30.2	1.0
Black only	42.4	1.2
Asian only	28.0	2.0
Women		
20 years and over, crude ²	33.4	1.0
20 years and over, age-adjusted ^{2,3}	29.7	0.8
Race and Hispanic origin ^{3,4}		
Hispanic or Latina	28.6	1.1
Not Hispanic or Latina:		
White only	28.0	0.8
Black only	44.0	1.4
Asian only	25.0	1.0

¹Hypertension is having measured high blood pressure (systolic pressure of at least 140 mm Hg or diastolic pressure of at least 90 mm Hg) and/or respondent report of taking antihypertensive medication. Excludes pregnant women.

²Includes all persons not shown separately.

³Estimates are age-adjusted to the year 2000 standard population using five age groups: 20–34 years, 35–44 years, 45–54 years, 55–64 years, and 65 years and over. Age-adjusted estimates in this table may differ from other age-adjusted estimates based on the same data and presented elsewhere if different age groups are used in the adjustment procedure. See Appendix II, Age adjustment.

⁴Persons of Mexican or Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The three non-Hispanic race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group. Data for Hispanic adults became available in 2007–2008 and for Asian adults starting in 2011–2012. See Appendix II, Hispanic origin; Race.

NOTE: Also see Table 54.

SOURCE: CDC/NCHS, National Health and Nutrition Examination Survey. See Appendix I, National Health and Nutrition Examination Survey (NHANES).

Data table for Figure 24 (page 1 of 3). Current cigarette smoking among adults aged 18 and over, by sex and race and Hispanic origin: United States, 1999–2014

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig24>

Sex, age, and year	Race and Hispanic origin ¹					
	Total ²		Not Hispanic or Latino ³			
	Crude	Age-adjusted ³	Hispanic or Latino ³	White only	Black only	Asian only
Men, 18 years and over			Percent current smokers ⁴			
1999	25.7	25.2	22.7	25.4	28.4	23.5
2000	25.6	25.2	23.5	25.8	25.7	19.5
2001	25.1	24.6	20.7	25.4	27.7	18.7
2002	25.1	24.6	21.4	25.5	26.7	17.5
2003	24.1	23.7	21.2	24.6	25.2	16.6
2004	23.4	23.0	17.9	24.2	23.5	16.7
2005	23.9	23.4	19.6	24.2	26.1	20.4
2006	23.9	23.6	19.3	24.6	26.8	15.6
2007	22.3	22.0	17.4	23.6	23.7	14.5
2008	23.1	22.8	19.1	23.9	24.9	15.1
2009	23.5	23.2	17.7	25.0	23.3	16.4
2010	21.5	21.2	15.2	23.0	23.7	14.6
2011	21.6	21.2	16.5	22.6	23.6	14.2
2012	20.5	20.6	16.9	22.0	22.1	15.6
2013	20.5	20.5	16.7	21.8	21.7	14.6
2014	18.8	19.0	13.8	20.1	22.0	14.0
Women, 18 years and over						
1999	21.5	21.6	11.9	23.8	20.5	*6.3
2000	20.9	21.1	12.9	23.1	20.8	7.4
2001	20.6	20.7	11.5	23.5	17.8	*6.0
2002	19.8	20.0	10.8	22.5	18.5	6.8
2003	19.2	19.4	10.4	22.1	18.1	6.0
2004	18.5	18.7	10.6	21.2	16.8	*4.9
2005	18.1	18.3	10.9	20.8	17.1	*5.9
2006	18.0	18.1	9.9	20.6	18.8	4.5
2007	17.4	17.5	8.4	20.7	15.5	*3.8
2008	18.3	18.5	10.4	21.5	17.4	4.7
2009	17.9	18.1	9.5	20.7	18.9	7.3
2010	17.3	17.5	9.1	20.4	16.8	4.3
2011	16.5	16.8	8.3	19.8	15.3	5.5
2012	15.8	15.9	7.5	19.2	14.3	5.4
2013	15.3	15.5	6.8	18.6	15.1	4.6
2014	14.8	15.1	7.4	18.3	13.7	5.1

See footnotes at end of table.

Data table for Figure 24 (page 2 of 3). Current cigarette smoking among adults aged 18 and over, by sex and race and Hispanic origin: United States, 1999–2014

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig24>

Sex, age, and year	Race and Hispanic origin ¹					
	Total ²		Not Hispanic or Latino ³			
	Crude	Age-adjusted ³	Hispanic or Latino ³	White only	Black only	Asian only
Men, 18 years and over			Standard error			
1999	0.5	0.5	1.1	0.6	1.5	2.9
2000	0.5	0.4	1.1	0.5	1.2	2.1
2001	0.4	0.4	1.0	0.5	1.3	2.4
2002	0.5	0.4	1.0	0.5	1.3	2.1
2003	0.4	0.4	1.1	0.5	1.2	2.1
2004	0.4	0.4	0.9	0.5	1.2	2.1
2005	0.5	0.5	0.9	0.6	1.3	2.3
2006	0.5	0.5	1.2	0.7	1.6	1.6
2007	0.6	0.5	1.3	0.7	1.4	1.6
2008	0.6	0.6	1.3	0.7	1.5	1.9
2009	0.5	0.5	0.9	0.7	1.2	1.5
2010	0.5	0.5	0.9	0.6	1.1	1.6
2011	0.4	0.4	0.9	0.6	1.1	1.3
2012	0.4	0.4	0.9	0.6	1.1	1.4
2013	0.5	0.5	1.0	0.6	1.2	1.6
2014	0.4	0.4	0.8	0.6	1.1	1.6
Women, 18 years and over						
1999	0.4	0.4	0.7	0.5	0.9	1.3
2000	0.4	0.4	0.8	0.5	0.9	1.4
2001	0.4	0.4	0.7	0.5	0.8	1.2
2002	0.4	0.4	0.7	0.5	0.9	1.3
2003	0.4	0.4	0.7	0.5	0.9	1.1
2004	0.4	0.4	0.7	0.5	0.9	1.1
2005	0.4	0.4	0.7	0.5	0.9	1.2
2006	0.4	0.4	0.8	0.6	1.0	0.8
2007	0.5	0.5	0.7	0.6	0.9	0.8
2008	0.5	0.5	0.8	0.7	1.0	0.8
2009	0.4	0.4	0.7	0.6	1.1	1.3
2010	0.4	0.4	0.6	0.6	0.9	0.7
2011	0.4	0.4	0.6	0.5	0.8	0.8
2012	0.4	0.4	0.6	0.5	0.7	0.8
2013	0.4	0.4	0.5	0.6	0.8	0.8
2014	0.4	0.4	0.5	0.7	0.8	0.9

See footnotes at end of table.

Data table for Figure 24 (page 3 of 3). Current cigarette smoking among adults aged 18 and over, by sex and race and Hispanic origin: United States, 1999–2014

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig24>

Sex, age, and year ⁵	Race and Hispanic origin ¹					
	Not Hispanic or Latino ³					
	Hispanic or Latino ³	White only	Black only	Asian only	American Indian or Alaska Native only	2 or more races
Men, 18 years and over			Percent current smokers ⁴			
1999–2001	22.2	25.5	27.2	20.1	32.6	35.2
2003–2005	19.5	24.3	24.9	18.1	36.4	31.4
2006–2008	18.6	24.0	25.1	15.2	39.2	24.8
2009–2011	16.4	23.5	23.5	15.0	31.1	28.9
2012–2014	15.7	21.3	21.9	14.8	28.0	29.8
Women, 18 years and over						
1999–2001	12.1	23.5	19.7	6.6	36.3	31.6
2003–2005	10.6	21.4	17.3	5.6	29.0	27.0
2006–2008	9.5	20.9	17.2	4.3	28.2	25.9
2009–2011	9.0	20.3	17.0	5.6	26.2	24.8
2012–2014	7.3	18.7	14.4	5.0	24.0	25.1
Men, 18 years and over			Standard error			
1999–2001	0.6	0.3	0.8	1.4	4.4	2.6
2003–2005	0.6	0.3	0.8	1.3	3.8	2.7
2006–2008	0.7	0.4	0.9	1.0	5.7	2.7
2009–2011	0.5	0.4	0.7	0.9	4.6	2.6
2012–2014	0.5	0.4	0.7	0.9	3.5	2.2
Women, 18 years and over						
1999–2001	0.4	0.3	0.5	0.7	3.5	2.5
2003–2005	0.4	0.3	0.5	0.7	3.6	2.4
2006–2008	0.4	0.4	0.6	0.5	3.8	2.4
2009–2011	0.4	0.4	0.6	0.6	3.4	2.0
2012–2014	0.3	0.3	0.5	0.5	3.2	2.1

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error of 20%–30%.

¹Persons of Hispanic origin may be of any race. Race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity*. The single-race categories plus multiple-race category shown in the table conform to the 1997 Standards. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

²Includes all persons not shown separately.

³Estimates are age-adjusted to the year 2000 standard population using five age groups: 18–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over. Age-adjusted estimates in this table may differ from other age-adjusted estimates based on the same data and presented elsewhere if different age groups are used in the adjustment procedure. See Appendix II, Age adjustment.

⁴Current cigarette smokers are defined as ever smoking 100 cigarettes in their lifetime and now smoke every day or some days. See Appendix II, Cigarette smoking.

⁵Three-year average annual estimates are shown in order to present estimates for the American Indian or Alaska Native and the multiple-race populations. Annual estimates are not stable for smaller population groups.

NOTES: Ties in highest and lowest rates were resolved by looking at additional decimal places. See Technical Notes. Also see Figure 7.

SOURCE: CDC/NCHS, National Health Interview Survey. Family core and sample adult questionnaires. See Appendix I, National Health Interview Survey (NHIS).

Data table for Figure 25 (page 1 of 3). Influenza vaccination among adults aged 18 and over, by age and race and Hispanic origin: United States, 1999–2014

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig25>

Age and year	Total ²	Race and Hispanic origin ¹			
		Hispanic or Latino	Not Hispanic or Latino		
			White only	Black only	Asian only
18–64 years		Percent with influenza vaccination in the past year ³			
1999	20.6	15.0	21.9	17.7	19.5
2000	21.4	14.6	23.2	16.2	23.8
2001	19.3	13.3	20.8	16.8	18.3
2002	20.8	14.3	22.2	18.4	21.7
2003	22.1	14.1	24.1	19.3	20.1
2004	22.7	14.3	25.3	17.0	23.3
2005	14.1	9.5	15.5	12.4	11.3
2006	20.5	13.5	22.4	18.1	21.1
2007	23.0	16.2	25.3	17.9	24.0
2008	25.8	17.6	28.2	22.3	25.8
2009	28.3	20.7	30.7	24.4	31.4
2010	30.1	23.8	32.2	24.6	35.0
2011	31.9	26.1	34.0	27.7	34.7
2012	31.5	25.2	33.4	27.5	37.5
2013	35.0	26.1	37.8	29.6	40.9
2014	35.8	27.9	38.5	30.8	41.3
65 years and over					
1999	65.7	55.1	67.9	49.7	71.7
2000	64.4	55.7	66.6	48.0	58.6
2001	63.1	51.8	65.4	47.9	58.6
2002	65.7	48.5	68.7	49.4	58.4
2003	65.5	45.4	68.6	47.8	63.3
2004	64.6	54.6	67.3	45.6	59.0
2005	59.7	41.7	63.2	39.6	58.9
2006	64.3	44.9	67.4	47.1	67.9
2007	66.7	52.3	69.4	55.4	63.4
2008	67.2	54.9	70.0	50.9	68.5
2009	66.8	57.0	69.0	52.9	68.9
2010	63.9	54.6	65.9	52.8	67.3
2011	66.9	57.3	69.0	53.4	69.5
2012	66.5	57.8	68.9	53.2	65.2
2013	67.9	57.2	70.2	55.9	70.0
2014	70.1	60.8	72.4	57.4	72.7

See footnotes at end of table.

Data table for Figure 25 (page 2 of 3). Influenza vaccination among adults aged 18 and over, by age and race and Hispanic origin: United States, 1999–2014

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig25>

Age and year	Total ²	Race and Hispanic origin ¹			
		Hispanic or Latino	Not Hispanic or Latino		
			White only	Black only	Asian only
Standard error					
18–64 years					
1999	0.3	0.7	0.4	0.8	1.7
2000	0.3	0.7	0.4	0.7	1.9
2001	0.3	0.6	0.4	0.8	1.5
2002	0.3	0.7	0.4	0.8	1.7
2003	0.3	0.6	0.4	0.8	1.6
2004	0.3	0.6	0.4	0.8	1.7
2005	0.3	0.5	0.3	0.7	1.3
2006	0.4	0.7	0.5	0.9	1.5
2007	0.4	0.8	0.6	0.9	1.8
2008	0.4	0.8	0.5	1.0	1.7
2009	0.4	0.8	0.5	1.1	1.8
2010	0.4	0.8	0.6	0.9	1.6
2011	0.4	0.8	0.5	0.9	1.4
2012	0.4	0.8	0.5	0.9	1.6
2013	0.4	0.8	0.5	1.0	1.6
2014	0.4	0.9	0.6	1.0	1.6
65 years and over					
1999	0.8	2.7	0.8	2.3	5.9
2000	0.7	2.6	0.8	2.3	5.9
2001	0.7	2.6	0.8	2.4	6.2
2002	0.7	2.7	0.8	2.3	5.5
2003	0.7	2.7	0.8	2.3	6.0
2004	0.7	2.7	0.8	2.4	6.0
2005	0.8	2.8	0.8	2.1	4.7
2006	0.9	2.9	1.1	2.3	3.9
2007	0.8	3.1	1.0	2.3	4.5
2008	0.9	3.1	1.0	2.5	3.7
2009	0.8	2.9	0.9	2.3	3.3
2010	0.8	2.6	0.9	2.1	3.5
2011	0.7	2.2	0.8	1.8	3.4
2012	0.8	2.3	0.9	2.1	3.2
2013	0.7	2.2	0.9	2.0	3.4
2014	0.7	2.2	0.8	2.0	2.8

See footnotes at end of table.

Data table for Figure 25 (page 3 of 3). Influenza vaccination among adults aged 18 and over, by age and race and Hispanic origin: United States, 1999–2014

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig25>

Age and year ⁴	Race and Hispanic origin ¹					
	Not Hispanic or Latino					
	Hispanic or Latino	White only	Black only	Asian only	American Indian or Alaska Native	2 or more races
Percent with influenza vaccination in the past year ³						
18–64 years						
1999–2001	14.3	22.0	16.9	20.5	24.1	20.2
2003–2005	12.6	21.6	16.2	18.1	20.9	20.8
2006–2008	15.8	25.3	19.4	23.6	26.8	24.5
2009–2011	23.6	32.3	25.5	33.7	33.4	24.8
2012–2014	26.4	36.5	29.3	39.9	43.0	30.8
65 years and over						
1999–2001	54.2	66.6	48.5	62.8	54.9	67.5
2003–2005	47.2	66.4	44.3	60.2	60.4	63.6
2006–2008	50.9	68.9	51.1	66.8	69.9	59.2
2009–2011	56.3	68.0	53.0	68.6	63.0	71.9
2012–2014	58.7	70.6	55.6	69.5	65.1	56.5
Standard error						
18–64 years						
1999–2001	0.4	0.2	0.4	1.0	2.2	1.7
2003–2005	0.4	0.2	0.4	0.9	2.4	1.7
2006–2008	0.5	0.4	0.5	1.0	4.5	2.1
2009–2011	0.5	0.3	0.5	0.9	3.1	1.7
2012–2014	0.5	0.3	0.5	0.9	3.1	1.7
65 years and over						
1999–2001	1.6	0.5	1.4	3.6	8.1	5.2
2003–2005	1.6	0.5	1.4	3.2	7.7	5.3
2006–2008	1.7	0.6	1.4	2.4	6.5	5.2
2009–2011	1.5	0.5	1.2	2.1	7.9	4.4
2012–2014	1.3	0.5	1.2	1.9	6.1	4.1

¹Persons of Hispanic origin may be of any race. Race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity*. The single-race categories plus multiple-race category shown in the table conform to the 1997 Standards. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

²Includes all persons not shown separately.

³Influenza vaccination is based on respondent report of receipt of a seasonal flu shot or influenza nasal spray (starting in 2005). Questions concerning use of influenza vaccination differed slightly on the National Health Interview Survey across the years for which data are shown. See Appendix II, Vaccination.

⁴Three-year average annual estimates are shown in order to present estimates for the American Indian or Alaska Native and the multiple-race populations. Annual estimates are not stable for smaller population groups.

NOTES: Also see Table 68. Ties in highest and lowest rates were resolved by looking at additional decimal places. See Technical Notes. Prevalence of influenza vaccination during the past 12 months is different from season-specific coverage, see: CDC. Surveillance of influenza vaccination coverage—United States, 2007–08 through 2011–12 influenza seasons. MMWR 2013;62(ss04):1–29. Available from: http://www.cdc.gov/mmwr/preview/mmwrhtml/ss6204a1.htm?s_cid=ss6204a1_w; and CDC. FluVaxView. Available from: <http://www.cdc.gov/flu/fluview/>. The recommendations of the Advisory Committee on Immunization Practices regarding who should receive an influenza vaccination have changed over the years, and changes in coverage estimates may reflect changes in recommendations. An influenza vaccine shortage occurred during the 2004–2005 influenza season. Delays in the availability of influenza shots also occurred in fall 2000 and, to a lesser extent, in fall 2001.

SOURCE: CDC/NCHS, National Health Interview Survey. Family core and sample adult questionnaires. See Appendix I, National Health Interview Survey (NHIS).

Data table for Figure 26 (page 1 of 3). No health insurance coverage among persons under age 65, by age and race and Hispanic origin: United States, 1999–June 2015 (preliminary data)

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig26>

Age and year	Total ²	Race and Hispanic origin ¹			
		Hispanic or Latino	Not Hispanic or Latino		
			White only	Black only	Asian only
Percent without health insurance coverage ³					
Under 18 years					
1999	11.9	26.7	8.1	11.9	10.4
2000	12.6	25.9	8.7	12.2	12.5
2001	11.2	24.6	7.2	10.6	12.6
2002	10.9	21.9	7.5	10.0	13.4
2003	9.8	20.2	6.4	8.9	12.0
2004	9.2	19.5	6.4	6.9	10.5
2005	9.3	17.5	6.5	8.9	11.4
2006	9.5	19.4	6.2	7.8	8.3
2007	9.0	15.3	7.1	6.2	8.0
2008	9.0	16.8	6.7	7.5	6.5
2009	8.2	14.7	6.0	6.6	7.5
2010	7.8	13.0	5.8	6.4	8.7
2011	7.0	12.3	4.8	5.5	7.8
2012	6.6	10.9	5.2	4.4	7.8
2013	6.6	11.8	4.7	5.1	5.9
2014	5.4	9.7	4.1	3.5	*4.3
2015, Jan–Jun ⁴	4.5	8.0	3.6	2.9	*
18–64 years					
1999	17.9	38.5	13.6	23.0	19.1
2000	18.9	41.4	13.9	23.6	19.5
2001	18.5	41.1	13.5	23.0	18.9
2002	19.3	40.8	14.4	23.2	18.7
2003	19.3	42.8	13.9	22.9	20.3
2004	19.3	42.9	14.0	22.7	18.6
2005	19.3	41.8	13.9	23.1	18.9
2006	20.0	43.8	14.6	22.2	16.5
2007	19.6	41.1	14.4	22.0	16.9
2008	19.9	42.6	14.4	22.9	15.7
2009	21.2	43.4	15.6	24.4	17.8
2010	22.3	43.3	16.3	27.1	19.4
2011	21.2	42.1	15.6	24.6	18.8
2012	20.9	41.3	15.1	23.6	19.1
2013	20.5	41.1	14.5	24.7	16.1
2014	16.3	34.1	11.5	17.6	12.1
2015, Jan–Jun ⁴	12.7	27.2	8.8	14.5	7.3

See footnotes at end of table.

Data table for Figure 26 (page 2 of 3). No health insurance coverage among persons under age 65, by age and race and Hispanic origin: United States, 1999–June 2015 (preliminary data)

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig26>

Age and year	Total ²	Race and Hispanic origin ¹			
		Hispanic or Latino	White only	Black only	Asian only
Standard error					
Under 18 years					
1999	0.3	0.9	0.4	0.8	1.5
2000	0.3	0.9	0.4	0.8	1.9
2001	0.4	0.9	0.4	0.9	2.1
2002	0.3	0.8	0.4	0.8	1.9
2003	0.3	0.8	0.4	0.7	2.1
2004	0.3	0.8	0.4	0.6	1.8
2005	0.3	0.7	0.4	0.7	1.9
2006	0.3	0.9	0.4	0.7	1.3
2007	0.4	0.8	0.5	0.6	1.4
2008	0.4	0.8	0.6	0.8	1.1
2009	0.4	0.8	0.5	0.6	1.1
2010	0.3	0.6	0.3	0.6	1.1
2011	0.3	0.6	0.3	0.6	1.1
2012	0.3	0.6	0.3	0.5	1.4
2013	0.3	0.6	0.3	0.6	1.0
2014	0.2	0.5	0.3	0.4	0.9
2015, Jan–Jun ⁴	0.4	0.7	0.5	0.6	*
18–64 years					
1999	0.3	0.8	0.3	0.7	1.4
2000	0.3	0.9	0.3	0.7	1.4
2001	0.3	0.8	0.3	0.7	1.3
2002	0.3	0.8	0.3	0.7	1.3
2003	0.3	0.8	0.3	0.7	1.4
2004	0.3	0.8	0.3	0.6	1.4
2005	0.3	0.7	0.3	0.7	1.2
2006	0.3	0.9	0.3	0.7	1.1
2007	0.3	0.8	0.3	0.7	1.1
2008	0.3	0.9	0.3	0.7	1.0
2009	0.3	0.9	0.3	0.7	1.1
2010	0.3	0.8	0.4	0.7	0.9
2011	0.3	0.7	0.3	0.6	0.9
2012	0.3	0.7	0.3	0.7	1.0
2013	0.3	0.8	0.3	0.6	0.8
2014	0.3	0.7	0.3	0.6	0.7
2015, Jan–Jun ⁴	0.3	0.9	0.3	0.7	0.6

See footnotes at end of table.

Data table for Figure 26 (page 3 of 3). No health insurance coverage among persons under age 65, by age and race and Hispanic origin: United States, 1999–June 2015 (preliminary data)

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig26>

Age and year ⁵	Race and Hispanic origin ¹					
	Hispanic or Latino	Not Hispanic or Latino				
		White only	Black only	Asian only	American Indian or Alaska Native only	2 or more races
Percent without health insurance coverage ³						
Under 18 years						
1999–2001	25.7	8.0	11.6	11.9	34.1	10.3
2003–2005	19.1	6.4	8.2	11.3	31.6	6.3
2006–2008	17.1	6.7	7.2	7.6	*24.2	9.2
2009–2011	13.3	5.5	6.2	8.0	*26.1	6.0
2012–2014	10.8	4.7	4.3	5.9	13.8	4.6
18–64 years						
1999–2001	40.4	13.7	23.2	19.2	39.8	21.1
2003–2005	42.5	13.9	22.9	19.2	36.9	21.1
2006–2008	42.5	14.5	22.4	16.4	39.1	22.4
2009–2011	42.9	15.8	25.4	18.7	39.9	25.3
2012–2014	38.8	13.7	22.0	15.7	33.2	19.4
Standard error						
Under 18 years						
1999–2001	0.6	0.2	0.5	1.1	4.0	1.1
2003–2005	0.5	0.2	0.4	1.1	4.0	0.9
2006–2008	0.5	0.3	0.4	0.8	7.2	1.0
2009–2011	0.4	0.2	0.4	0.7	7.3	0.7
2012–2014	0.4	0.2	0.3	0.7	2.6	0.5
18–64 years						
1999–2001	0.6	0.2	0.4	0.8	3.2	1.3
2003–2005	0.5	0.2	0.4	0.8	2.7	1.3
2006–2008	0.6	0.2	0.4	0.7	5.8	1.2
2009–2011	0.5	0.2	0.4	0.6	6.0	1.2
2012–2014	0.6	0.2	0.4	0.5	2.8	1.0

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.

¹Persons of Hispanic origin may be of any race. Race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity*. The single-race categories plus multiple-race category shown in the table conform to the 1997 Standards. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

²Includes all persons not shown separately.

³Persons not covered by private insurance, Medicaid, Children’s Health Insurance Program (CHIP), state-sponsored or other government-sponsored health plans (starting in 1997), Medicare, or military plans are considered to have no health insurance coverage. Persons with only Indian Health Service coverage are considered to have no health insurance coverage. Health insurance coverage is at the time of interview.

⁴Preliminary data based on the National Health Interview Survey’s Early Release program. Estimates based on the preliminary 6-month file may differ from estimates based on the final annual file and have larger standard errors associated with them than standard errors based on a final annual file. Available from: Martinez ME, Cohen RA. Health insurance coverage: Early release of estimates from the National Health Interview Survey, January–June 2015. NCHS. November 2015. Available from: <http://www.cdc.gov/nchs/nhis/releases.htm> and National Health Interview Survey, 2015 preliminary file. For more information, visit: <http://www.cdc.gov/nchs/nhis.htm>.

⁵Three-year average annual estimates are shown in order to present estimates for the American Indian or Alaska Native and the multiple-race populations. Annual estimates are not stable for smaller population groups.

NOTES: Also see Tables 102–105 and Figure 16. Ties in highest and lowest rates were resolved by looking at additional decimal places. See Technical Notes.

SOURCE: CDC/NCHS, National Health Interview Survey. Family core and sample adult questionnaires. See Appendix I, National Health Interview Survey (NHIS).

Data table for Figure 27 (page 1 of 2). Nonreceipt of needed dental care in the past 12 months due to cost among adults aged 18–64, by race and Hispanic origin: United States, 1999–2014

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig27>

Age and year	Total ²	Race and Hispanic origin ¹			
		Hispanic or Latino	Not Hispanic or Latino		
			White only	Black only	Asian only
Percent who did not receive needed dental care in the past 12 months due to cost ³					
18–64 years					
1999	9.3	8.5	9.3	10.4	4.5
2000	9.7	9.8	9.8	10.1	5.3
2001	10.4	12.0	10.2	10.3	5.1
2002	10.4	11.1	10.0	12.4	7.4
2003	11.5	13.5	11.1	12.8	5.3
2004	13.2	15.8	12.9	14.1	5.4
2005	13.0	15.5	12.3	15.3	6.8
2006	13.6	16.5	12.9	16.0	6.6
2007	13.0	16.0	12.5	13.6	7.5
2008	15.9	20.8	14.9	18.2	7.8
2009	16.8	22.2	15.5	19.0	9.1
2010	17.3	21.6	16.2	20.8	8.4
2011	16.4	21.1	15.4	18.2	9.4
2012	14.8	18.3	13.7	17.0	10.4
2013	14.3	18.4	13.3	16.3	8.8
2014	12.6	15.7	11.9	14.6	6.3
Standard error					
1999	0.2	0.5	0.3	0.7	0.9
2000	0.2	0.6	0.3	0.6	0.9
2001	0.2	0.6	0.3	0.6	0.9
2002	0.2	0.6	0.3	0.8	1.1
2003	0.3	0.7	0.3	0.7	0.9
2004	0.3	0.7	0.3	0.7	0.8
2005	0.3	0.6	0.3	0.8	1.0
2006	0.3	0.8	0.4	0.9	1.0
2007	0.3	0.8	0.4	0.8	0.9
2008	0.4	1.1	0.5	0.9	1.0
2009	0.4	1.0	0.5	0.9	1.0
2010	0.3	0.8	0.4	0.8	0.9
2011	0.3	0.7	0.4	0.8	0.9
2012	0.3	0.7	0.4	0.8	1.0
2013	0.3	0.7	0.4	0.7	0.9
2014	0.3	0.6	0.4	0.8	0.7

See footnotes at end of table.

Data table for Figure 27 (page 2 of 2). Nonreceipt of needed dental care in the past 12 months due to cost among adults aged 18–64, by race and Hispanic origin: United States, 1999–2014

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2015.htm#fig27>

Age and year ⁴	Race and Hispanic origin ¹					
	Not Hispanic or Latino					
	Hispanic or Latino	White only	Black only	Asian only	American Indian or Alaska Native only	2 or more races
Percent who did not receive needed dental care in the past 12 months due to cost ³						
18–64 years						
1999–2001	10.2	9.8	10.3	5.0	11.6	18.7
2003–2005	14.9	12.1	14.1	5.8	18.9	19.5
2006–2008	17.8	13.5	15.9	7.3	15.8	21.7
2009–2011	21.6	15.7	19.4	9.0	19.6	25.2
2012–2014	17.4	12.9	16.0	8.5	11.1	19.5
Standard error						
1999–2001	0.3	0.2	0.4	0.5	1.7	1.7
2003–2005	0.4	0.2	0.4	0.5	2.8	1.6
2006–2008	0.5	0.3	0.5	0.6	3.0	2.0
2009–2011	0.5	0.3	0.5	0.5	2.6	1.8
2012–2014	0.4	0.3	0.5	0.5	1.9	1.5

¹Persons of Hispanic origin may be of any race. Race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity*. The single-race categories plus multiple-race category shown in the table conform to the 1997 Standards. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

²Includes all persons not shown separately.

³Based on persons responding to the question, “During the past 12 months was there any time when [person] needed dental care (including checkups) but didn’t get it because [person] couldn’t afford it?”

⁴Three-year average annual estimates are shown in order to present estimates for the American Indian or Alaska Native and the multiple-race populations. Annual estimates are not stable for smaller population groups.

NOTE: Also see Table 63.

SOURCE: CDC/NCHS, National Health Interview Survey. Family core and sample adult questionnaires. See Appendix I, National Health Interview Survey (NHIS).

Data Sources

Data for the *Health, United States, 2015* Chartbook come from many surveys and data systems and cover a broad range of years. Detailed descriptions of the data sources included in the Chartbook are provided in Appendix I. Data Sources. Additional information clarifying and qualifying the data is included in the table notes and in Appendix II. Definitions and Methods.

Data Presentation

Many measures in the Chartbook are shown for people in specific age groups because of the strong effect of age on most health outcomes. Age-adjusted rates and age-adjusted percentages are computed to eliminate differences in observed rates that result from age differences in population composition (see Appendix II, Age adjustment). Age-adjusted rates and age-adjusted percentages are noted as such in the text; rates and percentages without this notation are crude rates and crude percentages. For some charts, data years are combined to increase sample size and the reliability of the estimates. Some charts present time trends, and others focus on differences in estimates among population subgroups for the most recent time point available. Figures 1–17 and the Highlights section generally present trends for the recent 10-year period. For some indicators, a slightly longer or shorter period may be shown due to design or data comparability issues. Trends are generally shown on a linear scale to emphasize absolute differences over time. The time trends for the overall mortality measures are shown on a logarithmic (log) scale to enable measures with large differences in magnitude to be shown on the same chart.

Point estimates and standard errors for Figures 1–17 are available either in the Trend Table and Excel spreadsheet specified in the note below the chart, or in the Chartbook tables section. For the Special Feature on racial and ethnic health disparities (Figures 18–27), data tables with point estimates and standard errors are contained in the Chartbook tables section. These data tables may include additional data that were not graphed because of space considerations.

Reliability of Estimates

Overall estimates generally have relatively small sampling errors, but estimates for certain population subgroups may be based on small numbers and have relatively large sampling errors. Numbers of deaths obtained from the National Vital Statistics System represent complete counts and therefore are not subject to sampling error. They are,

however, subject to random variation, which means that the number of events that actually occur in a given year may be considered as one of a large series of possible results that could have arisen under the same circumstances. When the number of events is small and the probability of such an event is small, considerable caution must be observed in interpreting the conditions described by the charts. Estimates that are unreliable because of large sampling errors or small numbers of events have been noted with an asterisk. The criteria used to designate or suppress unreliable estimates are indicated in the notes to the applicable tables or charts.

For NCHS surveys, point estimates and their corresponding variances were calculated using the SUDAAN software package, which takes into consideration the complex survey design (94). Standard errors for other surveys or data sets were computed using the methodology recommended by the programs providing the data, or were provided directly by those programs.

Statistical Testing

Data trends can be described in many ways. For most trend analyses presented in the Chartbook, increases or decreases in the estimates during the entire time period shown are assessed by the weighted least squares regression method in the National Cancer Institute's Joinpoint software (with Grid search and Bayesian Information Criterion (BIC) model selection). The default maximum number of joinpoints based on the number of available data points in the trend was used. Statistically significant changes in the trend were assessed at the 0.05 level. For more information on Joinpoint, see: <http://surveillance.cancer.gov/joinpoint>. Statistical significance of differences between regression coefficients at the 0.05 level was also taken into account to select a model with the fewest joinpoints or changes in trend. For some trend charts, there were too few observations for Joinpoint analysis. In those cases, either the difference between two points was assessed for statistical significance using z-tests or the statistical testing methods recommended by the data systems were used. Trend analyses using weighted least squares regression for Figures 1–17 were carried out on the log scale so that results provide estimates of percent change. However, as discussed below, trend analyses for figures in the Special Feature were carried out on the linear scale.

For analyses that show two time points, differences between the two points were assessed for statistical significance at the 0.05 level using two-sided significance tests (z-tests) without correction for multiple comparisons. Trend and data tables include point estimates and standard errors for users who would like to perform additional statistical tests.

Terms such as “similar,” “stable,” and “no difference” used in the text indicate that the statistics being compared were not significantly different. Lack of comment regarding the difference between statistics does not necessarily suggest that the difference was tested and found to be not significant. Because statistically significant differences or trends are partly a function of sample size (the larger the sample, the smaller the change that can be detected), they do not necessarily have public health significance (95).

Special Feature on Racial and Ethnic Health Disparities (Figures 18–27)

In general, the starting time period for trend analysis in the Special Feature is 1999. This is the earliest year for which National Health Interview Survey (NHIS) data were available for detailed racial and ethnic groups (see Appendix II, Race). Trend data on race and ethnicity are presented in the greatest detail possible after taking into account the quality of the data, the amount of missing data, and the number of observations. These issues significantly affect the availability of reportable data for certain populations, such as the Native Hawaiian or Other Pacific Islander population and the American Indian or Alaska Native population. Estimates for the Native Hawaiian or Other Pacific Islander population were unstable and are not presented. Three years of data were combined in order to present estimates for the American Indian or Alaska Native population in the data tables that accompany Figures 24–27.

There are various ways to quantify racial and ethnic differences in health and mortality, and different measures of disparity may lead to different conclusions (55–58). This Special Feature uses the maximal rate difference, one of three overall measures used in Healthy People 2020, to measure racial and ethnic disparities (59). The maximal rate difference is an overall measure of health disparities calculated as the absolute difference between the highest and lowest group rates in the population for a given characteristic, irrespective of other, intermediate rates (59). A decrease in the maximal rate difference does not capture whether the population health outcome overall is improving; rather it reflects progress toward eliminating disparities. As the absolute difference between the highest and lowest rates decreases toward 0, all the pairwise absolute differences between population subgroups will tend to 0. For determination of the highest and lowest group rates, estimates were ranked from highest to lowest based on the observed value to six decimal places, to avoid ties. Tests of statistical significance against other rates were not conducted. For consistency with the use of the absolute difference to measure disparity, all analyses in the Special Feature are carried out on the linear scale. For each figure in

the Special Feature that shows trends (Figures 19, 21, 23–27) the following analyses were carried out:

- (a) trend analysis of overall estimates;
- (b) trend analysis of estimates for each racial and ethnic group; and
- (c) trend analysis of the maximal rate difference.

These trend analyses provide information used to:

- (a) describe the trend in overall estimates as increasing, decreasing, or stable, and any changes in trend over the time period;
- (b) indicate whether the trend in estimates for different racial and ethnic groups is similar to the overall trend; and
- (c) describe the trend in disparity as measured by the maximal difference in rates as increasing, decreasing, or stable and any changes in trend over the time period.

In addition, a one-sided z-test was conducted to test whether the maximal difference in rates was 0 vs. >0 at the most recent time point (59). For figures in the Special Feature that only show estimates at a single time point, the maximal rate difference was calculated for that time point, and a one-sided z-test was conducted to test whether the maximal difference in rates was 0 vs. >0 .

References

1. Xu JQ, Murphy SL, Kochanek KD, Bastian BA. Deaths: Final data for 2013. National vital statistics reports; vol 64 no 2. Hyattsville, MD: NCHS; 2016.
2. Heron M. Deaths: Leading causes for 2013. National vital statistics reports; vol 65 no 2. Hyattsville, MD: NCHS; 2016.
3. HHS; National Action Alliance for Suicide Prevention. 2012 National strategy for suicide prevention: Goals and objectives for action: A report of the U.S. Surgeon General and of the National Action Alliance for Suicide Prevention. Washington, DC: HHS; 2012.
4. Institute of Medicine. Reducing suicide: A national imperative. Washington, DC: National Academies Press; 2002.
5. Rheingold AA, Zinzow H, Hawkins A, Saunders BE, Kilpatrick DG. Prevalence and mental health outcomes of homicide survivors in a representative U.S. sample of adolescents: Data from the 2005 National Survey of Adolescents. *J Child Psychol Psychiatry* 2012;53(6):687–94.
6. Smith EL, Cooper A. Homicide in the U.S. known to law enforcement, 2011. Washington, DC: U.S. Department of Justice, Office of Justice Programs; 2013.
7. Swanson JW, Bonnie RJ, Appelbaum PS. Getting serious about reducing suicide: More “how” and less “why.” *JAMA* 2015;314(21):2229–30.
8. Martin JA, Hamilton BE, Osterman MJK. Births in the United States, 2014. NCHS data brief, no 216. Hyattsville, MD: NCHS; 2015.
9. Ventura SJ, Hamilton BE, Mathews TJ. National and state patterns of teen births in the United States, 1940–2013. National vital statistics reports; vol 63 no 4. Hyattsville, MD: NCHS; 2014.
10. Adams D, Fullerton K, Jajosky R, et al. Summary of notifiable infectious diseases and conditions—United States, 2013. *MMWR* 2015;62(53):1–122.
11. CDC. National Notifiable Diseases Surveillance System (NNDSS): History and background. Atlanta, GA; 2015.
12. HHS. How tobacco smoke causes disease: The biology and behavioral basis for smoking-attributable disease: A report of the Surgeon General. Atlanta, GA: CDC, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 2010.
13. Freedman DS, Mei Z, Srinivasan SR, Berenson GS, Dietz WH. Cardiovascular risk factors and excess adiposity among overweight children and adolescents: The Bogalusa Heart Study. *J Pediatr* 2007;150(1):12–7.
14. Engeland A, Bjørge T, Tverdal A, Sjøgaard AJ. Obesity in adolescence and adulthood and the risk of adult mortality. *Epidemiology* 2004;15(1):79–85.
15. Kuczumski RJ, Ogden CL, Guo SS, et al. 2000 CDC growth charts for the United States: Methods and development. *NCHS Vital Health Stat* 11(246); 2002.
16. Barlow SE; Expert Committee. Expert Committee recommendations regarding the prevention, assessment, and treatment of child and adolescent overweight and obesity: Summary report. *Pediatrics* 2007;120(suppl 4):S164–92.
17. National Heart, Lung, and Blood Institute; National Institute of Diabetes and Digestive and Kidney Diseases. Clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults: The evidence report. NIH pub no 98–4083. Bethesda, MD: National Institutes of Health; 1998.
18. Jensen MD, Ryan DH, Apovian CM, Ard JD, Comuzzie AG, Donato KA, et al. 2013 AHA/ACC/TOS guideline for the management of overweight and obesity in adults: A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and the Obesity Society. *Circulation* 2014;129(25 suppl 2):S102–40.
19. HHS. The Surgeon General's vision for a healthy and fit nation. Rockville, MD: HHS, Office of the Surgeon General; 2010.
20. Flegal KM, Graubard BI, Williamson DF, Gail MH. Excess deaths associated with underweight, overweight, and obesity. *JAMA* 2005;293(15):1861–7.
21. Andrae B, Andersson TM, Lambert PC, Kemetli L, Silfverdal L, Strander B, et al. Screening and cervical cancer cure: Population based cohort study. *BMJ* 2012;344:e900.
22. Agency for Healthcare Research and Quality, U.S. Preventive Services Task Force. Final recommendation statement: Cervical cancer: Screening, March 2012. Rockville, MD: HHS, AHRQ; 2014.
23. Institute of Medicine. Hospital-based emergency care: At the breaking point. Washington, DC: National Academies Press; 2006.
24. DeLia D, Cantor J. Emergency department utilization and capacity. Research synthesis report no 17. Princeton, NJ: Robert Wood Johnson Foundation; 2009.
25. Taylor J. Don't bring me your tired, your poor: The crowded state of America's emergency departments. Issue brief no. 811. Washington, DC: National Health Policy Forum; 2006.
26. Kaiser Commission on Medicaid and the Uninsured. The uninsured and the difference health insurance makes. Menlo Park, CA: Kaiser Family Foundation; 2012.
27. Cunningham PJ, Felland LE. Falling behind: Americans' access to medical care deteriorates, 2003–2007. *Track Rep* 2008;19:1–5.
28. Blumenthal D, Tavenner M. The “meaningful use” regulation for electronic health records. *N Engl J Med* 2010;363(6):501–4.
29. Kern LM, Barrón Y, Dhopeswarkar RV, Edwards A, Kaushal R; HITEC Investigators. Electronic health records and ambulatory quality of care. *J Gen Intern Med* 2013;28(4):496–503.
30. Hsiao C-J, Hing E. Use and characteristics of electronic health record systems among office-based physician practices: United States, 2001–2013. NCHS data brief, no 143. Hyattsville, MD: NCHS; 2014.
31. Decker SL. In 2011 nearly one-third of physicians said they would not accept new Medicaid patients, but rising fees may help. *Health Aff (Millwood)* 2012;31(8):1673–9.
32. NCHS. National Electronic Health Records Survey 2013. Available from: http://www.cdc.gov/nchs/data/ahcd/2013_NEHRS_Questionnaire.pdf.
33. Hing E, Decker SL, Jamoom E. Acceptance of new patients with public and private insurance by office-based physicians: United States, 2013. NCHS data brief, no 195. Hyattsville, MD: NCHS; 2015.

34. Ingram DD, Franco SJ. 2013 NCHS urban–rural classification scheme for counties. *Vital Health Stat* 2(166); 2014.
35. Martin AB, Hartman M, Benson J, Catlin A; National Health Expenditure Accounts Team. National health spending in 2014: Faster growth driven by coverage expansion and prescription drug spending. *Health Aff (Millwood)* 2016;35(1):150–60.
36. Martinez ME, Cohen RA. Health insurance coverage: Early release of estimates from the National Health Interview Survey, January–June 2015. NCHS; 2015.
37. NCHS. National Health Interview Survey [unpublished analysis]. December 1, 2015.
38. Patient Protection and Affordable Care Act. Pub L No 111–148, 124 Stat 119, 132; 2010.
39. Cohen RA, Martinez ME. Health insurance coverage: Early release of estimates from the National Health Interview Survey, 2014. NCHS; 2015.
40. Kaiser Commission on Medicaid and the Uninsured. Medicaid eligibility for adults as of January 1, 2014. Menlo Park, CA: Kaiser Family Foundation; 2013.
41. Heckler MM. Report of the Secretary's Task Force on Black & Minority Health. Vol I, Executive summary. Washington, DC: HHS; 1985.
42. HHS; Offices of Minority Health. National Partnership for Action to End Health Disparities. Washington, DC; 2012.
43. Agency for Healthcare Research and Quality. National healthcare quality and disparity reports. Rockville, MD: HHS.
44. HHS. Healthy People 2010: Understanding and improving health. 2nd ed. Washington, DC: U.S. Government Printing Office; 2000.
45. HHS. HHS action plan to reduce racial and ethnic health disparities: A nation free of disparities in health and health care. Washington, DC.
46. Cooper RS, Kaufman JS, Ward R. Race and genomics. *N Engl J Med* 2003;348(12):1166–70.
47. Williams DR, Sternthal M. Understanding racial-ethnic disparities in health: Sociological contribution. *J Health Soc Behav* 2010;51(suppl):S15–27.
48. Office of Management and Budget. Race and ethnic standards for federal statistics and administrative reporting. Statistical Policy Directive 15; 1977.
49. Office of Management and Budget. Revisions to the standards for the classification of federal data on race and ethnicity. *Fed Regist* 1997;62(210):58782–90.
50. U.S. Census Bureau, Population Division. Annual estimates of the resident population by sex, race, and Hispanic origin for the United States, states, and counties: April 1, 2010 to July 1, 2014. Washington, DC; 2015.
51. DeNavas-Walt C, Proctor BD, U.S. Census Bureau. Income and poverty in the United States: 2014. Current Population Reports, P60–252. Washington, DC: U.S. Government Printing Office; 2015.
52. U.S. Census Bureau. Current Population Survey (CPS): CPS table creator, 2014 data from the CPS' annual social and economic supplement. Washington, DC.
53. Markides KS, Eschbach K. Aging, migration, and mortality: Current status of research on the Hispanic paradox. *J Gerontol B Psychol Sci Soc Sci* 2005;60(Spec No 2):68–75.
54. HHS. National partnership for action to end health disparities: Offices of minority health. Washington, DC; 2011. Available from: <http://minorityhealth.hhs.gov/npa/templates/browse.aspx?lvl=1&lvlid=34>.
55. Keppel K, Pamuk E, Lynch J, et al. Methodological issues in measuring health disparities. NCHS. *Vital Health Stat* 2(141); 2005.
56. Keppel KG, Pearcy JN, Klein RJ. Measuring progress in Healthy People 2010. Healthy People Statistical Notes, no 25. Hyattsville, MD: NCHS; 2004.
57. WHO. Handbook on health inequality monitoring with a special focus on low- and middle-income countries. Geneva, Switzerland; 2013.
58. Harper S, Lynch J. Methods for measuring cancer disparities: A review using data relevant to Healthy People 2010 cancer-related objectives. NCI Cancer Surveillance Monograph Series, no 6. Bethesda, MD: National Cancer Institute; 2005.
59. Talih M, Huang DT. Measuring progress toward target attainment and the elimination of health disparities in Healthy People 2020. Healthy People Statistical Notes, no 27. Hyattsville, MD: NCHS; 2016.
60. Kochanek KD, Murphy SL, Xu JQ. Deaths: Final data for 2011. National vital statistics reports; vol 63 no 3. Hyattsville, MD: NCHS; 2015.
61. Kochanek KD, Arias E, Anderson RN. Leading causes of death contributing to decrease in life expectancy gap between black and white populations: United States, 1999–2013. NCHS data brief, no 218. Hyattsville, MD: NCHS; 2015.
62. Singh GK, van Dyck PC. Infant mortality in the United States, 1935–2007: Over seven decades of progress and disparities. Rockville, MD: HHS, Health Resources and Services Administration, Maternal and Child Health Bureau; 2010.
63. Association of Maternal & Child Health Programs. Celebrating the legacy, shaping the future: 75 years of state and federal partnership to improve maternal and child health. Washington, DC; 2010.
64. Chang HH, Larson J, Blencowe H, Spong CY, Howson CP, Cairns-Smith S, et al. Preventing preterm births: Analysis of trends and potential reductions with interventions in 39 countries with very high human development index. *Lancet* 2013;381(9862):223–34.
65. Ancel PY, Goffinet F, EPIPAGE-2 Writing Group. Survival and morbidity of preterm children born at 22 through 34 weeks' gestation in France in 2011: Results of the EPIPAGE-2 cohort study. *JAMA Pediatr* 2015;169(3):230–8.
66. Adams JN, Feldman HM, Huffman LC, Loe IM. Sensory processing in preterm preschoolers and its association with executive function. *Early Hum Dev* 2015;91(3):227–33.
67. Brown L, Burns YR, Watter P, Gibbons KS, Gray PH. Motor performance, postural stability and behaviour of non-disabled extremely preterm or extremely low birth weight children at four to five years of age. *Early Hum Dev* 2015;91(5):309–15.
68. van der Pal-de Bruin KM, van der Pal SM, Verloove-Vanhorick SP, Walther FJ. Profiling the preterm or VLBW born adolescent; Implications of the Dutch POPS cohort follow-up studies. *Early Hum Dev* 2015;91(2):97–102.
69. Stoll BJ, Hansen NI, Bell EF, Shankaran S, Laptook AR, Walsh MC, et al. Neonatal outcomes of extremely preterm infants from the NICHD Neonatal Research Network. *Pediatrics* 2010;126(3):443–56.
70. Spong CY. Defining “term” pregnancy: Recommendations from the Defining “Term” Pregnancy Workgroup. *JAMA* 2013;309(23):2445–6.
71. Mathews TJ, MacDorman MF, Thoma ME. Infant mortality statistics from the 2013 period linked birth/infant death data set. National vital statistics reports; vol 64 no 9. Hyattsville, MD: NCHS; 2015.

72. MacDorman MF, Mathews TJ. Understanding racial and ethnic disparities in U.S. infant mortality rates. NCHS data brief, no 74. Hyattsville, MD: NCHS; 2011.
73. Martin JA, Osterman MJK, Kirmeyer SE, Gregory ECW. Measuring gestational age in vital statistics data: Transitioning to the obstetric estimate. National vital statistics reports; vol 64 no 5. Hyattsville, MD: NCHS; 2015.
74. Hamilton BE, Martin JA, Osterman MJK, et al. Births: Final data for 2014. National vital statistics reports; vol 64 no 12. Hyattsville, MD: NCHS; 2015.
75. Osterman MJK, Martin JA. Trends in low-risk cesarean delivery in the United States, 1990–2013. National vital statistics reports; vol 63 no 6. Hyattsville, MD: NCHS; 2014.
76. D'Alton ME, Hehir MP. Cesarean delivery rates: Revisiting a 3-decades-old dogma. *JAMA* 2015;314(21):2238–40.
77. Carroll MD, Navaneelan T, Bryan S, Ogden CL. Prevalence of obesity among children and adolescents in Canada and the United States. NCHS data brief, no 211. Hyattsville, MD: NCHS; 2015.
78. Organisation for Economic Co-operation and Development. Obesity update. Paris, France; 2014.
79. Ogden CL, Carroll MD, Fryar CD, Flegal KM. Prevalence of obesity among adults and youth: United States, 2011–2014. NCHS data brief, no 219. Hyattsville, MD: NCHS; 2015.
80. National Heart Lung and Blood Institute. National High Blood Pressure Education Program. Available from: <http://www.nhlbi.nih.gov/health/health-topics/topics/hbp>.
81. Nwankwo T, Yoon SS, Burt V, Gu Q. Hypertension among adults in the United States: National Health and Nutrition Examination Survey, 2011–2012. NCHS data brief, no 133. Hyattsville, MD: NCHS; 2013.
82. NCHS. National Health and Nutrition Examination Survey. Hyattsville, MD. [unpublished analysis].
83. CDC. Racial/ethnic disparities in the awareness, treatment, and control of hypertension—United States, 2003–2010. *MMWR* 2013;62(18):351–5.
84. Jamal A, Homa DM, O'Connor E, Babb SD, Caraballo RS, Singh T, et al. Current cigarette smoking among adults—United States, 2005–2014. *MMWR* 2015;64(44):1233–40.
85. HHS. The health consequences of smoking—50 years of progress: A Report of the Surgeon General, Chapter 12, 2014. Atlanta, GA: HHS, CDC, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 2014.
86. CDC. Experiences with obtaining influenza vaccination among persons in priority groups during a vaccine shortage—United States, October–November, 2004. *MMWR* 2004;53(49):1153–5.
87. Centers for Medicare & Medicaid Services. Children's Health Insurance Program (CHIP). Baltimore, MD.
88. Kaiser Family Foundation. A guide to the Supreme Court's decision on the ACA's Medicaid expansion. Menlo Park, CA; 2012.
89. HHS. Oral health in America: A Report of the Surgeon General. Rockville, MD: HHS, National Institute of Dental and Craniofacial Research, National Institutes of Health; 2000.
90. Bloom B, Cohen RA. Dental insurance for persons under age 65 years with private health insurance: United States, 2008. NCHS data brief, no 40. Hyattsville, MD: NCHS; 2010.
91. Institute of Medicine and National Research Council. Improving access to oral health care for vulnerable and underserved populations. Washington, DC: The National Academies Press; 2011.
92. National Association of Dental Plans. Who has dental benefits? Available from: http://www.nadp.org/Dental_Benefits_Basics/Dental_BB_1.aspx.
93. Agency for Healthcare Research and Quality. Medical Expenditure Panel Survey household component tables. Table 3: Dental services—Median and mean expenses per person with expense and distribution of expenses by source of payment: United States, 2012. Rockville, MD [generated interactively July 31, 2015].
94. SUDAAN, release 10.0.1 [computer software]. Research Triangle Park, NC: RTI International; 2009.
95. CDC. Youth Risk Behavior Surveillance System (YRBSS). Interpretation of YRBS trend data. Atlanta, GA; 2014. Available from: http://www.cdc.gov/HealthyYouth/yrbs/pdf/YRBS_trend_interpretation.pdf.

Table 1 (page 1 of 3). Resident population, by age, sex, race, and Hispanic origin: United States, selected years 1950–2014

 Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#001>.

[Data are based on the decennial census updated with data from multiple sources]

Sex, race, Hispanic origin, and year	Total resident population	Age										
		Under 1 year	1–4 years	5–14 years	15–24 years	25–34 years	35–44 years	45–54 years	55–64 years	65–74 years	75–84 years	85 years and over
All persons		Number, in thousands										
1950	150,697	3,147	13,017	24,319	22,098	23,759	21,450	17,343	13,370	8,340	3,278	577
1960	179,323	4,112	16,209	35,465	24,020	22,818	24,081	20,485	15,572	10,997	4,633	929
1970	203,212	3,485	13,669	40,746	35,441	24,907	23,088	23,220	18,590	12,435	6,119	1,511
1980	226,546	3,534	12,815	34,942	42,487	37,082	25,635	22,800	21,703	15,581	7,729	2,240
1990	248,710	3,946	14,812	35,095	37,013	43,161	37,435	25,057	21,113	18,045	10,012	3,021
2000	281,422	3,806	15,370	41,078	39,184	39,892	45,149	37,678	24,275	18,391	12,361	4,240
2010	308,746	3,944	16,257	41,026	43,626	41,064	41,071	45,007	36,483	21,713	13,061	5,493
2012	313,914	3,943	16,056	41,145	43,944	42,309	40,516	44,269	38,586	23,985	13,273	5,887
2013	316,129	3,942	15,926	41,221	43,954	42,845	40,453	43,768	39,316	25,217	13,447	6,041
2014	318,857	3,948	15,929	41,191	43,980	43,517	40,513	43,459	40,078	26,398	13,683	6,162
Male												
1950	74,833	1,602	6,634	12,375	10,918	11,597	10,588	8,655	6,697	4,024	1,507	237
1960	88,331	2,090	8,240	18,029	11,906	11,179	11,755	10,093	7,537	5,116	2,025	362
1970	98,912	1,778	6,968	20,759	17,551	12,217	11,231	11,199	8,793	5,437	2,436	542
1980	110,053	1,806	6,556	17,855	21,419	18,382	12,570	11,009	10,152	6,757	2,867	682
1990	121,239	2,018	7,581	17,971	18,915	21,564	18,510	12,232	9,955	7,907	3,745	841
2000	138,054	1,949	7,862	21,043	20,079	20,121	22,448	18,497	11,645	8,303	4,879	1,227
2010	151,781	2,014	8,305	20,970	22,318	20,632	20,436	22,142	17,601	10,097	5,477	1,790
2012	154,492	2,017	8,199	21,026	22,512	21,339	20,174	21,807	18,603	11,203	5,648	1,964
2013	155,652	2,017	8,136	21,061	22,525	21,641	20,145	21,569	18,957	11,798	5,761	2,042
2014	156,936	2,018	8,138	21,030	22,523	21,970	20,159	21,425	19,322	12,349	5,893	2,109
Female												
1950	75,864	1,545	6,383	11,944	11,181	12,162	10,863	8,688	6,672	4,316	1,771	340
1960	90,992	2,022	7,969	17,437	12,114	11,639	12,326	10,393	8,036	5,881	2,609	567
1970	104,300	1,707	6,701	19,986	17,890	12,690	11,857	12,021	9,797	6,998	3,683	969
1980	116,493	1,727	6,259	17,087	21,068	18,700	13,065	11,791	11,551	8,824	4,862	1,559
1990	127,471	1,928	7,231	17,124	18,098	21,596	18,925	12,824	11,158	10,139	6,267	2,180
2000	143,368	1,857	7,508	20,034	19,105	19,771	22,701	19,181	12,629	10,088	7,482	3,013
2010	156,964	1,930	7,952	20,056	21,309	20,432	20,635	22,864	18,882	11,617	7,584	3,704
2012	159,422	1,926	7,857	20,118	21,432	20,971	20,343	22,462	19,983	12,783	7,624	3,923
2013	160,477	1,925	7,791	20,160	21,429	21,203	20,307	22,198	20,360	13,419	7,686	3,999
2014	161,921	1,930	7,791	20,161	21,456	21,546	20,354	22,034	20,756	14,049	7,789	4,053
White male												
1950	67,129	1,400	5,845	10,860	9,689	10,430	9,529	7,836	6,180	3,736	1,406	218
1960	78,367	1,784	7,065	15,659	10,483	9,940	10,564	9,114	6,850	4,702	1,875	331
1970	86,721	1,501	5,873	17,667	15,232	10,775	9,979	10,090	7,958	4,916	2,243	487
1980	94,976	1,487	5,402	14,773	18,123	15,940	11,010	9,774	9,151	6,096	2,600	621
1990	102,143	1,604	6,071	14,467	15,389	18,071	15,819	10,624	8,813	7,127	3,397	760
2000	113,445	1,524	6,143	16,428	15,942	16,232	18,568	15,670	10,067	7,343	4,419	1,109
2010	121,403	1,518	6,281	16,043	17,069	16,139	16,208	18,096	14,840	8,726	4,866	1,617
2012	122,937	1,513	6,159	16,013	17,118	16,594	15,882	17,701	15,549	9,667	4,977	1,766
2013	123,559	1,509	6,101	16,000	17,093	16,764	15,813	17,447	15,787	10,160	5,056	1,830
2014	124,143	1,505	6,089	15,925	17,034	16,907	15,747	17,265	16,034	10,603	5,152	1,883
White female												
1950	67,813	1,341	5,599	10,431	9,821	10,851	9,719	7,868	6,168	4,031	1,669	314
1960	80,465	1,714	6,795	15,068	10,596	10,204	11,000	9,364	7,327	5,428	2,441	527
1970	91,028	1,434	5,615	16,912	15,420	11,004	10,349	10,756	8,853	6,366	3,429	890
1980	99,835	1,412	5,127	14,057	17,653	15,896	11,232	10,285	10,325	7,951	4,457	1,440
1990	106,561	1,524	5,762	13,706	14,599	17,757	15,834	10,946	9,698	9,048	5,687	2,001
2000	116,641	1,447	5,839	15,576	14,966	15,574	18,386	15,921	10,731	8,757	6,715	2,729
2010	124,020	1,451	5,993	15,270	16,153	15,552	15,941	18,311	15,586	9,846	6,601	3,314
2012	125,287	1,444	5,885	15,254	16,176	15,893	15,573	17,853	16,340	10,809	6,576	3,485
2013	125,785	1,440	5,826	15,247	16,146	16,021	15,489	17,569	16,588	11,319	6,603	3,537
2014	126,488	1,439	5,816	15,203	16,129	16,204	15,444	17,362	16,849	11,805	6,667	3,569

See footnotes at end of table.

Table 1 (page 2 of 3). Resident population, by age, sex, race, and Hispanic origin: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#001>.

[Data are based on the decennial census updated with data from multiple sources]

Sex, race, Hispanic origin, and year	Total resident population	Age										
		Under 1 year	1–4 years	5–14 years	15–24 years	25–34 years	35–44 years	45–54 years	55–64 years	65–74 years	75–84 years	85 years and over
Black or African American male												
Number, in thousands												
1950	7,300	---	¹ 944	1,442	1,162	1,105	1,003	772	459	299	² 113	---
1960	9,114	281	1,082	2,185	1,305	1,120	1,086	891	617	382	137	29
1970	10,748	245	975	2,784	2,041	1,226	1,084	979	739	461	169	46
1980	12,585	269	967	2,614	2,807	1,967	1,235	1,024	854	567	228	53
1990	14,420	322	1,164	2,700	2,669	2,592	1,962	1,175	878	614	277	66
2000	17,407	313	1,271	3,454	2,932	2,586	2,705	1,957	1,090	683	330	87
2010	20,101	341	1,388	3,408	3,591	2,801	2,639	2,708	1,832	886	396	110
2012	20,686	346	1,391	3,432	3,698	2,944	2,620	2,707	2,021	979	424	123
2013	20,935	346	1,381	3,447	3,720	3,019	2,629	2,693	2,092	1,038	439	129
2014	21,241	347	1,383	3,458	3,735	3,122	2,661	2,681	2,162	1,099	456	136
Black or African American female												
1950	7,745	---	¹ 941	1,446	1,300	1,260	1,112	796	443	322	² 125	---
1960	9,758	283	1,085	2,191	1,404	1,300	1,229	974	663	430	160	38
1970	11,832	243	970	2,773	2,196	1,456	1,309	1,134	868	582	230	71
1980	14,046	266	951	2,578	2,937	2,267	1,488	1,258	1,059	776	360	106
1990	16,063	316	1,137	2,641	2,700	2,905	2,279	1,416	1,135	884	495	156
2000	19,187	302	1,228	3,348	2,971	2,866	3,055	2,274	1,353	971	587	233
2010	21,965	330	1,343	3,292	3,568	3,066	2,962	3,056	2,197	1,192	675	282
2012	22,517	331	1,344	3,314	3,622	3,171	2,941	3,056	2,420	1,305	706	307
2013	22,762	331	1,333	3,333	3,630	3,224	2,950	3,043	2,499	1,378	723	318
2014	23,069	332	1,333	3,349	3,630	3,305	2,984	3,029	2,578	1,460	741	328
American Indian or Alaska Native male												
1980	702	17	59	153	161	114	75	53	37	22	9	2
1990	1,024	24	88	206	192	183	140	86	55	32	13	3
2000	1,488	28	109	301	271	229	229	165	88	45	18	5
2010	2,143	39	160	381	392	336	290	264	167	76	29	7
2012	2,210	39	158	386	396	350	296	270	186	89	33	9
2013	2,240	40	157	388	396	356	299	272	194	95	35	9
2014	2,269	40	156	389	395	362	302	274	201	102	37	10
American Indian or Alaska Native female												
1980	718	16	57	149	158	118	79	57	41	27	12	4
1990	1,041	24	85	200	178	186	148	92	61	41	21	6
2000	1,496	26	106	293	254	219	236	174	95	54	28	10
2010	2,121	38	156	370	364	316	282	273	179	87	41	14
2012	2,188	38	154	375	371	325	285	277	200	100	45	16
2013	2,217	39	153	378	374	329	288	277	209	107	48	18
2014	2,250	38	152	379	376	335	291	277	218	114	50	19
Asian or Pacific Islander male												
1980	1,814	35	130	321	334	366	252	159	110	72	30	6
1990	3,652	68	258	598	665	718	588	347	208	133	57	12
2000	5,713	84	339	861	934	1,073	947	705	399	231	112	27
2010	8,134	116	476	1,138	1,266	1,356	1,299	1,075	761	409	186	55
2012	8,658	119	492	1,195	1,301	1,451	1,376	1,128	847	469	214	66
2013	8,917	121	497	1,226	1,316	1,503	1,405	1,158	884	504	230	73
2014	9,284	127	510	1,257	1,359	1,580	1,449	1,205	925	545	248	80

See footnotes at end of table.

Table 1 (page 3 of 3). Resident population, by age, sex, race, and Hispanic origin: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#001>.

[Data are based on the decennial census updated with data from multiple sources]

Sex, race, Hispanic origin, and year	Total resident population	Age										
		Under 1 year	1–4 years	5–14 years	15–24 years	25–34 years	35–44 years	45–54 years	55–64 years	65–74 years	75–84 years	85 years and over
Number, in thousands												
Asian or Pacific Islander female												
1980	1,915	34	127	307	325	423	269	192	126	71	33	9
1990	3,805	65	247	578	621	749	664	371	264	166	65	17
2000	6,044	81	336	817	914	1,112	1,024	812	451	305	152	41
2010	8,859	110	460	1,124	1,223	1,498	1,450	1,223	920	491	267	93
2012	9,430	113	474	1,175	1,262	1,582	1,543	1,276	1,023	569	298	115
2013	9,713	116	479	1,202	1,279	1,628	1,580	1,309	1,064	615	313	127
2014	10,114	121	489	1,230	1,322	1,702	1,634	1,366	1,111	670	331	138
Hispanic or Latino male												
1980	7,280	187	661	1,530	1,646	1,256	761	570	364	200	86	19
1990	11,388	279	980	2,128	2,376	2,310	1,471	818	551	312	131	32
2000	18,162	395	1,506	3,469	3,564	3,494	2,653	1,551	804	474	203	50
2010	25,619	515	2,094	4,755	4,648	4,419	3,734	2,736	1,535	735	352	95
2012	26,930	523	2,105	4,959	4,784	4,579	3,919	2,967	1,747	840	389	116
2013	27,461	518	2,091	5,049	4,826	4,638	3,996	3,070	1,840	898	407	127
2014	28,018	517	2,097	5,107	4,868	4,666	4,083	3,194	1,955	964	428	140
Hispanic or Latina female												
1980	7,329	181	634	1,482	1,546	1,249	805	615	411	257	117	30
1990	10,966	268	939	2,039	2,028	2,073	1,448	868	632	403	209	59
2000	17,144	376	1,441	3,318	3,017	3,016	2,476	1,585	907	603	303	101
2010	24,859	497	2,008	4,561	4,206	4,016	3,564	2,728	1,679	914	510	176
2012	26,098	502	2,025	4,760	4,382	4,092	3,730	2,922	1,888	1,029	557	213
2013	26,610	496	2,013	4,844	4,449	4,124	3,794	3,011	1,979	1,089	580	230
2014	27,370	496	2,021	4,920	4,551	4,215	3,910	3,136	2,099	1,166	607	248
White, not Hispanic or Latino male												
1980	88,035	1,308	4,772	13,317	16,554	14,739	10,284	9,229	8,803	5,906	2,519	603
1990	91,743	1,351	5,181	12,525	13,219	15,967	14,481	9,875	8,303	6,837	3,275	729
2000	96,551	1,163	4,761	13,238	12,628	12,958	16,088	14,223	9,312	6,894	4,225	1,062
2010	98,386	1,067	4,438	11,817	12,930	12,171	12,813	15,606	13,434	8,045	4,536	1,528
2012	98,773	1,054	4,312	11,616	12,857	12,491	12,327	15,001	13,954	8,891	4,613	1,656
2013	98,937	1,058	4,270	11,529	12,794	12,612	12,193	14,654	14,108	9,332	4,677	1,711
2014	99,042	1,055	4,254	11,409	12,700	12,739	12,055	14,359	14,252	9,715	4,753	1,752
White, not Hispanic or Latina female												
1980	92,872	1,240	4,522	12,647	16,185	14,711	10,468	9,700	9,935	7,707	4,345	1,411
1990	96,557	1,280	4,909	11,846	12,749	15,872	14,520	10,153	9,116	8,674	5,491	1,945
2000	100,774	1,102	4,517	12,529	12,183	12,778	16,089	14,446	9,879	8,188	6,429	2,633
2010	101,741	1,016	4,225	11,219	12,426	11,972	12,718	15,839	14,049	9,000	6,125	3,150
2012	101,926	1,005	4,108	11,035	12,286	12,254	12,210	15,208	14,618	9,859	6,057	3,286
2013	101,982	1,007	4,063	10,958	12,194	12,358	12,073	14,844	14,785	10,314	6,064	3,321
2014	102,007	1,007	4,047	10,852	12,083	12,460	11,927	14,522	14,938	10,730	6,102	3,338

--- Data not available.

¹Population for age group under 5 years.

²Population for age group 75 years and over.

NOTES: The race groups, white, black, American Indian or Alaska Native, and Asian or Pacific Islander, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with *Health, United States, 2003*, population estimates for 1991–1999 are intercensal estimates based on the 1990 and 2000 censuses. Starting with *Health, United States, 2012*, population estimates for 2001–2009 are intercensal estimates based on the 2000 and 2010 censuses. Population estimates for 2011 and beyond are 2010-based postcensal estimates. Population figures are census counts as of April 1 for 1950, 1960, 1970, 1980, and 1990. For 2000 and 2010, population estimates are bridged-race April 1 census counts. Estimates for other years are as of July 1. See Appendix I, Population Census and Population Estimates. Populations for age groups may not sum to the total due to rounding. Unrounded population figures are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: U.S. Census Bureau: 1950 Nonwhite Population by Race. Special Report P-E, No. 3B. Washington, DC: U.S. Government Printing Office, 1951; U.S. Census of Population: 1960, Number of Inhabitants, PC(1)-A1, United States Summary, 1964; 1970, Number of Inhabitants, Final Report PC(1)-A1, United States Summary, 1971; U.S. population estimates, by age, sex, race, and Hispanic origin: 1980 to 1991. Current population reports, series P-25, no 1095. Washington, DC: U.S. Government Printing Office, Feb. 1993; NCHS. Estimates of the July 1, 1991–July 1, 1999; April 1, 2000; July 1, 2001–July 1, 2009; April 1, 2010; July 1, 2011–July 1, 2014 United States resident population by age, sex, race, and Hispanic origin, prepared under a collaborative arrangement with the U.S. Census Bureau, Population Estimates Program. Available from: http://www.cdc.gov/nchs/nvss/bridged_race.htm. See Appendix I, Population Census and Population Estimates.

Table 2 (page 1 of 2). Persons below poverty level, by selected characteristics, race, and Hispanic origin: United States, selected years 1973–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#002>.

[Data are based on household interviews of the civilian noninstitutionalized population]

<i>Selected characteristic, race, and Hispanic origin</i> ¹	1973	1980	1990	2000 ²	2010 ⁴	2012	2013(1) ⁵	2013(2) ⁵	2014 ⁶
All persons									
Percent below poverty									
All races	11.1	13.0	13.5	11.3	15.1	15.0	14.5	14.8	14.8
White only	8.4	10.2	10.7	9.5	13.0	12.7	12.3	12.9	12.7
Black or African American only	31.4	32.5	31.9	22.5	27.4	27.2	27.2	25.2	26.2
Asian only	---	---	12.2	9.9	12.2	11.7	10.5	13.1	12.0
Hispanic or Latino	21.9	25.7	28.1	21.5	26.5	25.6	23.5	24.7	23.6
Mexican	---	---	28.1	22.9	---	---	---	---	---
Puerto Rican	---	---	40.6	25.6	---	---	---	---	---
White only, not Hispanic or Latino	7.5	9.1	8.8	7.4	9.9	9.7	9.6	10.0	10.1
Related children under age 18 in families									
All races	14.2	17.9	19.9	15.6	21.5	21.3	19.5	20.9	20.7
White only	9.7	13.4	15.1	12.4	17.9	17.9	15.9	18.4	17.4
Black or African American only	40.6	42.1	44.2	30.9	39.0	37.5	38.0	33.8	37.1
Asian only	---	---	17.0	12.5	14.0	13.3	9.8	14.4	13.4
Hispanic or Latino	27.8	33.0	37.7	27.6	34.3	33.3	30.0	32.2	31.3
Mexican	---	---	35.5	29.5	---	---	---	---	---
Puerto Rican	---	---	56.7	32.1	---	---	---	---	---
White only, not Hispanic or Latino	---	11.3	11.6	8.5	11.7	11.8	10.1	12.7	11.9
Related children under age 18 in families with female householder and no husband present									
All races	---	50.8	53.4	40.1	46.6	47.2	45.8	47.4	46.5
White only	---	41.6	45.9	33.9	43.3	44.2	41.6	46.5	42.9
Black or African American only	---	64.8	64.7	49.3	53.2	53.3	54.0	49.6	52.8
Asian only	---	---	32.2	38.0	36.9	33.0	22.7	47.4	32.4
Hispanic or Latino	---	65.0	68.4	49.8	56.3	54.7	52.3	53.4	53.3
Mexican	---	---	62.4	51.4	---	---	---	---	---
Puerto Rican	---	---	82.7	55.3	---	---	---	---	---
White only, not Hispanic or Latino	---	---	39.6	28.0	34.7	36.5	33.6	39.5	35.8
All persons									
Number below poverty, in thousands									
All races	22,973	29,272	33,585	31,581	46,343	46,496	45,318	46,269	46,657
White only	15,142	19,699	22,326	21,645	31,083	30,816	29,936	31,287	31,089
Black or African American only	7,388	8,579	9,837	7,982	10,746	10,911	11,041	10,186	10,755
Asian only	---	---	858	1,258	1,899	1,921	1,785	2,255	2,137
Hispanic or Latino	2,366	3,491	6,006	7,747	13,522	13,616	12,744	13,356	13,104
Mexican	---	---	3,764	5,460	---	---	---	---	---
Puerto Rican	---	---	966	814	---	---	---	---	---
White only, not Hispanic or Latino	12,864	16,365	16,622	14,366	19,251	18,940	18,796	19,552	19,652
Related children under age 18 in families									
All races	9,453	11,114	12,715	11,005	15,598	15,437	14,142	15,116	14,987
White only	5,462	6,817	7,696	6,834	9,590	9,547	8,428	9,702	9,172
Black or African American only	3,822	3,906	4,412	3,495	4,271	4,097	4,153	3,678	4,036
Asian only	---	---	356	407	477	470	354	538	492
Hispanic or Latino	1,364	1,718	2,750	3,342	5,815	5,773	5,273	5,638	5,522
Mexican	---	---	1,733	2,537	---	---	---	---	---
Puerto Rican	---	---	490	329	---	---	---	---	---
White only, not Hispanic or Latino	---	5,174	5,106	3,715	4,544	4,510	3,833	4,784	4,440

See footnotes at end of table.

Table 2 (page 2 of 2). Persons below poverty level, by selected characteristics, race, and Hispanic origin: United States, selected years 1973–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#002>.

[Data are based on household interviews of the civilian noninstitutionalized population]

Selected characteristic, race, and Hispanic origin ¹	1973	1980	1990	2000 ²	2010 ⁴	2012	2013(1) ⁵	2013(2) ⁵	2014 ⁶
Related children under age 18 in families with female householder and no husband present	Number below poverty, in thousands								
All races	---	5,866	7,363	6,300	8,603	8,664	8,305	9,025	8,491
White only	---	2,813	3,597	3,090	4,495	4,598	4,316	5,155	4,426
Black or African American only	---	2,944	3,543	2,908	3,252	3,165	3,180	2,964	3,121
Asian only	---	---	80	162	141	128	89	159	136
Hispanic or Latino	---	809	1,314	1,407	2,707	2,809	2,763	3,069	2,739
Mexican	---	---	615	938	---	---	---	---	---
Puerto Rican	---	---	382	242	---	---	---	---	---
White only, not Hispanic or Latino	---	---	2,411	1,832	2,209	2,245	2,001	2,477	2,174

--- Data not available.

¹The race groups, white, black, and Asian, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 2002 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The three single-race categories shown in the table conform to the 1997 Standards. For 2002 and subsequent years, race-specific estimates are for persons who reported only one racial group. Estimates for single-race categories prior to 2002 are based on answers to the Current Population Survey question which asked respondents to choose only a single race. Prior to data year 2002, data were tabulated according to the 1977 Standards in which the Asian only category included Native Hawaiian or Other Pacific Islander. See Appendix II, Hispanic origin; Race.

²Estimates are consistent with 2001–2009 data through implementation of the 2000 census-based population controls and a 28,000-household sample expansion.

³Data for 2004 (shown in spreadsheet version) reflect a correction to the weights in the 2005 Annual Social and Economic (ASEC) Supplements of the Current Population Survey. See Appendix I, Current Population Survey (CPS).

⁴Data for 2010 and beyond reflect Census 2010-based population controls.

⁵For 2013 data, the CPS ASEC used a split panel to test a new set of income questions. Estimates for 2013 shown in the column labeled (1) are based on the approximately 68,000 addresses that received questions consistent with those for 2012 and earlier ASECs. Estimates for 2013 shown in the column labeled (2) are based on the approximately 30,000 addresses that received the new set of income questions. The vertical line in the table indicates the introduction of the new set of income questions into the ASEC estimates.

⁶Data for 2013(2) and beyond are based on a redesigned questionnaire that includes the new set of income questions; therefore data trends need to be interpreted with caution. For more information on the redesigned questionnaire and the impact of the new income questions on poverty estimates, see: <http://www.census.gov/content/dam/Census/library/publications/2015/demo/p60-252.pdf>.

NOTES: Estimates of poverty for 1991–1998 are based on 1990 postcensal population estimates. Estimates for 1999–2009 were based on Census 2000 population controls. Estimates for 2010 and beyond were based on Census 2010 population controls. Poverty level is based on family income and family size using U.S. Census Bureau poverty thresholds. See Appendix II, Poverty. Poverty estimates based on a supplemental poverty measure are available from the U.S. Census Bureau. In 2011–2013, an estimated 30.1% of American Indian or Alaska Native only persons (1,005,000 persons) were living below the poverty level, and an estimated 15.9% of Native Hawaiian or Other Pacific Islander only persons (176,000 persons) were living below the poverty level. Due to the redesign of the CPS ASEC income questions, 2013 is the last year that data were available to compute three-year estimates for the American Indian or Alaska Native only populations and the Native Hawaiian or Other Pacific Islander populations. Estimates for these groups will not be updated until 2016 estimates are available. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: U.S. Census Bureau, Current Population Survey, Annual Social and Economic Supplements; DeNavas-Walt C, Proctor BD. Income and poverty in the United States: 2014. Current Population Reports, P60–252. Washington, DC: U.S. Government Printing Office. 2015. Available from: <http://www.census.gov/content/dam/Census/library/publications/2015/demo/p60-252.pdf>. See Appendix I, Current Population Survey (CPS).

Table 3 (page 1 of 3). Crude birth rates, fertility rates, and birth rates, by age, race, and Hispanic origin of mother: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#003>.

[Data are based on birth certificates]

Race, Hispanic origin, and year	Crude birth rate ¹	Fertility rate ²	Age of mother									
			10–14 years	15–19 years			20–24 years	25–29 years	30–34 years	35–39 years	40–44 years	45–54 years ³
				Total	15–17 years	18–19 years						
All races												
Live births per 1,000 women												
1950	24.1	106.2	1.0	81.6	40.7	132.7	196.6	166.1	103.7	52.9	15.1	1.2
1960	23.7	118.0	0.8	89.1	43.9	166.7	258.1	197.4	112.7	56.2	15.5	0.9
1970	18.4	87.9	1.2	68.3	38.8	114.7	167.8	145.1	73.3	31.7	8.1	0.5
1980	15.9	68.4	1.1	53.0	32.5	82.1	115.1	112.9	61.9	19.8	3.9	0.2
1990	16.7	70.9	1.4	59.9	37.5	88.6	116.5	120.2	80.8	31.7	5.5	0.2
1995	14.6	64.6	1.3	56.0	35.5	87.7	107.5	108.8	81.1	34.0	6.6	0.3
2000	14.4	65.9	0.9	47.7	26.9	78.1	109.7	113.5	91.2	39.7	8.0	0.5
2004	14.0	66.4	0.6	40.5	21.8	68.7	101.5	116.5	96.2	45.5	9.0	0.5
2005	14.0	66.7	0.6	39.7	21.1	68.4	101.8	116.5	96.7	46.4	9.1	0.6
2010	13.0	64.1	0.4	34.2	17.3	58.2	90.0	108.3	96.5	45.9	10.2	0.7
2012	12.6	63.0	0.4	29.4	14.1	51.4	83.1	106.5	97.3	48.3	10.4	0.7
2013	12.4	62.5	0.3	26.5	12.3	47.1	80.7	105.5	98.0	49.3	10.4	0.8
2014	12.5	62.9	0.3	24.2	10.9	43.8	79.0	105.8	100.8	51.0	10.6	0.8
Race of child: ⁴ White												
1950	23.0	102.3	0.4	70.0	31.3	120.5	190.4	165.1	102.6	51.4	14.5	1.0
1960	22.7	113.2	0.4	79.4	35.5	154.6	252.8	194.9	109.6	54.0	14.7	0.8
1970	17.4	84.1	0.5	57.4	29.2	101.5	163.4	145.9	71.9	30.0	7.5	0.4
1980	14.9	64.7	0.6	44.7	25.2	72.1	109.5	112.4	60.4	18.5	3.4	0.2
Race of mother: ⁵ White												
1980	15.1	65.6	0.6	45.4	25.5	73.2	111.1	113.8	61.2	18.8	3.5	0.2
1990	15.8	68.3	0.7	50.8	29.5	78.0	109.8	120.7	81.7	31.5	5.2	0.2
1995	14.1	63.6	0.8	49.5	29.6	80.2	104.7	111.7	83.3	34.2	6.4	0.3
2000	13.9	65.3	0.6	43.2	23.3	72.3	106.6	116.7	94.6	40.2	7.9	0.4
2004	13.6	66.5	0.5	37.4	19.4	64.4	99.8	120.8	100.3	46.7	8.9	0.5
2005	13.6	66.8	0.5	36.7	18.8	64.0	99.9	120.7	100.7	47.6	9.0	0.6
2010	12.5	64.4	0.3	31.9	15.8	54.8	87.9	111.9	100.5	46.4	10.0	0.6
2012	12.1	63.0	0.3	27.4	13.0	48.3	80.8	109.2	100.2	48.5	10.0	0.7
2013	12.0	62.7	0.2	24.9	11.3	44.7	78.5	108.3	101.3	49.6	10.1	0.7
2014	12.0	63.2	0.2	23.0	10.2	42.0	77.3	108.6	103.9	51.2	10.2	0.7
Race of child: ⁴ Black or African American												
1960	31.9	153.5	4.3	156.1	---	---	295.4	218.6	137.1	73.9	21.9	1.1
1970	25.3	115.4	5.2	140.7	101.4	204.9	202.7	136.3	79.6	41.9	12.5	1.0
1980	22.1	88.1	4.3	100.0	73.6	138.8	146.3	109.1	62.9	24.5	5.8	0.3
Race of mother: ⁵ Black or African American												
1980	21.3	84.7	4.3	97.8	72.5	135.1	140.0	103.9	59.9	23.5	5.6	0.3
1990	22.4	86.8	4.9	112.8	82.3	152.9	160.2	115.5	68.7	28.1	5.5	0.3
1995	17.8	71.0	4.1	94.4	68.5	135.0	133.7	95.6	63.0	28.4	6.0	0.3
2000	17.0	70.0	2.3	77.4	49.0	118.8	141.3	100.3	65.4	31.5	7.2	0.4
2004	15.9	67.2	1.6	61.7	36.3	101.3	127.5	104.4	67.8	33.8	7.9	0.5
2005	16.1	68.5	1.6	60.1	34.5	101.2	129.5	107.0	70.2	35.1	8.4	0.5
2010	15.1	66.3	1.0	51.1	27.3	84.8	118.1	101.8	73.0	36.4	9.3	0.7
2012	14.7	65.1	0.8	44.0	22.0	74.4	108.7	101.7	75.1	39.2	9.7	0.7
2013	14.5	64.7	0.7	39.1	19.0	67.3	105.5	102.6	77.3	40.5	10.0	0.8
2014	14.5	64.6	0.6	35.1	16.7	61.9	102.6	103.1	79.4	42.6	10.2	0.8
American Indian or Alaska Native mother ⁵												
1980	20.7	82.7	1.9	82.2	51.5	129.5	143.7	106.6	61.8	28.1	8.2	*
1990	18.9	76.2	1.6	81.1	48.5	129.3	148.7	110.3	61.5	27.5	5.9	*
1995	15.3	63.0	1.6	72.9	44.6	122.2	123.1	91.6	56.5	24.3	5.5	*
2000	14.0	58.7	1.1	58.3	34.1	97.1	117.2	91.8	55.5	24.6	5.7	0.3
2004	12.8	54.2	0.8	47.2	26.7	79.9	105.4	87.1	51.9	23.9	5.6	0.2
2005	12.6	53.6	0.8	46.0	26.3	78.0	102.9	86.3	51.8	23.3	5.4	0.3
2010	11.0	48.6	0.5	38.7	20.1	66.1	91.0	74.4	48.4	22.3	5.2	0.3
2012	10.5	47.0	0.5	34.9	17.0	60.5	81.7	73.9	49.7	23.3	5.5	0.5
2013	10.3	46.4	0.4	31.1	15.9	53.3	78.9	75.6	50.4	24.7	5.5	0.3
2014	9.9	44.8	0.3	27.3	13.2	48.6	73.2	74.7	52.3	24.1	5.5	0.3

See footnotes at end of table.

Table 3 (page 2 of 3). Crude birth rates, fertility rates, and birth rates, by age, race, and Hispanic origin of mother: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#003>.

[Data are based on birth certificates]

Race, Hispanic origin, and year	Crude birth rate ¹	Fertility rate ²	Age of mother									
			10–14 years	15–19 years			20–24 years	25–29 years	30–34 years	35–39 years	40–44 years	45–54 years ³
				Total	15–17 years	18–19 years						
Live births per 1,000 women												
Asian or Pacific Islander mother ⁵												
1980	19.9	73.2	0.3	26.2	12.0	46.2	93.3	127.4	96.0	38.3	8.5	0.7
1990	19.0	69.6	0.7	26.4	16.0	40.2	79.2	126.3	106.5	49.6	10.7	1.1
1995	16.7	62.6	0.7	25.5	15.6	40.1	64.2	103.7	102.3	50.1	11.8	0.8
2000	17.1	65.8	0.3	20.5	11.6	32.6	60.3	108.4	116.5	59.0	12.6	0.8
2004	16.4	64.5	0.2	16.0	8.4	26.6	53.3	100.4	118.3	62.2	13.6	1.0
2005	15.9	63.0	0.2	15.4	7.7	26.4	52.9	96.6	115.3	61.8	13.7	1.0
2010	14.5	59.2	0.1	10.9	5.1	18.7	42.6	91.5	113.6	62.8	15.1	1.2
2012	15.1	62.2	0.1	9.7	4.1	17.7	41.4	95.8	121.3	68.1	16.1	1.4
2013	14.3	59.2	0.1	8.7	3.7	16.1	39.1	89.5	114.6	66.6	16.1	1.5
2014	14.6	60.7	0.1	7.7	3.3	13.9	37.5	90.0	121.3	68.9	16.1	1.5
Hispanic or Latina mother ^{5,6}												
1980	23.5	95.4	1.7	82.2	52.1	126.9	156.4	132.1	83.2	39.9	10.6	0.7
1990	26.7	107.7	2.4	100.3	65.9	147.7	181.0	153.0	98.3	45.3	10.9	0.7
1995	24.1	98.8	2.6	99.3	68.3	145.4	171.9	140.4	90.5	43.7	10.7	0.6
2000	23.1	95.9	1.7	87.3	55.5	132.6	161.3	139.9	97.1	46.6	11.5	0.6
2004	22.8	95.7	1.2	78.1	47.3	124.8	159.2	144.7	103.4	52.2	12.3	0.7
2005	22.9	96.4	1.3	76.5	45.8	124.4	161.1	147.0	105.6	53.3	12.8	0.8
2010	18.7	80.2	0.8	55.7	32.3	90.7	126.1	125.3	96.6	51.7	13.0	0.8
2012	17.1	74.4	0.6	46.3	25.5	77.2	111.5	119.6	94.3	51.6	13.2	0.8
2013	16.7	72.9	0.5	41.7	22.0	70.8	107.2	119.1	94.8	52.4	13.3	0.8
2014	16.5	72.1	0.4	38.0	19.3	66.1	104.5	118.7	96.5	53.6	13.5	0.9
White, not Hispanic or Latina mother ^{5,6}												
1980	14.2	62.4	0.4	41.2	22.4	67.7	105.5	110.6	59.9	17.7	3.0	0.1
1990	14.4	62.8	0.5	42.5	23.2	66.6	97.5	115.3	79.4	30.0	4.7	0.2
1995	12.5	57.5	0.4	39.3	22.0	66.2	90.2	105.1	81.5	32.8	5.9	0.3
2000	12.2	58.5	0.3	32.6	15.8	57.5	91.2	109.4	93.2	38.8	7.3	0.4
2004	11.7	58.9	0.2	26.7	12.0	48.6	83.0	112.1	98.3	45.1	8.3	0.5
2005	11.6	59.0	0.2	26.0	11.5	48.0	82.7	111.7	98.4	46.0	8.3	0.5
2010	10.9	58.7	0.2	23.5	10.0	42.5	74.9	105.8	99.9	44.1	9.2	0.6
2012	10.7	58.6	0.2	20.5	8.4	37.9	70.2	104.4	100.5	46.8	9.1	0.6
2013	10.7	58.7	0.1	18.6	7.4	35.0	68.3	103.5	101.9	48.0	9.1	0.7
2014	10.8	59.5	0.1	17.3	6.7	32.9	67.1	103.9	104.7	49.6	9.1	0.7
Black or African American, not Hispanic or Latina mother ^{5,6}												
1980	22.9	90.7	4.6	105.1	77.2	146.5	152.2	111.7	65.2	25.8	5.8	0.3
1990	23.0	89.0	5.0	116.2	84.9	157.5	165.1	118.4	70.2	28.7	5.6	0.3
1995	18.2	72.8	4.2	97.2	70.4	139.2	137.8	98.5	64.4	28.8	6.1	0.3
2000	17.3	71.4	2.4	79.2	50.1	121.9	145.4	102.8	66.5	31.8	7.2	0.4
2004	15.8	67.1	1.6	61.9	36.4	101.6	127.9	105.0	67.8	33.6	7.8	0.5
2005	15.8	67.2	1.6	59.4	34.1	100.2	127.9	105.5	68.8	34.2	8.2	0.5
2010	15.1	66.6	1.0	51.5	27.4	85.6	119.4	102.5	73.6	36.4	9.2	0.7
2012	14.6	65.0	0.8	43.9	21.9	74.1	109.0	101.7	75.1	38.9	9.6	0.7
2013	14.4	64.6	0.7	39.0	18.9	67.0	105.6	102.7	77.3	40.3	9.9	0.8
2014	14.4	64.5	0.6	34.9	16.6	61.5	102.8	103.3	79.6	42.5	10.1	0.9

See footnotes at end of table.

Table 3 (page 3 of 3). Crude birth rates, fertility rates, and birth rates, by age, race, and Hispanic origin of mother: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#003>.

[Data are based on birth certificates]

--- Data not available.

* Rates based on fewer than 20 births are considered unreliable and are not shown.

¹ Live births per 1,000 population.

² Total number of live births regardless of age of mother per 1,000 women aged 15–44.

³ Prior to 1997, data are for live births to mothers aged 45–49 per 1,000 women aged 45–49. In subsequent years, rates were computed by dividing the number of births to women aged 45 and over by the population of women aged 45–49. See Appendix II, Age.

⁴ Live births are tabulated by race of child. See Appendix II, Race.

⁵ Live births are tabulated by race and/or Hispanic origin of mother. See Appendix II, Race.

⁶ Prior to 1993, data from states that did not report Hispanic origin on the birth certificate were excluded. See Appendix II, Hispanic origin. Rates in 1985 were not calculated because estimates for the Hispanic and non-Hispanic populations were not available.

NOTES: Data are based on births adjusted for underregistration for 1950 and on registered births for all other years. Starting with 1970 data, births to persons who were not residents of the 50 states and the District of Columbia are excluded. Starting with *Health, United States, 2003*, rates for 1991–1999 were revised using intercensal population estimates based on the 1990 and 2000 censuses. Rates for 2000 were based on bridged-race April 1, 2000 census counts. Starting with *Health, United States, 2012*, rates for 2001–2009 were revised using intercensal population estimates based on the 2000 and 2010 censuses. Rates for 2010 were based on bridged-race April 1, 2010 census counts. Rates for 2011 and beyond were computed using 2010-based postcensal estimates. See Appendix I, Population Census and Population Estimates. The race groups, white, black, American Indian or Alaska Native, and Asian or Pacific Islander, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 2003 data, some states reported multiple-race data. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards, for comparability with other states. See Appendix II, Race. Interpretation of trend data for Hispanic women should take into consideration expansion of reporting areas. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Vital Statistics System, public-use Birth File. Hamilton BE, Martin JA, Osterman MJK, et al. Births: Final data for 2014. National vital statistics reports; vol 64 no 12. Hyattsville, MD: NCHS. 2015; Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr64/nvsr64_12.pdf. Ventura SJ. Births of Hispanic parentage, 1980 and 1985. Monthly vital statistics report; vol 32 no 6 and vol 36 no 11, suppl. Public Health Service. Hyattsville, MD. 1983 and 1988; Available from: http://www.cdc.gov/nchs/data/mvsr/supp/mv32_06sacc.pdf and http://www.cdc.gov/nchs/data/mvsr/supp/mv36_11s.pdf. Internet release of: Vital statistics of the United States, 2003, vol 1, Natality, Tables 1–1 and 1–7; available from: <http://www.cdc.gov/nchs/products/vsus.htm#electronic>. See Appendix I, National Vital Statistics System (NVSS).

Table 4. Nonmarital childbearing, by detailed race and Hispanic origin of mother, and maternal age: United States, selected years 1970–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#004>.

[Data are based on birth certificates]

Maternal race, Hispanic origin, and age	1970	1980	1990	1995	2000	2005	2010	2012	2013	2014
Live births per 1,000 unmarried women aged 15–44 ¹										
All races and origins	26.4	29.4	43.8	44.3	44.1	47.2	47.5	45.3	44.3	43.9
White ²	13.9	18.1	32.9	37.0	38.2	43.2	44.5	42.1	40.8	40.6
Black or African American ²	95.5	81.1	90.5	74.5	70.5	67.2	65.3	62.6	61.7	61.5
Asian or Pacific Islander	---	---	---	---	20.9	22.8	22.3	22.9	21.8	21.7
Hispanic or Latina ³	---	---	89.6	88.8	87.2	96.2	80.6	72.6	69.9	68.5
White, not Hispanic or Latina ³	---	---	24.4	28.1	28.0	30.4	32.9	32.1	31.7	31.8
Percent of live births to unmarried mothers										
All races and origins	10.7	18.4	28.0	32.2	33.2	36.9	40.8	40.7	40.6	40.2
White.	5.5	11.2	20.4	25.3	27.1	31.7	35.9	35.9	35.8	35.7
Black or African American	37.5	56.1	66.5	69.9	68.5	69.3	72.1	71.6	71.0	70.4
American Indian or Alaska Native	22.4	39.2	53.6	57.2	58.4	63.5	65.6	66.9	66.4	65.7
Asian or Pacific Islander ⁴	---	7.3	13.2	16.3	14.8	16.2	17.0	17.0	17.0	16.4
Hispanic or Latina ³	---	23.6	36.7	40.8	42.7	48.0	53.4	53.5	53.2	52.9
Mexican	---	20.3	33.3	38.1	40.7	46.7	52.0	52.1	51.9	51.6
Puerto Rican	---	46.3	55.9	60.0	59.6	61.7	65.2	65.1	64.6	63.9
Cuban	---	10.0	18.2	23.8	27.3	36.4	47.0	48.8	50.1	49.8
Central and South American	---	27.1	41.2	44.1	44.7	49.2	51.8	50.8	50.1	50.4
Other and unknown Hispanic or Latina	---	22.4	37.2	44.0	46.2	48.6	56.3	56.4	56.1	55.5
Not Hispanic or Latina: ³										
White	---	9.5	16.9	21.2	22.1	25.3	29.0	29.3	29.3	29.2
Black or African American	---	57.2	66.7	70.0	68.7	69.9	72.5	72.1	71.5	70.9
Number of live births, in thousands										
Live births to unmarried mothers	399	666	1,165	1,254	1,347	1,527	1,633	1,610	1,596	1,605
Percent distribution of live births to unmarried mothers										
Under 20 years.	50.1	40.8	30.9	30.9	28.0	23.1	20.1	17.1	15.4	13.9
20–24 years	31.8	35.6	34.7	34.5	37.4	38.3	36.8	36.9	36.8	36.1
25 years and over.	18.1	23.5	34.4	34.7	34.6	38.7	43.1	46.1	47.9	50.0

--- Data not available.

¹Rates computed by dividing births to unmarried mothers, regardless of age of mother, by the population of unmarried women aged 15–44. Population data for unmarried American Indian or Alaska Native women are not available for rate calculations. Prior to 2000, population data for unmarried Asian or Pacific Islander women were not available for rate calculations.

²For 1970 and 1975, birth rates are by race of child.

³Prior to 1993, data from states that did not report Hispanic origin on the birth certificate were excluded. See Appendix II, Hispanic origin. Data for non-Hispanic white and non-Hispanic black women for years prior to 1989 are not nationally representative and are provided solely for comparison with Hispanic data.

⁴Estimates are not available for Asian or Pacific Islander subgroups because not all states have adopted the 2003 revision of the U.S. Standard Certificate of Live Birth. See Appendix II, Race.

NOTES: National estimates for 1970 and 1975 for unmarried mothers are based on births occurring in states reporting marital status of mother. Changes in reporting procedures for marital status occurred in some states during the 1990s. Data for states in which marital status was not reported have been inferred and included with data from the remaining states. See Appendix II, Marital status. Interpretation of trend data for Hispanic births should take into consideration expansion of reporting areas. The race groups, white, black, American Indian or Alaska Native, and Asian or Pacific Islander, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 2003 data, some states reported multiple-race data. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards, for comparability with other states. See Appendix II, Race. Starting with *Health, United States, 2003*, rates for 1991–1999 were revised using intercensal population estimates based on the 1990 and 2000 censuses. Rates for 2000 were based on bridged-race April 1, 2000 census counts. Starting with *Health, United States, 2012*, rates for 2001–2009 were revised using intercensal population estimates based on the 2000 and 2010 censuses. Rates for 2010 were based on 2010 bridged-race April 2010 census counts. Rates for 2011 and beyond were computed using 2010-based postcensal estimates. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Vital Statistics System, public-use Birth File. Hamilton BE, Martin JA, Osterman MJK, et al. Births: Final data for 2014. National vital statistics reports; vol 64 no 12. Hyattsville, MD: NCHS. 2015; Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr64/nvsr64_12.pdf. Hamilton BE, Sutton PD, Ventura SJ. Revised birth and fertility rates for the 1990s and new rates for Hispanic populations, 2000 and 2001: United States. National vital statistics reports; vol 51 no 12. Hyattsville, MD: NCHS. 2003; Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr51/nvsr51_12.pdf. Births: Final data for each data year 1997–2007. National vital statistics reports. Hyattsville, MD; Final natality statistics for each data year 1993–1996. Monthly vital statistics report. Hyattsville, MD; Ventura SJ. Births to unmarried mothers: United States, 1980–1992. Vital Health Stat 21(53). 1995. See Appendix I, National Vital Statistics System (NVSS).

Table 5. Low birthweight live births, by detailed race and Hispanic origin of mother: United States, selected years 1970–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#005>.

[Data are based on birth certificates]

<i>Birthweight, maternal race, and Hispanic origin</i>	1970	1980	1990	2000	2005	2006	2010	2012	2013	2014
Low birthweight (less than 2,500 grams)					Percent of live births ¹					
All races	7.93	6.84	6.97	7.57	8.19	8.26	8.15	7.99	8.02	8.00
White	6.85	5.72	5.70	6.55	7.16	7.21	7.08	6.96	7.00	6.98
Black or African American	13.90	12.69	13.25	12.99	13.59	13.59	13.21	12.84	12.76	12.83
American Indian or Alaska Native	7.97	6.44	6.11	6.76	7.36	7.52	7.61	7.61	7.48	7.65
Asian or Pacific Islander ²	---	6.68	6.45	7.31	7.98	8.12	8.49	8.21	8.34	8.05
Hispanic or Latina ³	---	6.12	6.06	6.41	6.88	6.99	6.97	6.97	7.09	7.05
Mexican	---	5.62	5.55	6.01	6.49	6.58	6.49	6.48	6.62	6.58
Puerto Rican	---	8.95	8.99	9.30	9.92	10.14	9.55	9.40	9.38	9.54
Cuban	---	5.62	5.67	6.49	7.64	7.14	7.30	7.43	7.35	7.48
Central and South American	---	5.76	5.84	6.34	6.78	6.81	6.55	6.64	6.85	6.68
Other and unknown Hispanic or Latina	---	6.96	6.87	7.84	8.27	8.54	8.38	8.00	7.99	7.94
Not Hispanic or Latina: ³										
White	---	5.69	5.61	6.60	7.29	7.32	7.14	6.97	6.98	6.96
Black or African American	---	12.71	13.32	13.13	14.02	13.97	13.53	13.18	13.08	13.17
Very low birthweight (less than 1,500 grams)										
All races	1.17	1.15	1.27	1.43	1.49	1.49	1.45	1.42	1.41	1.40
White	0.95	0.90	0.95	1.14	1.20	1.20	1.17	1.15	1.14	1.14
Black or African American	2.40	2.48	2.92	3.07	3.15	3.05	2.90	2.85	2.82	2.79
American Indian or Alaska Native	0.98	0.92	1.01	1.16	1.17	1.28	1.28	1.33	1.32	1.27
Asian or Pacific Islander ²	---	0.92	0.87	1.05	1.14	1.12	1.17	1.13	1.18	1.15
Hispanic or Latina ³	---	0.98	1.03	1.14	1.20	1.19	1.20	1.22	1.21	1.23
Mexican	---	0.92	0.92	1.03	1.12	1.12	1.09	1.13	1.13	1.13
Puerto Rican	---	1.29	1.62	1.93	1.87	1.91	1.82	1.77	1.65	1.86
Cuban	---	1.02	1.20	1.21	1.50	1.28	1.42	1.55	1.27	1.45
Central and South American	---	0.99	1.05	1.20	1.19	1.13	1.09	1.13	1.15	1.12
Other and unknown Hispanic or Latina	---	1.01	1.09	1.42	1.36	1.36	1.46	1.38	1.37	1.38
Not Hispanic or Latina: ³										
White	---	0.87	0.93	1.14	1.21	1.20	1.16	1.13	1.11	1.10
Black or African American	---	2.47	2.93	3.10	3.27	3.15	2.98	2.94	2.90	2.87

--- Data not available.

¹Excludes live births with unknown birthweight. Percentage based on live births with known birthweight. See Appendix II, Birthweight.

²Estimates are not available for Asian or Pacific Islander subgroups because not all states have adopted the 2003 revision of the U.S. Standard Certificate of Live Birth. See Appendix II, Race.

³Prior to 1993, data from states that did not report Hispanic origin on the birth certificate were excluded. See Appendix II, Hispanic origin. Data for non-Hispanic white and non-Hispanic black women for years prior to 1989 are not nationally representative and are provided solely for comparison with Hispanic data.

NOTES: The race groups, white, black, American Indian or Alaska Native, and Asian or Pacific Islander, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 2003 data, some states reported multiple-race data. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards, for comparability with other states. See Appendix II, Race. Interpretation of trend data for Hispanic births should take into consideration expansion of reporting areas. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Vital Statistics System, public-use Birth File. Hamilton BE, Martin JA, Osterman MJK, et al. Births: Final data for 2014. National vital statistics reports; vol 64 no 12. Hyattsville, MD: NCHS. 2015; Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr64/nvsr64_12.pdf. See Appendix I, National Vital Statistics System (NVSS).

Table 6 (page 1 of 3). Low birthweight live births, by race and Hispanic origin of mother, state, and territory: United States and U.S. dependent areas, 2000–2002, 2003–2005, and 2012–2014

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2015.htm#006>.

[Data are based on birth certificates]

State and territory	Not Hispanic or Latina								
	All races			White			Black or African American		
	2000–2002	2003–2005	2012–2014	2000–2002	2003–2005	2012–2014	2000–2002	2003–2005	2012–2014
	Percent of live births weighing less than 2,500 grams ¹								
United States ²	7.69	8.07	8.01	6.75	7.18	6.97	13.19	13.77	13.14
Alabama	9.75	10.35	10.03	7.77	8.46	7.97	14.10	15.02	14.86
Alaska	5.71	6.02	5.77	4.84	5.34	5.25	10.70	11.74	8.88
Arizona	6.91	7.05	6.95	6.78	7.01	6.53	13.16	12.38	11.72
Arkansas	8.64	9.04	8.79	7.48	7.83	7.59	13.81	14.86	14.07
California	6.29	6.71	6.73	5.86	6.30	5.87	11.66	12.46	11.51
Colorado	8.60	9.04	8.80	8.24	8.81	8.27	14.59	15.20	13.59
Connecticut	7.52	7.74	7.76	6.48	6.60	6.56	12.28	12.88	12.08
Delaware	9.29	9.31	8.29	7.80	7.62	6.84	14.08	14.32	12.22
District of Columbia	11.85	11.06	9.62	6.35	6.28	5.98	14.60	13.96	12.41
Florida	8.18	8.59	8.59	6.98	7.38	7.18	12.58	13.28	12.97
Georgia	8.79	9.27	9.40	6.92	7.44	7.18	12.98	13.81	13.51
Hawaii	7.98	8.23	8.09	6.17	6.42	5.90	11.01	11.44	11.31
Idaho	6.41	6.65	6.59	6.29	6.60	6.46	*	*7.03	*7.23
Illinois	8.04	8.40	8.17	6.74	7.22	6.86	14.04	14.70	13.65
Indiana	7.54	8.10	7.93	6.95	7.54	7.30	12.89	13.46	12.89
Iowa	6.39	6.92	6.65	6.19	6.72	6.35	11.77	12.22	11.19
Kansas	6.96	7.28	7.06	6.66	6.97	6.54	12.37	13.42	12.84
Kentucky	8.38	8.86	8.71	7.84	8.50	8.33	13.84	13.52	13.37
Louisiana	10.40	11.02	10.74	7.56	8.12	8.06	14.44	15.33	15.22
Maine	6.12	6.58	7.11	6.13	6.57	6.96	*9.47	8.47	9.59
Maryland	8.88	9.17	8.62	6.79	7.19	6.64	13.00	13.13	12.17
Massachusetts	7.26	7.77	7.58	6.56	7.15	6.90	11.54	11.82	10.32
Michigan	7.94	8.28	8.34	6.55	7.00	6.99	14.24	14.43	13.60
Minnesota	6.23	6.43	6.52	5.80	5.93	5.94	10.54	10.71	9.56
Mississippi	10.82	11.62	11.49	7.97	8.67	8.23	14.48	15.60	15.96
Missouri	7.74	8.12	7.98	6.79	7.18	6.96	13.27	13.90	13.68
Montana	6.65	7.02	7.38	6.60	6.81	6.98	*	*15.58	*12.25
Nebraska	6.88	6.97	6.59	6.52	6.76	6.12	13.07	12.16	11.70
Nevada	7.44	8.11	8.09	7.19	7.78	7.34	13.40	13.98	13.21
New Hampshire	6.40	6.65	7.00	6.24	6.59	6.92	10.58	10.85	10.25
New Jersey	7.89	8.19	8.17	6.59	7.11	7.01	13.20	13.48	12.46
New Mexico	7.99	8.38	8.84	7.89	8.33	8.44	13.88	15.01	14.69
New York	7.76	8.11	7.91	6.48	6.82	6.60	12.02	12.78	12.16
North Carolina	8.90	9.07	8.82	7.49	7.73	7.37	13.83	14.33	13.46
North Dakota	6.28	6.49	6.26	6.13	6.37	6.06	*9.02	*9.43	9.30
Ohio	8.07	8.51	8.52	7.08	7.53	7.38	13.45	13.83	13.58
Oklahoma	7.75	7.92	7.99	7.35	7.63	7.60	13.57	13.62	13.44
Oregon	5.65	6.09	6.23	5.44	6.02	5.99	10.32	11.16	9.43
Pennsylvania	7.93	8.20	8.13	6.78	7.06	7.00	13.79	13.67	12.85
Rhode Island	7.47	8.12	7.34	6.75	7.39	6.47	12.32	11.22	10.90
South Carolina	9.74	10.15	9.56	7.40	7.82	7.41	14.29	15.19	14.36
South Dakota	6.58	6.71	6.33	6.37	6.62	5.92	*11.51	*7.27	9.37
Tennessee	9.20	9.35	9.09	7.95	8.26	7.95	14.23	14.51	13.98
Texas	7.54	8.07	8.25	6.81	7.43	7.31	12.82	13.91	13.36
Utah	6.48	6.68	6.94	6.28	6.45	6.66	13.09	12.05	9.82
Vermont	6.15	6.57	6.65	6.12	6.55	6.53	*	*	*8.49
Virginia	7.90	8.23	8.01	6.54	7.01	6.63	12.56	12.83	12.54
Washington	5.75	6.13	6.33	5.43	5.63	5.81	10.34	10.63	9.74
West Virginia	8.60	9.16	9.25	8.39	9.03	9.10	13.81	13.15	13.39
Wisconsin	6.58	6.93	7.16	5.83	6.18	6.36	13.25	13.59	13.39
Wyoming	8.35	8.71	8.78	8.12	8.74	8.62	*13.29	*	*11.90
American Samoa ³	3.51	3.75	4.40	---	---	---	---	---	---
Guam ³	7.88	8.81	8.36	*4.13	*4.01	*	*	*	*
Northern Marianas ³	8.05	7.55	7.16	---	---	*	---	---	*
Puerto Rico ³	11.14	11.92	10.99	---	---	10.73	---	---	14.13
Virgin Islands ³	10.21	11.14	---	*8.37	*5.90	---	9.89	12.51	---

See footnotes at end of table.

Table 6 (page 2 of 3). Low birthweight live births, by race and Hispanic origin of mother, state, and territory: United States and U.S. dependent areas, 2000–2002, 2003–2005, and 2012–2014

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2015.htm#006>.

[Data are based on birth certificates]

State and territory	Hispanic or Latina ⁴			American Indian or Alaska Native ⁵			Asian or Pacific Islander ⁵		
	2000–2002	2003–2005	2012–2014	2000–2002	2003–2005	2012–2014	2000–2002	2003–2005	2012–2014
	Percent of live births weighing less than 2,500 grams ¹								
United States ²	6.48	6.79	7.04	7.11	7.39	7.58	7.54	7.89	8.20
Alabama	6.95	6.92	6.70	9.68	10.53	9.66	7.38	8.02	8.92
Alaska	6.07	5.31	5.52	5.81	5.86	6.28	7.33	6.57	6.79
Arizona	6.56	6.69	6.72	6.85	7.11	6.84	7.95	7.92	7.74
Arkansas	5.79	6.54	6.60	8.11	8.86	7.63	7.73	6.74	9.60
California	5.66	6.10	6.27	6.21	6.49	6.63	7.15	7.42	7.66
Colorado	8.33	8.53	8.66	9.05	9.45	9.66	10.17	10.26	11.11
Connecticut	8.25	8.49	8.29	10.06	7.45	10.71	8.07	7.83	7.79
Delaware	6.81	7.03	6.36	*	*	*	9.89	9.33	7.82
District of Columbia	8.04	7.46	7.97	*	*	*	*7.00	8.97	7.72
Florida	6.61	6.98	7.27	7.11	7.38	8.24	8.35	8.73	8.54
Georgia	5.77	5.96	6.67	9.29	9.00	10.14	8.18	8.35	8.41
Hawaii	8.00	8.34	8.45	*4.99	*	*	8.45	8.84	8.84
Idaho	6.95	6.67	7.00	6.15	8.31	6.84	7.38	6.67	7.56
Illinois	6.31	6.60	6.89	8.60	9.46	8.75	8.49	8.28	8.84
Indiana	6.09	6.33	6.53	*7.74	*10.00	*9.88	7.41	7.87	7.78
Iowa	6.01	6.12	5.96	7.23	9.15	6.41	7.13	7.71	8.33
Kansas	5.93	6.09	6.38	6.20	7.09	6.83	6.69	7.34	8.84
Kentucky	7.73	6.85	6.49	*7.17	*8.54	*11.91	7.75	7.56	7.52
Louisiana	6.56	7.62	7.10	9.06	10.11	7.63	7.89	8.46	8.95
Maine	*6.03	*4.74	9.15	*	*	*7.16	*5.46	8.69	9.01
Maryland	6.73	7.18	7.00	9.74	10.87	7.12	7.42	7.93	8.17
Massachusetts	8.37	8.41	8.26	*7.11	*7.62	*7.69	7.57	7.63	7.83
Michigan	6.26	6.46	7.05	7.26	6.98	7.60	7.46	8.33	9.03
Minnesota	6.02	5.70	6.41	7.10	6.87	8.22	7.28	7.43	7.45
Mississippi	6.61	6.42	7.01	7.30	6.24	*5.12	6.83	8.06	8.65
Missouri	6.18	6.33	6.65	8.67	7.63	8.26	7.34	7.61	7.22
Montana	7.44	8.63	8.21	7.14	7.80	9.16	*5.95	*8.70	10.89
Nebraska	6.30	6.20	6.29	7.27	6.78	6.27	8.05	7.61	7.41
Nevada	6.34	6.74	6.96	6.80	7.58	6.78	7.56	10.35	9.76
New Hampshire	4.84	6.55	7.16	*	*	*	5.95	7.75	7.72
New Jersey	7.15	7.27	7.31	11.09	9.83	12.08	7.57	8.10	8.98
New Mexico	8.13	8.45	8.92	6.88	7.32	8.14	7.67	8.60	10.63
New York	7.38	7.59	7.67	7.81	7.31	7.20	7.33	7.89	8.01
North Carolina	6.13	6.27	6.66	10.30	11.01	10.83	8.20	7.77	7.96
North Dakota	*8.10	*5.84	4.95	6.62	6.78	7.04	*	*8.39	6.75
Ohio	7.20	7.13	7.80	8.86	10.22	9.06	7.86	8.27	8.67
Oklahoma	6.41	6.46	6.64	6.48	6.69	7.21	7.87	6.82	8.07
Oregon	5.54	5.43	6.08	7.23	7.34	7.39	6.78	7.00	7.44
Pennsylvania	8.97	9.00	8.67	9.15	10.95	10.15	7.48	7.99	8.35
Rhode Island	7.20	8.61	7.91	*10.32	13.66	*8.13	9.31	10.11	8.48
South Carolina	6.87	6.66	6.57	10.22	10.75	*7.42	8.02	8.13	8.14
South Dakota	6.89	5.94	7.90	6.84	7.04	7.09	*11.39	*9.50	8.33
Tennessee	6.28	6.04	6.43	*7.11	*6.63	7.95	8.60	7.76	8.18
Texas	6.88	7.23	7.56	6.67	7.33	6.67	7.78	8.33	9.40
Utah	7.20	7.26	7.37	6.37	7.46	8.17	7.23	8.20	8.87
Vermont	*	*	*	*	*	*	*	*8.08	*9.38
Virginia	6.07	6.28	6.56	*10.73	*9.20	9.66	7.50	7.71	8.17
Washington	5.31	5.93	6.18	7.08	7.31	7.16	6.37	6.90	7.48
West Virginia	*	*6.06	8.80	*	*	*	*9.16	*9.51	*5.92
Wisconsin	6.13	6.34	6.66	6.12	6.04	6.22	6.97	7.50	7.52
Wyoming	8.81	8.43	9.13	9.55	8.39	8.92	*12.04	*	*9.88
American Samoa ³	---	---	---	---	---	---	3.46	3.75	4.38
Guam ³	*	*	*	*	---	---	7.78	9.33	8.82
Northern Marianas ³	---	---	*	---	---	*	8.12	7.65	7.21
Puerto Rico ³	---	---	10.98	---	---	---	---	---	*
Virgin Islands ³	10.84	8.29	---	*12.50	---	---	*	*	---

See footnotes at end of table.

Table 6 (page 3 of 3). Low birthweight live births, by race and Hispanic origin of mother, state, and territory: United States and U.S. dependent areas, 2000–2002, 2003–2005, and 2012–2014

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2015.htm#006>.

[Data are based on birth certificates]

* Percentages preceded by an asterisk are based on fewer than 50 births. Percentages not shown are based on fewer than 20 births.

- - - Data not available.

– Quantity zero.

¹Excludes live births with unknown birthweight.

²Excludes data for American Samoa, Guam, Northern Marianas, Puerto Rico, and Virgin Islands.

³Comparable data were not available for all time periods and racial and ethnicity groups. Therefore, only selected low birthweight percentages are presented for the territories.

⁴Persons of Hispanic origin may be of any race. See Appendix II, Hispanic origin.

⁵Includes persons of Hispanic and non-Hispanic origin.

NOTES: For information on very low birthweight live births by state, see Table I–10 in Hamilton BE, Martin JA, Osterman MJK, et al. Births: Final data for 2014. National vital statistics reports; vol 64 no 12. Hyattsville, MD: NCHS. 2015; Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr64/nvsr64_12.pdf. Starting with 2003 data, some states and territories reported multiple-race data. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards, for comparability with other areas. See Appendix II, Race. Data for the territories are shown by race and ethnicity only if race-specific data are available for all years in the 3-year period. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Vital Statistics System, public-use and nonpublic-use Birth File. Hamilton BE, Martin JA, Osterman MJK, et al. Births: Final data for 2014. National vital statistics reports; vol 64 no 12. Hyattsville, MD: NCHS. 2015; Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr64/nvsr64_12.pdf. See Appendix I, National Vital Statistics System (NVSS).

Table 7. Legal abortions, legal abortion rates, and legal abortion ratios: United States and 47 continuous reporting areas, 2003–2012

Updated data when available, Excel, PDF: <http://www.cdc.gov/nchs/hus/contents2015.htm#007>.

[Data are based on reporting by state health departments and by hospitals and other medical facilities]

Data provider	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Number of legal abortions reported, in thousands										
Centers for Disease Control and Prevention (CDC) ¹	848	839	820	852	828	826	789	766	730	699
Guttmacher Institute ²	1,250	1,222	1,206	1,242	1,210	1,212	1,152	1,103	1,059	---
CDC 47 continuous reporting areas ³										
Number of legal abortions reported, in thousands	826	818	808	835	819	817	779	755	720	688
Percent of total legal abortions reported to CDC ⁴	97.4	97.5	98.5	97.9	99.0	98.9	98.7	98.6	98.5	98.4
Number of legal abortions per 1,000 women aged 15–44	16.1	15.9	15.7	16.2	15.8	15.8	15.0	14.6	13.9	13.2
Number of legal abortions per 1,000 live births	245	241	236	237	229	232	227	227	219	210

--- Data not available.

¹Overall trends presented in this table should be interpreted with caution because of the different numbers of reporting areas that provided data to CDC in different years. The following states did not report abortion data to CDC: California (2003–2012), Louisiana (2005), Maryland (2007–2012), New Hampshire (2003–2012), and West Virginia (2003–2004). For 2006, the number of legal abortions is greater than reported in the 2006 report because of numbers subsequently provided by Louisiana. For 2009, the number of legal abortions is greater than reported in the 2009 report because of numbers subsequently provided by Delaware.

²No surveys were conducted in 2003, 2006, or 2009. Data for those years were estimated by interpolation. See Appendix I, Guttmacher Institute Abortion Provider Census.

³Because overall trends in abortion data are affected by the number of reporting areas that provide data to CDC on an annual basis, CDC also presents estimates for the 47 reporting areas that provided data for the entire period from 2003 to 2012. The 47 continuous reporting areas include all states except California, Louisiana, Maryland, New Hampshire, and West Virginia. The District of Columbia and New York City are included in the 47 continuous reporting areas.

⁴Percentage of legal abortions that the 47 continuous reporting areas represented of the total number of legal abortions reported to CDC each year.

NOTES: Each year, CDC requests abortion data from the central health agencies of 52 reporting areas (the 50 states, the District of Columbia, and New York City). This information is provided voluntarily to CDC. See the annual Abortion Surveillance reports for more information on the characteristic-specific list of reporting areas. Available from: http://www.cdc.gov/reproductivehealth/Data_Stats/Abortion.htm. For methodological differences between CDC and the Guttmacher Institute Abortion Provider Census, see Appendix I, Abortion Surveillance System; Guttmacher Institute Abortion Provider Census. Some data were revised and differ from previous editions of *Health, United States*.

SOURCE: CDC, National Center for Chronic Disease Prevention and Health Promotion. CDC. Abortion surveillance—United States, 2012. *MMWR* 2015;64(SS10):1–40. Available from: http://www.cdc.gov/mmwr/preview/mmwrhtml/ss6410a1.htm?s_cid=ss6410a1_e. Guttmacher Institute Abortion Provider Survey. *Perspect Sex Reprod Health* 2014;46(1):3–14. Available from: <http://www.guttmacher.org/pubs/journals/psrh.46e0414.pdf>. See Appendix I, Abortion Surveillance System; Guttmacher Institute Abortion Provider Census.

Table 8 (page 1 of 6). Contraceptive use in the past month among women aged 15–44, by age, race and Hispanic origin, and method of contraception: United States, selected years 1982–2013

Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#008>.

[Data are based on household interviews of samples of women of childbearing age]

Race and Hispanic origin and year ¹	Age, in years				
	15–44	15–19	20–24	25–34	35–44
Number of women in population, in thousands					
All women: ²					
1982	54,099	9,521	10,629	19,644	14,305
1995	60,201	8,961	9,041	20,758	21,440
2002	61,561	9,834	9,840	19,522	22,365
2006–2010	61,755	10,478	10,365	19,722	21,190
2011–2013	60,887	9,547	10,338	20,790	20,212
Not Hispanic or Latina:					
White only:					
1982	41,279	7,010	8,081	14,945	11,243
1995	42,154	5,865	6,020	14,471	15,798
2002	39,498	6,069	5,938	12,073	15,418
2006–2010	37,384	6,034	6,173	11,953	13,224
2011–2013	34,674	4,889	5,606	12,086	12,094
Black or African American only:					
1982	6,825	1,383	1,456	2,392	1,593
1995	8,060	1,334	1,305	2,780	2,641
2002	8,250	1,409	1,396	2,587	2,857
2006–2010	8,451	1,566	1,493	2,621	2,771
2011–2013	8,491	1,432	1,572	2,794	2,693
Hispanic or Latina: ³					
1982	4,393	886	811	1,677	1,018
1995	6,702	1,150	1,163	2,450	1,940
2002	9,107	1,521	1,632	3,249	2,705
2006–2010	10,474	1,904	1,734	3,611	3,225
2011–2013	12,024	2,144	2,105	4,062	3,713
Percent of women in population using contraception					
All women: ²					
1982	55.7	24.2	55.8	66.7	61.6
1995	64.2	29.8	63.5	71.1	72.3
2002	61.9	31.5	60.7	68.6	69.9
2006–2010	62.2	30.5	58.3	67.3	74.9
2011–2013	61.7	33.3	60.4	67.4	70.0
Not Hispanic or Latina:					
White only:					
1982	57.3	23.6	58.7	67.8	63.5
1995	66.2	30.5	65.4	72.9	73.6
2002	64.6	35.0	66.3	69.9	71.4
2006–2010	65.6	35.1	62.7	69.7	77.2
2011–2013	65.3	40.1	60.9	69.2	73.7
Black or African American only:					
1982	51.6	29.8	52.3	63.5	52.0
1995	62.3	36.1	67.6	66.8	68.3
2002	57.6	32.9	50.8	67.9	63.8
2006–2010	54.2	25.5	50.0	60.9	66.2
2011–2013	57.9	21.0	64.9	65.2	65.9
Hispanic or Latina: ³					
1982	50.6	*	*36.8	67.2	59.0
1995	59.0	26.1	50.6	69.2	70.8
2002	59.0	20.4	57.4	66.2	72.9
2006–2010	59.7	22.3	54.0	66.0	77.7
2011–2013	57.3	26.3	52.7	67.8	66.5

See footnotes at end of table.

Table 8 (page 2 of 6). Contraceptive use in the past month among women aged 15–44, by age, race and Hispanic origin, and method of contraception: United States, selected years 1982–2013

Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#008>.

[Data are based on household interviews of samples of women of childbearing age]

Race and Hispanic origin and year ¹	Age, in years				
	15–44	15–19	20–24	25–34	35–44
Number of sexually active women in population, in thousands ⁴					
All women: ²					
1982	---	---	---	---	---
1995	41,796	3,341	6,272	15,687	16,495
2002	42,683	3,775	6,798	14,857	17,252
2006–2010	43,145	3,896	6,944	14,785	17,520
2011–2013	41,762	3,752	7,136	15,464	15,409
Not Hispanic or Latina:					
White only:					
1982	---	---	---	---	---
1995	29,994	2,202	4,276	11,194	12,322
2002	28,079	2,519	4,329	9,224	12,006
2006–2010	27,105	2,471	4,341	9,105	11,188
2011–2013	24,720	2,168	3,840	9,118	9,593
Black or African American only:					
1982	---	---	---	---	---
1995	5,579	598	967	2,039	1,975
2002	5,611	564	949	1,978	2,121
2006–2010	5,526	517	939	1,946	2,124
2011–2013	5,680	464	1,199	2,040	1,977
Hispanic or Latina: ³					
1982	---	---	---	---	---
1995	4,330	409	685	1,794	1,442
2002	6,075	405	1,070	2,462	2,138
2006–2010	6,978	563	1,076	2,656	2,683
2011–2013	7,853	723	1,335	3,076	2,720
Percent of sexually active women in population using contraception ⁴					
All women: ²					
1982	---	---	---	---	---
1995	92.5	80.2	91.7	94.0	93.9
2002	89.3	82.0	87.9	90.2	90.7
2006–2010	89.0	82.0	87.0	89.8	90.6
2011–2013	90.0	84.6	87.5	90.6	91.8
Not Hispanic or Latina:					
White only:					
1982	---	---	---	---	---
1995	93.0	81.7	93.0	93.9	94.2
2002	90.9	84.4	90.9	91.5	91.7
2006–2010	90.5	85.7	89.1	91.6	91.2
2011–2013	91.6	90.4	89.0	91.7	92.9
Black or African American only:					
1982	---	---	---	---	---
1995	90.0	80.0	91.3	91.6	90.9
2002	84.7	82.2	74.8	88.9	86.0
2006–2010	82.8	77.3	79.4	82.1	86.3
2011–2013	86.6	*	85.2	89.3	89.8
Hispanic or Latina: ³					
1982	---	---	---	---	---
1995	91.4	75.5	82.5	95.4	95.2
2002	88.4	76.4	87.5	87.4	92.3
2006–2010	89.6	75.5	87.0	89.7	93.4
2011–2013	87.8	78.0	83.1	89.5	90.8

See footnotes at end of table.

Table 8 (page 3 of 6). Contraceptive use in the past month among women aged 15–44, by age, race and Hispanic origin, and method of contraception: United States, selected years 1982–2013

Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#008>.

[Data are based on household interviews of samples of women of childbearing age]

Method of contraception and year	Age, in years				
	15–44	15–19	20–24	25–34	35–44
Female sterilization					
Percent of contracepting women					
1982	23.2	—	*4.5	22.1	43.5
1995	27.8	*	4.0	23.8	45.0
2002	27.0	—	3.6	21.6	45.8
2006–2010	26.6	*	*2.6	22.9	44.0
2011–2013	25.1	—	*2.3	21.7	44.2
Male sterilization					
1982	10.9	*	*3.6	10.1	19.9
1995	10.9	—	*	7.8	19.5
2002	10.2	—	*	7.2	18.2
2006–2010	10.8	*	*	7.1	19.8
2011–2013	9.0	—	*	4.2	19.6
Implant and other hormonal contraceptives ⁵					
1982
1995	1.3	*	3.7	*1.3	*
2002	1.0	*	*	*1.7	*
2006–2010	3.4	*4.7	6.4	4.4	*1.1
2011–2013	3.9	*	9.0	4.6	*
Injectable ⁶					
1982
1995	3.0	9.7	6.1	2.9	*0.8
2002	5.5	14.2	10.6	5.5	*1.9
2006–2010	3.9	11.4	5.9	4.2	*1.3
2011–2013	4.6	11.4	7.1	4.8	*
Birth control pill ⁷					
1982	28.0	63.9	55.1	25.7	*3.7
1995	27.0	43.8	52.1	33.4	8.7
2002	31.0	53.8	52.5	34.8	15.0
2006–2010	28.4	53.6	47.3	30.5	14.3
2011–2013	26.7	56.5	42.6	25.3	14.4
Intrauterine device					
1982	7.1	*	*4.2	9.7	6.9
1995	0.8	—	*	*0.8	1.1
2002	2.1	*	1.8	3.7	*1.3
2006–2010	5.6	*	5.6	7.2	4.9
2011–2013	10.7	*	10.2	15.1	8.4

See footnotes at end of table.

Table 8 (page 4 of 6). Contraceptive use in the past month among women aged 15–44, by age, race and Hispanic origin, and method of contraception: United States, selected years 1982–2013

Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#008>.

[Data are based on household interviews of samples of women of childbearing age]

Method of contraception and year	Age, in years				
	15–44	15–19	20–24	25–34	35–44
Diaphragm					
Percent of contracepting women					
1982	8.1	*6.0	10.2	10.3	4.0
1995	1.9	*	*	1.7	2.8
2002	*	–	*	*	*
2006–2010	*	–	–	*	*
2011–2013	*	–	–	*	–
Condom					
1982	12.0	20.8	10.7	11.4	11.3
1995	23.4	45.8	33.7	23.7	15.3
2002	23.8	44.6	36.0	23.1	15.6
2006–2010	23.1	34.7	39.6	25.2	12.8
2011–2013	22.8	34.1	38.5	23.8	12.4
Periodic abstinence-calendar rhythm					
1982	3.3	2.0	3.1	3.3	3.7
1995	3.3	*	*1.5	3.7	3.9
2002	2.0	*	*2.3	*1.7	*2.4
2006–2010	1.7	*	*	2.0	2.1
2011–2013	2.9	*	*	*2.7	*2.5
Periodic abstinence-natural family planning					
1982	0.6	–	*	0.9	*
1995	*0.5	–	*	*0.7	*
2002	*0.4	–	–	*	*
2006–2010	*	–	*	*	*
2011–2013	*	*	*	*	*
Withdrawal					
1982	2.0	2.9	3.0	1.8	1.3
1995	6.1	13.2	7.1	6.0	4.5
2002	8.8	15.0	11.9	10.7	4.7
2006–2010	10.1	14.5	15.1	10.2	7.3
2011–2013	12.6	17.9	16.8	15.3	6.9
Other methods ⁸					
1982	4.9	2.6	5.4	4.8	5.3
1995	3.2	*	3.2	3.1	3.4
2002	1.7	*	*0.9	*1.5	*1.8
2006–2010	0.6	*	*	*0.8	*
2011–2013	*	*	*	*	*

See footnotes at end of table.

Table 8 (page 5 of 6). Contraceptive use in the past month among women aged 15–44, by age, race and Hispanic origin, and method of contraception: United States, selected years 1982–2013

Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#008>.

[Data are based on household interviews of samples of women of childbearing age]

Method of contraception and year	Not Hispanic or Latina ¹		
	White only	Black or African American only	Hispanic or Latina ³
Female sterilization			
Percent of contracepting women			
1982	22.0	30.0	23.0
1995	24.5	39.9	36.6
2002	23.9	39.2	33.8
2006–2010	23.6	37.3	31.7
2011–2013	21.4	36.8	32.9
Male sterilization			
1982	13.0	*1.5	*
1995	13.7	*1.8	*4.0
2002	12.9	*	4.7
2006–2010	14.2	*	5.8
2011–2013	12.9	*1.9	4.0
Implant and other hormonal contraceptives ⁵			
1982
1995	*1.0	*2.4	*2.0
2002	*0.6	*	*2.6
2006–2010	3.0	4.7	3.3
2011–2013	*3.3	*	*4.1
Injectable ⁶			
1982
1995	2.4	5.4	4.7
2002	4.3	9.4	7.8
2006–2010	2.5	8.9	6.0
2011–2013	*3.1	10.1	*4.7
Birth control pill ⁷			
1982	26.4	37.9	30.2
1995	28.7	23.7	23.0
2002	34.9	23.1	22.0
2006–2010	33.1	18.7	20.2
2011–2013	30.2	17.9	19.2
Intrauterine device			
1982	5.8	9.3	19.2
1995	0.7	*	*
2002	1.7	*	5.3
2006–2010	5.6	5.0	6.8
2011–2013	11.0	6.5	13.5
Diaphragm			
1982	9.2	*3.2	*
1995	2.3	*	*
2002	*	*	—
2006–2010	*	*	*
2011–2013	—	—	—
Condom			
1982	13.1	6.3	*6.9
1995	22.5	24.9	21.2
2002	21.7	29.6	24.1
2006–2010	20.8	29.9	22.2
2011–2013	21.5	25.9	21.0
Periodic abstinence-calendar rhythm			
1982	3.2	2.9	3.9
1995	3.3	*1.7	3.2
2002	2.3	*	*
2006–2010	1.3	*	*2.7
2011–2013	*2.7	*2.2	*

See footnotes at end of table.

Table 8 (page 6 of 6). Contraceptive use in the past month among women aged 15–44, by age, race and Hispanic origin, and method of contraception: United States, selected years 1982–2013

Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#008>.

[Data are based on household interviews of samples of women of childbearing age]

Method of contraception and year	Not Hispanic or Latina ¹		
	White only	Black or African American only	Hispanic or Latina ³
Periodic abstinence-natural family planning			
Percent of contracepting women			
1982	0.7	0.3	—
1995	0.7	*	*
2002	*	*	*
2006–2010	*	*	*
2011–2013	*	—	*
Withdrawal			
1982	2.1	1.3	2.6
1995	6.4	3.3	5.7
2002	9.5	4.8	6.3
2006–2010	10.3	7.1	10.4
2011–2013	12.8	10.1	13.0
Other methods ⁸			
1982	4.6	7.3	5.0
1995	3.3	3.8	*2.2
2002	*1.7	*1.9	*1.2
2006–2010	0.6	*	*
2011–2013	*	*	*

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30% or based on fewer than 100 sample cases.

--- Data not available.

— Quantity zero.

. . . Data not applicable.

¹Starting with 1995 data, race-specific estimates are tabulated according to 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. Starting with 1995 data, race-specific estimates are for persons who reported only one racial group. Prior to data year 1995, data were tabulated according to the 1977 Standards. Estimates for single-race categories prior to 1995 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. See Appendix II, Race.

²Includes women of other or multiple race not shown separately.

³Persons of Hispanic origin may be of any race. See Appendix II, Hispanic origin.

⁴Had sexual (vaginal) intercourse in the past 3 months.

⁵Data collected starting with the 1995 survey. Includes data about the contraceptive patch, with data collection starting in the 2002 survey, and the contraceptive ring, with data collection starting in the 2006–2010 survey.

⁶Data collected starting with the 1995 survey.

⁷In 2011–2013, includes the oral contraceptive pill only. In previous surveys includes the oral contraceptive pill and emergency contraception/morning-after pill.

⁸In 2011–2013, includes emergency contraception, female condom/vaginal pouch, foam, cervical cap, Today sponge, suppository or insert, jelly or cream (without diaphragm), and other methods. See Appendix II, Contraception, for the list of other methods reported in previous surveys.

NOTES: Survey collects up to four methods of contraception used in the month of interview. Percents may not add to the total because more than one method could have been used in the month of interview. Standard errors for selected years are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Survey of Family Growth. See Appendix I, National Survey of Family Growth (NSFG).

Table 9. Breastfeeding among mothers aged 15–44, by year of baby's birth and selected characteristics of mother: United States, average annual 1986–1988 through 2008–2010

Updated data when available, Excel, PDF, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#009>.

[Data are based on household interviews of samples of women of childbearing age]

Maternal characteristic	1986–1988	1989–1991	1992–1994	1995–1998	1999–2001	2002–2004	2005–2007	2008–2010
Percent of babies breastfed								
Total	54.1	53.3	57.6	64.4	66.5	69.5	68.8	74.2
Age at baby's birth								
Under 20 years	28.4	34.7	41.0	49.5	47.3	60.0	50.7	65.5
20–24 years	48.2	44.3	50.0	55.9	59.3	61.4	64.3	64.2
25–29 years	58.2	56.4	57.4	68.1	63.5	71.1	70.6	75.8
30–44 years	68.6	66.0	70.2	72.8	80.0	77.1	76.2	82.9
Race and Hispanic origin ¹								
Not Hispanic or Latina:								
White only	59.1	58.4	61.7	66.5	68.7	73.8	72.3	77.4
Black or African American only	22.3	22.4	26.1	47.9	45.3	42.3	46.2	56.4
Hispanic or Latina	55.6	57.0	63.8	71.2	76.0	76.6	73.7	70.4
Education ²								
No high school diploma or GED	31.8	36.5	44.6	50.6	46.6	56.3	58.7	64.9
High school diploma or GED	47.4	45.5	51.1	55.9	61.6	61.2	55.4	63.4
Some college, no bachelor's degree	62.2	61.4	64.3	70.1	75.6	68.1	72.7	72.9
Bachelor's degree or higher	78.4	80.6	82.5	82.0	81.3	89.6	88.3	90.8
Geographic region ³								
Northeast	51.3	53.5	56.5	61.6	66.9	73.0	72.4	73.9
Midwest	52.3	49.6	51.7	61.7	61.9	66.0	66.2	69.8
South	44.6	43.6	48.6	58.1	60.9	62.2	62.6	69.3
West	71.4	69.5	77.3	78.1	78.9	83.3	79.0	83.1
Percent of babies breastfed 3 months or more								
Total	34.6	31.8	33.6	45.8	48.4	50.6	46.6	49.9
Age at baby's birth								
Under 20 years	18.5	*10.5	*11.7	30.0	30.0	37.6	26.6	36.2
20–24 years	26.1	24.1	25.1	36.6	41.8	38.0	38.6	36.3
25–29 years	36.9	32.3	35.6	46.3	43.7	50.2	49.0	54.3
30–44 years	50.1	46.8	46.7	57.5	62.4	63.9	56.3	60.1
Race and Hispanic origin ¹								
Not Hispanic or Latina:								
White only	37.7	35.2	36.6	47.8	49.7	54.5	49.5	51.3
Black or African American only	11.6	11.5	13.3	29.6	33.7	29.2	26.3	39.3
Hispanic or Latina	38.2	33.9	35.0	49.7	54.3	55.9	49.4	47.4
Education ²								
No high school diploma or GED	21.8	17.6	25.2	33.9	37.0	39.9	41.3	41.4
High school diploma or GED	28.2	28.0	27.4	36.9	43.1	41.9	36.8	41.8
Some college, no bachelor's degree	38.7	33.1	38.7	49.6	52.8	43.2	48.7	46.7
Bachelor's degree or higher	55.0	56.1	59.3	64.5	64.1	75.9	65.8	65.5
Geographic region ³								
Northeast	29.9	37.2	36.4	48.2	48.8	59.9	51.5	49.1
Midwest	30.3	31.5	30.1	42.0	42.8	46.8	41.6	43.4
South	27.7	20.1	26.2	38.9	44.4	42.7	40.5	41.5
West	52.4	42.9	45.3	58.2	59.2	62.6	57.8	64.7

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error of 20%–30%.

¹Starting with 1995 data, race-specific estimates are tabulated according to 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. Starting with 1995 data, race-specific estimates are for persons who reported only one racial group. Prior to data year 1995, data were tabulated according to the 1977 Standards. Estimates for single-race categories prior to 1995 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. See Appendix II, Race.

²Educational attainment is presented only for women aged 22–44. Education is as of date of interview. GED is General Educational Development high school equivalency diploma. See Appendix II, Education.

³See Appendix II, Geographic region.

NOTES: Data are based on single births to mothers aged 15–44 at interview, including those births that occurred when the mothers were younger than age 15. Data on breastfeeding for babies born in 1986–1994 are based on women interviewed for the 1995 National Survey of Family Growth (NSFG), also known as Cycle 5. Data for babies born in 1995–2001 are based on women interviewed for the 2002 NSFG, also known as Cycle 6. Data for babies born in 2002–2007 are based on women interviewed for the 2006–2010 NSFG, conducted after NSFG's transition from periodic to continuous interviewing. Data for babies born in 2008–2010 are based on women interviewed for the 2011–2013 NSFG. Standard errors are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Survey of Family Growth. See Appendix I, National Survey of Family Growth (NSFG).

Table 10 (page 1 of 2). Infant, neonatal, postneonatal, fetal, and perinatal mortality rates, by detailed race and Hispanic origin of mother: United States, selected years 1983–2013

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#010>.

[Data are based on linked birth and death certificates for infants and fetal death records]

<i>Maternal race and Hispanic origin</i>	1983 ¹	1985 ¹	1990 ¹	1995 ²	2000 ²	2005 ²	2010 ²	2012 ²	2013 ²
Infant ³ deaths per 1,000 live births									
All mothers	10.9	10.4	8.9	7.6	6.9	6.9	6.1	6.0	6.0
White	9.3	8.9	7.3	6.3	5.7	5.7	5.2	5.1	5.1
Black or African American	19.2	18.6	16.9	14.6	13.5	13.3	11.2	10.9	10.8
American Indian or Alaska Native	15.2	13.1	13.1	9.0	8.3	8.1	8.3	8.4	7.6
Asian or Pacific Islander ⁴	8.3	7.8	6.6	5.3	4.9	4.9	4.3	4.1	4.1
Hispanic or Latina ^{5,6}	9.5	8.8	7.5	6.3	5.6	5.6	5.3	5.1	5.0
Mexican	9.1	8.5	7.2	6.0	5.4	5.5	5.1	5.0	4.9
Puerto Rican	12.9	11.2	9.9	8.9	8.2	8.3	7.1	6.9	5.9
Cuban	7.5	8.5	7.2	5.3	4.6	4.4	3.8	5.0	3.0
Central and South American	8.5	8.0	6.8	5.5	4.6	4.7	4.4	4.1	4.3
Other and unknown Hispanic or Latina	10.6	9.5	8.0	7.4	6.9	6.4	6.1	5.6	5.9
Not Hispanic or Latina: ⁶									
White	9.2	8.6	7.2	6.3	5.7	5.8	5.2	5.0	5.1
Black or African American	19.1	18.3	16.9	14.7	13.6	13.6	11.5	11.2	11.1
Neonatal ³ deaths per 1,000 live births									
All mothers	7.1	6.8	5.7	4.9	4.6	4.5	4.0	4.0	4.0
White	6.1	5.8	4.6	4.1	3.8	3.8	3.5	3.4	3.4
Black or African American	12.5	12.3	11.1	9.6	9.1	8.9	7.3	7.3	7.3
American Indian or Alaska Native	7.5	6.1	6.1	4.0	4.4	4.0	4.3	4.9	4.1
Asian or Pacific Islander ⁴	5.2	4.8	3.9	3.4	3.4	3.4	3.0	2.9	3.0
Hispanic or Latina ^{5,6}	6.2	5.7	4.8	4.1	3.8	3.9	3.6	3.6	3.6
Mexican	5.9	5.4	4.5	3.9	3.6	3.8	3.5	3.6	3.5
Puerto Rican	8.7	7.6	6.9	6.1	5.8	5.9	4.8	4.9	4.2
Cuban	*5.0	6.2	5.3	*3.6	*3.2	*3.1	*2.9	3.9	*2.3
Central and South American	5.8	5.6	4.4	3.7	3.3	3.2	3.0	2.9	3.1
Other and unknown Hispanic or Latina	6.4	5.6	5.0	4.8	4.6	4.3	4.0	3.7	4.0
Not Hispanic or Latina: ⁶									
White	5.9	5.6	4.5	4.0	3.8	3.7	3.4	3.3	3.3
Black or African American	12.0	11.9	11.0	9.6	9.2	9.1	7.5	7.5	7.5
Postneonatal ³ deaths per 1,000 live births									
All mothers	3.8	3.6	3.2	2.6	2.3	2.3	2.1	2.0	1.9
White	3.2	3.1	2.7	2.2	1.9	2.0	1.8	1.7	1.6
Black or African American	6.7	6.3	5.9	5.0	4.3	4.3	3.9	3.6	3.5
American Indian or Alaska Native	7.7	7.0	7.0	5.1	3.9	4.0	4.0	3.5	3.5
Asian or Pacific Islander ⁴	3.1	2.9	2.7	1.9	1.4	1.5	1.3	1.2	1.1
Hispanic or Latina ^{5,6}	3.3	3.2	2.7	2.1	1.8	1.8	1.7	1.5	1.5
Mexican	3.2	3.2	2.7	2.1	1.8	1.7	1.6	1.5	1.4
Puerto Rican	4.2	3.5	3.0	2.8	2.4	2.4	2.3	2.0	1.7
Cuban	*2.5	*2.3	*1.9	*1.7	*	*1.4	*	*	*
Central and South American	2.6	2.4	2.4	1.9	1.4	1.5	1.4	1.3	1.2
Other and unknown Hispanic or Latina	4.2	3.9	3.0	2.6	2.3	2.1	2.1	1.9	1.9
Not Hispanic or Latina: ⁶									
White	3.2	3.0	2.7	2.2	1.9	2.1	1.8	1.7	1.7
Black or African American	7.0	6.4	5.9	5.0	4.4	4.5	4.0	3.7	3.7

See footnotes at end of table.

Table 10 (page 2 of 2). Infant, neonatal, postneonatal, fetal, and perinatal mortality rates, by detailed race and Hispanic origin of mother: United States, selected years 1983–2013

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#010>.

[Data are based on linked birth and death certificates for infants and fetal death records]

Maternal race and Hispanic origin	1983	1985	1990	1995	2000 ⁷	2005	2010	2012	2013
Fetal ⁸ deaths per 1,000 live births plus fetal deaths									
All mothers	---	---	---	7.0	6.6	6.2	6.0	6.1	6.0
Hispanic or Latina ⁵	---	---	---	---	5.8	5.4	5.2	5.3	5.2
Not Hispanic or Latina:									
White	---	---	---	---	5.3	4.8	4.8	4.9	4.9
Black or African American	---	---	---	---	12.0	11.1	10.8	10.7	10.5
Late fetal ⁹ deaths per 1,000 live births plus late fetal deaths									
All mothers	---	---	---	3.6	3.3	3.0	3.0	3.0	3.0
Hispanic or Latina ⁵	---	---	---	---	3.1	2.8	2.6	2.6	2.7
Not Hispanic or Latina:									
White	---	---	---	---	2.8	2.4	2.5	2.6	2.6
Black or African American	---	---	---	---	5.2	4.8	4.7	4.7	4.7
Perinatal ¹⁰ deaths per 1,000 live births plus late fetal deaths									
All mothers	---	---	---	7.6	7.0	6.6	6.2	6.2	6.2
Hispanic or Latina ⁵	---	---	---	---	6.1	5.9	5.5	5.6	5.6
Not Hispanic or Latina:									
White	---	---	---	---	5.7	5.4	5.1	5.2	5.3
Black or African American	---	---	---	---	12.6	12.2	10.6	10.8	10.7

* Estimates are considered unreliable. Rates preceded by an asterisk are based on fewer than 50 deaths in the numerator. Rates not shown are based on fewer than 20 deaths in the numerator.

--- Data not available.

¹Rates based on unweighted birth cohort data.

²Rates based on a period file using weighted data. See Appendix I, National Vital Statistics System (NVSS), Linked Birth/Infant Death Data Set.

³Infant (under 1 year of age), neonatal (under 28 days), and postneonatal (28 days–11 months).

⁴Estimates are not available for Asian or Pacific Islander subgroups because not all states have adopted the 2003 revision of the U.S. Standard Certificate of Live Birth. See Appendix II, Race.

⁵Persons of Hispanic origin may be of any race.

⁶Prior to 1995, data are shown only for states with an Hispanic-origin item on their birth certificates. See Appendix II, Hispanic origin.

⁷Rates for 1999–2004 (shown in spreadsheet version) exclude data from Oklahoma, which did not report Hispanic origin on the fetal death report in those years.

⁸Number of fetal deaths of 20 weeks or more gestation per 1,000 live births plus fetal deaths.

⁹Number of fetal deaths of 28 weeks or more gestation (late fetal deaths) per 1,000 live births plus late fetal deaths.

¹⁰Number of late fetal deaths plus infant deaths within 7 days of birth per 1,000 live births plus late fetal deaths.

NOTES: The race groups, white, black, American Indian or Alaska Native, and Asian or Pacific Islander include persons of Hispanic and non-Hispanic origin. Starting with 2003 data, some states reported multiple-race data. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards, for comparability with other states. See Appendix II, Race. National linked files do not exist for 1992–1994. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Vital Statistics System, public-use Linked Birth/Infant Death Data Set, public-use Fetal Death File, public-use Birth File. National Center for Health Statistics. Mathews TJ, MacDorman MF, Thoma ME. Infant mortality statistics from the 2013 period linked birth/infant death data set. National vital statistics reports; vol 64 no 9. Hyattsville, MD: NCHS. 2015 Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr64/nvsr64_09.pdf and MacDorman MF, Gregory ECW. Fetal and perinatal mortality: United States, 2013. National vital statistics reports; vol 64 no 8. Hyattsville, MD: NCHS; 2015. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr64/nvsr64_08.pdf. See Appendix I, National Vital Statistics System (NVSS).

Table 11. Infant mortality rates, by race: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#011>.

[Data are based on death certificates and birth certificates]

Race and year	Infant ¹	Neonatal ¹		Postneonatal ¹
		Under 28 days	Under 7 days	
All races				
Deaths per 1,000 live births				
1950 ²	29.2	20.5	17.8	8.7
1960 ²	26.0	18.7	16.7	7.3
1970	20.0	15.1	13.6	4.9
1980	12.6	8.5	7.1	4.1
1990	9.2	5.8	4.8	3.4
1995	7.6	4.9	4.0	2.7
2000	6.9	4.6	3.7	2.3
2004	6.8	4.5	3.6	2.3
2005	6.9	4.5	3.6	2.3
2010	6.1	4.0	3.2	2.1
2011	6.1	4.1	3.3	2.0
2012	6.0	4.0	3.3	2.0
2013	6.0	4.0	3.3	1.9
2014	5.8	3.9	3.2	1.9
Race of child: ³ White				
1950 ²	26.8	19.4	17.1	7.4
1960 ²	22.9	17.2	15.6	5.7
1970	17.8	13.8	12.5	4.0
1980	11.0	7.5	6.2	3.5
Race of mother: ⁴ White				
1980	10.9	7.4	6.1	3.5
1990	7.6	4.8	3.9	2.8
1995	6.3	4.1	3.3	2.2
2000	5.7	3.8	3.0	1.9
2004	5.7	3.8	3.0	1.9
2005	5.7	3.8	3.0	1.9
2010	5.2	3.5	2.7	1.7
2011	5.1	3.5	2.8	1.7
2012	5.1	3.5	2.8	1.6
2013	5.1	3.5	2.8	1.6
2014	4.9	3.4	2.7	1.6
Race of child: ³ Black or African American				
1950 ²	43.9	27.8	23.0	16.1
1960 ²	44.3	27.8	23.7	16.5
1970	32.6	22.8	20.3	9.9
1980	21.4	14.1	11.9	7.3
Race of mother: ⁴ Black or African American				
1980	22.2	14.6	12.3	7.6
1990	18.0	11.6	9.7	6.4
1995	15.1	9.8	8.2	5.3
2000	14.1	9.4	7.6	4.7
2004	13.8	9.1	7.3	4.7
2005	13.7	9.1	7.3	4.7
2010	11.6	7.5	6.0	4.1
2011	11.5	7.5	6.1	4.0
2012	11.2	7.3	6.0	3.9
2013	11.2	7.4	6.1	3.8
2014	11.0	7.3	6.0	3.7

¹Infant (under 1 year of age), neonatal (under 28 days), early neonatal (under 7 days), and postneonatal (28 days–11 months).

²Includes births and deaths of persons who were not residents of the 50 states and the District of Columbia.

³Infant deaths and live births are tabulated by race of infant. See Appendix II, Race.

⁴Infant deaths are tabulated by race of infant; live births are tabulated by race of mother. See Appendix II, Race.

NOTES: Infant mortality rates in this table are based on infant deaths from the mortality file (numerator) and live births from the natality file (denominator). Inconsistencies in reporting race for the same infant between the birth and death certificate can result in underestimated infant mortality rates for races other than white or black. Infant mortality rates for additional population groups are available from the Linked Birth/Infant Death Data Set and are presented in Table 10. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Vital Statistics System, public-use Mortality File, public-use Birth File; Deaths: Final data for 2014. National vital statistics reports (forthcoming). Available from: <http://www.cdc.gov/nchs/products/nvsr.htm>. See Appendix I, National Vital Statistics System (NVSS).

Table 12 (page 1 of 3). Infant mortality rates, by race and Hispanic origin of mother, state, and territory: United States and U.S. dependent areas, average annual 1989–1991, 2003–2005, and 2011–2013

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2015.htm#012>.

[Data are based on linked birth and death certificates for infants]

State and territory	Not Hispanic or Latina								
	All races			White			Black or African American		
	1989–1991 ¹	2003–2005 ²	2011–2013 ²	1989–1991 ¹	2003–2005 ²	2011–2013 ²	1989–1991 ¹	2003–2005 ²	2011–2013 ²
	Infant ³ deaths per 1,000 live births								
United States ⁴	9.0	6.8	6.0	7.3	5.7	5.1	17.2	13.6	11.3
Alabama	11.4	9.0	8.6	8.6	6.8	6.9	16.8	13.6	12.9
Alaska	9.2	6.5	4.9	7.2	5.3	3.6	*	*	*
Arizona	8.8	6.7	5.7	8.2	6.0	4.7	17.3	11.2	11.0
Arkansas	9.8	8.3	7.4	8.1	7.2	6.7	15.2	13.6	10.9
California	7.6	5.2	4.7	6.9	4.6	3.9	15.4	11.4	9.3
Colorado	8.7	6.3	5.1	8.0	5.2	4.2	16.7	16.3	9.6
Connecticut	7.9	5.5	5.1	5.9	3.9	3.7	17.0	12.7	10.2
Delaware	11.2	9.0	7.6	8.2	6.5	5.6	20.1	16.8	12.8
District of Columbia	20.3	12.2	7.3	*8.2	*3.4	*	23.9	17.2	11.1
Florida	9.4	7.2	6.2	7.2	5.8	5.0	16.2	12.9	10.8
Georgia	11.9	8.4	6.7	8.4	6.1	5.1	17.9	13.3	10.0
Hawaii	7.0	6.7	5.5	5.5	3.9	4.4	*13.6	*15.5	*
Idaho	8.9	6.1	5.4	8.9	6.1	5.0	*	*	*
Illinois	10.7	7.5	6.3	7.6	5.9	4.8	20.5	15.3	12.9
Indiana	9.4	7.9	7.2	8.4	7.1	6.5	17.3	15.1	12.9
Iowa	8.2	5.4	4.8	7.8	5.1	4.6	15.8	*11.0	10.7
Kansas	8.5	7.1	6.3	7.8	6.7	5.5	15.4	14.3	14.2
Kentucky	8.7	6.8	6.7	8.1	6.4	6.4	14.4	10.9	9.8
Louisiana ⁵	10.2	9.8	8.3	7.5	7.1	6.2	14.3	13.9	12.0
Maine	6.6	5.9	6.9	6.2	5.8	6.8	*	*	*
Maryland	9.1	8.0	6.6	6.3	5.2	4.1	15.0	13.7	11.1
Massachusetts	7.0	4.9	4.2	5.9	4.0	3.4	14.2	10.0	6.9
Michigan	10.5	8.0	6.8	7.7	6.2	5.3	20.7	16.4	13.1
Minnesota	7.3	4.8	5.0	6.4	4.3	4.3	18.5	8.9	8.8
Mississippi	11.5	10.7	9.2	7.9	7.0	6.8	15.2	15.6	12.4
Missouri	9.7	7.6	6.5	8.0	6.6	5.4	18.0	13.8	12.2
Montana	9.0	6.3	5.8	8.0	5.7	5.2	*	*	*
Nebraska	8.1	5.9	5.2	7.2	5.1	4.5	18.3	14.0	9.9
Nevada	8.6	5.9	5.3	7.8	5.6	5.1	16.9	12.2	9.5
New Hampshire ⁵	7.1	5.0	4.8	7.2	4.8	4.4	*	*	*
New Jersey	8.4	5.4	4.7	6.1	3.7	3.2	17.8	11.9	10.3
New Mexico	8.4	6.1	5.9	8.1	6.9	5.2	*17.2	*	*
New York	9.5	6.0	5.0	6.3	4.6	4.0	18.4	11.8	8.9
North Carolina	10.7	8.6	7.2	8.0	6.3	5.4	16.9	15.8	12.6
North Dakota	8.0	6.4	6.3	7.3	6.0	5.5	*	*	*
Ohio	9.0	7.8	7.6	7.7	6.4	6.3	16.2	15.6	13.6
Oklahoma ⁵	8.0	7.9	7.2	7.3	7.5	6.5	12.7	13.0	12.5
Oregon	8.0	5.7	5.0	7.4	5.5	4.7	21.3	*8.6	*8.3
Pennsylvania	9.2	7.3	6.7	7.2	5.8	5.2	19.1	13.6	12.7
Rhode Island	8.7	6.2	6.5	7.5	4.5	5.0	*13.6	*10.8	*9.4
South Carolina	11.8	9.0	7.2	8.4	6.4	5.3	17.2	14.2	11.5
South Dakota	9.5	7.2	7.0	7.5	6.2	5.7	*	*	*
Tennessee	10.2	8.9	7.2	7.8	7.0	6.1	18.2	16.3	11.7
Texas	7.9	6.5	5.8	6.9	5.9	5.1	14.1	12.4	10.7
Utah	7.0	4.9	5.2	6.8	4.5	4.8	*	*	*12.9
Vermont	6.6	5.4	4.5	6.3	5.3	4.4	*	*	*
Virginia	9.9	7.5	6.5	7.4	6.0	4.8	18.0	13.7	11.7
Washington	8.0	5.4	4.8	7.4	5.0	4.4	15.1	9.0	8.7
West Virginia	9.1	7.7	7.1	8.8	7.5	7.0	*15.7	*12.0	*12.0
Wisconsin	8.4	6.3	6.1	7.4	5.1	5.0	17.0	16.4	14.0
Wyoming	8.4	6.9	5.7	8.0	6.8	5.7	*	*	*
American Samoa ⁶	---	---	---	---	---	---	---	---	---
Guam ⁶	---	11.1	11.1	---	*	*	---	*	*
Northern Marianas ⁶	---	---	---	---	---	---	---	---	---
Puerto Rico ⁶	---	8.9	8.3	---	---	---	---	---	---
Virgin Islands ⁶	---	7.5	---	---	*	---	---	*	---

See footnotes at end of table.

Table 12 (page 2 of 3). Infant mortality rates, by race and Hispanic origin of mother, state, and territory: United States and U.S. dependent areas, average annual 1989–1991, 2003–2005, and 2011–2013

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2015.htm#012>.

[Data are based on linked birth and death certificates for infants]

State and territory	Hispanic or Latina ⁷			American Indian or Alaska Native ⁸			Asian or Pacific Islander ⁸		
	1989–1991 ¹	2003–2005 ²	2011–2013 ²	1989–1991 ¹	2003–2005 ²	2011–2013 ²	1989–1991 ¹	2003–2005 ²	2011–2013 ²
	Infant ³ deaths per 1,000 live births								
United States ⁴	7.5	5.6	5.1	12.6	8.4	8.1	6.6	4.8	4.2
Alabama	*	7.7	5.0	*	*	*	*	*	*
Alaska	*	*	*	15.7	9.2	8.1	*	*	*
Arizona	8.0	6.7	5.6	11.4	8.3	8.5	*8.5	6.7	5.1
Arkansas	*	6.0	6.2	*	*	*	*	*	*
California	7.0	5.0	4.6	11.0	6.2	5.9	6.4	4.2	3.8
Colorado	8.5	7.0	5.9	*16.5	*	*	*7.8	*5.7	*5.3
Connecticut	7.9	7.4	6.1	*	*	*	*	*	*
Delaware	*	*6.1	*5.1	*	*	*	*	*	*
District of Columbia	*8.8	*7.2	*5.5	*	*	*	*	*	*
Florida	7.1	5.2	4.6	*	*	*	*6.2	5.9	3.7
Georgia	9.0	5.5	4.7	*	*	*	*8.2	5.8	3.9
Hawaii	10.7	7.9	6.0	*	*	*	7.1	7.2	5.7
Idaho	*7.2	6.2	6.7	*	*	*	*	*	*
Illinois	9.2	6.2	5.3	*	*	*	6.0	4.5	4.7
Indiana	*7.2	6.8	6.1	*	*	*	*	*	*5.2
Iowa	*11.9	*5.2	*2.7	*	*	*	*	*	*
Kansas	8.7	6.2	6.8	*	*	*	*	*5.6	*
Kentucky	*	7.6	6.8	*	*	*	*	*	*
Louisiana ⁹	---	*5.7	4.8	*	*	*	*	*	*6.4
Maine	*	*	*	*	*	*	*	*	*
Maryland	7.2	5.8	5.2	*	*	*	7.5	4.3	4.5
Massachusetts	8.3	6.5	5.5	*	*	*	5.7	3.8	3.3
Michigan	7.9	7.6	6.0	*10.7	*	*8.9	*6.1	5.1	4.3
Minnesota	*8.4	4.3	5.4	17.3	*8.6	11.3	*5.1	3.8	4.6
Mississippi	*	*	*6.4	*	*	*	*	*	*
Missouri	*9.1	6.6	6.1	*	*	*	*9.1	*6.1	*4.2
Montana	*	*	*	16.7	*9.3	*9.8	*	*	*
Nebraska	*8.8	5.7	5.7	*18.2	*	*	*	*	*
Nevada	7.0	4.5	4.4	*	*	*	*	*5.8	*4.0
New Hampshire ⁹	---	*	*	*	*	*	*	*	*
New Jersey	7.5	5.2	4.4	*	*	*	5.6	5.0	3.8
New Mexico	7.8	5.3	6.1	9.8	7.6	5.9	*	*	*
New York	9.4	5.5	5.0	*15.2	*	*8.3	6.4	3.9	3.4
North Carolina	*7.5	6.6	5.6	12.2	10.2	10.6	*6.3	5.9	4.3
North Dakota	*	*	*	*13.8	*8.6	*13.2	*	*	*
Ohio	8.0	6.5	6.9	*	*	*	*4.8	*4.5	*4.2
Oklahoma ⁹	---	6.0	6.5	7.8	7.9	7.0	*	*	*7.6
Oregon	8.5	5.5	4.7	*15.7	*11.0	*10.2	*8.4	*5.8	*4.1
Pennsylvania	10.9	7.6	7.0	*	*	*	7.8	4.9	4.2
Rhode Island	*7.2	7.4	7.2	*	*	*	*	*	*
South Carolina	*	7.3	5.0	*	*	*	*	*	*
South Dakota	*	*	*	19.9	12.7	11.5	*	*	*
Tennessee	*	6.5	5.3	*	*	*	*	*8.1	*3.9
Texas	7.0	5.6	5.3	*	*	*	6.8	4.3	3.8
Utah	*7.0	5.8	5.1	*10.0	*	*	*10.7	*7.7	*7.5
Vermont	*	*	*	*	*	*	*	*	*
Virginia	7.6	5.4	5.8	*	*	*	6.0	4.5	5.0
Washington	7.6	4.9	4.2	19.6	9.5	8.7	6.2	4.8	4.4
West Virginia	*	*	*	*	*	*	*	*	*
Wisconsin	*7.3	6.1	5.2	*11.9	*8.2	*8.0	*6.7	*6.6	6.7
Wyoming	*	*	*	*	*	*	*	*	*
American Samoa ⁶	---	---	---	---	---	---	---	---	---
Guam ⁶	---	*	*	---	*	*	---	11.5	11.4
Northern Marianas ⁶	---	---	---	---	---	---	---	---	---
Puerto Rico ⁶	---	---	---	---	---	---	---	---	---
Virgin Islands ⁶	---	*	*	---	*	*	---	*	---

See footnotes at end of table.

Table 12 (page 3 of 3). Infant mortality rates, by race and Hispanic origin of mother, state, and territory: United States and U.S. dependent areas, average annual 1989–1991, 2003–2005, and 2011–2013

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2015.htm#012>.

[Data are based on linked birth and death certificates for infants]

* Estimates are considered unreliable. Rates preceded by an asterisk are based on fewer than 50 deaths in the numerator. Rates not shown are based on fewer than 20 deaths in the numerator.

- - - Data not available.

¹Rates based on unweighted birth cohort data.

²Rates based on period file using weighted data. See Appendix I, National Vital Statistics System (NVSS), Linked Birth/Infant Death Data Set.

³Under 1 year of age.

⁴Excludes data for American Samoa, Guam, Northern Marianas, Puerto Rico, and Virgin Islands.

⁵Rates for white and black are substituted for non-Hispanic white and non-Hispanic black for Louisiana for 1989, Oklahoma for 1989–1990, and New Hampshire for 1989–1991.

⁶Comparable data were not available for all time periods and for all racial and ethnicity groups. Therefore, only selected rates are presented for the territories. Linked birth/infant death data are not available for American Samoa and Northern Marianas.

⁷Persons of Hispanic origin may be of any race. See Appendix II, Hispanic origin.

⁸Includes persons of Hispanic origin.

⁹Rates for Hispanic origin exclude data from states not reporting Hispanic origin on the birth certificate for 1 or more years in a 3-year period.

NOTES: Starting with 2003 data, some states reported multiple-race data. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards, for comparability with other states. See Appendix II, Race. National linked files do not exist for 1992–1994.

SOURCE: CDC/NCHS, National Vital Statistics System, public-use and nonpublic-use Linked Birth/Infant Death Data Set. National Center for Health Statistics. Mathews TJ, MacDorman MF, Thoma ME. Infant mortality statistics from the 2013 period linked birth/infant death data set. National vital statistics reports; vol 64 no 9. Hyattsville, MD: NCHS; 2015. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr64/nvsr64_09.pdf. See Appendix I, National Vital Statistics System (NVSS).

Table 13. Infant mortality rates and international rankings: Organisation for Economic Co-operation and Development (OECD) countries, selected years 1960–2012

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#013>.

[Data are based on reporting by OECD countries]

Country ²	1960	1970	1980	1990	2000	2010	2011	2012	International rankings ¹	
									1960	2012
Infant ³ deaths per 1,000 live births										
Australia	20.2	17.9	10.7	8.2	5.2	4.1	3.8	3.3	5	11
Austria	37.5	25.9	14.3	7.8	4.8	3.9	3.6	3.2	19	10
Belgium	31.4	21.1	12.1	8.0	4.8	3.6	3.4	3.8	17	20
Canada	27.3	18.8	10.4	6.8	5.3	5.0	4.8	---	12	---
Chile	120.3	79.3	33.0	16.0	8.9	7.4	7.7	7.4	27	26
Czech Republic ⁴	20.0	20.2	16.9	10.8	4.1	2.7	2.7	2.6	4	4
Denmark	21.5	14.2	8.4	7.5	5.3	3.4	3.5	3.4	8	13
Finland	21.0	13.2	7.6	5.6	3.8	2.3	2.4	2.4	6	2
France	27.7	18.2	10.0	7.3	4.5	3.6	3.5	3.5	13	15
Germany ⁵	35.0	22.5	12.4	7.0	4.4	3.4	3.6	3.3	18	11
Greece	40.1	29.6	17.9	9.7	5.9	3.8	3.4	2.9	20	6
Hungary	47.6	35.9	23.2	14.8	9.2	5.3	4.9	4.9	23	23
Ireland	29.3	19.5	11.1	8.2	6.2	3.6	3.5	3.5	15	15
Israel ⁶	---	24.2	15.6	9.9	5.5	3.7	3.5	3.6	---	17
Italy	43.9	29.6	14.6	8.1	4.3	3.2	2.9	2.9	22	6
Japan	30.7	13.1	7.5	4.6	3.2	2.3	2.3	2.2	16	1
Korea	---	45.0	---	---	---	3.2	3.0	2.9	---	6
Mexico	92.3	---	52.6	32.5	20.8	14.1	13.7	13.3	26	28
Netherlands	16.5	12.7	8.6	7.1	5.1	3.8	3.6	3.7	2	19
New Zealand	22.6	16.7	13.0	8.4	6.3	5.5	5.2	---	10	---
Norway	16.0	11.3	8.1	6.9	3.8	2.8	2.4	2.5	1	3
Poland	56.1	36.4	25.4	19.4	8.1	5.0	4.7	4.6	24	22
Portugal	77.5	55.5	24.3	10.9	5.5	2.5	3.1	3.4	25	13
Slovak Republic ⁴	28.6	25.7	20.9	12.0	8.6	5.7	4.9	5.8	14	24
Spain	43.7	28.1	12.3	7.6	4.4	3.2	3.1	3.1	21	9
Sweden	16.6	11.0	6.9	6.0	3.4	2.5	2.1	2.6	3	4
Switzerland	21.1	15.1	9.1	6.8	4.9	3.8	3.8	3.6	7	17
Turkey	---	---	---	---	28.9	12.2	11.7	11.6	---	27
United Kingdom	22.5	18.5	12.1	7.9	5.6	4.2	4.2	4.1	9	21
United States	26.0	20.0	12.6	9.2	6.9	6.1	6.1	6.0	11	25

--- Data not available.

¹Break in series. See OECD website for additional information. Available from: <http://www.oecd.org/>.

¹Rankings are from lowest to highest infant mortality rates (IMR). Countries with the same IMR receive the same rank. The country with the next highest IMR is assigned the rank it would have received had the lower-ranked countries not been tied, i.e., skip a rank. The latest year's international rankings are based on 2012 data because that is the most current data year for which most countries have reported their final data to OECD. Countries without an estimate in the OECD database are omitted from this table. Relative rankings for individual countries may be affected if not all countries have reported data to OECD.

²Refers to countries, territories, cities, or geographic areas with at least 2.5 million population in 2000 (United Nations Department of Economic and Social Affairs/Population Division 172 World Population Prospects: The 2012 Revision, Volume 1: Comprehensive Tables. Available from: http://esa.un.org/wpp/Documentation/pdf/WPP2012_Volume-I_Comprehensive-Tables.pdf) and with complete counts of live births and infant deaths according to the United Nations Demographic Yearbook.

³The infant mortality rate is defined as the number of deaths of children under one year of age, expressed per 1,000 live births. Some of the international variation in infant mortality rates is due to variations among countries in registering practices for premature infants. See OECD website for additional information. Available at: <http://www.oecd.org/>.

⁴In 1993, Czechoslovakia was divided into two nations, the Czech Republic and Slovakia. Data for years prior to 1993 are from the Czech and Slovak regions of Czechoslovakia.

⁵Until 1990, estimates refer to the Federal Republic of Germany; from 1995 onward data refer to Germany after reunification.

⁶Statistical data for Israel are supplied by, and under the responsibility of, the relevant Israeli authorities. The use of such data by the OECD is without prejudice to the status of the Golan Heights, East Jerusalem, and Israeli settlements in the West Bank under the terms of international law.

NOTE: Some rates for selected countries and selected years were revised and differ from previous editions of *Health, United States*.

SOURCE: Organisation for Economic Co-operation and Development (OECD) Health Data 2015, incorporating revisions to the annual update. Available from: <http://www.oecd.org/>. See Appendix I, Organisation for Economic Co-operation and Development (OECD) Health Data.

Table 14 (page 1 of 2). Life expectancy at birth and at age 65, by sex: Organisation for Economic Co-operation and Development (OECD) countries, selected years 1980–2013

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#014>.

[Data are based on reporting by OECD countries]

Country	Male					Female				
	1980	1990	2000	2012	2013	1980	1990	2000	2012	2013
At birth	Life expectancy, in years									
Australia	71.0	73.9	76.6	79.9	80.1	78.1	80.1	82.0	84.3	84.3
Austria	69.0	72.3	75.2	78.4	78.6	76.1	79.0	81.2	83.6	83.8
Belgium	69.9	72.7	74.6	77.8	78.1	76.7	79.5	81.0	83.1	83.2
Canada	71.7	74.4	76.3	---	---	78.9	80.8	81.7	---	---
Chile	---	69.4	73.7	76.1	†76.3	---	76.5	80.0	81.3	†81.4
Czech Republic ¹	66.9	67.6	71.6	75.1	75.2	74.0	75.5	78.5	81.2	81.3
Denmark	71.2	72.0	74.5	78.1	78.3	77.3	77.8	79.2	82.1	82.4
Estonia	64.2	64.7	65.6	71.4	72.8	74.3	74.9	76.4	81.5	81.7
Finland	69.2	71.0	74.2	77.7	78.0	78.0	79.0	81.2	83.7	84.1
France	70.2	72.8	75.3	78.7	79.0	78.4	80.9	83.0	85.4	85.6
Germany ²	69.6	72.0	75.1	78.6	78.6	76.2	78.5	81.2	83.3	83.2
Greece	73.0	74.7	75.5	78.0	78.7	77.5	79.5	80.9	83.4	84.0
Hungary	65.5	65.2	67.5	††71.6	72.2	72.8	73.8	76.2	††78.7	79.1
Iceland	73.5	75.5	77.8	81.6	80.5	80.4	80.7	81.6	84.3	83.7
Ireland	70.1	72.1	74.0	78.7	79.0	75.6	77.7	79.2	83.2	83.1
Israel ³	72.1	74.9	76.7	79.9	80.3	75.7	78.4	80.9	83.6	83.9
Italy	70.6	73.8	76.9	79.8	80.3	77.4	80.3	82.8	84.8	85.2
Japan	73.4	75.9	77.7	79.9	80.2	78.8	81.9	84.6	86.4	86.6
Korea	61.8	67.3	72.3	77.9	78.5	70.0	75.5	79.6	84.6	85.1
Luxembourg	70.0	72.4	74.6	††79.1	79.8	75.6	78.7	81.3	††83.8	83.9
Mexico	64.1	67.0	70.5	71.4	71.7	70.2	74.0	76.1	77.3	77.4
Netherlands	72.5	73.8	75.6	79.3	79.5	79.2	80.2	80.7	83.0	83.2
New Zealand	70.1	72.5	75.9	79.3	79.5	76.2	78.4	80.8	83.0	83.2
Norway	72.4	73.4	76.0	79.5	79.8	79.3	79.9	81.5	83.5	83.8
Poland	66.0	66.3	††69.6	72.6	73.0	74.4	75.3	††78.0	81.1	81.2
Portugal	67.9	70.6	73.3	77.3	77.6	74.9	77.5	80.4	83.6	84.0
Slovak Republic ¹	66.7	66.7	69.2	72.5	72.9	74.4	75.7	77.5	79.9	80.1
Slovenia	---	69.8	72.2	77.1	77.2	---	77.8	79.9	83.3	83.6
Spain	72.3	73.4	75.8	79.5	80.2	78.4	80.6	82.9	85.5	86.1
Sweden	72.8	74.8	77.4	79.9	80.2	79.0	80.5	82.0	83.6	83.8
Switzerland	72.3	74.0	77.0	80.6	80.7	79.0	80.9	82.8	84.9	85.0
Turkey	55.8	††65.4	69.0	†72.0	††73.7	60.3	††69.5	73.1	†77.2	††79.4
United Kingdom	70.2	72.9	75.5	79.1	79.2	76.2	78.5	80.3	82.8	82.9
United States	70.0	71.8	74.1	76.4	76.4	77.4	78.8	79.3	81.2	81.2

See footnotes at end of table.

Table 14 (page 2 of 2). Life expectancy at birth and at age 65, by sex: Organisation for Economic Co-operation and Development (OECD) countries, selected years 1980–2013

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#014>.

[Data are based on reporting by OECD countries]

Country	Male					Female				
	1980	1990	2000	2012	2013	1980	1990	2000	2012	2013
At 65 years										
Life expectancy, in years										
Australia	13.7	15.2	16.9	19.1	19.2	17.9	19.0	20.4	22.0	22.1
Austria	12.9	14.4	16.0	18.1	18.2	16.3	18.1	19.6	21.3	21.5
Belgium	12.9	14.3	15.6	17.7	17.8	16.8	18.8	19.7	21.3	21.4
Canada	14.5	15.7	16.5	---	---	18.9	19.9	20.2	---	---
Chile	---	13.7	15.5	16.7	†16.8	---	17.2	19.3	19.8	†19.9
Czech Republic ¹	11.2	11.7	13.7	15.7	15.7	14.4	15.3	17.2	19.2	19.3
Denmark	13.6	14.0	15.2	17.5	17.7	17.7	17.9	18.3	20.2	20.4
Estonia	11.8	12.0	12.7	14.8	15.2	15.6	15.8	17.1	20.3	20.3
Finland	12.6	13.8	15.5	17.8	18.0	17.0	17.8	19.5	21.6	21.8
France	13.6	15.5	16.8	19.1	19.3	18.2	19.8	21.4	23.4	23.6
Germany ²	12.8	14.0	15.8	18.2	18.2	16.3	17.7	19.6	21.2	21.1
Greece	15.2	15.7	16.2	18.1	18.7	17.0	18.0	18.7	20.9	21.6
Hungary	11.6	12.1	13.0	††14.3	14.5	14.7	15.4	16.7	††18.1	18.4
Iceland	15.7	16.4	17.8	20.1	18.8	19.3	19.8	19.8	21.5	21.2
Ireland	12.6	13.3	14.6	18.0	18.1	15.7	17.0	18.0	21.1	20.8
Israel ³	---	15.7	17.0	18.8	19.2	---	17.8	19.0	21.0	21.3
Italy	13.3	15.2	16.7	18.5	18.9	17.1	18.9	20.7	22.1	22.6
Japan	14.6	16.2	17.5	18.9	19.1	17.7	20.0	22.4	23.8	24.0
Korea	10.5	12.4	14.3	17.5	18.0	15.1	16.3	18.2	22.0	22.4
Luxembourg	12.6	14.3	15.5	††18.4	19.1	16.5	18.5	20.1	††21.4	21.9
Mexico	15.4	16.0	16.5	16.7	16.7	17.0	18.0	18.4	18.6	18.6
Netherlands	13.7	14.4	15.4	18.0	18.2	18.0	19.1	19.3	21.0	21.2
New Zealand	13.2	14.6	16.4	18.8	18.9	17.0	18.3	19.8	21.2	21.3
Norway	14.3	14.6	16.1	18.3	18.5	18.2	18.7	19.9	21.0	21.4
Poland	12.0	12.4	††13.5	15.4	15.5	15.5	16.2	††17.5	19.8	19.9
Portugal	13.1	14.0	15.4	17.6	17.8	16.1	17.1	19.1	21.3	21.6
Slovak Republic ¹	12.0	12.3	12.9	14.6	14.7	15.2	16.0	16.7	18.5	18.8
Slovenia	---	13.3	14.2	17.1	17.2	---	17.1	18.7	21.1	21.4
Spain	14.6	15.5	16.7	18.7	19.2	17.8	19.3	20.8	22.8	23.4
Sweden	14.3	15.4	16.7	18.5	18.8	18.1	19.2	20.2	21.1	21.3
Switzerland	14.3	15.3	17.0	19.3	19.4	18.2	19.7	20.9	22.3	22.4
Turkey	11.7	††12.8	13.4	†14.1	††14.9	12.8	††14.3	15.1	†16.1	††18.5
United Kingdom	12.6	14.0	15.8	18.5	18.6	16.6	17.9	19.0	20.9	20.9
United States	14.1	15.1	16.0	17.9	17.9	18.3	18.9	19.0	20.5	20.5

--- Data not available.

¹Data are estimated. See OECD website for updated data and additional information. Available at: <http://www.oecd.org/>.

††Break in series. See OECD website for updated data and additional information. Available at: <http://www.oecd.org/>.

¹In 1993, Czechoslovakia was divided into two nations, the Czech Republic and Slovakia. Data for years prior to 1993 are from the Czech and Slovak regions of Czechoslovakia.

²Until 1990, estimates refer to the Federal Republic of Germany; from 1995 onward data refer to Germany after reunification.

³Statistical data for Israel are supplied by, and under the responsibility of, the relevant Israeli authorities. The use of such data by OECD is without prejudice to the status of the Golan Heights, East Jerusalem, and Israeli settlements in the West Bank under the terms of international law.

NOTES: Differences in life expectancy may reflect differences in reporting and calculation methods, which can vary by country, in addition to actual differences in mortality rates. Therefore, ranks are not presented and comparisons among countries should be made with caution. See Appendix II, Life expectancy. Some estimates for selected countries and selected years were revised and differ from previous editions of *Health, United States*.

SOURCE: Organisation for Economic Co-operation and Development (OECD) Health Data 2015, OECD. StatExtracts, available from: <http://www.oecd.org/>; CDC/NCHS. Vital statistics of the United States (selected years). Public Health Service. Washington, DC. See Appendix I, Organisation for Economic Co-operation and Development (OECD) Health Data.

Table 15 (page 1 of 2). Life expectancy at birth, at age 65, and at age 75, by sex, race, and Hispanic origin: United States, selected years 1900–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#015>.

[Data are based on death certificates]

Specified age and year	All races			White			Black or African American ¹		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
At birth									
Life expectancy, in years									
1900 ^{2,3}	47.3	46.3	48.3	47.6	46.6	48.7	33.0	32.5	33.5
1950 ³	68.2	65.6	71.1	69.1	66.5	72.2	60.8	59.1	62.9
1960 ³	69.7	66.6	73.1	70.6	67.4	74.1	63.6	61.1	66.3
1970	70.8	67.1	74.7	71.7	68.0	75.6	64.1	60.0	68.3
1980	73.7	70.0	77.4	74.4	70.7	78.1	68.1	63.8	72.5
1990	75.4	71.8	78.8	76.1	72.7	79.4	69.1	64.5	73.6
1995	75.8	72.5	78.9	76.5	73.4	79.6	69.6	65.2	73.9
2000	76.8	74.1	79.3	77.3	74.7	79.9	71.8	68.2	75.1
2001	77.0	74.3	79.5	77.5	74.9	80.0	72.0	68.5	75.3
2002	77.0	74.4	79.6	77.5	74.9	80.1	72.2	68.7	75.4
2003	77.2	74.5	79.7	77.7	75.1	80.2	72.4	68.9	75.7
2004	77.6	75.0	80.1	78.1	75.5	80.5	72.9	69.4	76.1
2005	77.6	75.0	80.1	78.0	75.5	80.5	73.0	69.5	76.2
2006	77.8	75.2	80.3	78.3	75.8	80.7	73.4	69.9	76.7
2007	78.1	75.5	80.6	78.5	76.0	80.9	73.8	70.3	77.0
2008	78.2	75.6	80.6	78.5	76.1	80.9	74.3	70.9	77.3
2009	78.5	76.0	80.9	78.8	76.4	81.2	74.7	71.4	77.7
2010	78.7	76.2	81.0	78.9	76.5	81.3	75.1	71.8	78.0
2011	78.7	76.3	81.1	79.0	76.6	81.3	75.3	72.2	78.2
2012	78.8	76.4	81.2	79.1	76.7	81.4	75.5	72.3	78.4
2013	78.8	76.4	81.2	79.1	76.7	81.4	75.5	72.3	78.4
2014	78.8	76.4	81.2	79.0	76.7	81.4	75.6	72.5	78.4
At 65 years									
1950 ³	13.9	12.8	15.0	14.1	12.8	15.1	13.9	12.9	14.9
1960 ³	14.3	12.8	15.8	14.4	12.9	15.9	13.9	12.7	15.1
1970	15.2	13.1	17.0	15.2	13.1	17.1	14.2	12.5	15.7
1980	16.4	14.1	18.3	16.5	14.2	18.4	15.1	13.0	16.8
1990	17.2	15.1	18.9	17.3	15.2	19.1	15.4	13.2	17.2
1995	17.4	15.6	18.9	17.6	15.7	19.1	15.6	13.6	17.1
2000	17.6	16.0	19.0	17.7	16.1	19.1	16.1	14.1	17.5
2001	17.9	16.2	19.2	18.0	16.3	19.3	16.2	14.2	17.7
2002	17.9	16.3	19.2	18.0	16.4	19.3	16.3	14.4	17.8
2003	18.1	16.5	19.3	18.2	16.6	19.4	16.5	14.5	18.0
2004	18.4	16.9	19.6	18.5	17.0	19.7	16.8	14.9	18.3
2005	18.4	16.9	19.6	18.5	17.0	19.7	16.9	15.0	18.3
2006	18.7	17.2	19.9	18.7	17.3	19.9	17.2	15.2	18.6
2007	18.8	17.4	20.0	18.9	17.4	20.1	17.3	15.4	18.8
2008	18.8	17.4	20.0	18.9	17.5	20.0	17.5	15.5	18.9
2009	19.1	17.7	20.3	19.2	17.7	20.3	17.8	15.9	19.2
2010	19.1	17.7	20.3	19.2	17.8	20.3	17.8	15.9	19.3
2011	19.2	17.8	20.3	19.2	17.8	20.4	18.0	16.2	19.4
2012	19.3	17.9	20.5	19.3	18.0	20.4	18.1	16.2	19.5
2013	19.3	17.9	20.5	19.3	18.0	20.5	18.1	16.3	19.5
2014	19.3	18.0	20.5	19.3	18.0	20.5	18.2	16.3	19.6
At 75 years									
1980	10.4	8.8	11.5	10.4	8.8	11.5	9.7	8.3	10.7
1990	10.9	9.4	12.0	11.0	9.4	12.0	10.2	8.6	11.2
1995	11.0	9.7	11.9	11.1	9.7	12.0	10.2	8.8	11.1
2000	11.0	9.8	11.8	11.0	9.8	11.9	10.4	9.0	11.3
2001	11.2	9.9	12.0	11.2	10.0	12.1	10.5	9.0	11.5
2002	11.2	10.0	12.0	11.2	10.0	12.1	10.5	9.1	11.5
2003	11.3	10.1	12.1	11.3	10.2	12.1	10.7	8.7	11.6
2004	11.5	10.4	12.4	11.6	10.4	12.4	10.9	9.4	11.2
2005	11.5	10.4	12.3	11.5	10.4	12.3	10.9	9.4	11.2
2006	11.7	10.6	12.5	11.1	10.6	12.5	11.1	9.1	12.0
2007	11.9	10.7	12.6	11.9	10.8	12.6	11.2	9.8	12.1
2008	11.8	10.7	12.6	11.8	10.7	12.6	11.3	9.8	12.2
2009	12.1	11.0	12.9	12.1	10.4	12.9	11.6	10.2	12.5
2010	12.1	11.0	12.9	12.1	11.0	12.8	11.6	10.2	12.5
2011	12.1	11.1	12.9	12.1	11.0	12.8	11.7	10.4	12.5
2012	12.2	11.2	12.9	12.1	11.1	12.9	11.8	10.4	12.7
2013	12.2	11.2	12.9	12.1	11.1	12.9	11.8	10.4	12.7
2014	12.2	11.2	13.0	12.2	11.2	12.9	11.8	10.5	12.7

See footnotes at end of table.

Table 15 (page 2 of 2). Life expectancy at birth, at age 65, and at age 75, by sex, race, and Hispanic origin: United States, selected years 1900–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#015>.

[Data are based on death certificates]

Specified age and year	White, not Hispanic			Black, not Hispanic			Hispanic ⁴		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
At birth									
Life expectancy, in years									
2006	78.2	75.7	80.6	73.1	69.5	76.4	80.3	77.5	82.9
2007	78.4	75.9	80.8	73.5	69.9	76.7	80.7	77.8	83.2
2008	78.4	76.0	80.7	73.9	70.5	77.0	80.8	78.0	83.3
2009	78.7	76.3	81.1	74.3	70.9	77.4	81.1	78.4	83.5
2010	78.8	76.4	81.1	74.7	71.4	77.7	81.2	78.5	83.8
2011	78.8	76.4	81.1	74.9	71.7	77.8	81.4	78.8	83.7
2012	78.9	76.6	81.2	75.1	71.8	78.1	81.6	79.1	83.9
2013	78.9	76.5	81.2	75.1	71.8	78.1	81.6	79.1	83.8
2014	78.8	76.5	81.1	75.2	72.0	78.1	81.8	79.2	84.0
At 65 years									
2006	18.7	17.2	19.9	17.1	15.1	18.5	20.2	18.5	21.5
2007	18.8	17.4	20.0	17.2	15.3	18.7	20.5	18.7	21.7
2008	18.8	17.4	20.0	17.4	15.4	18.8	20.4	18.7	21.6
2009	19.1	17.7	19.5	17.7	15.8	19.1	20.7	19.0	21.9
2010	19.1	17.7	20.3	17.7	15.8	19.1	20.6	18.8	22.0
2011	19.1	17.8	20.3	17.9	16.1	19.2	20.7	19.1	21.8
2012	19.3	17.9	20.4	18.0	16.1	19.4	21.0	19.5	22.1
2013	19.3	17.9	20.4	18.0	16.1	19.4	20.9	19.3	22.0
2014	19.3	18.0	20.5	18.1	16.2	19.5	21.1	19.6	22.2
At 75 years									
2006	11.7	10.6	12.5	11.1	9.6	12.0	13.0	11.7	13.7
2007	11.8	10.7	12.6	11.2	9.7	12.1	13.1	11.8	13.8
2008	11.8	10.7	12.6	11.3	9.8	12.2	13.0	11.7	13.8
2009	12.0	11.0	12.9	11.6	10.1	12.4	13.3	12.0	13.8
2010	12.0	11.0	12.8	11.6	10.1	12.5	13.2	11.7	14.1
2011	12.0	11.0	12.8	11.7	10.4	12.5	13.2	12.0	13.9
2012	12.1	11.1	12.9	11.7	10.4	12.6	13.5	12.3	14.2
2013	12.1	11.1	12.9	11.7	10.4	12.6	13.4	12.3	14.1
2014	12.1	11.1	12.9	11.8	10.4	12.6	13.6	12.4	14.3

¹Data shown for 1900–1960 are for the nonwhite population.

²Death registration area only. The death registration area increased from 10 states and the District of Columbia (D.C.) in 1900 to the coterminous United States in 1933. See Appendix II, Registration area.

³Includes deaths of persons who were not residents of the 50 states and D.C.

⁴Hispanic origin was added to the U.S. standard death certificate in 1989 and was adopted by every state in 1997. Life expectancies for the Hispanic population are adjusted for underreporting on the death certificate of Hispanic ethnicity, but are not adjusted to account for the potential effects of return migration. To address the effects of age misstatement at the oldest ages, the probability of death for Hispanic persons older than 80 years is estimated as a function of non-Hispanic white mortality with the use of the Brass relational logit model. See Appendix II, Hispanic origin. See Appendix II, Race, for a discussion of sources of bias in death rates by race and Hispanic origin.

NOTES: Populations for computing life expectancy for 1991–1999 are 1990-based postcensal estimates of the U.S. resident population. Starting with *Health, United States, 2012*, populations for computing life expectancy for 2001–2009 were based on intercensal population estimates of the U.S. resident population. Populations for computing life expectancy for 2010 were based on 2010 census counts. Life expectancy for 2011 and beyond was computed using 2010-based postcensal estimates. See Appendix I, Population Census and Population Estimates. In 1997, life table methodology was revised to construct complete life tables by single years of age that extend to age 100. (Anderson RN. Method for constructing complete annual U.S. life tables. NCHS. Vital Health Stat 2(129). 1999.) Previously, abridged life tables were constructed for 5-year age groups ending with 85 years and over. In 2000, the life table methodology was revised. The revised methodology is similar to that developed for the 1999–2001 decennial life tables. In 2008, the life table methodology was further refined. See Appendix II, Life expectancy. Starting with 2003 data, some states allowed the reporting of more than one race on the death certificate. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards, for comparability with other states. The race groups, white and black include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. See Appendix II, Race. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Vital Statistics System, public-use Mortality Files; Grove RD, Hetzel AM. Vital statistics rates in the United States, 1940–1960. Washington, DC: U.S. Government Printing Office, 1968; Arias E. United States life tables by Hispanic origin. Vital health statistics; vol 2 no 152. Hyattsville, MD: NCHS. 2010. Deaths: Final data for 2014. National vital statistics reports (forthcoming). Available from: <http://www.cdc.gov/nchs/products/nvsr.htm>. See Appendix I, National Vital Statistics System (NVSS).

Table 16 (page 1 of 2). Age-adjusted death rates, by race, Hispanic origin, state, and territory: United States and U.S. dependent areas, average annual 1979–1981, 1989–1991, and 2012–2014

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2015.htm#016>.

[Data are based on death certificates]

State and territory	All persons			White	Black or African American	American Indian or Alaska Native ¹	Asian or Pacific Islander ¹	Hispanic or Latino ¹	White, not Hispanic or Latino ¹
	1979–1981	1989–1991	2012–2014	2012–2014	2012–2014	2012–2014	2012–2014	2012–2014	2012–2014
	Age-adjusted death rate per 100,000 population ²								
United States ³	1,022.8	942.2	729.7	729.1	858.1	593.6	399.8	532.2	745.2
Alabama	1,091.2	1,037.9	920.2	905.9	998.2	332.2	253.0	336.2	914.4
Alaska	1,087.4	944.6	730.5	678.5	651.2	1,138.2	486.1	352.1	688.3
Arizona	951.5	873.5	672.6	667.4	769.5	837.6	382.2	598.5	675.2
Arkansas	1,017.0	996.3	891.6	881.8	1,005.4	235.2	508.4	318.3	891.2
California	975.5	911.0	621.8	652.8	808.4	371.0	398.1	517.2	686.6
Colorado	941.1	856.1	661.8	667.0	745.6	468.5	392.8	655.0	662.5
Connecticut	961.5	857.5	647.0	651.0	681.5	214.2	325.0	517.2	652.2
Delaware	1,069.7	1,001.9	735.4	727.8	816.7	*	374.4	405.5	734.2
District of Columbia	1,243.1	1,255.3	751.0	467.3	966.1	*	319.0	385.2	459.0
Florida	960.8	870.9	665.0	659.4	748.7	283.7	324.6	513.6	694.0
Georgia	1,094.3	1,037.4	805.4	793.9	872.7	190.9	394.0	299.0	813.9
Hawaii	801.2	752.2	588.6	649.7	516.8	*	568.5	802.7	640.7
Idaho	936.7	856.6	726.9	730.2	532.0	733.4	457.1	485.4	738.7
Illinois	1,063.7	973.8	726.3	711.2	918.8	153.5	376.9	456.8	727.7
Indiana	1,048.3	962.0	827.3	821.6	952.6	190.6	380.0	476.8	829.5
Iowa	919.9	848.2	721.7	719.5	912.3	542.5	496.5	396.6	723.4
Kansas	940.1	867.2	759.3	752.0	928.1	1,100.1	455.7	534.4	758.9
Kentucky	1,088.9	1,024.5	907.4	911.8	916.7	186.1	428.3	314.1	917.5
Louisiana	1,132.6	1,074.6	896.7	855.3	1,019.9	390.6	421.9	387.8	870.6
Maine	1,002.9	918.7	741.2	744.6	454.3	693.5	327.7	270.1	741.5
Maryland	1,063.3	985.2	706.3	693.3	800.4	241.6	342.1	323.6	707.7
Massachusetts	982.6	884.8	661.5	677.7	595.1	312.1	345.6	447.8	678.0
Michigan	1,050.2	966.0	780.1	759.0	963.9	820.8	345.3	617.8	759.6
Minnesota	892.9	825.2	649.1	642.2	727.9	1,048.3	483.4	394.9	644.6
Mississippi	1,108.7	1,071.4	946.6	908.1	1,037.1	716.8	485.5	260.4	916.6
Missouri	1,033.7	952.4	805.9	794.5	952.9	396.5	417.4	424.7	799.6
Montana	1,013.6	890.2	742.0	718.0	*	1,229.0	*	457.7	715.8
Nebraska	930.6	867.9	717.2	711.1	912.4	868.3	407.9	436.5	717.5
Nevada	1,077.4	1,017.4	764.1	793.2	817.8	531.4	429.7	461.9	848.8
New Hampshire	982.3	891.7	690.9	698.9	383.4	*	320.2	256.2	702.3
New Jersey	1,047.5	956.0	673.2	679.7	793.5	167.6	325.5	463.4	698.3
New Mexico	967.1	891.9	741.9	739.6	707.7	802.8	342.7	714.8	728.9
New York	1,051.8	973.7	645.9	663.7	666.9	178.0	355.5	501.6	662.6
North Carolina	1,050.4	986.0	779.8	760.3	884.2	765.8	360.2	301.2	772.9
North Dakota	922.4	818.4	701.2	678.8	337.4	1,353.8	*	518.0	677.5
Ohio	1,070.6	967.4	813.0	804.3	928.6	262.2	369.0	459.3	807.2
Oklahoma	1,025.6	961.4	900.0	888.6	1,017.9	981.6	464.0	536.6	900.5
Oregon	953.9	893.0	710.3	718.2	785.9	666.7	436.2	454.2	727.1
Pennsylvania	1,076.4	963.4	756.9	746.8	915.7	274.1	396.7	562.9	740.8
Rhode Island	990.8	889.6	699.0	715.0	453.2	507.0	389.6	390.7	722.7
South Carolina	1,104.6	1,030.0	833.9	805.7	942.7	377.1	376.4	356.5	816.0
South Dakota	941.9	846.4	700.3	664.6	266.9	1,274.7	395.2	258.0	668.0
Tennessee	1,045.5	1,011.8	880.6	872.8	966.9	257.1	404.8	330.6	879.5
Texas	1,014.9	947.6	750.0	750.4	891.5	150.3	393.4	631.6	787.3
Utah	924.9	823.2	706.7	709.6	615.2	731.5	549.0	533.0	718.7
Vermont	990.2	908.6	701.6	705.2	*	*	*	344.2	707.1
Virginia	1,054.0	963.1	724.1	714.7	856.6	263.0	374.2	353.6	724.8
Washington	947.7	869.4	677.8	691.2	727.4	852.2	417.5	448.1	699.6
West Virginia	1,100.3	1,031.5	930.7	933.4	983.6	*	281.3	215.8	938.2
Wisconsin	956.4	879.1	713.3	699.7	964.6	1,015.3	494.8	451.0	703.4
Wyoming	1,016.1	897.4	740.6	739.8	331.6	1,158.6	*	551.0	748.1
American Samoa ⁴	---	---	1,029.4	---	---	---	---	---	---
Guam ⁴	---	---	787.5	---	---	---	---	---	---
Northern Marianas ⁴	---	---	811.2	---	---	---	---	---	---
Puerto Rico ⁴	---	---	680.7	---	---	---	---	---	---
Virgin Islands ⁴	---	---	†644.0	---	---	---	---	---	---

See footnotes at end of table.

Table 16 (page 2 of 2). Age-adjusted death rates, by race, Hispanic origin, state, and territory: United States and U.S. dependent areas, average annual 1979–1981, 1989–1991, and 2012–2014

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2015.htm#016>.

[Data are based on death certificates]

-- Data not available.

¹Rate shown is for 2010–2012 because death data were not available for the Virgin Islands for 2013 and 2014.

* Prior to 2009–2011 (shown in spreadsheet file), data for states with populations under 10,000 in the middle year of a 3-year period, or fewer than 50 deaths for the 3-year period, are considered unreliable and are not shown. Starting with 2009–2011 estimates (shown in spreadsheet file), data for states with an average population for the 3-year period of under 10,000, or fewer than 50 deaths for the 3-year period, are considered unreliable and are not shown.

¹Death rates for Hispanic, American Indian or Alaska Native, and Asian or Pacific Islander persons should be interpreted with caution because of inconsistencies in reporting Hispanic origin or race on the death certificate (death rate numerators) compared with population figures (death rate denominators). The net effect of misclassification is an underestimation of deaths and death rates for races other than white and black. See Appendix II, Race, for a detailed discussion of sources of bias in death rates by race and Hispanic origin.

²Age-adjusted average annual death rates are calculated using the year 2000 standard population. Prior to 2001, age-adjusted rates were calculated using standard million proportions based on rounded population numbers. Starting with 2001 data, unrounded population numbers are used to calculate age-adjusted rates.

Age-adjusted rates for Puerto Rico, Virgin Islands, Guam, American Samoa, and Northern Marianas were computed by applying the age-specific death rates to the U.S. standard population combining the age groups for age 75 and over. For the territories, age groups were not available for those age 75 and over by age. See Appendix II, Age adjustment. Prior to 2009–2011 (shown in spreadsheet file), denominators for rates are resident population estimates for the middle year of each 3-year period, multiplied by 3. Starting with 2009–2011 estimates (shown in spreadsheet file), denominators for rates are the 3-year average population. See Appendix I, Population Census and Population Estimates.

³Excludes data for American Samoa, Guam, Northern Marianas, Puerto Rico, and Virgin Islands.

⁴Comparable population data were not available for all time periods and for all racial and ethnicity groups. Therefore, only selected rates are presented for the territories.

NOTES: The race groups, white, black, American Indian or Alaska Native, and Asian or Pacific Islander, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. United States, state, and territory rates for 2011 and beyond were calculated using 2010-based postcensal population estimates. Starting with 2003 data, some states allowed the reporting of more than one race on the death certificate. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards, for comparability with other states. See Appendix II, Race. Rates are rounded at the end of the calculation process. They may differ from rates based on the same data presented elsewhere if rounding is done earlier in the calculation process. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Vital Statistics System; numerator data from annual public-use and nonpublic-use Mortality Files; denominator data from state population estimates prepared by the U.S. Census Bureau 1980 from April 1, 1980 MARS Census File; 1990 from April 1, 1990 MARS Census File; 2011 and beyond from 2010-based postcensal bridged-race files. Available from: http://www.cdc.gov/nchs/nvss/bridged_race.htm. For the territories (except for Puerto Rico) populations are from the U.S. Census Bureau. International data base. 2010. Available from: <http://www.census.gov/population/international/>. For Puerto Rico, populations are from U.S. Census Bureau. Puerto Rico Commonwealth characteristics. Available from: http://www.census.gov/popest/data/puerto_rico/asrh/2014/index.html. See Appendix I, National Vital Statistics System (NVSS).

Table 17 (page 1 of 4). Age-adjusted death rates for selected causes of death, by sex, race, and Hispanic origin: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#017>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and cause of death ¹	1950 ^{2,3}	1960 ^{2,3}	1970 ³	1980 ³	1990 ³	2000 ⁴	2005 ⁴	2010 ⁴	2013 ⁴	2014 ⁴
All persons										
Age-adjusted death rate per 100,000 population ⁵										
All causes	1,446.0	1,339.2	1,222.6	1,039.1	938.7	869.0	815.0	747.0	731.9	724.6
Diseases of heart	588.8	559.0	492.7	412.1	321.8	257.6	216.8	179.1	169.8	167.0
Ischemic heart disease	---	---	---	345.2	249.6	186.8	148.2	113.6	102.6	98.8
Cerebrovascular diseases	180.7	177.9	147.7	96.2	65.3	60.9	48.0	39.1	36.2	36.5
Malignant neoplasms	193.9	193.9	198.6	207.9	216.0	199.6	185.1	172.8	163.2	161.2
Trachea, bronchus, and lung	15.0	24.1	37.1	49.9	59.3	56.1	52.7	47.6	43.4	42.1
Colon, rectum, and anus	---	30.3	28.9	27.4	24.5	20.8	17.7	15.8	14.6	14.3
Chronic lower respiratory diseases	---	---	---	28.3	37.2	44.2	43.9	42.2	42.1	40.5
Influenza and pneumonia	48.1	53.7	41.7	31.4	36.8	23.7	21.0	15.1	15.9	15.1
Chronic liver disease and cirrhosis	11.3	13.3	17.8	15.1	11.1	9.5	8.9	9.4	10.2	10.4
Diabetes mellitus ⁶	23.1	22.5	24.3	18.1	20.7	25.0	24.9	20.8	21.2	20.9
Alzheimer's disease	---	---	---	†	†	18.1	24.0	25.1	23.5	25.4
Human immunodeficiency virus (HIV) disease	---	---	---	---	10.2	5.2	4.2	2.6	2.1	2.0
Unintentional injuries	78.0	62.3	60.1	46.4	36.3	34.9	39.5	38.0	39.4	40.5
Motor vehicle-related injuries	24.6	23.1	27.6	22.3	18.5	15.4	15.2	11.3	10.9	10.8
Poisoning	2.5	1.7	2.8	1.9	2.3	4.5	8.0	10.6	12.2	13.1
Nephritis, nephrotic syndrome and nephrosis ⁶	---	---	---	9.1	9.3	13.5	14.7	15.3	13.2	13.2
Suicide ⁷	13.2	12.5	13.1	12.2	12.5	10.4	10.9	12.1	12.6	13.0
Homicide ⁷	5.1	5.0	8.8	10.4	9.4	5.9	6.1	5.3	5.2	5.1
Male										
All causes	1,674.2	1,609.0	1,542.1	1,348.1	1,202.8	1,053.8	971.9	887.1	863.6	855.1
Diseases of heart	699.0	687.6	634.0	538.9	412.4	320.0	268.2	225.1	214.5	210.9
Ischemic heart disease	---	---	---	459.7	328.2	241.4	192.3	151.3	138.2	133.5
Cerebrovascular diseases	186.4	186.1	157.4	102.2	68.5	62.4	48.4	39.3	36.7	36.9
Malignant neoplasms	208.1	225.1	247.6	271.2	280.4	248.9	227.2	209.9	196.0	192.9
Trachea, bronchus, and lung	24.6	43.6	67.5	85.2	91.1	76.7	69.1	60.3	53.7	51.7
Colon, rectum, and anus	---	31.8	32.3	32.8	30.4	25.1	21.2	19.0	17.4	16.9
Prostate	28.6	28.7	28.8	32.8	38.4	30.4	25.3	21.9	19.2	19.0
Chronic lower respiratory diseases	---	---	---	49.9	55.4	55.8	52.2	48.7	47.5	45.4
Influenza and pneumonia	55.0	65.8	54.0	42.1	47.8	28.9	24.9	18.2	18.6	17.8
Chronic liver disease and cirrhosis	15.0	18.5	24.8	21.3	15.9	13.4	12.4	12.9	13.8	14.1
Diabetes mellitus ⁶	18.8	19.9	23.0	18.1	21.7	27.8	28.8	24.9	25.6	25.6
Alzheimer's disease	---	---	---	†	†	15.2	19.5	21.0	19.3	20.6
Human immunodeficiency virus (HIV) disease	---	---	---	---	18.5	7.9	6.3	3.8	3.1	3.0
Unintentional injuries	101.8	85.5	87.4	69.0	52.9	49.3	55.0	51.5	53.1	54.6
Motor vehicle-related injuries	38.5	35.4	41.5	33.6	26.5	21.7	21.9	16.2	15.9	15.8
Poisoning	3.3	2.3	3.9	2.7	3.5	6.6	10.8	13.8	16.0	17.3
Nephritis, nephrotic syndrome and nephrosis ⁶	---	---	---	12.2	12.1	16.9	18.1	18.7	16.1	16.2
Suicide ⁷	21.2	20.0	19.8	19.9	21.5	17.7	18.1	19.8	20.3	20.7
Homicide ⁷	7.9	7.5	14.3	16.6	14.8	9.0	9.7	8.4	8.2	8.0
Female										
All causes	1,236.0	1,105.3	971.4	817.9	750.9	731.4	692.3	634.9	623.5	616.7
Diseases of heart	486.6	447.0	381.6	320.8	257.0	210.9	177.5	143.3	134.3	131.8
Ischemic heart disease	---	---	---	263.1	193.9	146.5	115.0	84.9	74.9	71.6
Cerebrovascular diseases	175.8	170.7	140.0	91.7	62.6	59.1	47.0	38.3	35.2	35.6
Malignant neoplasms	182.3	168.7	163.2	166.7	175.7	167.6	156.7	146.7	139.5	138.1
Trachea, bronchus, and lung	5.8	7.5	13.1	24.4	37.1	41.3	40.6	38.1	35.5	34.7
Colon, rectum, and anus	---	29.1	26.5	23.8	20.6	17.7	15.0	13.3	12.3	12.1
Breast	31.9	31.7	32.1	31.9	33.3	26.8	24.2	22.1	20.8	20.6
Chronic lower respiratory diseases	---	---	---	14.9	26.6	37.4	38.7	38.0	38.5	37.1
Influenza and pneumonia	41.9	43.8	32.7	25.1	30.5	20.7	18.6	13.1	14.0	13.2
Chronic liver disease and cirrhosis	7.8	8.7	11.9	9.9	7.1	6.2	5.8	6.2	6.8	7.1
Diabetes mellitus ⁶	27.0	24.7	25.1	18.0	19.9	23.0	21.9	17.6	17.6	17.2
Alzheimer's disease	---	---	---	†	†	19.3	26.2	27.3	25.9	28.3
Human immunodeficiency virus (HIV) disease	---	---	---	---	2.2	2.5	2.3	1.4	1.1	1.1
Unintentional injuries	54.0	40.0	35.1	26.1	21.5	22.0	25.3	25.6	26.6	27.3
Motor vehicle-related injuries	11.5	11.7	14.9	11.8	11.0	9.5	8.9	6.5	6.2	6.1
Poisoning	1.7	1.1	1.8	1.3	1.2	2.5	5.1	7.5	8.5	9.1
Nephritis, nephrotic syndrome and nephrosis ⁶	---	---	---	7.3	7.7	11.5	12.6	13.0	11.3	11.1
Suicide ⁷	5.6	5.6	7.4	5.7	4.8	4.0	4.4	5.0	5.5	5.8
Homicide ⁷	2.4	2.6	3.7	4.4	4.0	2.8	2.5	2.3	2.1	2.1

See footnotes at end of table.

Table 17 (page 2 of 4). Age-adjusted death rates for selected causes of death, by sex, race, and Hispanic origin: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#017>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and cause of death ¹	1950 ^{2,3}	1960 ^{2,3}	1970 ³	1980 ³	1990 ³	2000 ⁴	2005 ⁴	2010 ⁴	2013 ⁴	2014 ⁴
White ⁸										
Age-adjusted death rate per 100,000 population ⁵										
All causes	1,410.8	1,311.3	1,193.3	1,012.7	909.8	849.8	801.1	741.8	731.0	725.4
Diseases of heart	586.0	559.0	492.2	409.4	317.0	253.4	213.2	176.9	168.2	165.9
Ischemic heart disease	---	---	---	347.6	249.7	185.6	147.3	113.5	102.9	99.3
Cerebrovascular diseases	175.5	172.7	143.5	93.2	62.8	58.8	46.0	37.7	34.9	35.2
Malignant neoplasms	194.6	193.1	196.7	204.2	211.6	197.2	183.9	172.4	163.7	161.9
Trachea, bronchus, and lung	15.2	24.0	36.7	49.2	58.6	56.2	53.2	48.3	44.1	42.9
Colon, rectum, and anus	---	30.9	29.2	27.4	24.1	20.3	17.1	15.3	14.3	14.0
Chronic lower respiratory diseases	---	---	---	29.3	38.3	46.0	46.0	44.6	44.8	43.1
Influenza and pneumonia	44.8	50.4	39.8	30.9	36.4	23.5	20.9	14.9	15.8	15.1
Chronic liver disease and cirrhosis	11.5	13.2	16.6	13.9	10.5	9.6	9.2	9.9	10.7	11.2
Diabetes mellitus ⁶	22.9	21.7	22.9	16.7	18.8	22.8	22.8	19.0	19.4	19.3
Alzheimer's disease	---	---	---	†	†	18.8	24.7	26.0	24.4	26.4
Human immunodeficiency virus (HIV) disease	---	---	---	---	8.3	2.8	2.2	1.4	1.2	1.1
Unintentional injuries	77.0	60.4	57.8	45.3	35.5	35.1	40.7	40.3	41.9	43.1
Motor vehicle-related injuries	24.4	22.9	27.1	22.6	18.5	15.6	15.7	11.7	11.3	11.1
Poisoning	2.4	1.6	2.4	1.8	2.1	4.5	8.5	11.9	13.7	14.8
Nephritis, nephrotic syndrome and nephrosis ⁶	---	---	---	8.0	8.3	12.1	13.2	14.0	12.1	12.1
Suicide ⁷	13.9	13.1	13.8	13.0	13.4	11.3	12.1	13.6	14.2	14.7
Homicide ⁷	2.6	2.7	4.7	6.7	5.5	3.6	3.7	3.3	3.1	3.0
Black or African American ⁸										
All causes	1,722.1	1,577.5	1,518.1	1,314.8	1,250.3	1,121.4	1,035.1	898.2	860.8	849.3
Diseases of heart	588.7	548.3	512.0	455.3	391.5	324.8	278.0	224.9	210.4	206.3
Ischemic heart disease	---	---	---	334.5	267.0	218.3	175.7	131.2	117.5	112.8
Cerebrovascular diseases	233.6	235.2	197.1	129.1	91.6	81.9	67.0	53.0	49.0	49.7
Malignant neoplasms	176.4	199.1	225.3	256.4	279.5	248.5	223.5	203.8	189.2	185.6
Trachea, bronchus, and lung	11.1	23.7	41.3	59.7	72.4	64.0	58.1	51.4	46.8	44.5
Colon, rectum, and anus	---	22.8	26.1	28.3	30.6	28.2	25.1	21.8	19.4	18.6
Chronic lower respiratory diseases	---	---	---	19.2	28.1	31.6	31.1	29.0	29.5	28.4
Influenza and pneumonia	76.7	81.1	57.2	34.4	39.4	25.6	22.6	16.8	16.7	16.1
Chronic liver disease and cirrhosis	9.0	13.6	28.1	25.0	16.5	9.4	7.6	6.7	7.3	7.2
Diabetes mellitus ⁶	23.5	30.9	38.8	32.7	40.5	49.5	47.5	38.7	38.4	37.3
Alzheimer's disease	---	---	---	†	†	13.0	20.8	20.6	20.1	22.3
Human immunodeficiency virus (HIV) disease	---	---	---	---	26.7	23.3	19.2	11.6	8.9	8.3
Unintentional injuries	79.9	74.0	78.3	57.6	43.8	37.7	38.8	31.3	32.6	33.7
Motor vehicle-related injuries	26.0	24.2	31.1	20.2	18.8	15.7	14.4	10.9	10.9	11.1
Poisoning	2.8	2.9	5.8	3.1	4.1	6.0	8.1	7.3	8.9	9.6
Nephritis, nephrotic syndrome and nephrosis ⁶	---	---	---	20.9	19.8	28.7	30.3	29.3	25.0	24.6
Suicide ⁷	4.5	5.0	6.2	6.5	7.1	5.5	5.2	5.2	5.4	5.5
Homicide ⁷	28.3	26.0	44.0	39.0	36.3	20.5	21.1	17.7	17.8	17.2
American Indian or Alaska Native ⁸										
All causes	---	---	---	867.0	716.3	709.3	701.1	628.3	591.7	594.1
Diseases of heart	---	---	---	240.6	200.6	178.2	156.6	128.6	120.6	119.1
Ischemic heart disease	---	---	---	173.6	139.1	129.1	106.1	84.9	78.2	76.4
Cerebrovascular diseases	---	---	---	57.8	40.7	45.0	38.8	28.1	24.6	25.4
Malignant neoplasms	---	---	---	113.7	121.8	127.8	128.8	122.4	110.2	106.7
Trachea, bronchus, and lung	---	---	---	20.7	30.9	32.3	35.3	33.1	27.7	27.8
Colon, rectum, and anus	---	---	---	9.5	12.0	13.4	12.6	11.7	12.6	10.9
Chronic lower respiratory diseases	---	---	---	14.2	25.4	32.8	31.6	33.8	30.8	29.9
Influenza and pneumonia	---	---	---	44.4	36.1	22.3	23.6	15.9	15.0	15.1
Chronic liver disease and cirrhosis	---	---	---	45.3	24.1	24.3	21.6	22.8	24.8	24.2
Diabetes mellitus ⁶	---	---	---	29.6	34.1	41.5	44.1	36.4	34.1	31.3
Alzheimer's disease	---	---	---	†	†	9.1	15.0	17.2	12.7	15.2
Human immunodeficiency virus (HIV) disease	---	---	---	---	1.8	2.2	2.5	1.6	1.3	1.2
Unintentional injuries	---	---	---	99.0	62.6	51.3	51.3	46.9	47.1	49.5
Motor vehicle-related injuries	---	---	---	54.5	32.5	27.3	22.6	15.7	15.4	16.6
Poisoning	---	---	---	2.3	3.2	4.7	8.6	13.0	14.2	15.5
Nephritis, nephrotic syndrome and nephrosis ⁶	---	---	---	12.2	11.6	15.0	15.6	16.4	11.4	12.4
Suicide ⁷	---	---	---	11.9	11.7	9.8	10.7	10.8	11.7	10.9
Homicide ⁷	---	---	---	15.5	10.4	6.8	6.8	5.7	5.3	5.8

See footnotes at end of table.

Table 17 (page 3 of 4). Age-adjusted death rates for selected causes of death, by sex, race, and Hispanic origin: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#017>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and cause of death ¹	1950 ^{2,3}	1960 ^{2,3}	1970 ³	1980 ³	1990 ³	2000 ⁴	2005 ⁴	2010 ⁴	2013 ⁴	2014 ⁴
Asian or Pacific Islander ⁸										
Age-adjusted death rate per 100,000 population ⁵										
All causes	---	---	---	589.9	582.0	506.4	459.6	424.3	405.4	388.3
Diseases of heart	---	---	---	202.1	181.7	146.0	119.7	100.9	92.8	86.1
Ischemic heart disease	---	---	---	168.2	139.6	109.6	85.6	68.7	59.9	55.1
Cerebrovascular diseases	---	---	---	66.1	56.9	52.9	40.8	33.2	29.4	28.3
Malignant neoplasms	---	---	---	126.1	134.2	121.9	113.2	108.9	100.5	98.9
Trachea, bronchus, and lung	---	---	---	28.4	30.2	28.1	26.3	24.8	23.3	22.7
Colon, rectum, and anus	---	---	---	16.4	14.4	12.7	11.5	11.4	9.8	9.5
Chronic lower respiratory diseases	---	---	---	12.9	19.4	18.6	15.9	13.9	13.6	12.5
Influenza and pneumonia	---	---	---	24.0	31.4	19.7	16.8	14.4	15.0	12.9
Chronic liver disease and cirrhosis	---	---	---	6.1	5.2	3.5	3.6	3.2	3.3	3.5
Diabetes mellitus ⁶	---	---	---	12.6	14.6	16.4	17.3	15.5	15.8	15.0
Alzheimer's disease	---	---	---	†	†	5.5	8.5	10.9	11.1	12.1
Human immunodeficiency virus (HIV) disease	---	---	---	...	2.2	0.6	0.6	0.4	0.4	0.3
Unintentional injuries	---	---	---	27.0	23.9	17.9	18.1	15.0	15.2	15.1
Motor vehicle-related injuries	---	---	---	13.9	14.0	8.6	7.5	5.1	4.8	4.6
Poisoning	---	---	---	0.5	0.7	0.7	1.3	1.4	2.0	2.0
Nephritis, nephrotic syndrome and nephrosis ⁶	---	---	---	7.2	7.1	8.4	8.7	9.6	8.1	8.2
Suicide ⁷	---	---	---	7.8	6.7	5.5	5.1	6.2	5.9	6.0
Homicide ⁷	---	---	---	5.9	5.0	3.0	2.8	1.8	1.5	1.5
Hispanic or Latino ^{8,9}										
All causes	---	---	---	---	692.0	665.7	627.6	558.6	535.4	523.3
Diseases of heart	---	---	---	---	217.1	196.0	170.4	132.8	121.2	116.0
Ischemic heart disease	---	---	---	---	173.3	153.2	127.9	92.3	80.3	75.3
Cerebrovascular diseases	---	---	---	---	45.2	46.4	38.6	32.1	29.6	30.2
Malignant neoplasms	---	---	---	---	136.8	134.9	127.9	119.7	114.5	112.4
Trachea, bronchus, and lung	---	---	---	---	26.5	24.8	23.3	20.4	18.7	18.3
Colon, rectum, and anus	---	---	---	---	14.7	14.1	13.1	12.3	11.7	11.1
Chronic lower respiratory diseases	---	---	---	---	19.3	21.1	20.9	19.6	18.7	17.5
Influenza and pneumonia	---	---	---	---	29.7	20.6	18.5	13.7	13.2	12.8
Chronic liver disease and cirrhosis	---	---	---	---	18.3	16.5	14.1	13.7	14.0	14.5
Diabetes mellitus ⁶	---	---	---	---	28.2	36.9	35.4	27.1	26.3	25.1
Alzheimer's disease	---	---	---	---	†	10.4	15.6	18.5	17.7	19.8
Human immunodeficiency virus (HIV) disease	---	---	---	---	16.3	6.7	4.8	2.8	2.1	2.0
Unintentional injuries	---	---	---	---	34.6	30.1	31.8	25.8	26.9	26.8
Motor vehicle-related injuries	---	---	---	---	19.5	14.7	14.6	9.6	9.7	9.6
Poisoning	---	---	---	---	3.2	4.1	5.2	5.6	6.7	6.8
Nephritis, nephrotic syndrome and nephrosis ⁶	---	---	---	---	8.4	11.8	12.8	14.1	11.1	11.1
Suicide ⁷	---	---	---	---	7.8	5.9	5.6	5.9	5.7	6.3
Homicide ⁷	---	---	---	---	16.2	7.5	7.4	5.3	4.5	4.5
White, not Hispanic or Latino ⁹										
All causes	---	---	---	---	914.5	855.5	810.1	755.0	747.1	742.8
Diseases of heart	---	---	---	---	319.7	255.5	215.5	179.9	171.8	169.9
Ischemic heart disease	---	---	---	---	251.9	186.6	148.3	115.0	104.6	101.2
Cerebrovascular diseases	---	---	---	---	63.5	59.0	46.2	37.8	35.0	35.4
Malignant neoplasms	---	---	---	---	215.4	200.6	187.8	176.5	167.7	166.2
Trachea, bronchus, and lung	---	---	---	---	60.3	58.2	55.5	50.8	46.6	45.4
Colon, rectum, and anus	---	---	---	---	24.6	20.5	17.4	15.5	14.5	14.3
Chronic lower respiratory diseases	---	---	---	---	39.2	47.2	47.7	46.6	47.0	45.4
Influenza and pneumonia	---	---	---	---	36.5	23.5	21.0	14.9	15.9	15.1
Chronic liver disease and cirrhosis	---	---	---	---	9.9	9.0	8.7	9.4	10.3	10.6
Diabetes mellitus ⁶	---	---	---	---	18.3	21.8	21.8	18.2	18.6	18.6
Alzheimer's disease	---	---	---	---	†	19.1	25.1	26.4	24.8	26.8
Human immunodeficiency virus (HIV) disease	---	---	---	---	7.4	2.2	1.8	1.1	0.9	0.9
Unintentional injuries	---	---	---	---	35.0	35.3	41.5	42.4	44.2	45.8
Motor vehicle-related injuries	---	---	---	---	18.2	15.6	15.7	11.9	11.5	11.3
Poisoning	---	---	---	---	2.0	4.6	9.1	13.3	15.3	16.7
Nephritis, nephrotic syndrome and nephrosis ⁶	---	---	---	---	8.1	12.0	13.1	13.8	12.1	12.1
Suicide ⁷	---	---	---	---	13.8	12.0	13.0	15.0	15.9	16.4
Homicide ⁷	---	---	---	---	4.0	2.8	2.7	2.5	2.5	2.5

See footnotes at end of table.

Table 17 (page 4 of 4). Age-adjusted death rates for selected causes of death, by sex, race, and Hispanic origin: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#017>.

[Data are based on death certificates]

- - - Data not available.

¹Data for Alzheimer's disease are only presented for data years 1999 and beyond due to large differences in death rates caused by changes in the coding of the causes of death between ICD-9 and ICD-10. See Appendix II, Cause of death; Comparability ratio; Table IV; Table V.

. . . Category not applicable.

¹Underlying cause of death code numbers are based on the applicable revision of the *International Classification of Diseases (ICD)* for data years shown. See Appendix II, Cause of death; Table III; Table IV.

²Includes deaths of persons who were not residents of the 50 states and the District of Columbia (D.C.).

³Underlying cause of death was coded according to the 6th Revision of the ICD in 1950, 7th Revision in 1960, 8th Revision in 1970, and 9th Revision in 1980–1998. See Appendix II, Cause of death; Table III; Table IV.

⁴Starting with 1999 data, cause of death is coded according to ICD-10. See Appendix II, Cause of death; Comparability ratio; Table IV; Table V.

⁵Age-adjusted rates are calculated using the year 2000 standard population. Prior to 2001, age-adjusted rates were calculated using standard million proportions based on rounded population numbers. Starting with 2001 data, unrounded population numbers are used to calculate age-adjusted rates. See Appendix II, Age adjustment.

⁶Starting with 2011 data, the rules for selecting Renal failure as the underlying cause of death were changed, affecting the number of deaths in the Nephritis, nephrotic syndrome and nephrosis and Diabetes categories. These changes directly affect deaths with mention of Renal failure and other associated conditions, such as Diabetes mellitus with renal complications. The result is a decrease in the number of deaths for Nephritis, nephrotic syndrome and nephrosis and an increase in the number of deaths for Diabetes mellitus. Therefore, trend data for these two causes of death should be interpreted with caution. For more information, see Technical Notes in Deaths: Final data for 2011, available from: http://www.cdc.gov/nchs/data/nvsr/nvsr63/nvsr63_03.pdf.

⁷Figures for 2001 (in Excel spreadsheet on the Web) include September 11-related deaths for which death certificates were filed as of October 24, 2002. See Appendix II, Cause of death; Table IV for terrorism-related ICD-10 codes.

⁸The race groups, white, black, Asian or Pacific Islander, and American Indian or Alaska Native, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Death rates for Hispanic, American Indian or Alaska Native, and Asian or Pacific Islander persons should be interpreted with caution because of inconsistencies in reporting Hispanic origin or race on the death certificate (death rate numerators) compared with population figures (death rate denominators). The net effect of misclassification is an underestimation of deaths and death rates for races other than white and black. See Appendix II, Race, for a detailed discussion of sources of bias in death rates by race and Hispanic origin.

⁹Prior to 1997, data from states that did not report Hispanic origin on the death certificate were excluded. See Appendix II, Hispanic origin.

NOTES: Starting with *Health, United States, 2003*, rates for 1991–1999 were revised using intercensal population estimates based on the 1990 and 2000 censuses. For 2000, population estimates are bridged-race April 1 census counts. Starting with *Health, United States, 2012*, rates for 2001–2009 were revised using intercensal population estimates based on the 2000 and 2010 censuses. For 2010, population estimates are bridged-race April 1 census counts. Rates for 2011 and beyond were computed using 2010-based postcensal estimates. See Appendix I, Population Census and Population Estimates. Starting with 2003 data, some states allowed the reporting of more than one race on the death certificate. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards, for comparability with other states. See Appendix II, Race. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>. **Data have been revised and differ from previous editions of *Health, United States*.**

SOURCE: CDC/NCHS, National Vital Statistics System; Grove RD, Hetzel AM. Vital statistics rates in the United States, 1940–1960. Washington, DC: U.S. Government Printing Office. 1968; numerator data from National Vital Statistics System, annual public-use Mortality Files; denominator data from national population estimates for race groups from Table 1 and unpublished Hispanic population estimates for 1985–1996 prepared by the Housing and Household Economic Statistics Division, U.S. Census Bureau; Deaths: Final data for 2014. National vital statistics reports (forthcoming). Available from: <http://www.cdc.gov/nchs/products/nvsr.htm>. See Appendix I, National Vital Statistics System (NVSS).

Table 18 (page 1 of 4). Years of potential life lost before age 75 for selected causes of death, by sex, race, and Hispanic origin: United States, selected years 1980–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#018>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and cause of death ²	Crude	Age-adjusted ¹					
	2014 ³	1980 ²	1990 ²	2000 ³	2010 ³	2013 ³	2014 ³
All persons							
Years lost before age 75 per 100,000 population under age 75							
All causes	7,054.3	10,448.4	9,085.5	7,578.1	6,642.9	6,593.1	6,622.1
Diseases of heart	1,077.0	2,238.7	1,617.7	1,253.0	972.4	952.3	952.0
Ischemic heart disease	625.1	1,729.3	1,153.6	841.8	577.3	546.1	537.1
Cerebrovascular diseases	180.4	357.5	259.6	223.3	169.3	158.1	160.1
Malignant neoplasms	1,518.6	2,108.8	2,003.8	1,674.1	1,395.8	1,328.6	1,310.4
Trachea, bronchus, and lung	353.9	548.5	561.4	443.1	331.3	298.2	287.7
Colorectal	138.4	190.0	164.7	141.9	125.0	123.5	122.2
Prostate ⁴	59.0	84.9	96.8	63.6	52.2	47.5	47.6
Breast ⁵	274.1	463.2	451.6	332.6	262.4	250.0	245.9
Chronic lower respiratory diseases	211.2	169.1	187.4	188.1	172.4	176.6	174.1
Influenza and pneumonia	100.4	160.2	141.5	87.1	71.4	82.3	93.3
Chronic liver disease and cirrhosis	196.5	300.3	196.9	164.1	163.9	176.9	180.7
Diabetes mellitus ⁶	193.0	134.4	155.9	178.4	158.2	168.3	170.8
Alzheimer's disease	14.7	†	†	10.9	11.7	11.1	11.4
Human immunodeficiency virus (HIV) disease	54.6	...	383.8	174.6	76.6	58.1	55.0
Unintentional injuries	1,053.7	1,543.5	1,162.1	1,026.5	1,025.2	1,051.2	1,080.1
Motor vehicle-related injuries	374.4	912.9	716.4	574.3	400.6	386.6	383.0
Poisoning	449.5	68.0	81.2	163.6	379.7	430.9	465.8
Nephritis, nephrotic syndrome and nephrosis ⁶	75.8	---	50.4	70.7	73.1	65.7	66.7
Suicide ⁷	405.4	392.0	393.1	334.5	385.2	401.6	413.6
Homicide ⁷	216.1	425.5	417.4	266.5	239.0	229.8	224.5
Male							
All causes	8,733.9	13,777.2	11,973.5	9,572.2	8,329.5	8,249.5	8,276.4
Diseases of heart	1,471.7	3,352.1	2,356.0	1,766.0	1,370.8	1,338.2	1,326.9
Ischemic heart disease	904.0	2,715.1	1,766.3	1,255.4	864.8	816.2	797.6
Cerebrovascular diseases	204.2	396.7	286.6	244.6	190.7	182.1	184.9
Malignant neoplasms	1,603.3	2,360.8	2,214.6	1,810.8	1,500.8	1,415.9	1,396.9
Trachea, bronchus, and lung	397.7	821.1	764.8	554.9	390.5	345.1	331.8
Colorectal	160.6	214.9	194.3	167.3	148.0	146.8	144.3
Prostate	59.0	84.9	96.8	63.6	52.2	47.5	47.6
Chronic lower respiratory diseases	215.7	235.1	224.8	206.0	182.8	185.2	182.8
Influenza and pneumonia	110.7	202.5	180.0	102.8	82.6	94.1	103.7
Chronic liver disease and cirrhosis	261.4	415.0	283.9	236.9	226.9	242.1	242.6
Diabetes mellitus ⁶	236.7	140.4	170.4	203.8	194.8	208.6	214.0
Alzheimer's disease	12.6	†	†	10.6	10.7	10.2	10.3
Human immunodeficiency virus (HIV) disease	78.6	...	686.2	258.9	109.5	84.3	79.0
Unintentional injuries	1,483.6	2,342.7	1,715.1	1,475.6	1,432.1	1,463.5	1,509.6
Motor vehicle-related injuries	544.4	1,359.7	1,018.4	796.4	569.2	552.2	550.4
Poisoning	604.4	96.4	123.6	242.1	503.8	573.1	623.8
Nephritis, nephrotic syndrome and nephrosis ⁶	87.2	---	58.9	81.1	82.3	75.3	78.6
Suicide ⁷	626.6	605.6	634.8	539.1	607.0	619.8	635.1
Homicide ⁷	349.9	675.0	658.0	410.5	380.3	365.9	357.3
Female							
All causes	5,387.4	7,350.3	6,333.1	5,644.6	4,994.0	4,967.9	4,999.8
Diseases of heart	685.3	1,246.0	948.5	774.6	593.6	584.5	595.2
Ischemic heart disease	348.2	852.1	600.3	457.6	305.2	290.0	290.2
Cerebrovascular diseases	156.8	324.0	235.9	203.9	149.1	135.4	136.5
Malignant neoplasms	1,434.5	1,896.8	1,826.6	1,555.3	1,301.0	1,250.3	1,232.6
Trachea, bronchus, and lung	310.4	310.4	382.2	342.1	276.9	255.1	247.1
Colorectal	116.4	168.7	138.7	118.7	103.4	101.5	101.4
Breast	274.1	463.2	451.6	332.6	262.4	250.0	245.9
Chronic lower respiratory diseases	206.8	114.0	155.9	172.3	162.8	168.7	166.1
Influenza and pneumonia	90.2	122.0	106.2	72.3	60.7	70.9	83.4
Chronic liver disease and cirrhosis	132.1	194.5	115.1	94.5	103.5	114.4	121.5
Diabetes mellitus ⁶	149.6	128.5	142.3	154.4	123.5	129.8	129.6
Alzheimer's disease	16.8	†	†	11.1	12.6	12.0	12.5
Human immunodeficiency virus (HIV) disease	30.7	...	87.8	92.0	44.4	32.5	31.6
Unintentional injuries	627.1	755.3	607.4	573.2	616.4	635.7	647.5
Motor vehicle-related injuries	205.7	470.4	411.6	348.5	230.5	219.0	213.6
Poisoning	295.7	40.2	39.1	85.0	255.1	287.7	306.6
Nephritis, nephrotic syndrome and nephrosis ⁶	64.6	---	42.4	60.8	64.6	56.6	55.3
Suicide ⁷	185.8	184.2	153.3	129.1	163.7	182.5	191.2
Homicide ⁷	83.3	181.3	174.3	118.9	94.9	90.4	88.7

See footnotes at end of table.

Table 18 (page 2 of 4). Years of potential life lost before age 75 for selected causes of death, by sex, race, and Hispanic origin: United States, selected years 1980–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#018>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and cause of death ²	Crude		Age-adjusted ¹				
	2014 ³	1980 ²	1990 ²	2000 ³	2010 ³	2013 ³	2014 ³
White ⁸							
Years lost before age 75 per 100,000 population under age 75							
All causes	6,939.8	9,554.1	8,159.5	6,949.5	6,342.8	6,338.2	6,390.1
Diseases of heart	1,040.7	2,100.8	1,490.3	1,149.4	900.9	881.8	883.0
Ischemic heart disease	637.0	1,682.7	1,113.4	805.3	563.7	532.0	524.8
Cerebrovascular diseases	161.6	300.7	213.1	187.1	142.7	135.1	138.1
Malignant neoplasms	1,565.8	2,035.9	1,929.3	1,627.8	1,375.8	1,317.7	1,301.5
Trachea, bronchus, and lung	376.1	529.9	544.2	436.3	332.8	300.4	291.0
Colorectal	137.7	186.8	157.8	134.1	118.4	118.6	117.6
Prostate ⁴	55.1	74.8	86.6	54.3	45.3	41.9	41.8
Breast ⁵	265.5	460.2	441.7	315.6	245.0	236.0	230.3
Chronic lower respiratory diseases	230.1	165.4	182.3	185.3	176.1	180.1	178.7
Influenza and pneumonia	99.3	130.8	116.9	77.7	66.7	78.4	90.1
Chronic liver disease and cirrhosis	217.3	257.3	175.8	162.7	173.5	189.3	195.1
Diabetes mellitus ⁶	179.1	115.7	133.7	155.6	139.0	149.2	152.6
Alzheimer's disease	16.6	†	†	11.4	12.4	11.8	12.0
Human immunodeficiency virus (HIV) disease	30.2	...	309.0	94.7	39.9	31.5	30.2
Unintentional injuries	1,116.0	1,520.4	1,139.7	1,031.8	1,098.6	1,125.2	1,156.4
Motor vehicle-related injuries	381.3	939.9	726.7	586.1	419.0	401.2	395.9
Poisoning	510.0	64.9	74.4	167.2	435.4	493.0	534.7
Nephritis, nephrotic syndrome and nephrosis ⁶	63.6	---	37.0	52.5	57.4	52.7	53.6
Suicide ⁷	455.7	414.5	417.7	362.0	430.8	451.2	468.3
Homicide ⁷	118.0	271.7	234.9	156.6	138.7	127.7	125.7
Black or African American ⁸							
All causes	9,497.2	17,873.4	16,593.0	12,897.1	9,832.5	9,528.5	9,490.6
Diseases of heart	1,601.4	3,619.9	2,891.8	2,275.2	1,691.1	1,647.0	1,638.9
Ischemic heart disease	744.6	2,305.1	1,676.1	1,300.1	818.8	776.7	756.5
Cerebrovascular diseases	314.9	883.2	656.4	507.0	358.1	319.2	322.3
Malignant neoplasms	1,645.0	2,946.1	2,894.8	2,294.7	1,796.7	1,666.7	1,651.6
Trachea, bronchus, and lung	346.5	776.0	811.3	593.0	405.6	362.4	340.8
Colorectal	173.2	232.3	241.8	222.4	188.6	174.2	176.5
Prostate ⁴	102.1	200.3	223.5	171.0	127.3	109.9	109.0
Breast ⁵	382.9	524.2	592.9	500.0	420.8	389.8	386.7
Chronic lower respiratory diseases	193.9	203.7	240.6	232.7	187.7	200.1	193.6
Influenza and pneumonia	129.4	384.9	330.8	161.2	109.8	122.9	130.9
Chronic liver disease and cirrhosis	119.4	644.0	371.8	185.6	120.2	122.4	120.4
Diabetes mellitus ⁶	313.8	305.3	361.5	383.4	316.4	323.8	321.4
Alzheimer's disease	10.6	†	†	8.3	10.0	9.8	11.2
Human immunodeficiency virus (HIV) disease	210.5	...	1,014.7	763.3	329.5	238.4	222.9
Unintentional injuries	987.9	1,751.5	1,392.7	1,152.8	896.7	953.0	986.8
Motor vehicle-related injuries	411.0	750.2	699.5	580.8	393.4	402.6	404.6
Poisoning	276.7	99.4	144.3	196.6	218.9	265.6	289.7
Nephritis, nephrotic syndrome and nephrosis ⁶	163.6	---	160.9	216.9	193.2	165.7	167.1
Suicide ⁷	210.6	238.0	261.4	208.7	196.4	207.0	209.0
Homicide ⁷	817.0	1,580.8	1,612.9	941.6	821.2	813.5	785.1
American Indian or Alaska Native ⁸							
All causes	6,636.3	13,390.9	9,506.2	7,758.2	6,771.3	6,698.8	6,954.0
Diseases of heart	730.3	1,819.9	1,391.0	1,030.1	820.6	807.4	822.2
Ischemic heart disease	433.9	1,208.2	901.8	709.3	487.6	484.4	492.0
Cerebrovascular diseases	112.0	269.3	223.3	198.1	129.7	124.1	123.2
Malignant neoplasms	721.4	1,101.3	1,141.1	995.7	929.5	852.4	809.9
Trachea, bronchus, and lung	144.9	181.1	268.1	227.8	211.0	166.0	168.4
Colorectal	81.2	78.8	82.4	93.8	95.8	103.3	90.2
Prostate ⁴	32.4	66.7	42.0	44.5	36.8	32.6	39.0
Breast ⁵	99.3	205.5	213.4	174.1	145.0	108.5	110.9
Chronic lower respiratory diseases	125.5	89.3	129.0	151.8	154.5	135.6	140.5
Influenza and pneumonia	118.2	307.9	206.3	124.0	99.3	109.0	129.0
Chronic liver disease and cirrhosis	489.2	1,190.3	535.1	519.4	510.8	562.2	549.9
Diabetes mellitus ⁶	247.7	305.5	292.3	305.6	267.6	281.7	279.4
Alzheimer's disease	5.4	†	†	*	8.8	5.7	6.7
Human immunodeficiency virus (HIV) disease	26.5	...	70.1	68.4	46.1	33.8	29.5
Unintentional injuries	1,504.7	3,541.0	2,183.9	1,700.1	1,377.7	1,388.2	1,509.3
Motor vehicle-related injuries	625.6	2,102.4	1,301.5	1,032.2	570.6	575.7	605.4
Poisoning	511.7	92.9	119.5	180.1	449.6	487.3	537.8
Nephritis, nephrotic syndrome and nephrosis ⁶	71.1	---	88.5	102.0	81.7	73.7	80.5
Suicide ⁷	448.2	515.0	495.9	403.1	437.9	463.0	437.1
Homicide ⁷	248.1	628.9	434.2	278.5	256.4	227.8	239.0

See footnotes at end of table.

Table 18 (page 3 of 4). Years of potential life lost before age 75 for selected causes of death, by sex, race, and Hispanic origin: United States, selected years 1980–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#018>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and cause of death ²	Crude	Age-adjusted ¹					
	2014 ³	1980 ²	1990 ²	2000 ³	2010 ³	2013 ³	2014 ³
Asian or Pacific Islander ⁸							
Years lost before age 75 per 100,000 population under age 75							
All causes	2,987.8	5,378.4	4,705.2	3,811.1	3,061.2	3,050.9	2,954.4
Diseases of heart	412.0	952.8	702.2	567.9	400.1	413.1	402.1
Ischemic heart disease	246.8	697.7	486.6	381.1	250.6	250.4	239.8
Cerebrovascular diseases	124.9	266.9	233.5	199.4	148.3	134.3	121.8
Malignant neoplasms	825.3	1,218.6	1,166.4	1,033.8	874.7	830.7	799.4
Trachea, bronchus, and lung	141.7	238.2	204.7	185.8	148.2	136.7	136.5
Colorectal	81.9	115.9	105.1	91.6	87.6	87.0	79.1
Prostate ⁴	16.5	17.0	32.4	18.8	17.0	14.4	16.8
Breast ⁵	169.7	222.2	216.5	200.8	156.9	146.5	160.6
Chronic lower respiratory diseases	33.3	56.4	72.8	56.5	33.2	35.0	32.7
Influenza and pneumonia	43.4	79.3	74.0	48.6	38.4	36.4	42.8
Chronic liver disease and cirrhosis	43.0	85.6	72.4	44.8	41.7	42.9	41.6
Diabetes mellitus ⁶	76.9	83.1	74.0	77.0	69.5	74.0	73.8
Alzheimer's disease	2.6	†	†	3.5	3.2	3.6	2.6
Human immunodeficiency virus (HIV) disease	10.1	...	77.0	19.9	10.7	11.2	9.8
Unintentional injuries	316.2	742.7	636.6	425.7	303.0	308.5	307.0
Motor vehicle-related injuries	145.4	472.6	445.5	263.4	147.9	136.5	141.2
Poisoning	71.3	*	17.6	25.9	46.5	71.5	67.6
Nephritis, nephrotic syndrome and nephrosis ⁶	29.9	---	26.7	33.6	38.1	31.1	29.1
Suicide ⁷	210.4	217.1	200.6	168.6	199.7	206.0	203.1
Homicide ⁷	60.8	201.1	205.8	113.1	68.8	64.4	59.7
Hispanic or Latino ^{8,9}							
All causes	4,341.0	---	7,963.3	6,037.6	4,795.1	4,668.1	4,676.8
Diseases of heart	457.0	---	1,082.0	821.3	598.1	571.0	567.9
Ischemic heart disease	257.9	---	756.6	564.6	366.6	340.0	334.1
Cerebrovascular diseases	114.7	---	238.0	207.8	150.4	145.5	140.2
Malignant neoplasms	745.3	---	1,232.2	1,098.2	951.2	927.9	909.3
Trachea, bronchus, and lung	72.2	---	193.7	152.1	115.0	102.9	96.8
Colorectal	69.4	---	100.2	101.4	94.0	92.8	87.4
Prostate ⁴	23.6	---	47.7	42.9	38.2	35.9	36.5
Breast ⁵	152.6	---	299.3	230.7	180.0	181.2	183.4
Chronic lower respiratory diseases	42.1	---	78.8	68.5	59.6	56.2	52.7
Influenza and pneumonia	69.3	---	130.1	76.0	57.5	64.3	79.1
Chronic liver disease and cirrhosis	176.8	---	329.1	252.1	201.6	206.1	216.4
Diabetes mellitus ⁶	126.6	---	177.8	215.6	158.5	161.6	160.9
Alzheimer's disease	5.7	---	†	6.9	8.4	8.5	8.9
Human immunodeficiency virus (HIV) disease	43.5	---	600.1	209.4	74.9	52.5	49.5
Unintentional injuries	757.6	---	1,190.6	920.1	708.7	735.9	740.1
Motor vehicle-related injuries	371.0	---	740.8	540.2	340.3	344.8	348.3
Poisoning	227.7	---	121.9	145.9	191.2	224.3	234.1
Nephritis, nephrotic syndrome and nephrosis ⁶	44.8	---	54.4	62.0	67.7	57.5	56.3
Suicide ⁷	219.5	---	256.2	188.5	193.6	195.6	214.3
Homicide ⁷	213.0	---	720.8	335.1	238.0	199.3	197.6
White, not Hispanic or Latino ⁹							
All causes	7,498.1	---	8,022.5	6,960.5	6,545.3	6,593.2	6,659.4
Diseases of heart	1,176.6	---	1,504.0	1,175.1	943.2	930.5	932.4
Ischemic heart disease	725.5	---	1,127.2	824.7	590.8	560.9	553.6
Cerebrovascular diseases	170.7	---	210.1	183.0	139.1	130.3	134.9
Malignant neoplasms	1,757.6	---	1,974.1	1,668.4	1,421.5	1,362.2	1,349.1
Trachea, bronchus, and lung	452.4	---	566.8	460.3	359.1	327.2	318.6
Colorectal	153.4	---	162.1	136.2	121.2	122.1	122.3
Prostate ⁴	62.7	---	89.2	54.9	45.9	42.3	42.2
Breast ⁵	290.4	---	451.5	322.3	252.6	242.3	235.4
Chronic lower respiratory diseases	277.4	---	188.1	193.8	189.1	195.9	195.9
Influenza and pneumonia	105.1	---	112.3	76.4	67.8	80.5	91.1
Chronic liver disease and cirrhosis	222.5	---	162.4	150.9	166.9	183.9	188.6
Diabetes mellitus ⁶	188.8	---	131.2	150.2	136.7	147.0	150.6
Alzheimer's disease	19.3	---	†	11.7	12.7	12.1	12.3
Human immunodeficiency virus (HIV) disease	25.5	---	271.2	76.0	31.3	25.7	24.7
Unintentional injuries	1,189.3	---	1,114.7	1,041.4	1,183.0	1,214.6	1,253.8
Motor vehicle-related injuries	374.6	---	715.7	588.8	430.6	407.4	398.9
Poisoning	576.2	---	68.3	169.4	494.0	561.8	613.2
Nephritis, nephrotic syndrome and nephrosis ⁶	67.2	---	34.5	51.1	55.3	51.2	52.7
Suicide ⁷	511.4	---	433.0	389.2	483.8	512.9	531.3
Homicide ⁷	89.3	---	162.0	113.2	103.4	100.6	98.4

See footnotes at end of table.

Table 18 (page 4 of 4). Years of potential life lost before age 75 for selected causes of death, by sex, race, and Hispanic origin: United States, selected years 1980–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#018>.

[Data are based on death certificates]

. . . Category not applicable.

- - - Data not available.

† Data for Alzheimer's disease are only presented for data years 1999 and beyond due to large differences in death rates caused by changes in the coding of this cause of death between ICD-9 and ICD-10. See Appendix II, Cause of death; Comparability ratio; Table IV; Table V.

* Rates based on fewer than 20 deaths are considered unreliable and are not shown.

¹ Age-adjusted rates are calculated using the year 2000 standard population. Prior to 2001, age-adjusted rates were calculated using standard million proportions based on rounded population numbers. Starting with 2001 data, unrounded population numbers are used to calculate age-adjusted rates. See Appendix II, Age adjustment.

² Underlying cause of death was coded according to the 9th Revision of the *International Classification of Diseases* (ICD) in 1980–1998. See Appendix II, Cause of death; Table III; Table IV.

³ Starting with 1999 data, cause of death is coded according to ICD-10. See Appendix II, Cause of death; Comparability ratio; Table IV; Table V.

⁴ Rate for male population only.

⁵ Rate for female population only.

⁶ Starting with 2011 data, the rules for selecting Renal failure as the underlying cause of death were changed, affecting the number of deaths in the Nephritis, nephrotic syndrome and nephrosis and Diabetes categories. These changes directly affect deaths with mention of Renal failure and other associated conditions, such as Diabetes mellitus with renal complications. The result is a decrease in the number of deaths for Nephritis, nephrotic syndrome and nephrosis and an increase in the number of deaths for Diabetes mellitus. Therefore, trend data for these two causes of death should be interpreted with caution. For more information, see Technical Notes in Deaths: Preliminary data for 2011, available from: http://www.cdc.gov/nchs/data/nvsr/nvsr61/nvsr61_06.pdf.

⁷ Figures for 2001 (in Excel spreadsheet on the Web) include September 11-related deaths for which death certificates were filed as of October 24, 2002. See Appendix II, Cause of death; Table IV for terrorism-related ICD-10 codes.

⁸ The race groups, white, black, Asian or Pacific Islander, and American Indian or Alaska Native, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Death rates for Hispanic, American Indian or Alaska Native, and Asian or Pacific Islander persons should be interpreted with caution because of inconsistencies in reporting Hispanic origin or race on the death certificate (death rate numerators) compared with population figures (death rate denominators). The net effect of misclassification is an underestimation of deaths and death rates for races other than white and black. See Appendix II, Race, for a detailed discussion of sources of bias in death rates by race and Hispanic origin.

⁹ Prior to 1997, data from states that did not report Hispanic origin on the death certificate were excluded. See Appendix II, Hispanic origin.

NOTES: Starting with *Health, United States, 2003*, rates for 1991–1999 were revised using intercensal population estimates based on the 1990 and 2000 censuses. For 2000, population estimates are bridged-race April 1 census counts. Starting with *Health, United States, 2012*, rates for 2001–2009 were revised using intercensal population estimates based on the 2000 and 2010 censuses. For 2010, population estimates are bridged-race April 1 census counts. Rates for 2011 and beyond were computed using 2010-based postcensal estimates. See Appendix I, Population Census and Population Estimates. See Appendix II, Years of potential life lost (YPLL) for definition and method of calculation. Starting with 2003 data, some states allowed the reporting of more than one race on the death certificate. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards, for comparability with other states. See Appendix II, Race. Rates are rounded at the end of the calculation process. They may differ from rates based on the same data presented elsewhere if rounding is done earlier in the calculation process. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

Data have been revised and differ from previous editions of *Health, United States*.

SOURCE: CDC/NCHS, National vital statistics system; numerator data from annual public-use Mortality Files; denominator data from national population estimates for race groups from Table 1 and unpublished Hispanic population estimates for 1990–1996 prepared by the Housing and Household Economic Statistics Division, U.S. Census Bureau. See Appendix I, National Vital Statistics System (NVSS).

Table 19 (page 1 of 4). Leading causes of death and numbers of deaths, by sex, race, and Hispanic origin: United States, 1980 and 2014

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2015.htm#019>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and rank order	1980		2014	
	Cause of death	Deaths	Cause of death	Deaths
All persons				
Rank	All causes	1,989,841	All causes	2,626,418
1	Diseases of heart	761,085	Diseases of heart	614,348
2	Malignant neoplasms	416,509	Malignant neoplasms	591,700
3	Cerebrovascular diseases	170,225	Chronic lower respiratory diseases	147,101
4	Unintentional injuries	105,718	Unintentional injuries	135,928
5	Chronic obstructive pulmonary diseases	56,050	Cerebrovascular diseases	133,103
6	Pneumonia and influenza	54,619	Alzheimer's disease	93,541
7	Diabetes mellitus	34,851	Diabetes mellitus ¹	76,488
8	Chronic liver disease and cirrhosis	30,583	Influenza and pneumonia	55,227
9	Atherosclerosis	29,449	Nephritis, nephrotic syndrome and nephrosis ¹	48,146
10	Suicide	26,869	Suicide	42,826
Male				
Rank	All causes	1,075,078	All causes	1,328,241
1	Diseases of heart	405,661	Diseases of heart	325,077
2	Malignant neoplasms	225,948	Malignant neoplasms	311,296
3	Unintentional injuries	74,180	Unintentional injuries	85,340
4	Cerebrovascular diseases	69,973	Chronic lower respiratory diseases	69,456
5	Chronic obstructive pulmonary diseases	38,625	Cerebrovascular diseases	55,471
6	Pneumonia and influenza	27,574	Diabetes mellitus ¹	41,111
7	Suicide	20,505	Suicide	33,162
8	Chronic liver disease and cirrhosis	19,768	Alzheimer's disease	28,362
9	Homicide	18,779	Influenza and pneumonia	26,586
10	Diabetes mellitus	14,325	Chronic liver disease and cirrhosis	24,584
Female				
Rank	All causes	914,763	All causes	1,298,177
1	Diseases of heart	355,424	Diseases of heart	289,271
2	Malignant neoplasms	190,561	Malignant neoplasms	280,404
3	Cerebrovascular diseases	100,252	Chronic lower respiratory diseases	77,645
4	Unintentional injuries	31,538	Cerebrovascular diseases	77,632
5	Pneumonia and influenza	27,045	Alzheimer's disease	65,179
6	Diabetes mellitus	20,526	Unintentional injuries	50,588
7	Atherosclerosis	17,848	Diabetes mellitus ¹	35,377
8	Chronic obstructive pulmonary diseases	17,425	Influenza and pneumonia	28,641
9	Chronic liver disease and cirrhosis	10,815	Nephritis, nephrotic syndrome and nephrosis ¹	23,710
10	Certain conditions originating in the perinatal period	9,815	Septicemia	20,607
White				
Rank	All causes	1,738,607	All causes	2,237,880
1	Diseases of heart	683,347	Diseases of heart	524,695
2	Malignant neoplasms	368,162	Malignant neoplasms	502,933
3	Cerebrovascular diseases	148,734	Chronic lower respiratory diseases	134,541
4	Unintentional injuries	90,122	Unintentional injuries	117,151
5	Chronic obstructive pulmonary diseases	52,375	Cerebrovascular diseases	111,035
6	Pneumonia and influenza	48,369	Alzheimer's disease	84,990
7	Diabetes mellitus	28,868	Diabetes mellitus ¹	59,741
8	Atherosclerosis	27,069	Influenza and pneumonia	47,293
9	Chronic liver disease and cirrhosis	25,240	Suicide	38,723
10	Suicide	24,829	Nephritis, nephrotic syndrome and nephrosis ¹	37,976
Black or African American				
Rank	All causes	233,135	All causes	308,960
1	Diseases of heart	72,956	Diseases of heart	73,095
2	Malignant neoplasms	45,037	Malignant neoplasms	69,090
3	Cerebrovascular diseases	20,135	Cerebrovascular diseases	17,088
4	Unintentional injuries	13,480	Unintentional injuries	14,135
5	Homicide	10,172	Diabetes mellitus ¹	13,435
6	Certain conditions originating in the perinatal period	6,961	Chronic lower respiratory diseases	9,934
7	Pneumonia and influenza	5,648	Nephritis, nephrotic syndrome and nephrosis ¹	8,586
8	Diabetes mellitus	5,544	Homicide	7,903
9	Chronic liver disease and cirrhosis	4,790	Alzheimer's disease	6,567
10	Nephritis, nephrotic syndrome and nephrosis	3,416	Septicemia	6,386

See footnotes at end of table.

Table 19 (page 2 of 4). Leading causes of death and numbers of deaths, by sex, race, and Hispanic origin: United States, 1980 and 2014

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2015.htm#019>.

[Data are based on death certificates]

<i>Sex, race, Hispanic origin, and rank order</i>		1980		2014	
		<i>Cause of death</i>	<i>Deaths</i>	<i>Cause of death</i>	<i>Deaths</i>
American Indian or Alaska Native					
Rank	All causes		6,923	All causes	18,008
1	Diseases of heart		1,494	Diseases of heart	3,288
2	Unintentional injuries		1,290	Malignant neoplasms	3,153
3	Malignant neoplasms		770	Unintentional injuries	1,996
4	Chronic liver disease and cirrhosis		410	Chronic liver disease and cirrhosis	951
5	Cerebrovascular diseases		322	Diabetes mellitus ¹	945
6	Pneumonia and influenza		257	Chronic lower respiratory diseases	788
7	Homicide		217	Cerebrovascular diseases	649
8	Diabetes mellitus		210	Suicide	489
9	Certain conditions originating in the perinatal period		199	Influenza and pneumonia	412
10	Suicide		181	Nephritis, nephrotic syndrome and nephrosis ¹	338
Asian or Pacific Islander					
Rank	All causes		11,071	All causes	61,570
1	Diseases of heart		3,265	Malignant neoplasms	16,524
2	Malignant neoplasms		2,522	Diseases of heart	13,270
3	Cerebrovascular diseases		1,028	Cerebrovascular diseases	4,331
4	Unintentional injuries		810	Unintentional injuries	2,646
5	Pneumonia and influenza		342	Diabetes mellitus ¹	2,367
6	Suicide		249	Influenza and pneumonia	1,911
7	Certain conditions originating in the perinatal period		246	Chronic lower respiratory diseases	1,838
8	Diabetes mellitus		227	Alzheimer's disease	1,680
9	Homicide		211	Nephritis, nephrotic syndrome and nephrosis ¹	1,246
10	Chronic obstructive pulmonary diseases		207	Suicide	1,188
Hispanic or Latino					
Rank	---	---	---	All causes	169,387
1	---	---	---	Malignant neoplasms	36,447
2	---	---	---	Diseases of heart	34,021
3	---	---	---	Unintentional injuries	12,413
4	---	---	---	Cerebrovascular diseases	8,713
5	---	---	---	Diabetes mellitus ¹	7,795
6	---	---	---	Chronic liver disease and cirrhosis	5,658
7	---	---	---	Alzheimer's disease	4,934
8	---	---	---	Chronic lower respiratory diseases	4,795
9	---	---	---	Influenza and pneumonia	3,875
10	---	---	---	Nephritis, nephrotic syndrome and nephrosis ¹	3,273
White male					
Rank	All causes		933,878	All causes	1,128,993
1	Diseases of heart		364,679	Diseases of heart	277,921
2	Malignant neoplasms		198,188	Malignant neoplasms	266,137
3	Unintentional injuries		62,963	Unintentional injuries	72,807
4	Cerebrovascular diseases		60,095	Chronic lower respiratory diseases	62,989
5	Chronic obstructive pulmonary diseases		35,977	Cerebrovascular diseases	45,505
6	Pneumonia and influenza		23,810	Diabetes mellitus ¹	32,920
7	Suicide		18,901	Suicide	30,015
8	Chronic liver disease and cirrhosis		16,407	Alzheimer's disease	25,937
9	Diabetes mellitus		12,125	Influenza and pneumonia	22,643
10	Atherosclerosis		10,543	Chronic liver disease and cirrhosis	21,781
Black or African American male					
Rank	All causes		130,138	All causes	157,733
1	Diseases of heart		37,877	Diseases of heart	37,962
2	Malignant neoplasms		25,861	Malignant neoplasms	35,061
3	Unintentional injuries		9,701	Unintentional injuries	9,537
4	Cerebrovascular diseases		9,194	Cerebrovascular diseases	7,747
5	Homicide		8,274	Homicide	6,823
6	Certain conditions originating in the perinatal period		3,869	Diabetes mellitus ¹	6,452
7	Pneumonia and influenza		3,386	Chronic lower respiratory diseases	5,035
8	Chronic liver disease and cirrhosis		3,020	Nephritis, nephrotic syndrome and nephrosis ¹	4,034
9	Chronic obstructive pulmonary diseases		2,429	Septicemia	2,969
10	Diabetes mellitus		2,010	Influenza and pneumonia	2,736

See footnotes at end of table.

Table 19 (page 3 of 4). Leading causes of death and numbers of deaths, by sex, race, and Hispanic origin: United States, 1980 and 2014

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2015.htm#019>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and rank order	1980		2014	
	Cause of death	Deaths	Cause of death	Deaths
American Indian or Alaska Native male				
Rank	All causes	4,193	All causes	9,829
1	Unintentional injuries	946	Diseases of heart	1,915
2	Diseases of heart	917	Malignant neoplasms	1,727
3	Malignant neoplasms	408	Unintentional injuries	1,331
4	Chronic liver disease and cirrhosis	239	Diabetes mellitus ¹	504
5	Cerebrovascular diseases	163	Chronic liver disease and cirrhosis	481
6	Homicide	162	Suicide	363
7	Pneumonia and influenza	148	Chronic lower respiratory diseases	360
8	Suicide	147	Cerebrovascular diseases	296
9	Certain conditions originating in the perinatal period	107	Homicide	209
10	Diabetes mellitus	86	Influenza and pneumonia	195
Asian or Pacific Islander male				
Rank	All causes	6,809	All causes	31,686
1	Diseases of heart	2,174	Malignant neoplasms	8,371
2	Malignant neoplasms	1,485	Diseases of heart	7,279
3	Unintentional injuries	556	Cerebrovascular diseases	1,923
4	Cerebrovascular diseases	521	Unintentional injuries	1,665
5	Pneumonia and influenza	227	Diabetes mellitus ¹	1,235
6	Suicide	159	Chronic lower respiratory diseases	1,072
7	Chronic obstructive pulmonary diseases	158	Influenza and pneumonia	1,012
8	Homicide	151	Suicide	833
9	Certain conditions originating in the perinatal period	128	Nephritis, nephrotic syndrome and nephrosis ¹	652
10	Diabetes mellitus	103	Alzheimer's disease	508
Hispanic or Latino male				
Rank	---	---	All causes	92,474
1	---	---	Malignant neoplasms	19,040
2	---	---	Diseases of heart	18,884
3	---	---	Unintentional injuries	8,969
4	---	---	Diabetes mellitus ¹	4,149
5	---	---	Cerebrovascular diseases	4,092
6	---	---	Chronic liver disease and cirrhosis	3,911
7	---	---	Suicide	2,584
8	---	---	Chronic lower respiratory diseases	2,433
9	---	---	Homicide	2,122
10	---	---	Influenza and pneumonia	1,975
White female				
Rank	All causes	804,729	All causes	1,108,887
1	Diseases of heart	318,668	Diseases of heart	246,774
2	Malignant neoplasms	169,974	Malignant neoplasms	236,796
3	Cerebrovascular diseases	88,639	Chronic lower respiratory diseases	71,552
4	Unintentional injuries	27,159	Cerebrovascular diseases	65,530
5	Pneumonia and influenza	24,559	Alzheimer's disease	59,053
6	Diabetes mellitus	16,743	Unintentional injuries	44,344
7	Atherosclerosis	16,526	Diabetes mellitus ¹	26,821
8	Chronic obstructive pulmonary diseases	16,398	Influenza and pneumonia	24,650
9	Chronic liver disease and cirrhosis	8,833	Nephritis, nephrotic syndrome and nephrosis ¹	18,389
10	Certain conditions originating in the perinatal period	6,512	Septicemia	16,665
Black or African American female				
Rank	All causes	102,997	All causes	151,227
1	Diseases of heart	35,079	Diseases of heart	35,133
2	Malignant neoplasms	19,176	Malignant neoplasms	34,029
3	Cerebrovascular diseases	10,941	Cerebrovascular diseases	9,341
4	Unintentional injuries	3,779	Diabetes mellitus ¹	6,983
5	Diabetes mellitus	3,534	Chronic lower respiratory diseases	4,899
6	Certain conditions originating in the perinatal period	3,092	Alzheimer's disease	4,730
7	Pneumonia and influenza	2,262	Unintentional injuries	4,598
8	Homicide	1,898	Nephritis, nephrotic syndrome and nephrosis ¹	4,552
9	Chronic liver disease and cirrhosis	1,770	Septicemia	3,417
10	Nephritis, nephrotic syndrome and nephrosis	1,722	Essential hypertension and hypertensive renal disease	2,918

See footnotes at end of table.

Table 19 (page 4 of 4). Leading causes of death and numbers of deaths, by sex, race, and Hispanic origin: United States, 1980 and 2014

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2015.htm#019>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and rank order	1980		2014	
	Cause of death	Deaths	Cause of death	Deaths
American Indian or Alaska Native female				
Rank	All causes	2,730	All causes	8,179
1	Diseases of heart	577	Malignant neoplasms	1,426
2	Malignant neoplasms	362	Diseases of heart	1,373
3	Unintentional injuries	344	Unintentional injuries	665
4	Chronic liver disease and cirrhosis	171	Chronic liver disease and cirrhosis	470
5	Cerebrovascular diseases	159	Diabetes mellitus ¹	441
6	Diabetes mellitus	124	Chronic lower respiratory diseases	428
7	Pneumonia and influenza	109	Cerebrovascular diseases	353
8	Certain conditions originating in the perinatal period	92	Alzheimer's disease	224
9	Nephritis, nephrotic syndrome and nephrosis	56	Influenza and pneumonia	217
10	Homicide	55	Nephritis, nephrotic syndrome and nephrosis ¹	175
Asian or Pacific Islander female				
Rank	All causes	4,262	All causes	29,884
1	Diseases of heart	1,091	Malignant neoplasms	8,153
2	Malignant neoplasms	1,037	Diseases of heart	5,991
3	Cerebrovascular diseases	507	Cerebrovascular diseases	2,408
4	Unintentional injuries	254	Alzheimer's disease	1,172
5	Diabetes mellitus	124	Diabetes mellitus ¹	1,132
6	Certain conditions originating in the perinatal period	118	Unintentional injuries	981
7	Pneumonia and influenza	115	Influenza and pneumonia	899
8	Congenital anomalies	104	Chronic lower respiratory diseases	766
9	Suicide	90	Nephritis, nephrotic syndrome and nephrosis ¹	594
10	Homicide	60	Essential hypertension and hypertensive renal disease	572
Hispanic or Latina female				
Rank	---	---	All causes	76,913
1	---	---	Malignant neoplasms	17,407
2	---	---	Diseases of heart	15,137
3	---	---	Cerebrovascular diseases	4,621
4	---	---	Diabetes mellitus ¹	3,646
5	---	---	Unintentional injuries	3,444
6	---	---	Alzheimer's disease	3,334
7	---	---	Chronic lower respiratory diseases	2,362
8	---	---	Influenza and pneumonia	1,900
9	---	---	Chronic liver disease and cirrhosis	1,747
10	---	---	Nephritis, nephrotic syndrome and nephrosis ¹	1,573

--- Data not available. Complete coverage of all states for the Hispanic origin variable began in 1997.

¹Starting with 2011 data, the rules for selecting Renal failure as the underlying cause of death were changed, affecting the number of deaths in the Nephritis, nephrotic syndrome and nephrosis and Diabetes Categories. These changes directly affect deaths with mention of Renal failure and other associated conditions, such as Diabetes mellitus with renal complications. The result is a decrease in the number of deaths for Nephritis, nephrotic syndrome and nephrosis and an increase in the number of deaths for Diabetes mellitus. Therefore, trend data for these two causes of death should be interpreted with caution. For more information, see Technical Notes in Deaths: Final data for 2011, available from: http://www.cdc.gov/nchs/data/nvsr/nvsr63/nvsr63_03.pdf.

NOTES: For cause of death codes based on the *International Classification of Diseases, 9th Revision* (ICD-9) in 1980 and ICD-10 in 2014, see Appendix II, Cause of death; Cause-of-death ranking; Table III; Table IV. Starting with 2003 data, some states allowed the reporting of more than one race on the death certificate. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards, for comparability with other states. The race groups, white, black, Asian or Pacific Islander, and American Indian or Alaska Native, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. See Appendix II, Race; Hispanic origin. **Data have been revised and differ from previous editions of *Health, United States*.**

SOURCE: CDC/NCHS, National Vital Statistics System: Vital statistics of the United States, vol II, mortality, part A, 1980. Washington, DC: Public Health Service. 1985. Public-use 2014 Mortality File. Deaths: Final data for 2014. National vital statistics reports (forthcoming). Available from: <http://www.cdc.gov/nchs/products/nvsr.htm>. See Appendix I, National Vital Statistics System (NVSS).

Table 20 (page 1 of 2). Leading causes of death and numbers of deaths, by age: United States, 1980 and 2014

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2015.htm#020>.

[Data are based on death certificates]

Age and rank order	1980		2014	
	Cause of death	Deaths	Cause of death	Deaths
Under 1 year				
Rank	All causes	45,526	All causes	23,215
1	Congenital anomalies	9,220	Congenital malformations/deformations/ chromosomal abnormalities	4,746
2	Sudden infant death syndrome	5,510	Disorders related to short gestation and low birth weight	4,173
3	Respiratory distress syndrome	4,989	Newborn affected by maternal complications of pregnancy	1,574
4	Disorders relating to short gestation and unspecified low birthweight	3,648	Sudden infant death syndrome	1,545
5	Newborn affected by maternal complications of pregnancy	1,572	Unintentional injuries	1,160
6	Intrauterine hypoxia and birth asphyxia	1,497	Newborn affected by complications of placenta, cord and membranes	965
7	Unintentional injuries	1,166	Bacterial sepsis of newborn	544
8	Birth trauma	1,058	Respiratory distress of newborn	460
9	Pneumonia and influenza	1,012	Diseases of circulatory system	444
10	Newborn affected by complications of placenta, cord, and membranes	985	Neonatal hemorrhage	441
1–4 years				
Rank	All causes	8,187	All causes	3,830
1	Unintentional injuries	3,313	Unintentional injuries	1,216
2	Congenital anomalies	1,026	Congenital malformations/deformations/ chromosomal abnormalities	399
3	Malignant neoplasms	573	Homicide	364
4	Diseases of heart	338	Malignant neoplasms	321
5	Homicide	319	Diseases of heart	149
6	Pneumonia and influenza	267	Influenza and pneumonia	109
7	Meningitis	223	Septicemia	53
8	Meningococcal infection	110	Chronic lower respiratory diseases ¹	53
9	Certain conditions originating in the perinatal period	84	Benign neoplasms/neoplasms of uncertain/ unknown behavior	38
10	Septicemia	71	Conditions originating in perinatal period ¹	38
5–14 years				
Rank	All causes	10,689	All causes	5,250
1	Unintentional injuries	5,224	Unintentional injuries	1,480
2	Malignant neoplasms	1,497	Malignant neoplasms	852
3	Congenital anomalies	561	Suicide	428
4	Homicide	415	Congenital malformations/deformations/ chromosomal abnormalities	348
5	Diseases of heart	330	Homicide	279
6	Pneumonia and influenza	194	Diseases of heart	191
7	Suicide	142	Chronic lower respiratory diseases	139
8	Benign neoplasms	104	Influenza and pneumonia	98
9	Cerebrovascular diseases	95	Cerebrovascular diseases	88
10	Chronic obstructive pulmonary diseases	85	Benign neoplasms/neoplasms of uncertain/ unknown behavior	74
15–24 years				
Rank	All causes	49,027	All causes	28,791
1	Unintentional injuries	26,206	Unintentional injuries	11,797
2	Homicide	6,537	Suicide	5,090
3	Suicide	5,239	Homicide	4,171
4	Malignant neoplasms	2,683	Malignant neoplasms	1,569
5	Diseases of heart	1,223	Diseases of heart	953
6	Congenital anomalies	600	Congenital malformations/deformations/ chromosomal abnormalities	377
7	Cerebrovascular diseases	418	Influenza and pneumonia	199
8	Pneumonia and influenza	348	Diabetes mellitus ²	181
9	Chronic obstructive pulmonary diseases	141	Chronic lower respiratory diseases	178
10	Anemias	133	Cerebrovascular diseases	177

See footnotes at end of table.

Table 20 (page 2 of 2). Leading causes of death and numbers of deaths, by age: United States, 1980 and 2014

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2015.htm#020>.

[Data are based on death certificates]

Age and rank order	1980		2014	
	Cause of death	Deaths	Cause of death	Deaths
25–44 years				
Rank	All causes	108,658	All causes	118,173
1	Unintentional injuries	26,722	Unintentional injuries	33,366
2	Malignant neoplasms	17,551	Malignant neoplasms	14,891
3	Diseases of heart	14,513	Diseases of heart	13,709
4	Homicide	10,983	Suicide	13,289
5	Suicide	9,855	Homicide	6,769
6	Chronic liver disease and cirrhosis	4,782	Chronic liver disease and cirrhosis	3,307
7	Cerebrovascular diseases	3,154	Diabetes mellitus ²	2,708
8	Diabetes mellitus	1,472	Cerebrovascular diseases	2,324
9	Pneumonia and influenza	1,467	Human immunodeficiency virus (HIV) disease	1,757
10	Congenital anomalies	817	Influenza and pneumonia	1,674
45–64 years				
Rank	All causes	425,338	All causes	524,725
1	Diseases of heart	148,322	Malignant neoplasms	160,116
2	Malignant neoplasms	135,675	Diseases of heart	109,264
3	Cerebrovascular diseases	19,909	Unintentional injuries	38,607
4	Unintentional injuries	18,140	Chronic liver disease and cirrhosis	21,419
5	Chronic liver disease and cirrhosis	16,089	Chronic lower respiratory diseases	20,894
6	Chronic obstructive pulmonary diseases	11,514	Diabetes mellitus ²	19,404
7	Diabetes mellitus	7,977	Cerebrovascular diseases	17,076
8	Suicide	7,079	Suicide	16,313
9	Pneumonia and influenza	5,804	Septicemia	8,223
10	Homicide	4,019	Influenza and pneumonia	8,121
65 years and over				
Rank	All causes	1,341,848	All causes	1,922,271
1	Diseases of heart	595,406	Diseases of heart	489,722
2	Malignant neoplasms	258,389	Malignant neoplasms	413,886
3	Cerebrovascular diseases	146,417	Chronic lower respiratory diseases	124,693
4	Pneumonia and influenza	45,512	Cerebrovascular diseases	113,308
5	Chronic obstructive pulmonary diseases	43,587	Alzheimer's disease	92,604
6	Atherosclerosis	28,081	Diabetes mellitus ²	54,161
7	Diabetes mellitus	25,216	Unintentional injuries	48,282
8	Unintentional injuries	24,844	Influenza and pneumonia	44,836
9	Nephritis, nephrotic syndrome, and nephrosis	12,968	Nephritis, nephrotic syndrome and nephrosis ²	39,957
10	Chronic liver disease and cirrhosis	9,519	Septicemia	29,124

¹Chronic lower respiratory diseases is tied with Septicemia for the 7th rank in 2014 and Certain conditions originating in the perinatal period is tied with In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior for the 9th rank in 2014.

²Starting with 2011 data, the rules for selecting Renal failure as the underlying cause of death were changed, affecting the number of deaths in the Nephritis, nephrotic syndrome and nephrosis and Diabetes categories. These changes directly affect deaths with mention of Renal failure and other associated conditions, such as Diabetes mellitus with renal complications. The result is a decrease in the number of deaths for Nephritis, nephrotic syndrome and nephrosis and an increase in the number of deaths for Diabetes mellitus. Therefore, trend data for these two causes of death should be interpreted with caution. For more information, see Technical Notes in Deaths: Final data for 2011, available from: http://www.cdc.gov/nchs/data/nvsr/nvsr63/nvsr63_03.pdf.

NOTES: For cause of death codes based on the *International Classification of Diseases, 9th Revision (ICD–9)* in 1980 and ICD–10 in 2014, see Appendix II, Cause of death; Cause-of-death ranking; Table III; Table IV. **Data have been revised and differ from previous editions of *Health, United States*.**

SOURCE: CDC/NCHS, National Vital Statistics System: Vital statistics of the United States, vol II, mortality, part A, 1980. Washington, DC: Public Health Service. 1985. Public-use 2014 Mortality File. Deaths: Final data for 2014. National vital statistics reports (forthcoming). Available from: <http://www.cdc.gov/nchs/products/nvsr.htm>. See Appendix I, National Vital Statistics System (NVSS).

Table 21 (page 1 of 4). Death rates for all causes, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#021>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and age	1950 ¹	1960 ¹	1970	1980	1990	2000	2013	2014
Deaths per 100,000 resident population								
All persons								
All ages, age-adjusted ²	1,446.0	1,339.2	1,222.6	1,039.1	938.7	869.0	731.9	724.6
All ages, crude	963.8	954.7	945.3	878.3	863.8	854.0	821.5	823.7
Under 1 year	3,299.2	2,696.4	2,142.4	1,288.3	971.9	736.7	594.7	588.0
1–4 years	139.4	109.1	84.5	63.9	46.8	32.4	25.5	24.0
5–14 years	60.1	46.6	41.3	30.6	24.0	18.0	13.0	12.7
15–24 years	128.1	106.3	127.7	115.4	99.2	79.9	64.8	65.5
25–34 years	178.7	146.4	157.4	135.5	139.2	101.4	106.1	108.4
35–44 years	358.7	299.4	314.5	227.9	223.2	198.9	172.0	175.2
45–54 years	853.9	756.0	730.0	584.0	473.4	425.6	406.1	404.8
55–64 years	1,901.0	1,735.1	1,658.8	1,346.3	1,196.9	992.2	860.0	870.3
65–74 years	4,104.3	3,822.1	3,582.7	2,994.9	2,648.6	2,399.1	1,802.1	1,786.3
75–84 years	9,331.1	8,745.2	8,004.4	6,692.6	6,007.2	5,666.5	4,648.1	4,564.2
85 years and over	20,196.9	19,857.5	16,344.9	15,980.3	15,327.4	15,524.4	13,660.4	13,407.9
Male								
All ages, age-adjusted ²	1,674.2	1,609.0	1,542.1	1,348.1	1,202.8	1,053.8	863.6	855.1
All ages, crude	1,106.1	1,104.5	1,090.3	976.9	918.4	853.0	839.1	846.4
Under 1 year	3,728.0	3,059.3	2,410.0	1,428.5	1,082.8	806.5	650.5	638.6
1–4 years	151.7	119.5	93.2	72.6	52.4	35.9	28.6	26.7
5–14 years	70.9	55.7	50.5	36.7	28.5	20.9	14.6	14.9
15–24 years	167.9	152.1	188.5	172.3	147.4	114.9	92.6	93.8
25–34 years	216.5	187.9	215.3	196.1	204.3	138.6	145.4	148.8
35–44 years	428.8	372.8	402.6	299.2	310.4	255.2	213.8	216.7
45–54 years	1,067.1	992.2	958.5	767.3	610.3	542.8	500.7	496.5
55–64 years	2,395.3	2,309.5	2,282.7	1,815.1	1,553.4	1,230.7	1,088.4	1,098.2
65–74 years	4,931.4	4,914.4	4,873.8	4,105.2	3,491.5	2,979.6	2,186.0	2,175.5
75–84 years	10,426.0	10,178.4	10,010.2	8,816.7	7,888.6	6,972.6	5,474.2	5,369.2
85 years and over	21,636.0	21,186.3	17,821.5	18,801.1	18,056.6	17,501.4	14,911.6	14,642.2
Female								
All ages, age-adjusted ²	1,236.0	1,105.3	971.4	817.9	750.9	731.4	623.5	616.7
All ages, crude	823.5	809.2	807.8	785.3	812.0	855.0	804.4	801.7
Under 1 year	2,854.6	2,321.3	1,863.7	1,141.7	855.7	663.4	536.1	535.0
1–4 years	126.7	98.4	75.4	54.7	41.0	28.7	22.4	21.3
5–14 years	48.9	37.3	31.8	24.2	19.3	15.0	11.2	10.5
15–24 years	89.1	61.3	68.1	57.5	49.0	43.1	35.6	35.8
25–34 years	142.7	106.6	101.6	75.9	74.2	63.5	66.0	67.2
35–44 years	290.3	229.4	231.1	159.3	137.9	143.2	130.5	134.1
45–54 years	641.5	526.7	517.2	412.9	342.7	312.5	314.1	315.6
55–64 years	1,404.8	1,196.4	1,098.9	934.3	878.8	772.2	647.4	658.2
65–74 years	3,333.2	2,871.8	2,579.7	2,144.7	1,991.2	1,921.2	1,464.6	1,444.2
75–84 years	8,399.6	7,633.1	6,677.6	5,440.1	4,883.1	4,814.7	4,029.1	3,955.1
85 years and over	19,194.7	19,008.4	15,518.0	14,746.9	14,274.3	14,719.2	13,021.6	12,765.7
White male ³								
All ages, age-adjusted ²	1,642.5	1,586.0	1,513.7	1,317.6	1,165.9	1,029.4	859.2	853.4
All ages, crude	1,089.5	1,098.5	1,086.7	983.3	930.9	887.8	899.1	909.4
Under 1 year	3,400.5	2,694.1	2,113.2	1,230.3	896.1	667.6	566.4	551.3
1–4 years	135.5	104.9	83.6	66.1	45.9	32.6	26.2	23.8
5–14 years	67.2	52.7	48.0	35.0	26.4	19.8	13.9	14.0
15–24 years	152.4	143.7	170.8	167.0	131.3	105.8	87.1	88.3
25–34 years	185.3	163.2	176.6	171.3	176.1	124.1	139.8	145.3
35–44 years	380.9	332.6	343.5	257.4	268.2	233.6	208.0	212.6
45–54 years	984.5	932.2	882.9	698.9	548.7	496.9	491.7	488.9
55–64 years	2,304.4	2,225.2	2,202.6	1,728.5	1,467.2	1,163.3	1,050.7	1,063.8
65–74 years	4,864.9	4,848.4	4,810.1	4,035.7	3,397.7	2,905.7	2,152.0	2,143.3
75–84 years	10,526.3	10,299.6	10,098.8	8,829.8	7,844.9	6,933.1	5,507.2	5,419.1
85 years and over	22,116.3	21,750.0	18,551.7	19,097.3	18,268.3	17,716.4	15,220.4	15,000.3

See footnotes at end of table.

Table 21 (page 2 of 4). Death rates for all causes, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#021>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and age	1950 ¹	1960 ¹	1970	1980	1990	2000	2013	2014
Deaths per 100,000 resident population								
Black or African American male ³								
All ages, age-adjusted ²	1,909.1	1,811.1	1,873.9	1,697.8	1,644.5	1,403.5	1,052.8	1,034.0
All ages, crude	1,257.7	1,181.7	1,186.6	1,034.1	1,008.0	834.1	739.3	742.6
Under 1 year	---	5,306.8	4,298.9	2,586.7	2,112.4	1,567.6	1,120.1	1,125.4
1–4 years ⁴	1,412.6	208.5	150.5	110.5	85.8	54.5	40.6	42.2
5–14 years	95.1	75.1	67.1	47.4	41.2	28.2	19.6	21.0
15–24 years	289.7	212.0	320.6	209.1	252.2	181.4	135.5	135.4
25–34 years	503.5	402.5	559.5	407.3	430.8	261.0	218.6	212.1
35–44 years	878.1	762.0	956.6	689.8	699.6	453.0	312.4	308.5
45–54 years	1,905.0	1,624.8	1,777.5	1,479.9	1,261.0	1,017.7	678.8	671.8
55–64 years	3,773.2	3,316.4	3,256.9	2,873.0	2,618.4	2,080.1	1,628.0	1,611.5
65–74 years	5,310.3	5,798.7	5,803.2	5,131.1	4,946.1	4,253.5	3,064.4	3,047.4
75–84 years ⁵	10,101.9	8,605.1	9,454.9	9,231.6	9,129.5	8,486.0	6,362.4	6,172.6
85 years and over	---	14,844.8	12,222.3	16,098.8	16,954.9	16,791.0	13,657.1	13,291.7
American Indian or Alaska Native male ³								
All ages, age-adjusted ²	---	---	---	1,111.5	916.2	841.5	689.2	685.4
All ages, crude	---	---	---	597.1	476.4	415.6	416.5	433.2
Under 1 year	---	---	---	1,598.1	1,056.6	700.2	493.4	509.7
1–4 years	---	---	---	82.7	77.4	44.9	41.5	41.0
5–14 years	---	---	---	43.7	33.4	20.2	12.1	12.3
15–24 years	---	---	---	311.1	219.8	136.2	98.0	103.8
25–34 years	---	---	---	360.6	256.1	179.1	172.7	179.1
35–44 years	---	---	---	556.8	365.4	295.2	244.4	264.4
45–54 years	---	---	---	871.3	619.9	520.0	506.4	508.5
55–64 years	---	---	---	1,547.5	1,211.3	1,090.4	937.8	984.7
65–74 years	---	---	---	2,968.4	2,461.7	2,478.3	1,845.9	1,830.2
75–84 years	---	---	---	5,607.0	5,389.2	5,351.2	4,224.5	4,097.9
85 years and over	---	---	---	12,635.2	11,243.9	10,725.8	9,034.3	8,610.4
Asian or Pacific Islander male ³								
All ages, age-adjusted ²	---	---	---	786.5	716.4	624.2	487.8	462.0
All ages, crude	---	---	---	375.3	334.3	332.9	347.4	341.3
Under 1 year	---	---	---	816.5	605.3	529.4	408.9	384.3
1–4 years	---	---	---	50.9	45.0	23.3	19.3	14.3
5–14 years	---	---	---	23.4	20.7	12.9	11.1	9.9
15–24 years	---	---	---	80.8	76.0	55.2	42.2	44.7
25–34 years	---	---	---	83.5	79.6	55.0	54.2	54.1
35–44 years	---	---	---	128.3	130.8	104.9	88.5	83.0
45–54 years	---	---	---	342.3	287.1	249.7	220.4	212.3
55–64 years	---	---	---	881.1	789.1	642.4	517.6	519.9
65–74 years	---	---	---	2,236.1	2,041.4	1,661.0	1,126.4	1,107.1
75–84 years	---	---	---	5,389.5	5,008.6	4,328.2	3,239.9	3,047.8
85 years and over	---	---	---	13,753.6	12,446.3	12,125.3	10,142.8	9,263.1
Hispanic or Latino male ^{3,6}								
All ages, age-adjusted ²	---	---	---	---	886.4	818.1	639.8	626.8
All ages, crude	---	---	---	---	411.6	331.3	323.7	330.1
Under 1 year	---	---	---	---	921.8	637.1	501.1	508.3
1–4 years	---	---	---	---	53.8	31.5	23.1	20.1
5–14 years	---	---	---	---	26.0	17.9	11.7	12.5
15–24 years	---	---	---	---	159.3	107.7	72.9	75.4
25–34 years	---	---	---	---	234.0	120.2	99.9	103.6
35–44 years	---	---	---	---	341.8	211.0	143.3	149.4
45–54 years	---	---	---	---	533.9	439.0	348.0	341.0
55–64 years	---	---	---	---	1,123.7	965.7	797.4	787.7
65–74 years	---	---	---	---	2,368.2	2,287.9	1,710.4	1,655.1
75–84 years	---	---	---	---	5,369.1	5,395.3	4,218.1	4,103.3
85 years and over	---	---	---	---	12,272.1	13,086.2	10,596.0	10,318.0

See footnotes at end of table.

Table 21 (page 3 of 4). Death rates for all causes, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#021>.

Data are based on death certificates]

Sex, race, Hispanic origin, and age	1950 ¹	1960 ¹	1970	1980	1990	2000	2013	2014
Deaths per 100,000 resident population								
White, not Hispanic or Latino male ⁶								
All ages, age-adjusted ²	---	---	---	---	1,170.9	1,035.4	876.8	872.3
All ages, crude	---	---	---	---	985.9	978.5	1,032.1	1,045.4
Under 1 year	---	---	---	---	865.4	658.7	571.5	549.9
1–4 years	---	---	---	---	43.8	32.4	26.9	24.9
5–14 years	---	---	---	---	25.7	20.0	14.4	14.2
15–24 years	---	---	---	---	123.4	103.5	89.6	90.6
25–34 years	---	---	---	---	165.3	123.0	149.9	155.8
35–44 years	---	---	---	---	257.1	233.9	223.1	227.4
45–54 years	---	---	---	---	544.5	497.7	511.5	511.2
55–64 years	---	---	---	---	1,479.7	1,170.9	1,069.0	1,085.3
65–74 years	---	---	---	---	3,434.5	2,930.5	2,174.3	2,170.0
75–84 years	---	---	---	---	7,920.4	6,977.8	5,582.7	5,499.0
85 years and over	---	---	---	---	18,505.4	17,853.2	15,485.9	15,286.4
White female ³								
All ages, age-adjusted ²	1,198.0	1,074.4	944.0	796.1	728.8	715.3	623.6	617.6
All ages, crude	803.3	800.9	812.6	806.1	846.9	912.3	879.4	876.7
Under 1 year	2,566.8	2,007.7	1,614.6	962.5	690.0	550.5	456.8	457.6
1–4 years	112.2	85.2	66.1	49.3	36.1	25.5	20.1	19.6
5–14 years	45.1	34.7	29.9	22.9	17.9	14.1	10.6	10.0
15–24 years	71.5	54.9	61.6	55.5	45.9	41.1	35.5	35.5
25–34 years	112.8	85.0	84.1	65.4	61.5	55.1	64.6	66.9
35–44 years	235.8	191.1	193.3	138.2	117.4	125.7	126.8	130.7
45–54 years	546.4	458.8	462.9	372.7	309.3	281.4	303.4	305.8
55–64 years	1,293.8	1,078.9	1,014.9	876.2	822.7	730.9	623.4	635.1
65–74 years	3,242.8	2,779.3	2,470.7	2,066.6	1,923.5	1,868.3	1,453.2	1,433.1
75–84 years	8,481.5	7,696.6	6,698.7	5,401.7	4,839.1	4,785.3	4,072.0	4,001.3
85 years and over	19,679.5	19,477.7	15,980.2	14,979.6	14,400.6	14,890.7	13,316.1	13,079.1
Black or African American female ³								
All ages, age-adjusted ²	1,545.5	1,369.7	1,228.7	1,033.3	975.1	927.6	720.6	713.3
All ages, crude	1,002.0	905.0	829.2	733.3	747.9	733.0	651.1	655.5
Under 1 year	---	4,162.2	3,368.8	2,123.7	1,735.5	1,279.8	980.7	956.3
1–4 years ⁴	1,139.3	173.3	129.4	84.4	67.6	45.3	33.4	31.9
5–14 years	72.8	53.8	43.8	30.5	27.5	20.0	14.8	14.3
15–24 years	213.1	107.5	111.9	70.5	68.7	58.3	41.2	42.5
25–34 years	393.3	273.2	231.0	150.0	159.5	121.8	91.0	88.6
35–44 years	758.1	568.5	533.0	323.9	298.6	271.9	187.9	193.9
45–54 years	1,576.4	1,177.0	1,043.9	768.2	639.4	588.3	454.4	455.8
55–64 years	3,089.4	2,510.9	1,986.2	1,561.0	1,452.6	1,227.2	962.2	974.8
65–74 years	4,000.2	4,064.2	3,860.9	3,057.4	2,865.7	2,689.6	1,909.2	1,880.2
75–84 years ⁵	8,347.0	6,730.0	6,691.5	6,212.1	5,688.3	5,696.5	4,418.0	4,356.8
85 years and over	---	13,052.6	10,706.6	12,367.2	13,309.5	13,941.3	11,929.2	11,656.8
American Indian or Alaska Native female ³								
All ages, age-adjusted ²	---	---	---	662.4	561.8	604.5	508.3	514.1
All ages, crude	---	---	---	380.1	330.4	346.1	348.2	363.5
Under 1 year	---	---	---	1,352.6	688.7	492.2	305.9	412.5
1–4 years	---	---	---	87.5	37.8	39.8	25.5	20.4
5–14 years	---	---	---	33.5	25.5	17.7	10.9	10.8
15–24 years	---	---	---	90.3	69.0	58.9	42.8	46.8
25–34 years	---	---	---	178.5	102.3	84.8	93.5	88.0
35–44 years	---	---	---	286.0	156.4	171.9	172.4	179.0
45–54 years	---	---	---	491.4	380.9	284.9	334.5	354.3
55–64 years	---	---	---	837.1	805.9	772.1	616.6	606.2
65–74 years	---	---	---	1,765.5	1,679.4	1,899.8	1,314.7	1,368.6
75–84 years	---	---	---	3,612.9	3,073.2	3,850.0	3,221.9	3,226.6
85 years and over	---	---	---	8,567.4	8,201.1	9,118.2	8,008.4	7,893.5

See footnotes at end of table.

Table 21 (page 4 of 4). Death rates for all causes, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#021>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and age	1950 ¹	1960 ¹	1970	1980	1990	2000	2013	2014
Deaths per 100,000 resident population								
Asian or Pacific Islander female ³								
All ages, age-adjusted ²	---	---	---	425.9	469.3	416.8	343.0	331.1
All ages, crude	---	---	---	222.5	234.3	262.3	297.4	295.5
Under 1 year	---	---	---	755.8	518.2	434.3	329.3	338.6
1–4 years	---	---	---	35.4	32.0	20.0	18.4	12.5
5–14 years	---	---	---	21.5	13.0	11.7	8.8	6.5
15–24 years	---	---	---	32.3	28.8	22.4	17.7	17.8
25–34 years	---	---	---	45.4	37.5	27.6	25.7	24.8
35–44 years	---	---	---	89.7	69.9	65.6	51.5	49.1
45–54 years	---	---	---	214.1	182.7	155.5	128.0	122.3
55–64 years	---	---	---	440.8	483.4	390.9	287.2	283.2
65–74 years	---	---	---	1,027.7	1,089.2	996.4	703.9	700.5
75–84 years	---	---	---	2,833.6	3,127.9	2,882.4	2,346.7	2,237.5
85 years and over	---	---	---	7,923.3	10,254.0	9,052.2	8,240.4	7,945.3
Hispanic or Latina female ^{3,6}								
All ages, age-adjusted ²	---	---	---	---	537.1	546.0	448.6	437.5
All ages, crude	---	---	---	---	285.4	274.6	279.4	281.0
Under 1 year	---	---	---	---	746.6	553.6	433.7	432.1
1–4 years	---	---	---	---	42.1	27.5	18.4	17.2
5–14 years	---	---	---	---	17.3	13.4	9.9	9.7
15–24 years	---	---	---	---	40.6	31.7	25.6	27.2
25–34 years	---	---	---	---	62.9	43.4	41.4	41.3
35–44 years	---	---	---	---	109.3	100.5	78.3	78.5
45–54 years	---	---	---	---	253.3	223.8	189.6	185.3
55–64 years	---	---	---	---	607.5	548.4	437.2	443.3
65–74 years	---	---	---	---	1,453.8	1,423.2	1,039.9	1,036.7
75–84 years	---	---	---	---	3,351.3	3,624.5	3,037.8	2,931.3
85 years and over	---	---	---	---	10,098.7	11,202.8	9,651.3	9,250.2
White, not Hispanic or Latina female ⁶								
All ages, age-adjusted ²	---	---	---	---	734.6	721.5	638.4	633.8
All ages, crude	---	---	---	---	903.6	1,007.3	1,011.5	1,011.3
Under 1 year	---	---	---	---	655.3	530.9	448.5	451.0
1–4 years	---	---	---	---	34.0	24.4	20.4	20.2
5–14 years	---	---	---	---	17.6	13.9	10.6	9.8
15–24 years	---	---	---	---	46.0	42.6	38.0	37.5
25–34 years	---	---	---	---	60.6	56.8	70.2	73.2
35–44 years	---	---	---	---	116.8	128.1	138.6	143.8
45–54 years	---	---	---	---	312.1	285.0	320.5	325.5
55–64 years	---	---	---	---	834.5	742.1	640.5	653.5
65–74 years	---	---	---	---	1,940.2	1,891.0	1,483.9	1,462.9
75–84 years	---	---	---	---	4,887.3	4,819.3	4,142.2	4,078.9
85 years and over	---	---	---	---	14,533.1	14,971.7	13,502.5	13,290.4

--- Data not available.

¹Includes deaths of persons who were not residents of the 50 states and the District of Columbia (D.C.).

²Age-adjusted rates are calculated using the year 2000 standard population. Prior to 2001, age-adjusted rates were calculated using standard million proportions based on rounded population numbers. Starting with 2001 data, unrounded population numbers are used to calculate age-adjusted rates. See Appendix II, Age adjustment.

³The race groups, white, black, Asian or Pacific Islander, and American Indian or Alaska Native, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Death rates for Hispanic, American Indian or Alaska Native, and Asian or Pacific Islander persons should be interpreted with caution because of inconsistencies in reporting Hispanic origin or race on the death certificate (death rate numerators) compared with population figures (death rate denominators). The net effect of misclassification is an underestimation of deaths and death rates for races other than white and black. See Appendix II, Race, for a detailed discussion of sources of bias in death rates by race and Hispanic origin.

⁴In 1950, rate is for the age group under 5 years.

⁵In 1950, rate is for the age group 75 years and over.

⁶Prior to 1997, data from states that did not report Hispanic origin on the death certificate were excluded. See Appendix II, Hispanic origin.

NOTES: Starting with *Health, United States, 2003*, rates for 1991–1999 were revised using intercensal population estimates based on the 1990 and 2000 censuses. For 2000, population estimates are bridged-race April 1 census counts. Starting with *Health, United States, 2012*, rates for 2001–2009 were revised using intercensal population estimates based on the 2000 and 2010 censuses. For 2010, population estimates are bridged-race April 1 census counts. Rates for 2011 and beyond were computed using 2010-based postcensal estimates. See Appendix I, Population Census and Population Estimates. Starting with 2003 data, some states allowed the reporting of more than one race on the death certificate. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards, for comparability with other states. See Appendix II, Race. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Vital Statistics System; Grove RD, Hetzel AM. Vital statistics rates in the United States, 1940–1960. Washington, DC: U.S. Government Printing Office, 1968; numerator data from National Vital Statistics System, annual public-use Mortality Files; denominator data from national population estimates for race groups from Table 1 and unpublished Hispanic population estimates for 1985–1996 prepared by the Housing and Household Economic Statistics Division, U.S. Census Bureau; Deaths: Final data for 2014. National vital statistics reports (forthcoming). Available from: <http://www.cdc.gov/nchs/products/nvsr.htm>. See Appendix I, National Vital Statistics System (NVSS).

Table 22 (page 1 of 3). Death rates for diseases of heart, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#022>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and age	1950 ^{1,2}	1960 ^{1,2}	1970 ²	1980 ²	1990 ²	2000 ³	2013 ³	2014 ³
Deaths per 100,000 resident population								
All persons								
All ages, age-adjusted ⁴	588.8	559.0	492.7	412.1	321.8	257.6	169.8	167.0
All ages, crude	356.8	369.0	362.0	336.0	289.5	252.6	193.3	192.7
Under 1 year	4.1	6.6	13.1	22.8	20.1	13.0	7.8	8.0
1–4 years	1.6	1.3	1.7	2.6	1.9	1.2	1.1	0.9
5–14 years	3.9	1.3	0.8	0.9	0.9	0.7	0.4	0.5
15–24 years	8.2	4.0	3.0	2.9	2.5	2.6	2.1	2.2
25–34 years	20.9	15.6	11.4	8.3	7.6	7.4	7.6	7.7
35–44 years	88.3	74.6	66.7	44.6	31.4	29.2	25.6	25.6
45–54 years	309.2	271.8	238.4	180.2	120.5	94.2	80.3	80.1
55–64 years	804.3	737.9	652.3	494.1	367.3	261.2	184.6	185.8
65–74 years	1,857.2	1,740.5	1,558.2	1,218.6	894.3	665.6	390.3	385.2
75–84 years	4,311.0	4,089.4	3,683.8	2,993.1	2,295.7	1,780.3	1,095.1	1,070.2
85 years and over	9,152.5	9,317.8	7,891.3	7,777.1	6,739.9	5,926.1	4,013.9	3,920.9
Male								
All ages, age-adjusted ⁴	699.0	687.6	634.0	538.9	412.4	320.0	214.5	210.9
All ages, crude	424.7	439.5	422.5	368.6	297.6	249.8	206.5	207.1
Under 1 year	4.7	7.8	15.1	25.5	21.9	13.3	8.7	8.6
1–4 years	1.7	1.4	1.9	2.8	1.9	1.4	1.2	0.9
5–14 years	3.5	1.4	0.9	1.0	0.9	0.8	0.4	0.5
15–24 years	8.3	4.2	3.7	3.7	3.1	3.2	2.8	2.7
25–34 years	24.4	20.1	15.2	11.4	10.3	9.6	10.2	10.3
35–44 years	120.4	112.7	103.2	68.7	48.1	41.4	35.5	34.8
45–54 years	441.2	420.4	376.4	282.6	183.0	140.2	115.1	113.5
55–64 years	1,100.5	1,066.9	987.2	746.8	537.3	371.7	267.3	268.1
65–74 years	2,310.2	2,291.3	2,170.3	1,728.0	1,250.0	898.3	530.9	525.7
75–84 years	4,825.8	4,742.4	4,534.8	3,834.3	2,968.2	2,248.1	1,382.4	1,354.8
85 years and over	9,661.4	9,788.9	8,426.2	8,752.7	7,418.4	6,430.0	4,564.2	4,453.4
Female								
All ages, age-adjusted ⁴	486.6	447.0	381.6	320.8	257.0	210.9	134.3	131.8
All ages, crude	289.7	300.6	304.5	305.1	281.8	255.3	180.6	178.6
Under 1 year	3.4	5.4	10.9	20.0	18.3	12.5	6.9	7.4
1–4 years	1.6	1.1	1.6	2.5	1.9	1.0	0.9	0.9
5–14 years	4.3	1.2	0.8	0.9	0.8	0.5	0.4	0.4
15–24 years	8.2	3.7	2.3	2.1	1.8	2.1	1.5	1.6
25–34 years	17.6	11.3	7.7	5.3	5.0	5.2	4.9	5.0
35–44 years	57.0	38.2	32.2	21.4	15.1	17.2	15.7	16.4
45–54 years	177.8	127.5	109.9	84.5	61.0	49.8	46.6	47.5
55–64 years	507.0	429.4	351.6	272.1	215.7	159.3	107.5	109.3
65–74 years	1,434.9	1,261.3	1,082.7	828.6	616.8	474.0	266.8	261.7
75–84 years	3,873.0	3,582.7	3,120.8	2,497.0	1,893.8	1,475.1	879.8	854.8
85 years and over	8,798.1	9,016.8	7,591.8	7,350.5	6,478.1	5,720.9	3,732.9	3,643.8
White male ⁵								
All ages, age-adjusted ⁴	701.4	694.5	640.2	539.6	409.2	316.7	213.1	210.0
All ages, crude	434.2	454.6	438.3	384.0	312.7	265.8	222.6	223.9
45–54 years	424.1	413.2	365.7	269.8	170.6	130.7	109.8	107.7
55–64 years	1,082.6	1,056.0	979.3	730.6	516.7	351.8	254.0	255.9
65–74 years	2,309.4	2,297.9	2,177.2	1,729.7	1,230.5	877.8	516.0	512.7
75–84 years	4,908.0	4,839.9	4,617.6	3,883.2	2,983.4	2,247.0	1,389.0	1,361.5
85 years and over	9,952.3	10,135.8	8,818.0	8,958.0	7,558.7	6,560.8	4,701.6	4,603.9
Black or African American male ⁵								
All ages, age-adjusted ⁴	641.5	615.2	607.3	561.4	485.4	392.5	262.8	259.5
All ages, crude	348.4	330.6	330.3	301.0	256.8	211.1	177.2	178.7
45–54 years	624.1	514.0	512.8	433.4	328.9	247.2	178.3	179.5
55–64 years	1,434.0	1,236.8	1,135.4	987.2	824.0	631.2	430.0	425.5
65–74 years	2,140.1	2,281.4	2,237.8	1,847.2	1,632.9	1,268.8	807.5	794.6
75–84 years ⁶	4,107.9	3,533.6	3,783.4	3,578.8	3,107.1	2,597.6	1,660.0	1,651.4
85 years and over	---	6,037.9	5,367.6	6,819.5	6,479.6	5,633.5	3,754.5	3,693.0

See footnotes at end of table.

Table 22 (page 2 of 3). Death rates for diseases of heart, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#022>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and age	1950 ^{1,2}	1960 ^{1,2}	1970 ²	1980 ²	1990 ²	2000 ³	2013 ³	2014 ³
Deaths per 100,000 resident population								
American Indian or Alaska Native male⁵								
All ages, age-adjusted ⁴	---	---	---	320.5	264.1	222.2	152.3	149.7
All ages, crude	---	---	---	130.6	108.0	90.1	82.3	84.4
45–54 years	---	---	---	238.1	173.8	108.5	101.6	95.9
55–64 years	---	---	---	496.3	411.0	285.0	206.2	217.2
65–74 years	---	---	---	1,009.4	839.1	748.2	465.2	444.0
75–84 years	---	---	---	2,062.2	1,788.8	1,655.7	1,071.8	978.2
85 years and over	---	---	---	4,413.7	3,860.3	3,318.3	2,269.1	2,415.9
Asian or Pacific Islander male⁵								
All ages, age-adjusted ⁴	---	---	---	286.9	220.7	185.5	118.4	109.1
All ages, crude	---	---	---	119.8	88.7	90.6	81.9	78.4
45–54 years	---	---	---	112.0	70.4	61.1	50.9	53.4
55–64 years	---	---	---	306.7	226.1	182.6	133.3	121.4
65–74 years	---	---	---	852.4	623.5	482.5	274.9	250.5
75–84 years	---	---	---	2,010.9	1,642.2	1,354.7	754.6	726.6
85 years and over	---	---	---	5,923.0	4,617.8	4,154.2	2,848.5	2,459.7
Hispanic or Latino male^{5,7}								
All ages, age-adjusted ⁴	---	---	---	---	270.0	238.2	151.5	145.7
All ages, crude	---	---	---	---	91.0	74.7	66.9	67.4
45–54 years	---	---	---	---	116.4	84.3	63.5	64.0
55–64 years	---	---	---	---	363.0	264.8	178.0	169.5
65–74 years	---	---	---	---	829.9	684.8	416.6	391.6
75–84 years	---	---	---	---	1,971.3	1,733.2	1,057.5	1,019.7
85 years and over	---	---	---	---	4,711.9	4,897.5	3,106.2	2,987.5
White, not Hispanic or Latino male⁷								
All ages, age-adjusted ⁴	---	---	---	---	413.6	319.9	217.9	215.2
All ages, crude	---	---	---	---	336.5	297.5	259.0	261.0
45–54 years	---	---	---	---	172.8	134.3	117.2	115.0
55–64 years	---	---	---	---	521.3	356.3	260.2	263.6
65–74 years	---	---	---	---	1,243.4	885.1	520.2	518.9
75–84 years	---	---	---	---	3,007.7	2,261.9	1,408.1	1,382.1
85 years and over	---	---	---	---	7,663.4	6,606.6	4,794.2	4,703.8
White female⁵								
All ages, age-adjusted ⁴	479.2	441.7	376.7	315.9	250.9	205.6	132.0	130.0
All ages, crude	290.5	306.5	313.8	319.2	298.4	274.5	196.8	195.1
45–54 years	142.4	103.4	91.4	71.2	50.2	40.9	41.0	42.2
55–64 years	460.7	383.0	317.7	248.1	192.4	141.3	95.9	98.8
65–74 years	1,401.6	1,229.8	1,044.0	796.7	583.6	445.2	252.5	247.6
75–84 years	3,926.2	3,629.7	3,143.5	2,493.6	1,874.3	1,452.4	871.8	849.0
85 years and over	9,086.9	9,280.8	7,839.9	7,501.6	6,563.4	5,801.4	3,821.0	3,746.8
Black or African American female⁵								
All ages, age-adjusted ⁴	538.9	488.9	435.6	378.6	327.5	277.6	172.1	167.7
All ages, crude	289.9	268.5	261.0	249.7	237.0	212.6	153.4	152.3
45–54 years	526.8	360.7	290.9	202.4	155.3	125.0	94.2	94.9
55–64 years	1,210.7	952.3	710.5	530.1	442.0	332.8	213.0	211.0
65–74 years	1,659.4	1,680.5	1,553.2	1,210.3	1,017.5	815.2	450.8	437.9
75–84 years ⁶	3,499.3	2,926.9	2,964.1	2,707.2	2,250.9	1,913.1	1,122.2	1,085.7
85 years and over	---	5,650.0	5,003.8	5,796.5	5,766.1	5,298.7	3,425.3	3,269.8

See footnotes at end of table.

Table 22 (page 3 of 3). Death rates for diseases of heart, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#022>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and age	1950 ^{1,2}	1960 ^{1,2}	1970 ²	1980 ²	1990 ²	2000 ³	2013 ³	2014 ³
Deaths per 100,000 resident population								
American Indian or Alaska Native female⁵								
All ages, age-adjusted ⁴	---	---	---	175.4	153.1	143.6	93.9	94.0
All ages, crude	---	---	---	80.3	77.5	71.9	58.4	61.0
45–54 years	---	---	---	65.2	62.0	40.2	37.3	40.8
55–64 years	---	---	---	193.5	197.0	149.4	100.5	90.4
65–74 years	---	---	---	577.2	492.8	391.8	220.1	269.0
75–84 years	---	---	---	1,364.3	1,050.3	1,044.1	664.6	657.3
85 years and over	---	---	---	2,893.3	2,868.7	3,146.3	2,089.2	1,931.7
Asian or Pacific Islander female⁵								
All ages, age-adjusted ⁴	---	---	---	132.3	149.2	115.7	73.3	68.2
All ages, crude	---	---	---	57.0	62.0	65.0	61.8	59.2
45–54 years	---	---	---	28.6	17.5	15.9	12.5	11.9
55–64 years	---	---	---	92.9	99.0	68.8	41.7	34.9
65–74 years	---	---	---	313.3	323.9	229.6	124.7	124.8
75–84 years	---	---	---	1,053.2	1,130.9	866.2	521.0	485.4
85 years and over	---	---	---	3,211.0	4,161.2	3,367.2	2,272.4	2,097.2
Hispanic or Latina female^{5,7}								
All ages, age-adjusted ⁴	---	---	---	---	177.2	163.7	97.0	92.4
All ages, crude	---	---	---	---	79.4	71.5	55.9	55.3
45–54 years	---	---	---	---	43.5	28.2	22.2	21.3
55–64 years	---	---	---	---	153.2	111.2	63.1	70.2
65–74 years	---	---	---	---	460.4	366.3	195.8	190.0
75–84 years	---	---	---	---	1,259.7	1,169.4	702.3	657.5
85 years and over	---	---	---	---	4,440.3	4,605.8	2,738.8	2,558.9
White, not Hispanic or Latina female⁷								
All ages, age-adjusted ⁴	---	---	---	---	252.6	206.8	134.6	133.0
All ages, crude	---	---	---	---	320.0	304.9	228.0	226.9
45–54 years	---	---	---	---	50.2	41.9	44.0	45.8
55–64 years	---	---	---	---	193.6	142.9	99.0	101.4
65–74 years	---	---	---	---	584.7	448.5	256.3	251.3
75–84 years	---	---	---	---	1,890.2	1,458.9	881.3	862.1
85 years and over	---	---	---	---	6,615.2	5,822.7	3,876.5	3,813.8

--- Data not available.

¹Includes deaths of persons who were not residents of the 50 states and the District of Columbia (D.C.).

²Underlying cause of death was coded according to the 6th Revision of the *International Classification of Diseases* (ICD) in 1950, 7th Revision in 1960, 8th Revision in 1970, and 9th Revision in 1980–1998. See Appendix II, Cause of death; Table III; Table IV.

³Starting with 1999 data, cause of death is coded according to ICD–10. See Appendix II, Cause of death; Comparability ratio; Table IV; Table V.

⁴Age-adjusted rates are calculated using the year 2000 standard population. Prior to 2001, age-adjusted rates were calculated using standard million proportions based on rounded population numbers. Starting with 2001 data, unrounded population numbers are used to calculate age-adjusted rates. See Appendix II, Age adjustment.

⁵The race groups, white, black, Asian or Pacific Islander, and American Indian or Alaska Native, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Death rates for Hispanic, American Indian or Alaska Native, and Asian or Pacific Islander persons should be interpreted with caution because of inconsistencies in reporting Hispanic origin or race on the death certificate (death rate numerators) compared with population figures (death rate denominators). The net effect of misclassification is an underestimation of deaths and death rates for races other than white and black. See Appendix II, Race, for a detailed discussion of sources of bias in death rates by race and Hispanic origin.

⁶In 1950, rate is for the age group 75 years and over.

⁷Prior to 1997, data from states that did not report Hispanic origin on the death certificate were excluded. See Appendix II, Hispanic origin.

NOTES: Starting with *Health, United States, 2003*, rates for 1991–1999 were revised using intercensal population estimates based on the 1990 and 2000 censuses. For 2000, population estimates are bridged-race April 1 census counts. Starting with *Health, United States, 2012*, rates for 2001–2009 were revised using intercensal population estimates based on the 2000 and 2010 censuses. For 2010, population estimates are bridged-race April 1 census counts. Rates for 2011 and beyond were computed using 2010-based postcensal estimates. See Appendix I, Population Census and Population Estimates. Age groups were selected to minimize the presentation of unstable age-specific death rates based on small numbers of deaths and for consistency among comparison groups. Starting with 2003 data, some states allowed the reporting of more than one race on the death certificate. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards, for comparability with other states. See Appendix II, Race. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Vital Statistics System; numerator data from National Vital Statistics System, annual public-use Mortality Files; denominator data from national population estimates for race groups from Table 1 and unpublished Hispanic population estimates for 1985–1996 prepared by the Housing and Household Economic Statistics Division, U.S. Census Bureau; Deaths: Final data for 2014. National vital statistics reports (forthcoming). Available from: <http://www.cdc.gov/nchs/products/nvsr.htm>. See Appendix I, National Vital Statistics System (NVSS).

Table 23 (page 1 of 3). Death rates for cerebrovascular diseases, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#023>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and age	1950 ^{1,2}	1960 ^{1,2}	1970 ²	1980 ²	1990 ²	2000 ³	2013 ³	2014 ³
Deaths per 100,000 resident population								
All persons								
All ages, age-adjusted ⁴	180.7	177.9	147.7	96.2	65.3	60.9	36.2	36.5
All ages, crude	104.0	108.0	101.9	75.0	57.8	59.6	40.8	41.7
Under 1 year	5.1	4.1	5.0	4.4	3.8	3.3	2.7	2.4
1–4 years	0.9	0.8	1.0	0.5	0.3	0.3	0.2	0.2
5–14 years	0.5	0.7	0.7	0.3	0.2	0.2	0.2	0.2
15–24 years	1.6	1.8	1.6	1.0	0.6	0.5	0.3	0.4
25–34 years	4.2	4.7	4.5	2.6	2.2	1.5	1.2	1.3
35–44 years	18.7	14.7	15.6	8.5	6.4	5.8	4.2	4.3
45–54 years	70.4	49.2	41.6	25.2	18.7	16.0	12.4	12.3
55–64 years	194.2	147.3	115.8	65.1	47.9	41.0	28.9	29.3
65–74 years	554.7	469.2	384.1	219.0	144.2	128.6	74.2	74.5
75–84 years	1,499.6	1,491.3	1,254.2	786.9	498.0	461.3	268.9	265.7
85 years and over	2,990.1	3,680.5	3,014.3	2,283.7	1,628.9	1,589.2	906.0	929.7
Male								
All ages, age-adjusted ⁴	186.4	186.1	157.4	102.2	68.5	62.4	36.7	36.9
All ages, crude	102.5	104.5	94.5	63.4	46.7	46.9	34.5	35.3
Under 1 year	6.4	5.0	5.8	5.0	4.4	3.8	3.0	2.6
1–4 years	1.1	0.9	1.2	0.4	0.3	*	0.3	*
5–14 years	0.5	0.7	0.8	0.3	0.2	0.2	0.2	0.2
15–24 years	1.8	1.9	1.8	1.1	0.7	0.5	0.4	0.5
25–34 years	4.2	4.5	4.4	2.6	2.1	1.5	1.3	1.5
35–44 years	17.5	14.6	15.7	8.7	6.8	5.8	4.7	5.0
45–54 years	67.9	52.2	44.4	27.2	20.5	17.5	14.2	14.0
55–64 years	205.2	163.8	138.7	74.6	54.3	47.2	35.1	35.2
65–74 years	589.6	530.7	449.5	258.6	166.6	145.0	85.0	85.1
75–84 years	1,543.6	1,555.9	1,361.6	866.3	551.1	490.8	277.9	272.8
85 years and over	3,048.6	3,643.1	2,895.2	2,193.6	1,528.5	1,484.3	808.4	832.0
Female								
All ages, age-adjusted ⁴	175.8	170.7	140.0	91.7	62.6	59.1	35.2	35.6
All ages, crude	105.6	111.4	109.0	85.9	68.4	71.8	46.9	47.9
Under 1 year	3.7	3.2	4.0	3.8	3.1	2.7	2.5	2.1
1–4 years	0.7	0.7	0.7	0.5	0.3	0.4	*	*
5–14 years	0.4	0.6	0.6	0.3	0.2	0.2	0.2	0.2
15–24 years	1.5	1.6	1.4	0.8	0.6	0.5	0.3	0.3
25–34 years	4.3	4.9	4.7	2.6	2.2	1.5	1.1	1.2
35–44 years	19.9	14.8	15.6	8.4	6.1	5.7	3.7	3.7
45–54 years	72.9	46.3	39.0	23.3	17.0	14.5	10.6	10.7
55–64 years	183.1	131.8	95.3	56.8	42.2	35.3	23.1	23.8
65–74 years	522.1	415.7	333.3	188.7	126.7	115.1	64.8	65.2
75–84 years	1,462.2	1,441.1	1,183.1	740.1	466.2	442.1	262.1	260.3
85 years and over	2,949.4	3,704.4	3,081.0	2,323.1	1,667.6	1,632.0	955.8	980.6
White male ⁵								
All ages, age-adjusted ⁴	182.1	181.6	153.7	98.7	65.5	59.8	35.0	35.2
All ages, crude	100.5	102.7	93.5	63.1	46.9	48.4	35.8	36.7
45–54 years	53.7	40.9	35.6	21.7	15.4	13.6	11.7	11.9
55–64 years	182.2	139.0	119.9	64.0	45.7	39.7	29.9	29.6
65–74 years	569.7	501.0	420.0	239.8	152.9	133.8	77.6	77.3
75–84 years	1,556.3	1,564.8	1,361.6	852.7	539.2	480.0	271.1	265.3
85 years and over	3,127.1	3,734.8	3,018.1	2,230.8	1,545.4	1,490.7	815.2	841.4
Black or African American male ⁵								
All ages, age-adjusted ⁴	228.8	238.5	206.4	142.0	102.2	89.6	54.1	55.1
All ages, crude	122.0	122.9	108.8	73.0	53.0	46.1	35.1	36.5
45–54 years	211.9	166.1	136.1	82.1	68.4	49.5	30.7	29.6
55–64 years	522.8	439.9	343.4	189.7	141.7	115.4	78.1	79.4
65–74 years	783.6	899.2	780.1	472.3	326.9	268.5	165.0	169.8
75–84 years ⁶	1,504.9	1,475.2	1,445.7	1,066.3	721.5	659.2	387.4	393.2
85 years and over	---	2,700.0	1,963.1	1,873.2	1,421.5	1,458.8	814.1	827.6

See footnotes at end of table.

Table 23 (page 2 of 3). Death rates for cerebrovascular diseases, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#023>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and age	1950 ^{1,2}	1960 ^{1,2}	1970 ²	1980 ²	1990 ²	2000 ³	2013 ³	2014 ³
Deaths per 100,000 resident population								
American Indian or Alaska Native male⁵								
All ages, age-adjusted ⁴	---	---	---	66.4	44.3	46.1	22.7	25.3
All ages, crude	---	---	---	23.1	16.0	16.8	11.3	13.0
45–54 years	---	---	---	*	*	13.3	9.2	8.4
55–64 years	---	---	---	72.0	39.8	48.6	23.8	29.8
65–74 years	---	---	---	170.5	120.3	144.7	72.5	74.7
75–84 years	---	---	---	523.9	325.9	373.3	191.0	193.0
85 years and over	---	---	---	1,384.7	949.8	834.9	348.3	463.9
Asian or Pacific Islander male⁵								
All ages, age-adjusted ⁴	---	---	---	71.4	59.1	58.0	31.2	29.4
All ages, crude	---	---	---	28.7	23.3	27.2	21.3	20.7
45–54 years	---	---	---	17.0	15.6	15.0	15.7	11.0
55–64 years	---	---	---	59.9	51.8	49.3	27.7	30.0
65–74 years	---	---	---	197.9	167.9	135.6	73.0	66.4
75–84 years	---	---	---	619.5	483.9	438.7	230.5	217.2
85 years and over	---	---	---	1,399.0	1,196.6	1,415.6	686.7	666.1
Hispanic or Latino male^{5,7}								
All ages, age-adjusted ⁴	---	---	---	---	46.5	50.5	31.8	32.1
All ages, crude	---	---	---	---	15.6	15.8	14.0	14.6
45–54 years	---	---	---	---	20.0	18.1	13.8	13.3
55–64 years	---	---	---	---	49.2	48.8	33.9	33.2
65–74 years	---	---	---	---	126.4	136.1	82.1	77.0
75–84 years	---	---	---	---	356.6	392.9	242.1	245.8
85 years and over	---	---	---	---	866.3	1,029.9	620.6	656.6
White, not Hispanic or Latino male⁷								
All ages, age-adjusted ⁴	---	---	---	---	66.3	59.9	34.9	35.1
All ages, crude	---	---	---	---	50.6	53.9	40.8	41.8
45–54 years	---	---	---	---	14.9	13.0	11.0	11.3
55–64 years	---	---	---	---	45.1	38.7	29.0	28.7
65–74 years	---	---	---	---	154.5	133.1	76.6	76.7
75–84 years	---	---	---	---	547.3	482.3	272.0	265.8
85 years and over	---	---	---	---	1,578.7	1,505.9	825.6	851.2
White female⁵								
All ages, age-adjusted ⁴	169.7	165.0	135.5	89.0	60.3	57.3	34.2	34.7
All ages, crude	103.3	110.1	109.8	88.6	71.6	76.9	50.6	51.8
45–54 years	55.0	33.8	30.5	18.6	13.5	11.2	9.0	8.8
55–64 years	156.9	103.0	78.1	48.6	35.8	30.2	19.8	20.4
65–74 years	498.1	383.3	303.2	172.5	116.1	107.3	60.0	60.2
75–84 years	1,471.3	1,444.7	1,176.8	728.8	456.5	434.2	257.7	257.1
85 years and over	3,017.9	3,795.7	3,167.6	2,362.7	1,685.9	1,646.7	970.4	995.7
Black or African American female⁵								
All ages, age-adjusted ⁴	238.4	232.5	189.3	119.6	84.0	76.2	44.7	45.2
All ages, crude	128.3	127.7	112.2	77.8	60.7	58.3	39.2	40.5
45–54 years	248.9	166.2	119.4	61.8	44.1	38.1	21.1	22.4
55–64 years	567.7	452.0	272.4	138.4	96.9	76.4	47.7	48.4
65–74 years	754.4	830.5	673.5	361.7	236.7	190.9	109.1	112.6
75–84 years ⁶	1,496.7	1,413.1	1,338.3	917.5	595.0	549.2	333.0	322.6
85 years and over	---	2,578.9	2,210.5	1,891.6	1,495.2	1,556.5	894.1	934.0

See footnotes at end of table.

Table 23 (page 3 of 3). Death rates for cerebrovascular diseases, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#023>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and age	1950 ^{1,2}	1960 ^{1,2}	1970 ²	1980 ²	1990 ²	2000 ³	2013 ³	2014 ³
Deaths per 100,000 resident population								
American Indian or Alaska Native female⁵								
All ages, age-adjusted ⁴	---	---	---	51.2	38.4	43.7	25.5	25.0
All ages, crude	---	---	---	22.0	19.3	21.5	15.4	15.7
45–54 years	---	---	---	*	*	14.4	9.0	11.2
55–64 years	---	---	---	*	40.7	37.9	19.1	21.6
65–74 years	---	---	---	128.3	100.5	79.5	60.6	52.4
75–84 years	---	---	---	404.2	282.0	391.1	187.2	183.2
85 years and over	---	---	---	1,095.5	776.2	931.5	593.6	618.8
Asian or Pacific Islander female⁵								
All ages, age-adjusted ⁴	---	---	---	60.8	54.9	49.1	27.9	27.2
All ages, crude	---	---	---	26.4	24.3	28.7	23.8	23.8
45–54 years	---	---	---	20.3	19.7	13.3	8.6	8.6
55–64 years	---	---	---	43.7	42.1	33.3	18.2	18.4
65–74 years	---	---	---	136.1	124.0	102.8	54.0	52.3
75–84 years	---	---	---	446.6	396.6	386.0	203.0	197.9
85 years and over	---	---	---	1,545.2	1,395.0	1,246.6	754.6	747.4
Hispanic or Latina female^{5,7}								
All ages, age-adjusted ⁴	---	---	---	---	43.7	43.0	27.6	28.3
All ages, crude	---	---	---	---	20.1	19.4	16.1	16.9
45–54 years	---	---	---	---	15.2	12.4	9.1	7.7
55–64 years	---	---	---	---	38.5	31.9	19.7	17.2
65–74 years	---	---	---	---	102.6	95.2	53.6	54.7
75–84 years	---	---	---	---	308.5	311.3	211.4	219.7
85 years and over	---	---	---	---	1,055.3	1,108.9	698.4	744.6
White, not Hispanic or Latina female⁷								
All ages, age-adjusted ⁴	---	---	---	---	61.0	57.6	34.5	35.0
All ages, crude	---	---	---	---	77.2	85.5	58.3	59.7
45–54 years	---	---	---	---	13.2	10.9	8.8	8.9
55–64 years	---	---	---	---	35.7	29.9	19.6	20.6
65–74 years	---	---	---	---	116.9	107.6	60.2	60.3
75–84 years	---	---	---	---	461.9	438.3	260.3	259.0
85 years and over	---	---	---	---	1,714.7	1,661.6	984.3	1,009.2

* Rates based on fewer than 20 deaths are considered unreliable and are not shown.

--- Data not available.

¹Includes deaths of persons who were not residents of the 50 states and the District of Columbia (D.C.).

²Underlying cause of death was coded according to the 6th Revision of the *International Classification of Diseases* (ICD) in 1950, 7th Revision in 1960, 8th Revision in 1970, and 9th Revision in 1980–1998. See Appendix II, Cause of death; Table III; Table IV.

³Starting with 1999 data, cause of death is coded according to ICD–10. See Appendix II, Cause of death; Comparability ratio; Table IV; Table V.

⁴Age-adjusted rates are calculated using the year 2000 standard population. Prior to 2001, age-adjusted rates were calculated using standard million proportions based on rounded population numbers. Starting with 2001 data, unrounded population numbers are used to calculate age-adjusted rates. See Appendix II, Age adjustment.

⁵The race groups, white, black, Asian or Pacific Islander, and American Indian or Alaska Native, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Death rates for Hispanic, American Indian or Alaska Native, and Asian or Pacific Islander persons should be interpreted with caution because of inconsistencies in reporting Hispanic origin or race on the death certificate (death rate numerators) compared with population figures (death rate denominators). The net effect of misclassification is an underestimation of deaths and death rates for races other than white and black. See Appendix II, Race, for a detailed discussion of sources of bias in death rates by race and Hispanic origin.

⁶In 1950, rate is for the age group 75 years and over.

⁷Prior to 1997, data from states that did not report Hispanic origin on the birth certificate were excluded. See Appendix II, Hispanic origin.

NOTES: Starting with *Health, United States, 2003*, rates for 1991–1999 were revised using intercensal population estimates based on the 1990 and 2000 censuses. For 2000, population estimates are bridged-race April 1 census counts. Starting with *Health, United States, 2012*, rates for 2001–2009 were revised using intercensal population estimates based on the 2000 and 2010 censuses. For 2010, population estimates are bridged-race April 1 census counts. Rates for 2011 and beyond were computed using 2010-based postcensal estimates. See Appendix I, Population Census and Population Estimates. Age groups were selected to minimize the presentation of unstable age-specific death rates based on small numbers of deaths and for consistency among comparison groups. Starting with 2003 data, some states allowed the reporting of more than one race on the death certificate. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards, for comparability with other states. See Appendix II, Race. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Vital Statistics System; Grove RD, Hetzel AM. Vital statistics rates in the United States, 1940–1960. Washington, DC: U.S. Government Printing Office. 1968; numerator data from National Vital Statistics System, annual public-use Mortality Files; denominator data from national population estimates for race groups from Table 1 and unpublished Hispanic population estimates for 1985–1996 prepared by the Housing and Household Economic Statistics Division, U.S. Census Bureau; Deaths: Final data for 2014. National vital statistics reports (forthcoming). Available from: <http://www.cdc.gov/nchs/products/nvsr.htm>. See Appendix I, National Vital Statistics System (NVSS).

Table 24 (page 1 of 4). Death rates for malignant neoplasms, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#024>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and age	1950 ^{1,2}	1960 ^{1,2}	1970 ²	1980 ²	1990 ²	2000 ³	2013 ³	2014 ³
Deaths per 100,000 resident population								
All persons								
All ages, age-adjusted ⁴	193.9	193.9	198.6	207.9	216.0	199.6	163.2	161.2
All ages, crude	139.8	149.2	162.8	183.9	203.2	196.5	185.0	185.6
Under 1 year	8.7	7.2	4.7	3.2	2.3	2.4	1.6	1.3
1–4 years	11.7	10.9	7.5	4.5	3.5	2.7	2.1	2.0
5–14 years	6.7	6.8	6.0	4.3	3.1	2.5	2.2	2.1
15–24 years	8.6	8.3	8.3	6.3	4.9	4.4	3.4	3.6
25–34 years	20.0	19.5	16.5	13.7	12.6	9.8	8.6	8.3
35–44 years	62.7	59.7	59.5	48.6	43.3	36.6	28.1	27.8
45–54 years	175.1	177.0	182.5	180.0	158.9	127.5	105.5	103.2
55–64 years	390.7	396.8	423.0	436.1	449.6	366.7	288.2	287.6
65–74 years	698.8	713.9	754.2	817.9	872.3	816.3	616.9	603.1
75–84 years	1,153.3	1,127.4	1,169.2	1,232.3	1,348.5	1,335.6	1,139.4	1,125.9
85 years and over	1,451.0	1,450.0	1,320.7	1,594.6	1,752.9	1,819.4	1,635.4	1,632.9
Male								
All ages, age-adjusted ⁴	208.1	225.1	247.6	271.2	280.4	248.9	196.0	192.9
All ages, crude	142.9	162.5	182.1	205.3	221.3	207.2	197.6	198.4
Under 1 year	9.7	7.7	4.4	3.7	2.4	2.6	1.5	1.1
1–4 years	12.5	12.4	8.3	5.2	3.7	3.0	2.2	2.2
5–14 years	7.4	7.6	6.7	4.9	3.5	2.7	2.2	2.3
15–24 years	9.7	10.2	10.4	7.8	5.7	5.1	3.8	4.2
25–34 years	17.7	18.8	16.3	13.4	12.6	9.2	8.6	8.2
35–44 years	45.6	48.9	53.0	44.0	38.5	32.7	24.0	24.0
45–54 years	156.2	170.8	183.5	188.7	162.5	130.9	106.5	102.9
55–64 years	413.1	459.9	511.8	520.8	532.9	415.8	331.3	330.3
65–74 years	791.5	890.5	1,006.8	1,093.2	1,122.2	1,001.9	726.2	711.9
75–84 years	1,332.6	1,389.4	1,588.3	1,790.5	1,914.4	1,760.6	1,414.5	1,387.5
85 years and over	1,668.3	1,741.2	1,720.8	2,369.5	2,739.9	2,710.7	2,272.6	2,250.4
Female								
All ages, age-adjusted ⁴	182.3	168.7	163.2	166.7	175.7	167.6	139.5	138.1
All ages, crude	136.8	136.4	144.4	163.6	186.0	186.2	172.8	173.2
Under 1 year	7.6	6.8	5.0	2.7	2.2	2.3	1.8	1.6
1–4 years	10.8	9.3	6.7	3.7	3.2	2.5	1.9	1.8
5–14 years	6.0	6.0	5.2	3.6	2.8	2.2	2.1	1.8
15–24 years	7.6	6.5	6.2	4.8	4.1	3.6	3.0	2.9
25–34 years	22.2	20.1	16.7	14.0	12.6	10.4	8.6	8.5
35–44 years	79.3	70.0	65.6	53.1	48.1	40.4	32.1	31.6
45–54 years	194.0	183.0	181.5	171.8	155.5	124.2	104.6	103.4
55–64 years	368.2	337.7	343.2	361.7	375.2	321.3	248.1	247.9
65–74 years	612.3	560.2	557.9	607.1	677.4	663.6	520.8	507.5
75–84 years	1,000.7	924.1	891.9	903.1	1,010.3	1,058.5	933.3	928.0
85 years and over	1,299.7	1,263.9	1,096.7	1,255.7	1,372.1	1,456.4	1,310.1	1,311.7
White male ⁵								
All ages, age-adjusted ⁴	210.0	224.7	244.8	265.1	272.2	243.9	195.5	193.0
All ages, crude	147.2	166.1	185.1	208.7	227.7	218.1	213.0	214.4
25–34 years	17.7	18.8	16.2	13.6	12.3	9.2	8.5	8.4
35–44 years	44.5	46.3	50.1	41.1	35.8	30.9	23.9	24.0
45–54 years	150.8	164.1	172.0	175.4	149.9	123.5	105.7	102.3
55–64 years	409.4	450.9	498.1	497.4	508.2	401.9	323.1	323.5
65–74 years	798.7	887.3	997.0	1,070.7	1,090.7	984.3	723.3	707.6
75–84 years	1,367.6	1,413.7	1,592.7	1,779.7	1,883.2	1,736.0	1,421.7	1,400.4
85 years and over	1,732.7	1,791.4	1,772.2	2,375.6	2,715.1	2,693.7	2,290.7	2,279.7
Black or African American male ⁵								
All ages, age-adjusted ⁴	178.9	227.6	291.9	353.4	397.9	340.3	238.7	231.9
All ages, crude	106.6	136.7	171.6	205.5	221.9	188.5	165.6	165.1
25–34 years	18.0	18.4	18.8	14.1	15.7	10.1	10.2	8.3
35–44 years	55.7	72.9	81.3	73.8	64.3	48.4	27.3	28.8
45–54 years	211.7	244.7	311.2	333.0	302.6	214.2	132.4	128.8
55–64 years	490.8	579.7	689.2	812.5	859.2	626.4	464.0	450.5
65–74 years	636.5	938.5	1,168.9	1,417.2	1,613.9	1,363.8	941.5	925.8
75–84 years ⁶	853.5	1,053.3	1,624.8	2,029.6	2,478.3	2,351.8	1,633.2	1,569.4
85 years and over	---	1,155.2	1,387.0	2,393.9	3,238.3	3,264.8	2,465.6	2,378.6

See footnotes at end of table.

Table 24 (page 2 of 4). Death rates for malignant neoplasms, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#024>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and age	1950 ^{1,2}	1960 ^{1,2}	1970 ²	1980 ²	1990 ²	2000 ³	2013 ³	2014 ³
Deaths per 100,000 resident population								
American Indian or Alaska Native male ⁵								
All ages, age-adjusted ⁴	---	---	---	140.5	145.8	155.8	132.3	130.4
All ages, crude	---	---	---	58.1	61.4	67.0	73.6	76.1
25–34 years	---	---	---	*	*	*	6.2	*
35–44 years	---	---	---	*	22.8	21.4	13.1	15.2
45–54 years	---	---	---	86.9	86.9	70.3	61.9	58.7
55–64 years	---	---	---	213.4	246.2	255.6	233.5	221.2
65–74 years	---	---	---	613.0	530.6	648.0	477.8	520.7
75–84 years	---	---	---	936.4	1,038.4	1,152.5	1,009.1	1,015.8
85 years and over	---	---	---	1,471.2	1,654.4	1,584.2	1,488.1	1,256.3
Asian or Pacific Islander male ⁵								
All ages, age-adjusted ⁴	---	---	---	165.2	172.5	150.8	120.9	116.4
All ages, crude	---	---	---	81.9	82.7	85.2	90.5	90.2
25–34 years	---	---	---	6.3	9.2	7.4	6.5	6.0
35–44 years	---	---	---	29.4	27.7	26.1	20.3	15.9
45–54 years	---	---	---	108.2	92.6	78.5	67.6	64.6
55–64 years	---	---	---	298.5	274.6	229.2	186.4	190.7
65–74 years	---	---	---	581.2	687.2	559.4	387.5	399.6
75–84 years	---	---	---	1,147.6	1,229.9	1,086.1	899.5	839.8
85 years and over	---	---	---	1,798.7	1,837.0	1,823.2	1,575.3	1,468.0
Hispanic or Latino male ^{5,7}								
All ages, age-adjusted ⁴	---	---	---	---	174.7	171.7	138.8	135.9
All ages, crude	---	---	---	---	65.5	61.3	66.9	68.0
25–34 years	---	---	---	---	8.0	6.9	7.8	7.1
35–44 years	---	---	---	---	22.5	20.1	16.4	17.2
45–54 years	---	---	---	---	96.6	79.4	66.4	65.1
55–64 years	---	---	---	---	294.0	253.1	217.4	207.8
65–74 years	---	---	---	---	655.5	651.2	515.6	497.9
75–84 years	---	---	---	---	1,233.4	1,306.4	1,051.3	1,026.5
85 years and over	---	---	---	---	2,019.4	2,049.7	1,606.6	1,624.9
White, not Hispanic or Latino male ⁷								
All ages, age-adjusted ⁴	---	---	---	---	276.7	247.7	200.0	197.7
All ages, crude	---	---	---	---	246.2	244.4	247.3	249.3
25–34 years	---	---	---	---	12.8	9.7	8.5	8.6
35–44 years	---	---	---	---	36.8	32.3	25.6	25.6
45–54 years	---	---	---	---	153.9	127.2	111.9	108.5
55–64 years	---	---	---	---	520.6	412.0	332.9	335.0
65–74 years	---	---	---	---	1,109.0	1,002.1	737.2	722.1
75–84 years	---	---	---	---	1,906.6	1,750.2	1,444.8	1,424.1
85 years and over	---	---	---	---	2,744.4	2,714.1	2,330.5	2,319.1
White female ⁵								
All ages, age-adjusted ⁴	182.0	167.7	162.5	165.2	174.0	166.9	140.2	138.8
All ages, crude	139.9	139.8	149.4	170.3	196.1	199.4	186.8	187.2
25–34 years	20.9	18.8	16.3	13.5	11.9	10.1	8.5	8.3
35–44 years	74.5	66.6	62.4	50.9	46.2	38.2	31.8	31.2
45–54 years	185.8	175.7	177.3	166.4	150.9	120.1	103.0	101.8
55–64 years	362.5	329.0	338.6	355.5	368.5	319.7	245.4	244.9
65–74 years	616.5	562.1	554.7	605.2	675.1	665.6	526.4	512.4
75–84 years	1,026.6	939.3	903.5	905.4	1,011.8	1,063.4	945.4	943.2
85 years and over	1,348.3	1,304.9	1,126.6	1,266.8	1,372.3	1,459.1	1,326.5	1,329.0

See footnotes at end of table.

Table 24 (page 3 of 4). Death rates for malignant neoplasms, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#024>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and age	1950 ^{1,2}	1960 ^{1,2}	1970 ²	1980 ²	1990 ²	2000 ³	2013 ³	2014 ³
Deaths per 100,000 resident population								
Black or African American female⁵								
All ages, age-adjusted ⁴	174.1	174.3	173.4	189.5	205.9	193.8	158.5	156.8
All ages, crude	111.8	113.8	117.3	136.5	156.1	151.8	146.2	147.5
25–34 years	34.3	31.0	20.9	18.3	18.7	13.5	10.4	11.2
35–44 years	119.8	102.4	94.6	73.5	67.4	58.9	40.7	40.5
45–54 years	277.0	254.8	228.6	230.2	209.9	173.9	134.8	133.4
55–64 years	484.6	442.7	404.8	450.4	482.4	391.0	320.0	321.9
65–74 years	477.3	541.6	615.8	662.4	773.2	753.1	593.4	583.7
75–84 years ⁶	605.3	696.3	763.3	923.9	1,059.9	1,124.0	984.3	960.8
85 years and over	---	728.9	791.5	1,159.9	1,431.3	1,527.7	1,306.5	1,297.5
American Indian or Alaska Native female⁵								
All ages, age-adjusted ⁴	---	---	---	94.0	106.9	108.3	94.3	88.5
All ages, crude	---	---	---	50.4	62.1	61.3	65.8	63.4
25–34 years	---	---	---	*	*	*	6.4	*
35–44 years	---	---	---	36.9	31.0	23.7	18.8	17.5
45–54 years	---	---	---	96.9	104.5	59.7	69.1	67.1
55–64 years	---	---	---	198.4	213.3	200.9	155.1	139.5
65–74 years	---	---	---	350.8	438.9	458.3	373.0	359.8
75–84 years	---	---	---	446.4	554.3	714.0	656.2	611.5
85 years and over	---	---	---	786.5	843.7	983.2	844.8	817.9
Asian or Pacific Islander female⁵								
All ages, age-adjusted ⁴	---	---	---	93.0	103.0	100.7	86.0	86.2
All ages, crude	---	---	---	54.1	60.5	72.1	78.6	80.6
25–34 years	---	---	---	9.5	7.3	8.1	6.4	6.2
35–44 years	---	---	---	38.7	29.8	28.9	22.1	21.7
45–54 years	---	---	---	99.8	93.9	78.2	63.6	64.1
55–64 years	---	---	---	174.7	196.2	176.5	139.7	143.8
65–74 years	---	---	---	301.9	346.2	357.4	280.7	278.8
75–84 years	---	---	---	522.1	641.4	650.1	601.5	597.1
85 years and over	---	---	---	800.0	971.7	988.5	925.7	963.7
Hispanic or Latina female^{5,7}								
All ages, age-adjusted ⁴	---	---	---	---	111.9	110.8	97.3	95.7
All ages, crude	---	---	---	---	60.7	58.5	63.0	63.6
25–34 years	---	---	---	---	9.7	7.8	8.8	7.2
35–44 years	---	---	---	---	34.8	30.7	25.4	25.4
45–54 years	---	---	---	---	100.5	84.7	69.1	68.6
55–64 years	---	---	---	---	205.4	192.5	164.7	163.2
65–74 years	---	---	---	---	404.8	410.0	342.0	335.9
75–84 years	---	---	---	---	663.0	716.5	656.7	636.6
85 years and over	---	---	---	---	1,022.7	1,056.5	996.6	1,004.3
White, not Hispanic or Latina female⁷								
All ages, age-adjusted ⁴	---	---	---	---	177.5	170.0	143.9	142.7
All ages, crude	---	---	---	---	210.6	220.6	214.0	215.1
25–34 years	---	---	---	---	11.9	10.5	8.1	8.4
35–44 years	---	---	---	---	47.0	38.9	32.9	32.3
45–54 years	---	---	---	---	154.9	123.0	108.0	107.1
55–64 years	---	---	---	---	379.5	328.9	253.3	253.3
65–74 years	---	---	---	---	688.5	681.0	541.4	527.0
75–84 years	---	---	---	---	1,027.2	1,075.3	966.6	967.1
85 years and over	---	---	---	---	1,385.7	1,468.7	1,343.5	1,345.8

See footnotes at end of table.

Table 24 (page 4 of 4). Death rates for malignant neoplasms, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#024>.

[Data are based on death certificates]

-- Data not available.

* Rates based on fewer than 20 deaths are considered unreliable and are not shown.

¹Includes deaths of persons who were not residents of the 50 states and the District of Columbia (D.C.).

²Underlying cause of death was coded according to the 6th Revision of the *International Classification of Diseases* (ICD) in 1950, 7th Revision in 1960, 8th Revision in 1970, and 9th Revision in 1980–1998. See Appendix II, Cause of death; Tables III; and IV.

³Starting with 1999 data, cause of death is coded according to ICD–10. See Appendix II, Cause of death; Comparability ratio; Table IV; Table V.

⁴Age-adjusted rates are calculated using the year 2000 standard population. Prior to 2001, age-adjusted rates were calculated using standard million proportions based on rounded population numbers. Starting with 2001 data, unrounded population numbers are used to calculate age-adjusted rates. See Appendix II, Age adjustment.

⁵The race groups, white, black, Asian or Pacific Islander, and American Indian or Alaska Native, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Death rates for Hispanic, American Indian or Alaska Native, and Asian or Pacific Islander persons should be interpreted with caution because of inconsistencies in reporting Hispanic origin or race on the death certificate (death rate numerators) compared with population figures (death rate denominators). The net effect of misclassification is an underestimation of deaths and death rates for races other than white and black. See Appendix II, Race, for a detailed discussion of sources of bias in death rates by race and Hispanic origin.

⁶In 1950, rate is for the age group 75 years and over.

⁷Prior to 1997, data from states that did not report Hispanic origin on the death certificate were excluded. See Appendix II, Hispanic origin.

NOTES: Starting with *Health, United States, 2003*, rates for 1991–1999 were revised using intercensal population estimates based on the 1990 and 2000 censuses. For 2000, population estimates are bridged-race April 1 census counts. Starting with *Health, United States, 2012*, rates for 2001–2009 were revised using intercensal population estimates based on the 2000 and 2010 censuses. For 2010, population estimates are bridged-race April 1 census counts. Rates for 2011 and beyond were computed using 2010-based postcensal estimates. See Appendix I, Population Census and Population Estimates. Age groups were selected to minimize the presentation of unstable age-specific death rates based on small numbers of deaths and for consistency among comparison groups. Starting with 2003 data, some states allowed the reporting of more than one race on the death certificate. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards, for comparability with other states. See Appendix II, Race. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Vital Statistics System; GROVE RD, Hetzel AM. Vital statistics rates in the United States, 1940–1960. Washington, DC: U.S. Government Printing Office, 1968; numerator data from National Vital Statistics System, annual public-use Mortality Files; denominator data from national population estimates for race groups from Table 1 and unpublished Hispanic population estimates for 1985–1996 prepared by the Housing and Household Economic Statistics Division, U.S. Census Bureau; Deaths: Final data for 2014. National vital statistics reports (forthcoming). Available from: <http://www.cdc.gov/nchs/products/nvsr.htm>. See Appendix I, National Vital Statistics System (NVSS).

Table 25 (page 1 of 3). Death rates for malignant neoplasms of trachea, bronchus, and lung, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#025>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and age	1950 ^{1,2}	1960 ^{1,2}	1970 ²	1980 ²	1990 ²	2000 ³	2013 ³	2014 ³
Deaths per 100,000 resident population								
All persons								
All ages, age-adjusted ⁴	15.0	24.1	37.1	49.9	59.3	56.1	43.4	42.1
All ages, crude	12.2	20.3	32.1	45.8	56.8	55.3	49.4	48.8
Under 25 years	0.1	0.0	0.1	0.0	0.0	0.0	0.0	0.0
25–34 years	0.8	1.0	0.9	0.6	0.7	0.5	0.3	0.3
35–44 years	4.5	6.8	11.0	9.2	6.8	6.1	2.8	2.7
45–54 years	20.4	29.6	43.4	54.1	46.8	31.6	23.9	22.4
55–64 years	48.7	75.3	109.1	138.2	160.6	122.4	79.9	78.5
65–74 years	59.7	108.1	164.5	233.3	288.4	284.2	196.6	189.4
75–84 years	55.8	91.5	163.2	240.5	333.3	370.8	328.5	321.2
85 years and over	42.3	65.6	101.7	176.0	242.5	302.1	320.0	311.7
Male								
All ages, age-adjusted ⁴	24.6	43.6	67.5	85.2	91.1	76.7	53.7	51.7
All ages, crude	19.9	35.4	53.4	68.6	75.1	65.5	55.1	54.1
Under 25 years	0.0	0.0	0.1	0.1	0.0	*	*	*
25–34 years	1.1	1.4	1.3	0.8	0.9	0.5	0.3	0.3
35–44 years	7.1	10.5	16.1	11.9	8.5	6.9	2.8	2.9
45–54 years	35.0	50.6	67.5	76.0	59.7	38.5	25.9	23.8
55–64 years	83.8	139.3	189.7	213.6	222.9	154.0	96.9	93.8
65–74 years	98.7	204.3	320.8	403.9	430.4	377.9	236.0	229.5
75–84 years	82.6	167.1	330.8	488.8	572.9	532.2	413.4	395.7
85 years and over	62.5	107.7	194.0	368.1	513.2	521.2	463.0	442.8
Female								
All ages, age-adjusted ⁴	5.8	7.5	13.1	24.4	37.1	41.3	35.5	34.7
All ages, crude	4.5	6.4	11.9	24.3	39.4	45.4	44.0	43.7
Under 25 years	0.1	0.0	0.0	*	*	*	*	*
25–34 years	0.5	0.5	0.5	0.5	0.5	0.5	0.3	0.3
35–44 years	1.9	3.2	6.1	6.5	5.2	5.3	2.9	2.6
45–54 years	5.8	9.2	21.0	33.7	34.5	25.0	21.9	21.0
55–64 years	13.6	15.4	36.8	72.0	105.0	93.3	64.2	64.2
65–74 years	23.3	24.4	43.1	102.7	177.6	206.9	161.9	154.2
75–84 years	32.9	32.8	52.4	94.1	190.1	265.6	264.9	264.8
85 years and over	28.2	38.8	50.0	91.9	138.1	212.8	246.9	243.5
White male ⁵								
All ages, age-adjusted ⁴	25.1	43.6	67.1	83.8	89.0	75.7	53.7	51.8
All ages, crude	20.8	36.4	54.6	70.2	77.8	69.4	59.5	58.7
45–54 years	35.1	49.2	63.3	70.9	55.2	35.7	25.6	23.7
55–64 years	85.4	139.2	186.8	205.6	213.7	150.8	94.6	92.1
65–74 years	101.5	207.5	325.0	401.0	422.1	374.9	236.5	230.2
75–84 years	85.5	170.4	336.7	493.5	572.2	529.9	416.8	400.4
85 years and over	67.4	109.4	199.6	374.1	516.3	522.4	466.8	446.6
Black or African American male ⁵								
All ages, age-adjusted ⁴	17.8	42.6	75.4	107.6	125.4	101.1	65.7	61.9
All ages, crude	12.1	28.1	47.7	66.6	73.7	58.3	46.8	45.0
45–54 years	34.4	68.4	115.4	133.8	114.9	70.7	35.2	31.0
55–64 years	68.3	146.8	234.3	321.1	358.6	223.5	141.0	131.4
65–74 years	53.8	168.3	300.5	472.3	585.4	488.8	302.6	287.7
75–84 years ⁶	36.2	107.3	271.6	472.9	645.4	642.5	465.7	435.1
85 years and over	---	82.8	137.0	311.3	499.5	562.8	458.5	456.0
American Indian or Alaska Native male ⁵								
All ages, age-adjusted ⁴	---	---	---	31.7	47.5	42.9	34.4	34.1
All ages, crude	---	---	---	14.2	20.0	18.1	18.2	19.8
45–54 years	---	---	---	*	26.6	14.5	10.7	8.8
55–64 years	---	---	---	72.0	97.8	86.0	55.8	53.2
65–74 years	---	---	---	202.8	194.3	184.8	131.3	177.8
75–84 years	---	---	---	*	356.2	367.9	299.3	292.1
85 years and over	---	---	---	*	*	*	369.4	193.3

See footnotes at end of table.

Table 25 (page 2 of 3). Death rates for malignant neoplasms of trachea, bronchus, and lung, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#025>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and age	1950 ^{1,2}	1960 ^{1,2}	1970 ²	1980 ²	1990 ²	2000 ³	2013 ³	2014 ³
Deaths per 100,000 resident population								
Asian or Pacific Islander male ⁵								
All ages, age-adjusted ⁴	---	---	---	43.3	44.2	40.9	30.6	29.6
All ages, crude	---	---	---	22.1	20.7	22.7	22.3	22.4
45–54 years	---	---	---	33.3	18.8	17.2	12.7	13.1
55–64 years	---	---	---	94.4	74.4	61.4	42.5	44.7
65–74 years	---	---	---	174.3	215.8	183.2	110.3	108.9
75–84 years	---	---	---	301.3	307.5	323.2	257.5	239.8
85 years and over	---	---	---	*	421.3	378.0	387.3	363.2
Hispanic or Latino male ^{5,7}								
All ages, age-adjusted ⁴	---	---	---	---	44.1	39.0	26.2	25.0
All ages, crude	---	---	---	---	16.2	13.3	11.8	11.6
45–54 years	---	---	---	---	21.5	14.8	8.0	7.2
55–64 years	---	---	---	---	80.7	58.6	35.6	31.4
65–74 years	---	---	---	---	195.5	167.3	110.8	107.0
75–84 years	---	---	---	---	313.4	327.5	225.4	216.1
85 years and over	---	---	---	---	420.7	368.8	282.7	276.0
White, not Hispanic or Latino male ⁷								
All ages, age-adjusted ⁴	---	---	---	---	91.1	77.9	56.3	54.4
All ages, crude	---	---	---	---	84.7	78.9	71.0	70.2
45–54 years	---	---	---	---	57.8	37.7	28.7	26.9
55–64 years	---	---	---	---	221.0	157.7	101.0	99.0
65–74 years	---	---	---	---	431.4	387.3	246.6	240.3
75–84 years	---	---	---	---	580.4	537.7	430.8	414.3
85 years and over	---	---	---	---	520.9	527.3	478.0	457.5
White female ⁵								
All ages, age-adjusted ⁴	5.9	6.8	13.1	24.5	37.6	42.3	36.7	36.0
All ages, crude	4.7	5.9	12.3	25.6	42.4	49.9	48.9	48.7
45–54 years	5.7	9.0	20.9	33.0	34.6	24.8	22.6	21.7
55–64 years	13.7	15.1	37.2	71.9	105.7	96.1	66.0	66.3
65–74 years	23.7	24.8	42.9	104.6	181.3	213.2	169.0	161.4
75–84 years	34.0	32.7	52.6	95.2	194.6	272.7	274.1	275.3
85 years and over	29.3	39.1	50.6	92.4	138.3	215.9	250.9	247.7
Black or African American female ⁵								
All ages, age-adjusted ⁴	4.5	6.8	13.7	24.8	36.8	39.8	34.1	32.7
All ages, crude	2.8	4.3	9.4	18.3	28.1	30.8	31.3	30.7
45–54 years	7.5	11.3	23.9	43.4	41.3	32.9	24.0	23.2
55–64 years	12.9	17.9	33.5	79.9	117.9	95.3	71.9	69.8
65–74 years	14.0	18.1	46.1	88.0	164.3	194.1	151.0	142.3
75–84 years ⁶	*	31.3	49.1	79.4	148.1	224.3	235.8	227.5
85 years and over	---	34.2	44.8	85.8	134.9	185.9	224.9	220.8
American Indian or Alaska Native female ⁵								
All ages, age-adjusted ⁴	---	---	---	11.7	19.3	24.8	22.9	22.7
All ages, crude	---	---	---	6.0	11.2	14.0	15.6	15.8
45–54 years	---	---	---	*	22.9	12.1	12.3	14.4
55–64 years	---	---	---	*	53.7	52.6	31.1	28.0
65–74 years	---	---	---	*	78.5	151.5	124.9	114.4
75–84 years	---	---	---	*	111.8	136.3	174.6	179.3
85 years and over	---	---	---	*	*	*	137.0	161.4

See footnotes at end of table.

Table 25 (page 3 of 3). Death rates for malignant neoplasms of trachea, bronchus, and lung, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#025>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and age	1950 ^{1,2}	1960 ^{1,2}	1970 ²	1980 ²	1990 ²	2000 ³	2013 ³	2014 ³
Deaths per 100,000 resident population								
Asian or Pacific Islander female ⁵								
All ages, age-adjusted ⁴	---	---	---	15.4	18.9	18.4	18.0	17.6
All ages, crude	---	---	---	8.4	10.5	12.6	16.0	16.0
45–54 years	---	---	---	13.5	11.3	9.9	9.3	9.0
55–64 years	---	---	---	24.6	38.3	30.4	24.3	25.2
65–74 years	---	---	---	62.4	71.6	77.0	63.9	59.3
75–84 years	---	---	---	117.7	137.9	135.0	150.2	149.3
85 years and over	---	---	---	*	172.9	175.3	207.4	199.6
Hispanic or Latina female ^{5,7}								
All ages, age-adjusted ⁴	---	---	---	---	14.1	14.7	13.2	13.3
All ages, crude	---	---	---	---	7.2	7.2	8.0	8.3
45–54 years	---	---	---	---	8.7	7.1	5.4	5.1
55–64 years	---	---	---	---	25.1	22.2	18.9	19.3
65–74 years	---	---	---	---	66.8	66.0	52.9	53.0
75–84 years	---	---	---	---	94.3	112.3	107.2	109.7
85 years and over	---	---	---	---	118.2	137.5	146.7	151.9
White, not Hispanic or Latina female ⁷								
All ages, age-adjusted ⁴	---	---	---	---	39.0	44.1	39.0	38.4
All ages, crude	---	---	---	---	46.2	56.4	58.2	58.1
45–54 years	---	---	---	---	36.6	26.4	25.6	24.8
55–64 years	---	---	---	---	111.3	102.2	71.4	72.1
65–74 years	---	---	---	---	186.4	222.9	179.5	171.6
75–84 years	---	---	---	---	199.1	279.2	288.1	289.7
85 years and over	---	---	---	---	139.0	218.0	256.8	253.1

0.0 Quantity more than zero but less than 0.05.

* Rates based on fewer than 20 deaths are considered unreliable and are not shown.

--- Data not available.

¹Includes deaths of persons who were not residents of the 50 states and the District of Columbia (D.C.).

²Underlying cause of death was coded according to the 6th Revision of the *International Classification of Diseases* (ICD) in 1950, 7th Revision in 1960, 8th Revision in 1970, and 9th Revision in 1980–1998. See Appendix II, Cause of death; Table III; Table IV.

³Starting with 1999 data, cause of death is coded according to ICD–10. See Appendix II, Cause of death; Comparability ratio; Table IV; Table V.

⁴Age-adjusted rates are calculated using the year 2000 standard population. Prior to 2001, age-adjusted rates were calculated using standard million proportions based on rounded population numbers. Starting with 2001 data, unrounded population numbers are used to calculate age-adjusted rates. See Appendix II, Age adjustment.

⁵The race groups, white, black, Asian or Pacific Islander, and American Indian or Alaska Native, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Death rates for Hispanic, American Indian or Alaska Native, and Asian or Pacific Islander persons should be interpreted with caution because of inconsistencies in reporting Hispanic origin or race on the death certificate (death rate numerators) compared with population figures (death rate denominators). The net effect of misclassification is an underestimation of deaths and death rates for races other than white and black. See Appendix II, Race, for a detailed discussion of sources of bias in death rates by race and Hispanic origin.

⁶In 1950, rate is for the age group 75 years and over.

⁷Prior to 1997, data from states that did not report Hispanic origin on the death certificate were excluded. See Appendix II, Hispanic origin.

NOTES: Starting with *Health, United States, 2003*, rates for 1991–1999 were revised using intercensal population estimates based on the 1990 and 2000 censuses. For 2000, population estimates are bridged-race April 1 census counts. Starting with *Health, United States, 2012*, rates for 2001–2009 were revised using intercensal population estimates based on the 2000 and 2010 censuses. For 2010, population estimates are bridged-race April 1 census counts. Rates for 2011 and beyond were computed using 2010-based postcensal estimates. See Appendix I, Population Census and Population Estimates. Age groups were selected to minimize the presentation of unstable age-specific death rates based on small numbers of deaths and for consistency among comparison groups. Starting with 2003 data, some states allowed the reporting of more than one race on the death certificate. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards, for comparability with other states. See Appendix II, Race. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Vital Statistics System; Grove RD, Hetzel AM. Vital statistics rates in the United States, 1940–1960. Washington, DC: U.S. Government Printing Office. 1968; numerator data from National Vital Statistics System, annual public-use Mortality Files; denominator data from national population estimates for race groups from Table 1 and unpublished Hispanic population estimates for 1985–1996 prepared by the Housing and Household Economic Statistics Division, U.S. Census Bureau; Deaths: Final data for 2014. National vital statistics reports (forthcoming). Available from: <http://www.cdc.gov/nchs/products/nvsr.htm>. See Appendix I, National Vital Statistics System (NVSS).

Table 26 (page 1 of 2). Death rates for malignant neoplasm of breast among females, by race, Hispanic origin, and age: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#026>.

[Data are based on death certificates]

Race, Hispanic origin, and age	1950 ^{1,2}	1960 ^{1,2}	1970 ²	1980 ²	1990 ²	2000 ³	2010 ³	2013 ³	2014 ³
Deaths per 100,000 resident population									
All females									
All ages, age-adjusted ⁴	31.9	31.7	32.1	31.9	33.3	26.8	22.1	20.8	20.6
All ages, crude	24.7	26.1	28.4	30.6	34.0	29.2	26.1	25.5	25.5
Under 25 years	*	*	*	*	*	*	*	*	*
25–34 years	3.8	3.8	3.9	3.3	2.9	2.3	1.6	1.6	1.6
35–44 years	20.8	20.2	20.4	17.9	17.8	12.4	9.8	9.6	9.7
45–54 years	46.9	51.4	52.6	48.1	45.4	33.0	25.7	24.6	23.8
55–64 years	69.9	70.8	77.6	80.5	78.6	59.3	47.7	44.1	43.3
65–74 years	95.0	90.0	93.8	101.1	111.7	88.3	73.9	68.4	67.5
75–84 years	139.8	129.9	127.4	126.4	146.3	128.9	109.1	104.4	104.2
85 years and over	195.5	191.9	157.1	169.3	196.8	205.7	185.8	173.0	174.0
White ⁵									
All ages, age-adjusted ⁴	32.4	32.0	32.5	32.1	33.2	26.3	21.5	20.4	20.1
All ages, crude	25.7	27.2	29.9	32.3	35.9	30.7	27.3	26.7	26.6
35–44 years	20.8	19.7	20.2	17.3	17.1	11.3	8.8	8.8	8.9
45–54 years	47.1	51.2	53.0	48.1	44.3	31.2	23.9	23.0	22.0
55–64 years	70.9	71.8	79.3	81.3	78.5	57.9	45.9	42.5	41.4
65–74 years	96.3	91.6	95.9	103.7	113.3	89.3	73.3	68.4	67.4
75–84 years	143.6	132.8	129.6	128.4	148.2	130.2	110.2	105.0	105.2
85 years and over	204.2	199.7	161.9	171.7	198.0	205.5	186.8	174.9	175.9
Black or African American ⁵									
All ages, age-adjusted ⁴	25.3	27.9	28.9	31.7	38.1	34.5	30.3	28.3	28.1
All ages, crude	16.4	18.7	19.7	22.9	29.0	27.9	27.5	26.7	27.0
35–44 years	21.0	24.8	24.4	24.1	25.8	20.9	18.3	16.3	16.0
45–54 years	46.5	54.4	52.0	52.7	60.5	51.5	40.9	39.2	39.1
55–64 years	64.3	63.2	64.7	79.9	93.1	80.9	70.5	63.7	64.4
65–74 years	67.0	72.3	77.3	84.3	112.2	98.6	97.4	86.6	85.9
75–84 years ⁶	81.0	87.5	101.8	114.1	140.5	139.8	123.2	126.1	122.6
85 years and over	---	92.1	112.1	149.9	201.5	238.7	214.6	193.1	196.4
American Indian or Alaska Native ⁵									
All ages, age-adjusted ⁴	---	---	---	10.8	13.7	13.6	11.5	10.1	10.8
All ages, crude	---	---	---	6.1	8.6	8.7	8.0	7.5	8.0
35–44 years	---	---	---	*	*	*	*	*	*
45–54 years	---	---	---	*	23.9	14.4	13.2	11.2	9.4
55–64 years	---	---	---	*	*	40.0	25.2	20.6	19.3
65–74 years	---	---	---	*	*	42.5	34.3	38.2	40.2
75–84 years	---	---	---	*	*	71.8	61.1	50.5	61.7
85 years and over	---	---	---	*	*	*	*	*	*
Asian or Pacific Islander ⁵									
All ages, age-adjusted ⁴	---	---	---	11.9	13.7	12.3	11.9	11.1	11.6
All ages, crude	---	---	---	8.2	9.3	10.2	10.8	10.8	11.5
35–44 years	---	---	---	10.4	8.4	8.1	5.4	5.7	6.4
45–54 years	---	---	---	23.4	26.4	22.3	17.0	15.6	16.8
55–64 years	---	---	---	35.7	33.8	31.3	28.4	27.2	27.6
65–74 years	---	---	---	*	38.5	34.7	37.9	32.7	35.1
75–84 years	---	---	---	*	48.0	37.5	53.2	48.6	47.7
85 years and over	---	---	---	*	*	68.2	77.5	82.8	81.3
Hispanic or Latina ^{5,7}									
All ages, age-adjusted ⁴	---	---	---	---	19.5	16.9	14.4	14.6	14.5
All ages, crude	---	---	---	---	11.5	9.7	9.2	10.0	10.3
35–44 years	---	---	---	---	11.7	8.7	6.2	7.2	7.7
45–54 years	---	---	---	---	32.8	23.9	18.6	18.0	17.4
55–64 years	---	---	---	---	45.8	39.1	32.7	31.2	31.5
65–74 years	---	---	---	---	64.8	54.9	49.0	46.3	47.0
75–84 years	---	---	---	---	67.2	74.9	61.8	73.3	69.3
85 years and over	---	---	---	---	102.8	105.8	117.8	112.0	116.3

See footnotes at end of table.

Table 26 (page 2 of 2). Death rates for malignant neoplasm of breast among females, by race, Hispanic origin, and age: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#026>.

[Data are based on death certificates]

Race, Hispanic origin, and age	1950 ^{1,2}	1960 ^{1,2}	1970 ²	1980 ²	1990 ²	2000 ³	2010 ³	2013 ³	2014 ³
White, not Hispanic or Latina ⁷	Deaths per 100,000 resident population								
All ages, age-adjusted ⁴	---	---	---	---	33.9	26.8	22.1	20.9	20.6
All ages, crude	---	---	---	---	38.5	33.8	31.0	30.3	30.2
35–44 years	---	---	---	---	17.5	11.6	9.3	9.1	9.1
45–54 years	---	---	---	---	45.2	31.7	24.5	23.6	22.6
55–64 years	---	---	---	---	80.6	59.2	47.1	43.5	42.2
65–74 years	---	---	---	---	115.7	91.4	75.1	70.1	69.1
75–84 years	---	---	---	---	151.4	132.2	113.6	107.3	108.1
85 years and over	---	---	---	---	201.5	208.3	189.9	178.5	179.4

* Rates based on fewer than 20 deaths are considered unreliable and are not shown.

--- Data not available.

¹Includes deaths of persons who were not residents of the 50 states and the District of Columbia (D.C.).

²Underlying cause of death was coded according to the 6th Revision of the *International Classification of Diseases* (ICD) in 1950, 7th Revision in 1960, 8th Revision in 1970, and 9th Revision in 1980–1998. See Appendix II, Cause of death; Table III; Table IV.

³Starting with 1999 data, cause of death is coded according to ICD–10. See Appendix II, Cause of death; Comparability ratio; Table IV; Table V.

⁴Age-adjusted rates are calculated using the year 2000 standard population. Prior to 2001, age-adjusted rates were calculated using standard million proportions based on rounded population numbers. Starting with 2001 data, unrounded population numbers are used to calculate age-adjusted rates. See Appendix II, Age adjustment.

⁵The race groups, white, black, Asian or Pacific Islander, and American Indian or Alaska Native, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Death rates for Hispanic, American Indian or Alaska Native, and Asian or Pacific Islander persons should be interpreted with caution because of inconsistencies in reporting Hispanic origin or race on the death certificate (death rate numerators) compared with population figures (death rate denominators). The net effect of misclassification is an underestimation of deaths and death rates for races other than white and black. See Appendix II, Race, for a detailed discussion of sources of bias in death rates by race and Hispanic origin.

⁶In 1950, rate is for the age group 75 years and over.

⁷Prior to 1997, data from states that did not report Hispanic origin on the death certificate were excluded. See Appendix II, Hispanic origin.

NOTES: Starting with *Health, United States, 2003*, rates for 1991–1999 were revised using intercensal population estimates based on the 1990 and 2000 censuses. For 2000, population estimates are bridged-race April 1 census counts. Starting with *Health, United States, 2012*, rates for 2001–2009 were revised using intercensal population estimates based on the 2000 and 2010 censuses. For 2010, population estimates are bridged-race April 1 census counts. Rates for 2011 and beyond were computed using 2010-based postcensal estimates. See Appendix I, Population Census and Population Estimates. Age groups were selected to minimize the presentation of unstable age-specific death rates based on small numbers of deaths and for consistency among comparison groups. Starting with 2003 data, some states allowed the reporting of more than one race on the death certificate. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards, for comparability with other states. See Appendix II, Race. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Vital Statistics System; numerator data from National Vital Statistics System, annual public-use Mortality Files; denominator data from national population estimates for race groups from Table 1 and unpublished Hispanic population estimates for 1985–1996 prepared by the Housing and Household Economic Statistics Division, U.S. Census Bureau; Deaths: Final data for 2014. National vital statistics reports (forthcoming). Available from: <http://www.cdc.gov/nchs/products/nvsr.htm>. See Appendix I, National Vital Statistics System (NVSS).

Table 27 (page 1 of 4). Death rates for drug poisoning and drug poisoning involving opioid analgesics and heroin, by sex, age, race, and Hispanic origin: United States, selected years 1999–2014

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2015.htm#027>.

[Data are based on death certificates]

Sex, age, race, and Hispanic origin	1999	2000	2004	2005	2010	2011	2012	2013	2014
Drug poisoning deaths per 100,000 resident population ¹									
All persons									
All ages, age-adjusted ²	6.1	6.2	9.4	10.1	12.3	13.2	13.1	13.8	14.7
All ages, crude	6.0	6.2	9.4	10.1	12.4	13.3	13.2	13.9	14.8
Under 15 years	0.1	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.2
15–24 years	3.2	3.7	6.6	6.9	8.2	8.6	8.0	8.3	8.6
25–34 years	8.1	7.9	11.9	13.6	18.4	20.2	20.1	20.9	23.1
35–44 years	14.0	14.3	19.3	19.6	20.8	22.5	22.1	23.0	25.0
45–54 years	11.1	11.6	19.3	21.1	25.1	26.7	26.9	27.5	28.2
55–64 years	4.2	4.2	7.8	9.0	15.0	15.9	16.6	19.2	20.3
65–74 years	2.4	2.0	2.9	3.2	4.7	5.4	5.8	6.4	6.9
75–84 years	2.8	2.4	2.9	3.1	3.4	3.4	3.4	3.6	3.6
85 years and over	3.8	4.4	4.0	4.1	4.7	4.2	4.3	4.3	4.1
Male									
All ages, age-adjusted ²	8.2	8.3	11.8	12.8	15.0	16.1	16.1	17.0	18.3
All ages, crude	8.2	8.4	11.9	12.9	15.2	16.3	16.3	17.2	18.4
Under 15 years	0.1	0.2	0.2	0.2	0.3	0.2	0.2	0.2	0.2
15–24 years	4.5	5.3	9.6	10.0	11.6	12.4	11.4	11.7	12.1
25–34 years	11.5	11.3	16.6	18.7	25.0	27.5	27.0	28.6	31.9
35–44 years	19.2	19.5	23.8	24.4	24.9	26.8	27.1	28.1	30.8
45–54 years	15.2	15.7	23.8	25.8	28.5	30.4	30.4	31.5	32.9
55–64 years	4.9	4.4	8.6	10.6	17.3	18.5	19.4	22.7	23.5
65–74 years	2.7	2.1	2.9	3.3	4.5	5.4	6.2	6.9	7.3
75–84 years	2.5	2.5	2.8	3.4	3.6	3.4	3.2	3.7	3.8
85 years and over	4.4	5.9	4.8	5.2	5.1	4.3	5.3	5.9	4.3
Female									
All ages, age-adjusted ²	3.9	4.1	6.9	7.3	9.6	10.2	10.2	10.6	11.1
All ages, crude	3.9	4.1	6.9	7.4	9.8	10.3	10.3	10.7	11.3
Under 15 years	0.1	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.2
15–24 years	1.8	1.9	3.3	3.5	4.6	4.6	4.4	4.8	5.0
25–34 years	4.6	4.6	7.2	8.5	11.9	12.8	13.1	13.0	14.1
35–44 years	8.7	9.2	14.8	14.8	16.8	18.2	17.1	18.0	19.2
45–54 years	7.2	7.7	15.0	16.5	21.8	23.1	23.4	23.6	23.7
55–64 years	3.5	3.9	7.0	7.5	12.9	13.5	14.0	15.9	17.2
65–74 years	2.1	2.0	3.0	3.1	4.8	5.3	5.5	5.9	6.5
75–84 years	3.0	2.3	2.9	2.9	3.3	3.4	3.5	3.4	3.5
85 years and over	3.5	3.9	3.7	3.7	4.5	4.2	3.8	3.5	4.0
All ages, age-adjusted ^{2,3}									
Male:									
White	8.1	8.4	12.6	13.6	16.8	18.1	18.1	19.0	20.4
Black or African American	11.5	10.8	11.1	12.8	10.1	11.0	11.3	12.9	13.8
American Indian or Alaska Native	5.7	6.1	11.2	10.8	11.8	12.9	12.8	12.9	15.9
Asian or Pacific Islander	1.5	1.4	2.1	2.2	2.5	3.2	3.1	3.2	3.3
Hispanic or Latino	8.6	7.1	7.5	8.4	7.6	8.1	8.5	9.2	9.3
White, not Hispanic or Latino	8.0	8.6	13.7	14.7	19.0	20.5	20.4	21.4	23.2
Female:									
White	4.0	4.3	7.5	8.0	10.9	11.7	11.6	12.1	12.7
Black or African American	3.9	4.1	5.5	6.0	5.7	5.9	6.0	6.3	7.0
American Indian or Alaska Native	4.6	3.7	7.9	8.6	9.7	10.7	12.2	11.6	11.2
Asian or Pacific Islander	1.0	0.8	1.1	1.3	1.5	1.6	1.4	1.5	1.7
Hispanic or Latina	2.2	2.0	2.9	3.0	3.6	4.0	4.0	4.1	4.1
White, not Hispanic or Latina	4.3	4.5	8.3	8.8	12.5	13.3	13.2	13.8	14.6

See footnotes at end of table.

Table 27 (page 2 of 4). Death rates for drug poisoning and drug poisoning involving opioid analgesics and heroin, by sex, age, race, and Hispanic origin: United States, selected years 1999–2014

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2015.htm#027>.

[Data are based on death certificates]

Sex, age, race, and Hispanic origin	1999	2000	2004	2005	2010	2011	2012	2013	2014
Drug poisoning deaths involving opioid analgesics per 100,000 resident population ⁴									
All persons									
All ages, age-adjusted ²	1.4	1.5	3.4	3.7	5.4	5.4	5.1	5.1	5.9
All ages, crude	1.4	1.6	3.4	3.7	5.4	5.4	5.1	5.1	5.9
Under 15 years	*	0.0	0.1	0.1	0.1	0.1	0.1	0.1	0.1
15–24 years	0.7	0.8	2.7	2.7	3.9	3.6	2.8	2.6	3.1
25–34 years	1.9	1.9	4.4	5.3	8.5	8.5	7.7	7.5	9.0
35–44 years	3.5	3.7	6.8	6.9	9.1	9.3	8.8	8.6	10.3
45–54 years	2.9	3.2	7.1	7.9	10.9	11.2	10.6	10.6	11.7
55–64 years	1.0	1.1	2.6	3.1	6.2	6.3	6.6	7.5	8.5
65–74 years	0.4	0.4	0.8	1.0	1.5	1.8	2.0	2.3	2.7
75–84 years	0.3	0.2	0.5	0.6	0.7	0.7	0.9	0.8	0.9
85 years and over	*	*	0.5	0.9	1.1	0.8	0.8	0.9	0.9
Male									
All ages, age-adjusted ²	2.0	2.0	4.2	4.6	6.5	6.5	6.0	5.9	6.9
All ages, crude	2.0	2.1	4.2	4.6	6.6	6.5	6.0	5.9	7.0
Under 15 years	*	*	0.1	0.1	0.2	0.1	0.1	0.1	0.1
15–24 years	1.0	1.2	4.2	4.2	5.6	5.3	4.2	3.9	4.4
25–34 years	2.7	2.7	6.1	7.2	11.7	11.4	10.0	10.0	12.2
35–44 years	5.0	4.9	8.2	8.3	10.9	10.9	10.3	9.6	11.9
45–54 years	3.9	4.3	8.3	9.4	12.0	12.1	11.1	11.1	12.5
55–64 years	1.1	1.0	2.8	3.5	7.0	6.9	7.3	8.0	9.2
65–74 years	0.5	0.3	0.7	0.7	1.2	1.7	2.0	2.2	2.5
75–84 years	*	*	0.4	0.6	0.7	0.7	0.7	0.9	0.8
85 years and over	*	*	*	*	1.3	*	1.0	1.3	*
Female									
All ages, age-adjusted ²	0.9	1.1	2.5	2.8	4.2	4.3	4.2	4.3	4.9
All ages, crude	0.9	1.1	2.5	2.8	4.2	4.4	4.2	4.4	4.9
Under 15 years	*	*	0.1	*	0.1	0.1	0.1	0.1	0.1
15–24 years	0.3	0.4	1.1	1.2	2.1	1.9	1.5	1.4	1.7
25–34 years	1.1	1.2	2.8	3.4	5.3	5.5	5.3	5.0	5.7
35–44 years	2.1	2.5	5.4	5.6	7.3	7.8	7.3	7.6	8.7
45–54 years	1.9	2.2	5.9	6.5	9.8	10.2	10.1	10.1	10.9
55–64 years	0.8	1.1	2.4	2.8	5.5	5.7	6.0	6.9	7.8
65–74 years	0.3	0.4	0.9	1.2	1.7	1.8	2.0	2.4	2.9
75–84 years	0.4	*	0.6	0.6	0.7	0.7	0.9	0.7	1.0
85 years and over	*	*	*	0.8	1.1	0.8	0.7	0.8	1.0
All ages, age-adjusted ^{2,3}									
Male:									
White	2.2	2.3	4.8	5.3	7.7	7.6	7.0	6.8	8.0
Black or African American	1.2	1.2	1.8	2.1	2.2	2.4	2.3	2.7	3.9
American Indian or Alaska Native	*	1.9	4.5	4.4	5.3	5.5	5.8	4.8	6.4
Asian or Pacific Islander	*	*	0.4	0.5	0.8	1.0	0.7	0.9	0.9
Hispanic or Latino	2.9	1.7	2.1	2.2	2.4	2.6	2.5	2.7	2.7
White, not Hispanic or Latino	2.1	2.3	5.3	5.9	9.0	8.8	8.1	7.9	9.3
Female:									
White	1.0	1.2	2.9	3.2	4.8	5.1	4.9	5.0	5.6
Black or African American	0.6	0.6	1.2	1.4	2.0	2.0	2.0	2.2	2.6
American Indian or Alaska Native	*	*	2.7	3.8	4.9	4.6	5.4	5.4	4.6
Asian or Pacific Islander	*	*	*	0.4	0.5	0.4	0.4	0.3	0.5
Hispanic or Latina	0.5	0.5	1.0	1.0	1.3	1.4	1.5	1.5	1.6
White, not Hispanic or Latina	1.1	1.3	3.2	3.5	5.6	5.8	5.6	5.8	6.5

See footnotes at end of table.

Table 27 (page 3 of 4). Death rates for drug poisoning and drug poisoning involving opioid analgesics and heroin, by sex, age, race, and Hispanic origin: United States, selected years 1999–2014

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2015.htm#027>.

[Data are based on death certificates]

Sex, age, race, and Hispanic origin	1999	2000	2004	2005	2010	2011	2012	2013	2014
Drug poisoning deaths involving heroin per 100,000 resident population ⁴									
All persons									
All ages, age-adjusted ²	0.7	0.7	0.6	0.7	1.0	1.4	1.9	2.7	3.4
All ages, crude	0.7	0.7	0.6	0.7	1.0	1.4	1.9	2.6	3.3
Under 15 years	*	*	*	*	*	*	*	*	*
15–24 years	0.5	0.6	0.6	0.7	1.2	1.8	2.2	2.9	3.3
25–34 years	1.0	1.0	1.1	1.2	2.2	3.4	4.6	6.3	8.0
35–44 years	1.8	1.5	1.2	1.2	1.6	2.2	3.1	4.4	5.9
45–54 years	1.3	1.2	1.2	1.4	1.4	2.0	2.8	3.7	4.7
55–64 years	0.3	0.3	0.4	0.4	0.7	1.0	1.3	2.1	2.7
65–74 years	*	*	*	*	*	0.2	0.1	0.3	0.5
75–84 years	*	*	*	*	*	*	*	*	*
85 years and over	*	*	*	*	*	*	*	*	*
Male									
All ages, age-adjusted ²	1.2	1.1	1.1	1.1	1.6	2.3	3.1	4.2	5.2
All ages, crude	1.2	1.1	1.1	1.1	1.6	2.3	3.0	4.2	5.2
Under 15 years	*	*	*	*	*	*	*	*	*
15–24 years	0.8	0.9	1.0	1.0	1.9	2.8	3.2	4.2	4.8
25–34 years	1.6	1.7	1.9	2.0	3.5	5.4	7.1	9.9	12.3
35–44 years	3.0	2.6	2.0	1.9	2.8	3.6	5.1	6.9	9.2
45–54 years	2.3	2.2	2.0	2.3	2.4	3.2	4.5	6.0	7.2
55–64 years	0.5	0.4	0.6	0.7	1.1	1.7	2.3	3.6	4.4
65–74 years	*	*	*	*	*	0.3	0.3	0.5	0.9
75–84 years	*	*	*	*	*	*	*	*	*
85 years and over	*	*	*	*	*	*	*	*	*
Female									
All ages, age-adjusted ²	0.2	0.2	0.2	0.3	0.4	0.6	0.8	1.2	1.6
All ages, crude	0.2	0.2	0.2	0.3	0.4	0.6	0.8	1.1	1.5
Under 15 years	*	*	*	*	*	*	*	*	*
15–24 years	0.2	0.2	0.3	0.3	0.6	0.9	1.1	1.5	1.7
25–34 years	0.3	0.4	0.4	0.5	0.9	1.4	2.0	2.6	3.7
35–44 years	0.6	0.5	0.5	0.5	0.6	0.8	1.1	1.9	2.6
45–54 years	0.3	0.3	0.4	0.5	0.5	0.8	1.1	1.6	2.2
55–64 years	*	*	0.1	*	0.3	0.3	0.4	0.7	1.0
65–74 years	*	*	*	*	*	*	*	*	*
75–84 years	*	*	*	*	*	*	*	*	*
85 years and over	*	*	*	*	*	*	*	*	*
All ages, age-adjusted ^{2,3}									
Male:									
White	1.2	1.1	1.1	1.1	1.8	2.6	3.5	4.7	6.0
Black or African American	1.4	1.6	1.1	1.3	1.2	1.6	2.1	3.4	4.1
American Indian or Alaska Native	*	*	*	*	*	1.3	1.7	2.6	3.5
Asian or Pacific Islander	*	*	*	*	*	0.3	0.3	0.5	0.6
Hispanic or Latino	2.0	1.6	1.3	1.4	1.5	1.7	2.2	2.6	3.2
White, not Hispanic or Latino	1.1	1.0	1.1	1.1	1.9	2.9	3.9	5.3	6.7
Female:									
White	0.2	0.2	0.2	0.3	0.4	0.6	0.9	1.3	1.8
Black or African American	0.3	0.3	0.3	0.3	0.3	0.4	0.5	0.7	1.1
American Indian or Alaska Native	*	*	*	*	*	1.0	*	1.0	1.2
Asian or Pacific Islander	*	*	*	*	*	*	*	*	*
Hispanic or Latina	0.2	0.1	0.2	0.2	0.2	0.3	0.4	0.5	0.7
White, not Hispanic or Latina	0.2	0.2	0.3	0.3	0.5	0.7	1.1	1.5	2.1

See footnotes at end of table.

Table 27 (page 4 of 4). Death rates for drug poisoning and drug poisoning involving opioid analgesics and heroin, by sex, age, race, and Hispanic origin: United States, selected years 1999–2014

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2015.htm#027>.

[Data are based on death certificates]

* Rates based on fewer than 20 deaths are considered unreliable and are not shown.
0.0 Rate more than zero but less than 0.05.

¹Drug poisoning was coded using underlying cause of death according to the 10th Revision of the *International Classification of Diseases* (ICD–10). See Appendix II, Cause of death; Table IV. Drug poisoning deaths include those resulting from accidental or intentional overdoses of a drug, being given the wrong drug, taking the wrong drug in error, taking a drug inadvertently, or other misuses of drugs. These deaths are from all manners and intents, including unintentional, suicide, homicide, undetermined intent, legal intervention, and operations of war.

²Age-adjusted rates are calculated using the year 2000 standard population with unrounded population numbers. See Appendix II, Age adjustment.

³The race groups, white, black, Asian or Pacific Islander, and American Indian or Alaska Native, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Death rates for Hispanic, American Indian or Alaska Native, and Asian or Pacific Islander persons should be interpreted with caution because of inconsistencies in reporting Hispanic origin or race on the death certificate (death rate numerators) compared with population figures (death rate denominators). The net effect of misclassification is an underestimation of deaths and death rates for races other than white and black. See Appendix II, Race, for a detailed discussion of sources of bias in death rates by race and Hispanic origin.

⁴Opioid analgesics include opioids such as hydrocodone, codeine, and methadone, and synthetic narcotics such as fentanyl, tramadol, and propoxyphene (removed from the market in 2010). Drug poisoning deaths involving opioid analgesics include those with an underlying cause of drug poisoning and with an opioid analgesic mentioned in the ICD–10 multiple causes of death. Drug poisoning deaths involving heroin include those with an underlying cause of drug poisoning and with heroin mentioned in the ICD–10 multiple causes of death. See Appendix I, National Vital Statistics System (NVSS), Mortality Multiple Cause-of-Death File, See Appendix II, Cause of death; Table IV. Drug-poisoning deaths may involve multiple drugs. Deaths involving both opioid analgesics and heroin are included in the death rate for opioid analgesics and the death rate for heroin. Opioid analgesic death rates include deaths involving fentanyl, a synthetic opioid. A sharp increase in deaths involving synthetic opioids, other than methadone, in 2014 coincided with law enforcement reports of increased availability of illicitly manufactured, or non-pharmaceutical, fentanyl. Illicitly manufactured fentanyl cannot be distinguished from pharmaceutical fentanyl in death certificate data. For more information, see CDC health advisory: Increases in fentanyl drug confiscations and fentanyl-related overdose fatalities. Available from: <http://emergency.cdc.gov/han/han00384.asp>, and Rudd RA, Aleshire N, Zibbell JE, Gladden M. Increases in drug and opioid overdose deaths—United States, 2000–2014. *MMWR* 2016;64(50):1378–82. Available from: http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6450a3.htm?s_cid=mm6450a3_w. Metabolic breakdown of heroin into morphine in the body can make it difficult to distinguish between deaths from heroin and deaths from morphine based on the information on the death certificate. Some deaths reported to involve morphine could be deaths from heroin. This may result in an undercount of heroin-related deaths. For more information, see Hedegaard H, Chen LH, Warner M. Drug-poisoning deaths involving heroin: United States, 2000–2013. NCHS data brief, no 190. Hyattsville, MD: NCHS. 2015. Available from: <http://www.cdc.gov/nchs/data/databriefs/db190.htm>. In 1999–2014, 19%–25% of drug poisoning deaths did not include specific information on the death certificate on the type of drug that was involved. Some of these deaths could have potentially involved heroin or opioid analgesics. For more information, see NCHS Health E-Stat available from: http://www.cdc.gov/nchs/data/hestat/drug_poisoning/drug_poisoning.htm.

NOTES: Rates for 1999 were computed using intercensal population estimates based on the 1990 and 2000 censuses. Rates for 2000 were computed based on 2000 bridged-race April 1 census counts. Starting with *Health, United States, 2012*, rates for 2001–2009 were revised using intercensal population estimates based on the 2000 and 2010 censuses. Rates for 2010 were based on 2010 bridged-race April 1 census counts. Rates for 2011 and beyond were computed using 2010-based postcensal estimates. See Appendix I, Population Census and Population Estimates. Age groups were selected to minimize the presentation of unstable age-specific death rates based on small numbers of deaths and for consistency among comparison groups. For additional injury-related statistics, see the Web-based Injury Statistics Query and Reporting System, available from: <http://www.cdc.gov/injury/wisqars/index.html>. Starting with 2003 data, some states allowed the reporting of more than one race on the death certificate. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards, for comparability with other states. See Appendix II, Race. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Vital Statistics System; numerator data from National Vital Statistics System, annual public-use Mortality Files; denominator data from national population estimates for race groups from Table 1; Deaths: Final data for 2014. National vital statistics reports (forthcoming). Available from: <http://www.cdc.gov/nchs/products/nvsr.htm>. See Appendix I, National Vital Statistics System (NVSS).

Table 28 (page 1 of 4). Death rates for motor vehicle-related injuries, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#028>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and age	1950 ^{1,2}	1960 ^{1,2}	1970 ²	1980 ²	1990 ²	2000 ³	2010 ³	2013 ³	2014 ³
	Deaths per 100,000 resident population								
All persons									
All ages, age-adjusted ⁴	24.6	23.1	27.6	22.3	18.5	15.4	11.3	10.9	10.8
All ages, crude	23.1	21.3	26.9	23.5	18.8	15.4	11.4	11.2	11.1
Under 1 year	8.4	8.1	9.8	7.0	4.9	4.4	2.0	1.7	1.7
1–14 years	9.8	8.6	10.5	8.2	6.0	4.3	2.3	2.2	2.2
1–4 years	11.5	10.0	11.5	9.2	6.3	4.2	2.8	2.7	2.5
5–14 years	8.8	7.9	10.2	7.9	5.9	4.3	2.2	2.1	2.0
15–24 years	34.4	38.0	47.2	44.8	34.1	26.9	16.6	15.2	15.3
15–19 years	29.6	33.9	43.6	43.0	33.1	26.0	13.6	11.4	11.9
20–24 years	38.8	42.9	51.3	46.6	35.0	28.0	19.7	18.8	18.3
25–34 years	24.6	24.3	30.9	29.1	23.6	17.3	14.0	13.9	13.9
35–44 years	20.3	19.3	24.9	20.9	16.9	15.3	11.6	11.4	11.1
45–64 years	25.2	23.0	26.5	18.0	15.7	14.3	11.9	12.1	12.0
45–54 years	22.2	21.4	25.5	18.6	15.6	14.2	12.0	12.2	12.1
55–64 years	29.0	25.1	27.9	17.4	15.9	14.4	11.9	11.9	11.9
65 years and over	43.1	34.7	36.2	22.5	23.1	21.4	16.0	15.1	14.8
65–74 years	39.1	31.4	32.8	19.2	18.6	16.5	12.3	12.2	11.9
75–84 years	52.7	41.8	43.5	28.1	29.1	25.7	18.8	17.8	17.5
85 years and over	45.1	37.9	34.2	27.6	31.2	30.4	23.8	21.0	20.9
Male									
All ages, age-adjusted ⁴	38.5	35.4	41.5	33.6	26.5	21.7	16.2	15.9	15.8
All ages, crude	35.4	31.8	39.7	35.3	26.7	21.3	16.3	16.1	16.0
Under 1 year	9.1	8.6	9.3	7.3	5.0	4.6	2.2	1.7	1.9
1–14 years	12.3	10.7	13.0	10.0	7.0	4.9	2.7	2.5	2.5
1–4 years	13.0	11.5	12.9	10.2	6.9	4.7	3.0	3.0	2.8
5–14 years	11.9	10.4	13.1	9.9	7.0	5.0	2.5	2.4	2.4
15–24 years	56.7	61.2	73.2	68.4	49.5	37.4	23.1	21.2	21.5
15–19 years	46.3	51.7	64.1	62.6	45.5	33.9	17.8	14.7	16.1
20–24 years	66.7	73.2	84.4	74.3	53.3	41.2	28.5	27.3	26.5
25–34 years	40.8	40.1	49.4	46.3	35.7	25.5	21.0	20.8	20.8
35–44 years	32.5	29.9	37.7	31.7	24.7	22.0	16.9	16.9	16.4
45–64 years	37.7	33.3	38.9	26.5	21.9	20.2	17.9	18.2	18.1
45–54 years	33.6	31.6	37.2	27.6	22.0	20.4	17.9	18.3	18.2
55–64 years	43.1	35.6	40.9	25.4	21.7	19.8	17.8	18.1	18.1
65 years and over	66.6	52.1	54.4	33.9	32.1	29.5	22.2	21.5	21.0
65–74 years	59.1	45.8	47.3	27.3	24.2	21.7	17.1	17.5	17.0
75–84 years	85.0	66.0	68.2	44.3	41.2	35.6	25.9	24.9	24.8
85 years and over	78.1	62.7	63.1	56.1	64.5	57.5	40.2	35.3	34.0
Female									
All ages, age-adjusted ⁴	11.5	11.7	14.9	11.8	11.0	9.5	6.5	6.2	6.1
All ages, crude	10.9	11.0	14.7	12.3	11.3	9.7	6.8	6.4	6.3
Under 1 year	7.6	7.5	10.4	6.7	4.9	4.2	1.8	1.8	1.5
1–14 years	7.2	6.3	7.9	6.3	4.9	3.7	2.0	1.9	1.8
1–4 years	10.0	8.4	10.0	8.1	5.6	3.8	2.5	2.3	2.3
5–14 years	5.7	5.4	7.2	5.7	4.7	3.6	1.8	1.8	1.6
15–24 years	12.6	15.1	21.6	20.8	17.9	15.9	9.9	8.9	8.7
15–19 years	12.9	16.0	22.7	22.8	20.0	17.5	9.2	7.9	7.6
20–24 years	12.2	14.0	20.4	18.9	16.0	14.2	10.5	9.9	9.7
25–34 years	9.3	9.2	13.0	12.2	11.5	8.8	6.9	6.9	6.8
35–44 years	8.5	9.1	12.9	10.4	9.2	8.8	6.2	5.9	5.8
45–64 years	12.6	13.1	15.3	10.3	10.1	8.7	6.3	6.3	6.2
45–54 years	10.9	11.6	14.5	10.2	9.6	8.2	6.3	6.3	6.2
55–64 years	14.9	15.2	16.2	10.5	10.8	9.5	6.3	6.2	6.1
65 years and over	21.9	20.3	23.1	15.0	17.2	15.8	11.3	10.0	9.9
65–74 years	20.6	19.0	21.6	13.0	14.1	12.3	8.2	7.5	7.5
75–84 years	25.2	23.0	27.2	18.5	21.9	19.2	13.7	12.5	12.0
85 years and over	22.1	22.0	18.0	15.2	18.3	19.3	15.9	13.7	14.1
White male ⁵									
All ages, age-adjusted ⁴	37.9	34.8	40.4	33.8	26.3	21.8	16.7	16.3	16.1
All ages, crude	35.1	31.5	39.1	35.9	26.7	21.6	17.0	16.7	16.5
Under 1 year	9.1	8.8	9.1	7.0	4.8	4.2	2.0	1.7	1.8
1–14 years	12.4	10.6	12.5	9.8	6.6	4.8	2.7	2.4	2.4
15–24 years	58.3	62.7	75.2	73.8	52.5	39.6	24.6	22.9	23.0
25–34 years	39.1	38.6	47.0	46.6	35.4	25.1	21.4	21.1	20.8
35–44 years	30.9	28.4	35.2	30.7	23.7	21.8	17.4	17.3	16.6
45–64 years	36.2	31.7	36.5	25.2	20.6	19.7	18.3	18.5	18.3
65 years and over	67.1	52.1	54.2	32.7	31.4	29.4	22.7	22.1	21.5

See footnotes at end of table.

Table 28 (page 2 of 4). Death rates for motor vehicle-related injuries, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#028>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and age	1950 ^{1,2}	1960 ^{1,2}	1970 ²	1980 ²	1990 ²	2000 ³	2010 ³	2013 ³	2014 ³
Deaths per 100,000 resident population									
Black or African American male⁵									
All ages, age-adjusted ⁴	34.8	39.6	51.0	34.2	29.9	24.4	16.7	17.0	17.4
All ages, crude	37.2	33.1	44.3	31.1	28.1	22.5	15.9	16.4	16.8
Under 1 year	---	*	10.6	7.8	*	6.7	*	*	*
1–14 years ⁶	10.4	11.2	16.3	11.4	8.9	5.5	3.0	3.3	2.9
15–24 years	42.5	46.4	58.1	34.9	36.1	30.2	19.4	17.6	18.7
25–34 years	54.4	51.0	70.4	44.9	39.5	32.6	24.9	25.1	25.7
35–44 years	46.7	43.6	59.5	41.2	33.5	27.2	19.4	20.5	21.2
45–64 years	54.6	47.8	61.7	39.5	33.3	27.1	19.1	21.1	21.1
65 years and over	52.6	48.2	53.4	42.4	36.3	32.1	20.0	18.5	19.8
American Indian or Alaska Native male⁵									
All ages, age-adjusted ⁴	---	---	---	78.9	48.3	35.8	21.1	20.0	23.4
All ages, crude	---	---	---	74.6	47.6	33.6	19.8	19.0	22.2
1–14 years	---	---	---	15.1	11.6	7.8	*	3.9	4.2
15–24 years	---	---	---	126.1	75.2	56.8	31.9	26.8	27.3
25–34 years	---	---	---	107.0	78.2	49.8	23.8	30.3	31.8
35–44 years	---	---	---	82.8	57.0	36.3	24.5	19.8	24.2
45–64 years	---	---	---	77.4	45.9	32.0	23.2	20.4	30.5
65 years and over	---	---	---	97.0	43.0	48.5	26.6	25.8	26.8
Asian or Pacific Islander male⁵									
All ages, age-adjusted ⁴	---	---	---	19.0	17.9	10.6	6.5	6.4	6.2
All ages, crude	---	---	---	17.1	15.8	9.8	6.2	5.9	6.0
1–14 years	---	---	---	8.2	6.3	2.5	*	1.5	1.2
15–24 years	---	---	---	27.2	25.7	17.0	9.6	8.1	8.6
25–34 years	---	---	---	18.8	17.0	10.4	7.8	5.9	8.0
35–44 years	---	---	---	13.1	12.2	6.9	4.1	5.3	4.1
45–64 years	---	---	---	13.7	15.1	10.1	6.0	5.9	5.7
65 years and over	---	---	---	37.3	33.6	21.1	14.6	14.3	13.1
Hispanic or Latino male^{5,7}									
All ages, age-adjusted ⁴	---	---	---	---	29.5	21.3	14.0	14.2	14.3
All ages, crude	---	---	---	---	29.2	20.1	12.8	13.3	13.5
1–14 years	---	---	---	---	7.2	4.4	2.5	2.5	2.5
15–24 years	---	---	---	---	48.2	34.7	20.2	20.5	22.0
25–34 years	---	---	---	---	41.0	24.9	18.0	18.2	19.6
35–44 years	---	---	---	---	28.0	21.6	13.9	14.5	13.4
45–64 years	---	---	---	---	28.9	21.7	14.3	15.9	14.9
65 years and over	---	---	---	---	35.3	28.9	20.7	18.8	19.5
White, not Hispanic or Latino male⁷									
All ages, age-adjusted ⁴	---	---	---	---	25.7	21.7	17.1	16.5	16.2
All ages, crude	---	---	---	---	26.0	21.5	17.6	17.2	16.9
1–14 years	---	---	---	---	6.4	4.9	2.7	2.3	2.3
15–24 years	---	---	---	---	52.3	40.3	25.4	23.0	22.6
25–34 years	---	---	---	---	34.0	24.7	21.9	21.4	20.7
35–44 years	---	---	---	---	23.1	21.6	18.0	17.8	17.2
45–64 years	---	---	---	---	19.8	19.3	18.6	18.6	18.6
65 years and over	---	---	---	---	31.1	29.3	22.7	22.3	21.5
White female⁵									
All ages, age-adjusted ⁴	11.4	11.7	14.9	12.2	11.2	9.8	6.8	6.4	6.3
All ages, crude	10.9	11.2	14.8	12.8	11.6	10.0	7.1	6.7	6.6
Under 1 year	7.8	7.5	10.2	7.1	4.7	3.5	1.9	1.7	1.5
1–14 years	7.2	6.2	7.5	6.2	4.8	3.7	2.1	1.8	1.8
15–24 years	12.6	15.6	22.7	23.0	19.5	17.1	10.8	9.5	9.2
25–34 years	9.0	9.0	12.7	12.2	11.6	8.9	7.1	7.1	7.1
35–44 years	8.1	8.9	12.3	10.6	9.2	8.9	6.5	6.1	6.1
45–64 years	12.7	13.1	15.1	10.4	9.9	8.7	6.4	6.4	6.3
65 years and over	22.2	20.8	23.7	15.3	17.4	16.2	11.5	10.4	10.3

See footnotes at end of table.

Table 28 (page 3 of 4). Death rates for motor vehicle-related injuries, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#028>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and age	1950 ^{1,2}	1960 ^{1,2}	1970 ²	1980 ²	1990 ²	2000 ³	2010 ³	2013 ³	2014 ³
Deaths per 100,000 resident population									
Black or African American female⁵									
All ages, age-adjusted ⁴	9.3	10.4	14.1	8.5	9.6	8.4	5.9	5.7	5.6
All ages, crude	10.2	9.7	13.4	8.3	9.4	8.2	5.8	5.7	5.6
Under 1 year	---	8.1	11.9	*	7.0	*	*	*	*
1–14 years ⁶	7.2	6.9	10.2	6.3	5.3	3.9	2.0	2.3	2.3
15–24 years	11.6	9.9	13.4	8.0	9.9	11.7	7.8	7.7	7.7
25–34 years	10.8	9.8	13.3	10.6	11.1	9.4	6.8	7.2	6.9
35–44 years	11.1	11.0	16.1	8.3	9.4	8.2	5.8	6.4	5.7
45–64 years	11.8	12.7	16.7	9.2	10.7	9.0	6.3	6.2	6.2
65 years and over	14.3	13.2	15.7	9.5	13.5	10.4	8.6	5.9	6.4
American Indian or Alaska Native female⁵									
All ages, age-adjusted ⁴	---	---	---	32.0	17.5	19.5	10.6	11.0	10.1
All ages, crude	---	---	---	32.0	17.3	18.6	10.0	10.7	10.0
1–14 years	---	---	---	15.0	8.1	6.5	*	4.9	*
15–24 years	---	---	---	42.3	31.4	30.3	13.4	13.6	16.5
25–34 years	---	---	---	52.5	18.8	22.3	17.7	17.0	13.4
35–44 years	---	---	---	38.1	18.2	22.0	13.1	14.2	12.0
45–64 years	---	---	---	32.6	17.6	17.8	8.4	8.7	9.7
65 years and over	---	---	---	*	*	24.0	14.8	*	*
Asian or Pacific Islander female⁵									
All ages, age-adjusted ⁴	---	---	---	9.3	10.4	6.7	3.9	3.4	3.2
All ages, crude	---	---	---	8.2	9.0	5.9	3.6	3.4	3.2
1–14 years	---	---	---	7.4	3.6	2.3	*	*	*
15–24 years	---	---	---	7.4	11.4	6.0	3.3	3.0	3.0
25–34 years	---	---	---	7.3	7.3	4.5	3.1	2.5	2.5
35–44 years	---	---	---	8.6	7.5	4.9	2.0	*	2.0
45–64 years	---	---	---	8.5	11.8	6.4	4.3	4.1	3.3
65 years and over	---	---	---	18.6	24.3	18.5	12.2	10.8	9.4
Hispanic or Latina female^{5,7}									
All ages, age-adjusted ⁴	---	---	---	---	9.6	7.9	5.3	5.2	5.0
All ages, crude	---	---	---	---	8.9	7.2	4.9	4.8	4.7
1–14 years	---	---	---	---	4.8	3.9	2.0	1.7	1.9
15–24 years	---	---	---	---	11.6	10.6	7.7	7.4	7.5
25–34 years	---	---	---	---	9.4	6.5	5.0	5.1	5.3
35–44 years	---	---	---	---	8.0	7.3	4.5	4.7	4.1
45–64 years	---	---	---	---	11.4	8.3	5.6	5.3	5.3
65 years and over	---	---	---	---	14.9	13.4	9.4	9.0	8.1
White, not Hispanic or Latina female⁷									
All ages, age-adjusted ⁴	---	---	---	---	11.3	10.0	7.0	6.6	6.5
All ages, crude	---	---	---	---	11.7	10.3	7.5	7.1	7.0
1–14 years	---	---	---	---	4.7	3.5	2.0	1.8	1.7
15–24 years	---	---	---	---	20.4	18.4	11.4	10.0	9.6
25–34 years	---	---	---	---	11.7	9.3	7.6	7.5	7.5
35–44 years	---	---	---	---	9.3	9.0	6.9	6.4	6.6
45–64 years	---	---	---	---	9.7	8.7	6.4	6.5	6.4
65 years and over	---	---	---	---	17.5	16.3	11.6	10.5	10.4

See footnotes at end of table.

Table 28 (page 4 of 4). Death rates for motor vehicle-related injuries, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#028>.

[Data are based on death certificates]

-- Data not available.

* Rates based on fewer than 20 deaths are considered unreliable and are not shown.

¹Includes deaths of persons who were not residents of the 50 states and the District of Columbia (D.C.).

²Underlying cause of death was coded according to the 6th Revision of the *International Classification of Diseases* (ICD) in 1950, 7th Revision in 1960, 8th Revision in 1970, and 9th Revision in 1980–1998. See Appendix II, Cause of death; Table III; Table IV.

³Starting with 1999 data, cause of death is coded according to ICD–10. See Appendix II, Cause of death; Comparability ratio; Table IV; Table V.

⁴Age-adjusted rates are calculated using the year 2000 standard population. Prior to 2001, age-adjusted rates were calculated using standard million proportions based on rounded population numbers. Starting with 2001 data, unrounded population numbers are used to calculate age-adjusted rates. See Appendix II, Age adjustment.

⁵The race groups, white, black, Asian or Pacific Islander, and American Indian or Alaska Native, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Death rates for Hispanic, American Indian or Alaska Native, and Asian or Pacific Islander persons should be interpreted with caution because of inconsistencies in reporting Hispanic origin or race on the death certificate (death rate numerators) compared with population figures (death rate denominators). The net effect of misclassification is an underestimation of deaths and death rates for races other than white and black. See Appendix II, Race, for a detailed discussion of sources of bias in death rates by race and Hispanic origin.

⁶In 1950, rate is for the age group under 15 years.

⁷Prior to 1997, data from states that did not report Hispanic origin on the death certificate were excluded. See Appendix II, Hispanic origin.

NOTES: Starting with *Health, United States, 2003*, rates for 1991–1999 were revised using intercensal population estimates based on the 1990 and 2000 censuses. For 2000, population estimates are bridged-race April 1 census counts. Starting with *Health, United States, 2012*, rates for 2001–2009 were revised using intercensal population estimates based on the 2000 and 2010 censuses. For 2010, population estimates are bridged-race April 1 census counts. Rates for 2011 and beyond were computed using 2010-based postcensal estimates. See Appendix I, Population Census and Population Estimates. Age groups were selected to minimize the presentation of unstable age-specific death rates based on small numbers of deaths and for consistency among comparison groups. For additional injury-related statistics, see Web-based Injury Statistics Query and Reporting System (WISQARS), available from: <http://www.cdc.gov/injury/wisqars/index.html>. Starting with 2003 data, some states allowed the reporting of more than one race on the death certificate. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards, for comparability with other states. See Appendix II, Race. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Vital Statistics System; Grove RD, Hetzel AM. Vital statistics rates in the United States, 1940–1960. Washington, DC: U.S. Government Printing Office, 1968; numerator data from National Vital Statistics System, annual public-use Mortality Files; denominator data from national population estimates for race groups from Table 1 and unpublished Hispanic population estimates for 1985–1996 prepared by the Housing and Household Economic Statistics Division, U.S. Census Bureau; Deaths: Final data for 2014. National vital statistics reports (forthcoming). Available from: <http://www.cdc.gov/nchs/products/nvsr.htm>. See Appendix I, National Vital Statistics System (NVSS).

Table 29 (page 1 of 4). Death rates for homicide, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#029>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and age	1950 ^{1,2}	1960 ^{1,2}	1970 ²	1980 ²	1990 ²	2000 ³	2010 ³	2013 ³	2014 ³
Deaths per 100,000 resident population									
All persons									
All ages, age-adjusted ⁴	5.1	5.0	8.8	10.4	9.4	5.9	5.3	5.2	5.1
All ages, crude	5.0	4.6	8.1	10.6	9.9	6.0	5.3	5.1	5.0
Under 1 year	4.4	4.8	4.3	5.9	8.4	9.2	7.9	7.2	6.3
1–14 years	0.6	0.6	1.1	1.5	1.8	1.3	1.1	1.1	1.1
1–4 years	0.6	0.7	1.9	2.5	2.5	2.3	2.4	2.1	2.3
5–14 years	0.5	0.5	0.9	1.2	1.5	0.9	0.6	0.7	0.7
15–24 years	5.8	5.6	11.3	15.4	19.7	12.6	10.7	9.8	9.5
15–19 years	3.9	3.9	7.7	10.5	16.9	9.5	8.3	6.6	6.7
20–24 years	8.5	7.7	15.6	20.2	22.2	16.0	13.2	12.8	12.1
25–44 years	8.9	8.5	14.9	17.5	14.7	8.7	8.2	8.2	8.1
25–34 years	9.3	9.2	16.2	19.3	17.4	10.4	10.4	9.9	9.6
35–44 years	8.4	7.8	13.5	14.9	11.6	7.1	6.0	6.4	6.4
45–64 years	5.0	5.3	8.7	9.0	6.3	4.0	3.8	3.8	3.7
45–54 years	5.9	6.1	10.0	11.0	7.5	4.7	4.4	4.5	4.5
55–64 years	3.9	4.1	7.1	7.0	5.0	3.0	2.9	3.0	2.9
65 years and over	3.0	2.7	4.6	5.5	4.0	2.4	2.0	2.0	2.0
65–74 years	3.2	2.8	4.9	5.7	3.8	2.4	2.1	2.0	2.1
75–84 years	2.5	2.3	4.0	5.2	4.3	2.4	1.9	2.1	2.0
85 years and over	2.3	2.4	4.2	5.3	4.6	2.4	2.0	1.9	1.9
Male									
All ages, age-adjusted ⁴	7.9	7.5	14.3	16.6	14.8	9.0	8.4	8.2	8.0
All ages, crude	7.7	6.8	13.1	17.1	15.9	9.3	8.4	8.2	8.0
Under 1 year	4.5	4.7	4.5	6.3	8.8	10.4	8.8	8.7	7.1
1–14 years	0.6	0.6	1.2	1.6	2.0	1.5	1.4	1.2	1.2
1–4 years	0.5	0.7	1.9	2.7	2.7	2.5	2.8	2.3	2.4
5–14 years	0.6	0.5	1.0	1.2	1.7	1.1	0.8	0.8	0.8
15–24 years	8.6	8.4	18.2	24.0	32.5	20.9	18.2	16.7	16.1
15–19 years	5.5	5.7	12.1	15.9	27.8	15.5	14.0	11.4	11.2
20–24 years	13.5	11.8	25.6	32.2	36.9	26.7	22.6	21.6	20.6
25–44 years	13.8	12.8	24.4	28.9	23.5	13.3	13.3	13.4	13.2
25–34 years	14.4	13.9	26.8	31.9	27.7	16.7	17.3	16.4	15.9
35–44 years	13.2	11.7	21.7	24.5	18.6	10.3	9.2	10.1	10.3
45–64 years	8.1	8.1	14.8	15.2	10.2	6.0	5.6	5.7	5.7
45–54 years	9.5	9.4	16.8	18.4	11.9	6.9	6.7	6.9	6.9
55–64 years	6.3	6.4	12.1	11.8	8.0	4.6	4.3	4.3	4.3
65 years and over	4.8	4.3	7.7	8.8	5.8	3.3	2.6	2.8	2.7
65–74 years	5.2	4.6	8.5	9.2	5.8	3.4	2.9	2.9	2.9
75–84 years	3.9	3.7	5.9	8.1	5.7	3.2	2.1	2.5	2.2
85 years and over	2.5	3.6	7.4	7.5	6.7	3.3	2.2	2.9	2.2
Female									
All ages, age-adjusted ⁴	2.4	2.6	3.7	4.4	4.0	2.8	2.3	2.1	2.1
All ages, crude	2.4	2.4	3.4	4.5	4.2	2.8	2.2	2.1	2.1
Under 1 year	4.2	4.9	4.1	5.6	8.0	7.9	6.9	5.5	5.4
1–14 years	0.6	0.5	1.0	1.4	1.6	1.1	0.9	0.9	1.0
1–4 years	0.7	0.7	1.9	2.2	2.3	2.1	1.9	2.0	2.2
5–14 years	0.5	0.4	0.7	1.1	1.2	0.7	0.5	0.5	0.6
15–24 years	3.0	2.8	4.6	6.6	6.2	3.9	2.9	2.6	2.5
15–19 years	2.4	1.9	3.2	4.9	5.4	3.1	2.3	1.6	1.9
20–24 years	3.7	3.8	6.2	8.2	7.0	4.7	3.4	3.6	3.1
25–44 years	4.2	4.3	5.8	6.4	6.0	4.0	3.1	3.0	2.9
25–34 years	4.5	4.6	6.0	6.9	7.1	4.1	3.3	3.2	3.2
35–44 years	3.8	4.0	5.7	5.7	4.8	4.0	2.9	2.7	2.6
45–64 years	1.9	2.5	3.1	3.4	2.8	2.1	2.0	2.0	1.8
45–54 years	2.3	2.9	3.7	4.1	3.2	2.5	2.3	2.3	2.2
55–64 years	1.4	2.0	2.5	2.8	2.3	1.6	1.7	1.7	1.5
65 years and over	1.4	1.3	2.3	3.3	2.8	1.8	1.6	1.4	1.5
65–74 years	1.3	1.3	2.2	3.0	2.2	1.6	1.4	1.3	1.3
75–84 years	1.4	1.3	2.7	3.5	3.4	2.0	1.8	1.7	1.8
85 years and over	2.1	1.6	2.5	4.3	3.8	2.0	2.0	1.4	1.8

See footnotes at end of table.

Table 29 (page 2 of 4). Death rates for homicide, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#029>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and age	1950 ^{1,2}	1960 ^{1,2}	1970 ²	1980 ²	1990 ²	2000 ³	2010 ³	2013 ³	2014 ³
Deaths per 100,000 resident population									
White male ⁵									
All ages, age-adjusted ⁴	3.8	3.9	7.2	10.4	8.3	5.2	4.7	4.4	4.3
All ages, crude	3.6	3.6	6.6	10.7	8.8	5.2	4.7	4.4	4.3
Under 1 year	4.3	3.8	2.9	4.3	6.4	8.2	8.5	6.8	5.8
1–14 years	0.4	0.5	0.7	1.2	1.3	1.2	1.0	0.9	0.9
15–24 years	3.2	5.0	7.6	15.1	15.2	9.9	8.2	7.1	6.5
25–44 years	5.4	5.5	11.6	17.2	13.0	7.4	6.9	6.5	6.8
25–34 years	4.9	5.7	12.5	18.5	14.7	8.4	8.3	7.4	7.5
35–44 years	6.1	5.2	10.8	15.2	11.1	6.5	5.5	5.6	6.0
45–64 years	4.8	4.6	8.3	9.8	6.9	4.1	4.1	4.0	4.0
65 years and over	3.8	3.1	5.4	6.7	4.1	2.5	2.1	2.4	2.2
Black or African American male ⁵									
All ages, age-adjusted ⁴	47.0	42.3	78.2	69.4	63.1	35.4	31.5	31.6	30.6
All ages, crude	44.7	35.0	66.0	65.7	68.5	37.2	33.4	33.1	32.1
Under 1 year	---	10.3	14.3	18.6	21.4	23.3	12.3	17.9	13.9
1–14 years ⁶	1.8	1.5	4.4	4.1	5.8	3.1	3.4	2.6	3.1
15–24 years	53.8	43.2	98.3	82.6	137.1	85.3	71.0	66.6	65.0
25–44 years	92.8	80.5	140.2	130.0	105.4	55.8	55.9	57.9	54.8
25–34 years	104.3	86.4	154.5	142.9	123.7	73.9	76.1	73.2	67.7
35–44 years	80.0	74.4	124.0	109.3	81.2	38.5	34.5	40.4	39.7
45–64 years	46.0	44.6	82.3	70.6	41.4	21.9	17.6	18.4	18.2
65 years and over	16.5	17.3	33.3	30.9	25.7	12.8	8.0	7.3	8.4
American Indian or Alaska Native male ⁵									
All ages, age-adjusted ⁴	---	---	---	23.3	16.7	10.7	8.8	8.2	9.1
All ages, crude	---	---	---	23.1	16.6	10.7	9.5	8.3	9.2
15–24 years	---	---	---	35.4	25.1	17.0	17.6	7.1	12.1
25–44 years	---	---	---	39.2	25.7	17.0	14.8	15.4	13.6
45–64 years	---	---	---	22.1	14.8	*	6.5	8.0	9.7
Asian or Pacific Islander male ⁵									
All ages, age-adjusted ⁴	---	---	---	9.1	7.3	4.3	2.6	2.3	2.2
All ages, crude	---	---	---	8.3	7.9	4.4	2.7	2.3	2.3
15–24 years	---	---	---	9.3	14.9	7.8	4.0	3.5	3.4
25–44 years	---	---	---	11.3	9.6	4.6	3.3	2.7	3.0
45–64 years	---	---	---	10.4	7.0	6.1	3.1	2.5	2.5
Hispanic or Latino male ^{5,7}									
All ages, age-adjusted ⁴	---	---	---	---	27.4	11.8	8.7	7.3	7.2
All ages, crude	---	---	---	---	31.0	13.4	9.5	7.8	7.6
Under 1 year	---	---	---	---	8.7	6.6	7.0	6.8	4.5
1–14 years	---	---	---	---	3.1	1.7	1.1	0.9	0.7
15–24 years	---	---	---	---	55.4	28.5	19.7	15.3	14.2
25–44 years	---	---	---	---	46.4	17.2	13.2	11.0	11.6
25–34 years	---	---	---	---	50.9	19.9	16.8	13.3	13.1
35–44 years	---	---	---	---	39.3	13.5	8.9	8.4	9.9
45–64 years	---	---	---	---	20.5	9.1	6.9	6.1	5.6
65 years and over	---	---	---	---	9.4	4.4	3.2	2.9	3.3
White, not Hispanic or Latino male ⁷									
All ages, age-adjusted ⁴	---	---	---	---	5.6	3.6	3.3	3.4	3.3
All ages, crude	---	---	---	---	5.8	3.6	3.3	3.3	3.3
Under 1 year	---	---	---	---	5.4	8.3	8.7	6.5	6.5
1–14 years	---	---	---	---	0.9	1.0	0.9	0.9	0.9
15–24 years	---	---	---	---	7.5	4.7	4.1	3.9	3.5
25–44 years	---	---	---	---	8.7	5.2	4.7	4.9	4.9
25–34 years	---	---	---	---	9.3	5.2	5.0	5.2	5.3
35–44 years	---	---	---	---	8.0	5.2	4.4	4.6	4.6
45–64 years	---	---	---	---	5.7	3.6	3.6	3.6	3.6
65 years and over	---	---	---	---	3.7	2.3	2.0	2.3	2.0

See footnotes at end of table.

Table 29 (page 3 of 4). Death rates for homicide, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#029>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and age	1950 ^{1,2}	1960 ^{1,2}	1970 ²	1980 ²	1990 ²	2000 ³	2010 ³	2013 ³	2014 ³
Deaths per 100,000 resident population									
White female ⁵									
All ages, age-adjusted ⁴	1.4	1.5	2.3	3.2	2.7	2.1	1.8	1.7	1.7
All ages, crude	1.4	1.4	2.1	3.2	2.8	2.1	1.8	1.7	1.7
Under 1 year	3.9	3.5	2.9	4.3	5.1	5.0	5.8	4.3	4.5
1–14 years	0.4	0.4	0.7	1.1	1.0	0.8	0.7	0.7	0.8
15–24 years	1.3	1.5	2.7	4.7	4.0	2.7	2.0	1.8	1.7
25–44 years	2.0	2.1	3.3	4.2	3.8	2.9	2.4	2.3	2.3
45–64 years	1.5	1.7	2.1	2.6	2.3	1.8	1.7	1.7	1.5
65 years and over	1.2	1.2	1.9	2.9	2.2	1.6	1.6	1.4	1.5
Black or African American female ⁵									
All ages, age-adjusted ⁴	11.1	11.4	14.7	13.2	12.5	7.1	5.0	4.9	4.7
All ages, crude	11.5	10.4	13.2	13.5	13.4	7.2	5.1	4.9	4.7
Under 1 year	---	13.8	10.7	12.8	22.8	22.2	13.9	12.4	10.8
1–14 years ⁶	1.8	1.2	3.1	3.3	4.7	2.7	2.0	2.1	2.2
15–24 years	16.5	11.9	17.7	18.4	18.9	10.7	7.5	6.9	7.0
25–44 years	22.5	22.7	25.3	22.6	21.0	11.0	7.4	7.2	6.7
45–64 years	6.8	10.3	13.4	10.8	6.5	4.5	4.2	4.3	3.7
65 years and over	3.6	3.0	7.4	8.0	9.4	3.5	1.8	2.0	2.4
American Indian or Alaska Native female ⁵									
All ages, age-adjusted ⁴	---	---	---	8.1	4.6	3.0	2.5	2.4	2.5
All ages, crude	---	---	---	7.7	4.8	2.9	2.5	2.4	2.4
15–24 years	---	---	---	*	*	*	*	*	*
25–44 years	---	---	---	13.7	6.9	5.9	4.7	3.7	4.2
45–64 years	---	---	---	*	*	*	*	*	*
Asian or Pacific Islander female ⁵									
All ages, age-adjusted ⁴	---	---	---	3.1	2.8	1.7	1.2	0.9	1.0
All ages, crude	---	---	---	3.1	2.8	1.7	1.2	0.9	1.0
15–24 years	---	---	---	*	*	*	*	*	*
25–44 years	---	---	---	4.6	3.8	2.2	1.3	1.3	0.8
45–64 years	---	---	---	*	*	2.0	1.4	1.1	1.5
Hispanic or Latina female ^{5,7}									
All ages, age-adjusted ⁴	---	---	---	---	4.3	2.8	1.8	1.6	1.7
All ages, crude	---	---	---	---	4.7	2.8	1.8	1.6	1.8
Under 1 year	---	---	---	---	*	7.4	6.6	4.0	4.6
1–14 years	---	---	---	---	1.9	1.0	0.5	0.7	1.0
15–24 years	---	---	---	---	8.1	3.7	2.6	2.5	2.7
25–44 years	---	---	---	---	6.1	3.7	2.5	2.4	2.3
45–64 years	---	---	---	---	3.3	2.9	1.6	1.2	1.4
65 years and over	---	---	---	---	*	2.4	1.3	*	*
White, not Hispanic or Latina female ⁷									
All ages, age-adjusted ⁴	---	---	---	---	2.5	1.9	1.8	1.7	1.6
All ages, crude	---	---	---	---	2.5	1.9	1.7	1.7	1.6
Under 1 year	---	---	---	---	4.4	4.1	5.3	4.4	4.4
1–14 years	---	---	---	---	0.8	0.8	0.7	0.7	0.7
15–24 years	---	---	---	---	3.3	2.3	1.8	1.6	1.3
25–44 years	---	---	---	---	3.5	2.7	2.4	2.2	2.3
45–64 years	---	---	---	---	2.2	1.6	1.7	1.8	1.5
65 years and over	---	---	---	---	2.2	1.6	1.6	1.5	1.5

See footnotes at end of table.

Table 29 (page 4 of 4). Death rates for homicide, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#029>.

[Data are based on death certificates]

-- Data not available.

* Rates based on fewer than 20 deaths are considered unreliable and are not shown.

¹Includes deaths of persons who were not residents of the 50 states and the District of Columbia (D.C.).

²Underlying cause of death was coded according to the 6th Revision of the *International Classification of Diseases* (ICD) in 1950, 7th Revision in 1960, 8th Revision in 1970, and 9th Revision in 1980–1998. See Appendix II, Cause of death; Table III; Table IV.

³Starting with 1999 data, cause of death is coded according to ICD–10. See Appendix II, Cause of death; Comparability ratio; Table IV; Table V.

⁴Age-adjusted rates are calculated using the year 2000 standard population. Prior to 2001, age-adjusted rates were calculated using standard million proportions based on rounded population numbers. Starting with 2001 data, unrounded population numbers are used to calculate age-adjusted rates. See Appendix II, Age adjustment.

⁵The race groups, white, black, Asian or Pacific Islander, and American Indian or Alaska Native, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Death rates for Hispanic, American Indian or Alaska Native, and Asian or Pacific Islander persons should be interpreted with caution because of inconsistencies in reporting Hispanic origin or race on the death certificate (death rate numerators) compared with population figures (death rate denominators). The net effect of misclassification is an underestimation of deaths and death rates for races other than white and black. See Appendix II, Race, for a detailed discussion of sources of bias in death rates by race and Hispanic origin.

⁶In 1950, rate is for the age group under 15 years.

⁷Prior to 1997, data from states that did not report Hispanic origin on the death certificate were excluded. See Appendix II, Hispanic origin.

NOTES: Starting with *Health, United States, 2003*, rates for 1991–1999 were revised using intercensal population estimates based on the 1990 and 2000 censuses. For 2000, population estimates are bridged-race April 1 census counts. Starting with *Health, United States, 2012*, rates for 2001–2009 were revised using intercensal population estimates based on the 2000 and 2010 censuses. For 2010, population estimates are bridged-race April 1 census counts. Rates for 2011 and beyond were computed using 2010-based postcensal estimates. See Appendix I, Population Census and Population Estimates. Figures for 2001 include September 11-related deaths for which death certificates were filed as of October 24, 2002. See Appendix II, Cause of death; Table IV for terrorism-related ICD–10 codes. Age groups were selected to minimize the presentation of unstable age-specific death rates based on small numbers of deaths and for consistency among comparison groups. For additional injury-related statistics, see Web-based Injury Statistics Query and Reporting System (WISQARS), available from:

<http://www.cdc.gov/injury/wisqars/index.html>. Starting with 2003 data, some states allowed the reporting of more than one race on the death certificate. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards, for comparability with other states. See Appendix II, Race. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>. **Data have been revised and differ from previous editions of *Health, United States*.**

SOURCE: CDC/NCHS, National Vital Statistics System; Grove RD, Hetzel AM. Vital statistics rates in the United States, 1940–1960. Washington, DC: U.S. Government Printing Office, 1968; numerator data from National Vital Statistics System, annual public-use Mortality Files; denominator data from national population estimates for race groups from Table 1 and unpublished Hispanic population estimates for 1985–1996 prepared by the Housing and Household Economic Statistics Division, U.S. Census Bureau; Deaths: Final data for 2014. National vital statistics reports (forthcoming). Available from: <http://www.cdc.gov/nchs/products/nvsr.htm>. See Appendix I, National Vital Statistics System (NVSS).

Table 30 (page 1 of 3). Death rates for suicide, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#030>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and age	1950 ^{1,2}	1960 ^{1,2}	1970 ²	1980 ²	1990 ²	2000 ³	2010 ³	2013 ³	2014 ³
Deaths per 100,000 resident population									
All persons									
All ages, age-adjusted ⁴	13.2	12.5	13.1	12.2	12.5	10.4	12.1	12.6	13.0
All ages, crude	11.4	10.6	11.6	11.9	12.4	10.4	12.4	13.0	13.4
Under 1 year
1–4 years
5–14 years	0.2	0.3	0.3	0.4	0.8	0.7	0.7	1.0	1.0
15–24 years	4.5	5.2	8.8	12.3	13.2	10.2	10.5	11.1	11.6
15–19 years	2.7	3.6	5.9	8.5	11.1	8.0	7.5	8.3	8.7
20–24 years	6.2	7.1	12.2	16.1	15.1	12.5	13.6	13.7	14.2
25–44 years	11.6	12.2	15.4	15.6	15.2	13.4	15.0	15.5	15.8
25–34 years	9.1	10.0	14.1	16.0	15.2	12.0	14.0	14.8	15.1
35–44 years	14.3	14.2	16.9	15.4	15.3	14.5	16.0	16.2	16.6
45–64 years	23.5	22.0	20.6	15.9	15.3	13.5	18.6	19.0	19.5
45–54 years	20.9	20.7	20.0	15.9	14.8	14.4	19.6	19.7	20.2
55–64 years	26.8	23.7	21.4	15.9	16.0	12.1	17.5	18.1	18.8
65 years and over	30.0	24.5	20.8	17.6	20.5	15.2	14.9	16.1	16.7
65–74 years	29.6	23.0	20.8	16.9	17.9	12.5	13.7	15.0	15.6
75–84 years	31.1	27.9	21.2	19.1	24.9	17.6	15.7	17.1	17.5
85 years and over	28.8	26.0	19.0	19.2	22.2	19.6	17.6	18.6	19.3
Male									
All ages, age-adjusted ⁴	21.2	20.0	19.8	19.9	21.5	17.7	19.8	20.3	20.7
All ages, crude	17.8	16.5	16.8	18.6	20.4	17.1	19.9	20.6	21.1
Under 1 year
1–4 years
5–14 years	0.3	0.4	0.5	0.6	1.1	1.2	0.9	1.2	1.3
15–24 years	6.5	8.2	13.5	20.2	22.0	17.1	16.9	17.3	18.2
15–19 years	3.5	5.6	8.8	13.8	18.1	13.0	11.7	12.4	13.0
20–24 years	9.3	11.5	19.3	26.8	25.7	21.4	22.2	21.9	22.9
25–44 years	17.2	17.9	20.9	24.0	24.4	21.3	23.6	24.1	24.4
25–34 years	13.4	14.7	19.8	25.0	24.8	19.6	22.5	23.4	23.8
35–44 years	21.3	21.0	22.1	22.5	23.9	22.8	24.6	24.8	25.0
45–64 years	37.1	34.4	30.0	23.7	24.3	21.3	29.2	29.0	29.7
45–54 years	32.0	31.6	27.9	22.9	23.2	22.4	30.4	29.6	30.0
55–64 years	43.6	38.1	32.7	24.5	25.7	19.4	27.7	28.3	29.4
65 years and over	52.8	44.0	38.4	35.0	41.6	31.1	29.0	30.9	31.4
65–74 years	50.5	39.6	36.0	30.4	32.2	22.7	23.9	26.0	26.6
75–84 years	58.3	52.5	42.8	42.3	56.1	38.6	32.3	34.7	34.9
85 years and over	58.3	57.4	42.4	50.6	65.9	57.5	47.3	48.5	49.9
Female									
All ages, age-adjusted ⁴	5.6	5.6	7.4	5.7	4.8	4.0	5.0	5.5	5.8
All ages, crude	5.1	4.9	6.6	5.5	4.8	4.0	5.2	5.7	6.0
Under 1 year
1–4 years
5–14 years	0.1	0.1	0.2	0.2	0.4	0.3	0.4	0.7	0.7
15–24 years	2.6	2.2	4.2	4.3	3.9	3.0	3.9	4.5	4.6
15–19 years	1.8	1.6	2.9	3.0	3.7	2.7	3.1	3.9	4.2
20–24 years	3.3	2.9	5.7	5.5	4.1	3.2	4.7	5.2	5.0
25–44 years	6.2	6.6	10.2	7.7	6.2	5.4	6.4	6.8	7.2
25–34 years	4.9	5.5	8.6	7.1	5.6	4.3	5.3	6.1	6.3
35–44 years	7.5	7.7	11.9	8.5	6.8	6.4	7.5	7.6	8.2
45–64 years	9.9	10.2	12.0	8.9	7.1	6.2	8.6	9.4	9.8
45–54 years	9.9	10.2	12.6	9.4	6.9	6.7	9.0	10.0	10.7
55–64 years	9.9	10.2	11.4	8.4	7.3	5.4	8.0	8.7	8.9
65 years and over	9.4	8.4	8.1	6.1	6.4	4.0	4.2	4.6	5.0
65–74 years	10.1	8.4	9.0	6.5	6.7	4.0	4.8	5.4	5.9
75–84 years	8.1	8.9	7.0	5.5	6.3	4.0	3.7	3.9	4.3
85 years and over	8.2	6.0	5.9	5.5	5.4	4.2	3.3	3.3	3.4
White male⁵									
All ages, age-adjusted ⁴	22.3	21.1	20.8	20.9	22.8	19.1	22.0	22.6	23.3
All ages, crude	19.0	17.6	18.0	19.9	22.0	18.8	22.6	23.4	24.2
15–24 years	6.6	8.6	13.9	21.4	23.2	17.9	18.3	18.7	19.9
25–44 years	17.9	18.5	21.5	24.6	25.4	22.9	26.2	26.8	27.5
45–64 years	39.3	36.5	31.9	25.0	26.0	23.2	33.0	33.0	34.0
65 years and over	55.8	46.7	41.1	37.2	44.2	33.3	31.7	34.1	34.7
65–74 years	53.2	42.0	38.7	32.5	34.2	24.3	26.3	28.9	29.5
75–84 years	61.9	55.7	45.5	45.5	60.2	41.1	34.9	38.1	38.3
85 years and over	61.9	61.3	45.8	52.8	70.3	61.6	50.8	52.6	54.4

See footnotes at end of table.

Table 30 (page 2 of 3). Death rates for suicide, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#030>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and age	1950 ^{1,2}	1960 ^{1,2}	1970 ²	1980 ²	1990 ²	2000 ³	2010 ³	2013 ³	2014 ³
Deaths per 100,000 resident population									
Black or African American male ⁵									
All ages, age-adjusted ⁴	7.5	8.4	10.0	11.4	12.8	10.0	9.1	9.3	9.5
All ages, crude	6.3	6.4	8.0	10.3	12.0	9.4	8.7	9.0	9.2
15–24 years	4.9	4.1	10.5	12.3	15.1	14.2	11.1	11.5	12.0
25–44 years	9.8	12.6	16.1	19.2	19.6	14.3	14.5	14.4	14.4
45–64 years	12.7	13.0	12.4	11.8	13.1	9.9	9.5	10.0	9.9
65 years and over	9.0	9.9	8.7	11.4	14.9	11.5	8.3	8.4	8.9
65–74 years	10.0	11.3	8.7	11.1	14.7	11.1	7.6	7.8	7.7
75–84 years	*	*	*	10.5	14.4	12.1	9.9	9.1	11.0
85 years and over	---	*	*	*	*	*	*	*	*
American Indian or Alaska Native male ⁵									
All ages, age-adjusted ⁴	---	---	---	19.3	20.1	16.0	15.5	18.1	16.4
All ages, crude	---	---	---	20.9	20.9	15.9	16.1	17.9	16.0
15–24 years	---	---	---	45.3	49.1	26.2	30.6	29.1	23.5
25–44 years	---	---	---	31.2	27.8	24.5	20.9	26.6	26.2
45–64 years	---	---	---	*	*	15.4	17.8	18.3	15.1
65 years and over	---	---	---	*	*	*	*	*	13.4
Asian or Pacific Islander male ⁵									
All ages, age-adjusted ⁴	---	---	---	10.7	9.6	8.6	9.5	9.1	8.9
All ages, crude	---	---	---	8.8	8.7	7.9	9.3	9.2	9.0
15–24 years	---	---	---	10.8	13.5	9.1	10.9	11.9	12.9
25–44 years	---	---	---	11.0	10.6	9.9	10.6	11.9	9.7
45–64 years	---	---	---	13.0	9.7	9.7	12.8	10.9	12.1
65 years and over	---	---	---	18.6	16.8	15.4	14.9	10.9	11.6
Hispanic or Latino male ^{5,7}									
All ages, age-adjusted ⁴	---	---	---	---	13.7	10.3	9.9	9.3	10.3
All ages, crude	---	---	---	---	11.4	8.4	8.5	8.3	9.2
15–24 years	---	---	---	---	14.7	10.9	10.7	10.1	11.6
25–44 years	---	---	---	---	16.2	11.2	11.2	11.5	12.6
45–64 years	---	---	---	---	16.1	12.0	12.9	11.4	12.4
65 years and over	---	---	---	---	23.4	19.5	15.7	14.2	15.9
White, not Hispanic or Latino male ⁷									
All ages, age-adjusted ⁴	---	---	---	---	23.5	20.2	24.2	25.3	25.9
All ages, crude	---	---	---	---	23.1	20.4	25.7	26.9	27.7
15–24 years	---	---	---	---	24.4	19.5	20.4	21.4	22.4
25–44 years	---	---	---	---	26.4	25.1	30.3	31.2	31.8
45–64 years	---	---	---	---	26.8	24.0	35.4	36.1	37.2
65 years and over	---	---	---	---	45.4	33.9	32.7	35.6	36.1
White female ⁵									
All ages, age-adjusted ⁴	6.0	5.9	7.9	6.1	5.2	4.3	5.6	6.3	6.6
All ages, crude	5.5	5.3	7.1	5.9	5.3	4.4	5.9	6.5	6.9
15–24 years	2.7	2.3	4.2	4.6	4.2	3.1	4.2	5.0	5.0
25–44 years	6.6	7.0	11.0	8.1	6.6	6.0	7.3	7.9	8.4
45–64 years	10.6	10.9	13.0	9.6	7.7	6.9	9.9	10.9	11.5
65 years and over	9.9	8.8	8.5	6.4	6.8	4.3	4.5	5.1	5.5
Black or African American female ⁵									
All ages, age-adjusted ⁴	1.8	2.0	2.9	2.4	2.4	1.8	1.8	2.0	2.1
All ages, crude	1.5	1.6	2.6	2.2	2.3	1.7	1.8	2.0	2.1
15–24 years	1.8	*	3.8	2.3	2.3	2.2	2.0	2.6	2.6
25–44 years	2.3	3.0	4.8	4.3	3.8	2.6	2.8	2.9	2.9
45–64 years	2.7	3.1	2.9	2.5	2.9	2.1	2.1	2.7	2.7
65 years and over	*	*	2.6	*	1.9	1.3	*	*	1.3

See footnotes at end of table.

Table 30 (page 3 of 3). Death rates for suicide, by sex, race, Hispanic origin, and age: United States, selected years 1950–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#030>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and age	1950 ^{1,2}	1960 ^{1,2}	1970 ²	1980 ²	1990 ²	2000 ³	2010 ³	2013 ³	2014 ³
Deaths per 100,000 resident population									
American Indian or Alaska Native female⁵									
All ages, age-adjusted ⁴	---	---	---	4.7	3.6	3.8	6.1	5.3	5.5
All ages, crude	---	---	---	4.7	3.7	4.0	5.9	5.4	5.6
15–24 years	---	---	---	*	*	*	10.4	9.4	9.6
25–44 years	---	---	---	10.7	*	7.2	7.4	7.5	9.7
45–64 years	---	---	---	*	*	*	6.2	6.6	*
65 years and over	---	---	---	*	*	*	*	*	*
Asian or Pacific Islander female⁵									
All ages, age-adjusted ⁴	---	---	---	5.5	4.1	2.8	3.4	3.0	3.4
All ages, crude	---	---	---	4.7	3.4	2.7	3.4	3.1	3.5
15–24 years	---	---	---	*	3.9	2.7	3.5	3.6	4.1
25–44 years	---	---	---	5.4	3.8	3.3	4.1	3.7	4.0
45–64 years	---	---	---	7.9	5.0	3.2	4.7	3.9	4.2
65 years and over	---	---	---	*	8.5	5.2	4.3	3.3	5.2
Hispanic or Latina female^{5,7}									
All ages, age-adjusted ⁴	---	---	---	---	2.3	1.7	2.1	2.3	2.5
All ages, crude	---	---	---	---	2.2	1.5	2.0	2.2	2.4
15–24 years	---	---	---	---	3.1	2.0	3.1	3.3	3.4
25–44 years	---	---	---	---	3.1	2.1	2.4	2.8	3.0
45–64 years	---	---	---	---	2.5	2.5	2.8	3.2	3.5
65 years and over	---	---	---	---	*	*	2.2	1.2	2.1
White, not Hispanic or Latina female⁷									
All ages, age-adjusted ⁴	---	---	---	---	5.4	4.7	6.2	7.1	7.5
All ages, crude	---	---	---	---	5.6	4.9	6.7	7.5	7.9
15–24 years	---	---	---	---	4.3	3.3	4.4	5.4	5.4
25–44 years	---	---	---	---	7.0	6.7	8.6	9.3	9.8
45–64 years	---	---	---	---	8.0	7.3	10.7	12.0	12.6
65 years and over	---	---	---	---	7.0	4.4	4.7	5.4	5.8

... Category not applicable.

* Rates based on fewer than 20 deaths are considered unreliable and are not shown.

--- Data not available.

¹Includes deaths of persons who were not residents of the 50 states and the District of Columbia (D.C.).

²Underlying cause of death was coded according to the 6th Revision of the *International Classification of Diseases* (ICD) in 1950, 7th Revision in 1960, 8th Revision in 1970, and 9th Revision in 1980–1998. See Appendix II, Cause of death; Table III; Table IV.

³Starting with 1999 data, cause of death is coded according to ICD–10. See Appendix II, Cause of death; Comparability ratio; Table IV; Table V.

⁴Age-adjusted rates are calculated using the year 2000 standard population. Prior to 2001, age-adjusted rates were calculated using standard million proportions based on rounded population numbers. Starting with 2001 data, unrounded population numbers are used to calculate age-adjusted rates. See Appendix II, Age adjustment.

⁵The race groups, white, black, Asian or Pacific Islander, and American Indian or Alaska Native, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Death rates for Hispanic, American Indian or Alaska Native, and Asian or Pacific Islander persons should be interpreted with caution because of inconsistencies in reporting Hispanic origin or race on the death certificate (death rate numerators) compared with population figures (death rate denominators). The net effect of misclassification is an underestimation of deaths and death rates for races other than white and black. See Appendix II, Race, for a detailed discussion of sources of bias in death rates by race and Hispanic origin.

⁶In 1950, rate is for the age group 75 years and over.

⁷Prior to 1997, data from states that did not report Hispanic origin on the death certificate were excluded. See Appendix II, Hispanic origin.

NOTES: Starting with *Health, United States, 2003*, rates for 1991–1999 were revised using intercensal population estimates based on the 1990 and 2000 censuses. For 2000, population estimates are bridged-race April 1 census counts. Starting with *Health, United States, 2012*, rates for 2001–2009 were revised using intercensal population estimates based on the 2000 and 2010 censuses. For 2010, population estimates are bridged-race April 1 census counts. Rates for 2011 and beyond were computed using 2010-based postcensal estimates. See Appendix I, Population Census and Population Estimates. Figures for 2001 include September 11-related deaths for which death certificates were filed as of October 24, 2002. See Appendix II, Cause of death; Table IV for terrorism-related ICD–10 codes. Age groups were selected to minimize the presentation of unstable age-specific death rates based on small numbers of deaths and for consistency among comparison groups. For additional injury-related statistics, see Web-based Injury Statistics Query and Reporting System (WISQARS), available from: <http://www.cdc.gov/injury/wisqars/index.html>. Starting with 2003 data, some states allowed the reporting of more than one race on the death certificate. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards, for comparability with other states. See Appendix II, Race. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>. **Data have been revised and differ from previous editions of *Health, United States*.**

SOURCE: CDC/NCHS, National Vital Statistics System; Grove RD, Hetzel AM. Vital statistics rates in the United States, 1940–1960. Washington, DC: U.S. Government Printing Office, 1968; numerator data from National Vital Statistics System, annual public-use Mortality Files; denominator data from national population estimates for race groups from Table 1 and unpublished Hispanic population estimates for 1985–1996 prepared by the Housing and Household Economic Statistics Division, U.S. Census Bureau; Deaths: Final data for 2014. National vital statistics reports (forthcoming). Available from: <http://www.cdc.gov/nchs/products/nvsr.htm>. See Appendix I, National Vital Statistics System (NVSS).

Table 31 (page 1 of 3). Death rates for firearm-related injuries, by sex, race, Hispanic origin, and age: United States, selected years 1970–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#031>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and age	1970 ¹	1980 ¹	1990 ¹	1995 ¹	2000 ²	2010 ²	2013 ²	2014 ²
All persons								
All ages, age-adjusted ³	14.3	14.8	14.6	13.4	10.2	10.1	10.4	10.3
All ages, crude	13.1	14.9	14.9	13.5	10.2	10.3	10.6	10.5
Under 1 year	*	*	*	*	*	*	*	*
1–14 years	1.6	1.4	1.5	1.6	0.7	0.6	0.7	0.8
1–4 years	1.0	0.7	0.6	0.6	0.3	0.4	0.4	0.4
5–14 years	1.7	1.6	1.9	1.9	0.9	0.7	0.8	0.9
15–24 years	15.5	20.6	25.8	26.7	16.8	14.2	14.1	14.0
15–19 years	11.4	14.7	23.3	24.1	12.9	10.6	9.7	9.9
20–24 years	20.3	26.4	28.1	29.2	20.9	17.9	18.1	17.7
25–44 years	20.9	22.5	19.3	16.9	13.1	13.3	13.9	13.4
25–34 years	22.2	24.3	21.8	19.6	14.5	15.0	15.3	14.7
35–44 years	19.6	20.0	16.3	14.3	11.9	11.7	12.3	12.1
45–64 years	17.6	15.2	13.6	11.7	10.0	11.6	11.9	11.8
45–54 years	18.1	16.4	13.9	12.0	10.5	12.0	12.3	12.2
55–64 years	17.0	13.9	13.3	11.3	9.4	11.1	11.5	11.4
65 years and over	13.8	13.5	16.0	14.1	12.2	11.7	12.5	12.7
65–74 years	14.5	13.8	14.4	12.8	10.6	10.7	11.3	11.5
75–84 years	13.4	13.4	19.4	16.3	13.9	12.7	14.1	13.9
85 years and over	10.2	11.6	14.7	14.4	14.2	13.2	13.9	15.0
Male								
All ages, age-adjusted ³	24.8	25.9	26.1	23.8	18.1	17.9	18.3	18.0
All ages, crude	22.2	25.7	26.2	23.6	17.8	18.0	18.5	18.3
Under 1 year	*	*	*	*	*	*	*	*
1–14 years	2.3	2.0	2.2	2.3	1.1	1.0	1.0	1.1
1–4 years	1.2	0.9	0.7	0.8	0.4	0.6	0.6	0.5
5–14 years	2.7	2.5	2.9	2.9	1.4	1.1	1.2	1.3
15–24 years	26.4	34.8	44.7	46.5	29.4	25.0	24.4	24.3
15–19 years	19.2	24.5	40.1	41.6	22.4	18.4	17.0	17.2
20–24 years	35.1	45.2	49.1	51.5	37.0	31.8	31.2	30.8
25–44 years	34.1	38.1	32.6	28.4	22.0	22.9	23.7	22.8
25–34 years	36.5	41.4	37.0	33.2	24.9	26.4	26.3	25.2
35–44 years	31.6	33.2	27.4	23.6	19.4	19.3	20.8	20.1
45–64 years	31.0	25.9	23.4	20.0	17.1	19.9	20.1	19.8
45–54 years	30.7	27.3	23.2	20.1	17.6	20.3	20.4	19.9
55–64 years	31.3	24.5	23.7	19.8	16.3	19.3	19.8	19.8
65 years and over	29.7	29.7	35.3	30.7	26.4	24.1	25.3	25.7
65–74 years	29.5	27.8	28.2	25.1	20.3	20.0	20.9	21.5
75–84 years	31.0	33.0	46.9	37.8	32.2	27.5	29.8	29.2
85 years and over	26.2	34.9	49.3	47.1	44.7	37.4	38.3	40.7
Female								
All ages, age-adjusted ³	4.8	4.7	4.2	3.8	2.8	2.7	3.0	3.0
All ages, crude	4.4	4.7	4.3	3.8	2.8	2.7	3.0	3.0
Under 1 year	*	*	*	*	*	*	*	*
1–14 years	0.8	0.7	0.8	0.8	0.3	0.3	0.4	0.5
1–4 years	0.9	0.5	0.5	0.5	*	0.3	0.3	0.4
5–14 years	0.8	0.7	1.0	0.9	0.4	0.3	0.4	0.5
15–24 years	4.8	6.1	6.0	5.9	3.5	2.9	3.2	3.1
15–19 years	3.5	4.6	5.7	5.6	2.9	2.3	2.0	2.3
20–24 years	6.4	7.7	6.3	6.1	4.2	3.5	4.3	3.9
25–44 years	8.3	7.4	6.1	5.5	4.2	3.8	4.0	4.0
25–34 years	8.4	7.5	6.7	5.8	4.0	3.5	4.1	3.9
35–44 years	8.2	7.2	5.4	5.2	4.4	4.1	4.0	4.1
45–64 years	5.4	5.4	4.5	3.9	3.4	3.7	4.1	4.1
45–54 years	6.4	6.2	4.9	4.2	3.6	3.8	4.5	4.7
55–64 years	4.2	4.6	4.0	3.5	3.0	3.4	3.8	3.6
65 years and over	2.4	2.5	3.1	2.8	2.2	2.2	2.4	2.5
65–74 years	2.8	3.1	3.6	3.0	2.5	2.6	2.8	2.7
75–84 years	1.7	1.7	2.9	2.8	2.0	2.1	2.3	2.4
85 years and over	*	1.3	1.3	1.8	1.7	1.5	1.5	1.6
White male ⁴								
All ages, age-adjusted ³	19.7	22.1	22.0	20.1	15.9	16.1	16.5	16.3
All ages, crude	17.6	21.8	21.8	19.9	15.6	16.5	17.1	17.0
1–14 years	1.8	1.9	1.9	1.9	1.0	0.8	1.0	1.0
15–24 years	16.9	28.4	29.5	30.8	19.6	16.2	16.0	16.1
25–44 years	24.2	29.5	25.7	23.2	18.0	18.6	18.9	18.4
25–34 years	24.3	31.1	27.8	25.2	18.1	19.1	19.1	18.7
35–44 years	24.1	27.1	23.3	21.2	17.9	18.0	18.8	18.2
45–64 years	27.4	23.3	22.8	19.5	17.4	21.3	21.6	21.4
65 years and over	29.9	30.1	36.8	32.2	28.2	26.5	28.0	28.3

See footnotes at end of table.

Table 31 (page 2 of 3). Death rates for firearm-related injuries, by sex, race, Hispanic origin, and age: United States, selected years 1970–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#031>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and age	1970 ¹	1980 ¹	1990 ¹	1995 ¹	2000 ²	2010 ²	2013 ²	2014 ²
Deaths per 100,000 resident population								
Black or African American male ⁴								
All ages, age-adjusted ³	70.8	60.1	56.3	49.2	34.2	31.8	32.1	31.5
All ages, crude	60.8	57.7	61.9	52.9	36.1	33.4	33.5	32.9
1–14 years	5.3	3.0	4.4	4.4	1.8	1.9	1.4	1.9
15–24 years	97.3	77.9	138.0	138.7	89.3	73.2	69.9	68.9
25–44 years	126.2	114.1	90.3	70.2	54.1	57.3	60.2	57.4
25–34 years	145.6	128.4	108.6	92.3	74.8	78.2	76.6	71.3
35–44 years	104.2	92.3	66.1	46.3	34.3	35.2	41.5	41.0
45–64 years	71.1	55.6	34.5	28.3	18.4	16.5	16.7	16.9
65 years and over	30.6	29.7	23.9	21.8	13.8	9.4	8.8	10.9
American Indian or Alaska Native male ⁴								
All ages, age-adjusted ³	---	24.0	19.4	19.4	13.1	11.7	12.9	12.9
All ages, crude	---	27.5	20.5	20.9	13.2	12.5	12.5	12.4
15–24 years	---	55.3	49.1	40.9	26.9	26.0	18.4	17.7
25–44 years	---	43.9	25.4	31.2	16.6	16.9	18.8	19.3
45–64 years	---	*	*	14.2	12.2	11.1	13.5	12.6
65 years and over	---	*	*	*	*	*	*	*
Asian or Pacific Islander male ⁴								
All ages, age-adjusted ³	---	7.8	8.8	9.2	6.0	4.2	4.1	3.7
All ages, crude	---	8.2	9.4	10.0	6.2	4.4	4.3	3.8
15–24 years	---	10.8	21.0	24.3	9.3	6.8	5.9	6.3
25–44 years	---	12.8	10.9	10.6	8.1	6.0	6.4	4.8
45–64 years	---	10.4	8.1	8.2	7.4	4.4	4.6	4.1
65 years and over	---	*	*	*	*	3.9	*	3.4
Hispanic or Latino male ^{4,5}								
All ages, age-adjusted ³	---	---	27.6	23.8	13.6	10.5	9.4	9.4
All ages, crude	---	---	29.9	26.2	14.2	10.5	9.4	9.4
1–14 years	---	---	2.6	2.8	1.0	0.6	0.5	0.5
15–24 years	---	---	55.5	61.7	30.8	20.9	17.4	17.2
25–44 years	---	---	42.7	31.4	17.3	14.4	13.8	14.1
25–34 years	---	---	47.3	36.4	20.3	18.0	16.2	15.7
35–44 years	---	---	35.4	24.2	13.2	10.2	11.1	12.3
45–64 years	---	---	21.4	17.2	12.0	9.1	8.2	7.9
65 years and over	---	---	19.1	16.5	12.2	9.9	8.1	7.8
White, not Hispanic or Latino male ⁵								
All ages, age-adjusted ³	---	---	20.6	18.6	15.5	16.6	17.5	17.3
All ages, crude	---	---	20.4	18.5	15.7	17.6	18.8	18.7
1–14 years	---	---	1.6	1.6	1.0	0.9	1.2	1.2
15–24 years	---	---	24.1	23.5	16.2	14.2	15.1	15.4
25–44 years	---	---	23.3	21.4	17.9	19.4	20.2	19.4
25–34 years	---	---	24.7	22.5	17.2	18.9	19.7	19.2
35–44 years	---	---	21.6	20.4	18.4	19.9	20.8	19.7
45–64 years	---	---	22.7	19.5	17.8	22.8	23.5	23.4
65 years and over	---	---	37.4	32.5	29.0	27.6	29.6	29.9
White female ⁴								
All ages, age-adjusted ³	4.0	4.2	3.8	3.5	2.7	2.7	3.0	3.1
All ages, crude	3.7	4.1	3.8	3.5	2.7	2.8	3.1	3.2
15–24 years	3.4	5.1	4.8	4.5	2.8	2.3	2.7	2.6
25–44 years	6.9	6.2	5.3	4.9	3.9	3.7	3.9	4.0
45–64 years	5.0	5.1	4.5	4.0	3.5	4.1	4.7	4.7
65 years and over	2.2	2.5	3.1	2.8	2.4	2.5	2.7	2.7

See footnotes at end of table.

Table 31 (page 3 of 3). Death rates for firearm-related injuries, by sex, race, Hispanic origin, and age: United States, selected years 1970–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#031>.

[Data are based on death certificates]

Sex, race, Hispanic origin, and age	1970 ¹	1980 ¹	1990 ¹	1995 ¹	2000 ²	2010 ²	2013 ²	2014 ²
Deaths per 100,000 resident population								
Black or African American female ⁴								
All ages, age-adjusted ³	11.1	8.7	7.3	6.2	3.9	3.3	3.4	3.2
All ages, crude.	10.0	8.8	7.8	6.5	4.0	3.3	3.4	3.2
15–24 years	15.2	12.3	13.3	13.2	7.6	6.4	6.1	6.2
25–44 years	19.4	16.1	12.4	9.8	6.5	5.6	6.0	5.3
45–64 years	10.2	8.2	4.8	4.1	3.1	2.2	2.5	2.1
65 years and over	4.3	3.1	3.1	2.6	1.3	*	0.9	1.4
American Indian or Alaska Native female ⁴								
All ages, age-adjusted ³	---	5.8	3.3	3.8	2.9	2.6	2.0	2.4
All ages, crude.	---	5.8	3.4	4.1	2.9	2.4	2.0	2.4
15–24 years	---	*	*	*	*	*	*	*
25–44 years	---	10.2	*	7.0	5.5	3.7	*	5.6
45–64 years	---	*	*	*	*	*	*	*
65 years and over	---	*	*	*	*	*	*	*
Asian or Pacific Islander female ⁴								
All ages, age-adjusted ³	---	2.0	1.9	2.0	1.1	0.6	0.9	0.7
All ages, crude.	---	2.1	2.1	2.1	1.2	0.6	0.9	0.8
15–24 years	---	*	*	3.9	*	*	*	*
25–44 years	---	3.2	2.7	2.7	1.5	1.1	1.5	0.8
45–64 years	---	*	*	*	*	*	0.8	1.3
65 years and over	---	*	*	*	*	*	*	*
Hispanic or Latina female ^{4,5}								
All ages, age-adjusted ³	---	---	3.3	3.1	1.8	1.3	1.3	1.4
All ages, crude.	---	---	3.6	3.3	1.8	1.3	1.3	1.4
15–24 years	---	---	6.9	6.1	2.9	2.1	2.3	2.6
25–44 years	---	---	5.1	4.7	2.5	1.8	2.0	1.9
45–64 years	---	---	2.4	2.4	2.2	1.5	1.2	1.2
65 years and over	---	---	*	*	*	*	*	*
White, not Hispanic or Latina female ⁵								
All ages, age-adjusted ³	---	---	3.7	3.4	2.8	3.0	3.3	3.4
All ages, crude.	---	---	3.7	3.5	2.9	3.1	3.5	3.6
15–24 years	---	---	4.3	4.1	2.7	2.3	2.8	2.6
25–44 years	---	---	5.1	4.8	4.2	4.2	4.4	4.6
45–64 years	---	---	4.6	4.1	3.6	4.4	5.2	5.2
65 years and over	---	---	3.2	2.8	2.4	2.6	2.9	2.9

* Rates based on fewer than 20 deaths are considered unreliable and are not shown.

--- Data not available.

¹Underlying cause of death was coded according to the 8th Revision of the *International Classification of Diseases* (ICD) in 1970 and 9th Revision in 1980–1998. See Appendix II, Cause of death; Table III; Table IV.

²Starting with 1999 data, cause of death is coded according to ICD–10. See Appendix II, Cause of death; Comparability ratio; Table IV; Table V.

³Age-adjusted rates are calculated using the year 2000 standard population. Prior to 2001, age-adjusted rates were calculated using standard million proportions based on rounded population numbers. Starting with 2001 data, unrounded population numbers are used to calculate age-adjusted rates. See Appendix II, Age adjustment.

⁴The race groups, white, black, Asian or Pacific Islander, and American Indian or Alaska Native, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Death rates for Hispanic, American Indian or Alaska Native, and Asian or Pacific Islander persons should be interpreted with caution because of inconsistencies in reporting Hispanic origin or race on the death certificate (death rate numerators) compared with population figures (death rate denominators). The net effect of misclassification is an underestimation of deaths and death rates for races other than white and black. See Appendix II, Race, for a detailed discussion of sources of bias in death rates by race and Hispanic origin.

⁵Prior to 1997, data from states that did not report Hispanic origin on the death certificate were excluded. See Appendix II, Hispanic origin.

NOTES: Starting with *Health, United States, 2003*, rates for 1991–1999 were revised using intercensal population estimates based on the 1990 and 2000 censuses. For 2000, population estimates are bridged-race April 1 census counts. Starting with *Health, United States, 2012*, rates for 2001–2009 were revised using intercensal population estimates based on the 2000 and 2010 censuses. For 2010, population estimates are bridged-race April 1 census counts. Rates for 2011 and beyond were computed using 2010-based postcensal estimates. See Appendix I, Population Census and Population Estimates. Age groups were selected to minimize the presentation of unstable age-specific death rates based on small numbers of deaths and for consistency among comparison groups. For additional injury-related statistics, see Web-based Injury Statistics Query and Reporting System (WISQARS), available from: <http://www.cdc.gov/injury/wisqars/index.html>. Starting with 2003 data, some states allowed the reporting of more than one race on the death certificate. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards, for comparability with other states. See Appendix II, Race. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>. **Data have been revised and differ from previous editions of *Health, United States*.**

SOURCE: CDC/NCHS, National Vital Statistics System; numerator data from National Vital Statistics System, annual public-use Mortality Files; denominator data from national population estimates for race groups from Table 1 and unpublished Hispanic population estimates for 1985–1996 prepared by the Housing and Household Economic Statistics Division, U.S. Census Bureau; Deaths: Final data for 2014. National vital statistics reports (forthcoming). Available from: <http://www.cdc.gov/nchs/products/nvsr.htm>. See Appendix I, National Vital Statistics System (NVSS).

Table 32 (page 1 of 2). Occupational fatal injuries, by industry, sex, age, race, and Hispanic origin: United States, selected years 2003–2013

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2015.htm#032>.

[Data are compiled from various federal, state, and local administrative sources]

Characteristic	2003	2005	2009	2010	2011	2012	2013
	Number of deaths						
Total workforce	5,575	5,734	4,551	4,690	4,693	4,628	4,585
Sex							
Male	5,129	5,328	4,216	4,322	4,308	4,277	4,265
Female	446	406	335	368	385	351	319
Unspecified	0	0	0	0	0	0	1
Age							
Under 16 years	25	23	13	16	10	19	5
16–17 years	28	31	14	18	13	10	9
18–19 years	84	111	57	56	61	59	57
20–24 years	462	403	275	245	292	287	279
25–34 years	1,018	1,017	704	785	714	736	777
35–44 years	1,329	1,243	908	868	875	829	853
45–54 years	1,301	1,389	1,173	1,169	1,222	1,161	1,115
55–64 years	802	933	853	948	936	936	933
65 years and over	523	578	551	582	569	588	557
Unspecified	3	6	3	3	1	3	0
Race and Hispanic origin							
Hispanic or Latino	794	923	713	707	749	748	817
Not Hispanic or Latino	4,781	4,811	3,838	3,983	3,944	3,880	3,768
White	3,988	3,977	3,204	3,363	3,323	3,177	3,125
Black or African American	543	584	421	412	440	486	439
American Indian or Alaska Native	42	50	33	32	30	37	35
Asian	147	154	141	143	121	147	125
Native Hawaiian or Other Pacific Islander	11	9	7	6	3	7	7
Multiple races	3	*	7	8	15	5	12
Other races or not reported	47	35	25	19	12	21	25
Industry ¹							
Private sector	5,043	5,214	4,090	4,206	4,188	4,175	4,101
Agriculture, forestry, fishing, and hunting	709	715	575	621	566	509	500
Mining ²	141	159	99	172	155	181	155
Utilities	32	30	16	26	39	23	24
Construction	1,131	1,192	834	774	738	806	828
Manufacturing	420	393	319	329	327	327	312
Wholesale trade	191	209	190	191	190	204	201
Retail trade	344	400	307	311	268	273	263
Transportation and warehousing	808	885	633	661	749	741	733
Information	64	65	33	43	56	42	40
Finance and insurance	45	42	33	24	36	21	21
Real estate and rental and leasing	84	57	75	89	62	64	66
Professional and technical services	97	83	85	76	74	57	87
Management, administrative, and waste services ³	---	---	---	---	359	352	343
Educational services	41	46	27	30	37	34	32
Health care and social assistance	102	104	123	141	117	107	103
Arts, entertainment, and recreation	88	77	80	84	93	80	69
Accommodation and food services	187	136	151	154	138	152	138
Other services (except public administration)	194	210	173	192	183	199	186
Government ⁴	532	520	461	484	505	453	484

See footnotes at end of table.

Table 32 (page 2 of 2). Occupational fatal injuries, by industry, sex, age, race, and Hispanic origin: United States, selected years 2003–2013

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2015.htm#032>.

[Data are compiled from various federal, state, and local administrative sources]

-- Data not available.

* Estimates are unreliable or data do not meet publication criteria.

¹Industry data from 2003 to 2008 (shown in spreadsheet version) are based on the North American Industry Classification System (NAICS), 2002. Industry data from 2009 to the present are based on NAICS 2007. NAICS replaces the Standard Industrial Classification (SIC) system. Because of substantial differences between NAICS and SIC, industry data classified by these two systems are not comparable.

²Includes fatal injuries at all establishments categorized as Mining (Sector 21) in the NAICS, including establishments not governed by the Mine Safety and Health Administration (MSHA) rules and reporting, such as those in Oil and Gas Extraction.

³Starting with 2011 data, CFOI combined the categories "Management of companies and enterprises" and "Administrative and support and waste management and remediation services" into one category entitled "Management, administrative, and waste services."

⁴Includes fatal work injuries to workers employed by governmental organizations, regardless of industry.

NOTES: Fatal work injuries are based on revised data and may differ from originally published data from CFOI. See Appendix I, Census of Fatal Occupational Injuries (CFOI). Private sector totals include injuries with unknown industry. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: Department of Labor, Bureau of Labor Statistics, Census of Fatal Occupational Injuries. Revised annual data. See Appendix I, Census of Fatal Occupational Injuries (CFOI).

Table 33 (page 1 of 2). Selected notifiable disease rates and number of new cases: United States, selected years 1950–2013

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#033>.

[Data are based on reporting by state health departments]

Disease	1950	1960	1970	1980	1990	2000	2003	2010	2013
New cases per 100,000 population									
Acute hepatitis A viral infection	---	---	27.87	12.84	12.64	4.91	2.66	0.54	0.57
Acute hepatitis B viral infection	---	---	4.08	8.39	8.48	2.95	2.61	1.10	0.97
Diphtheria	3.83	0.51	0.21	0.00	0.00	0.00	0.00	—	—
<i>Haemophilus influenzae</i> , invasive	---	---	---	---	---	0.51	0.70	1.03	1.21
Lyme disease ¹	---	---	---	---	---	6.53	7.39	9.86	11.62
Measles (Rubeola)	211.01	245.42	23.23	5.96	11.17	0.03	0.02	0.02	0.06
Meningococcal disease	---	---	1.23	1.25	0.99	0.83	0.61	0.27	0.18
Mumps	---	---	55.55	3.86	2.17	0.13	0.08	0.85	0.19
Pertussis (whooping cough)	79.82	8.23	2.08	0.76	1.84	2.88	4.04	8.97	9.12
Poliomyelitis, paralytic ²	---	1.40	0.02	0.00	0.00	—	—	—	0.00
Rubella (German measles)	---	---	27.75	1.72	0.45	0.06	0.00	0.00	0.00
Salmonellosis, excluding typhoid fever	---	3.85	10.84	14.88	19.54	14.51	15.16	17.73	16.13
Shigellosis	15.45	6.94	6.79	8.41	10.89	8.41	8.19	4.82	4.06
Spotted fever rickettsiosis ³	---	---	0.19	0.52	0.26	0.18	0.38	0.65	1.08
Tuberculosis ⁴	---	30.83	18.28	12.25	10.33	6.01	5.17	3.64	3.05
Sexually transmitted diseases: ⁵									
Syphilis ⁶	146.02	68.78	44.80	30.30	54.32	11.20	11.79	14.93	17.87
Primary and secondary	16.73	9.06	10.80	12.00	20.26	2.12	2.47	4.49	5.50
Early latent	39.71	10.11	8.00	8.90	22.19	3.35	2.88	4.43	5.36
Late and late latent ⁷	70.22	45.91	24.70	9.20	10.32	5.53	6.30	5.89	6.90
Congenital ⁸	368.30	103.70	52.30	7.70	92.95	14.29	10.56	8.73	9.11
Chlamydia ⁹	---	---	---	---	160.19	251.38	301.74	426.01	443.46
Gonorrhea ¹⁰	192.50	145.40	294.20	442.10	276.43	128.67	115.23	100.76	105.34
Chancroid	3.34	0.94	0.70	0.30	1.69	0.03	0.02	0.01	0.00
Number of new cases									
Acute hepatitis A viral infection	---	---	56,797	29,087	31,441	13,397	7,653	1,670	1,781
Acute hepatitis B viral infection	---	---	8,310	19,015	21,102	8,036	7,526	3,374	3,050
Diphtheria	5,796	918	435	3	4	1	1	—	—
<i>Haemophilus influenzae</i> , invasive	---	---	---	---	---	1,398	2,013	3,151	3,792
Lyme disease ¹	---	---	---	---	---	17,730	21,273	30,158	36,307
Measles (Rubeola)	319,124	441,703	47,351	13,506	27,786	86	56	63	187
Meningococcal disease	---	---	2,505	2,840	2,451	2,256	1,756	833	556
Mumps	---	---	104,953	8,576	5,292	338	231	2,612	584
Pertussis (whooping cough)	120,718	14,809	4,249	1,730	4,570	7,867	11,647	27,550	28,639
Poliomyelitis, paralytic ²	---	2,525	31	4	6	—	—	—	1
Rubella (German measles)	---	---	56,552	3,904	1,125	176	7	5	9
Salmonellosis, excluding typhoid fever	---	6,929	22,096	33,715	48,603	39,574	43,657	54,424	50,634
Shigellosis	23,367	12,487	13,845	19,041	27,077	22,922	23,581	14,786	12,729
Spotted fever rickettsiosis ³	464	---	380	1,163	651	495	1,091	1,985	3,359
Tuberculosis ⁴	---	55,494	37,137	27,749	25,701	16,377	14,874	11,182	9,582
Sexually transmitted diseases: ⁵									
Syphilis ⁶	217,558	122,538	91,382	68,832	135,590	31,618	34,289	45,844	56,482
Primary and secondary	23,939	16,145	21,982	27,204	50,578	5,979	7,177	13,774	17,375
Early latent	59,256	18,017	16,311	20,297	55,397	9,465	8,361	13,604	16,929
Late and late latent ⁷	113,569	81,798	50,348	20,979	25,750	15,594	18,319	18,079	21,819
Congenital ⁸	13,377	4,416	1,953	277	3,865	580	432	387	359
Chlamydia ⁹	---	---	---	---	323,663	709,452	877,478	1,307,893	1,401,906
Gonorrhea ¹⁰	286,746	258,933	600,072	1,004,029	690,042	363,136	335,104	309,341	333,004
Chancroid	4,977	1,680	1,416	788	4,212	78	54	24	10

See footnotes at end of table.

Table 33 (page 2 of 2). Selected notifiable disease rates and number of new cases: United States, selected years 1950–2013

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#033>.

[Data are based on reporting by state health departments]

0.00 Rate more than zero but less than 0.005.

– Quantity zero.

- - - Data not available.

¹National surveillance case definition revised in 2008; probable cases not previously reported.

²Cases of vaccine-associated paralytic poliomyelitis caused by polio vaccine virus.

³Prior to 2010 data, cases of spotted fever rickettsiosis were reported as Rocky Mountain spotted fever (RMSF). Because serologic tests commonly used to diagnose RMSF exhibit cross-reactivity between spotted fever rickettsial pathogens, some cases reported as RMSF might actually be disease caused by other spotted fever rickettsial infections, and therefore are more correctly referred to as spotted fever rickettsiosis starting with 2010 data.

⁴Case reporting for tuberculosis began in 1953. Data prior to 1975 are not comparable with subsequent years because of changes in reporting criteria effective in 1975.

⁵For 1950, data for Alaska and Hawaii were not included. Starting with 1991, data include both civilian and military cases. Cases and rates shown do not include U.S. outlying areas of Guam, Puerto Rico, and the Virgin Islands.

⁶Includes stage of syphilis not stated.

⁷Includes cases of unknown duration.

⁸Rates include all cases of congenitally acquired syphilis per 100,000 live births. Cases of congenitally acquired syphilis were reported through 1994. Starting with 1995 data, only congenital syphilis for cases under 1 year of age were reported. See STD Surveillance Report for congenital syphilis rates.

⁹Prior to 1994, chlamydia was not notifiable. In 1994–1999, cases for New York were exclusively reported by New York City. Starting with 2000 data, cases for New York include the entire state.

¹⁰Data for 1994 do not include cases from Georgia.

NOTES: The total resident population was used to calculate all rates except for sexually transmitted disease rates prior to 1991; STD rates prior to 1991 used the civilian resident population. See Appendix I, Sexually Transmitted Disease (STD) Surveillance; Population Census and Population Estimates. Population data from states where diseases were not notifiable or not available were excluded from the rate calculation; see Appendix II, Notifiable disease. See Appendix I, National Notifiable Disease Surveillance System (NNDSS), for information on underreporting of notifiable diseases. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC, Office of Public Health Scientific Services (OPHSS); Center for Surveillance, Epidemiology and Laboratory Services (CELS); Division of Health Informatics and Surveillance (DHIS). MMWR 2015;62(53):1–119 and CDC. Available from: http://www.cdc.gov/mmwr/mmwr_nd/index.html. Sexually transmitted disease surveillance, 2014. Atlanta, GA: U.S. Department of Health and Human Services, 2015. Available from: <http://www.cdc.gov/std/stats/>. See Appendix I, National Notifiable Diseases Surveillance System (NNDSS); Sexually Transmitted Disease (STD) Surveillance.

Table 34 (page 1 of 2). Human immunodeficiency virus (HIV) diagnoses, by year of diagnosis and selected characteristics: United States, 2010–2014

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2015.htm#034>.

[Data are based on reporting by 50 states and the District of Columbia]

Sex, age at diagnosis, Hispanic origin and race, and region of residence	Year of diagnosis				
	2010 ¹	2011 ¹	2012 ¹	2013 ¹	2014 ¹
	Estimated number of HIV diagnoses ²				
All persons ³	44,940	43,510	43,165	42,566	44,073
Male, 13 years and over	34,871	34,146	34,259	34,034	35,571
Female, 13 years and over	9,831	9,166	8,656	8,340	8,328
Age at diagnosis					
Under 13 years	238	198	250	191	174
13–14 years	43	45	46	43	35
15–19 years	2,118	2,068	1,964	1,792	1,828
20–24 years	7,245	7,311	7,489	7,483	7,868
25–29 years	6,520	6,563	6,777	7,151	7,870
30–34 years	5,639	5,455	5,729	5,574	6,026
35–39 years	5,171	4,622	4,374	4,288	4,662
40–44 years	5,361	4,971	4,646	4,257	4,196
45–49 years	4,972	4,758	4,527	4,268	4,021
50–54 years	3,602	3,487	3,377	3,235	3,242
55–59 years	2,132	2,072	2,019	2,184	2,166
60–64 years	1,091	1,111	1,106	1,170	1,069
65 years and over	810	848	861	930	914
Hispanic origin and race ⁴					
Not Hispanic or Latino:					
White	12,135	11,738	11,752	11,581	12,025
Black or African American	20,987	20,064	19,581	19,252	19,540
American Indian or Alaska Native	177	163	193	186	222
Asian	727	802	848	859	1,046
Native Hawaiian or Other Pacific Islander	58	58	60	56	58
Multiple race	1,565	1,456	1,358	1,246	982
Hispanic or Latino ⁵	9,291	9,230	9,372	9,386	10,201
Region of residence					
Northeast	8,597	8,087	8,039	7,750	7,953
Midwest	5,664	5,580	5,717	5,654	5,529
South	22,550	22,079	21,480	21,508	22,196
West	8,129	7,764	7,929	7,654	8,395

See footnotes at end of table.

Table 34 (page 2 of 2). Human immunodeficiency virus (HIV) diagnoses, by year of diagnosis and selected characteristics: United States, 2010–2014

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2015.htm#034>.

[Data are based on reporting by 50 states and the District of Columbia]

Sex, age at diagnosis, Hispanic origin and race, and region of residence	Year of diagnosis				
	2010 ¹	2011 ¹	2012 ¹	2013 ¹	2014 ¹
Estimated number of HIV diagnoses per 100,000 resident population ²					
All persons	14.5	14.0	13.7	13.4	13.8
Male, 13 years and over	27.9	27.0	26.9	26.4	27.4
Female, 13 years and over	7.5	6.9	6.5	6.2	6.1
Age at diagnosis					
Under 13 years	0.4	0.4	0.5	0.4	0.3
13–14 years	0.5	0.5	0.6	0.5	0.4
15–19 years	9.6	9.5	9.2	8.5	8.7
20–24 years	33.4	33.0	33.1	32.7	34.3
25–29 years	30.8	30.8	31.7	33.1	35.8
30–34 years	28.1	26.6	27.4	26.2	28.0
35–39 years	25.8	23.6	22.4	21.8	23.4
40–44 years	25.6	23.6	22.1	20.4	20.4
45–49 years	22.0	21.5	20.8	20.1	19.3
50–54 years	16.1	15.4	14.9	14.3	14.4
55–59 years	10.8	10.2	9.7	10.3	10.1
60–64 years	6.4	6.2	6.2	6.5	5.8
65 years and over	2.0	2.0	2.0	2.1	2.0
Hispanic origin and race ⁴					
Not Hispanic or Latino:					
White	6.1	5.9	5.9	5.9	6.1
Black or African American	55.2	52.3	50.5	49.2	49.4
American Indian or Alaska Native	7.8	7.1	8.4	8.0	9.5
Asian	4.9	5.3	5.4	5.3	6.2
Native Hawaiian or Other Pacific Islander	11.7	11.4	11.6	10.6	10.6
Multiple race	27.7	25.0	22.6	20.1	15.4
Hispanic or Latino ⁵	18.3	17.8	17.7	17.3	18.4
Region of residence					
Northeast	15.5	14.5	14.4	13.8	14.2
Midwest	8.5	8.3	8.5	8.4	8.2
South	19.6	19.0	18.3	18.1	18.5
West	11.3	10.7	10.8	10.3	11.2

¹Based on diagnoses that occurred during 2010–2014 that were reported to CDC through July 31, 2015. The term diagnosis of HIV infection is defined as a diagnosis of HIV infection regardless of the stage of disease (stage 0, 1, 2, 3 [AIDS], or unknown) and refers to all persons with a diagnosis of HIV infection. In 2014, the criteria used to define HIV diagnoses changed. Cases diagnosed before 2014 were classified according to the 2008 HIV case definition and cases diagnosed in 2014 were classified according to the 2014 HIV case definition. Because of the change in case definition, HIV diagnoses prior to 2014 are not strictly comparable to HIV diagnoses for 2014. The vertical line in the table represents the discontinuity in the HIV diagnosis trend. See Appendix II, Human immunodeficiency virus (HIV) disease for discussion of HIV diagnoses reporting definitions and other issues affecting interpretation of trends.

²Numbers and rates are point estimates that result from statistical adjustments for reporting delays and missing risk factor information. The estimates do not include adjustments for incomplete reporting. See Appendix I, National HIV Surveillance System.

³All persons totals were calculated independent of values for subpopulations. Consequently, sums of subpopulations may not equal totals for all persons.

⁴Hispanic origin and race categories are mutually exclusive.

⁵Persons of Hispanic origin may be of any race. See Appendix II, Hispanic origin.

NOTES: Data shown are for the 50 states and the District of Columbia, and include newly diagnosed and reported cases. This table does not present HIV incidence or prevalence data. Rates for 2010–2014 were calculated using vintage 2014 population estimates from the U.S. Census Bureau. Variations in trends among subpopulations may be due to differences in testing behaviors, targeted HIV testing initiatives, or the numbers of new HIV infections in some subpopulations. Caution should be used when interpreting data on diagnoses of HIV infection. HIV surveillance reports may not be representative of all persons with HIV for several reasons: not all infected persons have been tested and diagnosed; results of anonymous tests are not reported to the name-based HIV registries of state and local health departments; testing patterns are influenced by the extent to which testing is routinely offered to specific groups; and surveillance and reporting practices among jurisdictions differ. The data presented here are estimates of the minimum number of persons for whom HIV infection has been diagnosed and reported to the surveillance system. Data have been revised and differ from previous editions of *Health, United States*.

SOURCE: CDC, National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention. Division of HIV/AIDS Prevention. HIV Surveillance Report. Diagnoses of HIV infection in the United States and Dependent Areas, 2014 (vol. 26). Atlanta, GA: U.S. Department of Health and Human Services, CDC. Available from: <http://www.cdc.gov/hiv/pdf/library/reports/surveillance/cdc-hiv-surveillance-report-us.pdf>. See Appendix I, National HIV Surveillance System.

Table 35 (page 1 of 5). Health conditions among children under age 18, by selected characteristics: United States, average annual, selected years 1997–1999 through 2012–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#035>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Current asthma ¹				Asthma attack in the past 12 months ²			
	1997–1999	2000–2002	2003–2005	2012–2014	1997–1999	2000–2002	2003–2005	2012–2014
Percent of children								
Under 18 years ³	---	---	8.7	8.7	5.4	5.7	5.4	4.9
Age								
0–4 years	---	---	6.1	4.6	4.3	4.7	4.2	3.0
5–17 years	---	---	9.6	10.3	5.7	6.1	5.8	5.6
5–9 years	---	---	9.1	9.9	5.6	6.3	6.1	6.2
10–17 years	---	---	9.9	10.5	5.8	5.9	5.7	5.3
Sex								
Male	---	---	9.9	9.8	6.2	6.6	6.3	5.7
Female	---	---	7.3	7.6	4.5	4.7	4.4	4.1
Race ⁴								
White only	---	---	7.7	7.7	5.0	5.2	4.9	4.4
Black or African American only	---	---	13.0	14.2	7.0	8.0	7.6	7.8
American Indian or Alaska Native only	---	---	12.2	10.2	6.4	*8.7	*6.1	*5.9
Asian only	---	---	4.8	5.1	4.3	4.7	3.3	2.9
Native Hawaiian or Other Pacific Islander only	---	---	*	*	---	*	*	*
2 or more races	---	---	13.5	11.5	---	7.3	8.8	6.3
Hispanic origin and race ⁴								
Hispanic or Latino	---	---	7.6	8.2	4.8	4.2	4.6	4.4
Not Hispanic or Latino	---	---	8.9	8.9	5.5	6.0	5.6	5.1
White only	---	---	7.9	7.7	5.1	5.5	5.0	4.4
Black or African American only	---	---	13.0	14.3	7.0	7.9	7.5	7.9
Percent of poverty level ⁵								
Below 100%	---	---	10.4	11.7	6.1	7.1	6.5	6.6
100%–199%	---	---	8.6	8.5	5.3	5.4	5.2	4.6
200%–399%	---	---	8.3	8.3	5.0	5.3	5.2	4.4
400% or more	---	---	7.9	6.9	5.2	5.5	4.9	4.2
Health insurance status at the time of interview ⁶								
Insured	---	---	9.0	8.9	5.6	5.9	5.6	5.0
Private	---	---	8.0	7.7	5.0	5.3	5.0	4.4
Medicaid	---	---	11.4	10.6	7.7	7.7	7.1	5.9
Uninsured	---	---	5.6	6.9	3.9	4.3	3.3	3.8

See footnotes at end of table.

Table 35 (page 2 of 5). Health conditions among children under age 18, by selected characteristics: United States, average annual, selected years 1997–1999 through 2012–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#035>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Attention-deficit/hyperactivity disorder ⁷				Serious emotional or behavioral difficulties ⁸			
	1997–1999	2000–2002	2003–2005	2012–2014	1997–1999	2000–2002	2003–2005	2012–2014
Age								
Percent of children								
5–17 years ³	6.5	7.5	7.6	10.2	---	---	5.1	5.4
5–9 years	4.8	5.2	5.6	7.7	---	---	4.3	5.1
10–17 years	7.6	9.0	8.9	11.8	---	---	5.6	5.6
Sex								
Male	9.6	10.8	10.7	14.1	---	---	6.1	6.9
Female	3.2	4.2	4.4	6.2	---	---	4.1	3.9
Race ⁴								
White only	7.1	8.1	7.8	10.7	---	---	5.1	5.5
Black or African American only	5.0	7.0	7.7	9.5	---	---	5.3	5.4
American Indian or Alaska Native only	*8.5	*	*9.4	*9.1	---	---	*	*
Asian only	*1.7	*	*1.6	2.1	---	---	*1.7	*1.8
Native Hawaiian or Other Pacific Islander only	---	*	*	*	---	---	*	*
2 or more races	---	7.4	9.7	13.8	---	---	8.2	8.7
Hispanic origin and race ⁴								
Hispanic or Latino	3.6	4.2	4.6	6.4	---	---	3.8	4.2
Not Hispanic or Latino	7.0	8.2	8.3	11.4	---	---	5.4	5.8
White only	7.7	9.0	8.8	12.5	---	---	5.6	6.1
Black or African American only	5.0	6.8	7.5	9.6	---	---	5.2	5.3
Percent of poverty level ⁵								
Below 100%	7.2	8.2	8.4	12.8	---	---	7.4	8.1
100%–199%	6.7	7.5	7.8	10.6	---	---	5.4	5.8
200%–399%	6.2	7.7	7.8	8.9	---	---	4.9	4.5
400% or more	6.1	7.1	6.9	9.3	---	---	3.7	4.0
Health insurance status at the time of interview ⁶								
Insured	6.7	7.8	7.8	10.6	---	---	5.2	5.6
Private	5.9	7.0	7.0	8.7	---	---	4.1	4.0
Medicaid	10.5	10.7	10.3	13.3	---	---	8.5	8.0
Uninsured	4.8	5.4	6.1	5.9	---	---	4.6	3.6

See footnotes at end of table.

Table 35 (page 3 of 5). Health conditions among children under age 18, by selected characteristics: United States, average annual, selected years 1997–1999 through 2012–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#035>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Food allergy ⁹				Skin allergy ¹⁰			
	1997–1999	2000–2002	2003–2005	2012–2014	1997–1999	2000–2002	2003–2005	2012–2014
	Percent of children							
Under 18 years ³	3.4	3.6	3.8	5.6	7.4	8.1	9.6	11.8
Age								
0–4 years	3.8	4.0	4.3	5.6	8.1	8.7	11.0	13.4
5–17 years	3.3	3.4	3.6	5.5	7.2	7.9	9.1	11.2
5–9 years	3.1	3.6	3.5	6.2	7.5	8.6	10.0	12.7
10–17 years	3.4	3.3	3.6	5.1	7.1	7.5	8.6	10.3
Sex								
Male	3.4	3.7	3.8	5.6	7.3	7.9	9.5	11.9
Female	3.5	3.4	3.8	5.5	7.6	8.4	9.8	11.7
Race ⁴								
White only	3.5	3.6	3.8	5.3	7.1	7.6	9.0	10.5
Black or African American only	3.1	3.0	3.7	6.4	9.0	10.4	12.4	17.3
American Indian or Alaska Native only	*	*4.8	*	*4.5	*4.1	*9.1	11.3	11.8
Asian only	3.9	4.4	4.3	5.4	8.0	8.4	7.5	11.1
Native Hawaiian or Other Pacific Islander only	---	*	*	*	---	*	*	*
2 or more races	---	5.2	4.6	7.6	---	10.9	14.0	17.1
Hispanic origin and race ⁴								
Hispanic or Latino	2.1	2.5	2.8	4.5	5.5	5.6	7.2	9.2
Not Hispanic or Latino	3.7	3.8	4.0	5.9	7.8	8.7	10.2	12.7
White only	3.8	3.9	4.1	5.7	7.5	8.2	9.7	11.2
Black or African American only	3.1	3.1	3.7	6.3	9.0	10.4	12.4	17.5
Percent of poverty level ⁵								
Below 100%	3.3	3.2	3.3	5.6	7.3	7.1	9.0	12.1
100%–199%	3.0	3.4	3.8	5.2	7.2	7.6	8.7	11.8
200%–399%	3.2	3.4	3.8	5.3	7.3	8.5	10.0	11.6
400% or more	4.2	4.0	4.1	6.1	7.9	8.8	10.5	11.8
Health insurance status at the time of interview ⁶								
Insured	3.5	3.7	3.9	5.6	7.7	8.5	10.0	12.0
Private	3.5	3.7	4.0	5.8	7.4	8.5	10.1	11.7
Medicaid	3.6	3.7	3.6	5.3	8.4	8.4	9.5	12.3
Uninsured	2.6	2.4	3.0	4.4	5.9	5.3	6.8	9.6

See footnotes at end of table.

Table 35 (page 4 of 5). Health conditions among children under age 18, by selected characteristics: United States, average annual, selected years 1997–1999 through 2012–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#035>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Hay fever or respiratory allergy ¹¹				Three or more ear infections ¹²			
	1997–1999	2000–2002	2003–2005	2012–2014	1997–1999	2000–2002	2003–2005	2012–2014
	Percent of children							
Under 18 years ³	17.5	17.7	17.3	16.0	7.1	6.7	5.8	5.2
Age								
0–4 years	10.7	10.4	10.1	9.6	13.7	12.8	11.0	9.7
5–17 years	19.9	20.3	20.0	18.4	4.8	4.5	3.8	3.6
5–9 years	17.3	18.1	17.9	16.7	7.1	6.9	5.7	5.7
10–17 years	21.6	21.7	21.2	19.4	3.2	2.9	2.7	2.3
Sex								
Male	18.6	18.8	18.9	17.7	7.3	6.9	5.9	5.7
Female	16.3	16.5	15.6	14.2	6.9	6.5	5.6	4.7
Race ⁴								
White only	17.9	18.5	17.8	16.2	7.4	7.2	6.3	5.6
Black or African American only	16.2	15.6	15.2	14.8	5.9	5.0	4.1	3.7
American Indian or Alaska Native only	15.2	16.4	16.5	16.1	*10.8	*6.3	*5.1	*8.4
Asian only	15.3	12.6	11.3	13.4	3.7	2.6	3.3	2.9
Native Hawaiian or Other Pacific Islander only	---	*	*	*	---	*	*	*
2 or more races	---	20.9	20.8	19.0	---	7.4	5.0	5.6
Hispanic origin and race ⁴								
Hispanic or Latino	12.4	12.4	12.8	12.4	6.1	6.7	6.2	5.5
Not Hispanic or Latino	18.4	18.8	18.3	17.1	7.3	6.7	5.7	5.1
White only	19.1	19.9	19.4	17.7	7.7	7.3	6.3	5.7
Black or African American only	16.3	15.5	15.1	15.1	5.9	4.9	4.0	3.6
Percent of poverty level ⁵								
Below 100%	14.3	14.0	14.2	13.2	8.3	7.9	6.7	6.8
100%–199%	15.4	15.6	16.0	14.2	7.1	6.8	5.7	5.6
200%–399%	18.5	18.1	17.7	17.0	6.8	6.5	5.6	4.6
400% or more	20.3	21.1	19.7	18.9	6.6	6.1	5.5	4.2
Health insurance status at the time of interview ⁶								
Insured	18.0	18.3	17.7	16.1	7.3	6.9	5.8	5.3
Private	18.8	19.2	18.5	17.7	6.6	6.4	5.2	4.4
Medicaid	15.0	16.0	16.1	13.8	10.2	8.7	7.4	6.5
Uninsured	14.3	12.6	13.5	14.8	5.9	4.9	5.4	4.4

See footnotes at end of table.

Table 35 (page 5 of 5). Health conditions among children under age 18, by selected characteristics: United States, average annual, selected years 1997–1999 through 2012–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#035>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

-- - Data not available.

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.

¹Based on parent or knowledgeable adult responding to both questions, “Has a doctor or other health professional ever told you that your child had asthma?” and “Does your child still have asthma?”

²Based on parent or knowledgeable adult responding to both questions, “Has a doctor or other health professional ever told you that your child had asthma?” and “During the past 12 months, did your child have an episode of asthma or an asthma attack?”

³Includes all other races not shown separately and unknown health insurance status.

⁴The race groups, white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The five single-race categories plus multiple-race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group; the category 2 or more races includes persons who reported more than one racial group. Prior to 1999, data were tabulated according to the 1977 Standards with four racial groups, and the Asian only category included Native Hawaiian or Other Pacific Islander. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

⁵Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Missing family income data were imputed for 1997 and beyond. See Appendix II, Family income; Poverty; Table VI.

⁶Health insurance categories are mutually exclusive. Persons who reported both Medicaid and private coverage are classified as having private coverage. Starting with 1997 data, state-sponsored health plan coverage is included as Medicaid coverage. Starting with 1999 data, coverage by the Children’s Health Insurance Program (CHIP) is included as Medicaid coverage. In addition to private and Medicaid, the insured category also includes military, other government, and Medicare coverage. Persons not covered by private insurance, Medicaid, CHIP, state-sponsored or other government-sponsored health plans, Medicare, or military plans are considered to have no health insurance coverage. Persons with only Indian Health Service coverage are considered to have no health insurance coverage. See Appendix II, Health insurance coverage.

⁷Based on parent or knowledgeable adult responding to the question, “Has a doctor or health professional ever told you that your child had attention-deficit/hyperactivity disorder (ADHD) or attention deficit disorder (ADD)?”

⁸Based on parent or knowledgeable adult responding to the question, “Overall, do you think that [child] has difficulties in any of the following areas: emotions, concentration, behavior, or being able to get along with other people?”

⁹Based on parent or knowledgeable adult responding to the question, “During the past 12 months, has your child had any kind of food or digestive allergy?”

¹⁰Based on parent or knowledgeable adult responding to the question, “During the past 12 months, has your child had any eczema or any kind of skin allergy?”

¹¹Based on parent or knowledgeable adult responding to the questions, “During the past 12 months, has your child had hay fever?” or to the question, “During the past 12 months, has your child had any kind of respiratory allergy?”

¹²Based on parent or knowledgeable adult responding to the question, “During the past 12 months, has your child had three or more ear infections?”

NOTES: Standard errors are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey, family core and sample child questionnaires. See Appendix I, National Health Interview Survey (NHIS).

Table 36 (page 1 of 4). Age-adjusted cancer incidence rates for selected cancer sites, by sex, race, and Hispanic origin: United States, selected geographic areas, selected years 1990–2012

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#036>.

[Data are based on the Surveillance, Epidemiology, and End Results (SEER) Program's 13 population-based cancer registries]

Site, sex, race, and Hispanic origin	1990	1995	2000	2002	2003	2005	2010	2011	2012	1990–2012 APC ¹
All sites										
Number of new cases per 100,000 population ²										
All persons	475.8	471.3	475.0	474.2	463.2	460.6	449.6	441.7	426.1	†–0.5
White	483.2	477.4	485.6	484.5	473.8	472.7	460.5	452.8	435.6	†–0.4
Black or African American	515.1	539.1	525.7	524.1	513.9	504.7	482.0	474.3	459.2	†–0.7
American Indian or Alaska Native ³	348.2	370.8	364.8	356.9	379.9	403.5	403.4	375.4	353.6	0.2
Asian or Pacific Islander	335.3	340.5	341.5	346.8	334.0	331.7	323.0	321.3	308.5	†–0.4
Hispanic or Latino ⁴	355.9	359.8	361.5	371.9	357.9	364.8	347.6	349.8	334.2	–0.3
White, not Hispanic or Latino ⁴	495.2	491.2	503.5	502.0	492.1	490.9	481.2	472.4	455.2	†–0.3
Male	584.1	564.7	565.4	558.3	545.7	534.6	511.1	496.4	464.2	†–1.0
White	591.1	563.4	569.4	563.8	550.5	543.1	517.5	503.7	469.8	†–0.9
Black or African American	688.3	742.4	707.1	688.5	668.9	636.4	593.5	567.7	540.8	†–1.4
American Indian or Alaska Native ³	396.3	424.0	373.1	381.5	447.2	425.2	451.2	401.6	344.7	–0.3
Asian or Pacific Islander	385.9	398.9	400.5	389.9	387.0	371.5	346.1	343.1	314.7	†–1.0
Hispanic or Latino ⁴	417.4	440.6	435.7	443.5	423.7	426.5	396.9	390.5	362.2	†–0.7
White, not Hispanic or Latino ⁴	606.7	577.3	588.6	581.4	568.9	561.3	538.4	523.8	489.9	†–0.8
Female	411.5	410.6	414.0	417.1	406.9	409.6	405.9	403.7	401.3	–0.1
White	421.2	423.2	430.0	431.4	422.1	424.6	420.3	417.7	414.3	0.0
Black or African American	406.5	403.1	403.8	414.6	409.8	415.9	404.7	409.9	403.2	0.0
American Indian or Alaska Native ³	315.6	337.0	365.5	337.7	335.5	389.0	376.2	361.3	367.7	†0.7
Asian or Pacific Islander	295.7	297.7	300.8	319.9	299.4	306.7	311.0	309.8	308.9	†0.2
Hispanic or Latina ⁴	322.6	310.5	317.6	328.0	317.4	326.3	317.5	326.1	320.1	0.1
White, not Hispanic or Latina ⁴	430.6	436.7	446.0	448.1	439.7	442.1	439.9	436.3	433.2	0.1
Lung and bronchus										
Male	95.0	86.9	77.8	75.7	75.5	71.5	62.9	60.7	58.3	†–2.1
White	94.1	85.0	76.3	74.9	74.3	70.8	62.7	60.1	57.4	†–2.0
Black or African American	134.2	137.0	111.7	109.1	112.1	98.0	82.3	82.3	80.6	†–2.5
American Indian or Alaska Native ³	77.3	82.8	62.6	46.2	74.2	67.4	65.3	55.3	44.1	†–1.2
Asian or Pacific Islander	64.4	60.1	63.9	57.9	58.7	58.2	50.9	49.9	48.3	†–1.2
Hispanic or Latino ⁴	59.4	52.4	45.2	49.0	46.5	44.0	34.3	35.6	33.6	†–2.1
White, not Hispanic or Latino ⁴	97.4	88.4	80.2	78.4	77.9	74.6	67.0	64.0	61.4	†–1.9
Female	47.2	49.3	48.6	49.4	49.8	50.0	46.0	44.8	43.9	†–0.3
White	48.5	51.7	50.7	51.5	52.4	51.9	48.0	46.5	45.5	–0.2
Black or African American	52.9	50.0	55.1	55.4	55.4	58.4	52.8	49.8	51.1	0.0
American Indian or Alaska Native ³	30.4	46.1	38.5	39.9	41.4	45.5	36.6	42.1	34.4	0.9
Asian or Pacific Islander	28.4	27.6	27.3	29.7	29.1	30.8	29.1	31.3	28.8	†0.3
Hispanic or Latina ⁴	26.2	25.1	24.3	24.9	24.4	22.9	24.6	23.9	22.6	†–0.5
White, not Hispanic or Latina ⁴	50.8	54.8	54.4	55.4	56.6	56.4	51.9	50.6	49.7	0.0
Colon and rectum										
Male	72.3	63.2	62.6	60.1	58.4	54.7	46.9	45.4	43.5	†–2.1
White	73.0	62.4	62.0	58.9	57.0	54.1	45.4	43.4	41.9	†–2.2
Black or African American	73.1	75.1	73.6	72.5	76.3	67.0	57.6	57.0	55.5	†–1.3
American Indian or Alaska Native ³	61.9	66.2	48.7	49.4	70.4	65.6	64.6	64.4	57.8	–0.1
Asian or Pacific Islander	60.9	58.8	58.0	59.0	53.3	47.7	45.5	45.7	42.1	†–1.6
Hispanic or Latino ⁴	47.6	46.0	50.5	46.0	47.4	47.3	42.0	43.0	40.0	†–0.7
White, not Hispanic or Latino ⁴	75.0	63.9	63.4	60.3	58.1	54.8	46.0	43.5	42.2	†–2.3
Female	50.2	45.9	46.1	45.2	43.5	41.4	36.0	34.7	33.4	†–1.5
White	49.7	45.4	45.6	44.1	42.9	40.3	34.5	33.7	32.7	†–1.6
Black or African American	61.3	54.9	58.1	56.3	55.4	54.1	46.3	42.2	41.0	†–1.2
American Indian or Alaska Native ³	45.8	47.9	39.1	50.9	45.1	48.3	41.9	46.0	45.1	–0.3
Asian or Pacific Islander	37.8	38.8	37.5	41.8	36.4	36.9	34.3	31.8	28.7	†–1.2
Hispanic or Latina ⁴	34.5	32.0	34.1	32.1	33.5	33.6	30.7	29.0	28.1	†–0.6
White, not Hispanic or Latina ⁴	50.9	46.7	46.7	45.5	44.0	41.3	34.9	34.5	33.5	†–1.6
Prostate										
Male	166.9	166.6	178.9	177.9	165.4	154.0	143.1	136.7	110.4	†–1.8
White	168.5	161.3	174.8	174.5	161.6	149.9	138.2	131.9	104.0	†–1.9
Black or African American	219.4	279.2	291.4	280.3	253.2	242.9	218.3	206.3	178.1	†–1.8
American Indian or Alaska Native ³	99.5	93.5	69.6	93.7	112.4	93.8	83.7	67.1	59.5	†–1.9
Asian or Pacific Islander	88.5	104.6	108.2	104.1	104.9	95.7	78.7	79.4	63.3	†–1.8
Hispanic or Latino ⁴	119.2	141.2	150.2	152.8	139.3	132.4	117.9	109.6	93.5	†–1.5
White, not Hispanic or Latino ⁴	172.2	163.6	178.6	177.6	165.1	152.5	142.2	136.3	106.7	†–1.9

See footnotes at end of table.

Table 36 (page 2 of 4). Age-adjusted cancer incidence rates for selected cancer sites, by sex, race, and Hispanic origin: United States, selected geographic areas, selected years 1990–2012

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#036>.

[Data are based on the Surveillance, Epidemiology, and End Results (SEER) Program's 13 population-based cancer registries]

Site, sex, race, and Hispanic origin	1990	1995	2000	2002	2003	2005	2010	2011	2012	1990–2012 APC ¹
Breast										
Number of new cases per 100,000 population ²										
Female	129.4	130.9	134.2	132.8	124.1	124.4	123.3	126.6	125.9	†–0.3
White	134.2	136.3	140.6	138.6	129.0	129.5	127.2	130.3	128.9	†–0.3
Black or African American	117.4	123.0	122.6	124.2	123.3	119.3	123.2	128.9	130.0	†0.3
American Indian or Alaska Native ³	69.6	96.0	100.2	81.3	93.5	105.6	91.3	105.2	96.9	0.4
Asian or Pacific Islander	88.1	88.4	94.7	101.0	91.9	95.6	100.3	102.7	103.9	†0.7
Hispanic or Latina ⁴	90.8	89.4	97.0	93.4	87.8	93.1	86.9	95.7	91.8	0.1
White, not Hispanic or Latina ⁴	138.6	141.8	147.0	145.8	135.7	136.2	135.4	137.2	136.5	–0.2
Cervix uteri										
Female	11.9	9.9	8.9	8.4	8.2	7.9	7.2	7.0	6.8	†–2.4
White	11.2	9.2	8.9	8.3	7.9	7.8	7.2	7.2	6.7	†–2.1
Black or African American	16.6	14.9	10.7	10.1	10.8	9.3	8.5	7.8	7.6	†–3.6
American Indian or Alaska Native ³	14.7	*	*	*	*	10.7	8.3	9.5	*	†–2.0
Asian or Pacific Islander	12.1	11.2	8.0	8.2	8.1	7.8	6.6	5.5	6.5	†–3.8
Hispanic or Latina ⁴	21.3	17.4	17.1	14.5	14.0	13.7	10.4	9.3	9.2	†–3.8
White, not Hispanic or Latina ⁴	9.7	7.8	7.1	7.0	6.5	6.4	6.2	6.5	5.9	†–1.9
Corpus and uterus, not otherwise specified										
Female	24.7	24.9	23.9	24.0	23.6	24.1	26.7	26.5	27.0	†0.4
White	26.4	26.4	25.6	24.8	25.0	25.4	27.6	27.2	27.9	0.2
Black or African American	17.0	17.9	17.4	22.4	20.3	21.5	24.6	27.7	25.0	†2.2
American Indian or Alaska Native ³	19.3	*	18.5	19.2	19.5	14.4	27.6	19.5	19.3	*
Asian or Pacific Islander	13.5	17.8	16.6	18.8	16.6	18.9	22.5	21.8	22.3	†1.8
Hispanic or Latina ⁴	17.7	16.5	15.8	17.2	17.8	19.3	20.5	22.1	23.7	†1.5
White, not Hispanic or Latina ⁴	27.1	27.5	26.8	25.8	26.0	26.2	28.6	27.7	28.3	0.1
Ovary										
Female	15.6	14.6	14.2	13.9	13.6	13.2	12.5	12.1	12.2	†–1.0
White	16.4	15.4	15.0	14.7	14.3	13.8	13.4	12.9	12.9	†–1.0
Black or African American	11.3	10.8	10.9	9.9	11.6	10.8	9.4	9.9	10.4	–0.4
American Indian or Alaska Native ³	21.9	*	18.9	*	13.4	12.2	11.9	10.8	15.7	*
Asian or Pacific Islander	11.1	10.4	10.4	12.2	10.2	10.9	9.7	9.6	9.5	†–0.5
Hispanic or Latina ⁴	12.2	11.7	10.6	13.8	11.6	11.7	11.8	11.0	12.1	–0.3
White, not Hispanic or Latina ⁴	16.8	15.9	15.6	14.7	14.7	14.1	13.4	13.1	12.9	†–1.1
Oral cavity and pharynx										
Male	18.5	16.5	15.8	15.7	15.2	15.1	15.5	15.9	15.4	†–0.7
White	18.0	16.4	15.6	15.9	15.2	15.5	16.0	16.8	16.2	†–0.4
Black or African American	25.4	22.3	19.4	18.1	17.5	15.8	13.8	13.8	14.0	†–2.8
American Indian or Alaska Native ³	*	*	*	*	16.3	10.4	21.3	15.9	15.3	*
Asian or Pacific Islander	14.9	11.8	13.7	12.9	11.9	11.5	11.7	10.9	10.6	†–1.1
Hispanic or Latino ⁴	10.7	12.3	9.0	9.6	8.8	9.7	8.8	10.9	9.9	–0.6
White, not Hispanic or Latino ⁴	18.8	16.9	16.6	16.9	16.3	16.4	17.5	18.0	17.4	–0.2
Female	7.3	7.0	6.2	6.5	5.9	6.1	6.1	6.1	6.3	†–0.8
White	7.4	7.1	6.2	6.5	5.8	6.0	6.3	6.3	6.5	†–0.7
Black or African American	6.4	6.7	5.4	6.3	6.7	6.9	5.3	5.2	4.5	†–1.5
American Indian or Alaska Native ³	*	*	*	*	*	*	*	*	7.4	*
Asian or Pacific Islander	6.1	5.2	6.3	6.0	5.2	5.9	5.2	4.7	6.0	†–0.8
Hispanic or Latina ⁴	4.0	3.7	3.7	3.7	3.8	3.5	4.2	3.7	3.8	–0.4
White, not Hispanic or Latina ⁴	7.8	7.5	6.6	7.0	6.2	6.5	6.7	6.9	7.0	†–0.6
Stomach										
Male	14.6	13.5	12.6	12.0	11.8	11.4	10.4	10.5	10.1	†–1.7
White	12.8	11.9	10.6	10.4	10.1	9.6	9.3	9.5	8.8	†–1.6
Black or African American	21.5	18.6	18.6	15.9	18.7	17.4	13.3	14.0	14.6	†–2.1
American Indian or Alaska Native ³	*	24.1	20.2	25.3	*	20.8	20.7	21.0	16.6	–0.5
Asian or Pacific Islander	26.8	24.5	22.9	20.5	19.2	20.2	15.2	14.2	14.8	†–2.9
Hispanic or Latino ⁴	20.2	19.3	16.1	16.4	16.2	15.5	14.9	14.2	13.1	†–1.7
White, not Hispanic or Latino ⁴	12.1	11.1	10.0	9.6	9.2	8.6	8.2	8.4	7.8	†–1.9

See footnotes at end of table.

Table 36 (page 3 of 4). Age-adjusted cancer incidence rates for selected cancer sites, by sex, race, and Hispanic origin: United States, selected geographic areas, selected years 1990–2012

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#036>.

[Data are based on the Surveillance, Epidemiology, and End Results (SEER) Program's 13 population-based cancer registries]

Site, sex, race, and Hispanic origin	1990	1995	2000	2002	2003	2005	2010	2011	2012	1990–2012 APC ¹
Stomach										
Number of new cases per 100,000 population ²										
Female	6.7	6.2	6.1	6.2	6.0	5.7	5.7	5.6	5.8	†–0.8
White	5.7	5.1	5.0	5.1	5.0	4.7	4.7	4.8	5.0	†–0.8
Black or African American	10.0	9.8	8.7	9.9	9.5	8.1	8.5	9.4	8.0	†–1.0
American Indian or Alaska Native ³	*	*	*	17.2	*	*	13.0	*	12.2	*
Asian or Pacific Islander	15.4	13.0	12.9	11.5	11.1	10.3	8.9	7.8	8.7	†–2.7
Hispanic or Latina ⁴	10.8	11.2	10.6	10.7	10.2	10.2	9.4	8.8	9.4	†–1.0
White, not Hispanic or Latina ⁴	5.1	4.5	4.2	4.2	4.1	3.8	3.6	3.9	3.9	†–1.5
Pancreas										
Male	13.0	12.7	12.8	12.8	12.5	13.6	13.7	14.2	13.9	†0.5
White	12.7	12.4	12.6	13.0	12.3	13.4	13.5	14.2	14.3	†0.7
Black or African American	19.3	19.1	18.2	13.9	17.4	18.3	18.5	16.8	15.8	–0.4
American Indian or Alaska Native ³	*	*	*	*	*	21.2	17.3	17.6	*	*
Asian or Pacific Islander	11.0	10.3	10.7	9.9	10.2	11.8	11.4	12.3	10.3	0.1
Hispanic or Latino ⁴	10.7	12.0	12.2	10.9	10.1	12.3	11.5	13.4	10.6	†0.8
White, not Hispanic or Latino ⁴	12.8	12.4	12.7	13.2	12.6	13.5	13.7	14.3	14.8	†0.7
Female	10.0	9.9	9.9	10.5	10.4	10.8	11.2	10.6	11.2	†0.6
White	9.7	9.6	9.6	10.1	10.2	10.6	10.9	10.3	10.9	†0.6
Black or African American	12.9	15.5	12.8	15.9	14.4	16.5	15.1	15.0	15.0	0.0
American Indian or Alaska Native ³	*	*	20.4	*	*	12.4	13.8	*	11.3	*
Asian or Pacific Islander	9.9	8.1	9.2	9.1	8.1	8.0	10.4	9.1	10.1	†1.0
Hispanic or Latina ⁴	9.9	9.0	9.1	11.1	8.6	11.1	9.9	10.1	10.1	0.1
White, not Hispanic or Latina ⁴	9.7	9.7	9.6	10.0	10.5	10.6	11.1	10.4	11.0	†0.7
Urinary bladder										
Male	37.2	35.4	36.8	35.8	36.9	37.0	35.8	33.8	34.0	†–0.2
White	40.7	38.8	40.8	39.3	40.7	40.9	39.8	37.9	37.8	†–0.2
Black or African American	19.7	19.7	20.2	20.9	23.2	23.1	22.3	20.7	23.1	†0.5
American Indian or Alaska Native ³	*	*	*	*	*	16.8	16.1	17.5	*	*
Asian or Pacific Islander	15.4	17.2	16.9	19.7	18.0	17.5	17.2	14.9	16.3	0.3
Hispanic or Latino ⁴	22.4	17.8	20.4	21.2	20.2	20.1	18.5	19.8	18.9	–0.3
White, not Hispanic or Latino ⁴	42.4	40.9	43.2	41.6	43.3	43.6	43.0	40.8	40.7	0.0
Female	9.5	9.3	9.1	9.1	9.2	9.0	8.6	8.3	8.3	†–0.5
White	10.0	10.1	10.0	10.0	10.0	9.7	9.4	9.1	9.1	†–0.4
Black or African American	8.7	7.3	7.8	8.5	7.8	7.9	7.0	7.1	6.2	–0.6
American Indian or Alaska Native ³	*	*	*	*	*	*	*	*	*	*
Asian or Pacific Islander	5.3	4.5	4.3	3.4	5.0	5.2	4.7	4.3	4.1	–0.2
Hispanic or Latino ⁴	5.8	5.3	5.8	6.4	4.6	5.9	4.7	5.6	4.9	–0.4
White, not Hispanic or Latino ⁴	10.3	10.6	10.5	10.5	10.8	10.3	10.2	9.8	9.9	†–0.2
Non-Hodgkin lymphoma										
Male	22.6	25.1	23.6	23.8	24.2	24.7	25.4	23.7	23.4	0.2
White	23.6	26.2	24.9	25.1	25.7	25.9	26.6	24.9	24.6	†0.2
Black or African American	17.7	21.7	17.6	18.1	19.4	19.7	21.3	17.5	17.1	–0.1
American Indian or Alaska Native ³	*	*	15.3	16.6	*	23.2	18.7	10.8	13.6	*
Asian or Pacific Islander	16.7	16.6	16.4	16.5	16.3	17.9	17.7	17.9	17.5	0.3
Hispanic or Latino ⁴	17.3	21.0	20.5	20.6	19.6	20.0	22.9	19.9	19.2	0.4
White, not Hispanic or Latino ⁴	24.3	26.7	25.5	25.7	26.5	26.9	27.2	25.6	25.7	†0.3
Female	14.5	15.2	16.0	16.5	17.2	16.5	17.0	15.8	16.0	†0.6
White	15.4	15.9	16.9	17.5	18.0	17.7	18.1	16.5	16.9	†0.7
Black or African American	10.4	10.3	11.9	11.8	13.4	13.3	12.6	12.9	13.5	†1.3
American Indian or Alaska Native ³	*	*	13.5	*	*	13.1	13.0	15.1	10.9	*
Asian or Pacific Islander	9.5	12.2	11.5	12.4	12.7	9.7	11.9	12.0	11.3	0.5
Hispanic or Latina ⁴	13.7	13.1	13.8	13.6	15.3	15.1	15.6	14.5	14.9	†0.8
White, not Hispanic or Latina ⁴	15.6	16.2	17.3	18.1	18.4	18.1	18.4	16.9	17.1	†0.7

See footnotes at end of table.

Table 36 (page 4 of 4). Age-adjusted cancer incidence rates for selected cancer sites, by sex, race, and Hispanic origin: United States, selected geographic areas, selected years 1990–2012

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#036>.

[Data are based on the Surveillance, Epidemiology, and End Results (SEER) Program's 13 population-based cancer registries]

Site, sex, race, and Hispanic origin	1990	1995	2000	2002	2003	2005	2010	2011	2012	1990–2012 APC ¹
Leukemia										
Number of new cases per 100,000 population ²										
Male	17.2	17.7	17.1	17.2	17.4	17.4	17.7	17.3	16.9	0.1
White	18.0	19.0	18.2	18.6	18.5	18.9	19.2	18.5	17.8	0.1
Black or African American	16.1	13.4	14.3	12.7	15.0	12.8	13.4	14.2	14.8	0.1
American Indian or Alaska Native ³	*	*	*	7.9	*	12.3	*	9.7	14.6	*
Asian or Pacific Islander	8.4	10.2	10.6	9.6	10.5	9.3	10.0	10.3	10.2	0.0
Hispanic or Latino ⁴	12.2	14.6	13.2	12.3	12.3	13.2	13.0	13.6	11.2	0.3
White, not Hispanic or Latino ⁴	18.3	19.3	18.7	19.2	19.0	19.3	19.7	19.1	18.6	†0.2
Female	9.9	10.3	10.4	10.1	10.1	10.1	10.6	10.3	10.4	†0.2
White	10.3	10.9	11.0	10.9	10.7	10.6	11.3	11.0	11.1	†0.3
Black or African American	8.5	8.4	9.8	7.5	9.1	9.5	9.0	9.1	9.1	0.1
American Indian or Alaska Native ³	*	*	*	*	*	*	8.7	*	6.6	*
Asian or Pacific Islander	5.7	6.4	6.4	6.4	6.5	6.4	6.0	6.3	6.5	0.0
Hispanic or Latina ⁴	8.5	8.2	7.8	8.7	7.2	8.2	8.6	8.5	8.8	0.4
White, not Hispanic or Latina ⁴	10.3	11.1	11.1	10.9	11.1	10.6	11.6	11.3	11.2	†0.4

† Annual percent change (APC) is significantly different from zero ($p < 0.05$).

0.0 APC is greater than -0.05 but less than 0.05 .

* Estimates are considered unreliable. Data not shown if the rate is based on fewer than 16 cases for the time interval. The trend is not shown if it is based on fewer than 10 cases for at least 1 year within the time interval.

¹ APC was calculated by fitting a linear regression model to the natural logarithm of the yearly rates from 1990–2012.

² Age-adjusted by 5-year age groups to the year 2000 U.S. standard population. Age-adjusted rates are based on at least 16 cases. See Appendix II, Age adjustment.

³ Estimates for the American Indian or Alaska Native populations are based on the Contract Health Service Delivery Area (CHSDA) counties within SEER areas.

⁴ Hispanic data exclude cases from Alaska. The race groups, white, black, Asian or Pacific Islander, and American Indian or Alaska Native, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. The North American Association of Central Cancer Registries (NAACCR) Hispanic Identification Algorithm was used on a combination of variables to classify cases as Hispanic for analytic purposes. See the report, NAACCR guideline for enhancing Hispanic-Latino identification, for more information. Available from: http://seer.cancer.gov/seerstat/variables/seer/yr1973_2006/race_ethnicity/. See Appendix II, Hispanic origin.

NOTES: See Appendix II, Incidence. Estimates are based on 13 SEER areas (November 2014 submission) and differ from published estimates based on 9 SEER areas or other submission dates. See Appendix I, Surveillance, Epidemiology, and End Results Program (SEER). The site variable distinguishes Kaposi Sarcoma and Mesothelioma as individual cancer sites. As a result, Kaposi Sarcoma and Mesothelioma cases do not contribute to other cancer sites. Estimates for 2001–2009 were computed using intercensal population estimates based on the 2000 and 2010 censuses. Data have been revised and differ from previous editions of *Health, United States*. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: National Institutes of Health, National Cancer Institute, Surveillance, Epidemiology, and End Results Program. Available from: <http://www.seer.cancer.gov>. See Appendix I, Surveillance, Epidemiology, and End Results Program (SEER).

Table 37. Five-year relative cancer survival rates for selected cancer sites, by race and sex: United States, selected geographic areas, selected years 1975–1977 through 2005–2011

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#037>.

[Data are based on the Surveillance, Epidemiology, and End Results (SEER) Program's nine population-based cancer registries]

Sex and site	White					Black or African American				
	1975–1977	1981–1983	1987–1989	1999–2001	2005–2011	1975–1977	1981–1983	1987–1989	1999–2001	2005–2011
Both sexes						Percent of patients				
All sites	49.8	51.3	56.6	67.1	69.7	39.1	38.8	43.0	57.9	62.2
Oral cavity and pharynx	54.0	54.0	55.9	62.2	68.1	36.0	30.9	34.0	44.6	44.9
Esophagus	5.5	7.3	10.5	18.8	21.0	3.5	4.3	6.6	12.8	13.6
Stomach	14.1	16.2	18.3	22.2	29.1	16.1	16.4	18.8	23.0	28.3
Colon	50.9	55.4	60.6	66.6	66.5	44.7	48.5	52.3	52.3	56.4
Rectum	48.2	52.0	58.7	66.4	68.1	44.4	40.1	52.3	59.1	64.9
Pancreas	2.5	2.5	3.2	4.9	7.8	2.3	3.6	5.5	5.6	7.2
Lung and bronchus	12.2	13.3	13.3	15.5	18.8	11.3	11.3	10.9	12.6	15.5
Urinary bladder	73.2	77.4	79.8	81.0	79.4	50.3	59.5	62.6	67.3	66.7
Non-Hodgkin lymphoma	46.8	50.7	51.3	64.9	73.0	48.6	49.4	45.9	55.7	64.0
Leukemia	34.5	38.0	43.8	51.6	62.5	33.2	33.9	35.0	43.2	55.4
Male										
All sites	42.7	46.5	52.8	67.5	70.2	32.7	34.2	38.9	61.0	65.4
Oral cavity and pharynx	53.7	52.8	54.0	62.0	68.0	29.7	25.3	30.0	39.4	42.1
Esophagus	4.8	6.5	11.0	18.6	21.2	2.0	3.7	5.3	11.1	12.6
Stomach	13.1	15.4	15.6	21.0	27.6	16.1	16.2	16.6	25.0	22.8
Colon	50.5	56.2	61.4	67.8	67.2	44.0	44.6	50.7	53.4	55.1
Rectum	47.3	50.9	58.9	66.5	67.9	41.4	37.7	47.7	60.0	60.2
Pancreas	2.7	2.1	3.1	5.4	7.9	2.5	3.7	5.1	3.7	8.5
Lung and bronchus	11.1	11.7	12.0	13.3	16.2	10.6	10.1	10.8	10.7	13.2
Prostate gland	68.5	73.0	84.3	99.7	99.7	60.7	62.8	71.2	97.3	97.6
Urinary bladder	74.3	78.5	81.9	81.5	80.8	56.5	64.9	67.5	71.5	71.3
Non-Hodgkin lymphoma	46.3	50.4	48.1	62.8	72.1	42.6	49.0	41.7	48.9	59.2
Leukemia	33.6	37.7	45.4	52.6	63.6	30.0	33.4	32.7	43.9	59.1
Female										
All sites	56.5	55.9	60.6	66.8	69.2	46.2	44.4	47.7	54.2	58.5
Colon	51.2	54.8	59.9	65.5	65.8	45.3	51.5	53.6	51.5	57.4
Rectum	49.3	53.2	58.4	66.3	68.2	46.8	42.4	56.9	58.2	69.6
Pancreas	2.3	3.0	3.3	4.4	7.7	1.9	3.2	5.8	7.4	6.1
Lung and bronchus	15.4	16.6	15.3	18.0	21.4	13.8	14.9	11.1	15.2	18.1
Melanoma of skin	86.2	87.2	91.3	94.6	95.0	*	*	89.5	76.1	78.1
Breast	75.6	77.1	85.1	90.8	92.0	62.1	63.4	71.1	78.7	80.5
Cervix uteri	69.7	67.8	72.5	73.3	71.0	64.6	59.2	57.0	66.0	59.6
Corpus and uterus, not otherwise specified	88.0	82.2	83.9	85.9	85.3	60.2	50.7	56.7	60.9	65.6
Ovary	35.3	38.4	38.1	43.5	45.6	41.9	37.5	33.7	35.4	37.7
Non-Hodgkin lymphoma	47.3	51.0	55.2	67.5	74.0	55.3	49.8	51.0	63.8	69.0

* Data for population groups with fewer than 25 cases are not shown because estimates are considered unreliable.

NOTES: Rates are based on follow-up of patients through 2012. The rate is the ratio of the observed survival rate for the patient group to the expected survival rate for persons in the general population similar to the patient group with respect to age, sex, race, and calendar year of observation. It estimates the chance of surviving the effects of cancer. The site variable distinguishes Kaposi Sarcoma and Mesothelioma as individual cancer sites. As a result, Kaposi Sarcoma and Mesothelioma cases are excluded from each of the sites shown except all sites combined. The race groups, white and black, include persons of Hispanic and non-Hispanic origin. Due to death certificate race-ethnicity classification and other methodological issues related to developing life tables, relative survival rates for race-ethnicity groups other than white and black are not calculated. Data have been revised and differ from previous editions of *Health, United States*. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: National Institutes of Health, National Cancer Institute, Surveillance, Epidemiology, and End Results Program. Available from: <http://www.seer.cancer.gov>. See Appendix I, Surveillance, Epidemiology, and End Results Program (SEER).

Table 38 (page 1 of 2). Respondent-reported prevalence of heart disease, cancer, and stroke among adults aged 18 and over, by selected characteristics: United States, average annual, selected years 1997–1998 through 2013–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#038>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Heart disease ¹				Cancer ²				Stroke ³			
	1997–1998	1999–2000	2010–2011	2013–2014	1997–1998	1999–2000	2010–2011	2013–2014	1997–1998	1999–2000	2010–2011	2013–2014
	Percent of adults											
18 years and over, age-adjusted ^{4,5}	12.0	11.1	11.1	10.7	4.9	5.1	6.0	5.9	2.3	2.2	2.6	2.5
18 years and over, crude ⁵	11.6	10.9	11.6	11.5	4.8	4.9	6.3	6.4	2.2	2.1	2.7	2.7
Age												
18–44 years	4.6	4.3	4.0	4.0	1.7	1.7	1.7	1.6	0.4	0.4	0.6	0.5
18–24 years	3.2	3.3	3.0	2.8	0.8	1.0	0.7	*0.6	*	*	*	*
25–44 years	5.0	4.6	4.4	4.5	2.0	1.9	2.0	2.0	0.4	0.5	0.7	0.6
45–64 years	13.5	12.6	13.0	12.0	5.4	5.2	6.9	6.7	2.3	2.0	2.9	2.9
45–54 years	10.9	10.0	9.6	8.6	4.0	4.0	4.9	4.3	1.4	1.3	2.1	1.8
55–64 years	17.4	16.6	17.1	15.8	7.4	7.2	9.3	9.3	3.8	3.1	3.9	4.2
65 years and over	31.8	29.6	30.5	29.4	14.1	15.2	18.5	18.2	8.1	8.1	8.2	7.9
65–74 years	27.8	25.8	25.6	25.1	12.4	13.1	15.9	15.6	6.7	6.2	6.3	6.0
75 years and over	37.0	34.3	36.5	35.4	16.2	17.7	21.7	21.8	9.8	10.3	10.6	10.5
Sex ⁴												
Male	12.3	11.9	12.4	11.9	4.1	4.4	5.5	5.3	2.6	2.4	2.6	2.6
Female	11.8	10.5	10.2	9.6	5.8	5.8	6.6	6.5	2.1	2.1	2.6	2.4
Sex and age												
Male:												
18–44 years	3.7	3.6	3.7	3.9	0.8	0.8	0.9	0.7	0.3	0.3	0.5	0.4
45–54 years	11.0	10.0	9.5	9.0	2.0	2.0	3.1	2.9	1.2	1.3	1.9	1.9
55–64 years	18.7	19.7	19.1	17.9	5.8	5.9	7.5	8.2	4.6	3.7	4.0	4.3
65–74 years	32.0	30.4	31.3	30.9	12.8	13.9	16.9	15.9	8.1	6.7	6.6	6.7
75 years and over	40.8	39.2	44.7	41.4	18.3	20.3	26.1	25.2	11.2	11.3	10.6	11.2
Female:												
18–44 years	5.5	4.9	4.3	4.1	2.6	2.5	2.5	2.4	0.4	0.4	0.6	0.6
45–54 years	10.8	9.9	9.6	8.2	6.0	5.9	6.6	5.7	1.5	1.4	2.2	1.7
55–64 years	16.2	13.8	15.3	14.0	8.8	8.4	10.9	10.4	3.2	2.6	3.9	4.0
65–74 years	24.5	22.0	20.6	20.0	12.1	12.5	15.0	15.3	5.5	5.8	6.1	5.3
75 years and over	34.6	31.2	30.9	31.1	14.9	16.1	18.7	19.4	9.0	9.6	10.6	10.1
Race ^{4,6}												
White only	12.2	11.3	11.2	10.9	5.2	5.4	6.3	6.2	2.2	2.1	2.3	2.3
Black or African American only	11.4	10.6	10.7	10.1	3.5	3.5	5.1	4.6	3.3	3.5	4.1	3.8
American Indian or Alaska Native only	18.6	14.7	12.5	10.8	*6.5	*5.7	6.5	*4.8	*5.0	*5.4	*4.7	*2.8
Asian only	6.9	6.3	7.2	6.0	2.4	*2.3	3.0	2.8	*1.2	*1.2	2.4	1.6
Native Hawaiian or Other Pacific Islander only	---	*	*	*	---	*	*	*	---	*	*	*
2 or more races	---	17.0	16.7	15.9	---	*4.7	7.9	7.8	---	*4.0	*3.9	3.3
Hispanic origin and race ^{4,6}												
Hispanic or Latino	8.7	8.0	8.4	8.0	2.9	3.0	3.4	4.0	2.1	1.9	2.7	2.5
Mexican	7.5	7.4	8.4	7.9	3.0	2.8	3.2	3.6	2.5	2.0	2.6	2.6
Not Hispanic or Latino	12.2	11.4	11.4	11.0	5.1	5.2	6.3	6.1	2.3	2.2	2.6	2.5
White only	12.5	11.6	11.7	11.4	5.4	5.5	6.7	6.5	2.2	2.1	2.3	2.3
Black or African American only	11.4	10.5	10.8	10.2	3.6	3.6	5.1	4.6	3.3	3.5	4.2	3.8
Education ^{7,8}												
No high school diploma or GED	15.1	13.8	14.6	13.7	5.3	5.5	5.8	5.6	3.9	3.8	4.4	4.1
High school diploma or GED	12.8	11.9	12.4	12.3	5.5	5.8	6.8	7.1	2.5	2.5	3.4	3.1
Some college or more	12.7	12.0	11.9	11.2	6.0	5.9	7.4	6.8	2.1	1.9	2.3	2.3

See footnotes at end of table.

Table 38 (page 2 of 2). Respondent-reported prevalence of heart disease, cancer, and stroke among adults aged 18 and over, by selected characteristics: United States, average annual, selected years 1997–1998 through 2013–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#038>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Heart disease ¹				Cancer ²				Stroke ³			
	1997–1998	1999–2000	2010–2011	2013–2014	1997–1998	1999–2000	2010–2011	2013–2014	1997–1998	1999–2000	2010–2011	2013–2014
Percent of poverty level ^{4,9}					Percent of adults							
Below 100%	15.3	13.6	13.9	13.7	4.9	4.9	5.3	5.2	4.3	3.7	4.6	4.3
100%–199%	13.2	12.0	12.3	11.9	4.8	5.3	5.9	5.7	3.1	3.2	3.7	3.6
200%–399%	11.5	11.0	11.3	10.6	4.9	5.1	6.2	6.1	2.1	2.1	2.5	2.4
400% or more	11.0	10.2	9.8	9.3	5.2	5.1	6.2	6.0	1.6	1.5	1.5	1.4
Hispanic origin and race and percent of poverty level ^{4,6,9}												
Hispanic or Latino:												
Below 100%	9.7	9.7	9.4	10.2	2.2	2.3	2.9	3.7	3.0	2.0	3.4	3.5
100%–199%	8.7	8.4	8.3	7.9	2.8	3.2	2.6	3.4	2.2	2.2	3.2	2.3
200%–399%	8.4	8.2	8.5	7.0	2.7	2.7	4.7	4.7	*1.8	*2.3	2.2	2.2
400% or more	8.4	5.6	7.5	7.6	*5.5	*4.5	3.3	4.7	*	*	*2.1	*1.9
Not Hispanic or Latino:												
White only:												
Below 100%	17.8	15.2	15.8	16.0	6.3	6.2	6.8	6.8	4.4	3.8	4.4	4.4
100%–199%	14.1	12.8	13.8	13.6	5.6	6.2	7.3	6.9	3.2	3.0	3.6	3.6
200%–399%	12.2	11.6	11.8	11.6	5.2	5.5	6.8	6.5	2.1	2.1	2.3	2.4
400% or more	11.3	10.6	10.2	9.8	5.4	5.3	6.6	6.5	1.6	1.5	1.4	1.4
Black or African American only:												
Below 100%	14.6	13.0	14.7	13.2	4.4	4.0	4.5	3.8	5.0	4.5	6.2	5.0
100%–199%	12.9	11.2	11.2	11.2	3.3	3.2	5.2	5.0	4.2	5.1	4.6	5.1
200%–399%	9.2	10.2	10.5	9.7	3.2	3.7	5.3	5.2	2.5	2.7	3.9	3.0
400% or more	9.5	8.9	7.3	7.0	4.0	4.3	5.6	3.8	*	*	*2.1	2.3
Geographic region ⁴												
Northeast	11.6	10.6	10.1	9.7	4.5	5.0	5.6	5.9	1.8	1.8	2.0	2.2
Midwest	12.1	11.4	11.5	11.8	5.1	5.2	6.7	6.1	2.3	2.2	2.6	2.5
South	12.5	11.5	12.1	11.2	5.0	5.0	6.2	6.0	2.6	2.5	2.9	2.7
West	11.1	10.4	9.9	9.4	5.1	5.0	5.5	5.5	2.1	2.0	2.4	2.2
Location of residence ^{4,10}												
Within MSA	11.7	10.7	10.8	10.3	4.9	5.0	5.9	5.8	2.2	2.1	2.4	2.3
Outside MSA	12.8	12.5	13.0	12.4	5.1	5.5	6.7	6.4	2.7	2.5	3.4	3.1

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.

-- Data not available.

¹Heart disease is based on self-reported responses to questions about whether respondents had ever been told by a doctor or other health professional that they had coronary heart disease, angina (angina pectoris), a heart attack (myocardial infarction), or any other kind of heart disease or heart condition.

²Cancer is based on self-reported responses to a question about whether respondents had ever been told by a doctor or other health professional that they had cancer or a malignancy of any kind. Excludes squamous cell and basal cell carcinomas.

³Stroke is based on self-reported responses to a question about whether respondents had ever been told by a doctor or other health professional that they had a stroke.

⁴Estimates are age-adjusted to the year 2000 standard population using five age groups: 18–44 years, 45–54 years, 55–64 years, 65–74 years, and 75 years and over. Age-adjusted estimates in this table may differ from other age-adjusted estimates based on the same data and presented elsewhere if different age groups are used in the adjustment procedure. See Appendix II, Age adjustment.

⁵Includes all other races not shown separately and unknown education level.

⁶The race groups, white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The five single-race categories plus multiple-race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group; the category 2 or more races includes persons who reported more than one racial group. Prior to 1999, data were tabulated according to the 1977 Standards with four racial groups, and the Asian only category included Native Hawaiian or Other Pacific Islander. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

⁷Estimates are for persons aged 25 and over and are age-adjusted to the year 2000 standard population using five age groups: 25–44 years, 45–54 years, 55–64 years, 65–74 years, and 75 years and over. See Appendix II, Age adjustment.

⁸GED is General Educational Development high school equivalency diploma. See Appendix II, Education.

⁹Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Missing family income data were imputed for 1997–1998 and beyond. See Appendix II, Family income; Poverty; Table VI.

¹⁰MSA is metropolitan statistical area. Starting with 2006 data, MSA status is determined using 2000 census data and the 2000 standards for defining MSAs. For data prior to 2006, see Appendix II, Metropolitan statistical area (MSA) for the applicable standards.

NOTES: Standard errors are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey, family core and sample adult questionnaires. See Appendix I, National Health Interview Survey (NHIS).

Table 39 (page 1 of 2). Number of respondent-reported chronic conditions from 10 selected conditions among adults aged 18 and over, by selected characteristics: United States, selected years 2002–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#039>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Number of respondent-reported chronic conditions from 10 selected conditions ¹											
	0–1 chronic conditions				2–3 chronic conditions				4 or more chronic conditions			
	2002	2010	2013	2014	2002	2010	2013	2014	2002	2010	2013	2014
	Percent distribution											
Total, age-adjusted ^{2,3}	78.5	75.4	76.7	76.9	17.9	19.9	19.0	19.1	3.6	4.6	4.3	3.9
Total, crude ²	78.5	74.0	74.5	74.5	17.9	21.1	20.8	21.0	3.6	4.9	4.7	4.5
Age												
18–64 years	85.1	81.3	82.4	82.5	12.9	16.0	14.9	15.2	2.0	2.8	2.6	2.3
18–44 years	93.3	92.2	93.2	92.8	6.2	7.2	6.3	6.8	0.5	0.6	0.4	0.4
18–24 years	96.4	96.6	97.1	96.9	3.5	3.3	2.8	2.9	*	*	*	*
25–44 years	92.3	90.7	91.8	91.3	7.1	8.6	7.6	8.2	0.7	0.8	0.6	0.5
45–64 years	71.4	66.2	67.7	68.5	24.2	28.1	26.6	26.7	4.4	5.7	5.6	4.8
45–54 years	78.4	74.1	76.2	77.2	18.6	22.3	20.1	20.6	3.1	3.6	3.7	2.1
55–64 years	60.9	56.4	58.4	59.1	32.7	35.3	33.8	33.3	6.4	8.3	7.8	7.6
65 years and over	44.6	37.9	39.1	39.9	43.4	46.5	46.8	46.0	12.0	15.6	14.1	14.1
65–74 years	47.6	41.2	42.6	43.4	41.4	45.5	45.2	44.3	11.0	13.2	12.2	12.3
75 years and over	41.1	33.8	34.4	34.9	45.8	47.7	48.9	48.4	13.2	18.6	16.7	16.7
Sex ³												
Male	79.7	76.2	77.6	77.8	16.5	19.3	18.2	18.4	3.8	4.5	4.3	3.8
Female	77.3	74.7	75.9	76.1	19.1	20.5	19.8	19.8	3.6	4.8	4.3	4.1
Race ^{3,4}												
White only	78.8	76.0	76.9	77.2	17.7	19.5	19.0	19.0	3.5	4.5	4.1	3.8
Black or African American only	74.3	70.4	72.9	72.4	21.3	23.8	21.4	22.1	4.4	5.8	5.7	5.6
American Indian or Alaska Native only	69.5	64.7	71.4	69.5	25.5	28.8	21.4	26.1	*	*	*7.2	*4.4
Asian only	86.5	82.9	84.6	85.2	11.3	14.4	12.8	12.7	*	2.8	2.7	2.1
Native Hawaiian or Other Pacific Islander only	*	*	*	*	*	*	*	*	*	*	*	*
2 or more races	72.3	66.6	70.3	72.0	20.2	23.2	21.4	21.5	*7.6	*10.2	8.3	6.6
Hispanic origin and race ^{3,4}												
Hispanic or Latino	82.2	78.0	80.0	79.2	14.8	17.9	16.1	17.0	3.0	4.1	3.9	3.8
Mexican	81.8	77.7	80.0	78.3	15.0	18.0	15.9	18.2	3.2	4.3	4.1	3.5
Not Hispanic or Latino	78.0	75.1	76.2	76.5	18.3	20.2	19.5	19.5	3.7	4.7	4.3	4.0
White only	78.3	75.6	76.2	76.7	18.1	19.8	19.6	19.5	3.6	4.5	4.2	3.8
Black or African American only	74.3	70.1	72.6	72.2	21.3	24.0	21.7	22.2	4.4	5.8	5.7	5.6
Percent of poverty level ^{3,5}												
Below 100%	71.9	69.2	69.7	70.1	21.3	22.7	21.9	23.3	6.8	8.1	8.4	6.6
100%–199%	76.4	72.6	73.6	73.1	18.6	21.0	20.7	21.5	5.0	6.4	5.7	5.4
200%–399%	77.8	75.6	76.9	76.5	18.9	19.9	18.9	19.6	3.3	4.5	4.2	3.9
400% or more	81.2	78.3	80.0	80.8	15.9	18.7	17.5	16.5	2.8	3.0	2.5	2.7
Health insurance status at the time of interview ^{6,7}												
18–64 years:												
Insured	85.1	82.5	84.1	84.3	13.0	15.0	13.5	13.8	1.9	2.5	2.4	1.9
Private	86.9	84.9	87.0	87.1	11.8	13.6	11.7	11.9	1.2	1.5	1.3	1.1
Medicaid	69.2	69.6	71.4	72.7	22.3	22.3	20.8	21.5	8.5	8.1	7.8	5.8
Uninsured	87.5	85.9	86.6	86.7	10.7	12.4	12.0	12.2	1.8	1.7	1.4	1.1

See footnotes at end of table.

Table 39 (page 2 of 2). Number of respondent-reported chronic conditions from 10 selected conditions among adults aged 18 and over, by selected characteristics: United States, selected years 2002–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#039>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Number of respondent-reported chronic conditions from 10 selected conditions ¹											
	0–1 chronic conditions				2–3 chronic conditions				4 or more chronic conditions			
	2002	2010	2013	2014	2002	2010	2013	2014	2002	2010	2013	2014
Geographic region ³						Percent distribution						
Northeast	79.4	77.7	78.3	79.1	17.4	18.1	17.4	17.7	3.1	4.1	4.3	3.2
Midwest	78.4	73.7	75.5	76.0	17.9	21.2	20.2	19.8	3.7	5.1	4.3	4.2
South	77.3	73.9	75.4	75.4	18.6	21.0	20.0	20.1	4.0	5.2	4.6	4.5
West.	79.7	77.6	78.9	78.6	17.0	18.5	17.4	18.0	3.3	3.9	3.6	3.4
Location of residence ^{3,8}												
Within MSA	79.1	76.1	77.5	77.8	17.5	19.5	18.5	18.4	3.4	4.4	4.0	3.8
Outside MSA.	76.0	71.7	72.3	72.0	19.6	22.6	22.2	23.2	4.4	5.7	5.6	4.8

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.

¹Adults were categorized as having 0–1, 2–3, or 4 or more of the following chronic conditions: hypertension, coronary heart disease, stroke, diabetes, cancer, arthritis, hepatitis, weak or failing kidneys, chronic obstructive pulmonary disease, or current asthma. Data from the National Health Interview Survey capture 10 of 20 chronic conditions used in a standardized approach for defining chronic conditions in the United States. Thus, these estimates are conservative in nature. For more information, see: Goodman RA, Posner SF, Huang ES, Parekh AK, Koh HK. Defining and measuring chronic conditions: imperatives for research, policy, program, and practice. *Prev Chronic Dis* 2013;10:120239. Available from: DOI: http://www.cdc.gov/pcd/issues/2013/12_0239.htm, and Ward BW, Schiller JS. Prevalence of multiple chronic conditions among US adults: estimates from the National Health Interview Survey, 2010. *Prev Chronic Dis* 2013;10:120203. Available from: DOI: http://www.cdc.gov/pcd/issues/2013/12_0203.htm.

²Includes all other races not shown separately and unknown health insurance status.

³Estimates are age-adjusted to the year 2000 standard population using five age groups: 18–44 years, 45–54 years, 55–64 years, 65–74 years, and 75 years and over. See Appendix II, Age adjustment.

⁴The race groups, white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity*. The five single-race categories plus multiple-race categories shown in the table conform to the 1997 Standards. Race-specific estimates are for persons who reported only one racial group; the category 2 or more races includes persons who reported more than one racial group. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

⁵Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Missing family income data were imputed. See Appendix II, Family income; Poverty; Table VI.

⁶Estimates are age-adjusted to the year 2000 standard population using three age groups: 18–44 years, 45–54 years, and 55–64 years. See Appendix II, Age adjustment.

⁷Health insurance categories are mutually exclusive. Persons who reported both Medicaid and private coverage are classified as having private coverage. State-sponsored health plan coverage is included as Medicaid coverage. Coverage by the Children’s Health Insurance Program (CHIP) is included with Medicaid coverage. In addition to private and Medicaid, the insured category also includes military plans, other government-sponsored health plans, and Medicare, not shown separately. Persons not covered by private insurance, Medicaid, CHIP, state-sponsored or other government-sponsored health plans (starting in 1997), Medicare, or military plans are considered to have no health insurance coverage. Persons with only Indian Health Service coverage are considered to have no health insurance coverage. See Appendix II, Health insurance coverage.

⁸MSA is metropolitan statistical area. Starting with 2006 data, MSA status is determined using 2000 census data and the 2000 standards for defining MSAs. For data prior to 2006, see Appendix II, Metropolitan statistical area (MSA) for the applicable standards.

NOTES: Standard errors are available in the spreadsheet version of this table. See <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey, family core and sample adult questionnaires. See Appendix I, National Health Interview Survey (NHIS).

Table 40 (page 1 of 3). Diabetes prevalence and glycemic control among adults aged 20 and over, by sex, age, and race and Hispanic origin: United States, selected years 1988–1994 through 2011–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#040>.

[Data are based on interviews and physical examinations of a sample of the civilian noninstitutionalized population]

Sex, age, and race and Hispanic origin ³	Physician-diagnosed and undiagnosed diabetes ^{1,2}				Physician-diagnosed diabetes ¹				Undiagnosed diabetes ²			
	1988– 1994	1999– 2002	2003– 2006	2011– 2014	1988– 1994	1999– 2002	2003– 2006	2011– 2014	1988– 1994	1999– 2002	2003– 2006	2011– 2014
20 years and over, age-adjusted ⁴	Percent of population											
All persons ⁵	8.8	9.9	10.6	11.9	5.2	6.6	7.7	9.0	3.6	3.2	3.0	2.9
Male	9.6	11.2	11.1	12.8	5.5	7.3	7.0	9.4	4.1	3.9	4.1	3.3
Female	8.2	8.6	10.2	11.2	5.1	5.9	8.2	8.7	3.2	2.7	2.0	2.4
Not Hispanic or Latino:												
White only	7.7	8.5	8.9	9.6	4.8	5.5	6.2	7.6	2.9	3.0	2.7	2.0
Black or African American only	16.3	14.0	16.7	18.0	9.1	9.2	13.0	13.4	7.2	4.8	3.8	4.6
Asian only	---	---	---	16.3	---	---	---	10.4	---	---	---	5.9
Hispanic or Latino	---	---	---	16.8	---	---	---	12.1	---	---	---	4.7
Mexican origin	15.6	13.9	17.1	18.0	10.7	10.8	12.9	13.0	5.0	3.1	4.2	5.1
Percent of poverty level: ⁶												
Below 100%	14.2	14.6	14.8	17.4	8.8	9.0	12.4	13.4	*5.4	5.6	*	3.9
100% or more	8.1	9.3	10.1	11.2	4.8	6.4	7.1	8.5	3.3	2.9	3.0	2.7
100%–199%	9.7	13.1	13.9	15.0	5.2	9.4	9.5	10.5	4.4	*3.6	4.4	4.5
200% or more	7.8	8.2	8.9	9.7	4.7	5.5	6.3	7.7	3.1	2.7	2.6	2.1
200%–399%	7.8	10.5	10.5	11.4	4.3	7.3	7.5	8.6	3.6	3.2	*3.1	2.8
400% or more	7.8	6.7	7.0	8.6	5.3	4.3	5.2	7.2	2.5	2.3	*	*1.4
20 years and over, crude												
All persons ⁵	8.3	9.8	10.9	12.6	4.9	6.6	7.9	9.6	3.4	3.2	3.0	3.0
Male	8.6	10.8	11.0	13.2	4.9	7.1	6.9	9.7	3.7	3.7	4.0	3.5
Female	8.0	8.9	10.8	12.1	5.0	6.1	8.7	9.5	3.1	2.8	2.1	2.5
Not Hispanic or Latino:												
White only	7.6	8.9	9.6	11.0	4.7	5.6	6.7	8.7	2.9	3.2	3.0	2.3
Black or African American only	13.3	12.5	15.8	17.5	7.2	8.3	12.3	13.0	6.1	4.2	3.5	4.6
Asian only	---	---	---	15.1	---	---	---	9.3	---	---	---	5.8
Hispanic or Latino	---	---	---	13.8	---	---	---	9.8	---	---	---	4.0
Mexican origin	10.4	9.3	12.6	14.3	6.3	7.2	8.8	10.1	4.1	2.0	*3.8	4.2
Percent of poverty level: ⁶												
Below 100%	11.6	13.4	12.7	15.0	7.2	8.4	10.6	11.2	4.4	5.1	*	3.7
100% or more	7.6	9.2	10.5	12.3	4.5	6.3	7.3	9.4	3.1	2.9	3.1	2.9
100%–199%	9.1	12.9	14.9	16.8	5.2	9.3	10.2	12.1	3.9	*3.6	4.7	4.7
200% or more	7.1	8.0	9.1	10.8	4.3	5.4	6.5	8.4	2.8	2.6	2.7	2.3
200%–399%	6.8	10.2	11.0	12.3	3.7	7.0	7.7	9.3	3.1	*3.1	*3.2	3.0
400% or more	7.6	6.4	7.4	9.5	5.2	4.1	5.3	7.7	*2.5	2.3	*2.2	*1.8
Age												
20–44 years	*2.1	4.4	3.9	4.0	*	3.2	2.8	2.6	1.1	*	*1.1	1.4
45–64 years	14.0	12.8	13.7	16.6	7.9	8.3	10.1	12.3	6.0	4.5	3.5	4.3
65 years and over	19.4	20.4	24.9	26.3	12.7	13.7	17.5	21.9	6.7	6.7	7.4	4.3

See footnotes at end of table.

Table 40 (page 2 of 3). Diabetes prevalence and glycemic control among adults aged 20 and over, by sex, age, and race and Hispanic origin: United States, selected years 1988–1994 through 2011–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#040>.

[Data are based on interviews and physical examinations of a sample of the civilian noninstitutionalized population]

Sex, age, and race and Hispanic origin ³	Poor glycemic control (A1c greater than 9%) among persons with physician-diagnosed diabetes			
	1988–1994	1999–2002	2003–2006	2011–2014
Percent of population with physician-diagnosed diabetes				
20 years and over, age-adjusted ⁴				
All persons ⁵	26.3	24.7	18.8	20.6
Male	22.4	27.7	20.7	24.1
Female	29.4	*20.3	17.1	18.0
Not Hispanic or Latino:				
White only	23.7	*22.9	*14.9	*16.6
Black or African American only	38.9	25.4	25.7	23.9
Asian only	---	---	---	*17.3
Hispanic or Latino	---	---	---	29.8
Mexican origin	29.8	28.0	*26.3	27.6
Percent of poverty level: ⁶				
Below 100%	37.2	30.6	*19.9	27.3
100% or more	22.8	*22.6	19.8	18.3
100%–199%	*	*	*19.2	21.7
200% or more	21.2	*25.6	20.8	*16.6
200%–399%	*24.2	*27.0	*19.1	*13.5
400% or more	*	*	*	*
20 years and over, crude				
All persons ⁵	23.3	18.4	13.0	15.6
Male	20.2	20.2	14.8	15.4
Female	25.8	16.7	11.5	15.7
Not Hispanic or Latino:				
White only	20.6	13.6	8.7	12.0
Black or African American only	34.2	25.4	21.0	19.0
Asian only	---	---	---	*12.8
Hispanic or Latino	---	---	---	25.5
Mexican origin	29.2	26.8	24.0	22.9
Percent of poverty level: ⁶				
Below 100%	30.2	25.6	17.6	23.2
100% or more	21.4	15.9	12.2	13.6
100%–199%	24.2	*14.9	*11.5	*13.9
200% or more	20.0	16.4	12.5	13.4
200%–399%	*21.2	*17.5	*10.7	13.1
400% or more	*18.3	*	14.8	*13.8
Age				
20–44 years	29.5	*32.7	25.2	26.2
45–64 years	26.0	19.9	16.6	17.8
65 years and over	18.0	*10.2	*4.1	9.2

See footnotes at end of table.

Table 40 (page 3 of 3). Diabetes prevalence and glycemic control among adults aged 20 and over, by sex, age, and race and Hispanic origin: United States, selected years 1988–1994 through 2011–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#040>.

[Data are based on interviews and physical examinations of a sample of the civilian noninstitutionalized population]

-- Data not available.

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.

¹Physician-diagnosed diabetes was obtained by self-report and excludes women who are pregnant.

²Undiagnosed diabetes is defined as a fasting plasma glucose (FPG) of at least 126 mg/dL or a hemoglobin A1c of at least 6.5% and no reported physician diagnosis. Respondents had fasted for at least 8 hours and less than 24 hours. Pregnant females are excluded. Estimates in some prior editions of *Health, United States* included data from respondents who had fasted for at least 9 hours and less than 24 hours. Starting in 2005–2006, testing was performed at a different laboratory and using different instruments than testing in earlier years. The National Health and Nutrition Examination Survey (NHANES) conducted crossover studies to evaluate the impact of these changes on FPG and A1c measurements and recommended adjustments to the FPG data. The adjustments recommended by NHANES were incorporated into the data presented here. For more information, see http://www.cdc.gov/nchs/hhanes/hhanes2005-2006/GLU_D.htm. Prior to *Health, United States, 2010*, the definition of undiagnosed diabetes did not consider hemoglobin A1c. The revised definition of undiagnosed diabetes was based on recommendations from the American Diabetes Association. For more information, see Standards of medical care in diabetes–2010. *Diabetes Care* 2010;33(suppl 1):S11–S61. To ensure data comparability, the revised definition of undiagnosed diabetes was applied to all data in this table. Also see Appendix II, Diabetes.

³Persons of Hispanic and Mexican origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The non-Hispanic race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group. Prior to data year 1999, estimates were tabulated according to the 1977 Standards. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. See Appendix II, Hispanic origin; Race.

⁴Estimates are age-adjusted to the year 2000 standard population using three age groups: 20–44 years, 45–64 years, and 65 years and over. Age-adjusted estimates in this table may differ from other age-adjusted estimates based on the same data and presented elsewhere if different age groups are used in the adjustment procedure. See Appendix II, Age adjustment.

⁵Includes persons of all other races and Hispanic origins not shown separately.

⁶Percent of poverty level was calculated by dividing family income by the U.S. Department of Health and Human Services' poverty guideline specific to family size, as well as the appropriate year, and state. Persons with unknown percent of poverty level are excluded (6% in 2011–2014). See Appendix II, Family income; Poverty.

NOTES: Excludes pregnant women. Fasting weights were used to obtain estimates of total, physician-diagnosed, and undiagnosed diabetes prevalence. Examination weights were used to obtain the poor glycemic control estimates. Estimates in this table may differ from other estimates based on the same data and presented elsewhere if different weights, age adjustment groups, definitions, or trend adjustments are used. Standard errors are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health and Nutrition Examination Survey. See Appendix I, National Health and Nutrition Examination Survey (NHANES).

Table 41 (page 1 of 3). Severe headache or migraine, low back pain, and neck pain among adults aged 18 and over, by selected characteristics: United States, selected years 1997–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#041>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Severe headache or migraine ¹			Low back pain ¹			Neck pain ¹		
	1997	2010	2014	1997	2010	2014	1997	2010	2014
Percent of adults with pain during the past 3 months									
18 years and over, age-adjusted ^{2,3}	15.8	16.6	15.3	28.2	28.4	28.1	14.7	15.4	14.6
18 years and over, crude ³	16.0	16.4	14.9	28.1	28.8	28.6	14.6	15.8	15.0
Age									
18–44 years	18.7	20.4	18.1	26.1	25.2	24.2	13.3	13.1	12.3
18–24 years	18.7	19.6	17.0	21.9	19.4	20.3	9.8	8.3	8.7
25–44 years	18.7	20.7	18.4	27.3	27.2	25.6	14.3	14.8	13.6
45–64 years	15.8	15.6	14.8	31.3	32.4	32.1	17.0	20.0	18.1
45–54 years	17.8	16.7	16.8	31.3	31.3	31.0	17.3	19.1	18.0
55–64 years	12.7	14.1	12.6	31.2	33.8	33.4	16.6	21.0	18.3
65 years and over	7.0	6.4	7.0	29.5	31.8	33.3	15.0	14.8	15.9
65–74 years	8.2	7.4	7.9	30.2	32.5	33.4	15.0	15.5	16.0
75 years and over	5.4	5.1	5.6	28.6	30.9	33.3	15.0	14.0	15.7
Sex ²									
Male	9.9	11.0	9.8	26.5	26.3	26.0	12.6	13.1	12.3
Female	21.4	22.1	20.5	29.6	30.3	30.0	16.6	17.6	16.8
Sex and age									
Male:									
18–44 years	11.9	13.5	11.3	24.8	23.2	21.6	11.6	11.0	10.6
45–54 years	10.3	10.4	10.9	29.4	29.6	30.8	13.9	16.3	13.7
55–64 years	8.8	9.6	8.4	30.7	32.8	32.6	14.6	17.6	15.4
65–74 years	5.0	5.5	5.8	29.0	28.4	29.7	13.6	12.8	13.6
75 years and over	*2.4	4.0	4.7	22.5	27.4	30.1	12.6	13.0	14.8
Female:									
18–44 years	25.4	27.3	24.7	27.3	27.1	26.7	14.9	15.2	13.9
45–54 years	24.9	22.9	22.5	33.1	33.0	31.1	20.6	21.8	22.0
55–64 years	16.3	18.2	16.5	31.7	34.7	34.2	18.4	24.1	21.0
65–74 years	10.7	9.1	9.8	31.1	36.1	36.5	16.1	17.8	18.2
75 years and over	7.4	5.8	6.3	32.4	33.2	35.6	16.5	14.6	16.4
Race ^{2,4}									
White only	15.9	16.7	15.6	28.7	29.1	28.7	15.1	16.0	15.3
Black or African American only	16.7	18.2	15.1	26.9	27.2	28.4	13.3	13.3	12.1
American Indian or Alaska Native only	18.9	18.8	19.2	33.3	33.6	28.9	16.2	16.9	16.9
Asian only	11.7	10.1	10.1	21.0	19.1	17.6	9.2	9.6	9.4
Native Hawaiian or Other Pacific Islander only	---	*	*	---	*	*	---	*	*
2 or more races	---	21.5	21.3	---	35.6	35.2	---	22.0	21.4
Hispanic origin and race ^{2,4}									
Hispanic or Latino	15.5	16.2	14.9	26.4	27.4	26.9	13.9	15.1	15.0
Mexican	14.6	15.7	14.7	25.2	26.5	26.5	12.9	14.7	14.8
Not Hispanic or Latino	15.9	16.8	15.5	28.4	28.7	28.4	14.9	15.5	14.8
White only	16.1	17.0	16.1	29.1	29.7	29.4	15.4	16.3	15.7
Black or African American only	16.8	18.4	15.1	26.9	27.1	28.0	13.3	13.3	11.8
Education ^{5,6}									
25 years and over:									
No high school diploma or GED	19.2	18.2	16.8	33.6	34.5	34.5	16.5	18.9	17.5
High school diploma or GED	16.0	17.4	15.6	30.2	31.9	32.3	15.5	16.8	15.5
Some college or more	13.8	15.1	14.3	26.9	28.0	26.9	14.6	15.8	15.1

See footnotes at end of table.

Table 41 (page 2 of 3). Severe headache or migraine, low back pain, and neck pain among adults aged 18 and over, by selected characteristics: United States, selected years 1997–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#041>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Severe headache or migraine ¹			Low back pain ¹			Neck pain ¹		
	1997	2010	2014	1997	2010	2014	1997	2010	2014
Percent of poverty level ^{2,7}									
Below 100%	23.3	22.7	21.7	35.4	34.9	36.8	18.6	20.2	19.2
100%–199%	18.9	19.5	17.8	30.8	32.5	31.9	16.1	17.7	17.8
200%–399%	15.5	16.6	15.1	27.9	28.5	28.2	14.8	15.2	14.2
400% or more	12.4	13.3	11.8	24.8	24.7	23.4	12.8	13.1	12.3
Hispanic origin and race and percent of poverty level ^{2,4,7}									
Hispanic or Latino:									
Below 100%	18.9	19.6	19.2	29.5	29.0	31.7	16.4	17.4	17.8
100%–199%	15.7	15.1	15.2	26.8	27.2	26.5	12.9	15.7	16.4
200%–399%	14.0	16.5	14.3	25.0	27.5	23.8	13.8	12.9	11.9
400% or more	13.0	14.0	9.6	21.6	25.6	25.2	12.1	15.3	14.0
Not Hispanic or Latino:									
White only:									
Below 100%	26.1	24.8	24.4	38.9	40.5	41.9	20.5	23.7	22.1
100%–199%	20.4	22.0	21.2	33.3	35.9	37.1	18.0	19.9	21.2
200%–399%	16.3	16.9	16.5	29.1	30.5	30.2	15.9	16.8	15.9
400% or more	12.5	13.8	12.4	25.4	25.2	24.3	13.1	13.6	12.9
Black or African American only:									
Below 100%	22.7	24.0	21.9	34.5	32.5	35.9	17.9	18.6	16.1
100%–199%	17.6	19.6	15.2	27.7	31.2	29.0	14.0	14.4	13.0
200%–399%	14.0	17.6	11.2	24.3	23.7	26.1	10.2	11.7	11.1
400% or more	12.9	12.2	11.8	21.5	21.0	20.4	11.9	8.5	6.8
Disability measure ^{2,8}									
Any basic actions difficulty or complex activity limitation									
Any basic actions difficulty	29.3	30.1	29.2	48.0	49.5	48.6	27.2	28.1	28.4
Any complex activity limitation	30.0	30.9	30.0	49.3	51.1	49.7	27.9	29.0	29.3
Any complex activity limitation	34.6	36.0	32.7	55.1	54.5	56.1	33.1	34.3	34.7
No disability	11.0	11.7	10.8	19.4	19.0	19.3	9.1	9.7	9.2
Geographic region ²									
Northeast	14.5	15.4	13.4	27.1	28.0	25.7	14.0	14.9	13.0
Midwest	15.6	16.8	17.2	28.7	28.1	30.2	15.3	16.0	16.3
South	17.1	18.2	14.8	27.5	28.3	27.2	13.9	14.6	13.3
West	15.3	15.1	15.5	30.0	29.3	29.1	16.1	16.5	16.4
Location of residence ^{2,9}									
Within MSA	15.2	16.3	14.9	27.0	27.5	27.3	14.2	14.9	14.3
Outside MSA	18.1	18.6	18.2	32.5	33.8	32.5	16.4	18.1	16.8

See footnotes at end of table.

Table 41 (page 3 of 3). Severe headache or migraine, low back pain, and neck pain among adults aged 18 and over, by selected characteristics: United States, selected years 1997–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#041>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

-- Data not available.

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.

¹In three separate questions, respondents were asked, "During the past 3 months, did you have a severe headache or migraine? ...low back pain? ...neck pain?" Respondents were instructed to report pain that had lasted a whole day or more, and not to report fleeting or minor aches or pains. Persons may be represented in more than one column.

²Estimates are age-adjusted to the year 2000 standard population using five age groups: 18–44 years, 45–54 years, 55–64 years, 65–74 years, and 75 years and over.

Age-adjusted estimates in this table may differ from other age-adjusted estimates based on the same data and presented elsewhere if different age groups are used in the adjustment procedure. See Appendix II, Age adjustment.

³Includes all other races not shown separately, unknown education level, and unknown disability status.

⁴The race groups, white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The five single-race categories plus multiple-race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group; the category 2 or more races includes persons who reported more than one racial group. Prior to 1999, data were tabulated according to the 1977 Standards with four racial groups, and the Asian only category included Native Hawaiian or Other Pacific Islander. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

⁵Estimates are for persons aged 25 and over and are age-adjusted to the year 2000 standard population using five age groups: 25–44 years, 45–54 years, 55–64 years, 65–74 years, and 75 years and over. See Appendix II, Age adjustment.

⁶GED is General Educational Development high school equivalency diploma. See Appendix II, Education.

⁷Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Missing family income data were imputed for 1997 and beyond. See Appendix II, Family income; Poverty; Table VI.

⁸Any basic actions difficulty or complex activity limitation is defined as having one or more of the following limitations or difficulties: movement difficulty, emotional difficulty, sensory (seeing or hearing) difficulty, cognitive difficulty, self-care (activities of daily living or instrumental activities of daily living) limitation, social limitation, or work limitation. For more information, see Appendix II, Basic actions difficulty; Complex activity limitation. Starting with 2007 data, the hearing question, a component of the basic actions difficulty measure, was revised. Consequently, data prior to 2007 are not comparable with data for 2007 and beyond. For more information on the impact of the revised hearing question, see Appendix II, Hearing trouble.

⁹MSA is metropolitan statistical area. Starting with 2006 data, MSA status is determined using 2000 census data and the 2000 standards for defining MSAs. For data prior to 2006, see Appendix II, Metropolitan statistical area (MSA) for the applicable standards.

NOTES: Standard errors are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey, sample adult questionnaire. See Appendix I, National Health Interview Survey (NHIS).

Table 42 (page 1 of 2). Disability measures among adults aged 18 and over, by selected characteristics: United States, selected years 1997–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#042>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	18 years and over				18–64 years				65 years and over			
	1997	2000	2010 ¹	2014 ¹	1997	2000	2010 ¹	2014 ¹	1997	2000	2010 ¹	2014 ¹
Number, in millions												
At least one basic actions difficulty or complex activity limitation ^{2,3}	60.9	59.0	73.7	74.8	41.3	39.3	50.7	48.7	19.6	19.7	23.0	26.1
At least one basic actions difficulty ²	56.7	55.2	69.2	70.6	38.1	36.4	47.2	45.5	18.6	18.7	22.0	25.0
At least one complex activity limitation ³	29.0	27.2	35.0	35.7	18.1	16.7	22.9	22.1	11.0	10.5	12.1	13.6
At least one basic actions difficulty or complex activity limitation ^{2,3}												
Percent												
Total, age-adjusted ^{4,5}	32.5	29.9	31.9	30.7
Total, crude ⁴	31.8	29.5	32.8	32.4	25.8	23.5	27.1	25.9	62.2	60.8	61.7	60.5
At least one basic actions difficulty ²												
Percent												
Total, age-adjusted ^{4,5}	30.1	27.9	29.9	28.9
Total, crude ⁴	29.4	27.5	30.8	30.5	23.6	21.7	25.1	24.2	58.8	58.1	59.3	58.2
Sex												
Male	25.6	23.8	26.3	26.3	20.7	18.9	21.4	20.8	54.5	53.4	53.8	52.3
Female	32.9	31.0	35.1	34.5	26.4	24.3	28.8	27.4	61.9	61.5	63.6	62.9
Race ⁶												
White only	29.6	28.1	31.2	30.9	23.5	21.8	25.1	24.2	58.5	58.0	59.2	57.7
Black or African American only	31.4	27.2	32.3	32.5	26.9	22.7	28.4	27.6	64.4	60.6	62.9	65.1
American Indian or Alaska Native only	43.8	36.8	41.6	40.3	41.9	34.1	38.5	38.0	66.0	70.2	74.0	55.4
Asian only	15.5	15.5	17.5	17.6	13.0	12.6	12.8	12.3	46.4	44.7	50.1	51.4
Native Hawaiian or Other Pacific Islander only	---	*	*	*	---	*	*	*	---	*	*	*
2 or more races	---	38.0	36.3	34.4	---	34.4	33.9	29.7	---	70.7	65.4	65.8
Hispanic origin and race ⁶												
Hispanic or Latino	23.8	19.6	24.7	24.1	21.0	16.6	21.2	20.5	54.6	57.5	61.5	58.8
Not Hispanic or Latino	30.0	28.5	31.8	31.7	23.9	22.4	25.9	24.9	59.0	58.2	59.1	58.1
White only	30.3	29.1	32.4	32.4	23.8	22.5	26.0	25.3	58.7	58.2	59.0	57.6
Black or African American only	31.5	27.3	32.6	33.0	27.0	22.9	28.6	28.0	64.4	60.4	63.2	65.0
Percent of poverty level ⁷												
Below 100%	41.9	38.4	40.6	41.8	36.2	31.9	36.3	37.3	74.1	71.6	72.7	73.9
100%–199%	38.2	37.1	38.7	37.7	29.2	26.5	30.5	29.3	66.6	69.4	69.5	67.2
200%–399%	28.4	28.2	31.1	31.0	22.0	22.1	24.1	23.1	56.1	53.9	58.9	60.9
400% or more	21.0	19.4	23.0	22.3	18.2	16.8	19.3	17.4	45.5	44.7	47.0	45.8
Location of residence ⁸												
Within MSA	27.7	25.9	29.2	29.0	22.3	20.3	23.6	23.0	56.6	56.7	59.2	57.4
Outside MSA	35.6	33.6	39.3	39.1	28.6	26.8	33.8	31.8	65.8	62.6	59.9	61.5

See footnotes at end of table.

Table 42 (page 2 of 2). Disability measures among adults aged 18 and over, by selected characteristics: United States, selected years 1997–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#042>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	18 years and over				18–64 years				65 years and over			
	1997	2000	2010 ¹	2014 ¹	1997	2000	2010 ¹	2014 ¹	1997	2000	2010 ¹	2014 ¹
At least one complex activity limitation ³												
Percent												
Total, age-adjusted ^{4,5}	15.6	13.7	14.9	14.1
Total, crude ⁴	15.1	13.4	15.5	15.0	11.2	9.8	12.1	11.4	35.1	32.0	32.3	31.0
Sex												
Male	13.7	12.0	14.0	13.5	10.6	9.4	11.3	10.6	31.9	28.1	30.1	27.8
Female	16.5	14.7	16.8	16.4	11.9	10.3	12.9	12.2	37.4	34.9	34.0	33.6
Race ⁶												
White only	15.0	13.6	15.2	14.9	10.9	9.8	11.7	11.2	34.3	31.5	31.7	30.1
Black or African American only	19.0	15.0	19.7	18.1	15.2	11.7	17.0	15.3	47.1	40.4	39.9	37.2
American Indian or Alaska Native only	23.7	20.6	15.4	22.0	22.1	17.8	14.5	19.0	*42.6	*54.9	*	42.7
Asian only	5.7	4.7	7.7	7.3	4.9	3.6	5.0	3.8	*14.8	*15.5	26.7	29.7
Native Hawaiian or Other Pacific Islander only	---	*	*	*	---	*	*	*	---	*	*	*
2 or more races	---	22.5	19.6	19.9	---	20.3	17.0	15.9	---	*42.2	53.6	47.7
Hispanic origin and race ⁶												
Hispanic or Latino	11.9	9.1	10.4	10.6	9.8	7.3	7.9	8.3	33.9	32.4	37.6	33.3
Not Hispanic or Latino	15.5	14.0	16.3	15.8	11.4	10.2	12.9	12.0	35.1	32.0	31.9	30.8
White only	15.4	14.1	16.1	15.9	11.1	10.1	12.5	11.9	34.4	31.5	31.1	30.0
Black or African American only	18.8	15.1	20.0	18.4	15.0	11.7	17.3	15.6	46.8	40.3	40.0	36.6
Percent of poverty level ⁷												
Below 100%	30.0	26.0	27.5	27.8	25.2	22.0	24.0	24.6	56.9	46.7	54.5	51.2
100%–199%	23.3	22.0	23.7	23.3	16.7	15.1	18.4	17.8	43.9	42.8	43.7	42.7
200%–399%	13.3	12.8	14.5	13.7	9.3	9.2	10.8	9.3	30.6	27.5	29.3	30.8
400% or more	7.3	6.4	7.7	7.3	5.8	5.0	5.8	4.9	20.2	19.6	19.8	18.7
Location of residence ⁸												
Within MSA	14.1	12.1	14.2	14.0	10.6	8.9	10.9	10.5	32.7	29.8	31.6	30.5
Outside MSA	19.0	18.2	22.2	21.3	13.6	13.4	18.8	17.3	42.8	38.8	35.2	33.3

... Category not applicable.

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.

--- Data not available.

¹Starting with 2007 data, the hearing question, a component of the basic actions difficulty measure, was revised. Consequently, data for basic actions difficulty prior to 2007 are not comparable with 2007 data and beyond. For more information on the impact of the revised hearing question, see Appendix II, Hearing trouble.

²A basic actions difficulty is defined as having one or more of the following difficulties: movement, emotional, sensory (seeing or hearing), or cognitive. For more information, see Appendix II, Basic actions difficulty. Starting with 2007 data, the hearing question, a component of basic actions difficulty, was revised. Consequently, data prior to 2007 are not comparable with data for 2007 and beyond. For more information on the impact of the revised hearing question, see Appendix II, Hearing trouble.

³A complex activity limitation is defined as having one or more of the following limitations: self-care (activities of daily living or instrumental activities of daily living), social, or work. For more information, see Appendix II, Complex activity limitation.

⁴Includes all other races not shown separately.

⁵Estimates are age-adjusted to the year 2000 standard population using five age groups: 18–44 years, 45–54 years, 55–64 years, 65–74 years, and 75 years and over. See Appendix II, Age adjustment.

⁶The race groups, white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The five single-race categories plus multiple-race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group; the category 2 or more races includes persons who reported more than one racial group. Prior to 1999, data were tabulated according to the 1977 Standards with four racial groups, and the Asian only category included Native Hawaiian or Other Pacific Islander. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

⁷Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Missing family income data were imputed for 1997 and beyond. See Appendix II, Family income; Poverty; Table VI.

⁸MSA is metropolitan statistical area. Starting with 2006 data, MSA status is determined using 2000 census data and the 2000 standards for defining MSAs. For data prior to 2006, see Appendix II, Metropolitan statistical area (MSA) for the applicable standards.

NOTES: Standard errors are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey, sample adult questionnaire. See Appendix I, National Health Interview Survey (NHIS).

Table 43 (page 1 of 2). Vision limitations among adults aged 18 and over, by selected characteristics: United States, selected years 1997–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#043>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Any trouble seeing, even with glasses or contacts ¹									
	1997	2000	2005	2007	2009	2010	2011	2012	2013	2014
	Percent of adults									
18 years and over, age-adjusted ^{2,3}	10.0	9.0	9.2	9.9	8.3	9.1	8.8	8.4	8.7	8.7
18 years and over, crude ³	9.8	8.9	9.3	10.0	8.6	9.4	9.2	8.8	9.1	9.1
Age										
18–44 years	6.2	5.3	5.5	6.9	5.3	6.2	5.5	5.4	5.5	5.6
18–24 years	5.4	4.2	5.0	6.9	4.8	5.8	5.2	5.1	5.5	4.9
25–44 years	6.5	5.7	5.7	6.8	5.6	6.3	5.6	5.5	5.5	5.9
45–64 years	12.0	10.7	11.2	12.2	10.8	11.6	12.0	11.3	11.1	11.3
45–54 years	12.2	10.9	11.0	12.3	10.5	10.7	11.7	11.2	10.6	11.5
55–64 years	11.6	10.5	11.5	12.1	11.2	12.7	12.4	11.5	11.7	11.0
65 years and over	18.1	17.4	17.4	15.3	13.1	13.9	13.6	12.7	14.3	13.5
65–74 years	14.2	13.6	13.2	12.9	10.3	12.2	12.2	11.0	11.5	11.5
75 years and over	23.1	21.9	22.0	17.9	16.5	16.1	15.2	14.9	18.0	16.5
Sex ²										
Male	8.8	7.9	7.9	8.5	7.2	7.9	7.6	7.1	7.5	7.6
Female	11.1	10.1	10.5	11.2	9.3	10.3	10.1	9.7	9.8	9.8
Sex and age										
Male:										
18–44 years	5.3	4.4	4.5	5.6	4.5	5.2	4.2	4.4	4.5	4.3
45–54 years	10.1	8.8	8.8	10.6	9.1	9.1	10.4	9.3	9.4	10.7
55–64 years	10.5	9.5	10.5	10.0	9.7	10.7	11.8	9.8	10.4	9.6
65–74 years	13.2	12.8	11.4	11.4	9.3	10.5	9.7	9.9	10.9	10.3
75 years and over	21.4	20.7	20.4	17.2	15.1	15.7	14.9	12.8	14.7	17.0
Female:										
18–44 years	7.1	6.2	6.5	8.1	6.2	7.1	6.9	6.4	6.5	7.0
45–54 years	14.2	12.8	13.2	13.9	11.9	12.3	13.0	12.9	11.8	12.2
55–64 years	12.6	11.5	12.4	14.2	12.6	14.6	13.0	13.1	12.9	12.3
65–74 years	15.0	14.4	14.8	14.2	11.2	13.6	14.5	11.9	12.1	12.5
75 years and over	24.2	22.7	23.0	18.4	17.4	16.4	15.4	16.4	20.3	16.1
Race ^{2,4}										
White only	9.7	8.8	9.1	9.9	8.1	8.8	8.6	8.4	8.7	8.5
Black or African American only	12.8	10.6	10.9	10.5	10.4	12.1	10.8	9.2	10.0	11.1
American Indian or Alaska Native only	19.2	16.6	*14.9	18.0	*12.3	15.0	15.0	13.0	13.7	16.9
Asian only	6.2	6.3	5.5	5.7	5.5	5.3	6.3	5.7	4.9	5.7
Native Hawaiian or Other Pacific Islander only	---	*	*	*	*	*	*	*	*	*
2 or more races	---	16.2	16.4	16.9	14.8	13.1	12.4	15.6	11.8	11.4
Hispanic origin and race ^{2,4}										
Hispanic or Latino	10.0	9.7	9.6	9.9	8.7	9.2	9.4	9.4	9.7	8.8
Mexican	10.2	8.3	9.9	10.1	8.7	9.0	10.4	9.3	10.9	9.3
Not Hispanic or Latino	10.0	9.1	9.2	10.0	8.3	9.2	8.8	8.4	8.6	8.8
White only	9.8	8.9	9.1	10.1	8.1	8.9	8.6	8.4	8.6	8.5
Black or African American only	12.8	10.6	10.9	10.6	10.5	12.2	10.7	9.3	10.1	11.3
Education ^{5,6}										
25 years of age and over:										
No high school diploma or GED	15.0	12.2	13.5	13.4	12.6	14.1	13.9	12.9	12.8	13.1
High school diploma or GED	10.6	9.5	10.3	10.9	9.2	10.5	10.4	9.3	10.0	9.3
Some college or more	8.9	8.9	8.6	9.2	7.6	8.0	7.9	7.9	8.0	8.4

See footnotes at end of table.

Table 43 (page 2 of 2). Vision limitations among adults aged 18 and over, by selected characteristics: United States, selected years 1997–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#043>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Any trouble seeing, even with glasses or contacts ¹									
	1997	2000	2005	2007	2009	2010	2011	2012	2013	2014
Percent of poverty level ^{2,7}										
						Percent of adults				
Below 100%	17.0	12.9	15.3	15.0	14.3	14.8	14.2	13.7	15.6	13.2
100%–199%	12.9	11.6	11.5	13.0	11.1	12.2	11.5	10.9	11.2	10.9
200%–399%	9.1	8.8	8.9	9.4	8.0	9.0	8.7	7.9	7.6	8.8
400% or more	7.3	7.1	6.9	7.8	5.7	6.4	6.0	6.1	6.4	6.4
Hispanic origin and race and percent of poverty level ^{2,4,7}										
Hispanic or Latino:										
Below 100%	12.8	11.0	13.6	13.4	12.2	10.8	13.9	13.1	13.1	10.2
100%–199%	11.2	9.4	8.8	11.1	8.1	10.8	9.6	10.0	10.6	9.3
200%–399%	8.1	9.2	8.2	7.2	9.0	8.9	8.3	6.8	7.4	9.2
400% or more	*8.1	10.5	8.0	10.6	*4.6	5.3	5.1	7.8	9.0	6.6
Not Hispanic or Latino:										
White only:										
Below 100%	17.9	13.1	16.2	16.3	13.4	16.8	14.4	14.5	17.7	14.6
100%–199%	13.1	12.0	12.7	14.2	12.1	12.6	12.3	11.7	11.8	11.3
200%–399%	9.2	9.2	9.0	10.3	8.3	8.8	9.0	8.5	7.8	8.7
400% or more	7.3	7.0	6.9	7.7	5.8	6.7	5.9	6.0	6.4	6.3
Black or African American only:										
Below 100%	17.9	13.6	16.0	15.1	17.8	15.8	15.5	13.7	15.3	14.2
100%–199%	16.0	12.9	11.3	14.0	11.7	14.9	12.3	11.3	11.4	12.1
200%–399%	9.3	7.7	9.7	7.3	8.1	12.0	8.5	6.8	8.0	10.8
400% or more	7.7	8.3	6.4	6.9	5.6	6.6	8.6	6.4	6.7	8.6
Geographic region ²										
Northeast	8.6	7.4	8.1	8.1	7.3	7.8	7.6	6.4	7.4	7.0
Midwest	9.5	9.6	9.7	10.3	8.2	9.1	8.7	8.7	9.0	9.0
South	11.4	9.2	9.8	10.1	8.7	10.6	9.4	9.1	8.9	9.1
West	9.7	9.9	8.6	10.5	8.6	8.0	9.1	8.9	9.1	9.3
Location of residence ^{2,8}										
Within MSA	9.5	8.5	8.6	9.6	8.2	8.6	8.6	8.2	8.4	8.4
Outside MSA	12.0	11.1	11.7	11.4	9.0	11.6	10.3	9.8	10.6	10.3

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.

--- Data not available.

¹ Respondents were asked, “Do you have any trouble seeing, even when wearing glasses or contact lenses?” Respondents were also asked, “Are you blind or unable to see at all?” In this analysis, any trouble seeing and blind are combined into one category.

² Estimates are age-adjusted to the year 2000 standard population using five age groups: 18–44 years, 45–54 years, 55–64 years, 65–74 years, and 75 years and over.

Age-adjusted estimates in this table may differ from other age-adjusted estimates based on the same data and presented elsewhere if different age groups are used in the adjustment procedure. See Appendix II, Age adjustment.

³ Includes all other races not shown separately and unknown education level.

⁴ The race groups, white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The five single-race categories plus multiple-race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group; the category 2 or more races includes persons who reported more than one racial group. Prior to 1999, data were tabulated according to the 1977 Standards with four racial groups, and the Asian only category included Native Hawaiian or Other Pacific Islander. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

⁵ Estimates are for persons aged 25 and over and are age-adjusted to the year 2000 standard population using five age groups: 25–44 years, 45–54 years, 55–64 years, 65–74 years, and 75 years and over. See Appendix II, Age adjustment.

⁶ GED is General Educational Development high school equivalency diploma. See Appendix II, Education.

⁷ Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Missing family income data were imputed for 1997 and beyond. See Appendix II, Family income; Poverty; Table VI.

⁸ MSA is metropolitan statistical area. Starting with 2006 data, MSA status is determined using 2000 census data and the 2000 standards for defining MSAs. For data prior to 2006, see Appendix II, Metropolitan statistical area (MSA) for the applicable standards.

NOTES: Standard errors are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey, sample adult questionnaire. See Appendix I, National Health Interview Survey (NHIS).

Table 44. (page 1 of 2). Hearing limitations among adults aged 18 and over, by selected characteristics: United States, selected years 2007–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#044>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Level of hearing trouble											
	Any hearing trouble (a little, moderate, a lot of trouble, or deaf) ¹				Moderate, a lot of trouble, or deaf ¹				A lot of trouble or deaf ¹			
	2007	2010	2013	2014	2007	2010	2013	2014	2007	2010	2013	2014
	Percent of adults											
18 years and over, age-adjusted ^{2,3}	14.7	15.6	14.3	15.6	5.6	5.7	5.4	5.8	2.3	2.1	2.0	2.3
18 years and over, crude ³	14.9	16.2	15.3	16.8	5.7	5.9	5.7	6.3	2.3	2.2	2.1	2.4
Age												
18–44 years	6.0	6.7	5.3	6.3	1.3	1.7	1.3	1.7	0.4	0.5	0.4	0.6
18–24 years	4.1	5.4	3.4	4.1	*	*1.2	*1.0	*	*	*	*	*
25–44 years	6.6	7.2	6.0	7.2	1.6	1.8	1.5	1.9	0.5	0.5	0.4	0.6
45–64 years	17.6	18.9	17.3	19.2	6.0	6.1	5.6	5.9	2.0	1.9	1.7	2.0
45–54 years	14.7	15.6	13.6	15.4	4.1	4.8	3.9	3.9	1.2	1.2	1.1	1.3
55–64 years	21.5	23.2	21.5	23.3	8.5	7.8	7.5	8.2	3.0	2.7	2.4	2.7
65 years and over	36.9	37.5	37.3	38.7	18.6	17.7	17.3	18.5	8.7	7.6	7.4	7.8
65–74 years	29.8	31.2	29.7	33.1	11.9	12.9	12.1	13.5	4.7	4.6	4.4	5.0
75 years and over	45.0	45.2	47.7	46.7	26.3	23.7	24.4	25.7	13.3	11.1	11.4	11.8
Sex ²												
Male	18.3	18.9	17.3	19.1	7.7	7.4	6.9	7.7	3.1	2.8	2.5	2.9
Female	11.5	12.7	11.7	12.6	3.9	4.3	4.1	4.3	1.6	1.6	1.6	1.7
Sex and age												
Male:												
18–44 years	6.8	7.7	6.1	7.4	1.6	1.9	1.6	2.0	*0.5	*0.7	*0.4	*0.6
45–54 years	18.7	18.2	16.4	18.1	5.3	5.7	5.0	4.9	1.5	*1.1	*1.4	1.4
55–64 years	28.4	30.1	27.4	30.2	12.9	11.5	10.1	10.9	4.7	3.9	3.1	3.8
65–74 years	39.4	41.0	38.7	41.9	17.7	17.9	16.5	19.3	7.0	6.7	6.0	6.9
75 years and over	54.6	53.1	54.8	57.0	34.8	29.7	30.6	33.2	16.9	14.5	14.1	15.5
Female:												
18–44 years	5.1	5.8	4.5	5.3	1.0	1.4	1.1	1.3	*0.3	*0.3	*0.4	*0.5
45–54 years	11.0	13.0	10.8	12.8	2.9	3.9	2.9	2.9	*1.0	*1.3	*0.9	*1.2
55–64 years	15.0	16.7	16.1	16.9	4.3	4.4	5.1	5.7	*1.3	1.6	1.7	1.7
65–74 years	21.8	22.8	21.8	25.5	6.9	8.6	8.3	8.4	2.8	2.9	3.1	3.3
75 years and over	38.8	39.9	42.9	39.5	20.9	19.7	20.2	20.4	11.1	8.9	9.5	9.2
Race ^{2,4}												
White only	15.6	16.5	15.3	16.7	6.0	6.1	5.8	6.3	2.4	2.3	2.2	2.4
Black or African American only	8.5	10.3	9.4	10.3	2.7	3.3	2.8	3.5	1.2	1.1	1.0	1.2
American Indian or Alaska Native only	17.9	21.1	18.8	13.5	*8.8	*7.1	*7.4	*8.2	*3.8	*	*	*
Asian only	8.0	8.0	8.2	9.5	2.8	2.6	2.6	2.7	*	*1.0	*1.1	*1.3
Native Hawaiian or Other Pacific Islander only	*	*	*	*	*	*	*	*	*	*	*	*
2 or more races	24.4	23.3	17.1	20.1	11.5	8.7	6.6	7.1	*4.9	*	*1.4	*
Hispanic origin and race ^{2,4}												
Hispanic or Latino	10.9	10.9	9.6	12.2	4.3	3.5	3.4	4.7	2.5	1.4	1.3	1.9
Mexican	11.8	11.5	10.8	13.8	4.4	3.5	3.8	6.2	2.5	*1.5	1.5	2.6
Not Hispanic or Latino	15.2	16.2	15.0	16.2	5.8	6.0	5.6	6.0	2.3	2.2	2.1	2.3
White only	16.5	17.5	16.2	17.6	6.3	6.5	6.2	6.5	2.5	2.4	2.3	2.5
Black or African American only	8.4	10.3	9.4	10.4	2.7	3.3	2.8	3.4	1.2	1.1	1.0	1.2
Education ^{5,6}												
25 years and over:												
No high school diploma or GED	17.9	19.7	17.8	17.2	7.6	8.1	6.7	6.7	4.1	3.2	3.0	3.3
High school diploma or GED	17.2	18.1	17.7	18.9	6.4	6.5	6.6	7.4	2.8	2.5	2.5	2.8
Some college or more	15.4	16.2	14.9	16.8	6.1	6.0	5.7	6.1	1.9	2.0	1.9	2.1

See footnotes at end of table.

Table 44 (page 2 of 2). Hearing limitations among adults aged 18 and over, by selected characteristics: United States, selected years 2007–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#044>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Level of hearing trouble											
	Any hearing trouble (a little, moderate, a lot of trouble, or deaf) ¹				Moderate, a lot of trouble, or deaf ¹				A lot of trouble or deaf ¹			
	2007	2010	2013	2014	2007	2010	2013	2014	2007	2010	2013	2014
Percent of poverty level ^{2,7}						Percent of adults						
Below 100%	16.2	16.7	16.3	16.5	6.8	6.8	6.3	6.3	3.4	2.7	2.5	2.8
100%–199%	15.5	17.2	15.2	16.6	5.8	6.6	5.5	6.3	2.8	2.5	2.4	2.4
200%–399%	15.1	15.7	14.6	16.0	5.8	5.6	5.5	6.2	2.4	2.1	2.1	2.3
400% or more	13.6	14.5	13.1	14.7	5.3	5.0	4.9	5.2	1.6	1.8	1.6	2.0
Hispanic origin and race and percent of poverty level ^{2,4,7}												
Hispanic or Latino:												
Below 100%	12.7	9.1	10.4	13.0	*6.1	*3.5	3.9	4.4	*	*	*	*
100%–199%	9.7	11.8	9.4	12.0	*3.1	4.3	3.1	4.6	*2.1	*2.3	*1.6	*2.1
200%–399%	9.8	10.3	9.0	12.2	*3.9	*2.6	4.0	5.0	*	*	*	*
400% or more	13.3	12.4	10.8	12.3	*5.6	*3.2	*2.9	*5.6	*	*	*	*
Not Hispanic or Latino:												
White only:												
Below 100%	20.9	21.7	20.6	19.9	8.8	9.2	8.1	8.2	4.3	3.7	3.2	3.9
100%–199%	18.8	20.8	18.3	21.0	7.2	8.3	7.0	7.8	3.3	3.0	2.9	2.9
200%–399%	17.2	17.9	17.0	18.0	6.4	6.5	6.3	7.0	2.6	2.3	2.4	2.5
400% or more	14.3	15.4	14.2	16.0	5.6	5.4	5.4	5.7	1.7	2.0	1.8	2.1
Black or African American only:												
Below 100%	9.3	11.6	13.1	13.9	*2.8	4.0	4.7	4.5	*	*1.5	*1.4	*1.9
100%–199%	9.8	11.1	12.2	11.3	*3.1	3.0	2.8	3.9	*	*0.7	*1.1	*1.1
200%–399%	7.8	10.4	7.0	9.4	*2.2	3.6	2.4	*2.9	*	*	*	*
400% or more	7.1	7.7	5.2	6.6	*	*2.8	*	*	*	*	*	*
Geographic region ²												
Northeast	13.3	13.9	13.1	12.3	5.2	4.4	5.0	4.5	1.7	1.4	1.8	1.6
Midwest	16.0	17.5	16.5	17.4	6.1	6.3	6.4	6.1	2.3	2.3	2.3	2.3
South	14.0	16.0	13.8	16.1	5.4	6.3	5.0	6.1	2.5	2.6	2.1	2.3
West	15.5	14.4	14.1	15.8	5.9	5.3	5.2	6.2	2.4	1.9	1.8	2.6
Location of residence ^{2,8}												
Within MSA	14.0	14.7	13.5	14.9	5.3	5.4	5.0	5.5	2.1	1.9	1.8	2.1
Outside MSA	18.0	20.1	18.4	19.8	7.2	7.5	7.3	7.3	3.3	3.0	2.9	2.9

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.

¹Starting in 2007, respondents were asked questions about their hearing WITHOUT the use of hearing aids or other listening devices. “Is your hearing excellent, good, a little trouble hearing, moderate trouble, a lot of trouble, or are you deaf?” See Appendix II, Hearing trouble.

²Estimates are age-adjusted to the year 2000 standard population using five age groups: 18–44 years, 45–54 years, 55–64 years, 65–74 years, and 75 years and over. Age-adjusted estimates in this table may differ from other age-adjusted estimates based on the same data and presented elsewhere if different age groups are used in the adjustment procedure. See Appendix II, Age adjustment.

³Includes all other races not shown separately and unknown education level.

⁴The race groups, white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The five single-race categories plus multiple-race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group; the category 2 or more races includes persons who reported more than one racial group. Prior to 1999, data were tabulated according to the 1977 Standards with four racial groups, and the Asian only category included Native Hawaiian or Other Pacific Islander. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

⁵Estimates are for persons aged 25 and over and are age-adjusted to the year 2000 standard population using five age groups: 25–44 years, 45–54 years, 55–64 years, 65–74 years, and 75 years and over. See Appendix II, Age adjustment.

⁶GED is General Educational Development high school equivalency diploma. See Appendix II, Education.

⁷Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Missing family income data were imputed for 1997 and beyond. See Appendix II, Family income; Poverty; Table VI.

⁸MSA is metropolitan statistical area. Starting with 2006 data, MSA status is determined using 2000 census data and the 2000 standards for defining MSAs. For data prior to 2006, see Appendix II, Metropolitan statistical area (MSA) for the applicable standards.

NOTES: Starting with *Health, United States, 2013*, the hearing measures shown in this table were revised to provide a consistent definition over time. For a longer trend, see *Health, United States, 2012*. Available from: <http://www.cdc.gov/nchs/hus.htm>. Standard errors are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey, sample adult questionnaire. See Appendix I, National Health Interview Survey (NHIS).

Table 45 (page 1 of 2). Respondent-assessed fair-poor health status, by selected characteristics: United States, selected years 1991–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#045>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	1991 ¹	1995 ¹	1997	2000	2005	2010	2013	2014
Percent of persons with fair or poor health ²								
All ages, age-adjusted ^{3,4}	10.4	10.6	9.2	9.0	9.2	9.6	9.4	8.9
All ages, crude ⁴	10.0	10.1	8.9	8.9	9.3	10.1	10.2	9.8
Age								
Under 18 years	2.6	2.6	2.1	1.7	1.8	2.0	1.7	1.6
Under 6 years	2.7	2.7	1.9	1.5	1.6	1.8	1.6	1.3
6–17 years	2.6	2.5	2.1	1.8	1.9	2.2	1.8	1.8
18–44 years	6.1	6.6	5.3	5.1	5.5	6.3	6.2	6.1
18–24 years	4.8	4.5	3.4	3.3	3.3	3.9	3.6	3.7
25–44 years	6.4	7.2	5.9	5.7	6.3	7.2	7.2	7.0
45–54 years	13.4	13.4	11.7	11.9	11.6	13.3	14.1	12.8
55–64 years	20.7	21.4	18.2	17.9	18.3	19.4	19.2	18.4
65 years and over	29.0	28.3	26.7	26.9	26.6	24.4	23.1	21.7
65–74 years	26.0	25.6	23.1	22.5	23.4	21.2	19.7	19.5
75 years and over	33.6	32.2	31.5	32.1	30.2	28.3	27.6	24.9
Sex ³								
Male	10.0	10.1	8.8	8.8	8.8	9.2	9.0	8.7
Female	10.8	11.1	9.7	9.3	9.5	10.0	9.8	9.2
Race ^{3,5}								
White only	9.6	9.7	8.3	8.2	8.6	8.8	8.7	8.3
Black or African American only	16.8	17.2	15.8	14.6	14.3	14.9	14.3	13.6
American Indian or Alaska Native only	18.3	18.7	17.3	17.2	13.2	17.8	15.3	14.1
Asian only	7.8	9.3	7.8	7.4	6.8	8.1	7.7	7.3
Native Hawaiian or Other Pacific Islander only	---	---	---	*	*	*	*	*
2 or more races	---	---	---	16.2	14.5	15.6	13.9	12.8
Black or African American; White	---	---	---	*14.5	8.3	*16.7	*19.0	11.9
American Indian or Alaska Native; White	---	---	---	18.7	17.2	19.0	16.2	17.6
Hispanic origin and race ^{3,5}								
Hispanic or Latino	15.6	15.1	13.0	12.8	13.3	13.1	12.7	12.2
Mexican	17.0	16.7	13.1	12.8	14.3	13.7	13.3	13.0
Not Hispanic or Latino	10.0	10.1	8.9	8.7	8.7	9.2	9.0	8.5
White only	9.1	9.1	8.0	7.9	8.0	8.2	8.2	7.7
Black or African American only	16.8	17.3	15.8	14.6	14.4	14.9	14.2	13.6
Percent of poverty level ^{3,6}								
Below 100%	22.8	23.7	20.8	19.6	20.4	20.9	21.8	19.8
100%–199%	14.7	15.5	13.9	14.1	14.4	15.2	14.4	14.2
200%–399%	7.9	7.9	8.2	8.4	8.3	8.3	8.2	7.9
400% or more	4.9	4.7	4.1	4.5	4.7	4.3	4.0	3.9
Hispanic origin and race and percent of poverty level ^{3,5,6}								
Hispanic or Latino:								
Below 100%	23.6	22.7	19.9	18.7	20.2	19.2	20.3	18.7
100%–199%	18.0	16.9	13.5	15.3	15.3	15.6	14.3	14.1
200%–399%	10.3	10.1	10.0	10.3	10.3	10.3	10.3	9.4
400% or more	6.6	4.0	5.7	5.5	7.6	6.4	5.3	5.5
Not Hispanic or Latino:								
White only:								
Below 100%	21.9	22.8	19.7	18.8	20.1	20.9	22.2	20.3
100%–199%	14.0	14.8	13.3	13.4	13.8	14.8	14.4	14.1
200%–399%	7.5	7.3	7.7	7.9	7.9	7.7	7.7	7.4
400% or more	4.7	4.6	3.9	4.2	4.3	4.0	3.8	3.5
Black or African American only:								
Below 100%	25.8	27.7	25.3	23.8	23.3	23.9	24.8	21.8
100%–199%	17.0	19.3	19.2	18.2	17.6	18.3	17.5	16.8
200%–399%	12.0	11.4	12.2	11.7	11.2	11.2	9.8	9.8
400% or more	5.9	6.5	6.1	7.3	7.1	6.8	5.0	5.7

See footnotes at end of table.

Table 45 (page 2 of 2). Respondent-assessed fair-poor health status, by selected characteristics: United States, selected years 1991–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#045>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	1991 ¹	1995 ¹	1997	2000	2005	2010	2013	2014
Disability measure among adults 18 years and over ^{3,7}								
Percent of persons with fair or poor health ²								
Any basic actions difficulty or complex activity limitation	---	---	27.0	27.6	28.5	28.7	30.5	28.8
Any basic actions difficulty	---	---	27.3	27.7	29.1	28.9	30.7	29.1
Any complex activity limitation	---	---	42.9	45.6	46.3	46.0	47.8	47.6
No disability	---	---	3.4	3.8	3.6	3.5	3.8	3.8
Geographic region ³								
Northeast	8.3	9.1	8.0	7.6	7.5	7.9	8.2	7.2
Midwest	9.1	9.7	8.1	8.0	8.3	9.0	8.8	8.5
South	13.1	12.3	10.8	10.7	11.0	11.1	10.6	10.2
West	9.7	10.1	8.8	8.8	8.6	9.2	9.1	8.6
Location of residence ^{3,8}								
Within MSA	9.9	10.1	8.7	8.5	8.7	9.2	9.1	8.5
Outside MSA	11.9	12.6	11.1	11.1	11.2	11.9	11.4	11.4

--- Data not available.

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.

¹Data prior to 1997 are not strictly comparable with data for later years due to the 1997 questionnaire redesign. See Appendix I, National Health Interview Survey (NHIS).

²See Appendix II, Health status, respondent-assessed.

³Estimates are age-adjusted to the year 2000 standard population using six age groups: under 18 years, 18–44 years, 45–54 years, 55–64 years, 65–74 years, and 75 years and over. The disability measure is age-adjusted using the five adult age groups. See Appendix II, Age adjustment.

⁴Includes all other races not shown separately and unknown disability status.

⁵The race groups, white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The five single-race categories plus multiple-race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group; the category 2 or more races includes persons who reported more than one racial group. Prior to 1999, data were tabulated according to the 1977 Standards with four racial groups, and the Asian only category included Native Hawaiian or Other Pacific Islander. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

⁶Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Missing family income data were imputed for 1991 and beyond. See Appendix II, Family income; Poverty; Table VI.

⁷Any basic actions difficulty or complex activity limitation is defined as having one or more of the following limitations or difficulties: movement difficulty, emotional difficulty, sensory (seeing or hearing) difficulty, cognitive difficulty, self-care (activities of daily living or instrumental activities of daily living) limitation, social limitation, or work limitation. For more information, see Appendix II, Basic actions difficulty; Complex activity limitation. Starting with 2007 data, the hearing question, a component of the basic actions difficulty measure, was revised. Consequently, data prior to 2007 are not comparable with data for 2007 and beyond. For more information on the impact of the revised hearing question, see Appendix II, Hearing trouble.

⁸MSA is metropolitan statistical area. Starting with 2006 data, MSA status is determined using 2000 census data and the 2000 standards for defining MSAs. For data prior to 2006, see Appendix II, Metropolitan statistical area (MSA) for the applicable standards.

NOTES: Standard errors for selected years are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey, family core and sample adult questionnaires. See Appendix I, National Health Interview Survey (NHIS).

Table 46 (page 1 of 2). Serious psychological distress in the past 30 days among adults aged 18 and over, by selected characteristics: United States, average annual, selected years 1997–1998 through 2013–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#046>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	1997–1998	1999–2000	2001–2002	2004–2005	2010–2011	2013–2014 ¹
Percent of adults with serious psychological distress ²						
18 years and over, age-adjusted ^{3,4}	3.2	2.6	3.1	3.0	3.3	3.4
18 years and over, crude ⁴	3.2	2.6	3.1	3.0	3.4	3.5
Age						
18–44 years	2.9	2.3	2.9	2.8	2.9	3.1
18–24 years	2.7	2.2	2.8	2.5	2.4	2.5
25–44 years	3.0	2.4	3.0	2.9	3.1	3.4
45–64 years	3.7	3.2	3.9	3.7	4.5	4.5
45–54 years	3.9	3.5	4.2	3.9	4.2	4.6
55–64 years	3.4	2.6	3.4	3.4	4.7	4.5
65 years and over	3.1	2.4	2.4	2.5	2.4	2.5
65–74 years	2.5	2.3	2.4	2.2	2.6	2.7
75 years and over	3.8	2.5	2.4	2.9	2.1	2.2
Sex ³						
Male	2.5	2.0	2.4	2.3	2.8	2.9
Female	3.8	3.1	3.8	3.7	3.7	3.9
Race ^{3,5}						
White only	3.1	2.5	3.0	2.9	3.2	3.4
Black or African American only	4.0	2.9	3.5	3.6	3.7	3.5
American Indian or Alaska Native only	7.8	*7.2	8.1	*3.5	5.6	*5.4
Asian only	2.0	*1.4	*1.8	1.7	1.7	1.9
Native Hawaiian or Other Pacific Islander only	---	*	*	*	*	*
2 or more races	---	4.8	5.0	7.9	5.6	8.6
Hispanic origin and race ^{3,5}						
Hispanic or Latino	5.0	3.5	4.0	3.7	4.0	4.5
Mexican	5.2	2.9	3.8	3.6	3.6	4.6
Not Hispanic or Latino	3.0	2.5	3.1	3.0	3.2	3.3
White only	2.9	2.4	3.0	2.9	3.2	3.2
Black or African American only	3.9	2.9	3.5	3.6	3.7	3.4
Percent of poverty level ^{3,6}						
Below 100%	9.1	6.8	8.4	8.6	8.2	9.1
100%–199%	5.0	4.4	5.2	5.0	5.0	5.5
200%–399%	2.5	2.3	2.8	2.5	2.9	2.6
400% or more	1.3	1.2	1.3	1.1	1.2	1.2
Hispanic origin and race and percent of poverty level ^{3,5,6}						
Hispanic or Latino:						
Below 100%	8.6	6.1	7.5	6.6	7.5	8.2
100%–199%	5.4	3.8	4.1	3.9	4.3	5.0
200%–399%	3.4	2.1	3.5	2.6	3.1	3.2
400% or more	*	2.3	*	*1.9	*1.4	*1.8
Not Hispanic or Latino:						
White only:						
Below 100%	9.6	7.8	9.2	10.2	9.6	10.7
100%–199%	5.2	4.9	5.9	5.6	5.6	6.5
200%–399%	2.5	2.3	2.9	2.6	3.2	2.7
400% or more	1.3	1.1	1.3	1.1	1.1	1.2
Black or African American only:						
Below 100%	8.7	6.0	7.2	7.6	7.7	7.4
100%–199%	4.3	3.6	4.9	4.8	4.4	3.5
200%–399%	2.2	*1.7	2.3	2.1	1.9	2.3
400% or more	*	*1.0	*	*	*1.5	*0.7

See footnotes at end of table.

Table 46 (page 2 of 2). Serious psychological distress in the past 30 days among adults aged 18 and over, by selected characteristics: United States, average annual, selected years 1997–1998 through 2013–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#046>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	1997–1998	1999–2000	2001–2002	2004–2005	2010–2011	2013–2014 ¹
Geographic region ³						
Percent of adults with serious psychological distress ²						
Northeast	2.7	1.9	2.8	2.5	3.0	2.8
Midwest	2.6	2.5	2.9	2.7	3.1	3.5
South	3.8	2.9	3.5	3.7	3.6	3.7
West	3.3	2.8	3.0	2.8	3.3	3.4
Location of residence ^{3,7}						
Within MSA	3.0	2.3	3.0	2.8	3.1	3.3
Outside MSA	3.9	3.5	3.8	4.0	4.0	4.3

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.

-- Data not available.

¹In 2013, the six psychological distress questions were moved to the adult selected items section of the sample adult questionnaire. Observed differences between the 2012 and earlier estimates and the 2013 and later estimates may be partially or fully attributable to this change in question placement within the sample adult questionnaire.

²Serious psychological distress is measured by a six-question scale that asks respondents how often they experienced each of the six symptoms of psychological distress in the past 30 days. Respondents must have answered all six questions to have a computed K6 score. Only those with K6 scores were included in this analysis. See Appendix II, Serious psychological distress.

³Estimates are age-adjusted to the year 2000 standard population using five age groups: 18–44 years, 45–54 years, 55–64 years, 65–74 years, and 75 years and over. See Appendix II, Age adjustment.

⁴Includes all other races not shown separately.

⁵The race groups, white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The five single-race categories plus multiple-race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group; the category 2 or more races includes persons who reported more than one racial group. Prior to 1999, data were tabulated according to the 1977 Standards with four racial groups, and the Asian only category included Native Hawaiian or Other Pacific Islander. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

⁶Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Missing family income data were imputed for 1997 and beyond. See Appendix II, Family income; Poverty; Table VI.

⁷MSA is metropolitan statistical area. Starting with 2006–2007 data (shown in spreadsheet), MSA status is determined using 2000 census data and the 2000 standards for defining MSAs. For data prior to 2006, see Appendix II, Metropolitan statistical area (MSA) for the applicable standards.

NOTES: Standard errors for selected years are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey, sample adult questionnaire. See Appendix I, National Health Interview Survey (NHIS).

Table 47 (page 1 of 2). Current cigarette smoking among adults aged 18 and over, by sex, race, and age: United States: selected years 1965–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#047>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

<i>Sex, race, and age</i>	1965 ¹	1979 ¹	1985 ¹	1990 ¹	2000	2004	2005	2010	2012	2013	2014
18 years and over, age-adjusted ²											
Percent of adults who were current cigarette smokers ³											
All persons	41.9	33.3	29.9	25.3	23.1	20.8	20.8	19.3	18.2	17.9	17.0
Male	51.2	37.0	32.2	28.0	25.2	23.0	23.4	21.2	20.6	20.5	19.0
Female	33.7	30.1	27.9	22.9	21.1	18.7	18.3	17.5	15.9	15.5	15.1
White male ⁴	50.4	36.4	31.3	27.6	25.4	23.0	23.3	21.4	20.7	20.5	18.8
Black or African American male ⁴	58.8	43.9	40.2	32.8	25.7	23.5	25.9	23.3	22.0	21.8	21.7
White female ⁴	33.9	30.3	27.9	23.5	22.0	19.5	19.1	18.3	16.9	16.3	16.0
Black or African American female ⁴	31.8	30.5	30.9	20.8	20.7	16.9	17.1	16.6	14.2	14.9	13.4
18 years and over, crude											
All persons	42.4	33.5	30.1	25.5	23.2	20.9	20.9	19.3	18.1	17.8	16.8
Male	51.9	37.5	32.6	28.4	25.6	23.4	23.9	21.5	20.5	20.5	18.8
Female	33.9	29.9	27.9	22.8	20.9	18.5	18.1	17.3	15.8	15.3	14.8
White male ⁴	51.1	36.8	31.7	28.0	25.7	23.2	23.6	21.4	20.3	20.3	18.5
Black or African American male ⁴	60.4	44.1	39.9	32.5	26.2	23.9	26.5	24.3	22.0	21.9	21.8
White female ⁴	34.0	30.1	27.7	23.4	21.4	19.1	18.7	17.9	16.6	15.9	15.5
Black or African American female ⁴	33.7	31.1	31.0	21.2	20.8	17.3	17.3	17.0	14.7	15.1	13.5
All males											
18–44 years	57.9	40.4	35.2	31.4	29.2	26.1	27.1	23.9	24.0	22.9	21.7
18–24 years	54.1	35.0	28.0	26.6	28.1	25.6	28.0	22.8	20.1	21.9	18.5
25–34 years	60.7	43.9	38.2	31.6	28.9	26.1	27.7	26.1	28.0	24.4	23.7
35–44 years	58.2	41.8	37.6	34.5	30.2	26.5	26.0	22.5	22.8	22.1	22.0
45–64 years	51.9	39.3	33.4	29.3	26.4	25.0	25.2	23.2	20.2	21.9	19.4
45–54 years	55.9	42.0	34.9	32.1	28.8	26.7	28.1	25.2	21.4	21.4	19.9
55–64 years	46.6	36.4	31.9	25.9	22.6	22.7	21.1	20.7	18.8	22.6	18.8
65 years and over	28.5	20.9	19.6	14.6	10.2	9.8	8.9	9.7	10.6	10.6	9.8
White male ⁴											
18–44 years	57.1	40.0	34.6	31.3	30.2	26.5	27.7	24.6	24.8	23.4	21.7
18–24 years	53.0	34.3	28.4	27.4	30.4	26.7	29.7	23.8	21.9	23.5	20.0
25–34 years	60.1	43.6	37.3	31.6	29.7	26.3	27.7	26.6	28.4	24.6	23.4
35–44 years	57.3	41.3	36.6	33.5	30.6	26.6	26.3	23.1	23.3	21.9	21.2
45–64 years	51.3	38.3	32.1	28.7	25.8	24.4	24.5	22.5	19.4	21.7	19.0
45–54 years	55.3	40.9	33.7	31.3	28.0	25.9	27.4	24.5	20.7	21.2	19.7
55–64 years	46.1	35.3	30.5	25.6	22.5	22.4	20.4	20.1	17.9	22.2	18.2
65 years and over	27.7	20.5	18.9	13.7	9.8	9.4	7.9	9.6	10.3	10.0	9.4
Black or African American male ⁴											
18–44 years	66.3	45.2	39.6	32.9	25.5	22.8	25.1	22.6	21.3	20.9	22.2
18–24 years	62.8	40.2	27.2	21.3	20.9	18.0	21.6	18.8	13.2	*13.2	*13.9
25–34 years	68.4	47.5	45.6	33.8	23.2	21.2	29.8	25.7	24.9	24.8	28.0
35–44 years	67.3	48.6	45.0	42.0	30.7	28.4	23.3	22.6	24.7	24.0	24.0
45–64 years	57.9	50.0	46.1	36.7	32.2	29.2	32.4	31.8	24.6	25.7	24.0
45–54 years	62.4	51.5	47.7	42.0	35.6	32.1	33.9	33.2	23.3	25.7	22.5
55–64 years	51.8	47.9	44.4	30.2	26.3	24.1	29.8	29.6	26.4	25.6	25.9
65 years and over	36.4	26.2	27.7	21.5	14.2	14.1	16.8	10.0	17.4	15.5	13.9
All females											
18–44 years	42.1	34.7	31.4	25.6	24.5	21.4	21.2	19.1	16.9	16.6	16.6
18–24 years	38.1	33.8	30.4	22.5	24.9	21.5	20.7	17.4	14.5	15.4	14.8
25–34 years	43.7	33.7	32.0	28.2	22.3	21.0	21.5	20.6	19.4	17.9	17.5
35–44 years	43.7	37.0	31.5	24.8	26.2	21.6	21.3	19.0	16.1	16.3	17.0
45–64 years	32.0	30.7	29.9	24.8	21.7	19.8	18.8	19.1	18.9	18.1	16.8
45–54 years	37.5	32.6	32.4	28.5	22.2	20.7	20.9	21.3	21.3	20.6	18.7
55–64 years	25.0	28.6	27.4	20.5	20.9	18.6	16.1	16.5	16.2	15.2	14.8
65 years and over	9.6	13.2	13.5	11.5	9.3	8.1	8.3	9.3	7.5	7.5	7.5

See footnotes at end of table.

Table 47 (page 2 of 2). Current cigarette smoking among adults aged 18 and over, by sex, race, and age: United States, selected years 1965–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#047>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Sex, race, and age	1965 ¹	1979 ¹	1985 ¹	1990 ¹	2000	2004	2005	2010	2012	2013	2014
Percent of adults who were current cigarette smokers ³											
White female ⁴											
18–44 years	42.2	35.1	31.6	26.5	26.5	22.7	22.6	20.5	18.6	17.8	17.8
18–24 years	38.4	34.5	31.8	25.4	28.5	22.9	22.6	18.4	16.9	17.0	16.5
25–34 years	43.4	34.1	32.0	28.5	24.9	22.6	23.1	22.0	20.7	19.2	18.6
35–44 years	43.9	37.2	31.0	25.0	26.6	22.7	22.2	20.5	17.6	17.0	18.0
45–64 years	32.7	30.6	29.7	25.4	21.4	20.1	18.9	19.5	19.4	18.4	17.6
45–54 years	38.2	32.5	32.4	29.1	21.9	21.4	21.0	22.4	22.7	21.2	19.9
55–64 years	25.7	28.5	27.2	21.2	20.6	18.4	16.2	15.9	15.8	15.5	15.3
65 years and over	9.8	13.8	13.3	11.5	9.1	8.2	8.4	9.4	7.5	7.9	7.6
Black or African American female ⁴											
18–44 years	42.9	34.7	33.5	22.8	20.8	17.8	16.9	17.1	12.3	15.1	13.9
18–24 years	37.1	31.8	23.7	10.0	14.2	15.6	14.2	14.2	*7.4	11.8	*9.3
25–34 years	47.8	35.2	36.2	29.1	15.5	18.3	16.9	19.3	17.3	16.4	15.1
35–44 years	42.8	37.7	40.2	25.5	30.2	18.9	19.0	17.2	11.2	16.4	16.4
45–64 years	25.7	34.2	33.4	22.6	25.6	20.9	21.0	19.8	20.4	18.8	15.0
45–54 years	32.3	36.2	36.4	26.5	26.5	20.3	22.2	20.4	20.1	22.2	15.7
55–64 years	16.5	31.9	29.8	17.6	24.2	22.0	19.1	18.9	20.8	14.8	14.2
65 years and over	7.1	*8.5	14.5	11.1	10.2	6.7	10.0	9.4	9.1	6.5	8.1

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error of 20%–30%.

¹Data prior to 1997 are not strictly comparable with data for later years due to the 1997 questionnaire redesign. See Appendix I, National Health Interview Survey (NHIS).

²Estimates are age-adjusted to the year 2000 standard population using five age groups: 18–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over. Age-adjusted estimates in this table may differ from other age-adjusted estimates based on the same data and presented elsewhere if different age groups are used in the adjustment procedure. See Appendix II, Age adjustment.

³Starting with 1993 data (shown in spreadsheet version), current cigarette smokers were defined as ever smoking 100 cigarettes in their lifetime and smoking now every day or some days. For previous definition, see Appendix II, Cigarette smoking.

⁴The race groups, white and black, include persons of Hispanic and non-Hispanic origin. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The single-race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group. Prior to 1999, data were tabulated according to the 1977 Standards. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

NOTES: Standard errors for selected years are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey. Data are from the core questionnaire (1965) and the following questionnaire supplements: hypertension (1974), smoking (1979), alcohol and health practices (1983), health promotion and disease prevention (1985, 1990–1991), cancer control and cancer epidemiology (1992), and year 2000 objectives (1993–1995). Starting with 1997, data are from the family core and sample adult questionnaires. See Appendix I, National Health Interview Survey (NHIS).

Table 48. Age-adjusted prevalence of current cigarette smoking among adults aged 25 and over, by sex, race, and education level: United States, selected years 1974–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#048>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Sex, race, and education level	1974 ¹	1979 ¹	1985 ¹	1990 ¹	1995 ¹	2000	2005	2010	2013	2014
25 years and over, age-adjusted ²										
Percent of adults who were current cigarette smokers ³										
All persons ⁴	36.9	33.1	30.0	25.4	24.5	22.6	20.3	19.2	17.8	17.1
No high school diploma or GED	43.7	40.7	40.8	36.7	35.6	31.6	28.2	26.9	25.8	24.4
High school diploma or GED	36.2	33.6	32.0	29.1	29.1	29.2	27.0	27.0	25.6	25.9
Some college, no bachelor's degree	35.9	33.2	29.5	23.4	22.6	21.7	21.8	21.3	19.5	18.6
Bachelor's degree or higher	27.2	22.6	18.5	13.9	13.6	10.9	9.1	8.3	7.7	7.0
All males ⁴	42.9	37.3	32.8	28.2	26.4	24.7	22.7	21.0	20.3	19.1
No high school diploma or GED	52.3	47.6	45.7	42.0	39.7	36.0	31.7	29.7	31.6	27.7
High school diploma or GED	42.4	38.9	35.5	33.1	32.7	32.1	29.9	29.3	28.8	28.2
Some college, no bachelor's degree	41.8	36.5	32.9	25.9	23.7	23.3	24.9	23.2	20.4	20.2
Bachelor's degree or higher	28.3	22.7	19.6	14.5	13.8	11.6	9.7	8.7	8.7	7.9
White males ^{4,5}	41.9	36.7	31.7	27.6	25.9	24.7	22.4	21.0	20.1	18.6
No high school diploma or GED	51.5	47.6	45.0	41.8	38.7	38.2	31.6	29.4	30.2	26.0
High school diploma or GED	42.0	38.5	34.8	32.9	32.9	32.4	30.0	29.6	28.4	27.9
Some college, no bachelor's degree	41.6	36.4	32.2	25.4	23.3	23.5	24.5	23.4	20.3	20.3
Bachelor's degree or higher	27.8	22.5	19.1	14.4	13.4	11.3	9.3	8.8	8.8	7.4
Black or African American males ^{4,5}	53.4	44.4	42.1	34.5	31.6	26.4	26.5	23.9	23.1	22.9
No high school diploma or GED	58.1	49.7	50.5	41.6	41.9	38.2	35.9	34.4	41.5	38.7
High school diploma or GED	*50.7	48.6	41.8	37.4	36.6	29.0	30.1	28.8	31.9	28.7
Some college, no bachelor's degree	*45.3	39.2	41.8	28.1	26.4	19.9	27.4	24.2	18.0	18.8
Bachelor's degree or higher	*41.4	*36.8	*32.0	*20.8	*17.3	14.6	10.0	8.1	7.7	9.8
All females ⁴	32.0	29.5	27.5	22.9	22.9	20.5	18.0	17.5	15.5	15.2
No high school diploma or GED	36.6	34.8	36.5	31.8	31.7	27.1	24.6	23.7	19.8	21.2
High school diploma or GED	32.2	29.8	29.5	26.1	26.4	26.6	24.1	24.9	22.0	23.5
Some college, no bachelor's degree	30.1	30.0	26.3	21.0	21.6	20.4	19.1	19.6	18.8	17.3
Bachelor's degree or higher	25.9	22.5	17.1	13.3	13.3	10.1	8.5	7.9	6.8	6.2
White females ^{4,5}	31.7	29.7	27.3	23.3	23.1	21.0	18.6	18.3	16.2	16.0
No high school diploma or GED	36.8	35.8	36.7	33.4	32.4	28.4	24.6	24.0	19.0	21.0
High school diploma or GED	31.9	29.9	29.4	26.5	26.8	27.8	25.9	25.8	23.5	25.4
Some college, no bachelor's degree	30.4	30.7	26.7	21.2	22.2	21.1	19.5	21.0	19.8	18.3
Bachelor's degree or higher	25.5	21.9	16.5	13.4	13.5	10.2	9.1	8.7	7.3	6.6
Black or African American females ^{4,5}	35.6	30.3	32.0	22.4	25.7	21.6	17.5	17.0	15.3	14.0
No high school diploma or GED	36.1	31.6	39.4	26.3	32.3	31.1	27.8	25.8	26.5	22.5
High school diploma or GED	40.9	32.6	32.1	24.1	27.8	25.4	18.2	22.9	17.0	15.9
Some college, no bachelor's degree	32.3	*28.9	23.9	22.7	20.8	20.4	17.5	15.0	15.3	14.4
Bachelor's degree or higher	*36.3	*43.3	26.6	17.0	17.3	10.8	*6.6	*6.6	7.3	6.9

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%.

¹Data prior to 1997 are not strictly comparable with data for later years due to the 1997 questionnaire redesign. See Appendix I, National Health Interview Survey (NHIS).

²Estimates are age-adjusted to the year 2000 standard population using four age groups: 25–34 years, 35–44 years, 45–64 years, and 65 years and over. See Appendix II, Age adjustment. For age groups where smoking was 0% or 100%, the age-adjustment procedure was modified to substitute the percentage smoking from the next lower education group.

³Starting with 1993 data (shown in spreadsheet version), current cigarette smokers were defined as ever smoking 100 cigarettes in their lifetime and smoking now every day or some days. For previous definition, see Appendix II, Cigarette smoking.

⁴Includes unknown education level. Education categories shown are for 1997 and subsequent years. GED is General Educational Development high school equivalency diploma. In 1974–1995 the following categories based on number of years of school completed were used: less than 12 years, 12 years, 13–15 years, 16 years or more. See Appendix II, Education.

⁵The race groups, white and black, include persons of Hispanic and non-Hispanic origin. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The single-race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group. Prior to 1999, data were tabulated according to the 1977 Standards. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

NOTES: Standard errors for selected years are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey. Data are from the following questionnaire supplements: hypertension (1974), smoking (1979), alcohol and health practices (1983), health promotion and disease prevention (1985, 1990–1991), cancer control and cancer epidemiology (1992), and year 2000 objectives (1993–1995). Starting with 1997, data are from the family core and sample adult questionnaires. See Appendix I, National Health Interview Survey (NHIS).

Table 49 (page 1 of 3). Current cigarette smoking among adults aged 18 and over, by sex, race, Hispanic origin, age, and education level: United States, average annual, selected years 1990–1992 through 2012–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#049>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Male			Female		
	1990–1992 ¹	1999–2001	2012–2014	1990–1992 ¹	1999–2001	2012–2014
18 years and over, age-adjusted ²						
Percent of adults who were current cigarette smokers ³						
All persons ⁴	27.9	25.0	20.0	23.7	21.1	15.5
Race ⁵						
White only	27.4	25.1	20.0	24.3	22.2	16.4
Black or African American only	33.9	27.2	21.8	23.1	19.7	14.2
American Indian or Alaska Native only	34.2	30.3	23.1	36.7	34.7	18.0
Asian only	24.8	20.3	14.6	6.3	6.7	5.2
Native Hawaiian or Other Pacific						
Islander only	---	*	*	---	*	*
2 or more races	---	34.4	29.2	---	30.7	23.4
American Indian or Alaska Native; White	---	38.7	36.6	---	38.9	30.5
Hispanic origin and race ⁵						
Hispanic or Latino	25.7	22.2	15.7	15.8	12.1	7.3
Mexican	26.2	21.9	15.4	14.8	10.6	6.5
Not Hispanic or Latino	28.1	25.5	21.0	24.4	22.3	17.1
White only	27.7	25.5	21.3	25.2	23.5	18.7
Black or African American only	33.9	27.2	21.9	23.2	19.7	14.4
18 years and over, crude						
All persons ⁴	28.4	25.5	19.9	23.6	21.0	15.3
Race ⁵						
White only	27.8	25.4	19.7	24.1	21.7	16.0
Black or African American only	33.2	27.5	21.9	23.3	19.8	14.4
American Indian or Alaska Native only	35.5	31.8	23.1	37.3	36.9	18.7
Asian only	24.9	21.4	15.3	6.3	6.9	5.4
Native Hawaiian or Other Pacific						
Islander only	---	*	*	---	*	*
2 or more races	---	35.9	29.7	---	31.5	22.5
American Indian or Alaska Native; White	---	41.1	32.9	---	40.1	30.9
Hispanic origin and race ⁵						
Hispanic or Latino	26.5	23.2	16.4	16.6	12.6	7.4
Mexican	27.1	22.8	15.8	15.0	11.0	6.5
Not Hispanic or Latino	28.5	25.8	20.6	24.2	21.9	16.7
White only	28.0	25.5	20.5	24.8	22.7	17.8
Black or African American only	33.3	27.5	22.0	23.3	19.8	14.6
Age and Hispanic origin and race ⁵						
18–24 years:						
Hispanic or Latino	19.3	22.6	14.4	12.8	12.9	5.9
Not Hispanic or Latino:						
White only	28.9	32.7	24.4	28.7	30.8	20.5
Black or African American only	17.7	21.9	13.4	10.8	13.0	9.7
25–34 years:						
Hispanic or Latino	29.9	23.2	20.4	19.2	12.5	8.0
Not Hispanic or Latino:						
White only	32.7	30.8	27.2	30.9	27.4	23.0
Black or African American only	34.6	23.3	26.6	29.2	16.9	16.6
35–44 years:						
Hispanic or Latino	32.1	25.3	15.7	19.9	14.1	7.3
Not Hispanic or Latino:						
White only	32.3	29.6	24.1	27.3	28.3	20.6
Black or African American only	44.1	32.0	24.6	31.3	27.5	15.2
45–64 years:						
Hispanic or Latino	26.6	24.7	16.2	17.1	13.5	9.4
Not Hispanic or Latino:						
White only	28.4	25.1	20.7	26.1	22.1	20.0
Black or African American only	38.0	34.0	24.5	26.1	23.6	18.3
65 years and over:						
Hispanic or Latino	16.1	12.6	10.9	6.6	5.9	3.6
Not Hispanic or Latino:						
White only	14.2	10.0	9.8	12.3	9.8	8.1
Black or African American only	25.2	17.6	15.5	10.7	11.0	7.8

See footnotes at end of table.

Table 49 (page 2 of 3). Current cigarette smoking among adults aged 18 and over, by sex, race, Hispanic origin, age, and education level: United States, average annual, selected years 1990–1992 through 2012–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#049>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Male			Female		
	1990–1992 ¹	1999–2001	2012–2014	1990–1992 ¹	1999–2001	2012–2014
Percent of poverty level ^{2,6}		Percent of adults who were current cigarette smokers ³				
Below 100%	40.5	36.5	33.1	30.7	29.1	24.4
100%–199%	35.0	32.8	27.1	26.9	25.6	20.3
200%–399%	26.5	27.3	20.7	22.6	22.3	15.6
400% or more	22.5	18.8	12.9	19.0	15.9	9.3
Hispanic origin and race and percent of poverty level ^{2,5,6}						
Hispanic or Latino:						
Below 100%	29.2	25.3	18.3	16.3	14.4	9.9
100%–199%	29.5	22.0	16.1	16.0	11.8	6.9
200%–399%	23.7	23.6	15.5	15.9	12.0	6.3
400% or more	19.7	18.1	13.5	13.6	9.4	6.4
Not Hispanic or Latino:						
White only:						
Below 100%	44.2	40.7	43.3	37.8	38.3	35.7
100%–199%	36.3	37.5	34.3	31.1	32.0	29.5
200%–399%	26.4	28.5	22.3	23.7	24.8	19.3
400% or more	22.5	19.1	13.3	19.5	17.1	10.5
Black or African American only:						
Below 100%	43.5	40.6	35.6	28.9	27.7	24.1
100%–199%	36.0	33.9	25.7	20.3	21.3	14.7
200%–399%	31.4	24.9	21.0	21.4	17.3	10.1
400% or more	24.3	17.9	11.1	19.2	12.6	6.8
Disability measure ^{2,7}						
Any basic actions difficulty or complex activity limitation	---	33.1	28.8	---	28.1	23.7
Any basic actions difficulty	---	33.2	29.2	---	28.2	23.7
Any complex activity limitation	---	37.6	32.0	---	30.6	28.4
No disability	---	22.8	17.3	---	18.8	12.3
Education, Hispanic origin, and race ^{5,8}						
25 years and over, age-adjusted ⁹						
No high school diploma or GED:						
Hispanic or Latino	30.2	24.3	17.6	15.8	12.1	7.0
Not Hispanic or Latino:						
White only	46.1	43.5	43.4	40.4	39.3	39.5
Black or African American only	45.4	40.0	39.2	31.3	29.4	26.2
High school diploma or GED:						
Hispanic or Latino	29.6	24.1	19.1	18.4	12.5	9.0
Not Hispanic or Latino:						
White only	32.9	31.8	31.2	28.4	29.2	29.3
Black or African American only	38.2	31.4	29.9	25.4	23.0	16.8
Some college or more:						
Hispanic or Latino	20.4	17.1	11.8	14.3	11.1	7.2
Not Hispanic or Latino:						
White only	19.3	17.6	14.3	18.1	16.7	13.6
Black or African American only	25.6	19.2	15.4	22.8	16.9	12.2

See footnotes at end of table.

Table 49 (page 3 of 3). Current cigarette smoking among adults aged 18 and over, by sex, race, Hispanic origin, age, and education level: United States, average annual, selected years 1990–1992 through 2012–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#049>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

-- Data not available.

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.

¹Data prior to 1997 are not strictly comparable with data for later years due to the 1997 questionnaire redesign. See Appendix I, National Health Interview Survey (NHIS).

²Estimates are age-adjusted to the year 2000 standard population using five age groups: 18–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over. See Appendix II, Age adjustment. For age groups where smoking is 0% or 100%, the age-adjustment procedure was modified to substitute the percentage smoking from the previous 3-year period.

³Starting with 1993 data, current cigarette smokers were defined as ever smoking 100 cigarettes in their lifetime and smoking now every day or some days. For previous definition, see Appendix II, Cigarette smoking.

⁴Includes all other races not shown separately, unknown education level, and unknown disability measure.

⁵The race groups, white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The five single-race categories plus multiple-race categories shown in the table conform to the 1997 Standards. Starting with 1999–2001 data, race-specific estimates are for persons who reported only one racial group; the category 2 or more races includes persons who reported more than one racial group. Prior to 1999, data were tabulated according to the 1977 Standards with four racial groups, and the Asian only category included Native Hawaiian or Other Pacific Islander. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

⁶Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Missing family income data were imputed for 1990 and beyond. See Appendix II, Family income; Poverty; Table VI.

⁷Any basic actions difficulty or complex activity limitation is defined as having one or more of the following limitations or difficulties: movement difficulty, emotional difficulty, sensory (seeing or hearing) difficulty, cognitive difficulty, self-care (activities of daily living or instrumental activities of daily living) limitation, social limitation, or work limitation. For more information, see Appendix II, Basic actions difficulty; Complex activity limitation. Starting with 2007 data, the hearing question, a component of the basic actions difficulty measure, was revised. Consequently, data prior to 2007 are not comparable with data for 2007 and beyond. For more information on the impact of the revised hearing question, see Appendix II, Hearing trouble.

⁸Education categories shown are for 1997 and subsequent years. GED is General Educational Development high school equivalency diploma. In years prior to 1997, the following categories based on number of years of school completed were used: less than 12 years, 12 years, 13 years or more. See Appendix II, Education.

⁹Estimates are age-adjusted to the year 2000 standard using four age groups: 25–34 years, 35–44 years, 45–64 years, and 65 years and over. See Appendix II, Age adjustment.

NOTES: Standard errors for selected years are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey. Data are from the following questionnaire supplements: health promotion and disease prevention (1990–1991), cancer control and cancer epidemiology (1992), and year 2000 objectives (1993–1995). Starting with 1997, data are from the family core and sample adult questionnaires. See Appendix I, National Health Interview Survey (NHIS).

Table 50 (page 1 of 2). Use of selected substances in the past month among persons aged 12 and over, by age, sex, race, and Hispanic origin: United States, selected years 2002–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hs/content2015.htm#050>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population aged 12 and over]

Age, sex, race, and Hispanic origin	Any illicit drug ¹			Marijuana			Nonmedical use of any psychotherapeutic drug ²		
	2002	2013	2014	2002	2013	2014	2002	2013	2014
Percent of population									
12 years and over	8.3	9.4	10.2	6.2	7.5	8.4	2.7	2.5	2.5
Age									
12–13 years	4.2	2.6	3.4	1.4	1.0	1.1	1.7	1.3	1.8
14–15 years	11.2	7.8	7.9	7.6	5.8	5.5	4.0	2.2	2.6
16–17 years	19.8	15.8	16.5	15.7	14.2	15.0	6.3	3.1	3.4
18–25 years	20.2	21.5	22.0	17.3	19.1	19.6	5.5	4.8	4.4
26–34 years	10.5	15.3	15.1	7.7	12.6	12.7	3.7	4.4	4.1
35 years and over	4.6	5.6	6.7	3.1	4.0	5.2	1.6	1.6	1.7
Sex									
Male	10.3	11.5	12.8	8.1	9.7	10.9	2.8	2.6	2.6
Female	6.4	7.3	7.7	4.4	5.6	6.0	2.6	2.3	2.3
Age and sex									
12–17 years	11.6	8.8	9.4	8.2	7.1	7.4	4.0	2.2	2.6
Male	12.3	9.6	9.6	9.1	7.9	7.9	3.6	2.0	2.3
Female	10.9	8.0	9.1	7.2	6.2	6.8	4.4	2.4	3.0
Hispanic origin and race ³									
Not Hispanic or Latino:									
White only	8.5	9.5	10.4	6.5	7.7	8.7	2.8	2.5	2.5
Black or African American only	9.7	10.5	12.4	7.4	8.7	10.3	2.0	2.3	2.7
American Indian or Alaska Native only	10.1	12.3	14.9	6.7	10.8	11.8	3.2	2.1	4.8
Native Hawaiian or Other Pacific Islander only	7.9	14.0	15.6	4.4	13.4	12.1	3.8	1.1	6.2
Asian only	3.5	3.1	4.1	1.8	2.2	2.8	0.7	0.8	1.1
2 or more races	11.4	17.4	15.0	9.0	16.0	12.4	3.5	2.5	3.2
Hispanic or Latino	7.2	8.8	8.9	4.3	6.5	6.7	2.9	2.9	2.5

Age, sex, race, and Hispanic origin	Alcohol use			Binge alcohol use ⁴			Heavy alcohol use ⁵		
	2002	2013	2014	2002	2013	2014	2002	2013	2014
Percent of population									
12 years and over	51.0	52.2	52.7	22.9	22.9	23.0	6.7	6.3	6.2
Age									
12–13 years	4.3	2.1	2.1	1.8	0.8	0.8	0.3	0.1	0.1
14–15 years	16.6	9.5	8.5	9.2	4.5	3.9	1.9	0.7	0.5
16–17 years	32.6	22.7	23.3	21.4	13.1	13.1	5.6	2.7	2.4
18–25 years	60.5	59.6	59.6	40.9	37.9	37.7	14.9	11.3	10.8
26–34 years	61.4	66.1	66.0	33.1	37.4	35.5	9.0	10.9	8.9
35 years and over	52.1	53.6	54.4	18.6	19.0	19.6	5.2	5.0	5.3
Sex									
Male	57.4	57.1	57.3	31.2	30.2	30.0	10.8	9.5	9.3
Female	44.9	47.5	48.4	15.1	16.0	16.4	3.0	3.3	3.2
Age and sex									
12–17 years	17.6	11.6	11.5	10.7	6.2	6.1	2.5	1.2	1.0
Male	17.4	11.2	10.8	11.4	6.6	6.4	3.1	1.4	1.2
Female	17.9	11.9	12.3	9.9	5.8	5.7	1.9	1.0	0.9
Hispanic origin and race ³									
Not Hispanic or Latino:									
White only	55.0	57.7	57.7	23.4	24.0	23.5	7.5	7.3	7.1
Black or African American only	39.9	43.6	44.2	21.0	20.1	21.6	4.4	4.5	4.5
American Indian or Alaska Native only	44.7	37.3	42.3	27.9	23.5	27.7	8.7	5.8	9.2
Native Hawaiian or Other Pacific Islander only	*	38.4	37.9	25.2	24.7	18.3	8.3	8.9	4.6
Asian only	37.1	34.5	38.7	12.4	12.4	14.5	2.6	2.0	2.0
2 or more races	49.9	47.4	49.5	19.8	19.6	21.0	7.5	7.5	5.8
Hispanic or Latino	42.8	43.0	44.4	24.8	24.1	24.7	5.9	4.8	5.1

See footnotes at end of table.

Table 50 (page 2 of 2). Use of selected substances in the past month among persons aged 12 and over, by age, sex, race, and Hispanic origin: United States, selected years 2002–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#050>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population aged 12 and over]

Age, sex, race, and Hispanic origin	Any tobacco ⁶			Cigarettes			Cigars		
	2002	2013	2014	2002	2013	2014	2002	2013	2014
Percent of population									
12 years and over	30.4	25.5	25.2	26.0	21.3	20.8	5.4	4.7	4.5
Age									
12–13 years	3.8	1.3	1.1	3.2	0.9	0.7	0.7	0.2	0.3
14–15 years	13.4	6.4	5.1	11.2	4.3	3.4	3.8	1.4	1.5
16–17 years	29.0	15.5	14.4	24.9	11.4	10.2	9.3	5.2	4.4
18–25 years	45.3	37.0	35.0	40.8	30.6	28.4	11.0	10.0	9.7
26–34 years	38.2	38.3	34.8	32.7	33.0	29.4	6.6	7.8	6.8
35 years and over	27.9	22.8	23.7	23.4	19.0	19.7	4.1	3.3	3.3
Sex									
Male	37.0	31.1	31.1	28.7	23.6	23.2	9.4	7.7	7.5
Female	24.3	20.2	19.7	23.4	19.0	18.6	1.7	2.0	1.7
Age and sex									
12–17 years	15.2	7.8	7.0	13.0	5.6	4.9	4.5	2.3	2.1
Male	16.0	9.1	8.2	12.3	5.7	5.1	6.2	3.2	2.7
Female	14.4	6.5	5.8	13.6	5.5	4.6	2.7	1.4	1.5
Hispanic origin and race ³									
Not Hispanic or Latino:									
White only	32.0	27.7	27.6	26.9	22.7	22.3	5.5	4.8	4.6
Black or African American only	28.8	27.1	26.6	25.3	23.0	22.5	6.8	6.9	6.5
American Indian or Alaska Native only	44.3	40.1	37.8	37.1	36.5	32.5	5.2	6.1	4.2
Native Hawaiian or Other Pacific Islander only	*	25.8	30.6	*	21.1	25.4	4.1	2.1	3.2
Asian only	18.6	10.1	10.2	17.7	8.5	9.2	1.1	2.0	1.2
2 or more races	38.1	31.2	29.5	35.0	27.1	24.4	5.5	5.5	6.5
Hispanic or Latino	25.2	18.8	18.8	23.0	16.8	16.7	5.0	3.7	3.7

* Estimates are considered unreliable. Data not shown if the relative standard error is greater than 17.5% of the log transformation of the proportion, the minimum effective sample size is less than 68, the minimum nominal sample size is less than 100, or the prevalence is close to 0% or 100%.

¹Any illicit drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic drug used nonmedically. See Appendix II, Illicit drug use.

²Nonmedical use of prescription-type psychotherapeutic drugs includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs. Special questions on methamphetamine were added in 2005 and 2006. Data for years prior to 2007 have been adjusted for comparability.

³Persons of Hispanic origin may be of any race. Data on race and Hispanic origin were collected using the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity*. Single-race categories shown include persons who reported only one racial group. The category 2 or more races includes persons who reported more than one racial group. See Appendix II, Hispanic origin; Race.

⁴Binge alcohol use is defined as drinking five or more drinks on the same occasion on at least 1 day in the past 30 days. Occasion is defined as at the same time or within a couple of hours of each other. See Appendix II, Alcohol consumption; Binge drinking.

⁵Heavy alcohol use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days. By definition, all heavy alcohol users are also binge alcohol users.

⁶Any tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. See Appendix II, Cigarette smoking.

NOTES: The National Survey on Drug Use & Health (NSDUH), formerly called the National Household Survey on Drug Abuse (NHSDA), began a new baseline in 2002 and cannot be compared with previous years. Starting with 2011 data, 2010-census based control totals were used in the weighting process. Because of methodological differences among the National Survey on Drug Use & Health, the Monitoring the Future (MTF) Study, and the Youth Risk Behavior Survey (YRBS), rates of substance use measured by these surveys are not directly comparable. See Appendix I, Monitoring the Future (MTF) Study; National Survey on Drug Use & Health (NSDUH); Youth Risk Behavior Survey (YRBS). See Appendix II, Substance use. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use & Health. Available from: <http://www.samhsa.gov/data/population-data-nsduh>. See Appendix I, National Survey on Drug Use & Health (NSDUH).

Table 51 (page 1 of 3). Use of selected substances in the past 30 days among 12th graders, 10th graders, and 8th graders, by sex and race: United States, selected years 1980–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#051>.

[Data are based on a survey of 12th graders, 10th graders, and 8th graders in the coterminous United States]

<i>Substance, grade in school, sex, and race</i>	<i>1980</i>	<i>1990</i>	<i>2000</i>	<i>2004</i>	<i>2005</i>	<i>2010</i>	<i>2011</i>	<i>2012</i>	<i>2013</i>	<i>2014</i>
Cigarettes										
Percent using substance in the past 30 days										
All 12th graders	30.5	29.4	31.4	25.0	23.2	19.2	18.7	17.1	16.3	13.6
Male	26.8	29.1	32.8	25.3	24.8	21.9	21.5	19.3	18.4	15.2
Female.	33.4	29.2	29.7	24.1	20.7	15.7	15.1	14.5	13.2	11.6
White	31.0	32.5	36.6	28.2	27.0	22.2	22.2	20.1	18.5	16.5
Black or African American	25.2	12.0	13.6	11.3	10.0	10.7	8.7	8.4	10.8	7.5
All 10th graders	---	---	23.9	16.0	14.9	13.6	11.8	10.8	9.1	7.2
Male	---	---	23.8	16.2	14.5	15.0	13.4	12.0	10.5	7.7
Female.	---	---	23.6	15.7	15.1	12.1	10.0	9.6	7.5	6.6
White	---	---	27.3	18.1	17.0	14.8	13.7	12.2	10.4	8.5
Black or African American	---	---	11.3	9.6	7.7	7.0	7.2	6.2	4.2	4.2
All 8th graders	---	---	14.6	9.2	9.3	7.1	6.1	4.9	4.5	4.0
Male	---	---	14.3	8.3	8.7	7.4	6.2	4.6	4.0	3.5
Female.	---	---	14.7	9.9	9.7	6.8	5.7	4.9	4.7	4.2
White	---	---	16.4	9.4	9.5	7.9	6.5	5.0	4.3	4.6
Black or African American	---	---	8.4	7.5	6.7	4.0	4.2	3.8	3.3	2.0
Marijuana										
All 12th graders	33.7	14.0	21.6	19.9	19.8	21.4	22.6	22.9	22.7	21.2
Male	37.8	16.1	24.7	23.0	23.6	25.2	26.4	26.5	26.4	24.3
Female.	29.1	11.5	18.3	16.6	15.8	16.9	18.4	18.8	18.7	17.9
White	34.2	15.6	22.0	21.5	21.7	21.6	22.9	22.3	21.3	21.4
Black or African American	26.5	5.2	17.5	14.2	15.1	19.7	22.2	22.4	25.7	20.7
All 10th graders	---	---	19.7	15.9	15.2	16.7	17.6	17.0	18.0	16.6
Male	---	---	23.3	17.4	16.7	20.1	20.8	19.8	20.6	17.4
Female.	---	---	16.2	14.2	13.4	13.3	14.5	14.4	15.3	15.7
White	---	---	20.1	15.8	15.7	15.9	16.9	16.6	16.5	15.8
Black or African American	---	---	17.0	17.2	13.5	15.9	20.0	17.6	20.7	17.4
All 8th graders	---	---	9.1	6.4	6.6	8.0	7.2	6.5	7.0	6.5
Male	---	---	10.2	6.3	7.6	9.2	8.5	7.0	6.7	6.9
Female.	---	---	7.8	6.3	5.7	6.8	5.7	6.0	7.2	5.9
White	---	---	8.3	5.5	6.0	7.1	5.9	4.7	4.8	4.7
Black or African American	---	---	8.5	8.1	8.2	8.2	8.0	7.1	9.3	6.5
Cocaine										
All 12th graders	5.2	1.9	2.1	2.3	2.3	1.3	1.1	1.1	1.1	1.0
Male	6.0	2.3	2.7	2.9	2.6	1.9	1.5	1.5	1.4	1.5
Female.	4.3	1.3	1.6	1.7	1.8	0.7	0.7	0.6	0.5	0.5
White	5.4	1.8	2.2	2.5	2.3	1.2	1.2	1.0	0.7	0.8
Black or African American	2.0	0.5	1.0	0.9	0.5	0.9	0.8	0.5	0.6	1.4
All 10th graders	---	---	1.8	1.7	1.5	0.9	0.7	0.8	0.8	0.6
Male	---	---	2.1	1.9	1.9	1.1	0.8	0.8	1.2	0.8
Female.	---	---	1.4	1.4	1.2	0.5	0.5	0.7	0.5	0.4
White	---	---	1.7	1.7	1.5	0.7	0.5	0.5	0.6	0.5
Black or African American	---	---	0.4	0.4	0.8	0.6	0.6	1.2	0.9	0.5
All 8th graders	---	---	1.2	0.9	1.0	0.6	0.8	0.5	0.5	0.5
Male	---	---	1.3	0.8	0.9	0.6	0.7	0.5	0.4	0.5
Female.	---	---	1.1	1.0	1.0	0.6	0.7	0.4	0.5	0.4
White	---	---	1.1	0.8	0.9	0.5	0.5	0.3	0.4	0.2
Black or African American	---	---	0.5	0.8	0.3	0.3	0.5	0.5	0.5	0.4

See footnotes at end of table.

Table 51 (page 2 of 3). Use of selected substances in the past 30 days among 12th graders, 10th graders, and 8th graders, by sex and race: United States, selected years 1980–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#051>.

[Data are based on a survey of 12th graders, 10th graders, and 8th graders in the coterminous United States]

<i>Substance, grade in school, sex, and race</i>	<i>1980</i>	<i>1990</i>	<i>2000</i>	<i>2004</i>	<i>2005</i>	<i>2010</i>	<i>2011</i>	<i>2012</i>	<i>2013</i>	<i>2014</i>
Inhalants										
Percent using substance in the past 30 days										
All 12th graders	1.4	2.7	2.2	1.5	2.0	1.4	1.0	0.9	1.0	0.7
Male	1.8	3.5	2.9	1.7	2.4	2.1	1.1	0.9	1.2	1.0
Female.	1.0	2.0	1.7	1.3	1.6	0.7	0.9	0.8	0.7	0.5
White	1.4	3.0	2.1	1.6	2.1	1.1	0.9	0.6	0.6	0.6
Black or African American	1.0	1.5	2.1	1.0	1.4	1.5	1.3	0.9	1.3	1.4
All 10th graders	---	---	2.6	2.4	2.2	2.0	1.7	1.4	1.3	1.1
Male	---	---	3.0	2.4	1.9	1.6	1.5	1.2	1.4	0.9
Female.	---	---	2.2	2.3	2.5	2.4	2.0	1.6	1.3	1.2
White	---	---	2.8	2.6	2.2	1.7	1.4	1.1	1.0	0.9
Black or African American	---	---	1.5	1.4	1.4	1.8	1.6	1.2	1.9	1.4
All 8th graders	---	---	4.5	4.5	4.2	3.6	3.2	2.7	2.3	2.2
Male	---	---	4.1	4.0	3.1	2.8	2.5	1.9	1.6	1.6
Female.	---	---	4.8	5.1	5.3	4.4	3.9	3.4	2.9	2.6
White	---	---	4.8	4.4	4.0	3.2	2.7	2.1	1.7	1.7
Black or African American	---	---	2.3	3.8	2.9	2.2	2.8	3.0	2.4	2.4
MDMA (Ecstasy)										
All 12th graders	---	---	3.6	1.2	1.0	1.4	2.3	0.9	1.5	1.4
Male	---	---	4.1	1.6	1.0	1.5	2.8	1.2	2.1	2.1
Female.	---	---	3.1	0.9	1.0	1.2	1.8	0.6	0.9	0.8
White	---	---	3.9	1.2	1.0	0.9	2.1	0.9	1.5	1.3
Black or African American	---	---	1.9	1.1	0.9	1.1	1.1	0.4	0.7	1.4
All 10th graders	---	---	2.6	0.8	1.0	1.9	1.6	1.0	1.2	0.8
Male	---	---	2.5	1.0	1.0	2.3	1.7	1.1	1.5	0.9
Female.	---	---	2.5	0.6	0.9	1.5	1.3	1.0	1.0	0.7
White	---	---	2.5	0.9	1.0	1.5	1.1	1.0	0.9	0.6
Black or African American	---	---	1.8	0.1	0.3	1.1	1.1	1.1	0.4	0.8
All 8th graders	---	---	1.4	0.8	0.6	1.1	0.6	0.5	0.5	0.4
Male	---	---	1.6	0.7	0.8	1.2	0.7	0.4	0.4	0.4
Female.	---	---	1.2	0.9	0.4	1.1	0.5	0.6	0.5	0.3
White	---	---	1.4	0.6	0.6	1.0	0.4	0.4	0.3	0.3
Black or African American	---	---	0.8	1.2	0.9	0.5	0.2	0.5	0.6	0.5
Alcohol ¹										
All 12th graders	72.0	57.1	50.0	48.0	47.0	41.2	40.0	41.5	39.2	37.4
Male	77.4	61.3	54.0	51.1	50.7	44.2	42.1	43.8	41.8	37.4
Female.	66.8	52.3	46.1	45.1	43.3	37.9	37.5	38.8	36.3	37.1
White	75.8	62.2	55.3	52.5	52.2	44.1	43.4	44.3	42.8	42.1
Black or African American	47.7	32.9	29.3	29.2	28.8	30.8	29.4	29.8	27.0	24.9
All 10th graders	---	---	41.0	35.2	33.2	28.9	27.2	27.6	25.7	23.5
Male	---	---	43.3	36.3	32.8	30.1	28.2	28.0	26.0	23.0
Female.	---	---	38.6	34.0	33.6	27.7	26.0	27.1	25.3	23.9
White	---	---	44.3	37.3	36.7	29.2	28.9	29.2	26.9	25.9
Black or African American	---	---	24.7	25.4	20.8	21.3	20.3	20.1	17.7	15.5
All 8th graders	---	---	22.4	18.6	17.1	13.8	12.7	11.0	10.2	9.0
Male	---	---	22.5	17.9	16.2	13.2	12.1	10.3	9.3	8.2
Female.	---	---	22.0	19.0	17.9	14.3	12.8	11.5	11.2	9.5
White	---	---	23.9	18.6	17.3	12.8	11.8	9.6	9.4	8.7
Black or African American	---	---	15.1	16.0	13.9	12.7	10.5	9.4	9.9	7.9

See footnotes at end of table.

Table 51 (page 3 of 3). Use of selected substances in the past 30 days among 12th graders, 10th graders, and 8th graders, by sex and race: United States, selected years 1980–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#051>.

[Data are based on a survey of 12th graders, 10th graders, and 8th graders in the coterminous United States]

Substance, grade in school, sex, and race	1980	1990	2000	2004	2005	2010	2011	2012	2013	2014
Binge drinking ² Percent in the last 2 weeks										
All 12th graders	41.2	32.2	30.0	29.2	27.1	23.2	21.6	23.7	22.1	19.4
Male	52.1	39.1	36.7	34.3	32.6	28.0	25.5	27.2	26.1	22.3
Female	30.5	24.4	23.5	24.2	21.6	18.4	17.6	19.7	18.1	16.6
White	44.6	36.2	34.4	33.1	31.8	26.5	25.3	26.2	25.0	22.5
Black or African American	17.0	11.6	11.0	11.7	10.9	12.6	10.0	13.0	12.0	10.6
All 10th graders	---	---	24.1	19.9	19.0	16.3	14.7	15.6	13.7	12.6
Male	---	---	27.6	21.6	19.9	17.9	16.5	16.4	14.7	13.1
Female	---	---	20.6	18.1	17.9	14.6	12.7	14.8	12.5	12.2
White	---	---	26.6	22.1	21.5	16.0	16.1	16.5	14.7	14.1
Black or African American	---	---	10.6	9.8	8.4	11.5	7.3	9.3	7.9	7.2
All 8th graders	---	---	11.7	9.4	8.4	7.2	6.4	5.1	5.1	4.1
Male	---	---	11.7	8.8	8.2	6.5	6.1	4.6	4.5	3.5
Female	---	---	11.3	9.9	8.6	7.8	6.5	5.5	5.7	4.6
White	---	---	12.5	9.5	8.4	6.7	5.8	3.9	4.6	3.7
Black or African American	---	---	6.2	6.4	5.8	5.9	4.4	4.2	4.8	4.1

--- Data not available.

¹In 1993, the alcohol question was changed to indicate that a drink meant more than a few sips. Data for 1993, available in the spreadsheet version of this table, are based on a half sample. See Appendix II, Alcohol consumption.

²Five or more alcoholic drinks in a row at least once in the prior 2-week period. See Appendix II, Binge drinking.

NOTES: Estimates for Hispanic students are not shown due to small sample size. For 2-year estimates for Hispanic students, see Johnston LD, O'Malley PM, Miech RA, Bachman JG, Schulenberg JE. (2015). Demographic subgroup trends among adolescents in the use of various licit and illicit drugs 1975–2014 (Monitoring the Future Occasional Paper No. 83). Ann Arbor, MI: Institute for Social Research, University of Michigan, 530 pp. Available at: <http://monitoringthefuture.org/pubs.html#papers>. Because of methodological differences among the National Survey on Drug Use & Health (NSDUH), the Monitoring the Future Study (MTF), and the Youth Risk Behavior Survey (YRBS), rates of substance use measured by these surveys are not directly comparable. See Appendix I, National Survey on Drug Use & Health (NSDUH); Monitoring the Future (MTF) Study; Youth Risk Behavior Survey (YRBS). See Appendix II, Cigarette smoking; Illicit drug use; Substance use. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: Monitoring the Future Study. Institute for Social Research, the University of Michigan. Supported by National Institutes of Health, National Institute on Drug Abuse. See Appendix I, Monitoring the Future (MTF) Study.

Table 52 (page 1 of 3). Health risk behaviors among students in grades 9–12, by sex, grade level, race, and Hispanic origin: United States, selected years 1991–2013

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#052>.

[Data are based on a national sample of high school students, grades 9–12]

Sex, grade level, race, and Hispanic origin	Seriously considered suicide ¹				In a physical fight ¹				Carried a weapon ^{2,3}			
	1991	2001	2011	2013	1991	2001	2011	2013	1991	2001	2011	2013
	Percent of students											
Total	29.0	19.0	15.8	17.0	42.5	33.2	32.8	24.7	26.1	17.4	16.6	17.9
Male												
Total	20.8	14.2	12.5	11.6	50.2	43.1	40.7	30.2	40.6	29.3	25.9	28.1
9th grade	17.6	14.7	12.9	9.9	57.8	50.0	46.0	33.2	44.4	33.7	26.6	26.4
10th grade	19.5	13.8	11.4	11.3	50.2	45.0	44.2	30.9	41.5	28.4	26.4	26.4
11th grade	25.3	14.1	14.3	14.0	51.0	38.0	36.3	31.6	44.0	28.1	25.9	30.5
12th grade	20.7	13.7	11.5	11.0	42.3	36.5	34.1	23.8	33.1	25.6	24.1	29.5
Not Hispanic or Latino:												
White	21.7	14.9	12.8	11.4	49.1	43.1	37.7	27.1	41.2	31.3	27.2	33.4
Black or African American	13.3	9.2	9.0	10.2	58.4	43.9	45.8	37.5	43.4	22.4	21.0	18.2
Hispanic or Latino	18.0	12.2	12.6	11.5	48.5	42.4	44.4	34.2	40.0	26.0	24.5	23.8
Female												
Total	37.2	23.6	19.3	22.4	34.4	23.9	24.4	19.2	10.9	6.2	6.8	7.9
9th grade	40.3	26.2	21.5	24.6	42.9	30.3	28.8	23.3	10.4	7.4	7.6	8.6
10th grade	39.7	24.1	22.3	23.4	35.4	24.9	25.5	21.9	11.1	5.4	6.1	9.2
11th grade	38.4	23.6	16.7	22.3	34.5	20.3	22.7	16.7	12.9	5.9	6.2	5.9
12th grade	30.7	18.9	15.8	18.7	25.4	16.9	19.4	13.9	9.5	5.3	7.1	7.5
Not Hispanic or Latina:												
White	38.6	24.2	18.4	21.1	32.2	21.7	20.4	14.6	7.5	5.1	6.2	8.3
Black or African American	29.4	17.2	17.4	18.6	43.8	29.6	32.3	32.1	23.6	8.6	7.5	7.2
Hispanic or Latina	34.6	26.5	21.0	26.0	34.8	29.3	28.7	22.8	12.9	7.4	7.5	7.7

Sex, grade level, race, and Hispanic origin	Rarely or never wore a seatbelt ⁴				Rode with a driver who had been drinking alcohol ^{2,4}				Drove while drinking alcohol ⁵			
	1991	2001	2011	2013	1991	2001	2011	2013	1991	2001	2011	2013
	Percent of students											
Total	25.9	14.1	7.7	7.6	39.9	30.7	24.1	21.9	16.7	13.3	8.2	10.0
Male												
Total	30.0	18.1	8.9	9.1	40.0	31.8	23.3	21.4	21.5	17.2	9.5	12.0
9th grade	30.0	19.4	10.3	9.8	33.9	29.2	20.7	18.1	8.6	9.9	6.1	9.6
10th grade	25.5	16.6	9.0	8.4	36.6	31.5	23.1	19.9	16.1	12.5	6.0	7.4
11th grade	29.5	17.5	7.0	9.7	45.0	32.8	22.4	23.4	26.4	22.1	10.4	14.0
12th grade	34.7	18.6	8.5	8.3	44.7	34.5	27.4	25.3	34.5	27.2	16.0	15.7
Not Hispanic or Latino:												
White	28.6	17.7	7.3	8.5	40.2	31.2	20.5	19.6	23.3	18.6	8.9	12.4
Black or African American	37.5	20.3	12.6	11.8	37.5	31.2	22.5	18.9	14.0	12.5	7.8	6.9
Hispanic or Latino	37.1	17.7	10.1	8.9	47.2	37.1	30.7	28.9	25.1	15.8	11.5	14.5
Female												
Total	21.6	10.2	6.3	6.1	39.8	29.6	24.9	22.4	11.7	9.5	6.7	7.8
9th grade	25.0	10.8	8.4	7.1	36.0	31.3	22.9	20.8	3.3	3.7	3.3	6.1
10th grade	20.4	10.3	5.9	5.7	38.8	29.9	23.5	23.8	7.3	8.4	5.2	4.6
11th grade	20.8	9.7	4.9	6.3	39.7	25.4	25.2	21.8	14.2	11.1	7.8	8.0
12th grade	20.2	9.4	5.5	5.1	44.8	31.3	28.0	23.2	21.7	17.3	11.2	10.5
Not Hispanic or Latina:												
White	18.7	9.7	5.1	4.7	40.9	29.4	23.8	19.9	13.6	10.9	7.0	8.2
Black or African American	31.9	12.2	8.0	7.1	33.8	24.2	23.2	24.8	6.2	3.3	4.0	5.4
Hispanic or Latina	25.9	11.3	8.4	8.7	46.7	39.3	30.7	29.2	9.5	10.5	7.8	8.4

See footnotes at end of table.

Table 52 (page 2 of 3). Health risk behaviors among students in grades 9–12, by sex, grade level, race, and Hispanic origin: United States, selected years 1991–2013

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#052>.

[Data are based on a national sample of high school students, grades 9–12]

Sex, grade level, race, and Hispanic origin	Ever had sexual intercourse				Did not use a condom at last sex ^{6,7}				Ever physically forced to have sex ⁸			
	1991	2001	2011	2013	1991	2001	2011	2013	1991	2001	2011	2013
Percent of students												
Total	54.1	45.6	47.4	46.8	53.8	42.1	39.8	40.9	---	7.7	8.0	7.3
Male												
Total	57.4	48.5	49.2	47.5	45.5	34.9	33.0	34.2	---	5.1	4.5	4.2
9th grade	45.6	40.5	37.8	32.0	44.1	31.1	33.0	30.5	---	5.9	3.5	3.8
10th grade	50.9	42.2	44.5	41.1	43.1	30.7	30.1	30.7	---	4.1	4.2	2.8
11th grade	64.5	54.0	54.5	54.3	43.2	34.7	33.0	29.4	---	4.3	5.2	4.7
12th grade	68.3	61.0	62.6	65.4	49.3	40.8	35.3	42.0	---	5.8	4.7	5.5
Not Hispanic or Latino:												
White	52.7	45.1	44.0	42.2	44.8	36.2	33.7	38.2	---	3.8	3.2	3.1
Black or African American	88.1	68.8	66.9	68.4	43.0	27.3	24.6	27.0	---	8.5	6.1	5.2
Hispanic or Latino	64.1	53.0	53.0	51.7	53.0	40.9	36.6	33.5	---	6.2	5.4	5.2
Female												
Total	50.8	42.9	45.6	46.0	62.0	48.7	46.4	46.9	---	10.3	11.8	10.5
9th grade	32.2	29.1	27.8	28.1	49.7	33.4	43.7	43.5	---	8.6	8.2	8.3
10th grade	45.3	39.3	43.0	41.7	63.6	47.8	43.3	44.5	---	10.7	12.2	11.8
11th grade	60.2	49.7	51.9	53.9	59.3	47.3	44.5	45.2	---	9.9	12.7	10.5
12th grade	65.1	60.1	63.6	62.8	67.4	58.8	51.1	51.6	---	12.2	14.5	11.2
Not Hispanic or Latina:												
White	47.1	41.3	44.5	45.3	62.0	49.0	46.6	46.8	---	9.8	12.0	9.1
Black or African American	75.9	53.4	53.6	53.4	60.6	39.3	46.2	44.7	---	10.6	11.0	11.5
Hispanic or Latina	43.3	44.0	43.9	46.9	73.1	52.4	47.0	49.3	---	11.6	11.2	12.2

Sex, grade level, race, and Hispanic origin	Watched television 3 or more hours ⁹				Not physically active at least 60 minutes every day ^{7,10}				Got fewer than 8 hours of sleep ^{7,11}			
	1991	2001	2011	2013	1991	2001	2011	2013	1991	2001	2011	2013
Percent of students												
Total	---	38.3	32.4	32.5	---	---	71.3	72.9	---	---	68.6	68.3
Male												
Total	---	41.8	33.3	32.8	---	---	61.7	63.4	---	---	66.4	65.5
9th grade	---	51.4	33.9	34.6	---	---	61.2	59.5	---	---	56.9	55.0
10th grade	---	42.3	35.3	32.4	---	---	57.4	65.4	---	---	64.1	62.9
11th grade	---	36.8	32.3	32.3	---	---	63.8	63.0	---	---	71.3	70.6
12th grade	---	33.5	30.9	31.9	---	---	65.1	66.5	---	---	75.2	75.7
Not Hispanic or Latino:												
White	---	35.7	27.3	25.7	---	---	59.6	62.5	---	---	65.0	64.6
Black or African American	---	69.1	54.4	55.3	---	---	64.8	62.8	---	---	72.1	71.2
Hispanic or Latino	---	49.7	38.4	36.5	---	---	64.4	66.1	---	---	66.3	64.6
Female												
Total	---	35.0	31.6	32.2	---	---	81.5	82.3	---	---	70.9	71.1
9th grade	---	39.6	33.8	35.3	---	---	77.8	79.9	---	---	63.2	65.2
10th grade	---	36.2	31.7	32.2	---	---	81.9	79.5	---	---	69.2	70.1
11th grade	---	32.5	30.4	30.4	---	---	82.0	85.6	---	---	75.5	72.4
12th grade	---	29.2	29.9	30.6	---	---	85.1	84.7	---	---	77.2	77.6
Not Hispanic or Latina:												
White	---	26.5	23.9	24.3	---	---	80.3	81.3	---	---	69.8	70.6
Black or African American	---	68.6	54.9	52.2	---	---	83.1	84.0	---	---	72.0	72.4
Hispanic or Latina	---	46.0	37.2	39.0	---	---	83.1	82.6	---	---	72.3	69.8

See footnotes at end of table.

Table 52 (page 3 of 3). Health risk behaviors among students in grades 9–12, by sex, grade level, race, and Hispanic origin: United States, selected years 1991–2013

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#052>.

[Data are based on a national sample of high school students, grades 9–12]

-- Data not available.

¹During the past 12 months.

²During the past 30 days.

³Such as a gun, knife, or club.

⁴When riding in a car driven by someone else.

⁵Among students who drove a vehicle during the past 30 days.

⁶Among students who had sexual intercourse during the past 3 months.

⁷Percent is 100 minus percent presented in MMWR Youth Risk Behavior Surveillance Summaries. See Surveillance Summaries at <http://www.cdc.gov/healthyyouth/yrbs/cdcreports.htm>.

⁸Data prior to 1999 are not available.

⁹On an average school day. Data prior to 1999 are not available.

¹⁰During the past 7 days. Data prior to 2011 are not available.

¹¹On an average school night. Data prior to 2007 are not available.

NOTES: Only youths attending school participated in the survey. YRBS is conducted biennially. Persons of Hispanic origin may be of any race. See Appendix II, Hispanic origin; Race; Suicidal ideation. Standard errors for selected years are available in the spreadsheet version of this table. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/National Center for HIV, Hepatitis, STD, and TB Prevention, Youth Risk Behavior Survey. See Youth Online website at <http://nccd.cdc.gov/youthonline>. See Appendix I, Youth Risk Behavior Survey (YRBS).

Table 53 (page 1 of 2). Selected health conditions and risk factors, by age: United States, selected years 1988–1994 through 2013–2014

Updated data when available, Excel, PDF, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#053>.

[Data are based on interviews and physical examinations of a sample of the civilian noninstitutionalized population]

Health condition	1988–1994	1999–2000	2001–2002	2003–2004	2005–2006	2007–2008	2009–2010	2011–2012	2013–2014
Diabetes ¹ Percent of adults aged 20 and over									
Total, age-adjusted ²	8.8	9.0	10.6	10.9	10.4	11.4	11.5	11.9	11.9
Total, crude	8.3	8.6	10.3	10.9	10.9	11.9	12.1	12.5	12.7
Hypercholesterolemia ³									
Total, age-adjusted ⁴	22.8	25.5	24.6	27.9	27.4	27.6	27.2	28.2	27.4
Total, crude	21.5	24.5	24.2	27.9	28.1	28.8	28.6	30.4	29.3
High cholesterol ⁵									
Total, age-adjusted ⁴	20.8	18.3	16.5	16.9	15.6	14.2	13.2	12.7	11.1
Total, crude	19.6	17.7	16.4	17.0	15.9	14.6	13.6	13.1	11.1
Hypertension ⁶									
Total, age-adjusted ⁴	25.5	30.0	29.7	32.1	30.5	31.2	30.0	30.0	30.8
Total, crude	24.1	28.9	28.9	32.5	31.7	32.6	31.9	32.5	33.5
Uncontrolled high blood pressure among persons with hypertension ⁷									
Total, age-adjusted ⁴	77.2	71.9	68.3	63.8	63.0	56.2	55.7	54.6	51.3
Total, crude	73.9	69.1	65.4	60.8	56.6	51.8	46.7	48.0	46.1
Overweight (includes obesity) ⁸									
Total, age-adjusted ⁴	56.0	64.5	65.6	66.4	66.9	68.1	68.8	68.6	70.4
Total, crude	54.9	64.1	65.6	66.5	67.3	68.3	69.2	69.0	70.7
Obesity ⁹									
Total, age-adjusted ⁴	22.9	30.5	30.5	32.3	34.4	33.7	35.7	34.9	37.8
Total, crude	22.3	30.3	30.6	32.3	34.7	33.9	35.9	35.1	37.9
Untreated dental caries ¹⁰									
Total, age-adjusted ⁴	27.7	24.4	21.3	29.8	24.4	21.7	---	25.5	---
Total, crude	28.2	25.0	21.7	30.2	24.5	21.8	---	25.5	---
Obesity ¹¹ Percent of persons under age 20									
2–5 years	7.2	10.3	10.6	14.0	11.0	10.1	12.1	8.4	9.4
6–11 years	11.3	15.1	16.3	18.8	15.1	19.6	18.0	17.7	17.4
12–19 years	10.5	14.8	16.7	17.4	17.8	18.1	18.4	20.5	20.6
Untreated dental caries ¹⁰									
5–19 years	24.3	23.6	21.2	25.6	16.2	16.9	14.6	17.5	---

See footnotes at end of table.

Table 53 (page 2 of 2). Selected health conditions and risk factors, by age: United States, selected years 1988–1994 through 2013–2014

Updated data when available, Excel, PDF, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#053>.

[Data are based on interviews and physical examinations of a sample of the civilian noninstitutionalized population]

-- Data not available.

¹Includes physician-diagnosed and undiagnosed diabetes. Estimates were obtained using fasting weights. Physician-diagnosed diabetes was obtained by self-report and excludes women who reported having diabetes only during pregnancy. Undiagnosed diabetes is defined as a fasting plasma glucose (FPG) of at least 126 mg/dL or a hemoglobin A1c of at least 6.5% and no reported physician diagnosis. Pregnant women were excluded. Adjustments to FPG recommended by NHANES for trend analysis were incorporated into the data presented here. For more information, see http://www.cdc.gov/nchs/nhanes/nhanes2005-2006/GLU_D.htm. See Appendix II, Diabetes. See related Table 40.

²Estimates are age-adjusted to the year 2000 standard population using three age groups: 20–44 years, 45–64 years, and 65 years and over. Age-adjusted estimates in this table may differ from other age-adjusted estimates based on the same data presented elsewhere if different age groups are used in the adjustment procedure. See Appendix II, Age adjustment.

³Hypercholesterolemia is defined as measured serum total cholesterol greater than or equal to 240 mg/dL or reporting taking cholesterol-lowering medication. Respondents were asked, "Are you now following this advice [from a doctor or health professional] to take prescribed medicine [to lower your cholesterol]?" See Appendix II, Cholesterol. See related Table 55.

⁴Estimates are age-adjusted to the year 2000 standard population using five age groups: 20–34 years, 35–44 years, 45–54 years, 55–64 years, and 65 years and over. Age-adjusted estimates in this table may differ from other age-adjusted estimates based on the same data and presented elsewhere if different age groups are used in the adjustment procedure. See Appendix II, Age adjustment.

⁵High cholesterol is defined as greater than or equal to 240 mg/dL (6.20 mmol/L). This second measure of cholesterol presented in *Health, United States* is based solely on measured high serum total cholesterol. See Appendix II, Cholesterol. See related Table 55.

⁶Hypertension is defined as having measured high blood pressure and/or taking antihypertensive medication. High blood pressure is defined as having measured systolic pressure of at least 140 mm Hg or diastolic pressure of at least 90 mm Hg. Those with high blood pressure also may be taking prescribed medicine for high blood pressure. For antihypertensive medication use, respondents were asked, "Are you now taking prescribed medicine for your high blood pressure?" Pregnant women are excluded. See Appendix II, Blood pressure, high. See related Table 54.

⁷Uncontrolled high blood pressure among persons with hypertension is defined as measured systolic pressure of at least 140 mm Hg or diastolic pressure of at least 90 mm Hg, among those with measured high blood pressure or reporting taking antihypertensive medication. Pregnant women are excluded. See Appendix II, Blood pressure, high. See related Table 54.

⁸Overweight is defined as body mass index (BMI) greater than or equal to 25, based on the NHANES variable, Body Mass Index. Excludes pregnant women. See Appendix II, Body mass index (BMI). See related Table 58.

⁹Obesity is defined as body mass index (BMI) greater than or equal to 30, based on the NHANES variable, Body Mass Index. Excludes pregnant women. See Appendix II, Body mass index (BMI). See related Table 58.

¹⁰Untreated dental caries refers to decay on the crown or enamel surface of a tooth (i.e., coronal caries) that has not been treated or filled. The presence of caries was evaluated in primary and permanent teeth for persons aged 5 and older. The third molars were not included. Persons without at least one natural tooth (primary or permanent) were excluded. Over time, there have been changes in the NHANES oral health examination process, ages examined, and methodology. For more information, see Appendix II, Dental caries. See related Table 60.

¹¹Obesity is defined as body mass index (BMI) at or above the sex- and age-specific 95th percentile BMI (based on the variable BMXBMI) using cutoff points from the 2000 CDC growth charts for the United States: Methods and development. NCHS. Vital Health Stat 11(246). 2002. Available at: http://www.cdc.gov/nchs/data/series/sr_11/sr11_246.pdf. Excludes pregnant girls. See related Table 59.

NOTES: Standard errors are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health and Nutrition Examination Survey. See Appendix I, National Health and Nutrition Examination Survey (NHANES).

Table 54 (page 1 of 2). Hypertension among adults aged 20 and over, by selected characteristics: United States, selected years 1988–1994 through 2011–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#054>.

[Data are based on interviews and physical examinations of a sample of the civilian noninstitutionalized population]

Sex, age, race and Hispanic origin ¹ , and percent of poverty level	Hypertension ^{2,3} (measured high blood pressure and/or taking antihypertensive medication)				Uncontrolled high blood pressure among persons with hypertension ⁴			
	1988–1994	1999–2002	2003–2006	2011–2014	1988–1994	1999–2002	2003–2006	2011–2014
20 years and over, age-adjusted ⁵	Percent of population							
Both sexes ⁶	25.5	30.0	31.3	30.4	77.2	70.6	63.3	52.8
Male	26.4	28.8	31.8	31.0	83.2	73.3	65.0	58.1
Female	24.4	30.6	30.3	29.7	68.5	61.8	53.6	45.5
Not Hispanic or Latino:								
White only	24.4	28.3	29.9	29.1	76.6	69.1	62.3	48.4
White only, male	25.6	27.6	31.2	30.2	82.6	70.3	63.3	53.5
White only, female	23.0	28.5	28.3	28.0	67.0	63.6	47.5	40.2
Black or African American only	38.1	42.3	43.4	43.3	77.5	71.5	64.8	58.7
Black or African American only, male	37.5	40.6	42.2	42.4	84.0	74.3	70.2	66.2
Black or African American only, female	38.3	43.5	44.1	44.0	71.1	67.2	59.0	52.8
Asian only	---	---	---	26.5	---	---	---	64.7
Asian only, male	---	---	---	28.0	---	---	---	68.6
Asian only, female	---	---	---	25.0	---	---	---	58.7
Hispanic or Latino	---	---	---	28.2	---	---	---	57.5
Hispanic or Latino, male	---	---	---	27.7	---	---	---	69.6
Hispanic or Latina, female	---	---	---	28.6	---	---	---	43.4
Mexican origin	26.1	27.6	27.0	28.4	85.7	84.1	72.8	59.3
Mexican origin, male	26.9	26.8	24.8	27.5	87.9	89.5	70.7	71.4
Mexican origin, female	25.0	27.9	28.6	29.4	77.6	71.5	66.1	40.7
Percent of poverty level: ⁷								
Below 100%	31.7	33.9	35.0	34.1	75.0	71.2	69.8	54.8
100%–199%	26.6	33.5	34.1	33.4	76.0	73.4	68.2	56.4
200%–399%	24.7	30.2	31.9	30.0	76.2	67.8	63.9	51.5
400% or more	22.6	26.4	28.9	27.7	81.5	70.3	56.8	53.5
20 years and over, crude								
Both sexes ⁶	24.1	30.2	32.1	33.0	73.9	67.3	58.6	47.0
Male	23.8	27.6	31.3	32.6	79.3	67.1	58.4	49.9
Female	24.4	32.7	32.9	33.4	68.8	67.4	58.8	44.3
Not Hispanic or Latino:								
White only	24.4	30.6	32.9	34.5	72.7	65.5	57.2	44.3
White only, male	24.3	28.3	32.4	34.6	78.0	64.0	56.2	46.0
White only, female	24.6	32.8	33.4	34.5	67.8	66.9	58.2	42.5
Black or African American only	31.8	39.1	41.0	43.0	75.9	69.1	60.1	52.6
Black or African American only, male	31.1	35.9	38.8	40.9	83.3	71.3	65.9	58.3
Black or African American only, female	32.5	41.9	42.8	44.8	70.0	67.5	55.5	48.1
Asian only	---	---	---	25.0	---	---	---	56.7
Asian only, male	---	---	---	25.7	---	---	---	61.2
Asian only, female	---	---	---	24.3	---	---	---	52.4
Hispanic or Latino	---	---	---	22.0	---	---	---	53.2
Hispanic or Latino, male	---	---	---	20.4	---	---	---	59.6
Hispanic or Latina, female	---	---	---	23.6	---	---	---	47.3
Mexican origin	16.2	17.5	18.1	21.3	83.8	80.9	67.7	55.4
Mexican origin, male	16.4	16.5	16.6	20.2	86.5	86.9	66.9	62.6
Mexican origin, female	15.9	18.8	20.0	22.5	80.6	74.5	68.6	47.8
Percent of poverty level: ⁷								
Below 100%	25.7	30.3	28.8	29.1	74.0	71.3	67.3	52.4
100%–199%	26.7	34.8	36.8	36.5	75.1	70.7	63.2	48.3
200%–399%	22.4	29.9	33.1	33.2	73.4	64.4	58.0	46.7
400% or more	22.0	26.8	29.2	32.2	74.3	63.8	53.4	44.2

See footnotes at end of table.

Table 54 (page 2 of 2). Hypertension among adults aged 20 and over, by selected characteristics: United States, selected years 1988–1994 through 2011–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#054>.

[Data are based on interviews and physical examinations of a sample of the civilian noninstitutionalized population]

Sex, age, race and Hispanic origin ¹ , and percent of poverty level	Hypertension ^{2,3} (measured high blood pressure and/or taking antihypertensive medication)				Uncontrolled high blood pressure among persons with hypertension ⁴			
	1988–1994	1999–2002	2003–2006	2011–2014	1988–1994	1999–2002	2003–2006	2011–2014
Male	Percent of population							
20–44 years	10.9	12.1	14.2	11.5	90.5	79.7	71.1	67.8
20–34 years	7.1	*8.1	9.2	6.9	92.6	89.9	83.1	74.1
35–44 years	17.1	17.1	21.1	18.8	89.0	73.3	63.6	64.2
45–64 years	34.2	36.4	41.2	43.8	73.1	61.4	57.0	49.7
45–54 years	29.2	31.0	36.2	33.0	76.2	66.4	59.3	46.3
55–64 years	40.6	45.0	50.2	54.9	70.3	55.9	53.9	51.7
65–74 years	54.4	59.6	64.1	63.4	74.3	59.1	45.9	38.2
75 years and over	60.4	69.0	65.0	72.3	82.5	74.3	59.7	46.5
Female								
20–44 years	6.5	8.3	6.9	10.2	63.4	58.3	49.1	41.9
20–34 years	2.9	*2.7	*2.2	4.3	82.2	56.9	*47.9	55.5
35–44 years	11.2	15.1	12.6	18.6	56.8	58.6	49.4	37.7
45–64 years	32.8	40.0	43.4	39.5	62.1	60.5	55.5	36.8
45–54 years	23.9	31.8	36.2	28.1	58.5	61.1	57.4	40.3
55–64 years	42.6	53.9	54.4	52.1	64.3	60.0	53.6	34.8
65–74 years	56.2	72.7	70.8	64.3	68.7	73.5	58.5	44.5
75 years and over	73.6	83.1	80.2	79.9	81.9	78.1	70.3	61.5

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error of 20%–30%.

--- Data not available.

¹Persons of Hispanic and Mexican origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The non-Hispanic race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group. Prior to data year 1999, estimates were tabulated according to the 1977 Standards. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. See Appendix II, Hispanic origin; Race.

²Hypertension is defined as having measured high blood pressure and/or taking antihypertensive medication. High blood pressure is defined as having measured systolic pressure of at least 140 mm Hg or diastolic pressure of at least 90 mm Hg. Those with high blood pressure also may be taking prescribed medicine for high blood pressure. Those taking antihypertensive medication may not have measured high blood pressure but are still classified as having hypertension. See Appendix II, Blood pressure, high.

³Respondents were asked, “Are you now taking prescribed medicine for your high blood pressure?”

⁴Uncontrolled high blood pressure among persons with hypertension is defined as measured systolic pressure of at least 140 mm Hg or diastolic pressure of at least 90 mm Hg, among those with measured high blood pressure or reporting taking antihypertensive medication. See Appendix II, Blood pressure, high.

⁵Estimates are age-adjusted to the year 2000 standard population using five age groups: 20–34 years, 35–44 years, 45–54 years, 55–64 years, and 65 years and over. Age-adjusted estimates in this table may differ from other age-adjusted estimates based on the same data and presented elsewhere if different age groups are used in the adjustment procedure. See Appendix II, Age adjustment.

⁶Includes persons of all races and Hispanic origins, not just those shown separately.

⁷Percent of poverty level was calculated by dividing family income by the U.S. Department of Health and Human Services’ poverty guideline specific to family size, as well as the appropriate year, and state. Persons with unknown percent of poverty level are excluded (6% in 2011–2014). See Appendix II, Family income; Poverty.

NOTES: Percentages are based on the average of blood pressure measurements taken. In 2011–2014, 85% of participants had three systolic or diastolic blood pressure readings. Excludes pregnant women. Standard errors are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health and Nutrition Examination Survey. See Appendix I, National Health and Nutrition Examination Survey (NHANES).

Table 55 (page 1 of 4). Cholesterol among adults aged 20 and over, by selected characteristics: United States, selected years 1988–1994 through 2011–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#055>.

[Data are based on interviews and laboratory data of a sample of the civilian noninstitutionalized population]

<i>Sex, age, race and Hispanic origin¹, and percent of poverty level</i>	1988–1994	1999–2002	2003–2006	2007–2010	2011–2014
Percent of population with hypercholesterolemia (serum total cholesterol greater than or equal to 240 mg/dL or taking cholesterol-lowering medications) ³					
20 years and over, age-adjusted ²					
Both sexes ⁴	22.8	25.0	27.7	27.4	27.8
Male	21.1	25.3	27.7	28.0	28.4
Female	24.0	24.3	27.4	26.7	27.3
Not Hispanic or Latino:					
White only	22.9	25.8	28.5	27.8	28.7
White only, male	21.1	26.0	28.7	28.1	29.4
White only, female	24.2	25.1	28.2	27.4	28.0
Black or African American only	21.3	21.3	23.2	25.6	25.2
Black or African American only, male	18.6	20.1	22.8	25.4	24.5
Black or African American only, female	23.1	22.0	23.3	25.6	25.7
Asian only	---	---	---	---	26.0
Asian only, male	---	---	---	---	27.4
Asian only, female	---	---	---	---	24.6
Hispanic or Latino					
Hispanic or Latino, male	---	---	---	27.3	26.3
Hispanic or Latina, female	---	---	---	29.1	26.6
Hispanic or Latina, female	---	---	---	25.2	25.8
Mexican origin	20.0	20.6	24.2	27.4	24.8
Mexican origin, male	19.9	21.6	24.2	28.6	26.6
Mexican origin, female	19.8	19.3	24.1	25.5	22.7
Percent of poverty level: ⁵					
Below 100%	23.0	25.0	27.9	26.5	29.2
100%–199%	22.1	25.9	27.6	27.6	25.4
200%–399%	23.1	26.5	27.5	28.9	29.0
400% or more	21.7	23.1	27.9	26.6	28.1
20 years and over, crude					
Both sexes ⁴	21.5	25.0	28.0	28.7	29.8
Male	19.6	25.1	27.5	28.7	29.5
Female	23.2	24.8	28.5	28.7	30.1
Not Hispanic or Latino:					
White only	22.3	26.9	30.3	30.9	33.1
White only, male	20.0	26.8	29.7	30.4	32.6
White only, female	24.5	27.0	30.8	31.4	33.5
Black or African American only	18.1	19.3	21.7	24.4	24.8
Black or African American only, male	16.0	18.5	21.3	24.1	24.0
Black or African American only, female	19.7	19.9	21.9	24.7	25.4
Asian only	---	---	---	---	24.7
Asian only, male	---	---	---	---	25.9
Asian only, female	---	---	---	---	23.7
Hispanic or Latino					
Hispanic or Latino, male	---	---	---	22.3	21.2
Hispanic or Latina, female	---	---	---	23.7	21.3
Hispanic or Latina, female	---	---	---	20.7	21.1
Mexican origin	15.6	15.5	19.0	22.4	19.1
Mexican origin, male	16.2	17.0	19.3	23.7	21.2
Mexican origin, female	14.9	13.8	18.7	21.0	16.8
Percent of poverty level: ⁵					
Below 100%	19.4	21.6	24.1	22.3	25.3
100%–199%	21.3	25.4	28.3	28.7	27.4
200%–399%	21.3	26.2	28.1	30.6	31.6
400% or more	21.9	24.2	28.7	29.6	32.2
Male					
20–44 years	13.1	16.1	16.5	14.3	12.3
20–34 years	8.2	10.4	10.2	8.5	6.7
35–44 years	21.0	23.1	25.2	22.5	21.1
45–64 years	30.1	36.0	35.7	39.0	39.3
45–54 years	29.6	34.1	32.4	34.0	32.9
55–64 years	30.8	39.1	41.6	46.2	46.0
65–74 years	27.4	36.3	49.4	48.9	55.8
75 years and over	24.4	29.0	37.1	45.2	54.4

See footnotes at end of table.

Table 55 (page 2 of 4). Cholesterol among adults aged 20 and over, by selected characteristics: United States, selected years 1988–1994 through 2011–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#055>.

[Data are based on interviews and laboratory data of a sample of the civilian noninstitutionalized population]

<i>Sex, age, race and Hispanic origin¹, and percent of poverty level</i>	1988–1994	1999–2002	2003–2006	2007–2010	2011–2014
Percent of population with hypercholesterolemia (serum total cholesterol greater than or equal to 240 mg/dL or taking cholesterol-lowering medications) ³					
Female					
20–44 years	9.9	11.4	12.9	10.6	9.0
20–34 years	7.3	9.1	10.8	6.8	6.1
35–44 years	13.5	14.4	15.8	15.7	13.2
45–64 years	36.4	31.7	37.3	39.1	40.6
45–54 years	28.2	27.2	29.6	29.1	31.2
55–64 years	45.8	39.2	49.2	51.4	51.2
65–74 years	46.9	51.9	55.3	53.3	58.1
75 years and over	41.2	44.0	47.3	52.5	59.1
Percent of population with high cholesterol (serum total cholesterol greater than or equal to 240 mg/dL) ⁶					
20 years and over, age-adjusted ²					
Both sexes ⁴	20.8	17.3	16.3	13.7	11.9
Male	19.0	16.4	15.1	12.6	10.8
Female	22.0	17.8	17.1	14.4	12.7
Not Hispanic or Latino:					
White only	20.8	17.5	16.9	13.9	12.4
White only, male	18.8	16.5	15.5	12.2	11.2
White only, female	22.2	18.1	18.0	15.3	13.3
Black or African American only	19.5	15.5	12.2	11.3	8.6
Black or African American only, male	16.9	12.4	10.9	10.8	7.7
Black or African American only, female	21.4	17.7	13.3	11.5	9.4
Asian only	---	---	---	---	10.9
Asian only, male	---	---	---	---	10.6
Asian only, female	---	---	---	---	11.0
Hispanic or Latino	---	---	---	14.7	13.0
Hispanic or Latino, male	---	---	---	15.5	13.1
Hispanic or Latina, female	---	---	---	13.7	12.7
Mexican origin	18.7	15.8	16.1	14.6	11.2
Mexican origin, male	18.5	17.4	17.6	15.1	12.8
Mexican origin, female	18.7	13.8	14.4	13.6	9.3
Percent of poverty level: ⁵					
Below 100%	20.6	18.3	18.1	14.4	12.3
100%–199%	20.6	19.1	16.7	15.0	11.3
200%–399%	20.8	18.9	15.8	14.4	13.0
400% or more	19.5	14.4	15.9	12.3	11.5
20 years and over, crude					
Both sexes ⁴	19.6	17.3	16.4	14.1	12.1
Male	17.7	16.5	15.2	12.9	10.7
Female	21.3	18.0	17.5	15.2	13.5
Not Hispanic or Latino:					
White only	20.3	18.0	17.4	14.7	12.9
White only, male	18.0	16.9	15.7	12.6	10.8
White only, female	22.5	19.1	18.9	16.7	14.9
Black or African American only	16.7	14.4	11.7	11.1	8.6
Black or African American only, male	14.7	12.2	10.8	10.9	7.4
Black or African American only, female	18.2	16.1	12.5	11.3	9.5
Asian only	---	---	---	---	10.9
Asian only, male	---	---	---	---	10.9
Asian only, female	---	---	---	---	10.9
Hispanic or Latino	---	---	---	13.5	12.1
Hispanic or Latino, male	---	---	---	14.7	12.7
Hispanic or Latina, female	---	---	---	12.2	11.6
Mexican origin	14.9	12.9	14.2	13.6	10.6
Mexican origin, male	15.4	15.0	15.7	14.7	12.3
Mexican origin, female	14.3	10.7	12.6	12.3	8.8
Percent of poverty level: ⁵					
Below 100%	17.6	16.4	16.8	12.8	11.3
100%–199%	19.8	18.2	16.0	14.6	11.1
200%–399%	19.3	18.7	15.8	14.6	13.4
400% or more	19.9	15.5	17.1	13.7	12.5

See footnotes at end of table.

Table 55 (page 3 of 4). Cholesterol among adults aged 20 and over, by selected characteristics: United States, selected years 1988–1994 through 2011–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#055>.

[Data are based on interviews and laboratory data of a sample of the civilian noninstitutionalized population]

<i>Sex, age, race and Hispanic origin¹, and percent of poverty level</i>	1988–1994	1999–2002	2003–2006	2007–2010	2011–2014
	Percent of population with high cholesterol (serum total cholesterol greater than or equal to 240 mg/dL) ⁶				
Male					
20–44 years	12.5	14.2	14.1	11.1	10.0
20–34 years	8.2	9.8	9.5	7.6	6.0
35–44 years	19.4	19.7	20.5	16.2	16.2
45–64 years	27.2	22.2	19.1	17.7	13.6
45–54 years	26.6	23.6	20.8	18.7	15.7
55–64 years	28.0	19.9	16.0	16.3	11.5
65–74 years	21.9	13.7	10.9	7.5	7.6
75 years and over	20.4	10.2	9.6	6.8	*3.6
Female					
20–44 years	9.4	10.4	11.3	8.4	7.2
20–34 years	7.3	8.9	10.3	5.8	5.4
35–44 years	12.3	12.4	12.7	11.9	9.8
45–64 years	33.4	23.0	23.9	21.3	19.9
45–54 years	26.7	21.4	19.7	17.7	18.0
55–64 years	40.9	25.6	30.5	25.6	22.1
65–74 years	41.3	32.3	24.2	20.6	15.8
75 years and over	38.2	26.5	18.6	20.2	16.1
20 years and over, age-adjusted ²					
Both sexes ⁴	206	203	200	196	192
Male	204	202	198	194	189
Female	207	204	202	198	195
Not Hispanic or Latino:					
White only	206	204	201	196	193
White only, male	205	202	198	193	189
White only, female	208	205	203	199	196
Black or African American only	205	199	194	192	186
Black or African American only, male	202	195	193	191	183
Black or African American only, female	207	202	195	192	189
Asian only	---	---	---	---	191
Asian only, male	---	---	---	---	189
Asian only, female	---	---	---	---	192
Hispanic or Latino	---	---	---	198	194
Hispanic or Latino, male	---	---	---	199	193
Hispanic or Latina, female	---	---	---	197	195
Mexican origin	206	202	202	198	192
Mexican origin, male	206	204	203	200	194
Mexican origin, female	206	199	200	196	191
Percent of poverty level: ⁵					
Below 100%	205	201	203	196	191
100%–199%	205	204	201	198	191
200%–399%	207	205	199	196	193
400% or more	205	202	201	195	194

See footnotes at end of table.

Table 55 (page 4 of 4). Cholesterol among adults aged 20 and over, by selected characteristics: United States, selected years 1988–1994 through 2011–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#055>.

[Data are based on interviews and laboratory data of a sample of the civilian noninstitutionalized population]

<i>Sex, age, race and Hispanic origin¹, and percent of poverty level</i>	1988–1994	1999–2002	2003–2006	2007–2010	2011–2014
20 years and over, crude		Mean serum total cholesterol level, mg/dL			
Both sexes ⁴	204	203	200	197	192
Male	202	202	198	194	188
Female	206	204	202	199	196
Not Hispanic or Latino:					
White only	206	205	202	198	194
White only, male	203	203	198	193	188
White only, female	208	206	205	201	199
Black or African American only	200	197	193	191	186
Black or African American only, male	198	194	192	191	183
Black or African American only, female	201	199	194	191	189
Asian only	---	---	---	---	191
Asian only, male	---	---	---	---	190
Asian only, female	---	---	---	---	192
Hispanic or Latino					
Hispanic or Latino, male	---	---	---	197	193
Hispanic or Latina, female	---	---	---	199	193
Mexican origin	199	197	198	198	192
Mexican origin, male	199	200	200	200	194
Mexican origin, female	198	194	196	195	189
Percent of poverty level: ⁵					
Below 100%	200	198	200	194	189
100%–199%	202	202	199	197	190
200%–399%	205	204	199	197	193
400% or more	206	204	203	198	196
Male					
20–44 years	194	196	196	194	188
20–34 years	186	188	186	186	179
35–44 years	206	207	209	205	202
45–64 years	216	213	206	202	196
45–54 years	216	215	208	204	200
55–64 years	216	212	202	199	192
65–74 years	212	202	191	182	180
75 years and over	205	195	187	176	168
Female					
20–44 years	189	191	192	187	184
20–34 years	184	185	188	181	179
35–44 years	195	198	197	195	193
45–64 years	225	215	213	211	209
45–54 years	217	211	208	208	207
55–64 years	235	221	219	214	210
65–74 years	233	224	214	207	200
75 years and over	229	217	206	203	199

--- Data not available.

¹Persons of Hispanic and Mexican origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The non-Hispanic race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group. Prior to data year 1999, estimates were tabulated according to the 1977 Standards. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. See Appendix II, Hispanic origin; Race.

²Estimates are age-adjusted to the year 2000 standard population using five age groups: 20–34 years, 35–44 years, 45–54 years, 55–64 years, and 65 years and over. Age-adjusted estimates may differ from other age-adjusted estimates based on the same data and presented elsewhere if different age groups are used in the adjustment procedure. See Appendix II, Age adjustment.

³Hypercholesterolemia is defined as measured serum total cholesterol greater than or equal to 240 mg/dL or reporting taking cholesterol-lowering medications. Respondents were asked, "Are you now following this advice [from a doctor or health professional] to take prescribed medicine [to lower your cholesterol]?"

⁴Includes persons of all races and Hispanic origins, not just those shown separately.

⁵Percent of poverty level was calculated by dividing family income by the U.S. Department of Health and Human Services' poverty guideline specific to family size, as well as the appropriate year, and state. Persons with unknown percent of poverty level are excluded (6% in 2011–2014). See Appendix II, Family income; Poverty.

⁶High cholesterol is defined as serum total cholesterol greater than or equal to 240 mg/dL (6.20 mmol/L), regardless of whether the respondent reported taking cholesterol-lowering medications.

NOTES: See Appendix II, Cholesterol. Standard errors for selected years are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health and Nutrition Examination Survey. See Appendix I, National Health and Nutrition Examination Survey (NHANES).

Table 56 (page 1 of 2). Mean macronutrient intake among adults aged 20 and over, by sex and age: United States, selected years 1988–1994 through 2009–2012

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#056>.

[Data are based on dietary recall interviews of a sample of the civilian noninstitutionalized population]

Sex and age	1988–1994	1999–2002	2003–2006	2009–2012
Percent kcal from carbohydrates				
Both sexes, age-adjusted ¹	49.8	50.7	48.9	49.5
Both sexes, crude	49.8	50.7	48.9	49.4
20–44 years	49.2	51.3	49.3	49.9
45–64 years	49.7	49.3	47.5	48.5
65–74 years	51.1	50.5	49.2	49.0
75 years and over	53.0	52.6	51.5	51.0
Male, age-adjusted ¹	48.5	49.5	47.8	48.1
Male, crude	48.4	49.4	47.7	48.0
20–44 years	48.1	50.2	48.4	48.5
45–64 years	48.3	48.0	46.3	47.2
65–74 years	49.4	49.4	47.6	47.0
75 years and over	50.9	51.0	50.3	50.3
Female, age-adjusted ¹	51.0	51.9	49.9	50.8
Female, crude	51.0	51.9	49.9	50.7
20–44 years	50.3	52.5	50.2	51.3
45–64 years	51.0	50.6	48.7	49.8
65–74 years	52.5	51.4	50.6	50.9
75 years and over	54.2	53.7	52.4	51.6
Percent kcal from protein				
Both sexes, age-adjusted ¹	15.5	15.3	15.6	15.7
Both sexes, crude	15.4	15.3	15.6	15.7
20–44 years	15.0	14.9	15.3	15.6
45–64 years	15.9	15.6	16.0	15.8
65–74 years	16.2	16.3	15.9	16.4
75 years and over	16.0	15.4	15.6	15.8
Male, age-adjusted ¹	15.5	15.4	15.6	16.0
Male, crude	15.4	15.4	15.6	16.0
20–44 years	15.0	15.0	15.4	15.8
45–64 years	15.9	15.7	15.8	16.0
65–74 years	15.9	16.3	16.0	16.6
75 years and over	16.3	15.7	15.8	16.0
Female, age-adjusted ¹	15.5	15.2	15.6	15.5
Female, crude	15.4	15.2	15.6	15.5
20–44 years	14.9	14.8	15.2	15.3
45–64 years	15.9	15.5	16.1	15.5
65–74 years	16.5	16.3	15.9	16.2
75 years and over	15.9	15.3	15.5	15.6
Percent kcal from total fat				
Both sexes, age-adjusted ¹	33.5	33.0	33.7	32.9
Both sexes, crude	33.5	33.0	33.7	33.0
20–44 years	34.0	32.4	33.1	32.3
45–64 years	33.4	33.9	34.6	33.5
65–74 years	32.3	33.4	34.3	33.7
75 years and over	32.0	32.8	33.1	33.3
Male, age-adjusted ¹	33.8	33.0	33.5	33.0
Male, crude	33.9	33.0	33.6	33.0
20–44 years	34.1	32.2	32.6	32.2
45–64 years	33.9	34.0	34.8	33.8
65–74 years	33.0	33.4	34.5	34.1
75 years and over	33.0	33.2	33.3	33.1
Female, age-adjusted ¹	33.2	33.1	33.8	32.8
Female, crude	33.2	33.1	33.9	32.9
20–44 years	33.9	32.6	33.6	32.4
45–64 years	32.9	33.9	34.4	33.2
65–74 years	31.6	33.3	34.1	33.3
75 years and over	31.5	32.6	32.9	33.5

See footnotes at end of table.

Table 56 (page 2 of 2). Mean macronutrient intake among adults aged 20 and over, by sex and age: United States, selected years 1988–1994 through 2009–2012

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#056>.

[Data are based on dietary recall interviews of a sample of the civilian noninstitutionalized population]

Sex and age	1988–1994	1999–2002	2003–2006	2009–2012
		Percent kcal from saturated fat		
Both sexes, age-adjusted ¹	11.2	10.7	11.2	10.6
Both sexes, crude	11.2	10.7	11.2	10.6
20–44 years	11.5	10.8	11.1	10.5
45–64 years	11.1	10.8	11.4	10.8
65–74 years	10.7	10.5	11.2	10.7
75 years and over	10.7	10.3	11.0	10.8
Male, age-adjusted ¹	11.3	10.7	11.1	10.6
Male, crude	11.4	10.7	11.1	10.6
20–44 years	11.5	10.8	11.0	10.4
45–64 years	11.2	10.7	11.3	10.9
65–74 years	10.9	10.6	11.2	10.8
75 years and over	11.2	10.7	11.2	10.7
Female, age-adjusted ¹	11.1	10.7	11.2	10.6
Female, crude	11.1	10.7	11.3	10.6
20–44 years	11.4	10.8	11.2	10.5
45–64 years	10.9	10.9	11.5	10.6
65–74 years	10.4	10.4	11.3	10.7
75 years and over	10.5	10.1	10.8	10.9

¹Estimates are age-adjusted to the year 2000 standard population using four age groups: 20–44 years, 45–64 years, 65–74 years, and 75 years and over. Age-adjusted estimates in this table may differ from other age-adjusted estimates based on the same data and presented elsewhere if different age groups are used in the adjustment procedure. See Appendix II, Age adjustment.

NOTES: Starting in 2001, 24-hour dietary recall data were collected in the mobile examination center (day 1 file) and on a second day by telephone interview (day 2 file). For comparability across survey years, this table is based on day 1 data only. It is recognized that usual intake of macronutrients based on 2 or more days of dietary data would be more precise (Freedman LS, Guenther PM, Dodd KW, Krebs-Smith SM, Midthune D. The population distribution of ratios of usual intakes of dietary components that are consumed every day can be estimated from repeated 24-hour recalls. *J Nutr* 2010 Jan;140(1):111–6.) Two days of data are available only in later years of the continuous NHANES survey. Thus, in order to present trends, macronutrient intake estimates on a given day are presented in this table. This table excludes individuals who reported no energy intake. Energy intake included kilocalories from all foods and beverages, including alcoholic beverages, consumed during the previous 24-hour period. Macronutrients (carbohydrates, protein, and fat) do not sum to 100% because information for alcohol is not shown in the table. See *Health, United States, 2013*, Table 67, for earlier data years. Standard errors are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health and Nutrition Examination Survey. U.S. Department of Agriculture, Agriculture Research Service. Beltsville Human Nutrition Research Center, Food Surveys Research Group, What We Eat in America. See Appendix I, National Health and Nutrition Examination Survey (NHANES).

Table 57 (page 1 of 5). Participation in leisure-time aerobic and muscle-strengthening activities that meet the federal 2008 Physical Activity Guidelines for Americans among adults aged 18 and over, by selected characteristics: United States, selected years 1998–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#057>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	2008 Physical Activity Guidelines for Americans ¹									
	Met both aerobic activity and muscle-strengthening guidelines					Met neither aerobic activity nor muscle-strengthening guideline				
	1998	2000	2010	2013	2014	1998	2000	2010	2013	2014
	Percent									
18 years and over, age-adjusted ^{2,3}	14.3	15.0	20.7	21.0	21.5	56.6	54.7	49.1	46.5	46.8
18 years and over, crude ³	14.5	15.1	20.4	20.4	20.9	56.3	54.6	49.5	47.2	47.5
Age										
18–44 years	18.9	18.9	25.7	25.7	26.7	50.7	49.1	43.1	40.3	40.8
18–24 years	23.8	23.8	29.6	30.3	31.1	46.5	44.5	39.4	35.5	38.0
25–44 years	17.4	17.3	24.3	24.0	25.1	51.9	50.6	44.4	42.0	41.8
45–64 years	11.4	12.8	17.7	17.8	17.8	58.8	57.6	51.0	50.2	50.6
45–54 years	13.2	14.5	19.2	20.1	19.3	56.9	55.4	48.9	48.4	49.2
55–64 years	8.6	10.1	15.9	15.3	16.1	61.8	61.0	53.7	52.1	52.1
65 years and over	5.5	6.8	10.4	11.7	11.7	71.0	67.0	64.6	59.4	58.7
65–74 years	7.0	8.4	13.6	14.7	14.4	65.6	60.3	59.9	54.0	53.1
75 years and over	3.5	4.9	6.4	7.6	7.9	77.8	75.0	70.3	66.8	66.7
Sex ²										
Male	17.5	17.9	25.1	25.0	25.5	50.8	49.6	43.8	42.0	43.4
Female	11.4	12.3	16.5	17.2	17.7	61.9	59.4	54.0	50.7	50.0
Sex and age										
Male:										
18–44 years	23.0	23.0	31.8	31.7	32.3	44.3	43.0	37.1	34.7	36.2
45–54 years	16.1	16.0	20.9	22.3	21.5	52.9	52.7	45.2	46.3	48.0
55–64 years	9.4	11.3	19.1	17.6	17.6	58.2	58.7	50.1	49.6	52.1
65–74 years	9.5	9.4	16.6	15.9	17.5	58.9	55.3	55.6	49.7	50.5
75 years and over	4.9	7.1	9.1	7.8	10.7	69.5	66.7	62.8	60.6	60.3
Female:										
18–44 years	14.9	15.0	19.6	19.8	21.3	56.9	55.0	49.0	45.7	45.2
45–54 years	10.5	13.1	17.5	18.0	17.3	60.8	57.9	52.4	50.5	50.3
55–64 years	7.8	9.0	13.1	13.2	14.8	65.0	63.1	57.0	54.4	52.2
65–74 years	5.1	7.7	11.0	13.6	11.8	70.9	64.3	63.6	57.7	55.4
75 years and over	2.6	3.6	4.6	7.5	5.9	83.0	80.0	75.3	71.0	71.3
Race ^{2,4}										
White only	14.8	15.7	21.4	21.7	22.1	55.2	53.1	47.6	45.2	45.6
Black or African American only	11.7	12.2	17.2	17.7	19.9	65.7	64.6	58.5	54.7	53.4
American Indian or Alaska Native only	16.0	*10.6	*12.7	16.8	24.1	57.6	67.1	54.0	50.8	51.5
Asian only	13.5	14.1	17.8	18.3	17.0	59.1	55.0	51.7	47.6	49.9
Native Hawaiian or Other Pacific Islander only	---	*	*	*	*	---	*	*	*	*
2 or more races	---	19.0	25.9	22.4	21.0	---	52.8	45.0	44.4	46.1
Hispanic origin and race ^{2,4}										
Hispanic or Latino	9.4	9.2	14.4	16.6	15.3	67.7	66.5	60.2	53.8	55.2
Mexican	8.7	8.1	13.2	15.0	14.3	69.5	67.0	60.7	53.4	55.8
Not Hispanic or Latino	14.9	15.8	21.9	21.9	22.7	55.3	53.2	47.2	45.2	45.2
White only	15.5	16.5	22.9	22.9	23.7	53.6	51.4	45.0	43.1	43.3
Black or African American only	11.7	12.2	17.4	17.8	20.1	65.8	64.6	58.4	54.7	53.3
Education ^{5,6}										
No high school diploma or GED	4.6	4.3	7.7	8.0	7.1	76.3	74.0	69.8	66.6	66.7
High school diploma or GED	8.6	9.5	12.7	13.8	13.1	64.6	61.7	59.0	57.0	57.2
Some college or more	18.2	18.9	25.0	24.4	25.4	48.0	47.1	42.1	40.6	40.4

See footnotes at end of table.

Table 57 (page 2 of 5). Participation in leisure-time aerobic and muscle-strengthening activities that meet the federal 2008 Physical Activity Guidelines for Americans among adults aged 18 and over, by selected characteristics: United States, selected years 1998–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#057>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	2008 Physical Activity Guidelines for Americans ¹									
	Met both aerobic activity and muscle-strengthening guidelines					Met neither aerobic activity nor muscle-strengthening guideline				
	1998	2000	2010	2013	2014	1998	2000	2010	2013	2014
Percent of poverty level ^{2,7}						Percent				
Below 100%	8.0	9.3	12.0	12.6	13.1	71.3	68.0	63.9	59.5	60.2
100%–199%	9.0	9.0	12.7	14.3	14.1	67.1	65.5	60.6	56.7	56.6
200%–399%	12.6	13.2	19.2	19.2	19.7	58.0	56.8	50.6	48.0	49.0
400% or more	20.2	20.5	29.1	29.1	29.9	46.2	45.0	36.9	35.6	35.6
Hispanic origin and race and percent of poverty level ^{2,4,7}										
Hispanic or Latino:										
Below 100%	4.6	4.4	8.9	8.7	9.0	78.0	75.2	68.6	65.1	65.0
100%–199%	7.0	5.0	9.3	12.2	12.8	71.2	72.2	66.7	60.1	57.5
200%–399%	11.1	10.2	15.7	20.6	17.2	63.8	63.1	57.6	48.5	52.2
400% or more	17.4	19.6	28.1	27.6	26.1	55.6	52.8	42.5	39.5	41.3
Not Hispanic or Latino:										
White only:										
Below 100%	9.9	11.7	13.7	14.9	16.1	66.9	63.5	60.5	56.2	57.4
100%–199%	9.6	10.3	14.1	15.7	14.7	65.1	62.6	56.4	53.8	54.8
200%–399%	13.1	13.9	20.0	19.0	19.8	56.1	54.7	48.6	46.8	47.8
400% or more	20.2	21.0	29.9	30.0	31.0	45.2	43.7	35.2	33.9	33.6
Black or African American only:										
Below 100%	7.1	9.5	11.3	11.6	12.9	74.6	72.1	66.9	63.5	60.5
100%–199%	8.8	9.5	11.7	13.8	15.0	69.8	69.2	67.0	62.0	59.6
200%–399%	10.6	11.8	20.8	17.1	23.2	64.5	64.3	53.3	52.7	50.8
400% or more	21.2	17.6	26.1	28.5	30.4	54.2	54.9	47.7	39.4	41.7
Disability measure ^{2,8}										
Any basic actions difficulty or complex activity limitation										
Any basic actions difficulty	10.2	10.3	13.6	14.1	14.3	64.4	62.2	59.1	56.9	58.3
Any complex activity limitation	9.8	10.3	13.8	13.8	14.2	64.8	62.1	59.2	57.3	58.6
No disability	7.7	7.2	8.9	10.8	10.7	71.9	71.2	67.2	65.1	66.3
	16.0	17.0	24.2	24.2	24.5	52.5	50.6	43.3	40.2	41.2
Geographic region ²										
Northeast	14.2	17.0	20.2	22.2	22.3	57.0	51.8	49.1	47.1	48.8
Midwest	15.0	16.4	20.7	21.7	20.8	54.9	53.4	49.7	46.3	47.7
South	11.8	12.1	18.8	18.2	20.1	61.4	59.7	51.8	49.5	48.2
West	18.5	16.7	24.0	23.7	24.0	49.5	50.1	44.5	41.5	42.3
Location of residence ^{2,9}										
Within MSA	14.9	15.7	21.8	21.9	22.4	55.8	54.1	47.8	45.3	45.9
Outside MSA	12.2	12.3	14.5	15.7	16.0	59.7	56.9	56.9	53.5	52.4

See footnotes at end of table.

Table 57 (page 3 of 5). Participation in leisure-time aerobic and muscle-strengthening activities that meet the federal 2008 Physical Activity Guidelines for Americans among adults aged 18 and over, by selected characteristics: United States, selected years 1998–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#057>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	2008 Physical Activity Guidelines for Americans ¹									
	Met aerobic activity guideline					Met muscle-strengthening guideline				
	1998	2000	2010	2013	2014	1998	2000	2010	2013	2014
	Percent									
18 years and over, age-adjusted ^{2,3}	40.0	42.2	47.3	50.1	50.0	17.7	18.0	24.4	24.4	24.6
18 years and over, crude ³	40.3	42.4	46.9	49.4	49.3	17.9	18.1	24.0	23.9	24.0
Age										
18–44 years	45.7	47.7	53.8	56.9	56.7	22.5	22.1	28.8	28.5	29.3
18–24 years	49.3	52.2	57.2	61.5	59.5	28.0	27.2	32.8	33.3	33.8
25–44 years	44.6	46.3	52.5	55.1	55.7	20.8	20.5	27.4	26.8	27.7
45–64 years	38.2	39.7	45.2	46.4	46.2	14.4	15.5	21.5	21.3	21.0
45–54 years	40.1	42.1	47.6	48.4	47.8	16.2	17.0	22.6	23.3	22.4
55–64 years	35.3	36.1	42.1	44.1	44.4	11.5	13.1	20.1	19.0	19.4
65 years and over	26.0	30.1	30.5	35.8	36.5	8.6	9.8	15.4	16.7	16.5
65–74 years	31.7	36.8	35.9	41.8	42.4	9.7	11.3	17.9	19.2	19.0
75 years and over	18.7	22.1	23.9	27.6	28.2	7.2	8.0	12.3	13.4	13.0
Sex ²										
Male	45.4	47.4	52.1	54.3	53.3	21.2	20.8	29.1	28.7	28.8
Female	35.1	37.6	42.7	46.2	47.0	14.4	15.4	19.8	20.2	20.7
Sex and age										
Male:										
18–44 years	51.5	53.6	59.0	61.8	60.8	27.2	26.3	35.6	35.2	35.4
45–54 years	44.3	45.2	50.7	50.4	48.9	18.8	18.0	24.8	25.5	24.5
55–64 years	38.3	38.9	46.0	46.7	44.7	12.9	13.8	22.9	21.1	20.7
65–74 years	38.5	41.8	40.7	46.5	45.6	12.0	12.2	20.6	20.0	21.5
75 years and over	26.1	30.7	32.3	33.2	35.2	9.5	10.1	14.5	14.3	15.3
Female:										
18–44 years	40.0	42.0	48.5	52.0	52.7	17.9	17.9	22.1	22.0	23.4
45–54 years	36.1	39.1	44.7	46.4	46.7	13.7	16.1	20.4	21.2	20.4
55–64 years	32.5	33.5	38.6	41.7	44.2	10.3	12.4	17.5	17.1	18.3
65–74 years	26.2	32.6	31.8	37.7	39.7	7.8	10.5	15.6	18.5	16.8
75 years and over	14.0	16.8	18.3	23.7	23.2	5.7	6.7	10.8	12.7	11.4
Race ^{2,4}										
White only	41.5	44.1	48.9	51.5	51.3	18.0	18.5	24.8	25.0	25.2
Black or African American only	30.4	31.7	37.3	41.4	43.6	15.6	16.0	21.4	21.7	23.1
American Indian or Alaska Native only	39.7	29.7	42.0	47.4	44.1	18.2	13.9	16.7	20.0	27.9
Asian only	37.1	41.7	44.2	49.5	47.5	17.2	17.2	21.9	21.0	19.5
Native Hawaiian or Other Pacific Islander only	---	*	*	*	*	---	*	*	*	*
2 or more races	---	43.9	50.2	51.6	50.5	---	22.2	30.4	26.7	24.8
Hispanic origin and race ^{2,4}										
Hispanic or Latino	29.1	30.8	36.2	42.9	41.3	12.7	11.9	18.1	19.8	19.0
Mexican	27.4	30.0	35.9	43.5	40.6	11.9	11.3	16.7	18.1	18.1
Not Hispanic or Latino	41.3	43.7	49.1	51.5	51.7	18.3	18.8	25.5	25.2	25.8
White only	43.1	45.7	51.5	53.6	53.7	18.7	19.3	26.3	26.2	26.7
Black or African American only	30.4	31.7	37.3	41.4	43.6	15.6	16.0	21.6	21.8	23.3
Education ^{5,6}										
No high school diploma or GED	21.4	23.9	27.1	30.7	31.2	7.0	6.6	10.9	10.9	9.3
High school diploma or GED	32.6	35.7	37.3	39.9	39.7	11.4	12.1	16.2	16.9	16.2
Some college or more	48.1	49.4	53.9	55.7	56.1	22.1	22.4	28.9	28.1	28.9

See footnotes at end of table.

Table 57 (page 4 of 5). Participation in leisure-time aerobic and muscle-strengthening activities that meet the federal 2008 Physical Activity Guidelines for Americans among adults aged 18 and over, by selected characteristics: United States, selected years 1998–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#057>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	2008 Physical Activity Guidelines for Americans ¹									
	Met aerobic activity guideline					Met muscle-strengthening guideline				
	1998	2000	2010	2013	2014	1998	2000	2010	2013	2014
Percent of poverty level ^{2,7}						Percent				
Below 100%	25.9	29.3	32.2	37.2	36.7	10.8	12.3	15.8	15.9	16.2
100%–199%	29.9	32.0	36.0	39.5	40.4	12.0	11.5	16.1	18.0	17.3
200%–399%	38.8	39.9	45.5	48.8	47.8	15.9	16.5	23.1	22.4	23.0
400% or more	50.0	52.0	59.3	61.1	61.4	24.0	23.4	32.8	32.5	32.9
Hispanic origin and race and percent of poverty level ^{2,4,7}										
Hispanic or Latino:										
Below 100%	19.5	22.1	27.8	32.2	31.2	7.1	7.2	12.4	11.6	13.0
100%–199%	25.6	25.8	30.1	36.4	39.0	10.2	7.1	12.6	15.6	16.3
200%–399%	33.1	33.0	38.8	48.6	43.5	14.6	14.0	19.5	23.5	21.8
400% or more	40.6	45.1	53.4	56.8	56.2	21.1	21.7	32.1	31.2	28.7
Not Hispanic or Latino:										
White only:										
Below 100%	30.2	34.0	35.5	40.0	39.7	12.8	14.7	17.5	18.5	18.9
100%–199%	32.2	34.8	40.6	42.7	42.5	12.5	12.9	17.0	19.0	17.6
200%–399%	40.8	42.3	47.8	50.1	49.0	16.2	16.9	23.6	22.0	22.9
400% or more	51.0	53.4	61.0	62.7	63.3	24.0	23.8	33.5	33.3	34.0
Black or African American only:										
Below 100%	22.7	25.4	29.3	33.4	37.0	10.0	12.1	15.3	14.8	15.7
100%–199%	26.9	28.0	28.5	33.0	37.4	12.1	12.3	16.0	18.9	18.1
200%–399%	30.6	31.4	41.9	42.9	45.6	15.5	16.2	25.7	21.6	26.9
400% or more	41.7	40.3	48.5	57.0	54.8	25.4	22.4	29.8	32.8	34.0
Disability measure ^{2,8}										
Any basic actions difficulty or complex activity limitation										
activity limitation	31.8	34.2	36.4	38.6	38.1	13.9	14.0	18.0	18.5	18.0
Any basic actions difficulty	31.3	34.0	36.6	38.3	37.7	13.6	14.2	18.1	18.2	17.9
Any complex activity limitation	24.4	24.9	27.9	29.5	29.2	11.5	11.3	13.9	16.2	15.1
No disability	44.3	46.6	53.4	56.7	56.0	19.3	19.8	27.4	27.4	27.3
Geographic region ²										
Northeast	39.6	45.3	46.9	49.2	48.6	17.5	20.0	24.3	26.0	24.9
Midwest	42.0	43.5	46.1	50.0	48.9	18.2	19.3	24.7	25.3	24.3
South	35.3	37.3	45.0	47.3	48.7	15.0	15.1	22.0	21.5	23.2
West	46.7	46.9	52.0	55.4	54.5	22.3	19.7	27.5	26.9	27.1
Location of residence ^{2,9}										
Within MSA	40.8	42.9	48.7	51.3	51.0	18.3	18.6	25.4	25.4	25.5
Outside MSA	37.1	39.9	39.1	43.4	44.6	15.4	15.5	18.5	18.7	19.0

See footnotes at end of table.

Table 57 (page 5 of 5). Participation in leisure-time aerobic and muscle-strengthening activities that meet the federal 2008 Physical Activity Guidelines for Americans among adults aged 18 and over, by selected characteristics: United States, selected years 1998–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#057>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.
-- Data not available.

¹Starting with *Health, United States, 2010*, measures of physical activity shown in this table changed to reflect the federal 2008 Physical Activity Guidelines for Americans (available from: <http://www.health.gov/PAGuidelines/>). This table presents four measures of physical activity that are of interest to the public health community: the percentage of adults who met the federal 2008 guidelines for both aerobic activity and muscle strengthening; the percentage who met neither the aerobic activity guideline nor the muscle-strengthening guideline; the percentage who met the aerobic activity guideline; and the percentage who met the muscle-strengthening guideline. Persons who met neither the aerobic activity nor the muscle-strengthening guideline were unable to be active, were completely inactive, or had some aerobic or muscle-strengthening activities but amounts were insufficient to meet the guidelines. The percentage of persons who met the aerobic activity guideline includes those who may or may not have also met the muscle-strengthening guideline. Similarly, the percentage of persons who met the muscle-strengthening guideline includes those who may or may not have also met the aerobic activity guideline. The federal 2008 guidelines recommend that for substantial health benefits adults perform at least 150 minutes (2 hours and 30 minutes) a week of moderate-intensity, or 75 minutes (1 hour and 15 minutes) a week of vigorous-intensity aerobic physical activity, or an equivalent combination of moderate- and vigorous-intensity aerobic activity. Aerobic activity should be performed in episodes of at least 10 minutes, and preferably should be spread throughout the week. The 2008 guidelines also recommend that adults perform muscle-strengthening activities that are moderate or high intensity and involve all major muscle groups on 2 or more days a week, because these activities provide additional health benefits. See Appendix II, Physical activity, leisure-time.

²Estimates are age-adjusted to the year 2000 standard population using five age groups: 18–44 years, 45–54 years, 55–64 years, 65–74 years, and 75 years and over. Age-adjusted estimates in this table may differ from other age-adjusted estimates based on the same data and presented elsewhere if different age groups are used in the adjustment procedure. See Appendix II, Age adjustment.

³Includes all other races not shown separately, unknown education level, and unknown disability status.

⁴The race groups, white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The five single-race categories plus multiple-race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group; the category 2 or more races includes persons who reported more than one racial group. Prior to 1999, data were tabulated according to the 1977 Standards with four racial groups, and the Asian only category included Native Hawaiian or Other Pacific Islander. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

⁵Estimates are for persons aged 25 and over and are age-adjusted to the year 2000 standard population using five age groups: 25–44 years, 45–54 years, 55–64 years, 65–74 years, and 75 years and over. See Appendix II, Age adjustment.

⁶GED is General Educational Development high school equivalency diploma. See Appendix II, Education.

⁷Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Missing family income data were imputed for 1997 and beyond. See Appendix II, Family income; Poverty; Table VI.

⁸Any basic actions difficulty or complex activity limitation is defined as having one or more of the following limitations or difficulties: movement difficulty, emotional difficulty, sensory (seeing or hearing) difficulty, cognitive difficulty, self-care (activities of daily living or instrumental activities of daily living) limitation, social limitation, or work limitation. For more information, see Appendix II, Basic actions difficulty; Complex activity limitation. Starting with 2007 data, the hearing question, a component of the basic actions difficulty measure, was revised. Consequently, data prior to 2007 are not comparable with data for 2007 and beyond. For more information on the impact of the revised hearing question, see Appendix II, Hearing trouble.

⁹MSA is metropolitan statistical area. Starting with 2006 data, MSA status is determined using 2000 census data and the 2000 standards for defining MSAs. For data prior to 2006, see Appendix II, Metropolitan statistical area (MSA) for the applicable standards.

NOTES: Standard errors are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey, family core and sample adult questionnaires. See Appendix I, National Health Interview Survey (NHIS).

Table 58 (page 1 of 7). Normal weight, overweight, and obesity among adults aged 20 and over, by selected characteristics: United States, selected years 1988–1994 through 2011–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#058>.

[Data are based on measured height and weight of a sample of the civilian noninstitutionalized population]

Sex, age, race and Hispanic origin ¹ , and percent of poverty level	Normal weight (BMI from 18.5 to 24.9) ²				
	1988–1994	1999–2002	2003–2006	2007–2010	2011–2014
20 years and over, age-adjusted ³					
Percent of population					
Both sexes ⁴	41.6	33.0	31.6	29.8	28.9
Male	37.9	30.2	26.6	25.7	26.0
Female	45.0	35.7	36.5	33.7	31.7
Not Hispanic or Latino:					
White only	43.1	34.6	33.2	31.4	30.0
White only, male	37.3	29.6	26.8	25.5	25.6
White only, female	48.7	39.5	39.6	36.9	34.3
Black or African American only	33.9	27.6	22.7	22.7	22.0
Black or African American only, male	40.1	34.7	27.0	28.5	29.0
Black or African American only, female	29.2	21.6	19.2	17.9	16.0
Asian only	---	---	---	---	55.7
Asian only, male	---	---	---	---	50.2
Asian only, female	---	---	---	---	60.5
Hispanic or Latino	---	---	---	21.1	20.9
Hispanic or Latino, male	---	---	---	19.0	19.5
Hispanic or Latina, female	---	---	---	23.5	22.3
Mexican origin	30.1	26.9	24.4	19.6	17.7
Mexican origin, male	30.2	26.5	23.8	18.5	16.5
Mexican origin, female	29.8	27.5	25.1	21.3	19.1
Percent of poverty level: ⁵					
Below 100%	37.5	32.7	32.1	27.3	28.1
100%–199%	39.3	30.5	31.3	27.6	24.6
200%–399%	41.8	29.6	29.7	29.7	27.5
400% or more	45.5	36.5	33.7	32.1	33.4
20 years and over, crude					
Both sexes ⁴	42.6	32.9	31.4	29.6	28.6
Male	39.4	30.4	26.6	25.8	26.2
Female	45.7	35.4	35.9	33.2	31.0
Not Hispanic or Latino:					
White only	43.6	34.0	32.3	30.4	29.0
White only, male	38.2	29.2	26.2	24.8	25.1
White only, female	48.8	38.7	38.2	35.7	32.8
Black or African American only	35.9	28.2	22.7	23.0	21.9
Black or African American only, male	41.5	35.9	27.1	29.4	29.3
Black or African American only, female	31.2	21.8	19.2	17.6	15.8
Asian only	---	---	---	---	55.8
Asian only, male	---	---	---	---	50.7
Asian only, female	---	---	---	---	60.3
Hispanic or Latino	---	---	---	22.3	21.7
Hispanic or Latino, male	---	---	---	20.3	20.2
Hispanic or Latina, female	---	---	---	24.6	23.1
Mexican origin	34.0	29.4	25.5	20.8	18.4
Mexican origin, male	35.2	29.4	25.2	19.5	17.0
Mexican origin, female	32.5	29.5	25.8	22.3	19.9
Percent of poverty level: ⁵					
Below 100%	39.8	34.5	33.2	29.2	29.8
100%–199%	41.5	31.5	31.7	28.0	25.2
200%–399%	42.9	29.7	29.6	29.5	27.4
400% or more	44.6	35.3	32.1	30.5	31.3
Male					
20–34 years	51.1	40.3	35.9	37.5	37.6
35–44 years	33.4	29.0	24.1	19.8	20.5
45–54 years	33.6	24.0	20.8	21.8	18.7
55–64 years	28.6	23.8	19.3	19.4	22.7
65–74 years	30.1	22.8	21.2	21.6	22.2
75 years and over	40.9	32.0	33.1	25.4	28.1
Female					
20–34 years	57.9	42.5	45.1	41.1	38.1
35–44 years	47.1	37.1	37.6	34.4	32.5
45–54 years	37.2	33.1	31.1	30.7	27.8
55–64 years	31.5	27.6	29.5	26.7	24.2
65–74 years	37.0	26.4	28.5	23.9	26.9
75 years and over	43.0	36.9	35.4	35.4	32.9

See footnotes at end of table.

Table 58 (page 2 of 7). Normal weight, overweight, and obesity among adults aged 20 and over, by selected characteristics: United States, selected years 1988–1994 through 2011–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#058>.

[Data are based on measured height and weight of a sample of the civilian noninstitutionalized population]

Sex, age, race and Hispanic origin ¹ , and percent of poverty level	Overweight or obese (BMI greater than or equal to 25.0) ²				
	1988–1994	1999–2002	2003–2006	2007–2010	2011–2014
20 years and over, age-adjusted ³					
Percent of population					
Both sexes ⁴	56.0	65.1	66.7	68.5	69.5
Male	60.9	68.8	72.1	73.3	73.0
Female	51.4	61.6	61.3	63.9	66.2
Not Hispanic or Latino:					
White only	54.4	63.3	64.8	66.8	68.5
White only, male	61.6	69.4	71.8	73.6	73.7
White only, female	47.5	57.2	57.9	60.3	63.5
Black or African American only	63.7	70.5	76.1	75.5	76.3
Black or African American only, male	57.8	62.6	71.6	70.0	69.6
Black or African American only, female	68.2	77.2	79.8	80.0	82.0
Asian only	---	---	---	---	40.3
Asian only, male	---	---	---	---	46.9
Asian only, female	---	---	---	---	34.4
Hispanic or Latino	---	---	---	78.4	78.4
Hispanic or Latino, male	---	---	---	80.6	79.6
Hispanic or Latina, female	---	---	---	75.7	77.1
Mexican origin	68.9	72.3	75.0	79.9	81.6
Mexican origin, male	68.9	73.2	75.8	81.3	82.7
Mexican origin, female	68.9	71.2	73.9	78.0	80.3
Percent of poverty level: ⁵					
Below 100%	59.6	64.7	65.7	69.7	69.1
100%–199%	58.0	67.3	66.5	70.5	73.9
200%–399%	56.0	68.6	69.0	68.6	71.6
400% or more	52.4	62.2	64.7	66.9	65.6
20 years and over, crude					
Both sexes ⁴	54.9	65.2	66.9	68.7	69.8
Male	59.4	68.6	72.1	73.2	72.8
Female	50.7	62.0	61.9	64.5	67.0
Not Hispanic or Latino:					
White only	53.8	63.9	65.8	67.8	69.6
White only, male	60.6	69.9	72.5	74.2	74.2
White only, female	47.4	58.2	59.4	61.7	65.2
Black or African American only	61.8	70.0	76.0	75.2	76.3
Black or African American only, male	56.7	61.7	71.6	69.1	69.1
Black or African American only, female	66.0	76.9	79.7	80.2	82.3
Asian only	---	---	---	---	40.0
Asian only, male	---	---	---	---	46.3
Asian only, female	---	---	---	---	34.5
Hispanic or Latino	---	---	---	77.1	77.5
Hispanic or Latino, male	---	---	---	79.3	78.8
Hispanic or Latina, female	---	---	---	74.6	76.3
Mexican origin	64.8	69.8	73.9	78.8	81.0
Mexican origin, male	63.9	70.1	74.6	80.2	82.3
Mexican origin, female	65.9	69.3	73.0	77.1	79.5
Percent of poverty level: ⁵					
Below 100%	56.8	62.5	64.4	67.8	67.1
100%–199%	55.7	66.2	66.0	70.1	73.1
200%–399%	54.9	68.5	69.0	68.8	71.8
400% or more	53.3	63.7	66.5	68.5	67.7
Male					
20–34 years	47.5	57.4	61.6	61.1	60.4
35–44 years	65.5	70.5	75.2	80.2	79.3
45–54 years	66.1	75.7	78.5	76.8	80.8
55–64 years	70.5	75.4	79.7	79.8	76.7
65–74 years	68.5	76.2	78.0	77.5	76.1
75 years and over	56.5	67.4	65.8	73.2	71.0
Female					
20–34 years	37.0	52.9	50.9	55.4	58.5
35–44 years	49.6	60.6	60.7	63.9	65.6
45–54 years	60.3	65.1	67.3	66.2	71.4
55–64 years	66.3	72.2	69.6	72.2	74.3
65–74 years	60.3	70.9	70.5	74.2	71.2
75 years and over	52.3	59.9	62.6	63.2	64.6

See footnotes at end of table.

Table 58 (page 3 of 7). Normal weight, overweight, and obesity among adults aged 20 and over, by selected characteristics: United States, selected years 1988–1994 through 2011–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#058>.

[Data are based on measured height and weight of a sample of the civilian noninstitutionalized population]

<i>Sex, age, race and Hispanic origin¹, and percent of poverty level</i>	<i>Obesity (BMI greater than or equal to 30.0)²</i>				
	<i>1988–1994</i>	<i>1999–2002</i>	<i>2003–2006</i>	<i>2007–2010</i>	<i>2011–2014</i>
20 years and over, age-adjusted ³					
Percent of population					
Both sexes ⁴	22.9	30.4	33.4	34.7	36.4
Male	20.2	27.5	32.4	33.9	34.5
Female	25.5	33.2	34.3	35.5	38.1
Not Hispanic or Latino:					
White only	21.6	29.4	32.0	33.3	34.6
White only, male	20.2	28.0	32.4	34.1	34.0
White only, female	22.9	30.7	31.6	32.5	35.3
Black or African American only	30.7	39.2	45.5	47.0	48.0
Black or African American only, male	20.9	27.8	35.7	38.3	37.9
Black or African American only, female	38.3	48.6	53.4	54.0	56.5
Asian only	---	---	---	---	11.8
Asian only, male	---	---	---	---	11.3
Asian only, female	---	---	---	---	11.9
Hispanic or Latino	---	---	---	38.9	42.6
Hispanic or Latino, male	---	---	---	35.7	39.1
Hispanic or Latina, female	---	---	---	42.1	45.6
Mexican origin	29.3	32.7	35.3	40.3	46.4
Mexican origin, male	23.8	27.8	29.5	36.3	43.3
Mexican origin, female	35.2	38.0	41.8	44.6	49.6
Percent of poverty level: ⁵					
Below 100%	28.1	34.7	35.0	37.2	39.2
100%–199%	26.1	34.1	35.9	37.3	42.6
200%–399%	22.7	32.1	35.7	36.8	38.8
400% or more	18.7	25.5	28.9	31.3	29.7
20 years and over, crude					
Both sexes ⁴	22.3	30.5	33.5	34.9	36.5
Male	19.5	27.5	32.4	33.9	34.5
Female	25.0	33.4	34.6	35.9	38.5
Not Hispanic or Latino:					
White only	21.3	29.8	32.4	33.8	35.3
White only, male	19.8	28.4	32.6	34.4	34.3
White only, female	22.7	31.3	32.2	33.2	36.2
Black or African American only	29.5	39.1	45.3	46.9	48.2
Black or African American only, male	20.7	27.5	35.8	38.1	37.6
Black or African American only, female	36.7	48.7	53.2	54.2	56.9
Asian only	---	---	---	---	11.8
Asian only, male	---	---	---	---	11.7
Asian only, female	---	---	---	---	11.9
Hispanic or Latino	---	---	---	38.0	42.3
Hispanic or Latino, male	---	---	---	34.8	39.7
Hispanic or Latina, female	---	---	---	41.5	45.0
Mexican origin	26.4	31.0	34.5	39.5	45.9
Mexican origin, male	20.6	26.0	29.0	35.6	43.5
Mexican origin, female	33.3	37.0	41.2	44.2	48.6
Percent of poverty level: ⁵					
Below 100%	25.9	33.0	34.6	36.5	38.1
100%–199%	24.3	32.8	35.0	36.8	41.9
200%–399%	22.0	31.8	35.5	36.8	38.8
400% or more	19.3	27.2	30.7	32.4	31.1
Male					
20–34 years	14.1	21.7	26.2	27.1	28.5
35–44 years	21.3	28.5	37.0	37.2	39.8
45–54 years	23.2	30.6	34.6	36.6	36.6
55–64 years	27.2	35.5	39.3	37.3	38.1
65–74 years	24.1	31.9	33.0	41.5	36.2
75 years and over	13.2	18.0	24.0	26.6	26.8
Female					
20–34 years	18.5	28.3	28.4	30.4	33.4
35–44 years	25.5	32.1	36.1	37.1	39.1
45–54 years	32.4	36.9	40.0	36.9	41.7
55–64 years	33.7	42.1	41.0	43.4	44.4
65–74 years	26.9	39.3	36.4	40.3	40.7
75 years and over	19.2	23.6	24.2	28.7	30.5

See footnotes at end of table.

Table 58 (page 4 of 7). Normal weight, overweight, and obesity among adults aged 20 and over, by selected characteristics: United States, selected years 1988–1994 through 2011–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#058>.

[Data are based on measured height and weight of a sample of the civilian noninstitutionalized population]

Sex, age, race and Hispanic origin ¹ , and percent of poverty level	Grade 1 Obesity (BMI from 30.0 to 34.9) ²				
	1988–1994	1999–2002	2003–2006	2007–2010	2011–2014
20 years and over, age-adjusted ³					
Percent of population					
Both sexes ⁴	14.8	17.9	19.8	19.9	20.6
Male	14.9	18.2	21.8	22.3	22.0
Female	14.7	17.6	17.9	17.6	19.3
Not Hispanic or Latino:					
White only	14.0	17.6	19.3	19.2	19.6
White only, male	14.9	18.9	21.6	22.7	21.4
White only, female	13.2	16.2	17.0	15.9	17.9
Black or African American only	17.3	19.0	23.1	23.0	23.5
Black or African American only, male	14.2	16.1	22.4	20.8	22.0
Black or African American only, female	19.6	21.6	23.8	24.8	24.9
Asian only	---	---	---	---	9.3
Asian only, male	---	---	---	---	9.4
Asian only, female	---	---	---	---	8.9
Hispanic or Latino	---	---	---	23.7	25.6
Hispanic or Latino, male	---	---	---	23.9	27.2
Hispanic or Latina, female	---	---	---	23.5	24.1
Mexican origin	20.3	21.1	22.8	24.8	26.9
Mexican origin, male	18.9	19.5	22.0	24.7	29.7
Mexican origin, female	22.0	22.9	23.6	24.9	23.8
Percent of poverty level: ⁵					
Below 100%	16.6	17.3	19.3	19.8	20.7
100%–199%	16.1	17.7	20.6	19.8	21.7
200%–399%	14.5	19.8	21.6	20.2	22.7
400% or more	13.3	16.6	18.0	19.4	18.2
20 years and over, crude					
Both sexes ⁴	14.4	17.9	19.8	20.0	20.7
Male	14.3	18.1	21.8	22.3	22.0
Female	14.5	17.7	18.0	17.9	19.4
Not Hispanic or Latino:					
White only	13.8	17.8	19.5	19.6	20.2
White only, male	14.6	19.1	21.8	22.8	22.0
White only, female	13.2	16.6	17.3	16.6	18.5
Black or African American only	16.6	19.0	22.9	22.8	23.3
Black or African American only, male	14.0	15.8	22.2	20.6	21.4
Black or African American only, female	18.7	21.7	23.5	24.6	24.9
Asian only	---	---	---	---	9.4
Asian only, male	---	---	---	---	9.8
Asian only, female	---	---	---	---	9.0
Hispanic or Latino	---	---	---	22.9	25.4
Hispanic or Latino, male	---	---	---	22.9	27.2
Hispanic or Latina, female	---	---	---	22.9	23.7
Mexican origin	18.1	20.2	22.2	24.1	26.9
Mexican origin, male	15.8	18.2	21.6	23.8	29.4
Mexican origin, female	20.7	22.4	22.9	24.5	24.1
Percent of poverty level: ⁵					
Below 100%	15.2	16.4	19.1	19.2	19.5
100%–199%	15.2	17.5	20.4	19.8	21.5
200%–399%	14.0	19.6	21.5	20.3	22.6
400% or more	13.5	17.4	18.6	19.9	19.0
Male					
20–34 years	9.8	13.7	18.1	19.0	16.7
35–44 years	14.7	19.3	24.9	23.2	23.9
45–54 years	17.3	17.8	22.4	22.6	25.8
55–64 years	20.6	25.3	27.0	25.2	25.1
65–74 years	19.4	22.1	20.5	26.1	22.5
75 years and over	10.9	15.7	18.5	20.6	20.3
Female					
20–34 years	10.8	15.9	14.2	14.0	14.7
35–44 years	13.9	14.8	19.7	17.0	21.9
45–54 years	17.5	19.4	18.4	18.6	19.7
55–64 years	20.0	21.6	19.8	22.5	21.7
65–74 years	16.0	23.4	20.3	19.4	21.9
75 years and over	14.4	14.1	18.2	19.8	19.9

See footnotes at end of table.

Table 58 (page 5 of 7). Normal weight, overweight, and obesity among adults aged 20 and over, by selected characteristics: United States, selected years 1988–1994 through 2011–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#058>.

[Data are based on measured height and weight of a sample of the civilian noninstitutionalized population]

Sex, age, race and Hispanic origin ¹ , and percent of poverty level	Grade 2 Obesity (BMI from 35.0 to 39.9) ²				
	1988–1994	1999–2002	2003–2006	2007–2010	2011–2014
20 years and over, age-adjusted ³					
Percent of population					
Both sexes ⁴	5.2	7.6	8.2	8.9	8.8
Male	3.5	5.9	7.1	7.4	7.6
Female	6.8	9.2	9.3	10.3	9.9
Not Hispanic or Latino:					
White only	4.9	7.4	7.8	8.6	8.5
White only, male	3.5	5.8	7.2	7.3	7.8
White only, female	6.3	9.0	8.4	9.9	9.1
Black or African American only	7.8	11.2	12.0	12.0	12.3
Black or African American only, male	4.1	8.3	7.6	10.2	8.9
Black or African American only, female	10.7	13.6	15.4	13.4	15.1
Asian only	---	---	---	---	1.9
Asian only, male	---	---	---	---	*
Asian only, female	---	---	---	---	*2.1
Hispanic or Latino	---	---	---	9.8	10.6
Hispanic or Latino, male	---	---	---	7.8	7.3
Hispanic or Latina, female	---	---	---	12.0	13.5
Mexican origin	6.1	7.4	8.0	9.8	12.4
Mexican origin, male	3.8	5.4	5.1	7.2	8.3
Mexican origin, female	8.4	9.4	11.2	12.9	16.6
Percent of poverty level: ⁵					
Below 100%	6.8	9.6	8.6	10.0	10.1
100%–199%	6.5	9.7	9.0	9.4	11.0
200%–399%	5.2	7.5	8.8	10.3	9.4
400% or more	3.6	5.7	6.7	7.6	6.5
20 years and over, crude					
Both sexes ⁴	5.1	7.7	8.2	8.8	8.8
Male	3.5	6.0	7.0	7.3	7.5
Female	6.6	9.3	9.4	10.3	10.1
Not Hispanic or Latino:					
White only	4.8	7.5	8.0	8.7	8.5
White only, male	3.4	5.9	7.4	7.4	7.6
White only, female	6.2	9.1	8.5	9.9	9.4
Black or African American only	7.6	11.1	11.7	11.9	12.5
Black or African American only, male	4.2	8.2	7.5	10.2	9.1
Black or African American only, female	10.4	13.5	15.3	13.3	15.4
Asian only	---	---	---	---	1.9
Asian only, male	---	---	---	---	*
Asian only, female	---	---	---	---	*2.1
Hispanic or Latino	---	---	---	9.7	10.3
Hispanic or Latino, male	---	---	---	7.6	7.6
Hispanic or Latina, female	---	---	---	12.0	13.1
Mexican origin	5.6	6.8	7.6	9.7	11.7
Mexican origin, male	3.7	5.1	4.7	7.0	8.6
Mexican origin, female	7.9	8.8	11.2	13.0	15.2
Percent of poverty level: ⁵					
Below 100%	6.3	9.5	8.4	9.7	10.1
100%–199%	6.2	8.9	8.7	9.2	10.8
200%–399%	5.1	7.5	8.8	10.1	9.4
400% or more	3.8	6.4	7.4	7.9	6.5
Male					
20–34 years	2.9	4.1	4.5	4.7	7.0
35–44 years	*3.5	5.9	7.9	8.8	10.3
45–54 years	*3.5	8.5	8.3	8.9	6.1
55–64 years	5.5	*7.4	8.4	6.7	6.8
65–74 years	*3.8	6.9	10.3	11.8	8.9
75 years and over	*	*	*3.9	4.6	*4.9
Female					
20–34 years	5.1	8.0	7.9	8.6	9.6
35–44 years	7.1	9.4	9.2	12.6	8.1
45–54 years	8.4	10.4	12.4	10.6	11.5
55–64 years	9.4	10.9	11.4	11.5	11.2
65–74 years	6.7	9.8	9.6	11.7	12.4
75 years and over	3.7	7.2	*3.9	5.5	7.4

See footnotes at end of table.

Table 58 (page 6 of 7). Normal weight, overweight, and obesity among adults aged 20 and over, by selected characteristics: United States, selected years 1988–1994 through 2011–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#058>.

[Data are based on measured height and weight of a sample of the civilian noninstitutionalized population]

Sex, age, race and Hispanic origin ¹ , and percent of poverty level	Grade 3 Obesity (BMI greater than or equal to 40.0) ²				
	1988–1994	1999–2002	2003–2006	2007–2010	2011–2014
20 years and over, age-adjusted ³					
Percent of population					
Both sexes ⁴	2.9	4.9	5.4	6.0	6.9
Male	1.7	3.3	3.5	4.2	4.9
Female	4.0	6.4	7.2	7.6	8.9
Not Hispanic or Latino:					
White only	2.7	4.4	4.9	5.4	6.5
White only, male	*1.8	3.3	3.5	4.0	4.7
White only, female	3.5	5.5	6.3	6.7	8.2
Black or African American only	5.6	8.9	10.4	11.9	12.1
Black or African American only, male	2.5	3.4	5.6	7.3	7.0
Black or African American only, female	8.0	13.4	14.2	15.8	16.5
Asian only	---	---	---	---	*
Asian only, male	---	---	---	---	*
Asian only, female	---	---	---	---	*
Hispanic or Latino	---	---	---	5.3	6.4
Hispanic or Latino, male	---	---	---	4.0	4.7
Hispanic or Latina, female	---	---	---	6.6	8.1
Mexican origin	2.9	4.2	4.6	5.6	7.1
Mexican origin, male	*	*2.9	*2.4	4.4	5.3
Mexican origin, female	4.9	5.7	6.9	6.8	9.2
Percent of poverty level: ⁵					
Below 100%	4.7	7.8	7.0	7.5	8.5
100%–199%	3.6	6.7	6.3	8.1	9.8
200%–399%	3.0	4.8	5.2	6.3	6.7
400% or more	1.9	3.2	4.2	4.4	5.0
20 years and over, crude					
Both sexes ⁴	2.8	4.9	5.4	6.0	7.0
Male	1.7	3.4	3.5	4.3	4.9
Female	3.8	6.4	7.2	7.7	9.0
Not Hispanic or Latino:					
White only	2.6	4.5	4.9	5.5	6.6
White only, male	*1.8	3.4	3.5	4.1	4.7
White only, female	3.3	5.6	6.3	6.8	8.3
Black or African American only	5.3	8.9	10.6	12.2	12.3
Black or African American only, male	2.6	3.5	6.1	7.2	7.1
Black or African American only, female	7.6	13.4	14.4	16.3	16.7
Asian only	---	---	---	---	*
Asian only, male	---	---	---	---	*
Asian only, female	---	---	---	---	*
Hispanic or Latino	---	---	---	5.4	6.6
Hispanic or Latino, male	---	---	---	4.3	5.0
Hispanic or Latina, female	---	---	---	6.5	8.1
Mexican origin	2.7	4.1	4.7	5.7	7.3
Mexican origin, male	*1.1	*2.7	*2.7	4.9	*5.6
Mexican origin, female	4.7	5.7	7.0	6.6	9.3
Percent of poverty level: ⁵					
Below 100%	4.3	7.1	7.1	7.5	8.5
100%–199%	3.0	6.4	5.9	7.9	9.6
200%–399%	2.9	4.7	5.2	6.3	6.7
400% or more	2.0	3.5	4.7	4.6	5.5
Male					
20–34 years	*1.3	3.9	3.6	3.4	4.7
35–44 years	*	*3.2	4.2	5.2	5.6
45–54 years	*	*4.2	*3.9	5.1	*4.6
55–64 years	*	*2.8	3.9	5.4	*6.2
65–74 years	*	*	*2.1	*3.6	*4.7
75 years and over	*	*	*	*	*
Female					
20–34 years	2.6	4.5	6.3	7.7	9.1
35–44 years	4.5	7.9	7.2	7.5	9.0
45–54 years	6.4	7.2	9.2	7.7	10.5
55–64 years	4.2	9.5	9.8	9.4	11.5
65–74 years	4.2	6.2	*6.4	9.2	6.5
75 years and over	*	*	*2.1	*3.4	*3.3

See footnotes at end of table.

Table 58 (page 7 of 7). Normal weight, overweight, and obesity among adults aged 20 and over, by selected characteristics: United States, selected years 1988–1994 through 2011–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#058>.

[Data are based on measured height and weight of a sample of the civilian noninstitutionalized population]

-- Data not available.

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.

¹Persons of Hispanic and Mexican origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The non-Hispanic race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group. Prior to data year 1999, estimates were tabulated according to the 1977 Standards. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. See Appendix II, Hispanic origin; Race.

²Body mass index (BMI) equals weight in kilograms divided by height in meters squared. In *Health, United States* the NHANES variable, Body Mass Index, is used to assign persons to BMI categories. See Appendix II, Body mass index (BMI).

³Estimates are age-adjusted to the year 2000 standard population using five age groups: 20–34 years, 35–44 years, 45–54 years, 55–64 years, and 65 years and over.

Age-adjusted estimates in this table may differ from other age-adjusted estimates based on the same data and presented elsewhere if different age groups are used in the adjustment procedure. See Appendix II, Age adjustment.

⁴Includes persons of all races and Hispanic origins, not just those shown separately.

⁵Percent of poverty level was calculated by dividing family income by the U.S. Department of Health and Human Services' poverty guideline specific to family size, as well as the appropriate year, and state. Persons with unknown percent of poverty level are excluded (6% in 2011–2014). See Appendix II, Family income; Poverty.

NOTES: Percents do not sum to 100 because the percentage of persons with BMI less than normal weight (18.5 kilograms per meters squared) is not shown and the percentage of persons with obesity is a subset of the percentage with overweight. Height was measured without shoes. Excludes pregnant women. See *Health, United States, 2013*, Table 69, for earlier data years. Standard errors for selected years are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health and Nutrition Examination Survey. See Appendix I, National Health and Nutrition Examination Survey (NHANES).

Table 59 (page 1 of 2). Obesity among children and adolescents aged 2–19 years, by selected characteristics: United States, selected years 1988–1994 through 2011–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#059>.

[Data are based on measured height and weight of a sample of the civilian noninstitutionalized population]

<i>Sex, age, race and Hispanic origin¹, and percent of poverty level</i>	1988–1994	1999–2002	2003–2006	2007–2010	2011–2014
2–5 years					
Percent of population					
Both sexes ²	7.2	10.3	12.5	11.1	8.9
Not Hispanic or Latino:					
White only	5.2	8.7	10.8	9.0	*5.2
Black or African American only	7.8	8.8	14.9	15.0	10.4
Asian only	---	---	---	---	*
Hispanic or Latino	---	---	---	15.3	15.6
Mexican origin	12.3	13.1	16.7	14.6	15.3
Boys	6.2	10.0	12.8	11.9	9.2
Not Hispanic or Latino:					
White only	*4.5	*8.2	11.1	8.8	*
Black or African American only	7.9	*8.0	13.3	15.7	*9.0
Asian only	---	---	---	---	*
Hispanic or Latino	---	---	---	17.7	16.7
Mexican origin	12.4	14.1	18.8	19.1	*14.5
Girls	8.2	10.6	12.2	10.2	8.6
Not Hispanic or Latina:					
White only	5.9	*9.0	10.4	*9.2	*4.4
Black or African American only	7.6	9.6	16.6	*14.2	11.9
Asian only	---	---	---	---	*
Hispanic or Latina	---	---	---	12.7	14.6
Mexican origin	12.3	*12.2	14.5	*9.9	*16.1
Percent of poverty level: ³					
Below 100%	9.7	10.9	14.3	13.2	11.6
100%–199%	7.3	*13.8	12.7	11.8	10.2
200%–399%	5.6	*7.6	11.9	13.9	*7.7
400% or more	*	*	*10.0	*5.8	*
6–11 years					
Both sexes ²	11.3	15.9	17.0	18.8	17.5
Not Hispanic or Latino:					
White only	10.2	13.6	15.0	16.4	13.6
Black or African American only	14.6	19.8	21.3	23.9	21.4
Asian only	---	---	---	---	*9.8
Hispanic or Latino	---	---	---	23.8	25.0
Mexican origin	16.4	21.8	23.7	23.3	25.3
Boys	11.6	16.9	18.0	20.7	17.6
Not Hispanic or Latino:					
White only	10.7	14.0	15.5	18.6	13.0
Black or African American only	12.3	17.0	18.6	23.3	21.2
Asian only	---	---	---	---	*14.7
Hispanic or Latino	---	---	---	26.0	25.8
Mexican origin	17.5	26.5	27.5	24.3	25.3
Girls	11.0	14.7	15.8	16.9	17.5
Not Hispanic or Latina:					
White only	*9.8	13.1	14.4	14.0	14.4
Black or African American only	17.0	22.8	24.0	24.5	21.6
Asian only	---	---	---	---	*
Hispanic or Latina	---	---	---	21.5	24.1
Mexican origin	15.3	17.1	19.7	22.4	25.3
Percent of poverty level: ³					
Below 100%	11.4	19.1	22.0	22.2	21.5
100%–199%	11.1	16.4	19.2	20.7	20.4
200%–399%	11.7	15.3	16.7	18.9	15.7
400% or more	*	*12.9	*9.2	*12.5	*12.2

See footnotes at end of table.

Table 59 (page 2 of 2). Obesity among children and adolescents 2–19 years of age, by selected characteristics: United States, selected years 1988–1994 through 2011–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#059>.

[Data are based on physical examinations of a sample of the civilian noninstitutionalized population]

<i>Sex, age, race and Hispanic origin¹, and percent of poverty level</i>	<i>1988–1994</i>	<i>1999–2002</i>	<i>2003–2006</i>	<i>2007–2010</i>	<i>2011–2014</i>
12–19 years					
Percent of population					
Both sexes ²	10.5	16.0	17.6	18.2	20.5
Not Hispanic or Latino:					
White only	10.3	13.7	16.0	15.9	19.6
Black or African American only	13.4	21.1	22.9	24.1	22.6
Asian only	---	---	---	---	9.4
Hispanic or Latino	---	---	---	22.5	22.8
Mexican origin	13.8	22.3	21.1	23.1	23.5
Boys	11.3	16.7	18.2	19.4	20.1
Not Hispanic or Latino:					
White only	11.6	14.6	17.3	17.1	18.7
Black or African American only	10.7	18.8	18.4	21.2	20.9
Asian only	---	---	---	---	12.9
Hispanic or Latino	---	---	---	26.0	22.7
Mexican origin	14.1	24.7	22.1	27.9	22.8
Girls	9.7	15.3	16.8	16.9	21.0
Not Hispanic or Latina:					
White only	8.9	12.6	14.5	14.6	20.4
Black or African American only	16.3	23.5	27.7	27.1	24.4
Asian only	---	---	---	---	*5.7
Hispanic or Latina	---	---	---	18.7	22.8
Mexican origin	*13.4	19.6	19.9	18.0	24.2
Percent of poverty level: ³					
Below 100%	15.8	19.8	19.3	24.3	22.4
100%–199%	11.2	15.1	18.4	20.1	25.7
200%–399%	9.4	15.7	19.3	16.3	19.7
400% or more	*	13.9	12.6	14.0	*13.7

--- Data not available.

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.

¹Persons of Hispanic and Mexican origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The non-Hispanic race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group. Prior to data year 1999, estimates were tabulated according to the 1977 Standards. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. See Appendix II, Hispanic origin; Race.

²Includes persons of all races and Hispanic origins, not just those shown separately.

³Percent of poverty level was calculated by dividing family income by the U.S. Department of Health and Human Services' poverty guideline specific to family size, as well as the appropriate year, and state. Persons with unknown percent of poverty level are excluded (6% in 2011–2014). See Appendix II, Family income; Poverty.

NOTES: Obesity is defined as body mass index (BMI) at or above the sex- and age-specific 95th percentile from the 2000 CDC Growth Charts: United States. Kuczmarski RJ, Ogden CL, Guo SS, Grummer-Strawn LM, Flegal KM, Mei Z, Wei R, Curtin LR, Roche AF, Johnson CL. 2000 CDC Growth Charts for the United States: methods and development. *Vital Health Stat 11*. 2002 May;(246):1–190. Available at: http://www.cdc.gov/nchs/data/series/sr_11/sr11_246.pdf. In *Health, United States*, the NHANES variable, Body Mass Index, is used to assign persons to BMI categories. Age is at time of examination at the mobile examination center. Crude rates, not age-adjusted rates, are shown. Height was measured without shoes. Excludes pregnant females. See *Health, United States, 2013*, Table 70, for earlier data years. Standard errors for selected years are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>. Data for 1988–1994 have been revised and differ from previous editions.

SOURCE: CDC/NCHS, National Health and Nutrition Examination Survey. See Appendix I, National Health and Nutrition Examination Survey (NHANES).

Table 60 (page 1 of 2). Untreated dental caries, by selected characteristics: United States, selected years 1988–1994 through 2011–2012

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#060>.

[Data are based on dental examinations of a sample of the civilian noninstitutionalized population]

Sex, race and Hispanic origin ¹ , and percent of poverty level	Age 5–19 years				Age 20–44 years			
	1988–1994	1999–2002	2005–2008	2011–2012	1988–1994	1999–2002	2005–2008	2011–2012
Percent of persons with untreated dental caries								
Total ²	24.3	22.5	16.6	17.5	29.5	26.0	25.1	27.4
Sex								
Male	23.6	23.7	17.6	18.9	32.8	27.0	28.4	29.0
Female	25.0	21.3	15.5	16.1	26.4	24.9	21.8	25.9
Race and Hispanic origin								
Not Hispanic or Latino:								
White only	19.4	18.5	13.3	14.5	24.8	20.7	21.1	22.1
Black or African American only	33.9	29.2	22.6	23.2	49.2	43.4	36.7	41.4
Asian only	---	---	---	*15.2	---	---	---	17.7
Hispanic or Latino								
Mexican origin	37.9	33.9	22.4	24.2	40.0	35.6	35.1	38.0
Percent of poverty level: ³								
Below 100%	39.0	31.9	25.4	24.6	47.8	42.4	39.8	40.2
100%–199%	29.6	29.7	19.3	21.6	43.7	36.7	37.7	38.0
200%–399%	16.6	18.0	14.7	15.4	24.5	24.9	22.3	22.0
400% or more	*10.4	8.9	9.3	*4.8	12.5	9.8	12.4	*11.3
Race and Hispanic origin, and percent of poverty level ³								
Not Hispanic or Latino:								
White only:								
Below 100% of poverty level	33.8	28.0	25.0	21.7	42.9	36.9	37.7	*39.0
100% or more of poverty level	17.3	16.5	11.6	12.3	22.7	18.5	18.6	17.8
Black or African American only:								
Below 100% of poverty level	37.4	36.7	27.3	29.2	60.0	55.3	48.7	48.0
100% or more of poverty level	31.2	25.0	19.5	18.0	44.8	37.6	33.8	37.8
Asian only:								
Below 100% of poverty level	---	---	---	*	---	---	---	*30.0
100% or more of poverty level	---	---	---	*13.7	---	---	---	14.6
Hispanic or Latino:								
Below 100% of poverty level	---	---	---	24.5	---	---	---	38.3
100% or more of poverty level	---	---	---	19.4	---	---	---	32.1
Mexican origin:								
Below 100% of poverty level	47.5	40.2	25.9	24.2	52.7	42.2	42.1	38.0
100% or more of poverty level	28.0	27.0	20.5	22.8	31.2	32.5	31.0	35.1

See footnotes at end of table.

Table 60 (page 2 of 2). Untreated dental caries, by selected characteristics: United States, selected years 1988–1994 through 2011–2012

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#060>.

[Data are based on dental examinations of a sample of the civilian noninstitutionalized population]

Sex, race and Hispanic origin ¹ , and percent of poverty level	Age 45–64 years				Age 65 years and over			
	1988–1994	1999–2002	2005–2008	2011–2012	1988–1994	1999–2002	2005–2008	2011–2012
Percent of persons with untreated dental caries								
Total ²	25.4	20.1	21.6	25.8	27.1	18.4	19.9	18.9
Sex								
Male	28.5	24.0	25.4	31.7	31.2	21.8	25.1	20.8
Female	22.6	16.5	18.0	20.3	24.1	15.7	15.6	17.3
Race and Hispanic origin								
Not Hispanic or Latino:								
White only	21.7	15.6	17.1	22.1	24.6	16.0	17.8	15.5
Black or African American only	46.2	39.8	44.4	43.2	51.2	38.6	35.8	40.9
Asian only	---	---	---	16.0	---	---	---	27.3
Hispanic or Latino								
Mexican origin	41.4	34.3	35.4	44.1	46.3	37.9	36.4	48.3
Percent of poverty level: ³								
Below 100%	49.5	39.3	47.6	51.6	46.8	30.5	41.3	39.5
100%–199%	42.5	35.9	37.7	41.0	37.6	25.3	22.5	33.1
200%–399%	25.0	24.8	27.6	28.1	24.1	15.6	16.6	*12.2
400% or more	13.0	9.6	10.0	11.7	15.6	11.1	13.1	*
Race and Hispanic origin, and percent of poverty level ³								
Not Hispanic or Latino:								
White only:								
Below 100% of poverty level	47.6	31.2	45.4	*51.6	38.5	*27.1	*35.6	46.1
100% or more of poverty level	19.7	14.2	15.4	19.8	23.9	15.4	16.1	14.1
Black or African American only:								
Below 100% of poverty level	62.9	52.5	61.0	69.7	56.3	40.1	55.7	*44.5
100% or more of poverty level	41.2	37.7	41.4	35.2	47.8	40.3	30.6	39.1
Asian only:								
Below 100% of poverty level	---	---	---	*31.3	---	---	---	*
100% or more of poverty level	---	---	---	*12.7	---	---	---	*23.8
Hispanic or Latino:								
Below 100% of poverty level	---	---	---	36.1	---	---	---	*
100% or more of poverty level	---	---	---	35.9	---	---	---	*25.0
Mexican origin:								
Below 100% of poverty level	52.7	49.0	51.6	*45.7	62.8	46.8	56.3	*56.7
100% or more of poverty level	34.5	31.1	30.9	43.4	35.4	36.2	26.2	*45.0

--- Data not available.

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE of greater than 30%.

¹Persons of Hispanic and Mexican origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The non-Hispanic race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group. Prior to data year 1999, estimates were tabulated according to the 1977 Standards. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. See Appendix II, Hispanic origin; Race.

²Includes persons of all races and Hispanic origins, not just those shown separately, and those with unknown percent of poverty level.

³Percent of poverty level was calculated by dividing family income by the U.S. Department of Health and Human Services' poverty guideline specific to family size, as well as the appropriate year, and state. Persons with unknown percent of poverty level are excluded (6% in 2011–2012). See Appendix II, Family income; Poverty.

NOTES: Untreated dental caries refers to decay on the crown or enamel surface of a tooth (i.e., coronal caries) that has not been treated or filled. Decay in the root (i.e., root caries) was not included. The presence of caries was evaluated in primary and permanent teeth for persons aged 5 and older. The third molars were not included. Persons without at least one natural tooth (primary or permanent) were classified as edentulous (without any teeth) and were excluded. The majority of edentulous persons are aged 65 and over. Estimates of edentulism among persons aged 65 and over are 33% in 1988–1994, 23% in 2005–2008, and 19% in 2011–2012. Over time, there have been changes in the NHANES oral health examination process, ages examined, and methodology. Therefore, data trends need to be interpreted with caution. For more information on the methodology changes, see Appendix II, Dental caries. Standard errors are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health and Nutrition Examination Survey. See Appendix I, National Health and Nutrition Examination Survey (NHANES).

Table 61 (page 1 of 2). No usual source of health care among children under age 18, by selected characteristics: United States, average annual, selected years 1993–1994 through 2013–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#061>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Under 18 years			Under 6 years			6–17 years		
	1993–1994 ¹	1999–2000	2013–2014	1993–1994 ¹	1999–2000	2013–2014	1993–1994 ¹	1999–2000	2013–2014
	Percent of children without a usual source of health care ²								
All children ³	7.7	6.9	3.8	5.2	4.6	2.6	9.0	8.0	4.4
Sex									
Male	8.1	6.7	3.9	5.3	4.5	2.5	9.6	7.8	4.6
Female	7.3	7.1	3.7	5.0	4.7	2.8	8.5	8.2	4.2
Race ⁴									
White only	7.0	6.3	3.7	4.7	4.4	2.5	8.3	7.2	4.3
Black or African American only	10.3	7.7	3.9	7.6	4.4	*2.4	11.9	9.1	4.7
American Indian or Alaska Native only	*9.3	*9.4	*4.8	*	*	*	*8.7	*9.4	*
Asian only	9.7	10.0	4.6	*3.4	*5.8	*3.8	13.5	12.2	5.1
Native Hawaiian or Other Pacific Islander only	---	*	*	---	*	*	---	*	*
2 or more races	---	*4.9	4.0	---	*	*	---	*7.2	4.6
Hispanic origin and race ⁴									
Hispanic or Latino	14.3	14.2	6.5	9.3	9.0	4.2	17.7	17.2	7.8
Not Hispanic or Latino	6.7	5.5	3.0	4.4	3.6	2.1	7.8	6.3	3.4
White only	5.7	4.7	2.5	3.7	3.3	1.7	6.7	5.4	2.9
Black or African American only	10.2	7.6	3.8	7.7	4.5	*2.5	11.6	9.0	4.4
Percent of poverty level ⁵									
Below 100%	13.9	13.1	6.5	9.4	7.6	4.7	16.8	16.2	7.5
100%–199%	9.8	10.6	5.2	6.7	7.5	3.6	11.6	12.2	5.9
200%–399%	3.7	4.8	2.7	1.9	3.2	1.4	4.5	5.6	3.3
400% or more	3.7	2.6	1.5	*1.6	1.5	*1.0	5.0	3.0	1.8
Hispanic origin and race and percent of poverty level ^{4,5}									
Hispanic or Latino:									
Below 100%	19.6	19.4	8.5	12.7	11.6	5.9	24.8	24.5	10.1
100%–199%	15.3	17.1	7.3	9.9	11.3	*3.4	18.9	20.4	9.3
200%–399%	5.2	8.3	4.0	*	*5.0	*3.4	6.7	10.1	4.4
400% or more	*	*3.8	*2.0	*	*	*	*	*5.0	*
Not Hispanic or Latino:									
White only:									
Below 100%	10.2	10.7	5.1	6.5	*6.3	*	12.7	13.1	5.6
100%–199%	8.7	7.8	3.6	6.3	5.7	*3.2	10.1	8.8	3.8
200%–399%	3.3	4.0	2.3	1.6	2.7	*	4.0	4.6	3.0
400% or more	4.0	2.3	1.3	*1.7	*1.5	*0.7	5.4	2.6	1.6
Black or African American only:									
Below 100%	13.7	9.4	4.7	10.9	*4.7	*3.3	15.5	11.8	5.4
100%–199%	9.1	9.7	3.6	*6.0	*6.4	*	10.8	11.2	*4.0
200%–399%	5.0	5.0	*	*	*	*	6.2	5.7	*
400% or more	*	*3.5	*	*	*	*	*	*4.0	*
Health insurance status at the time of interview ⁶									
Insured	5.0	3.9	2.4	3.3	2.6	1.8	5.9	4.5	2.8
Private	3.8	3.4	1.9	1.9	2.2	1.1	4.6	3.9	2.3
Medicaid	8.9	5.3	3.2	6.4	3.5	2.6	11.3	6.7	3.5
Uninsured	23.5	29.3	25.8	18.0	20.8	20.3	26.0	32.9	27.7
Health insurance status prior to interview ⁶									
Insured continuously all 12 months	4.6	3.6	2.2	3.1	2.3	1.6	5.5	4.2	2.5
Uninsured for any period up to 12 months	15.3	15.0	13.2	10.9	12.5	11.0	18.1	16.4	14.3
Uninsured more than 12 months	27.6	35.8	31.8	21.4	26.8	*22.0	30.0	39.1	34.4

See footnotes at end of table.

Table 61 (page 2 of 2). No usual source of health care among children under age 18, by selected characteristics: United States, average annual, selected years 1993–1994 through 2013–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#061>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Under 18 years			Under 6 years			6–17 years		
	1993–1994 ¹	1999–2000	2013–2014	1993–1994 ¹	1999–2000	2013–2014	1993–1994 ¹	1999–2000	2013–2014
Percent of poverty level and health insurance status prior to interview ^{5,6}									
Percent of children without a usual source of health care ²									
Below 100%:									
Insured continuously all 12 months	8.6	5.7	3.4	5.8	*2.7	2.8	10.7	7.5	3.7
Uninsured for any period up to 12 months . . .	21.7	19.8	21.3	18.0	*16.0	*19.8	23.7	21.9	22.1
Uninsured more than 12 months	31.2	42.7	45.2	25.5	31.0	*	33.4	47.1	48.8
100%–199%:									
Insured continuously all 12 months	5.6	5.2	2.5	3.7	3.7	*1.9	6.7	6.0	2.8
Uninsured for any period up to 12 months . . .	14.5	15.4	13.9	*9.7	*14.4	*12.3	18.0	15.9	14.8
Uninsured more than 12 months	27.6	34.4	31.2	21.4	26.4	*	30.2	37.4	32.3
200%–399%:									
Insured continuously all 12 months	2.8	3.2	1.8	1.5	2.1	*1.0	3.4	3.7	2.2
Uninsured for any period up to 12 months . . .	9.1	11.1	8.2	*	*8.4	*	11.6	12.7	*10.1
Uninsured more than 12 months	18.2	27.1	*18.7	*9.7	*20.3	*	21.0	29.4	*21.5
400% or more:									
Insured continuously all 12 months	3.1	2.0	1.3	*	*1.2	*0.8	4.3	2.4	1.6
Uninsured for any period up to 12 months . . .	*	*10.3	*	*	*	*	*	*	*
Uninsured more than 12 months	*	*30.0	*	*	*	*	*	*33.3	*
Geographic region									
Northeast	4.1	2.8	1.7	2.9	2.3	*1.4	4.8	3.0	1.8
Midwest	5.2	5.3	2.9	4.1	3.7	2.4	5.9	6.0	3.1
South	10.9	8.5	4.5	7.3	5.8	2.7	12.7	9.8	5.4
West	8.6	9.7	5.0	5.3	5.7	3.5	10.6	11.7	5.7
Location of residence ⁷									
Within MSA	7.7	6.8	3.8	5.0	4.7	2.7	9.2	7.8	4.3
Outside MSA	7.8	7.4	3.9	6.0	4.2	*2.1	8.7	8.7	4.7

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.

--- Data not available.

¹Data prior to 1997 are not strictly comparable with data for later years due to the 1997 questionnaire redesign. See Appendix I, National Health Interview Survey (NHIS).

²Persons who report the emergency department as their usual source of care are defined as having no usual source of care. See Appendix II, Usual source of care.

³Includes all other races not shown separately and unknown health insurance status.

⁴The race groups, white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The five single-race categories plus multiple-race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group; the category 2 or more races includes persons who reported more than one racial group. Prior to 1999, data were tabulated according to the 1977 Standards with four racial groups, and the Asian only category included Native Hawaiian or Other Pacific Islander. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

⁵Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Missing family income data were imputed starting in 1993. See Appendix II, Family income; Poverty; Table VI.

⁶Health insurance categories are mutually exclusive. Persons who reported both Medicaid and private coverage are classified as having private coverage. Medicaid includes other public assistance through 1996. Starting with 1997 data, state-sponsored health plan coverage is included as Medicaid coverage. Starting with 1999 data, coverage by the Children's Health Insurance Program (CHIP) is included with Medicaid coverage. In addition to private and Medicaid, the insured category also includes military, other government, and Medicare coverage. Persons not covered by private insurance, Medicaid, CHIP, public assistance (through 1996), state-sponsored or other government-sponsored health plans (starting in 1997), Medicare, or military plans are considered to have no health insurance coverage. Persons with only Indian Health Service coverage are considered to have no health insurance coverage. Health insurance status was unknown for 8%–9% of children in 1993–1996 and about 1% in 1997–2014. See Appendix II, Health insurance coverage.

⁷MSA is metropolitan statistical area. Starting with 2005–2006 data, MSA status is determined using 2000 census data and the 2000 standards for defining MSAs. For data prior to 2005, see Appendix II, Metropolitan statistical area (MSA) for the applicable standards.

NOTES: Standard errors are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey, access to care and health insurance supplements (1993–1996). Starting in 1997, data are from the family core and sample child questionnaires. See Appendix I, National Health Interview Survey (NHIS).

Table 62 (page 1 of 2). No usual source of health care among adults aged 18–64, by selected characteristics: United States, average annual, selected years 1993–1994 through 2013–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#062>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	1993–1994 ¹	1999–2000	2003–2004	2005–2006	2007–2008	2009–2010	2011–2012	2013–2014
Percent of adults without a usual source of health care ²								
18–64 years ³	18.9	17.8	17.3	18.4	18.5	20.3	19.5	18.2
Age								
18–44 years	21.7	21.6	21.7	23.5	23.6	26.0	25.0	23.8
18–24 years	26.6	27.2	28.0	29.8	28.6	29.8	27.8	26.3
19–25 years	28.0	29.0	29.7	31.8	30.0	33.1	30.3	28.5
25–44 years	20.3	19.9	19.5	21.3	21.8	24.7	23.9	22.8
45–64 years	12.8	10.9	10.4	10.7	11.0	12.3	12.0	10.8
45–54 years	14.1	12.0	11.7	12.3	13.1	14.7	14.2	12.8
55–64 years	11.1	9.2	8.7	8.4	8.3	9.3	9.6	8.6
Sex								
Male	23.9	24.1	22.5	23.9	23.9	25.9	24.4	23.1
Female	14.1	11.8	12.4	13.0	13.1	14.8	14.8	13.6
Race ⁴								
White only	18.4	16.7	17.0	18.1	18.0	19.7	18.9	17.8
Black or African American only	20.0	19.2	18.4	19.8	20.5	22.4	21.9	19.9
American Indian or Alaska Native only	19.7	19.2	21.5	21.9	24.4	26.7	23.6	23.4
Asian only	24.8	22.1	19.3	17.9	17.8	20.8	20.8	18.1
Native Hawaiian or Other Pacific Islander only	---	*	*	*	*	*	*	*
2 or more races	---	21.0	18.4	20.9	21.4	27.5	22.3	23.9
American Indian or Alaska Native; White	---	25.8	17.8	21.4	20.9	27.1	19.0	24.2
Hispanic origin and race ⁴								
Hispanic or Latino	30.3	32.6	32.9	35.1	32.5	33.3	33.6	28.9
Mexican	32.4	36.5	36.4	39.3	36.6	35.7	35.6	30.6
Not Hispanic or Latino	17.7	15.8	14.9	15.6	16.0	17.9	16.8	16.1
White only	17.1	14.9	14.0	14.8	15.1	16.8	15.5	15.0
Black or African American only	19.7	19.2	18.1	19.2	20.2	22.2	21.6	19.6
Percent of poverty level ⁵								
Below 100%	29.5	29.6	28.9	32.1	30.4	33.8	32.1	30.6
100%–199%	25.4	27.1	26.6	27.8	29.1	30.5	30.2	26.6
200%–399%	15.6	17.2	17.3	17.8	18.9	20.5	19.3	17.9
400% or more	13.4	11.6	10.1	10.4	10.2	10.8	9.6	9.7
Hispanic origin and race and percent of poverty level ^{4,5}								
Hispanic or Latino:								
Below 100%	40.0	44.4	42.8	46.7	43.7	45.5	42.9	40.1
100%–199%	36.9	40.6	39.7	41.8	40.6	39.7	40.0	34.3
200%–399%	20.7	26.9	28.2	31.2	28.0	29.1	29.4	24.2
400% or more	13.8	16.1	16.4	16.4	16.9	14.0	15.4	13.2
Not Hispanic or Latino:								
White only:								
Below 100%	28.2	24.2	23.0	26.2	25.2	28.8	27.0	26.9
100%–199%	23.3	23.0	22.0	23.5	24.9	26.6	25.7	23.4
200%–399%	14.8	15.3	15.4	15.3	16.7	18.6	16.9	16.3
400% or more	13.4	11.2	9.4	9.8	9.5	10.3	8.8	9.1
Black or African American only:								
Below 100%	24.7	23.7	24.3	29.5	27.1	30.1	29.9	27.4
100%–199%	22.3	24.4	22.8	22.6	25.7	28.5	28.2	23.7
200%–399%	16.5	18.2	16.3	16.2	19.7	20.1	18.5	17.3
400% or more	11.7	12.0	11.3	10.3	10.2	10.5	10.1	9.5
Health insurance status at the time of interview ⁶								
Insured	13.3	10.9	9.4	9.7	10.1	10.6	10.5	10.5
Private	13.1	11.1	9.5	9.6	10.0	10.6	10.1	10.5
Medicaid	16.3	9.9	9.9	11.6	11.7	12.5	13.1	11.9
Uninsured	43.1	49.2	50.2	53.0	52.1	55.6	54.1	52.5
Health insurance status prior to interview ⁶								
Insured continuously all 12 months	12.7	10.3	8.7	8.9	9.1	9.8	9.6	9.5
Uninsured for any period up to 12 months	30.9	31.2	32.1	33.4	35.1	36.5	33.2	32.5
Uninsured more than 12 months	46.9	54.8	55.0	58.0	56.1	59.5	57.8	56.4

See footnotes at end of table.

Table 62 (page 2 of 2). No usual source of health care among adults aged 18–64, by selected characteristics: United States, average annual, selected years 1993–1994 through 2013–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#062>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	1993–1994 ¹	1999–2000	2003–2004	2005–2006	2007–2008	2009–2010	2011–2012	2013–2014
Percent of poverty level and health insurance status prior to interview ^{5,6}		Percent of adults without a usual source of health care ²						
Below 100%:								
Insured continuously all 12 months	16.7	11.6	11.2	12.0	12.7	13.0	14.0	13.4
Uninsured for any period up to 12 months . .	33.6	31.9	36.2	36.5	37.4	37.8	35.6	38.3
Uninsured more than 12 months	50.1	57.1	57.2	63.2	61.1	65.3	61.3	58.8
100%–199%:								
Insured continuously all 12 months	14.7	12.3	10.5	10.4	11.9	12.5	12.8	11.9
Uninsured for any period up to 12 months . .	30.9	34.6	34.2	37.8	35.9	38.1	35.9	30.0
Uninsured more than 12 months	47.6	54.9	55.1	57.0	56.8	58.5	57.9	57.4
200%–399%:								
Insured continuously all 12 months	11.7	10.6	9.4	9.4	9.4	10.6	10.0	9.7
Uninsured for any period up to 12 months . .	29.2	29.0	30.9	31.3	36.3	37.6	33.2	32.8
Uninsured more than 12 months	44.5	53.6	54.2	55.5	54.2	56.6	55.3	55.1
400% or more:								
Insured continuously all 12 months	11.8	9.3	7.5	7.7	7.5	7.9	7.4	7.7
Uninsured for any period up to 12 months . .	31.5	30.2	27.5	28.6	30.3	31.2	25.6	27.5
Uninsured more than 12 months	36.5	51.8	51.6	54.2	47.9	53.8	52.9	47.4
Disability measure ⁷								
Any basic actions difficulty or complex activity limitation	---	14.1	14.3	15.2	16.6	16.8	16.5	15.0
Any basic actions difficulty	---	14.1	14.5	15.4	16.5	16.7	16.5	14.9
Any complex activity limitation	---	11.6	10.7	11.1	13.6	13.5	13.5	11.6
No disability	---	18.8	18.2	19.4	19.1	21.5	20.5	19.3
Geographic region								
Northeast	14.7	12.8	12.1	12.2	12.5	14.0	13.1	12.2
Midwest	16.2	17.0	14.7	15.8	16.6	17.5	17.1	16.5
South	21.8	19.7	19.7	21.4	21.4	23.5	22.2	20.4
West	21.1	20.1	21.0	21.1	20.0	22.9	22.8	20.9
Location of residence ⁸								
Within MSA	19.3	18.1	17.6	18.7	18.7	20.3	19.8	18.4
Outside MSA	17.5	16.8	16.2	16.7	16.9	20.4	17.8	17.1

--- Data not available.

* Estimates are considered unreliable. Data not shown have a relative standard error greater than 30%.

¹Data prior to 1997 are not strictly comparable with data for later years due to the 1997 questionnaire redesign. See Appendix I, National Health Interview Survey (NHIS).

²Persons who report the emergency department as their usual source of care are defined as having no usual source of care. See Appendix II, Usual source of care.

³Includes all other races not shown separately, unknown health insurance status, and unknown disability status.

⁴The race groups, white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity and are not strictly comparable with estimates for earlier years. The five single-race categories plus multiple-race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group; the category 2 or more races includes persons who reported more than one racial group. Prior to 1999, data were tabulated according to the 1977 Standards with four racial groups, and the Asian only category included Native Hawaiian or Other Pacific Islander. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

⁵Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Missing family income data were imputed starting in 1993. See Appendix II, Family income; Poverty; Table VI.

⁶Health insurance categories are mutually exclusive. Persons who reported both Medicaid and private coverage are classified as having private coverage. Medicaid includes other public assistance through 1996. Starting with 1997 data, state-sponsored health plan coverage is included as Medicaid coverage. Starting with 1999 data, coverage by the Children's Health Insurance Program (CHIP) is included with Medicaid coverage. In addition to private and Medicaid, the insured category also includes military, other government, and Medicare coverage. Persons not covered by private insurance, Medicaid, CHIP, public assistance (through 1996), state-sponsored or other government-sponsored health plans (starting in 1997), Medicare, or military plans are considered to have no health insurance coverage. Persons with only Indian Health Service coverage are considered to have no health insurance coverage. In 1993–1996, health insurance status was unknown for 8%–9% of adults in the sample. In 1997–2014, health insurance status was unknown for about 1% of adults aged 18–64. See Appendix II, Health insurance coverage.

⁷Any basic actions difficulty or complex activity limitation is defined as having one or more of the following limitations or difficulties: movement difficulty, emotional difficulty, sensory (seeing or hearing) difficulty, cognitive difficulty, self-care (activities of daily living or instrumental activities of daily living) limitation, social limitation, or work limitation. For more information, see Appendix II, Basic actions difficulty; Complex activity limitation. Starting with 2007 data, the hearing question, a component of the basic actions difficulty measure, was revised. Consequently, data prior to 2007 are not comparable with data for 2007 and beyond. For more information on the impact of the revised hearing question, see Appendix II, Hearing trouble.

⁸MSA is metropolitan statistical area. Starting with 2005–2006 data, MSA status is determined using 2000 census data and the 2000 standards for defining MSAs. For data prior to 2005, see Appendix II, Metropolitan statistical area (MSA) for the applicable standards.

NOTES: Standard errors are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey, access to care and health insurance supplements (1993–1996). Starting in 1997, data are from the family core and sample adult questionnaires. See Appendix I, National Health Interview Survey (NHIS).

Table 63 (page 1 of 3). Delay or nonreceipt of needed medical care, nonreceipt of needed prescription drugs, or nonreceipt of needed dental care during the past 12 months due to cost, by selected characteristics: United States, selected years 1997–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#063>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Delay or nonreceipt of needed medical care due to cost ¹				Nonreceipt of needed prescription drugs due to cost ²				Nonreceipt of needed dental care due to cost ³			
	1997	2004	2010	2014	1997	2004	2010	2014	1997	2004	2010	2014
Total ⁴	8.3	8.8	10.9	8.2	4.8	7.1	8.3	5.6	8.6	10.7	13.5	10.0
	Percent											
Age												
Under 19 years	4.5	4.5	4.5	2.9	2.1	2.9	2.8	1.6	6.0	6.6	6.6	4.3
Under 18 years	4.4	4.4	4.4	2.8	2.2	2.8	2.7	1.6	6.0	6.6	6.6	4.2
Under 6 years	3.3	3.7	3.7	2.1	1.6	2.2	2.5	1.3	3.9	3.8	3.9	2.8
6–17 years	4.9	4.7	4.8	3.2	2.4	3.1	2.8	1.7	6.8	7.5	7.5	4.7
18–64 years	10.7	11.4	14.7	11.2	6.3	9.2	11.2	7.6	10.6	13.2	17.3	12.6
18–44 years	11.0	11.4	14.5	10.7	6.9	9.4	11.2	7.1	11.7	14.5	17.9	12.2
18–24 years	10.2	11.4	13.5	8.2	6.7	9.3	9.7	5.9	11.6	14.8	17.4	9.2
25–34 years	11.4	11.8	15.3	12.3	6.9	9.2	12.0	7.5	12.3	14.7	18.3	14.2
35–44 years	11.0	11.1	14.4	11.0	7.1	9.5	11.3	7.7	11.2	14.1	17.8	12.3
19–25 years	11.1	12.6	14.8	9.1	7.7	10.1	10.9	6.5	13.1	16.2	18.9	10.2
45–64 years	10.1	11.2	14.9	11.7	5.1	8.8	11.3	8.3	8.4	11.3	16.5	13.3
45–54 years	10.6	11.6	15.0	11.8	5.6	9.7	11.5	8.3	9.4	12.6	17.8	13.2
55–64 years	9.3	10.7	14.6	11.7	4.2	7.7	11.0	8.2	7.0	9.4	14.9	13.3
65 years and over	4.6	4.7	5.0	4.3	2.8	5.4	4.7	3.7	3.5	5.5	6.9	7.2
65–74 years	5.0	5.5	6.3	5.6	3.4	6.2	6.3	4.7	4.2	6.7	9.0	8.3
75 years and over	4.1	3.9	3.4	2.6	2.0	4.6	2.8	2.3	2.6	4.2	4.3	5.6
18–64 years												
Sex												
Male	9.3	10.3	13.5	10.1	5.1	7.3	8.8	5.8	8.8	11.4	15.2	10.5
Female	12.0	12.4	15.7	12.2	7.4	11.0	13.5	9.3	12.4	15.0	19.4	14.7
Race ⁵												
White only	10.8	11.5	14.5	11.1	5.9	8.9	10.8	7.2	10.6	13.3	17.1	12.6
Black or African American only	10.8	11.7	17.4	13.3	9.5	12.3	15.6	10.9	10.8	14.2	20.7	14.8
American Indian or Alaska Native only	14.5	13.7	*15.7	12.0	*10.1	*11.9	18.6	*12.0	18.8	20.7	23.1	*13.3
Asian only	6.3	6.3	8.0	6.2	*2.8	*3.0	4.2	*3.3	7.8	5.7	8.7	6.6
Native Hawaiian or Other Pacific Islander only	---	*	*	*	---	*	*	*	---	*	*	*
2 or more races	---	16.6	24.0	16.8	---	12.2	16.6	13.5	---	18.9	25.6	19.2
Hispanic origin and race ⁵												
Hispanic or Latino	10.5	11.0	15.4	11.4	6.7	10.4	13.0	8.4	11.5	15.8	21.6	15.7
Mexican	9.7	10.6	15.6	11.3	6.5	9.7	13.5	7.7	11.3	15.8	22.0	16.9
Not Hispanic or Latino	10.7	11.4	14.5	11.1	6.3	9.0	10.9	7.5	10.5	12.8	16.6	12.0
White only	10.9	11.6	14.3	11.0	5.9	8.7	10.3	7.1	10.5	12.9	16.2	11.9
Black or African American only	10.8	11.7	17.5	13.4	9.5	12.3	15.6	10.8	10.8	14.1	20.8	14.6
Education ⁶												
No high school diploma or GED	16.2	15.8	20.6	16.6	11.5	16.0	18.1	13.3	14.5	19.7	26.3	20.6
High school diploma or GED	11.1	12.0	16.1	13.0	7.0	9.8	13.8	10.1	11.4	14.1	20.1	16.2
Some college or more	9.2	10.3	13.4	10.4	4.3	7.2	9.2	6.1	8.8	10.9	14.4	10.8
Percent of poverty level ⁷												
Below 100%	19.6	20.4	23.4	20.2	14.8	17.6	21.5	16.1	19.4	22.8	30.4	23.7
100%–199%	17.9	19.3	24.0	17.9	11.6	16.8	18.4	12.5	18.3	22.9	29.2	20.6
200%–399%	10.5	12.1	15.2	11.5	5.5	9.2	11.4	7.0	10.2	14.3	17.3	12.6
400% or more	4.6	5.2	6.8	4.3	1.7	3.5	3.9	2.5	4.5	5.7	7.0	4.5

See footnotes at end of table.

Table 63 (page 2 of 3). Delay or nonreceipt of needed medical care, nonreceipt of needed prescription drugs, or nonreceipt of needed dental care during the past 12 months due to cost, by selected characteristics: United States, selected years 1997–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#063>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Delay or nonreceipt of needed medical care due to cost ¹				Nonreceipt of needed prescription drugs due to cost ²				Nonreceipt of needed dental care due to cost ³			
	1997	2004	2010	2014	1997	2004	2010	2014	1997	2004	2010	2014
Hispanic origin and race and percent of poverty level ^{5,7}												
Percent												
Hispanic or Latino:												
Below 100%	14.6	14.0	19.0	16.2	10.6	13.2	18.9	13.1	16.1	19.6	30.5	22.6
100%–199%	12.2	14.4	18.6	13.2	8.1	12.5	14.7	9.8	13.5	19.4	25.2	19.8
200%–399%	8.0	9.2	13.9	9.6	4.4	9.7	11.5	6.1	9.2	13.7	18.1	11.8
400% or more	5.1	4.6	7.7	5.1	*	4.2	4.6	*3.1	4.5	8.2	9.1	5.7
Not Hispanic or Latino:												
White only:												
Below 100%	24.3	25.3	26.1	24.7	17.3	19.8	24.6	17.9	23.4	25.2	31.8	25.3
100%–199%	20.9	23.0	27.6	21.3	12.4	19.1	19.9	14.6	20.6	26.1	31.7	22.8
200%–399%	11.4	13.3	16.0	12.3	5.4	9.4	11.3	7.0	10.6	15.4	18.0	13.7
400% or more	4.6	5.3	6.9	4.3	1.7	3.4	3.8	2.5	4.5	5.7	6.9	4.5
Black or African American only:												
Below 100%	16.1	19.3	24.4	18.0	14.9	20.8	21.1	18.0	14.8	23.4	29.7	24.1
100%–199%	14.3	16.2	22.9	17.3	13.9	18.2	21.3	13.2	16.4	20.0	28.2	17.2
200%–399%	8.8	9.6	14.6	12.6	7.0	9.0	13.7	8.6	8.6	11.2	16.1	11.4
400% or more	4.6	4.6	8.1	5.0	*2.9	*3.9	5.6	*	4.3	4.8	9.1	*4.3
Health insurance status at the time of interview ⁸												
Insured	6.8	6.9	9.1	7.5	3.7	5.9	7.3	5.7	7.2	8.7	11.8	9.6
Private	6.0	6.2	8.2	6.4	2.9	4.8	6.0	4.1	6.2	7.3	9.2	6.9
Medicaid	11.9	11.9	12.5	11.6	11.1	13.2	13.5	11.6	14.8	18.9	24.2	20.3
Uninsured	27.6	30.2	34.5	30.4	18.0	22.9	25.7	17.6	26.1	32.3	37.7	28.2
Health insurance status prior to interview ⁸												
Insured continuously all 12 months	5.5	5.7	7.6	5.7	2.8	4.9	6.2	4.4	6.0	7.7	10.5	7.8
Uninsured for any period up to 12 months	28.7	32.3	35.1	30.7	17.7	23.4	25.1	20.1	25.2	28.7	33.6	29.9
Uninsured more than 12 months	30.6	31.8	35.9	31.7	18.9	23.7	26.2	17.4	28.0	34.2	39.4	28.2
Percent of poverty level and health insurance status prior to interview ^{7,8}												
Below 100%:												
Insured continuously all 12 months	9.4	9.6	10.1	9.3	8.1	10.0	11.4	9.5	10.7	13.9	20.7	15.9
Uninsured for any period up to 12 months	31.9	36.5	36.7	37.8	25.5	28.1	35.7	27.6	31.6	33.4	39.0	37.7
Uninsured more than 12 months	32.4	33.6	38.5	35.5	21.6	26.7	31.5	23.0	29.4	34.0	42.3	31.7
100%–199%:												
Insured continuously all 12 months	9.5	9.9	12.5	9.9	6.0	10.8	11.9	8.7	11.0	14.7	19.7	14.1
Uninsured for any period up to 12 months	33.6	35.9	38.5	31.3	20.5	27.6	26.5	21.4	28.2	32.4	38.9	32.4
Uninsured more than 12 months	30.0	32.3	37.4	31.3	19.5	24.0	26.1	17.2	29.3	35.7	40.7	30.1
200%–399%:												
Insured continuously all 12 months	6.1	6.9	9.5	6.9	2.9	5.4	7.4	4.5	6.8	9.3	11.6	8.4
Uninsured for any period up to 12 months	27.1	31.2	33.7	30.6	14.0	22.1	23.2	18.0	21.6	27.1	32.5	28.3
Uninsured more than 12 months	31.3	30.5	32.4	28.2	17.3	22.6	23.7	13.8	26.5	34.1	36.1	26.0
400% or more:												
Insured continuously all 12 months	3.1	3.2	4.6	2.8	0.8	2.3	2.9	1.9	3.1	3.9	5.2	3.6
Uninsured for any period up to 12 months	20.8	25.9	30.7	21.2	10.7	15.9	14.0	11.5	19.3	22.3	21.6	17.7
Uninsured more than 12 months	25.5	29.5	31.8	29.5	13.5	18.3	16.3	*	23.6	30.0	34.6	*14.8

See footnotes at end of table.

Table 63 (page 3 of 3). Delay or nonreceipt of needed medical care, nonreceipt of needed prescription drugs, or nonreceipt of needed dental care during the past 12 months due to cost, by selected characteristics: United States, selected years 1997–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#063>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Delay or nonreceipt of needed medical care due to cost ¹				Nonreceipt of needed prescription drugs due to cost ²				Nonreceipt of needed dental care due to cost ³				
	1997	2004	2010	2014	1997	2004	2010	2014	1997	2004	2010	2014	
Disability measure ⁹													Percent
Any basic actions difficulty or complex activity limitation	23.3	25.2	28.9	24.4	14.8	20.5	22.6	17.8	19.8	23.7	28.8	24.9	
Any basic actions difficulty	24.2	25.5	28.9	25.9	15.3	21.1	23.3	18.3	20.1	23.7	29.2	25.2	
Any complex activity limitation	25.7	27.8	30.8	25.2	19.4	23.8	27.3	23.0	23.2	27.3	33.7	30.1	
No disability	9.0	9.8	13.2	9.8	3.4	5.5	7.0	4.1	7.5	9.9	13.1	8.5	
Geographic region													
Northeast	8.8	8.7	10.2	8.5	4.9	7.3	7.7	5.6	8.9	11.1	12.9	9.9	
Midwest	10.5	10.9	14.8	11.6	5.9	7.6	11.6	8.6	9.7	11.6	16.0	12.4	
South	11.8	12.5	16.5	12.2	7.3	11.6	13.5	8.5	10.9	14.7	19.6	12.7	
West	10.8	12.2	15.1	11.1	6.3	8.3	10.0	6.7	13.1	14.4	18.4	14.7	
Location of residence ¹⁰													
Within MSA	10.2	10.8	14.2	10.8	5.9	8.8	10.8	7.2	10.0	12.9	17.0	12.3	
Outside MSA	12.5	13.6	17.4	13.4	7.9	10.8	13.6	10.4	12.9	14.5	19.1	15.0	

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.

--- Data not available.

¹Based on persons responding to the questions, “During the past 12 months was there any time when person needed medical care but did not get it because person couldn’t afford it?” and “During the past 12 months has medical care been delayed because of worry about the cost?”

²Based on persons responding to the question, “During the past 12 months was there any time when person needed prescription medicine but didn’t get it because person couldn’t afford it?”

³Based on persons responding to the question, “During the past 12 months was there any time when person needed dental care (including checkups) but didn’t get it because person couldn’t afford it?”

⁴Includes all other races not shown separately, unknown health insurance status, unknown education level, and unknown disability status.

⁵The race groups, white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The five single-race categories plus multiple-race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group; the category 2 or more races includes persons who reported more than one racial group. Prior to 1999, data were tabulated according to the 1977 Standards with four racial groups, and the Asian only category included Native Hawaiian or Other Pacific Islander. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

⁶Estimates are for persons aged 25–64. GED is General Educational Development high school equivalency diploma. See Appendix II, Education.

⁷Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Missing family income data were imputed for 1997 and beyond. See Appendix II, Family income; Poverty; Table VI.

⁸For information on the health insurance categories, see Appendix II, Health insurance coverage.

⁹Any basic actions difficulty or complex activity limitation is defined as having one or more of the following limitations or difficulties: movement difficulty, emotional difficulty, sensory (seeing or hearing) difficulty, cognitive difficulty, self-care (activities of daily living or instrumental activities of daily living) limitation, social limitation, or work limitation. For more information, see Appendix II, Basic actions difficulty; Complex activity limitation. Starting with 2007 data, the hearing question, a component of the basic actions difficulty measure, was revised. Consequently, data prior to 2007 are not comparable with data for 2007 and beyond. For more information on the impact of the revised hearing question, see Appendix II, Hearing trouble.

¹⁰MSA is metropolitan statistical area. Starting with 2006 data, MSA status is determined using 2000 census data and the 2000 standards for defining MSAs. For data prior to 2006, see Appendix II, Metropolitan statistical area (MSA) for the applicable standards.

NOTES: Standard errors and additional data years are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey, family core, sample child, and sample adult questionnaires. See Appendix I, National Health Interview Survey (NHIS).

Table 64 (page 1 of 2). No health care visits to an office or clinic within the past 12 months among children under age 18, by selected characteristics: United States, average annual, selected years 1997–1998 through 2013–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#064>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Under 18 years			Under 6 years			6–17 years		
	1997–1998	2001–2002	2013–2014	1997–1998	2001–2002	2013–2014	1997–1998	2001–2002	2013–2014
	Percent of children without a health care visit ¹								
All children ²	12.8	12.1	8.7	5.7	6.3	5.1	16.3	14.9	10.4
Sex									
Male	12.9	12.3	9.1	4.9	6.4	5.4	16.8	15.1	10.9
Female	12.7	11.9	8.3	6.5	6.1	4.8	15.8	14.6	9.9
Race ³									
White only	12.2	11.5	8.7	5.5	6.4	5.2	15.5	13.9	10.4
Black or African American only	14.3	13.3	8.7	6.5	5.9	5.6	18.1	16.8	10.2
American Indian or Alaska Native only	13.8	*18.6	*11.6	*	*	*	*17.6	*23.0	*15.6
Asian only	16.3	15.6	9.4	*5.6	*6.8	*4.6	22.1	20.5	11.8
Native Hawaiian or Other Pacific Islander only	---	*	*	---	*	*	---	*	*
2 or more races	---	8.3	6.4	---	*3.3	*3.5	---	12.4	8.3
Hispanic origin and race ³									
Hispanic or Latino	19.3	18.8	11.6	9.7	9.6	6.4	25.3	24.0	14.4
Not Hispanic or Latino	11.6	10.6	7.8	4.8	5.4	4.7	14.9	13.0	9.2
White only	10.7	9.7	7.4	4.3	5.3	4.5	13.7	11.7	8.8
Black or African American only	14.5	13.4	8.6	6.5	6.0	5.8	18.3	16.8	9.9
Percent of poverty level ⁴									
Below 100%	17.6	17.3	10.7	8.1	9.1	8.1	23.6	21.8	12.1
100%–199%	16.2	14.8	11.0	7.2	7.4	5.3	20.8	18.7	13.9
200%–399%	11.7	11.2	8.5	4.9	5.4	4.3	14.8	13.8	10.5
400% or more	7.4	7.7	5.1	3.0	4.1	2.8	9.5	9.3	6.2
Hispanic origin and race and percent of poverty level ^{3,4}									
Hispanic or Latino:									
Below 100%	23.2	22.1	12.1	11.7	10.4	9.3	31.1	29.4	13.7
100%–199%	20.9	21.3	13.7	9.7	12.3	4.8	28.1	26.2	18.0
200%–399%	15.7	15.5	10.3	8.0	*7.3	*4.7	19.7	20.0	13.3
400% or more	7.8	9.7	5.8	*	*	*	9.3	12.5	6.9
Not Hispanic or Latino:									
White only:									
Below 100%	14.0	13.2	9.6	*5.6	*8.6	*7.1	19.7	15.6	11.0
100%–199%	14.1	11.8	9.1	6.0	*6.0	*5.6	18.0	14.8	10.9
200%–399%	10.9	10.2	8.1	4.3	4.8	4.6	13.9	12.5	9.8
400% or more	7.2	7.4	5.2	*2.8	4.2	2.6	9.1	8.6	6.3
Black or African American only:									
Below 100%	15.8	16.1	9.3	7.6	*7.8	*7.6	20.5	20.3	10.2
100%–199%	16.4	13.3	10.3	*7.7	*4.4	*5.8	20.4	17.5	12.3
200%–399%	13.3	12.2	8.5	*4.9	*6.5	*	16.7	14.6	10.1
400% or more	8.3	8.9	*2.7	*	*	*	10.7	11.5	*3.1
Health insurance status at the time of interview ⁵									
Insured	10.4	9.8	7.5	4.5	4.7	4.7	13.4	12.3	8.9
Private	10.4	9.5	7.1	4.3	4.3	3.5	13.1	11.8	8.5
Medicaid	10.1	10.3	8.0	5.0	5.5	5.9	14.4	13.3	9.3
Uninsured	28.8	31.9	27.1	14.6	21.0	13.3	34.9	36.3	32.0
Health insurance status prior to interview ⁵									
Insured continuously all 12 months	10.3	9.5	7.3	4.4	4.6	4.6	13.2	12.0	8.6
Uninsured for any period up to 12 months	15.9	17.7	14.2	7.7	10.3	7.7	20.9	21.9	17.6
Uninsured more than 12 months	34.9	41.4	34.7	19.9	30.2	*16.8	40.2	45.3	39.3

See footnotes at end of table.

Table 64 (page 2 of 2). No health care visits to an office or clinic within the past 12 months among children under age 18, by selected characteristics: United States, average annual, selected years 1997–1998 through 2013–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#064>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Under 18 years			Under 6 years			6–17 years		
	1997–1998	2001–2002	2013–2014	1997–1998	2001–2002	2013–2014	1997–1998	2001–2002	2013–2014
Percent of poverty level and health insurance status prior to interview ^{4,5}									
Percent of children without a health care visit ¹									
Below 100%:									
Insured continuously all 12 months	12.6	11.7	8.9	5.7	6.1	7.6	17.6	14.9	9.7
Uninsured for any period up to 12 months . .	19.9	21.8	15.7	*9.9	*14.4	*11.9	26.1	26.6	17.9
Uninsured more than 12 months	39.9	48.2	35.6	24.9	*28.0	*	45.2	55.7	40.6
100%–199%:									
Insured continuously all 12 months	12.6	10.9	8.4	4.8	4.2	4.4	16.7	14.5	10.5
Uninsured for any period up to 12 months . .	15.6	18.9	16.1	*8.7	*10.7	*7.8	20.2	23.2	20.2
Uninsured more than 12 months	33.7	41.3	38.6	21.3	35.4	*	37.9	43.6	42.2
200%–399%:									
Insured continuously all 12 months	10.5	10.0	7.6	4.5	4.6	4.0	13.2	12.4	9.3
Uninsured for any period up to 12 months . .	12.8	14.5	11.2	*	*7.1	*	17.2	18.7	14.1
Uninsured more than 12 months	29.9	30.8	27.4	*11.8	*24.2	*	36.5	32.9	31.9
400% or more:									
Insured continuously all 12 months	7.0	7.2	4.8	2.9	3.9	2.6	8.8	8.7	5.7
Uninsured for any period up to 12 months . .	*10.8	*11.4	*12.8	*	*	*	*15.1	*14.1	*18.7
Uninsured more than 12 months	*28.8	*38.4	*36.0	*	*	*	*37.7	*40.3	*46.5
Geographic region									
Northeast	7.0	6.0	5.7	3.1	3.9	3.5	8.9	6.9	6.8
Midwest	12.2	10.3	7.6	5.9	5.1	4.2	15.3	12.8	9.2
South	14.3	14.0	9.3	5.6	7.0	6.4	18.5	17.4	10.7
West	16.3	16.0	10.6	7.9	8.1	4.9	20.7	20.0	13.4
Location of residence ⁶									
Within MSA	12.3	11.7	8.2	5.4	6.1	4.9	15.9	14.5	9.8
Outside MSA	14.6	13.5	11.7	6.9	6.9	*6.6	17.9	16.3	14.2

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.

--- Data not available.

¹ Respondents were asked how many times a doctor or other health care professional was seen in the past 12 months at a doctor's office, clinic, or some other place. Excluded are visits to emergency rooms, hospitalizations, home visits, and telephone calls. Starting with 2000 data, dental visits were also excluded. See Appendix II, Health care contact.

² Includes all other races not shown separately and unknown health insurance status.

³ The race groups, white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The five single-race categories plus multiple-race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group; the category 2 or more races includes persons who reported more than one racial group. Prior to 1999, data were tabulated according to the 1977 Standards with four racial groups, and the Asian only category included Native Hawaiian or Other Pacific Islander. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

⁴ Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Missing family income data were imputed starting in 1997. See Appendix II, Family income; Poverty; Table VI.

⁵ Health insurance categories are mutually exclusive. Persons who reported both Medicaid and private coverage are classified as having private coverage. Starting with 1997 data, state-sponsored health plan coverage is included as Medicaid coverage. Starting with 1999 data, coverage by the Children's Health Insurance Program (CHIP) is included with Medicaid coverage. In addition to private and Medicaid, the insured category also includes military, other government, and Medicare coverage. Persons not covered by private insurance, Medicaid, CHIP, state-sponsored or other government-sponsored health plans (starting in 1997), Medicare, or military plans are considered to have no health insurance coverage. Persons with only Indian Health Service coverage are considered to have no health insurance coverage. See Appendix II, Health insurance coverage.

⁶ MSA is metropolitan statistical area. Starting with 2005–2006 data, MSA status is determined using 2000 census data and the 2000 standards for defining MSAs. For data prior to 2005, see Appendix II, Metropolitan statistical area (MSA) for the applicable standards.

NOTES: Standard errors for selected years are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey, family core and sample child questionnaires. See Appendix I, National Health Interview Survey (NHIS).

Table 65 (page 1 of 3). Health care visits to doctor offices, emergency departments, and home visits within the past 12 months, by selected characteristics: United States, selected years 1997–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#065>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Number of health care visits ¹											
	None			1–3 visits			4–9 visits			10 or more visits		
	1997	2010	2014	1997	2010	2014	1997	2010	2014	1997	2010	2014
	Percent distribution											
Total, age-adjusted ^{2,3}	16.5	15.6	15.3	46.2	45.4	50.4	23.6	25.8	22.8	13.7	13.2	11.5
Total, crude ²	16.5	15.4	14.9	46.5	45.2	49.8	23.5	26.0	23.3	13.5	13.5	11.9
Age												
Under 18 years	11.8	8.1	7.9	54.1	55.6	62.8	25.2	28.2	23.1	8.9	8.2	6.2
Under 6 years	5.0	3.7	4.3	44.9	48.9	55.6	37.0	36.8	32.4	13.0	10.6	7.7
6–17 years	15.3	10.4	9.7	58.7	59.1	66.3	19.3	23.6	18.5	6.8	6.9	5.5
18–44 years	21.7	24.2	23.2	46.7	43.9	48.7	19.0	20.6	17.9	12.6	11.3	10.2
18–24 years	22.0	25.9	23.7	46.8	43.4	48.7	20.0	21.1	19.3	11.2	9.6	8.4
25–44 years	21.6	23.6	23.1	46.7	44.1	48.6	18.7	20.5	17.4	13.0	11.9	10.8
45–64 years	16.9	14.8	15.0	42.9	42.8	46.8	24.7	26.1	24.2	15.5	16.4	14.0
45–54 years	17.9	17.6	18.1	43.9	43.5	48.8	23.4	23.9	21.2	14.8	15.0	11.9
55–64 years	15.3	11.1	11.6	41.3	41.9	44.7	26.7	28.8	27.4	16.7	18.2	16.3
65 years and over	8.9	5.3	5.6	34.7	33.8	36.9	32.5	36.7	35.6	23.8	24.2	21.9
65–74 years	9.8	6.3	6.4	36.9	36.1	39.2	31.6	35.7	35.0	21.6	21.9	19.4
75 years and over	7.7	4.1	4.4	31.8	31.0	33.7	33.8	38.0	36.4	26.6	27.0	25.5
Sex ³												
Male	21.3	20.4	19.7	47.1	46.4	51.0	20.6	22.7	20.1	11.0	10.5	9.3
Female	11.8	10.9	11.1	45.4	44.4	49.9	26.5	28.8	25.4	16.3	15.9	13.6
Race ^{3,4}												
White only	16.0	15.3	15.2	46.1	44.9	49.6	23.9	26.1	23.3	14.0	13.7	11.9
Black or African American only	16.8	15.7	14.8	46.1	47.2	52.1	23.2	24.7	22.8	13.9	12.4	10.3
American Indian or Alaska Native only	17.1	19.4	19.7	38.0	40.3	39.5	24.2	28.1	26.2	20.7	12.2	14.5
Asian only	22.8	20.4	17.3	49.1	49.9	59.7	19.7	22.1	15.8	8.3	7.6	7.2
Native Hawaiian or Other Pacific Islander only	---	*	*	---	*	*	---	*	*	---	*	*
2 or more races	---	13.9	17.0	---	42.3	47.0	---	25.2	22.2	---	18.6	13.8
Hispanic origin and race ^{3,4}												
Hispanic or Latino	24.9	23.5	21.8	42.3	43.2	47.6	20.3	22.6	20.9	12.5	10.7	9.6
Mexican	28.9	25.2	23.1	40.8	43.3	46.8	18.5	21.4	20.4	11.8	10.1	9.7
Not Hispanic or Latino	15.4	14.0	13.8	46.7	45.8	51.0	24.0	26.5	23.3	13.9	13.7	11.9
White only	14.7	13.2	13.2	46.6	45.3	50.1	24.4	27.1	24.0	14.3	14.4	12.7
Black or African American only	16.9	15.6	14.6	46.1	47.3	52.2	23.1	24.9	23.0	13.8	12.2	10.2
Percent of poverty level ^{3,5}												
Below 100%	20.6	20.4	18.9	37.8	37.5	42.5	22.7	25.1	22.9	18.9	17.0	15.7
100%–199%	20.1	20.8	19.2	43.3	42.1	45.9	21.7	23.1	22.3	14.9	13.9	12.6
200%–399%	16.4	16.2	16.2	47.2	46.3	51.3	23.6	25.4	21.8	12.8	12.1	10.6
400% or more	12.8	10.2	10.7	49.8	49.4	55.1	24.9	27.6	23.8	12.5	12.7	10.4

See footnotes at end of table.

Table 65 (page 2 of 3). Health care visits to doctor offices, emergency departments, and home visits within the past 12 months, by selected characteristics: United States, selected years 1997–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#065>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Number of health care visits ¹											
	None			1–3 visits			4–9 visits			10 or more visits		
	1997	2010	2014	1997	2010	2014	1997	2010	2014	1997	2010	2014
Percent distribution												
Hispanic origin and race and percent of poverty level ^{3,4,5}												
Hispanic or Latino:												
Below 100%	30.2	28.7	25.4	34.8	36.5	41.1	19.9	22.5	21.2	15.0	12.3	12.3
100%–199%	28.7	27.7	26.2	39.7	42.7	44.7	20.4	19.9	19.7	11.2	9.8	9.4
200%–399%	20.7	21.6	20.1	47.4	45.0	51.1	19.8	23.1	19.9	12.1	10.3	8.9
400% or more	15.2	11.3	11.5	50.4	51.1	56.6	22.6	26.1	23.6	11.8	11.5	8.3
Not Hispanic or Latino:												
White only:												
Below 100%	17.0	15.0	15.5	38.3	37.0	40.8	23.9	27.4	24.1	20.9	20.6	19.7
100%–199%	17.3	18.4	15.7	44.1	40.4	45.8	22.2	24.7	23.4	16.3	16.5	15.2
200%–399%	15.4	14.7	15.2	46.9	46.0	50.0	24.3	26.3	23.1	13.4	13.0	11.7
400% or more	12.5	9.9	10.4	49.1	48.2	53.6	25.5	28.4	24.7	13.0	13.5	11.3
Black or African American only:												
Below 100%	17.4	18.4	16.3	38.5	39.8	46.0	23.4	25.0	24.2	20.7	16.8	13.6
100%–199%	18.8	17.6	17.1	43.7	45.7	45.3	22.9	24.3	25.2	14.5	12.5	12.4
200%–399%	16.6	15.1	14.9	49.7	49.0	57.5	22.9	25.7	19.4	10.8	10.2	8.1
400% or more	14.0	10.0	8.8	54.3	58.2	59.8	22.7	22.5	23.8	9.0	9.3	7.7
Health insurance status at the time of interview ^{6,7}												
Under 65 years:												
Insured	14.3	12.3	13.0	49.0	48.5	54.0	23.6	26.1	22.2	13.1	13.1	10.8
Private	14.7	12.4	13.3	50.6	51.0	56.5	23.1	25.5	21.2	11.6	11.1	9.0
Medicaid	9.8	10.9	11.9	35.5	38.2	43.8	26.5	28.0	25.8	28.2	23.0	18.4
Uninsured	33.7	37.2	39.0	42.8	42.2	44.3	15.3	15.2	12.3	8.2	5.4	4.4
Health insurance status prior to interview ^{6,7}												
Under 65 years:												
Insured continuously all 12 months	14.1	12.1	12.6	49.2	48.6	54.3	23.6	26.2	22.3	13.0	13.0	10.8
Uninsured for any period up to 12 months	18.9	18.5	20.8	46.0	47.8	50.7	20.8	22.0	19.3	14.4	11.6	9.3
Uninsured more than 12 months	39.0	43.8	44.2	41.4	39.7	42.1	13.2	12.6	10.6	6.4	3.9	3.0
Percent of poverty level and health insurance status prior to interview ^{5,6,7}												
Under 65 years:												
Below 100%:												
Insured continuously all 12 months	13.8	12.7	11.8	39.7	39.5	46.0	25.2	27.5	24.1	21.4	20.3	18.1
Uninsured for any period up to 12 months	19.7	16.9	22.5	37.6	43.0	46.7	21.9	25.0	19.2	20.9	15.1	11.5
Uninsured more than 12 months	41.2	45.0	44.2	39.9	38.1	39.9	12.2	13.6	12.4	6.6	3.3	3.5
100%–199%:												
Insured continuously all 12 months	16.0	14.8	13.9	46.4	44.4	49.4	21.9	24.8	23.0	15.8	16.0	13.7
Uninsured for any period up to 12 months	18.8	21.0	19.3	45.1	46.0	48.6	21.0	20.6	22.9	15.0	12.4	9.1
Uninsured more than 12 months	38.7	43.2	47.0	41.0	39.4	40.4	14.0	12.4	9.6	6.3	5.0	3.0
200%–399%:												
Insured continuously all 12 months	15.1	13.6	14.3	49.4	49.4	54.6	23.4	25.3	21.2	12.1	11.7	9.9
Uninsured for any period up to 12 months	17.9	18.8	21.3	49.3	49.7	53.0	20.0	19.7	16.6	12.8	11.8	9.1
Uninsured more than 12 months	37.0	43.8	42.4	43.8	40.7	46.3	12.6	13.3	8.6	6.6	*2.2	2.7
400% or more:												
Insured continuously all 12 months	12.4	9.7	10.6	52.2	51.8	57.9	23.9	26.8	22.3	11.5	11.6	9.2
Uninsured for any period up to 12 months	17.2	16.6	19.4	50.0	53.5	56.7	24.2	23.9	17.5	*8.5	*6.0	6.4
Uninsured more than 12 months	35.1	39.2	39.3	44.1	46.0	42.6	15.1	*8.8	*14.9	*5.7	*	3.2
Respondent-assessed health status ³												
Fair or poor	7.8	8.4	9.4	23.3	24.0	27.8	29.0	30.2	28.5	39.9	37.3	34.3
Good to excellent	17.2	16.3	15.9	48.4	47.5	52.7	23.3	25.5	22.3	11.1	10.7	9.1

See footnotes at end of table.

Table 65 (page 3 of 3). Health care visits to doctor offices, emergency departments, and home visits within the past 12 months, by selected characteristics: United States, selected years 1997–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#065>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Number of health care visits ¹											
	None			1–3 visits			4–9 visits			10 or more visits		
	1997	2010	2014	1997	2010	2014	1997	2010	2014	1997	2010	2014
Disability measure among adults 18 years of age and over ^{3,8}												
Percent distribution												
Any basic actions difficulty or complex activity limitation	11.1	11.5	12.1	32.0	30.9	32.3	27.9	29.3	28.4	29.1	28.3	27.2
Any basic actions difficulty	11.1	11.5	11.9	31.9	30.3	32.1	27.5	29.2	28.9	29.4	29.0	27.0
Any complex activity limitation	7.1	6.9	7.5	23.7	23.0	22.4	27.5	29.1	27.0	41.7	41.0	43.0
No disability	20.9	20.5	19.9	49.6	47.5	52.5	20.8	23.4	20.5	8.7	8.5	7.1
Geographic region ³												
Northeast	13.2	12.6	13.3	45.9	46.3	51.6	26.0	26.4	23.1	14.9	14.7	12.0
Midwest	15.9	13.4	13.6	47.7	46.8	50.9	22.8	26.4	23.3	13.6	13.3	12.2
South	17.2	16.1	16.0	46.1	44.2	49.7	23.3	26.6	23.1	13.5	13.2	11.2
West	19.1	19.1	17.2	44.8	45.2	50.5	22.8	23.5	21.4	13.3	12.2	10.8
Location of residence ^{3,9}												
Within MSA	16.2	15.6	15.3	46.4	45.8	51.0	23.7	25.6	22.5	13.7	13.0	11.3
Outside MSA	17.3	15.9	15.3	45.4	42.7	47.2	23.3	27.0	24.6	13.9	14.4	12.9

--- Data not available.

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.

¹This table presents a summary measure of the number of visits to hospital emergency departments, home visits by a nurse or other health care professional, and visits to doctor offices, clinics, or some other place during a 12-month period. See Appendix II, Emergency department or emergency room visit; Health care contact; Home visit.

²Includes all other races not shown separately, unknown health insurance status, and unknown disability status.

³Estimates are age-adjusted to the year 2000 standard population using six age groups: Under 18 years, 18–44 years, 45–54 years, 55–64 years, 65–74 years, and 75 years and over. The disability measure is age-adjusted using the five adult age groups. See Appendix II, Age adjustment.

⁴The race groups, white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity and are not strictly comparable with estimates for earlier years. The five single-race categories plus multiple-race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group; the category 2 or more races includes persons who reported more than one racial group. Prior to 1999, data were tabulated according to the 1977 Standards with four racial groups, and the Asian only category included Native Hawaiian or Other Pacific Islander. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

⁵Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Missing family income data were imputed for 1997 and beyond. See Appendix II, Family income; Poverty; Table VI.

⁶Estimates for persons under age 65 are age-adjusted to the year 2000 standard population using four age groups: Under 18 years, 18–44 years, 45–54 years, and 55–64 years. See Appendix II, Age adjustment.

⁷Health insurance categories are mutually exclusive. Persons who reported both Medicaid and private coverage are classified as having private coverage. Starting with 1997 data, state-sponsored health plan coverage is included as Medicaid coverage. Starting with 1999 data, coverage by the Children's Health Insurance Program (CHIP) is included with Medicaid coverage. In addition to private and Medicaid, the insured category also includes military plans, other government-sponsored health plans, and Medicare, not shown separately. Persons not covered by private insurance, Medicaid, CHIP, state-sponsored or other government-sponsored health plans (starting in 1997), Medicare, or military plans are considered to have no health insurance coverage. Persons with only Indian Health Service coverage are considered to have no health insurance coverage. See Appendix II, Health insurance coverage.

⁸Any basic actions difficulty or complex activity limitation is defined as having one or more of the following limitations or difficulties: movement difficulty, emotional difficulty, sensory (seeing or hearing) difficulty, cognitive difficulty, self-care (activities of daily living or instrumental activities of daily living) limitation, social limitation, or work limitation. For more information, see Appendix II, Basic actions difficulty; Complex activity limitation. Starting with 2007 data, the hearing question, a component of the basic actions difficulty measure, was revised. Consequently, data prior to 2007 are not comparable with data for 2007 and beyond. For more information on the impact of the revised hearing question, see Appendix II, Hearing trouble.

⁹MSA is metropolitan statistical area. Starting with 2006 data, MSA status is determined using 2000 census data and the 2000 standards for defining MSAs. For data prior to 2006, see Appendix II, Metropolitan statistical area (MSA) for the applicable standards.

NOTES: Standard errors are available in the spreadsheet version of this table. See <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey, family core and sample adult questionnaires. See Appendix I, National Health Interview Survey (NHIS).

Table 66 (page 1 of 3). Vaccination coverage for selected diseases among children aged 19–35 months, by race, Hispanic origin, poverty level, and location of residence in metropolitan statistical area: United States, selected years 1998–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#066>.

[Data are based on telephone interviews of a sample of the civilian noninstitutionalized population, supplemented by a survey of interview participants' immunization providers]

Vaccination and year	Race and Hispanic origin ¹						Poverty level ²		Location of residence				
	Not Hispanic or Latino						Below poverty level	At or above poverty level	Inside MSA ³				
	All	White only	Black or African American only	American Indian or Alaska Native only	Asian only ⁴	Native Hawaiian or Other Pacific Islander only ⁴			2 or more races or Latino	Central city	Remaining area	Outside MSA ³	
Percent of children aged 19–35 months													
Combined 7-vaccine series: ⁵													
2009	44.3	45.2	39.6	*	38.6	*	40.7	45.9	41.3	45.7	44.8	44.6	42.4
2010	56.6	56.9	54.5	64.1	59.3	*	61.3	55.5	52.8	58.7	56.5	57.2	55.2
2011	68.5	68.8	63.7	65.9	70.8	*	70.9	69.5	63.6	71.6	69.5	67.9	67.4
2012	68.4	69.3	64.8	*	71.6	*	71.5	67.8	63.4	71.6	67.6	69.4	68.0
2013	70.4	72.1	65.0	70.1	72.7	*	71.8	69.3	64.4	73.8	68.8	72.5	69.1
2014	71.6	72.6	65.4	*	69.5	*	68.5	74.3	65.7	75.4	70.8	72.7	71.2
DTP/DT/DTaP (4 doses or more): ⁶													
2000	81.7	84.4	76.1	77.8	84.5	*	81.5	78.6	76.2	83.5	79.9	82.8	82.9
2005	85.7	87.1	84.0	*	88.8	*	86.3	83.6	81.8	87.4	84.8	87.0	84.7
2006	85.2	86.6	81.2	82.7	86.0	86.2	83.8	84.5	81.0	86.8	84.4	86.3	84.5
2007	84.5	85.3	82.3	86.4	87.5	*	84.2	83.8	81.1	85.9	84.5	85.1	82.9
2008	84.6	85.0	80.1	82.0	92.3	*	87.6	84.9	79.9	86.8	85.0	85.3	81.8
2009	83.9	85.8	78.6	82.1	86.6	93.1	81.8	82.9	80.1	85.7	83.8	84.2	84.2
2010	84.4	84.5	83.7	81.8	88.3	*	82.8	84.4	80.8	86.1	84.0	85.0	83.7
2011	84.6	85.0	81.3	72.7	92.0	93.0	87.1	84.1	81.0	86.8	86.1	83.8	82.2
2012	82.5	83.6	79.6	88.2	88.1	*	85.6	80.8	78.5	85.0	82.4	83.4	80.5
2013	83.1	85.3	74.7	78.1	89.0	*	83.1	82.3	77.8	86.0	81.8	84.7	82.4
2014	84.2	85.5	79.1	*	87.4	*	79.6	85.4	79.1	87.4	83.6	85.3	83.1
Polio (3 doses or more):													
2000	89.5	90.6	86.6	90.8	92.7	91.2	91.2	87.9	86.9	89.9	88.1	90.1	91.1
2005	91.7	91.4	91.0	*	92.9	*	93.8	92.3	89.7	92.4	90.6	92.6	92.2
2006	92.8	93.3	90.4	91.0	92.4	96.0	91.7	93.3	92.1	93.1	92.6	93.1	93.2
2007	92.6	92.6	91.1	94.8	95.0	87.4	92.3	93.0	91.9	92.8	92.0	92.7	93.8
2008	93.6	93.6	91.5	90.6	96.5	*	94.3	94.3	91.8	94.4	93.7	94.0	92.5
2009	92.8	93.3	90.9	92.2	94.0	97.3	92.8	92.5	92.0	93.3	93.5	92.1	92.1
2010	93.3	93.2	94.0	94.6	92.8	95.1	90.2	93.8	92.4	93.6	92.7	94.1	93.1
2011	93.9	93.9	93.9	88.1	96.5	96.6	93.5	93.8	93.6	94.2	94.3	93.4	94.2
2012	92.8	93.0	92.9	95.2	92.3	*	93.3	92.5	91.8	93.4	92.6	92.9	92.8
2013	92.7	93.7	91.2	92.2	95.5	*	90.8	91.6	89.2	94.4	91.9	93.2	93.4
2014	93.3	93.3	92.0	93.8	93.2	93.8	94.0	93.8	92.0	94.5	92.7	94.2	92.7
Measles, Mumps, Rubella:													
2000	90.5	91.6	87.7	89.4	89.3	94.5	88.1	90.0	88.9	90.9	89.7	91.0	90.8
2005	91.5	91.4	91.9	89.7	91.9	90.3	93.7	91.1	89.3	92.1	91.6	91.8	90.4
2006	92.3	92.8	90.9	89.3	94.7	94.3	91.0	92.0	91.1	93.1	92.5	92.5	91.5
2007	92.3	92.1	91.5	96.2	93.9	87.6	94.6	92.6	91.3	92.6	91.8	92.8	92.3
2008	92.1	91.3	92.0	95.8	94.7	97.0	94.0	92.8	92.3	92.0	92.6	92.3	90.4
2009	90.0	90.8	88.2	94.9	90.7	96.9	88.5	89.3	88.8	90.6	91.1	88.6	88.6
2010	91.5	90.6	92.1	93.4	91.7	96.9	89.7	92.9	91.3	91.4	92.4	90.5	91.4
2011	91.6	91.1	90.8	94.8	93.9	98.7	91.1	92.4	91.3	91.7	92.0	91.2	91.5
2012	90.8	90.9	90.9	92.0	89.8	*	92.3	90.7	89.9	91.4	90.1	91.0	92.4
2013	91.9	91.5	90.9	96.3	96.7	90.4	91.5	92.1	90.5	92.5	91.5	92.4	91.3
2014	91.5	91.2	90.3	96.5	95.7	95.7	90.5	91.9	89.5	92.8	91.9	91.2	91.2
Hib (full series): ⁷													
2009	54.8	55.3	51.2	*	54.6	*	53.7	55.4	51.4	56.5	55.5	54.9	53.0
2010	66.8	67.5	65.4	77.1	69.5	*	70.1	64.8	61.3	69.7	66.5	68.4	63.4
2011	80.4	81.0	74.6	73.7	83.5	*	82.0	81.6	75.5	83.4	81.4	80.3	77.8
2012	80.9	82.2	77.5	84.7	86.1	*	82.5	79.5	76.4	84.0	80.5	81.8	79.9
2013	82.0	84.2	74.9	82.9	82.0	*	84.9	80.9	75.8	85.3	80.6	84.3	79.7
2014	82.0	83.8	75.2	83.8	83.1	*	78.7	82.8	76.3	85.5	81.4	82.7	81.6

See footnotes at end of table.

Table 66 (page 2 of 3). Vaccination coverage for selected diseases among children aged 19–35 months, by race, Hispanic origin, poverty level, and location of residence in metropolitan statistical area: United States, selected years 1998–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#066>.

[Data are based on telephone interviews of a sample of the civilian noninstitutionalized population, supplemented by a survey of interview participants' immunization providers]

Vaccination and year	Race and Hispanic origin ¹							Poverty level ²		Location of residence			
	Not Hispanic or Latino							2 or more races or Latino	Below poverty level	At or above poverty level	Inside MSA ³		
	All	White only	Black or African American only	American Indian or Alaska Native only	Asian only ⁴	Native Hawaiian or Other Pacific Islander only ⁴	Central city				Remaining area	Outside MSA ³	
Percent of children aged 19–35 months													
Hepatitis A (2 doses or more):													
2008	40.4	---	---	---	---	---	---	---	---	---	---	---	---
2009	46.6	46.2	41.3	33.2	50.9	*	47.8	49.3	47.3	46.2	48.2	46.9	42.0
2010	49.7	45.8	48.6	*	50.8	*	49.8	57.0	51.0	49.1	52.4	48.8	45.1
2011	52.2	50.0	50.9	*	56.9	*	50.2	56.3	50.7	53.4	55.0	50.9	47.6
2012	53.0	52.6	52.0	*	57.5	*	49.4	54.4	49.4	55.4	54.7	53.0	48.2
2013	54.7	53.4	49.1	*	67.3	*	57.8	56.6	53.5	56.1	55.5	55.2	50.1
2014	57.5	55.4	56.7	*	67.7	*	53.7	61.6	54.0	59.2	58.9	58.1	51.2
Hepatitis B (3 doses or more):													
2000	90.3	91.4	88.8	91.9	89.5	93.1	92.6	88.2	87.3	91.4	89.4	90.3	92.3
2005	92.9	93.1	92.7	90.1	92.7	*	94.4	92.7	91.4	93.5	91.8	93.9	93.4
2006	93.3	93.8	91.5	95.1	91.5	97.0	91.7	93.6	92.9	93.5	92.9	94.0	92.9
2007	92.7	92.5	91.2	96.7	93.8	*	92.1	93.6	92.1	92.9	92.2	92.8	93.5
2008	93.5	93.4	92.1	91.5	97.5	*	94.9	93.7	91.4	94.4	93.4	94.1	92.6
2009	92.4	92.3	91.6	92.5	93.1	96.2	93.3	92.6	92.3	92.7	92.8	91.8	91.8
2010	91.8	91.4	92.1	97.2	91.7	96.7	89.9	92.5	91.5	92.0	91.2	92.0	92.7
2011	91.1	90.3	92.1	92.6	95.5	91.1	90.7	91.5	91.8	91.2	91.0	90.7	92.5
2012	89.7	89.3	89.7	94.0	93.2	*	92.2	89.4	89.4	89.8	89.5	89.6	90.7
2013	90.8	91.0	91.1	96.1	92.0	94.9	90.7	89.7	88.3	92.0	89.6	91.8	91.4
2014	91.6	90.7	92.3	98.5	92.9	95.2	92.9	91.9	91.3	92.0	90.5	92.5	91.9
Varicella:⁸													
1998	43.2	41.9	42.4	28.0	52.6	---	---	46.9	40.5	44.1	45.1	45.2	34.3
2000	67.8	66.3	67.6	65.8	76.3	*	69.7	70.2	63.5	69.2	69.0	69.8	60.2
2005	87.9	86.1	90.6	82.2	91.9	*	90.1	89.2	87.3	87.7	88.4	88.2	85.7
2006	89.2	88.7	89.1	85.4	92.7	90.4	91.2	89.6	88.6	90.0	90.0	90.0	86.1
2007	90.0	89.2	89.8	94.9	93.7	88.6	91.6	90.6	89.2	90.1	90.1	90.4	88.7
2008	90.7	89.8	90.4	93.8	94.2	92.3	90.9	91.8	90.1	91.1	91.9	90.4	88.4
2009	89.6	89.2	88.2	89.2	89.5	97.5	90.6	90.7	89.0	90.2	90.6	88.5	88.5
2010	90.4	88.9	91.5	95.7	92.5	92.7	88.9	92.3	89.6	90.6	90.8	90.1	90.0
2011	90.8	89.6	91.2	90.1	93.5	99.0	91.9	92.0	90.2	90.9	90.9	91.0	89.8
2012	90.2	89.8	90.4	92.5	91.9	*	90.9	90.9	89.7	90.6	90.1	90.0	91.3
2013	91.2	90.0	92.1	95.4	96.0	88.7	91.0	92.0	90.3	91.6	91.1	91.6	90.3
2014	91.0	90.3	90.1	95.7	95.3	94.9	90.0	92.1	89.9	91.9	91.4	91.1	89.8
PCV (4 doses or more):⁹													
2005	53.7	57.3	46.2	*	56.2	*	54.2	50.5	44.6	57.1	51.7	57.7	48.4
2006	68.4	70.9	60.5	62.7	64.8	*	71.3	67.4	61.8	71.1	68.9	70.5	61.8
2007	75.3	76.6	70.3	80.4	75.0	*	74.1	75.4	72.8	76.3	74.8	77.3	71.0
2008	80.1	81.4	76.4	70.6	82.3	*	85.4	78.6	74.2	82.8	80.7	81.4	75.1
2009	80.4	83.4	73.2	76.2	72.5	*	73.1	80.6	74.8	83.2	79.7	81.8	81.8
2010	83.3	84.2	79.7	85.3	78.9	*	83.0	83.9	78.7	85.6	82.6	84.3	82.6
2011	84.4	85.3	81.3	75.3	84.9	93.1	84.0	84.6	80.6	86.9	85.0	84.6	82.3
2012	81.9	83.5	77.1	*	80.7	*	84.1	82.1	76.7	85.3	80.4	84.0	80.8
2013	82.0	84.1	76.1	79.0	85.6	*	83.0	80.4	74.5	86.1	80.7	84.1	79.9
2014	82.9	84.5	78.0	*	80.9	93.1	82.1	83.2	76.9	86.9	81.4	84.5	82.9
Rotavirus vaccine:¹⁰													
2009	43.9	46.4	38.0	*	41.7	*	38.4	43.7	37.7	47.1	44.6	46.6	35.6
2010	59.2	60.2	52.7	*	62.6	*	57.7	60.5	51.5	62.9	59.2	62.2	51.6
2011	67.3	68.3	62.5	57.7	66.9	*	67.8	68.3	61.1	71.1	68.9	67.4	62.7
2012	68.6	70.5	60.4	*	69.9	*	69.3	70.0	63.0	72.5	68.8	70.5	62.5
2013	72.6	74.8	62.1	*	74.9	*	72.8	73.7	64.3	76.9	72.4	74.7	66.7
2014	71.7	74.8	61.6	*	72.4	*	73.9	71.3	62.8	76.9	71.2	73.2	68.4

See footnotes at end of table.

Table 66 (page 3 of 3). Vaccination coverage for selected diseases among children aged 19–35 months, by race, Hispanic origin, poverty level, and location of residence in metropolitan statistical area: United States, selected years 1998–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#066>.

[Data are based on telephone interviews of a sample of the civilian noninstitutionalized population, supplemented by a survey of interview participants' immunization providers]

Vaccination and year	Not Hispanic or Latino					
	White only		Black or African American only		Hispanic or Latino only	
	Below poverty level ²	At or above poverty level ²	Below poverty level ²	At or above poverty level ²	Below poverty level ²	At or above poverty level ²
	Percent of children aged 19–35 months					
Combined 7-vaccine series: ⁵						
2009	43.2	45.6	37.8	43.5	43.5	48.5
2010	48.7	59.0	53.4	56.3	55.0	55.2
2011	59.8	71.8	61.0	68.0	67.9	71.1
2012	58.2	72.1	62.7	68.5	68.1	68.3
2013	61.3	74.9	60.4	69.1	68.6	70.2
2014	61.2	75.4	61.5	71.0	71.8	79.4

--- Data not available.

* Estimates are considered unreliable. For data prior to 2007, percents not shown if the unweighted sample size for the numerator was less than 30, or the confidence interval half-width divided by the estimate was greater than 50%, or the confidence interval half-width was greater than 10. Starting with 2007 data, percents not shown if the unweighted sample size for the denominator was less than 30, or the confidence interval half-width divided by the estimate was greater than 60%, or the confidence interval half-width was greater than 10.

¹Persons of Hispanic origin may be of any race. Starting with 2000 data, estimates were tabulated using the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity*. Estimates for earlier years were tabulated using the 1977 Standards on Race and Ethnicity. See Appendix II, Hispanic origin; Race.

²Poverty level is based on family income and family size using U.S. Census Bureau poverty thresholds. In 2014, 3.3% of the 14,893 children with provider-reported vaccination history data, 4.7% of Hispanic, 2.4% of non-Hispanic white, and 5.3% of non-Hispanic black children, were missing information about poverty level and were omitted from the estimates of vaccination coverage by poverty level (unweighted percentages). See Appendix II, Poverty. See Appendix I, National Immunization Survey (NIS).

³MSA is metropolitan statistical area. See Appendix II, Metropolitan statistical area (MSA).

⁴Prior to data year 2000, the category Asian included Native Hawaiian and Other Pacific Islander.

⁵The combined 7-vaccine series consists of 4 or more doses of diphtheria and tetanus toxoids and pertussis vaccine (DTP), diphtheria and tetanus toxoids vaccine (DT), or diphtheria and tetanus toxoids and acellular pertussis vaccine (DTaP); 3 or more doses of any poliovirus vaccine; 1 or more doses of a measles-containing vaccine (MCV); 3 or more doses or 4 or more doses of *Haemophilus influenzae* type b vaccine (Hib) depending on Hib vaccine product type (full series Hib); 3 or more doses of hepatitis B vaccine; 1 or more doses of varicella vaccine; and 4 or more doses of pneumococcal conjugate vaccine (PCV). The vaccine shortage that ended in September 2004 might have reduced coverage with the fourth dose of PCV among children in the 2007 National Immunization Survey (NIS) cohort. Also see footnote 7 for additional information on (Hib) vaccination.

⁶Diphtheria and tetanus toxoids and pertussis vaccine (DTP), diphtheria and tetanus toxoids (DT), and diphtheria and tetanus toxoids vaccine and acellular pertussis vaccine (DTaP).

⁷*Haemophilus influenzae* type b vaccine (Hib) full series includes primary series plus the booster dose. Before January 2009, NIS did not distinguish between Hib vaccine product types; therefore, children who received 3 doses of a vaccine product that requires 4 doses were misclassified as fully vaccinated. In addition, there was a Hib vaccine shortage during December 2007–September 2009. For more information, see Changes in measurement of *Haemophilus influenzae* serotype b (Hib) vaccination coverage—National Immunization Survey, United States, 2009. MMWR 59(33);1069–72. Available from: http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5933a3.htm?s_cid=mm5933a3_e%0d%0a.

⁸Recommended in 1996. Data collection for varicella began in July 1996.

⁹PCV is pneumococcal conjugate vaccine. Recommended in 2000. Data collection for PCV began in July 2001. Data for 4 doses of PCV are not available prior to 2005.

¹⁰Rotavirus vaccine includes 2 or more or 3 or more doses, depending on the product type received. Recommended in 2006. Data collection for rotavirus began in 2009.

NOTES: Final estimates from the National Immunization Survey include an adjustment for children with missing immunization provider data. Additional information on childhood immunizations is available from: <http://www.cdc.gov/vaccines/schedules/index.html>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>. Data have been revised and differ from previous editions of *Health, United States*.

SOURCE: CDC/NCHS and National Center for Immunization and Respiratory Diseases, National Immunization Survey. Available from: <http://www.cdc.gov/vaccines/imz-managers/coverage/imz-coverage.html> and <http://www.cdc.gov/nchs/nis.htm>. See Appendix I, National Immunization Survey (NIS).

Table 67. Vaccination coverage for selected diseases among adolescents aged 13–17, by selected characteristics: United States, 2008–2014

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2015.htm#067>.

[Data are based on telephone interviews of a sample of the civilian noninstitutionalized population, supplemented by a survey of interview participants' immunization providers]

Vaccination coverage	2008	2009	2010	2011	2012	2013 ¹	2014 ¹			
Percent of adolescents aged 13–17										
Measles, mumps, rubella (2 doses or more) . . .	89.3	89.1	90.5	91.1	91.4	89.6	90.7			
Hepatitis B (3 doses or more)	87.9	89.9	91.6	92.3	92.8	91.3	91.4			
History of varicella or received varicella vaccine (2 doses or more) ²	73.5	75.7	76.8	79.9	82.6	82.7	85.0			
Tdap (1 dose or more) ³	40.8	55.6	68.7	78.2	84.6	84.7	87.6			
Meningococcal conjugate vaccine (MenACWY) (1 dose or more) ⁴	41.8	53.6	62.7	70.5	74.0	76.6	79.3			
Human papillomavirus (HPV) (3 doses or more among females) ⁵	17.9	26.7	32.0	34.8	33.4	36.8	39.7			
Human papillomavirus (HPV) (3 doses or more among males) ⁵	1.3	6.8	13.4	21.6			
	<i>Race and Hispanic origin⁶</i>				<i>Poverty level⁷</i>		<i>Location of residence</i>			
	<i>Not Hispanic or Latino</i>							<i>Inside MSA⁸</i>		
	<i>White only</i>	<i>Black or African American only</i>	<i>American Indian or Alaska Native only</i>	<i>Asian only</i>	<i>Hispanic or Latino</i>	<i>Below poverty level</i>	<i>At or above poverty level</i>	<i>Central city</i>	<i>Remaining area</i>	<i>Outside MSA⁸</i>
	Percent of adolescents aged 13–17									
Measles, mumps, rubella (2 doses or more) . . .	91.0	91.1	94.1	85.8	90.5	90.5	90.8	91.1	90.1	91.2
Hepatitis B (3 doses or more)	92.2	91.4	93.9	85.5	90.5	90.3	91.9	91.7	91.0	91.9
No history of varicella disease and received varicella (2 doses or more) ²	80.0	84.6	84.7	82.3	82.5	82.7	80.8	82.4	82.1	73.0
Tdap (1 dose or more) ³	88.6	87.6	86.1	85.2	86.7	85.8	88.4	88.5	87.8	84.6
Meningococcal conjugate vaccine (MenACWY) (1 dose or more) ⁴	78.2	80.3	73.5	82.5	82.1	79.0	79.5	81.2	81.1	68.1
Human papillomavirus (HPV) (3 doses or more among females) ⁵	37.5	39.0	39.4	35.7	46.9	44.7	37.9	43.6	37.1	36.5
Human papillomavirus (HPV) (3 doses or more among males) ⁵	18.8	20.4	26.3	26.6	27.8	27.2	20.2	23.1	21.9	16.7

... Category not applicable.

¹Starting in 2014, NIS–Teen implemented a new definition of adequate provider data. Data for 2013 shown in this table were revised based on the 2014 definition. In general, 2013 NIS–Teen vaccination coverage estimates using the revised adequate provider data definition were different, and generally lower, than original 2013 NIS–Teen estimates. Thus, data for 2013 and beyond are not directly comparable with data for 2006–2012. For more information on the new criteria and its effect on coverage levels, see Appendix I, National Immunization Survey (NIS).

²Varicella is chickenpox.

³Tdap refers to tetanus toxoid-diphtheria vaccine (Td) or tetanus toxoid, reduced diphtheria toxoid, and acellular pertussis vaccine (Tdap) or tetanus-unknown type vaccine received at or after the age of 10 years.

⁴Includes persons receiving MenACWY or meningococcal-unknown type vaccine.

⁵For 2008, refers to HPV vaccine quadrivalent; for 2009 and beyond, refers to HPV vaccine quadrivalent or bivalent.

⁶Persons of Hispanic origin may be of any race. Estimates were tabulated using the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity*. Data for Native Hawaiian or Other Pacific Islander persons and persons of multiple races were not included because of small sample sizes. See Appendix II, Hispanic origin; Race.

⁷Poverty level is based on family income and family size using U.S. Census Bureau poverty thresholds. In 2014, 3.4% (unweighted) of adolescents with provider-reported vaccination data were missing information about poverty level and were not included in the estimates of vaccination coverage by poverty level. See Appendix II, Poverty.

⁸MSA is metropolitan statistical area. See Appendix II, Metropolitan statistical area (MSA).

NOTES: Vaccination coverage estimates are based on provider-verified responses from parents who live in households with telephones. Complex statistical methods are used to adjust vaccination estimates to account for adolescents whose parents refuse to participate in the survey, for adolescents who live in households without telephones, or for adolescents whose vaccination histories cannot be verified through their providers. Starting in 2011, the NIS sampling frame was expanded from a single-landline frame to dual-landline and cellular telephone sampling frames. See Appendix I, National Immunization Survey (NIS). Detailed vaccination data among adolescents, by race and Hispanic origin, percent of poverty level, and MSA were not available prior to 2008. Interpretation of vaccination data needs to take into account when specific vaccines were licensed and recommended for use among adolescents. Quadrivalent HPV vaccine was licensed by the U.S. Food and Drug Administration (FDA) in June 2006. For the initial recommendations on HPV vaccination, see: CDC. Quadrivalent human papillomavirus vaccine: Recommendations of the Advisory Committee on Immunization Practices. MMWR 2007;56(RR–02):1–24. Available from: http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5602a1.htm?s_cid=rr5602a1_e; HPV vaccine was recommended for males in October 2011. CDC. Recommendations on the use of quadrivalent human papillomavirus vaccine in males—Advisory Committee on Immunization Practices (ACIP), 2011. MMWR 2011;60(50):1705–8. Available from: <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6050a3.htm>. Meningococcal vaccine was licensed for use by the FDA in January 2005. For the initial recommendations on meningococcal vaccination, see: CDC. Prevention and control of meningococcal disease: Recommendations of the Advisory Committee on Immunization Practices. MMWR 2005;54(RR–07):1–21. Available from: <http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5407a1.htm>. Tdap vaccines were licensed by the FDA in May and June of 2005. For the initial recommendations on Tdap vaccination, see: CDC. Preventing tetanus, diphtheria, and pertussis among adolescents: Use of tetanus toxoid, reduced diphtheria toxoid and acellular pertussis vaccines. Recommendations of the Advisory Committee on Immunization Practices. MMWR 2006;55(RR–03):1–34. Available from: <http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5503a1.htm>. See Appendix I, National Immunization Survey (NIS). Additional information on the recommended schedule for adolescent vaccination is available from: <http://www.cdc.gov/vaccines/schedules/index.html>.

SOURCE: CDC/NCHS and National Center for Immunization and Respiratory Diseases, National Immunization Survey–Teen. Available from: <http://www.cdc.gov/vaccines/imz-managers/coverage/imz-coverage.html>. See Appendix I, National Immunization Survey (NIS).

Table 68 (page 1 of 2). Influenza vaccination among adults aged 18 and over, by selected characteristics: United States, selected years 1989–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#068>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	1989	1995	2000	2004	2010	2011	2012	2013	2014
Percent receiving influenza vaccination during past 12 months ¹									
18 years and over, age-adjusted ^{2,3}	9.6	23.7	28.7	29.5	35.3	37.2	36.8	39.9	41.0
18 years and over, crude ³	9.1	23.0	28.4	29.4	35.8	37.9	37.7	41.0	42.2
Age									
18–44 years	3.3	12.0	15.6	16.8	24.6	26.0	25.6	28.5	30.2
45–64 years	8.8	24.5	31.6	32.1	37.8	40.0	39.4	43.7	43.3
65 years and over	30.4	58.2	64.4	64.6	63.9	66.9	66.5	67.9	70.1
65–74 years	28.0	54.9	61.1	60.1	60.5	63.0	62.6	64.4	67.1
75 years and over	34.2	63.0	68.4	69.7	68.2	71.9	71.7	72.8	74.3
18 years and over									
Sex									
Male	8.5	21.5	26.7	27.0	31.3	34.1	33.4	37.0	37.9
Female	9.7	24.4	30.0	31.7	40.0	41.5	41.7	44.7	46.1
Race ⁴									
White only	9.6	23.7	30.1	30.9	36.9	39.0	38.7	42.2	43.4
Black or African American only	6.4	19.0	19.8	20.2	28.1	30.8	30.6	33.0	34.1
American Indian or Alaska Native only	10.9	*16.5	31.1	25.0	36.3	38.1	37.2	37.9	42.6
Asian only	4.3	20.0	27.0	26.5	38.6	39.1	40.9	43.9	45.0
Native Hawaiian or Other Pacific Islander only	---	---	*	*	*	*	*	*	*
2 or more races	---	---	25.3	28.5	28.9	30.8	30.1	34.7	34.3
Hispanic origin and race ⁴									
Hispanic or Latino	5.9	16.1	17.7	17.4	26.5	28.9	28.1	28.9	31.0
Mexican	5.2	16.0	16.6	16.9	25.1	27.6	26.8	29.2	31.2
Not Hispanic or Latino	9.4	23.7	29.8	31.2	37.3	39.4	39.4	43.1	44.2
White only	9.9	24.6	31.4	33.1	38.8	41.0	40.9	44.9	46.0
Black or African American only	6.4	19.2	19.9	20.3	28.0	30.9	30.9	33.2	34.4
Percent of poverty level ⁵									
Below 100%	8.9	20.6	23.1	22.4	25.0	28.1	29.2	30.1	32.0
100%–199%	11.5	23.4	28.1	28.3	31.3	32.7	32.3	35.5	36.7
200%–399%	8.0	22.8	29.6	29.9	34.8	38.0	36.8	40.1	40.8
400% or more	9.0	24.3	29.2	31.8	42.7	44.1	44.5	48.6	49.7
Hispanic origin and race and percent of poverty level ^{4,5}									
Hispanic or Latino:									
Below 100%	5.1	13.3	14.5	14.7	21.9	25.3	25.4	25.8	26.8
100%–199%	7.5	17.8	15.8	15.8	23.3	27.2	25.7	25.4	29.6
200%–399%	6.3	15.7	19.3	17.6	27.5	29.3	28.6	29.4	30.8
400% or more	6.0	19.8	22.2	23.7	36.4	36.0	35.5	37.6	39.8
Not Hispanic or Latino:									
White only:									
Below 100%	10.6	23.9	27.8	27.0	26.5	29.3	31.3	32.4	35.1
100%–199%	13.2	25.6	33.1	34.7	35.0	34.9	35.8	40.7	41.2
200%–399%	8.3	23.8	32.5	33.7	37.5	40.7	38.9	42.9	44.0
400% or more	9.4	25.5	30.5	33.3	43.8	45.9	46.2	50.4	51.3
Black or African American only:									
Below 100%	7.5	19.8	20.0	20.0	24.0	27.5	27.7	28.3	29.4
100%–199%	6.4	18.8	21.3	21.7	28.6	30.7	29.9	30.9	33.0
200%–399%	6.9	20.4	19.5	19.1	27.4	31.6	31.8	35.2	34.6
400% or more	5.6	15.6	19.2	20.9	32.9	33.9	34.2	38.2	41.6

See footnotes at end of table.

Table 68 (page 2 of 2). Influenza vaccination among adults aged 18 and over, by selected characteristics: United States, selected years 1989–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#068>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	1989	1995	2000	2004	2010	2011	2012	2013	2014
Disability measure ⁶ Percent receiving influenza vaccination during past 12 months ¹									
Any basic actions difficulty or complex activity limitation	---	---	40.8	42.3	44.6	47.6	47.3	50.6	51.4
Any basic actions difficulty	---	---	41.0	42.7	45.1	48.1	47.8	51.3	51.8
Any complex activity limitation	---	---	44.7	46.0	47.7	50.8	50.9	52.3	54.7
No disability	---	---	23.2	23.7	31.6	33.2	33.2	36.7	37.9
Geographic region									
Northeast	8.6	21.2	28.0	31.1	39.1	40.8	40.1	44.5	45.0
Midwest	8.8	22.7	28.5	30.8	37.6	37.9	38.4	41.4	43.6
South	9.5	24.7	28.7	28.9	35.1	38.2	37.2	40.1	41.5
West	9.4	22.3	28.1	27.3	32.6	35.4	36.0	39.4	39.7
Location of residence ⁷									
Within MSA	8.5	22.3	27.5	28.7	35.7	37.7	37.7	40.8	41.9
Outside MSA	11.4	25.7	31.8	32.4	36.1	39.2	37.9	42.2	44.1

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.
 --- Data not available.

¹Questions concerning use of influenza vaccination differed slightly on the National Health Interview Survey across the years for which data are shown. See Appendix II, Vaccination. Data prior to 1997 are not strictly comparable with data for later years due to the 1997 questionnaire redesign. See Appendix I, National Health Interview Survey (NHIS).

²Estimates are age-adjusted to the year 2000 standard population using four age groups: 18–44 years, 45–64 years, 65–74 years, and 75 years and over. See Appendix II, Age adjustment.

³Includes all other races not shown separately, unknown disability status, and unknown poverty level in 1989.

⁴The race groups, white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The five single-race categories plus multiple-race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group; the category 2 or more races includes persons who reported more than one racial group. Prior to 1999, data were tabulated according to the 1977 Standards with four racial groups, and the Asian only category included Native Hawaiian or Other Pacific Islander. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

⁵Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Poverty level was unknown for 11% of persons aged 18 and over in 1989. Missing family income data were imputed for 1991 and beyond. See Appendix II, Family income; Poverty; Table VI.

⁶Any basic actions difficulty or complex activity limitation is defined as having one or more of the following limitations or difficulties: movement difficulty, emotional difficulty, sensory (seeing or hearing) difficulty, cognitive difficulty, self-care (activities of daily living or instrumental activities of daily living) limitation, social limitation, or work limitation. For more information, see Appendix II, Basic actions difficulty; Complex activity limitation. Starting with 2007 data, the hearing question, a component of the basic actions difficulty measure, was revised. Consequently, data prior to 2007 are not comparable with data for 2007 and beyond. For more information on the impact of the revised hearing question, see Appendix II, Hearing trouble.

⁷MSA is metropolitan statistical area. Starting with 2006 data, MSA status is determined using 2000 census data and the 2000 standards for defining MSAs. For data prior to 2006, see Appendix II, Metropolitan statistical area (MSA) for the applicable standards.

NOTES: Interpretation of vaccination data needs to take into account when age-specific universal recommendations were issued. Medicare payment for the costs of the vaccine and its administration began in 1993. In 2000, CDC's Advisory Committee on Immunization Practices (ACIP) recommended universal influenza vaccination, with rare exceptions, for persons aged 50 and over. See, *Health, United States, 2014*, Table 74 for historical data for adults age 50 and over. In 2010, ACIP recommended universal influenza vaccination, with rare exceptions, for persons aged 6 months and over. For current ACIP vaccination recommendations, see:

<http://www.cdc.gov/flu/professionals/acip/index.htm>. Standard errors for selected years are available in the spreadsheet version of this table. Available from:

<http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at:

<http://www.cdc.gov/nchs/hus.htm>. Some data were revised and differ from previous editions of *Health, United States*.

SOURCE: CDC/NCHS, National Health Interview Survey. Data are from the following questionnaire supplements: immunization (1981), health promotion and disease prevention (1991), and the year 2000 objectives (1993–1995). Starting in 1997, data are from the sample adult questionnaire. See Appendix I, National Health Interview Survey (NHIS).

Table 69 (page 1 of 2). Pneumococcal vaccination among adults aged 18 and over, by selected characteristics: United States, selected years 1989–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#069>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	1989	1995	2000	2004	2010	2011	2012	2013	2014
Percent of adults ever receiving pneumococcal vaccination ¹									
18 years and over, age-adjusted ^{2,3}	4.6	12.0	15.4	16.8	19.2	20.6	19.9	19.9	20.5
18 years and over, crude ³	4.4	11.7	15.1	16.5	19.6	21.1	20.7	21.0	21.8
Age									
18–44 years	2.1	6.6	5.1	5.4	6.9	8.3	7.9	7.5	8.3
45–64 years	3.7	8.8	12.2	14.2	17.7	18.4	18.0	18.8	18.4
65 years and over	14.1	34.0	53.1	56.8	59.7	62.3	59.9	59.7	61.3
65–74 years	13.1	31.4	48.2	50.4	54.6	56.0	55.0	54.4	55.8
75 years and over	15.7	37.8	59.1	64.2	66.0	70.0	66.4	67.1	69.3
High-risk group ⁴									
Total, 18–64 years	---	---	18.3	20.9	18.3	20.0	19.9	21.0	20.2
18–44 years	---	---	11.3	11.4	9.8	12.4	11.3	11.7	11.3
45–64 years	---	---	23.3	27.0	26.7	27.3	28.0	29.4	28.3
65 years and over									
Sex									
Male	13.9	34.6	52.1	54.3	57.6	59.5	55.8	57.1	58.4
Female	14.3	33.6	53.9	58.7	61.3	64.5	63.1	61.8	63.7
Race ⁵									
White only	14.8	35.3	55.6	59.1	61.6	64.7	62.3	61.7	63.1
Black or African American only	6.4	21.9	30.6	38.6	45.5	47.5	46.0	48.4	49.2
American Indian or Alaska Native only	31.2	*	70.1	*42.0	*48.5	53.0	*36.3	52.9	57.1
Asian only	*	*23.4	40.9	35.1	47.9	40.3	41.1	45.0	47.7
Native Hawaiian or Other Pacific Islander only	---	---	*	*	*	*	*	*	*
2 or more races	---	---	55.6	*48.8	65.5	77.1	45.4	50.8	71.2
Hispanic origin and race ⁵									
Hispanic or Latino	9.8	23.2	30.4	33.7	39.0	43.1	43.4	39.2	45.2
Mexican	12.9	*18.8	32.0	33.3	41.4	47.1	45.5	47.4	47.8
Not Hispanic or Latino	14.3	34.5	54.4	58.3	61.3	63.8	61.2	61.4	62.7
White only	15.0	35.9	56.8	60.9	63.5	66.5	64.0	63.6	64.7
Black or African American only	6.2	21.8	30.6	38.6	46.2	47.6	46.1	48.7	49.8
Percent of poverty level ⁶									
Below 100%	11.2	28.7	40.6	42.5	42.6	49.6	39.5	50.5	47.3
100%–199%	15.1	30.7	51.4	56.1	57.2	60.3	59.8	58.0	59.5
200%–399%	15.1	36.1	55.8	60.5	62.2	63.4	63.6	61.7	64.5
400% or more	15.5	39.5	56.9	58.5	64.0	66.4	61.4	61.6	63.2
Hispanic origin and race and percent of poverty level ^{5,6}									
Hispanic or Latino:									
Below 100%	*	*14.1	23.8	31.8	30.2	34.8	30.9	35.3	34.1
100%–199%	*11.0	*15.6	32.3	29.0	36.9	49.3	42.0	39.1	44.4
200%–399%	*11.1	*34.4	37.6	42.7	45.8	39.2	54.5	36.1	52.1
400% or more	*	*55.1	*26.4	*	43.0	49.1	46.4	49.1	54.0
Not Hispanic or Latino:									
White only:									
Below 100%	13.3	32.5	47.9	50.6	51.1	60.3	46.5	59.1	55.4
100%–199%	16.0	33.5	56.1	61.9	61.3	64.6	66.1	63.3	64.1
200%–399%	15.7	37.1	57.6	62.9	64.9	66.9	65.9	65.2	66.9
400% or more	15.9	39.3	59.5	60.4	66.0	68.6	63.5	63.2	64.5
Black or African American only:									
Below 100%	*5.0	*22.6	28.8	27.0	34.9	39.5	36.1	48.9	46.0
100%–199%	7.8	*20.9	28.1	36.4	46.4	45.6	44.5	46.9	49.1
200%–399%	*5.9	*21.7	35.5	51.3	51.8	54.2	54.1	49.4	47.9
400% or more	*	*	*32.6	45.0	50.1	49.1	45.4	50.3	56.0

See footnotes at end of table.

Table 69 (page 2 of 2). Pneumococcal vaccination among adults aged 18 and over, by selected characteristics: United States, selected years 1989–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#069>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	1989	1995	2000	2004	2010	2011	2012	2013	2014
Percent of adults ever receiving pneumococcal vaccination ¹									
Any basic actions difficulty or complex activity limitation ⁷									
Any basic actions difficulty or complex activity limitation	---	---	56.6	61.2	63.9	67.0	65.4	64.4	66.7
Any basic actions difficulty	---	---	56.8	61.8	64.2	67.3	66.0	64.9	66.7
Any complex activity limitation	---	---	58.0	62.1	65.2	66.7	65.7	66.1	67.6
No disability	---	---	47.9	50.1	53.3	55.6	53.2	53.1	53.7
Geographic region									
Northeast	10.4	28.2	51.2	56.0	56.7	60.0	58.0	59.1	59.6
Midwest	13.7	31.0	52.6	59.5	61.2	65.6	63.8	62.3	65.4
South	14.9	35.9	51.3	57.2	60.9	63.2	59.5	59.3	60.9
West	17.9	41.1	59.7	53.7	58.9	59.5	58.2	58.3	59.3
Location of residence ⁸									
Within MSA	13.1	33.8	52.4	56.7	58.8	61.7	59.3	59.0	60.7
Outside MSA	16.9	34.7	55.4	57.3	63.3	64.6	62.4	62.8	64.0

--- Data not available.

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.

¹Questions concerning receipt of pneumococcal vaccination differed slightly on the National Health Interview Survey across the years for which data are shown. See Appendix II, Vaccination. Data prior to 1997 are not strictly comparable with data for later years due to the 1997 questionnaire redesign. See Appendix I, National Health Interview Survey (NHIS).

²Estimates are age-adjusted to the year 2000 standard population using four age groups: 18–44 years, 45–64 years, 65–74 years, and 75 years and over. See Appendix II, Age adjustment.

³Includes all other races not shown separately, unknown poverty level in 1989, and unknown disability status.

⁴High-risk group membership is based on recommendations of CDC’s Advisory Committee on Immunization Practices (ACIP). The high-risk group includes persons who reported diabetes, cancer, heart, lung, liver, or kidney disease. Starting in 2009, this definition was expanded to also include persons who reported asthma or cigarette smoking, to be consistent with the revised ACIP recommendation. Starting with data year 2012, the survey questionnaire changed and now asks respondents if a health professional had ever told them they had chronic obstructive pulmonary disease (COPD), and this information was added to the list of lung diseases used to construct the high-risk category. For more information on high-risk groups, see the 2009 ACIP recommendation. Available from: <http://www.cdc.gov/mmwr/pdf/wk/mm5934.pdf>.

⁵The race groups, white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The five single-race categories plus multiple-race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group; the category 2 or more races includes persons who reported more than one racial group. Prior to 1999, data were tabulated according to the 1977 Standards with four racial groups, and the Asian only category included Native Hawaiian or Other Pacific Islander. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

⁶Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Poverty level was unknown for 11% of persons aged 18 and over in 1989. Missing family income data were imputed for 1991 and beyond. See Appendix II, Family income; Poverty; Table VI.

⁷Any basic actions difficulty or complex activity limitation is defined as having one or more of the following limitations or difficulties: movement difficulty, emotional difficulty, sensory (seeing or hearing) difficulty, cognitive difficulty, self-care (activities of daily living or instrumental activities of daily living) limitation, social limitation, or work limitation. For more information, see Appendix II, Basic actions difficulty; Complex activity limitation. Starting with 2007 data, the hearing question, a component of the basic actions difficulty measure, was revised. Consequently, data prior to 2007 are not comparable with data for 2007 and beyond. For more information on the impact of the revised hearing question, see Appendix II, Hearing trouble.

⁸MSA is metropolitan statistical area. Starting with 2006 data, MSA status is determined using 2000 census data and the 2000 standards for defining MSAs. For data prior to 2006, see Appendix II, Metropolitan statistical area (MSA) for the applicable standards.

NOTES: In 1997, CDC’s Advisory Committee on Immunization Practices (ACIP) recommended universal pneumonia vaccination for adults aged 65 and over. A pneumococcal polysaccharide vaccine was first licensed in 1977. Medicare payment for the costs of the vaccine and its administration began in 1981. CDC. Prevention of pneumococcal disease: Recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR* 1997;46(RR-08);1–24. Available from: <http://www.cdc.gov/mmwr/preview/mmwrhtml/00047135.htm>. For more information on the adult vaccination schedule, see: <http://www.cdc.gov/vaccines/schedules/index.html>. Standard errors for selected years are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>. Some estimates have been revised and differ from previous editions of *Health, United States*.

SOURCE: CDC/NCHS, National Health Interview Survey. Data are from the following questionnaire supplements: immunization (1981), health promotion and disease prevention (1991), and the year 2000 objectives (1993–1995). Starting in 1997, data are from the sample adult questionnaire. See Appendix I, National Health Interview Survey (NHIS).

Table 70 (page 1 of 3). Use of mammography among women aged 40 and over, by selected characteristics: United States, selected years 1987–2013

Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#070>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	1987	1993	1994	2000	2003	2005	2008	2010	2013
Percent of women having a mammogram within the past 2 years ¹									
40 years and over, age-adjusted ^{2,3}	29.0	59.7	61.0	70.4	69.5	66.6	67.1	66.5	65.7
40 years and over, crude ²	28.7	59.7	60.9	70.4	69.7	66.8	67.6	67.1	66.8
50 years and over, age-adjusted ^{2,3}	27.3	59.7	60.9	73.7	72.4	68.2	70.3	68.8	69.1
50 years and over, crude ²	27.4	59.7	60.6	73.6	72.4	68.4	70.5	69.2	69.5
Age									
40–49 years	31.9	59.9	61.3	64.3	64.4	63.5	61.5	62.3	59.6
50–64 years	31.7	65.1	66.5	78.7	76.2	71.8	74.2	72.6	71.4
65 years and over	22.8	54.2	55.0	67.9	67.7	63.8	65.5	64.4	66.9
65–74 years	26.6	64.2	63.0	74.0	74.6	72.5	72.6	71.9	75.3
75 years and over	17.3	41.0	44.6	61.3	60.6	54.7	57.9	55.7	56.5
Race ⁴									
40 years and over, crude:									
White only	29.6	60.0	60.6	71.4	70.1	67.4	67.9	67.4	66.8
Black or African American only	24.0	59.1	64.3	67.8	70.4	64.9	68.0	67.9	67.1
American Indian or Alaska Native only	*	49.8	65.8	47.4	63.1	72.8	62.7	71.2	62.6
Asian only	*	55.1	55.8	53.5	57.6	54.6	66.1	62.4	66.6
Native Hawaiian or Other Pacific Islander only	---	---	---	*	*	*	*	*	*
2 or more races	---	---	---	69.2	65.3	63.7	55.2	51.4	65.4
Hispanic origin and race ⁴									
40 years and over, crude:									
Hispanic or Latina	18.3	50.9	51.9	61.2	65.0	58.8	61.2	64.2	61.4
Not Hispanic or Latina	29.4	60.3	61.5	71.1	70.1	67.5	68.3	67.4	67.5
White only	30.3	60.6	61.3	72.2	70.5	68.3	68.7	67.8	67.6
Black or African American only	23.8	59.2	64.4	67.9	70.5	65.2	68.3	67.4	67.2
Age and Hispanic origin and race ⁴									
40–49 years:									
Hispanic or Latina	*15.3	52.6	47.5	54.1	59.4	54.2	54.1	59.8	56.4
Not Hispanic or Latina:									
White only	34.3	61.6	62.0	67.2	65.2	65.5	64.1	62.6	60.3
Black or African American only	27.8	55.6	67.2	60.9	68.2	62.1	59.5	63.5	59.4
50–64 years:									
Hispanic or Latina	23.0	59.2	60.1	66.5	69.4	61.5	71.3	68.6	65.6
Not Hispanic or Latina:									
White only	33.6	66.2	67.5	80.6	77.2	73.5	74.1	73.5	72.1
Black or African American only	26.4	65.5	63.6	77.7	76.2	71.6	76.7	74.0	71.7
65 years and over:									
Hispanic or Latina	*	*35.7	48.0	68.3	69.5	63.8	59.0	65.2	63.2
Not Hispanic or Latina:									
White only	24.0	54.7	54.9	68.3	68.1	64.7	66.1	65.0	67.3
Black or African American only	14.1	56.3	61.0	65.5	65.4	60.5	66.4	60.9	68.8
Age and percent of poverty level ⁵									
40 years and over, crude:									
Below 100%	14.6	41.1	44.2	54.8	55.4	48.5	51.4	51.4	49.9
100%–199%	20.9	47.5	48.6	58.1	60.8	55.3	55.8	53.8	56.7
200%–399%	29.7	63.2	65.0	68.8	69.9	67.2	64.4	66.2	66.0
400% or more	42.9	74.1	74.1	81.5	77.7	76.6	79.0	78.1	77.2
40–49 years:									
Below 100%	18.6	36.1	43.0	47.4	50.6	42.5	46.6	48.1	43.3
100%–199%	18.4	47.8	47.6	43.6	54.0	49.8	46.5	46.2	52.0
200%–399%	31.2	63.0	64.5	60.2	63.0	61.8	56.8	59.2	58.5
400% or more	44.1	69.6	69.9	75.8	71.6	73.6	72.5	73.6	69.0
50–64 years:									
Below 100%	14.6	47.3	46.2	61.7	58.3	50.4	57.5	54.7	55.0
100%–199%	24.2	47.0	49.0	68.3	64.0	58.8	58.9	57.3	57.2
200%–399%	29.7	66.1	69.6	75.1	74.1	70.7	69.8	70.7	69.5
400% or more	44.7	78.7	78.0	86.9	84.9	80.6	84.3	82.8	80.9
65 years and over:									
Below 100%	13.1	40.4	43.9	54.8	57.0	52.3	49.1	50.6	49.8
100%–199%	19.9	47.6	48.8	60.3	62.8	56.1	59.4	55.5	59.3
200%–399%	27.7	60.3	61.0	71.1	72.3	68.6	65.0	67.2	68.1
400% or more	34.7	71.3	73.0	81.9	73.0	72.6	78.3	74.5	79.0

See footnotes at end of table.

Table 70 (page 2 of 3). Use of mammography among women aged 40 and over, by selected characteristics: United States, selected years 1987–2013

Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#070>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	1987	1993	1994	2000	2003	2005	2008	2010	2013
Health insurance status at the time of interview ⁶									
Percent of women having a mammogram within the past 2 years ¹									
40–64 years:									
Insured	---	66.2	68.3	76.0	75.1	72.5	73.4	74.1	72.1
Private	---	67.1	69.4	77.1	76.3	74.5	74.2	75.6	73.4
Medicaid	---	51.9	54.5	61.7	63.5	55.6	64.2	64.4	63.5
Uninsured	---	36.0	34.0	40.7	41.5	38.1	39.7	36.0	37.3
Health insurance status prior to interview ⁶									
40–64 years:									
Insured continuously all 12 months	---	66.6	68.6	76.8	75.6	73.1	74.1	74.7	72.7
Uninsured for any period up to 12 months	---	49.4	49.9	53.0	56.0	51.3	55.3	57.3	54.5
Uninsured more than 12 months	---	28.4	26.6	34.0	37.0	32.9	34.6	30.0	32.8
Age and education ⁷									
40 years and over, crude:									
No high school diploma or GED	17.8	46.4	48.2	57.7	58.1	52.8	53.8	53.0	53.6
High school diploma or GED	31.3	59.0	61.0	69.7	67.8	64.9	65.2	64.4	63.4
Some college or more	37.7	69.5	69.7	76.2	75.1	72.7	73.4	72.1	71.6
40–49 years:									
No high school diploma or GED	15.1	43.6	50.4	46.8	53.3	51.2	46.9	44.9	46.9
High school diploma or GED	32.6	56.6	55.8	59.0	60.8	58.8	57.2	58.4	51.8
Some college or more	39.2	66.1	68.7	70.6	68.1	68.3	66.3	66.5	64.3
50–64 years:									
No high school diploma or GED	21.2	51.4	51.6	66.5	63.4	56.9	64.9	56.7	58.2
High school diploma or GED	33.8	62.4	67.8	76.6	71.8	70.1	70.4	69.9	66.9
Some college or more	40.5	78.5	74.7	84.2	82.7	77.0	78.5	77.0	75.7
65 years and over:									
No high school diploma or GED	16.5	44.2	45.6	57.4	56.9	50.7	49.2	54.1	53.4
High school diploma or GED	25.9	57.4	59.1	71.8	69.7	64.3	65.7	62.5	66.5
Some college or more	32.3	64.8	64.3	74.1	75.1	73.0	75.6	70.9	73.6
Disability measure ⁸									
40 years and over, crude:									
Any basic actions difficulty or complex activity limitation	---	---	---	67.8	67.2	63.5	63.9	63.3	63.5
Any basic actions difficulty	---	---	---	67.9	67.3	63.5	63.9	63.3	63.8
Any complex activity limitation	---	---	---	64.1	62.3	59.9	60.2	58.2	58.4
No disability	---	---	---	72.6	71.8	69.8	71.1	70.8	69.8

See footnotes at end of table.

Table 70 (page 3 of 3). Use of mammography among women aged 40 and over, by selected characteristics: United States, selected years 1987–2013

Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#070>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.
-- - Data not available.

¹Questions concerning use of mammography differed slightly on the National Health Interview Survey across the years for which data are shown. See Appendix II, Mammography.

²Includes all other races not shown separately, unknown poverty level in 1987, unknown health insurance status, unknown education level, and unknown disability status.

³Estimates for women aged 40 and over are age-adjusted to the year 2000 standard population using four age groups: 40–49 years, 50–64 years, 65–74 years, and 75 years and over. Estimates for women 50 years of age and over are age-adjusted using three age groups. See Appendix II, Age adjustment.

⁴The race groups, white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The five single-race categories plus multiple-race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group; the category 2 or more races includes persons who reported more than one racial group. Prior to 1999, data were tabulated according to the 1977 Standards with four racial groups, and the Asian only category included Native Hawaiian or Other Pacific Islander. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

⁵Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Poverty level was unknown for 11% of women aged 40 and over in 1987. Missing family income data were imputed for 1997 and beyond. See Appendix II, Family income; Poverty; Table VI.

⁶Health insurance categories are mutually exclusive. Persons who reported both Medicaid and private coverage are classified as having private coverage. Starting with 1997 data, state-sponsored health plan coverage is included as Medicaid coverage. Starting with 1999 data, coverage by the Children's Health Insurance Program (CHIP) is included with Medicaid coverage. In addition to private and Medicaid, the insured category also includes military plans, other government-sponsored health plans, and Medicare, not shown separately. Persons not covered by private insurance, Medicaid, CHIP, public assistance (through 1996), state-sponsored or other government-sponsored health plans (starting in 1997), Medicare, or military plans are considered to have no health insurance coverage. Persons with only Indian Health Service coverage are considered to have no health insurance coverage. See Appendix II, Health insurance coverage.

⁷Education categories shown are for 1998 and subsequent years. GED is General Educational Development high school equivalency diploma. In years prior to 1998, the following categories based on number of years of school completed were used: less than 12 years, 12 years, 13 years or more. See Appendix II, Education.

⁸Any basic actions difficulty or complex activity limitation is defined as having one or more of the following limitations or difficulties: movement difficulty, emotional difficulty, sensory (seeing or hearing) difficulty, cognitive difficulty, self-care (activity of daily living or instrumental activities of daily living) limitation, social limitation, or work limitation. For more information, see Appendix II, Basic actions difficulty; Complex activity limitation. Starting with 2007 data, the hearing question, a component of the basic actions difficulty measure, was revised. Consequently, data prior to 2007 are not comparable with 2007 data and beyond. For more information on the impact of the revised hearing question, see Appendix II, Hearing trouble.

NOTES: See Appendix II, Mammography, for a discussion of the U.S. Preventive Services Task Force recommendations for mammography screening. Standard errors are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data starting in 1997 are not strictly comparable with data for earlier years due to the 1997 questionnaire redesign. See Appendix I, National Health Interview Survey (NHIS). Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey. Data are from the following supplements: cancer control (1987), health promotion and disease prevention (1990–1991), and year 2000 objectives (1993–1994). Starting in 1998, data are from the family core and sample adult questionnaires. See Appendix I, National Health Interview Survey (NHIS).

Table 71 (page 1 of 5). Use of Pap smears among women aged 18 and over, by selected characteristics: United States, selected years 1987–2013

Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#071>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	1987	1993	1999	2000	2003	2005	2008	2010	2013
Percent of women having a Pap smear within the past 3 years ¹									
18 years and over, age-adjusted ^{2,3}	74.1	77.7	80.8	81.3	79.2	77.9	75.6	73.7	70.4
18 years and over, crude ²	74.4	77.7	80.8	81.2	79.0	77.7	75.1	73.2	69.4
Age									
18–44 years	83.3	84.6	86.8	84.9	83.9	83.6	81.8	80.4	77.2
18–20 years	59.4	66.8	65.3	59.8	63.7	61.1	57.5	52.0	38.6
21–44 years	86.1	86.2	89.2	87.8	86.1	86.3	84.8	84.0	81.6
21–24 years	85.3	86.1	85.3	84.1	82.7	84.0	80.2	81.1	74.6
25–44 years	86.3	86.3	89.9	88.5	86.8	86.8	85.7	84.6	83.2
45–64 years	70.5	77.2	81.7	84.6	81.3	80.6	78.8	76.9	73.9
45–54 years	75.7	82.1	83.8	86.3	83.6	83.4	81.0	79.9	78.6
55–64 years	65.2	70.6	78.4	82.0	77.8	76.8	76.0	73.2	68.6
65 years and over	50.8	57.6	61.0	64.5	60.8	54.9	50.0	47.1	42.7
65–74 years	57.9	64.7	70.0	71.6	70.1	66.3	61.6	58.0	54.5
75 years and over	40.4	48.0	50.8	56.7	51.1	42.7	37.5	34.6	27.9
Race ⁴									
18 years and over, crude:									
White only	74.1	77.3	80.6	81.3	78.7	77.7	74.9	72.8	68.7
Black or African American only	80.7	82.7	85.7	85.1	84.0	81.1	80.1	77.9	75.3
American Indian or Alaska Native only	85.4	78.1	92.2	76.8	84.8	75.2	69.4	73.4	70.1
Asian only	51.9	68.8	64.4	66.4	68.3	64.1	65.1	68.0	65.3
Native Hawaiian or Other Pacific Islander only	---	---	*	*	*	*	*	*	*
2 or more races	---	---	86.9	80.0	81.6	86.2	77.1	70.8	70.8
Hispanic origin and race ⁴									
18 years and over, crude:									
Hispanic or Latina	67.6	77.2	76.3	77.0	75.4	75.5	75.4	73.6	70.5
Not Hispanic or Latina	74.9	77.8	81.3	81.7	79.5	78.0	75.1	73.1	69.2
White only	74.7	77.3	81.0	81.8	79.3	78.1	74.9	72.8	68.4
Black or African American only	80.9	82.7	86.0	85.1	83.8	81.2	80.0	77.4	75.1
Age, Hispanic origin, and race ⁴									
18–44 years:									
Hispanic or Latina	73.9	80.9	77.0	78.1	75.9	76.5	77.9	75.9	72.3
Not Hispanic or Latina:									
White only	84.5	85.3	88.7	86.6	85.8	85.8	83.8	82.1	79.0
Black or African American only	89.1	88.0	90.8	88.5	88.6	86.4	83.5	84.2	82.8
45–64 years:									
Hispanic or Latina	57.7	75.8	79.5	77.8	77.9	78.4	78.2	75.4	74.4
Not Hispanic or Latina:									
White only	71.2	77.2	81.9	85.9	81.4	81.4	79.0	77.2	73.6
Black or African American only	76.2	80.3	84.6	85.7	84.7	80.5	82.1	78.2	76.0
65 years and over:									
Hispanic or Latina	41.7	57.1	63.7	66.8	64.6	60.0	52.6	54.2	49.4
Not Hispanic or Latina:									
White only	51.8	57.1	60.5	64.2	60.7	54.1	49.0	46.5	41.4
Black or African American only	44.8	61.2	64.5	67.2	59.6	60.1	58.7	48.0	45.8
Age and percent of poverty level ⁵									
18 years and over, crude:									
Below 100%	64.3	70.3	73.6	72.0	70.5	68.7	68.9	65.1	60.6
100%–199%	68.2	71.2	72.5	73.4	71.4	69.0	65.0	64.3	59.8
200%–399%	77.6	80.6	80.6	80.2	78.6	77.9	72.5	71.3	68.5
400% or more	83.6	85.1	87.6	89.1	86.6	85.7	84.4	83.1	79.4
18–44 years:									
Below 100%	77.1	77.0	79.7	77.1	77.1	76.2	76.5	73.0	69.2
100%–199%	80.4	81.9	84.0	79.4	79.5	78.1	75.5	75.7	72.6
200%–399%	84.8	86.6	86.7	86.1	84.0	85.5	82.6	79.8	78.6
400% or more	88.9	91.3	91.1	89.8	89.5	88.7	87.0	88.9	84.5
45–64 years:									
Below 100%	53.6	66.5	73.1	73.6	66.0	65.9	66.2	61.7	54.9
100%–199%	60.4	64.8	70.4	76.1	71.4	69.6	65.6	63.2	61.2
200%–399%	71.0	79.5	79.9	80.0	80.8	79.3	75.3	75.2	73.7
400% or more	79.1	83.9	87.4	91.5	87.5	87.4	87.1	85.7	82.8
65 years and over:									
Below 100%	33.2	47.4	51.9	53.7	52.6	44.4	41.6	35.1	34.1
100%–199%	50.4	55.7	54.7	61.0	55.4	49.5	43.5	40.7	33.0
200%–399%	58.0	59.7	64.0	65.1	62.4	56.8	45.8	47.1	39.6
400% or more	65.2	67.5	70.4	75.4	70.2	64.6	65.7	57.7	58.1

See footnotes at end of table.

Table 71 (page 2 of 5). Use of Pap smears among women aged 18 and over, by selected characteristics: United States, selected years 1987–2013

Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#071>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	1987	1993	1999	2000	2003	2005	2008	2010	2013
Health insurance status at the time of interview ⁶									
Percent of women having a Pap smear within the past 3 years ¹									
18–64 years, crude:									
Insured	---	84.7	87.2	87.8	86.4	85.6	83.4	82.8	80.0
Private	---	84.8	87.5	88.0	87.0	86.5	84.2	84.2	81.3
Medicaid	---	82.7	84.2	85.8	82.8	80.9	80.3	78.0	75.7
Uninsured	---	69.4	73.3	70.4	66.6	67.7	67.1	61.9	57.6
Health insurance status prior to interview ⁶									
18–64 years, crude:									
Insured continuously all 12 months	---	84.8	87.3	88.0	86.6	85.8	83.7	83.2	80.4
Uninsured for any period up to 12 months	---	81.8	83.5	83.7	81.8	81.3	78.9	78.3	72.3
Uninsured more than 12 months	---	65.1	68.8	65.1	60.2	62.0	62.1	55.2	52.7
Age and education ⁷									
25 years and over, crude:									
No high school diploma or GED	57.1	61.9	66.1	69.9	64.9	64.1	60.6	56.7	56.2
High school diploma or GED	76.4	78.2	79.3	79.8	75.9	73.8	69.5	66.8	62.0
Some college or more	84.0	84.4	87.8	88.0	86.2	84.6	82.6	80.7	77.1
25–44 years:									
No high school diploma or GED	75.1	73.6	79.0	79.6	71.7	75.5	76.2	69.1	71.7
High school diploma or GED	85.6	85.4	87.6	86.2	84.3	83.1	80.0	79.0	79.5
Some college or more	90.1	89.8	93.0	91.4	90.8	90.5	89.3	89.0	86.1
45–64 years:									
No high school diploma or GED	58.0	65.6	71.6	75.7	71.4	69.7	70.4	63.4	63.0
High school diploma or GED	72.3	77.6	79.8	81.8	77.6	79.0	73.9	72.4	67.0
Some college or more	80.1	83.0	85.7	89.1	86.2	84.1	83.0	81.5	78.7
65 years and over:									
No high school diploma or GED	44.0	50.7	51.8	56.6	52.5	46.0	36.7	37.7	33.8
High school diploma or GED	55.4	61.6	63.7	66.9	61.2	52.5	49.3	42.6	38.8
Some college or more	59.4	62.3	68.8	69.8	67.8	63.8	59.0	54.9	49.7
Disability measure ⁸									
18 years and over, crude:									
Any basic actions difficulty or complex activity limitation	---	---	74.4	75.4	72.7	69.1	66.1	63.8	59.3
Any basic actions difficulty	---	---	74.3	75.1	72.6	69.1	66.2	63.6	59.2
Any complex activity limitation	---	---	69.3	71.0	67.6	62.2	60.1	58.5	52.8
No disability	---	---	83.8	84.1	82.5	82.6	80.4	78.9	75.2

See footnotes at end of table.

Table 71 (page 3 of 5). Use of Pap smears among women aged 18 and over, by selected characteristics: United States, selected years 1987–2013

Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#071>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	1987	1993	1999	2000	2003	2005	2008	2010	2013
Percent of women having a Pap smear within the past 3 years, among those who have not had a hysterectomy ⁹									
18 years and over, age-adjusted ^{2,3}	77.3	78.7	81.6	82.7	---	79.5	78.1	76.2	72.7
18 years and over, crude ²	77.8	80.0	82.6	83.3	---	80.7	79.3	77.3	73.9
Age									
18–44 years	85.1	84.7	86.3	84.9	---	83.8	81.8	80.3	77.2
18–20 years	62.1	67.3	63.3	59.8	---	61.3	57.6	52.0	38.6
21–44 years	87.7	86.4	89.0	88.0	---	86.6	85.0	84.0	81.7
21–24 years	85.9	86.0	84.8	84.3	---	84.0	80.2	81.1	74.5
25–44 years	88.1	86.5	89.7	88.7	---	87.2	86.0	84.7	83.5
45–64 years	75.8	79.2	83.8	86.9	---	83.3	83.7	81.6	79.2
45–54 years	80.9	82.9	85.5	87.6	---	85.5	83.8	83.1	81.7
55–64 years	70.5	73.6	80.6	85.5	---	79.6	83.6	79.4	75.9
65 years and over	55.4	59.7	63.7	68.6	---	59.1	56.1	54.1	48.8
65–74 years	62.8	67.9	71.9	75.9	---	72.1	69.9	66.9	63.0
75 years and over	44.4	49.9	54.7	60.9	---	46.2	41.9	39.3	29.9
Race ⁴									
18 years and over, crude:									
White only	77.8	79.9	82.8	83.7	---	81.1	79.6	77.4	73.8
Black or African American only	82.3	83.3	87.2	86.8	---	82.1	82.5	80.8	77.4
American Indian or Alaska Native only	85.9	78.2	94.1	77.7	---	75.6	74.8	78.9	74.3
Asian only	52.5	69.6	63.4	66.9	---	64.6	65.3	69.7	66.9
Native Hawaiian or Other Pacific Islander only	---	---	*	*	---	*	*	*	*
2 or more races	---	---	87.5	82.2	---	88.8	81.6	72.5	74.6
Hispanic origin and race ⁴									
18 years and over, crude:									
Hispanic or Latina	69.8	77.3	75.1	78.0	---	75.9	77.3	74.7	71.6
Not Hispanic or Latina	78.5	80.2	83.5	84.0	---	81.4	79.6	77.8	74.3
White only	78.6	80.2	83.6	84.4	---	82.1	80.2	78.1	74.4
Black or African American only	82.4	83.4	87.5	86.8	---	82.3	82.4	80.4	77.2
Age, Hispanic origin, and race ⁴									
18–44 years:									
Hispanic or Latina	75.1	80.2	76.0	77.9	---	76.5	78.3	75.6	72.0
Not Hispanic or Latina:									
White only	86.5	85.7	88.3	86.6	---	86.2	83.9	82.1	79.3
Black or African American only	90.3	87.6	90.6	88.7	---	86.1	83.3	84.0	82.5
45–64 years:									
Hispanic or Latina	62.4	75.3	77.8	81.0	---	78.6	81.0	77.7	77.3
Not Hispanic or Latina:									
White only	77.0	79.3	84.7	88.5	---	85.0	84.7	82.7	80.0
Black or African American only	78.0	81.1	86.6	87.4	---	80.7	85.6	81.7	77.6
65 years and over:									
Hispanic or Latina	43.8	58.9	60.9	71.2	---	60.0	53.7	56.4	49.4
Not Hispanic or Latina:									
White only	56.8	60.0	63.8	68.0	---	59.2	56.2	54.4	48.7
Black or African American only	46.3	55.8	65.1	72.1	---	59.3	64.1	52.7	45.6
Age and percent of poverty level ⁵									
18 years and over, crude:									
Below 100%	67.5	71.7	74.8	73.8	---	70.3	72.3	67.6	62.7
100%–199%	71.6	73.7	75.2	75.7	---	72.6	69.6	69.3	64.7
200%–399%	81.0	83.0	82.5	83.0	---	81.4	77.3	76.0	73.9
400% or more	87.0	87.8	88.9	90.5	---	88.2	87.8	87.1	84.0
18–44 years:									
Below 100%	79.3	77.2	79.0	76.8	---	76.1	76.6	73.0	69.3
100%–199%	81.8	82.1	83.7	79.2	---	78.1	75.4	75.6	72.3
200%–399%	86.6	86.5	86.2	86.0	---	86.1	82.4	79.7	78.4
400% or more	90.2	91.9	90.6	90.0	---	88.8	87.3	88.9	84.7
45–64 years:									
Below 100%	58.0	65.8	74.7	75.6	---	64.8	70.7	63.7	54.9
100%–199%	66.1	64.2	72.2	78.2	---	71.3	70.0	67.8	65.6
200%–399%	76.9	82.2	81.2	81.7	---	81.7	79.5	79.5	81.0
400% or more	84.4	86.6	89.7	93.7	---	90.9	92.4	90.8	88.0
65 years and over:									
Below 100%	36.4	47.5	53.5	55.9	---	43.7	44.7	36.5	36.8
100%–199%	54.6	56.6	56.3	63.3	---	54.4	48.7	48.1	37.7
200%–399%	62.8	63.5	68.3	71.8	---	61.4	53.3	56.1	43.8
400% or more	73.0	71.7	72.9	78.6	---	70.1	70.9	63.7	67.6

See footnotes at end of table.

Table 71 (page 4 of 5). Use of Pap smears among women aged 18 and over, by selected characteristics: United States, selected years 1987–2013

Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#071>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	1987	1993	1999	2000	2003	2005	2008	2010	2013
Health insurance status at the time of interview ⁶									
Percent of women having a Pap smear within the past 3 years, among those who have not had a hysterectomy ⁹									
18–64 years, crude:									
Insured	---	85.9	87.8	88.7	---	87.1	85.8	85.1	82.3
Private	---	86.0	88.1	88.8	---	87.9	86.6	86.2	83.5
Medicaid	---	83.9	84.2	86.9	---	82.6	82.4	79.7	77.6
Uninsured	---	70.2	74.3	70.8	---	68.0	67.9	63.1	59.6
Health insurance status prior to interview ⁶									
18–64 years, crude:									
Insured continuously all 12 months	---	86.1	88.0	88.9	---	87.2	86.1	85.4	82.9
Uninsured for any period up to 12 months	---	81.7	84.4	84.4	---	82.7	80.9	79.7	74.5
Uninsured more than 12 months	---	66.5	69.9	65.5	---	62.7	62.4	56.6	54.6
Age and education ⁷									
25 years and over, crude:									
No high school diploma or GED	61.7	63.2	68.3	72.5	---	66.9	67.5	61.0	60.2
High school diploma or GED	80.0	80.2	81.2	82.7	---	77.1	73.6	71.5	68.3
Some college or more	86.7	86.7	89.9	90.1	---	88.2	86.8	85.3	82.3
25–44 years:									
No high school diploma or GED	77.3	73.1	78.4	78.6	---	74.7	76.5	69.0	71.6
High school diploma or GED	87.6	85.6	87.4	86.2	---	83.4	79.5	78.8	79.6
Some college or more	91.5	90.0	92.9	91.7	---	91.1	89.7	89.2	86.5
45–64 years:									
No high school diploma or GED	63.9	65.5	73.2	77.5	---	70.5	74.8	66.8	65.7
High school diploma or GED	77.0	78.8	81.6	84.1	---	80.1	77.9	75.8	71.4
Some college or more	85.5	86.2	87.7	91.0	---	87.9	87.9	86.4	84.5
65 years and over:									
No high school diploma or GED	48.4	51.3	52.7	59.7	---	49.2	43.0	40.6	34.5
High school diploma or GED	60.4	63.8	65.0	71.3	---	56.5	53.6	48.7	45.3
Some college or more	63.6	65.7	75.6	74.9	---	69.9	66.1	64.0	57.6
Disability measure ⁸									
18 years and over, crude:									
Any basic actions difficulty or complex activity limitation	---	---	77.8	78.6	---	73.7	73.4	70.6	66.1
Any basic actions difficulty	---	---	77.8	78.5	---	73.9	73.8	70.6	66.2
Any complex activity limitation	---	---	73.9	73.9	---	67.4	68.1	65.9	59.1
No disability	---	---	84.5	85.1	---	84.0	82.1	80.8	77.4

See footnotes at end of table.

Table 71 (page 5 of 5). Use of Pap smears among women aged 18 and over, by selected characteristics: United States, selected years 1987–2013

Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#071>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

- - - Data not available.

* Estimates are considered unreliable. Data not shown have a relative standard error greater than 30%.

¹Includes all women aged 18 and over who had a Pap smear within past 3 years, including women who reported having had a hysterectomy. Questions concerning use of Pap smears differed slightly on the National Health Interview Survey across the years for which data are shown. See Appendix II, Pap smear. Data prior to 1997 are not strictly comparable with data for later years due to the 1997 questionnaire redesign. See Appendix I, National Health Interview Survey (NHIS).

²Includes all other races not shown separately, unknown poverty level in 1987, unknown health insurance status, unknown education level, and unknown disability status.

³Estimates are age-adjusted to the year 2000 standard population using five age groups: 18–44 years, 45–54 years, 55–64 years, 65–74 years, and 75 years and over. Age-adjusted estimates in this table may differ from other age-adjusted estimates based on the same data and presented elsewhere if different age groups are used in the adjustment procedure. See Appendix II, Age adjustment.

⁴The race groups, white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The five single-race categories plus multiple-race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group; the category 2 or more races includes persons who reported more than one racial group. Prior to 1999, data were tabulated according to the 1977 Standards with four racial groups, and the Asian only category included Native Hawaiian or Other Pacific Islander. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

⁵Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Poverty level was unknown for 9% of women aged 18 and over in 1987. Missing family income data were imputed for 1993 and beyond. See Appendix II, Family income; Poverty; Table VI.

⁶Health insurance categories are mutually exclusive. Persons who reported both Medicaid and private coverage are classified as having private coverage. Starting with 1997 data, state-sponsored health plan coverage is included as Medicaid coverage. Starting with 1999 data, coverage by the Children's Health Insurance Program (CHIP) is included with Medicaid coverage. In addition to private and Medicaid, the insured category also includes military plans, other government-sponsored health plans, and Medicare, not shown separately. Persons not covered by private insurance, Medicaid, CHIP, public assistance (through 1996), state-sponsored or other government-sponsored health plans (starting in 1997), Medicare, or military plans are considered to have no health insurance coverage. Persons with only Indian Health Service coverage are considered to have no health insurance coverage. See Appendix II, Health insurance coverage.

⁷Education categories shown are for 1998 and subsequent years. GED is General Educational Development high school equivalency diploma. In years prior to 1998, the following categories based on number of years of school completed were used: less than 12 years, 12 years, 13 years or more. See Appendix II, Education.

⁸Any basic actions difficulty or complex activity limitation is defined as having one or more of the following limitations or difficulties: movement difficulty, emotional difficulty, sensory (seeing or hearing) difficulty, cognitive difficulty, self-care (activities of daily living or instrumental activities of daily living) limitation, social limitation, or work limitation. For more information, see Appendix II, Basic actions difficulty; Complex activity limitation. Starting with 2007 data, the hearing question, a component of the basic actions difficulty measure, was revised. Consequently, data prior to 2007 are not comparable with data for 2007 and beyond. For more information on the impact of the revised hearing question, see Appendix II, Hearing trouble.

⁹The U.S. Preventive Services Task Force recommends against routine Pap smear screening in women who have had a total hysterectomy for benign disease. Therefore, Pap smear screening estimates are presented among women who have not had a hysterectomy, in addition to the estimates among all women, although it is not known, from National Health Interview Survey (NHIS) data, if the hysterectomy was for benign disease. Questions concerning hysterectomy differed slightly on NHIS across the years for which data are shown. See Appendix II, Pap smear.

NOTES: Currently, the U.S. Preventive Services Task Force (USPSTF) recommends Pap smears every three years for women aged 21 to 65, although the USPSTF recommendations have changed over time. See Appendix II, Pap smear. Standard errors are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey. Data are from the following supplements: cancer control (1987) and year 2000 objectives (1993–1994). Starting in 1998, data are from the family core and sample adult questionnaires. See Appendix I, National Health Interview Survey (NHIS).

Table 72 (page 1 of 2). Use of colorectal tests or procedures among adults aged 50–75, by selected characteristics: United States, selected years 2000–2013

Excel, PDF, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#072>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Any colorectal test or procedure ^{1,2}					Colonoscopy ^{2,3}				
	2000	2005	2008	2010	2013	2000	2005	2008	2010	2013
Percent of adults aged 50–75										
All adults 50–75 years ⁴	33.9	44.3	51.6	58.7	57.8	19.1	37.6	46.7	54.9	54.5
Sex										
Male	33.1	44.4	51.4	58.5	56.7	19.5	37.9	46.9	54.7	53.4
Female	34.5	44.2	51.9	58.8	58.9	18.8	37.4	46.6	55.1	55.5
Race ⁵										
White only	34.9	45.6	52.8	59.8	58.4	19.7	38.9	47.8	56.0	55.3
Black or African American only	29.6	38.1	46.9	55.2	58.0	17.4	32.2	43.1	51.8	54.1
American Indian or Alaska Native only	*35.2	*33.9	28.5	48.9	49.3	*	*	*26.7	46.7	45.8
Asian only	20.4	30.8	47.1	47.1	49.8	*8.6	24.4	39.3	43.6	43.7
Native Hawaiian or Other Pacific Islander only	*	*	*	*	*	*	*	*	*	*
2 or more races	37.5	33.8	38.4	51.9	50.5	*25.1	29.6	37.4	48.4	48.4
Hispanic origin and race ⁵										
Hispanic or Latino	21.7	28.5	34.0	46.5	41.5	13.3	23.1	29.3	43.9	37.5
Mexican	19.3	24.6	27.5	44.6	39.2	11.2	18.2	21.2	41.3	35.2
Not Hispanic or Latino	34.7	45.6	53.3	59.9	59.6	19.5	38.9	48.4	56.0	56.3
White only	35.7	47.4	54.8	61.3	60.4	20.0	40.5	49.8	57.3	57.4
Black or African American only	29.7	38.0	47.4	55.3	58.2	17.5	32.0	43.5	52.0	54.6
Percent of poverty level ⁶										
Below 100%	26.5	28.7	33.9	37.9	43.7	16.3	23.6	28.5	34.8	40.5
100%–199%	29.4	38.4	42.7	47.9	48.4	17.7	31.5	38.0	43.3	44.8
200%–399%	33.7	43.6	49.9	58.0	55.8	18.6	37.0	44.3	54.6	52.0
400% or more	37.1	49.6	58.9	67.3	65.6	20.5	42.8	54.5	63.6	62.7
Hispanic origin and race and percent of poverty level ^{5,6}										
Hispanic or Latino:										
Below 100%	15.3	19.3	21.1	33.7	35.7	*9.3	13.1	17.9	32.1	32.0
100%–199%	16.8	24.6	27.7	39.6	35.1	8.6	19.4	24.4	36.3	31.2
200%–399%	23.6	28.3	39.3	47.5	41.5	*13.7	21.6	33.8	46.0	37.3
400% or more	31.1	42.1	43.9	63.3	53.0	22.4	39.3	37.6	59.5	48.8
Not Hispanic or Latino:										
White only:										
Below 100%	29.6	30.6	39.8	40.4	46.8	19.3	26.8	33.2	36.4	44.0
100%–199%	32.1	42.4	46.0	50.0	51.9	19.7	35.0	40.7	44.5	48.3
200%–399%	35.2	47.3	51.6	59.7	57.6	19.3	40.2	45.8	56.3	54.0
400% or more	37.9	50.6	60.5	68.0	66.2	20.7	43.8	56.3	64.3	63.6
Black or African American only:										
Below 100%	27.5	29.0	35.1	39.2	45.5	14.5	23.5	30.1	36.4	41.2
100%–199%	28.7	36.2	46.7	49.0	51.4	17.2	30.3	43.2	46.5	47.3
200%–399%	27.7	35.8	48.5	60.5	61.3	16.5	31.8	44.7	56.2	57.9
400% or more	33.9	48.9	54.3	68.1	70.5	20.7	40.2	50.6	64.6	67.5
Education ⁷										
No high school diploma or GED	25.9	34.5	36.2	44.6	43.5	14.9	29.0	31.8	41.5	39.9
High school diploma or GED	33.1	42.1	48.5	53.7	53.4	19.0	35.7	44.6	50.8	50.4
Some college or more	37.8	48.7	57.5	64.7	63.1	20.9	41.6	52.1	60.4	59.6

See footnotes at end of table.

Table 72 (page 2 of 2). Use of colorectal tests or procedures among adults aged 50–75, by selected characteristics: United States, selected years 2000–2013

Excel, PDF, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#072>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Any colorectal test or procedure ^{1,2}					Colonoscopy ^{2,3}				
	2000	2005	2008	2010	2013	2000	2005	2008	2010	2013
Disability measure ⁸										
Percent of adults aged 50–75										
Any basic actions difficulty or complex activity limitation	37.8	47.7	54.2	59.5	61.0	22.1	40.1	48.5	55.5	57.6
Any basic actions difficulty	38.1	47.9	54.6	59.7	61.5	22.5	40.6	48.9	55.8	58.0
Any complex activity limitation	37.4	48.1	52.4	59.4	59.9	22.6	39.7	46.7	55.1	55.7
No disability	30.9	41.6	50.0	58.5	55.5	16.6	35.6	45.8	54.9	52.2
Geographic region										
Northeast	34.4	50.9	54.7	64.3	61.0	19.1	44.8	51.0	61.7	59.4
Midwest	35.2	43.5	52.5	58.4	59.5	19.8	36.6	47.8	55.2	57.3
South	32.5	43.9	51.6	57.4	56.4	20.0	38.1	47.4	54.4	53.8
West	34.1	39.6	48.2	56.3	55.9	16.3	31.3	41.1	49.7	48.7
Location of residence ⁹										
Within MSA	34.1	44.7	52.4	59.6	58.3	19.0	37.9	47.6	55.8	54.8
Outside MSA	33.2	42.7	48.5	54.4	55.6	19.6	36.7	43.3	50.9	52.8

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.
¹Includes reports of home fecal occult blood test (FOBT) in the past year, sigmoidoscopy procedure in the past 5 years with FOBT in the past 3 years, or colonoscopy in the past 10 years. Colorectal procedures are performed for diagnostic and screening purposes.
²Questions differed slightly on the National Health Interview Survey across the years for which data are shown. See Appendix II, Colorectal tests or procedures.
³Includes any colonoscopy in the past 10 years, alone or in addition to another type of colorectal test or procedure.
⁴Includes all other races not shown separately, unknown disability status, and unknown education level.
⁵The race groups, white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. The five single-race and multiple-race categories shown in the table conform to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity*. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.
⁶Based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Missing family income data were imputed. See Appendix II, Family income; Poverty; Table VI.
⁷GED is General Educational Development high school equivalency diploma. See Appendix II, Education.
⁸Any basic actions difficulty or complex activity limitation is defined as having one or more of the following limitations or difficulties: movement difficulty, emotional difficulty, sensory (seeing or hearing) difficulty, cognitive difficulty, self-care (activities of daily living or instrumental activities of daily living) limitation, social limitation, or work limitation. For more information, see Appendix II, Basic actions difficulty; Complex activity limitation. Starting with 2007 data, the hearing question, a component of the basic actions difficulty measure, was revised. Consequently, data prior to 2007 are not comparable with data for 2007 and beyond. For more information on the impact of the revised hearing question, see Appendix II, Hearing trouble.
⁹MSA is metropolitan statistical area. Starting with 2006 data, MSA status is determined using 2000 census data and the 2000 standards for defining MSAs. For data prior to 2006, see Appendix II, Metropolitan statistical area (MSA) for the applicable standards.

NOTES: In 2008, the U.S. Preventive Services Task Force (USPSTF) recommended screening for colorectal cancer annually using FOBT, every 5 years using sigmoidoscopy with FOBT every 3 years, or every 10 years using colonoscopy, in adults, beginning at age 50 and continuing until age 75. See: <http://www.uspreventiveservicestaskforce.org/uspstf08/colocancer/colors.htm> for more information. Colonoscopy is one of the three modalities currently recommended by USPSTF for colorectal cancer screening. USPSTF does not recommend one screening method over another, and the risks and benefits of these screening methods vary. Colonoscopy estimates are shown separately because of the recent large increase in its utilization. The American College of Gastroenterology recommends that African American persons start routine testing for colorectal cancer at age 45. See: <http://www.acg.gi.org/patients/ccrk/> for more information. Standard errors for selected years are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey. Family core and sample adult questionnaires. See Appendix I, National Health Interview Survey (NHIS).

Table 73 (page 1 of 4). Emergency department visits within the past 12 months among children under age 18, by selected characteristics: United States, selected years 1997–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#073>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Under 18 years			Under 6 years			6–17 years		
	1997	2010	2014	1997	2010	2014	1997	2010	2014
	Percent of children with one or more emergency department visits ¹								
All children ²	19.9	22.1	16.7	24.3	27.8	22.6	17.7	19.1	13.8
Sex									
Male	21.5	23.3	16.9	25.2	29.3	22.1	19.6	20.1	14.4
Female	18.3	20.9	16.4	23.3	26.3	23.2	15.7	18.2	13.2
Race ³									
White only	19.4	21.2	15.9	22.6	26.6	21.5	17.8	18.4	13.2
Black or African American only	24.0	27.6	21.3	33.1	34.0	30.0	19.4	24.2	17.1
American Indian or Alaska Native only	*24.1	20.9	28.7	*24.3	*35.4	*31.3	*24.0	*	*27.5
Asian only	12.6	15.0	10.4	20.8	18.4	13.2	8.6	13.3	8.8
Native Hawaiian or Other Pacific Islander only	---	*	*	---	*	*	---	*	*
2 or more races	---	27.2	19.4	---	34.9	25.3	---	21.6	15.6
Hispanic origin and race ³									
Hispanic or Latino	21.1	23.6	17.1	25.7	30.2	21.8	18.1	19.4	14.6
Not Hispanic or Latino	19.7	21.7	16.6	24.0	27.0	22.9	17.6	19.0	13.6
White only	19.2	20.4	15.8	22.2	25.1	21.8	17.7	18.2	13.0
Black or African American only	23.6	27.2	20.7	32.7	34.4	30.1	19.2	23.3	16.4
Percent of poverty level ⁴									
Below 100%	25.1	30.6	24.7	29.5	35.4	31.2	22.2	27.6	21.1
100%–199%	22.0	25.7	18.5	28.0	31.6	23.4	19.0	22.3	16.0
200%–399%	18.0	18.4	13.8	21.4	22.7	20.1	16.4	16.4	10.7
400% or more	16.3	15.9	11.4	19.1	21.7	16.0	15.1	13.3	9.3
Hispanic origin and race and percent of poverty level ^{3,4}									
Hispanic or Latino:									
Percent of poverty level:									
Below 100%	21.9	27.0	21.0	25.0	32.0	26.3	19.6	23.4	17.8
100%–199%	20.8	23.3	16.6	28.8	31.6	21.7	15.6	18.0	14.1
200%–399%	21.4	19.5	14.4	24.6	25.2	16.5	19.6	16.1	13.4
400% or more	17.7	21.4	11.3	*20.2	28.6	*16.2	16.4	18.0	*8.8
Not Hispanic or Latino:									
White only:									
Percent of poverty level:									
Below 100%	25.5	33.7	28.8	27.2	37.4	32.2	24.4	31.6	27.2
100%–199%	22.3	26.3	19.0	25.8	29.2	23.2	20.7	24.7	16.9
200%–399%	17.8	17.6	13.8	20.9	21.2	20.9	16.3	15.9	10.1
400% or more	16.5	15.5	11.4	19.0	21.0	17.5	15.4	13.2	9.0
Black or African American only:									
Percent of poverty level:									
Below 100%	29.3	32.4	24.8	39.5	41.6	37.0	23.0	26.6	18.2
100%–199%	22.5	27.5	20.6	31.7	34.5	28.6	18.5	23.7	16.7
200%–399%	18.5	22.3	14.6	23.9	24.6	*19.5	16.3	21.4	13.2
400% or more	16.1	18.9	17.1	*18.8	*24.1	*19.5	15.2	16.1	*16.0
Health insurance status at the time of interview ⁵									
Insured	19.8	22.3	16.9	24.4	28.1	22.7	17.5	19.2	14.0
Private	17.5	17.1	12.4	20.9	21.8	17.6	15.9	14.9	10.1
Medicaid	28.2	30.0	22.9	33.0	35.5	28.5	24.1	26.4	19.6
Uninsured	20.2	19.4	14.7	23.0	24.0	22.9	18.9	17.6	12.0
Health insurance status prior to interview ⁵									
Insured continuously all 12 months	19.6	22.2	16.8	24.1	28.1	22.8	17.3	19.1	13.9
Uninsured for any period up to 12 months	24.0	23.7	18.0	27.1	28.0	20.7	21.9	21.3	16.6
Uninsured more than 12 months	18.4	17.6	11.3	19.3	*21.3	*	18.1	16.7	*9.0

See footnotes at end of table.

Table 73 (page 2 of 4). Emergency department visits within the past 12 months among children under age 18, by selected characteristics: United States, selected years 1997–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#073>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Under 18 years			Under 6 years			6–17 years		
	1997	2010	2014	1997	2010	2014	1997	2010	2014
Percent of poverty level and health insurance status prior to interview ^{4,5}									
Percent of children with one or more emergency department visits ¹									
Below 100%:									
Insured continuously all 12 months	26.3	31.7	25.5	30.9	36.3	31.4	22.8	28.7	22.1
Uninsured for any period up to 12 months	26.5	30.3	21.6	29.7	34.7	*28.5	24.4	27.5	*18.0
Uninsured more than 12 months	17.5	*19.6	*	*16.0	*	*	18.0	*16.0	*
100%–199%:									
Insured continuously all 12 months	21.8	26.2	18.6	28.0	32.4	23.6	18.6	22.4	16.0
Uninsured for any period up to 12 months	24.5	28.4	21.8	29.7	30.9	*26.1	21.0	27.0	*19.6
Uninsured more than 12 months	19.5	17.6	*13.4	*22.5	*	*	18.6	*17.2	*13.8
200%–399%:									
Insured continuously all 12 months	17.7	18.4	13.9	21.2	22.8	20.7	16.1	16.3	10.5
Uninsured for any period up to 12 months	21.1	16.2	*15.3	*19.5	*22.7	*	22.1	*12.6	*15.8
Uninsured more than 12 months	19.2	*17.4	*9.1	*22.7	*	*	17.6	*18.7	*
400% or more:									
Insured continuously all 12 months	16.2	16.1	11.5	18.9	22.0	16.0	15.1	13.5	9.5
Uninsured for any period up to 12 months	*19.2	*	*	*	*	*	*	*	*
Uninsured more than 12 months	*	*	*	*	*	*	*	*	*
Geographic region									
Northeast	18.5	22.3	17.6	20.7	27.8	22.2	17.4	19.6	15.5
Midwest	19.5	23.3	18.7	26.0	28.8	24.8	16.4	20.7	15.9
South	21.8	23.4	16.7	25.6	30.4	23.9	19.9	19.5	13.2
West	18.5	19.1	14.2	23.5	23.3	18.9	15.9	16.8	11.8
Location of residence ⁶									
Within MSA	19.7	21.8	16.5	23.9	27.7	22.4	17.4	18.6	13.7
Outside MSA	20.8	24.2	17.8	26.2	28.6	24.5	18.6	22.1	14.7
Percent of children with two or more emergency department visits ¹									
All children ²	7.1	8.4	4.8	9.6	10.8	6.8	5.8	7.2	3.9
Sex									
Male	7.3	8.5	5.0	9.9	11.3	7.1	6.0	7.0	4.0
Female	6.9	8.3	4.6	9.4	10.3	6.5	5.7	7.3	3.7
Race ³									
White only	6.6	7.6	4.3	8.4	10.1	5.7	5.7	6.3	3.6
Black or African American only	9.6	12.6	7.2	14.9	15.7	12.6	6.9	11.0	4.7
American Indian or Alaska Native only	*	*	*12.6	*	*	*	*	*	*10.4
Asian only	*5.7	7.3	*2.4	*12.9	*	*	*	*7.1	*
Native Hawaiian or Other Pacific Islander only	---	*	*	---	*	*	---	*	*
2 or more races	---	10.3	6.2	---	*11.7	*6.7	---	*9.2	*5.9
Hispanic origin and race ³									
Hispanic or Latino	8.9	8.6	5.0	11.8	11.7	6.6	7.0	6.6	4.2
Not Hispanic or Latino	6.8	8.4	4.8	9.2	10.5	6.9	5.7	7.3	3.8
White only	6.2	7.4	4.1	7.8	9.3	5.5	5.5	6.4	3.5
Black or African American only	9.3	12.3	7.1	14.6	15.8	12.7	6.8	10.4	4.5
Percent of poverty level ⁴									
Below 100%	11.1	13.4	8.9	14.5	15.3	12.2	8.9	12.1	7.1
100%–199%	8.3	10.3	5.6	12.2	13.4	7.0	6.3	8.4	4.9
200%–399%	6.2	6.3	3.4	7.4	7.3	5.4	5.6	5.9	2.5
400% or more	4.0	5.0	2.2	5.0	7.3	*2.8	3.6	3.9	1.9

See footnotes at end of table.

Table 73 (page 3 of 4). Emergency department visits within the past 12 months among children under age 18, by selected characteristics: United States, selected years 1997–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#073>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Under 18 years			Under 6 years			6–17 years		
	1997	2010	2014	1997	2010	2014	1997	2010	2014
Hispanic origin and race and percent of poverty level ^{3,4}									
Percent of children with two or more emergency department visits ¹									
Hispanic or Latino:									
Percent of poverty level:									
Below 100%	10.4	9.9	7.3	13.9	10.9	9.6	8.0	9.2	5.9
100%–199%	8.2	9.4	4.5	12.0	15.4	5.8	5.7	5.5	*3.8
200%–399%	8.5	5.9	*3.5	10.0	*8.0	*	*7.6	*4.6	*
400% or more	*5.0	*6.5	*	*	*	*	*	*5.2	*
Not Hispanic or Latino:									
White only:									
Percent of poverty level:									
Below 100%	10.7	14.0	11.6	12.2	15.5	*13.7	9.8	13.1	*10.6
100%–199%	8.0	10.4	*4.5	11.2	12.3	*3.9	6.4	9.4	*4.8
200%–399%	6.0	5.7	3.1	6.7	*6.5	*5.1	5.6	5.4	*2.1
400% or more	3.7	5.0	2.2	4.6	7.6	*	3.3	3.9	*1.8
Black or African American only:									
Percent of poverty level:									
Below 100%	12.7	16.1	8.6	19.1	22.1	14.8	8.8	12.4	*5.2
100%–199%	9.2	12.4	*7.6	*13.5	*14.6	*14.3	*7.2	11.1	*4.2
200%–399%	5.8	9.9	*5.3	*8.9	*10.2	*	*4.5	*9.8	*
400% or more	*	*3.7	*	*	*	*	*	*	*
Health insurance status at the time of interview ⁵									
Insured	7.0	8.5	4.8	9.6	11.0	6.8	5.7	7.1	3.8
Private	5.2	5.5	2.6	6.8	7.4	3.6	4.5	4.6	2.1
Medicaid	13.1	12.8	7.7	16.2	15.3	10.2	10.4	11.2	6.3
Uninsured	7.7	8.0	*5.9	9.8	*8.5	*	6.8	7.8	*4.8
Health insurance status prior to interview ⁵									
Insured continuously all 12 months	6.9	8.4	4.8	9.4	10.8	6.8	5.7	7.1	3.8
Uninsured for any period up to 12 months	8.5	10.1	*5.6	11.5	13.3	*8.4	6.6	8.4	*4.2
Uninsured more than 12 months	6.8	7.8	*4.5	*8.6	*	*	6.2	*7.9	*5.0
Geographic region									
Northeast	6.2	7.8	5.1	7.6	10.3	7.7	5.4	6.6	3.9
Midwest	6.6	9.1	5.7	10.4	11.4	7.9	4.8	8.0	4.6
South	8.0	9.1	4.9	10.1	12.9	6.9	6.9	7.1	3.9
West	7.1	7.2	3.8	10.0	7.6	5.3	5.6	7.0	3.1
Location of residence ⁶									
Within MSA	7.2	8.3	4.6	9.6	10.6	6.6	5.9	7.0	3.7
Outside MSA	6.8	9.3	6.3	9.7	12.2	8.7	5.6	7.9	*5.2

See footnotes at end of table.

Table 73 (page 4 of 4). Emergency department visits within the past 12 months among children under age 18, by selected characteristics: United States, selected years 1997–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#073>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.

-- Data not available.

¹See Appendix II, Emergency department or emergency room visit.

²Includes all other races not shown separately and unknown health insurance status.

³The race groups, white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The five single-race categories plus multiple-race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group; the category 2 or more races includes persons who reported more than one racial group. Prior to 1999, data were tabulated according to the 1977 Standards with four racial groups, and the Asian only category included Native Hawaiian or Other Pacific Islander. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

⁴Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Missing family income data were imputed for 1997 and beyond. See Appendix II, Family income; Poverty; Table VI.

⁵Health insurance categories are mutually exclusive. Persons who reported both Medicaid and private coverage are classified as having private coverage. Starting with 1997 data, state-sponsored health plan coverage is included as Medicaid coverage. Starting with 1999 data, coverage by the Children's Health Insurance Program (CHIP) is included with Medicaid coverage. In addition to private and Medicaid, the insured category also includes military, other government, and Medicare coverage. Persons not covered by private insurance, Medicaid, CHIP, state-sponsored or other government-sponsored health plans (starting in 1997), Medicare, or military plans are considered to have no health insurance coverage. Persons with only Indian Health Service coverage are considered to have no health insurance coverage. See Appendix II, Health insurance coverage.

⁶MSA is metropolitan statistical area. Starting with 2006 data, MSA status is determined using 2000 census data and the 2000 standards for defining MSAs. For data prior to 2006, see Appendix II, Metropolitan statistical area (MSA) for the applicable standards.

NOTES: Standard errors are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey, family core and sample child questionnaires. See Appendix I, National Health Interview Survey (NHIS).

Table 74 (page 1 of 3). Emergency department visits within the past 12 months among adults aged 18 and over, by selected characteristics: United States, selected years 1997–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#074>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	One or more emergency department visits				Two or more emergency department visits			
	1997	2000	2010	2014	1997	2000	2010	2014
Percent of adults with emergency department visits ¹								
18 years and over, age-adjusted ^{2,3}	19.6	20.2	21.4	18.6	6.7	6.9	7.8	6.7
18 years and over, crude ²	19.6	20.1	21.3	18.6	6.7	6.8	7.7	6.7
Age								
18–44 years	20.7	20.5	22.0	18.4	6.8	7.0	8.4	6.7
18–24 years	26.3	25.7	25.4	20.9	9.1	8.8	9.6	8.2
25–44 years	19.0	18.8	20.7	17.5	6.2	6.4	8.0	6.1
45–64 years	16.2	17.6	19.2	17.5	5.6	5.6	6.7	6.4
45–54 years	15.7	17.9	18.6	16.2	5.5	5.8	6.6	5.8
55–64 years	16.9	17.0	19.8	18.9	5.7	5.3	6.8	7.0
65 years and over	22.0	23.7	23.7	21.2	8.1	8.6	7.7	7.1
65–74 years	20.3	21.6	20.7	18.9	7.1	7.4	6.4	6.1
75 years and over	24.3	26.2	27.4	24.4	9.3	10.0	9.4	8.6
Sex ³								
Male	19.1	18.7	18.5	16.9	5.9	5.7	6.0	5.3
Female	20.2	21.6	24.3	20.3	7.5	7.9	9.6	8.1
Race ^{3,4}								
White only	19.0	19.4	20.7	17.7	6.2	6.4	7.2	6.1
Black or African American only	25.9	26.5	28.6	26.3	11.1	10.8	12.6	11.1
American Indian or Alaska Native only	24.8	30.3	22.6	31.1	13.1	*12.6	*11.8	18.6
Asian only	11.6	13.6	13.3	11.0	*2.9	*3.8	3.3	2.8
Native Hawaiian or Other Pacific Islander only	---	*	*	*	---	*	*	*
2 or more races	---	32.5	29.7	23.8	---	11.3	11.1	8.1
American Indian or Alaska Native; White	---	33.9	31.1	25.5	---	*9.4	*15.2	*9.1
Hispanic origin and race ^{3,4}								
Hispanic or Latino	19.2	18.3	19.8	17.0	7.4	7.0	6.9	6.0
Mexican	17.8	17.4	18.1	17.4	6.4	7.1	6.1	6.1
Not Hispanic or Latino	19.7	20.6	21.9	19.0	6.7	6.9	8.1	6.9
White only	19.1	19.8	21.1	18.1	6.2	6.4	7.4	6.3
Black or African American only	25.9	26.5	29.0	26.6	11.0	10.8	12.7	11.2
Percent of poverty level ^{3,5}								
Below 100%	28.1	29.0	30.6	28.6	12.8	13.3	14.9	12.9
100%–199%	23.8	23.9	25.6	23.4	9.3	9.6	10.5	9.5
200%–399%	18.3	19.8	20.4	17.2	5.9	6.3	6.8	6.2
400% or more	15.9	16.8	17.0	13.5	3.9	4.5	4.7	3.4
Hispanic origin and race and percent of poverty level ^{3,4,5}								
Hispanic or Latino:								
Below 100%	22.1	22.4	23.6	22.6	9.8	9.7	11.5	10.2
100%–199%	19.2	18.1	19.9	17.0	8.1	6.7	6.3	5.7
200%–399%	18.5	17.3	18.1	14.7	6.0	7.4	5.2	4.9
400% or more	14.6	16.4	18.8	13.6	*3.8	*4.3	*5.5	*3.4
Not Hispanic or Latino:								
White only:								
Below 100%	29.5	30.1	33.3	30.7	13.0	13.9	15.5	13.4
100%–199%	24.3	25.5	26.8	25.4	9.1	10.4	11.2	10.8
200%–399%	18.1	20.1	20.3	17.1	5.8	6.3	6.5	6.1
400% or more	15.8	16.3	16.9	13.5	3.8	4.1	4.9	3.3
Black or African American only:								
Below 100%	34.6	35.4	36.9	35.3	17.5	17.4	20.2	17.7
100%–199%	29.2	28.5	33.5	30.4	12.8	12.2	15.9	12.9
200%–399%	20.8	23.2	25.7	22.0	8.1	8.0	10.2	8.8
400% or more	18.2	22.6	18.8	18.2	5.9	8.8	*4.0	5.6

See footnotes at end of table.

Table 74 (page 2 of 3). Emergency department visits within the past 12 months among adults aged 18 and over, by selected characteristics: United States, selected years 1997–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#074>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	One or more emergency department visits				Two or more emergency department visits			
	1997	2000	2010	2014	1997	2000	2010	2014
Health insurance status at the time of interview ^{6,7}								
Percent of adults with emergency department visits ¹								
18–64 years:								
Insured	18.8	19.5	20.8	18.2	6.1	6.4	7.5	6.7
Private	16.9	17.6	17.4	14.5	4.7	5.1	5.2	4.2
Medicaid	37.6	42.2	40.2	34.9	19.7	21.0	21.1	18.4
Uninsured	20.0	19.3	21.3	16.5	7.5	6.9	8.9	5.9
Health insurance status prior to interview ^{6,7}								
18–64 years:								
Insured continuously all 12 months	18.3	19.0	20.2	17.6	5.8	6.1	7.1	6.4
Uninsured for any period up to 12 months	25.5	28.2	26.0	22.9	9.4	10.3	12.5	9.3
Uninsured more than 12 months	18.9	17.3	20.6	15.4	7.1	6.4	8.1	5.4
Percent of poverty level and health insurance status prior to interview ^{5,6,7}								
18–64 years:								
Below 100%:								
Insured continuously all 12 months	30.2	31.6	35.2	32.7	14.7	15.4	18.3	16.0
Uninsured for any period up to 12 months	34.1	43.7	34.2	29.6	16.1	18.1	16.5	14.5
Uninsured more than 12 months	20.8	20.5	23.4	20.2	8.1	9.1	11.7	6.9
100%–199%:								
Insured continuously all 12 months	24.5	25.5	26.1	25.0	8.9	10.2	10.8	11.0
Uninsured for any period up to 12 months	28.7	27.7	29.7	29.0	12.3	11.7	15.6	11.6
Uninsured more than 12 months	19.0	17.4	21.2	14.2	8.3	6.4	7.8	5.8
200%–399%:								
Insured continuously all 12 months	17.5	19.5	19.6	16.3	5.3	6.3	6.0	6.0
Uninsured for any period up to 12 months	21.6	24.6	25.4	19.9	6.6	7.3	12.2	7.2
Uninsured more than 12 months	16.8	15.6	17.6	12.6	5.9	4.5	5.7	*4.2
400% or more:								
Insured continuously all 12 months	14.9	15.5	15.9	12.5	3.7	3.7	4.5	2.8
Uninsured for any period up to 12 months	18.0	20.1	12.5	9.2	*3.1	6.4	*	*
Uninsured more than 12 months	19.1	15.8	19.4	*10.7	*	*5.2	*	*
Disability measure ^{3,8}								
Any basic actions difficulty or complex activity limitation	30.8	32.0	34.9	32.2	13.5	14.6	16.8	14.8
Any basic actions difficulty	30.5	32.4	35.0	32.6	13.5	14.9	17.2	15.0
Any complex activity limitation	39.7	41.5	43.8	41.5	19.9	21.2	24.5	21.7
No disability	14.5	15.3	16.1	13.2	3.7	3.9	4.4	3.6
Geographic region ³								
Northeast	19.5	20.0	22.6	18.3	6.9	6.2	8.4	6.4
Midwest	19.3	20.1	22.3	20.6	6.2	6.9	8.2	7.7
South	20.9	21.2	22.1	19.1	7.3	7.6	8.0	6.9
West	17.7	18.6	18.9	16.0	6.0	6.3	6.7	5.4
Location of residence ^{3,9}								
Within MSA	19.1	19.6	20.8	18.1	6.4	6.6	7.5	6.4
Outside MSA	21.5	22.5	25.5	22.0	7.8	7.8	9.8	8.9

See footnotes at end of table.

Table 74 (page 3 of 3). Emergency department visits within the past 12 months among adults aged 18 and over, by selected characteristics: United States, selected years 1997–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#074>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.
- - - Data not available.

¹See Appendix II, Emergency department or emergency room visit.

²Includes all other races not shown separately, unknown health insurance status, and unknown disability status.

³Estimates are for persons aged 18 and over and are age-adjusted to the year 2000 standard population using five age groups: 18–44 years, 45–54 years, 55–64 years, 65–74 years, and 75 years and over. See Appendix II, Age adjustment.

⁴The race groups, white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The five single-race categories plus multiple-race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group; the category 2 or more races includes persons who reported more than one racial group. Prior to 1999, data were tabulated according to the 1977 Standards with four racial groups, and the Asian only category included Native Hawaiian or Other Pacific Islander. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

⁵Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Missing family income data were imputed for 1997 and beyond. See Appendix II, Family income; Poverty; Table VI.

⁶Estimates for persons aged 18–64 are age-adjusted to the year 2000 standard population using three age groups: 18–44 years, 45–54 years, and 55–64 years. See Appendix II, Age adjustment.

⁷Health insurance categories are mutually exclusive. Persons who reported both Medicaid and private coverage are classified as having private coverage. Starting with 1997 data, state-sponsored health plan coverage is included as Medicaid coverage. Starting with 1999 data, coverage by the Children's Health Insurance Program (CHIP) is included with Medicaid coverage. In addition to private and Medicaid, the insured category also includes military plans, other government-sponsored health plans, and Medicare, not shown separately. Persons not covered by private insurance, Medicaid, CHIP, state-sponsored or other government-sponsored health plans (starting in 1997), Medicare, or military plans are considered to have no health insurance coverage. Persons with only Indian Health Service coverage are considered to have no health insurance coverage. See Appendix II, Health insurance coverage.

⁸Any basic actions difficulty or complex activity limitation is defined as having one or more of the following limitations or difficulties: movement difficulty, emotional difficulty, sensory (seeing or hearing) difficulty, cognitive difficulty, self-care (activities of daily living or instrumental activities of daily living) limitation, social limitation, or work limitation. For more information, see Appendix II, Basic actions difficulty; Complex activity limitation. Starting with 2007 data, the hearing question, a component of the basic actions difficulty measure, was revised. Consequently, data prior to 2007 are not comparable with data for 2007 and beyond. For more information on the impact of the revised hearing question, see Appendix II, Hearing trouble.

⁹MSA is metropolitan statistical area. Starting with 2006 data, MSA status is determined using 2000 census data and the 2000 standards for defining MSAs. For data prior to 2006, see Appendix II, Metropolitan statistical area (MSA) for the applicable standards.

NOTES: Standard errors are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey, family core and sample adult questionnaires. See Appendix I, National Health Interview Survey (NHIS).

Table 75 (page 1 of 2). Initial injury-related visits to hospital emergency departments, by sex, age, and intent and mechanism of injury: United States, average annual, selected years 2005–2006 through 2010–2011

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#075>.

[Data are based on reporting by a sample of hospital emergency departments]

<i>Sex, age, and intent and mechanism of injury</i> ¹	<i>2005–2006</i>	<i>2008–2009</i>	<i>2010–2011</i>	<i>2005–2006</i>	<i>2008–2009</i>	<i>2010–2011</i>
Both sexes	Initial injury-related visits, in thousands			Initial injury-related visits per 10,000 persons		
All ages, age-adjusted ^{2,3}	31,706	31,328	33,007	1,076.4	1,040.8	1,084.0
All ages, crude ²	31,706	31,328	33,007	1,068.6	1,029.4	1,067.9
Unintentional injuries ⁴	25,658	25,725	27,275	864.7	845.3	882.4
Falls	8,100	8,900	9,932	273.0	292.4	321.3
Struck by or against objects or persons	2,935	2,916	3,166	98.9	95.8	102.4
Motor vehicle traffic	3,714	3,508	3,557	125.2	115.3	115.1
Cut or pierce	2,145	2,008	1,922	72.3	66.0	62.2
Intentional injuries	1,977	2,313	2,446	66.6	76.0	79.1
Male						
All ages, age-adjusted ^{2,3}	16,966	16,640	17,483	1,166.1	1,118.0	1,164.5
All ages, crude ²	16,966	16,640	17,483	1,164.2	1,111.8	1,150.5
Unintentional injuries ⁴	13,736	13,590	14,451	942.5	908.0	951.0
Falls	3,685	3,944	4,689	252.9	263.5	308.6
Struck by or against objects or persons	1,833	1,863	2,008	125.8	124.4	132.2
Motor vehicle traffic	1,733	1,734	1,710	118.9	115.8	112.5
Cut or pierce	1,392	1,263	1,236	95.5	84.4	81.4
Intentional injuries	1,135	1,266	1,396	77.8	84.6	91.8
Under 18 years ²	5,072	5,132	5,309	1,346.6	1,351.1	1,397.8
Unintentional injuries ⁴	4,391	4,509	4,724	1,165.8	1,187.1	1,243.9
Falls	1,362	1,512	1,737	361.5	398.1	457.4
Struck by or against objects or persons	816	909	997	216.6	239.2	262.6
Motor vehicle traffic	357	305	301	94.8	80.3	79.1
Cut or pierce	291	284	238	77.3	74.8	62.7
Intentional injuries	190	194	167	50.4	51.1	44.1
18–24 years ²	2,552	2,562	2,511	1,729.5	1,695.5	1,612.1
Unintentional injuries ⁴	1,985	1,947	1,890	1,345.4	1,288.6	1,213.7
Falls	318	366	390	215.2	242.4	250.4
Struck by or against objects or persons	290	283	259	196.9	187.4	166.6
Motor vehicle traffic	386	373	357	261.6	247.0	229.3
Cut or pierce	265	215	192	179.5	142.6	123.5
Intentional injuries	273	381	403	185.2	252.2	258.7
25–44 years ²	5,199	4,611	4,850	1,243.6	1,109.5	1,184.3
Unintentional injuries ⁴	4,001	3,540	3,690	957.1	851.8	901.1
Falls	763	703	815	182.4	169.2	199.1
Struck by or against objects or persons	472	401	452	112.9	96.4	110.4
Motor vehicle traffic	629	578	591	150.5	139.1	144.3
Cut or pierce	480	401	423	114.8	96.5	103.2
Intentional injuries	436	495	589	104.4	119.2	143.8
45–64 years ²	2,842	2,996	3,270	790.0	780.7	822.7
Unintentional injuries ⁴	2,275	2,437	2,741	632.5	635.1	689.6
Falls	599	669	909	166.6	174.2	228.6
Struck by or against objects or persons	208	216	204	57.9	56.4	51.4
Motor vehicle traffic	262	375	334	72.9	97.7	84.0
Cut or pierce	285	306	294	79.2	79.7	73.9
Intentional injuries	205	168	219	57.1	43.9	55.2
65 years and over ²	1,301	1,340	1,544	837.5	805.1	871.6
Unintentional injuries ⁴	1,082	1,157	1,406	696.8	695.2	793.5
Falls	644	694	838	414.5	416.7	473.0
Struck by or against objects or persons	46	*54	95	29.8	*32.2	53.6
Motor vehicle traffic	98	103	128	63.4	61.7	72.1
Cut or pierce	70	*57	90	45.3	*34.0	50.6
Intentional injuries	*	*	*	*	*	*

See footnotes at end of table.

Table 75 (page 2 of 2). Initial injury-related visits to hospital emergency departments, by sex, age, and intent and mechanism of injury: United States, average annual, selected years 2005–2006 through 2010–2011

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#075>.

[Data are based on reporting by a sample of hospital emergency departments]

<i>Sex, age, and intent and mechanism of injury</i> ¹	2005–2006	2008–2009	2010–2011	2005–2006	2008–2009	2010–2011
Female	Initial injury-related visits, in thousands			Initial injury-related visits per 10,000 persons		
All ages, age-adjusted ^{2,3}	14,740	14,688	15,524	980.5	955.6	997.2
All ages, crude ²	14,740	14,688	15,524	976.3	949.7	988.0
Unintentional injuries ⁴	11,922	12,134	12,824	789.7	784.6	816.1
Falls	4,415	4,956	5,243	292.4	320.4	333.6
Struck by or against objects or persons	1,102	1,053	1,158	73.0	68.1	73.7
Motor vehicle traffic	1,981	1,774	1,847	131.2	114.7	117.6
Cut or pierce	753	745	685	49.9	48.2	43.6
Intentional injuries	843	1,048	1,050	55.8	67.7	66.8
Under 18 years ²	3,625	3,508	3,673	1,008.7	967.5	1,013.2
Unintentional injuries ⁴	3,058	3,008	3,120	851.1	829.5	860.7
Falls	1,039	1,096	1,138	289.1	302.3	314.0
Struck by or against objects or persons	419	439	425	116.7	121.1	117.2
Motor vehicle traffic	367	249	302	102.1	68.6	83.4
Cut or pierce	160	154	158	44.4	42.4	43.7
Intentional injuries	188	222	196	52.3	61.4	54.1
18–24 years ²	1,882	1,736	1,936	1,329.3	1,194.5	1,297.1
Unintentional injuries ⁴	1,431	1,325	1,530	1,010.5	911.7	1,025.0
Falls	290	307	305	205.0	210.9	204.5
Struck by or against objects or persons	146	110	171	103.4	75.4	114.7
Motor vehicle traffic	397	360	460	280.6	247.5	308.1
Cut or pierce	116	77	*94	82.2	53.2	*63.3
Intentional injuries	176	232	251	124.2	159.7	168.4
25–44 years ²	4,173	4,087	4,233	1,004.2	996.6	1,034.6
Unintentional injuries ⁴	3,266	3,179	3,308	785.8	775.1	808.5
Falls	873	1,004	941	210.1	244.7	229.9
Struck by or against objects or persons	309	198	284	74.3	48.3	69.4
Motor vehicle traffic	719	621	616	173.1	151.3	150.5
Cut or pierce	269	270	219	64.7	65.9	53.6
Intentional injuries	313	396	408	75.4	96.5	99.8
45–64 years ²	2,904	3,061	3,101	767.8	760.0	741.9
Unintentional injuries ⁴	2,278	2,539	2,519	602.2	630.4	602.7
Falls	865	1,012	1,075	228.7	251.2	257.1
Struck by or against objects or persons	160	216	197	42.2	53.5	47.2
Motor vehicle traffic	359	399	345	94.8	99.0	82.6
Cut or pierce	158	190	157	41.7	47.2	37.6
Intentional injuries	149	161	182	39.4	39.9	43.5
65 years and over ²	2,155	2,294	2,582	1,002.9	1,016.3	1,110.7
Unintentional injuries ⁴	1,889	2,083	2,348	879.1	922.8	1,009.8
Falls	1,347	1,538	1,784	626.9	681.2	767.2
Struck by or against objects or persons	69	91	81	31.9	40.4	34.7
Motor vehicle traffic	139	146	124	64.5	64.7	53.5
Cut or pierce	*50	*54	*56	*23.3	*23.9	*24.2
Intentional injuries	*	*	*	*	*	*

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.

¹Intent and mechanism of injury are based on the first-listed external cause of injury code (E code). Intentional injuries include suicide attempts and assaults. See Appendix II, External cause of injury; Injury; Injury-related visit; Table IX for a listing of E codes.

²Includes all injury-related visits not shown separately in table, including those with undetermined intent (1% in 2010–2011) and insufficient or no information to code cause of injury (9% in 2010–2011).

³Rates are age-adjusted to the year 2000 standard population using six age groups: under 18 years, 18–24 years, 25–44 years, 45–64 years, 65–74 years, and 75 years and over. See Appendix II, Age adjustment.

⁴Includes unintentional injury-related visits with mechanism of injury not shown in table.

NOTES: An emergency department visit was considered injury-related if the first-listed diagnosis was injury-related (ICD–9–CM 800–909.2, 909.4, 909.9–994.9, 995.50–995.59, and 995.80–995.85) or the first-listed external cause code (E code) was injury-related (ICD–9–CM E800–E869, E880–E929, and E950–E999). See: http://www.cdc.gov/nchs/injury/injury_tools.htm for code used to classify injury-related visits in this table. Visits with a first-listed diagnosis or first-listed E code describing a complication or adverse effect of medical care were not considered injury related. For more information on injury-related visits, see Bergen G, Chen LH, Warner M, Fingerhut LA. Injury in the United States: 2007 Chartbook. Hyattsville, MD: NCHS. 2008. Available from: <http://www.cdc.gov/nchs/data/misc/injury2007.pdf>. Estimates for first-listed injury-related visits were further limited to those visits that were initial visits for the injury. This was determined using an imputed variable in 2005–2006; for 2007 and beyond this was determined by using the initial visit episode of care information collected on the questionnaire. Limiting the estimates to initial visits decreases the total number of injury-related visits by 9% in 2005–2006, 14% in 2007–2008, 10%–12% in 2008–2009 and 2009–2010 (shown in spreadsheet version), and 10% in 2010–2011. Rates were calculated using estimates of the civilian population of the United States including institutionalized persons. Population data are from unpublished tabulations provided by the U.S. Census Bureau. Rates for 2005–2010 were calculated using postcensal population estimates based on the 2000 census. Rates for 2011 and beyond were calculated using postcensal population estimates based on the 2010 census. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Hospital Ambulatory Medical Care Survey. See Appendix I, National Hospital Ambulatory Medical Care Survey (NHAMCS).

Table 76 (page 1 of 3). Visits to physician offices, hospital outpatient departments, and hospital emergency departments, by age, sex, and race: United States, selected years 2000–2012

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#076>.

[Data are based on reporting by a sample of office-based physicians, hospital outpatient departments, and hospital emergency departments]

Age, sex, and race	All places ¹				Physician offices			
	2000	2010	2011	2012 ²	2000	2010	2011	2012 ²
Age								
Number of visits, in thousands								
Total	1,014,848	1,239,387	1,249,047	---	823,542	1,008,802	987,029	928,630
Under 18 years	212,165	246,228	263,387	---	163,459	191,500	206,285	171,045
18–44 years	315,774	342,797	333,427	---	243,011	261,941	239,224	234,645
45–64 years	255,894	352,001	353,591	---	216,783	296,385	285,784	275,307
45–54 years	142,233	171,039	173,334	---	119,474	140,819	136,429	129,816
55–64 years	113,661	180,962	180,258	---	97,309	155,566	149,355	145,491
65 years and over	231,014	298,362	298,642	---	200,289	258,976	255,736	247,634
65–74 years	116,505	151,075	151,970	---	102,447	132,201	131,233	126,436
75 years and over	114,510	147,287	146,672	---	97,842	126,775	124,503	121,197
Number of visits per 100 persons								
Total, age-adjusted ³	374	401	400	---	304	325	314	292
Total, crude	370	408	408	---	300	332	322	301
Under 18 years	293	331	357	---	226	257	280	232
18–44 years	291	310	302	---	224	237	216	211
45–64 years	422	441	431	---	358	371	349	335
45–54 years	385	388	392	---	323	320	309	297
55–64 years	481	505	477	---	412	434	395	380
65 years and over	706	767	745	---	612	666	638	592
65–74 years	656	713	683	---	577	624	590	532
75 years and over	766	831	822	---	654	715	698	670
Sex and age								
Male, age-adjusted ³	325	350	354	---	261	283	280	254
Male, crude	314	350	356	---	251	283	281	258
Under 18 years	302	340	372	---	231	262	294	236
18–44 years	203	205	208	---	148	151	145	140
45–54 years	316	324	322	---	260	265	250	251
55–64 years	428	460	430	---	367	396	351	343
65–74 years	614	680	655	---	539	597	566	505
75 years and over	771	871	869	---	670	760	758	685
Female, age-adjusted ³	420	452	444	---	345	367	348	328
Female, crude	424	464	457	---	348	379	361	342
Under 18 years	285	322	341	---	221	252	265	229
18–44 years	377	415	393	---	298	323	286	281
45–54 years	451	450	459	---	384	372	364	340
55–64 years	529	546	520	---	453	469	436	413
65–74 years	692	741	707	---	609	647	611	556
75 years and over	763	804	790	---	645	685	657	660
Race and age ⁴								
White, age-adjusted ³	380	408	411	---	315	336	333	315
White, crude	381	421	424	---	316	349	345	330
Under 18 years	306	341	397	---	243	270	324	264
18–44 years	301	319	312	---	239	249	233	231
45–54 years	386	389	378	---	330	326	305	313
55–64 years	480	505	474	---	416	440	400	400
65–74 years	641	727	677	---	568	642	594	555
75 years and over	764	838	815	---	658	723	698	693
Black or African American, age-adjusted ³	353	439	430	---	239	316	277	229
Black or African American, crude	324	425	416	---	214	303	266	221
Under 18 years	264	351	275	---	167	241	153	168
18–44 years	257	339	350	---	149	222	199	168
45–54 years	383	466	550	---	269	339	378	252
55–64 years	495	617	566	---	373	481	412	305
65–74 years	656	715	733	---	512	565	539	421
75 years and over	745	845	821	---	568	682	605	522

See footnotes at end of table.

Table 76 (page 2 of 3). Visits to physician offices, hospital outpatient departments, and hospital emergency departments, by age, sex, and race: United States, selected years 2000–2012

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hs/content2015.htm#076>.

[Data are based on reporting by a sample of office-based physicians, hospital outpatient departments, and hospital emergency departments]

Age, sex, and race	Hospital outpatient departments				Hospital emergency departments				
	2000	2010	2011	2012	2000	2010	2011	2012	
Age		Number of visits, in thousands							
Total	83,289	100,742	125,721	---	108,017	129,843	136,296	---	
Under 18 years	21,076	24,913	27,651	---	27,630	29,815	29,451	---	
18–44 years	26,947	28,159	37,557	---	45,816	52,697	56,646	---	
45–64 years	20,772	27,739	37,980	---	18,339	27,877	29,828	---	
45–54 years	11,558	13,639	19,310	---	11,201	16,581	17,595	---	
55–64 years	9,214	14,100	18,670	---	7,138	11,296	12,232	---	
65 years and over	14,494	19,932	22,534	---	16,232	19,454	20,372	---	
65–74 years	7,515	10,675	12,529	---	6,543	8,199	8,208	---	
75 years and over	6,979	9,257	10,005	---	9,690	11,255	12,163	---	
		Number of visits per 100 persons							
Total, age-adjusted ³	31	33	40	---	40	43	45	---	
Total, crude	30	33	41	---	39	43	44	---	
Under 18 years	29	33	37	---	38	40	40	---	
18–44 years	25	25	34	---	42	48	51	---	
45–64 years	34	35	46	---	30	35	36	---	
45–54 years	31	31	44	---	30	38	40	---	
55–64 years	39	39	49	---	30	32	32	---	
65 years and over	44	51	56	---	50	50	51	---	
65–74 years	42	50	56	---	37	39	37	---	
75 years and over	47	52	56	---	65	64	68	---	
Sex and age									
Male, age-adjusted ³	26	27	32	---	38	40	42	---	
Male, crude	25	27	33	---	38	39	41	---	
Under 18 years	29	34	37	---	41	43	41	---	
18–44 years	17	16	20	---	38	38	43	---	
45–54 years	26	24	34	---	30	35	38	---	
55–64 years	32	32	45	---	30	32	34	---	
65–74 years	38	47	52	---	36	37	37	---	
75 years and over	42	50	49	---	59	60	62	---	
Female, age-adjusted ³	35	38	48	---	41	47	48	---	
Female, crude	35	39	49	---	41	46	48	---	
Under 18 years	29	33	38	---	35	37	39	---	
18–44 years	33	35	47	---	46	57	59	---	
45–54 years	36	37	53	---	31	40	41	---	
55–64 years	45	46	54	---	31	31	31	---	
65–74 years	46	54	60	---	37	40	37	---	
75 years and over	49	53	61	---	69	66	72	---	
Race and age⁴									
White, age-adjusted ³	28	31	37	---	37	41	42	---	
White, crude	28	32	38	---	37	40	41	---	
Under 18 years	27	33	37	---	36	39	37	---	
18–44 years	23	25	31	---	39	45	47	---	
45–54 years	28	28	37	---	28	34	35	---	
55–64 years	36	36	44	---	28	29	30	---	
65–74 years	38	48	49	---	35	37	34	---	
75 years and over	44	52	52	---	63	62	65	---	
Black or African American, age-adjusted ³	51	51	69	---	62	73	85	---	
Black or African American, crude	48	50	68	---	62	72	83	---	
Under 18 years	40	48	*50	---	57	62	72	---	
18–44 years	40	37	55	---	68	81	96	---	
45–54 years	61	54	89	---	53	73	83	---	
55–64 years	70	73	94	---	52	62	60	---	
65–74 years	85	*85	*121	---	59	66	73	---	
75 years and over	85	*74	*98	---	92	89	118	---	

See footnotes at end of table.

Table 76 (page 3 of 3). Visits to physician offices, hospital outpatient departments, and hospital emergency departments, by age, sex, and race: United States, selected years 2000–2012

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#076>.

[Data are based on reporting by a sample of office-based physicians, hospital outpatient departments, and hospital emergency departments]

-- Data not available.

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.

¹All places includes visits to physician offices and hospital outpatient and emergency departments. See Appendix II, Emergency department; Emergency department or emergency room visit; Office visit; Outpatient department; Outpatient visit.

²In 2012, data for all places and physician offices exclude visits to community health centers; in 2006–2011, data for all places and physician offices include visits to community health centers (2%–3% of visits to physician offices in 2006–2011 were to community health centers). Prior to 2006, visits to community health centers were not included in the survey.

³Estimates are age-adjusted to the year 2000 standard population using six age groups: under 18 years, 18–44 years, 45–54 years, 55–64 years, 65–74 years, and 75 years and over. See Appendix II, Age adjustment.

⁴Estimates by racial group should be used with caution because information on race was collected from medical records and race is imputed for records missing that information. Information on the race imputation process used in each data year is available in the public-use file documentation. Available from:

<http://www.cdc.gov/nchs/ahcd.htm>. Starting with 1999 data, the instruction for the race item on the Patient Record Form was changed so that more than one race could be recorded. In previous years only one race could be recorded. Estimates for race in this table are for visits where only one race was recorded. Because of the small number of responses with more than one racial group recorded, estimates for visits with multiple races recorded are unreliable and are not presented.

NOTES: Rates for 1995–2000 were computed using 1990-based postcensal estimates of the civilian noninstitutionalized population as of July 1, adjusted for net underenumeration using the 1990 National Population Adjustment Matrix from the U.S. Census Bureau. For 2001–2010 data, rates were computed using 2000-based postcensal estimates of the civilian noninstitutionalized population as of July 1. For 2011 data and beyond, rates were computed using 2010-based postcensal estimates of the civilian noninstitutionalized population as of July 1. Rates using the civilian noninstitutionalized population will be overestimated to the extent that visits by institutionalized persons are counted in the numerator (for example, hospital emergency department visits by nursing home residents) but institutionalized persons are omitted from the denominator (the civilian noninstitutionalized population). Starting with *Health, United States, 2005*, data for physician offices for 2001 and beyond use a revised weighting scheme. See Appendix I, National Ambulatory Medical Care Survey (NAMCS); National Hospital Ambulatory Medical Care Survey (NHAMCS). Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Ambulatory Medical Care Survey and National Hospital Ambulatory Medical Care Survey. See Appendix I, National Ambulatory Medical Care Survey (NAMCS); National Hospital Ambulatory Medical Care Survey (NHAMCS).

Table 77 (page 1 of 2). Visits to primary care generalist and specialty care physicians, by selected characteristics and type of physician: United States, selected years 1980–2012

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#077>.

[Data are based on reporting by a sample of office-based physicians]

Age, sex, and race	Type of primary care generalist physician ¹											
	All primary care generalists				General and family practice				Internal medicine			
	1980	2000	2010	2012 ²	1980	2000	2010	2012 ²	1980	2000	2010	2012 ²
Age						Percent distribution						
Total	66.2	58.9	55.2	53.2	33.5	24.1	21.1	20.3	12.1	15.3	13.9	12.7
Under 18 years	77.8	79.7	80.9	80.7	26.1	19.9	15.3	12.4	2.0	*	*	*1.3
18–44 years	65.3	62.1	62.7	57.4	34.3	28.2	27.8	24.2	8.6	12.7	11.6	10.2
45–64 years	60.2	51.2	46.7	44.6	36.3	26.4	23.1	23.0	19.5	20.1	18.5	16.3
45–54 years	60.2	52.3	48.7	47.0	37.4	27.8	26.2	24.4	17.1	18.7	15.7	15.2
55–64 years	60.2	49.9	44.8	42.5	35.4	24.7	20.4	21.7	21.8	21.7	21.0	17.2
65 years and over	61.6	46.5	38.3	39.7	37.5	20.2	16.4	19.0	22.7	24.5	20.5	19.0
65–74 years	61.2	46.6	37.3	39.8	37.4	19.7	17.5	20.0	22.1	24.5	18.2	17.6
75 years and over	62.3	46.4	39.2	39.6	37.6	20.8	15.4	18.0	23.5	24.5	22.8	20.4
Sex and age												
Male:												
Under 18 years	77.3	77.7	80.1	78.6	25.6	18.3	15.7	12.3	2.0	*	*	*1.0
18–44 years	50.8	51.5	51.7	46.1	38.0	34.2	33.7	29.9	11.5	14.4	16.4	14.3
45–64 years	55.6	49.4	43.7	43.5	34.4	28.7	24.4	25.2	20.5	19.8	19.1	18.2
65 years and over	58.2	43.1	36.6	37.5	35.6	19.3	16.2	18.9	22.3	23.8	20.3	18.5
Female:												
Under 18 years	78.5	82.0	81.7	82.9	26.6	21.7	14.9	12.6	2.0	*	*	*1.7
18–44 years	72.1	67.2	67.9	62.8	32.5	25.3	25.0	21.5	7.3	11.9	9.4	8.2
45–64 years	63.4	52.5	48.9	45.4	37.7	24.9	22.2	21.3	18.9	20.2	18.1	14.8
65 years and over	63.9	48.9	39.6	41.3	38.7	20.9	16.7	19.1	22.9	25.0	20.5	19.3
Race and age ³												
White:												
Under 18 years	77.6	78.5	79.6	80.5	26.4	21.2	15.6	13.0	2.0	*	*	*1.4
18–44 years	64.8	61.4	61.2	56.4	34.5	29.2	27.9	24.8	8.6	11.0	11.1	9.6
45–64 years	59.6	49.3	45.2	43.7	36.0	27.3	22.8	23.4	19.2	17.1	17.5	14.9
65 years and over	61.4	45.1	37.6	39.4	36.6	20.3	16.6	19.4	23.3	23.0	19.7	18.2
Black or African American:												
Under 18 years	79.9	87.3	88.0	81.0	23.7	*	*16.5	10.2	*2.2	*	*	*
18–44 years	68.5	65.0	72.6	61.1	31.7	22.0	29.4	23.1	9.0	20.9	*14.0	12.2
45–64 years	66.1	61.7	57.0	48.7	38.6	23.3	26.7	24.7	22.6	35.9	24.5	18.7
65 years and over	64.6	52.8	45.2	41.5	49.0	*18.5	*18.6	18.2	14.2	33.4	*25.4	21.8

See footnotes at end of table.

Table 77 (page 2 of 2). Visits to primary care generalist and specialty care physicians, by selected characteristics and type of physician: United States, selected years 1980–2012

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#077>.

[Data are based on reporting by a sample of office-based physicians]

Age, sex, and race	Type of primary care generalist physician ¹								Specialty care physicians			
	Obstetrics and gynecology				Pediatrics				1980	2000	2010	2012 ²
	1980	2000	2010	2012 ²	1980	2000	2010	2012 ²				
Age					Percent distribution							
Total	9.6	7.8	7.8	7.6	10.9	11.7	12.4	12.5	33.8	41.1	44.8	46.8
Under 18 years	1.3	*1.1	*1.3	0.8	48.5	57.3	63.4	66.1	22.2	20.3	19.1	19.3
18–44 years	21.7	20.4	22.3	21.6	0.7	*0.9	1.0	1.3	34.7	37.9	37.3	42.6
45–64 years	4.2	4.5	4.9	5.3	*	*	*	*	39.8	48.8	53.3	55.4
45–54 years	5.6	5.6	6.7	7.2	*	*	*	*	39.8	47.7	51.3	53.0
55–64 years	2.9	3.3	3.3	3.5	*	*	*	*	39.8	50.1	55.2	57.5
65 years and over	1.4	1.5	1.3	1.7	*	*	*	*	38.4	53.5	61.7	60.3
65–74 years	1.7	2.0	1.7	2.1	*	*	*	*	38.8	53.4	62.7	60.2
75 years and over	1.0	*1.0	*1.0	1.2	*	*	*	*	37.7	53.6	60.8	60.4
Sex and age												
Male:												
Under 18 years	49.4	58.0	63.7	65.2	22.7	22.3	19.9	21.4
18–44 years	1.0	*1.7	*1.4	1.8	49.2	48.5	48.3	53.9
45–64 years	*	*	*	*	44.4	50.6	56.3	56.5
65 years and over	*	*	*	*	41.8	56.9	63.4	62.5
Female:												
Under 18 years	2.5	2.1	*2.8	1.6	47.4	56.5	63.1	67.0	21.5	18.0	18.3	17.1
18–44 years	31.7	29.6	32.5	32.0	0.6	*	*0.9	1.1	27.9	32.8	32.1	37.2
45–64 years	6.7	7.3	8.5	9.0	*	*	*	*	36.6	47.5	51.1	54.6
65 years and over	2.1	2.6	2.4	2.9	*	*	*	*	36.1	51.1	60.4	58.7
Race and age ³												
White:												
Under 18 years	1.1	*1.2	*1.3	0.8	48.2	54.7	61.7	65.3	22.4	21.5	20.4	19.5
18–44 years	21.0	20.4	21.1	20.6	0.7	*0.8	*1.1	1.4	35.2	38.6	38.8	43.6
45–64 years	4.1	4.7	4.7	5.2	*	*	*	*	40.4	50.7	54.8	56.3
65 years and over	1.4	1.5	*1.3	1.7	*	*	*	*	38.6	54.9	62.4	60.6
Black or African American:												
Under 18 years	2.8	*	*	*	51.2	75.0	70.2	67.9	20.1	*12.7	*12.0	19.0
18–44 years	27.1	20.7	28.4	24.9	*	*	*	*	31.5	35.0	27.4	38.9
45–64 years	4.8	*2.4	*5.6	5.3	*	*	*	*	33.9	38.3	43.0	51.3
65 years and over	*	*	*1.2	*	*	*	*	0.0	35.4	47.2	54.8	58.5

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have a RSE greater than 30%. ... Category not applicable.

¹Type of physician is based on physician's self-designated primary area of practice. Primary care generalist physicians are defined as practitioners in the fields of general and family practice, general internal medicine, general obstetrics and gynecology, and general pediatrics and exclude primary care specialists. Primary care generalists in general and family practice exclude primary care specialties, such as sports medicine and geriatrics. Primary care internal medicine physicians exclude internal medicine specialists, such as allergists, cardiologists, and endocrinologists. Primary care obstetrics and gynecology physicians exclude obstetrics and gynecology specialties, such as gynecological oncology, maternal and fetal medicine, obstetrics and gynecology critical care medicine, and reproductive endocrinology. Primary care pediatricians exclude pediatric specialists, such as adolescent medicine specialists, neonatologists, pediatric allergists, and pediatric cardiologists. See Appendix II, Physician specialty.

²In 2012, data exclude visits to community health centers; in 2006–2011, data include visits to community health centers (2%–3% of visits to physician offices in 2006–2011 were to community health centers). Prior to 2006, visits to community health centers were not included in the survey.

³Estimates by racial group should be used with caution because information on race was collected from medical records. In 2012, race data were missing and imputed for 34% of visits. Information on the race imputation process used in each data year is available in the public-use file documentation. Available from: <http://www.cdc.gov/nchs/ahcd.htm>. Starting with 1999 data, the instruction for the race item on the Patient Record Form was changed so that more than one race could be recorded. In previous years only one racial category could be checked. Estimates for racial groups presented in this table are for visits where only one race was recorded. Because of the small number of responses with more than one racial group checked, estimates for visits with multiple races checked are unreliable and are not presented.

NOTES: This table presents data on visits to physician offices and excludes visits to other sites, such as hospital outpatient and emergency departments. See Appendix II, Office visit. In 1980, the survey excluded Alaska and Hawaii. Data for all other years include all 50 states and the District of Columbia. Visits with specialty of physician unknown are excluded. Starting with *Health, United States, 2005*, data for 2001 and later years for physician offices use a revised weighting scheme. See Appendix I, National Ambulatory Medical Care Survey (NAMCS). Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Ambulatory Medical Care Survey. See Appendix I, National Ambulatory Medical Care Survey (NAMCS).

Table 78 (page 1 of 2). Dental visits in the past year, by selected characteristics: United States, selected years 1997–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#078>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	2 years and over			2–17 years			18–64 years			65 years and over ¹		
	1997	2010	2014	1997	2010	2014	1997	2010	2014	1997	2010	2014
Percent of persons with a dental visit in the past year ²												
Total ³	65.1	64.7	66.6	72.7	78.9	83.0	64.1	61.1	62.0	54.8	57.7	62.4
Sex												
Male	62.9	61.7	64.1	72.3	78.3	82.3	60.4	56.8	58.1	55.4	56.2	62.2
Female	67.1	67.5	68.9	73.0	79.6	83.8	67.7	65.4	65.8	54.4	58.9	62.5
Race ⁴												
White only	66.4	65.6	67.7	74.0	79.2	83.4	65.7	62.4	63.3	56.8	59.3	64.9
Black or African American only	58.9	58.8	60.7	68.8	79.0	83.0	57.0	53.1	54.8	35.4	40.6	42.7
American Indian or Alaska Native only	55.1	57.4	59.4	66.8	73.2	88.6	49.9	49.8	46.8	*	72.2	51.8
Asian only	62.5	66.5	66.5	69.9	74.8	78.9	60.3	64.6	64.8	53.9	61.9	55.0
Native Hawaiian or Other Pacific Islander only	---	*	*	---	*	*	---	*	*	---	*	*
2 or more races	---	65.2	66.4	---	77.9	79.2	---	54.7	56.8	---	48.1	53.8
Black or African American; White	---	72.5	72.3	---	78.4	77.6	---	62.1	62.3	---	*	78.4
American Indian or Alaska Native; White	---	54.7	54.9	---	70.0	81.2	---	49.0	45.5	---	*54.5	45.1
Hispanic origin and race ⁴												
Hispanic or Latino	54.0	56.5	59.7	61.0	74.8	81.4	50.8	48.5	50.2	47.8	42.1	51.3
Not Hispanic or Latino	66.4	66.2	68.0	74.7	80.1	83.5	65.7	63.4	64.5	55.2	59.0	63.3
White only	68.0	67.6	69.7	76.4	80.9	84.4	67.5	65.4	66.5	57.2	60.9	66.2
Black or African American only	58.8	58.7	60.6	68.8	79.2	83.0	56.9	53.1	55.1	35.3	40.5	42.6
Percent of poverty level ⁵												
Below 100%	50.5	50.6	52.5	62.0	73.2	78.2	46.9	41.0	41.9	31.5	32.8	35.1
100%–199%	50.8	51.6	54.4	62.5	73.4	78.2	48.3	44.1	47.1	40.8	43.8	44.1
200%–399%	66.2	63.5	65.8	76.1	79.0	83.6	63.4	59.6	60.9	60.7	57.9	62.1
400% or more	78.9	79.3	80.4	85.7	88.0	90.5	77.7	77.5	77.8	74.7	77.2	81.5
Hispanic origin and race and percent of poverty level ^{4,5}												
Hispanic or Latino:												
Below 100%	45.7	50.8	54.2	55.9	74.3	80.3	39.2	34.7	37.6	33.6	32.4	39.3
100%–199%	47.2	50.8	55.1	53.8	71.1	78.3	43.5	40.2	43.7	47.9	39.5	46.8
200%–399%	61.2	59.1	62.7	70.5	76.5	83.9	57.5	54.1	55.5	57.0	46.0	55.2
400% or more	73.0	73.3	74.0	82.4	84.2	89.4	70.8	71.6	69.6	64.9	54.3	74.3
Not Hispanic or Latino:												
White only:												
Below 100%	51.7	49.3	51.0	64.4	69.1	73.8	50.6	44.4	45.4	32.0	36.4	36.0
100%–199%	52.4	52.7	53.1	66.1	75.3	78.2	50.4	47.2	47.4	42.2	45.4	45.7
200%–399%	67.5	64.7	67.1	77.1	79.6	83.7	65.0	61.4	62.7	61.9	59.8	64.5
400% or more	79.7	79.8	81.8	86.8	88.6	91.6	78.5	77.9	79.3	75.5	78.8	83.2
Black or African American only:												
Below 100%	52.8	52.0	52.5	66.1	78.0	82.0	46.2	39.7	39.2	27.7	20.9	27.4
100%–199%	48.7	50.0	57.2	61.2	75.9	81.5	46.3	41.5	51.7	26.9	33.6	32.9
200%–399%	63.3	61.2	61.5	75.0	81.2	84.0	60.7	57.2	57.3	41.5	45.3	43.6
400% or more	74.6	77.2	75.6	81.8	87.2	88.8	73.4	75.9	74.8	66.1	69.8	66.8

See footnotes at end of table.

Table 78 (page 2 of 2). Dental visits in the past year, by selected characteristics: United States, selected years 1997–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#078>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	2 years and over			2–17 years			18–64 years			65 years and over ¹		
	1997	2010	2014	1997	2010	2014	1997	2010	2014	1997	2010	2014
Disability measure ⁶ Percent of persons with a dental visit in the past year ²												
Any basic actions difficulty or complex activity limitation	55.1	53.5	52.4	49.0	50.7	56.1
Any basic actions difficulty	54.7	53.2	52.6	48.7	50.5	56.0
Any complex activity limitation	51.0	47.4	48.3	44.6	43.1	49.4
No disability	67.4	64.2	65.3	64.2	68.8	71.7
Geographic region												
Northeast	69.6	70.1	71.9	77.5	83.8	84.6	69.6	67.9	69.2	55.5	61.5	67.7
Midwest	68.4	67.3	68.1	76.4	80.8	82.5	67.4	64.3	64.4	57.6	58.2	63.3
South	60.2	60.9	62.6	68.0	77.4	82.4	59.4	56.5	56.7	49.0	54.1	58.7
West	65.0	63.9	67.7	71.5	76.1	83.3	62.9	60.2	63.1	61.9	59.8	62.9
Location of residence ⁷												
Within MSA	66.7	65.9	67.7	73.6	79.3	83.4	65.7	62.4	63.2	57.6	59.4	64.2
Outside MSA	59.1	58.4	59.8	69.3	76.4	80.6	58.0	53.8	54.4	46.1	51.3	54.8

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.

--- Data not available.

... Category not applicable.

¹Based on the 1997–2014 National Health Interview Surveys, about 21%–30% of persons aged 65 and over were edentulous (having lost all their natural teeth). In 1997–2014, about 69%–73% of older dentate persons, compared with 17%–24% of older edentate persons, had a dental visit in the past year.

²Respondents were asked, “About how long has it been since you last saw or talked to a dentist?” See Appendix II, Dental visit.

³Includes all other races not shown separately and unknown disability status.

⁴The race groups, white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The five single-race categories plus multiple-race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group; the category 2 or more races includes persons who reported more than one racial group. Prior to 1999, data were tabulated according to the 1977 Standards with four racial groups, and the Asian only category included Native Hawaiian or Other Pacific Islander. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

⁵Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Missing family income data were imputed for 1997 and beyond. See Appendix II, Family income; Poverty; Table VI.

⁶Any basic actions difficulty or complex activity limitation is defined as having one or more of the following limitations or difficulties: movement difficulty, emotional difficulty, sensory (seeing or hearing) difficulty, cognitive difficulty, self-care (activities of daily living or instrumental activities of daily living) limitation, social limitation, or work limitation. For more information, see Appendix II, Basic actions difficulty; Complex activity limitation. Starting with 2007 data, the hearing question, a component of the basic actions difficulty measure, was revised. Consequently, data prior to 2007 are not comparable with data for 2007 and beyond. For more information on the impact of the revised hearing question, see Appendix II, Hearing trouble.

⁷MSA is metropolitan statistical area. Starting with 2006 data, MSA status is determined using 2000 census data and the 2000 standards for defining MSAs. For data prior to 2006, see Appendix II, Metropolitan statistical area (MSA) for the applicable standards.

NOTES: Standard errors for selected years are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey, sample child and sample adult questionnaires. See Appendix I, National Health Interview Survey (NHIS).

Table 79 (page 1 of 2). Prescription drug use in the past 30 days, by sex, race and Hispanic origin, and age: United States, selected years 1988–1994 through 2009–2012

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#079>.

[Data are based on a sample of the civilian noninstitutionalized population]

Sex, race and Hispanic origin ¹ , and age	At least one prescription drug in past 30 days				Three or more prescription drugs in past 30 days				Five or more prescription drugs in past 30 days			
	1988– 1994	1999– 2002	2003– 2006	2009– 2012	1988– 1994	1999– 2002	2003– 2006	2009– 2012	1988– 1994	1999– 2002	2003– 2006	2009– 2012
All ages, age-adjusted ²												
Percent of population												
Both sexes ³	39.1	45.2	46.9	47.3	11.8	17.8	21.0	20.6	4.0	7.5	10.0	10.1
Male	32.7	39.8	41.7	42.7	9.4	14.8	18.0	19.1	2.9	6.1	8.4	9.3
Female	45.0	50.3	51.9	51.8	13.9	20.4	23.8	22.0	4.9	8.7	11.4	10.8
Not Hispanic or Latino:												
White only	41.1	48.7	50.7	52.4	12.4	18.9	22.7	22.0	4.2	7.8	10.5	10.6
White only, male	34.2	43.0	45.1	47.2	9.9	15.9	19.2	20.2	3.1	6.3	8.7	9.6
White only, female	47.6	54.3	56.3	57.6	14.6	21.8	26.1	23.8	5.1	9.2	12.3	11.5
Black or African American only	36.9	40.1	41.8	43.7	12.6	16.5	19.2	21.9	3.8	7.7	10.3	10.9
Black or African American only, male	31.1	35.4	37.0	37.5	10.2	14.5	17.2	19.2	2.9	6.4	9.4	9.5
Black or African American only, female	41.4	43.8	45.7	48.8	14.3	18.1	20.7	24.0	4.5	8.7	11.1	12.0
Hispanic or Latino	---	---	---	35.4	---	---	---	15.9	---	---	---	8.4
Hispanic or Latino, male	---	---	---	31.7	---	---	---	14.7	---	---	---	7.8
Hispanic or Latina, female	---	---	---	39.2	---	---	---	17.1	---	---	---	8.9
Mexican origin	31.7	31.7	33.3	34.0	9.0	11.2	14.6	15.4	2.9	4.4	6.3	8.3
Mexican origin, male	27.5	25.8	28.4	30.7	7.0	9.5	11.3	14.0	2.0	3.5	4.7	7.6
Mexican origin, female	36.0	37.8	38.2	37.6	11.0	12.8	17.7	16.8	3.7	5.2	7.8	8.9
All ages, crude												
Both sexes ³	37.8	45.0	47.3	48.7	11.0	17.6	21.3	21.8	3.6	7.4	10.1	10.7
Male	30.6	38.6	41.1	43.4	8.3	13.9	17.4	19.4	2.5	5.6	8.0	9.3
Female	44.6	51.1	53.2	53.9	13.6	21.1	25.1	24.1	4.7	9.1	12.2	12.0
Not Hispanic or Latino:												
White only	41.4	50.7	53.5	56.5	12.5	20.6	25.3	25.9	4.2	8.7	12.0	12.8
White only, male	33.5	43.8	46.8	50.7	9.5	16.5	20.7	23.1	2.9	6.6	9.5	11.1
White only, female	48.9	57.5	59.9	62.1	15.4	24.5	29.7	28.5	5.4	10.8	14.5	14.4
Black or African American only	31.2	36.0	38.7	42.0	9.2	13.5	16.6	20.2	2.6	6.2	8.7	9.8
Black or African American only, male	25.5	30.7	32.9	35.1	7.0	10.9	14.0	16.9	1.8	4.8	7.3	8.1
Black or African American only, female	36.2	40.6	43.7	48.0	11.1	15.7	18.8	23.1	3.3	7.4	9.9	11.3
Hispanic or Latino	---	---	---	28.9	---	---	---	10.6	---	---	---	5.0
Hispanic or Latino, male	---	---	---	25.0	---	---	---	9.0	---	---	---	4.4
Hispanic or Latina, female	---	---	---	33.1	---	---	---	12.2	---	---	---	5.7
Mexican origin	24.0	23.6	25.2	26.4	4.8	6.1	8.5	9.2	1.4	2.1	3.3	4.4
Mexican origin, male	20.1	18.8	21.1	23.5	3.4	4.8	6.1	8.3	0.9	1.6	2.2	4.1
Mexican origin, female	28.1	28.9	29.8	29.5	6.4	7.5	11.1	10.2	1.9	2.7	4.4	*4.6
Both sexes												
Under 18 years	20.5	23.8	24.7	23.5	2.4	4.1	4.0	3.6	*	*0.8	*0.9	0.8
18–44 years	31.3	35.9	37.4	38.1	5.7	8.4	10.6	9.6	1.2	2.3	3.5	3.3
45–64 years	54.8	64.1	65.2	67.2	20.0	30.8	35.3	34.7	7.4	13.3	17.1	16.3
65 years and over	73.6	84.7	89.4	89.8	35.3	51.8	62.9	64.8	13.8	27.1	36.1	39.1
Male												
Under 18 years	20.4	25.7	25.3	23.1	2.6	4.3	4.0	4.1	*	*	*1.0	0.9
18–44 years	21.5	27.1	28.0	29.6	3.6	6.7	6.9	7.5	*0.8	1.7	2.1	2.8
45–64 years	47.2	55.6	58.4	63.1	15.1	23.6	29.8	31.4	4.8	9.5	13.4	14.2
65 years and over	67.2	80.1	88.2	87.7	31.3	46.3	60.5	64.6	11.3	24.7	34.5	37.9
Female												
Under 18 years	20.6	21.7	24.0	23.8	2.3	3.9	4.0	3.1	*	*0.8	*0.7	*0.6
18–44 years	40.7	44.6	46.6	46.4	7.6	10.2	14.3	11.8	1.7	2.8	4.9	3.9
45–64 years	62.0	72.0	71.6	71.1	24.7	37.5	40.6	37.8	9.7	16.8	20.7	18.3
65 years and over	78.3	88.1	90.3	91.4	38.2	55.9	64.7	64.9	15.6	28.9	37.4	40.0

See footnotes at end of table.

Table 79 (page 2 of 2). Prescription drug use in the past 30 days, by sex, race and Hispanic origin, and age: United States, selected years 1988–1994 through 2009–2012

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#079>.

[Data are based on a sample of the civilian noninstitutionalized population]

- - - Data not available.

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.

¹Persons of Hispanic and Mexican origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The non-Hispanic race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group. Prior to data year 1999, estimates were tabulated according to the 1977 Standards. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. See Appendix II, Hispanic origin; Race.

²Estimates are age-adjusted to the year 2000 standard population using four age groups: under 18 years, 18–44 years, 45–64 years, and 65 years and over. Age-adjusted estimates in this table may differ from other age-adjusted estimates based on the same data and presented elsewhere if different age groups are used in the adjustment procedure. See Appendix II, Age adjustment.

³Includes persons of all races and Hispanic origins, not just those shown separately.

NOTES: See Appendix II, Drug. Standard errors are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health and Nutrition Examination Survey. See Appendix I, National Health and Nutrition Examination Survey (NHANES).

Table 80 (page 1 of 3). Selected prescription drug classes used in the past 30 days, by sex and age: United States, selected years 1988–1994 through 2009–2012

Updated data when available, Excel, PDF, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#080>.

[Data are based on a sample of the civilian noninstitutionalized population]

Age group and Multum Lexicon Plus therapeutic class ¹ (common indications for use)	Total			Male			Female		
	1988–1994	1999–2002	2009–2012	1988–1994	1999–2002	2009–2012	1988–1994	1999–2002	2009–2012
All ages									
Percent of population with at least one prescription drug in drug class in past 30 days									
Antihyperlipidemic agents (high cholesterol)	1.7	6.5	13.1	1.5	7.1	13.9	1.8	5.8	12.3
Analgesics (pain relief)	7.2	9.4	8.8	5.4	7.3	7.5	9.0	11.3	10.0
Antidepressants (depression and related disorders)	1.8	6.4	9.0	1.2	4.4	6.0	2.3	8.3	11.8
Proton pump inhibitors or H2 antagonists (gastric reflux, ulcers) ²	2.8	5.3	8.2	2.4	4.7	7.3	3.0	5.9	9.1
Beta-adrenergic blocking agents (high blood pressure, heart disease)	3.1	4.4	7.7	2.7	4.1	7.3	3.5	4.6	8.1
ACE inhibitors (high blood pressure, heart disease)	2.4	4.6	6.8	2.4	4.7	7.6	2.4	4.5	6.1
Antidiabetic agents (diabetes)	2.6	3.7	5.9	2.5	3.7	6.2	2.6	3.8	5.6
Diuretics (high blood pressure, heart disease, kidney disease) ³	3.4	4.1	5.6	2.3	3.1	4.5	4.4	5.1	6.6
Thyroid hormones (hypothyroidism)	2.3	3.9	5.0	0.8	1.5	2.3	3.7	6.2	7.5
Bronchodilators (asthma, breathing)	2.6	3.5	5.0	2.5	3.1	4.6	2.7	3.8	5.3
Sex hormones (contraceptives, menopause, hot flashes) ⁴	9.8	15.2	8.6
Anxiolytics, sedatives, and hypnotics (anxiety, insomnia, and related disorders)	2.8	3.3	4.6	1.9	2.6	3.8	3.6	4.0	5.4
Antihypertensive combinations (high blood pressure)	2.4	2.9	4.3	1.4	1.9	3.7	3.3	3.8	4.8
Anticonvulsants (epilepsy, seizure, and related disorders)	1.4	2.4	4.1	1.2	2.1	3.5	1.6	2.7	4.6
Calcium channel blocking agents (high blood pressure, heart disease)	3.6	4.2	4.0	3.4	3.5	3.9	3.8	4.8	4.1
Under 18 years									
Bronchodilators (asthma, breathing)	3.0	4.0	5.1	3.3	4.4	5.7	2.7	3.6	4.4
CNS stimulants (attention-deficit/hyperactivity disorder)	*0.8	2.9	3.5	*1.2	4.4	5.0	*	1.4	1.9
Penicillins (bacterial infections)	6.1	5.1	3.3	5.9	5.2	2.7	6.4	5.0	3.8
Leukotriene modifiers (asthma, allergies)	0.7	2.0	...	*0.9	2.3	...	*	1.7
Antihistamines (allergies)	2.0	4.4	1.7	2.1	4.9	1.9	1.9	3.9	1.6
Respiratory inhalant products (asthma, chronic obstructive pulmonary disease, and related disorders)	*0.7	1.5	2.0	*	1.7	2.4	*	1.3	1.5
Adrenal cortical steroids (anti-inflammatory)	*0.5	0.8	1.4	*	*0.7	1.5	*0.5	0.9	*1.2
Nasal preparations (nose symptoms)	*	1.1	1.6	*	*1.3	1.9	*	1.0	*1.3
Antidepressants (depression and related disorders)	*	1.8	*1.2	*	2.2	*	*	*1.5	*
Upper respiratory combinations (cough and cold, congestion)	2.3	2.3	*0.8	2.6	*2.4	*0.9	2.0	*2.2	*
Analgesics (pain relief)	1.2	1.4	1.2	*1.2	1.3	*1.2	1.4	1.6	*1.1
Dermatological agents (skin symptoms)	0.7	1.1	1.4	*	1.1	*1.2	*1.0	*1.1	1.7
18–44 years									
Analgesics (pain relief)	7.2	8.0	7.6	5.1	6.0	5.8	9.1	9.9	9.5
Antidepressants (depression and related disorders)	1.6	6.0	8.4	*1.0	3.6	5.9	2.3	8.5	10.8
Sex hormones (contraceptives, menopause, hot flashes) ⁴	11.5	13.5	14.0
Proton pump inhibitors or H2 antagonists (gastric reflux, ulcers) ²	2.0	3.0	4.8	1.6	3.0	5.1	2.4	3.0	4.4
Anxiolytics, sedatives, and hypnotics (anxiety, insomnia, and related disorders)	1.4	2.1	4.0	*1.0	*1.7	3.2	1.9	2.5	4.7
Anticonvulsants (epilepsy, seizure, and related disorders)	0.8	1.6	3.2	*0.6	1.6	2.9	1.0	*1.5	3.6
Bronchodilators (asthma, breathing)	1.4	2.2	3.5	*1.1	1.6	2.7	*1.8	2.8	4.2
Antihyperlipidemic agents (high cholesterol)	*0.4	1.3	2.3	*	2.0	*2.8	*	*	1.8
Antihistamines (allergies)	2.5	3.9	1.9	1.8	3.6	*1.5	3.2	4.2	2.2
Thyroid hormones (hypothyroidism)	1.3	1.6	1.9	*	*	*0.8	2.1	2.8	3.1
ACE inhibitors (high blood pressure, heart disease)	0.7	1.4	1.8	*0.9	1.5	2.1	*0.6	*1.2	1.6
Antidiabetic agents (diabetes)	*1.0	1.5	2.1	*	*1.5	2.0	*1.0	*1.6	2.1
Muscle relaxants (muscle spasm and related disorders)	1.0	1.3	1.7	*1.3	*1.1	*1.5	*0.7	*1.4	1.9
Beta-adrenergic blocking agents (high blood pressure, heart disease)	1.1	*1.2	1.7	*0.9	*1.3	1.3	1.3	*	2.1
Nasal preparations (nose symptoms)	*0.6	1.5	1.5	*	*1.2	*1.1	*0.7	1.7	1.8

See footnotes at end of table.

Table 80 (page 2 of 3). Selected prescription drug classes used in the past 30 days, by sex and age: United States, selected years 1988–1994 through 2009–2012

Updated data when available, Excel, PDF, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#080>.

[Data are based on a sample of the civilian noninstitutionalized population]

Age group and Multum Lexicon Plus therapeutic class ¹ (common indications for use)	Total			Male			Female		
	1988–1994	1999–2002	2009–2012	1988–1994	1999–2002	2009–2012	1988–1994	1999–2002	2009–2012
45–64 years									
Percent of population with at least one prescription drug in drug class in past 30 days									
Antihyperlipidemic agents (high cholesterol)	4.3	13.8	23.8	4.4	17.2	26.3	4.2	10.7	21.4
Proton pump inhibitors or H2 antagonists (gastric reflux, ulcers) ²	5.2	9.9	13.2	5.3	8.4	11.7	5.2	11.3	14.6
Antidepressants (depression and related disorders)	3.5	10.5	14.3	*2.3	7.0	9.6	4.6	13.8	18.6
Sex hormones (contraceptives, menopause, hot flashes) ⁴	19.9	30.3	9.3
Analgesics (pain relief)	11.9	16.0	14.2	9.2	13.5	13.2	14.3	18.3	15.2
Beta-adrenergic blocking agents (high blood pressure, heart disease)	6.6	8.7	11.5	7.0	7.8	11.6	6.2	9.5	11.4
ACE inhibitors (high blood pressure, heart disease)	5.2	8.8	11.8	5.7	9.8	13.1	4.6	7.9	10.6
Antidiabetic agents (diabetes)	5.5	7.0	10.0	5.9	7.8	11.0	5.1	6.3	9.1
Thyroid hormones (hypothyroidism)	4.7	6.6	8.5	*1.2	*2.7	3.6	8.1	10.1	13.0
Antihypertensive combinations (high blood pressure)	5.3	5.6	8.0	3.3	*3.7	8.1	7.1	7.3	7.9
Anxiolytics, sedatives, and hypnotics (anxiety, insomnia, and related disorders)	6.0	6.2	7.4	4.3	4.9	6.8	7.5	7.4	7.9
Diuretics (high blood pressure, heart disease, kidney disease) ³	6.1	6.6	8.8	4.8	4.8	7.6	7.3	8.3	9.8
Anticonvulsants (epilepsy, seizure, and related disorders)	2.7	4.3	6.3	*2.5	3.5	5.5	2.9	5.1	7.0
Bronchodilators (asthma, breathing)	3.4	3.8	5.8	2.9	3.1	4.8	3.8	4.5	6.7
Calcium channel blocking agents (high blood pressure, heart disease)	7.0	6.7	5.4	8.2	5.9	5.9	5.9	7.5	5.0
65 years and over									
Antihyperlipidemic agents (high cholesterol)	5.9	23.4	46.8	5.3	24.3	51.9	6.4	22.7	42.8
Beta-adrenergic blocking agents (high blood pressure, heart disease)	11.8	15.9	31.2	10.4	17.5	32.6	12.8	14.8	30.0
Diuretics (high blood pressure, heart disease, kidney disease) ³	16.2	19.2	21.6	12.2	17.1	19.4	19.1	20.7	23.3
ACE inhibitors (high blood pressure, heart disease)	9.5	16.9	23.3	9.8	18.0	28.8	9.3	16.1	19.0
Proton pump inhibitors or H2 antagonists (gastric reflux, ulcers) ²	7.5	14.6	21.6	7.2	14.1	18.9	7.7	15.0	23.8
Antidiabetic agents (diabetes)	9.0	12.4	19.0	9.0	12.9	22.0	9.0	12.0	16.7
Anticoagulants or antiplatelet agents (blood clot prevention) ⁵	6.1	9.1	16.3	6.8	11.5	20.3	5.6	7.4	13.0
Analgesics (pain relief)	13.8	18.4	15.8	11.4	15.0	14.5	15.6	20.9	16.9
Calcium channel blocking agents (high blood pressure, heart disease)	16.1	19.1	17.3	14.5	17.4	16.6	17.3	20.4	17.8
Thyroid hormones (hypothyroidism)	7.0	14.3	15.6	3.3	6.7	9.2	9.7	19.8	20.8
Antihypertensive combinations (high blood pressure)	9.6	9.8	13.1	6.0	7.4	9.2	12.2	11.6	16.3
Antidepressants (depression and related disorders)	3.0	9.3	15.0	*2.3	7.2	10.3	3.5	10.8	18.7
Angiotensin II inhibitors (high blood pressure, heart disease)	4.8	12.0	...	4.1	11.6	...	5.3	12.3
Antiarrhythmic agents (heart rhythm irregularities)	23.1	16.6	9.3	21.6	17.9	9.1	24.3	15.6	9.5
65–74 years									
Antihyperlipidemic agents (high cholesterol)	7.3	26.2	45.8	6.2	26.6	49.4	8.1	25.9	42.6
Beta-adrenergic blocking agents (high blood pressure, heart disease)	11.3	14.8	26.2	10.6	16.0	29.4	11.9	13.9	23.3
ACE inhibitors (high blood pressure, heart disease)	9.6	17.2	21.9	10.6	18.1	28.5	8.9	16.4	16.0
Proton pump inhibitors or H2 antagonists (gastric reflux, ulcers) ²	7.0	14.7	19.4	6.3	13.4	16.8	7.5	15.8	21.7
Antidiabetic agents (diabetes)	8.8	12.9	19.5	8.0	13.8	21.4	9.4	12.0	17.9
Diuretics (high blood pressure, heart disease, kidney disease) ³	14.2	15.9	18.7	10.8	14.6	16.1	17.0	16.9	21.0
Analgesics (pain relief)	13.0	18.5	15.8	10.5	14.9	14.7	15.0	21.4	16.9
Antihypertensive combinations (high blood pressure)	8.1	8.0	13.1	4.8	*6.7	9.4	10.8	9.0	16.3
Anticoagulants or antiplatelet agents (blood clot prevention) ⁵	5.4	6.7	11.3	6.3	9.8	16.0	4.6	*4.2	7.2
Antidepressants (depression and related disorders)	2.8	9.3	15.2	*2.3	5.8	10.8	3.1	12.1	19.1
Calcium channel blocking agents (high blood pressure, heart disease)	15.0	16.1	13.6	14.0	15.3	14.9	15.8	16.8	12.5
Thyroid hormones (hypothyroidism)	6.4	13.0	14.7	*3.4	*5.0	*8.5	8.9	19.7	20.2
Angiotensin II inhibitors (high blood pressure, heart disease)	4.2	11.2	...	*3.5	9.7	...	4.9	12.5
Antiarrhythmic agents (heart rhythm irregularities)	20.2	13.0	6.7	19.0	15.5	7.4	21.1	10.8	6.1

See footnotes at end of table.

Table 80 (page 3 of 3). Selected prescription drug classes used in the past 30 days, by sex and age: United States, selected years 1988–1994 through 2009–2012

Updated data when available, Excel, PDF, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#080>.

[Data are based on a sample of the civilian noninstitutionalized population]

Age group and Multum Lexicon Plus therapeutic class ¹ (common indications for use)	Total			Male			Female		
	1988–1994	1999–2002	2009–2012	1988–1994	1999–2002	2009–2012	1988–1994	1999–2002	2009–2012
75 years and over	Percent of population with at least one prescription drug in drug class in past 30 days								
Antihyperlipidemic agents (high cholesterol)	3.8	19.9	48.2	*3.5	21.1	55.7	4.0	19.2	43.1
Beta-adrenergic blocking agents (high blood pressure, heart disease)	12.5	17.3	37.9	9.8	19.6	37.6	14.1	15.8	38.1
Diuretics (high blood pressure, heart disease, kidney disease) ³	19.2	23.2	25.4	14.7	20.5	24.5	21.9	24.9	26.1
ACE inhibitors (high blood pressure, heart disease)	9.3	16.4	25.3	8.5	17.7	29.2	9.8	15.6	22.6
Anticoagulants or antiplatelet agents (blood clot prevention) ⁵	7.2	12.0	22.8	7.8	13.9	27.0	6.9	10.9	20.0
Proton pump inhibitors or H2 antagonists (gastric reflux, ulcers) ²	8.3	14.6	24.6	9.0	15.3	22.1	7.9	14.2	26.4
Calcium channel blocking agents (high blood pressure, heart disease)	17.8	22.8	22.2	15.3	20.5	19.3	19.2	24.2	24.1
Thyroid hormones (hypothyroidism)	7.9	15.8	16.9	3.0	9.2	10.3	10.9	20.0	21.4
Analgesics (pain relief)	15.1	18.4	15.8	13.0	15.1	14.2	16.3	20.4	16.9
Antidiabetic agents (diabetes)	9.3	11.8	18.3	10.7	11.5	22.9	8.5	12.0	15.2
Antihypertensive combinations (high blood pressure)	11.9	12.0	13.2	8.3	*8.2	8.9	14.0	14.4	16.2
Antiarrhythmic agents (heart rhythm irregularities)	27.7	21.0	12.9	26.3	21.3	11.8	28.6	20.7	13.7
Angiotensin II inhibitors (high blood pressure, heart disease)	5.4	13.2	...	*4.9	14.6	...	5.8	12.2
Antidepressants (depression and related disorders)	3.4	9.3	14.7	*2.3	9.2	9.6	4.0	9.4	18.3

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.
 ... Category not applicable.

¹The drug therapeutic class is based on the December 2012 Lexicon Plus, a proprietary database of Cerner Multum, Inc. Lexicon Plus is a comprehensive database of all prescription and some nonprescription drug products available in the U.S. drug market. Data on prescription drug use are collected by the National Health and Nutrition Examination Survey. Respondents were asked if they had taken a prescription drug in the past 30 days. Those who answered “yes” were asked to show the interviewer the medication containers for all prescriptions. If no container was available, the respondent was asked to verbally report the name of the medication. Each drug’s complete name was recorded and classified. Data presented here are based on the second level classification of prescription drugs. Up to four classes are assigned to each drug. Drugs classified into more than one class were counted in each class. For more information, see http://www.cdc.gov/nchs/nhanes/nhanes1999-2000/RXQ_DRUG.htm. See Appendix II, Multum Lexicon Plus therapeutic class.

²The drugs classes proton pump inhibitors (272) and H2 antagonists (94) have been combined because of their similar indications for use.

³This category includes carbonic anhydrase inhibitors which are primarily used to treat glaucoma.

⁴Although sex hormones may be used by males, most are used by females. Therefore, data for sex hormones are only presented for females.

⁵The drugs classes anticoagulants (82) and antiplatelet agents (83) have been combined because of their similar indications for use.

NOTES: Some drug classes were not available in 1988–1994 and are coded as not applicable. See Appendix II, Drug. Standard errors are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health and Nutrition Examination Survey. See Appendix I, National Health and Nutrition Examination Survey (NHANES).

Table 81 (page 1 of 4). Persons with hospital stays in the past year, by selected characteristics: United States, selected years 1997–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#081>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	One or more hospital stays ¹					Two or more hospital stays ¹				
	1997	2000	2010	2013	2014	1997	2000	2010	2013	2014
	Percent									
1 year and over, age-adjusted ^{2,3}	7.8	7.6	7.0	6.7	6.4	1.8	1.8	1.8	1.7	1.6
1 year and over, crude ²	7.7	7.5	7.2	6.9	6.8	1.7	1.8	1.9	1.8	1.7
Age										
1–17 years	2.8	2.5	2.4	2.1	2.0	0.5	0.4	0.5	0.4	0.4
1–5 years	3.9	3.8	3.4	3.4	3.0	0.7	0.7	0.6	0.6	0.7
6–17 years	2.3	1.9	1.9	1.6	1.6	0.4	0.3	0.5	0.3	0.3
18–44 years	7.4	7.0	6.3	6.1	5.8	1.2	1.1	1.3	1.1	1.1
18–24 years	7.9	7.0	5.7	5.3	4.6	1.3	1.1	1.1	1.0	0.9
25–44 years	7.3	7.0	6.6	6.4	6.2	1.2	1.2	1.3	1.2	1.1
45–64 years	8.2	8.4	8.3	7.8	7.4	2.2	2.2	2.5	2.2	2.3
45–54 years	6.9	7.3	7.3	6.2	6.1	1.7	1.8	2.1	1.8	1.9
55–64 years	10.2	10.0	9.5	9.7	8.7	2.9	2.8	2.9	2.6	2.6
65 years and over	18.0	18.2	16.1	15.3	15.3	5.4	5.8	4.9	4.7	4.3
65–74 years	16.1	16.1	13.6	12.6	13.8	4.8	4.9	3.8	3.4	4.0
75 years and over	20.4	20.7	19.0	19.0	17.5	6.2	6.8	6.2	6.5	4.8
75–84 years	19.8	20.1	18.3	17.6	16.1	6.1	6.2	6.1	6.0	4.2
85 years and over	22.8	23.4	20.8	22.4	20.9	6.2	9.0	6.6	7.9	6.3
1–64 years										
Total, 1–64 years ^{2,4}	6.3	6.1	5.7	5.3	5.1	1.3	1.2	1.3	1.2	1.2
Sex										
Male, crude	4.4	4.2	4.2	4.1	3.9	0.9	1.0	1.1	1.0	1.1
1–17 years	2.9	2.4	2.4	2.1	2.0	0.6	0.4	0.5	0.4	0.5
18–44 years	3.6	3.1	2.9	2.9	2.6	0.6	0.6	0.7	0.7	0.6
45–54 years	6.0	7.0	6.4	5.9	5.7	1.4	1.8	1.9	1.7	1.8
55–64 years	11.1	10.2	9.3	9.8	9.0	3.0	3.0	2.8	2.4	2.8
Female, crude	8.0	7.9	7.6	7.0	6.7	1.6	1.5	1.7	1.5	1.4
1–17 years	2.6	2.5	2.3	2.1	2.1	0.5	0.4	0.5	0.4	0.4
18–44 years	11.2	10.8	9.8	9.2	8.9	1.8	1.7	1.9	1.6	1.5
45–54 years	7.6	7.6	8.3	6.5	6.6	2.0	1.9	2.3	2.0	2.0
55–64 years	9.4	9.8	9.7	9.6	8.5	2.9	2.7	2.9	2.7	2.5
Race ^{4,5}										
White only	6.2	5.9	5.6	5.2	5.0	1.2	1.1	1.3	1.1	1.1
Black or African American only	7.6	7.4	6.7	6.2	6.2	1.9	1.9	1.9	1.8	2.1
American Indian or Alaska Native only	7.6	7.0	*7.6	8.9	6.6	*	*	*2.4	*2.6	*
Asian only	3.9	3.9	3.6	3.6	3.1	*0.5	*0.6	*0.4	0.6	0.5
Native Hawaiian or Other Pacific Islander only	---	*	*	*	*	---	*	*	*	*
2 or more races	---	8.8	7.7	7.6	5.9	---	*1.6	*2.4	*1.7	*1.4
Hispanic origin and race ^{4,5}										
Hispanic or Latino	6.8	5.5	5.2	4.8	4.5	1.3	0.9	1.1	1.1	1.1
Not Hispanic or Latino	6.2	6.1	5.8	5.4	5.2	1.3	1.3	1.4	1.2	1.2
White only	6.1	6.0	5.7	5.4	5.2	1.2	1.2	1.3	1.1	1.1
Black or African American only	7.5	7.4	6.7	6.2	6.2	1.9	1.9	1.9	1.8	2.1
Percent of poverty level ^{4,6}										
Below 100%	10.3	9.1	8.3	8.6	8.1	2.8	2.6	2.7	2.7	2.7
100%–199%	7.3	7.3	7.0	6.0	6.1	1.7	1.9	1.9	1.6	1.7
200%–399%	6.0	6.0	5.2	4.8	4.7	1.2	1.1	1.1	1.1	1.0
400% or more	4.7	5.0	4.5	4.2	3.8	0.7	0.8	0.8	0.6	0.6

See footnotes at end of table.

Table 81 (page 2 of 4). Persons with hospital stays in the past year, by selected characteristics: United States, selected years 1997–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#081>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	One or more hospital stays ¹					Two or more hospital stays ¹				
	1997	2000	2010	2013	2014	1997	2000	2010	2013	2014
Hispanic origin and race and percent of poverty level ^{4,5,6}										
Percent										
Hispanic or Latino:										
Below 100%	9.1	7.4	7.3	6.4	6.6	2.0	1.6	2.0	2.0	2.0
100%–199%	5.9	5.4	4.8	4.4	4.4	1.0	0.8	1.1	1.0	1.1
200%–399%	5.9	4.6	4.3	4.3	3.8	1.1	0.7	0.7	0.8	0.8
400% or more	5.5	4.7	4.4	4.2	2.7	*1.1	*0.6	*0.8	*0.6	*
Not Hispanic or Latino:										
White only:										
Below 100%	10.7	9.6	8.8	9.5	8.8	3.2	2.7	2.9	2.7	2.9
100%–199%	7.7	7.8	7.8	6.9	6.8	1.8	2.2	2.2	1.9	1.7
200%–399%	6.1	6.1	5.5	5.1	5.1	1.2	1.1	1.2	1.1	1.0
400% or more	4.7	5.0	4.6	4.2	4.0	0.7	0.8	0.8	0.6	0.6
Black or African American only:										
Below 100%	11.4	10.8	9.4	9.4	8.8	3.3	3.4	3.1	3.6	3.6
100%–199%	8.0	8.5	7.7	6.5	7.2	2.1	2.3	2.3	1.6	2.6
200%–399%	6.2	6.1	5.3	4.5	4.2	1.5	1.3	1.4	1.2	1.3
400% or more	4.7	5.8	4.5	4.7	4.7	*0.9	*1.3	*1.0	*1.0	*0.9
Health insurance status at the time of interview ^{4,7}										
Insured	6.6	6.4	6.2	5.7	5.4	1.3	1.3	1.4	1.2	1.2
Private	5.6	5.5	5.0	4.5	4.2	1.0	1.0	0.9	0.8	0.7
Medicaid	16.1	15.9	12.7	11.5	10.4	4.9	4.7	4.5	3.8	3.6
Uninsured	4.8	4.5	4.0	4.0	3.5	1.0	0.9	0.9	1.0	0.8
Health insurance status prior to interview ^{4,7}										
Insured continuously all 12 months	6.5	6.3	6.0	5.5	5.3	1.3	1.2	1.4	1.2	1.2
Uninsured for any period up to 12 months	8.5	8.4	7.9	7.5	6.5	1.8	1.9	1.9	1.9	1.6
Uninsured more than 12 months	3.8	3.5	3.0	3.3	2.7	0.8	0.8	0.8	0.9	0.7
Percent of poverty level and health insurance status prior to interview ^{4,6,7}										
Below 100%:										
Insured continuously all 12 months	12.4	10.7	10.4	10.3	9.8	3.7	3.1	3.4	3.3	3.4
Uninsured for any period up to 12 months	13.7	13.4	10.4	12.1	10.2	3.4	*3.4	3.0	4.0	3.1
Uninsured more than 12 months	4.9	5.0	4.0	4.7	3.3	1.0	*1.6	1.3	1.3	*1.1
100%–199%:										
Insured continuously all 12 months	8.5	8.6	8.5	7.0	7.1	2.0	2.3	2.5	1.8	2.0
Uninsured for any period up to 12 months	9.3	9.1	10.1	8.5	7.3	*1.9	*2.2	1.9	*1.8	1.7
Uninsured more than 12 months	3.8	3.2	2.7	3.2	2.8	*0.7	*0.7	*0.5	*1.0	*0.8
200%–399%:										
Insured continuously all 12 months	6.3	6.4	5.6	5.2	5.0	1.3	1.2	1.2	1.1	1.0
Uninsured for any period up to 12 months	7.0	6.6	6.1	5.5	5.0	*1.5	*1.3	*1.6	*1.4	*1.0
Uninsured more than 12 months	3.3	2.8	2.6	2.2	2.0	*0.7	*0.4	*0.7	*0.7	*
400% or more:										
Insured continuously all 12 months	4.9	5.1	4.7	4.3	3.9	0.7	0.8	0.8	0.6	0.6
Uninsured for any period up to 12 months	3.9	6.0	4.1	*4.4	3.2	*	*	*	*	*
Uninsured more than 12 months	*	*2.1	*1.8	*	*2.5	*	*	*	*	*
Disability measure among adults 18–64 years ^{4,8}										
Any basic actions difficulty or complex activity limitation	14.1	15.1	14.3	14.2	13.4	4.1	4.4	5.2	4.6	4.6
Any basic actions difficulty	13.9	15.1	14.2	14.2	13.6	4.1	4.4	5.1	4.8	4.7
Any complex activity limitation	21.5	22.6	21.2	20.5	21.5	7.7	8.8	8.6	8.2	8.4
No disability	5.8	5.6	5.4	4.8	4.5	0.6	0.7	0.8	0.6	0.5

See footnotes at end of table.

Table 81 (page 3 of 4). Persons with hospital stays in the past year, by selected characteristics: United States, selected years 1997–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#081>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	One or more hospital stays ¹					Two or more hospital stays ¹				
	1997	2000	2010	2013	2014	1997	2000	2010	2013	2014
Geographic region ⁴						Percent				
Northeast	6.0	5.5	5.2	5.3	5.1	1.2	1.0	1.2	1.2	1.1
Midwest	6.5	6.3	6.3	5.6	5.3	1.5	1.3	1.5	1.3	1.3
South	6.8	6.6	6.0	5.7	5.2	1.4	1.5	1.5	1.3	1.3
West	5.4	5.2	4.9	4.5	4.6	0.8	0.9	1.1	0.8	1.0
Location of residence ^{4,9}										
Within MSA	6.1	5.8	5.5	5.2	5.0	1.2	1.1	1.3	1.1	1.1
Outside MSA	7.0	6.9	6.9	6.3	5.8	1.6	1.5	1.6	1.7	1.4
65 years and over										
Total 65 years and over ^{2,10}	18.1	18.3	16.2	15.7	15.6	5.4	5.8	4.9	4.9	4.4
65–74 years	16.1	16.1	13.6	12.6	13.8	4.8	4.9	3.8	3.4	4.0
75 years and over	20.4	20.7	19.0	19.0	17.5	6.2	6.8	6.2	6.5	4.8
Sex ¹⁰										
Male	19.0	19.5	16.2	16.1	16.6	5.8	5.8	5.4	4.8	4.7
Female	17.5	17.4	16.2	15.4	14.8	5.1	5.7	4.6	5.0	4.1
Hispanic origin and race ^{5,10}										
Hispanic or Latino	17.3	16.6	13.9	15.1	14.7	6.2	6.4	5.0	4.9	4.2
Not Hispanic or Latino	18.2	18.4	16.4	15.7	15.6	5.4	5.8	4.9	4.9	4.4
White only	18.3	18.4	16.5	15.8	15.7	5.4	5.7	4.9	4.9	4.3
Black or African American only	18.9	19.8	16.9	18.2	16.7	5.5	7.5	5.5	5.8	5.5
Percent of poverty level ^{6,10}										
Below 100%	20.9	20.9	18.8	18.3	17.5	6.4	7.5	5.1	6.8	6.1
100%–199%	19.6	19.2	17.2	18.3	17.8	6.5	6.6	5.2	6.0	5.3
200%–399%	17.3	18.1	16.0	15.4	15.5	4.9	5.8	5.5	4.5	4.5
400% or more	16.6	16.0	15.0	13.4	13.9	4.7	4.2	4.1	3.9	3.4
Disability measure ^{8,10}										
Any basic actions difficulty or complex activity limitation	22.6	24.7	20.2	20.2	19.9	7.2	8.6	6.4	7.1	6.2
Any basic actions difficulty	22.7	24.7	20.4	20.4	20.0	7.2	8.7	6.6	7.2	6.3
Any complex activity limitation	29.0	31.5	25.4	26.6	24.8	10.8	12.2	9.2	10.8	9.5
No disability	7.8	9.7	10.6	8.8	8.1	1.1	1.9	*1.6	*1.6	*1.4
Geographic region ¹⁰										
Northeast	17.2	16.6	16.5	15.2	15.6	5.1	4.5	6.1	5.1	4.4
Midwest	18.2	19.5	16.4	16.2	17.6	5.6	7.2	4.7	4.6	4.0
South	19.4	19.5	16.4	16.4	15.0	6.1	6.3	4.7	5.3	4.9
West	16.5	16.4	15.3	14.5	14.3	4.4	4.4	4.5	4.3	3.9
Location of residence ^{9,10}										
Within MSA	17.8	17.8	15.9	15.6	15.4	5.2	5.4	4.8	4.9	4.5
Outside MSA	19.1	19.6	17.3	16.0	16.2	6.3	6.9	5.6	5.0	3.9

See footnotes at end of table.

Table 81 (page 4 of 4). Persons with hospital stays in the past year, by selected characteristics: United States, selected years 1997–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#081>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.

-- Data not available.

¹These estimates exclude hospitalizations for institutionalized persons and those who died while hospitalized, because they are outside the scope of this survey. See Appendix II, Hospital utilization.

²Includes all other races not shown separately, unknown health insurance status, and unknown disability status.

³Estimates are for persons 1 year of age and over and are age-adjusted to the year 2000 standard population using six age groups: 1–17 years, 18–44 years, 45–54 years, 55–64 years, 65–74 years, and 75 years and over. See Appendix II, Age adjustment.

⁴Estimates are for persons aged 1–64 and are age-adjusted to the year 2000 standard population using four age groups: 1–17 years, 18–44 years, 45–54 years, and 55–64 years. The disability measure is age-adjusted using the three adult age groups. See Appendix II, Age adjustment.

⁵The race groups, white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The five single-race categories plus multiple-race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group; the category 2 or more races includes persons who reported more than one racial group. Prior to 1999, data were tabulated according to the 1977 Standards with four racial groups, and the Asian only category included Native Hawaiian or Other Pacific Islander. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

⁶Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Missing family income data were imputed for 1997 and beyond. See Appendix II, Family income; Poverty; Table VI.

⁷Health insurance categories are mutually exclusive. Persons who reported both Medicaid and private coverage are classified as having private coverage. Starting with 1997 data, state-sponsored health plan coverage is included as Medicaid coverage. Starting with 1999 data, coverage by the Children's Health Insurance Program (CHIP) is included with Medicaid coverage. In addition to private and Medicaid, the insured category also includes military, other government, and Medicare coverage. Persons not covered by private insurance, Medicaid, CHIP, state-sponsored or other government-sponsored health plans (starting in 1997), Medicare, or military plans are considered to have no health insurance coverage. Persons with only Indian Health Service coverage are considered to have no health insurance coverage. See Appendix II, Health insurance coverage.

⁸Any basic actions difficulty or complex activity limitation is defined as having one or more of the following limitations or difficulties: movement difficulty, emotional difficulty, sensory (seeing or hearing) difficulty, cognitive difficulty, self-care (activities of daily living or instrumental activities of daily living) limitation, social limitation, or work limitation. For more information, see Appendix II, Basic actions difficulty; Complex activity limitation. Starting with 2007 data, the hearing question, a component of the basic actions difficulty measure, was revised. Consequently, data prior to 2007 are not comparable with data for 2007 and beyond. For more information on the impact of the revised hearing question, see Appendix II, Hearing trouble.

⁹MSA is metropolitan statistical area. Starting with 2006 data, MSA status is determined using 2000 census data and the 2000 standards for defining MSAs. For data prior to 2006, see Appendix II, Metropolitan statistical area (MSA) for the applicable standards.

¹⁰Estimates are for persons aged 65 and over and are age-adjusted to the year 2000 standard population using two age groups: 65–74 years and 75 years and over. See Appendix II, Age adjustment.

NOTES: Standard errors are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey, family core and sample adult questionnaires. See Appendix I, National Health Interview Survey (NHIS).

Table 82. Hospital admissions, average length of stay, outpatient visits, and outpatient surgery, by type of ownership and size of hospital: United States, selected years 1975–2013

[Data are based on reporting by a census of hospitals]

Type of ownership and size of hospital	1975	1980	1990	2000	2010	2011	2012	2013
Admissions								
Number, in thousands								
All hospitals	36,157	38,892	33,774	34,891	36,915	36,565	36,156	35,416
Federal	1,913	2,044	1,759	1,034	911	892	901	949
Nonfederal ¹	34,243	36,848	32,015	33,946	36,004	35,673	35,256	34,467
Community ²	33,435	36,143	31,181	33,089	35,149	34,843	34,422	33,609
Nonprofit	23,722	25,566	22,878	24,453	25,532	25,185	24,751	24,319
For profit	2,646	3,165	3,066	4,141	4,925	5,060	5,224	5,052
State-local government	7,067	7,413	5,236	4,496	4,693	4,598	4,447	4,238
6–24 beds	174	159	95	141	199	197	197	189
25–49 beds	1,431	1,254	870	995	1,169	1,173	1,128	1,087
50–99 beds	3,675	3,700	2,474	2,355	2,173	2,104	2,017	2,021
100–199 beds	7,017	7,162	5,833	6,735	6,125	6,022	5,920	5,754
200–299 beds	6,174	6,596	6,333	6,702	6,569	6,464	6,298	6,156
300–399 beds	4,739	5,358	5,091	5,135	5,835	5,851	5,660	5,344
400–499 beds	3,689	4,401	3,644	3,617	3,869	3,863	3,966	3,750
500 beds or more	6,537	7,513	6,840	7,410	9,210	9,169	9,235	9,307
Average length of stay ³								
Number of days								
All hospitals	11.4	10.0	9.1	6.8	6.2	6.1	6.1	6.1
Federal	20.3	16.8	14.9	12.8	11.8	10.8	9.9	9.6
Nonfederal ¹	10.8	9.6	8.8	6.6	6.1	6.0	6.0	6.0
Community ²	7.7	7.6	7.2	5.8	5.4	5.4	5.4	5.4
Nonprofit	7.8	7.7	7.3	5.7	5.3	5.2	5.2	5.3
For profit	6.6	6.5	6.4	5.4	5.3	5.3	5.3	5.5
State-local government	7.6	7.3	7.7	6.7	6.2	6.2	6.3	6.3
6–24 beds	5.6	5.3	5.4	4.3	4.3	4.5	4.4	4.6
25–49 beds	6.0	5.8	6.1	5.1	5.2	5.2	5.3	5.5
50–99 beds	6.8	6.7	7.2	6.5	6.4	6.5	6.8	6.7
100–199 beds	7.1	7.0	7.1	5.7	5.3	5.2	5.2	5.2
200–299 beds	7.5	7.4	6.9	5.7	5.1	5.1	5.1	5.1
300–399 beds	7.8	7.6	7.0	5.5	5.1	5.1	5.1	5.1
400–499 beds	8.1	7.9	7.3	5.6	5.3	5.3	5.2	5.3
500 beds or more	9.1	8.7	8.1	6.3	5.7	5.7	5.7	5.7
Outpatient visits ⁴								
Number, in thousands								
All hospitals	254,844	262,951	368,184	592,673	750,408	754,454	777,961	787,422
Federal	51,957	50,566	58,527	63,402	90,134	87,975	92,891	98,676
Nonfederal ¹	202,887	212,385	309,657	531,972	660,274	666,479	685,070	688,746
Community ²	190,672	202,310	301,329	521,405	651,424	656,079	674,971	677,951
Nonprofit	131,435	142,156	221,073	393,168	494,178	496,643	512,237	516,162
For profit	7,713	9,696	20,110	43,378	48,201	50,013	53,854	53,191
State-local government	51,525	50,459	60,146	84,858	109,045	109,423	108,880	108,599
6–24 beds	915	1,155	1,471	4,555	9,934	10,531	10,628	10,888
25–49 beds	5,855	6,227	10,812	27,007	43,099	45,098	46,693	47,453
50–99 beds	16,303	17,976	27,582	49,385	57,701	56,126	56,800	58,123
100–199 beds	35,156	36,453	58,940	114,183	120,902	120,555	123,765	123,562
200–299 beds	32,772	36,073	60,561	99,248	110,661	109,901	111,664	112,921
300–399 beds	29,169	30,495	43,699	73,444	90,515	95,282	93,787	89,747
400–499 beds	22,127	25,501	33,394	52,205	65,543	66,428	72,413	71,359
500 beds or more	48,375	48,430	64,870	101,378	153,067	152,158	159,222	163,897
Outpatient surgery								
Percent of total surgeries ⁵								
Community hospitals ²	---	16.3	50.5	62.7	63.6	64.2	64.5	65.6

--- Data not available.

¹The category of nonfederal hospitals comprises psychiatric hospitals, tuberculosis and other respiratory diseases hospitals, and long-term and short-term general and other special hospitals. See Appendix II, Hospital.

²Community hospitals are nonfederal short-term general and special hospitals whose facilities and services are available to the public. The types of facilities included in the community hospitals category have changed over time. See Appendix II, Hospital.

³Average length of stay is the number of inpatient days divided by the number of admissions. See Appendix II, Average length of stay.

⁴Outpatient visits include visits to the emergency department, outpatient department, referred visits (pharmacy, EKG, radiology), and outpatient surgery. See Appendix II, Outpatient visit.

⁵Total surgeries is a measure of patients with at least one surgical procedure. Persons with multiple surgical procedures during the same outpatient visit or inpatient stay are counted only once. See Appendix II, Outpatient surgery.

SOURCE: American Hospital Association (AHA). Annual Survey of Hospitals. Hospital Statistics, 1976, 1981, 1991–92, 2002, 2012, 2013, 2014, and 2015 editions. Chicago, IL. (Reprinted from AHA Hospital Statistics by permission, Copyright 1976, 1981, 1991–92, 2002, 2012, 2013, 2014, and 2015 by Health Forum, LLC, an American Hospital Association Company.) See Appendix I, American Hospital Association (AHA) Annual Survey of Hospitals.

Table 83. Active physicians and physicians in patient care, by state: United States, selected years 1975–2013

[Data are based on reporting by physicians]

State	Active physicians ^{1,2}						Physicians in patient care ^{1,2,3}					
	1975	1985	2000 ⁴	2010	2012	2013	1975	1985	2000 ⁴	2010	2012	2013
	Number per 10,000 civilian population											
United States	15.3	20.7	25.8	27.2	28.3	29.4	13.5	18.0	22.7	24.0	26.9	27.6
Alabama	9.2	14.2	19.8	21.4	21.8	22.4	8.6	13.1	18.2	20.6	21.1	21.5
Alaska	8.4	13.0	18.5	24.3	24.2	25.0	7.8	12.1	16.3	23.3	23.2	23.7
Arizona	16.7	20.2	20.9	22.6	24.2	25.5	14.1	17.1	17.6	21.6	23.2	24.0
Arkansas	9.1	13.8	18.8	20.2	20.9	21.5	8.5	12.8	17.3	19.4	20.2	20.6
California	18.8	23.7	23.8	26.1	26.9	27.8	17.3	21.5	21.6	24.7	25.6	26.2
Colorado	17.3	20.7	24.0	26.9	27.6	29.1	15.0	17.7	20.9	25.5	26.3	27.4
Connecticut	19.8	27.6	33.7	36.0	37.6	38.4	17.7	24.3	30.3	33.6	35.2	35.8
Delaware	14.3	19.7	24.7	26.3	26.4	27.4	12.7	17.1	21.0	25.2	25.3	25.8
District of Columbia	39.6	55.3	62.5	76.9	73.8	74.7	34.6	45.6	54.5	68.8	65.9	66.1
Florida	15.2	20.2	24.1	26.0	26.5	27.2	13.4	17.8	21.2	25.0	25.5	25.7
Georgia	11.5	16.2	20.4	21.3	22.3	23.4	10.6	14.7	18.6	20.2	21.2	22.0
Hawaii	16.2	21.5	26.4	31.3	29.7	30.8	14.7	19.8	24.0	29.6	28.2	29.0
Idaho	9.5	12.1	15.8	18.4	18.4	19.2	8.9	11.4	14.4	17.9	18.0	18.6
Illinois	14.5	20.5	26.1	27.9	28.7	30.1	13.1	18.2	23.1	26.6	27.5	28.5
Indiana	10.6	14.7	20.0	22.2	22.6	23.3	9.6	13.2	18.0	21.3	21.7	22.2
Iowa	11.4	15.6	19.8	21.8	22.0	23.2	9.4	12.4	15.5	20.8	21.0	21.6
Kansas	12.8	17.3	21.8	24.0	24.5	25.4	11.2	15.1	18.8	23.1	23.6	24.1
Kentucky	10.9	15.1	20.6	23.1	23.3	24.6	10.1	13.9	19.1	22.2	22.5	23.6
Louisiana	11.4	17.3	23.8	25.4	26.8	27.2	10.5	16.1	22.4	24.5	25.9	26.3
Maine	12.8	18.7	26.8	31.8	32.0	33.7	10.7	15.6	21.7	30.2	30.5	31.7
Maryland	18.6	30.4	35.4	39.1	39.5	40.9	16.5	24.9	31.1	34.9	35.5	36.5
Massachusetts	20.8	30.2	38.6	43.4	44.6	47.0	18.3	25.4	34.4	40.0	41.3	43.0
Michigan	15.4	20.8	26.3	28.9	30.1	31.5	12.0	16.0	20.2	27.6	28.8	29.4
Minnesota	14.9	20.5	24.9	30.1	30.3	31.1	13.7	18.5	23.0	28.2	28.9	29.5
Mississippi	8.4	11.8	16.6	18.3	18.6	19.5	8.0	11.1	15.2	17.6	18.0	18.7
Missouri	15.0	20.5	24.7	26.3	27.4	28.9	11.6	16.3	20.2	25.1	26.2	26.9
Montana	10.6	14.0	20.4	22.5	22.4	23.1	10.1	13.2	18.8	21.8	21.7	22.3
Nebraska	12.1	15.7	21.7	24.5	24.8	26.0	10.9	14.4	20.1	23.4	23.8	24.8
Nevada	11.9	16.0	18.0	19.8	19.6	20.3	10.9	14.5	15.9	19.2	19.0	19.3
New Hampshire	14.3	18.1	23.8	29.5	30.6	32.0	13.1	16.7	21.7	28.2	29.3	30.4
New Jersey	16.2	23.4	31.1	31.8	32.5	33.5	14.0	19.8	26.2	30.1	30.9	31.5
New Mexico	12.2	17.0	20.9	23.8	24.1	25.2	10.1	14.7	18.5	22.5	22.9	23.6
New York	22.7	29.0	36.2	36.4	38.3	39.4	20.2	25.2	32.3	34.2	36.2	36.8
North Carolina	11.7	16.9	22.3	25.0	25.4	26.4	10.6	15.0	20.5	23.7	24.1	24.9
North Dakota	9.7	15.8	19.2	25.0	25.0	25.3	9.2	14.9	19.8	24.1	24.2	24.4
Ohio	14.1	19.9	25.4	28.5	29.5	31.4	12.2	16.8	21.3	27.3	28.3	29.5
Oklahoma	11.6	16.1	19.4	21.0	21.5	22.3	9.4	12.9	14.8	20.2	20.7	21.0
Oregon	15.6	19.7	22.9	28.3	29.1	30.7	13.8	17.6	20.5	26.9	27.8	29.0
Pennsylvania	16.6	23.6	31.6	32.6	33.1	35.1	13.9	19.2	25.4	30.7	31.2	32.5
Rhode Island	17.8	23.3	32.5	37.1	38.2	40.2	16.1	20.2	28.8	35.2	36.3	37.8
South Carolina	10.0	14.7	21.0	23.3	23.4	24.1	9.3	13.6	19.4	22.4	22.5	23.1
South Dakota	8.2	13.4	19.2	23.0	23.8	24.6	7.7	12.3	17.7	22.2	22.8	23.6
Tennessee	12.4	17.7	23.6	26.0	26.8	27.7	11.3	16.2	21.8	24.8	25.6	26.3
Texas	12.5	16.8	20.3	21.5	22.3	23.2	11.0	14.7	17.9	20.6	21.5	22.1
Utah	14.1	17.2	19.6	21.0	21.9	22.6	13.0	15.5	17.8	20.0	21.0	21.5
Vermont	18.2	23.8	32.0	35.7	36.2	38.2	15.5	20.3	28.8	33.4	34.0	35.6
Virginia	12.9	19.5	23.9	27.0	27.2	28.1	11.9	17.8	22.0	25.7	26.0	26.6
Washington	15.3	20.2	23.7	27.1	27.4	28.4	13.6	17.9	21.2	25.5	25.8	26.5
West Virginia	11.0	16.3	23.5	25.5	26.0	27.1	10.0	14.6	19.5	24.5	24.9	25.4
Wisconsin	12.5	17.7	23.1	26.8	27.3	27.9	11.4	15.9	20.9	25.6	26.2	26.6
Wyoming	9.5	12.9	17.3	19.7	19.2	19.5	8.9	12.0	15.7	19.1	18.8	18.9

¹Includes active doctors of medicine (MDs) and active doctors of osteopathy (DOs). See Appendix II, Physician.

²Starting with 2003 data, federal and nonfederal physicians are included. Data prior to 2003 included nonfederal physicians only.

³Prior to 2006, excludes DOs. Excludes physicians in medical teaching, administration, research, and other nonpatient care activities. Includes residents.

⁴Data for DOs are as of January 2001.

NOTES: Data for MDs are as of December 31. Data for DOs are as of May 31, unless otherwise specified. Starting with *Health, United States, 2012*, data for DOs for 2009 and beyond are from the American Medical Association (AMA). Prior to 2009, data for DOs are from the American Osteopathic Association (AOA).

SOURCE: American Medical Association (AMA): Physician distribution and medical licensure in the U.S., 1975; Physician characteristics and distribution in the U.S., 1986, 2002–2003, 2012, 2014, and 2015 editions; Department of Physician Practice and Communications Information, Division of Survey and Data Resources, AMA. (Copyright 1976, 1986, 2003, 2012, 2014, and 2015: Used with permission of the AMA); American Osteopathic Association: 1975–1976 Yearbook and Directory of Osteopathic Physicians, 1985–1986 Yearbook and Directory of Osteopathic Physicians. See Appendix I, American Medical Association (AMA) Physician Masterfile; American Osteopathic Association (AOA).

Table 84. Doctors of medicine, by place of medical education and activity: United States and outlying U.S. areas, selected years 1975–2013

[Data are based on reporting by physicians]

Place of medical education and activity	1975	1985	1995	2000	2005	2010	2012	2013
Number of doctors of medicine								
Total doctors of medicine	393,742	552,716	720,325	813,770	902,053	985,375	1,026,788	1,045,910
Active doctors of medicine ¹	340,280	497,140	625,443	692,368	762,438	794,862	826,001	854,698
Place of medical education:								
U.S. medical graduates	---	392,007	481,137	527,931	571,798	595,908	615,100	636,707
International medical graduates ²	---	105,133	144,306	164,437	190,640	198,954	210,901	217,991
Activity:								
Patient care ^{3,4}	287,837	431,527	564,074	631,431	718,473	752,572	784,633	809,845
Office-based practice	213,334	329,041	427,275	490,398	563,225	565,024	585,933	600,863
General and family practice	46,347	53,862	59,932	67,534	74,999	77,098	78,935	80,240
Cardiovascular diseases	5,046	9,054	13,739	16,300	17,519	17,454	17,512	17,657
Dermatology	3,442	5,325	6,959	7,969	8,795	9,272	9,669	9,910
Gastroenterology	1,696	4,135	7,300	8,515	9,742	10,466	10,985	11,322
Internal medicine	28,188	52,712	72,612	88,699	107,028	110,612	116,937	120,439
Pediatrics	12,687	22,392	33,890	42,215	51,854	53,054	56,692	58,719
Pulmonary diseases	1,166	3,035	4,964	6,095	7,321	7,846	8,365	8,870
General surgery	19,710	24,708	24,086	24,475	26,079	24,327	24,448	25,024
Obstetrics and gynecology	15,613	23,525	29,111	31,726	34,659	34,083	34,570	34,780
Ophthalmology	8,795	12,212	14,596	15,598	16,580	15,723	16,002	16,331
Orthopedic surgery	8,148	13,033	17,136	17,367	19,115	19,325	19,581	20,013
Otolaryngology	4,297	5,751	7,139	7,581	8,206	7,964	8,021	8,136
Plastic surgery	1,706	3,299	4,612	5,308	6,011	6,180	6,322	6,414
Urological surgery	5,025	7,081	7,991	8,460	8,955	8,606	8,558	8,563
Anesthesiology	8,970	15,285	23,770	27,624	31,887	31,819	32,604	33,218
Diagnostic radiology	1,978	7,735	12,751	14,622	17,618	17,503	17,916	18,203
Emergency medicine	---	---	11,700	14,541	20,173	20,654	22,223	23,414
Neurology	1,862	4,691	7,623	8,559	10,400	10,547	11,249	11,762
Pathology, anatomical/clinical	4,195	6,877	9,031	10,267	11,747	10,688	10,648	10,481
Psychiatry	12,173	18,521	23,334	24,955	27,638	25,690	26,171	26,696
Radiology	6,970	7,355	5,994	6,674	7,049	7,032	7,228	7,527
Other specialty	15,320	28,453	29,005	35,314	39,850	39,081	41,297	43,144
Hospital-based practice	74,503	102,486	136,799	141,033	155,248	187,548	198,700	208,982
Residents and interns ⁵	53,527	72,159	93,650	95,125	95,391	108,142	116,460	117,203
Full-time hospital staff	20,976	30,327	43,149	45,908	59,857	79,406	82,240	91,779
Other professional activity ⁶	24,252	44,046	40,290	41,556	43,965	42,290	41,368	44,853
Inactive	21,449	38,646	72,326	75,168	99,823	125,928	142,716	147,676
Not classified	26,145	13,950	20,579	45,136	39,304	64,153	57,649	43,536
Unknown address	5,868	2,980	1,977	1,098	488	432	422	---

--- Data not available.

¹Doctors of medicine who are inactive, have unknown address, or primary specialty not classified are excluded. See Appendix II, Physician.

²International medical graduates received their medical education in schools outside of the United States and Canada.

³Specialty information is based on the physician's self-designated primary area of practice. Categories include generalists and specialists. See Appendix II, Physician specialty.

⁴Starting with 2003 data, federal and nonfederal doctors of medicine are included. Data prior to 2003 included nonfederal doctors of medicine only.

⁵Starting with 1990 data, clinical fellows are included in this category. In prior years, clinical fellows were included in the other professional activity category.

⁶Includes doctors of medicine in medical teaching, administration, research, and other nonpatient care activities. Prior to 1990, this category also included clinical fellows.

NOTES: Data for doctors of medicine are as of December 31. Outlying areas include Puerto Rico, the U.S. Virgin Islands, and the U.S. Pacific islands.

SOURCE: American Medical Association (AMA). Physician distribution and medical licensure in the U.S., 1975; Physician characteristics and distribution in the U.S., 1986, 1996–1997, 2002–2003, 2007, 2012, 2014, and 2015 editions, Department of Physician Practice and Communications Information, Division of Survey and Data Resources, AMA. (Copyright 1976, 1986, 1997, 2003, 2007, 2012, 2014, and 2015: Used with permission of the AMA.) See Appendix I, American Medical Association (AMA) Physician Masterfile.

Table 85. Doctors of medicine in primary care, by specialty: United States and outlying U.S. areas, selected years 1949–2013

[Data are based on reporting by physicians]

Specialty	1949 ¹	1960 ¹	1970	1980	1990	2000	2010	2012	2013
	Number								
Total doctors of medicine ²	201,277	260,484	334,028	467,679	615,421	813,770	985,375	1,026,788	1,045,910
Active doctors of medicine ³	191,577	247,257	310,845	414,916	547,310	692,368	794,862	826,001	854,698
General primary care specialists	113,222	125,359	134,354	170,705	213,514	274,653	304,687	313,793	319,881
General practice/family medicine	95,980	88,023	57,948	60,049	70,480	86,312	94,746	96,552	98,298
Internal medicine	12,453	26,209	39,924	58,462	76,295	101,353	113,591	118,504	121,127
Obstetrics/Gynecology	---	---	18,532	24,612	30,220	35,922	38,520	39,324	40,045
Pediatrics	4,789	11,127	17,950	27,582	36,519	51,066	57,830	59,413	60,411
Primary care subspecialists	---	---	3,161	16,642	30,911	52,294	76,122	83,532	90,147
Family medicine	---	---	---	---	---	483	1,445	1,764	1,991
Internal medicine	---	---	1,948	13,069	22,054	34,831	50,730	55,357	59,256
Obstetrics/Gynecology	---	---	344	1,693	3,477	4,319	4,277	4,186	4,141
Pediatrics	---	---	869	1,880	5,380	12,661	19,670	22,225	24,759
	Percent of active doctors of medicine								
General primary care specialists	59.1	50.7	43.2	41.1	39.0	39.7	38.3	38.0	37.4
General practice/family medicine	50.1	35.6	18.6	14.5	12.9	12.5	11.9	11.7	11.5
Internal medicine	6.5	10.6	12.8	14.1	13.9	14.6	14.3	14.3	14.2
Obstetrics/Gynecology	---	---	6.0	5.9	5.5	5.2	4.8	4.8	4.7
Pediatrics	2.5	4.5	5.8	6.6	6.7	7.4	7.3	7.2	7.1
Primary care subspecialists	---	---	1.0	4.0	5.6	7.6	9.6	10.1	10.5
Family medicine	---	---	---	---	---	0.1	0.2	0.2	0.2
Internal medicine	---	---	0.6	3.1	4.0	5.0	6.4	6.7	6.9
Obstetrics/Gynecology	---	---	0.1	0.4	0.6	0.6	0.5	0.5	0.5
Pediatrics	---	---	0.3	0.5	1.0	1.8	2.5	2.7	2.9

--- Data not available.

0.0 Percentage greater than zero but less than 0.05.

¹Estimated by the Bureau of Health Professions, Health Resources and Services Administration. Active doctors of medicine (MDs) include those with address unknown and primary specialty not classified.

²Data on federal and nonfederal doctors of medicine engaged in office- or hospital-based patient care and other professional activities.

³Starting with 1970 data, MDs who are inactive, have unknown address, or primary specialty not classified are excluded. See Appendix II, Physician.

NOTES: See Appendix II, Physician specialty. Data are as of December 31 except for 1990–1994 data, which are as of January 1, and 1949 data, which are as of midyear. Outlying areas include Puerto Rico, the U.S. Virgin Islands, and the U.S. Pacific islands.

SOURCE: Health Manpower Source Book: Medical Specialists, USDHEW, 1962; American Medical Association (AMA). Distribution of physicians in the United States, 1970; Physician characteristics and distribution in the U.S., 1981, 1992, 2002–2003, 2012, 2014, and 2015 editions, Department of Physician Practice and Communications Information, Division of Survey and Data Resources, AMA. (Copyright 1971, 1982, 1992, 2003, 2012, 2014, and 2015: Used with permission of the AMA.) See Appendix I, American Medical Association (AMA) Physician Masterfile.

Table 86. Active dentists, by state: United States, selected years 2001–2013

[Data are based on reporting by dentists]

State	2001	2006	2011	2012	2013	2001	2006	2011	2012	2013
	Number of dentists					Number of dentists per 100,000 civilian population				
United States	163,345	172,603	186,025	188,820	191,347	57.32	57.85	59.68	60.11	60.46
Alabama	1,880	1,921	2,056	2,107	2,128	42.08	41.50	42.82	43.74	44.02
Alaska	457	489	555	571	577	72.11	72.41	76.81	78.10	78.26
Arizona	2,374	3,061	3,465	3,515	3,617	45.02	50.77	53.53	53.61	54.51
Arkansas	1,047	1,114	1,161	1,187	1,210	38.90	39.48	39.51	40.25	40.90
California	22,709	26,388	28,680	29,119	29,425	65.86	73.26	76.07	76.50	76.57
Colorado	2,844	3,098	3,486	3,563	3,623	64.26	65.63	68.09	68.63	68.72
Connecticut	2,590	2,587	2,721	2,695	2,742	75.45	73.55	75.78	74.98	76.18
Delaware	352	383	409	422	420	44.24	44.57	45.05	46.03	45.39
District of Columbia	603	533	564	568	579	104.96	93.40	90.91	89.44	89.20
Florida	8,158	8,754	9,583	9,774	9,947	49.87	48.19	50.15	50.50	50.75
Georgia	3,614	4,115	4,574	4,629	4,701	43.14	44.94	46.61	46.67	47.03
Hawaii	1,022	1,009	1,047	1,048	1,060	83.36	77.04	75.97	75.25	75.23
Idaho	690	864	927	913	932	52.27	58.83	58.53	57.22	57.79
Illinois	8,154	7,994	8,416	8,476	8,599	65.29	63.22	65.45	65.84	66.71
Indiana	2,870	2,842	3,074	3,066	3,116	46.84	44.88	47.17	46.90	47.42
Iowa	1,516	1,526	1,601	1,601	1,604	51.71	51.16	52.24	52.05	51.87
Kansas	1,314	1,347	1,437	1,442	1,461	48.63	48.75	50.07	49.97	50.45
Kentucky	2,256	2,287	2,452	2,427	2,488	55.46	54.20	56.11	55.37	56.55
Louisiana	2,058	2,017	2,189	2,240	2,221	45.96	46.88	47.84	48.65	47.98
Maine	598	642	650	660	693	46.51	48.50	48.95	49.68	52.16
Maryland	3,955	3,989	4,160	4,248	4,268	73.59	70.89	71.19	72.10	71.87
Massachusetts	4,898	4,797	5,181	5,234	5,232	76.56	74.84	78.35	78.64	77.99
Michigan	5,783	5,928	6,072	6,036	6,075	57.88	59.07	60.47	61.06	61.37
Minnesota	2,880	3,105	3,210	3,254	3,284	57.80	60.13	60.02	60.48	60.57
Mississippi	1,117	1,140	1,220	1,253	1,275	39.15	39.24	40.96	41.96	42.61
Missouri	2,634	2,666	2,846	2,897	2,900	46.69	45.63	47.35	48.08	47.97
Montana	511	525	613	616	598	56.34	55.11	61.44	61.28	58.92
Nebraska	1,103	1,117	1,166	1,201	1,203	64.13	63.01	63.29	64.73	64.37
Nevada	846	1,177	1,396	1,432	1,448	40.32	46.66	51.35	51.97	51.87
New Hampshire	735	815	833	840	847	58.54	62.29	63.20	63.57	64.04
New Jersey	6,054	6,922	7,181	7,264	7,238	71.28	79.92	81.21	81.84	81.22
New Mexico	814	861	1,023	1,055	1,062	44.44	43.88	49.22	50.61	50.89
New York	14,309	14,062	14,211	14,262	14,468	74.98	73.61	72.80	72.74	73.46
North Carolina	3,474	4,016	4,588	4,674	4,719	42.31	45.04	47.54	47.95	47.91
North Dakota	305	311	370	383	394	47.73	47.89	54.00	54.58	54.43
Ohio	5,929	5,797	5,932	6,012	6,003	52.07	50.49	51.38	52.05	51.88
Oklahoma	1,664	1,749	1,912	1,924	1,943	47.99	48.66	50.49	50.41	50.43
Oregon	2,197	2,431	2,643	2,643	2,708	63.35	66.22	68.34	67.79	68.94
Pennsylvania	7,595	7,454	7,529	7,676	7,698	61.75	59.58	59.08	60.11	60.23
Rhode Island	588	576	581	582	566	55.62	54.18	55.23	55.29	53.73
South Carolina	1,839	1,958	2,228	2,267	2,288	45.24	44.93	47.68	48.00	47.95
South Dakota	348	382	443	440	457	45.91	48.78	53.75	52.73	54.05
Tennessee	2,912	2,947	3,131	3,195	3,246	50.64	48.40	48.93	49.50	49.96
Texas	9,642	10,365	12,451	12,861	13,391	45.23	44.37	48.53	49.29	50.52
Utah	1,409	1,559	1,852	1,878	1,892	61.70	61.73	65.78	65.77	65.18
Vermont	354	343	357	361	365	57.82	55.07	56.99	57.66	58.23
Virginia	4,189	4,367	4,854	5,068	5,194	58.19	56.91	59.85	61.85	62.80
Washington	3,957	4,312	4,856	4,917	4,951	66.11	67.68	71.18	71.30	70.99
West Virginia	863	835	873	874	890	47.91	45.68	47.06	47.08	48.01
Wisconsin	3,069	2,860	3,072	3,137	3,215	56.76	51.28	53.81	54.80	55.98
Wyoming	266	266	294	306	309	53.77	50.89	51.79	53.04	52.98

NOTES: Data include professionally active dentists only. Professionally active dentists include those whose primary occupation is one of the following: private practice (full- or part-time), dental school/faculty staff member, armed forces, other federal services (i.e., Veterans' Affairs, Public Health Service), state or local government employee, hospital staff dentist, graduate student/intern/resident, or other health/dental organization staff member. U.S. totals include dentists with unknown state of practice not shown separately and may include missing data.

SOURCE: American Dental Association, Health Policy Institute, Dentist Supply in the US: 2001–2013, Tables 1 and 3 (Copyright 2015 American Dental Association. Reprinted with permission. All rights reserved.) Any form of reproduction is strictly prohibited without prior written permission of the American Dental Association. See Appendix I, American Dental Association (ADA).

Table 87. Healthcare employment and wages, by selected occupations: United States, selected years 2000–2014Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#087>.

[Data are based on a semiannual mail survey of nonfarm establishments]

Occupation title	2000	2010	2013	2014	2000	2010	2013	2014
Healthcare practitioners and technical occupations								
	Employment ¹				Mean hourly wage ²			
Audiologists	11,530	12,860	11,550	12,250	\$22.92	\$33.58	\$35.75	\$36.92
Cardiovascular technologists and technicians	40,080	48,720	51,010	51,080	16.81	24.38	25.95	26.54
Dental hygienists	148,460	177,520	192,330	196,520	24.99	33.02	34.39	34.60
Diagnostic medical sonographers	31,760	53,010	58,250	59,760	22.03	31.20	32.29	32.88
Dietetic technicians	28,010	23,890	26,420	28,690	10.98	13.86	13.74	13.75
Dietitians and nutritionists	43,030	53,510	59,530	59,490	18.76	26.13	27.07	27.62
Emergency medical technicians and paramedics	165,530	221,760	237,660	235,760	11.89	16.01	16.77	16.88
Licensed practical and licensed vocational nurses	679,470	730,290	705,200	695,610	14.65	19.88	20.63	20.87
Magnetic resonance imaging technologists	---	---	32,000	33,130	---	---	31.71	32.36
Medical and clinical laboratory technicians	146,060	156,480	157,080	160,460	13.93	18.36	19.35	19.59
Medical and clinical laboratory technologists	144,530	164,430	162,630	161,710	19.84	27.34	28.59	29.12
Medical records and health information technicians	143,870	176,090	180,760	184,740	11.74	16.83	18.13	18.68
Nuclear medicine technologists	18,030	21,600	20,020	20,320	21.56	33.20	34.60	35.21
Nurse anesthetists	---	---	35,430	36,590	---	---	75.81	76.40
Nurse midwives	---	---	5,460	5,110	---	---	44.34	46.97
Nurse practitioners	---	---	113,370	122,050	---	---	45.71	47.11
Occupational therapists	75,150	100,300	108,410	110,520	24.10	35.28	37.45	38.46
Opticians, dispensing	66,580	62,200	68,390	73,110	12.67	16.73	17.17	17.43
Pharmacists	212,660	268,030	287,420	290,780	33.39	52.59	56.01	56.96
Pharmacy technicians	190,940	333,500	362,690	368,760	10.38	14.10	14.83	14.95
Physical therapists	120,410	180,280	195,670	200,670	27.62	37.50	39.51	40.35
Physician assistants	55,490	81,420	88,110	91,670	29.17	41.89	45.36	46.77
Psychiatric technicians	53,350	72,650	66,760	64,540	12.53	15.15	16.09	16.91
Radiation therapists	13,100	16,590	16,950	16,380	25.59	37.64	39.30	40.25
Radiologic technologists ³	172,080	216,730	194,000	193,400	17.93	26.80	27.29	27.65
Recreational therapists	26,940	20,830	18,640	17,950	14.23	19.92	21.88	22.14
Registered nurses ⁴	2,189,670	2,655,020	2,661,890	2,687,310	22.31	32.56	33.13	33.55
Respiratory therapists	82,670	109,270	118,640	119,410	18.37	26.54	27.83	28.12
Respiratory therapy technicians	28,230	13,570	12,070	10,610	16.46	22.28	23.01	23.46
Speech-language pathologists	82,850	112,530	125,050	126,500	23.31	33.60	35.56	36.01
Healthcare support occupations								
Dental assistants	250,870	294,030	309,540	314,330	12.86	16.41	17.13	17.43
Home health aides	561,120	982,840	806,710	799,080	8.71	10.46	10.60	10.77
Massage therapists	24,620	60,040	79,040	87,670	15.51	19.12	19.42	20.09
Medical assistants	330,830	523,260	571,690	584,970	11.46	14.31	14.80	15.01
Medical equipment preparers	32,760	47,310	51,300	50,550	10.68	14.59	16.02	16.28
Medical transcriptionists	97,330	78,780	68,350	61,210	12.37	16.12	16.95	17.11
Nursing assistants ⁵	1,273,460	1,451,090	1,427,830	1,427,740	9.18	12.09	12.51	12.62
Occupational therapy aides	8,890	7,180	8,710	8,570	11.21	14.95	13.90	13.96
Occupational therapy assistants	15,910	27,720	30,450	32,230	16.76	24.66	26.56	27.53
Pharmacy aides	59,890	49,580	42,250	41,240	9.10	10.98	11.78	12.28
Physical therapist aides	34,620	45,900	48,630	48,730	10.06	12.02	12.50	12.82
Physical therapist assistants	44,120	65,960	72,640	76,910	16.52	23.95	25.63	26.12
Psychiatric aides	57,680	64,730	75,340	72,860	10.79	12.84	12.98	13.67

--- Data not available.

¹Employment is the number of filled positions. This table includes both full-time and part-time wage and salary positions. Estimates do not include the self-employed, owners and partners in unincorporated firms, household workers, or unpaid family workers. Estimates were rounded to the nearest 10.²The mean hourly wage rate for an occupation is the total wages that all workers in the occupation earn in an hour, divided by the total number of employees in the occupation. More information is available from: http://www.bls.gov/oes/current/oes_tec.htm.³2012–2014 data are not comparable with earlier data. Starting with 2012 data, the radiologic technologists and technicians occupation category was split into two occupations as part of the 2010 Standard Occupational Classification (SOC) revision: Radiologic technologists (29–2034) and Magnetic resonance imaging technologists (29–2035).⁴2012–2014 data are not comparable to earlier data. Starting with 2012 data, the registered nurses occupation category was split into four occupations as part of the 2010 SOC revision: Registered nurses (29–1141), plus three advanced practice nursing occupations: Nurse anesthetists (29–1151), Nurse midwives (29–1161), and Nurse practitioners (29–1171).⁵2012–2014 data are not comparable to earlier data. Starting with 2012 data, the nursing aides, orderlies, and attendants occupation category was split into two occupations as part of the 2010 SOC revision: Nursing assistants (31–1014) and Orderlies (31–1015).

NOTES: This table excludes occupations such as dentists, physicians, and chiropractors, which have a large percentage of workers who are self-employed. Challenges in using Occupational Employment Statistics (OES) data as a time series include changes in the occupational, industrial, and geographical classification systems; changes in the way data are collected; changes in the survey reference period; and changes in mean wage estimation methodology, as well as permanent features of the methodology. See Appendix I, Occupational Employment Statistics (OES).

SOURCE: U.S. Department of Labor, Bureau of Labor Statistics. Occupational Employment Statistics. Available from: http://www.bls.gov/oes/current/oes_nat.htm#29-0000. See Appendix I, Occupational Employment Statistics (OES).

Table 88. First-year enrollment and graduates of health professions schools, and number of schools, by selected profession: United States, selected academic years 1980–1981 through 2013–2014

[Data are based on reporting by health professions associations]

Profession	Academic years						
	1980–1981	1990–1991	2000–2001	2010–2011	2011–2012	2012–2013	2013–2014
First-year enrollment				Number			
Dentistry	6,030	4,001	4,327	5,170	5,493	5,697	5,904
Medicine (Allopathic) ^{1,2}	17,186	16,876	16,699	19,082	19,947	20,279	20,803
Medicine (Osteopathic) ³	1,496	1,950	2,927	5,428	5,788	5,986	6,636
Optometry ¹	1,174	1,245	1,384	1,661	1,674	1,760	1,818
Pharmacy ^{1,4}	7,377	8,267	8,382	13,077	13,464	14,011	14,008
Podiatry	695	561	475	671	672	687	671
Public Health: ^{1,5,6}							
Schools and programs	---	---	---	---	---	---	13,766
Schools	3,348	4,087	5,840	11,205	11,345	11,588	12,486
Graduates							
Dentistry	5,256	5,550	4,367	5,070	5,229	5,351	5,491
Medicine (Allopathic) ¹	15,632	15,427	15,796	17,363	17,341	18,157	18,078
Medicine (Osteopathic)	1,151	1,534	2,510	4,159	4,458	4,806	4,997
Optometry ^{1,7}	1,092	1,224	1,310	1,308	1,383	1,545	1,541
Pharmacy ¹	7,323	7,122	7,000	11,931	12,719	13,207	13,838
Podiatry	597	591	531	543	537	572	564
Public Health: ^{1,6}							
Schools and programs	---	---	---	---	---	---	11,932
Schools	3,168	3,995	5,747	9,717	9,969	10,477	11,052
Schools							
Dentistry	60	56	55	58	61	62	65
Medicine (Allopathic) ^{1,8}	125	125	124	135	138	141	141
Medicine (Osteopathic)	14	15	19	34	34	37	40
Optometry ¹	13	17	17	20	20	21	21
Pharmacy ¹	72	74	82	123	129	130	133
Podiatry	5	7	7	9	9	9	9
Public Health: ^{1,6,9}							
Schools and programs	---	---	---	---	---	---	91
Schools	21	24	28	46	49	50	56

¹Includes data from schools in Puerto Rico.

²Includes new entrants and those repeating the initial year.

³May also include persons enrolled in first-year classes for data years 1980–1981.

⁴Starting with 2005–2006 data, first-year enrollment for pharmacy schools include Pharm.D.1 enrollments only. Prior to 2005, first-year enrollment data include both Pharm.D.1, B.S. Pharmacy, and B.Pharm. enrollments. Includes second from last year for baccalaureate and third from last year for Pharm.D.1 and does not include first-year enrollees in accelerated programs.

⁵Starting with 2008–2009 data, first-year enrollment data for public health schools include spring, summer, and fall enrollment. All other data years include fall enrollment only and are not directly comparable.

⁶Includes data from a school of public health in Mexico as of 2007 school year. 2013–2014 data reported for 53 schools and 24 programs of public health.

⁷Excludes graduates of “special” optometry degree programs which include, but are not limited to, accelerated programs for those entering optometry schools with a doctoral degree or graduates of foreign optometry programs and modified extended programs for those returning to schools after an absence, changing professional fields or taking a reduced course load for personal reasons.

⁸Includes schools with preliminary and provisional accreditation, in addition to fully accredited schools.

⁹Includes programs of public health as Association of Schools & Programs of Public Health members as of 2013 school year.

NOTE: Data on the number of schools and first-year enrollments are reported as of the beginning of the academic year, while data on the number of graduates are reported as of the end of the academic year.

SOURCE: American Dental Association: 2014–2015 Survey of Dental Education Series, Report 1: Academic Programs, Enrollment and Graduates. Available from: <http://www.ada.org/en/science-research/health-policy-institute/data-center/dental-education> (Copyright 2016 American Dental Association. Reprinted with permission. All rights reserved.) Any form of reproduction is strictly prohibited without prior written permission of the American Dental Association; Association of American Medical Colleges (AAMC): AAMC Data Book 2015 – Medical Schools and Teaching Hospitals by the Numbers, Washington, DC. 2015. Table A1 (number of schools) and Table B1 (number of first-year enrollment students and number of graduates). Used with permission of the AAMC; American Association of Colleges of Osteopathic Medicine: Trends in Osteopathic Medical School Applicants, Enrollment and Graduates, 2016. Chevy Chase, MD. 2016. Available from: <http://www.aacom.org/reports-programs-initiatives/aacom-reports>. Reprinted with permission from AACOM, All rights reserved; Association of Schools and Colleges of Optometry: Annual Student Data Report Academic Years 2014–2015 and unpublished data. Available from: <http://www.opted.org/data-surveys/>; American Association of Colleges of Pharmacy: Fall 2014 Profile of Pharmacy Students. Available from: <http://www.aacp.org/resources/research/institutionalresearch/Pages/StudentApplications,EnrollmentsandDegreesConferred.aspx>; American Association of Colleges of Podiatric Medicine: Applicant, Matriculant, and Graduate Statistics, 2006 through 2014 and unpublished data. Available from: <http://www.aacpm.org>. Used with permission of the AACPM; Association of Schools & Programs of Public Health: unpublished data. Washington, DC. Used with permission of the ASPPH; Bureau of Health Professions: United States Health Personnel FACTBOOK. Health Resources and Services Administration. Rockville, MD. 2003. See Appendix I, American Dental Association (ADA); Association of American Medical Colleges (AAMC); American Association of Colleges of Osteopathic Medicine (AACOM); Association of Schools and Colleges of Optometry (ASCO); American Association of Colleges of Pharmacy (AACCP); American Association of Colleges of Podiatric Medicine (AACPM); Association of Schools & Programs of Public Health (ASPPH).

Table 89. Hospitals, beds, and occupancy rates, by type of ownership and size of hospital: United States, selected years 1975–2013

[Data are based on reporting by a census of hospitals]

Type of ownership and size of hospital	1975	1980	1990	2000	2005	2010	2012	2013
Hospitals								
	Number							
All hospitals	7,156	6,965	6,649	5,810	5,756	5,754	5,723	5,686
Federal	382	359	337	245	226	213	211	213
Nonfederal ¹	6,774	6,606	6,312	5,565	5,530	5,541	5,512	5,473
Community ²	5,875	5,830	5,384	4,915	4,936	4,985	4,999	4,974
Nonprofit	3,339	3,322	3,191	3,003	2,958	2,904	2,894	2,904
For profit	775	730	749	749	868	1,013	1,068	1,060
State-local government	1,761	1,778	1,444	1,163	1,110	1,068	1,037	1,010
6–24 beds	299	259	226	288	370	424	462	469
25–49 beds	1,155	1,029	935	910	1,032	1,167	1,192	1,186
50–99 beds	1,481	1,462	1,263	1,055	1,001	970	954	959
100–199 beds	1,363	1,370	1,306	1,236	1,129	1,029	1,012	995
200–299 beds	678	715	739	656	619	585	570	571
300–399 beds	378	412	408	341	368	352	348	334
400–499 beds	230	266	222	182	173	185	189	183
500 beds or more	291	317	285	247	244	273	272	277
Beds								
All hospitals	1,465,828	1,364,516	1,213,327	983,628	946,997	941,995	920,829	914,513
Federal	131,946	117,328	98,255	53,067	45,837	44,940	38,557	38,747
Nonfederal ¹	1,333,882	1,247,188	1,115,072	930,561	901,160	897,055	882,272	875,766
Community ²	941,844	988,387	927,360	823,560	802,311	804,943	800,566	795,603
Nonprofit	658,195	692,459	656,755	582,988	561,106	555,768	545,287	543,929
For profit	73,495	87,033	101,377	109,883	113,510	124,652	135,008	134,643
State-local government	210,154	208,895	169,228	130,689	127,695	124,523	120,271	117,031
6–24 beds	5,615	4,932	4,427	5,156	6,316	7,261	7,791	7,763
25–49 beds	41,783	37,478	35,420	33,333	33,726	37,446	38,338	38,039
50–99 beds	106,776	105,278	90,394	75,865	71,737	69,470	67,879	67,892
100–199 beds	192,438	192,892	183,867	175,778	161,593	148,090	145,556	143,760
200–299 beds	164,405	172,390	179,670	159,807	151,290	142,616	139,212	140,113
300–399 beds	127,728	139,434	138,938	117,220	126,899	121,749	120,554	115,511
400–499 beds	101,278	117,724	98,833	80,763	76,894	82,071	84,007	81,148
500 beds or more	201,821	218,259	195,811	175,638	173,856	196,240	197,229	201,377
Occupancy rate³								
	Percent							
All hospitals	76.7	77.7	69.5	66.1	69.3	66.6	65.2	64.7
Federal	80.7	80.1	72.9	68.2	66.0	65.3	63.5	64.5
Nonfederal ¹	76.3	77.4	69.2	65.9	69.5	66.6	65.3	64.7
Community ²	75.0	75.2	66.8	63.9	67.3	64.5	63.4	62.9
Nonprofit	77.5	78.2	69.3	65.5	69.1	66.2	64.9	64.5
For profit	65.9	65.2	52.8	55.9	59.6	57.1	56.8	56.2
State-local government	70.4	71.1	65.3	63.2	66.7	64.4	63.8	62.9
6–24 beds	48.0	46.8	32.3	31.7	33.5	32.3	30.8	30.5
25–49 beds	56.7	52.8	41.3	41.3	47.1	44.8	43.1	42.7
50–99 beds	64.7	64.2	53.8	54.8	59.0	55.1	55.2	55.1
100–199 beds	71.2	71.4	61.5	60.0	63.2	60.4	58.1	57.6
200–299 beds	77.1	77.4	67.1	65.0	67.7	64.0	63.2	61.6
300–399 beds	79.7	79.7	70.0	65.7	70.1	67.4	65.1	64.9
400–499 beds	81.1	81.2	73.5	69.1	71.2	68.5	67.5	67.6
500 beds or more	80.9	82.1	77.3	72.2	75.9	73.0	72.6	72.1

¹The category of nonfederal hospitals comprises psychiatric hospitals, tuberculosis and other respiratory diseases hospitals, and long-term and short-term general and other special hospitals. See Appendix II, Hospital.

²Community hospitals are nonfederal short-term general and special hospitals whose facilities and services are available to the public. The types of facilities included in the community hospitals category have changed over time. See Appendix II, Hospital.

³Estimated percentage of staffed beds that are occupied. Occupancy rate is calculated as the average daily census (from the American Hospital Association) divided by the number of hospital beds. See Appendix II, Occupancy rate.

SOURCE: American Hospital Association (AHA). Annual Survey of Hospitals. Hospital Statistics, 1976, 1981, 1991–92, 2002, 2012, 2014, and 2015 editions. Chicago, IL. (Reprinted from AHA Hospital Statistics by permission, Copyright 1976, 1981, 1991–92, 2002, 2012, 2014, and 2015 by Health Forum, LLC, an American Hospital Association Company.) See Appendix I, American Hospital Association (AHA) Annual Survey of Hospitals.

Table 90. Community hospital beds and average annual percent change, by state: United States, selected years 1980–2013

[Data are based on reporting by a census of hospitals]

State	1980	1990	2000	2010	2012	2013	1980–1990	1990–2000	2000–2010	2010–2013
	Beds per 1,000 resident population						Average annual percent change ¹			
United States	4.5	3.7	2.9	2.6	2.6	2.5	-1.9	-2.4	-1.1	-1.3
Alabama	5.1	4.6	3.7	3.2	3.1	3.1	-1.0	-2.2	-1.4	-1.1
Alaska	2.7	2.3	2.3	2.2	2.1	2.1	-1.6	-	-0.4	-1.5
Arizona	3.6	2.7	2.1	2.0	2.1	2.0	-2.8	-2.5	-0.5	-
Arkansas	5.0	4.6	3.7	3.2	3.2	3.2	-0.8	-2.2	-1.4	-
California	3.6	2.7	2.1	1.9	1.9	1.8	-2.8	-2.5	-1.0	-1.8
Colorado	4.2	3.2	2.2	2.0	2.0	2.0	-2.7	-3.7	-0.9	-
Connecticut	3.5	2.9	2.3	2.3	2.3	2.2	-1.9	-2.3	-	-1.5
Delaware	3.6	3.0	2.3	2.4	2.2	2.2	-1.8	-2.6	0.4	-2.9
District of Columbia	7.3	7.6	5.8	5.7	5.7	5.6	0.4	-2.7	-0.2	-0.6
Florida	5.1	3.9	3.2	2.9	2.8	2.7	-2.6	-2.0	-1.0	-2.4
Georgia	4.6	4.0	2.9	2.6	2.5	2.5	-1.4	-3.2	-1.1	-1.3
Hawaii	3.1	2.7	2.5	2.4	2.0	2.0	-1.4	-0.8	-0.4	-5.9
Idaho	3.7	3.2	2.7	2.2	2.1	2.1	-1.4	-1.7	-2.0	-1.5
Illinois	5.1	4.0	3.0	2.6	2.5	2.5	-2.4	-2.8	-1.4	-1.3
Indiana	4.5	3.9	3.2	2.8	2.7	2.6	-1.4	-2.0	-1.3	-2.4
Iowa	5.7	5.1	4.0	3.3	3.2	3.2	-1.1	-2.4	-1.9	-1.0
Kansas	5.8	4.8	4.0	3.5	3.5	3.5	-1.9	-1.8	-1.3	-
Kentucky	4.5	4.3	3.7	3.3	3.2	3.2	-0.5	-1.5	-1.1	-1.0
Louisiana	4.8	4.6	3.9	3.4	3.3	3.4	-0.4	-1.6	-1.4	-
Maine	4.7	3.7	2.9	2.7	2.7	2.6	-2.4	-2.4	-0.7	-1.3
Maryland	3.6	2.8	2.1	2.0	2.1	2.1	-2.5	-2.8	-0.5	1.6
Massachusetts	4.4	3.6	2.6	2.4	2.4	2.5	-2.0	-3.2	-0.8	1.4
Michigan	4.4	3.7	2.6	2.6	2.5	2.5	-1.7	-3.5	-	-1.3
Minnesota	5.7	4.4	3.4	2.9	2.8	2.7	-2.6	-2.5	-1.6	-2.4
Mississippi	5.3	5.0	4.8	4.4	4.3	4.3	-0.6	-0.4	-0.9	-0.8
Missouri	5.7	4.8	3.6	3.1	3.2	3.1	-1.7	-2.8	-1.5	-
Montana	5.9	5.8	4.7	3.8	3.7	3.7	-0.2	-2.1	-2.1	-0.9
Nebraska	6.0	5.5	4.8	4.0	3.8	3.6	-0.9	-1.4	-1.8	-3.5
Nevada	4.2	2.8	1.9	2.0	2.0	2.0	-4.0	-3.8	0.5	-
New Hampshire	3.9	3.1	2.3	2.2	2.1	2.1	-2.3	-2.9	-0.4	-1.5
New Jersey	4.2	3.7	3.0	2.4	2.4	2.4	-1.3	-2.1	-2.2	-
New Mexico	3.1	2.8	1.9	2.0	2.0	1.8	-1.0	-3.8	0.5	-3.5
New York	4.5	4.1	3.5	3.0	2.9	2.9	-0.9	-1.6	-1.5	-1.1
North Carolina	4.2	3.3	2.9	2.4	2.3	2.3	-2.4	-1.3	-1.9	-1.4
North Dakota	7.4	7.0	6.0	5.1	4.7	4.0	-0.6	-1.5	-1.6	-7.8
Ohio	4.7	4.0	3.0	3.0	2.9	2.9	-1.6	-2.8	-	-1.1
Oklahoma	4.6	4.0	3.2	3.0	3.0	3.0	-1.4	-2.2	-0.6	-
Oregon	3.5	2.8	1.9	1.7	1.7	1.7	-2.2	-3.8	-1.1	-
Pennsylvania	4.8	4.4	3.4	3.2	3.1	3.1	-0.9	-2.5	-0.6	-1.1
Rhode Island	3.8	3.2	2.3	2.3	2.3	2.1	-1.7	-3.2	-	-3.0
South Carolina	3.9	3.3	2.9	2.7	2.7	2.7	-1.7	-1.3	-0.7	-
South Dakota	5.5	6.1	5.7	5.0	5.0	4.9	1.0	-0.7	-1.3	-0.7
Tennessee	5.5	4.8	3.6	3.3	3.1	3.1	-1.4	-2.8	-0.9	-2.1
Texas	4.7	3.5	2.7	2.4	2.4	2.3	-2.9	-2.6	-1.2	-1.4
Utah	3.1	2.6	1.9	1.8	1.8	1.8	-1.7	-3.1	-0.5	-
Vermont	4.4	3.0	2.7	2.1	2.0	1.9	-3.8	-1.0	-2.5	-3.3
Virginia	4.1	3.3	2.4	2.2	2.2	2.2	-2.1	-3.1	-0.9	-
Washington	3.1	2.5	1.9	1.7	1.8	1.7	-2.1	-2.7	-1.1	-
West Virginia	5.5	4.7	4.4	4.0	3.9	3.8	-1.6	-0.7	-0.9	-1.7
Wisconsin	4.9	3.8	2.9	2.4	2.3	2.2	-2.5	-2.7	-1.9	-2.9
Wyoming	3.6	4.8	3.9	3.6	3.3	3.3	2.9	-2.1	-0.8	-2.9

- Quantity zero.

¹See Appendix II, Average annual rate of change (percent change).

NOTES: Community hospitals are nonfederal short-term general and special hospitals whose facilities and services are available to the public. The types of facilities included in the community hospitals category have changed over time. See Appendix II, Hospital. See *Health, United States, 2013*, Table 108, for 1970 hospital data.

SOURCE: American Hospital Association (AHA). Annual Survey of Hospitals. Hospital Statistics, 1981, 1991–92, 2002, 2012, 2014, and 2015 editions. Chicago, IL. (Reprinted from AHA Hospital Statistics by permission, Copyright 1981, 1991–92, 2002, 2012, 2014, and 2015 by Health Forum, LLC, an American Hospital Association Company.) See Appendix I, American Hospital Association (AHA) Annual Survey of Hospitals.

Table 91. Occupancy rates in community hospitals and average annual percent change, by state: United States, selected years 1980–2013

[Data are based on reporting by a census of hospitals]

State	1980	1990	2000	2010	2013	1980–1990	1990–2000	2000–2010	2010–2013
	Occupancy rate ¹					Average annual percent change ²			
United States	75	67	64	65	63	-1.1	-0.5	0.2	-1.0
Alabama	73	63	60	61	61	-1.5	-0.5	0.2	-
Alaska	58	50	57	61	66	-1.5	1.3	0.7	2.7
Arizona	74	62	63	65	59	-1.8	0.2	0.3	-3.2
Arkansas	70	62	59	55	53	-1.2	-0.5	-0.7	-1.2
California	69	64	66	68	64	-0.7	0.3	0.3	-2.0
Colorado	72	64	58	60	57	-1.2	-1.0	0.3	-1.7
Connecticut	80	77	75	78	75	-0.4	-0.3	0.4	-1.3
Delaware	82	77	75	74	73	-0.6	-0.3	-0.1	-0.5
District of Columbia	83	75	74	73	72	-1.0	-0.1	-0.1	-0.5
Florida	72	62	61	63	62	-1.5	-0.2	0.3	-0.5
Georgia	70	66	63	66	67	-0.6	-0.5	0.5	0.5
Hawaii	75	85	76	72	74	1.3	-1.1	-0.5	0.9
Idaho	65	56	53	51	50	-1.5	-0.5	-0.4	-0.7
Illinois	75	66	60	62	59	-1.3	-0.9	0.3	-1.6
Indiana	78	61	56	58	58	-2.4	-0.9	0.4	-
Iowa	69	62	58	56	56	-1.1	-0.7	-0.4	-
Kansas	69	56	53	54	52	-2.1	-0.5	0.2	-1.3
Kentucky	77	62	62	60	59	-2.1	-	-0.3	-0.6
Louisiana	70	57	56	59	54	-2.0	-0.2	0.5	-2.9
Maine	75	72	64	62	62	-0.4	-1.2	-0.3	-
Maryland	84	79	73	74	70	-0.6	-0.8	0.1	-1.8
Massachusetts	82	74	71	73	69	-1.0	-0.4	0.3	-1.9
Michigan	78	66	65	66	65	-1.7	-0.2	0.2	-0.5
Minnesota	74	67	67	64	63	-1.0	-	-0.5	-0.5
Mississippi	71	59	59	54	53	-1.8	-	-0.9	-0.6
Missouri	75	62	58	61	59	-1.9	-0.7	0.5	-1.1
Montana	66	61	67	63	60	-0.8	0.9	-0.6	-1.6
Nebraska	67	58	59	55	54	-1.4	0.2	-0.7	-0.6
Nevada	69	60	71	68	67	-1.4	1.7	-0.4	-0.5
New Hampshire	73	67	59	60	61	-0.9	-1.3	0.2	0.6
New Jersey	83	80	69	71	66	-0.4	-1.5	0.3	-2.4
New Mexico	66	58	58	57	56	-1.3	-	-0.2	-0.6
New York	86	86	79	79	78	-	-0.8	-	-0.4
North Carolina	78	73	70	70	69	-0.7	-0.4	0.0	-0.5
North Dakota	69	64	60	59	62	-0.7	-0.6	-0.2	1.7
Ohio	79	65	61	61	59	-1.9	-0.6	-	-1.1
Oklahoma	68	58	56	57	54	-1.6	-0.4	0.2	-1.8
Oregon	69	57	59	59	59	-1.9	0.3	-	-
Pennsylvania	80	73	68	67	64	-0.9	-0.7	-0.1	-1.5
Rhode Island	86	79	72	69	70	-0.8	-0.9	-0.4	0.5
South Carolina	77	71	69	66	62	-0.8	-0.3	-0.4	-2.1
South Dakota	61	62	65	62	62	0.2	0.5	-0.5	-
Tennessee	76	64	56	60	61	-1.7	-1.3	0.7	0.6
Texas	70	57	59	60	59	-2.0	0.3	0.2	-0.6
Utah	70	59	56	53	51	-1.7	-0.5	-0.5	-1.3
Vermont	74	67	67	65	69	-1.0	-	-0.3	2.0
Virginia	78	67	68	67	68	-1.5	0.1	-0.1	0.5
Washington	72	63	60	63	62	-1.3	-0.5	0.5	-0.5
West Virginia	76	63	61	61	60	-1.9	-0.3	-	-0.5
Wisconsin	74	65	60	60	60	-1.3	-0.8	-	-
Wyoming	57	54	56	56	53	-0.5	0.4	-	-1.8

- Quantity zero.

¹Estimated percent of staffed beds that are occupied. Occupancy rate is calculated as the average daily census (inpatient days divided by 365) divided by the number of hospital beds. See Appendix II, Occupancy rate.

²See Appendix II, Average annual rate of change (percent change).

NOTES: Community hospitals are nonfederal short-term general and special hospitals whose facilities and services are available to the public. The types of facilities included in the community hospitals category have changed over time. See Appendix II, Hospital. See *Health, United States, 2013*, Table 109, for 1970 hospital data.

SOURCE: American Hospital Association (AHA). Annual Survey of Hospitals. Hospital Statistics, 1981, 1991–92, 2002, 2012, and 2015 editions. Chicago, IL. (Reprinted from AHA Hospital Statistics by permission, Copyright 1981, 1991–92, 2002, 2012, and 2015 by Health Forum, LLC, an American Hospital Association Company.) See Appendix I, American Hospital Association (AHA) Annual Survey of Hospitals.

Table 92 (page 1 of 2). Nursing homes, beds, residents, and occupancy rates, by state: United States, selected years 1995–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#092>.

[Data are based on a census of certified nursing facilities]

State	Nursing homes				Beds			
	1995	2000	2013	2014	1995	2000	2013	2014
	Number							
United States	16,389	16,886	15,663	15,643	1,751,302	1,795,388	1,697,484	1,693,943
Alabama	221	225	228	226	23,353	25,248	26,685	26,388
Alaska	15	15	17	18	814	821	779	693
Arizona	152	150	146	147	16,162	17,458	16,607	16,605
Arkansas	256	255	230	229	29,952	25,715	24,546	24,558
California	1,382	1,369	1,226	1,217	140,203	131,762	121,381	119,866
Colorado	219	225	211	214	19,912	20,240	20,371	20,431
Connecticut	267	259	231	229	32,827	32,433	27,841	27,673
Delaware	42	43	46	46	4,739	4,906	4,986	4,876
District of Columbia	19	20	19	19	3,206	3,078	2,766	2,766
Florida	627	732	687	689	72,656	83,365	83,178	83,545
Georgia	352	363	358	357	38,097	39,817	39,883	39,975
Hawaii	34	45	47	46	2,513	4,006	4,215	4,213
Idaho	76	84	77	78	5,747	6,181	5,930	5,951
Illinois	827	869	769	761	103,230	110,766	98,883	98,348
Indiana	556	564	515	528	59,538	56,762	58,764	59,555
Iowa	419	467	444	443	39,959	37,034	32,183	31,950
Kansas	429	392	345	345	30,016	27,067	25,653	25,730
Kentucky	288	307	283	287	23,221	25,341	26,170	26,300
Louisiana	337	337	280	280	37,769	39,430	35,189	35,066
Maine	132	126	107	105	9,243	8,248	7,020	6,953
Maryland	218	255	230	228	28,394	31,495	28,487	28,115
Massachusetts	550	526	421	416	54,532	56,030	48,660	48,320
Michigan	432	439	432	433	49,473	50,696	46,970	46,521
Minnesota	432	433	380	377	43,865	42,149	30,405	30,319
Mississippi	183	190	205	205	16,059	17,068	18,550	18,434
Missouri	546	551	513	512	52,679	54,829	55,106	55,273
Montana	100	104	83	83	7,210	7,667	6,713	6,732
Nebraska	231	236	217	219	18,169	17,877	15,855	16,005
Nevada	42	51	51	52	3,998	5,547	5,979	6,040
New Hampshire	74	83	76	76	7,412	7,837	7,510	7,501
New Jersey	300	361	365	361	43,967	52,195	52,417	52,051
New Mexico	83	80	71	71	6,969	7,289	6,716	6,869
New York	624	665	631	628	107,750	120,514	116,448	117,131
North Carolina	391	410	421	422	38,322	41,376	44,598	45,088
North Dakota	87	88	81	81	7,125	6,954	6,138	6,131
Ohio	943	1,009	955	954	106,884	105,038	91,563	90,653
Oklahoma	405	392	311	309	33,918	33,903	29,396	28,962
Oregon	161	150	138	137	13,885	13,500	12,276	12,210
Pennsylvania	726	770	703	699	92,625	95,063	88,284	88,236
Rhode Island	94	99	84	84	9,612	10,271	8,715	8,720
South Carolina	166	178	189	188	16,682	18,102	19,689	19,631
South Dakota	114	114	111	111	8,296	7,844	6,909	6,945
Tennessee	322	349	320	321	37,074	38,593	37,140	37,268
Texas	1,266	1,215	1,205	1,212	123,056	125,052	135,350	136,000
Utah	91	93	98	99	7,101	7,651	8,500	8,577
Vermont	23	44	38	37	1,862	3,743	3,199	3,174
Virginia	271	278	286	288	30,070	30,595	32,638	32,497
Washington	285	277	225	221	28,464	25,905	21,641	21,286
West Virginia	129	139	126	127	10,903	11,413	10,888	10,888
Wisconsin	413	420	392	389	48,754	46,395	34,730	33,959
Wyoming	37	40	39	39	3,035	3,119	2,984	2,965

See footnotes at end of table.

Table 92 (page 2 of 2). Nursing homes, beds, residents, and occupancy rates, by state: United States, selected years 1995–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#092>.

[Data are based on a census of certified nursing facilities]

State	Residents				Occupancy rate ¹			
	1995	2000	2013	2014	1995	2000	2013	2014
	Number							
United States	1,479,550	1,480,076	1,371,926	1,368,667	84.5	82.4	80.8	80.8
Alabama	21,691	23,089	22,764	22,731	92.9	91.4	85.3	86.1
Alaska	634	595	498	612	77.9	72.5	63.9	88.3
Arizona	12,382	13,253	11,344	11,428	76.6	75.9	68.3	68.8
Arkansas	20,823	19,317	17,774	17,688	69.5	75.1	72.4	72.0
California	109,805	106,460	102,324	102,245	78.3	80.8	84.3	85.3
Colorado	17,055	17,045	15,957	16,309	85.7	84.2	78.3	79.8
Connecticut	29,948	29,657	24,610	24,250	91.2	91.4	88.4	87.6
Delaware	3,819	3,900	4,217	4,314	80.6	79.5	84.6	88.5
District of Columbia	2,576	2,858	2,569	2,539	80.3	92.9	92.9	91.8
Florida	61,845	69,050	72,679	73,487	85.1	82.8	87.4	88.0
Georgia	35,933	36,559	33,889	33,930	94.3	91.8	85.0	84.9
Hawaii	2,413	3,558	3,714	3,663	96.0	88.8	88.1	86.9
Idaho	4,697	4,640	3,909	3,841	81.7	75.1	65.9	64.5
Illinois	83,696	83,604	72,856	72,563	81.1	75.5	73.7	73.8
Indiana	44,328	42,328	38,649	38,893	74.5	74.6	65.8	65.3
Iowa	27,506	29,204	24,980	24,859	68.8	78.9	77.6	77.8
Kansas	25,140	22,230	18,400	18,337	83.8	82.1	71.7	71.3
Kentucky	20,696	22,730	22,818	23,008	89.1	89.7	87.2	87.5
Louisiana	32,493	30,735	25,600	25,854	86.0	77.9	72.8	73.7
Maine	8,587	7,298	6,342	6,239	92.9	88.5	90.3	89.7
Maryland	24,716	25,629	24,360	24,430	87.0	81.4	85.5	86.9
Massachusetts	49,765	49,805	41,595	41,255	91.3	88.9	85.5	85.4
Michigan	43,271	42,615	39,288	39,374	87.5	84.1	83.6	84.6
Minnesota	41,163	38,813	27,201	26,695	93.8	92.1	89.5	88.0
Mississippi	15,247	15,815	16,165	16,129	94.9	92.7	87.1	87.5
Missouri	39,891	38,586	37,828	38,326	75.7	70.4	68.7	69.3
Montana	6,415	5,973	4,689	4,619	89.0	77.9	69.9	68.6
Nebraska	16,166	14,989	12,070	12,043	89.0	83.8	76.1	75.2
Nevada	3,645	3,657	4,749	4,821	91.2	65.9	79.4	79.8
New Hampshire	6,877	7,158	6,813	6,767	92.8	91.3	90.7	90.2
New Jersey	40,397	45,837	45,450	45,185	91.9	87.8	86.7	86.8
New Mexico	6,051	6,503	5,531	5,439	86.8	89.2	82.4	79.2
New York	103,409	112,957	105,965	105,390	96.0	93.7	91.0	90.0
North Carolina	35,511	36,658	36,908	37,058	92.7	88.6	82.8	82.2
North Dakota	6,868	6,343	5,702	5,664	96.4	91.2	92.9	92.4
Ohio	79,026	81,946	77,129	76,325	73.9	78.0	84.2	84.2
Oklahoma	26,377	23,833	19,376	19,108	77.8	70.3	65.9	66.0
Oregon	11,673	9,990	7,373	7,343	84.1	74.0	60.1	60.1
Pennsylvania	84,843	83,880	79,554	79,598	91.6	88.2	90.1	90.2
Rhode Island	8,823	9,041	7,986	8,011	91.8	88.0	91.6	91.9
South Carolina	14,568	15,739	16,744	16,773	87.3	86.9	85.0	85.4
South Dakota	7,926	7,059	6,335	6,381	95.5	90.0	91.7	91.9
Tennessee	33,929	34,714	29,990	28,897	91.5	89.9	80.7	77.5
Texas	89,354	85,275	93,712	93,170	72.6	68.2	69.2	68.5
Utah	5,832	5,703	5,383	5,515	82.1	74.5	63.3	64.3
Vermont	1,792	3,349	2,726	2,686	96.2	89.5	85.2	84.6
Virginia	28,119	27,091	28,249	28,486	93.5	88.5	86.6	87.7
Washington	24,954	21,158	17,199	17,005	87.7	81.7	79.5	79.9
West Virginia	10,216	10,334	9,524	9,535	93.7	90.5	87.5	87.6
Wisconsin	43,998	38,911	28,062	27,485	90.2	83.9	80.8	80.9
Wyoming	2,661	2,605	2,377	2,364	87.7	83.5	79.7	79.7

--- Data not available.

¹Percentage of beds occupied (number of nursing home residents per 100 nursing home beds).

NOTES: Annual numbers of nursing homes, beds, and residents are based on the Centers for Medicare & Medicaid Services' reporting cycle. Starting with 2013 data, a new editing rule was used for number of beds. For the U.S., the number of beds decreased by less than 1%. For most states, this caused little or no change in the data. The change in the number of beds also caused a change in some occupancy rates. Because of the methodology change, trends should be interpreted with caution. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: Cowles CM ed., 2014 Nursing Home Statistical Yearbook. Anacortes, WA: Cowles Research Group, 2015 and previous editions; and Cowles Research Group, unpublished data. Based on data from the Centers for Medicare & Medicaid Services' Quality Improvement Evaluation System (QIES) and its predecessor, the Online Survey Certification and Reporting Database (OSCAR). See Appendix I, Quality Improvement Evaluation System (QIES).

Table 93 (page 1 of 2). Gross domestic product, national health expenditures, per capita amounts, percent distribution, and average annual percent change: United States, selected years 1960–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#093>.

[Data are compiled from various sources by the Centers for Medicare & Medicaid Services]

<i>Gross domestic product and national health expenditures</i>	1960	1970	1980	1990	2000	2009	2012	2013	2014
Amount, in billions									
Gross domestic product (GDP)	\$543	\$1,076	\$2,863	\$5,980	\$10,285	\$14,419	\$16,155	\$16,663	\$17,348
Deflator (2009 = 100.0)									
Price deflator for GDP ¹	17.5	22.8	44.5	66.8	81.9	100.0	105.2	106.9	108.7
Amount, in billions									
National health expenditures	\$27.2	\$74.6	\$255.3	\$721.4	\$1,369.7	\$2,496.4	\$2,799.0	\$2,879.9	\$3,031.3
Health consumption expenditures	24.7	67.0	235.5	674.1	1,286.4	2,357.5	2,645.8	2,727.4	2,877.4
Personal health care	23.3	63.1	217.0	615.3	1,162.0	2,115.9	2,371.8	2,441.3	2,563.6
Administration and net cost of private health insurance	1.1	2.6	12.1	38.7	81.3	167.5	197.9	209.5	234.8
Public health	0.4	1.4	6.4	20.0	43.0	74.1	76.0	76.6	79.0
Investment ²	2.5	7.5	19.9	47.3	83.3	139.0	153.2	152.5	153.9
Deflator (2009 = 100.0)									
Chain-weighted national health expenditure deflator ¹	---	---	---	---	---	100.0	106.9	108.3	110.2
Per capita amount, in dollars									
National health expenditures	\$146	\$355	\$1,108	\$2,843	\$4,857	\$8,147	\$8,927	\$9,115	\$9,523
Health consumption expenditures	133	319	1,022	2,657	4,562	7,693	8,438	8,632	9,040
Personal health care	125	300	942	2,425	4,121	6,905	7,564	7,727	8,054
Administration and net cost of private health insurance	6	13	52	153	288	546	631	663	738
Public health	2	6	28	79	153	242	243	242	248
Investment ²	13	36	86	187	295	453	489	483	483
Percent									
National health expenditures as percent of GDP	5.0	6.9	8.9	12.1	13.3	17.3	17.3	17.3	17.5
Percent distribution									
National health expenditures	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Health consumption expenditures	90.8	89.9	92.2	93.4	93.9	94.4	94.5	94.7	94.9
Personal health care	85.5	84.6	85.0	85.3	84.8	84.8	84.7	84.8	84.6
Administration and net cost of private health insurance	3.9	3.5	4.7	5.4	5.9	6.7	7.1	7.3	7.7
Public health	1.4	1.8	2.5	2.8	3.1	3.0	2.7	2.7	2.6
Investment ²	9.2	10.1	7.8	6.6	6.1	5.6	5.5	5.3	5.1
Average annual percent change from previous year shown ³									
GDP	7.1	10.3	7.6	5.6	3.8	3.9	3.1	4.1
National health expenditures	10.6	13.1	10.9	6.6	6.9	3.9	2.9	5.3
Health consumption expenditures	10.5	13.4	11.1	6.7	7.0	3.9	3.1	5.5
Personal health care	10.5	13.2	11.0	6.6	6.9	3.9	2.9	5.0
Administration and net cost of private health insurance	9.4	16.4	12.4	7.7	8.4	5.7	5.8	12.1
Public health	13.8	16.9	12.0	8.0	6.2	0.8	0.7	3.1
Investment ²	11.6	10.2	9.1	5.8	5.9	3.3	-0.5	0.9
National health expenditures, per capita	9.3	12.1	9.9	5.5	5.9	3.1	2.1	4.5
Health consumption expenditures	9.1	12.3	10.0	5.6	6.0	3.1	2.3	4.7
Personal health care	9.1	12.1	9.9	5.4	5.9	3.1	2.2	4.2
Administration and net cost of private health insurance	8.0	14.9	11.4	6.5	7.4	4.9	5.1	11.3
Public health	11.6	16.7	10.9	6.8	5.2	0.1	-0.4	2.5
Investment ²	10.7	9.1	8.1	4.7	4.9	2.6	-1.2	0.0

See footnotes at end of table.

Table 93 (page 2 of 2). Gross domestic product, national health expenditures, per capita amounts, percent distribution, and average annual percent change: United States, selected years 1960–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#093>.

[Data are compiled from various sources by the Centers for Medicare & Medicaid Services]

-- Data not available.

... Category not applicable.

¹Year 2009 = 100. For more information on the detailed price series recommended for deflating each category of spending see the National Health Expenditure Accounts Methodology Paper, 2014 and NHE Deflator Methodology paper. Available from:

<https://www.cms.gov/Research-Statistics-Data-and-Systems/Statistics-Trends-and-Reports/NationalHealthExpendData/NationalHealthAccountsHistorical.html>.

²Investment consists of research and structures and equipment.

³See Appendix II, Average annual rate of change (percent change).

NOTES: Dollar amounts shown are in current dollars. See Appendix II, Gross domestic product (GDP); Health expenditures, national. Percents are calculated using unrounded data. Estimates may not add to totals because of rounding. Census resident-based population less armed forces overseas and population of outlying areas used to calculate per capita. For more information on NHE categories, sources, and methods, see the National Health Expenditure Accounts Methodology Paper, 2014. Available from:

<https://www.cms.gov/Research-Statistics-Data-and-Systems/Statistics-Trends-and-Reports/NationalHealthExpendData/NationalHealthAccountsHistorical.html>. See Appendix I, National Health Expenditure Accounts (NHEA). Data have been revised and differ from previous editions of *Health, United States*.

SOURCE: Centers for Medicare & Medicaid Services, Office of the Actuary, National Health Statistics Group, National Health Expenditure Accounts, National health expenditures aggregate. Available from:

<https://www.cms.gov/Research-Statistics-Data-and-Systems/Statistics-Trends-and-Reports/NationalHealthExpendData/NationalHealthAccountsHistorical.html>, accessed on December 12, 2015. U.S. Department of Commerce Bureau of Economic Analysis, National Economic Accounts, National Income and Product Accounts, Table 1.1.4, accessed on December 10, 2015. Available from: <http://www.bea.gov/iTable/iTable.cfm?ReqID=9&step=1>. See Appendix I, National Health Expenditure Accounts (NHEA); National Income and Product Accounts (NIPA).

Table 94 (page 1 of 2). National health expenditures, average annual percent change, and percent distribution, by type of expenditure: United States, selected years 1960–2014

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2015.htm#094>.

[Data are compiled from various sources by the Centers for Medicare & Medicaid Services]

Type of national health expenditure	1960	1970	1980	1990	2000	2012	2013	2014
Amount, in billions								
National health expenditures	\$27.2	\$74.6	\$255.3	\$721.4	\$1,369.7	\$2,799.0	\$2,879.9	\$3,031.3
Health consumption expenditures	24.7	67.0	235.5	674.1	1,286.4	2,645.8	2,727.4	2,877.4
Personal health care	23.3	63.1	217.0	615.3	1,162.0	2,371.8	2,441.3	2,563.6
Hospital care	9.0	27.2	100.5	250.4	415.5	902.7	933.9	971.8
Professional services	7.9	19.8	64.5	207.3	387.4	749.5	767.5	801.6
Physician and clinical services	5.6	14.3	47.7	158.4	288.7	563.0	576.8	603.7
Other professional services	0.4	0.7	3.5	17.3	36.6	77.6	80.3	84.4
Dental services	2.0	4.7	13.3	31.6	62.1	108.9	110.4	113.5
Other health, residential, and personal care	0.4	1.3	8.4	23.8	63.9	137.9	144.5	150.4
Home health care ¹	0.1	0.2	2.4	12.5	32.3	76.9	79.4	83.2
Nursing care facilities and continuing care retirement communities ¹	0.8	4.0	15.3	44.7	85.0	148.3	150.2	155.6
Retail outlet sales of medical products	5.0	10.6	25.9	76.5	177.8	356.5	365.8	401.0
Prescription drugs	2.7	5.5	12.0	40.3	121.0	259.1	265.3	297.7
Durable medical equipment	0.7	1.7	4.1	13.8	25.2	43.7	44.9	46.4
Other nondurable medical products	1.6	3.3	9.8	22.4	31.6	53.7	55.6	56.9
Government administration ²	0.1	0.7	2.8	7.2	17.1	33.5	36.3	40.2
Net cost of health insurance ³	1.0	1.9	9.3	31.6	64.2	164.4	173.2	194.6
Government public health activities ⁴	0.4	1.4	6.4	20.0	43.1	76.0	76.6	79.0
Investment	2.5	7.5	19.9	47.3	83.3	153.2	152.5	153.9
Research ⁵	0.7	2.0	5.4	12.7	25.5	48.4	46.5	45.5
Structures and equipment	1.8	5.6	14.4	34.6	57.8	104.8	106.0	108.3
Average annual percent change from previous year shown ⁶								
National health expenditures	...	10.6	13.1	10.9	6.6	6.1	2.9	5.3
Health consumption expenditures	...	10.5	13.4	11.1	6.7	6.2	3.1	5.5
Personal health care	...	10.5	13.2	11.0	6.6	6.1	2.9	5.0
Hospital care	...	11.7	14.0	9.6	5.2	6.7	3.5	4.1
Professional services	...	9.6	12.6	12.4	6.5	5.7	2.4	4.4
Physician and clinical services	...	9.9	12.8	12.7	6.2	5.7	2.5	4.6
Other professional services	...	6.3	17.0	17.4	7.8	6.4	3.5	5.2
Dental services	...	9.0	11.0	9.0	7.0	4.8	1.5	2.8
Other health, residential, and personal care	...	11.5	20.5	11.0	10.4	6.6	4.7	4.1
Home health care ¹	...	14.5	26.9	18.1	9.9	7.5	3.3	4.8
Nursing care facilities and continuing care retirement communities ¹	...	17.4	14.2	11.4	6.6	4.7	1.3	3.6
Retail outlet sales of medical products	...	7.7	9.4	11.4	8.8	6.0	2.6	9.6
Prescription drugs	...	7.5	8.2	12.8	11.6	6.5	2.4	12.2
Durable medical equipment	...	9.0	8.8	13.0	6.2	4.7	2.8	3.2
Other nondurable medical products	...	7.4	11.4	8.6	3.5	4.5	3.5	2.4
Government administration ²	...	30.0	14.1	10.0	9.0	5.8	8.5	10.7
Net cost of health insurance ³	...	6.4	17.2	13.0	7.4	8.1	5.3	12.4
Government public health activities ⁴	...	13.8	16.9	12.0	8.0	4.9	0.7	3.1
Investment	...	11.6	10.2	9.1	5.8	5.2	-0.5	0.9
Research ⁵	...	10.9	10.8	8.9	7.2	5.5	-4.1	-2.0
Structures and equipment	...	11.9	10.0	9.2	5.3	5.1	1.2	2.2
Percent distribution								
National health expenditures	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Health consumption expenditures	90.8	89.9	92.2	93.4	93.9	94.5	94.7	94.9
Personal health care	85.5	84.6	85.0	85.3	84.8	84.7	84.8	84.6
Hospital care	33.0	36.4	39.4	34.7	30.3	32.3	32.4	32.1
Professional services	29.1	26.5	25.3	28.7	28.3	26.8	26.7	26.4
Physician and clinical services	20.4	19.2	18.7	22.0	21.1	20.1	20.0	19.9
Other professional services	1.4	1.0	1.4	2.4	2.7	2.8	2.8	2.8
Dental services	7.3	6.3	5.2	4.4	4.5	3.9	3.8	3.7
Other health, residential, and personal care	1.6	1.7	3.3	3.3	4.7	4.9	5.0	5.0
Home health care ¹	0.2	0.3	0.9	1.7	2.4	2.7	2.8	2.7
Nursing care facilities and continuing care retirement communities ¹	3.0	5.4	6.0	6.2	6.2	5.3	5.2	5.1
Retail outlet sales of medical products	18.5	14.2	10.1	10.6	13.0	12.7	12.7	13.2
Prescription drugs	9.8	7.4	4.7	5.6	8.8	9.3	9.2	9.8
Durable medical equipment	2.7	2.3	1.6	1.9	1.8	1.6	1.6	1.5
Other nondurable medical products	6.0	4.5	3.8	3.1	2.3	1.9	1.9	1.9
Government administration ²	0.2	1.0	1.1	1.0	1.2	1.2	1.3	1.3
Net cost of health insurance ³	3.7	2.5	3.6	4.4	4.7	5.9	6.0	6.4
Government public health activities ⁴	1.4	1.8	2.5	2.8	3.1	2.7	2.7	2.6
Investment	9.2	10.1	7.8	6.6	6.1	5.5	5.3	5.1
Research ⁵	2.6	2.6	2.1	1.8	1.9	1.7	1.6	1.5
Structures and equipment	6.7	7.5	5.7	4.8	4.2	3.7	3.7	3.6

See footnotes at end of table.

Table 94 (page 2 of 2). National health expenditures, average annual percent change, and percent distribution, by type of expenditure: United States, selected years 1960–2014

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2015.htm#094>.

[Data are compiled from various sources by the Centers for Medicare & Medicaid Services]

Type of national health expenditure	1960	1970	1980	1990	2000	2012	2013	2014
	Percent distribution							
Personal health care	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Hospital care	38.6	43.1	46.3	40.7	35.8	38.1	38.3	37.9
Professional services	34.1	31.3	29.7	33.7	33.3	31.6	31.4	31.3
Physician and clinical services	23.9	22.7	22.0	25.7	24.8	23.7	23.6	23.5
Other professional services	1.7	1.2	1.6	2.8	3.2	3.3	3.3	3.3
Dental services	8.5	7.5	6.1	5.1	5.3	4.6	4.5	4.4
Other health, residential, and personal care	1.9	2.1	3.9	3.9	5.5	5.8	5.9	5.9
Home health care ¹	0.2	0.3	1.1	2.0	2.8	3.2	3.3	3.2
Nursing care facilities and continuing care retirement communities ¹	3.5	6.4	7.0	7.3	7.3	6.3	6.2	6.1
Retail outlet sales of medical products	21.7	16.8	11.9	12.4	15.3	15.0	15.0	15.6
Prescription drugs	11.5	8.7	5.6	6.5	10.4	10.9	10.9	11.6
Durable medical equipment	3.2	2.8	1.9	2.2	2.2	1.8	1.8	1.8
Other nondurable medical products	7.0	5.3	4.5	3.6	2.7	2.3	2.3	2.2

. . . Category not applicable.

¹Includes expenditures for care in freestanding facilities only. Additional services of this type are provided in hospital-based facilities and are considered hospital care.

²Includes all administrative costs (federal and state and local employees' salaries; contracted employees, including fiscal intermediaries; rent and building costs; computer systems and programs; other materials and supplies; and other miscellaneous expenses) associated with insuring individuals enrolled in the following health insurance programs: Medicare, Medicaid, Children's Health Insurance Program, Department of Defense, Department of Veterans Affairs, Indian Health Service, workers' compensation, maternal and child health, vocational rehabilitation, Substance Abuse and Mental Health Services Administration, and other federal programs.

³Net cost of health insurance is calculated as the difference between calendar year incurred premiums earned and benefits incurred for private health insurance. This includes administrative costs, and in some cases additions to reserves, rate credits and dividends, premium taxes, and net underwriting gains or losses. Also included in this category is the difference between premiums earned and benefits incurred for the private health insurance companies that insure the enrollees of the following programs: Medicare, Medicaid, Children's Health Insurance Program, and workers' compensation (health portion only).

⁴Includes health care services delivered by government public health agencies.

⁵Research and development expenditures of drug companies and other manufacturers and providers of medical equipment and supplies are excluded. These are included in the expenditure class in which the product falls because such expenditures are covered by the payment received for that product.

⁶See Appendix II, Average annual rate of change (percent change).

NOTES: Percents and average annual percent change are calculated using unrounded data. For more information on NHE categories, sources, and methods, see the National Health Expenditure Accounts Methodology Paper, 2014. Available from:

<https://www.cms.gov/Research-Statistics-Data-and-Systems/Statistics-Trends-and-Reports/NationalHealthExpendData/NationalHealthAccountsHistorical.html>. See Appendix I, National Health Expenditure Accounts (NHEA). Data have been revised and differ from previous editions of *Health, United States*.

SOURCE: Centers for Medicare & Medicaid Services, Office of the Actuary, National Health Statistics Group, National Health Expenditure Accounts, National health expenditures. Available from:

<https://www.cms.gov/Research-Statistics-Data-and-Systems/Statistics-Trends-and-Reports/NationalHealthExpendData/NationalHealthAccountsHistorical.html>, accessed on December 12, 2015. See Appendix I, National Health Expenditure Accounts (NHEA).

Table 95 (page 1 of 3). Personal health care expenditures, by source of funds and type of expenditure: United States, selected years 1960–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#095>.

[Data are compiled from various sources by the Centers for Medicare & Medicaid Services]

Type of personal health care expenditure and source of funds	1960	1970	1980	1990	2000	2009	2012	2013	2014
Per capita	\$125	\$300	\$942	\$2,425	\$4,121	\$6,905	\$7,564	\$7,727	\$8,054
	Amount								
	Amount, in billions								
All personal health care expenditures ¹	\$23.3	\$63.1	\$217.0	\$615.3	\$1,162.0	\$2,115.9	\$2,371.8	\$2,441.3	\$2,563.6
Out-of-pocket payments	12.9	25.0	58.1	137.9	199.0	294.6	318.7	325.5	329.8
Health insurance	6.6	29.7	132.1	402.9	844.2	1,636.8	1,841.6	1,893.7	2,000.3
Private health insurance	5.0	14.1	61.5	204.8	406.1	734.6	822.0	834.6	868.8
Medicare	7.3	36.3	107.3	216.3	470.3	534.8	551.2	580.7
Medicaid	5.0	24.7	69.7	186.9	346.2	387.8	407.7	444.9
Federal	2.7	13.7	40.3	109.3	230.6	223.5	235.5	273.6
State and local	2.3	11.0	29.4	77.6	115.6	164.3	172.3	171.3
CHIP ²	2.5	9.5	10.7	11.4	10.8
Federal	1.8	6.7	7.4	7.9	7.6
State and local	0.8	2.8	3.3	3.5	3.3
Other health insurance programs ³	1.7	3.3	9.6	21.2	32.3	76.2	86.4	88.9	95.0
Other third-party payers and programs ⁴	3.7	8.5	26.7	74.5	118.9	184.5	211.5	222.1	233.5
	Deflator (2009 = 100.0)								
Chain-weighted personal health care deflator ⁵	9.3	13.5	28.5	56.3	75.7	100.0	106.8	108.4	109.9
	Percent distribution								
All sources of funds	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Out-of-pocket payments	55.7	39.6	26.8	22.4	17.1	13.9	13.4	13.3	12.9
Health insurance	28.5	47.0	60.9	65.5	72.6	77.4	77.6	77.6	78.0
Private health insurance	21.3	22.3	28.4	33.3	34.9	34.7	34.7	34.2	33.9
Medicare	11.5	16.7	17.4	18.6	22.2	22.5	22.6	22.7
Medicaid	8.0	11.4	11.3	16.1	16.4	16.3	16.7	17.4
Federal	4.3	6.3	6.6	9.4	10.9	9.4	9.6	10.7
State and local	3.7	5.1	4.8	6.7	5.5	6.9	7.1	6.7
CHIP ²	0.2	0.5	0.5	0.5	0.4
Federal	0.2	0.3	0.3	0.3	0.3
State and local	0.1	0.1	0.1	0.1	0.1
Other health insurance programs ³	7.2	5.2	4.4	3.4	2.8	3.6	3.6	3.6	3.7
Other third-party payers and programs ⁴	15.8	13.4	12.3	12.1	10.2	8.7	8.9	9.1	9.1
	Amount, in billions								
Hospital expenditures ⁶	\$9.0	\$27.2	\$100.5	\$250.4	\$415.5	\$778.1	\$902.7	\$933.9	\$971.8
	Percent distribution								
All sources of funds	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Out-of-pocket payments	20.6	9.0	5.4	4.5	3.2	3.3	3.5	3.5	3.2
Health insurance	50.7	71.5	79.7	82.6	86.2	87.5	86.8	86.3	86.5
Private health insurance	35.6	32.5	36.6	38.5	33.9	36.6	37.7	37.5	37.3
Medicare	19.7	26.1	26.9	29.7	27.6	26.4	26.1	25.8
Medicaid	9.7	9.2	10.6	17.1	17.2	16.6	16.7	17.3
Federal	5.2	5.0	6.3	10.3	11.3	9.6	9.6	10.7
State and local	4.5	4.2	4.3	6.8	5.8	7.0	7.1	6.5
CHIP ²	0.2	0.4	0.4	0.4	0.3
Federal	0.2	0.3	0.3	0.3	0.2
State and local	0.1	0.1	0.1	0.1	0.1
Other health insurance programs ³	15.2	9.6	7.8	6.5	5.3	5.8	5.7	5.7	5.9
Other third-party payers and programs ⁴	28.6	19.5	15.0	12.9	10.6	9.2	9.8	10.2	10.3
	Amount, in billions								
Physician and clinical expenditures	\$5.6	\$14.3	\$47.7	\$158.4	\$288.7	\$500.5	\$563.0	\$576.8	\$603.7
	Percent distribution								
All sources of funds	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Out-of-pocket payments	59.5	45.1	29.8	18.9	11.1	9.1	9.3	9.3	9.0
Health insurance	33.1	48.8	59.8	67.8	76.4	80.4	80.0	79.7	79.9
Private health insurance	28.7	29.4	34.8	42.1	47.1	45.9	44.7	43.6	42.2
Medicare	11.5	17.4	19.2	20.3	22.4	22.7	22.8	22.9
Medicaid	4.5	5.1	4.4	6.7	8.0	8.3	9.0	10.6
Federal	2.4	2.9	2.6	3.9	5.5	5.1	5.6	7.4
State and local	2.1	2.2	1.8	2.7	2.5	3.3	3.4	3.2
CHIP ²	0.3	0.6	0.6	0.6	0.5
Federal	0.2	0.4	0.4	0.4	0.4
State and local	0.1	0.2	0.2	0.2	0.2
Other health insurance programs ³	4.4	3.4	2.4	2.1	2.1	3.5	3.6	3.6	3.7
Other third-party payers and programs ⁴	7.4	6.1	10.4	13.3	12.5	10.5	10.7	11.0	11.2

See footnotes at end of table.

Table 95 (page 2 of 3). Personal health care expenditures, by source of funds and type of expenditure: United States, selected years 1960–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#095>.

[Data are compiled from various sources by the Centers for Medicare & Medicaid Services]

Type of personal health care expenditure and source of funds	1960	1970	1980	1990	2000	2009	2012	2013	2014
Amount, in billions									
Dental services expenditures	\$2.0	\$4.7	\$13.3	\$31.6	\$62.1	\$102.3	\$108.9	\$110.4	\$113.5
Percent distribution									
All sources of funds	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Out-of-pocket payments	96.0	90.0	65.8	48.3	44.2	41.6	41.5	41.3	40.3
Health insurance	3.2	9.5	33.4	51.2	55.3	57.9	58.0	58.2	59.3
Private health insurance	1.9	4.5	28.4	47.9	50.3	48.3	47.4	47.3	47.6
Medicare	0.0	0.1	0.3	0.3	0.4	0.4
Medicaid	3.4	3.8	2.4	3.9	7.5	7.9	8.0	8.9
Federal	1.8	2.1	1.3	2.2	5.1	4.6	4.6	5.6
State and local	1.6	1.7	1.0	1.7	2.4	3.3	3.4	3.3
CHIP ²	0.4	0.7	1.2	1.4	1.3
Federal	0.3	0.5	0.8	0.9	0.9
State and local	0.1	0.2	0.4	0.4	0.4
Other health insurance programs ³	1.3	1.6	1.2	0.9	0.6	1.1	1.2	1.1	1.2
Other third-party payers and programs ⁴	0.8	0.4	0.8	0.6	0.6	0.5	0.5	0.4	0.4
Amount, in billions									
Home health care expenditures ⁷	\$0.1	\$0.2	\$2.4	\$12.5	\$32.3	\$67.4	\$76.9	\$79.4	\$83.2
Percent distribution									
All sources of funds	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Out-of-pocket payments	12.3	9.5	15.3	17.8	19.2	8.3	8.5	8.7	8.9
Health insurance	5.3	37.7	53.7	66.2	71.8	88.4	88.2	87.9	87.9
Private health insurance	1.8	3.2	14.7	22.8	24.0	7.4	8.1	8.9	9.9
Medicare	26.8	26.8	26.0	26.5	45.0	43.5	42.3	41.7
Medicaid	6.8	11.6	17.1	20.9	35.6	36.1	36.1	35.6
Federal	3.2	6.2	9.2	11.3	23.2	19.8	19.9	20.1
State and local	3.2	5.4	8.0	9.6	12.4	16.3	16.2	15.5
CHIP ²	0.0	0.0	0.0	0.0	0.0
Federal	0.0	0.0	0.0	0.0	0.0
State and local	0.0	0.0	0.0	0.0	0.0
Other health insurance programs ³	3.5	1.4	0.5	0.3	0.3	0.4	0.5	0.5	0.5
Other third-party payers and programs ⁴	80.7	52.7	31.1	16.0	9.0	3.3	3.3	3.4	3.3
Amount, in billions									
Nursing care facilities and continuing care retirement communities expenditures ⁸	\$0.8	\$4.0	\$15.3	\$44.7	\$85.0	\$136.9	\$148.3	\$150.2	\$155.6
Percent distribution									
All sources of funds	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Out-of-pocket payments	74.4	49.2	40.5	40.3	31.9	27.9	27.1	27.2	26.5
Health insurance	28.5	51.9	49.0	61.2	65.5	66.0	65.9	66.3
Private health insurance	0.2	1.3	6.2	8.8	7.4	8.0	8.0	8.4
Medicare	3.5	2.0	3.8	12.7	22.0	22.9	22.8	22.9
Medicaid	23.3	46.2	36.7	37.5	33.2	32.1	32.0	31.9
Federal	12.5	26.1	20.7	21.8	22.1	18.4	18.3	18.3
State and local	10.8	20.1	16.1	15.7	11.1	13.7	13.8	13.6
CHIP ²	0.0	0.0	0.0	0.0	0.0
Federal	0.0	0.0	0.0	0.0	0.0
State and local	0.0	0.0	0.0	0.0	0.0
Other health insurance programs ³	0.0	1.5	2.4	2.2	2.2	2.9	3.0	3.0	3.0
Other third-party payers and programs ⁴	25.5	22.3	7.6	10.8	6.9	6.6	6.9	7.0	7.3
Amount, in billions									
Prescription drug expenditures	\$2.7	\$5.5	\$12.0	\$40.3	\$121.0	\$252.7	\$259.1	\$265.3	\$297.7
Percent distribution									
All sources of funds	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Out-of-pocket payments	96.0	82.4	71.3	56.8	27.8	19.4	17.5	16.4	15.0
Health insurance	1.5	16.5	26.9	40.3	70.3	79.2	81.5	82.7	84.3
Private health insurance	1.3	8.8	15.0	27.0	50.5	46.0	43.7	43.1	42.8
Medicare	0.5	1.7	21.6	26.0	27.8	29.0
Medicaid	7.6	11.7	12.6	16.3	8.0	8.1	8.3	9.2
Federal	4.1	6.8	7.2	9.3	5.4	4.5	4.6	5.9
State and local	3.5	4.9	5.4	7.0	2.7	3.7	3.7	3.2
CHIP ²	0.2	0.5	0.6	0.5	0.5
Federal	0.2	0.4	0.4	0.4	0.3
State and local	0.1	0.2	0.2	0.2	0.1
Other health insurance programs ³	0.1	0.1	0.2	0.2	1.5	3.1	3.1	2.9	2.9
Other third-party payers and programs ⁴	2.5	1.1	1.8	3.0	1.9	1.4	1.0	0.9	0.7

See footnotes at end of table.

Table 95 (page 3 of 3). Personal health care expenditures, by source of funds and type of expenditure: United States, selected years 1960–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#095>.

[Data are compiled from various sources by the Centers for Medicare & Medicaid Services]

Type of personal health care expenditure and source of funds	1960	1970	1980	1990	2000	2009	2012	2013	2014
Amount, in billions									
All other personal health care expenditures ⁹	\$3.2	\$7.1	\$25.7	\$77.3	\$157.3	\$278.1	\$312.9	\$325.2	\$338.1
Percent distribution									
All sources of funds	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Out-of-pocket payments	84.6	74.2	56.9	49.7	37.6	31.6	31.3	31.4	31.2
Health insurance	4.1	9.1	25.7	33.7	45.0	52.2	53.6	53.9	54.0
Private health insurance	3.1	4.6	7.8	13.0	13.7	14.2	14.9	14.6	14.6
Medicare	1.0	2.8	5.5	8.1	10.3	10.6	10.4	10.3
Medicaid	3.0	14.8	15.0	22.8	27.0	27.4	28.3	28.5
Federal	1.6	8.1	8.5	13.0	18.0	15.6	16.1	16.5
State and local	1.4	6.7	6.4	9.8	9.0	11.8	12.1	12.0
CHIP ²	0.2	0.4	0.4	0.4	0.4
Federal	0.1	0.3	0.3	0.3	0.3
State and local	0.1	0.1	0.1	0.1	0.1
Other health insurance programs ³	1.1	0.5	0.3	0.2	0.2	0.2	0.3	0.3	0.3
Other third-party payers and programs ⁴	11.2	16.7	17.4	16.6	17.4	16.2	15.1	14.7	14.8

... Category not applicable.

0.0 Quantity more than zero but less than 0.05.

¹Includes all expenditures other than expenses for government administration, net cost of health insurance, public health activities, research, and structures and equipment.

²The Children's Health Insurance Program (CHIP) including Medicaid CHIP expansions.

³Includes Department of Defense and Department of Veterans Affairs.

⁴Includes worksite health care, other private revenues, Indian Health Service, workers' compensation, general assistance, maternal and child health, vocational rehabilitation, other federal programs, Substance Abuse and Mental Health Services Administration, other state and local programs, and school health.

⁵The personal health care deflator is calculated as a chain-weighted price index using the Producer Price Indexes for hospitals, offices of physicians, medical and diagnostic laboratories, home health care services, and nursing care facilities; and Consumer Price Indices specific to each of the remaining personal health care components. For more information on the detailed price series recommended for deflating each category of spending see the National Health Expenditure Accounts Methodology Paper, 2014 and NHE Deflator Methodology paper. Available from:

<https://www.cms.gov/Research-Statistics-Data-and-Systems/Statistics-Trends-and-Reports/NationalHealthExpendData/NationalHealthAccountsHistorical.html>.

⁶Includes expenditures for hospital-based nursing home and home health agency care.

⁷Includes expenditures for care in freestanding facilities only. Additional services of this type are provided in hospital-based facilities and are considered hospital care.

⁸Includes expenditures for care in freestanding nursing homes. Expenditures for care in hospital-based nursing homes are included with hospital care.

⁹Includes expenditures for other professional services, other nondurable medical products, durable medical equipment, and other health, residential, and personal care, not shown separately. See Appendix II, Health expenditures, national.

NOTES: Percents may not add to totals because of rounding. Census resident-based population less armed forces overseas and population of outlying areas used to calculate per capita. The Medicare and Medicaid programs began coverage in 1965. The Children's Health Insurance Program began coverage in 1997. For more information on NHE sources and methods, see the National Health Expenditure Accounts Methodology Paper, 2014. Available from:

<https://www.cms.gov/Research-Statistics-Data-and-Systems/Statistics-Trends-and-Reports/NationalHealthExpendData/Downloads/DSM-14.pdf>. See Appendix I, National Health Expenditure Accounts (NHEA). Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at:

<http://www.cdc.gov/nchs/hus.htm>. Data have been revised and differ from previous editions of *Health, United States*.

SOURCE: Centers for Medicare & Medicaid Services, Office of the Actuary, National Health Statistics Group, National Health Expenditure Accounts, National health expenditures. Available from:

<https://www.cms.gov/Research-Statistics-Data-and-Systems/Statistics-Trends-and-Reports/NationalHealthExpendData/NationalHealthAccountsHistorical.html>, accessed on December 12, 2015. Martin AB, Hartman M, Benson J, Catlin A. National health spending in 2014: Faster growth driven by coverage expansion and prescription drug spending. *Health Aff* 2015;35(1):1–11. See Appendix I, National Health Expenditure Accounts (NHEA).

Table 96 (page 1 of 3). Cost of hospital discharges with common hospital operating room procedures in nonfederal community hospitals, by age and selected principal procedure: United States, selected years 2000–2013

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2015.htm#096>.

[National estimates are compiled by the Agency for Healthcare Research and Quality using discharge data from participating states]

Age and principal operating room procedure ¹	Mean inflation-adjusted cost per hospitalization: 2013 dollars ²			Number of discharges with operating room principal procedure			Total inflation-adjusted national costs: 2013 dollars (in millions) ²		
	2000	2005	2013	2000	2005	2013	2000	2005	2013
All ages									
Hospital discharges with an operating room principal procedure ³	\$13,541	\$16,248	\$18,780	8,743,631	9,964,151	8,414,072	\$117,663	\$162,038	\$157,891
Laminectomy (back surgery)	8,345	9,448	14,064	285,636	248,800	167,380	2,393	2,352	2,363
Heart valve procedures	43,829	53,711	51,415	79,719	93,802	102,425	3,488	5,058	5,264
Coronary artery bypass graft (CABG)	31,952	38,907	41,274	337,972	221,325	157,720	10,836	8,625	6,510
Percutaneous coronary angioplasty (PTCA) (balloon angioplasty of heart)	15,314	18,871	20,924	581,183	727,912	403,550	8,899	13,743	8,454
Insertion, revision, replacement, removal of cardiac pacemaker or cardioverter/defibrillator	28,254	36,181	35,074	66,286	160,629	87,465	1,887	5,804	3,067
Colorectal resection (removal of part of the bowel)	19,887	23,171	23,446	253,780	274,599	237,285	5,152	6,371	5,562
Appendectomy	7,483	8,685	10,190	269,089	298,829	201,700	1,991	2,594	2,061
Cholecystectomy (gall bladder removal)	10,622	12,356	13,090	389,079	376,158	318,145	4,097	4,649	4,168
Hysterectomy	6,671	7,423	10,122	580,019	550,659	211,050	3,842	4,094	2,138
Cesarean section	5,547	5,579	6,054	898,859	1,258,990	1,136,704	4,869	7,028	6,887
Treatment, fracture or dislocation of hip and femur	12,844	15,610	17,177	237,615	251,071	241,510	3,103	3,917	4,148
Arthroplasty knee (knee replacement)	14,106	15,954	16,497	318,854	533,216	700,740	4,471	8,510	11,563
Hip replacement	15,296	17,459	17,337	295,940	369,634	457,195	4,582	6,447	7,927
Spinal fusion	17,802	25,373	28,696	204,320	322,610	405,245	3,551	8,194	11,635
Under 18 years									
Hospital discharges with an operating room principal procedure ³	13,589	19,768	26,307	382,455	535,472	312,630	5,022	10,539	8,185
Incision and excision of CNS (a type of brain surgery)	29,501	35,818	44,842	6,352	11,436	7,385	181	411	328
Tonsillectomy and/or adenoidectomy	4,517	5,844	7,381	12,045	16,288	9,830	56	96	73
Small bowel resection (removal of part of the small bowel)	37,050	51,668	51,450	1,712	2,993	1,735	62	153	89
Appendectomy	6,713	8,297	9,894	75,481	85,790	52,160	493	712	516
Cesarean section	6,159	5,852	6,173	23,690	28,609	13,735	132	167	85
Spinal fusion	29,930	47,106	56,920	7,463	12,880	10,580	221	601	599
18–44 years									
Hospital discharges with an operating room principal procedure ³	8,948	10,127	11,969	2,806,078	3,101,461	2,392,184	24,568	31,439	28,611
Incision and excision of CNS (a type of brain surgery)	26,014	31,206	36,661	19,510	18,229	17,505	488	572	641
Laminectomy (back surgery)	7,509	8,799	13,129	95,687	67,321	31,755	723	593	419
Appendectomy	6,892	7,902	9,146	133,662	135,504	84,790	908	1,071	778
Cholecystectomy (gall bladder removal)	8,700	9,617	10,671	132,538	128,987	112,165	1,107	1,242	1,200
Oophorectomy (removal of one or both ovaries)	6,486	7,552	10,650	38,252	33,384	19,015	251	252	202
Ligation of fallopian tubes (“tying” of fallopian tubes)	4,796	4,620	7,150	75,221	74,449	34,365	340	344	243
Hysterectomy	6,186	6,740	9,231	291,704	255,025	83,310	1,779	1,721	769
Cesarean section	5,530	5,569	6,045	873,231	1,226,170	1,119,309	4,725	6,833	
Treatment, fracture or dislocation of lower extremity (other than hip or femur)	9,533	12,103	16,582	68,015	59,527	46,080	637	720	763
Spinal fusion	16,735	23,501	26,766	73,228	87,276	67,645	1,185	2,052	1,811

See footnotes at end of table.

Table 96 (page 2 of 3). Cost of hospital discharges with common hospital operating room procedures in nonfederal community hospitals, by age and selected principal procedure: United States, selected years 2000–2013

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2015.htm#096>.

[National estimates are compiled by the Agency for Healthcare Research and Quality using discharge data from participating states]

Age and principal operating room procedure ¹	Mean inflation-adjusted cost per hospitalization: 2013 dollars ²			Number of discharges with operating room principal procedure			Total inflation-adjusted national costs: 2013 dollars (in millions) ²		
	2000	2005	2013	2000	2005	2013	2000	2005	2013
45–64 years									
Hospital discharges with an operating room principal procedure ³	\$14,812	\$17,822	\$20,923	2,435,212	2,909,930	2,653,754	\$35,913	\$51,931	\$55,450
Laminectomy (back surgery)	8,417	9,328	14,619	107,720	96,084	66,415	908	897	974
Heart valve procedures.	41,166	49,216	49,346	22,849	26,650	26,270	935	1,319	1,297
Coronary artery bypass graft (CABG).	29,854	35,640	39,557	139,897	94,742	65,095	4,196	3,384	2,575
Percutaneous coronary angioplasty (PTCA) (balloon angioplasty of heart)	14,822	18,223	20,386	252,151	318,811	179,935	3,730	5,814	3,675
Insertion, revision, replacement, removal of cardiac pacemaker or cardioverter/defibrillator.	34,603	39,265	37,353	15,957	44,029	25,785	548	1,728	962
Colorectal resection (removal of part of the bowel)	17,889	20,639	22,075	76,604	95,108	89,920	1,398	1,966	1,985
Cholecystectomy (gall bladder removal)	10,019	11,897	12,981	117,432	117,636	103,040	1,177	1,402	1,339
Oophorectomy	7,765	8,885	12,913	21,232	22,459	23,285	165	199	300
Hysterectomy	6,805	7,568	10,170	231,498	242,030	99,070	1,573	1,835	1,008
Arthroplasty knee (knee replacement)	14,430	16,008	16,680	95,902	199,682	288,955	1,377	3,197	4,820
Hip replacement.	15,897	17,707	17,229	65,118	105,138	161,640	1,043	1,859	2,785
Spinal fusion	17,087	23,584	27,182	87,388	150,313	192,890	1,454	3,548	5,246
65–74 years									
Hospital discharges with an operating room principal procedure ³	16,594	19,819	21,637	1,511,467	1,602,345	1,614,725	25,212	31,808	34,922
Laminectomy (back surgery)	8,818	9,418	13,755	45,976	45,731	39,185	406	431	541
Heart valve procedures.	44,858	54,620	50,611	23,236	24,774	27,265	1,033	1,357	1,380
Coronary artery bypass graft (CABG).	32,514	39,683	40,825	112,652	70,404	55,040	3,661	2,796	2,247
Percutaneous coronary angioplasty (PTCA) (balloon angioplasty of heart)	15,247	18,693	20,981	166,497	196,855	108,310	2,534	3,682	2,274
Insertion, revision, replacement, removal of cardiac pacemaker or cardioverter/defibrillator.	30,868	37,338	36,361	19,096	44,900	23,365	592	1,675	848
Endarterectomy (plaque removal from artery lining brain, head, neck)	8,886	9,534	10,707	51,292	40,715	30,710	467	389	329
Colorectal resection (removal of part of the bowel)	19,999	23,474	23,221	63,693	62,337	59,650	1,313	1,466	1,384
Cholecystectomy (gall bladder removal)	11,736	14,096	14,976	65,953	55,566	48,440	787	783	726
Arthroplasty knee (knee replacement)	14,367	15,905	16,219	110,961	177,306	255,495	1,579	2,823	4,146
Hip replacement.	15,237	17,168	17,156	71,986	86,918	125,935	1,113	1,491	2,160
Spinal fusion	18,817	27,237	29,430	23,419	47,031	93,200	438	1,282	2,744
75–84 years									
Hospital discharges with an operating room principal procedure ³	16,844	20,484	21,837	1,224,573	1,360,428	1,036,865	20,927	27,902	22,647
Laminectomy (back surgery)	9,514	10,333	13,278	31,059	31,945	23,195	298	330	310
Heart valve procedures.	46,270	57,554	51,773	21,004	25,070	28,635	984	1,449	1,482
Coronary artery bypass graft (CABG).	35,253	44,112	45,346	68,750	45,187	30,025	2,443	1,995	1,361
Percutaneous coronary angioplasty (PTCA) (balloon angioplasty of heart)	16,143	19,976	22,065	111,169	144,906	70,560	1,806	2,895	1,558
Insertion, revision, replacement, removal of cardiac pacemaker or cardioverter/defibrillator.	25,320	35,091	34,053	19,975	48,557	22,500	515	1,701	767
Endarterectomy (plaque removal from artery lining brain, head, neck)	9,230	9,884	11,111	45,337	38,073	23,750	432	377	264
Colorectal resection (removal of part of the bowel)	21,810	26,029	25,187	62,096	61,238	41,135	1,391	1,594	1,035
Cholecystectomy (gall bladder removal)	13,450	16,626	16,666	52,448	49,823	35,420	720	827	589
Treatment, fracture or dislocation of hip and femur	12,095	14,547	16,064	73,332	72,878	63,090	913	1,062	1,014
Arthroplasty knee (knee replacement)	14,360	16,007	16,381	79,138	121,865	125,790	1,138	1,951	2,061
Hip replacement.	15,053	17,326	17,378	92,715	105,580	97,705	1,418	1,828	1,698
Spinal fusion	19,579	28,418	30,094	11,770	22,902	36,895	229	651	1,110

See footnotes at end of table.

Table 96 (page 3 of 3). Cost of hospital discharges with common hospital operating room procedures in nonfederal community hospitals, by age and selected principal procedure: United States, selected years 2000–2013

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2015.htm#096>.

[National estimates are compiled by the Agency for Healthcare Research and Quality using discharge data from participating states]

Age and principal operating room procedure ¹	Mean inflation-adjusted cost per hospitalization: 2013 dollars ²			Number of discharges with operating room principal procedure			Total inflation-adjusted national costs: 2013 dollars (in millions) ²		
	2000	2005	2013	2000	2005	2013	2000	2005	2013
85 years and over									
Hospital discharges with an operating room principal procedure ³	\$15,444	\$18,846	\$19,999	382,341	434,936	402,235	\$5,997	\$8,204	\$8,054
Heart valve procedures	48,641	61,736	52,006	2,985	3,957	10,650	145	244	553
Coronary artery bypass graft (CABG)	39,659	50,802	47,898	5,280	4,174	3,345	207	213	160
Percutaneous coronary angioplasty (PTCA) (balloon angioplasty of heart)	18,321	22,279	22,408	16,682	28,942	21,740	303	644	487
Insertion, revision, replacement, removal of cardiac pacemaker or cardioverter/defibrillator.	15,027	25,544	26,001	7,071	13,675	9,175	109	348	240
Colorectal resection (removal of part of the bowel)	23,568	27,864	25,578	20,729	20,456	15,935	499	570	408
Cholecystectomy (gall bladder removal)	16,368	18,461	17,606	15,698	16,732	14,080	260	308	248
Treatment, fracture or dislocation of hip and femur	11,777	13,968	15,627	76,900	77,692	76,795	932	1,087	1,202
Arthroplasty knee	14,559	16,813	17,611	10,122	15,784	16,880	148	265	297
Hip replacement	14,638	17,397	17,737	50,005	53,975	54,380	743	938	965
Amputation of lower extremity (amputation of leg, foot or toe)	13,511	17,465	17,250	12,855	10,029	7,535	176	175	130

¹Data are based on valid operating room procedures. Operating room procedures were identified using the Centers for Medicare & Medicaid Services' Diagnosis Related Groups (DRGs). For DRGs, physician panels identified *International Classification of Diseases* (ICD–9-CM) procedure codes that would be performed in operating rooms in most hospitals. Operating room procedures, as defined by DRGs, are classified by the Clinical Classifications Software (CCS) into 1 of 231 clinically meaningful categories. Mean costs per hospitalization are based on the principal procedure as determined by the CCS. The number of discharges is based on the first-listed (principal) major procedure. See Appendix II, Procedure.

²Charges (the amount billed by the hospital) were converted to costs using cost-charge ratios from the Centers for Medicare & Medicaid Services. Costs are for the entire hospitalization including the principal procedure. Costs were adjusted for inflation to 2013 dollars using the gross domestic product deflator Table 1.1.4. Price Indexes downloaded from <https://www.bea.gov/iTable/iTable.cfm?ReqID=9&step=1> on October, 13, 2015. See Appendix II, Cost-charge ratio.

³Includes discharges for operating room principal procedures not shown separately.

NOTES: Excludes newborn infants. The number of states participating in the sample varied over time from 28 states in 2000 to 46 in 2011, 44 in 2012, and 43 states and D.C. in 2013. See Appendix I, Healthcare Cost and Utilization Project (HCUP), National (Nationwide) Inpatient Sample, for a list of states available in each year. In 2012, the HCUP-NIS was redesigned and changed from a sample of hospitals to a sample of discharges from all participating community hospitals. For this report, the statistics for years prior to 2012 were regenerated using new trend weights taking into account the 2012 redesign. For more information on the 2012 redesign, see: Houchens R, Ross D, Elixhauser A, Jiang J. Nationwide Inpatient Sample (NIS) redesign final report. 2014. HCUP methods series report # 2014–04 ONLINE. April 4, 2014. U.S. Agency for Healthcare Research and Quality. Available from: <http://www.hcup-us.ahrq.gov/reports/methods/2014-04.pdf>. The estimates are weighted to provide national estimates. Standard errors are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>. Data have been revised and differ from previous editions of *Health, United States*.

SOURCE: Agency for Healthcare Research and Quality, Healthcare Cost and Utilization Project, National (Nationwide) Inpatient Sample. See Appendix I, Healthcare Cost and Utilization Project (HCUP), National (Nationwide) Inpatient Sample.

Table 97 (page 1 of 3). Expenses for health care and prescribed medicine, by selected population characteristics: United States, selected years 1987–2012

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hs/content2015.htm#097>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population and a sample of medical providers]

Characteristic	Total expenses ¹										
	Population in millions ²			Percent of persons with expense				Mean annual expense per person with expense ³			
	1997	2000	2012	1987	1997	2000	2012	1987	1997	2000	2012
All ages	271.3	278.4	313.5	84.5	84.1	83.5	84.7	\$3,147	\$3,467	\$3,600	\$5,089
Under 65 years:											
Total	237.1	243.6	268.2	83.2	82.5	81.8	82.7	2,446	2,630	2,836	4,186
Under 6 years	23.8	24.1	24.1	88.9	88.0	86.7	88.8	2,087	1,227	1,498	2,145
6–17 years	48.1	48.4	49.8	80.2	81.7	80.0	85.7	1,360	1,378	1,490	2,064
18–44 years	108.9	109.0	112.0	81.5	78.3	77.7	75.5	2,149	2,383	2,540	3,512
45–64 years	56.3	62.1	82.3	87.0	89.2	88.5	88.9	4,170	4,614	4,750	6,799
Sex											
Male	118.0	120.9	133.1	78.8	77.6	76.6	77.6	2,311	2,376	2,715	4,117
Female	119.1	122.7	135.2	87.5	87.4	87.0	87.8	2,563	2,852	2,941	4,247
Hispanic origin and race ⁴											
Hispanic or Latino	29.4	32.0	50.2	71.0	69.5	69.0	72.5	1,957	2,188	1,931	3,240
Not Hispanic or Latino:											
White	166.2	169.2	162.5	86.9	87.2	86.6	87.6	2,452	2,820	2,967	4,585
Black or African American	31.3	32.1	33.9	72.2	72.1	71.3	77.5	2,973	2,108	3,012	3,752
Asian	14.4	75.2	3,221
American Indian, Alaska Native, Native Hawaiian, Other Pacific Islander, and Multiple Race	7.2	83.2	4,104
Insurance status ⁵											
Any private insurance	174.0	181.6	174.5	86.5	86.5	85.9	87.8	2,345	2,679	2,702	4,424
Public insurance only	29.8	29.7	54.2	82.4	83.3	83.6	86.0	3,948	3,196	4,305	4,347
Uninsured all year	33.3	32.3	39.6	61.8	61.1	57.3	55.8	1,507	1,570	2,000	2,198
65 years and over:											
Total	34.2	34.8	45.3	93.7	95.2	95.5	96.3	7,787	8,507	8,187	9,678
Sex											
Male	14.6	15.0	20.1	92.0	94.5	93.4	96.1	7,968	9,560	8,778	9,703
Female	19.6	19.8	25.1	94.9	95.7	97.1	96.5	7,664	7,732	7,756	9,658
Hispanic origin and race ⁴											
Hispanic or Latino	1.7	1.9	3.3	82.5	94.2	92.5	93.4	7,407	8,901	7,346	7,224
Not Hispanic or Latino:											
White	28.8	28.9	35.4	94.9	95.9	95.9	97.2	7,665	8,550	8,311	10,022
Black or African American	2.8	2.9	4.0	88.5	92.2	94.0	94.9	9,413	8,378	7,873	9,108
Asian	1.7	90.6	7,677
American Indian, Alaska Native, Native Hawaiian, Other Pacific Islander, and Multiple Race	*	*	*
Insurance status ⁶											
Medicare only	8.8	12.0	16.9	85.9	92.1	94.8	95.3	6,132	7,837	7,028	8,322
Medicare and private insurance	21.7	19.2	21.4	95.4	97.0	96.0	97.4	7,705	8,297	8,394	9,818
Medicare and other public insurance	3.2	3.2	6.3	94.4	93.2	96.3	97.0	11,978	11,991	11,233	13,299

See footnotes at end of table.

Table 97 (page 2 of 3). Expenses for health care and prescribed medicine, by selected population characteristics: United States, selected years 1987–2012

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#097>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population and a sample of medical providers]

Characteristic	Prescribed medicine expenses ⁷							
	Percent of persons with expense				Mean annual out-of-pocket expenses for prescribed medicine per person with prescribed medicine expenses ³			
	1987	1997	2000	2012	1987	1997	2000	2012
All ages	57.3	62.1	62.3	62.4	\$186	\$289	\$365	\$287
Under 65 years:								
Total	54.0	58.7	58.5	57.6	137	205	265	243
Under 6 years	61.8	61.3	56.9	51.1	48	50	49	31
6–17 years	44.3	48.2	46.2	46.0	92	77	93	102
18–44 years	51.3	55.9	56.0	51.8	107	174	202	190
45–64 years	65.3	71.8	73.3	74.3	261	380	500	389
Sex								
Male	46.5	51.5	51.3	51.0	127	182	233	250
Female	61.4	65.8	65.6	64.1	145	222	290	238
Hispanic origin and race ⁴								
Hispanic or Latino	41.6	47.7	45.0	45.1	100	136	195	147
Not Hispanic or Latino:								
White	57.7	63.1	63.8	63.7	144	221	285	273
Black or African American	44.1	50.0	47.6	52.7	122	165	218	188
Asian	42.7	203
American Indian, Alaska Native, Native Hawaiian, Other Pacific Islander, and Multiple Race	58.7	323
Insurance status ⁵								
Any private insurance	56.5	61.6	61.6	61.4	142	194	227	261
Public insurance only	56.5	62.0	62.4	60.0	95	201	380	136
Uninsured all year	35.1	40.2	37.6	37.3	151	295	440	352
65 years and over:								
Total	81.6	86.0	88.3	90.8	428	691	831	451
Sex								
Male	78.0	82.8	83.9	90.7	399	622	623	468
Female	84.0	88.3	91.5	90.9	447	739	975	437
Hispanic origin and race ⁴								
Hispanic or Latino	74.7	87.5	83.9	87.3	567	563	699	262
Not Hispanic or Latino:								
White	82.3	86.7	89.0	92.0	436	714	862	482
Black or African American	79.5	85.3	85.3	89.2	335	574	710	360
Asian	78.4	372
American Indian, Alaska Native, Native Hawaiian, Other Pacific Islander, and Multiple Race	*	*
Insurance status ⁶								
Medicare only	70.6	82.1	87.7	88.7	473	799	992	461
Medicare and private insurance	83.4	88.1	89.0	92.6	445	701	768	502
Medicare and other public coverage	88.2	85.0	88.5	92.4	161	387	658	263

See footnotes at end of table.

Table 97 (page 3 of 3). Expenses for health care and prescribed medicine, by selected population characteristics: United States, selected years 1987–2012

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#097>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population and a sample of medical providers]

... Category not applicable.

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error equal to or greater than 30%. Data not shown if based on fewer than 100 sample cases.

¹Includes expenses for inpatient hospital and physician services, ambulatory physician and nonphysician services, prescribed medicines, home health services, dental services, and other medical equipment, supplies, and services that were purchased or rented during the year. Excludes expenses for over-the-counter medications, phone contacts with health providers, and premiums for health insurance.

²Includes persons in the civilian noninstitutionalized population for all or part of the year. Expenditures for persons in this population for only part of the year are restricted to those incurred during periods of eligibility (e.g., expenses incurred during periods of institutionalization and military service are not included in estimates).

³Estimates of expenses were converted to 2012 dollars using the Consumer Price Index (all items). See Appendix II, Consumer Price Index (CPI).

⁴Persons of Hispanic origin may be of any race. Estimates for Asian persons as well as for American Indian, Alaska Native, Native Hawaiian, Other Pacific Islander, and Multiple Race persons are not available for years prior to 2002 because Asian persons could not be distinguished separately and multiple race information was not collected.

⁵Any private insurance includes individuals with insurance that provided coverage for hospital and physician care at any time during the year, other than Medicare, Medicaid, or other public coverage for hospital or physician services. Public insurance only includes individuals who were not covered by private insurance at any time during the year but were covered by Medicare, Medicaid, other public coverage for hospital or physician services, and/or CHAMPUS/CHAMPVA (TRICARE) at any point during the year. Uninsured includes persons not covered by either private or public insurance throughout the entire year or period of eligibility for the survey. Individuals with Indian Health Service coverage only are considered uninsured.

⁶Populations do not add to total because uninsured persons and persons with unknown insurance status were excluded.

⁷Includes expenses for all prescribed medications that were purchased or refilled during the survey year.

NOTES: Estimates for 1987 are based on the National Medical Expenditure Survey (NMES); estimates for other years are based on the Medical Expenditure Panel Survey (MEPS). Because expenditures in NMES were based primarily on charges and those for MEPS were based on payments, NMES data were adjusted to be more comparable with MEPS by using estimated charge-to-payment ratios for 1987. Overall, this resulted in an approximate 11% reduction from the unadjusted 1987 NMES expenditure estimates. For a detailed explanation of this adjustment, see Zuvekas S, Cohen J. A guide to comparing health care expenditures in the 1996 MEPS to the 1987 NMES. *Inquiry* 2002;39(1):76–86. See Appendix I, Medical Expenditure Panel Survey (MEPS). Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>. Data have been revised and differ from previous editions of *Health, United States*.

SOURCE: Agency for Healthcare Research and Quality, Center for Financing, Access, and Cost Trends. 1987 National Medical Expenditure Survey and 1996–2012 Medical Expenditure Panel Surveys. See Appendix I, Medical Expenditure Panel Survey (MEPS).

Table 98 (page 1 of 3). Sources of payment for health care, by selected population characteristics: United States, selected years 1987–2012

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hs/content2015.htm#098>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population and a sample of medical providers]

Characteristic	All sources	Source of payment for health care							
		Out of pocket				Private insurance ¹			
		1987	1997	2000	2012	1987	1997	2000	2012
		Percent distribution							
All ages	100.0	24.8	19.4	19.4	14.1	36.6	40.3	40.3	41.5
Under 65 years:									
Total	100.0	26.2	21.1	20.3	14.7	46.6	53.1	52.5	53.9
Under 6 years	100.0	18.5	14.2	10.3	6.2	39.5	49.3	51.2	46.2
6–17 years	100.0	35.7	29.0	27.7	19.6	47.3	53.2	48.8	51.5
18–44 years	100.0	27.4	21.1	19.9	15.7	46.8	52.9	51.2	55.4
45–64 years	100.0	24.0	20.1	20.2	14.1	47.8	53.6	54.5	54.2
Sex									
Male	100.0	24.5	21.3	18.1	13.6	44.6	50.3	52.2	52.8
Female	100.0	27.5	21.0	22.1	15.7	48.1	55.1	52.7	54.8
Hispanic origin and race ²									
Hispanic or Latino	100.0	22.0	18.8	20.5	11.4	36.1	42.3	45.8	41.7
Not Hispanic or Latino:									
White	100.0	28.2	21.8	21.7	16.0	50.1	55.8	55.1	58.3
Black or African American	100.0	15.5	17.1	11.8	10.4	30.0	42.3	40.5	41.9
Asian	100.0	15.2	54.4
American Indian, Alaska Native, Native Hawaiian, Other Pacific Islander, and Multiple Race	100.0	15.0	43.1
Insurance status ³									
Any private insurance	100.0	29.0	21.6	21.2	15.9	60.0	67.6	70.2	73.8
Public insurance only	100.0	8.9	10.6	9.8	5.5
Uninsured all year	100.0	40.6	41.3	40.4	37.3
65 years and over	100.0	22.0	16.3	17.5	12.8	15.8	16.5	14.9	14.1
Sex									
Male	100.0	21.7	14.2	14.2	11.7	17.6	20.1	16.8	14.3
Female	100.0	22.2	18.1	20.2	13.7	14.4	13.2	13.3	13.9
Hispanic origin and race ²									
Hispanic or Latino	100.0	*13.5	13.6	13.9	6.6	*4.7	5.9	8.4	6.0
Not Hispanic or Latino:									
White	100.0	23.7	17.0	18.3	13.8	16.7	17.9	15.2	14.6
Black or African American	100.0	11.2	11.4	13.6	9.9	*11.9	8.8	9.3	12.6
Asian	100.0	9.7	22.4
American Indian, Alaska Native, Native Hawaiian, Other Pacific Islander, and Multiple Race	100.0	*	*
Insurance status									
Medicare only	100.0	29.8	19.8	22.2	15.3
Medicare and private insurance	100.0	23.4	17.3	17.0	14.5	18.9	25.7	25.3	28.3
Medicare and other public coverage	100.0	*6.2	5.2	9.1	4.6

See footnotes at end of table.

Table 98 (page 2 of 3). Sources of payment for health care, by selected population characteristics: United States, selected years 1987–2012

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hs/content2015.htm#098>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population and a sample of medical providers]

Characteristic	Source of payment for health care							
	Public sources ⁴				Other ⁵			
	1987	1997	2000	2012	1987	1997	2000	2012
	Percent distribution							
All ages	34.1	34.4	35.4	39.8	4.5	5.9	5.0	4.6
Under 65 years:								
Total	21.3	18.1	21.3	26.1	6.0	7.7	6.0	5.2
Under 6 years	35.8	25.4	33.6	40.8	6.2	11.2	4.9	6.8
6–17 years	11.8	14.1	20.1	27.2	5.2	3.7	3.4	1.7
18–44 years	19.4	15.7	21.1	22.9	6.4	10.3	7.8	6.0
45–64 years	22.4	20.3	20.2	26.5	5.8	6.0	5.2	5.2
Sex								
Male	23.9	19.5	23.5	28.8	7.1	8.9	6.3	4.8
Female	19.2	17.0	19.5	23.8	5.2	6.8	5.7	5.6
Hispanic origin and race ²								
Hispanic or Latino	35.8	28.9	27.5	39.4	6.0	10.0	6.2	7.6
Not Hispanic or Latino:								
White	15.9	15.3	18.0	21.0	5.8	7.1	5.2	4.7
Black or African American	47.2	30.7	38.8	41.1	7.3	9.9	8.8	6.5
Asian	*26.0	*4.4
American Indian, Alaska Native, Native Hawaiian, Other Pacific Islander, and Multiple Race	38.5	*3.4
Insurance status ³								
Any private insurance	6.2	6.6	5.3	8.1	4.8	4.2	3.3	2.2
Public insurance only	87.2	80.7	84.4	90.4	3.9	8.7	5.8	*4.2
Uninsured all year	28.6	7.5	*21.2	9.5	30.9	51.1	38.4	53.3
65 years and over	60.8	64.8	64.7	70.0	1.5	2.5	2.9	3.1
Sex								
Male	58.8	63.4	66.9	71.7	*1.9	2.3	2.2	2.2
Female	62.3	65.9	63.0	68.5	1.1	2.7	3.5	3.9
Hispanic origin and race ²								
Hispanic or Latino	80.2	77.8	75.6	84.7	*1.6	*2.7	*2.2	2.7
Not Hispanic or Latino:								
White	58.0	62.6	64.1	68.3	1.6	2.5	2.4	3.3
Black or African American	76.3	77.6	68.3	74.8	0.6	2.2	*8.9	2.7
Asian	66.9	0.9
American Indian, Alaska Native, Native Hawaiian, Other Pacific Islander, and Multiple Race	*	*
Insurance status								
Medicare only	68.8	72.4	72.2	77.1	1.4	7.7	5.7	7.7
Medicare and private insurance	56.1	56.3	57.1	56.5	1.6	0.6	*0.6	*0.6
Medicare and other public coverage	92.9	92.7	87.3	93.6	1.0	*2.3	*4.0	1.9

See footnotes at end of table.

Table 98 (page 3 of 3). Sources of payment for health care, by selected population characteristics: United States, selected years 1987–2012

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#098>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population and a sample of medical providers]

... Category not applicable.

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error equal to or greater than 30%. Data not shown if based on fewer than 100 sample cases.

¹Private insurance includes any type of private insurance payments reported for people with private health insurance coverage during the year.

²Persons of Hispanic origin may be of any race. Estimates for Asian persons as well as for American Indian, Alaska Native, Native Hawaiian, Other Pacific Islander, and Multiple Race persons are not available for years prior to 2002 because Asian persons could not be distinguished separately and multiple race information was not collected.

³Any private insurance includes individuals with insurance that provided coverage for hospital and physician care at any time during the year, other than Medicare, Medicaid, or other public coverage for hospital or physician services. Public insurance only includes individuals who were not covered by private insurance at any time during the year but were covered by Medicare, Medicaid, other public coverage for hospital or physician services, and/or CHAMPUS/CHAMPVA (TRICARE) at any point during the year. Uninsured includes persons not covered by either private or public insurance throughout the entire year or period of eligibility for the survey. However, some expenses for the uninsured were paid by sources that were not defined as health insurance coverage, such as the Department of Veterans Affairs, community and neighborhood clinics, the Indian Health Service, state and local health departments, state programs other than Medicaid, workers' compensation, and other unclassified sources (e.g., automobile, home, or liability insurance). Individuals with Indian Health Service coverage only are considered uninsured.

⁴Public sources include payments made by Medicare, Medicaid, the Department of Veterans Affairs, other federal sources (e.g., Indian Health Service, military treatment facilities, and other care provided by the federal government), CHAMPUS/CHAMPVA (TRICARE), and various state and local sources (e.g., community and neighborhood clinics, state and local health departments, and state programs other than Medicaid).

⁵Other sources includes workers' compensation, unclassified sources (automobile, home, or liability insurance, and other miscellaneous or unknown sources), Medicaid payments reported for people who were not enrolled in the program at any time during the year, and any type of private insurance payments reported for people without private health insurance coverage during the year.

NOTES: Estimates for 1987 are based on the National Medical Expenditure Survey (NMES); estimates for other years are based on the Medical Expenditure Panel Survey (MEPS). Because expenditures in NMES were based primarily on charges and those for MEPS were based on payments, NMES data were adjusted to be more comparable with MEPS using estimated charge-to-payment ratios for 1987. Overall, this resulted in an approximate 11% reduction from the unadjusted 1987 NMES expenditure estimates. For a detailed explanation of this adjustment, see Zuvekas S, Cohen J. A guide to comparing health care expenditures in the 1996 MEPS to the 1987 NMES. *Inquiry* 2002;39(1):76–86. Percents sum to 100 across sources within years. See Appendix I, Medical Expenditure Panel Survey (MEPS). Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>. Data have been revised and differ from previous editions of *Health, United States*.

SOURCE: Agency for Healthcare Research and Quality, Center for Financing, Access, and Cost Trends. 1987 National Medical Expenditure Survey and 1996–2012 Medical Expenditure Panel Surveys. See Appendix I, Medical Expenditure Panel Survey (MEPS).

Table 99. Out-of-pocket health care expenses among persons with medical expenses, by age: United States, selected years 1987–2012

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hs/content2015.htm#099>.

[Data are based on household interviews for a sample of the civilian noninstitutionalized population and a sample of medical providers]

Age and year	Percent of persons with expenses	Amount paid out of pocket among persons with expenses ¹						
		Total	\$0	\$1–\$99	\$100–\$499	\$500–\$999	\$1,000–\$1,999	\$2,000 or more
All ages		Percent distribution						
1987	84.5	100.0	10.4	18.7	36.3	15.6	10.6	8.5
1997	84.1	100.0	8.5	25.0	34.6	14.5	10.0	7.3
2000	83.5	100.0	6.9	25.5	33.8	15.1	10.2	8.5
2005	84.7	100.0	8.7	20.5	30.7	16.1	12.3	11.7
2011	84.6	100.0	12.1	21.9	31.7	14.9	11.0	8.4
2012	84.7	100.0	12.2	23.5	31.5	13.7	10.5	8.5
Under 6 years								
1987	88.9	100.0	19.2	26.8	39.4	9.6	3.0	2.1
1997	88.0	100.0	20.0	43.2	29.5	4.2	2.3	0.8
2000	86.7	100.0	16.7	49.3	27.4	4.7	1.3	*0.6
2005	88.9	100.0	27.2	35.6	27.8	6.6	2.0	0.9
2011	89.6	100.0	40.1	31.0	22.8	4.0	1.3	*0.7
2012	88.8	100.0	38.4	35.3	19.8	3.7	2.4	*0.4
6–17 years								
1987	80.2	100.0	15.5	25.9	37.9	9.2	6.0	5.4
1997	81.7	100.0	16.5	34.8	32.5	7.7	4.2	4.2
2000	80.0	100.0	14.7	35.6	33.4	7.4	4.3	4.7
2005	83.0	100.0	18.6	31.3	31.2	9.6	5.0	4.2
2011	84.9	100.0	28.5	29.8	26.3	7.1	3.8	4.4
2012	85.7	100.0	29.0	28.4	26.8	6.4	3.5	6.0
18–44 years								
1987	81.5	100.0	10.1	20.5	39.1	15.4	9.0	6.0
1997	78.3	100.0	7.3	27.2	39.1	14.5	7.6	4.4
2000	77.7	100.0	5.8	28.0	39.1	14.7	7.6	4.8
2005	77.1	100.0	7.0	24.0	36.7	15.7	9.4	7.1
2011	75.6	100.0	8.7	26.3	37.5	13.3	8.3	5.8
2012	75.5	100.0	9.6	28.8	35.2	12.2	8.2	5.9
45–64 years								
1987	87.0	100.0	5.6	11.7	34.8	20.7	15.6	11.6
1997	89.2	100.0	3.4	15.9	34.8	20.0	15.8	10.1
2000	88.5	100.0	2.6	15.0	33.8	21.0	15.7	11.9
2005	89.7	100.0	2.4	12.3	28.6	21.2	19.2	16.2
2011	88.8	100.0	3.7	16.3	32.0	19.6	16.2	12.3
2012	88.9	100.0	3.8	18.0	33.5	17.7	15.2	11.8
65–74 years								
1987	92.8	100.0	5.3	9.0	26.3	21.9	19.7	17.7
1997	94.6	100.0	3.2	10.4	30.3	22.4	17.8	15.9
2000	94.7	100.0	1.5	9.5	26.2	21.8	20.6	20.5
2005	95.9	100.0	1.7	6.2	23.3	20.9	21.8	26.1
2011	96.2	100.0	2.6	9.4	29.8	23.9	20.2	14.1
2012	95.5	100.0	1.9	12.3	32.2	21.5	17.9	14.2
75 years and over								
1987	95.1	100.0	5.6	7.2	23.5	19.5	19.7	24.4
1997	95.8	100.0	2.4	9.1	26.4	19.6	21.6	20.9
2000	96.5	100.0	2.6	9.2	24.0	21.3	19.1	23.7
2005	97.4	100.0	1.6	6.1	20.0	19.5	19.6	33.2
2011	96.7	100.0	2.4	10.8	29.0	22.6	19.4	15.8
2012	97.4	100.0	2.3	11.7	30.4	22.7	18.1	14.9

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error equal to or greater than 30%.

¹Estimates of expenses were converted to 2012 dollars using the Consumer Price Index (all items). See Appendix II, Consumer Price Index (CPI).

NOTES: Includes persons in the civilian noninstitutionalized population for all or part of the year. Expenses for persons in this population for only part of the year are restricted to those incurred during periods of eligibility (e.g., expenses incurred during periods of institutionalization and military service are not included in estimates). Out-of-pocket expenses include expenditures for inpatient hospital and physician services, ambulatory physician and nonphysician services, prescribed medicines, home health services, dental services, and various other medical equipment, supplies, and services that were purchased or rented during the year. Out-of-pocket expenses for over-the-counter medications, phone contacts with health providers, and premiums for health insurance policies are not included in these estimates. Estimates for 1987 are based on the National Medical Expenditure Survey (NMES); estimates for other years are based on the Medical Expenditure Panel Survey (MEPS). Because expenditures in NMES were based primarily on charges and those for MEPS were based on payments, NMES data were adjusted to be more comparable with MEPS using estimated charge to payment ratios for 1987. Overall, this resulted in an approximate 11% reduction from the unadjusted 1987 NMES expenditure estimates. For a detailed explanation of this adjustment, see Zuvekas S, Cohen J. A guide to comparing health care expenditures in the 1996 MEPS to the 1987 NMES. *Inquiry* 2002;39(1):76–86. See Appendix I, Medical Expenditure Panel Survey (MEPS). Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hs.htm>. Data have been revised and differ from previous editions of *Health, United States*.

SOURCE: Agency for Healthcare Research and Quality, Center for Financing, Access, and Cost Trends. 1987 National Medical Expenditure Survey and 1997–2012 Medical Expenditure Panel Surveys. See Appendix I, Medical Expenditure Panel Survey (MEPS).

Table 100 (page 1 of 2). National health expenditures and percent distribution, by sponsor: United States, selected years 1987–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#100>.

[Data are compiled from various sources by the Centers for Medicare & Medicaid Services]

Type of sponsor	1987	1990	1995	2000	2010	2012	2013	2014
Amount, in billions								
National health expenditures	\$516.5	\$721.4	\$1,021.6	\$1,369.7	\$2,595.7	\$2,799.0	\$2,879.9	\$3,031.3
Businesses, households, and other private revenue	353.6	488.2	640.5	883.6	1,444.6	1,581.0	1,618.3	1,672.6
Private business	115.9	170.6	233.7	335.8	518.8	571.9	581.9	606.4
Private business contribution to employer-sponsored private health insurance premiums ¹	78.1	121.8	166.3	244.1	399.5	437.9	442.6	459.0
Employer Medicare hospital insurance trust fund payroll taxes ²	24.4	29.4	43.1	62.3	79.6	88.3	91.1	95.8
Workers' compensation and temporary disability insurance	11.7	17.1	21.4	25.9	34.9	40.5	42.7	45.9
Worksite health care	1.7	2.2	2.9	3.5	4.7	5.2	5.4	5.7
Household	196.6	261.1	328.1	443.6	751.2	811.7	827.4	844.0
Household contribution to employer-sponsored private health insurance premiums ³	29.4	48.9	74.1	102.3	219.4	235.9	237.4	234.7
Household contribution to direct purchase insurance	11.5	15.7	20.8	25.8	42.6	48.6	50.2	51.3
Medical portion of property and casualty insurance ⁴	10.5	12.8	16.1	17.6	28.5	29.3	30.6	32.4
Employee and self-employment payroll taxes and voluntary premiums paid to Medicare hospital insurance trust fund ⁵	29.4	35.7	56.0	82.6	112.0	125.8	124.4	132.9
Premiums paid by individuals to Medicare supplementary medical insurance trust fund and the pre-existing condition insurance plan ⁶	6.2	10.2	16.4	16.3	49.2	53.4	59.3	62.9
Out-of-pocket health spending	109.7	137.9	144.8	199.0	299.5	318.7	325.5	329.8
Other private revenues ⁷	41.1	56.5	78.6	104.2	174.6	197.4	209.1	222.2
Governments	162.9	233.2	381.2	486.1	1,151.1	1,218.0	1,261.6	1,358.7
Federal government	85.2	124.1	215.4	260.5	731.1	730.0	755.5	843.7
Federal government contribution to employer-sponsored private health insurance premiums	4.9	9.9	11.4	14.3	28.5	31.0	32.4	33.2
Employer Medicare hospital insurance trust fund payroll taxes	1.7	2.0	2.3	2.7	4.1	4.1	4.0	4.0
Marketplace tax credits and subsidies ⁸	18.5
Federal general revenue and Medicare net trust fund expenditures ⁹	17.7	27.7	57.6	49.2	247.1	264.8	273.5	288.5
Federal portion of Medicaid payments	27.9	42.6	85.9	116.8	266.3	242.8	257.7	305.1
Medicare buy-in premiums ¹⁰	0.3	0.7	1.9	2.5	9.2	7.8	8.3	8.4
Retiree drug subsidy payments to employer-sponsored health insurance plans	3.9	3.0	1.8	1.6
Other federal health insurance and programs ¹¹	32.7	41.3	56.2	75.1	172.0	176.2	177.8	184.3
State and local government	77.7	109.1	165.7	225.5	420.0	488.0	506.0	515.0
State and local government contribution to employer-sponsored private health insurance premiums ¹²	14.8	24.8	36.7	54.4	138.0	148.1	154.1	159.8
Employer Medicare hospital insurance trust fund payroll taxes	3.1	4.1	5.6	7.5	11.2	11.3	11.5	11.8
State portion of Medicaid payments	22.5	31.1	58.9	83.5	130.8	179.2	189.0	190.6
Medicare buy-in premiums ¹⁰	0.2	0.5	1.4	1.8	4.0	5.4	5.7	5.8
Other programs ¹³	37.0	48.6	63.1	78.4	135.8	144.0	145.7	147.0
Percent distribution								
National health expenditures	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Businesses, households, and other private revenue	68.5	67.7	62.7	64.5	55.7	56.5	56.2	55.2
Private business	22.4	23.6	22.9	24.5	20.0	20.4	20.2	20.0
Private business contribution to employer-sponsored private health insurance premiums ¹	15.1	16.9	16.3	17.8	15.4	15.6	15.4	15.1
Employer Medicare hospital insurance trust fund payroll taxes ²	4.7	4.1	4.2	4.5	3.1	3.2	3.2	3.2
Workers' compensation and temporary disability insurance	2.3	2.4	2.1	1.9	1.3	1.4	1.5	1.5
Worksite health care	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2
Household	38.1	36.2	32.1	32.4	28.9	29.0	28.7	27.8
Household contribution to employer-sponsored private health insurance premiums ³	5.7	6.8	7.3	7.5	8.5	8.4	8.2	7.7
Household contribution to direct purchase insurance	2.2	2.2	2.0	1.9	1.6	1.7	1.7	1.7
Medical portion of property and casualty insurance ⁴	2.0	1.8	1.6	1.3	1.1	1.0	1.1	1.1
Employee and self-employment payroll taxes and voluntary premiums paid to Medicare hospital insurance trust fund ⁵	5.7	4.9	5.5	6.0	4.3	4.5	4.3	4.4
Premiums paid by individuals to Medicare supplementary medical insurance trust fund and the pre-existing condition insurance plan ⁶	1.2	1.4	1.6	1.2	1.9	1.9	2.1	2.1
Out-of-pocket health spending	21.2	19.1	14.2	14.5	11.5	11.4	11.3	10.9
Other private revenues ⁷	8.0	7.8	7.7	7.6	6.7	7.1	7.3	7.3

See footnotes at end of table.

Table 100 (page 2 of 2). National health expenditures and percent distribution, by sponsor: United States, selected years 1987–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#100>.

[Data are compiled from various sources by the Centers for Medicare & Medicaid Services]

Type of sponsor	1987	1990	1995	2000	2010	2012	2013	2014
	Percent distribution							
Governments	31.5	32.3	37.3	35.5	44.3	43.5	43.8	44.8
Federal government	16.5	17.2	21.1	19.0	28.2	26.1	26.2	27.8
Federal government contribution to employer-sponsored private health insurance premiums	0.9	1.4	1.1	1.0	1.1	1.1	1.1	1.1
Employer Medicare hospital insurance trust fund payroll taxes	0.3	0.3	0.2	0.2	0.2	0.1	0.1	0.1
Marketplace tax credits and subsidies ⁸	0.6
Federal general revenue and Medicare net trust fund expenditures ⁹	3.4	3.8	5.6	3.6	9.5	9.5	9.5	9.5
Federal portion of Medicaid payments	5.4	5.9	8.4	8.5	10.3	8.7	8.9	10.1
Medicare buy-in premiums ¹⁰	0.1	0.1	0.2	0.2	0.4	0.3	0.3	0.3
Retiree drug subsidy payments to employer-sponsored health insurance plans	0.2	0.1	0.1	0.1
Other federal health insurance and programs ¹¹	6.3	5.7	5.5	5.5	6.6	6.3	6.2	6.1
State and local government	15.0	15.1	16.2	16.5	16.2	17.4	17.6	17.0
State and local government contribution to employer-sponsored private health insurance premiums ¹²	2.9	3.4	3.6	4.0	5.3	5.3	5.4	5.3
Employer Medicare hospital insurance trust fund payroll taxes	0.6	0.6	0.5	0.5	0.4	0.4	0.4	0.4
State portion of Medicaid payments	4.4	4.3	5.8	6.1	5.0	6.4	6.6	6.3
Medicare buy-in premiums ¹⁰	0.0	0.1	0.1	0.1	0.2	0.2	0.2	0.2
Other programs ¹³	7.2	6.7	6.2	5.7	5.2	5.1	5.1	4.8

... Category not applicable.

¹Excludes Medicare Retiree Drug Subsidy (RDS) payments to private plans beginning in 2006, small-business tax credits beginning in 2010, and Early Retirement Reinsurance Program (ERRP) payments for 2010–2011.

²Includes one-half of self-employment contribution to the Medicare hospital insurance (HI) trust fund.

³Excludes government-subsidized Consolidated Omnibus Budget Reconciliation Act (COBRA) payments in 2009–2011.

⁴Includes property and casualty insurance premium portions that are used to pay medical claims for automobile, homeowners, or other liability insurance.

⁵Includes one-half of self-employment contributions to the Medicare hospital insurance trust fund and income taxation of Social Security benefits.

⁶Includes premiums paid for the Pre-Existing Condition Insurance Plan (PCIP) in 2010–2014.

⁷Includes health-related philanthropic support, nonoperating revenue, investment income, and privately funded structures and equipment.

⁸Includes Affordable Care Act (ACA) health insurance premium tax credits and cost-sharing subsidies beginning in 2014.

⁹Excludes Medicare hospital trust fund payroll taxes and premiums, Medicare supplementary medical insurance premiums, Part D state phase-down payments to Medicare beginning in 2006, Medicare premium buy-in programs by Medicaid for people eligible for both Medicaid and Medicare (dual eligibles), and trust fund revenues from the income taxation of Social Security benefits.

¹⁰Medicare premium buy-in programs are for people eligible for both Medicaid and Medicare (dual eligibles).

¹¹Includes maternal and child health, vocational rehabilitation, Substance Abuse and Mental Health Services Administration, Indian Health Service, federal workers' compensation, and other federal programs, public health activities, Department of Defense, Department of Veterans Affairs, Children's Health Insurance Program (CHIP), and investment (research, structures and equipment). Also includes government-subsidized COBRA payments in 2009–2011, small business tax credits beginning in 2010, and ERRP payments in 2010–2011. Excludes premiums paid for the Pre-Existing Condition Insurance Plan (PCIP) premiums in 2010–2014.

¹²Excludes Medicare RDS payments to state and local government employer plans beginning in 2006 and ERRP payments in 2010–2011.

¹³Includes maternal and child health, vocational rehabilitation, general assistance, school health, CHIP, public health activities, other state and local programs, investment (research, structures and equipment). Also includes Part D state phase-down payments to Medicare beginning in 2006. See Appendix II, Health expenditures, national.

NOTES: This table disaggregates health expenditures according to five classes of sponsors: businesses, households (individuals), federal government, state and local governments, and nonpatient revenue sources such as philanthropy. Where businesses or households pay dedicated funds into government health programs (for example, Medicare) or employers and employees share in the cost of health premiums, these costs are assigned to businesses or households accordingly. This results in a lower share of expenditures being assigned to the federal government than for tabulations of expenditures by source of funds. Estimates of national health expenditure by source of funds aim to track government-sponsored health programs over time and do not delineate the role of business employers in paying for health care. Some of the sponsor categories were revised or added in 2014 to account for changes in the health care system. See Appendix I, National Health Expenditure Accounts (NHEA). Estimates may not sum to totals because of rounding. For more information on NHE categories, sources, and methods, see the National Health Expenditure Accounts Methodology Paper, 2014. Available from: <https://www.cms.gov/Research-Statistics-Data-and-Systems/Statistics-Trends-and-Reports/NationalHealthExpendData/NationalHealthAccountsHistorical.html>. Data have been revised and differ from previous editions of *Health, United States*.

SOURCE: Centers for Medicare & Medicaid Services, Office of the Actuary, National Health Statistics Group. Businesses, Households, and Governments. National Health Expenditure Accounts, National health expenditures. Available from: <https://www.cms.gov/Research-Statistics-Data-and-Systems/Statistics-Trends-and-Reports/NationalHealthExpendData/NationalHealthAccountsHistorical.html>, accessed on December 12, 2015. See Appendix I, National Health Expenditure Accounts (NHEA).

Table 101. Employers' costs per employee-hour worked for total compensation, wages and salaries, and health insurance, by selected characteristics: United States, selected years 1999–2015

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#101>.

[Data are based on surveys of a sample of employers]

Characteristic	1999	2000	2005	2008	2010	2011	2013	2014	2015
Total compensation per employee-hour worked									
State and local government	\$28.00	\$29.05	\$35.50	\$37.84	\$39.81	\$40.54	\$42.12	\$43.10	\$44.25
Total private industry	19.00	19.85	24.17	26.76	27.73	28.10	29.13	29.99	31.65
Census region:									
Northeast	20.94	22.67	27.09	30.56	32.13	32.16	33.43	34.79	38.93
Midwest	18.36	19.22	24.23	25.98	26.75	27.47	27.93	28.71	29.08
South	16.97	17.81	21.36	23.90	24.72	24.93	26.60	27.14	29.04
West	20.74	20.88	25.98	28.70	29.52	29.95	30.54	31.59	32.23
Union status:									
Union	24.75	25.88	33.17	36.28	37.16	37.68	40.43	43.84	46.62
Nonunion	18.20	19.07	23.09	25.64	26.67	27.08	28.02	28.63	30.18
Establishment employment size:									
1–99 employees	16.27	17.16	20.22	22.23	22.84	23.21	23.92	25.03	26.45
100 or more	21.88	22.81	28.94	31.68	33.33	33.69	35.25	35.76	37.78
100–499	18.14	19.30	24.44	26.80	28.55	28.69	29.71	29.92	32.07
500 or more	26.37	26.93	34.59	37.60	39.76	40.53	43.05	44.04	46.19
Wages and salaries as a percent of total compensation									
State and local government	70.6	70.8	68.3	65.9	65.9	65.5	64.8	64.4	64.0
Total private industry	73.0	73.0	71.0	70.6	70.6	70.7	70.3	69.9	69.3
Census region:									
Northeast	72.0	72.2	70.4	69.8	69.0	69.5	68.8	68.4	66.9
Midwest	71.9	72.4	70.1	69.8	70.0	69.8	69.5	69.5	69.2
South	74.0	73.5	72.1	71.8	71.8	71.9	71.6	71.2	70.8
West	74.1	74.0	70.9	70.8	71.1	71.0	70.6	70.0	69.9
Union status:									
Union	65.5	65.2	62.6	61.9	61.6	61.1	59.8	60.0	59.8
Nonunion	74.4	74.4	72.4	72.1	72.0	72.1	71.8	71.4	70.8
Establishment employment size:									
1–99 employees	75.5	75.5	73.9	73.8	73.6	74.0	74.0	73.5	72.5
100 or more	71.0	71.0	68.5	68.2	68.2	68.0	67.3	66.9	66.7
100–499	72.6	72.8	70.2	69.8	70.0	69.9	69.1	68.8	68.5
500 or more	69.7	69.4	67.0	66.9	66.5	66.2	65.6	65.1	64.9
Health insurance as a percent of total compensation									
State and local government	7.6	7.8	10.2	11.0	11.4	11.7	11.7	11.7	11.6
Total private industry	5.4	5.5	6.8	7.2	7.5	7.5	7.8	7.9	7.7
Census region:									
Northeast	5.7	5.6	6.8	6.9	7.5	7.8	8.1	8.2	7.7
Midwest	5.8	5.8	7.3	7.9	8.3	8.3	8.6	8.6	8.5
South	5.2	5.4	6.6	6.9	7.2	7.2	7.2	7.3	7.1
West	4.8	5.0	6.3	6.9	7.1	7.1	7.4	7.5	7.7
Union status:									
Union	8.2	8.4	10.3	10.9	11.8	12.3	12.9	12.6	12.1
Nonunion	4.9	5.0	6.2	6.5	6.8	6.8	7.0	7.1	7.0
Establishment employment size:									
1–99 employees	4.7	4.8	5.9	6.1	6.4	6.3	6.5	6.6	6.4
100 or more	5.9	6.0	7.5	8.0	8.4	8.6	8.8	8.9	8.7
100–499	5.6	5.6	7.5	7.9	8.3	8.4	8.7	8.7	8.5
500 or more	6.2	6.4	7.6	8.0	8.5	8.7	8.9	9.1	8.9

NOTES: Costs are calculated annually from March survey data. Total compensation includes wages, salaries and benefits. See Appendix II, Employer costs for employee compensation. See *Health, United States, 2013*, Table 121 for prior years of data. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: U.S. Department of Labor, Bureau of Labor Statistics, National Compensation Survey: Employer Costs for Employee Compensation Annual, 1999–2001; Quarterly, 2002–2003; March release, 2004–2015. Available from: <http://www.bls.gov/nchs/ect/>. See Appendix I, National Compensation Survey (NCS).

Table 102 (page 1 of 3). Private health insurance coverage among persons under age 65, by selected characteristics: United States, selected years 1984–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#102>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Private health insurance ¹								
	1984 ²	1997	2000 ³	2004	2005	2010	2012	2013	2014
	Number, in millions								
Total ⁴	157.5	165.8	174.0	174.5	174.7	163.9	164.9	165.3	170.7
	Percent of population								
Total ⁴	76.8	70.7	71.5	68.8	68.2	61.7	61.8	61.8	63.7
Age									
Under 19 years	72.6	66.1	66.7	63.3	62.3	54.3	53.6	53.5	54.1
Under 6 years	68.1	61.3	62.7	58.1	56.6	48.3	48.4	47.3	50.2
6–18 years	74.8	68.4	68.5	65.5	64.9	57.2	56.0	56.3	55.9
Under 18 years	72.6	66.1	66.6	63.2	62.1	54.1	53.4	53.2	53.7
6–17 years	74.9	68.5	68.5	65.6	64.7	57.2	55.8	56.0	55.4
18–64 years	78.6	72.7	73.5	71.1	70.7	64.7	65.1	65.1	67.4
18–44 years	76.5	69.4	70.5	67.3	66.6	60.0	61.4	61.8	64.3
18–24 years	67.4	59.3	60.3	58.2	58.0	52.3	58.1	59.0	62.0
19–25 years	67.4	58.3	59.1	57.3	56.3	51.8	58.1	58.9	62.2
25–34 years	77.4	68.1	70.1	65.5	65.1	58.7	58.7	59.0	62.0
35–44 years	83.9	76.4	77.0	74.8	73.7	66.9	66.7	67.0	68.6
45–64 years	83.3	79.0	78.7	77.1	76.9	71.3	70.0	69.5	71.7
45–54 years	83.3	80.4	80.0	77.8	77.4	70.9	69.6	69.8	71.6
55–64 years	83.3	76.9	76.7	76.1	76.2	71.8	70.4	69.1	71.7
Sex									
Male	77.3	70.9	71.6	68.7	68.0	61.1	61.8	61.9	63.8
Female	76.2	70.5	71.3	68.9	68.4	62.4	61.9	61.7	63.5
Sex and marital status ⁵									
Male:									
Married	85.0	81.6	81.5	80.0	79.6	75.1	74.9	74.8	77.1
Divorced, separated, widowed	65.5	59.9	62.2	59.0	56.7	50.6	51.0	50.9	54.0
Never married	71.3	63.3	63.8	60.4	60.2	52.5	54.7	55.8	58.2
Female:									
Married	83.8	81.0	81.0	79.7	79.3	75.6	75.0	74.3	75.9
Divorced, separated, widowed	63.1	59.1	63.2	58.6	59.9	53.9	51.8	52.1	55.1
Never married	72.2	63.8	64.2	62.2	61.5	54.1	56.2	56.2	58.4
Race ⁶									
White only	79.9	74.2	75.7	71.4	70.9	64.9	64.8	64.7	66.6
Black or African American only	58.1	54.7	55.9	53.9	52.9	44.8	45.8	45.4	47.1
American Indian or Alaska Native only	49.1	39.4	43.7	44.7	43.0	31.7	34.9	36.0	34.7
Asian only	69.9	68.0	72.1	71.6	72.2	68.1	67.6	69.4	72.5
Native Hawaiian or Other Pacific Islander only	---	---	*	*	*	*	*	*	*
2 or more races	---	---	61.4	62.0	57.6	52.4	52.9	50.0	55.4
Hispanic origin and race ⁶									
Hispanic or Latino	55.7	46.4	47.8	41.7	42.4	36.8	36.7	37.3	41.2
Mexican	53.3	42.3	45.4	39.1	39.7	33.4	34.1	34.9	39.0
Puerto Rican	48.4	47.0	51.1	47.3	48.5	46.0	43.7	42.1	46.8
Cuban	72.5	71.0	63.9	57.9	58.1	53.8	49.1	45.3	56.6
Other Hispanic or Latino	61.6	49.9	50.7	45.1	45.6	40.9	39.5	41.2	43.1
Not Hispanic or Latino	78.7	74.0	75.2	73.7	73.0	67.0	67.5	67.4	68.9
White only	82.4	78.1	79.5	77.9	77.3	72.0	72.6	72.4	73.7
Black or African American only	58.2	54.9	56.0	54.6	53.1	45.1	46.4	45.7	48.0
Age and percent of poverty level ⁷									
Under 65 years:									
Below 100%	32.2	23.3	25.2	21.8	21.4	16.0	16.5	15.5	17.4
100%–199%	70.3	53.5	50.1	45.7	44.7	34.8	36.7	35.1	38.2
100%–133%	59.4	39.7	39.3	36.2	36.0	24.4	26.9	25.3	26.5
134%–199%	75.2	60.1	55.3	50.3	49.4	40.3	42.4	40.8	45.1
200%–399%	89.3	80.8	78.1	75.8	74.8	70.7	71.3	71.3	73.6
400% or more	95.4	91.8	91.9	90.8	90.6	89.9	90.6	90.4	91.5

See footnotes at end of table.

Table 102 (page 2 of 3). Private health insurance coverage among persons under age 65, by selected characteristics: United States, selected years 1984–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#102>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Private health insurance ¹								
	1984 ²	1997	2000 ³	2004	2005	2010	2012	2013	2014
	Percent of population								
Under 19 years:									
Below 100%	29.6	19.3	20.3	15.1	15.0	9.8	10.0	9.3	9.5
100%–199%	73.6	54.7	49.5	43.6	41.6	31.5	32.4	29.2	30.4
100%–133%	63.8	39.3	37.1	33.3	32.6	20.1	22.3	18.3	18.2
134%–199%	78.4	62.4	56.1	48.7	47.0	38.1	38.4	36.2	38.6
200%–399%	91.1	83.5	80.8	78.2	76.6	72.6	72.5	71.8	73.5
400% or more	96.2	93.3	93.0	92.4	92.5	91.2	91.2	92.3	92.5
Under 18 years:									
Below 100%	28.5	18.3	19.5	14.2	14.2	9.2	9.1	8.4	8.6
100%–199%	73.9	54.7	49.4	43.6	41.4	31.5	32.1	28.5	30.2
100%–133%	63.9	38.7	36.8	33.2	32.0	19.9	21.6	17.8	18.2
134%–199%	78.6	62.8	56.2	48.8	47.0	38.3	38.4	35.3	38.3
200%–399%	91.3	83.7	81.1	78.4	76.6	72.6	72.5	71.9	73.2
400% or more	96.1	93.5	93.1	92.6	92.5	91.4	91.4	92.2	92.6
18–64 years:									
Below 100%	35.0	26.8	29.1	26.5	25.9	20.4	20.9	19.9	22.7
100%–199%	68.3	52.8	50.5	46.9	46.5	36.4	38.9	38.2	42.1
100%–133%	56.6	40.3	40.9	37.9	38.3	26.9	29.6	29.2	31.1
134%–199%	73.3	58.6	54.9	51.2	50.7	41.3	44.2	43.2	48.2
200%–399%	88.3	79.4	76.7	74.7	74.0	70.0	70.8	71.1	73.7
400% or more	95.2	91.3	91.6	90.3	90.1	89.5	90.4	89.9	91.2
Disability measure among adults 18–64 years ⁸									
Any basic actions difficulty or complex activity limitation	---	61.6	63.1	58.9	58.1	53.0	50.8	48.6	51.1
Any basic actions difficulty	---	62.3	63.9	59.4	58.8	53.8	51.7	49.2	51.8
Any complex activity limitation	---	47.9	48.4	45.9	44.0	38.6	36.0	34.8	34.7
No disability	---	77.4	77.2	74.5	73.7	69.3	70.2	70.7	72.5
Geographic region									
Northeast	80.5	74.2	76.3	74.0	74.0	68.2	67.2	66.1	67.7
Midwest	80.6	77.1	78.8	76.3	74.6	66.7	68.4	68.0	68.7
South	74.3	67.3	66.8	64.1	62.5	57.5	57.3	57.4	59.4
West	71.9	65.4	66.5	64.1	65.6	58.9	58.5	59.6	62.9
Location of residence ⁹									
Within MSA	77.5	71.2	72.3	69.6	69.0	62.9	63.0	63.0	64.8
Outside MSA	75.2	68.4	67.8	65.5	64.6	55.1	55.3	54.7	56.2

See footnotes at end of table.

Table 102 (page 3 of 3). Private health insurance coverage among persons under age 65, by selected characteristics: United States, selected years 1984–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#102>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

-- Data not available.

* Estimates are considered unreliable. Data not shown have a relative standard error greater than 30%.

¹Any private health insurance coverage (both individual and insurance obtained through the workplace) at the time of interview; includes those who also had another type of coverage.

²Data prior to 1997 are not strictly comparable with data for later years due to the 1997 questionnaire redesign. See Appendix I, National Health Interview Survey (NHIS) and Appendix II, Health insurance coverage.

³Estimates for 2000–2002 were calculated using 2000-based sample weights and may differ from estimates in other reports that used 1990-based sample weights for 2000–2002 estimates.

⁴Includes all other races not shown separately, those with unknown marital status, unknown disability status, and, in 1984 and 1989, persons with unknown poverty level.

⁵Includes persons aged 14–64.

⁶The race groups, white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The five single-race categories plus multiple-race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group; the category 2 or more races includes persons who reported more than one racial group. Prior to 1999, data were tabulated according to the 1977 Standards with four racial groups, and the Asian only category including Native Hawaiian or Other Pacific Islander. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

⁷Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Poverty level was unknown for 10%–11% of persons under age 65 in 1984 and 1989. Missing family income data were imputed for 1995 and beyond. See Appendix II, Family income; Poverty; Table VI.

⁸Any basic actions difficulty or complex activity limitation is defined as having one or more of the following limitations or difficulties: movement difficulty, emotional difficulty, sensory (seeing or hearing) difficulty, cognitive difficulty, self-care (activities of daily living or instrumental activities of daily living) limitation, social limitation, or work limitation. For more information, see Appendix II, Basic actions difficulty; Complex activity limitation. Starting with 2007 data, the hearing question, a component of the basic actions difficulty measure, was revised. Consequently, data prior to 2007 are not comparable with data for 2007 and beyond. For more information on the impact of the revised hearing question, see Appendix II, Hearing trouble.

⁹MSA is metropolitan statistical area. Starting with 2006 data, MSA status is determined using 2000 census data and the 2000 standards for defining MSAs. For data prior to 2006, see Appendix II, Metropolitan statistical area (MSA) for the applicable standards.

NOTES: This table includes persons who had private coverage through the workplace in addition to other types of health insurance coverage. Private health insurance coverage is at the time of interview. The number of persons with private coverage was calculated by multiplying the percentage with private coverage by the number of persons under age 65 in the civilian noninstitutionalized U.S. population, which was determined from the post-stratification Census control total for each survey year. Percentages of persons with private coverage were calculated with unknown values excluded from denominators. See Appendix II, Health insurance coverage. Standard errors are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey, health insurance supplements (1984, 1989, 1994–1996). Starting with 1997, data are from the family core and the sample adult questionnaires. See Appendix I, National Health Interview Survey (NHIS).

Table 103 (page 1 of 3). Private health insurance coverage obtained through the workplace among persons under age 65, by selected characteristics: United States, selected years 1984–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#103>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Private insurance obtained through workplace ¹								
	1984 ²	1997	2000 ³	2004	2005	2010	2012	2013	2014
	Number, in millions								
Total ⁴	141.8	153.6	160.8	159.5	160.1	147.6	148.6	148.3	146.4
	Percent of population								
Total ⁴	69.1	66.4	67.1	64.0	63.6	56.6	56.9	56.6	56.8
Age									
Under 19 years	66.4	62.8	63.1	59.6	58.7	50.9	50.1	49.6	49.8
Under 6 years	62.1	58.3	58.9	54.8	53.4	44.9	45.0	44.2	46.0
6–18 years	68.4	64.9	64.9	61.7	61.1	53.8	52.4	52.0	51.5
Under 18 years	66.5	62.8	63.0	59.6	58.6	50.7	49.9	49.3	49.5
6–17 years	68.7	65.1	65.0	61.9	61.1	53.8	52.3	51.8	51.2
18–64 years	70.3	68.0	68.8	65.8	65.7	58.9	59.6	59.5	59.6
18–44 years	69.6	65.7	66.5	62.6	62.2	54.6	56.7	56.9	57.4
18–24 years	58.7	54.9	55.5	52.2	52.1	45.3	52.7	53.1	54.3
19–25 years	59.0	53.7	54.2	51.2	50.6	44.1	52.7	53.0	54.4
25–34 years	71.2	64.6	66.4	61.0	61.1	53.3	53.8	53.8	54.6
35–44 years	77.4	72.7	73.2	70.7	69.9	62.8	62.7	63.1	62.6
45–64 years	71.8	72.8	72.9	70.8	70.9	64.8	63.6	62.9	62.6
45–54 years	74.6	75.6	75.6	72.4	72.6	65.9	64.4	64.3	64.3
55–64 years	69.0	68.4	68.6	68.5	68.6	63.4	62.6	61.3	60.9
Sex									
Male	69.8	66.7	67.3	64.1	63.6	56.1	57.1	56.9	57.2
Female	68.4	66.2	66.9	63.9	63.6	57.1	56.8	56.4	56.5
Sex and marital status ⁵									
Male:									
Married	77.9	77.4	77.5	75.3	75.3	70.1	69.9	69.8	70.0
Divorced, separated, widowed	58.0	55.2	57.4	53.8	51.9	45.3	46.2	45.3	46.1
Never married	61.5	58.4	58.8	54.9	54.9	46.2	49.6	50.0	50.4
Female:									
Married	76.1	76.4	76.3	74.5	74.2	69.8	69.3	68.5	68.1
Divorced, separated, widowed	51.9	53.8	57.8	53.2	54.3	48.1	46.3	46.3	46.3
Never married	63.5	59.6	60.1	56.7	56.3	48.2	50.9	50.6	51.1
Race ⁶									
White only	72.0	69.7	71.0	66.2	66.1	59.3	59.6	59.2	59.5
Black or African American only	52.4	52.6	53.4	51.4	50.6	42.3	43.2	42.9	42.4
American Indian or Alaska Native only	45.8	37.2	41.7	42.2	39.9	*29.4	34.0	34.2	31.0
Asian only	59.0	61.7	65.8	65.6	64.4	60.6	60.1	61.4	62.3
Native Hawaiian or Other Pacific Islander only	---	---	*	*	*	*	*	*	*
2 or more races	---	---	59.8	58.2	54.8	49.5	48.8	46.9	51.2
Hispanic origin and race ⁶									
Hispanic or Latino	52.0	43.9	45.3	39.2	40.0	34.6	34.6	34.9	36.8
Mexican	50.5	40.8	43.6	37.2	37.6	31.6	32.5	32.5	35.3
Puerto Rican	45.9	45.1	49.4	44.3	46.2	43.6	41.6	40.8	43.2
Cuban	57.4	58.4	53.6	51.2	53.5	47.4	42.8	41.2	44.7
Other Hispanic or Latino	57.4	47.0	47.3	41.7	42.6	37.8	36.7	38.6	37.7
Not Hispanic or Latino	70.7	69.5	70.6	68.5	68.0	61.3	62.0	61.7	61.6
White only	74.0	73.3	74.5	72.1	71.9	65.7	66.6	66.1	65.9
Black or African American only	52.5	52.9	53.6	52.2	50.9	42.6	43.6	43.2	43.3
Age and percent of poverty level ⁷									
Under 65 years:									
Below 100%	24.1	20.0	21.0	18.2	17.8	12.4	13.6	12.2	12.3
100%–199%	61.7	48.9	45.4	41.2	40.1	30.2	32.2	31.0	31.4
100%–133%	50.0	35.4	35.0	32.0	31.3	20.6	23.0	21.7	20.7
134%–199%	66.9	55.4	50.5	45.7	44.8	35.3	37.5	36.3	37.8
200%–399%	82.8	76.5	73.4	70.7	69.8	65.3	65.9	65.6	65.9
400% or more	88.8	87.4	87.9	85.7	86.1	84.2	85.1	84.6	84.8

See footnotes at end of table.

Table 103 (page 2 of 3). Private health insurance coverage obtained through the workplace among persons under age 65, by selected characteristics: United States, selected years 1984–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#103>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Private insurance obtained through workplace ¹								
	1984 ²	1997	2000 ³	2004	2005	2010	2012	2013	2014
Percent of population									
Under 19 years:									
Below 100%	23.6	17.0	17.1	13.6	13.3	8.2	8.7	7.8	7.2
100%–199%	67.0	51.2	45.8	40.5	38.3	28.8	29.7	26.6	27.1
100%–133%	56.1	35.8	33.6	30.8	29.1	17.9	20.5	16.4	15.7
134%–199%	72.3	59.0	52.2	45.3	43.7	35.1	35.2	33.1	34.7
200%–399%	85.7	80.0	76.9	73.9	72.4	68.7	68.0	66.7	67.9
400% or more	90.8	89.7	89.5	87.6	88.3	86.5	86.3	86.9	87.2
Under 18 years:									
Below 100%	23.0	16.2	16.6	12.8	12.5	7.8	8.1	7.2	6.7
100%–199%	67.5	51.2	45.8	40.7	38.2	28.8	29.4	26.0	26.9
100%–133%	56.3	35.2	33.5	30.8	28.6	17.8	19.8	16.0	15.7
134%–199%	72.8	59.4	52.4	45.6	43.9	35.2	35.2	32.4	34.6
200%–399%	85.9	80.2	77.1	74.2	72.4	68.7	68.1	66.8	67.8
400% or more	90.7	89.8	89.7	87.8	88.5	86.6	86.4	86.8	87.3
18–64 years:									
Below 100%	24.8	22.7	24.0	21.6	21.2	15.4	16.9	15.2	15.7
100%–199%	58.3	47.6	45.2	41.5	41.1	30.9	33.6	33.3	33.5
100%–133%	46.0	35.5	35.9	32.7	32.9	22.1	24.6	24.7	23.4
134%–199%	63.6	53.2	49.5	45.7	45.3	35.3	38.7	38.0	39.2
200%–399%	81.4	74.7	71.7	69.2	68.7	63.9	65.0	65.2	65.2
400% or more	88.5	86.8	87.5	85.1	85.4	83.6	84.7	84.0	84.1
Disability measure among adults 18–64 years ⁸									
Any basic actions difficulty or complex activity limitation	---	57.3	58.5	54.1	53.3	48.0	45.8	44.0	43.3
Any basic actions difficulty	---	58.0	59.1	54.7	54.0	48.9	46.7	44.6	44.2
Any complex activity limitation	---	43.3	43.5	41.1	38.9	32.8	30.5	29.6	26.3
No disability	---	72.5	72.5	69.0	68.5	63.5	64.7	64.7	64.8
Geographic region									
Northeast	74.0	71.0	72.5	70.1	70.6	64.4	63.4	62.3	62.0
Midwest	72.0	72.6	74.9	71.9	70.1	61.8	63.8	62.6	61.8
South	66.2	62.9	62.5	59.6	58.0	52.2	52.2	52.4	52.5
West	64.7	60.7	61.1	57.5	59.7	52.7	52.8	53.6	55.2
Location of residence ⁹									
Within MSA	70.9	67.3	68.2	64.9	64.5	57.9	58.1	57.8	58.2
Outside MSA	65.3	62.8	62.6	60.1	59.6	49.4	50.3	49.4	48.3

See footnotes at end of table.

Table 103 (page 3 of 3). Private health insurance coverage obtained through the workplace among persons under age 65, by selected characteristics: United States, selected years 1984–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#103>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

-- Data not available.

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.

¹Any private insurance at the time of interview that was originally obtained through a present or former employer or union, or, starting with 1997 data, through the workplace, self-employment, or a professional association; includes those who also had another type of coverage.

²Data prior to 1997 are not strictly comparable with data for later years due to the 1997 questionnaire redesign. See Appendix I, National Health Interview Survey (NHIS) and Appendix II, Health insurance coverage.

³Estimates for 2000–2002 were calculated using 2000-based sample weights and may differ from estimates in other reports that used 1990-based sample weights for 2000–2002 estimates.

⁴Includes all other races not shown separately, those with unknown marital status, unknown disability status, and, in 1984 and 1989, persons with unknown poverty level.

⁵Includes persons aged 14–64.

⁶The race groups, white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The five single-race categories plus multiple-race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group; the category 2 or more races includes persons who reported more than one racial group. Prior to 1999, data were tabulated according to the 1977 Standards with four racial groups, and the Asian only category included Native Hawaiian or Other Pacific Islander. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

⁷Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Poverty level was unknown for 10%–11% of persons under age 65 in 1984 and 1989. Missing family income data were imputed for 1995 and beyond. See Appendix II, Family income; Poverty; Table VI.

⁸Any basic actions difficulty or complex activity limitation is defined as having one or more of the following limitations or difficulties: movement difficulty, emotional difficulty, sensory (seeing or hearing) difficulty, cognitive difficulty, self-care (activities of daily living or instrumental activities of daily living) limitation, social limitation, or work limitation. For more information, see Appendix II, Basic actions difficulty; Complex activity limitation. Starting with 2007 data, the hearing question, a component of the basic actions difficulty measure, was revised. Consequently, data prior to 2007 are not comparable with data for 2007 and beyond. For more information on the impact of the revised hearing question, see Appendix II, Hearing trouble.

⁹MSA is metropolitan statistical area. Starting with 2006 data, MSA status is determined using 2000 census data and the 2000 standards for defining MSAs. For data prior to 2006, see Appendix II, Metropolitan statistical area (MSA) for the applicable standards.

NOTES: This table includes persons who had private coverage through the workplace in addition to other types of health insurance coverage. Private coverage through the workplace is at the time of interview. The number of persons with private coverage through the workplace was calculated by multiplying the percentage with private coverage through the workplace by the number of persons under age 65 in the civilian noninstitutionalized U.S. population, which was determined from the post-stratification Census control total for each survey year. Percentages of persons with private coverage obtained through the workplace were calculated with unknown values excluded from denominators. See Appendix II, Health insurance coverage. Standard errors are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey, health insurance supplements (1984, 1989, 1994–1996). Starting with 1997, data are from the family core and the sample adult questionnaires. See Appendix I, National Health Interview Survey (NHIS).

Table 104 (page 1 of 3). Medicaid coverage among persons under age 65, by selected characteristics: United States, selected years 1984–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#104>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	1984 ¹	1997	2000 ²	2004(1) ³	2004(2) ³	2010 ³	2012 ³	2013 ³	2014 ³
Number, in millions									
Total ⁴	14.0	22.9	23.2	31.1	31.6	44.8	48.1	48.5	52.6
Percent of population									
Total ⁴	6.8	9.7	9.5	12.3	12.5	16.9	18.0	18.1	19.6
Age									
Under 19 years	11.7	18.0	19.2	25.4	25.8	35.7	38.1	38.1	38.6
Under 6 years	15.5	24.7	24.7	31.8	32.4	43.7	45.7	45.9	43.8
6–18 years	9.8	14.9	16.8	22.5	22.9	31.8	34.7	34.6	36.3
Under 18 years	11.9	18.4	19.6	25.9	26.4	36.4	38.9	38.9	39.4
6–17 years	10.1	15.2	17.2	23.1	23.4	32.5	35.5	35.5	37.3
18–64 years	4.5	5.9	5.2	6.7	6.8	9.2	10.0	10.2	12.1
18–44 years	5.1	6.6	5.6	7.5	7.7	10.9	11.6	11.6	13.8
18–24 years	6.4	8.8	8.1	10.3	10.4	14.5	15.4	14.2	17.9
19–25 years	6.3	8.5	7.3	9.0	9.1	12.6	13.4	12.1	16.1
25–34 years	5.3	6.8	5.5	7.6	7.8	11.1	11.4	11.7	13.3
35–44 years	3.5	5.2	4.3	5.7	5.8	8.1	8.8	9.6	11.3
45–64 years	3.4	4.6	4.5	5.4	5.5	6.8	8.0	8.4	9.9
45–54 years	3.2	4.0	4.2	5.4	5.5	7.0	8.2	8.6	9.8
55–64 years	3.6	5.6	4.9	5.4	5.5	6.6	7.7	8.2	9.9
Sex									
Male	5.4	8.4	8.2	10.8	11.0	15.2	16.3	16.5	17.8
Female	8.1	11.1	10.8	13.7	13.9	18.5	19.7	19.8	21.4
Sex and marital status ⁵									
Male:									
Married	1.9	2.5	2.2	2.9	3.0	4.0	4.8	5.3	6.2
Divorced, separated, widowed	4.9	5.7	6.1	6.7	6.8	9.3	9.7	10.3	12.0
Never married	4.8	7.0	7.2	10.2	10.4	13.5	15.1	14.8	17.6
Female:									
Married	2.6	3.5	3.1	4.2	4.3	5.7	6.2	6.9	8.1
Divorced, separated, widowed	16.0	14.7	12.7	14.9	15.2	17.6	18.8	18.8	21.6
Never married	10.7	14.2	13.2	16.9	17.1	22.2	22.6	22.2	24.9
Race ⁶									
White only	4.6	7.4	7.1	10.2	10.4	14.5	15.5	15.6	16.9
Black or African American only	20.5	22.4	21.2	24.5	24.9	30.4	31.6	31.6	34.1
American Indian or Alaska Native only	*28.2	19.6	15.1	18.0	18.4	21.6	36.5	32.0	35.5
Asian only	*8.7	9.6	7.5	9.6	9.8	12.0	13.0	13.2	14.7
Native Hawaiian or Other Pacific Islander only	---	---	*	*	*	*	*	*	*
2 or more races	---	---	19.1	19.0	19.3	27.4	29.1	30.4	30.2
Hispanic origin and race ⁶									
Hispanic or Latino	13.3	17.6	15.5	21.9	22.5	28.6	30.5	29.5	31.3
Mexican	12.2	17.2	14.0	21.9	22.4	29.5	31.0	29.8	32.1
Puerto Rican	31.5	31.0	29.4	28.5	29.1	35.7	35.3	36.9	35.7
Cuban	*4.8	7.3	9.2	17.9	17.9	17.3	22.9	23.3	22.4
Other Hispanic or Latino	7.9	15.3	14.5	19.9	20.8	24.5	28.3	27.0	28.6
Not Hispanic or Latino	6.2	8.7	8.5	10.5	10.7	14.4	15.2	15.5	16.9
White only	3.7	6.1	6.1	7.8	7.9	11.0	11.5	11.9	13.0
Black or African American only	20.7	22.1	21.0	24.1	24.6	30.0	31.3	31.3	33.4
Age and percent of poverty level ⁷									
Under 65 years:									
Below 100%	33.0	40.5	38.4	44.2	45.0	50.8	52.5	53.7	56.5
100%–199%	5.3	13.0	16.2	21.6	22.0	28.5	30.1	30.8	34.0
100%–133%	8.7	20.1	22.4	28.5	29.1	36.3	38.0	38.8	43.9
134%–199%	3.7	9.5	13.1	18.2	18.6	24.4	25.5	26.2	28.1
200%–399%	0.8	2.7	4.0	6.1	6.1	8.4	9.0	9.0	9.9
400% or more	0.2	0.8	0.9	1.5	1.5	2.0	1.7	1.9	2.2

See footnotes at end of table.

Table 104 (page 2 of 3). Medicaid coverage among persons under age 65, by selected characteristics: United States, selected years 1984–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#104>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	1984 ¹	1997	2000 ²	2004(1) ³	2004(2) ³	2010 ³	2012 ³	2013 ³	2014 ³
Percent of population									
Under 19 years:									
Below 100%	42.0	56.4	56.9	67.5	68.9	78.4	82.1	82.1	83.3
100%–199%	6.5	20.3	27.8	38.7	39.5	53.5	56.3	58.9	59.1
100%–133%	10.3	31.1	36.4	48.3	49.2	63.5	68.8	70.3	71.7
134%–199%	4.7	14.8	23.3	33.9	34.6	47.7	48.8	51.6	50.6
200%–399%	1.0	4.4	7.6	12.1	12.2	17.7	18.8	19.1	18.5
400% or more	*	1.3	2.1	3.2	3.2	4.3	3.6	3.2	3.8
Under 18 years:									
Below 100%	43.3	58.0	58.5	69.2	70.7	79.8	83.7	83.9	84.7
100%–199%	6.6	20.8	28.4	39.5	40.2	54.3	57.3	60.1	60.0
100%–133%	10.4	32.0	36.9	48.9	49.8	64.6	70.1	71.2	72.3
134%–199%	4.8	15.1	23.8	34.7	35.4	48.2	49.6	52.9	51.6
200%–399%	1.0	4.5	7.6	12.2	12.3	18.0	19.1	19.5	18.9
400% or more	*	1.3	2.2	3.3	3.3	4.3	3.6	3.3	4.0
18–64 years:									
Below 100%	25.3	28.0	24.9	28.6	28.9	32.4	34.0	35.4	39.7
100%–199%	4.5	8.6	9.1	11.9	12.2	15.7	16.8	17.1	21.4
100%–133%	7.6	13.0	13.2	17.0	17.4	21.0	21.8	22.0	28.3
134%–199%	3.1	6.5	7.2	9.5	9.7	13.0	13.9	14.4	17.6
200%–399%	0.7	1.9	2.4	3.4	3.4	4.8	5.1	5.1	6.7
400% or more	0.2	0.7	0.6	1.0	1.0	1.3	1.2	1.6	1.7
Disability measure among adults 18–64 years ⁸									
Any basic actions difficulty or complex activity limitation	---	13.2	12.8	14.7	14.9	17.8	19.3	21.1	22.9
Any basic actions difficulty	---	12.7	12.2	14.0	14.2	16.7	18.4	20.6	22.0
Any complex activity limitation	---	22.9	23.2	23.9	24.1	30.0	30.8	32.3	35.6
No disability	---	3.5	3.0	4.5	4.7	6.8	7.0	6.8	8.7
Geographic region									
Northeast	8.6	11.3	10.6	12.8	13.0	17.9	19.3	20.8	21.4
Midwest	7.4	8.4	8.0	10.2	10.4	17.3	16.3	16.9	18.6
South	5.1	8.7	9.4	12.2	12.4	16.0	17.8	17.8	18.7
West	7.0	11.7	10.4	14.2	14.4	17.1	19.1	18.0	20.9
Location of residence ⁹									
Within MSA	7.1	9.7	8.9	11.7	11.9	16.1	17.4	17.4	18.9
Outside MSA	6.1	10.1	11.9	14.8	15.0	21.4	21.4	22.5	24.4

See footnotes at end of table.

Table 104 (page 3 of 3). Medicaid coverage among persons under age 65, by selected characteristics: United States, selected years 1984–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#104>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

-- Data not available.

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.

¹Data prior to 1997 are not strictly comparable with data for later years due to the 1997 questionnaire redesign. See Appendix I, National Health Interview Survey (NHIS) and Appendix II, Health insurance coverage.

²Estimates for 2000–2002 were calculated using 2000-based sample weights and may differ from estimates in other reports that used 1990-based sample weights for 2000–2002 estimates.

³Beginning in quarter 3 of the 2004 NHIS, persons under age 65 with no reported coverage were asked explicitly about Medicaid coverage. Estimates were calculated without and with the additional information from this question in the columns labeled 2004(1) and 2004(2), respectively, and estimates were calculated with the additional information starting with 2005 data.

⁴Includes all other races not shown separately, those with unknown marital status, unknown disability status, and, in 1984 and 1989, persons with unknown poverty level. Includes persons aged 14–64.

⁵The race groups, white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The five single-race categories plus multiple-race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group; the category 2 or more races includes persons who reported more than one racial group. Prior to 1999, data were tabulated according to the 1977 Standards with four racial groups, and the Asian only category included Native Hawaiian or Other Pacific Islander. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

⁷Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Poverty level was unknown for 10%–11% of persons under age 65 in 1984 and 1989. Missing family income data were imputed for 1995 and beyond. See Appendix II, Family income; Poverty; Table VI.

⁸Any basic actions difficulty or complex activity limitation is defined as having one or more of the following limitations or difficulties: movement difficulty, emotional difficulty, sensory (seeing or hearing) difficulty, cognitive difficulty, self-care (activities of daily living or instrumental activities of daily living) limitation, social limitation, or work limitation. For more information, see Appendix II, Basic actions difficulty; Complex activity limitation. Starting with 2007 data, the hearing question, a component of the basic actions difficulty measure, was revised. Consequently, data prior to 2007 are not comparable with data for 2007 and beyond. For more information on the impact of the revised hearing question, see Appendix II, Hearing trouble.

⁹MSA is metropolitan statistical area. Starting with 2006 data, MSA status is determined using 2000 census data and the 2000 standards for defining MSAs. For data prior to 2006, see Appendix II, Metropolitan statistical area (MSA) for the applicable standards.

NOTES: The category Medicaid coverage includes persons who had any of the following at the time of interview: Medicaid, other public assistance through 1996, state-sponsored health plan starting in 1997, or Children's Health Insurance Program (CHIP) starting in 1999; it includes those who also had another type of coverage in addition to one of these. In 2014, 17.0% of persons under age 65 reported being covered by Medicaid, 1.4% by state-sponsored health plans, and 1.3% by CHIP. Estimates may not sum to total because of rounding. The number of persons with Medicaid coverage was calculated by multiplying the percentage with Medicaid coverage by the number of persons under age 65 in the civilian noninstitutionalized U.S. population, which was determined from the post-stratification Census control total for each survey year. Percentages of persons with Medicaid coverage were calculated with unknown values excluded from denominators. See Appendix II, Health insurance coverage; Medicaid. Standard errors are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey, health insurance supplements (1984, 1989, 1994–1996). Starting with 1997, data are from the family core and the sample adult questionnaires. See Appendix I, National Health Interview Survey (NHIS).

Table 105 (page 1 of 3). No health insurance coverage among persons under age 65, by selected characteristics: United States, selected years 1984–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#105>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	1984 ¹	1997	2000 ²	2004(1) ³	2004(2) ³	2010 ³	2012 ³	2013 ³	2014 ³
Number, in millions									
Total ⁴	29.8	41.0	41.4	42.1	41.6	48.3	45.2	44.6	35.7
Percent of population									
Total ⁴	14.5	17.5	17.0	16.6	16.4	18.2	16.9	16.7	13.3
Age									
Under 19 years	14.1	14.4	12.9	10.1	9.6	8.3	7.0	7.1	5.7
Under 6 years	14.9	12.5	11.8	8.9	8.2	6.3	4.6	5.0	4.1
6–18 years	13.8	15.2	13.4	10.6	10.3	9.2	8.1	8.0	6.5
Under 18 years	13.9	14.0	12.6	9.7	9.2	7.8	6.6	6.6	5.4
6–17 years	13.4	14.7	13.0	10.0	9.7	8.6	7.6	7.4	6.1
18–64 years	14.8	19.0	18.9	19.4	19.3	22.3	20.9	20.5	16.3
18–44 years	17.1	22.4	22.4	23.6	23.5	27.1	24.8	24.2	19.7
18–24 years	25.0	30.1	30.4	30.1	30.0	31.4	24.5	24.6	18.1
19–25 years	25.1	31.5	32.3	32.3	32.2	33.8	26.3	26.7	19.7
25–34 years	16.2	23.8	23.3	25.7	25.5	28.3	28.1	27.1	22.7
35–44 years	11.2	16.7	16.9	17.6	17.5	22.6	21.7	21.0	17.7
45–64 years	9.6	12.4	12.6	12.9	12.8	15.7	15.6	15.4	11.8
45–54 years	10.5	12.8	12.8	13.7	13.6	17.9	17.7	17.1	13.7
55–64 years	8.7	11.8	12.4	11.7	11.6	12.8	13.2	13.5	9.7
Sex									
Male	15.3	18.7	18.1	18.1	17.9	20.3	18.5	18.1	14.7
Female	13.8	16.3	15.9	15.2	14.9	16.1	15.4	15.2	11.9
Sex and marital status ⁵									
Male:									
Married	11.1	13.9	14.1	14.5	14.4	17.2	16.2	15.9	12.6
Divorced, separated, widowed	24.9	28.8	25.8	27.1	27.0	31.4	29.3	28.1	23.2
Never married	22.4	27.9	27.2	27.6	27.5	31.1	27.5	26.9	21.9
Female:									
Married	11.2	13.0	13.3	13.2	13.1	14.7	14.6	14.6	11.6
Divorced, separated, widowed	19.2	23.2	21.3	23.3	23.0	23.6	24.2	22.8	17.7
Never married	16.3	20.5	21.1	19.6	19.3	21.9	19.6	19.6	15.1
Race ⁶									
White only	13.6	16.4	15.4	16.3	16.1	17.6	16.7	16.3	13.3
Black or African American only	19.9	20.1	19.5	18.1	17.6	20.6	18.0	18.9	13.7
American Indian or Alaska Native only	22.5	38.1	38.4	35.0	34.6	44.0	27.0	29.4	28.3
Asian only	18.5	19.5	17.6	16.7	16.5	17.1	16.8	14.2	10.8
Native Hawaiian or Other Pacific Islander only	---	---	*	*	*	*	*	*	*
2 or more races	---	---	16.8	12.6	12.3	15.8	14.5	15.3	10.1
Hispanic origin and race ⁶									
Hispanic or Latino	29.5	34.5	35.6	35.1	34.4	32.0	30.4	30.7	25.5
Mexican	33.8	39.4	39.9	38.1	37.6	34.8	33.2	33.4	27.2
Puerto Rican	18.3	19.0	16.4	21.0	20.4	13.7	14.4	15.6	13.0
Cuban	21.6	21.1	25.4	22.8	22.8	26.5	24.3	26.6	19.4
Other Hispanic or Latino	27.4	33.0	33.4	33.3	32.3	32.4	30.1	28.8	26.2
Not Hispanic or Latino	13.2	15.2	14.0	13.3	13.2	15.2	13.9	13.4	10.5
White only	11.9	13.8	12.5	12.1	12.0	13.7	12.7	12.2	9.7
Black or African American only	19.7	20.0	19.5	17.8	17.3	20.7	17.8	18.8	13.5
Age and percent of poverty level ⁷									
Under 65 years:									
Below 100%	33.9	33.7	34.2	31.8	31.0	30.3	28.2	28.0	23.0
100%–199%	21.8	30.6	31.0	29.4	29.0	32.4	29.3	29.3	23.4
100%–133%	28.8	36.6	35.7	32.3	31.7	34.9	31.1	30.4	25.0
134%–199%	18.7	27.7	28.7	28.0	27.6	31.0	28.4	28.6	22.4
200%–399%	7.6	14.2	15.4	15.7	15.6	17.4	16.2	16.1	12.6
400% or more	3.2	6.1	5.9	5.9	5.9	5.6	4.9	4.8	3.8

See footnotes at end of table.

Table 105 (page 2 of 3). No health insurance coverage among persons under age 65, by selected characteristics: United States, selected years 1984–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#105>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	1984 ¹	1997	2000 ²	2004(1) ³	2004(2) ³	2010 ³	2012 ³	2013 ³	2014 ³
Percent of population									
Under 19 years:									
Below 100%	29.0	23.8	22.6	17.2	15.7	11.3	8.3	8.9	6.7
100%–199%	18.0	23.7	22.1	16.5	15.8	13.5	11.1	11.7	9.3
100%–133%	24.4	28.2	26.5	18.4	17.6	15.9	9.6	11.6	9.4
134%–199%	14.9	21.4	19.7	15.5	14.9	12.0	12.0	11.7	9.3
200%–399%	5.1	9.7	9.6	8.1	8.0	7.4	6.8	6.7	5.7
400% or more	1.8	4.0	3.5	2.8	2.8	2.3	2.2	1.9	1.7
Under 18 years:									
Below 100%	28.9	23.2	22.0	16.5	15.0	10.6	7.6	8.2	6.4
100%–199%	17.5	23.2	21.7	15.8	15.1	12.7	10.4	11.1	8.7
100%–133%	24.0	28.1	26.4	17.9	17.1	15.1	9.0	11.2	8.9
134%–199%	14.4	20.7	19.1	14.7	14.1	11.3	11.3	11.1	8.5
200%–399%	4.9	9.4	9.3	7.7	7.6	7.0	6.7	6.3	5.5
400% or more	1.8	3.9	3.3	2.6	2.6	2.1	2.1	1.8	*1.6
18–64 years:									
Below 100%	37.6	41.2	42.4	41.4	41.0	42.7	40.5	40.0	32.9
100%–199%	24.4	34.7	36.4	36.7	36.5	42.1	38.6	37.8	30.5
100%–133%	31.9	41.7	41.7	40.4	40.0	45.7	42.2	40.4	33.9
134%–199%	21.1	31.5	34.0	35.0	34.8	40.3	36.5	36.4	28.7
200%–399%	8.9	16.4	18.2	19.1	19.1	21.3	19.8	19.7	15.3
400% or more	3.4	6.7	6.6	6.8	6.8	6.5	5.6	5.6	4.3
Disability measure among adults 18–64 years ⁸									
Any basic actions difficulty or complex activity limitation	---	20.1	17.6	19.8	19.6	20.8	20.4	20.4	16.2
Any basic actions difficulty	---	20.1	17.6	20.0	19.8	20.9	20.3	20.4	16.3
Any complex activity limitation	---	20.2	16.1	18.1	17.9	17.2	18.3	17.1	12.5
No disability	---	17.6	18.5	19.3	19.2	21.6	20.4	19.9	16.3
Geographic region									
Northeast	10.2	13.5	12.2	11.9	11.8	12.4	11.5	11.2	9.3
Midwest	11.3	13.2	12.3	12.6	12.4	14.1	13.6	13.1	10.3
South	17.7	20.9	20.5	20.2	19.9	21.9	20.3	19.9	16.9
West	18.2	20.6	20.7	19.1	18.9	20.6	19.0	18.9	13.3
Location of residence ⁹									
Within MSA	13.6	16.9	16.6	16.4	16.2	17.8	16.4	16.2	13.0
Outside MSA	16.6	19.8	18.6	17.4	17.2	20.4	19.9	19.3	15.2

See footnotes at end of table.

Table 105 (page 3 of 3). No health insurance coverage among persons under age 65, by selected characteristics: United States, selected years 1984–2014

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#105>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

-- Data not available.

* Estimates are considered unreliable. Data not shown have a relative standard error greater than 30%.

¹Data prior to 1997 are not strictly comparable with data for later years due to the 1997 questionnaire redesign. See Appendix I, National Health Interview Survey (NHIS) and Appendix II, Health insurance coverage.

²Estimates for 2000–2002 were calculated using 2000-based sample weights and may differ from estimates in other reports that used 1990-based sample weights for 2000–2002 estimates.

³Beginning in quarter 3 of the 2004 NHIS, persons under age 65 with no reported coverage were asked explicitly about Medicaid coverage. Estimates were calculated without and with the additional information from this question in the columns labeled 2004(1) and 2004(2), respectively, and estimates were calculated with the additional information starting with 2005 data.

⁴Includes all other races not shown separately, those with unknown marital status, unknown disability status, and, in 1984 and 1989, persons with unknown poverty level.

⁵Includes persons aged 14–64.

⁶The race groups, white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The five single-race categories plus multiple-race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group; the category 2 or more races includes persons who reported more than one racial group. Prior to 1999, data were tabulated according to the 1977 Standards with four racial groups, and the Asian only category included Native Hawaiian or Other Pacific Islander. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

⁷Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Poverty level was unknown for 10%–11% of persons under age 65 in 1984 and 1989. Missing family income data were imputed for 1995 and beyond. See Appendix II, Family income; Poverty; Table VI.

⁸Any basic actions difficulty or complex activity limitation is defined as having one or more of the following limitations or difficulties: movement difficulty, emotional difficulty, sensory (seeing or hearing) difficulty, cognitive difficulty, self-care (activities of daily living or instrumental activities of daily living) limitation, social limitation, or work limitation. For more information, see Appendix II, Basic actions difficulty; Complex activity limitation. Starting with 2007 data, the hearing question, a component of the basic actions difficulty measure, was revised. Consequently, data prior to 2007 are not comparable with data for 2007 and beyond. For more information on the impact of the revised hearing question, see Appendix II, Hearing trouble.

⁹MSA is metropolitan statistical area. Starting with 2006 data, MSA status is determined using 2000 census data and the 2000 standards for defining MSAs. For data prior to 2006, see Appendix II, Metropolitan statistical area (MSA) for the applicable standards.

NOTES: Persons not covered by private insurance, Medicaid, Children's Health Insurance Program (CHIP), public assistance (through 1996), state-sponsored or other government-sponsored health plans (starting in 1997), Medicare, or military plans are considered to have no health insurance coverage. Persons with only Indian Health Service coverage are considered to have no health insurance coverage. Health insurance coverage is at the time of interview. The number of persons with no health insurance coverage was calculated by multiplying the percentage with no coverage by the number of persons under age 65 in the civilian noninstitutionalized U.S. population, which was determined from the post-stratification Census control total for each survey year. Percentages of persons without coverage were calculated with unknown values excluded from denominators. See Appendix II, Children's Health Insurance Program (CHIP); Health insurance coverage; Medicaid. Standard errors are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey, health insurance supplements (1984, 1989, 1994–1996). Starting with 1997, data are from the family core and the sample adult questionnaires. See Appendix I, National Health Interview Survey (NHIS).

Table 106 (page 1 of 2). Health insurance coverage of noninstitutionalized Medicare beneficiaries aged 65 and over, by type of coverage and selected characteristics: United States, selected years 1992–2013

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#106>.

[Data are based on household interviews of a sample of noninstitutionalized Medicare beneficiaries]

Characteristic	Medicare Advantage plan ¹					Medicaid ²				
	1992	1995	2000	2012	2013	1992	1995	2000	2012	2013
Age										
Number, in millions										
65 years and over	1.1	2.6	5.9	12.4	13.8	2.7	2.8	2.7	3.6	3.6
Percent distribution										
65 years and over	3.9	8.9	19.3	29.1	31.2	9.4	9.6	9.0	8.5	8.1
65–74 years	4.2	9.5	20.6	29.1	30.3	7.9	8.8	8.5	7.6	7.2
75–84 years	3.7	8.3	18.5	30.1	33.8	10.6	9.6	8.9	9.2	8.6
85 years and over	*	7.3	16.3	26.3	29.4	16.6	13.6	11.2	10.3	11.4
Sex										
Male	4.6	9.2	19.3	28.9	30.0	6.3	6.2	6.3	5.6	5.7
Female	3.4	8.6	19.3	29.2	32.3	11.6	12.0	10.9	10.7	10.1
Race and Hispanic origin										
White, not Hispanic or Latino . . .	3.6	8.4	18.4	26.9	29.1	5.6	5.4	5.1	5.1	4.9
Black, not Hispanic or Latino . . .	*	7.9	20.7	30.3	33.4	28.5	30.3	23.6	19.0	17.8
Hispanic	*	15.5	27.5	45.5	46.7	39.0	40.5	28.7	20.2	20.1
Percent of poverty level ³										
Below 100%	3.6	7.7	18.4	---	---	22.3	17.2	15.9	---	---
100%–less than 200%	3.7	9.5	23.4	---	---	6.7	6.3	8.4	---	---
200% or more	4.2	10.1	18.0	---	---	*	*	*	---	---
Marital status										
Married	4.6	9.5	18.7	28.6	30.2	4.0	4.3	4.3	3.9	3.5
Widowed	2.3	7.7	19.4	28.6	31.2	14.9	15.0	13.6	13.8	12.5
Divorced	*	9.7	24.4	32.4	35.2	23.4	24.5	20.2	15.7	16.0
Never married	*	*	15.8	27.4	32.1	19.2	19.0	17.0	14.5	17.9
Employer-sponsored plan ⁴										
Medigap ⁵										
Characteristic	1992	1995	2000	2012	2013	1992	1995	2000	2012	2013
Age										
Number, in millions										
65 years and over	12.5	11.3	10.7	12.0	12.1	9.9	9.5	7.6	8.0	8.2
Percent distribution										
65 years and over	42.8	38.6	35.2	28.1	27.4	33.9	32.5	25.0	18.9	18.7
65–74 years	46.9	41.1	36.6	29.8	29.5	31.4	29.9	21.7	17.5	17.5
75–84 years	38.2	37.1	35.0	25.9	24.4	37.5	35.2	27.8	20.2	19.8
85 years and over	31.6	30.2	29.4	26.1	24.9	38.3	37.6	31.1	22.1	21.1
Sex										
Male	46.3	42.1	37.7	30.2	29.1	30.6	30.0	23.4	17.2	17.2
Female	40.4	36.0	33.4	26.4	25.9	36.2	34.4	26.2	20.3	19.9
Race and Hispanic origin										
White, not Hispanic or Latino . . .	45.9	41.3	38.6	30.6	29.5	37.2	36.2	28.3	22.4	22.1
Black, not Hispanic or Latino . . .	25.9	26.7	22.0	27.4	26.7	13.6	10.2	7.5	5.7	5.3
Hispanic	20.7	16.9	15.8	13.1	13.9	15.8	10.1	11.3	6.3	5.6
Percent of poverty level ³										
Below 100%	29.0	32.1	28.1	---	---	30.8	29.8	22.6	---	---
100%–less than 200%	37.5	32.0	27.0	---	---	39.3	39.1	28.4	---	---
200% or more	58.4	52.8	49.0	---	---	32.8	32.2	26.2	---	---
Marital status										
Married	49.9	44.6	41.0	32.9	33.0	33.0	32.6	25.6	20.1	19.6
Widowed	34.1	30.3	28.7	22.6	21.3	37.5	35.2	26.7	19.9	19.9
Divorced	27.3	26.6	22.4	19.7	17.2	27.9	24.1	16.9	13.0	14.8
Never married	38.0	35.1	28.5	24.3	21.5	29.1	26.2	21.9	16.7	13.1

See footnotes at end of table.

Table 106 (page 2 of 2). Health insurance coverage of noninstitutionalized Medicare beneficiaries aged 65 and over, by type of coverage and selected characteristics: United States, selected years 1992–2013

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#106>.

[Data are based on household interviews of a sample of noninstitutionalized Medicare beneficiaries]

Characteristic	Medicare fee-for-service only or Other ⁶				
	1992	1995	2000	2012	2013
Age					
Number, in millions					
65 years and over	2.9	3.1	3.5	6.6	6.4
Percent distribution					
65 years and over	9.9	10.5	11.5	15.5	14.6
65–74 years	9.7	10.7	12.6	16.0	15.5
75–84 years	10.1	9.9	9.9	14.6	13.5
85 years and over	10.8	11.3	12.1	15.1	13.3
Sex					
Male	12.2	12.6	13.3	18.0	18.0
Female	8.3	8.9	10.2	13.4	11.9
Race and Hispanic origin					
White, not Hispanic or Latino . . .	7.7	8.7	9.6	15.1	14.3
Black, not Hispanic or Latino . . .	26.7	25.0	26.1	17.6	16.8
Hispanic	18.3	17.1	16.7	14.9	13.7
Percent of poverty level ³					
Below 100%	14.3	13.3	15.1	---	---
100%–less than 200%	12.9	13.1	12.7	---	---
200% or more	4.0	4.5	6.3	---	---
Marital status					
Married	8.5	9.0	10.5	14.6	13.8
Widowed	11.2	11.9	11.6	15.1	15.2
Divorced	15.7	15.1	16.1	19.3	16.9
Never married	*	13.1	16.8	17.2	15.4

* Estimates are considered unreliable if the sample cell size is 50 or fewer.

--- Data not available.

¹Enrollee has a Medicare Advantage plan regardless of other insurance. Medicare Advantage plans include health maintenance organizations, preferred provider organizations, private fee-for-service plans, special needs plans, and Medicare medical savings account plans. Starting with 2013 data, the term Medicare Risk Health Maintenance Organization was replaced with Medicare Advantage plan. See Appendix II, Managed care.

²Enrolled in Medicaid and not enrolled in a Medicare Advantage plan. See Appendix II, Managed care.

³Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. See Appendix II, Family income; Poverty.

⁴Private insurance plans purchased through employers (own, current, or former employer, family business, union, or former employer or union of spouse) and not enrolled in a Medicare Advantage plan or Medicaid.

⁵Supplemental insurance purchased privately or through organizations such as American Association of Retired Persons or professional organizations, and not enrolled in a Medicare Advantage plan, Medicaid, or employer-sponsored plan.

⁶Medicare fee-for-service only or other public plans (except Medicaid).

NOTES: Data for noninstitutionalized Medicare beneficiaries. Insurance categories are mutually exclusive. Persons with more than one type of coverage are categorized according to the order in which the health insurance categories appear in the table. See Appendix I, Medicare Current Beneficiary Survey (MCBS). Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: Centers for Medicare & Medicaid Services, Medicare Current Beneficiary Survey, Access to Care file. See Appendix I, Medicare Current Beneficiary Survey (MCBS).

Table 107 (page 1 of 2). Medicare enrollees and expenditures and percent distribution, by Medicare program and type of service: United States and other areas, selected years 1970–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#107>.

[Data are compiled from various sources by the Centers for Medicare & Medicaid Services]

Medicare program and type of service	1970	1980	1990	1995	2000	2005	2010	2011	2012	2013	2014 ¹
Enrollees											
Number, in millions											
Total Medicare ²	20.4	28.4	34.3	37.6	39.7	42.6	47.7	48.9	50.9	52.5	53.8
Hospital insurance	20.1	28.0	33.7	37.2	39.3	42.2	47.4	48.5	50.5	52.1	53.5
Supplementary medical insurance (SMI) ³	19.5	27.3	32.6	35.6	37.3	---	---	---	---	---	---
Part B	19.5	27.3	32.6	35.6	37.3	39.8	43.9	44.9	46.5	47.9	49.3
Part D ⁴	---	---	---	---	---	1.8	34.8	35.7	37.4	39.1	40.5
Expenditures											
Amount, in billions											
Total Medicare	\$7.5	\$36.8	\$111.0	\$184.2	\$221.8	\$336.4	\$522.9	\$549.1	\$574.2	\$582.9	\$613.3
Total hospital insurance (HI)	5.3	25.6	67.0	117.6	131.1	182.9	247.9	256.7	266.8	266.2	269.3
HI payments to managed care organizations ⁵	---	0.0	2.7	6.7	21.4	24.9	60.7	64.6	70.2	73.1	74.0
HI payments for fee-for-service utilization	5.1	25.0	63.4	109.5	105.1	156.6	183.3	187.0	189.2	184.7	186.4
Inpatient hospital	4.8	24.1	56.9	82.3	87.1	123.3	135.9	133.9	138.9	134.1	135.6
Skilled nursing facility	0.2	0.4	2.5	9.1	11.1	19.3	27.1	31.9	28.4	28.4	28.8
Home health agency	0.1	0.5	3.7	16.2	4.0	6.0	7.2	7.1	6.8	6.9	6.6
Hospice	---	---	0.3	1.9	2.9	8.0	13.1	14.0	15.0	15.3	15.5
Other programs ⁶	---	---	---	---	---	---	---	0.9	2.8	3.5	3.7
Home health agency transfer ⁷	---	---	---	---	1.7	---	---	---	---	---	---
Medicare Advantage premiums ⁸	---	---	---	---	---	---	0.2	0.2	0.2	0.3	0.3
Accounting error (CY 2005–2008) ⁹	---	---	---	---	---	-1.9	---	---	---	---	---
Administrative expenses ¹⁰	0.2	0.5	0.9	1.4	2.9	3.3	3.8	4.0	4.3	4.7	4.9
Total supplementary medical insurance (SMI) ³	2.2	11.2	44.0	66.6	90.7	153.5	274.9	292.5	307.4	316.7	344.0
Total Part B	2.2	11.2	44.0	66.6	90.7	152.4	212.9	225.3	240.5	247.1	265.9
Part B payments to managed care organizations ⁵	0.0	0.2	2.8	6.6	18.4	22.0	55.2	59.1	66.0	72.7	85.7
Part B payments for fee-for-service utilization ¹¹	1.9	10.4	39.6	58.4	72.2	125.0	154.3	162.3	170.3	170.8	175.8
Physician/supplier ¹²	1.8	8.2	29.6	---	---	---	---	---	---	---	---
Outpatient hospital ¹³	0.1	1.9	8.5	---	---	---	---	---	---	---	---
Independent laboratory ¹⁴	0.0	0.1	1.5	---	---	---	---	---	---	---	---
Physician fee schedule	---	---	---	31.7	37.0	57.7	64.0	67.5	69.5	68.6	69.2
Durable medical equipment	---	---	---	3.7	4.7	8.0	8.3	8.2	8.2	7.2	6.3
Laboratory ¹⁵	---	---	---	4.3	4.4	6.9	8.4	8.4	9.2	9.1	8.2
Other ¹⁶	---	---	---	9.9	13.6	26.7	34.1	36.0	38.3	38.3	39.6
Hospital ¹⁷	---	---	---	8.7	8.1	18.7	27.6	30.2	33.6	36.1	41.4
Home health agency	0.0	0.2	0.1	0.2	4.5	7.1	12.0	12.1	11.4	11.6	11.2
Home health agency transfer ⁷	---	---	---	---	-1.7	---	---	---	---	---	---
Medicare Advantage premiums ⁸	---	---	---	---	---	0.2	0.2	0.2	0.2	0.3	0.3
Accounting error (CY 2005–2008) ⁹	---	---	---	---	---	1.9	---	---	---	---	---
Administrative expenses ¹⁰	0.2	0.6	1.5	1.6	1.8	2.8	3.2	3.7	4.0	3.4	4.1
Part D start-up costs ¹⁸	---	---	---	---	---	0.7	---	---	---	---	---
Total Part D ⁴	---	---	---	---	---	1.1	62.1	67.1	66.9	69.7	78.1
Percent distribution of expenditures											
Total hospital insurance (HI)	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
HI payments to managed care organizations ⁵	---	0.0	4.0	5.7	16.3	13.6	24.5	25.2	26.3	27.5	27.5
HI payments for fee-for-service utilization	97.0	97.9	94.6	93.1	80.2	85.6	73.9	72.8	70.9	69.4	69.2
Inpatient hospital	91.4	94.3	85.0	70.0	66.4	67.4	54.8	52.2	52.1	50.4	50.3
Skilled nursing facility	4.7	1.5	3.7	7.8	8.5	10.6	10.9	12.4	10.7	10.7	10.7
Home health agency	1.0	2.1	5.5	13.8	3.1	3.3	2.9	2.8	2.6	2.6	2.5
Hospice	---	---	0.5	1.6	2.2	4.4	5.3	5.5	5.6	5.7	5.7
Other programs ⁶	---	---	---	---	---	---	---	0.3	1.1	1.3	1.4
Home health agency transfer ⁷	---	---	---	---	1.3	---	---	---	---	---	---
Medicare Advantage premiums ⁸	---	---	---	---	---	---	0.1	0.1	0.1	0.1	0.1
Accounting error (CY 2005–2008) ⁹	---	---	---	---	---	-1.0	---	---	---	---	---
Administrative expenses ¹⁰	3.0	2.1	1.4	1.2	2.2	1.8	1.5	1.6	1.6	1.8	1.8

See footnotes at end of table.

Table 107 (page 2 of 2). Medicare enrollees and expenditures and percent distribution, by Medicare program and type of service: United States and other areas, selected years 1970–2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#107>.

[Data are compiled from various sources by the Centers for Medicare & Medicaid Services]

Medicare program and type of service	1970	1980	1990	1995	2000	2005	2010	2011	2012	2013	2014 ¹
Percent distribution of expenditures											
Total supplementary medical insurance (SMI) ³	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Total Part B	100.0	100.0	100.0	100.0	100.0	99.3	77.4	77.1	78.2	78.0	77.3
Part B payments to managed care organizations ⁵	1.2	1.8	6.4	9.9	20.2	14.3	20.1	20.2	21.5	22.9	24.9
Part B payments for fee-for-service utilization ¹¹	88.1	92.8	90.1	87.6	79.6	81.5	56.1	55.5	55.4	53.9	51.1
Physician/supplier ¹²	80.9	72.8	67.3	---	---	---	---	---	---	---	---
Outpatient hospital ¹³	5.2	16.9	19.3	---	---	---	---	---	---	---	---
Independent laboratory ¹⁴	0.5	1.0	3.4	---	---	---	---	---	---	---	---
Physician fee schedule	---	---	---	47.5	40.8	37.6	23.3	23.1	22.6	21.7	20.1
Durable medical equipment	---	---	---	5.5	5.2	5.2	3.0	2.8	2.7	2.3	1.8
Laboratory ¹⁵	---	---	---	6.4	4.8	4.5	3.1	2.9	3.0	2.9	2.4
Other ¹⁶	---	---	---	14.8	15.0	17.4	12.4	12.3	12.5	12.1	11.5
Hospital ¹⁷	---	---	---	13.0	8.9	12.2	10.0	10.3	10.9	11.4	12.0
Home health agency	1.5	2.1	0.2	0.3	4.9	4.6	4.4	4.1	3.7	3.7	3.3
Home health agency transfer ⁷	---	---	---	---	-1.9	---	---	---	---	---	---
Medicare Advantage premiums ⁸	---	---	---	---	---	---	0.1	0.1	0.1	0.1	0.1
Accounting error (CY 2005–2008) ⁹	---	---	---	---	---	1.2	---	---	---	---	---
Administrative expenses ¹⁰	10.7	5.4	3.5	2.4	2.0	1.8	1.2	1.3	1.3	1.1	1.2
Part D start-up costs ¹⁸	---	---	---	---	---	0.4	---	---	---	---	---
Total Part D ⁴	---	---	---	---	---	0.7	22.6	22.9	21.8	22.0	22.7

--- Category not applicable or data not available.

0.0 Quantity more than zero but less than 0.05.

¹Preliminary estimates.

²Average number enrolled in the hospital insurance (HI) and/or supplementary medical insurance (SMI) programs for the period. See Appendix II, Medicare.

³Starting with 2004 data, the SMI trust fund consists of two separate accounts: Part B (which pays for a portion of the costs of physicians' services, outpatient hospital services, and other related medical and health services for voluntarily enrolled individuals) and Part D (Medicare Prescription Drug Account, which pays private plans to provide prescription drug coverage).

⁴The Medicare Modernization Act, enacted December 8, 2003, established within SMI two Part D accounts related to prescription drug benefits: the Medicare Prescription Drug Account and the Transitional Assistance Account. The Medicare Prescription Drug Account is used in conjunction with the broad, voluntary prescription drug benefits that began in 2006. The Transitional Assistance Account was used to provide transitional assistance benefits, beginning in 2004 and extending through 2005, for certain low-income beneficiaries prior to the start of the new prescription drug benefit. The amounts shown for Total Part D expenditures—and thus for total SMI expenditures and total Medicare expenditures—for 2006 and later years include estimated amounts for premiums paid directly from Part D beneficiaries to Part D prescription drug plans.

⁵Medicare-approved managed care organizations. See Appendix II, Managed care.

⁶Includes Community-Based Care Transitions Program (\$0.1 billion in each of 2011–2014) and Electronic Health Records Incentive Program (\$0.7 billion in 2011, \$2.7 billion in 2012, \$3.4 billion in 2013, and \$3.6 billion in 2014).

⁷For 1998 to 2003, data reflects annual home health HI to SMI transfer amounts.

⁸When a beneficiary chooses a Medicare Advantage plan whose monthly premium exceeds the benchmark amount, the additional premiums (that is, amounts beyond those paid by Medicare to the plan) are the responsibility of the beneficiary. Beneficiaries subject to such premiums may choose to either reimburse the plans directly or have the additional premiums deducted from their Social Security checks. The amounts shown here are only those additional premiums deducted from Social Security checks. These amounts are transferred to the HI trust and SMI trust funds and then transferred from the trust funds to the plans.

⁹Represents misallocation of benefit payments between the HI trust fund and the Part B account of the SMI trust fund from May 2005 to September 2007, and the transfer made in June 2008 to correct the misallocation.

¹⁰Includes expenditures for research, experiments and demonstration projects, peer review activity (performed by Peer Review Organizations from 1983 to 2001 and by Quality Review Organizations from 2002 to present), and to combat and prevent fraud and abuse.

¹¹Type-of-service reporting categories for fee-for-service reimbursement differ before and after 1991.

¹²Includes payment for physicians, practitioners, durable medical equipment, and all suppliers other than independent laboratory through 1990. Starting with 1991 data, physician services subject to the physician fee schedule are shown. Payments for laboratory services paid under the laboratory fee schedule and performed in a physician office are included under Laboratory beginning in 1991. Payments for durable medical equipment are shown separately beginning in 1991. The remaining services from the Physician/supplier category are included in Other.

¹³Includes payments for hospital outpatient department services, skilled nursing facility outpatient services, Part B services received as an inpatient in a hospital or skilled nursing facility setting, and other types of outpatient facilities. Starting with 1991 data, payments for hospital outpatient department services, except for laboratory services, are listed under Hospital. Hospital outpatient laboratory services are included in the Laboratory line.

¹⁴Starting with 1991 data, those independent laboratory services that were paid under the laboratory fee schedule (most of the independent laboratory category) are included in the Laboratory line; the remaining services are included in the Physician fee schedule and Other lines.

¹⁵Payments for laboratory services paid under the laboratory fee schedule performed in a physician office, independent laboratory, or in a hospital outpatient department.

¹⁶Includes payments for physician-administered drugs; freestanding ambulatory surgical center facility services; ambulance services; supplies; freestanding end-stage renal disease (ESRD) dialysis facility services; rural health clinics; outpatient rehabilitation facilities; psychiatric hospitals; and federally qualified health centers.

¹⁷Includes the hospital facility costs for Medicare Part B services that are predominantly in the outpatient department, with the exception of hospital outpatient laboratory services, which are included on the Laboratory line. Physician reimbursement is included on the Physician fee schedule line.

¹⁸Part D start-up costs were funded through the SMI Part B account in 2004–2008.

NOTES: Estimates are subject to change as more recent data become available. Totals may not equal the sum of the components because of rounding. Estimates are for Medicare-covered services furnished to Medicare enrollees residing in the United States, Puerto Rico, Virgin Islands, Guam, other outlying areas, foreign countries, and unknown residence. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>. Estimates in this table have been revised and differ from previous editions of *Health, United States*.

SOURCE: Centers for Medicare & Medicaid Services (CMS), Office of the Actuary, Medicare and Medicaid Cost Estimates Group. Estimates are based on unpublished data from CMS, the Office of the Actuary, and Treasury Department financial statements.

Table 108 (page 1 of 2). Medicare beneficiaries, by race, Hispanic origin, and selected characteristics: United States, selected years 1992–2012

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#108>.

[Data are based on household interviews of a sample of Medicare beneficiaries and Medicare administrative records]

Characteristic	Not Hispanic or Latino											
	All			White			Black or African American			Hispanic or Latino		
	1992	2011	2012	1992	2011	2012	1992	2011	2012	1992	2011	2012
	Number of beneficiaries, in millions											
All Medicare beneficiaries	36.8	50.0	52.0	30.9	38.2	39.4	3.3	4.7	4.9	1.9	4.5	5.0
	Percent distribution of beneficiaries											
All Medicare beneficiaries	100.0	100.0	100.0	84.2	76.3	75.6	8.9	9.4	9.4	5.2	9.1	9.7
	Medical care use											
	Percent of beneficiaries with at least one service											
All Medicare beneficiaries:												
Long-term care facility stay	7.7	8.5	8.4	8.0	9.1	9.2	6.2	7.5	6.8	4.2	5.9	5.2
Community-only residents:												
Inpatient hospital	17.9	17.1	15.4	18.1	17.2	15.4	18.4	19.5	16.2	16.6	15.3	14.4
Outpatient hospital	57.9	73.5	68.8	57.8	74.5	70.4	61.1	73.5	70.0	53.1	67.6	58.7
Physician/supplier ¹	92.4	96.5	96.4	93.0	97.0	96.9	89.1	96.3	94.5	87.9	94.1	94.9
Dental	40.4	45.0	48.4	43.1	49.4	52.9	23.5	27.8	30.2	29.1	32.9	39.3
Prescription medicine	85.2	94.7	94.7	85.5	95.0	94.6	83.1	94.4	95.1	84.6	93.5	94.5
	Expenditures											
	Expenditures per beneficiary											
All Medicare beneficiaries:												
Total health care ²	\$6,716	\$16,383	\$17,197	\$6,816	\$16,195	\$17,064	\$7,043	\$20,061	\$19,975	\$5,784	\$15,976	\$16,656
Long-term care facility ³	1,581	2,763	2,767	1,674	2,929	3,053	1,255	2,964	2,424	*758	2,024	1,823
Community-only residents:												
Total personal health care	5,054	12,186	12,880	4,988	11,757	12,556	5,530	15,179	15,887	4,938	13,156	13,218
Inpatient hospital	2,098	2,330	2,128	2,058	2,223	2,035	2,493	3,220	2,781	1,999	2,412	2,385
Outpatient hospital	504	1,500	1,568	478	1,425	1,545	668	2,409	2,297	511	1,280	1,024
Physician/supplier ¹	1,524	3,135	3,553	1,525	3,219	3,672	1,398	3,105	3,605	1,587	2,642	2,939
Dental	142	408	455	153	447	499	70	219	298	97	279	375
Prescription medicine	468	2,987	3,261	481	2,916	3,155	417	3,421	3,755	389	3,139	3,461
Long-term care facility residents only:												
Long-term care facility ⁴	23,054	46,413	47,332	23,177	45,795	46,684	21,272	56,121	50,600	*25,026	*43,043	*51,241
	Sex											
	Percent distribution of beneficiaries											
Both sexes	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Male	42.9	45.3	45.5	42.7	45.3	45.1	42.0	42.6	45.5	46.7	48.8	48.6
Female	57.1	54.7	54.5	57.3	54.7	54.9	58.0	57.4	54.5	53.3	51.2	51.4
	Eligibility criteria and age											
All Medicare beneficiaries ⁵	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Disabled	10.2	16.7	16.7	8.6	13.9	13.7	19.1	31.5	31.6	16.5	23.1	22.1
Under 45 years	3.5	3.7	3.6	2.9	3.0	2.8	7.6	8.4	7.7	6.9	4.8	4.4
45–64 years	6.5	13.0	13.1	5.8	10.9	10.9	11.5	23.1	23.9	9.6	18.2	17.7
Aged	89.8	83.3	83.2	91.4	86.1	86.2	81.0	68.5	68.4	83.5	76.9	77.9
65–74 years	51.5	45.5	46.6	52.0	46.0	47.7	48.0	40.2	38.6	49.4	46.7	46.0
75–84 years	28.8	25.7	24.9	29.5	26.8	25.7	24.0	19.8	21.8	27.1	22.9	23.2
85 years and over	9.7	12.0	11.7	9.9	13.3	12.8	9.0	8.5	8.0	6.9	7.4	8.7
	Living arrangement ⁶											
All living arrangements	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Alone	27.0	28.8	28.1	27.5	29.5	28.8	27.7	32.4	30.9	20.2	22.0	23.6
With spouse	51.2	48.6	48.5	53.3	51.0	50.6	33.3	29.6	31.7	50.4	48.0	45.9
With children	9.1	10.3	10.4	7.7	7.9	8.2	16.8	20.4	19.1	16.6	17.4	17.0
With others	7.6	8.7	9.5	6.2	7.7	8.4	18.1	14.3	15.1	10.8	10.3	11.2
Long-term care facility	5.1	3.6	3.6	5.3	3.9	3.9	4.0	3.2	3.3	*2.0	*2.3	*2.3

See footnotes at end of table.

Table 108 (page 2 of 2). Medicare beneficiaries, by race, Hispanic origin, and selected characteristics: United States, selected years 1992–2012

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#108>.

[Data are based on household interviews of a sample of Medicare beneficiaries and Medicare administrative records]

Characteristic	Not Hispanic or Latino											
	All			White			Black or African American			Hispanic or Latino		
	1992	2011	2012	1992	2011	2012	1992	2011	2012	1992	2011	2012
Age and limitation of activity ⁷												
Percent distribution of beneficiaries												
Disabled, under age 65	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
None	22.7	26.0	24.0	21.8	23.9	23.1	26.2	35.2	30.3	21.2	24.8	22.4
IADL only	39.0	34.9	32.7	38.9	36.0	33.7	35.8	31.9	32.4	46.1	36.1	29.3
1 or 2 ADLs	21.2	23.8	25.4	21.5	24.7	26.2	21.2	20.5	20.5	*20.9	*22.6	*25.9
3–5 ADLs	17.2	15.3	17.9	17.9	15.4	17.0	*16.8	*12.3	16.8	*11.9	*16.4	22.4
65–74 years	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
None	67.0	71.9	68.5	68.7	73.6	70.1	55.1	65.7	62.9	59.2	68.5	65.0
IADL only	17.8	14.7	15.1	17.0	14.1	14.8	22.9	18.8	18.9	*20.9	15.0	14.5
1 or 2 ADLs	10.4	8.9	11.3	9.6	8.4	10.4	14.4	*9.5	*13.4	*15.7	*9.8	*12.1
3–5 ADLs	4.8	4.5	5.1	4.6	4.0	4.7	*7.6	*6.0	*4.8	*4.2	*6.7	*8.4
75–84 years	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
None	46.6	53.9	52.9	47.5	55.4	53.4	42.0	52.2	55.2	44.3	47.6	46.1
IADL only	23.9	22.4	21.6	23.6	21.6	22.1	26.7	20.0	*15.0	*27.8	27.9	22.5
1 or 2 ADLs	16.5	14.8	16.0	16.8	14.5	15.5	15.3	*17.0	19.0	*14.9	*14.2	17.2
3–5 ADLs	13.0	8.9	9.5	12.2	8.4	8.9	*15.9	*10.8	*10.9	*13.0	*10.3	*14.3
85 years and over	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
None	19.9	27.5	25.7	20.2	28.7	27.0	*19.6	*24.3	*18.7	*19.7	*17.5	*16.1
IADL only	20.9	27.6	25.6	20.2	28.3	26.2	*22.1	*18.7	*25.5	*24.7	*25.3	*19.9
1 or 2 ADLs	23.5	20.2	22.5	23.5	20.4	23.0	*24.3	*23.6	*18.4	*23.7	*15.7	*26.3
3–5 ADLs	35.8	24.8	26.1	36.1	22.6	23.9	*34.0	*33.3	*37.4	*31.8	*41.6	*37.7

* Estimates are based on 50 persons or fewer or have a relative standard error of 30% or higher and are considered unreliable.

¹Physician/supplier services include medical and osteopathic doctor and health practitioner visits, diagnostic laboratory and radiology services, medical and surgical services, and durable medical equipment and nondurable medical supplies.

²Total health care expenditures by Medicare beneficiaries, including expenses paid by Medicare and all other sources of payment for the following services: inpatient hospital, outpatient hospital, physician/supplier, dental, prescription medicine, home health, and hospice and long-term care facility care. Excluded are health insurance premiums.

³Expenditures for long-term care in facilities for all beneficiaries include facility room and board expenses for beneficiaries who resided in a facility for the full year, for beneficiaries who resided in a facility for part of the year and in the community for part of the year, and expenditures for short-term facility stays for full-year or part-year community residents. See Appendix II, Long-term care facility.

⁴Expenditures for facility-based long-term care for facility-based beneficiaries include facility room and board expenses for beneficiaries who resided in a facility for the full year and for beneficiaries who resided in a facility for part of the year and in the community for part of the year. Excluded are expenditures for short-term facility stays for full-year community residents. See Appendix II, Long-term care facility.

⁵Medicare beneficiaries with end-stage renal disease (ESRD) are included within the subgroups Aged and Disabled. In 2012, less than 1% of Medicare beneficiaries qualified because of ESRD. See Appendix II, Medicare.

⁶In 2012, less than 1% of Medicare beneficiaries had an unknown living arrangement.

⁷IADL is instrumental activities of daily living; ADL is activities of daily living. Includes data for both community and long-term care facility residents. See Appendix II, Activities of daily living (ADL); Instrumental activities of daily living (IADL).

NOTES: Percentages and percent distributions are calculated using unrounded numbers. Expenditures include expenses for Medicare beneficiaries paid by Medicare and all other sources of payment. Estimates include individuals enrolled in the hospital insurance (HI) and/or supplementary medical insurance (SMI) programs at any time during the calendar year. A new imputation methodology was used for 2012 estimates; therefore some utilization estimates may not be comparable to previous years.

SOURCE: Centers for Medicare & Medicaid Services, Medicare Current Beneficiary Survey, Cost and Use file, Health and Health Care of the Medicare Population. Available from: <http://www.cms.hhs.gov/mcbs> and unpublished data. See Appendix I, Medicare Current Beneficiary Survey (MCBS).

Table 109 (page 1 of 2). Medicaid and Children's Health Insurance Program beneficiaries and payments, by basis of eligibility, and race and Hispanic origin: United States, selected fiscal years 1999–2012

Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#109>.

[Data are compiled by the Centers for Medicare & Medicaid Services from the Medicaid Data System]

<i>Basis of eligibility and race and Hispanic origin</i>	1999	2000	2005	2007	2008	2009	2010	2011 ¹	2012 ¹
Beneficiaries²									
Number, in millions									
All beneficiaries	40.1	42.8	57.7	56.8	58.8	62.6	65.7	71.4	71.6
Percent of beneficiaries									
Basis of eligibility:									
Aged (65 years and over)	9.4	8.7	7.6	7.1	7.1	6.7	6.5	5.7	5.7
Blind and disabled	16.7	16.1	14.2	14.8	14.8	14.4	14.3	13.1	13.2
Adults in families with dependent children ³	18.7	20.5	21.8	21.8	22.0	23.1	23.7	18.9	19.5
Children under age 21 ⁴	46.9	46.1	47.2	48.4	47.8	47.7	48.3	40.5	40.6
Other Title XIX ⁵	8.4	8.6	9.1	7.8	8.4	8.1	7.2	9.6	8.7
Separate CHIP ⁶	---	---	---	---	---	---	---	12.2	12.3
Race and Hispanic origin:⁷									
White	---	---	39.3	38.6	38.1	38.2	38.9	37.2	37.2
Black or African American	---	---	21.5	21.6	21.1	20.7	20.6	19.8	20.1
American Indian or Alaska Native	---	---	1.2	1.2	1.3	1.2	1.2	1.1	1.1
Asian or Pacific Islander	---	---	3.5	3.5	3.5	3.6	3.6	3.6	3.8
Asian	---	---	2.5	2.6	2.6	2.7	2.7	2.8	2.9
Pacific Islander	---	---	0.9	0.9	0.9	0.9	0.9	0.9	0.8
Hispanic or Latino	---	---	20.6	21.6	21.7	22.3	22.3	17.4	17.4
Multiple race or unknown	---	---	13.9	13.5	14.3	14.0	13.3	20.9	20.4
Payments⁸									
Amount, in billions									
All payments	\$153.5	\$168.3	\$274.9	\$276.2	\$296.8	\$326.0	\$339.0	\$368.6	\$363.9
Percent distribution									
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Basis of eligibility:									
Aged (65 years and over)	27.7	26.4	23.1	20.7	20.6	19.7	19.4	16.6	16.7
Blind and disabled	42.9	43.2	43.4	43.3	43.5	43.4	43.4	40.8	41.2
Adults in families with dependent children ³	10.3	10.6	11.8	12.4	12.7	13.9	14.2	12.6	13.3
Children under age 21 ⁴	15.7	15.9	17.2	19.4	19.2	19.6	19.8	18.1	18.2
Other Title XIX ⁵	3.4	3.9	4.6	4.2	4.0	3.3	3.1	4.5	3.5
Separate CHIP ⁶	---	---	---	---	---	---	---	7.4	7.2
Race and Hispanic origin:⁷									
White	---	---	53.0	50.7	50.2	50.0	50.2	48.1	48.3
Black or African American	---	---	19.8	20.8	20.6	20.7	20.5	20.3	20.8
American Indian or Alaska Native	---	---	1.2	1.2	1.3	1.2	1.3	1.2	1.3
Asian or Pacific Islander	---	---	2.7	2.8	2.9	3.1	3.0	3.1	3.2
Asian	---	---	1.9	2.0	2.1	2.3	2.3	2.4	2.5
Pacific Islander	---	---	0.8	0.8	0.8	0.8	0.7	0.7	0.7
Hispanic or Latino	---	---	12.2	13.1	13.7	14.2	14.2	9.9	9.7
Multiple race or unknown	---	---	11.1	11.4	11.4	10.8	10.8	17.4	16.7
Payments per beneficiary⁸									
Amount									
All beneficiaries	\$3,819	\$3,936	\$4,768	\$4,862	\$5,051	\$5,209	\$5,160	\$5,159	\$5,082
Basis of eligibility:									
Aged (65 years and over)	11,268	11,929	14,427	14,141	14,742	15,337	15,286	15,073	14,862
Blind and disabled	9,832	10,559	14,531	14,194	14,843	15,670	15,695	16,104	15,825
Adults in families with dependent children ³	2,104	2,030	2,583	2,753	2,912	3,144	3,095	3,443	3,460
Children under age 21 ⁴	1,282	1,358	1,732	1,951	2,035	2,145	2,122	2,300	2,281
Other Title XIX ⁵	1,532	1,778	2,380	2,622	2,407	2,104	2,219	2,402	2,030
Separate CHIP ⁶	---	---	---	---	---	---	---	3,125	2,979
Race and Hispanic origin:⁷									
White	---	---	6,422	6,390	6,657	6,809	6,663	6,677	6,598
Black or African American	---	---	4,397	4,669	4,928	5,216	5,142	5,308	5,266
American Indian or Alaska Native	---	---	4,626	4,826	5,218	5,382	5,421	5,461	5,649
Asian or Pacific Islander	---	---	3,710	3,863	4,133	4,402	4,300	4,483	4,365
Asian	---	---	3,624	3,847	4,123	4,386	4,307	4,482	4,383
Pacific Islander	---	---	3,947	3,907	4,161	4,448	4,275	4,484	4,302
Hispanic or Latino	---	---	2,822	2,960	3,175	3,322	3,276	2,944	2,821
Multiple race or unknown	---	---	3,816	4,106	4,014	4,025	4,173	4,298	4,161

See footnotes at end of table.

Table 109 (page 2 of 2). Medicaid and Children's Health Insurance Program beneficiaries and payments, by basis of eligibility, and race and Hispanic origin: United States, selected fiscal years 1999–2012

Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#109>.

[Data are compiled by the Centers for Medicare & Medicaid Services from the Medicaid Data System]

-- Data not available.

¹Starting with 2011, a new tabular methodology was used. Therefore, estimates may not be comparable to earlier data and caution should be used with trend analysis.

²Beneficiaries include those who received services through Medicaid or the Children's Health Insurance Program (CHIP). Beneficiary counts for 2011 and 2012 data were derived from MSIS claims files. Separate CHIP beneficiaries are included for 2011 and 2012.

³Includes adults who meet the requirements for the Aid to Families with Dependent Children (AFDC) program that were in effect in their state on July 16, 1996, or, at state option, more liberal criteria (with some exceptions). Includes adults in the Temporary Assistance for Needy Families (TANF) program. Starting with 2001 data, includes women in the Breast and Cervical Cancer Prevention and Treatment Program and unemployed adults. For more information on the eligibility requirements, see Appendix II, Medicaid.

⁴Includes children (including those in the foster care system) in the TANF program. For more information on the eligibility requirements, see Appendix II, Medicaid.

⁵Includes some participants in the Supplemental Security Income program and other people deemed medically needy in participating states. Prior to 2001, includes unemployed adults. Excludes foster care children and includes unknown eligibility.

⁶CHIP is Children's Health Insurance Program. CHIP provides federal funds for states to provide health care coverage to eligible low-income, uninsured children who do not qualify for Medicaid. Some states use CHIP funds to expand Medicaid. For 2012 data, all states except Colorado and Idaho had separate CHIP beneficiaries. See Appendix II, Children's Health Insurance Program (CHIP).

⁷Race and Hispanic origin are as determined on initial Medicaid application. Categories are mutually exclusive. Starting with 2001 data, the Hispanic category included Hispanic persons, regardless of race. Persons indicating more than one race were included in the multiple race category.

⁸Payments for 2011 and 2012 data were derived from MSIS claims files. Medicaid payment data for 2010 and earlier excluded disproportionate share hospital (DSH) payments (\$14.7 billion in FY2010) and DSH mental health facility payments (\$2.9 billion in FY2010).

NOTES: Data are for fiscal year ending September 30. See Appendix II, Medicaid; Medicaid payments. For more information, see: <http://www.medicaid.gov>. Colorado and Idaho had not reported 2012 data as of the date accessed. Starting with 2011, a new tabular methodology was used. Therefore, estimates may not be comparable to earlier data and caution should be used with trend analysis. For more information on data and analytic issues, see:

<https://www.cms.gov/Research-Statistics-Data-and-Systems/Computer-Data-and-Systems/MedicaidDataSourcesGenInfo/MSIS-Tables.html>. Estimates for 2011 have been revised and differ from previous editions of *Health, United States*.

SOURCE: Centers for Medicare & Medicaid Services, Center for Medicaid and CHIP Services, Medicaid Statistical Information System (MSIS), granular file. MSIS data for 2011 and 2012 were accessed January 7, 2016. See Appendix I, Medicaid Statistical Information System (MSIS).

Table 110 (page 1 of 2). Medicaid and Children's Health Insurance Program beneficiaries and payments, by type of service: United States, selected fiscal years 1999–2012

Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#110>.

[Data are compiled by the Centers for Medicare & Medicaid Services from the Medicaid Data System]

Type of service	1999	2000	2005	2007	2008	2009	2010	2011 ¹	2012 ¹
Beneficiaries²									
Number, in millions									
All beneficiaries	40.2	42.8	57.7	56.8	58.8	62.6	65.7	71.4	71.6
Percent of beneficiaries									
Inpatient hospital	11.2	11.5	9.5	9.0	8.9	8.7	6.9	11.4	11.8
Mental health facility	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Intermediate care facility for individuals with intellectual disabilities ³	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.1	0.1
Nursing facility	4.0	4.0	3.0	2.9	2.7	2.6	2.4	2.5	2.3
Physician	45.7	44.7	42.0	38.8	36.9	36.9	36.9	62.6	64.4
Dental	14.0	13.8	16.2	16.8	16.7	17.8	19.1	26.6	27.4
Other practitioner	9.9	11.1	10.2	9.5	8.8	8.8	9.2	13.6	14.2
Outpatient hospital	30.9	30.9	28.2	26.2	25.2	26.4	24.2	38.8	39.5
Clinic	16.8	17.9	20.7	20.6	20.2	20.6	20.7	24.4	23.6
Laboratory and radiological	25.4	26.6	27.7	27.8	26.6	26.2	25.8	41.6	42.0
Home health	2.0	2.3	2.1	2.1	1.9	1.7	1.7	2.4	2.5
Prescribed drugs	49.4	48.0	49.2	42.1	41.8	42.6	44.7	56.8	57.2
Capitated care	51.5	49.7	58.1	64.5	64.9	66.6	70.8	93.6	86.9
Primary care case management	9.7	13.0	15.1	12.5	14.9	13.1	13.3	13.3	13.6
Personal support	10.1	10.6	11.8	11.6	10.8	10.7	11.0	1.7	1.7
Other care ⁴	21.6	21.4	21.9	21.5	21.3	20.6	19.9	47.6	47.9
Payments⁵									
Amount, in billions									
All payments	\$153.5	\$168.3	\$274.9	\$276.2	\$296.8	\$326.0	\$339.0	\$368.6	\$363.9
Percent distribution									
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Inpatient hospital	14.5	14.4	12.8	13.4	12.5	11.8	9.9	10.7	10.3
Mental health facility	1.1	1.1	0.8	0.9	0.8	0.8	0.7	0.7	0.6
Intermediate care facility for individuals with intellectual disabilities ³	6.1	5.6	4.3	4.3	4.2	3.9	3.7	3.6	3.3
Nursing facility	21.7	20.5	16.3	16.8	16.1	14.9	14.4	13.1	13.0
Physician	4.3	4.0	4.1	3.6	3.5	3.5	3.5	3.3	3.0
Dental	0.8	0.8	1.1	1.2	1.3	1.4	1.6	1.6	1.3
Other practitioner	0.3	0.4	0.4	0.3	0.3	0.3	0.3	0.5	0.5
Outpatient hospital	4.0	4.2	3.6	3.7	3.7	3.7	3.8	3.5	3.5
Clinic	3.8	3.7	3.2	3.1	3.1	3.1	3.2	3.4	3.2
Laboratory and radiological	0.8	0.8	1.1	1.1	1.0	1.0	1.0	1.0	0.8
Home health	1.9	1.9	2.0	2.3	2.2	2.2	2.1	2.0	1.8
Prescribed drugs	10.8	11.9	15.6	8.0	7.9	7.8	8.0	7.9	6.1
Capitated care	14.0	14.5	16.9	21.2	23.0	25.5	27.2	29.5	33.6
Primary care case management	0.3	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Personal support	6.9	6.9	7.5	8.4	8.3	8.0	7.7	3.3	3.3
Other care ⁴	8.6	8.8	10.2	11.6	12.0	11.9	12.7	15.9	15.6

See footnotes at end of table.

Table 110 (page 2 of 2). Medicaid and Children's Health Insurance Program beneficiaries and payments, by type of service: United States, selected fiscal years 1999–2012

Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#110>.

[Data are compiled by the Centers for Medicare & Medicaid Services from the Medicaid Data System]

Type of service	1999	2000	2005	2007	2008	2009	2010	2011 ¹	2012 ¹
Payments per beneficiary ⁵									
	Amount								
Total payment per beneficiary	\$3,819	\$3,936	\$4,768	\$4,862	\$5,051	\$5,209	\$5,160	\$5,159	\$5,082
Inpatient hospital	4,943	4,919	6,411	7,191	7,083	7,070	7,347	4,858	4,437
Mental health facility	18,094	17,800	19,252	21,407	21,975	21,404	20,782	14,557	13,221
Intermediate care facility for individuals with intellectual disabilities ³	76,443	79,330	107,028	113,735	123,053	127,837	125,851	129,806	119,903
Nursing facility	20,568	20,220	26,185	28,282	29,533	29,551	31,617	26,995	28,060
Physician	357	356	465	457	485	496	492	271	238
Dental	214	238	326	340	389	423	432	301	245
Other practitioner	118	139	200	170	171	171	190	186	183
Outpatient hospital	491	533	617	695	736	735	803	470	454
Clinic	860	805	749	741	772	792	791	712	680
Laboratory and radiological	114	113	183	185	188	198	205	119	102
Home health	3,571	3,135	4,487	5,334	5,789	6,628	6,375	4,286	3,565
Prescribed drugs	837	975	1,509	926	957	951	926	719	540
Capitated care	1,040	1,148	1,386	1,598	1,786	1,991	1,983	1,627	1,964
Primary care case management	119	30	27	33	32	41	49	45	47
Personal support	2,583	2,543	3,035	3,534	3,852	3,903	3,593	9,959	9,619
Other care ⁴	1,508	1,600	2,228	2,611	2,856	3,015	3,289	1,724	1,653

¹Starting with 2011, a new tabular methodology was used. Therefore, estimates may not be comparable to earlier data and caution should be used with trend analysis.

²Beneficiaries include those who received services through Medicaid or the Children's Health Insurance Program (CHIP). Separate CHIP beneficiaries are included for 2011 and 2012.

³This category was previously known as Intermediate care facility for the mentally retarded. This is a change in terminology only and not measurement.

⁴Estimates for 2010 and earlier include unknown services and payments with Other care.

⁵Medicaid payment data for 2010 and earlier exclude disproportionate share hospital (DSH) payments (\$14.7 billion in FY2010) and DSH mental health facility payments (\$2.9 billion in FY2010).

NOTES: Data are for fiscal year ending September 30. See Appendix II, Medicaid; Medicaid payments. Beneficiaries receiving more than one type of service are included in each category. For more information, see: <http://www.medicare.gov>. Colorado and Idaho had not reported 2012 data as of the date accessed. Starting with 2011, a new tabular methodology was used. Therefore, estimates may not be comparable to earlier data and caution should be used with trend analysis. For more information on data and analytic issues, see: <https://www.cms.gov/Research-Statistics-Data-and-Systems/Computer-Data-and-Systems/MedicaidDataSourcesGenInfo/MSIS-Tables.html>. Estimates for 2011 have been revised and differ from previous editions of *Health, United States*.

SOURCE: Centers for Medicare & Medicaid Services, Center for Medicaid and CHIP Services, Medicaid Statistical Information System (MSIS), granular file. MSIS data for 2011 and 2012 were accessed January 7, 2016. See Appendix I, Medicaid Statistical Information System (MSIS).

Table 111. Department of Veterans Affairs health care expenditures and use, and persons treated, by selected characteristics: United States, selected fiscal years 2005–2014

Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#111>.

[Data are compiled from patient records, enrollment information, and budgetary data by the Department of Veterans Affairs]

Type of expenditure and use	2005	2007	2008	2009	2010	2011	2012	2013	2014
Health care expenditures									
Amount, in millions									
All expenditures ¹	\$30,291	\$34,025	\$38,282	\$42,955	\$47,280	\$50,575	\$51,880	\$54,738	\$58,010
Percent distribution									
All services	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Inpatient hospital	24.3	24.0	23.5	22.7	21.4	20.6	20.1	19.8	19.8
Outpatient care	53.4	53.5	53.2	53.5	52.5	52.6	53.8	53.2	55.5
Nursing home care	8.4	8.3	8.1	7.8	7.4	7.2	7.3	7.0	7.0
All other ²	13.9	14.2	15.2	16.0	18.8	19.6	18.8	20.0	17.7
Health care use									
Number, in thousands									
Inpatient hospital discharges ³	614	607	622	640	656	653	646	632	619
Outpatient visits ⁴	57,169	62,234	66,484	73,969	79,457	83,146	87,370	90,226	93,852
Nursing home discharges ⁵	61	63	64	65	67	63	67	69	68
Inpatients ⁶									
Total	488	477	492	512	532	540	546	545	558
Percent distribution									
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veterans with service-connected disability	37.6	39.9	41.1	42.6	43.5	44.9	46.5	48.3	50.0
Veterans without service-connected disability	61.5	59.1	58.0	56.4	55.6	54.3	52.6	50.8	49.1
Low income	39.9	36.9	35.4	34.8	34.6	33.4	32.1	30.4	28.8
Veterans receiving aid and attendance or housebound benefits or who are catastrophically disabled	12.1	11.3	11.1	10.5	10.1	9.8	9.6	9.4	9.0
Veterans receiving medical care subject to copayments ⁷	8.6	9.8	10.0	9.5	9.3	9.3	9.2	9.3	9.6
Other and unknown ⁸	1.0	1.0	1.6	1.6	1.6	1.7	1.7	1.7	1.6
Nonveterans	0.9	0.9	0.9	1.0	0.9	0.9	0.9	0.9	0.9
Outpatients ⁶									
Number, in thousands									
Total	5,077	5,221	5,291	5,439	5,631	5,789	5,903	6,009	6,176
Percent distribution									
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Veterans with service-connected disability	31.6	33.8	34.7	37.1	38.6	39.8	41.7	44.0	45.8
Veterans without service-connected disability	62.7	60.8	59.7	57.2	56.4	55.1	53.3	51.1	49.0
Low income	31.8	28.9	27.2	25.9	25.7	24.9	24.0	22.6	21.3
Veterans receiving aid and attendance or housebound benefits or who are catastrophically disabled	3.5	3.5	3.5	3.4	3.4	3.3	3.2	3.2	3.1
Veterans receiving medical care subject to copayments ⁷	25.4	25.5	25.2	23.8	23.0	22.3	21.4	20.7	20.1
Other and unknown ⁸	2.0	3.0	3.8	4.0	4.3	4.6	4.6	4.6	4.5
Nonveterans	5.7	5.4	5.7	5.7	5.1	5.1	5.1	4.9	5.3

¹Health care expenditures exclude construction, medical administration, and miscellaneous operating expenses at Department of Veterans Affairs (VA) headquarters.

²Includes expenditures for miscellaneous benefits and services, contract hospitals, education and training, subsidies to state veterans hospitals, nursing homes and residential rehabilitation treatment programs (formerly domiciliaries), and the Civilian Health and Medical Program of the Department of Veterans Affairs.

³Discharges from medicine, surgery, psychiatry, rehabilitation medicine, spinal cord, and neurology units, and residential rehabilitation treatment programs (formerly, domiciliary care). Does not include long-term stays.

⁴Hospital outpatient care. Includes the following services: physicians, laboratory tests, home-based primary care, or outpatient fee-basis care.

⁵Includes VA-covered state nursing home veteran patients.

⁶Individuals receiving services. Individuals with multiple discharges or visits are only counted once in the inpatient or outpatient category. The inpatient and outpatient totals are not additive because most inpatients are also treated as outpatients.

⁷Includes veterans who receive medical care subject to copayments according to income level, based on financial means testing.

⁸Includes expenditures for services for veterans who were prisoners of war, exposed to Agent Orange, and other. Veterans reporting Agent Orange exposure but not treated for it were means tested and placed in the low income or other group depending on income.

NOTES: Some veterans have multiple sources of health coverage, including Medicare or private insurance. Estimates in this table relate only to health care use paid for by the Veteran's Administration. At the end of FY2014, the veteran population was estimated at 22.0 million, with 45% aged 65 and over. Of all living veterans, 5% had served during World War II, 9% during the Korean conflict, 33% during the Vietnam era, 32% during the Persian Gulf War (service from August 2, 1990 to present), and 25% during peacetime. Percentages sum to more than 100% because some veterans serve during more than one war. See Appendix I, Department of Veterans Affairs National Enrollment and Patient Databases. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: Department of Veterans Affairs (VA), Office of the Assistant Deputy Under Secretary for Health, National Patient Care Database, National Enrollment Database, budgetary data, and unpublished data. Veteran population estimates were provided by the VA's Office of the Actuary. See Appendix I, Department of Veterans Affairs National Enrollment and Patient Databases.

Table 112 (page 1 of 2). Medicare enrollees, enrollees in managed care, payment per fee-for-service enrollee, and short-stay hospital utilization, by state: United States, 1994 and 2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#112>.

[Data are compiled by the Centers for Medicare & Medicaid Services]

State	Enrollment, in thousands ¹		Percent of enrollees in managed care ²		Average payment per fee-for-service enrollee		Short-stay hospital utilization for fee-for-service enrollees			
	1994	2014	1994	2014	1994	2014	Discharges per 1,000 enrollees ³		Average length of stay, in days ³	
							1994	2014	1994	2014
United States ⁴	36,190	52,846	7.9	29.9	\$4,375	\$9,423	345	283	7.5	5.4
Alabama	633	947	0.8	24.0	4,454	8,433	413	312	7.0	5.5
Alaska	33	80	0.6	0.9	3,687	8,153	269	182	6.3	5.8
Arizona	578	1,096	24.8	37.6	4,442	8,731	292	233	5.9	4.8
Arkansas	416	584	0.2	19.4	3,719	8,272	366	288	7.0	5.3
California	3,582	5,476	30.0	39.3	5,219	9,908	366	240	6.1	5.3
Colorado	413	755	17.2	36.2	3,935	7,967	302	221	6.0	4.8
Connecticut	497	619	2.6	24.2	4,426	10,307	287	294	8.1	5.8
Delaware	99	175	0.2	7.9	4,712	9,606	326	288	8.1	5.5
District of Columbia	80	87	3.9	11.9	5,655	9,884	376	331	10.1	6.4
Florida	2,584	3,900	13.8	37.9	5,027	10,754	326	328	7.1	5.3
Georgia	819	1,469	0.4	28.9	4,402	8,818	378	277	6.9	5.4
Hawaii	146	237	29.8	46.4	3,069	5,990	301	160	9.1	6.3
Idaho	146	271	2.5	31.6	3,045	7,768	274	180	5.2	4.6
Illinois	1,605	2,026	5.5	18.4	4,324	9,817	374	317	7.3	5.2
Indiana	805	1,124	2.6	22.5	3,945	9,128	345	294	6.9	5.1
Iowa	470	561	3.1	14.4	3,080	7,970	322	232	6.6	5.2
Kansas	378	477	3.3	13.5	3,847	8,458	348	257	6.5	5.0
Kentucky	578	846	2.3	24.7	3,862	8,728	396	314	7.2	5.1
Louisiana	572	772	0.4	28.3	5,468	10,158	399	307	7.2	5.4
Maine	198	299	0.1	20.5	3,464	8,168	322	218	7.6	5.2
Maryland	596	905	1.4	8.8	4,997	10,684	362	301	7.5	5.4
Massachusetts	924	1,189	6.1	20.5	5,147	10,167	350	291	7.6	5.3
Michigan	1,331	1,853	0.7	30.5	4,307	9,998	328	335	7.6	5.2
Minnesota	625	888	19.6	51.8	3,394	11,783	334	384	5.7	4.8
Mississippi	391	549	0.1	13.5	4,189	9,482	423	315	7.4	5.7
Missouri	821	1,113	3.4	26.5	4,191	8,776	349	305	7.3	5.1
Montana	128	195	0.4	17.3	3,114	6,781	306	184	5.9	4.9
Nebraska	247	306	2.2	12.1	2,926	8,499	281	229	6.3	5.1
Nevada	187	435	19.0	32.3	4,306	9,473	291	245	7.0	5.8
New Hampshire	152	259	0.2	7.2	3,414	8,166	281	217	7.6	5.4
New Jersey	1,158	1,462	2.6	15.5	4,531	10,499	354	297	10.2	5.8
New Mexico	205	361	13.6	30.8	3,110	7,624	301	211	6.0	4.9
New York	2,601	3,278	6.2	35.9	4,855	10,309	334	293	11.2	6.7
North Carolina	1,001	1,719	0.5	28.5	3,465	8,672	314	279	8.0	5.3
North Dakota	101	117	0.6	15.0	3,218	8,662	327	239	6.3	5.2
Ohio	1,649	2,108	2.4	38.8	3,982	9,742	350	323	7.1	5.0
Oklahoma	481	665	2.5	16.5	4,098	8,793	355	290	7.0	5.3
Oregon	469	728	27.7	43.2	3,285	7,505	305	190	5.2	4.7
Pennsylvania	2,053	2,489	3.3	39.6	5,212	9,353	379	306	8.0	5.4
Rhode Island	166	200	7.0	35.8	4,148	8,816	312	291	8.1	5.4
South Carolina	497	910	0.1	22.2	3,777	8,750	319	267	8.3	5.5
South Dakota	114	152	0.1	16.6	2,952	8,447	356	248	6.1	4.9
Tennessee	754	1,206	0.3	32.3	4,441	8,686	375	302	7.1	5.3
Texas	2,029	3,516	4.1	29.2	4,703	10,340	333	288	7.2	5.4
Utah	182	333	9.4	33.4	3,443	8,080	238	207	5.4	4.3
Vermont	82	128	0.1	7.3	3,182	7,858	283	172	7.6	5.6
Virginia	803	1,311	1.5	16.7	3,748	8,251	348	275	7.3	5.2
Washington	676	1,146	12.5	29.7	3,401	7,632	269	209	5.3	4.9
West Virginia	326	411	8.3	26.3	3,798	8,536	420	306	7.1	5.4
Wisconsin	752	1,023	2.0	35.4	3,246	8,437	310	255	6.8	4.9
Wyoming	58	92	3.3	3.9	3,537	7,676	315	202	5.6	4.8

See footnotes at end of table.

Table 112 (page 2 of 2). Medicare enrollees, enrollees in managed care, payment per fee-for-service enrollee, and short-stay hospital utilization, by state: United States, 1994 and 2014

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#112>.

[Data are compiled by the Centers for Medicare & Medicaid Services]

¹Total persons enrolled in the hospital insurance (Part A) program, supplementary medical insurance (Part B) program, or both, as of July 1. Includes fee-for-service and managed care enrollees.

²See Appendix II, Managed care.

³Data are for fee-for-service enrollees only.

⁴Includes residents of the 50 states and the District of Columbia.

NOTES: In 1994, 92% of Medicare enrollees were in fee-for-service; in 2014, 70% of enrollees were in fee-for-service. See Appendix II, Medicare; Fee-for-service health insurance. Prior to 2004, enrollment and percentage of enrollees in managed care were based on a 5% annual Denominator File derived from the Centers for Medicare & Medicaid Services' (CMS) Enrollment Database. Starting with 2004 data, the enrollee counts were pulled from the 100% Denominator File. Payments per fee-for-service enrollee are based on fee-for-service billing reimbursement for a 5% sample of Medicare beneficiaries as recorded in CMS' National Claims History File. Prior to 2011, short-stay hospital utilization is based on the Medicare Provider Analysis and Review (MedPAR) stay records for a 20% sample of Medicare beneficiaries. Beginning in 2011, short-stay hospital utilization is based on the MedPAR stay records for 100% of Medicare beneficiaries. Estimates may not sum to totals because of rounding. State based on residence of the beneficiary. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: Centers for Medicare & Medicaid Services; Office of Research, Development, and Information. Health Care Financing Review: Medicare and Medicaid Statistical Supplements for publication years 1996 to 2010; Center for Strategic Planning. Medicare & Medicaid Research Review: Medicare and Medicaid Statistical Supplement for publication year 2011; Office of Information Products and Data Analytics. Medicare and Medicaid Statistical Supplements for publication year 2012. Data for 2013 (shown in spreadsheet version) and 2014 are unpublished. See Appendix I, Medicare Administrative Data.

Table 113. Medicaid and Children's Health Insurance Program beneficiaries and payments per beneficiary, by state: United States, selected fiscal years 2000–2012

Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hsu/contents2015.htm#113>.

[Data are compiled by the Centers for Medicare & Medicaid Services from the Medicaid Data System]

State	Beneficiaries, in thousands ¹				Payments per beneficiary ²			
	2000	2010	2011 ³	2012 ³	2000	2010	2011 ³	2012 ³
United States	42,763	65,700	71,449	71,604	\$3,936	\$5,160	\$5,159	\$5,082
Alabama	619	931	945	964	3,860	4,342	4,427	4,261
Alaska	96	127	137	138	4,876	9,520	9,578	9,638
Arizona	681	1,805	2,005	1,698	3,100	5,270	4,707	4,843
Arkansas	489	773	803	810	3,086	4,916	4,548	4,425
California	7,915	11,212	11,798	10,987	2,155	3,094	3,184	3,232
Colorado	381	682	757	---	4,747	4,840	4,597	---
Connecticut	420	664	749	774	6,762	8,120	7,796	7,603
Delaware	115	210	234	246	4,584	6,380	6,298	6,383
District of Columbia	139	211	239	250	5,715	8,577	8,895	6,509
Florida	2,360	3,656	3,949	4,052	3,114	4,412	4,893	4,657
Georgia	1,290	1,875	2,194	2,296	2,774	3,717	3,817	3,963
Hawaii	204	288	328	335	2,626	4,692	4,475	4,466
Idaho	131	430	263	---	4,530	2,871	5,493	---
Illinois	1,516	2,758	2,940	3,089	5,150	4,222	4,401	4,349
Indiana	705	1,177	1,278	1,342	4,224	4,889	4,509	4,882
Iowa	314	508	548	562	4,707	5,920	5,970	6,093
Kansas	263	364	481	435	4,670	6,309	5,335	5,883
Kentucky	771	959	1,124	1,108	3,780	5,532	5,014	5,114
Louisiana	761	1,237	1,327	1,343	3,456	4,439	4,149	4,318
Maine	192	330	339	370	6,820	4,451	4,669	5,248
Maryland	665	940	1,073	1,127	5,396	7,273	6,694	6,611
Massachusetts	1,047	1,637	1,733	1,542	5,153	6,760	6,537	7,009
Michigan	1,352	2,219	2,390	2,355	3,611	5,127	4,867	4,996
Minnesota	559	851	1,036	1,106	5,857	8,390	7,673	7,825
Mississippi	605	801	834	897	2,987	4,197	4,398	4,160
Missouri	890	1,141	1,207	1,206	3,673	5,429	5,212	5,466
Montana	104	126	137	141	4,173	6,023	5,818	5,856
Nebraska	229	269	304	311	4,185	5,890	5,306	5,441
Nevada	138	334	376	390	3,733	3,899	3,716	3,526
New Hampshire	97	148	155	165	6,712	6,805	6,676	6,387
New Jersey	822	1,229	1,410	1,596	5,724	6,963	6,306	5,896
New Mexico	376	557	636	633	3,325	4,971	4,058	4,009
New York	3,420	5,011	5,758	6,076	7,646	8,526	8,893	7,954
North Carolina	1,209	1,876	1,911	2,071	3,996	5,111	5,033	4,829
North Dakota	61	83	89	92	5,852	8,261	8,222	8,474
Ohio	1,305	2,319	2,537	2,574	5,434	6,231	6,237	6,289
Oklahoma	507	853	972	997	3,163	4,355	3,891	3,894
Oregon	542	644	771	787	3,135	4,948	4,642	4,776
Pennsylvania	1,492	2,326	2,544	2,578	4,266	6,834	6,958	6,901
Rhode Island	179	214	229	234	5,982	7,367	7,005	6,769
South Carolina	685	953	983	1,005	3,900	5,339	5,241	4,739
South Dakota	102	142	135	136	3,935	5,479	5,670	5,626
Tennessee	1,568	1,532	1,548	1,683	2,226	5,914	7,240	7,305
Texas	2,603	4,745	5,266	6,021	3,487	4,367	4,257	3,673
Utah	224	369	448	443	4,277	5,404	4,805	5,344
Vermont	139	181	188	189	3,451	5,525	5,592	5,690
Virginia	627	969	1,057	1,077	3,960	6,045	5,781	5,605
Washington	895	1,330	1,431	1,487	2,717	4,744	4,382	4,208
West Virginia	335	397	420	411	4,154	6,774	6,965	7,426
Wisconsin	577	1,230	1,356	1,398	5,039	4,393	4,185	4,070
Wyoming	46	76	78	78	4,609	7,540	7,380	7,425

--- Data not available.

¹Beneficiaries include those who received services through Medicaid or the Children's Health Insurance Program (CHIP). Separate CHIP beneficiaries are included for 2011 and 2012.

²Medicaid payment data for 2010 and earlier exclude disproportionate share hospital (DSH) payments (\$14.7 billion in FY2010) and DSH mental health facility payments (\$2.9 billion in FY2010).

³Starting with 2011, a new tabular methodology was used. Therefore, estimates may not be comparable to earlier data and caution should be used with trend analysis.

NOTES: See Appendix II, Children's Health Insurance Program (CHIP); Medicaid; Medicaid payments. Colorado and Idaho had not reported 2012 data as of the date accessed. Starting with 2011, a new tabular methodology was used. Therefore, estimates may not be comparable to earlier data and caution should be used with trend analysis. For more information on data and analytic issues, see: <https://www.cms.gov/Research-Statistics-Data-and-Systems/Computer-Data-and-Systems/MedicaidDataSourcesGenInfo/MSIS-Tables.html>. Estimates for 2011 have been revised and differ from previous editions of *Health, United States*.

SOURCE: Centers for Medicare & Medicaid Services, Center for Medicaid and CHIP Services, Medicaid Statistical Information System (MSIS), granular file. MSIS data for 2011 and 2012 were accessed January 7, 2016. See Appendix I, Medicaid Statistical Information System (MSIS).

Table 114 (page 1 of 3). Persons under age 65 without health insurance coverage, by age, state, and territory: United States and Puerto Rico, 2009–2014

Updated data when available, Excel, PDF, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#114>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population in the United States and Puerto Rico]

Age, state, and territory	2009	2010	2011	2012	2013	2014
Under 65 years						
				Percent		
United States ¹	17.2	17.6	17.2	16.9	16.7	13.4
Alabama	16.0	16.9	16.3	15.4	15.9	14.0
Alaska	22.0	19.4	21.7	22.0	20.3	19.2
Arizona	19.8	19.5	19.9	20.3	20.3	16.0
Arkansas	19.3	20.2	19.7	19.0	18.8	13.8
California	20.1	20.7	20.3	20.0	19.3	14.0
Colorado	17.4	17.6	17.0	16.3	15.7	11.9
Connecticut	10.1	10.3	9.8	10.6	10.8	8.0
Delaware	11.9	11.5	10.2	9.9	11.7	8.7
District of Columbia	7.8	8.4	8.3	6.1	7.1	6.1
Florida	24.8	25.4	25.0	24.0	24.2	20.1
Georgia	21.0	21.9	21.8	20.8	21.1	17.8
Hawaii	7.9	8.7	8.4	7.7	8.3	5.7
Idaho	19.1	20.3	17.8	18.3	18.5	15.4
Illinois	14.9	15.7	14.5	14.6	14.4	11.2
Indiana	15.8	17.0	16.6	16.6	16.2	13.8
Iowa	10.1	10.8	10.4	9.7	10.2	6.7
Kansas	14.5	15.6	14.4	14.6	14.3	12.2
Kentucky	16.5	17.5	16.6	15.8	16.8	10.2
Louisiana	19.5	20.1	19.9	19.0	19.2	16.6
Maine	11.9	12.5	13.1	12.4	13.3	11.8
Maryland	12.5	12.7	11.5	11.5	11.4	9.0
Massachusetts	4.9	4.9	4.8	4.5	4.4	3.9
Michigan	14.1	14.3	13.6	13.4	12.8	9.8
Minnesota	10.2	10.1	9.9	9.4	9.5	6.8
Mississippi	20.0	20.7	20.2	19.7	19.6	16.8
Missouri	15.3	15.2	15.7	16.1	15.2	13.5
Montana	21.9	19.6	21.8	21.4	19.7	16.2
Nebraska	13.1	13.4	13.6	12.7	12.2	10.8
Nevada	24.6	25.4	24.7	25.0	23.4	17.5
New Hampshire	12.0	12.7	11.4	12.8	12.6	10.9
New Jersey	14.2	15.0	14.7	14.5	15.3	12.6
New Mexico	23.1	22.8	22.8	21.6	22.2	17.3
New York	13.0	13.5	13.0	12.5	12.4	9.9
North Carolina	18.2	19.2	18.7	18.9	18.0	15.2
North Dakota	11.4	11.5	11.5	12.4	11.9	9.2
Ohio	14.0	14.1	13.9	13.4	12.8	9.7
Oklahoma	21.3	22.0	21.3	21.1	20.4	17.7
Oregon	19.8	19.7	18.0	16.9	17.4	11.5
Pennsylvania	11.3	11.9	11.8	11.3	11.3	10.1
Rhode Island	12.8	13.7	12.6	13.3	13.9	8.3
South Carolina	19.2	20.2	19.4	19.3	18.4	16.0
South Dakota	15.6	13.6	13.4	12.3	14.3	11.6
Tennessee	16.3	16.4	17.0	16.1	16.2	13.9
Texas	26.3	26.3	25.5	24.8	24.5	21.2
Utah	15.7	17.0	16.5	15.3	14.7	13.6
Vermont	9.9	9.0	8.3	7.7	8.1	5.4
Virginia	13.3	14.5	14.1	14.2	14.1	12.4
Washington	15.3	16.1	16.0	15.7	16.1	10.5
West Virginia	16.7	17.2	18.2	16.9	16.1	10.7
Wisconsin	10.4	10.9	10.6	10.6	10.4	8.7
Wyoming	17.5	16.6	17.5	18.3	14.6	14.1
Puerto Rico ²	9.4	9.4	8.9	8.5	7.7	7.2

See footnotes at end of table.

Table 114 (page 2 of 3). Persons under age 65 without health insurance coverage, by age, state, and territory: United States and Puerto Rico, 2009–2014

Updated data when available, Excel, PDF, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#114>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population in the United States and Puerto Rico]

Age, state, and territory	2009	2010	2011	2012	2013	2014
Under 18 years	Percent					
United States ¹	8.5	8.0	7.5	7.1	7.1	6.0
Alabama	6.1	5.9	5.2	4.0	4.5	3.7
Alaska	12.8	9.3	13.9	13.3	11.7	12.3
Arizona	12.1	13.0	12.8	12.8	12.1	10.0
Arkansas	6.2	6.3	5.5	5.4	5.7	4.5
California	9.4	9.0	8.0	8.0	7.3	5.4
Colorado	9.8	9.8	9.3	8.1	8.4	6.0
Connecticut	3.9	2.9	2.5	3.7	4.1	3.9
Delaware	5.7	5.6	3.5	3.6	5.1	*5.0
District of Columbia	*	*2.0	*4.1	*	*2.2	*2.6
Florida	14.7	12.7	11.9	10.8	11.0	9.2
Georgia	10.7	9.8	9.5	8.9	9.5	7.5
Hawaii	2.4	3.7	3.9	2.9	3.2	2.0
Idaho	10.7	10.6	8.5	7.6	8.3	7.2
Illinois	4.4	4.8	3.4	3.2	4.3	3.8
Indiana	8.5	8.9	8.4	8.0	8.4	7.2
Iowa	4.4	4.3	4.6	4.3	4.8	3.2
Kansas	8.1	7.7	6.1	6.9	6.6	6.1
Kentucky	5.8	5.8	5.9	5.9	5.9	4.3
Louisiana	6.4	5.6	5.7	5.3	5.6	5.0
Maine	5.7	3.8	5.5	4.2	5.1	6.1
Maryland	4.7	4.9	4.5	3.8	4.3	3.4
Massachusetts	1.7	1.4	1.6	1.3	1.5	1.6
Michigan	4.3	4.2	3.9	4.2	4.2	3.6
Minnesota	7.0	6.3	6.1	5.7	6.1	3.5
Mississippi	10.1	8.2	7.4	7.2	7.3	5.4
Missouri	7.2	6.3	6.7	7.2	7.3	6.8
Montana	13.4	12.7	12.8	10.9	10.4	8.6
Nebraska	6.3	5.2	7.3	5.4	5.9	5.0
Nevada	18.0	17.9	16.1	16.5	13.9	9.7
New Hampshire	4.6	4.9	3.2	4.2	3.5	5.2
New Jersey	6.2	6.0	5.2	5.1	5.7	4.5
New Mexico	12.0	9.9	9.1	8.1	9.0	7.6
New York	4.8	4.8	4.4	4.0	4.1	3.4
North Carolina	7.9	8.1	7.8	7.3	5.9	5.3
North Dakota	6.3	6.6	7.6	7.4	7.7	6.7
Ohio	6.4	5.9	6.1	5.4	5.1	4.9
Oklahoma	11.1	10.4	10.9	9.9	10.5	8.7
Oregon	10.8	8.8	7.0	5.6	6.3	4.3
Pennsylvania	5.0	5.2	5.4	5.1	5.0	5.4
Rhode Island	4.9	4.8	3.9	5.1	6.0	3.3
South Carolina	9.5	9.8	8.7	7.8	7.0	5.2
South Dakota	6.8	7.2	5.6	3.9	7.2	7.2
Tennessee	5.7	5.3	5.8	5.6	5.7	5.2
Texas	16.3	14.7	13.3	12.3	12.5	11.2
Utah	10.2	11.1	11.1	9.3	9.0	9.2
Vermont	*3.3	*2.7	*	*3.0	*	*
Virginia	6.7	6.4	5.8	5.6	5.7	5.9
Washington	7.0	6.4	6.1	5.5	6.3	4.4
West Virginia	5.4	4.7	5.0	3.9	4.0	3.0
Wisconsin	4.6	5.2	4.6	4.7	4.4	4.9
Wyoming	9.0	7.4	8.7	9.9	6.3	6.7
Puerto Rico ²	4.2	4.5	4.0	4.3	3.5	3.1

See footnotes at end of table.

Table 114 (page 3 of 3). Persons under age 65 without health insurance coverage, by age, state, and territory: United States and Puerto Rico, 2009–2014

Updated data when available, Excel, PDF, and standard errors: <http://www.cdc.gov/nchs/hus/contents2015.htm#114>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population in the United States and Puerto Rico]

Age, state, and territory	2009	2010	2011	2012	2013	2014
18–64 years	Percent					
United States ¹	20.6	21.4	21.0	20.6	20.3	16.2
Alabama	19.8	21.2	20.5	19.7	20.2	17.9
Alaska	25.8	23.6	24.8	25.5	23.9	22.0
Arizona	23.3	22.3	22.8	23.4	23.7	18.5
Arkansas	24.6	25.8	25.4	24.5	24.0	17.6
California	24.5	25.3	25.1	24.7	23.8	17.3
Colorado	20.3	20.5	19.9	19.4	18.5	14.0
Connecticut	12.4	13.0	12.4	13.1	13.1	9.4
Delaware	14.3	13.7	12.7	12.2	14.1	10.1
District of Columbia	9.2	9.9	9.3	7.1	8.3	7.0
Florida	28.6	29.9	29.5	28.6	28.8	23.8
Georgia	25.4	26.9	26.8	25.6	25.8	21.9
Hawaii	9.9	10.5	10.1	9.4	10.1	7.1
Idaho	22.8	24.7	22.0	23.1	23.1	19.2
Illinois	19.0	20.0	18.8	19.0	18.1	13.9
Indiana	18.8	20.3	19.8	20.0	19.2	16.4
Iowa	12.2	13.3	12.6	11.8	12.3	8.0
Kansas	17.1	18.9	17.9	17.8	17.4	14.8
Kentucky	20.6	22.0	20.7	19.5	20.8	12.4
Louisiana	24.8	25.8	25.6	24.4	24.5	21.2
Maine	14.0	15.4	15.5	15.1	15.8	13.7
Maryland	15.4	15.5	14.0	14.3	14.0	11.0
Massachusetts	5.9	6.1	5.8	5.5	5.3	4.6
Michigan	17.8	18.1	17.3	16.8	16.0	12.1
Minnesota	11.4	11.6	11.3	10.9	10.8	8.1
Mississippi	24.3	26.0	25.6	25.0	24.6	21.4
Missouri	18.4	18.7	19.2	19.5	18.2	16.0
Montana	25.0	22.1	25.0	25.1	23.0	18.9
Nebraska	15.9	16.8	16.2	15.7	14.8	13.1
Nevada	27.3	28.4	28.1	28.4	27.1	20.4
New Hampshire	14.4	15.3	14.1	15.6	15.6	12.6
New Jersey	17.2	18.3	18.2	17.9	18.7	15.5
New Mexico	27.8	28.2	28.5	27.2	27.6	21.1
New York	16.0	16.6	16.0	15.4	15.2	12.1
North Carolina	22.2	23.5	22.9	23.3	22.6	19.0
North Dakota	13.2	13.3	12.9	14.1	13.4	10.1
Ohio	16.9	17.2	16.8	16.4	15.6	11.5
Oklahoma	25.6	26.8	25.6	25.6	24.5	21.4
Oregon	23.1	23.6	21.9	20.9	21.2	14.0
Pennsylvania	13.6	14.2	14.0	13.4	13.5	11.7
Rhode Island	15.5	16.7	15.5	16.0	16.4	9.8
South Carolina	23.0	24.2	23.5	23.6	22.5	19.9
South Dakota	19.1	16.3	16.6	15.7	17.2	13.4
Tennessee	20.4	20.7	21.2	20.0	20.1	17.1
Texas	30.9	31.5	30.9	30.4	29.7	25.6
Utah	18.6	20.1	19.3	18.4	17.7	16.0
Vermont	12.0	11.1	10.3	9.2	9.6	6.8
Virginia	15.8	17.4	17.1	17.3	17.1	14.7
Washington	18.4	19.7	19.6	19.4	19.8	12.7
West Virginia	20.5	21.5	22.7	21.2	20.2	13.3
Wisconsin	12.5	13.1	12.8	12.8	12.7	10.1
Wyoming	20.7	20.2	20.7	21.5	17.8	16.9
Puerto Rico ²	11.5	11.3	10.9	10.0	9.2	8.7

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error of 20%–30%. Data not shown have an RSE greater than 30%.

¹Excludes data for Puerto Rico.

²Data for Puerto Rico are collected in the Puerto Rico Community Survey. Data are not collected for the other territories.

NOTES: Health insurance estimates are shown for the civilian noninstitutionalized population. Data for 2009 use Census 2000 population controls, and data for 2010 and beyond use Census 2010 population controls. Questions on health insurance coverage ask about current coverage as of the day of American Community Survey (ACS) interview. Persons were considered uninsured if they were not covered by private health insurance, Medicare, Medicaid, Medical Assistance, TRICARE or other military health care, veteran's coverage through the Veteran's Administration, or other government coverage. People with Indian Health Service coverage only are considered uninsured by ACS. Standard errors for selected years are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Standard errors were computed with replicate weights using 80 balanced repeated replicate weights (BRR) with a Fay-modified BRR adjustment factor of 0.5.

SOURCE: U.S. Census Bureau, American Community Survey, public-use microdata sample. See Appendix I, American Community Survey (ACS).

Appendix Contents

Appendix I. Data Sources	349		
Government Sources	350		
Abortion Surveillance System	350		
American Community Survey (ACS)	350		
Census of Fatal Occupational Injuries (CFOI)	351		
Current Population Survey (CPS)	352		
Department of Veterans Affairs National Enrollment and Patient Databases	353		
Employee Benefits Survey—See Appendix I, National Compensation Survey (NCS).			
Healthcare Cost and Utilization Project (HCUP), National (Nationwide) Inpatient Sample	353		
Medicaid Statistical Information System (MSIS)	354		
Medical Expenditure Panel Survey (MEPS)	355		
Medicare Administrative Data	355		
Medicare Current Beneficiary Survey (MCBS)	356		
Monitoring the Future (MTF) Study	357		
National Ambulatory Medical Care Survey (NAMCS)	358		
National Compensation Survey (NCS)	359		
National Health and Nutrition Examination Survey (NHANES)	360		
National Health Expenditure Accounts (NHEA)	361		
National Health Interview Survey (NHIS)	362		
National HIV Surveillance System	364		
National Hospital Ambulatory Medical Care Survey (NHAMCS)	364		
National Immunization Survey (NIS)	365		
National Income and Product Accounts (NIPA)	366		
National Medical Expenditure Survey (NMES)—See Appendix I, Medical Expenditure Panel Survey (MEPS).			
National Notifiable Diseases Surveillance System (NNDSS)	367		
National Survey of Family Growth (NSFG)	368		
National Survey on Drug Use & Health (NSDUH)	369		
National Vital Statistics System (NVSS)	370		
Birth File	371		
Fetal Death Data Set	371		
Mortality Multiple Cause-of-Death File	372		
Linked Birth/Infant Death Data Set	374		
Occupational Employment Statistics (OES)	374		
Population Census and Population Estimates	375		
Decennial Census	375		
Race Data on the 1990 Census	375		
Race Data on the 2000 Census	375		
Race Data on the 2010 Census	375		
Modified Decennial Census Files	376		
Postcensal Population Estimates	376		
Intercensal Population Estimates	376		
Bridged-race Population Estimates	376		
Quality Improvement Evaluation System (QIES)	377		
Sexually Transmitted Disease (STD) Surveillance	378		
Surveillance, Epidemiology, and End Results Program (SEER)	378		
Youth Risk Behavior Survey (YRBS)	379		
Private and Global Sources	380		
American Association of Colleges of Osteopathic Medicine (AACOM)	380		
American Association of Colleges of Pharmacy (AACP)	380		
American Association of Colleges of Podiatric Medicine (AACPM)	380		
American Dental Association (ADA)	380		
American Hospital Association (AHA) Annual Survey of Hospitals	381		
American Medical Association (AMA) Physician Masterfile	381		
American Osteopathic Association (AOA)	381		
Association of American Medical Colleges (AAMC)	381		
Association of Schools and Colleges of Optometry (ASCO)	382		
Association of Schools & Programs of Public Health (ASPPH)	382		
Guttmacher Institute Abortion Provider Census	382		
Organisation for Economic Co-operation and Development (OECD) Health Data	382		
Appendix II. Definitions and Methods	384		
Acquired immunodeficiency syndrome (AIDS)	384		
Active physician—See Appendix II, Physician.			
Activities of daily living (ADL)	384		
Admission	384		
Age	384		
Age adjustment	385		
AIDS—See Appendix II, Acquired immunodeficiency syndrome (AIDS).			
Alcohol consumption	387		
Any-listed diagnosis—See Appendix II, Diagnosis.			
Average annual rate of change (percent change)	387		
Average length of stay	387		
Basic actions difficulty	387		
Bed, health facility	388		
Binge drinking	388		
Birth cohort	388		
Birth rate—See Appendix II, Rate: Birth and related rates.			
Birthweight	388		
Blood pressure, high	388		

Body mass index (BMI)	389	Health maintenance organization (HMO)	405
Cause of death	389	Health services and supplies expenditures—See	
Cause-of-death ranking	390	Appendix II, Health expenditures, national.	
Children's Health Insurance Program (CHIP)	390	Health status, respondent-assessed	405
Cholesterol	390	Hearing trouble	405
Cigarette smoking	393	Hispanic origin	405
Civilian noninstitutionalized population; Civilian		HIV—See Appendix II, Human immunodeficiency	
population—See Appendix II, Population.		virus (HIV) disease.	
Colorectal tests or procedures.	393	Home visit.	408
Community hospital—See Appendix II, Hospital.		Hospital.	408
Comparability ratio	394	Hospital-based physician—See Appendix II, Physician.	
Compensation—See Appendix II, Employer costs		Hospital day—See Appendix II, Days of care.	
for employee compensation.		Hospital utilization	409
Complex activity limitation	395	Human immunodeficiency virus (HIV) disease.	409
Consumer Price Index (CPI)	396	Hypercholesterolemia—See Appendix II, Cholesterol.	
Contraception	396	Hypertension—See Appendix II, Blood pressure, high.	
Cost-charge ratio	396	ICD; ICD codes—See Appendix II, Cause of death;	
Critical access hospital—See Appendix II, Hospital.		<i>International Classification of Diseases (ICD)</i> .	
Crude birth rate; Crude death rate—See Appendix II,		Illicit drug use.	410
Rate: Birth and related rates; Rate: Death and		Immunization—See Appendix II, Vaccination.	
related rates.		Incidence	411
Days of care	396	Income—See Appendix II, Family income.	
Death rate—See Appendix II, Rate: Death and		Individual practice association (IPA)—See Appendix II,	
related rates.		Health maintenance organization (HMO).	
Dental caries.	396	Industry of employment	411
Dental visit	397	Infant death	411
Diabetes	397	Injury.	411
Diagnosis	397	Injury-related visit	412
Diagnostic and other nonsurgical procedure—See		Inpatient	412
Appendix II, Procedure.		Inpatient care—See Appendix II, Hospital utilization.	
Discharge	398	Inpatient day—See Appendix II, Days of care.	
Domiciliary care home—See Appendix II, Long-term		Instrumental activities of daily living (IADL)	412
care facility; Nursing home.		Insurance—See Appendix II, Health insurance	
Drug	398	coverage.	
Drug abuse—See Appendix II, Illicit drug use.		Intermediate care facility—See Appendix II, Nursing	
Education	398	home.	
Emergency department.	399	<i>International Classification of Diseases (ICD)</i>	412
Emergency department or emergency room		<i>International Classification of Diseases, 9th Revision,</i>	
visit	399	<i>Clinical Modification (ICD-9-CM)</i>	413
Employer costs for employee compensation	399	<i>International Classification of Diseases, 10th Revision,</i>	
Ethnicity—See Appendix II, Hispanic origin.		<i>Clinical Modification/Procedure Coding System</i>	
Exercise—See Appendix II, Physical activity, leisure-		<i>(ICD-10-CM/PCS)</i>	413
time.		Late fetal death rate—See Appendix II, Rate: Death	
Expenditures—See Appendix II, Health expenditures,		and related rates.	
national.		Leading causes of death—See Appendix II, Cause-of-	
External cause of injury	399	death ranking.	
Family income	399	Length of stay—See Appendix II, Average length of	
Federal hospital—See Appendix II, Hospital.		stay.	
Fee-for-service health insurance	401	Life expectancy	413
Fertility rate—See Appendix II, Rate: Birth and		Limitation of activity	414
related rates.		Long-term care facility.	414
General hospital—See Appendix II, Hospital.		Low birthweight—See Appendix II, Birthweight.	
Geographic region	401	Mammography.	414
Gestation	401	Managed care.	415
Gross domestic product (GDP)	401	Marital status	416
Health care contact	401	Maternal age—See Appendix II, Age.	
Health expenditures, national	402	Medicaid	417
Health insurance coverage	403	Medicaid payments	419

Medical specialty—See Appendix II, Physician specialty.		Registered hospital—See Appendix II, Hospital.	
Medicare	419	Registration area	432
Metropolitan statistical area (MSA)	420	Relative standard error (RSE)	432
Micropolitan statistical area.	420	Relative survival rate	432
Multum Lexicon Plus therapeutic class.	420	Reporting area	432
Neonatal mortality rate—See Appendix II, Rate: Death and related rates.		Resident, health facility	432
Nonprofit hospital—See Appendix II, Hospital.		Resident population—See Appendix II, Population.	
North American Industry Classification System (NAICS)—See Appendix II, Industry of employment.		Rural—See Appendix II, Urbanization.	
Notifiable disease.	421	Self-assessment of health—See Appendix II, Health status, respondent-assessed.	
Nursing home.	421	Serious psychological distress.	432
Nursing home expenditures—See Appendix II, Health expenditures, national.		Short-stay hospital—See Appendix II, Hospital.	
Obesity—See Appendix II, Body mass index (BMI).		Skilled nursing facility—See Appendix II, Nursing home.	
Occupancy rate	421	Smoker—See Appendix II, Cigarette smoking.	
Office-based physician—See Appendix II, Physician.		Special hospital—See Appendix II, Hospital.	
Office visit.	421	Substance use	433
Operation—See Appendix II, Procedure.		Suicidal ideation.	433
Outpatient department	422	Surgery—See Appendix II, Outpatient surgery; Procedure.	
Outpatient surgery.	422	Surgical specialty—See Appendix II, Physician specialty.	
Outpatient visit.	422	Tobacco use—See Appendix II, Cigarette smoking.	
Overweight—See Appendix II, Body mass index (BMI).		Uninsured	433
Pap smear	422	Urbanization.	434
Patient—See Appendix II, Inpatient; Office visit; Outpatient visit.		Usual source of care	434
Percent change/percentage change—See Appendix II, Average annual rate of change (percent change).		Vaccination	434
Perinatal mortality rate; ratio—See Appendix II, Rate: Death and related rates.		Wages and salaries—See Appendix II, Employer costs for employee compensation.	
Personal care home with or without nursing—See Appendix II, Nursing home.		Years of potential life lost (YPLL)	435
Personal health care expenditures—See Appendix II, Health expenditures, national.			
Physical activity, leisure-time.	423		
Physician.	423		
Physician specialty	424		
Population	424		
Postneonatal mortality rate—See Appendix II, Rate: Death and related rates.			
Poverty	425		
Preferred provider organization (PPO)	425		
Prevalence	425		
Primary care specialty—See Appendix II, Physician specialty.			
Private expenditures—See Appendix II, Health expenditures, national.			
Procedure	425		
Proprietary hospital—See Appendix II, Hospital.			
Public expenditures—See Appendix II, Health expenditures, national.			
Purchasing power parities (PPPs)	426		
Race	426		
Rate.	431		
Region—See Appendix II, Geographic region.			

Appendix II: Tables

Table I. United States projected year 2000 standard population and age groups used to age-adjust data.	385
Table II. United States projected year 2000 standard population and proportion distribution, by age, for age-adjusting death rates prior to 2001	386
Table III. Revision of the <i>International Classification of Diseases</i> (ICD), by year of conference in which adopted and years in use in the United States	390
Table IV. Cause-of-death codes, by applicable revision of the <i>International Classification of Diseases</i> (ICD)	391
Table V. Comparability of selected causes of death between the 9th and 10th revisions of the <i>International Classification of Diseases</i> (ICD)	395
Table VI. Imputed family income percentages in the National Health Interview Survey, by selected characteristics: United States, 1990–2014	402
Table VII. Percentage of persons under age 65 with Medicaid or who are uninsured, by selected demographic characteristics, using Method 1 and Method 2 estimation procedures: United States, 2004	406

Table VIII. Codes for industries, based on the North American Industry Classification System (NAICS)	412
Table IX. Codes for external causes of injury, from the <i>International Classification of Diseases, 9th Revision, Clinical Modification</i>	413
Table X. Codes for procedure categories for Healthcare Cost and Utilization Project data, from the <i>International Classification of Diseases, 9th Revision, Clinical Modification</i>	415
Table XI. Current cigarette smoking among adults aged 18 and over, by race and Hispanic origin under the 1997 and 1977 Standards for federal data on race and ethnicity: United States, average annual, 1993–1995	428
Table XII. Private health care coverage among persons under age 65, by race and Hispanic origin under the 1997 and 1977 Standards for federal data on race and ethnicity: United States, average annual, 1993–1995.	429
Appendix II: Figure	
Figure I. U.S. Census Bureau: Four geographic regions and nine divisions of the United States	403

Appendix I. Data Sources

Health, United States consolidates the most current data on the health of the population of the United States, the availability and use of health care resources, and health care expenditures. Information was obtained from the data files and published reports of many federal government, private, and global agencies and organizations. In each case, the sponsoring agency or organization collected data using its own methods and procedures. Therefore, data in this report may vary considerably with respect to source, method of collection, definitions, and reference period.

Although a detailed description and comprehensive evaluation of each data source are beyond the scope of this appendix, readers should be aware of the general strengths and weaknesses of the different data collection systems shown in *Health, United States*. For example, population-based surveys are able to collect socioeconomic data and information on the impact of an illness, such as limitation of activity. These data are limited by the amount of information a respondent remembers or is willing to report. For example, a respondent may not know detailed medical information, such as a precise diagnosis or the type of medical procedure performed, and therefore cannot report that information. In contrast, records-based surveys, which collect data from physician and hospital records, usually contain good diagnostic information but little or no information about the socioeconomic characteristics of individuals or the impact of illnesses on individuals.

Different data collection systems may cover different populations, and understanding these differences is critical to interpreting the resulting data. Data on vital statistics and national expenditures cover the entire population. However, most data on morbidity cover only the civilian noninstitutionalized population and thus may not include data for military personnel, who are usually young; for institutionalized people, including the prison population, who may be of any age; or for nursing home residents, who are usually older.

All data collection systems are subject to error, and records may be incomplete or contain inaccurate information. Respondents may not remember essential information, a question may not mean the same thing to different respondents, and some institutions or individuals may not respond at all. It is not always possible to measure the magnitude of these errors or their effect on the data. Where possible, table notes describe the universe and method of data collection, to assist users in evaluating data quality.

Some information is collected in more than one survey, and estimates of the same statistic may vary among surveys because of different survey methodologies, sampling frames, questionnaires, definitions, and tabulation

categories. For example, cigarette use is measured by the National Health Interview Survey, the National Survey on Drug Use & Health, the Monitoring the Future Study, and the Youth Risk Behavior Survey. These surveys use slightly different questions, cover persons of differing ages, and interview in diverse settings (e.g., at school compared with at home), so estimates may differ.

Overall estimates generally have relatively small sampling errors, but estimates for certain population subgroups may be based on a small sample size and have relatively large sampling errors. Numbers of births and deaths from the National Vital Statistics System represent complete counts (except for births in those states where data are based on a 50% sample for certain years). Therefore, these data are not subject to sampling error. However, when the figures are used for analytical purposes, such as the comparison of rates over a period, the number of events that actually occurred may be considered as one of a large series of possible results that could have arisen under the same circumstances. When the number of events is small and the probability of such an event is rare, estimates may be unstable, and considerable caution must be used in interpreting the statistics. Estimates that are unreliable because of large sampling errors or small numbers of events are noted with asterisks in tables, and the criteria used to determine unreliable estimates are indicated in an accompanying footnote.

In this appendix, government data sources are listed alphabetically by data set name, and private and global sources are listed separately. To the extent possible, government data systems are described using a standard format. The *Overview* is a brief, general statement about the purpose or objectives of the data system. The *Coverage* section describes the population or events that the data system covers: for example, residents of the United States, the noninstitutionalized population, persons in specific population groups, or other entities that are included in the survey or data system. The *Methodology* section presents a short description of the methods used to collect the data. The *Sample Size and Response Rate* section provides these statistics for surveys. The *Issues Affecting Interpretation* section describes major changes in the data collection methodology or other factors that must be considered when analyzing trends shown in *Health, United States*: for example, a major survey redesign that may introduce a discontinuity in the trend. For additional information about the methodology, data files, and history of a data source, consult the *References* and *For More Information* sections that follow each summary.

Government Sources

Abortion Surveillance System

CDC/National Center for Chronic Disease Prevention and Health Promotion (NCCDPHP)

Overview. The Abortion Surveillance System documents the number and characteristics of women obtaining legal induced abortions in the United States.

Coverage. The system includes women of all ages, including adolescents, who obtain legal induced abortions.

Methodology. Each year, CDC requests tabulated data from the central health agencies of 52 reporting areas (the 50 states, the District of Columbia [D.C.], and New York City) to document the number and characteristics of women obtaining abortions in the United States. For the purpose of surveillance, a legal induced abortion is defined as an intervention performed by a licensed clinician (e.g., a physician, nurse-midwife, nurse practitioner, or physician assistant) that is intended to terminate a suspected or known ongoing intrauterine pregnancy and produce a nonviable fetus at any gestational age.

In most states, collection of abortion data is facilitated by the legal requirement for hospitals, facilities, and physicians to report abortions to a central health agency. These central health agencies voluntarily report abortion data to CDC and provide only the aggregate numbers for the abortion data they have collected through their independent surveillance systems. Although reporting to CDC is voluntary, most reporting areas provide aggregate abortion numbers; during 2003–2012, a total of 47 reporting areas provided CDC a continuous annual record of abortion numbers.

Issues Affecting Interpretation. Because reporting areas establish their own reporting requirements for abortion and send their data to CDC voluntarily, CDC is unable to obtain the total number of abortions performed in the United States. Although most states legally require medical providers to submit a report for all the abortions they perform, enforcement of this requirement varies. Additionally, although most reporting areas collect and send abortion data to CDC, during 2003–2012, 5 of the 52 reporting areas did not provide CDC with data on a consistent annual basis (the five states that did not report continuously for the period 2003–2012 were California, Louisiana, Maryland, New Hampshire, and West Virginia). Because of these limitations, during the period covered by this report the total annual number of abortions recorded by CDC was consistently approximately 70% of the number recorded by the Guttmacher Institute, which uses numerous active follow-up techniques to increase the completeness of the data obtained through its periodic national census of abortion providers.

Reference

Pazol K, Creanga AA, Jamieson DJ. Abortion surveillance—United States, 2012. *MMWR Surveill Summ* 2015;64(SS–10):1–40. Available from: http://www.cdc.gov/mmwr/preview/mmwrhtml/ss6410a1.htm?s_cid=ss6410a1_e.

For More Information. See the NCCDPHP surveillance and research website at: http://www.cdc.gov/reproductivehealth/Data_Stats/index.htm.

American Community Survey (ACS)

U.S. Census Bureau

Overview. ACS provides annual estimates of income, education, employment, health insurance coverage, and housing costs and conditions for U.S. residents. Estimates from ACS complement population data collected by the U.S. Census Bureau during the decennial census. Topics currently included on an annual basis in ACS were previously collected once a decade through the decennial census long form.

Coverage. ACS covers U.S. residents residing in all 3,141 counties in the 50 states and D.C., and all 78 municipalities in Puerto Rico. ACS began data collection for U.S. residents residing in housing units in January 2005 and for residents residing in group quarters facilities in January 2006. Annual ACS estimates are available every year for states and for specific geographic areas with populations of 65,000 or more.

Methodology. Starting with 2013 data, the ACS data collection operation uses up to four modes to collect information: Internet, mail, telephone, and personal visit interviews. The first mode includes a mailed request to respond to the ACS questionnaire over the Internet, followed later by an option to complete a paper questionnaire and return it by mail. If neither an Internet nor mail questionnaire is received, a follow-up interview by phone or personal visit is attempted for a sample of nonrespondents. Prior to 2013, Internet collection was not used, and only three modes of collection were used. Each month, a sample of housing unit addresses and residents of group quarters facilities receive questionnaires. Housing units include a house, apartment, mobile home or trailer, a group of rooms, or a single room occupied as separate living quarters, or if vacant, intended for occupancy as separate living quarters. Group quarters are places where people live or stay that are normally owned or managed by an entity or organization providing housing and services for the residents. These services may include custodial or medical care as well as other types of assistance, and residency is commonly restricted to persons receiving these services. The group quarters population comprises both the institutional and noninstitutional group quarters populations. The institutional group quarters population

includes residents under formally authorized supervised care, such as those in skilled nursing facilities, adult correctional facilities, and psychiatric hospitals. The noninstitutional group quarters population includes residents of colleges or university housing, military barracks, and group homes.

ACS creates two sets of weights: a weight to each sample person record (both household and group quarters persons) and a weight to each sample housing unit record. For information on the weighting procedure, see the ACS methodology website at: <https://www.census.gov/programs-surveys/acs/methodology.html>.

Sample Size and Response Rate. Each year from 2005 through 2010, approximately 2.9 million housing unit addresses in the U.S. and 36,000 in Puerto Rico were selected to participate in ACS. Starting in 2011, the housing unit sample was increased to 3.54 million addresses per year. For 2005–2012, the housing unit response rate was 97%–98%; in 2013, the housing unit response rate was 90%; and in 2014 it was 97%. Beginning in 2006, the ACS sample was expanded to include 2.5% of the population living in group quarters, which included approximately 20,000 group quarters facilities and 195,000 residents of group quarters in the United States and Puerto Rico. In 2013, the group quarters sample for college dormitories was restricted to the nonsummer months. The group quarters response rate ranged between 95% and 98% for 2005–2014. For year-specific response rates, see: <http://www.census.gov/acs/www/methodology/sample-size-and-data-quality/response-rates/index.php>.

Issues Affecting Interpretation. Several changes were made to the ACS questionnaire at the beginning of 2008, including the introduction of new questions on health insurance coverage. Health insurance coverage estimates are methodologically consistent for data year 2009 and subsequent years (O'Hara and Medalia). In addition, the methodology for weighting the group quarters survey changed starting in 2011.

References

Torrieri N, Program Staff. American Community Survey design and methodology (January 2014). Washington, DC: U.S. Census Bureau; 2014. Available from: http://www2.census.gov/programs-surveys/acs/methodology/design_and_methodology/acs_design_methodology_report_2014.pdf.

O'Hara B, Medalia C. CPS and ACS health insurance estimates: Consistent trends from 2009–2012. SEHSD working paper 2014–29. Washington, DC: U.S. Census Bureau, Social, Economic, and Housing Statistics Division; 2014. Available from: http://www.census.gov/hhes/www/hlthins/data/incpovhlth/2013/CPS_ACS_Trends.pdf.

For More Information. See the ACS website at: <http://www.census.gov/programs-surveys/acs/>.

Census of Fatal Occupational Injuries (CFOI)

Bureau of Labor Statistics (BLS)

Overview. CFOI compiles comprehensive and timely information on fatal work injuries, to monitor workplace safety and to inform private and public health efforts to improve workplace safety.

Coverage. The data cover all 50 states and D.C. In selected years, data are available for Puerto Rico, the Virgin Islands, and Guam but are not included in *Health, United States* because of data comparability issues.

Methodology. CFOI is administered by BLS, in conjunction with participating state agencies, to compile counts that are as complete as possible to identify, verify, and profile fatal work injuries. Key information about each workplace fatal injury (occupation and other worker characteristics, equipment or machinery involved, and circumstances of the event) is obtained by cross-referencing source documents. For a fatal occupational injury to be included in the census, the decedent must have been employed (i.e., self-employed, working for pay, or volunteering) at the time of the event, engaged in a legal work activity, or present at the site of the incident as a requirement of his or her job. These criteria are generally broader than those used by federal and state agencies administering specific laws and regulations. Fatal work injuries that occur during a person's commute to or from work are excluded from the census counts. Fatal work injuries to volunteer workers who are exposed to the same work hazards and perform the same duties or functions as paid employees and who meet the CFOI work relationship criteria are included.

Data for CFOI are compiled from various federal, state, and local administrative sources, including death certificates, workers' compensation reports and claims, reports to various regulatory agencies, medical examiner reports, police reports, and news reports. Diverse sources are used because studies have shown that no single source captures all job-related fatal injuries. Source documents are matched so that each fatal work injury is counted only once. To ensure that a fatal work injury occurred while the decedent was at work, information is verified from two or more independent source documents or from a source document and a follow-up questionnaire.

Issues Affecting Interpretation. The numbers of fatal occupational injuries are revised once after the initial preliminary release. States have up to 8 months to update their initial published counts and may identify additional fatal work injuries after data collection has closed for a reference year. Fatal work injuries initially excluded from the published count because of insufficient information to determine work relationship may subsequently be verified as work-related and included in the revised counts. Increases in the published counts over the last 5 years based on

additional information have averaged 173 fatal occupational injuries per year, or less than 4% of the annual total.

Beginning with 2003 data, CFOI began using the 2002 North American Industry Classification System (NAICS). Starting with 2009 data, CFOI began using the 2007 NAICS to classify industries. In *Health, United States*, industry data are presented at the two-digit level. Most of the differences between the 2002 and 2007 NAICS are at a more detailed level. Therefore, the adoption of the 2007 NAICS for CFOI is unlikely to affect the trend presented in *Health, United States*. (See Appendix II, Industry of employment.)

Reference

Bureau of Labor Statistics. Revisions to the 2013 Census of Fatal Occupational Injuries (CFOI) counts. Washington, DC: U.S. Department of Labor; 2015. Available from: http://www.bls.gov/iif/oshwc/cfoi/cfoi_revised13.pdf.

For More Information. See the CFOI website at: <http://www.bls.gov/iif/oshcfoi1.htm> and the CFOI section of the *BLS Handbook of Methods* at: <http://www.bls.gov/opub/hom/pdf/homch9.pdf>.

Current Population Survey (CPS)

Bureau of Labor Statistics (BLS) and U.S. Census Bureau

Overview. CPS provides current estimates and trends in employment, unemployment, poverty, and other characteristics of the general labor force, the population as a whole, and various population subgroups.

Coverage. The Census 2000-based CPS sample, referred to as basic CPS, was introduced in April 2004, and implementation was completed by July 2005 with coverage in every state and D.C. For CPS labor force data, the adult universe (i.e., the population of marriageable age) is composed of persons aged 15 and over in the civilian noninstitutionalized population. The sample for the March CPS supplement included members of the Armed Forces who are living in a household with at least one civilian adult, as well as additional Hispanic households that are not included in the monthly labor force estimates.

Methodology. The CPS interview is divided into three parts: (a) household and demographic information, (b) labor force information, and (c) supplement information for months that include supplements. Comprehensive work experience information is gathered on the employment status, occupation, and industry of persons interviewed.

The basic CPS sample is selected from multiple frames using multiple stages of selection. Each unit is selected with a known probability to represent similar units in the universe. The sample design is state-based, with the sample in each state being independent of the others. One person generally responds for all eligible members of a household.

Estimates of poverty presented in *Health, United States* from CPS are derived from the Annual Social and Economic Supplement (ASEC), formerly called the Annual Demographic Supplement (ADS) and commonly called the March Supplement. ASEC collects data on family characteristics, household composition, marital status, migration, income from all sources, weeks worked, time spent looking for work or on layoff from a job, occupation and industry classification of the job held longest during the year, and receipt of noncash benefits (such as food stamps, school lunch program, employer-provided group health insurance plan, personal health insurance, Medicaid, Medicare, TRICARE or military health care, and energy assistance).

The additional Hispanic sample is from the previous November's basic CPS sample. If a person is identified as being of Hispanic origin from the November interview and is still residing at the same address in March, that housing unit is eligible for the March survey. This amounts to a near-doubling of the Hispanic sample because there is no overlap of housing units between the basic CPS samples in November and March.

The ASEC sample weight is an adjusted version of the final CPS sample weight. The final CPS sample weight is the product of the basic weight, the adjustments for special weighting, the noninterview adjustment, the first-stage ratio adjustment factor, and the second-stage ratio adjustment factor. Due to differences in the questionnaire, sample, and data uses for the ASEC supplement, the ASEC sample weight should be used for poverty estimates.

Sample Size and Response Rate. Beginning with 2001, the Children's Health Insurance Program (CHIP) sample expansion was introduced. This included an increase in the basic CPS sample to 60,000 households per month. Prior to 2001, estimates were based on 50,000 households per month. The expansion also included an additional 12,000 households that were allocated differentially across states, based on prior information of the number of uninsured children in each state, to produce statistically reliable current state data on the number of low-income children who do not have health insurance coverage. In an average month, the nonresponse rate for the basic CPS is about 7%–8%.

Issues Affecting Interpretation. Over the years, the number of income questions has expanded, questions on work experience and other characteristics have been added, and the month of interview was moved to March. In 2002, an ASEC (March Supplement) sample increase was implemented, requiring more time for data collection. Thus, additional ASEC interviews are now taking place in February and April. However, even with this sample increase, most of the data collection still occurs in March.

In 1994, major changes were introduced that included a complete redesign of the questionnaire and the introduction of computer-assisted interviewing for the entire survey. In addition, some of the labor force concepts

and definitions were revised. Prior to the redesign, CPS data were primarily collected using a paper-and-pencil form. Beginning in 1994, population controls were based on the 1990 census and adjusted for the estimated population undercount. Starting with *Health, United States, 2003*, poverty estimates for data years 2000 and beyond were recalculated based on the expanded CHIP sample, and Census 2000-based population controls were implemented. Starting with 2002 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. Starting with *Health, United States, 2012*, Census 2010-based population controls were implemented for poverty estimates for 2010 and beyond. For a discussion of the impact of the implementation of the Census 2010-based controls on poverty estimate trends, see: DeNavas-Walt, Proctor, and Smith (2012).

For 2013 data, the CPS ASEC used a split panel to test a new set of income questions. Starting with *Health, United States, 2015*, estimates for 2013 are presented two ways: using questions consistent with previous ASEC surveys and using the new set of income questions. Because data for 2013 (using the new income questions) and data for 2014 are based on the new set of income questions from the redesigned questionnaire, data trends need to be interpreted with caution.

References

U.S. Census Bureau. Current Population Survey: Design and methodology. Technical paper no 66. Washington, DC: U.S. Census Bureau; 2006. Available from: <http://www.census.gov/prod/2006pubs/tp-66.pdf>.

DeNavas-Walt C, Proctor BD, Smith JC. Income, poverty, and health insurance coverage in the United States: 2011. Current Population Reports, P-60-243. Washington, DC: U.S. Government Printing Office; 2012. Available from: <https://www.census.gov/prod/2012pubs/p60-243.pdf>.

DeNavas-Walt C, Proctor BD. Income and poverty in the United States: 2014. Current Population Reports, P-60-252. Washington, DC: U.S. Government Printing Office; 2015. Available from: <http://www.census.gov/content/dam/Census/library/publications/2015/demo/p60-252.pdf>.

For More Information. See the CPS website at: <http://www.census.gov/cps>.

Department of Veterans Affairs National Enrollment and Patient Databases

Department of Veterans Affairs (VA)

Overview. The VA compiles and analyzes multiple data sets on the health and health care of its clients and other

veterans to monitor access and quality of care and to conduct program and policy evaluations. The VA maintains nationwide systems that contain a statistical record for each episode of care provided under VA auspices, in VA and non-VA hospitals, nursing homes, VA residential rehabilitation treatment programs (formerly called domiciliaries), and VA outpatient clinics. The VA also maintains enrollment information for each veteran enrolled in the VA health care system.

Coverage. U.S. veterans who receive services within the VA medical system are included. Data are available for some nonveterans who receive care at VA facilities.

Methodology. Encounter data from VA clinical information systems are collected locally at each VA medical center and transmitted electronically to the VA's Austin Automation Center for use in providing nationwide statistics, reports, and comparisons.

Issues Affecting Interpretation. The databases include users of the VA health care system. VA eligibility is a hierarchy based on service-connected disabilities, income, age, and availability of services. Therefore, different VA programs may serve populations with different sociodemographic characteristics than those served by other health care systems.

For More Information. See the VA Information Resource Center website at: <http://www.virec.research.va.gov/Index.asp>.

Employee Benefits Survey—See Appendix I, National Compensation Survey (NCS).

Healthcare Cost and Utilization Project (HCUP), National (Nationwide) Inpatient Sample

Agency for Healthcare Research and Quality (AHRQ)

Overview. HCUP is a family of health care databases and related software tools developed through a federal-state-industry partnership to build a multistate health data system for health care research and decision making. The National (Nationwide) Inpatient Sample (HCUP-NIS), a component of HCUP, is the largest all-payer inpatient care database that is publicly available in the United States.

HCUP-NIS contains a core set of clinical and nonclinical information found in a typical discharge abstract, including all-listed diagnoses and procedures, discharge status, patient demographics, and charges for all patients regardless of payer (e.g., persons covered by Medicare, Medicaid, and private insurance, as well as those without insurance coverage).

Coverage. In 2013, HCUP–NIS covered more than 94% of all U.S. community hospital discharges (excluding discharges from rehabilitation or long-term acute care hospitals) from 43 states and D.C. Community hospitals are defined by the American Hospital Association as nonfederal, short-term, general, and other specialty hospitals, excluding hospital units of institutions.

The number of states participating in HCUP–NIS has generally increased each year. In the years of data presented in *Health, United States*, the number of states participating was 28 in 2000, 37 in 2005, 45 in 2010, 46 in 2011, 44 in 2012, and 43 states and D.C. in 2013. In 2013, all states except Alabama, Alaska, Delaware, Idaho, Maine, Mississippi, and New Hampshire were included.

Methodology. In 2012, HCUP–NIS was redesigned to improve national estimates. To highlight the design change, beginning with 2012 data, AHRQ renamed HCUP–NIS from the “Nationwide Inpatient Sample” to the “National Inpatient Sample.” The redesigned HCUP–NIS is now a sample of discharge records from all HCUP-participating hospitals. It approximates a 20% stratified sample of discharges from U.S. community hospitals, excluding rehabilitation and long-term acute care hospitals. The information abstracted from hospital discharge records is translated into a uniform format to facilitate both multistate and national-state comparisons and analyses.

Prior to 2012, HCUP–NIS was designed to approximate a 20% stratified sample of U.S. community hospitals, rather than a sample of discharges. The pre-2012 HCUP–NIS was a stratified probability sample of hospitals in the frame, with sampling probabilities proportional to the number of U.S. community hospitals in each stratum (ownership and control, bed size, teaching status, urban or rural location, and U.S. region). Discharge records for all patients in the sampled hospitals were included in the pre-2012 HCUP–NIS. To permit longitudinal analysis, the statistics for years prior to 2012 presented in *Health, United States* were regenerated using new trend weights taking into account the redesign.

Hospital costs are derived from total hospital charges using hospital-specific cost-to-charge ratios based on hospital cost reports from the Centers for Medicare & Medicaid Services. Hospital charges reflect the amount the hospital billed for the entire hospital stay and do not include professional (physician) fees. Costs will tend to reflect the actual costs to produce hospital services, whereas charges represent what the hospital billed for the care. Costs are adjusted for economy-wide inflation by removing increases that reflect the effect of changing average prices for the same goods and services. The U.S. Bureau of Economic Analysis Gross Domestic Product Price Index is used to remove economy-wide inflation. Additional inflation that is specific to the hospital sector is not removed in this calculation.

Sample Size and Response Rate. The 2013 HCUP–NIS contains data from 7.1 million hospital stays sampled from 4,924 hospitals.

Issues Affecting Interpretation. Weights are produced to create national estimates, but because the number of participating states has increased over time, estimates from earlier years may be biased if omitted states have substantially different hospitalization patterns than states that provided data. In 2012, the survey was redesigned. HCUP–NIS is now a sample of discharge records from all HCUP-participating hospitals, rather than a sample of hospitals from which all discharges were retained. The statistics for years prior to 2012 presented in *Health, United States* were regenerated using new trend weights taking into account the redesign.

References

Agency for Healthcare Research and Quality. Introduction to the HCUP National Inpatient Sample (NIS), 2013. In: Healthcare Cost and Utilization Project—HCUP: A federal-state-industry partnership in health data. Rockville, MD: AHRQ; 2015. Available from: <https://www.hcup-us.ahrq.gov/db/nation/nis/NISIntroduction2013.pdf>.

Houchens R, Ross D, Elixhauser A, Jiang J. Nationwide Inpatient Sample (NIS) redesign final report; 2014. HCUP Methods Series Report #2014–04 ONLINE. April 4, 2014. U.S. Agency for Healthcare Research and Quality. Available from: <https://www.hcup-us.ahrq.gov/reports/methods/2014-04.pdf>.

For More Information. See the HCUP website at: <http://www.hcup-us.ahrq.gov/>.

Medicaid Statistical Information System (MSIS)

Centers for Medicare & Medicaid Services (CMS)

Overview. CMS works with its state partners to collect data on each person served by the Medicaid program, in order to monitor and evaluate access to and quality of care, trends in program eligibility, characteristics of enrollees, changes in payment policy, and other program-related issues. MSIS is the primary data source for Medicaid statistical information. Data collected include claims for services and their associated payments for each Medicaid beneficiary, by type of service. MSIS also collects information on the characteristics of every Medicaid-eligible individual, including eligibility and demographic information.

Coverage. Medicaid data for all 50 states and D.C. are available starting in 1999. The data include information about all individuals enrolled in the Medicaid program, the services they receive, and the payments made for those services.

Methodology. Beginning in FY 1999, as a result of legislation enacted from the Balanced Budget Act of 1997, states were required to submit individual eligibility and claims data

tapes to CMS quarterly, through MSIS. Prior to FY 1999, states were required to submit an annual HCFA–2082 report, designed to collect aggregated statistical data on eligibles, recipients, services, and expenditures during a federal fiscal year (October 1 through September 30) or, at state option, to submit eligibility data and claims through MSIS. The claims data reflect bills adjudicated or processed during the year, rather than services used during the year.

Issues Affecting Interpretation. Starting with 2011 data, estimates were derived from Medicaid claims files and a new methodology was used to obtain estimates. Therefore, caution should be used when comparing data for 2010 and earlier with more recent data. For more information on data and analytic issues, see: <https://www.cms.gov/Research-Statistics-Data-and-Systems/Computer-Data-and-Systems/MedicaidDataSourcesGenInfo/MSIS-Tables.html>.

For More Information. See the CMS websites at: <http://www.cms.hhs.gov/home/medicaid.asp> and <http://www.medicare.gov/Medicare-CHIP-Program-Information/By-Topics/Data-and-Systems/Data-and-Systems.html> and the Research Data Assistance Center (ResDAC) website at: <http://cms.gov/Research-Statistics-Data-and-Systems/Research/ResearchGenInfo/ResearchDataAssistanceCenter.html>. (Also see Appendix II, Medicaid.)

Medical Expenditure Panel Survey (MEPS)

Agency for Healthcare Research and Quality (AHRQ)

Overview. MEPS produces nationally representative estimates of health care use, expenditures, sources of payment, insurance coverage, and quality of care. MEPS consists of three components: the Household Component (HC), the Medical Provider Component (MPC), and the Insurance Component (IC). Data from MEPS–HC and MEPS–MPC are used in *Health, United States*.

Coverage. The U.S. civilian noninstitutionalized population is the primary population represented.

Methodology. MEPS–HC is a national probability survey conducted on an annual basis since 1996. The panel design of the survey features five rounds of interviewing covering 2 full calendar years. The HC is a nationally representative survey of the civilian noninstitutionalized population drawn from a subsample of households that participated in the prior year's National Health Interview Survey. Missing expenditure data in the HC are imputed largely from data collected in the MPC.

The MPC collects data from hospitals, physicians, home health care providers, and pharmacies that were reported in the HC as providing care to MEPS sample persons. Data are collected in the MPC to improve the accuracy of the expenditure estimates that would be obtained if derived solely from the HC. The MPC is particularly useful in obtaining expenditure information for persons enrolled in

managed care plans and Medicaid recipients. Sample sizes for the MPC vary from year to year depending on the HC sample size and the MPC sampling rates for providers.

The MEPS predecessor, the 1987 National Medical Expenditure Survey (NMES), consisted of two components: the Household Survey (HS) and the Medical Provider Survey (MPS). The NMES–HS component was designed to provide nationally representative estimates for the U.S. civilian noninstitutionalized population for the calendar year 1987. Data from the NMES–MPS component were used in conjunction with HS data to produce estimates of health care expenditures. The NMES–HS consisted of four rounds of household interviews. Income information was collected in a special supplement administered early in 1988. Events under the scope of the NMES–MPS included medical services provided by or under the direction of a physician, all hospital events, and home health care.

Sample Size and Response Rate. In the 2012 MEPS, there were 14,763 families covered, and 37,182 respondents over the course of the year. For the same year, the overall annual response rate was 56.3%, reflecting nonresponse to the National Health Interview Survey from which the MEPS sample was selected, as well as nonresponse and attrition in MEPS.

Issues Affecting Interpretation. The 1987 estimates are based on NMES, and 1996 and later years' estimates are based on MEPS. Because expenditures in NMES were based primarily on charges, whereas those for MEPS were based on payments, data for NMES were adjusted to be more comparable with MEPS by using estimated charge-to-payment ratios for 1987. For a detailed explanation of this adjustment, see Zuvekas and Cohen (2002).

References

Ezzati-Rice TM, Rohde F, Greenblatt J. Sample design of the Medical Expenditure Panel Survey Household Component, 1998–2007. Methodology report no 22. Rockville, MD: Agency for Healthcare Research and Quality; 2008. Available from: http://www.meps.ahrq.gov/mepsweb/data_files/publications/mr22/mr22.shtml.

Zuvekas SH, Cohen JW. A guide to comparing health care expenditures in the 1996 MEPS to the 1987 NMES. *Inquiry* 2002;39(1):76–86.

For More Information. See the MEPS website at: <http://www.meps.ahrq.gov/mepsweb/>.

Medicare Administrative Data

Centers for Medicare & Medicaid Services (CMS)

Overview. CMS collects and synthesizes Medicare enrollment, spending, and claims data to monitor and evaluate access to and quality of care, trends in utilization, changes in payment policy, and other program-related issues. Data include claims information for services furnished

to Medicare fee-for-service beneficiaries and Medicare enrollment data. Claims data include type of service, procedures, diagnoses, dates of service, charge amounts, and payment amounts. Enrollment data include date of birth, sex, race, ethnicity, and reason for entitlement.

Coverage. Enrollment data are for all persons enrolled in the Medicare program. Claims data include data for Medicare fee-for-service beneficiaries who received services and for whom claims were filed. In general, claims data are not included for beneficiaries enrolled in managed care plans.

Methodology. The claims and utilization data files contain extensive utilization information at various levels of summarization for a variety of providers and services. There are many types and levels of these files: National Claims History (NCH) files, Standard Analytic files (SAFs), Medicare Provider and Analysis Review (MedPAR) files, Medicare enrollment files, and various other files.

The NCH files contain all institutional and noninstitutional claims submitted during a calendar year, including adjustment claims. SAFs contain “final action” claims data in which all adjustments have been resolved. Both the NCH and SAF files contain information collected by Medicare to pay for health care services provided to a Medicare beneficiary. SAFs are available for each institutional (inpatient, outpatient, skilled nursing facility, hospice, or home health agency) and noninstitutional (physician and durable medical equipment providers) claim type. The record unit of SAFs is the claim (some episodes of care may have more than one claim).

MedPAR files contain inpatient hospital and skilled nursing facility (SNF) final action stay records. Each MedPAR record represents a stay in an inpatient hospital or SNF. An inpatient stay record summarizes all services rendered to a beneficiary from the time of admission to a facility, through discharge. Each MedPAR record may represent one claim or multiple claims, depending on the length of a beneficiary's stay and the amount of inpatient services used throughout the stay.

The Denominator file contains demographic and enrollment information about each beneficiary enrolled in Medicare during a calendar year. The information in the Denominator file is frozen in March of the following calendar year. Some of the information contained in this file includes the beneficiary unique identifier, state and county codes, ZIP code, date of birth, date of death, sex, race, age, monthly entitlement indicators (for Medicare Part A, Medicare Part B, or Part A and Part B), reasons for entitlement, state buy-in indicators, and monthly managed care indicators (yes or no). The Denominator file is used to determine beneficiary demographic characteristics, entitlement, and beneficiary participation in Medicare managed care organizations (MCOs).

Issues Affecting Interpretation. Because Medicare MCOs might not file claims, files based only on claims data will exclude care for persons enrolled in Medicare MCOs. In addition, to maintain a manageable file size, some files are based on a

sample of enrollees rather than on all Medicare enrollees. Coding and the interpretation of Medicare coverage rules have also changed over the life of the Medicare program.

For More Information. See the CMS Research Data Assistance Center (ResDAC) website at: <http://www.resdac.org> and the CMS website at: [http://www.cms.gov/Research-Statistics-Data-and-Systems.html](http://www.cms.gov/Research-Statistics-Data-and-Systems/Research-Statistics-Data-and-Systems.html). (Also see Appendix II, Medicare.)

Medicare Current Beneficiary Survey (MCBS)

Centers for Medicare & Medicaid Services (CMS)

Overview. MCBS produces nationally representative estimates of health and functional status, health care use and expenditures, health insurance coverage, and socioeconomic and demographic characteristics of Medicare beneficiaries. It is used to estimate expenditures and sources of payment for all services used by Medicare beneficiaries, including copayments, deductibles, and noncovered services; to ascertain all types of health insurance coverage and relate coverage to sources of payment; and to trace processes over time, such as changes in health status and the effects of program changes.

Coverage. MCBS is a continuous survey of a nationally representative sample of aged, institutionalized, and disabled Medicare beneficiaries.

Methodology. The overlapping panel design of the survey allows each sample person (or their proxies) to be interviewed three times a year for 4 years, whether he or she resides in the community or a facility or moves between the two settings, using the version of the questionnaire appropriate to the setting. Sampled people are interviewed using computer-assisted personal interviewing (CAPI) survey instruments. Because residents of long-term care facilities often are in poor health, information about institutionalized residents is collected from proxy respondents such as nurses and other primary caregivers affiliated with the facility. The sample is selected from the Medicare enrollment files, with oversampling among disabled persons under age 65 and among persons aged 85 and over.

MCBS has two components: the Cost and Use file and the Access to Care file. Medicare claims are linked to survey-reported events to produce the Cost and Use file, which provides complete expenditure and source-of-payment data on all health care services, including those not covered by Medicare. The Access to Care file contains information on beneficiaries' access to health care, satisfaction with care, and usual source of care. The sample for this file represents the always-enrolled population—those who participated in the Medicare program for the entire year. In contrast, the Cost and Use file represents the ever-enrolled population, including those who entered Medicare and those who died during the year.

Sample Size and Response Rate. Each fall, about one-third of the MCBS sample is retired and roughly 6,000 new sample persons are included in the survey; the exact number chosen is based on projections of target samples of 12,000 persons with 3 years of cost and use information distributed appropriately across the sample cells. In the community, response rates for initial interviews are approximately 80%; once respondents have completed the first interview, their participation in subsequent rounds is 95% or more. In recent rounds, data have been collected from approximately 16,000 beneficiaries. Roughly 90% of the sample is made up of persons who live in the community, with the remaining persons living in long-term care facilities. Response rates for facility interviews approach 100%.

Issues Affecting Interpretation. Because only Medicare beneficiaries are included in MCBS, the survey excludes a small proportion of persons aged 65 and over who are not enrolled in Medicare. This should be noted when using MCBS to make estimates of the entire population aged 65 and over in the United States. The 2012 Cost and Use file estimates were created with a new imputation methodology; therefore some utilization estimates may not be comparable with previous years.

References

Adler GS. A profile of the Medicare Current Beneficiary Survey. *Health Care Financ Rev* 1994;15(4):153–63.

Lo A, Chu A, Apodaca R. Redesign of the Medicare Current Beneficiary Survey sample. Rockville, MD: Westat, Inc.; 2003. Available from: <http://www.amstat.org/sections/srms/Proceedings/y2002/Files/JSM2002-000662.pdf>.

For More Information. See the MCBS website at: <http://www.cms.hhs.gov/MCBS>.

Monitoring the Future (MTF) Study

University of Michigan, supported by the National Institute on Drug Abuse (NIDA)

Overview. MTF is an ongoing study that uses annual surveys to track the behaviors, attitudes, and values of U.S. secondary school students, college students, and adults through age 55. Data collected include lifetime, annual, and 30-day prevalence of use of many illegal drugs, inhalants, tobacco, and alcohol.

Coverage. MTF surveys a sample of 12th, 10th, and 8th graders in public and private high schools in the coterminous United States. Follow-up questionnaires are mailed to a sample of each graduating class for a number of years after their initial participation, to gather information on college students, young adults, and older adults.

Methodology. The survey design is a multistage random sample, with stage 1 being the selection of particular geographic areas, stage 2 the selection of one or more schools in each area, and stage 3 the selection of students within each school. Data are collected using self-administered questionnaires conducted in the classroom by representatives of the University of Michigan's Institute for Social Research. Dropouts and students who are absent on the day of the survey are excluded. Recognizing that the dropout population is at higher risk for drug use, MTF was expanded in 1991 to include similar nationally representative samples of 8th and 10th graders, who have lower dropout rates than seniors and include future high-risk 12th grade dropouts. For more information on MTF adjustments for absentees and dropouts, see Johnston et al. (2014 and preceding); and Miech et al. (2015).

Sample Size and Response Rate. In 2014, a total of 41,551 students in 377 public and private schools in the coterminous United States participated. The annual senior samples comprised 13,015 12th graders in 122 public and private high schools nationwide. The 10th-grade samples involved 13,341 students in 114 schools, and the 8th-grade samples had 15,195 students in 141 schools. Student response rates were 90% for grade 8, 88% for grade 10, and 82% for grade 12 and have been relatively constant across time. Absentees constitute virtually all of the nonresponding students.

Issues Affecting Interpretation. Estimates of substance use among youth based on the National Survey on Drug Use & Health (NSDUH) are not directly comparable with estimates based on MTF and the Youth Risk Behavior Survey (YRBS). In addition to the fact that MTF excludes dropouts and absentees, rates are not directly comparable across these surveys because of differences in populations covered, sample design, questionnaires, interview setting, and data cleaning procedures. NSDUH collects data in residences, whereas MTF and YRBS collect data in school classrooms. In addition, NSDUH estimates are tabulated by age, whereas MTF and YRBS estimates are tabulated by grade, representing different ages as well as different populations.

References

Miech RA, Johnston LD, O'Malley PM, Bachman JG, Schulenberg JE. Monitoring the Future National Survey results on drug use: 1975–2014. Vol I. Secondary school students. Ann Arbor, MI: Institute for Social Research, The University of Michigan; 2015. Available from: http://www.monitoringthefuture.org/pubs/monographs/mtf-vol1_2014.pdf.

Cowan CD. Coverage, sample design, and weighting in three federal surveys. *J Drug Issues* 2001;31(3):599–614.

Johnston LD, O'Malley PM, Bachman JG, Schulenberg JE, Miech, RA. Monitoring the Future National Survey results on drug use, 1975–2013: Vol I, Secondary school

students. Ann Arbor, MI: Institute for Social Research, The University of Michigan; 2014. Available from: http://www.monitoringthefuture.org/pubs/monographs/mtf-vol1_2013.pdf.

For More Information. See the NIDA website at: <http://www.nida.nih.gov/Infobox/HSYouthtrends.html> and the MTF website at: <http://www.monitoringthefuture.org>.

National Ambulatory Medical Care Survey (NAMCS)

CDC/NCHS

Overview. NAMCS provides national data about the provision and use of medical care services in office-based physician practices in the United States, using information collected from medical records. Data are collected on type of providers seen; reason for visit; diagnoses; drugs ordered, provided, or continued; and selected procedures and tests ordered or performed during the visit. Patient data include age, sex, race, and expected source of payment. Data are also collected on selected characteristics of physician practices, including the adoption and use of electronic health record (EHR) systems.

Coverage. NAMCS covers patient encounters in the offices of nonfederally employed physicians classified by the American Medical Association (AMA) or American Osteopathic Association (AOA) as office-based patient care physicians in the United States. Patient encounters with physicians engaged in prepaid practices (health maintenance organizations [HMOs], independent practice organizations [IPAs], and other prepaid practices) are included in NAMCS. Excluded are visits to hospital-based physicians; visits to specialists in anesthesiology, pathology, or radiology; and visits to physicians who are principally engaged in teaching, research, or administration. Telephone contacts and nonoffice visits are also excluded. Starting in 2006, NAMCS includes visits to a separate sample of community health centers (CHCs). In 2012, the NAMCS survey sample size was increased fivefold to allow for state-level estimates in the 34 most populous states and the U.S. Census Bureau divisions.

Methodology. A multistage probability design is employed. Beginning in 1989–2011, the first-stage sample consisted of 112 primary sampling units (PSUs), which were selected from about 1,900 such units into which the United States had been divided. In each sample PSU, a sample of practicing nonfederal, office-based physicians was selected from master files maintained by AMA and AOA. The final stage involved systematic random samples of office visits during randomly assigned 7-day reporting periods. In 2012, the sampling design was changed to a list sample of physicians, instead of an area sample, to ensure adequate representation for state-level estimates. Starting in 1989, the survey included all 50 states and D.C.

Starting in 2006–2011, a dual-sampling procedure was used to select CHC physicians and nonphysician clinicians. First, the traditional NAMCS sample was selected using the methods described above. Second, information from the Health Resources and Services Administration and the Indian Health Service was used to select a sample of CHCs. Within CHCs, a maximum of three health care providers were selected, including physicians, physician assistants, nurse practitioners, or nurse midwives. After selection, CHC providers followed traditional NAMCS methods for selecting patient visits. Another major change for 2012 was the mode of data collection—from in-person interviews with a paper questionnaire to obtain physician practice information to laptop-assisted data collection using automated survey instruments. Over time, interviewer abstraction from visit records has been increasing. In 2012, medical abstraction by interviewers was the predominant method of data collection.

Since 2008, a supplemental mail survey on EHR systems has been conducted in addition to the core NAMCS. This supplement is known as the National Ambulatory Medical Care Survey–National Electronic Health Records Survey (NEHRS). Starting in 2010, the mail NEHRS sample size was increased fivefold to allow for state-level estimates without needing to combine NEHRS with the core NAMCS. Survey questions have been added since the introduction of NEHRS.

The U.S. Census Bureau acts as the data collection agent for NAMCS. Starting in 2012, Census field representatives have used laptops containing an automated version of each survey instrument to (a) conduct induction interviews with the physician or his or her representative to obtain information about the practice and ensure that it is within the scope of the survey; (b) determine which visits to sample; and (c) abstract and record data from medical charts. Prior to 2012, physicians were asked to perform their own visit sampling and record abstraction using a paper-and-pencil mode of data collection, but Census field representatives were available to perform these tasks if needed. Beginning in 2012, abstraction by field representatives became the preferred mode of data collection, accounting for 98% of records collected.

Sample data are weighted to produce national estimates. The estimation procedure used in NAMCS has four basic components: inflation by the reciprocal of the probability of selection, adjustment for nonresponse, ratio adjustment to fixed totals, and weight smoothing.

Sample Size and Response Rate. In 2011, a sample of 3,819 physicians was selected: 2,555 were in-scope and 1,400 participated, for an unweighted response rate of 54% (54% weighted). Data were provided for 30,872 visits. In 2012, a sample of 15,740 physicians was selected: 9,574 were in-scope and 3,010 participated, for an unweighted response rate of 39% (39% weighted). Data were provided for 76,330 visits. The response rates have been modified to accommodate the mixture of one- and two-stage samples

of providers. The 2013 NAMCS–NEHRS had a sample of 10,302 physicians. The unweighted response rate was 70% (67% weighted).

Issues Affecting Interpretation. The NAMCS patient record form is modified approximately every 2–4 years to reflect changes in physician practice characteristics, patterns of care, and technological innovations. Examples of recent changes include increasing the number of drugs recorded on the patient record form and adding checkboxes for specific tests or procedures performed. Sample sizes vary by survey year. For some years it is suggested that analysts combine two or more years of data if they wish to examine relatively rare populations or events. Starting with *Health, United States, 2005*, data for survey years 2001–2002 were revised to be consistent with the weighting scheme introduced in the 2003 NAMCS data. For more information on the new weighting scheme, see Hing et al. (2005). The 2012 sampling design change may affect trending 2012 with earlier data. For more information on the new sampling design, see Hing et al. (2016).

References

Hing E, Cherry DK, Woodwell DA. National Ambulatory Medical Care Survey: 2003 summary. Advance data from vital and health statistics; no 365. Hyattsville, MD: NCHS; 2005. Available from: <http://www.cdc.gov/nchs/data/ad/ad365.pdf>.

Hing E, Shimizu IM, Talwalkar A. Nonresponse bias in estimates from the 2012 National Ambulatory Medical Care Survey. *Vital Health Stat* 2(171). Hyattsville, MD: NCHS; 2016. Available from: http://www.cdc.gov/nchs/data/series/sr_02/sr02_171.pdf.

For More Information. See the National Health Care Surveys website at: <http://www.cdc.gov/nchs/dhcs.htm> and the Ambulatory Health Care Data website at: <http://www.cdc.gov/nchs/ahcd.htm>.

National Compensation Survey (NCS)

Bureau of Labor Statistics (BLS)

Overview. NCS provides comprehensive measures of occupational earnings, compensation cost trends, benefit incidence, and detailed plan provisions based on surveys of a sample of employers.

Coverage. NCS provides information for the nation for the nine census divisions and for 152 selected areas (combined statistical areas, metropolitan statistical areas, micropolitan statistical areas, and county clusters). NCS includes both full- and part-time workers who are paid a wage or salary and includes data for the civilian economy, including both private industry and state and local government. It excludes agriculture, private household workers, the self-employed, and the federal government.

Methodology. NCS is conducted quarterly by BLS' Office of Compensation and Working Conditions. The sample consists of approximately 152 areas that represent the nation's metropolitan statistical areas and micropolitan statistical areas (as defined by the Office of Management and Budget [OMB]) and the remaining portions of the 50 states. The sample is selected using a three-stage design. The first stage involves the selection of geographic areas for the state and local government sample and the private industry sample. In the second stage, establishments are selected systematically, with the probability of selection proportionate to their relative employment size within sampled areas. Use of this technique means that the larger an establishment's employment, the greater its chance of selection. The third stage of sampling is a probability sample of occupations within a sampled establishment. This step is performed by the BLS field economist during an interview with the respondent establishment in which selection of an occupation is based on probability of selection proportionate to employment in the establishment, and each occupation is classified under its corresponding major occupational group.

Data collection is conducted by BLS field economists. Data are gathered from each establishment on the primary business activity of the establishment; types of occupations; number of employees; wages, salaries, and benefits; hours of work; and duties and responsibilities. Data are collected for the pay period including the 12th day of the survey months of March, June, September, and December.

Sample Size and Response Rate. The March 2015 sample consists of about 8,600 establishments in private industry and about 1,500 establishments in state and local government.

Issues Affecting Interpretation. Prior to 1999, estimates were based on multiple surveys that were replaced by NCS; therefore, trend analyses based on estimates prior to 1999 should be interpreted with care.

The state and local government sample is revised every 10 years and was replaced in its entirety in December 2007. As a result of this replacement, the number of state and local government occupations and establishments increased substantially. The private industry sample is rotated approximately every 5 years, which makes the sample more representative of the economy and reduces respondent burden. The sample is replaced on a cross-area, cross-establishment basis.

Compensation cost levels in state and local government should not be directly compared with levels in private industry. Differences between these sectors stem from factors such as variation in work activities and occupational structures.

References

Bureau of Labor Statistics. Employer costs for employee compensation—March 2015 [press release USDL–14–

1075]. Washington, DC: U.S. Department of Labor; 2015 June 10. Available from: http://www.bls.gov/news.release/archives/ecec_06102015.htm.

Wiatrowski WJ. The National Compensation Survey: Compensation statistics for the 21st century. Washington, DC: U.S. Department of Labor, Bureau of Labor Statistics. Compensation and Working Conditions (CWC) Online 2000; Winter:5–14. Available from: <http://www.bls.gov/opub/mlr/cwc/the-national-compensation-survey-compensation-statistics-for-the-21st-century.pdf>.

U.S. Bureau of Labor Statistics. BLS handbook of methods, Ch. 8: National compensation measures; 2007. Available from: <http://www.bls.gov/opub/hom/pdf/homch8.pdf>.

For More Information. See the NCS website at: <http://www.bls.gov/ncs/>.

National Health and Nutrition Examination Survey (NHANES)

CDC/NCHS

Overview. NHANES is designed to assess the health and nutritional status of adults and children in the United States. The survey is unique in that it combines interviews and physical examinations. NHANES collects data on the prevalence of chronic diseases and conditions (including undiagnosed conditions) and on risk factors such as obesity, elevated serum cholesterol levels, hypertension, diet and nutritional status, and numerous other measures.

Coverage. NHANES III, conducted during 1988–1994, and the continuous NHANES, begun in 1999, target the civilian noninstitutionalized U.S. population.

Methodology. NHANES includes clinical examinations, selected medical and laboratory tests, and self-reported data. NHANES interviews persons in their homes and conducts medical examinations in a mobile examination center (MEC), including laboratory analysis of blood, urine, and other tissue samples. Medical examinations and laboratory tests follow very specific protocols and are standardized as much as possible to ensure comparability across sites and providers. In 1999–2002, as a substitute for the MEC examinations, a small number of survey participants received an abbreviated health examination in their homes if they were unable to come to the MEC.

The survey for NHANES III was conducted from 1988 to 1994 using a stratified, multistage probability design to sample the civilian U.S. population living in households. About 40,000 persons aged 2 months and over were selected and asked to complete an extensive interview and a physical examination. Participants were selected from households in 81 survey units across the United States. Children aged 2 months to 5 years, persons aged 60 and over, black

persons, and persons of Mexican origin were oversampled to provide precise descriptive information on the health status of selected population groups in the United States.

Beginning in 1999, NHANES became a continuous annual survey, collecting data every year from a representative sample of the civilian noninstitutionalized U.S. population, newborns and older, through in-home personal interviews and physical examinations in the MEC. The sample design is a complex, multistage, clustered design using unequal probabilities of selection. The first-stage sample frame for continuous NHANES during 1999–2001 was the list of primary sampling units (PSUs) selected for the design of the National Health Interview Survey. Typically, an NHANES PSU is a county. For 2002, an independent sample of PSUs (based on current census data) was selected. This independent design was used for the period 2002–2006. In 2007–2010 and 2011–2014, the sample was redesigned. For 1999, because of a delay in the start of data collection, 12 distinct PSUs were in the annual sample. For each year in 2000–2014, 15 PSUs were selected. The within-PSU design involves forming secondary sampling units that are nested within census tracts, selecting dwelling units within secondary units, and then selecting sample persons within dwelling units. Selection of the final sample person involves differential probabilities of selection according to the demographic variables of sex (male or female), race and ethnicity, and age. Because of the differential probabilities of selection, dwelling units are screened for potential sample persons.

Beginning in 1999, NHANES oversampled low-income persons, adolescents aged 12–19, persons aged 60 and over, African American persons, and persons of Mexican origin. The sample for data years 1999–2006 was not designed to give a nationally representative sample for the total Hispanic population residing in the United States. Starting with 2007–2010 data collection, all Hispanic persons were oversampled, not just persons of Mexican origin, and adolescents were no longer oversampled. In 2011–2014, the sampling design was changed and the following groups were oversampled: Hispanic persons; non-Hispanic black persons; non-Hispanic Asian persons; non-Hispanic white and other persons at or below 130% of poverty; and non-Hispanic white and other persons aged 80 and over. For more information on the sample design for 1999–2006, see: http://www.cdc.gov/nchs/data/series/sr_02/sr02_155.pdf; for 2007–2010, see: http://www.cdc.gov/nchs/data/series/sr_02/sr02_160.pdf; and for 2011–2014, see: http://www.cdc.gov/nchs/data/series/sr_02/sr02_162.pdf.

The estimation procedure used to produce national statistics for all NHANES involved inflation by the reciprocal of the probability of selection, adjustment for nonresponse, and poststratified ratio adjustment to population totals. Sampling errors also were estimated, to measure the reliability of the statistics.

Sample Size and Response Rate. Over the 6-year survey period of NHANES III, 39,695 persons were selected; the household

interview response rate was 86% (33,994); and the medical examination response rate was 78% (30,818). For NHANES 2011–2012, a total of 13,431 persons were eligible, of which 73% (9,756) were interviewed and 70% (9,338) completed the health examination component. For NHANES 2013–2014, a total of 14,332 persons were eligible, of which 71% (10,175) were interviewed and 68% (9,813) completed the health examination component. For more information on unweighted NHANES response rates and response weights using sample size weighted to Current Population Survey population totals, see: http://www.cdc.gov/nchs/nhanes/response_rates_CPS.htm.

Issues Affecting Interpretation. Data elements, laboratory tests performed, and the technological sophistication of medical examination and laboratory equipment have changed over time. Therefore, trend analyses should carefully examine how specific data elements were collected across the various survey years. Data files are revised periodically. If the file changes are minor and the impact on estimates small, then the data are not revised in *Health, United States*. Major data changes are incorporated.

Periodically, NHANES changes its sampling design to oversample different groups. Because the total sample size in any year is fixed due to operational constraints, sample sizes for the other oversampled groups (including Hispanic persons and non-low-income white and other persons) were decreased. Therefore, trend analyses on demographic subpopulations should be carefully evaluated to determine if the sample sizes meet the NHANES Analytic Guidelines. In general, any 2-year data cycle in NHANES can be combined with adjacent 2-year data cycles to create analytic data files based on 4 or more years of data, in order to improve precision. However, because of the sample design change for 2011–2012, the data user should be aware of the implications if these data are combined with data from earlier survey cycles. Users are advised to examine their estimates carefully to see if the 4-year estimates (and sampling errors) are consistent with each set of 2-year estimates.

References

Ezzati TM, Massey JT, Waksberg J, et al. Sample design: Third National Health and Nutrition Examination Survey. NCHS. Vital Health Stat 1992;2(113). Available from: http://www.cdc.gov/nchs/data/series/sr_02/sr02_113.pdf.

NCHS. Plan and operation of the Third National Health and Nutrition Examination Survey, 1988–94. Vital Health Stat 1994;1(32). Available from: http://www.cdc.gov/nchs/data/series/sr_01/sr01_032.pdf.

Johnson CL, Paulose-Ram R, Ogden CL, et al. National Health and Nutrition Examination Survey: Analytic guidelines, 1999–2010. NCHS. Vital Health Stat 2013;2(161). Available from: http://www.cdc.gov/nchs/data/series/sr_02/sr02_161.pdf.

Johnson CL, Dohrmann SM, Burt VL, Mohadjer LK. National Health and Nutrition Examination Survey: Sample design, 2011–2014. Vital Health Stat 2014;2(162). Available from: http://www.cdc.gov/nchs/data/series/sr_02/sr02_162.pdf.

For More Information. See the NHANES website at: <http://www.cdc.gov/nchs/nhanes.htm>.

National Health Expenditure Accounts (NHEA)

Centers for Medicare & Medicaid Services (CMS)

Overview. NHEA provide estimates of aggregate health care expenditures in the United States from 1960 onward. NHEA contain all of the main components of the health care system within a unified, mutually exclusive and exhaustive structure. The accounts measure spending for health care in the United States by type of good or service delivered (e.g., hospital care, physician and clinical services, or retail prescription drugs) and by the source of funds that pay for that care (e.g., private health insurance, Medicare, Medicaid, or out-of-pocket). A consistent set of definitions is used for health care goods and services and for sources of funds that finance health care expenditures, allowing for comparisons over time.

Methodology. The primary data sources used to estimate hospital care spending are the American Hospital Association's (AHA) Annual Survey and the U.S. Census Bureau's Services Annual Survey (SAS). These sources are supplemented by data on federal hospital spending. Expenditures for physician and clinical services are estimated using data from SAS and the U.S. Census Bureau's quinquennial Economic Census. Expenditures for nursing care facilities and continuing care retirement communities, home health care, dentists, and the services of health care professionals (e.g., chiropractors, private duty nurses, therapists, and podiatrists) are estimated using data from SAS and the quinquennial Economic Census. The estimate of retail spending for prescription drugs is based on prescription drug data from the U.S. Census Bureau's Census of Retail Trade and data from IMS Health (Parsippany, NJ), an organization that collects data on retail sales of prescription drugs.

Expenditures for durable and other nondurable medical products purchased in retail outlets are based on input-output and personal consumption expenditure data (Bureau of Economic Analysis), the Economic Census and Annual Retail Trade Survey (ARTS) data (U.S. Census Bureau), Consumer Expenditure Survey data (Bureau of Labor Statistics [BLS]), Medical Expenditure Panel Surveys (MEPS) data (Agency for Healthcare Research and Quality [AHRQ]), and over-the-counter sales data from Kline and Company, Inc. Durable and nondurable products provided to inpatients in hospitals or nursing homes, and those provided by licensed health professionals or through home health

care agencies, are excluded from NHEA estimates of durable and nondurable medical products but are included with the expenditure estimates for the provider service category.

The Structures and Equipment component of NHEA includes estimates of the value of new construction put in place and new capital equipment (including software) purchased by the medical sector during the year. These estimates are based on a variety of data from the U.S. Census Bureau and the Bureau of Economic Analysis, including the Annual Capital Expenditures Survey, the C-30 Survey, and data from the National Income and Product Accounts.

Expenditures for noncommercial research are included in the Investment category of NHEA and are developed primarily from information gathered by the National Institutes of Health and the National Science Foundation. The cost of commercial research (such as by drug companies) is assumed to be embedded in the price charged for the product and therefore is not included in the noncommercial research category.

Private health insurance spending for health care goods and services is derived using data from the U.S. Census Bureau, the American Medical Association (AMA), the American Hospital Association (AHA), and IMS Health, as well as household data from surveys such as the National Medical Care Expenditure Survey (National Center for Health Services Research, 1987) and later, MEPS (AHRQ, 1996–2014). The net cost of private health insurance (which includes administrative costs, additions to reserves, rate credits and dividends, premium taxes, and net underwriting gains or losses) is estimated using data from A.M. Best (Oldwick, NJ), the National Association of Insurance Commissioners, BLS surveys on the cost of employer-sponsored health insurance and consumer expenditures, MEPS data for self-insured plans, data from privately funded surveys, and numerous consulting firms and private health insurance trade organizations.

Estimates of federal health care program spending (e.g., Medicare, Medicaid, and Department of Defense) were developed using administrative records maintained by the servicing agencies. Out-of-pocket spending (direct spending by consumers for copayments, coinsurance, deductibles, and payments for goods and services not covered by insurance) was estimated using data from SAS (U.S. Census Bureau), the Consumer Expenditure Survey (BLS), MEPS (AHRQ), the AHA Annual Survey, and IMS Health.

Issues Affecting Interpretation. Every 5 years, NHEA undergoes a comprehensive revision that includes the incorporation of newly available source data, methodological and definitional changes, and benchmark estimates from the Economic Census. During these comprehensive revisions, the entire NHEA time series is opened for revision.

References

Martin AB, Hartman M, Benson J, Catlin A; the National Health Expenditure Accounts Team. National health

spending in 2014: Faster growth driven by coverage expansion and prescription drug spending. *Health Aff (Millwood)* 2015;35(1):1–11.

Centers for Medicare & Medicaid Services. National Health Expenditure Accounts: Methodology paper, 2014: Definitions, sources, and methods. Baltimore, MD: CMS; 2015. Available from: <https://www.cms.gov/Research-Statistics-Data-and-Systems/Statistics-Trends-and-Reports/NationalHealthExpendData/Downloads/DSM-14.pdf>.

Centers for Medicare & Medicaid Services. Summary of 2014 comprehensive revision to National Health Expenditure Accounts. Baltimore, MD: CMS; 2015. Available from: <https://www.cms.gov/Research-Statistics-Data-and-Systems/Statistics-Trends-and-Reports/NationalHealthExpendData/Downloads/benchmark2014.pdf>.

For More Information. See the CMS National Health Expenditure Accounts website at: <http://www.cms.gov/Research-Statistics-Data-and-Systems/Statistics-Trends-and-Reports/NationalHealthExpendData/NationalHealthAccountsHistorical.html>.

National Health Interview Survey (NHIS)

CDC/NCHS

Overview. NHIS monitors the health of the U.S. population through the collection and analysis of data on a broad range of health topics. A major strength of this survey lies in the ability to analyze health measures by many demographic and socioeconomic characteristics. During household interviews, NHIS obtains information on activity limitation, illnesses, injuries, chronic conditions, health insurance coverage (or lack thereof), utilization of health care, and other health topics.

Coverage. The survey covers the civilian noninstitutionalized population of the United States. Among those excluded are patients in long-term care facilities, persons on active duty with the Armed Forces (although their dependents are included), incarcerated persons, and U.S. nationals living in foreign countries.

Methodology. NHIS is a cross-sectional household interview survey. Sampling and interviewing are continuous throughout each year. The sampling plan follows a multistage area probability design that permits the representative sampling of households. Traditionally, the sample for NHIS is redesigned and redrawn about every 10 years to better measure the changing U.S. population and to meet new survey objectives. A new sample design was implemented in the 2006 survey and will be used until 2016. The fundamental structure of the 2006 design is very similar to the previous design for the 1995–2005 surveys. Only the current sampling plan covering design years 2006–2015 is addressed here. The first stage of the current sampling plan

consists of a sample of 428 primary sampling units (PSUs) drawn from approximately 1,900 geographically defined PSUs that cover the 50 states and D.C. A PSU consists of a county, a small group of contiguous counties, or a metropolitan statistical area.

Within a PSU, two types of second-stage units are used: area segments and permit segments. Area segments are defined geographically and contain an expected 8, 12, or 16 addresses. Permit segments cover housing units built after the 2000 census. The permit segments are defined using updated lists of building permits issued in the PSU since 2000 and contain an expected four addresses. Within each segment, all occupied households at the sample addresses are targeted for interview.

The total NHIS sample of PSUs is subdivided into four separate panels, or subdesigns, such that each panel is a representative sample of the U.S. population. This design feature has a number of advantages, including flexibility for the total sample size. The households selected for interview each week in NHIS are a probability sample representative of the target population.

Oversampling of the black and Hispanic populations was retained in the 2006–2015 design to allow for more precise estimation of health characteristics in these populations. The current sample design also oversamples the Asian population. In addition, the sample adult selection process was revised so that when black, Hispanic, or Asian persons aged 65 and over are present, they have an increased chance of being selected as the sample adult.

The current NHIS questionnaire, implemented in 1997, has two basic parts: a Basic Module or Core and one or more supplements that vary by year. The Core remains largely unchanged from year to year and allows for trend analysis and for data from more than 1 year to be pooled to increase the sample size for analytic purposes. The Core contains three components: the Family, the Sample Adult, and the Sample Child. The Family component collects information on everyone in the family. From each family in NHIS, one sample adult is randomly selected to participate in the Sample Adult questionnaire. For families with children under age 18, one sample child is randomly selected to participate in the Sample Child questionnaire. For children, information is provided by a knowledgeable family member aged 18 or over residing in the household. Because some health issues are different for children and adults, these two questionnaires differ in some items, but both collect basic information on health status, use of health care services, health conditions, and health behaviors.

Sample Size and Response Rate. The NHIS sample size varies from year to year. It may be reduced for budgetary reasons or may be augmented if supplementary funding is available. Between 1997 and 2005, the sample numbered about 100,000 persons annually, with about 30,000–36,000 persons participating in the Sample Adult and about 12,000–14,000 in the Sample Child questionnaires. In the

2006–2015 redesign, the NHIS sample was reduced by 13% compared with the 1995–2005 design. With four sample panels and no sample cuts or augmentations, the expected annual NHIS sample size (completed interviews) during survey years 2006–2010 was on average 37,000 households containing about 81,000 persons.

In 2011–2014, the NHIS sample size was augmented in 32 states and D.C. The main goal of the augmentation was to increase the number of states for which reliable state-level estimates can be made. In 2011, the sample size was augmented by approximately 13%; in 2012, by approximately 21%; in 2013, by approximately 18%; and in 2014, by approximately 28%. In 2014, the sample numbered 112,053 persons, with 36,697 persons participating in the Sample Adult and 13,380 in the Sample Child questionnaires. In 2014 the total household response rate was 74%. The final response rate was 59% for the Sample Adult file and 67% for the Sample Child file.

Issues Affecting Interpretation. In 1997, the questionnaire was redesigned: some basic concepts were changed, and other concepts were measured in different ways. For some questions there was a change in the reference period. Also in 1997, the collection methodology changed from paper-and-pencil questionnaires to computer-assisted personal interviewing (CAPI). Because of the major redesign of the questionnaire in 1997, most NHIS trend tables in *Health, United States* begin with 1997 data. Starting with *Health, United States, 2005*, estimates for 2000–2002 were revised to use 2000-based weights and differ from previous editions of *Health, United States* that used 1990-based weights for those data years. The weights available on the public-use NHIS files for 2000–2002 are 1990-based. Data for 2003–2011 use weights derived from the 2000 census. Data for 2012 and beyond use weights derived from the 2010 census. In 2006–2010, the sample size was reduced, and this is associated with slightly larger variance estimates than in other years when a larger sample was fielded. Starting in 2010, a geographic nonresponse adjustment was made to both the sample adult weight and the sample child weight. See Moriarity (2009).

References

Moriarity C. 2009 National Health Interview Survey sample adult and sample child nonresponse bias analysis. Hyattsville, MD: NCHS; 2010. Available from: http://www.cdc.gov/nchs/data/nhis/nr_bias_analysis_report_2009_NHIS.pdf.

Parsons VL, Moriarity C, Jonas K, et al. Design and estimation for the National Health Interview Survey, 2006–2015. NCHS. Vital Health Stat 2(165); 2014.

For More Information. See the NHIS website at: <http://www.cdc.gov/nchs/nhis.htm>.

National HIV Surveillance System

CDC/National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention (NCHHSTP)

Overview. Human immunodeficiency virus (HIV) surveillance data are used to detect and monitor cases of HIV infection in the United States, identify epidemiologic trends, identify unusual cases requiring follow-up, and inform public health efforts to prevent and control the disease. Data collected on persons diagnosed with HIV infection include age, sex, race, ethnicity, mode of exposure, and geographic region.

Coverage. All 50 states, D.C., and six U.S. dependent areas (American Samoa, Guam, Northern Mariana Islands, Puerto Rico, Republic of Palau, and the U.S. Virgin Islands) report confirmed diagnoses of HIV infection to CDC using a uniform surveillance case definition and case report form. As of April 2008, all reporting areas had implemented confidential, name-based HIV infection reporting and agreed to participate in CDC's National HIV Surveillance System. *Health, United States* only presents data for the 50 states and D.C.

Methodology. HIV surveillance is conducted by health departments in each state or dependent area and D.C. Using a standard confidential case report form, the health departments collect information that is then transmitted electronically, without personal identifiers, to CDC.

The statistical adjustment of data on diagnoses of HIV infection (including stage 3, AIDS) is based on estimates of reporting-delay distributions, which are calculated by using a modified semiparametric life table statistical procedure. This procedure takes into account differences in reporting delays due to sex, race/ethnicity, and HIV transmission categories; reporting city, state, or territory; geographic region; size of the metropolitan statistical area; and type of facility where the diagnosis was made. HIV surveillance data are provisional and are updated annually.

Issues Affecting Interpretation. Although the completeness of reporting of cases of HIV infection to state and local health departments differs by geographic region and patient population, studies conducted by state and local health departments indicate that the reporting of cases of HIV infection in most areas of the United States is more than 80% complete. To assess trends in cases of HIV infection and deaths, it is preferable to use case data adjusted for reporting delays and presented by year of diagnosis, rather than straight counts of cases presented by year of report.

Reference

CDC. HIV surveillance report. Atlanta, GA; [published annually]. Available from: <http://www.cdc.gov/hiv/library/reports/surveillance/index.html>.

For More Information. See the NCHHSTP website at: <http://www.cdc.gov/nchhstp>.

National Hospital Ambulatory Medical Care Survey (NHAMCS)

CDC/NCHS

Overview. NHAMCS provides national data on the provision and use of medical care services in hospital emergency and outpatient departments, using information collected from medical records. Data are collected on types of providers seen; reason for visit; diagnoses; drugs ordered, provided, or continued; and selected procedures and tests performed during the visit. Patient data include age, sex, race, and expected source of payment. Data are also collected on selected characteristics of the hospitals included in the survey.

Coverage. NHAMCS covers visits to emergency departments (EDs) and outpatient departments (OPDs) of nonfederal, short-stay, or general hospitals in the United States. Telephone contacts are excluded. Starting in 2009, the survey includes visits to hospital-based ambulatory surgery centers (ASCs). Starting in 2010, visits to freestanding ASCs are included.

Methodology. The four-stage probability sample design used in NHAMCS involves samples of (a) geographically defined primary sampling units (PSUs), (b) hospitals within PSUs, (c) clinics or emergency service areas within OPDs or EDs, and (d) patient visits within clinics or emergency service areas. EDs are treated as their own stratum, and all service areas within EDs are included. The first-stage sample of NHAMCS consists of 112 PSUs selected from 1,900 such units that make up the United States. Within PSUs, 600 general and short-stay hospitals were sampled and assigned to 1 of 16 panels. In any given year, 13 panels are included. Each panel is assigned to a 4-week reporting period during the survey year.

In the NHAMCS OPD, a clinic is defined as an administrative unit of the OPD in which ambulatory medical care is provided under the supervision of a physician. Clinics where only ancillary services (e.g., radiology, laboratory services, physical rehabilitation, renal dialysis, and pharmacy) are provided, or other settings in which physician services are not typically provided, are considered out of scope. If a hospital OPD has five or fewer in-scope clinics, all are included in the sample. If an OPD has more than five clinics, the clinics are assigned to one of six specialty groups: general medicine, surgery, pediatrics, obstetrics and gynecology, substance abuse, and other. Within these specialty groups, clinics are grouped into clinic sampling units (SUs). A clinic SU is generally one clinic, except when a clinic expects fewer than 30 visits. In that case, it is grouped with one or more other clinics to form a clinic SU. If the grouped SU is selected, all clinics included in that SU are included in the sample. Prior to 2001, generally a sample of five clinic SUs was selected per hospital, based on probability proportional to the total expected number of patient visits to the clinic during the assigned 4-week

reporting period. Starting in 2001, clinic sampling within each hospital was stratified. If an OPD had more than five clinics, two clinic SUs were selected from each of the six specialty groups with a probability proportional to the total expected number of visits to the clinic. The change was made to ensure that at least two SUs were sampled from each of the specialty group strata.

The U.S. Census Bureau acts as the data collection agent for NHAMCS. Census field representatives contact sample hospitals to determine whether they have a 24-hour ED or an OPD that offers physician services. Visits to eligible EDs and OPDs are systematically sampled over the 4-week reporting period such that about 100 ED encounters and about 150–200 OPD encounters are selected. Hospital staff are asked to complete patient record forms (PRFs) for each sampled visit, but census field representatives typically abstract data for approximately two-thirds of these visits.

Sample data are weighted to produce national estimates. The estimation procedure used in NHAMCS has three basic components: inflation by the reciprocal of the probability of selection, adjustment for nonresponse, and population weighting ratio adjustment.

Sample Size and Response Rate. In any given year, the hospital sample consists of approximately 500 hospitals, of which 80% have EDs and about one-half have eligible OPDs. Typically, about 1,000 clinics are selected from participating hospital OPDs.

In 2011, the number of PRFs completed for EDs was 31,084 and for OPDs was 32,233, and the hospital response rate was 80% for EDs and 67% for OPDs.

Issues Affecting Interpretation. The NHAMCS PRF is modified approximately every 2 to 4 years to reflect changes in physician practice characteristics, patterns of care, and technological innovations. Examples of recent changes include an increase in the number of drugs recorded on the PRF and adding checkboxes for specific tests or procedures performed.

Reference

McCaig LF, McLemore T. Plan and operation of the National Hospital Ambulatory Medical Care Survey. NCHS. Vital Health Stat 1994;1(34). Available from: http://www.cdc.gov/nchs/data/series/sr_01/sr01_034acc.pdf.

For More Information. See the National Health Care Surveys website at: <http://www.cdc.gov/nchs/dhcs.htm> and the Ambulatory Health Care Data website at: <http://www.cdc.gov/nchs/ahcd.htm>.

National Immunization Survey (NIS)

CDC/National Center for Immunization and Respiratory Diseases (NCIRD) and NCHS

Overview. NIS is a continuing nationwide telephone sample survey to monitor vaccination coverage rates among children aged 19–35 months and among teenagers (NIS–Teen) aged 13–17. Data collection for children aged 19–35 months started in 1994, and data collection for teenagers aged 13–17 started in 2006.

Coverage. Children aged 19–35 months and adolescents aged 13–17 in the civilian noninstitutionalized population are represented in this survey. Estimates of vaccine-specific coverage are available for the nation, states, and selected local areas.

Methodology. NIS is a nationwide telephone sample survey of households with age-eligible children. The survey uses a two-phase sample design. First, a random-digit-dialing sample of telephone numbers is drawn. When households with age-eligible children are contacted, the interviewer collects information on the vaccinations received by all age-eligible children and obtains permission to contact the children's vaccination providers. Second, identified providers are sent vaccination history questionnaires by mail. Final weighted estimates are adjusted for households without telephones and for nonresponse. All vaccination coverage estimates are based on provider-reported vaccination histories. NIS–Teen followed the same sample design and data collection procedures as NIS except that only one age-eligible adolescent was selected from each screened household for data collection.

Starting in 2011, the NIS sampling frame was expanded from a single-landline frame to dual-landline and cellular telephone sampling frames. This change increased the representativeness of the sample characteristics but had little effect on the final 2011 NIS and NIS–Teen national estimates of vaccination coverage overall and when stratified by poverty status. See details of the dual-frame sample design in the annual NIS Data User's Guide on the NIS Website. Available from: http://www.cdc.gov/nchs/nis/data_files.htm.

Sample Size and Response Rate. In 2014, the Council of American Survey Research Organizations (CASRO) response rate for the NIS landline sample was 62.6%, and 33.5% for the cellular telephone sample. Of the 11,198 age-eligible children with completed household interviews from the landline sample, 7,093 (63.3%) had adequate provider data. From the cellular telephone sample, 7,800 (58.9%) of the 13,233 eligible children with completed household interviews had adequate provider data.

The CASRO response rate for the 2014 NIS–Teen landline sample was 60.3%, and 31.2% for the cellular telephone sample. Of the 19,705 age-eligible adolescents with completed household interviews from the landline sample,

11,243 (57.1%) had adequate provider data. From the cellular telephone sample, 9,584 (52.3%) of the 18,342 eligible adolescents with completed household interviews had adequate provider data.

Issues Affecting Interpretation. Starting with *Health, United States, 2015*, estimates are from the NIS website and may differ slightly from estimates published previously in *Morbidity and Mortality Weekly Report* (MMWR) articles.

The findings in recent years are subject to several limitations. Data year 2011 was the first year that NIS and NIS–Teen used a dual-frame sampling scheme that included landline and cellular telephone households. Estimates from 2011 and subsequent years might not be comparable with those from prior to 2011 when surveys were conducted via landline telephone only. NIS is a telephone survey, and statistical adjustments might not compensate fully for nonresponse and for households without landline telephones prior to 2011. Underestimates of vaccination coverage might have resulted in exclusive use of provider-reported vaccination histories because completeness of records is unknown. Finally, although national coverage estimates are precise, annual estimates and trends for state and local areas should be interpreted with caution because of smaller sample sizes and wider confidence intervals.

Before January 2009, NIS did not distinguish between Hib vaccine production types; therefore, children who received three doses of a vaccine product that requires four doses were misclassified as fully vaccinated. For more information, see “Changes in measurement of *Haemophilus influenzae* serotype b (Hib) vaccination coverage—National Immunization Survey, United States, 2009. MMWR 2010;59:1069–72.”

Starting in 2014, NIS–Teen defined an adolescent's vaccination record as having adequate provider data if that adolescent had vaccination history data from one or more of the named vaccination providers, or if the parent reported that the adolescent was completely unvaccinated. Prior to 2014, the adequate provider data definition had more criteria, and it was based on a comparison of provider report of vaccination history with parental report of vaccination history, either by shot card report or recall.

To assess the effect of the change in the adequate provider definition criteria on vaccination coverage estimates, NIS recomputed estimates from the 2006–2013 survey. In general, 2013 NIS–Teen vaccination coverage estimates using the revised adequate provider data definition were different, and generally lower, than original 2013 NIS–Teen estimates. Differences between revised and original 2013 national vaccination estimates ranged from –0.1 percentage point to –2.2 percentage points. For more information on the revised adequate provider data criteria, see: <http://www.cdc.gov/vaccines/imz-managers/coverage/nis/teen/apd-report.html>, and for revised 2013 estimates based on the 2014 criteria, see: CDC. National, regional, state, and selected local area vaccination coverage among

adolescents aged 13–17 years—United States, 2014. MMWR 2015;64(29):784–92. Available from: <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6429a3.htm>. Because of the revision in the adequate provider definition, NIS–Teen vaccination coverage estimates for 2013 and beyond cannot be directly compared with previously published 2006–2013 NIS–Teen survey vaccination coverage estimates based on the previous adequate provider definition.

References

CDC. National, state, and selected local area vaccination coverage among children aged 19–35 months—United States, 2014. MMWR 2015;64(33):889–96. Available from: <http://www.cdc.gov/mmwr/pdf/wk/mm6433.pdf>.

CDC. National, regional, state, and selected local area vaccination coverage among adolescents aged 13–17 years—United States, 2014. MMWR 2015;64(29):784–92. Available from: <http://www.cdc.gov/mmwr/pdf/wk/mm6429.pdf>.

Smith PJ, Hoaglin DC, Battaglia MP, et al. Statistical methodology of the National Immunization Survey, 1994–2002. NCHS. Vital Health Stat 2005;2(138). Available from: http://www.cdc.gov/nchs/data/series/sr_02/sr02_138.pdf.

CDC. Announcement: Addition of households with only cellular telephone service to the National Immunization Survey, 2011. Available from: http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6134a5.htm?s_cid=mm6134a5_e%0d%0a.

CDC. Changes in measurement of *Haemophilus influenzae* serotype b (Hib) vaccination coverage—National Immunization Survey, United States, 2009. MMWR 2010;59(33):1069–72. Available from: http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5933a3.htm?s_cid=mm5933a3_e%0d%0a.

For More Information. See the NIS website at: <http://www.cdc.gov/nchs/nis.htm>.

National Income and Product Accounts (NIPA)

Bureau of Economic Analysis (BEA)

Overview. NIPA are a set of economic accounts that provide detailed measures of the value and composition of national output and the incomes generated in the production of that output. Essentially, NIPA provide a detailed snapshot of the myriad transactions that make up the economy—buying and selling goods and services, hiring of labor, investing, renting property, paying taxes, and the like. NIPA estimates show U.S. production, distribution, consumption, investment, and saving.

The best-known NIPA measure is the gross domestic product (GDP), which is defined as the market value of the goods

and services produced by labor and property located in the United States. NIPA calculate GDP as the sum of familiar final expenditure components: personal consumption expenditures, private investment, government spending (consumption and investment), and net exports. However, GDP is just one of many economic measures presented in NIPA. Another key NIPA indicator presented in *Health, United States* is the implicit price deflator for GDP.

The conceptual framework of NIPA is illustrated by seven summary accounts: the domestic income and product account, the private enterprise income account, the personal income and outlay account, the government receipts and expenditures account, the foreign transactions current account, the domestic capital account, and the foreign transactions capital account. These summary accounts record a use (or expenditure) in one account for one sector and a corresponding source (or receipt) in an account of another sector or of the same sector. This integrated system provides a comprehensive measure of economic activity in a consistently defined framework without double counting.

Coverage. Source data for NIPA domestic estimates cover all 50 states and D.C. The U.S. national income and product statistics were first presented as part of a complete and consistent double-entry accounting system in the summer of 1947.

Methodology. NIPA estimates are revised on a quarterly, annual, and quinquennial basis. For GDP and most other NIPA series, a set of three current quarterly estimates is released each year. Quarterly estimates provide the first look at the path of U.S. economic activity. Annual revisions of NIPA are usually carried out each summer. These revisions incorporate source data that are based on more extensive annual surveys, on annual data from other sources, and on later revisions to the monthly and quarterly source data, and they generally cover the three previous calendar years. Comprehensive revisions are carried out at about 5-year intervals and may result in revisions that extend back many years. These estimates incorporate all of the best available source data, such as data from the quinquennial U.S. Economic Census.

NIPA measures are built up from a wide range of source data using a variety of estimating methods. To ensure consistency and accuracy, NIPA use various adjustment and estimation techniques to estimate data. Three general types of adjustments are made to the source data that are incorporated into the NIPA estimates. The first consists of adjustments that are needed so that the data conform to appropriate NIPA concepts and definitions. The second type of adjustment involves filling gaps in coverage. The third type of adjustment involves time of recording and valuation. Source data must occasionally be adjusted to account for special circumstances that affect the accuracy of the data. For example, quarterly and monthly NIPA estimates are seasonally adjusted at the detailed-series level when the series demonstrate statistically significant seasonal patterns.

Source data may also be used as indicators to extrapolate annual estimates. For more information, see “An introduction to the National Income and Product Accounts methodology papers: U.S. National Income and Product Accounts,” available from: http://www.bea.gov/scb/pdf/national/nipa/methpap/mpi1_0907.pdf; and “Concepts and methods of the U.S. National Income and Product Accounts,” available from: <http://www.bea.gov/national/pdf/NIPAhandbookch1-4.pdf>.

Issues Affecting Interpretation. NIPA source data and estimates are revised frequently. Data are released at different times, estimates are updated as they become available, new concepts and definitions are incorporated, and source data may change due to improvements in collection and new methodologies. As a result, major estimates such as GDP and its major components undergo frequent revision, and historical data are changed. For more information, see the BEA (NIPA) website at: http://www.bea.gov/scb/pdf/2013/03%20March/0313_nipa_comprehensive_revision_preview.pdf.

Reference

U.S. Bureau of Economic Analysis. A guide to the National Income and Product Accounts of the United States. Washington, DC: BEA; 2006. Available from: <http://www.bea.gov/national/pdf/nipaguid.pdf>.

For More Information. See the BEA (NIPA) website at: <http://www.bea.gov/national/index.htm>.

National Medical Expenditure Survey (NMES)—See Appendix I, Medical Expenditure Panel Survey (MEPS).

National Notifiable Diseases Surveillance System (NNDSS)

CDC

Overview. The CDC National Notifiable Diseases Surveillance System (NNDSS) is a nationwide collaboration that enables all levels of public health (local, state, territorial, federal, and international) to share health information to monitor, control, and prevent the occurrence and spread of state-reportable and nationally notifiable infectious and some noninfectious diseases and conditions. NNDSS is a multifaceted program that includes the surveillance system for collection, analysis, and sharing of health data, resources, and information about policies and standards, at the local, state, and national levels. NNDSS provides weekly provisional and annual finalized information on the occurrence of diseases defined as notifiable by the Council of State and Territorial Epidemiologists (CSTE). Data include incidence of reportable diseases, which are nationally notifiable using uniform surveillance case definitions.

Coverage. Notifiable disease reports are received from health departments in the 50 states, five territories, D.C., and New York City. Policies for reporting notifiable disease cases can vary by disease or reporting jurisdiction, depending on case status classification (i.e., confirmed, probable, or suspect).

Methodology. CDC, in partnership with CSTE, administers NNDSS. Reportable disease surveillance is conducted by public health practitioners at local, state, and national levels to support disease prevention and control. Data on a subset of reportable conditions that have been designated nationally notifiable are then submitted to CDC without personal identifiers. The system also provides annual summaries of the finalized data. CSTE and CDC annually review the status of national infectious disease surveillance and recommend additions or deletions to the list of nationally notifiable diseases, based on the need to respond to emerging priorities. For example, Q fever and tularemia became nationally notifiable in 2000. However, reporting nationally notifiable diseases to CDC is voluntary. Because reporting is currently mandated by law or regulation only at the local and state levels, the list of diseases that are considered reportable varies by state. For example, reporting of cyclosporiasis to CDC is not done by some states in which this disease is not reportable to local or state authorities.

State epidemiologists report cases of nationally notifiable diseases to CDC, which tabulates and publishes these data in *Morbidity and Mortality Weekly Report* (MMWR) and in *Summary of Notifiable Diseases, United States* (before 1985, titled *Annual Summary*).

Issues Affecting Interpretation. NNDSS data must be interpreted in light of reporting practices. Some diseases that cause severe clinical illness (for example, plague and rabies) are likely reported accurately if diagnosed by a clinician. However, persons who have diseases that are clinically mild and infrequently associated with serious consequences (e.g., salmonellosis) may not seek medical care from a health care provider. Even if these less severe diseases are diagnosed, they are less likely to be reported.

The degree of completeness of data reporting is also influenced by the diagnostic facilities available, the control measures in effect, public awareness of a specific disease, and the interests, resources, and priorities of state and local officials responsible for disease control and public health surveillance. Finally, factors such as changes in case definitions for public health surveillance, introduction of new diagnostic tests, or discovery of new disease entities can cause changes in disease reporting that are independent of the true incidence of disease.

Reference

CDC. Summary of notifiable diseases—United States, 2013. MMWR 2015;62(53):1–119. Available from: http://www.cdc.gov/mmwr/mmwr_nd/index.html.

For More Information. See the NNDSS website at: <http://wwwn.cdc.gov/nndss/>.

National Survey of Family Growth (NSFG)

CDC/NCHS

Overview. NSFG gathers information on family life, marriage and divorce, pregnancy, infertility, use of contraception, and men's and women's health. NSFG provides national data on factors affecting birth and pregnancy rates, adoption, and maternal and infant health. Data collected include sexual activity, marriage, divorce and remarriage, unmarried cohabitation, forced sexual intercourse, contraception and sterilization, infertility, breastfeeding, pregnancy loss, low birthweight, and use of medical care for family planning and infertility.

Coverage. NSFG in 1982, 1988, and 1995 included women aged 15–44 in the civilian noninstitutionalized population of the U.S. The 2002, 2006–2010, and 2011–2013 NSFG included both men and women aged 15–44 in the household population of the United States. The household population of the United States refers to the civilian noninstitutionalized population and active-duty military personnel who are not living on military bases.

Methodology. The 2006–2010 and 2011–2013 NSFG sample design consisted of five stages of selection: primary sampling units (PSUs), blocks or segments, housing units, one eligible person per housing unit, and housing units or persons for phase 2 data collection (intended to raise response rates and correct any bias due to nonresponse). Samples of 110 PSUs were drawn for 2006–2010, and 117 PSUs were drawn for 2011–2015. Both PSU samples were divided into four fully representative national samples. Interviewing was done for 1 year in each subsample; the entire 110-PSU design was completed in the 4-year period, 2006–2010. The 2011–2013 file contained interviews from 65 PSUs, equal to two of the four subsamples drawn for the 2011–2015 NSFG. The interviews were administered in person by trained female interviewers using a laptop or notebook computer with computer-assisted personal interviewing (CAPI) or audio computer-assisted self-interview (ACASI) programs.

In all survey years, black women were sampled at higher rates than white women so that more reliable statistics could be produced for black women. In both the 1995 and 2002 surveys, Hispanic persons were also oversampled. In the 2006–2010 and 2011–2013 NSFG, black and Hispanic adults and all 15- to 19-year-olds were oversampled.

To produce national estimates from the sample for the millions of women aged 15–44 in the United States, data for the interviewed sample women were (a) inflated by the reciprocal of the probability of selection at each stage of sampling (for example, if there was a 1 in 5,000 chance that a woman would be selected for the sample, her sampling weight was 5,000); (b) adjusted for nonresponse; and (c) poststratified, or aligned with benchmark population sizes based on data from the U.S. Census Bureau.

Sample Size and Response Rate. For the 1982, 1995, 2002, and 2006–2010 surveys, the response rate ranged from 78%–80%. The response for the 2011–2013 survey was 73%. Sample sizes have varied over the surveys: in 1982 the sample size was 7,969; in 1995 it was 10,847; in 2002 it was 7,643; in 2006–2010 it was 12,279; and in 2011–2013 it was 5,601.

References

Bachrach CA, Horn MC, Mosher WD, Shimizu I. National Survey of Family Growth, cycle III: Sample design, weighting, and variance estimation. *NCHS. Vital Health Stat* 1985;2(98). Available from: http://www.cdc.gov/nchs/data/series/sr_02/sr02_098.pdf.

Judkins DR, Mosher WD, Botman S. National Survey of Family Growth: Design, estimation, and inference. *NCHS. Vital Health Stat* 1991;2(109). Available from: http://www.cdc.gov/nchs/data/series/sr_02/sr02_109.pdf.

Kelly JE, Mosher WD, Duffer AP, Kinsey SH. Plan and operation of the 1995 National Survey of Family Growth. *NCHS. Vital Health Stat* 1997;1(36). Available from: http://www.cdc.gov/nchs/data/series/sr_01/sr01_036.pdf.

Potter FJ, Iannacchione VG, Mosher WD, et al. Sample design, sampling weights, imputation, and variance estimation in the 1995 National Survey of Family Growth. *NCHS. Vital Health Stat* 1998;2(124). Available from: http://www.cdc.gov/nchs/data/series/sr_02/sr02_124.pdf.

Groves RM, Benson G, Mosher WD, et al. Plan and operation of cycle 6 of the National Survey of Family Growth. *NCHS. Vital Health Stat* 2005;1(42). Available from: http://www.cdc.gov/nchs/data/series/sr_01/sr01_042.pdf.

Groves RM, Mosher WD, Lepkowski J, Kirgis NG. Planning and development of the continuous National Survey of Family Growth. *NCHS. Vital Health Stat* 2009;1(48). Available from: http://www.cdc.gov/nchs/data/series/sr_01/sr01_048.pdf.

Lepkowski JM, Mosher WD, Davis KE, et al. The 2006–2010 National Survey of Family Growth: Sample design and analysis of a continuous survey. *NCHS. Vital Health Stat* 2010;2(150). Available from: http://www.cdc.gov/nchs/data/series/sr_02/sr02_150.pdf.

Lepkowski JM, Mosher WD, Groves RM, et al. Responsive design, weighting, and variance estimation in the 2006–2010 National Survey of Family Growth. *NCHS. Vital Health Stat* 2013;2(158). Available from: http://www.cdc.gov/nchs/data/series/sr_02/sr02_158.pdf.

For More Information. See the NSFG website at: <http://www.cdc.gov/nchs/nsfg.htm>.

National Survey on Drug Use & Health (NSDUH)

Substance Abuse and Mental Health Services Administration (SAMHSA)

Overview. NSDUH, formerly called the National Household Survey on Drug Abuse (NHSDA), collects data on substance use, abuse, and dependence; mental health problems; and receipt of substance abuse and mental health treatment. NSDUH reports on the prevalence, incidence, and patterns of drug and alcohol use and abuse in the general U.S. civilian noninstitutionalized population aged 12 and over. NSDUH also reports on substance use disorders, substance use treatment, health care, mental health disorders, and mental health service utilization.

Coverage. NSDUH is representative of persons aged 12 and over in the civilian noninstitutionalized population of the United States, and in each state and D.C.

The survey covers residents of households (including those living in houses, townhouses, apartments, and condominiums), persons in noninstitutional group quarters (including those in shelters, boarding houses, college dormitories, migratory work camps, and halfway houses), and civilians living on military bases. Persons excluded from the survey include homeless people who do not use shelters, active military personnel, and residents of institutional group quarters such as jails and hospitals.

Methodology. The data collection method is in-person interviews conducted with a sample of individuals at their place of residence. Computer-assisted interviewing (CAI) methods, including audio computer-assisted self-interviewing (ACASI), are used to provide a private and confidential setting to complete the interview.

NSDUH uses a 50-state (and D.C.) sample design that is revised periodically. In 2014, NSDUH introduced an independent multistage area probability sample within each state and D.C. States are the first level of stratification. Each state was stratified into approximately equally populated state sampling regions (SSRs), and then census tracts within each SSR were selected, census block groups within census tracts, and area segments (i.e., a collection of census blocks) within census block groups. Finally, dwelling units (DUs) were selected within segments, and within each selected DU, up to two residents who were at least 12 years old were selected for the interview.

In addition, in 2014, changes were made in the sample sizes allocated to each state and to different age groups, in order to increase the precision of national and many state estimates as well as estimates for older adults. In particular, samples sizes were increased in the 12 most populous states. States with sample increases will have more precise estimates than in previous years, whereas states with smaller sample sizes will have some reductions in precision. However, all states will still have reasonable levels of

precision. This allocation of sample to states is also thought to be more cost-efficient. Starting in 2014, the sample size was redistributed by age group so that 25% of the sample is allocated to those aged 12–17, 25% to those aged 18–25, and 50% to those aged 26 or older. Although the sample sizes for age groups 12–17 and 18–25 were reduced, these two groups are still considered to be oversampled since they represent approximately 10% and 13% of the total population, respectively.

Sample Size and Response Rate. Nationally, 127,605 household addresses were successfully screened for the 2014 survey, conducted from January to December 2014. In these screened households, a total of 91,640 sample persons were selected, from which 67,901 completed interviews were obtained, including 17,046 interviews from adolescents aged 12–17 and 50,855 interviews from adults aged 18 or over. Weighted response rates were 82% for household screening and 71% for interviewing.

Issues Affecting Interpretation. Several improvements to the survey were implemented in 2002, when the survey was redesigned as NSDUH. In addition to the name change, respondents were offered a \$30 incentive payment for participation in the survey starting in 2002, and quality control procedures for data collection were enhanced in 2001 and 2002. Because of these improvements and modifications, estimates from NSDUH completed in 2002 and later should not be compared with estimates from the 2001 or earlier versions of the survey. The data collected in 2002 represent a new baseline for tracking trends in substance use and other measures. Special questions on methamphetamine were added in 2005 and 2006. Data for years prior to 2007 were adjusted for comparability. Starting with 2011 data, 2010-census based control totals were used in the weighting process. Analysis weights in the 2002 through 2010 NSDUHs were derived from the 2000 census data. This reweighting to the 2010 census data could affect comparisons between estimates for 2011 and subsequent years and those from prior years. An analysis of the impact of reweighting showed that the percentages of substance users were largely unaffected. For more information, see: <http://www.samhsa.gov/data/NSDUH/NSDUHCensusEffects/Index.aspx>.

Estimates of substance use for youth based on NSDUH are not directly comparable with estimates based on the Monitoring the Future (MTF) Study and the Youth Risk Behavior Survey (YRBS). In addition to the fact that MTF excludes dropouts and absentees, rates are not directly comparable across these surveys because of differences in the populations covered, sample design, questionnaires, and interview setting. NSDUH collects data in residences, whereas MTF and YRBS collect data in school classrooms. Further, NSDUH estimates are tabulated by age, whereas MTF and YRBS estimates are tabulated by grade, representing different ages as well as different populations.

References

Substance Abuse and Mental Health Services Administration. 2014 National Survey on Drug Use and Health: Methodological summary and definitions. Rockville, MD: SAMHSA; 2015. Available from: <http://www.samhsa.gov/data/sites/default/files/NSDUH-MethodSummDefs2014/NSDUH-MethodSummDefs2014.pdf>.

Substance Abuse and Mental Health Services Administration. Results from the 2014 National Survey on Drug Use and Health: Detailed tables. Rockville, MD: SAMHSA; 2015. Available from: <http://www.samhsa.gov/data/sites/default/files/NSDUH-DetTabs2014/NSDUH-DetTabs2014.htm>.

For More Information. See the NSDUH website at: <http://www.samhsa.gov/data/population-data-nsduh> and the Center for Behavioral Health Statistics and Quality (the data collection agency) website at: <http://www.samhsa.gov/about-us/who-we-are/offices-centers/cbhsq>.

National Vital Statistics System (NVSS)

CDC/NCHS

Overview. NVSS collects and publishes official national statistics on births, deaths, fetal deaths, and, prior to 1996, marriages and divorces occurring in the United States, based on U.S. Standard Certificates. Fetal deaths are classified and tabulated separately from other deaths. The vital statistics files—Birth, Fetal Death, Mortality Multiple Cause-of-Death, Linked Birth/Infant Death, and Compressed Mortality—are described in detail below.

Coverage. NVSS collects and presents U.S. resident data for the aggregate of 50 states, New York City, and D.C., as well as for each individual state and D.C. and the U.S. dependent areas of Puerto Rico, Virgin Islands, Guam, American Samoa, and Northern Marianas. Vital events occurring in the United States to non-U.S. residents, and vital events occurring abroad to U.S. residents, are excluded. Starting with *Health, United States, 2013*, information on vital events for Puerto Rico, Virgin Islands, Guam, American Samoa, and Northern Marianas is shown in selected tables that show data by state, but are not included in U.S. totals.

Methodology. NCHS' Division of Vital Statistics obtains information on births and deaths from the registration offices of each of the 50 states, New York City, D.C., Puerto Rico, Virgin Islands, Guam, American Samoa, and Northern Marianas. Until 1972, microfilm copies of all death certificates and a 50% sample of birth certificates were received from all registration areas and processed by NCHS. In 1972, some states began sending their data to NCHS through the Cooperative Health Statistics System (CHSS). States that participated in the CHSS program processed 100% of their death and birth records and sent the entire data file to NCHS on computer tapes. Currently, data are sent

to NCHS through the Vital Statistics Cooperative Program (VSCP), following the same procedures as with CHSS. The number of participating states grew from 6 in 1972 to 46 in 1984. Starting in 1985, all 50 states and D.C. participated in VSCP.

U.S. Standard Certificates. U.S. Standard Certificates of Live Birth and Death and Fetal Death Reports are revised periodically, allowing evaluation and addition, modification, and deletion of items. Beginning with 1989, revised Standard Certificates replaced the 1978 versions. The 1989 revision of the birth certificate included items to identify the Hispanic parentage of newborns and to expand information about maternal and infant health characteristics. The 1989 revision of the death certificate included items on educational attainment and Hispanic origin of decedents, as well as changes to improve the medical certification of cause of death. Standard Certificates recommended by NCHS are modified in each registration area to serve the area's needs. However, most certificates conform closely in content and arrangement to the Standard Certificate, and all certificates contain a minimum data set specified by NCHS. The 2003 revision of vital records went into effect in some states and territories beginning in 2003, but full implementation in all states and territories will be phased in over several years. The 2003 revision of the birth certificate included changes in ascertainment of education level, prenatal care, and tobacco use during pregnancy. The 2003 revision of the death certificate included changes in the ascertainment of multiple races, education level, tobacco use, and maternal mortality.

Birth File

Overview. Vital statistics natality data are a fundamental source of demographic, geographic, and medical and health information on all births occurring in the United States. This is one of the few sources of comparable health-related data for small geographic areas over an extended time period. The data are used to present the characteristics of babies and their mothers, track trends such as birth rates for teenagers, and compare natality trends with those in other countries.

The Birth file includes characteristics of the baby, such as sex, birthweight, and weeks of gestation; demographic information about the parents, such as age, race, Hispanic origin, parity, educational attainment, marital status, and state of residence; medical and health information, such as prenatal care, based on hospital records; and behavioral risk factors for the birth, such as mother's tobacco use during pregnancy.

Coverage. Birth data presented in *Health, United States* are based on reporting from all 50 states and D.C. Data for Puerto Rico, Virgin Islands, Guam, American Samoa, and Northern Marianas are shown in selected state tables but are not included in U.S. totals. Beginning with 1970, births to nonresidents of the United States are excluded.

Methodology. In the United States, state laws require birth certificates to be completed for all births. The registration of births is the responsibility of the professional attendant at birth, generally a physician or midwife. The birth certificate must be filed with the local registrar of the district in which the birth occurs. Each birth must be reported promptly; the reporting requirements vary from state to state, ranging from 24 hours to as much as 10 days after the birth.

Federal law mandates national collection and publication of birth and other vital statistics data. NVSS is the result of cooperation between NCHS and the states to provide access to statistical information from birth certificates. Standard forms for the collection of the data, and model procedures for the uniform registration of the events, are developed and recommended for state use through cooperative activities of the states and NCHS. NCHS shares the costs incurred by the states in providing vital statistics data for national use.

Issues Affecting Interpretation. Two-thirds (66%) of all births in 2009, 76% in 2010, 83% in 2011, 86% in 2012, 90% in 2013, and 96% in 2014 were reported using the 2003 revision of the U.S. Standard Certificate of Live Birth. Interpretation of trend data should take into consideration changes to reporting areas. For methodological and reporting area changes for the following birth certificate items, see Appendix II, Age; Hispanic origin; Marital status; Race.

Reference

Hamilton BE, Martin JA, Osterman MJK, et al. Births: Final data for 2014. National vital statistics reports; vol 64 no 12. Hyattsville, MD: NCHS; 2015. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr64/nvsr64_12.pdf.

For More Information. See the Birth Data website at: <http://www.cdc.gov/nchs/births.htm>, and Vitalstats at: http://www.cdc.gov/nchs/data_access/Vitalstatsonline.htm.

Fetal Death Data Set

Overview. Fetal mortality refers to the intrauterine death of a fetus at any gestational age. In *Health, United States*, data are presented for fetal deaths at 20 weeks or more. The Fetal Death data set includes characteristics of the fetus, such as sex, birthweight, and weeks of gestation; demographic information about the mother, such as age, race, Hispanic origin, live-birth order, and marital status; and medical and health information, such as maternal diabetes and hypertension.

Coverage. Data presented in *Health, United States* are based on reporting from all 50 states and D.C. Data for Puerto Rico, Virgin Islands, Guam, American Samoa, and Northern Marianas are not included in U.S. totals but are included in the Fetal Death User Guide available from the NCHS website at: http://www.cdc.gov/nchs/data_access/VitalStatsOnline.htm, and in periodic reports.

Methodology. Fetal death means the death of a fetus prior to delivery from the mother, irrespective of the duration of pregnancy. Fetal deaths do not include induced terminations of pregnancy. This definition of fetal death, adopted by NCHS as the nationally recommended standard, is based on the definition published by the World Health Organization in 1950 and revised in 1988. The term fetal death encompasses other commonly used terms, including stillbirth, spontaneous abortion, and miscarriage. All U.S. states and registration areas have definitions similar to the standard definition, except for Puerto Rico and Wisconsin, which have no formal definition.

State laws require the reporting of fetal deaths, and federal law mandates national collection and publication of fetal death data. States and reporting areas submit fetal mortality data to NCHS as part of a cooperative agreement. Standard forms and procedures for the collection of the data are developed and recommended for state use through cooperative activities of the states and NCHS. NCHS shares the costs incurred by the states in providing vital statistics data for national use.

In addition to fetal mortality rates, perinatal mortality rates are also presented in *Health, United States*. Perinatal mortality includes both late fetal deaths (of at least 28 weeks of gestation) and early infant (neonatal) deaths (within 7 days of birth). Data on early infant deaths come from the Linked Birth/Infant Death data set.

Issues Affecting Interpretation. Reporting requirements for fetal deaths vary by state, and these differences have important implications for comparisons of fetal mortality rates by state. The majority of states require reporting of fetal deaths at 20 weeks of gestation or more, or a minimum of 350 grams birthweight (roughly equivalent to 20 weeks), or some combination of the two. However, seven states require reporting of fetal deaths at all periods of gestation, and one state requires reporting beginning at 16 weeks of gestation. Further, two states require the reporting of fetal deaths with birthweights of 500 grams or more (roughly equivalent to 22 weeks of gestation).

There is substantial evidence that not all fetal deaths for which reporting is required are, in fact, reported. Underreporting of fetal deaths is most likely to occur in the earlier part of the required reporting period for each state. For example, in 2013, for those states requiring reporting of fetal deaths at all periods of gestation, 56.4% of fetal deaths at 20 weeks of gestation or more were at 20–27 weeks, whereas for states requiring reporting of fetal deaths of 500 grams or more, only 33.8% were at 20–27 weeks, thus indicating substantial underreporting of early fetal deaths in some states.

References

MacDorman MF, Gregory ECW. Fetal and perinatal mortality: United States, 2013. National vital statistics reports; vol 64 no 8. Hyattsville, MD: NCHS; 2015.

Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr64/nvsr64_08.pdf.

Gregory ECW, MacDorman MF, Martin JA. Trends in fetal and perinatal mortality in the United States, 2006–2012. NCHS data brief, no 169. Hyattsville, MD: NCHS; 2014. Available from: <http://www.cdc.gov/nchs/data/databriefs/db169.htm>.

MacDorman MF, Kirmeyer SE, Wilson EC. Fetal and perinatal mortality, United States, 2006. National vital statistics report; vol 60 no 8. Hyattsville, MD: NCHS; 2012. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr60/nvsr60_08.pdf.

For More Information. See the NCHS Fetal Deaths data website at: http://www.cdc.gov/nchs/fetal_death.htm.

Mortality Multiple Cause-of-Death File

Overview. Vital statistics mortality data are a fundamental source of demographic, geographic, and underlying and multiple cause-of-death information. Multiple cause-of-death data reflect all medical information reported on death certificates and complement traditional underlying cause-of-death data. Multiple-cause data give information on diseases that are a factor in death, whether or not they are the underlying cause of death; on associations among diseases; and on injuries leading to death.

The Mortality multiple cause-of-death file includes demographic information on age, sex, race, Hispanic origin, state of residence, and educational attainment, as well as medical information on causes of death. This data set is one of the few sources of comparable health-related data for small geographic areas over an extended time period. The data are used to present the characteristics of those dying in the United States, to determine life expectancy, and to compare mortality trends with those in other countries.

Coverage. Mortality data presented in *Health, United States* are based on reporting from all 50 states and D.C. Data for Puerto Rico, Virgin Islands, Guam, American Samoa, and Northern Marianas are shown in selected state tables, but are not included in U.S. totals. Beginning with 1970, mortality statistics for the U.S. exclude deaths of nonresidents of the U.S. Mortality statistics for Puerto Rico, Virgin Islands, American Samoa, and Northern Marianas excluded deaths of nonresidents for each area. For Guam, mortality statistics exclude deaths that occurred to a resident of any place other than Guam or the U.S. (50 states and D.C.).

Methodology. By law, the registration of deaths is the responsibility of the funeral director. The funeral director obtains demographic data for the death certificate from an informant. The physician in attendance at the death is required to certify the cause of death. Where death is from other than natural causes, a coroner or medical examiner may be required to examine the body and certify the cause of death.

NCHS is responsible for compiling and publishing annual national statistics on causes of death. In carrying out this responsibility, NCHS adheres to the World Health Organization (WHO) Nomenclature Regulations. These regulations require (a) that cause of death be coded in accordance with the applicable revision of the *International Classification of Diseases* (ICD) (see Appendix II, *International Classification of Diseases* [ICD]; Table III); and (b) that underlying cause of death be selected in accordance with international rules. Traditionally, national mortality statistics have been based on a count of deaths, with one underlying cause assigned for each death.

Prior to 1968, mortality medical data were based on manual coding of an underlying cause of death for each certificate, in accordance with WHO rules. Starting with 1968, NCHS converted to computerized coding of the underlying cause and manual coding of all causes (multiple causes) on the death certificate. In this system, called Automated Classification of Medical Entities (ACME), multiple-cause codes serve as inputs to the computer software, which employs WHO rules to select the underlying cause. ACME is used to select the underlying cause of death for all death certificates in the United States, and cause-of-death data in *Health, United States* are coded using ACME.

In addition, NCHS has developed two computer systems as inputs to ACME. Beginning with 1990 data, the Mortality Medical Indexing, Classification, and Retrieval system (MICAR) was introduced to automate coding of multiple causes of death. MICAR provides more detailed information on the conditions reported on death certificates than is available through the ICD code structure. Then, beginning with data year 1993, SuperMICAR, an enhancement of MICAR, was introduced. SuperMICAR allows for literal entry of the multiple cause-of-death text as reported by the certifier. This information is then processed automatically by the MICAR and ACME computer systems. Records that cannot be processed automatically by MICAR or SuperMICAR are multiple-cause-coded manually and then further processed through ACME. Starting in 2003, SuperMICAR was used to process all of the nation's death records.

Data for the entire United States refer to events occurring within the United States; data for geographic areas are by place of residence. For methodological and reporting area changes for the following death certificate items, see Appendix II, Hispanic origin; Race.

Issues Affecting Interpretation. The ICD by which cause of death is coded and classified is revised approximately every 10–20 years. Because revisions of the ICD may cause discontinuities in trend data by cause of death, comparison of death rates by cause of death across ICD revisions should be done with caution and with reference to the comparability ratio. (See Appendix II, Comparability ratio.) Prior to 1999, modifications to the ICD were made only when a new revision of the ICD was implemented. A process for updating the ICD was introduced with the 10th revision

(ICD–10) that allows for midrevision changes. These changes, however, may affect comparability of data between years for select causes of death. Minor changes may be implemented every year, whereas major changes may be implemented every 3 years (e.g., 2003 data year). In data year 2006, major changes were implemented, including the addition and deletion of several ICD codes. For more information, see Heron et al. (2009).

Multiple-cause data were obtained from all certificates for 1968–1971, 1973–1980, and 1983–present. Data were obtained from a 50% sample of certificates for 1972. Multiple-cause data for 1981 and 1982 were obtained from a 50% sample of certificates from 19 registration areas. For the other states, data were obtained from all certificates.

The death certificate has been revised periodically. A revised U.S. Standard Certificate of Death was recommended for state use beginning January 1, 1989. Among the changes were the addition of a new item on educational attainment and Hispanic origin of the decedent and changes to improve the medical certification of cause of death. The U.S. Standard Certificate of Death was revised again in 2003; states are adopting this new certificate on a rolling basis.

The 2003 revision permits reporting of more than one race (multiple races). This change was implemented to reflect the increasing diversity of the U.S. population and to be consistent with the decennial census. Some states, however, are still using the 1989 revision of the U.S. Standard Certificate of Death, which allows only a single race to be reported. Until all states adopt the new death certificate, the race data reported using the 2003 revision are “bridged” for those for whom more than one race was reported (multiple race) to one single race, to provide comparability with race data reported on the 1989 revision. For more information on the impact of the 2003 certificate revisions on mortality data presented in *Health, United States*, see Appendix II, Race.

References

NCHS. Deaths: Final data for 2014. [Forthcoming]. Available from: <http://www.cdc.gov/nchs/products/nvsr.htm>.

Xu JQ, Murphy SL, Kochanek KD, Bastian BA. Deaths: Final data for 2013. National vital statistics reports; vol 64 no 2. Hyattsville, MD: NCHS; 2016. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr64/nvsr64_02.pdf.

Heron M, Hoyert DL, Murphy SL, et al. Deaths: Final data for 2006. National vital statistics reports; vol 57 no 14. Hyattsville, MD: NCHS; 2009. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr57/nvsr57_14.pdf.

NCHS. Multiple causes of death in the United States. Monthly vital statistics report; vol 32 no 10 suppl 2. Hyattsville, MD: NCHS; 1984. Available from: http://www.cdc.gov/nchs/data/mvsvr/supp/mv32_10s2.pdf.

For More Information. See the Mortality Data website at: <http://www.cdc.gov/nchs/deaths.htm>.

Linked Birth/Infant Death Data Set

Overview. National linked files of live births and infant deaths are used for research on infant mortality. The Linked Birth/Infant Death data set links information from the birth certificate to information from the death certificate for each infant death in the United States. The purpose of the linkage is to use the many additional variables from the birth certificate, including the more accurate race and ethnicity data, for more detailed analyses of infant mortality patterns. The Linked Birth/Infant Death data set includes all variables on the natality (Birth) file, including racial and ethnic information, birthweight, and maternal smoking, as well as variables on the Mortality file, including cause of death and age at death.

Coverage. To be included in the U.S. linked file, both the birth and death must have occurred in the 50 states, D.C., Puerto Rico, Virgin Islands, or Guam. Data for Puerto Rico, Virgin Islands, and Guam are shown in selected state tables but are not included in U.S. totals. Linked birth/infant death data are not available for American Samoa and Northern Marianas.

Methodology. Infant deaths are defined as a death before the infant's first birthday. About 98%–99% of infant death records can be linked to their corresponding birth certificates. The linkage makes available extensive information from the birth certificate about the pregnancy, maternal risk factors, infant characteristics, and health items at birth that can be used for more detailed analyses of infant mortality. The linked file is used for calculating infant mortality rates by race and ethnicity, which are more accurately measured from the birth certificate.

Starting with 1995 data, linked birth/infant death data files are available in two different formats: period data and birth cohort data. The numerator for the period linked file consists of all infant deaths occurring in a given data year linked to their corresponding birth certificates, whether the birth occurred in that year or the previous year. The numerator for the birth cohort linked file consists of deaths to infants born in a given year. In both cases, the denominator is all births occurring in the year. For example, the 2013 period linked file contains a numerator file that consists of all infant deaths occurring in 2013 that have been linked to their corresponding birth certificates, whether the birth occurred in 2012 or 2013. In contrast, the 2013 birth cohort linked file will contain a numerator file that consists of all infant deaths to babies born in 2013, whether the death occurred in 2013 or 2014. Although the birth cohort format has methodological advantages, it creates substantial delays in data availability because it is necessary to wait until the close of the following data year to include all infant deaths in the birth cohort. Starting with 1995 data, period linked files are used for infant mortality rate tables in *Health, United States*.

Other changes to the data set starting with 1995 include the addition of record weights to compensate for the 1%–2% of infant death records that could not be linked to their

corresponding birth records. In addition, not-stated birthweight was imputed if the period of gestation was known. This imputation was done to improve the accuracy of birthweight-specific infant mortality rates because the percentage of records with not-stated birthweight is generally higher for infant deaths (4.2% in 2013) than for live births (0.1% in 2013). In 2013, not-stated birthweight was imputed for 0.10% of births.

Issues Affecting Interpretation. Period linked file data starting with 1995 are not strictly comparable with birth cohort data for 1983–1991. A new revision of the birth certificate was introduced in 2003 and is being adopted by states on a voluntary, rolling basis.

Reference

Mathews TJ, MacDorman MF, Thoma ME. Infant mortality statistics from the 2013 period Linked Birth/Infant Death data set. National vital statistics report; vol 64 no 9. Hyattsville, MD: NCHS; 2015. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr64/nvsr64_09.pdf.

For More Information. See the NCHS Linked Birth and Infant Death Data website at: <http://www.cdc.gov/nchs/linked.htm>.

Occupational Employment Statistics (OES)

Bureau of Labor Statistics (BLS)

Overview. The OES program conducts a semiannual survey designed to produce estimates of employment and wages for specific occupations. The program collects data on wage and salary workers in nonfarm establishments in order to produce employment and wage estimates for about 800 occupations. The OES program produces these occupational estimates for all industries combined at different geographic levels—for the nation; the 50 states and D.C.; metropolitan and nonmetropolitan areas; and Guam, Puerto Rico, and the U.S. Virgin Islands. National occupational employment and wage estimates are also available by industry for more than 430 industry aggregations, and by public/private ownership across all industries and for schools and hospitals.

Coverage. The OES survey covers all full-time and part-time wage and salary workers in nonfarm establishments. The survey does not cover the self-employed, owners and partners in unincorporated firms, household workers, or unpaid family workers.

Methodology. The OES program surveys approximately 200,000 establishments per panel (every 6 months), taking 3 years to fully collect the sample of 1.2 million establishments. The estimates for occupations in nonfarm establishments are based on OES data collected for the reference months of May and November. May 2014 employment and wage estimates are based on all data

collected from establishments sampled in the May 2014, November 2013, May 2013, November 2012, May 2012, and November 2011 semiannual panels. The overall national response rate for the six panels is 74.3% based on establishments and 70.5% based on employment. The OES survey is a federal-state cooperative program between BLS and state workforce agencies (SWAs). BLS provides the procedures and technical support, draws the sample, and produces the survey materials, while SWAs collect most of the data. SWAs from all 50 states plus D.C., Puerto Rico, Guam, and the U.S. Virgin Islands participate in the survey. Occupational employment and wage rate estimates at the national level are produced by BLS using data from the 50 states and D.C. Employers who respond to states' requests to participate in the OES survey make these estimates possible.

Issues Affecting Interpretation. Over time, OES data have had changes in the occupational, industrial, and geographical classification systems; data collection methods; survey reference period; and mean wage estimation methodology. Because of these changes as well as permanent features of the OES methodology, caution should be used in trend analysis.

OES occupational estimates are based on the Office of Management and Budget's Standard Occupational Classification (SOC) system. The OES survey classifies workers into more than 800 detailed occupations; these detailed occupations are aggregated into 23 SOC major groups. Only 22 SOC major groups are included in OES; Major group 55, Military Specific Occupations, is not included.

OES estimates for 1999 through 2009 classified occupations according to the 2000 SOC system. OES estimates for 2010 and 2011 were based on a hybrid structure using both the 2000 and 2010 SOC systems. For more information about the hybrid structure, see http://www.bls.gov/oes/oes_ques.htm. OES estimates for 2012 to 2014 classified occupations according to the 2010 SOC system.

Reference

Bureau of Labor Statistics. Occupational employment and wages, May 2014. Washington, DC: U.S. Department of Labor; 2015. Available from: <http://www.bls.gov/oes/home.htm>.

For More Information. See the OES website at: <http://www.bls.gov/OES>.

Population Census and Population Estimates

U.S. Census Bureau

Decennial Census

The census of population (decennial census) has been held in the United States every 10 years since 1790. Since 1930, it

has enumerated the resident population as of April 1 of the census year. Data on sex, race, Hispanic origin, age, and marital status are collected from 100% of the enumerated population.

Race Data on the 1990 Census

The question on race on the 1990 census was based on the Office of Management and Budget's (OMB) 1977 *Race and Ethnic Standards for Federal Statistics and Administrative Reporting* (Statistical Policy Directive 15). This document specified rules for the collection, tabulation, and reporting of racial and ethnic data within the federal statistical system. The 1977 Standards required federal agencies to report race-specific tabulations using four single-race categories: American Indian or Alaska Native, Asian or Pacific Islander, black, and white. Under the 1977 Standards, race and ethnicity were considered to be two separate and distinct concepts. Thus, persons of Hispanic origin may be of any race.

Race Data on the 2000 Census

The question on race on the 2000 census was based on OMB's 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* (Fed Regist 1997 October 30;62:58781–90). (Also see Appendix II, Race.) The 1997 Standards incorporated two major changes in the collection, tabulation, and presentation of race data. First, the 1997 Standards increased from four to five the minimum set of categories to be used by federal agencies for identification of race: American Indian or Alaska Native, Asian, black or African American, Native Hawaiian or Other Pacific Islander, and white. Second, the 1997 Standards included the requirement that federal data collection programs allow respondents to select one or more race categories when responding to a query on their racial identity. This provision means that there are potentially 31 race groups, depending on whether an individual selects one, two, three, four, or all five of the race categories. The 1997 Standards continue to call for use, when possible, of a separate question on Hispanic or Latino ethnicity and specify that the ethnicity question should appear before the question on race. Thus, under the 1997 Standards, as under the 1977 Standards, persons of Hispanic origin may be of any race.

Race Data on the 2010 Census

Similar to race data on the 2000 census, the question on race on the 2010 census was based on OMB's 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* (Fed Regist 1997 October 30;62:58781–90). (Also see Appendix II, Race.) The 1997 Standards required a minimum set of categories to be used by federal agencies for identification of race: American Indian or Alaska Native, Asian, black or African American, Native Hawaiian or Other Pacific Islander, and white and require that federal data collection programs allow respondents to select one or

more race categories when responding to a query on their racial identity. The 1997 Standards continue to call for use, when possible, of a separate question on Hispanic or Latino ethnicity and specify that the ethnicity question should appear before the question on race. Thus, under the 1997 Standards, as under the 1977 Standards, persons of Hispanic origin may be of any race.

Modified Decennial Census Files

For several decades, the U.S. Census Bureau has produced Modified Decennial Census files. These modified files incorporate adjustments to the 100% April 1 count data for (a) errors in the census data discovered subsequent to publication, (b) misreported age data, and (c) nonspecified race.

For the 1990 census, the U.S. Census Bureau modified the age, race, and sex data on the census and produced the Modified Age-Race-Sex (MARS) file. The differences between the population counts in the original census file and the MARS file are primarily due to modification of the race data. Of the 248.7 million persons enumerated in 1990, 9.8 million did not specify their race (over 95% were of Hispanic origin). For the 1990 MARS file, these persons were assigned the race reported by a nearby person with an identical response to the Hispanic origin question.

For the 2000 and 2010 censuses, the U.S. Census Bureau modified the race data and produced the Modified Race Data Summary files. For these files, persons who did not report a race (reported only the category Some Other Race) as part of their race response were assigned by imputation to one of the 31 race groups, which are the single- and multiple-race combinations of the five race categories specified in the 1997 OMB race and ethnicity standards. For the 2000 census, 97% of the 15.4 million persons who did not report a race were of Hispanic origin. Because a large proportion of those identifying their race as Some Other Race are Hispanic, for the 2010 census, a new instruction was added that, for the census, Hispanic origins are not races. For the 2010 census, 97% of the 19.1 million persons who did not report a race (reported only the category Some Other Race) were of Hispanic origin.

Postcensal Population Estimates

Postcensal population estimates are estimates made for the years following a census, before the next census has been taken. Postcensal population estimates are derived annually by updating the resident population enumerated in the decennial census using a components-of-population-change approach. Each annual series includes estimates for the current data year and revised estimates for the earlier years in the decade. The following formula is used to derive national estimates for a given year from those for the previous year, starting with the decennial census enumerated resident population as the base:

Resident population estimate
+ births to U.S. resident women
– deaths to U.S. residents
+ net international migration.

The postcensal estimates are consistent with official decennial census figures and do not reflect estimated decennial census underenumeration.

Estimates for the earlier years in a given series are revised to reflect changes in the components-of-change data sets (for example, births to U.S. resident women from a preliminary natality file are replaced with counts from a final natality file). To help users keep track of which postcensal estimate is being used, each annual series is referred to as a “vintage,” and the last year in the series is used to name the series. For example, both the Vintage 2011 and the Vintage 2012 postcensal series have revised estimates for July 1, 2011, but the estimates for July 1, 2011, from the Vintage 2011 and Vintage 2012 postcensal series differ.

The U.S. Census Bureau also produces postcensal estimates of the resident population of states and counties, using the components-of-population-change method. An additional component of population change—net internal migration—is involved.

Intercensal Population Estimates

Intercensal population estimates are estimates made for the years between two decennial censuses and are produced once the census at the end of the decade has been completed. They replace the postcensal estimates produced prior to the completion of the census at the end of the decade. Intercensal estimates are more accurate than postcensal estimates because they are based on both the census at the beginning and the census at the end of the decade. They are derived by adjusting the final postcensal estimates for the decade to correct for the error of closure (the difference between the estimated population at the end of the decade and the census count for that date). The patterns of population change observed over the decade are preserved. The intercensal estimates for the 1990s were produced using the same methodology used to generate the intercensal estimates for the 1980s. The revised intercensal population estimates for 2000–2009 were produced using a modified version of the methodology used previously. Vital rates calculated using postcensal population estimates are routinely revised when intercensal estimates become available.

Bridged-race Population Estimates

Race data on the 2000 and 2010 censuses are not comparable with race data on other data systems that are continuing to collect data using the 1977 OMB Standards on race and ethnicity during the transition to full implementation of the 1997 OMB Standards. For example,

states are implementing the revised birth and death certificates—which have race and ethnicity items that are compliant with the 1997 OMB Standards—at different times, and to date some states are still using the 1989 certificates that collect race and ethnicity data in accordance with the 1977 OMB Standards. Thus, population estimates for 1990 and beyond with race categories comparable with the 1977 OMB categories are needed so that race-specific birth and death rates can be calculated. To meet this need, NCHS, in collaboration with the U.S. Census Bureau, developed methodology to bridge the 31 race groups in Census 2000 and Census 2010 to the four single-race categories specified under the 1977 OMB Standards.

The bridging methodology was developed using information from the 1997–2000 National Health Interview Survey (NHIS). NHIS provides a unique opportunity to investigate multiple-race groups because, since 1982, it has allowed respondents to choose more than one race but has also asked respondents reporting multiple races to choose a primary race. The bridging methodology developed by NCHS involved the application of regression models relating person-level and county-level covariates to the selection of a particular primary race by the multiple-race respondents. The bridging proportions derived from these models have been applied by the U.S. Census Bureau to various unbridged resident population files. These applications have resulted in bridged-race population estimates for each of the four single-race categories: American Indian or Alaska Native, Asian or Pacific Islander, black, and white.

In *Health, United States*, vital rates for 1991–1999 were calculated using the July 1, 1991–July 1, 1999 bridged-race intercensal estimates. Vital rates for 2000 were calculated using the bridged-race April 1, 2000, census counts, and those for 2010 were calculated using the bridged-race April 1, 2010, census counts. Starting with *Health, United States, 2012*, vital rates for 2001–2009 have been recalculated using the July 1, 2001–July 1, 2009, revised intercensal bridged-race population estimates. Vital rates for 2011 and beyond will be calculated using bridged-race estimates of the July 1 population from the corresponding postcensal vintage.

Reference

Ingram DD, Parker JD, Schenker N, et al. United States Census 2000 population with bridged race categories. NCHS. Vital Health Stat 2003;2(135). Available from: http://www.cdc.gov/nchs/data/series/sr_02/sr02_135.pdf.

For More Information. See the U.S. Census Bureau website at: <http://www.census.gov> and the NCHS website for U.S. Census populations with bridged race categories at: http://www.cdc.gov/nchs/nvss/bridged_race.htm.

Quality Improvement Evaluation System (QIES)

Centers for Medicare & Medicaid Services (CMS)

Overview. This administrative database, referred to in *Health, United States* as QIES, is created from the Certification and Survey Provider Enhanced Reporting (CASPER) and QIES systems. QIES is a CMS database that contains information from the standard annual facility survey data submitted by state survey agencies to CMS for certification to participate in the Medicare and Medicaid programs in the United States and territories. (Data for the territories are not shown in *Health, United States*.) The purpose of the facility survey certification process is to ensure that facilities meet current CMS care requirements and thus can be paid for services furnished to Medicare and Medicaid beneficiaries. In 2012, QIES replaced the Online Survey Certification and Reporting Database (OSCAR). QIES (and its predecessor OSCAR) contain information on facility and patient characteristics and health deficiencies issued by the government during the survey process.

Coverage. Facilities in the United States that are certified to receive Medicare or Medicaid payments are included.

Methodology. QIES data are compiled by the state survey agency and a facility representative. The data are reviewed during the survey process and then submitted electronically to CMS. The information provided can be audited at any time.

All certified facilities are inspected periodically by representatives of the state survey agency (generally the department of health). Some facilities are inspected twice, or more often, during any given reporting cycle. To avoid overcounting, the data must be edited and duplicates removed. Data editing and compilation of nursing home data were performed by Cowles Research Group (CRG; Anacortes, WA.) and published in the group's *Nursing Home Statistical Yearbook* series.

References

Cowles CM, ed. Nursing home statistical yearbooks for 2003–2014. Anacortes, WA: CRG; published 2004–2015, respectively.

Centers for Medicare & Medicaid Services. Certification and compliance. Baltimore, MD: CMS; 2005. Available from: <http://www.cms.gov/Research-Statistics-Data-and-Systems/Files-for-Order/NonIdentifiableDataFiles/Index.html>.

For More Information. See the CMS website at: <http://www.cms.gov/Research-Statistics-Data-and-Systems/Files-for-Order/NonIdentifiableDataFiles/index.html> and the CRG website at: <http://www.longtermcareinfo.com/index.html>.

Sexually Transmitted Disease (STD) Surveillance

CDC/National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention (NCHHSTP)

Overview. Surveillance information on the incidence and prevalence of STDs is used to inform public and private health efforts to control these diseases. Case reporting data are available for nationally notifiable chancroid, chlamydia, gonorrhea, and syphilis. Enhanced surveillance of these conditions and surveillance of other STDs, such as genital herpes simplex virus, genital warts or other human papillomavirus infections, and trichomoniasis use data collected from other sources, including data from sentinel surveillance and national surveys.

Coverage. Case reports of STDs are reported to CDC by STD surveillance systems operated by state and local STD control programs and health departments in 50 states, D.C., selected cities, 3,142 U.S. counties, and outlying areas consisting of U.S. dependencies, possessions, and independent nations in free association with the United States. Data from outlying areas are not included in *Health, United States*.

Methodology. Information is obtained from the following data sources: (a) notifiable disease reporting from state and local STD programs; (b) projects that monitor STD positivity and prevalence in various settings, including the National Job Training Program, the STD Surveillance Network, and the Gonococcal Isolate Surveillance Project; and (c) national sample surveys implemented by federal and private organizations. STD data are submitted to CDC on a variety of hard-copy summary reporting forms (monthly, quarterly, and annually) and in electronic summary or individual case-specific (line-listed) formats through the National Electronic Telecommunications System for Surveillance.

Issues Affecting Interpretation. Because of incomplete diagnosis and reporting, the number of STD cases reported to CDC undercounts the actual number of infections occurring among the U.S. population.

Reference

CDC. Sexually transmitted disease surveillance 2014. Atlanta, GA: CDC; 2015. Available from: <http://www.cdc.gov/std/stats14/default.htm>.

For More Information. See the STD Data and Statistics website at: <http://www.cdc.gov/std/stats> and the STD Diseases & Related Conditions website at: <http://www.cdc.gov/std/default.htm>.

Surveillance, Epidemiology, and End Results Program (SEER)

National Cancer Institute (NCI)

Overview. SEER tracks the incidence of new cancers each year and collects follow-up information on all previously diagnosed patients until their death. For each cancer, SEER registries routinely collect data on patient demographics, primary tumor site, morphology, stage at diagnosis, first course of treatment, and follow-up for vital status.

Coverage. The SEER 9 registries (Atlanta, Connecticut, Detroit, Hawaii, Iowa, New Mexico, San Francisco-Oakland, Seattle-Puget Sound, and Utah) have been part of the program continuously since 1975. The SEER 13 registries (the SEER 9 registries plus Los Angeles, San Jose-Monterey, rural Georgia, and the Alaska Native Tumor Registry) have been part of the program continuously since 1992. The SEER 18 registries (the SEER 13 plus Greater Georgia, Kentucky, Greater California, New Jersey, and Louisiana) have been part of the program continuously since 2000. SEER currently collects and publishes cancer incidence and survival data from 18 population-based cancer registries covering approximately 28% of the U.S. population.

Methodology. A cancer registry collects and stores data on cancers diagnosed in a specific hospital or medical facility (hospital-based registry) or in a defined geographic area (population-based registry). A population-based registry includes, but is not limited to, a number of hospital-based registries. In SEER registry areas, trained coders abstract medical records using the *International Classification of Diseases for Oncology, 3rd edition (ICD-O-3)* to classify site and tumor morphology. The ICD-O-3 coding also includes updates for hematopoietic codes based on *WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues* (2008). All SEER data in this report were collected with or converted to ICD-O-3.

NCI obtains population counts from the U.S. Census Bureau and uses them to calculate incidence rates. It also uses estimation procedures as needed to obtain estimates for years and races not included in data provided by the Census Bureau. Life tables used to determine general population life expectancy when calculating relative survival rates were obtained from NCHS and in-house calculations. Separate life tables are used for each race-sex-specific group included in SEER.

Issues Affecting Interpretation. Because of the addition of registries over time, analysis of long-term incidence and survival trends is limited to those registries that have been in SEER for similar lengths of time. Analysis of Hispanic, and American Indian or Alaska Native data is limited to shorter trends. Starting with *Health, United States, 2006*, the North American Association of Central Cancer Registries (NAACCR) Hispanic Identification Algorithm was used on a combination of variables to classify cases as Hispanic for

analytic purposes. Starting with *Health, United States, 2007*, Hispanic incidence data exclude data for Alaska. Earlier editions of *Health, United States* also excluded Hispanic data for Hawaii and Seattle. Starting with *Health, United States, 2007*, incidence estimates for the American Indian or Alaska Native population are limited to contract health service delivery area (CHSDA) counties within SEER reporting areas. This change is believed to produce estimates that more accurately reflect the incidence rates for this population group. More information on CHSDA is available from: http://www.ihs.gov/chs/index.cfm?module=chs_requirements_chsda. For more information on SEER estimates by race and ethnicity, see: http://seer.cancer.gov/seerstat/variables/seer/race_ethnicity/index.html. Rates presented in this report may differ somewhat from those reported previously due to changes in population estimates and the addition and deletion of small numbers of incidence cases.

Reference

Howlader N, Noone AM, Krapcho M, Garshell J, Miller D, Altekruse SF, et al. (eds). SEER cancer statistics review, 1975–2012. Bethesda, MD: National Cancer Institute. Based on November 2014 SEER data submission, posted to the SEER website, April 2015. Available from: http://seer.cancer.gov/csr/1975_2012/.

For More Information. See the SEER website at: <http://seer.cancer.gov>.

Youth Risk Behavior Survey (YRBS)

CDC/National Center for HIV, Hepatitis, STD, and TB Prevention (NCHHSTP)

Overview. YRBS monitors health risk behaviors among students in grades 9–12 that contribute to morbidity and mortality in both adolescence and adulthood. The six areas monitored are behaviors that contribute to unintentional injuries and violence; tobacco use; alcohol and other drug use; sexual behaviors that contribute to unintended pregnancy and sexually transmitted diseases (STDs), including human immunodeficiency virus (HIV) infection; unhealthy dietary behaviors; and physical inactivity. In addition, YRBS monitors the prevalence of obesity, asthma, and sleep behaviors.

Coverage. National data are representative of high school students in public and private schools in the United States.

Methodology. The national YRBS school-based surveys have been conducted biennially since 1991. A three-stage cluster sample design is used to produce a nationally representative sample of students in grades 9–12 attending public and private schools. In 2013, the first-stage sampling frame comprised primary sampling units (PSUs) consisting of counties, subareas of large counties, or groups of smaller, adjacent counties. PSUs were categorized into strata

according to their metropolitan statistical area (MSA) status (e.g., urban city) and the percentages of black and Hispanic students in the PSUs. PSUs were sampled with probability proportional to overall school enrollment size for the PSU. In the second stage of sampling, schools with any of grades 9–12 were sampled with probability proportional to school enrollment size. The third stage of sampling consisted of random sampling in each of grades 9–12, one or two classrooms from either a required subject (e.g., English or Social Studies) or a required period (e.g., Homeroom or second period).

All students in sampled classes were eligible to participate. Schools, classes, and students that refused to participate were not replaced. To enable a separate analysis of data for black and Hispanic students, two classes per grade, rather than one, were sampled in schools with a high enrollment of black and Hispanic students. Prior to 2013, three strategies were used to oversample black and Hispanic students: (a) larger sampling rates were used to select PSUs that were in high-black and high-Hispanic strata; (b) a modified measure of size was used to increase the probability of sampling schools with a disproportionately high minority enrollment; and (c) two classes per grade, rather than one, were sampled in schools with a high enrollment of black and Hispanic students. A weighting factor is applied to each student record to adjust for nonresponse and for the varying probabilities of selection, including those resulting from the oversampling of black and Hispanic students.

Sample Size and Response Rate. The sample size for the 2013 YRBS was 13,583 students in 148 schools. The school response rate was 77%, and the student response rate was 88%, for an overall response rate of 68%.

Issues Affecting Interpretation. National YRBS data are subject to at least two limitations. First, these data apply only to adolescents who attend regular high school, including some charter, public alternative, special education, and vocational schools. These students may not be representative of all persons in this age group because those who have dropped out of high school are not surveyed. Second, the extent of underreporting or overreporting cannot be determined, although the survey questions demonstrate good test-retest reliability.

Estimates of substance use for youth based on YRBS differ from the National Survey on Drug Use & Health (NSDUH) and the Monitoring the Future (MTF) Study. Rates are not directly comparable across these surveys because of differences in populations covered, sample designs, questionnaires, and interview settings. NSDUH collects data in residences, whereas MTF and YRBS collect data in school classrooms. In addition, NSDUH estimates are tabulated by age, whereas MTF and YRBS estimates are tabulated by grade, representing different ages as well as different populations. All YRBS data collection is anonymous.

References

Brener ND, Kann L, Shanklin SL, Kinchen S, Eaton DK, Hawkins J, et al. Methodology of the Youth Risk Behavior Surveillance System—2013. *MMWR* 2013;62(RR-1):1–23. Available from: <http://www.cdc.gov/mmwr/preview/mmwrhtml/rr6201a1.htm>.

Kann L, Kinchen S, Shanklin SL, Flint KH, Hawkins J, Harris WA, et al. Youth Risk Behavior Surveillance—United States, 2013. *MMWR Surveill Summ* 2014;63(SS-4):1–172. Available from: <http://www.cdc.gov/mmwr/pdf/ss/ss6304.pdf>.

Cowan CD. Coverage, sample design, and weighting in three federal surveys. *J Drug Issues* 2001;31(3):599–614.

For More Information. See the YRBS website at: <http://www.cdc.gov/yrebs>.

Private and Global Sources

American Association of Colleges of Osteopathic Medicine (AACOM)

AACOM compiles data on various aspects of osteopathic medical education for distribution to the profession, the government, and the public. Questionnaires are sent annually to schools of osteopathic medicine requesting information on characteristics of applicants, students and graduates, faculty, curriculum, contract and grant activity, revenues and expenditures, and clinical facilities.

Reference

American Association of Colleges of Osteopathic Medicine. Trends in osteopathic medical school applicants, enrollment and graduates 2016. Chevy Chase, MD: AACOM; 2016.

For More Information. See the AACOM website at: <http://www.aacom.org>.

American Association of Colleges of Pharmacy (AACP)

AACP compiles data on colleges and schools of pharmacy, including information on student enrollment and types of degrees conferred. Data are collected through an annual survey. In 2013–2014, the response rate was 99.2%.

Reference

American Association of Colleges of Pharmacy. Fall 2014 profile of pharmacy students. Available from: <http://www.aacp.org/resources/research/institutionalresearch/Pages/StudentApplications,EnrollmentsandDegreesConferred.aspx>.

For More Information. See the AACP website at: <http://www.aacp.org>.

American Association of Colleges of Podiatric Medicine (AACPM)

AACPM compiles data on colleges of podiatric medicine, including information on the schools and enrollment. Data are collected annually through written questionnaires. The response rate is 100%.

Reference

American Association of Colleges of Podiatric Medicine. Applicant, matriculant, and graduate statistics. Available from: <http://www.aacpm.org>.

For More Information. See the AACPM website at: <http://www.aacpm.org>.

American Dental Association (ADA)

The ADA Masterfile contains the most up-to-date information on dentists in the United States. The Masterfile is a database of all dentists, practicing and nonpracticing, in the United States. It is updated through a variety of methods including reconciliation with state licensure databases, death records, and various surveys and censuses of dentists carried out by ADA.

ADA's Health Policy Institute conducts annual surveys of predoctoral dental educational institutions. A questionnaire, mailed to all dental schools, collects information on academic programs, admissions, enrollment, attrition, graduates, educational expenses and financial assistance, patient care, advanced dental education, and faculty positions.

References

American Dental Association, Survey Center. Supply of dentists in the U.S.: 2001–2013, Tables 1 and 3. Available from: <http://www.ada.org/en/science-research/health-policy-institute/data-center/supply-of-dentists>.

American Dental Association. 2014–2015 survey of dental education series. Report 1: Academic programs, enrollment and graduates. Chicago, IL: ADA; 2015. Available from: <http://www.ada.org/en/science-research/health-policy-institute/data-center/dental-education>.

For More Information. See the ADA website at: <http://www.ada.org>.

American Hospital Association (AHA) Annual Survey of Hospitals

Data from AHA's annual survey are based on questionnaires sent to all AHA-registered and nonregistered hospitals in the United States and its associated areas: American Samoa, Guam, the Marshall Islands, Puerto Rico, and the Virgin Islands. U.S. government hospitals located outside the United States are excluded. Overall, the average response rate over the past 5 years has been approximately 83%. For nonreporting hospitals and for the survey questionnaires of reporting hospitals on which some information was missing, estimates are made for all data except those on beds, bassinets, facilities, and services. Data for beds and bassinets of nonreporting hospitals are based on the most recent information available from those hospitals. Data for facilities and services are based only on reporting hospitals. Estimates of other types of missing data are based on data reported the previous year, if available. When unavailable, estimates are based on data furnished by reporting hospitals similar in size, control, major service provided, length of stay, and geographic and demographic characteristics.

Reference

American Hospital Association, Annual survey of hospitals. Hospital statistics, 2015. Chicago, IL: AHA; 2015.

For More Information. See the AHA website at: <http://www.aha.org>.

American Medical Association (AMA) Physician Masterfile

A master file of physicians has been maintained by AMA since 1906. The Physician Masterfile contains data on all physicians in the United States, both members and nonmembers of AMA, and on those graduates of American medical schools temporarily practicing overseas. The file also includes information on international medical graduates (IMGs) who are graduates of foreign medical schools, who reside in the United States, and who meet U.S. educational standards for primary recognition as physicians.

A file is initiated on each individual upon entry into medical school or, in the case of IMGs, upon entry into the United States. Between 1969 and 1985, a mail questionnaire survey was conducted every 4 years to update the file information on professional activities, self-designated area of specialization, and present employment status. Between 1985 and 2006, approximately one-third to one-fourth of all physicians were surveyed each year. Since then, AMA has employed a more diversified survey approach in which more than 500,000 active physicians are targeted each year through mail, telephone, and web-based surveys.

Reference

American Medical Association. Physician characteristics and distribution in the U.S., 2015. Chicago, IL: AMA Division of Survey and Data Resources; 2015.

For More Information. See the AMA website at: <http://www.ama-assn.org>.

American Osteopathic Association (AOA)

AOA was established to promote the public health, to encourage scientific research, and to maintain and improve high standards of medical education in osteopathic colleges. Among its activities, AOA compiles the number of osteopathic physicians (DOs); the number of active DOs by gender, age, and specialty and by 50 states and D.C.; and the number of osteopathic medical students, by selected characteristics.

Reference

American Osteopathic Association. 2015 osteopathic medical profession report. Chicago, IL: AOA; 2015. Available from: <http://www.osteopathic.org/inside-aoa/about/aoa-annual-statistics/Pages/default.aspx>.

For More Information. See the AOA website at: <http://www.osteopathic.org>.

Association of American Medical Colleges (AAMC)

As part of its mission to serve and lead the academic medicine community to improve the health of all, AAMC collects information on student enrollment in medical schools through a variety of sources. Among the data services and sources offered are the Medical College Admission Test (MCAT), the American Medical College Application Service (AMCAS), the Electronic Residency Application Service (ERAS), and the Student Records System. The AAMC Data Warehouse stores data relevant to both applicants and students, and from these two source files the association derives summary statistics about applicants, accepted applicants, matriculants, enrollees, and graduates. AAMC has developed policies and procedures to ensure that the privacy of individual and institutional data are protected and meet federal, state, AAMC, and professional standards. Applicant, enrollment, and graduate statistical data are arranged by academic year, which begins July 1 and ends June 30.

Reference

Association of American Medical Colleges. AAMC data book: Medical schools and teaching hospitals by the numbers, 2015. Washington, DC: AAMC; 2015.

For More Information. See the AAMC website at: <http://www.aamc.org>.

Association of Schools and Colleges of Optometry (ASCO)

ASCO compiles data on various aspects of optometric education, including data on schools and enrollment. Schools and colleges complete an annual questionnaire. The response rate is 100%.

Reference

Association of Schools and Colleges of Optometry. Annual student data report: Academic year 2014–2015. Rockville, MD: ASCO; 2015. Available from: <http://www.opted.org/student-data-reports/>.

For More Information. See the ASCO website at: <http://www.opted.org>.

Association of Schools & Programs of Public Health (ASPPH)

ASPPH compiles data on Council on Education for Public Health-accredited schools and programs of public health in the United States, Puerto Rico, Mexico, and Canada. Unlike health professional schools that emphasize specific clinical occupations, schools and programs of public health offer study in specialty areas such as biostatistics, epidemiology, environmental health, occupational health, health administration, health planning, nutrition, maternal and child health, social and behavioral sciences, and other population-based sciences. Data collection is conducted annually from all ASPPH member schools and programs. The response rate in 2013–2014 was 85%.

Reference

Association of Schools and Programs of Public Health. [unpublished data]. Washington, DC: ASPPH; 2014.

For More Information. See the ASPPH website at: <http://www.aspph.org>.

Guttmacher Institute Abortion Provider Census

The Guttmacher Institute (previously called the Alan Guttmacher Institute, or AGI) is a not-for-profit organization for reproductive health research, policy analysis, and public education. Guttmacher has collected or estimated national abortion data since 1973 by conducting surveys every 3–4 years and extrapolating estimates for the intervening years. Guttmacher reports the number of legal induced abortions and the number, types, and locations of abortion providers by state and region.

The abortion data reported to Guttmacher contain data on women of all ages, including adolescents who obtain legal induced abortions, and includes both surgical and

medication (e.g., using mifepristone, misoprostol, or methotrexate) abortion procedures. Data are collected from three major categories of providers that were identified as potential providers of abortion services: clinics, physicians, and hospitals.

Questionnaires are mailed to all potential providers, with two additional mailings and telephone follow-up for nonresponse. All questionnaires ask the number of induced abortions performed at the provider's location. State health statistics agencies are also contacted, requesting all available data reported by providers to each state health agency on the number of abortions performed in the survey year. For states that provide data to Guttmacher, the health agency figures are used for providers who do not respond to the survey. Estimates of the number of abortions performed by some providers are ascertained from knowledgeable sources, including other providers of reproductive health services.

In the 2012–2013 survey, respondents were asked to report the number of induced abortions performed in their facilities during 2010 and 2011. Of the 2,288 potential providers surveyed between April 2012 and May 2013, 1,222 responded directly or in follow-up; health department data were used for 470 providers; 71 facilities had closed or stopped offering abortion services during the survey period; knowledgeable sources were used for 51 providers; and Guttmacher made its own estimates for 474 facilities, usually relying on prior abortion census results. The level of internal estimation was higher than in the 2008 survey.

Between 2003 and 2011, the total number of abortions reported to CDC has been about one-third less than the total estimated by Guttmacher. (See Appendix I, Abortion Surveillance System.)

Reference

Jones RK, Jerman J. Abortion incidence and service availability in the United States, 2011. *Perspect Sex Reprod Health* 2014;46(1):3–14. Available from: <http://www.guttmacher.org/pubs/journals/psrh.46e0414.pdf>.

For More Information. See The Guttmacher Institute website at: <http://www.guttmacher.org>.

Organisation for Economic Co-operation and Development (OECD) Health Data

OECD provides annual data on statistical indicators for health and health systems collected from 34 member countries, with some time series going back to 1960.

OECD was established in 1961 with a mandate to promote policies to achieve the highest sustainable economic growth and a rising standard of living among member countries. The organization now comprises 34 member countries: Australia, Austria, Belgium, Canada, Chile, the Czech Republic, Denmark, Estonia, Finland, France, Germany,

Greece, Hungary, Iceland, Ireland, Israel, Italy, Japan, Korea, Luxembourg, Mexico, the Netherlands, New Zealand, Norway, Poland, Portugal, the Slovak Republic, Slovenia, Spain, Sweden, Switzerland, Turkey, the United Kingdom, and the United States.

Each year, OECD compiles cross-country data in the OECD Health Data database, one of the most comprehensive sources of comparable health-related statistics.

For More Information. See the OECD website at:

<http://www.oecd.org/health>.

Appendix II. Definitions and Methods

This appendix contains an alphabetical listing of terms used in *Health, United States*, and these definitions are specific to the data presented in this report. The methods used for calculating age-adjusted rates, average annual rates of change, relative standard errors, birth rates, death rates, and years of potential life lost are described. Included are standard populations used for age adjustment (Tables I and II), the years when the revisions for *International Classification of Diseases* (ICD) codes were in effect (Table III), codes for cause of death from the 6th through 10th revisions of ICD (Table IV), and comparability ratios between the 9th and 10th revisions (ICD-9 and ICD-10) for selected causes (Table V), imputed family income percentages from the National Health Interview Survey (NHIS) (Table VI), an analysis of the effect of added probe questions for Medicare and Medicaid coverage on health insurance rates in NHIS (Table VII), industry codes from the North American Industry Classification System (NAICS) (Table VIII), and ICD-9 Clinical Modification (ICD-9-CM) codes for external causes of injury and procedure categories (Tables IX and X). Standards for presenting federal data on race and ethnicity are described, and sample tabulations of NHIS data comparing the 1977 and 1997 Office of Management and Budget standards for the classification of federal data on race and ethnicity are presented in Tables XI and XII.

Acquired immunodeficiency syndrome (AIDS)—Human immunodeficiency virus (HIV) is the pathogen that causes AIDS, and HIV disease is the term that encompasses all of the condition's stages—from infection to the deterioration of the immune system and the onset of opportunistic diseases. However, AIDS is still the term most people use to refer to the immune deficiency caused by HIV. An AIDS diagnosis indicates that the person has reached the late stages of the disease and is given to people with HIV who have been diagnosed with at least one of a set of opportunistic diseases or whose laboratory values indicate advanced disease. All 50 states, the District of Columbia (D.C.), and six U.S. dependent areas (American Samoa, Guam, Northern Mariana Islands, Puerto Rico, Republic of Palau, and U.S. Virgin Islands) report confirmed diagnoses of HIV infection and AIDS cases to CDC using a uniform surveillance case definition and case report form. The case reporting definitions have changed over time to incorporate a broader range of AIDS-indicator diseases and conditions and use HIV diagnostic tests to improve the sensitivity and specificity of the definition. Because of these case definition changes, caution should be used when interpreting AIDS trends. (Also see Appendix II, Human immunodeficiency virus [HIV] disease.)

Active physician—See Appendix II, Physician.

Activities of daily living (ADL)—ADLs are activities related to personal care and include bathing or showering, dressing, getting into or out of bed or a chair, getting around inside the home, using the toilet, and eating. In the National Health Interview Survey, respondents were asked whether they or family members need the help of another person with personal care because of a physical, mental, or emotional problem.

In the Medicare Current Beneficiary Survey, if a sample person had any difficulty performing an activity by him- or herself and without special equipment, or did not perform the activity at all because of health problems, the person was categorized as having a limitation in that activity. The limitation may have been temporary or chronic at the time of interview. Sampled people who were administered a community interview answered questions about health status and functioning themselves, if able to do so. If the sample person was not able to respond, a proxy answered the questions. For persons in a long-term care facility, a proxy such as a nurse answered questions about the sample person's health status and functioning. Starting in 1997, interview questions for people residing in long-term care facilities were changed slightly from those administered to people living in the community, in order to differentiate residents who were independent from those who received supervision or assistance with transferring, locomotion on unit, dressing, eating, toilet use, and bathing. (Also see Appendix II, Basic actions difficulty; Complex activity limitation; Instrumental activities of daily living [IADL]; Limitation of activity.)

Admission—The American Hospital Association defines admissions as persons, excluding newborns, accepted for inpatient services during the survey reporting period. (Also see Appendix II, Days of care; Discharge; Inpatient.)

Age—Age is reported as age at last birthday (i.e., age in completed years), often calculated by subtracting the date of birth from the reference date, with the reference date being the date of the examination, interview, or other contact with an individual.

Mother's (maternal) age is reported on the birth certificate by all states. Birth statistics are presented for mothers aged 10–49 through 1996 and aged 10–54 starting in 1997, based on mother's date of birth or age as reported on the birth certificate. The age of the mother is edited for upper and lower limits. When the age of the mother is computed to be under 10 or 55 and over (50 and over in 1964–1996), it is considered not stated and is imputed according to the age of the mother from the previous birth record of the same race and total birth order (total of fetal deaths and live births). Before 1963, not-stated ages were distributed in proportion to the known ages for each racial group.

Table I. United States projected year 2000 standard population and age groups used to age-adjust data

<i>Data system and age</i>	<i>Population</i>
DVS mortality data	
Total	274,633,642
Under 75 years	258,059,676
Under 1 year	3,794,901
1–4 years	15,191,619
5–14 years	39,976,619
15–24 years	38,076,743
25–34 years	37,233,437
35–44 years	44,659,185
45–54 years	37,030,152
55–64 years	23,961,506
65–74 years	18,135,514
75–84 years	12,314,793
85 years and over	4,259,173
DVS (Table 18)	
Under 75 years	258,059,676
Under 1 year	3,794,901
1–14 years	55,168,238
15–24 years	38,076,743
25–34 years	37,233,437
35–44 years	44,659,185
45–54 years	37,030,152
55–64 years	23,961,506
65–74 years	18,135,514
NHIS, NAMCS, and NHAMCS	
All ages	274,633,642
18 years and over	203,852,188
25 years and over	177,593,760
40 years and over	118,180,367
65 years and over	34,709,480
Under 18 years	70,781,454
2–17 years	63,227,991
18–44 years	108,151,050
18–24 years	26,258,428
25–34 years	37,233,437
35–44 years	44,659,185
45–64 years	60,991,658
45–54 years	37,030,152
55–64 years	23,961,506
65–74 years	18,135,514
75 years and over	16,573,966
18–49 years	127,956,843
40–64 years:	
40–49 years	42,285,022
50–64 years	41,185,865

See footnotes at end of table.

Table I. United States projected year 2000 standard population and age groups used to age-adjust data—Con.

<i>Data system and age</i>	<i>Population</i>
NHANES	
20 years and over	195,850,985
20–34 years	55,490,662
35–44 years	44,659,185
45–54 years	37,030,152
55–64 years	23,961,506
65 years and over	34,709,480
NHANES (Tables 38 and 53)	
20–44 years	100,149,847
45–64 years	60,991,658
65 years and over	34,709,480
NHANES (Table 56)	
20–44 years	100,149,847
45–64 years	60,991,658
65–74 years	18,135,514
75 years and over	16,573,966
NHANES (Table 79)	
Under 18 years	70,781,454
18–44 years	108,151,050
45–64 years	60,991,658
65 years and over	34,709,480

NOTES: DVS is Division of Vital Statistics. NHIS is National Health Interview Survey. NAMCS is National Ambulatory Medical Care Survey. NHAMCS is National Hospital Ambulatory Medical Care Survey. NHANES is National Health and Nutrition Examination Survey. SOURCE: National Institutes of Health, National Cancer Institute. Surveillance, Epidemiology, and End Results (SEER). Standard populations—single ages. Available from: <http://seer.cancer.gov/stdpopulations>.

Beginning in 1997, the birth rate for the maternal age group 45–49 has included data for mothers aged 50–54 in the numerator and has been based on the population of women aged 45–49 in the denominator. Beginning with 2003 data, age of mother is imputed for stated ages 8 and under and 65 and over, for births occurring in states using the 2003 revision of the birth certificate. Starting with 2007 data, age of mother is imputed for all births for stated ages 8 and under and 65 and over, regardless of the birth certificate version used. As with data for earlier years, age is imputed according to the age of mother from the previous record with the same race and total birth order.

Age adjustment—Age adjustment is used to compare risks for two or more populations at one point in time or for one population at two or more points in time. Age-adjusted rates are computed by the direct method by applying age-specific rates in a population of interest to a standardized age distribution, to eliminate differences in observed rates that result from age differences in population composition.

Table II. United States projected year 2000 standard population and proportion distribution, by age, for age-adjusting death rates prior to 2001

Age	Population	Proportion distribution (weight)	Standard million
Total	274,634,000	1.000000	1,000,000
Under 1 year	3,795,000	0.013818	13,818
1–4 years	15,192,000	0.055317	55,317
5–14 years	39,977,000	0.145565	145,565
15–24 years	38,077,000	0.138646	138,646
25–34 years	37,233,000	0.135573	135,573
35–44 years	44,659,000	0.162613	162,613
45–54 years	37,030,000	0.134834	134,834
55–64 years	23,961,000	0.087247	87,247
65–74 years	18,136,000	0.066037	66,037
75–84 years	12,315,000	¹ 0.044842	44,842
85 years and over	4,259,000	0.015508	15,508

¹Figure is rounded up instead of down to force total to 1.0.

SOURCE: CDC/NCHS. Anderson RN, Rosenberg HM. Age standardization of death rates: Implementation of the year 2000 standard. National vital statistics reports; vol 47 no 3. Hyattsville, MD: NCHS; 1998. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr47/nvs47_03.pdf.

Age-adjusted rates should be viewed as relative indexes rather than actual measures of risk.

Age-adjusted rates are calculated by the direct method, as follows:

$$\sum_{i=1}^n r_i \times (p_i / P)$$

where

r_i = rate in age group i in the population of interest

p_i = standard population in age group i

$$P = \sum_{i=1}^n p_i$$

n = total number of age groups over the age range of the age-adjusted rate.

Age adjustment by the direct method requires the use of a standard age distribution. The standard for age-adjusting death rates and estimates from surveys in *Health, United States* is the projected year 2000 U.S. resident population. Starting with *Health, United States, 2000*, the projected year 2000 U.S. standard population replaced the 1970 civilian noninstitutionalized population for age-adjusting estimates from most NCHS surveys; and starting with *Health, United States, 2001*, it was used uniformly and replaced the 1940 U.S. population for age-adjusting mortality statistics and the 1980 U.S. resident population, which previously had been used for age-adjusting estimates from the National Health and Nutrition Examination Survey.

Changing the standard population has implications for racial and ethnic differentials in mortality. For example, the mortality ratio for the black to white populations is reduced

from 1.6 using the 1940 standard to 1.4 using the 2000 standard, reflecting the greater weight the 2000 standard gives to the older population, in which race differentials in mortality are smaller.

Age-adjusted estimates from any data source presented in *Health, United States* that use the projected year 2000 U.S. resident population may differ from age-adjusted estimates based on the same data presented in other reports if different age groups are used in the adjustment procedure.

For more information on implementing the 2000 population standard for age-adjusting death rates, see: Anderson RN, Rosenberg HM. Age standardization of death rates: Implementation of the year 2000 standard. National vital statistics reports; vol 47 no 3. Hyattsville, MD: NCHS; 1998. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr47/nvs47_03.pdf. For more information on the derivation of age-adjustment weights for use with NCHS survey data, see: Klein RJ, Schoenborn CA. Age adjustment using the 2000 projected U.S. population. Healthy People 2010 statistical notes, no 20. Hyattsville, MD: NCHS; 2001. Available from: <http://www.cdc.gov/nchs/data/statnt/statnt20.pdf>. The projected year 2000 U.S. standard population is available from the National Cancer Institute's Surveillance, Epidemiology, and End Results (SEER) Program: <http://seer.cancer.gov/stdpopulations/stdpop.singleages.html>.

Mortality data—Death rates are age-adjusted to the projected year 2000 U.S. standard population (Table I). Prior to 2001 data, age-adjusted rates were calculated using standard million proportions based on rounded population numbers (Table II). Starting with 2001 data, unrounded population numbers are used to age-adjust. Adjustment is based on 11 age groups, with two exceptions. First, age-adjusted death rates for black males and black females in 1950 are based on nine age

groups, with under 1 and 1–4 combined as one group, and 75–84 and 85 and over combined as one group. Second, age-adjusted rates for years of potential life lost before age 75 also use the projected year 2000 standard population and are based on eight age groups: under 1, 1–14, 15–24, and 10-year age groups through 65–74.

National Health and Nutrition Examination Survey (NHANES)—Estimates based on the National Health Examination Survey and NHANES are generally age-adjusted to the projected year 2000 U.S. standard population by using five age groups: 20–34, 35–44, 45–54, 55–64, and 65–74 or 65 and over (Table I). Prior to *Health, United States, 2001*, these estimates were age-adjusted to the 1980 U.S. resident population.

National Health Care Surveys—Estimates based on the National Ambulatory Medical Care Survey, and the National Hospital Ambulatory Medical Care Survey are age-adjusted to the projected year 2000 U.S. standard population (Table I). Information on the age groups used in the age-adjustment procedure is contained in the footnotes to the specific tables.

National Health Interview Survey (NHIS)—Estimates based on NHIS are age-adjusted to the projected year 2000 U.S. standard population (Table I). Prior to *Health, United States, 2000*, NHIS estimates were age-adjusted to the 1970 civilian noninstitutionalized population. Information on the age groups used in the age-adjustment procedure is contained in the footnotes to the specific tables.

AIDS—See Appendix II, Acquired immunodeficiency syndrome (AIDS).

Alcohol consumption—Alcohol consumption is measured differently in the following data systems. (Also see Appendix II, Binge drinking.)

Monitoring the Future (MTF) Study—This school-based survey of secondary school students collects information on alcohol use by using self-completed questionnaires. To determine whether they have tried alcohol in their lifetime, students are asked a preliminary alcohol consumption (defined as beer, wine, liquor, and any other beverage that contains alcohol) screening question: “Have you ever had any alcoholic beverage to drink—more than just a few sips?” Students who reply in the affirmative are then asked additional questions about their alcohol consumption over different time frames: “On how many occasions (if any) have you had alcohol to drink—more than just a few sips... in your lifetime, ...in the last 12 months, ...in the last 30 days?” A subsequent question asks, “Think back over the last two weeks. How many times have you had five or more drinks in a row?” A drink is defined as a bottle of beer, a glass of wine, a shot glass of liquor, a mixed drink, etc.

National Survey on Drug Use & Health (NSDUH)—Starting in 1999, NSDUH information about the frequency of the

consumption of alcoholic beverages in the past 30 days has been obtained for all persons surveyed who are aged 12 and over. An extensive list of examples of the kinds of beverages covered is given to respondents prior to question administration. A drink is defined as a can or bottle of beer, a glass of wine or a wine cooler, a shot of liquor, or a mixed drink with liquor in it. Those times when the respondent had only a sip or two from a drink are not considered consumption. Alcohol use is based on the following questions: “During the past 30 days, on how many days did you drink one or more drinks of an alcoholic beverage?”, “On the days that you drank during the past 30 days, how many drinks did you usually have?”, and “During the past 30 days, on how many days did you have five or more drinks on the same occasion? By ‘occasion,’ we mean at the same time or within a couple of hours of each other.”

Any-listed diagnosis—See Appendix II, Diagnosis.

Average annual rate of change (percent change)—In *Health, United States*, average annual rates of change, or growth rates, are calculated as follows:

$$[(P_n / P_o)^{1/N} - 1] \times 100$$

where

P_n = later time period

P_o = earlier time period

N = number of years in interval.

This geometric rate of change assumes that a variable increases or decreases at the same rate during each year between the two time periods.

Average length of stay—The American Hospital Association computes average length of stay by dividing the number of inpatient days by the number of admissions. (Also see Appendix II, Days of care; Discharge; Inpatient.)

Basic actions difficulty—Basic actions difficulty is a composite measure of disability designed to capture limitations or difficulties in movement, emotional, sensory, or cognitive functioning associated with a health problem. Persons with more than one of these difficulties are counted only once in the estimates. The full range of functional areas cannot be assessed on the basis of National Health Interview Survey (NHIS) questions; however, the available questions can identify difficulty in the following core areas of functioning:

- Movement (walking, standing, sitting, bending or kneeling, reaching overhead, grasping objects with fingers, and lifting).
- Selected elements of emotional functioning, in particular, feelings that interfere with accomplishing daily activities. Respondents were classified based on responses to a series of questions that measure psychological distress.

- Sensory functioning, based on difficulties seeing or hearing.
- Selected elements in cognitive functioning, specifically difficulties with remembering, or experiencing confusion.

For many measures of disability, only disabilities resulting from an underlying condition that is chronic (based on nature and duration) are considered. However, whether the underlying conditions related to the core areas of basic actions difficulty were chronic was not a requirement in classifying persons. In *Health, United States*, respondents missing responses in a series of questions were classified as missing for that component. Respondents reporting that they “do not do this activity” were classified as missing for that activity. For hearing, respondents reporting that they were “deaf” or had “a lot of trouble” hearing without the use of hearing aids or other listening devices were coded as having a hearing limitation. For more information on how this measure was constructed using NHIS data, including the specific questions asked, see: Altman B, Bernstein A. Disability and health in the United States, 2001–2005. Hyattsville, MD: NCHS; 2008. Available from: <http://www.cdc.gov/nchs/data/misc/disability2001-2005.pdf>. (Also see Appendix II, Complex activity limitation; Hearing trouble.)

Bed, health facility—The American Hospital Association defines bed count as the number of beds, cribs, and pediatric bassinets that are set up and staffed for use by inpatients on the last day of the reporting period. In the Centers for Medicare & Medicaid Service’s Quality Improvement Evaluation System (QIES) (formerly the Online Survey Certification and Reporting [OSCAR]) database, all beds in certified facilities are counted on the day of certification inspection. (Also see Appendix II, Hospital; Occupancy rate.)

Binge drinking—Binge drinking is measured in the following data systems. (Also see Appendix II, Alcohol consumption.)

Monitoring the Future (MTF) Study—This school-based survey of secondary school students collects information on alcohol use by using self-completed questionnaires. To determine whether they have tried alcohol, students are asked a preliminary screening question: “Have you ever had any alcoholic beverage to drink—more than just a few sips?” Students who reply in the affirmative are then asked additional questions about their alcohol consumption, including one on binge drinking: “Think back over the last two weeks. How many times have you had five or more drinks in a row?” A drink is defined as a bottle of beer, a glass of wine, a shot glass of liquor, a mixed drink, etc. Information on binge drinking is obtained for 12th graders (starting in 1975) and for 8th and 10th graders (starting in 1991).

National Survey on Drug Use & Health (NSDUH)—In NSDUH, binge alcohol use is defined as “Five or more

drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) at least once in the past 30 days.” Heavy alcohol use is defined as “Five or more drinks on the same occasion (binge drinking) on at least 5 different days in the past 30 days.” (Also see Appendix II, Alcohol consumption.)

Birth cohort—A birth cohort consists of all persons born within a given period of time, such as a calendar year.

Birth rate—See Appendix II, Rate: Birth and related rates.

Birthweight—Birthweight is the first weight of the newborn obtained after birth. Low birthweight is defined as weighing less than 2,500 grams (5 lb 8 oz). Very low birthweight is defined as weighing less than 1,500 grams (3 lb 4 oz). Prior to 1979, low birthweight was defined as weighing 2,500 grams or less, and very low birthweight as weighing 1,500 grams or less.

Blood pressure, high—In *Health, United States*, a person is considered to have hypertension if they have measured high blood pressure (i.e., average measured systolic blood pressure of at least 140 mm Hg or diastolic pressure of at least 90 mm Hg) and/or if they report that they are taking a prescription medicine for high blood pressure (respondents were asked, “Are you now taking prescribed medicine for your high blood pressure?”). Uncontrolled high blood pressure is defined as having an average measured systolic blood pressure of at least 140 mm Hg or diastolic pressure of at least 90 mm Hg, among those with hypertension. Those with uncontrolled high blood pressure also may be taking prescribed medicine for high blood pressure. These blood pressure definitions are consistent with the following: National Heart, Lung, and Blood Institute. Seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. NIH pub no 04–5230. Bethesda, MD: National Institutes of Health; 2004. Available from: <http://www.nhlbi.nih.gov/guidelines/hypertension/jnc7full.pdf>; and Go AS, Bauman M, King SMC, Fonarow GC, Lawrence W, Williams KA, et al. AHA/ACC/CDC. An effective approach to high blood pressure control: A science advisory from the American Heart Association, the American College of Cardiology, and the Centers for Disease Control and Prevention. Hypertension 2013. Available from: <http://hyper.ahajournals.org/content/early/2013/11/14/HYP.0000000000000003.citation>.

Blood pressure data presented in *Health, United States* are from the National Health and Nutrition Examination Survey (NHANES). Blood pressure is measured by averaging up to three blood pressure readings taken for an NHANES participant. Blood pressure readings of 0 mm Hg are assumed to be in error and are not included in the estimates. The methods used to measure the blood pressure of participants have changed over the different NHANES survey years. Changes include the following:

- Number of blood pressure measurements taken (increased from one to four).

- Equipment maintenance procedures.
- Training of persons taking readings (physician, nurse, or interviewer).
- Proportion zero end-digits for systolic and diastolic readings.
- Published diastolic definition.
- Location where the measurements were taken (mobile examination center [MEC] or home).

In 1999 and subsequent years, blood pressure has been measured in the NHANES MEC by one of the MEC physicians. For people aged 8 and over, three consecutive blood pressure readings are obtained using the same arm. If a blood pressure measurement was interrupted or the measurer was unable to get one or more of the readings, a fourth attempt may be made. Both systolic and diastolic measurements are recorded to the nearest even number.

In NHANES III, three sets of blood pressure measurements were taken in the MEC for examinees aged 5 and over. Blood pressure measurements were also taken by trained interviewers during the household interview, on sample persons aged 17 and over. Systolic and diastolic average blood pressures were computed as the arithmetic mean of six or fewer measurements obtained at the household interview (maximum of three) and the MEC examination (maximum of three). If the examinee did not have blood pressure measurements taken in the MEC, this variable was calculated from measurements taken at the household interview. Both systolic and diastolic measurements were recorded to the nearest even number.

For more information on changes in blood pressure measurement in NHANES up to 1991, see: Burt VL, Cutler JA, Higgins M, Horan MJ, Labarthe D, Whelton P, et al. Trends in the prevalence, awareness, treatment, and control of hypertension in the adult U.S. population: Data from the health examination surveys, 1960 to 1991. *Hypertension* 1995;26(1):60–9.

Body mass index (BMI)—BMI is a measure that adjusts body weight for height. It is calculated as weight in kilograms divided by height in meters squared. Normal weight for adults is defined as a BMI of 18.5 to less than 25.0; overweight or obese is greater than or equal to 25.0; and obesity is greater than or equal to 30.0. Within the obesity category, Grade 1 obesity is defined as a BMI of 30.0 to less than 35.0; Grade 2 is 35.0 to less than 40.0; and Grade 3 is 40.0 or greater. Prior to assigning a person to a BMI category, BMI is rounded to one decimal place. In *Health, United States*, the NHANES variable Body Mass Index is used to assign persons to BMI categories. BMI cut points are defined in the following: National Heart, Lung, and Blood Institute. Managing overweight and obesity in adults: Systematic evidence review from the Obesity Expert Panel. Bethesda, MD: National Institutes of Health; 2013. Available from: <https://www.nhlbi.nih.gov/health-pro/guidelines/in-develop/obesity-evidence-review>; Jensen MD, Ryan DH,

Apovian CM, Ard JD, Comuzzie AG, Donato KA, et al. 2013 AHA/ACC/TOS guideline for the management of overweight and obesity in adults: A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and the Obesity Society. *Circulation*; 2013. Available from: <http://circ.ahajournals.org/content/early/2013/11/11/01.cir.0000437739.71477.ee.citation>; and HHS. Healthy People 2020: Nutrition, physical activity, and obesity; 2012. Available from: <http://www.healthypeople.gov/2020/Leading-Health-Indicators>.

Obesity for children and adolescents is defined as a BMI at or above the sex- and age-specific 95th percentile from the 2000 CDC Growth Charts (<http://www.cdc.gov/growthcharts/>). The age used is age in months from the age at time of examination. Also see, Kuczmarski RJ, Ogden CL, Guo SS, et al. 2000 CDC Growth Charts for the United States: methods and development. *Vital Health Stat* 11. 2002 May; (246):1–190. Available at: http://www.cdc.gov/nchs/data/series/sr_11/sr11_246.pdf. Starting with *Health, United States, 2010*, the terminology describing excess weight among children changed from previous editions. The term obesity now refers to children who were formerly labeled as overweight. This is a change in terminology only and not a change in measurement. For more information, see: Ogden CL, Flegal KM. Changes in terminology for childhood overweight and obesity. *National health statistics report*; no 25. Hyattsville, MD: NCHS; 2010. Available from: <http://www.cdc.gov/nchs/data/nhsr/nhsr025.pdf>.

Cause of death—For the purpose of national mortality statistics, every death is attributed to one underlying condition, based on information reported on the death certificate and using the international rules for selecting the underlying cause of death from the conditions stated on the certificate. The underlying cause is defined by the World Health Organization (WHO) as “the disease or injury that initiated the train of events leading directly to death, or the circumstances of the accident or violence that produced the fatal injury.” Generally, more medical information is reported on death certificates than is directly reflected in the underlying cause of death. Conditions that are not selected as the underlying cause of death constitute the nonunderlying causes of death, also known as multiple cause of death.

Cause of death is coded according to the appropriate revision of the *International Classification of Diseases* (ICD) (Table III). Effective with deaths occurring in 1999, the United States began using the 10th revision of the ICD (ICD–10); during the period 1979–1998, causes of death were coded and classified according to the 9th revision (ICD–9). Table IV lists ICD codes for the 6th through 10th revisions for causes of death shown in *Health, United States*. In *Health, United States*, common terms are sometimes used in the text in place of medical terminology. Examples include “cancer” for “malignant neoplasm” and “kidney disease” for “nephritis, nephrotic syndrome and nephrosis.”

Table III. Revision of the *International Classification of Diseases (ICD)*, by year of conference in which adopted and years in use in the United States

ICD revision	Year of conference in which adopted	Years in use in United States
1st	1900	1900–1909
2nd	1909	1910–1920
3rd	1920	1921–1929
4th	1929	1930–1938
5th	1938	1939–1948
6th	1948	1949–1957
7th	1955	1958–1967
8th	1965	1968–1978
9th	1975	1979–1998
10th	1990	1999–present

SOURCE: CDC/NCHS. Available from: <http://www.cdc.gov/nchs/icd.htm>.

Each ICD revision has produced discontinuities in cause-of-death trends. These discontinuities are measured by using comparability ratios that are essential to the interpretation of mortality trends. For further discussion, see: http://www.cdc.gov/nchs/nvss/mortality/comparability_icd.htm. (Also see Appendix II, Comparability ratio; *International Classification of Diseases* [ICD]; and Appendix I, National Vital Statistics System [NVSS]; Multiple Cause-of-Death File.)

Cause-of-death ranking—Selected causes of death of public health and medical importance are compiled into tabulation lists and are ranked according to the number of deaths assigned to these causes. The top-ranking causes determine the leading causes of death. Certain causes on the tabulation lists are not ranked if, for example, the category title represents a group title (such as “Major cardiovascular diseases” and “Symptoms, signs, and abnormal clinical and laboratory findings, not elsewhere classified”) or the category title begins with the words “Other” or “All other.” In addition, when one of the titles that represents a subtotal (such as “Malignant neoplasms”) is ranked, its component parts are not ranked. The tabulation lists used for ranking in the 10th revision of the *International Classification of Diseases* (ICD–10) include the List of 113 Selected Causes of Death, which replaces the ICD–9 List of 72 Selected Causes, HIV Infection and Alzheimer’s Disease; and the ICD–10 List of 130 Selected Causes of Infant Death, which replaces the ICD–9 List of 60 Selected Causes of Infant Death and HIV Infection. Causes that are tied receive the same rank; the next cause is assigned the rank it would have received had the lower-ranked causes not been tied, that is, a rank is skipped. For more information, see the annual series of “Deaths: Final Data” and “Deaths: Leading Causes” reports, available from: <http://www.cdc.gov/nchs/products/nvsr.htm>. (Also see Appendix II, *International Classification of Diseases* [ICD].)

Children’s Health Insurance Program (CHIP)—Title XXI of the Social Security Act, often referred to as the Children’s Health Insurance Program (CHIP), is a program originally

enacted by the Balanced Budget Act of 1997. The Children’s Health Insurance Program Reauthorization Act of 2009 (CHIPRA, P.L. 111–3) reauthorized CHIP and appropriated funding for CHIP through FY 2013. The Affordable Care Act of 2010 (ACA, P.L. 111–148) extends CHIP funding through FY 2015, and the Medicare Access and CHIP Reauthorization Act of 2015 (P.L. 114–10) extended funding with no programmatic changes for CHIP through 2017. CHIP provides federal funds for states to provide health care coverage to eligible low-income, uninsured children whose income is too high to qualify for Medicaid. Generally CHIP is only available through age 18. CHIP gives states broad flexibility in program design within a federal framework that includes important beneficiary protections. Funds from CHIP may be used for a separate child health program or to expand Medicaid. Although CHIP is not part of Medicaid, in some instances in *Health, United States*, data on CHIP and Medicaid are presented together, and those instances are discussed in the footnotes of the respective tables. For more information, see: <https://www.medicare.gov/chip/chip-program-information.html>. (Also see Appendix II, Health insurance coverage; Medicaid.)

Cholesterol—Serum total cholesterol is a combination of high-density lipoprotein (HDL) cholesterol, low-density lipoprotein (LDL) cholesterol, and very low-density lipoprotein (VLDL) cholesterol and is highly correlated with LDL cholesterol. High serum total cholesterol is a risk factor for cardiovascular disease (see Wilson PW, D’Agostino RB, Levy D, Belanger AM, Silbershatz H, Kannel WB. Prediction of coronary heart disease using risk factor categories. *Circulation* 97(18):1837–47. 1998). In its 2002 report on high blood cholesterol, the National Cholesterol Education Program Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III, or ATP III) considered a serum total cholesterol value greater than or equal to 240 mg/dL (6.20 mmol/L) as high. A more recent set of guidelines—the result of a collaboration among the National Heart, Lung, and Blood Institute; the American College of Cardiology; and the American Heart Association—focused on which groups of people could benefit from statin use, based on their risk factors. Because *Health, United States* focuses on providing population-level prevalence data rather than individual-level estimates, three broad indicators of cholesterol are presented based on measured serum total cholesterol level and the reported use of cholesterol-lowering medications. Cholesterol levels are determined using the NHANES T_CHOL file. For more information on the current cholesterol guidelines, see: Management of blood cholesterol in adults: Systematic evidence review from the Cholesterol Expert Panel. Bethesda, MD: National Institutes of Health, National Heart, Lung, and Blood Institute; 2013. Available from: <http://www.nhlbi.nih.gov/health-pro/guidelines/in-develop/cholesterol-in-adults>; and Stone NJ, Robinson JG, Lichtenstein AH, Merz CNB, Blum CB, Eckel RH, et al. 2013 ACC/AHA guideline on the treatment of blood cholesterol to reduce atherosclerotic cardiovascular risk in

Table IV. Cause-of-death codes, by applicable revision of the *International Classification of Diseases (ICD)*

<i>Cause of death (10th Revision titles)</i>	<i>6th and 7th Revisions</i>	<i>8th Revision</i>	<i>9th Revision</i>	<i>10th Revision</i>
Communicable diseases	001–139, 460–466, 480–487, 771.3	A00–B99, J00–J22
Chronic and noncommunicable diseases	140–459, 470–478, 490–799	C00–I99, J30–R99
Meningococcal infection	036	A39
Septicemia	038	A40–A41
Human immunodeficiency virus (HIV) disease ¹	*042–*044	B20–B24
Malignant neoplasms	140–205	140–209	140–208	C00–C97
Colon, rectum, and anus	153–154	153–154	153–154	C18–C21
Trachea, bronchus, and lung	162–163	162	162	C33–C34
Breast	170	174	174–175	C50
Prostate	177	185	185	C61
In situ neoplasms, Benign neoplasms, and Neoplasms of uncertain or unknown behavior	210–239	210–239	210–239	D00–D48
Diabetes mellitus	260	250	250	E10–E14
Anemias	280–285	D50–D64
Meningitis	320–322	G00, G03
Alzheimer's disease	331.0	G30
Diseases of heart	400–402, 410–443	390–398, 402, 404, 410–429	390–398, 402, 404, 410–429	I00–I09, I11, I13, I20–I51
Ischemic heart disease	410–414, 429.2	I20–I25
Essential hypertension and hypertensive renal disease	I10, I12, I15
Cerebrovascular diseases	330–334	430–438	430–434, 436–438	I60–I69
Atherosclerosis	440	I70
Influenza and pneumonia ²	480–483, 490–493	470–474, 480–486	480–487	J09–J18
Chronic lower respiratory diseases	241, 501, 502, 527.1	490–493, 519.3	490–494, 496	J40–J47
Chronic liver disease and cirrhosis	581	571	571	K70, K73–K74
Nephritis, nephrotic syndrome, and nephrosis	580–589	N00–N07, N17–N19, N25–N27
Pregnancy, childbirth, and the puerperium	640–689	630–678	630–676	O00–O99
Congenital malformations, deformations, and chromosomal abnormalities	740–759	Q00–Q99
Certain conditions originating in the perinatal period	760–779	P00–P96
Newborn affected by maternal complications of pregnancy	761	P01
Newborn affected by complications of placenta, cord, and membranes	762	P02
Disorders related to short gestation and low birthweight, not elsewhere classified	765	P07
Birth trauma	767	P10–P15
Intrauterine hypoxia and birth asphyxia	768	P20–P21
Respiratory distress of newborn	769	P22
Bacterial sepsis of newborn	P36
Necrotizing enterocolitis of newborn	777.5	P77
Sudden infant death syndrome	798.0	R95

See footnotes at end of table.

Table IV. Cause-of-death codes, by applicable revision of the *International Classification of Diseases (ICD)*—Con.

<i>Cause of death (10th Revision titles)</i>	<i>6th and 7th Revisions</i>	<i>8th Revision</i>	<i>9th Revision</i>	<i>10th Revision</i>
Occupational diseases:				
Angiosarcoma of liver	C22.3
Malignant mesothelioma	158.8, 158.9, 163	C45
Pneumoconiosis	500–505	J60–J66
Coal workers' pneumoconiosis	500	J60
Asbestosis	501	J61
Silicosis	502	J62
Other (including unspecified)	503–505	J63–J66
Injuries ²	E800–E869, E880–E929, E950–E999	*U01–*U03, V01–Y36, Y85–Y87, Y89
Unintentional injuries ³	E800–E936, E960–E965	E800–E929, E940–E946	E800–E869, E880–E929	V01–X59, Y85–Y86
Motor vehicle-related injuries ³	E810–E835	E810–E823	E810–E825	V02–V04, V09.0, V09.2, V12–V14, V19.0–V19.2, V19.4–V19.6, V20–V79, V80.3–V80.5, V81.0– V81.1, V82.0–V82.1, V83–V86, V87.0–V87.8, V88.0–V88.8, V89.0, V89.2
Poisoning	E870–E888, E890–E895	E850–E877	E850–E869	X40–X49
Suicide ²	E963, E970– E979	E950–E959	E950–E959	*U03, X60–X84, Y87.0
Homicide ²	E964, E980– E983	E960–E969	E960–E969	*U01–*U02, X85–Y09, Y87.1
Firearm-related injury	E922, E955, E965, E970, E985	E922, E955.0– E955.4, E965.0–E965.4, E970, E985.0– E985.4	*U01.4, W32–W34, X72–X74, X93–X95, Y22–Y24, Y35.0
Injury by drug poisoning	X40–X44, X60–X64, X85, Y10–Y14
Heroin	X40–X44, X60–X64, X85, Y10–Y14 (underlying cause) and T40.1 (multiple cause)
Opioid analgesics	X40–X44, X60–X64, X85, Y10–Y14 (underlying cause) and T40.2–T40.4 (multiple cause)

... Cause-of-death codes are not provided for causes not shown in *Health, United States*.

¹Categories for coding human immunodeficiency virus (HIV) infection were introduced in 1987. The asterisk (*) indicates codes that are not part of ICD–9.

²Starting with 2001 data, NCHS introduced categories *U01–*U03 for classifying and coding deaths due to acts of terrorism. The asterisk (*) indicates codes that are not part of ICD–10. Starting with 2007 data, NCHS introduced the category J09 for coding avian influenza virus. In 2009, the title for the ICD–10 code J09 was changed from Influenza due to identified avian Influenza virus to Influenza due to certain identified influenza virus. This change was made to accommodate deaths from influenza A (H1N1) virus in the ICD–10 code J09 for data years 2009 and beyond.

³In the public health community, the term unintentional injuries is preferred to accidents, and the term motor vehicle-related injuries is preferred to motor vehicle accidents.

SOURCE: CDC/NCHS. Advance report: Final mortality statistics, 1974. Monthly vital statistics report; vol 24 no 11 suppl. Hyattsville, MD: NCHS; 1976. Available from: http://www.cdc.gov/nchs/data/mvsr/supp/mv24_11sacc.pdf.

Hoyert DL, Kochanek KD, Murphy SL. Deaths: Final data for 1997. National vital statistics reports; vol 47 no 19. Hyattsville, MD: NCHS; 1999. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr47/nvs47_19.pdf.

Hoyert DL, Heron MP, Murphy SL, Kung H-C. Deaths: Final data for 2003. National vital statistics reports; vol 54 no 13. Hyattsville, MD: NCHS; 2006. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr54/nvsr54_13.pdf.

Murphy SL, Xu JQ, Kochanek KD. Deaths: Final data for 2010. National vital statistics reports; vol 61 no 4. Hyattsville, MD: NCHS; 2013. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr61/nvsr61_04.pdf.

Xu JQ, Murphy SL, Kochanek KD, et al. Deaths: Final data for 2013. National vital statistics reports; vol 64 no 2. Hyattsville, MD: NCHS; 2015. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr64/nvsr64_02.pdf.

adults: A report of the American College of Cardiology/ American Heart Association Task Force on Practice Guidelines. *Circulation*. 2014;129:S1–45. Available from: http://circ.ahajournals.org/content/129/25_suppl_2/S1.full.

In *Health, United States*, three measures of total cholesterol are presented: hypercholesterolemia, high serum total cholesterol, and mean serum total cholesterol. Hypercholesterolemia is based on both laboratory testing and self-reported medication use. It is defined as measured serum total cholesterol greater than or equal to 240 mg/dL or reporting taking cholesterol-lowering medications. Respondents who were told by a doctor or health professional that their cholesterol was high, and were told by a doctor to take cholesterol-lowering medication and who answered “yes” to the question, “Are you now following this advice?” were classified as taking cholesterol-lowering medication. High serum total cholesterol is defined as measured serum total cholesterol greater than or equal to 240 mg/dL (6.20 mmol/L). Both high serum cholesterol and mean serum total cholesterol are based on serum samples collected during the National Health and Nutrition Examination Survey (NHANES) examination.

Venous blood serum samples collected from NHANES participants at mobile examination centers were frozen and shipped on dry ice to the laboratory conducting the lipid analyses. Serum total cholesterol was measured on all examined adults regardless of whether they had fasted, and data were analyzed regardless of fasting status. Cholesterol measurements are standardized according to the criteria of CDC—and later the CDC–National Heart, Lung, and Blood Institute Cholesterol Standardization Program—to ensure comparable and accurate measurements. For more information, see: Myers GL, Cooper GR, Winn CL, Smith SJ. The Centers for Disease Control–National Heart, Lung, and Blood Institute Lipid Standardization Program: An approach to accurate and precise lipid measurements. *Clin Lab Med* 1989;9(1):105–35. A detailed summary of the procedures used for measurement of total cholesterol in the earlier NHANES survey years has been published in: Carroll MD, Kit BK, Lacher DA, Shero ST, Mussolino ME. Trends in lipids and lipoproteins in US adults, 1988–2010. *JAMA* 2012;308(15):1545–54. A description of the laboratory procedures for the total cholesterol measurement for different NHANES survey years is published by NCHS and is available from: <http://www.cdc.gov/nchs/nhanes.htm>.

Cigarette smoking—Cigarette smoking and related tobacco use are measured in the following data systems.

Monitoring the Future (MTF) Study—Information on current cigarette smoking was obtained for 12th graders (starting in 1975) and for 8th and 10th graders (starting in 1991), based on the following question: “How frequently have you smoked cigarettes during the past 30 days?”

National Health Interview Survey (NHIS)—Information about cigarette smoking is obtained for adults aged 18

and over. Starting in 1993, current smokers are identified by asking the following two questions: “Have you smoked at least 100 cigarettes in your entire life?” and “Do you now smoke cigarettes every day, some days, or not at all?” Persons who smoked 100 cigarettes and who now smoke every day or some days were defined as current smokers. Before 1992, current smokers were identified based on positive responses to the following two questions: “Have you smoked 100 cigarettes in your entire life?” and “Do you smoke now?” (traditional definition). In 1992, the definition of current smoker in NHIS was modified to separately identify persons who smoked every day and those who smoked on some days (revised definition). In 1992, cigarette smoking data were collected for a half-sample, with one-half of respondents (one-quarter sample) answering the traditional smoking questions and the other one-half (one-quarter sample) answering the revised smoking question, “Do you smoke every day, some days, or not at all?” An unpublished analysis of the 1992 traditional smoking measure revealed that the crude percentage of current smokers aged 18 and over remained the same as for 1991. The estimates for 1992 shown in *Health, United States* combine data collected using both the traditional and revised questions. Estimates for 1993 and beyond use the revised questions.

In 1993–1995, estimates of cigarette smoking prevalence were based on a half-sample. Smoking data were not collected in 1996. Starting in 1997, smoking data were collected in the Sample Adult questionnaire. Starting in 2014, a question was added to the survey on the use of electronic cigarettes, often referred to as e-cigarettes. Electronic cigarette use was not considered in the definition of current cigarette smoking. For more information on e-cigarette use, see: Schoenborn CA, Gindi RM. Electronic cigarette use among adults: United States, 2014. NCHS data brief, no 217. Hyattsville, MD: NCHS. 2015. Available from: <http://www.cdc.gov/nchs/data/databriefs/db217.htm>. For more information on survey methodology and sample sizes pertaining to NHIS cigarette smoking data, see the NHIS Adult Tobacco Use Information website at: <http://www.cdc.gov/nchs/nhis/tobacco.htm>.

National Survey on Drug Use & Health (NSDUH)—Information on current cigarette smoking is obtained for all persons surveyed who are aged 12 and over, based on the following question: “Now think about the past 30 days, that is, from [DATE] up to and including today. During the past 30 days, have you smoked part or all of a cigarette?” Electronic cigarette use was not considered in the definition of current cigarette smoking.

Civilian noninstitutionalized population; Civilian population—See Appendix II, Population.

Colorectal tests or procedures—Colorectal tests or procedures are used to detect polyps, abnormal cell growth,

lesions, and other gastrointestinal conditions, including colon cancer. These tests may include home fecal occult blood tests, sigmoidoscopy, or colonoscopy. The time interval between screenings varies, depending on the type of test as well as individual risk factors and prior screening history.

In the National Health Interview Survey, questions about colorectal tests or procedures were asked of respondents aged 40 and over on an intermittent schedule, and the questions varied over time.

In 2000, 2003, 2005, and 2008, respondents were asked, "Have you ever had a sigmoidoscopy, colonoscopy, or proctoscopy?" In 2010 and 2013, respondents were asked two separate questions: "Have you ever had a colonoscopy?" and "Have you ever had a sigmoidoscopy?" An additional question about colorectal testing, "Have you ever had a blood stool test using a home testing kit?" was asked in all of these survey years.

Respondents who replied that they had a colorectal test or procedure were asked subsequent questions about the month, year, and time since their most recent test or procedure. In 2000 and 2003, if respondents did not provide the year of, or the time since, their most recent colorectal exam, they were asked about the time frame of their most recent exam (i.e., whether they had the exam a year ago or less, more than 1 year ago but not more than 2 years ago, more than 2 years ago but not more than 3 years ago, more than 3 years ago but not more than 5 years ago, more than 5 years ago but not more than 10 years ago, or over 10 years ago). For adults who provided the year, but not the month, of their most recent exam, the exam date was coded as July 15 of the provided year.

In 2005, 2008, 2010, and 2013, the questionnaire skip pattern was modified so that respondents giving an incomplete or partial date (missing month or year) of their most recent colorectal exam were asked a follow-up question about the time since their most recent exam (i.e., whether they had the exam a year ago or less, more than 1 year ago but not more than 2 years ago, more than 2 years ago but not more than 3 years ago, more than 3 years ago but not more than 5 years ago, more than 5 years ago but not more than 10 years ago, or over 10 years ago). In 2010 only, additional questions on the use of virtual or CT colonoscopy were included in the questionnaire, but these questions were not used to determine whether respondents had a colorectal test or procedure.

Colorectal screening tests and procedures may be used for diagnostic or screening purposes. The current recommendation, made by the U.S. Preventive Services Task Force in 2008, is for the use of fecal occult blood testing, sigmoidoscopy, or colonoscopy in adults aged 50 to 75. However, these recommendations were currently undergoing review by the U.S. Preventive Services Task Force at the time this report was prepared. For a summary of current colorectal screening recommendations and the status of the

review, see: <http://www.uspreventiveservicestaskforce.org/uspstf/uspscoco.htm>.

In *Health, United States*, estimates of colorectal tests are presented for adults aged 50–75 who had any colorectal test or procedure (defined as reporting a home fecal occult blood test [FOBT] in the past year, a sigmoidoscopy procedure in the past 5 years with FOBT in the past 3 years, or a colonoscopy in the past 10 years).

Community hospital—See Appendix II, Hospital.

Comparability ratio—About every 10 to 20 years, the *International Classification of Diseases* (ICD) is revised to stay abreast of advances in medical science and changes in medical terminology. Each of these revisions produces breaks in the continuity of cause-of-death statistics because of changes in classification and in the rules for selecting an underlying cause of death. Classification and rule changes affect cause-of-death trend data by shifting deaths away from some cause-of-death categories and into others. Comparability ratios measure the effect of changes in classification and coding rules. For the causes shown in Table V, comparability ratios range between 0.6974 and 1.5812. Influenza and pneumonia had the lowest comparability ratio (0.6974), indicating that this cause is about 30% less likely to be selected as the underlying cause of death under ICD–10 than under ICD–9. Alzheimer's disease had the highest comparability ratio (1.5812), indicating that Alzheimer's disease is 58% more likely to be selected as the underlying cause when ICD–10 coding is used.

For selected causes of death, the ICD–9 codes used to calculate death rates for 1980–1998 differ from the ICD–9 codes most nearly comparable with the corresponding ICD–10 cause-of-death category, which also affects the ability to compare death rates across ICD revisions. Examples of these causes are Ischemic heart disease; Cerebrovascular diseases; Trachea, bronchus, and lung cancer; Unintentional injuries; and Homicide. To address this source of discontinuity, mortality trends for 1980–1998 were recalculated using ICD–9 codes that are more comparable with codes for corresponding ICD–10 categories. Table IV shows the ICD–9 codes used for these causes. This modification may lessen the discontinuity between the 9th and 10th revisions, but the effect on the discontinuity between the 8th and 9th revisions is not measured.

Comparability ratios shown in Table V are based on a comparability study in which the same deaths were coded using both the 9th and 10th revisions. The comparability ratio was calculated by dividing the number of deaths classified by ICD–10 by the number of deaths classified by ICD–9. The resulting ratios represent the net effect of the 10th revision on cause-of-death statistics and can be used to adjust mortality statistics for causes of death classified by the 9th revision to be comparable with cause-specific mortality statistics classified by the 10th revision.

Table V. Comparability of selected causes of death between the 9th and 10th revisions of the *International Classification of Diseases (ICD)*

<i>Cause of death</i> ¹	<i>Final comparability ratio</i> ²
Human immunodeficiency virus (HIV) disease	1.0821
Malignant neoplasms	1.0093
Colon, rectum, and anus	0.9988
Trachea, bronchus, and lung	0.9844
Breast	1.0073
Prostate	1.0144
Diabetes mellitus	1.0193
Alzheimer's disease	1.5812
Diseases of heart	0.9852
Ischemic heart diseases	1.0006
Essential (primary) hypertension and hypertensive renal disease	1.1162
Cerebrovascular diseases	1.0502
Influenza and pneumonia	0.6974
Chronic lower respiratory diseases	1.0411
Chronic liver disease and cirrhosis	1.0321
Nephritis, nephrotic syndrome and nephrosis	1.2555
Pregnancy, childbirth, and the puerperium	1.1404
Unintentional injuries	1.0251
Motor vehicle-related injuries	0.9527
Poisoning	1.0365
Suicide	1.0022
Homicide	1.0020
Firearm-related injury	1.0012
Chronic and noncommunicable diseases	1.0100
Injuries	1.0159

¹See Table IV for ICD–9 and ICD–10 cause-of-death codes.

²Ratio of number of deaths classified by ICD–10 to number of deaths classified by ICD–9.

SOURCE: CDC/NCHS. Final comparability ratios for 113 selected causes of death. Available from: ftp://ftp.cdc.gov/pub/Health_Statistics/NCHS/Datasets/Comparability/icd9_icd10/Comparability_Ratio_tables.xls.

Miniño M, Anderson RN, Fingerhut LA, Boudreaux MA, Warner M. Deaths: Injuries, 2002. National vital statistics reports; vol 54 no 10. Hyattsville, MD: NCHS; 2006. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr54/nvsr54_10.pdf.

The application of comparability ratios to mortality statistics helps make the analysis of change between 1998 and 1999 more accurate and complete. The 1998 comparability-modified death rate is calculated by multiplying the comparability ratio by the 1998 death rate. Comparability-modified rates should be used to estimate mortality change between 1998 and 1999.

Caution should be used when applying the comparability ratios presented in Table V to age-, race-, and sex-specific mortality data. Demographic subgroups may sometimes differ with regard to their cause-of-death distribution, and this would result in demographic variation in cause-specific comparability ratios.

For more information, see: Anderson RN, Miniño M, Hoyert DL, Rosenberg HM. Comparability of cause of death between ICD–9 and ICD–10: Preliminary estimates. National vital statistics reports; vol 49 no 2. Hyattsville, MD: NCHS; 2001. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr49/nvsr49_02.pdf; Kochanek KD, Smith BL, Anderson RN. Deaths: Preliminary data for 1999. National vital statistics reports; vol 49 no 3. Hyattsville, MD: NCHS; 2001. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr49/nvsr49_03.pdf; Final ratios for 113 selected causes of death, available from: ftp://ftp.cdc.gov/pub/Health_Statistics/NCHS/Datasets/Comparability/icd9_icd10/; and the ICD comparability ratio website at: http://www.cdc.gov/nchs/nvss/mortality/comparability_icd.htm. (Also see Appendix II, Cause of death; *International Classification of Diseases* [ICD].)

Compensation—See Appendix II, Employer costs for employee compensation.

Complex activity limitation—Complex activity limitation is a composite measure of disability constructed to measure disability as defined by the inability to function successfully in certain social roles. Complex activities consist of the tasks and organized activity that make up numerous social roles such as working, maintaining a household, living independently, or participating in community activities. Complex activity performance requires the execution of a combination of core areas of functioning. Complex activities include the following:

- Maintaining independence, including self-care and the ability to carry out activities associated with maintaining a household, such as shopping, cooking, and taking care of bills (measures are based on questions commonly known as activities of daily living [ADLs] and instrumental activities of daily living [IADLs]). Limitations in these activities usually reflect severe restrictions and are associated with limitations in other complex activities.
- Difficulties experienced with social and leisure activities—represented in this measure by using questions about attending movies or sporting events, visiting friends, or pursuing hobbies or relaxation activities.
- Perceived limitation in the ability to work (a core aspect of social participation for the majority of the U.S. population)—represented by the respondent's self-defined limitation in the kind or amount of work they can do or their inability to work at a job or business.

For many measures of disability, only disabilities resulting from an underlying condition that is chronic (based on nature and duration) are considered. However, whether the underlying conditions related to the complex activities were chronic was not a requirement in classifying persons as having a complex activity limitation. In *Health, United States*, respondents missing responses in a series of questions were classified as missing for that component. Respondents

reporting that they “do not do this activity” were classified as missing for that activity. For more information on how this measure was constructed using data from the National Health Interview Survey, including the specific questions asked, see: Altman B, Bernstein A. Disability and health in the United States, 2001–2005. Hyattsville, MD: NCHS; 2008. Available from: <http://www.cdc.gov/nchs/data/misc/disability2001-2005.pdf>. (Also see Appendix II, Activities of daily living [ADL]; Basic actions difficulty; Instrumental activities of daily living [IADL].)

Consumer Price Index (CPI)—The CPI, prepared by the U.S. Bureau of Labor Statistics, is a monthly measure of the average change in prices of goods and services purchased by urban households. The medical care component of the CPI shows trends in medical care prices based on specific indicators of hospital, medical, and drug prices. A revised definition of the CPI has been in use since January 1988. (Also see Appendix II, Gross domestic product [GDP]; and Health expenditures, national.)

Contraception—The National Survey of Family Growth collects information on contraceptive use as reported by women aged 15–44. To determine current contraceptive use, women were asked to identify up to 4, out of 21, contraceptive methods they had used during the month of interview. Contraceptive methods listed as “other methods” in 2011–2013 included emergency contraception, contraceptive ring, female condom/vaginal pouch, foam, cervical cap, Today-brand sponge, suppository or insert, jelly or cream (without diaphragm), and other methods. Previously, contraceptive methods listed as “other methods” included the following: for 2006–2010, the contraceptive ring, female condom/vaginal pouch, foam, cervical cap, Today-brand sponge, suppository or insert, jelly or cream (without diaphragm), and other methods; for 2002, the female condom, foam, cervical cap, Today sponge, suppository or insert, jelly or cream (without diaphragm), or other method; for 1995, the female condom or vaginal pouch, foam, cervical cap, Today sponge, suppository or insert, jelly or cream, or other method; for 1988, foam, douche, Today sponge, suppository or insert, jelly or cream, or other method; and for 1982, foam, douche, suppository or insert, or other method.

Cost-charge ratio—The Agency for Healthcare Research and Quality’s Healthcare Cost and Utilization Project (HCUP) contains data on total charges per discharge as reported on the hospital discharge record. This charge information represents the amount the hospital billed for services but does not reflect how much hospital services actually cost or the specific amounts that hospitals received in payment. Data on costs may be of more interest to some users. The HCUP Cost-to-Charge ratio files convert charges to costs. Each file contains hospital-specific cost-to-charge ratios based on all-payer inpatient cost for nearly every hospital in HCUP. Cost information was obtained from hospital cost reports collected by the Centers for Medicare & Medicaid

Services. Some imputations for missing values were necessary. These files are unique by year.

Critical access hospital—See Appendix II, Hospital.

Crude birth rate; Crude death rate—See Appendix II, Rate: Birth and related rates; Rate: Death and related rates.

Days of care—Days of care is defined by the American Hospital Association as the number of adult and pediatric days of care rendered during the entire reporting period. Days of care for newborns are excluded. (Also see Appendix II, Admission; Average length of stay; Discharge; Hospital; Hospital utilization; Inpatient.)

Death rate—See Appendix II, Rate: Death and related rates.

Dental caries—Dental caries is evidence of decay on the crown or enamel surface of a tooth (i.e., coronal caries) and includes treated and untreated caries. Untreated dental caries refers to decay on the crown or enamel surface of a tooth (i.e., coronal caries) that has not been treated or filled. Decay in the root (i.e., root caries) was not included.

In *Health, United States*, estimates on the presence of caries are based on evaluation of primary and permanent teeth for persons aged 5 and over. The third molars were not included. Persons without at least one natural tooth (primary or permanent) were classified as edentulous (without any teeth) and were excluded. The majority of edentulous persons are aged 65 and over. Estimates of edentulism among persons aged 65 and over are 33% in 1988–1994, 23% in 2005–2008, and 19% in 2011–2012.

Dental caries was identified by an oral examination as part of the National Health and Nutrition Examination Survey (NHANES). Over time, there have been changes in the NHANES oral health examination process, ages examined, and methodology. During 1988–1994, a full-mouth complete oral health exam was conducted by a trained dentist on those aged 1 and over. During 1999–2004, a full-mouth complete oral health exam was conducted by a trained dentist on those aged 2 and over. During 2005–2008, data were collected for those aged 5 and over by a trained health technologist using the Basic Screening Examination (BSE), a simplified screening process to collect information on untreated caries, dental restorations, and dental sealants. During 2009–2010, the BSE was conducted by a trained dental hygienist on those aged 3–19. During 2005–2008 and 2009–2010, the use of the BSE does not allow us to determine if untreated decay was found in permanent teeth or primary teeth. For 2011–2012 data, a full-mouth complete oral health exam was conducted by a trained dentist on those aged 1 and over.

For more information, see: Dye BA, Barker LK, Li X, Lewis BG, Beltrán-Aguilar ED. Overview and quality assurance for the oral health component of the National Health and Nutrition Examination Survey (NHANES), 2005–08. *J Public Health Dent* 2011;71(1):54–61; and the following NHANES

resources: http://www.cdc.gov/nchs/nhanes/nhanes2005-2006/OHX_D.htm, http://www.cdc.gov/nchs/nhanes/nhanes2007-2008/OHX_E.htm, http://www.cdc.gov/nchs/nhanes/nhanes2009-2010/OHXDEN_F.htm, and http://www.cdc.gov/nchs/nhanes/2011-2012/OHXDEN_G.htm.

Dental visit—Starting in 1997, National Health Interview Survey respondents were asked, “About how long has it been since you last saw or talked to a dentist? Include all types of dentists, such as orthodontists, oral surgeons, and all other dental specialists as well as hygienists.” Starting in 2001, the question was modified slightly to ask respondents how long it had been since they last saw a dentist. Questions about dental visits were not asked for children under age 2 for years 1997–1999 and under age 1 for years 2000 and beyond. Starting with 1997 data, estimates are presented for people with a dental visit in the past year.

Diabetes—Diabetes is a group of conditions in which insulin is not adequately secreted or utilized. Diabetes is a leading cause of disease and death in the United States. Using data from the National Health and Nutrition Examination Survey (NHANES), three measures of diabetes are presented in *Health, United States*: physician-diagnosed diabetes, undiagnosed diabetes, and total diabetes. Physician-diagnosed diabetes data were obtained by self-report. Respondents who answered “yes” to the question, “Other than during pregnancy, have you ever been told by a doctor or health professional that you have diabetes or sugar diabetes?” were classified as having physician-diagnosed diabetes.

Only respondents who were not classified as having physician-diagnosed diabetes were evaluated to determine if they had undiagnosed diabetes. Undiagnosed diabetes was based on the results of laboratory testing of whole blood and blood plasma samples collected from NHANES participants at mobile examination centers. Undiagnosed diabetes was defined as a fasting plasma glucose (FPG) of at least 126 mg/dL or a hemoglobin A1c of at least 6.5% and no reported physician diagnosis. Respondents had fasted for at least 8 hours and less than 24 hours at the time of the blood draw. Fasting is not necessary to measure hemoglobin A1c. However, to be consistent with the subsample of fasting respondents used for FPG, assessment of undiagnosed diabetes in *Health, United States* is limited to the fasting subsample. Total diabetes includes those who were classified as having either physician-diagnosed or undiagnosed diabetes. Fasting weights were used to obtain prevalence estimates, and pregnant women were excluded.

Starting with *Health, United States, 2010*, an elevated hemoglobin A1c (greater than or equal to 6.5%) was included as a component of the definition of undiagnosed diabetes, along with FPG. Previous editions of *Health, United States* did not evaluate hemoglobin A1c to classify respondents as having undiagnosed diabetes; undiagnosed diabetes was based solely on elevated FPG (greater than or equal to 126 mg/dL) among those without physician-

diagnosed diabetes. The revised definition of undiagnosed diabetes was based on recommendations from the American Diabetes Association (ADA). Hemoglobin A1c was recommended as a component in diagnosing diabetes because recent improvements in assay standardization make A1c results more reliable. In addition, research has provided evidence linking elevated A1c levels with diabetic complications, thus allowing for a threshold to be set above which patients would be diagnosed as having diabetes. Although the ADA recommends using hemoglobin A1c greater than or equal to 6.5% as an indicator of undiagnosed diabetes, it cautions that A1c may be misleading in individuals with certain blood disorders (including sickle cell trait), which may have specific ethnic or geographic distributions. Therefore, clinicians may use other criteria and tests to diagnose a specific patient. For more information, see: Diagnosis and classification of diabetes mellitus. *Diabetes Care* 2015;38(suppl 1):S8–S16; Standards of medical care in diabetes—2010. *Diabetes Care* 2010;33(suppl 1):S11–S61; and International Expert Committee Report on the role of the A1c assay in the diagnosis of diabetes. *Diabetes Care* 2009;32(7):1327–34. To ensure data comparability over time, the revised definition of undiagnosed diabetes was applied to all estimates shown in *Health, United States*. As expected, this revised definition increased the percentage of respondents classified as having undiagnosed diabetes.

Periodically, NHANES laboratory testing is performed at different laboratories and using different instruments than testing in earlier years. In those instances, NHANES conducts crossover studies to evaluate the impact of these changes on laboratory measurements, and thus their impact on the evaluation of data over time. Crossover studies have been conducted to evaluate the impact of laboratory changes on both FPG and A1c. The recommended adjustments to FPG to account for laboratory changes from 2005–2006 to present have been incorporated in estimates presented in *Health, United States* so that these estimates are compatible with those from earlier years. NHANES does not recommend any adjustments to the A1c data.

Estimates presented in *Health, United States* may differ from other estimates based on the same data and presented elsewhere if different weights, age-adjustment groups, definitions, or trend adjustments are used.

For more information, see: https://www.cdc.gov/Nchs/Nhanes/2013-2014/GHB_H.htm and https://www.cdc.gov/Nchs/Nhanes/2013-2014/GLU_H.htm.

Diagnosis—Diagnosis is the act or process of identifying or determining the nature and cause of a disease or injury through evaluation of patient history, examination, and review of laboratory data. Diagnoses in the National Ambulatory Medical Care Survey and the National Hospital Ambulatory Medical Care Survey are abstracted from medical records and are currently coded to the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Starting with 2016 data, diagnosis data will be

classified using *International Classification of Diseases, 10th Revision, Clinical Modification/Procedure Coding System (ICD-10-CM/PCS)*.

For a given medical care encounter, the first-listed diagnosis can be used to categorize the visit, or if more than one diagnosis is recorded on the medical record, the visit can be categorized based on all diagnoses recorded. Analyzing first-listed diagnoses avoids double-counting events such as visits or hospitalizations; the first-listed diagnosis is often, but not always, considered the most important or dominant condition among all comorbid conditions. However, the choice of the first-listed diagnosis by the medical facility may be influenced by reimbursement or other factors. (Also see Appendix II, External cause of injury; Injury; Injury-related visit.)

Diagnostic and other nonsurgical procedure—See Appendix II, Procedure.

Discharge—The National Health Interview Survey defines a hospital discharge as the completion of any continuous period of stay of one night or more in a hospital as an inpatient. According to the Healthcare Cost and Utilization Project—National (Nationwide) Inpatient Sample, a discharge is a completed inpatient hospitalization. A hospitalization may be completed by death or by release of the patient to the customary place of residence, a nursing home, another hospital, or other locations. (Also see Appendix II, Admission; Average length of stay; Days of care; Hospital utilization; Inpatient.)

Domiciliary care home—See Appendix II, Long-term care facility; Nursing home.

Drug—Drugs are pharmaceutical agents, by any route of administration, for the prevention, diagnosis, or treatment of medical conditions or diseases. Data on specific drug use are collected in several NCHS surveys. (Also see Appendix II, Multum Lexicon Plus therapeutic class.)

National Health and Nutrition Examination Survey (NHANES)—Drug information from NHANES III and from NHANES for 1999 and subsequent years was collected during in-person interviews conducted in participants' homes. Starting with 2001 data, participants were asked whether they had taken a medication in the past 30 days for which they needed a prescription. For 1999–2000 and 1988–1994 data, the question wording differed slightly; participants were asked whether they had taken a prescription medication in the past month. For all survey years, those who answered “yes” were asked to provide the prescription medication containers for the interviewer. For each medication reported, the interviewer entered the product's complete name from the container. If no container was available, the interviewer asked the participant to verbally report the name of the medication. In addition, participants were asked how long they had been taking the medication and the main reason for use.

All reported medication names were converted to their standard generic ingredient name. For multi-ingredient products, the ingredients were listed in alphabetical order and counted as one drug (e.g., Tylenol #3 was listed as acetaminophen; codeine). No trade or proprietary names were provided on the data file.

Drug data from NHANES provide a snapshot of all prescribed drugs reported by a sample of the civilian noninstitutionalized population for a 30-day period (or past month, for earlier survey years). Drugs taken on an irregular basis, such as every other day, once per week, or for a 10-day period, were captured in the 30-day recall period. Data shown in *Health, United States* for the percentage of the population reporting multiple prescription drugs during the past 30 days include a range of drug utilization patterns; for example, persons who took three or more drugs daily during the past 30 days or persons who took a different drug three separate times would be classified as taking three or more drugs in the past 30 days, as long as at least three different drugs were taken at some time during the past 30 days.

For more information on prescription drug data collection and coding in NHANES, see: http://www.cdc.gov/nchs/nhanes/nhanes1999-2000/RXQ_DRUG.htm.

For more information on NHANES III prescription drug data collection and coding, see: ftp://ftp.cdc.gov/pub/Health_Statistics/NCHS/nhanes/nhanes3/2A/pupremed.pdf. (Also see Appendix I, National Health and Nutrition Examination Survey [NHANES].)

Drug abuse—See Appendix II, Illicit drug use.

Education—Several approaches to defining educational categories are used in *Health, United States*.

National Health Interview Survey (NHIS)—Starting in 1997, the NHIS questionnaire was changed to ask, “What is the highest level of school [person] has completed or the highest degree received?” Responses were used to categorize adults according to educational credentials (i.e., no high school diploma or general educational development high school equivalency diploma [GED]; high school diploma or GED; some college, no bachelor's degree; bachelor's degree or higher).

Prior to 1997, the education variable in NHIS was measured by asking, “What is the highest grade or year of regular school [person] has ever attended?” and “Did [person] finish the grade/year?” Responses were used to categorize adults according to years of education completed (i.e., less than 12, 12, 13–15, or 16 years or more).

Data from the 1996 and 1997 NHIS were used to compare distributions of educational attainment for adults aged 25 and over, using categories based on educational credentials (1997) and categories based on

years of education completed (1996). A larger percentage of persons reported some college than 13–15 years of education, and a correspondingly smaller percentage reported a high school diploma or GED than 12 years of education. In 1997, 19% of adults reported no high school diploma, 31% a high school diploma or GED, 26% some college, and 24% a bachelor's degree or higher. In 1996, 18% of adults reported less than 12 years of education, 37% reported 12 years, 20% reported 13–15 years, and 25% reported 16 or more years of education.

National Health and Nutrition Examination Survey (NHANES)—In 1988–1994 (NHANES III) the questionnaire asked, “What is the highest grade or year of regular school [person] has completed?” Responses were used to categorize adults according to educational credentials (i.e., no high school diploma or GED; high school diploma or GED; some college, no bachelor's degree; bachelor's degree or higher). Starting with 1999–2000 data, the questionnaire was changed to ask, “What is the highest grade or level of school [you have/(person) has] completed or the highest degree [you have/(person) has] received?” For data on children, education is based on the level of education completed by the head of the household. The question asked is, “What is the highest grade or level of school [you have/(person) has] completed or the highest degree [you have/(person) has] received?”

Emergency department—According to the National Hospital Ambulatory Medical Care Survey, an emergency department is a hospital facility that is staffed 24 hours a day and provides unscheduled outpatient services to patients whose condition requires immediate care. Emergency services provided under the “hospital as landlord” arrangement were also eligible. An emergency department was in scope if it was staffed 24 hours a day. If an in-scope emergency department had an emergency service area that was open less than 24 hours a day, then that area was included under the emergency department. If a hospital had an emergency department that was staffed less than 24 hours a day, that department was considered an outpatient clinic. (Also see Appendix II, Emergency department or emergency room visit; Outpatient department.)

Emergency department or emergency room visit—Starting with the 1997 National Health Interview Survey, respondents to the Sample Adult questionnaire and the Sample Child questionnaire (generally a parent) were asked about the number of visits to hospital emergency rooms during the past 12 months, including visits that resulted in hospitalization. In the National Hospital Ambulatory Medical Care Survey, an emergency department visit is a direct personal exchange between a patient and a physician or other health care provider working under the physician's supervision, for the purpose of seeking care and receiving personal health services. (Also see Appendix II, Emergency department; Injury-related visit.)

Employer costs for employee compensation—Employer costs for employee compensation is a measure of the average cost, per employee hour worked, to employers for wages, salaries, and benefits. Wages and salaries are defined as the hourly straight-time wage rate or, for workers not paid on an hourly basis, straight-time earnings divided by the corresponding hours. Straight-time wage and salary rates are total earnings before payroll deductions, excluding premium pay for work in addition to the regular work schedule (e.g., overtime, weekends, and holidays), shift differentials, and nonproduction bonuses such as discretionary holiday bonuses and lump-sum payments provided in lieu of wage increases. Production bonuses, incentive earnings, commission payments, and cost-of-living adjustments are included in straight-time wage and salary rates. Benefits covered are paid leave (paid vacations, holidays, sick leave, and other leave), supplemental pay (premium pay for overtime, weekends, or holidays), shift differentials, nonproduction bonuses, insurance benefits (life, health, and short- and long-term disability), retirement and savings benefits (pension and other retirement plans and savings and thrift plans), and legally required benefits (Social Security, Medicare, federal and state unemployment insurance, and workers' compensation). (Also see Appendix I, National Compensation Survey [NCS].)

Ethnicity—See Appendix II, Hispanic origin.

Exercise—See Appendix II, Physical activity, leisure-time.

Expenditures—See Appendix II, Health expenditures, national. (Also see Appendix I, National Health Expenditure Accounts [NHEA].)

External cause of injury—The external cause of injury is used for classifying the circumstances in which injuries occur. The *International Classification of Diseases, 9th Revision (ICD-9)*, External Cause of Injury Matrix, is a two-dimensional array describing both the mechanism or external cause of the injury (e.g., fall, motor-vehicle traffic) and the manner or intent of the injury (e.g., unintentional, self-inflicted, or assault). Although this matrix was originally developed for mortality, it has been adapted for use with the ICD-9 Clinical Modification (ICD-9-CM) and will be used in *Health, United States* until 2016 data are available. Data for 2016 and beyond will be classified using the *International Classification of Diseases, 10th Revision, Clinical Modification/Procedure Coding System (ICD-10-CM/PCS)*. For more information, see the NCHS website at: http://www.cdc.gov/nchs/injury/injury_tools.htm; and see: Bergen G, Chen LH, Warner M, Fingerhut LA. Injury in the United States: 2007 chartbook. Hyattsville, MD: NCHS; 2008. Available from: <http://www.cdc.gov/nchs/data/misc/injury2007.pdf>.

Family income—For the National Health Interview Survey and the National Health and Nutrition Examination Survey, all people within a household who are related to each other by blood, marriage or cohabitation, or adoption constitute a family. Each member of a family is classified according to the

total income of the family. Unrelated individuals are classified according to their own income.

National Health Interview Survey (NHIS)—Prior to 1997, family income was the total income received by members of a family (or by an unrelated individual) in the 12 months before interview. Family income included wages, salaries, rents from property, interest, dividends, profits and fees from their own businesses, pensions, and help from relatives. Starting in 1997, NHIS collected family income data for the calendar year prior to interview (e.g., 2014 family income data were based on calendar year 2013 information). The 1997–2006 instrument allowed the respondent to supply a specific dollar amount (up to \$999,995). Any family income responses greater than \$999,995 were entered as \$999,996. Respondents who did not know or refused to give a dollar amount in response to this question were asked if their total combined family income for the previous year was \$20,000 or more, or less than \$20,000. If the respondent answered this question, he or she was then given one of two flash cards and asked to indicate which income group listed on the card best represented the family's combined income during the previous calendar year. One flash card listed incomes that were \$20,000 or more, and the other flash card listed incomes that were less than \$20,000. Starting with the 2007 NHIS, the income amount follow-up questions that had been in place since 1997 were replaced with a series of unfolding bracket questions. The unfolding bracket method asked a series of closed-ended income range questions (e.g., “Is it less than \$50,000?”) if the respondent did not provide an answer to the exact income amount question. The closed-ended income range questions were constructed so that each successive question establishes a smaller range for the amount of the family's income. In 2011, several new unfolding-bracket income questions were added to NHIS to improve the assignment of poverty status. Additional questions focused on assessing whether a family's income was less than 200% of the poverty threshold or 200% or more of the poverty threshold. The question wording varied according to family size. In addition, a question was added for respondents who answered that their family's income was \$100,000 or more as to whether their family's income was less than \$150,000, or \$150,000 or more. For more information on this series of family income questions, see: 2014 NHIS public-use data release. NCHS. 2015. Available from: <http://www.cdc.gov/nchs/nhis/2014imputedincome.htm>.

Also see: Pleis JR, Cohen RA. Impact of income bracketing on poverty measures used in the National Health Interview Survey's Early Release Program: Preliminary data from the 2007 NHIS. Hyattsville, MD: NCHS. 2007. Available from: <http://www.cdc.gov/nchs/data/nhis/income.pdf>.

For NHIS respondents, family income data are used in the computation of a poverty measure. Starting with *Health, United States, 2004*, a new methodology for imputing family income data for NHIS was implemented for data years 1997 and beyond. Multiple imputations were performed for survey years 1997 and beyond, with five sets of imputed values created to allow for the assessment of variability caused by imputation. A detailed description of the multiple imputation procedure, and data files for 1997 and beyond, are available from: http://www.cdc.gov/nchs/nhis/quest_data_related_1997_forward.htm, through the Data Release or the Imputed Income Files link under that year. For data years 1990–1996, about 16%–18% of persons had missing data for family income. In those years, missing values were imputed for family income by using a sequential hot deck within matrix cells imputation approach. A detailed description of the imputation procedure and data files, with imputed annual family income for 1990–1996, is available from: ftp://ftp.cdc.gov/pub/Health_Statistics/NCHS/Datasets/NHIS/1990-96_Family_Income/. (Also see Appendix II, Poverty; Table VI.)

National Health and Nutrition Examination Survey (NHANES)—In NHANES 1999 and onward, family income is asked in a series of questions about possible sources of income, including wages, salaries, interest and dividends, federal programs, child support, rents, royalties, and other possible sources. After the information about sources of income was obtained in the family interview income section of the questionnaire, the respondent was asked to report total combined family income for him- or herself and the other members of their family, in dollars. If the respondent did not provide an answer or did not know the total combined family income, he or she was asked if the total family income was less than \$20,000 or \$20,000 or more. If the respondent answered, a follow-up question asked the respondent to select an income range from a list on a printed flash card. The midpoint of the income range was then used as the total family income value. Family income values are used to calculate a poverty measure. NHANES II (1976–1980) included questions on components of income; NHANES III (1988–1994) did not ask the detailed components-of-income questions but asked respondents to identify their income based on a set of ranges provided on a flash card. Family income was not imputed for individuals or families with no reported income information in any of the NHANES survey years. (Also see Appendix II, Poverty.)

National Immunization Survey (NIS)—Prior to 1998, family income was the total income received by all family members in the past 12 months at the time of interview. Following the changes in the NHIS income questions, NIS changed the reference period for 1998 onward and collected income received by all family members for the calendar year prior to the interview year for households with age-eligible children (e.g., 2014 NIS

family income data are based on calendar year 2013 income). Family income is the combined total income received by all members of a family before taxes. For the family income questions, the household respondent is asked to include income received from jobs, Social Security, retirement income, unemployment payments, public assistance, interest, dividends, net income from business, farm, rent, or any other sources. Respondents who answered “don't know” or refused to give a dollar amount for the total family income were asked a cascading sequence of income questions (a total of 15 cascading questions that attempt to place the family income into one of 15 income intervals ranging from less than or equal to \$7,500 to greater than or equal to \$75,000). The initial question asks if the family income for the prior year was more or less than \$20,000. Subsequent sets of income range questions are asked so that each successive question establishes a smaller income range. The midpoint of the income range is used as the total family income value for respondents who answered “don't know” or refused to give a dollar amount. A family income variable is constructed from the total family income question and the cascading income questions. If an exact income is given, family income is set to this amount; otherwise it is set to the midpoint of the tightest bounds established by the cascading income questions. The values of total family income are used to calculate a poverty measure. For NIS, this ratio is calculated only for households with age-eligible children, using the actual family income value or the midpoint of the interval from the series of cascading questions in the numerator and the poverty threshold provided by the Census Bureau for the size of the family and the number of related children in the household in the denominator. Details of the income questions and computation of the income-to-poverty ratio for each data collection year can be found in the NIS data documentation (Data User's Guide and Household Interview Questionnaire) provided on the NIS website at: <http://www.cdc.gov/vaccines/imz-managers/nis/data-tables.html>.

For more information, see: Battaglia MP, Hoaglin DC, Izrael D, Khare M, Mokdad A. Improving income imputation by using partial income information and ecological variables. Presented at the American Statistical Association–Joint Statistical Meeting; 2002 Aug 11–15, New York, NY. Available from: http://www.cdc.gov/nchs/data/nis/estimation_weighting/Battaglia2002.pdf.

Federal hospital—See Appendix II, Hospital.

Fee-for-service health insurance—Fee-for-service health insurance is private (commercial) health insurance that reimburses health care providers on the basis of a fee for each health service provided to the insured person. It is also known as indemnity health insurance. In addition, “fee-for-service” is a term often applied to original Medicare, to distinguish it from Medicare managed-care plans and other

new payment systems. (Also see Appendix II, Health insurance coverage; Managed care; Medicare.)

Fertility rate—See Appendix II, Rate: Birth and related rates.

General hospital—See Appendix II, Hospital.

Geographic region—The U.S. Census Bureau groups the 50 states and D.C., for statistical purposes, into four geographic regions (Northeast, Midwest, South, and West) and nine divisions based on geographic proximity. (See Figure I.)

Gestation—For the National Vital Statistics System and CDC's Abortion Surveillance System, the period of gestation is defined as beginning with the first day of the last normal menstrual period and ending with the day of birth or day of termination of pregnancy. Data on gestational age are subject to error for several reasons, including imperfect maternal recall or misidentification of the last menstrual period because of postconception bleeding, delayed ovulation, or intervening early miscarriage.

Gross domestic product (GDP)—The GDP is the market value of the goods and services produced by labor and property located in the United States. As long as the labor and property are located in the United States, the suppliers (i.e., the workers and, for property, the owners) may be U.S. residents or residents of other countries. (Also see Appendix II, Consumer Price Index [CPI]; Health expenditures, national.)

Health care contact—Starting in 1997, the National Health Interview Survey has collected information on health care contacts with doctors and other health care professionals by using the following series of questions: “During the past 12 months, how many times have you gone to a hospital emergency room about your own health?”, “During the past 12 months, did you receive care at home from a nurse or other health care professional? What was the total number of home visits received?”, and “During the past 12 months, how many times have you seen a doctor or other health care professional about your own health at a doctor's office, a clinic, or some other place? Do not include times you were hospitalized overnight, visits to hospital emergency rooms, home visits, or telephone calls.” Starting with 2000 data, this question was amended to specifically exclude dental visits.

For 1997–1999, for each question, respondents were shown a flash card with response categories of 0, 1, 2–3, 4–9, 10–12, or 13 or more visits. For tabulation of the 1997–1999 data, responses of 2–3 were recoded to 2, responses of 4–9 were recoded to 6, responses of 10–12 were recoded to 11, and 13 or more visits were recoded to 13. The recoded values for the three types of visits were then added to yield an estimate of total health care contacts. Starting with 2000 data, response categories were expanded to 0, 1, 2–3, 4–5, 6–7, 8–9, 10–12, 13–15, or 16 or more. For 2000 and more recent data, these response categories were recoded to the midpoint of the range. The category of 16 or more was recoded to 16. The recoded values for the three types of visits were then added

Table VI. Imputed family income percentages in the National Health Interview Survey, by selected characteristics: United States, 1990–2014

Year	All ages	Under 18 years	18 years and over	18–64 years	Under 65 years	1–64 years	65 years and over	Females 18 years and over	Females 40 years and over	2 years and over	45 years and over
	Percent										
1990	16	14	18	16	15	15	24	18	21	17	22
1991	18	15	19	17	17	17	26	19	23	18	23
1992	18	16	19	18	17	17	27	20	23	18	23
1993	16	14	17	16	15	15	23	17	19	16	20
1994	17	15	18	17	16	16	25	18	21	17	21
1995	16	14	16	15	15	15	22	17	19	16	19
1996	17	14	17	16	16	16	24	18	20	17	20
1997	24	21	26	24	23	23	34	26	30	25	30
1998	29	25	30	28	27	27	39	30	34	29	34
1999	31	27	32	30	29	29	43	33	37	31	37
2000	32	28	33	31	30	31	45	34	38	32	38
2001	32	27	33	31	30	30	44	34	37	32	38
2002	32	28	33	31	30	30	44	33	37	32	37
2003	33	30	35	33	32	32	44	35	38	34	38
2004	33	29	34	32	31	31	41	34	36	33	37
2005	33	29	34	32	31	31	44	35	37	33	38
2006	34	31	35	33	33	33	45	36	39	34	39
2007	33	29	34	32	31	31	43	35	38	33	37
2008	30	27	31	29	29	29	40	32	34	30	34
2009	25	21	26	24	23	23	34	26	29	25	29
2010	25	20	26	24	23	23	36	27	30	25	30
2011	22	19	23	22	21	21	31	24	26	23	26
2012	23	19	24	22	21	21	32	24	27	23	27
2013	23	19	24	23	22	22	31	25	27	23	27
2014	23	20	24	23	22	22	31	25	27	23	27

NOTES: Percentages are weighted. See Appendix II, Family income.

SOURCE: CDC/NCHS, National Health Interview Survey. See Appendix I, National Health Interview Survey (NHIS).

to yield an estimate of the summary measure of health care contacts (including doctor's visits, hospital emergency room visits, and home visits). After summing the three component visit variables, respondents with values on the edge of the categories presented in *Health, United States* were rounded down to provide a more conservative estimate of the number of visits. For example, a respondent with 3.5 health care contacts was included in the 1–3 visits category, and a respondent with 9.5 health care contacts was included in the 4–9 visits category. Respondents were included in this analysis only if they were known on all three visit variables.


Analyses of the percentage of children without a health care visit are based on the following question: "During the past 12 months, how many times has [person] seen a doctor or other health care professional about [his/her] health at a doctor's office, a clinic, or some other place? Do not include times [person] was hospitalized overnight, visits to hospital emergency rooms, home visits, or telephone calls." Starting with 2000 data, this question was amended to specifically exclude dental visits. (Also see Appendix II, Emergency department or emergency room visit; Home visit.)

Health expenditures, national—National health expenditures are estimated by the Centers for Medicare & Medicaid Services (CMS) and measure calendar year spending for health care in the United States by type of service delivered (e.g., hospital care, physician services, nursing home care) and source of funding for those services (e.g., private health insurance, Medicare, Medicaid, out-of-pocket spending). CMS produces both historical and projected estimates of health expenditures by category. (Also see Appendix I, National Health Expenditure Accounts [NHEA]; Appendix II, Gross domestic product [GDP].) Types of national health expenditures include:

Health consumption expenditures are outlays for goods and services relating directly to patient care, plus expenses for administering health insurance programs, the net cost of health insurance, and public health activities. This category is equivalent to total national health expenditures minus expenditures for investment in noncommercial research and structures and equipment.

Personal health care expenditures are outlays for goods and services relating directly to patient care. These

Figure I. U.S. Census Bureau: Four geographic regions and nine divisions of the United States



expenditures are total national health expenditures minus expenditures for investment, health insurance program administration and the net cost of insurance, and public health activities.

Business, household, and other private expenditures are outlays for services paid for by nongovernmental sources, such as consumers, private industry, and philanthropic and other non-patient-care sources.

Government expenditures are outlays for services paid for by federal, state, and local government agencies or expenditures required by governmental mandate (such as workers' compensation insurance payments).

Health insurance coverage—Health insurance is broadly defined to include both public and private payers who cover medical expenditures incurred by a defined population in a variety of settings. Estimates of health insurance are available from several different government surveys. Because of differences in methodology, question wording, and recall period, estimates from different sources may vary and are not directly comparable. For more information, see: Health insurance measurement: Differences by data source. Available from: http://www.census.gov/content/dam/Census/library/infographics/health_insurance_measurement.pdf.

American Community Survey (ACS)—For point-in-time health insurance estimates, ACS respondents were asked about their coverage at the time of interview. Respondents were asked: “Is this person CURRENTLY covered by any of the following types of health insurance or health coverage plans? Mark yes or no for each type of coverage: Insurance through a current or former employer or union [of this person or another family member]; Insurance purchased directly from an insurance company [by this person or another family member]; Medicare, for people 65 and older, or people with certain disabilities; Medicaid, Medical Assistance, or any kind of government-assistance plan for those with low incomes or a disability; TRICARE or other military health care; VA (including those who have ever used or enrolled for VA health care); Indian Health Service; Any other type of health insurance or health coverage plan [specify plan].” In ACS, persons were considered uninsured if they were not covered by private health insurance, Medicare, Medicaid, Medical Assistance, TRICARE or other military health care, veteran’s coverage through the Veteran’s Administration, or other government coverage. People with Indian Health Service coverage only were considered uninsured in ACS.

National Health Interview Survey (NHIS)—For point-in-time health insurance estimates, NHIS respondents were asked about their coverage at the time of interview. For 1993–1996, respondents were asked about their coverage in the previous month. Questions on health insurance coverage were expanded starting in 1993, compared with previous years. In 1997, the entire questionnaire was redesigned and data were collected using a computer-assisted personal interview (CAPI). In 2007, questions on health insurance coverage were expanded again to include three new questions on high-deductible health plans, health savings accounts, and flexible spending accounts.

Respondents were considered to be covered by private health insurance if they indicated private health insurance or, prior to 1997, if they were covered by a single-service hospital plan. Private health insurance includes managed care such as health maintenance organizations (HMOs).

Private insurance obtained through the workplace was defined as any private insurance that was originally obtained through a present or former employer or union or, starting in 1997, through the workplace, self-employment, or a professional association. Starting in 2011, respondents were also asked whether health insurance coverage was obtained through parents or another relative. Coverage obtained through parents or another relative was not included as workplace coverage.

Until 1996, persons were defined as having Medicaid or other public assistance coverage if they indicated that they had either Medicaid or other public assistance or if they reported receiving Aid to Families with Dependent Children (AFDC) or Supplemental Security Income (SSI). After welfare reform in late 1996, Medicaid was delinked from AFDC and SSI. Starting in 1997, persons were considered to be covered by Medicaid if they reported Medicaid or a state-sponsored health program. Starting in 1999, persons were considered covered by Medicaid if they reported coverage by the Children's Health Insurance Program (CHIP). Medicare or military health plan coverage was also determined in the interview and, starting in 1997, other government-sponsored program coverage was determined as well.

If respondents did not report coverage under one of the above types of plans and they had unknown coverage under either private health insurance or Medicaid, they were considered to have unknown coverage.

The remaining respondents without any indicated coverage were considered uninsured. The uninsured were persons who did not have coverage under private health insurance, Medicare, Medicaid, public assistance, a state-sponsored health plan, other government-sponsored programs, or a military health plan. Persons with only Indian Health Service (IHS) coverage were considered uninsured. Although NHIS respondents who

report IHS coverage as their only source of coverage are currently recoded to being uninsured, IHS provides a comprehensive health service delivery system for approximately 2.2 million American Indian or Alaska Native persons. See: <http://www.ihs.gov/newsroom/factsheets/ihsyear2015profile/>. Estimates of the percentage of persons who were uninsured based on NHIS may differ slightly from those based on other sources because of differences in survey questions, recall period, and other aspects of survey methodology.

In NHIS, on average less than 2% of people aged 65 and over reported no current health insurance coverage, but the small sample size precludes the presentation of separate estimates for this population. Therefore, the term “uninsured” refers only to the population under age 65.

Two additional questions were added to the health insurance section of NHIS beginning with the third quarter of 2004 (Table VII). One question was asked of persons aged 65 and over who had not indicated that they had Medicare: “People covered by Medicare have a card which looks like this. [Are/Is] [person] covered by Medicare?” The other question was asked of persons under age 65 who had not indicated any type of coverage: “There is a program called Medicaid that pays for health care for persons in need. In this state it is also called [state name]. [Are/Is] [person] covered by Medicaid?” Respondents who originally classified themselves as uninsured, but whose classification was changed to Medicare or Medicaid on the basis of a “yes” response to either question, subsequently received appropriate follow-up questions concerning periods of noncoverage for insured respondents. Of the 892 people (unweighted) who were eligible to receive the Medicare probe question in the third and fourth quarters of 2004, 55% indicated that they were covered by Medicare. Of the 9,146 people (unweighted) who were eligible to receive the Medicaid probe question in the third and fourth quarters of 2004, 3% indicated that they were covered by Medicaid. Estimates in *Health, United States* were calculated using the responses to the two additional probe questions. For a complete discussion of the effect of the addition of these two probe questions on the estimates for insurance coverage, see: Cohen RA, Martinez ME. Impact of Medicare and Medicaid probe questions on health insurance estimates from the National Health Interview Survey, 2004. Health E-Stats. NCHS; 2005. Available from: <http://www.cdc.gov/nchs/data/hestat/impact04/impact04.htm>.

Survey respondents may be covered by health insurance at the time of interview but may have experienced one or more lapses in coverage during the 12 months prior to interview. Starting with *Health, United States, 2006*, NHIS estimates have been presented for the following three exhaustive categories: (a) people with health insurance continuously for the full 12 months prior to interview,

(b) those who had a period of up to 12 months prior to interview without coverage, and (c) those who were uninsured for more than 12 months prior to interview. This stub variable has been added to selected tables. Two additional NHIS questions were used to determine the appropriate category for the survey respondents: (a) all persons without a known comprehensive health insurance plan were asked, "About how long has it been since [person] last had health care coverage?"; and (b) all persons with known health insurance coverage were asked, "In the past 12 months, was there any time when [person] did NOT have ANY health insurance coverage?"

(Also see Appendix II, Children's Health Insurance Program [CHIP]; Fee-for-service health insurance; Health maintenance organization [HMO]; Managed care; Medicaid; Medicare; Uninsured.)

Health maintenance organization (HMO)—An HMO is a health care system that assumes or shares both the financial risks and the delivery risks associated with providing comprehensive medical services to a voluntarily enrolled population in a particular geographic area, usually in return for a fixed, prepaid fee. Pure HMO enrollees use only the prepaid, capitated health services of the HMO panel of medical care providers. Open-ended HMO enrollees use the prepaid HMO health services but may also receive medical care from providers who are not part of the HMO panel. There is usually a substantial deductible, copayment, or coinsurance associated with use of nonpanel providers. HMO model types are as follows:

Group model HMO is an HMO that contracts with a single multispecialty medical group to provide care to the HMO's membership. The group practice may work exclusively with the HMO, or it may provide services to non-HMO patients as well. The HMO pays the medical group a negotiated per capita rate, which the group distributes among its physicians, usually on a salaried basis.

Staff model HMO is a closed-panel HMO (where patients can receive services only through a limited number of providers) in which physicians are HMO employees. The providers see members in the HMO's own facilities.

Network model HMO is an HMO that contracts with multiple physician groups to provide services to HMO members. It may include single or multispecialty groups.

Individual practice association (IPA) is a health care provider organization composed of a group of independent practicing physicians who maintain their own offices and band together for the purpose of contracting their services to HMOs, preferred provider organizations, and insurance companies. An IPA may contract with and provide services to both HMO and non-HMO plan participants.

Mixed model HMO is an HMO that combines features of more than one HMO model.

(Also see Appendix II, Managed care; Preferred provider organization [PPO].)

Health services and supplies expenditures—See Appendix II, Health expenditures, national.

Health status, respondent-assessed—Health status was measured in the National Health Interview Survey by asking the family respondent about his or her health or the health of a family member: "Would you say [person's] health in general is excellent, very good, good, fair, or poor?"

Hearing trouble—In the National Health Interview Survey, information about hearing trouble is obtained by asking respondents how well they hear without the use of hearing aids. Prior to 2007 data, respondents were asked, "Which statement best describes your hearing without a hearing aid: good, a little trouble, a lot of trouble, or deaf?" In the 2007 data, the question was revised to expand the response categories. Respondents were asked, "These next questions are about your hearing WITHOUT the use of hearing aids or other listening devices. Is your hearing excellent, good, [do you have] a little trouble hearing, moderate trouble, a lot of trouble, or are you deaf?" Starting with 2008 data, respondents were asked, "WITHOUT the use of hearing aids or other listening devices, is your hearing excellent, good, [do you have] a little trouble hearing, moderate trouble, a lot of trouble, or are you deaf?" Because of the expanded response categories, 2007 and subsequent data are not strictly comparable with earlier years and caution is urged when interpreting trends. For example, in 2006, 3.5% of adults (aged 18 and over) were classified as having hearing difficulty (response categories: a lot of trouble or deaf). In 2007, 2.3% of adults (aged 18 and over) were classified as having hearing difficulty (response categories: a lot of trouble or deaf). This more than 30% decline from 2006 to 2007 in the estimate of those with hearing trouble is likely attributable to the addition of the moderate trouble response category, rather than changes in the prevalence of hearing trouble. Although all age groups saw a decline in the percentage reporting hearing trouble between 2006 and 2007, the amount of the decline varied. There was a 50% decline in reported hearing trouble among adults aged 18–44 (from 0.8% in 2006 to 0.4% in 2007). Among adults aged 45–64, the percentage that reported hearing trouble declined 43%, from 3.5% in 2006 to 2.0% in 2007. Among adults aged 65 and over, reported hearing trouble declined 24%, from 11.4% in 2006 to 8.7% in 2007.

For more information, see: Pleis JR, Lucas JW. Summary health statistics for U.S. adults: National Health Interview Survey, 2007. NCHS. Vital Health Stat 2009;10(240). Available from: http://www.cdc.gov/nchs/data/series/sr_10/sr10_240.pdf. (Also see Appendix II, Basic actions difficulty.)

Hispanic origin—Hispanic or Latino origin includes persons of Mexican, Puerto Rican, Cuban, Central and South American, and other or unknown Latin American or Spanish origin. Persons of Hispanic origin may be of any race.

Table VII. Percentage of persons under age 65 with Medicaid or who are uninsured, by selected demographic characteristics, using Method 1 and Method 2 estimation procedures: United States, 2004

Characteristic	Medicaid ¹		Uninsured ²	
	Method 2 ³	Method 1 ³	Method 2 ³	Method 1 ³
	Percent (standard error)			
Age				
Under 65 years	12.0 (0.24)	11.8 (0.24)	16.4 (0.23)	16.6 (0.23)
Under 18 years	25.4 (0.49)	24.9 (0.49)	9.2 (0.30)	9.7 (0.29)
18–64 years	6.6 (0.17)	6.5 (0.17)	19.3 (0.26)	19.4 (0.26)
Percent of poverty level ⁴				
Below 100%	47.5 (1.03)	46.6 (1.03)	29.6 (0.89)	30.5 (0.92)
100%–less than 200%	22.0 (0.59)	21.5 (0.60)	28.9 (0.66)	29.4 (0.66)
200% or more	2.9 (0.13)	2.8 (0.13)	9.4 (0.23)	9.5 (0.23)
Age and percent of poverty level ⁴				
Under 18 years:				
Below 100%	71.9 (1.35)	70.2 (1.35)	14.5 (1.15)	16.2 (1.22)
100%–less than 200%	39.2 (1.13)	38.4 (1.14)	15.0 (0.81)	15.8 (0.82)
200% or more	6.2 (0.33)	6.1 (0.33)	4.9 (0.30)	4.9 (0.30)
18–64 years:				
Below 100%	31.2 (1.02)	30.8 (1.02)	39.7 (1.09)	40.1 (1.09)
100%–less than 200%	12.0 (0.48)	11.8 (0.48)	37.0 (0.72)	37.2 (0.72)
200% or more	1.7 (0.11)	1.7 (0.10)	11.0 (0.26)	11.1 (0.26)
Hispanic origin and race ⁵				
Hispanic or Latino	22.2 (0.55)	21.5 (0.55)	34.4 (0.64)	35.1 (0.65)
Mexican	22.0 (0.63)	21.5 (0.63)	37.6 (0.82)	38.1 (0.83)
Not Hispanic or Latino	10.2 (0.25)	10.1 (0.25)	13.2 (0.23)	13.3 (0.23)
White only	7.4 (0.26)	7.4 (0.26)	12.0 (0.25)	12.1 (0.25)
Black or African American only	23.9 (0.80)	23.5 (0.79)	17.3 (0.58)	17.8 (0.58)

¹Includes persons who do not have private coverage but who have Medicaid or other state-sponsored health plans, including the Children's Health Insurance Program (CHIP).

²Includes persons who have not indicated that they are covered at the time of interview under private health insurance, Medicare, Medicaid, CHIP, a state-sponsored health plan, other government programs, or military health plan (includes VA, TRICARE, and CHAMP–VA). This category includes persons who are only covered by Indian Health Service or only have a plan that pays for one type of service, such as accidents or dental care.

³Starting with the third quarter of 2004, two additional questions were added to the National Health Interview Survey (NHIS) insurance section to reduce potential errors in reporting of Medicare and Medicaid status. Persons aged 65 and over not reporting Medicare coverage were asked explicitly about Medicare coverage, and persons under age 65 with no reported coverage were asked explicitly about Medicaid coverage. Estimates calculated without using the additional information from these questions are noted as Method 1. Estimates calculated using the additional information from these questions are noted as Method 2.

⁴Based on family income and family size and composition, using the U.S. Census Bureau's poverty thresholds. The percentage of respondents with unknown poverty level was 28.2% in 2004. See the *NHIS Survey Description* for 2004. Available from: <http://www.cdc.gov/nchs/data/nhis/srvydesc.pdf>.

⁵Persons of Hispanic origin may be of any race or combination of races. Similarly, the category Not Hispanic or Latino refers to all persons who are not of Hispanic or Latino origin, regardless of race.

SOURCE: CDC/NCHS, National Health Interview Survey, 2004, Family Core Component. Data are based on household interviews of a sample of the civilian noninstitutionalized population. Available from: <http://www.cdc.gov/nchs/data/hestat/impact04/impact04.htm>. See Appendix I, National Health Interview Survey (NHIS).

Birth file—The reporting area for an Hispanic-origin item on the birth certificate expanded between 1980 and 1993 (when the Hispanic item was included on the birth certificate in all states and D.C.). Trend data on births of Hispanic and non-Hispanic parentage in *Health, United States* are affected by expansion of the reporting areas, which affects numbers of events, composition of the Hispanic population, and maternal and infant health characteristics.

In 1980 and 1981, information on births of Hispanic parentage was reported on the birth certificate by the following 22 states: Arizona, Arkansas, California, Colorado, Florida, Georgia, Hawaii, Illinois, Indiana, Kansas, Maine, Mississippi, Nebraska, Nevada, New Jersey, New Mexico, New York, North Dakota, Ohio, Texas, Utah, and Wyoming. In 1982 Tennessee, and in 1983 D.C., began reporting this information. Between 1983 and 1987, information on births of Hispanic parentage was available for 23 states and D.C. In 1988, this information became available for Alabama, Connecticut, Kentucky, Massachusetts, Montana, North Carolina, and Washington state, increasing the number of states reporting information on births of Hispanic parentage to 30 states and D.C. In 1989, this information became available from an additional 17 states, increasing the number of Hispanic-reporting states to 47 and D.C. In 1989, only Louisiana, New Hampshire, and Oklahoma did not report Hispanic parentage on the birth certificate. With the inclusion of Louisiana in 1989 and Oklahoma in 1990 as Hispanic-reporting states, 99% of birth records included information on mother's origin. Hispanic origin of the mother was reported on the birth certificates of 49 states and D.C. in 1991 and 1992; only New Hampshire did not provide this information. Starting in 1993, Hispanic origin of mother was reported by all 50 states and D.C.

Starting with 2003 data, some states began using the 2003 revision of the U.S. Standard Certificate of Live Birth. Hispanic origin and race are collected separately on the birth certificate. The Hispanic origin question on the 2003 revision of the birth certificate asks respondents to select only one response. Occasionally, more than one Hispanic origin response is given; that is, a specified Hispanic origin group (Mexican, Puerto Rican, Cuban, or Central and South American) in combination with one or more other specified Hispanic origin groups. From 2003 through 2012, respondents who selected more than one Hispanic origin on the birth certificate were classified as other Hispanic. In 2012, 0.4% of births in the revised-state reporting area, plus Massachusetts (unrevised state that also reported more than one Hispanic origin response), were to women reporting more than one Hispanic origin. Beginning with 2013 data, respondents who select more than one Hispanic origin are randomly assigned to a single Hispanic origin. The Hispanic origin question on the 1989 revision of the birth certificate also offers the opportunity to report more than one origin;

however, NCHS processing guidelines for unrevised data allow for coding only the first Hispanic origin listed.

Linked birth/Infant death file—The linked birth/infant death file is particularly useful for computing accurate infant mortality rates by race and Hispanic origin because the race and Hispanic origin of the mother from the birth certificate are used in both the numerator and denominator of the linked birth/infant death infant mortality rate. In contrast, infant mortality rates based on the vital statistics mortality file use for the numerator race and Hispanic origin as reported on the death certificate and for the denominator the race and Hispanic origin of the mother as reported on the birth certificate. Race and Hispanic origin information from the birth certificate, which is reported by the mother, is considered more reliable than race and Hispanic origin information from the death certificate, which is reported by the funeral director based on information provided by an informant or by observation. See Appendix II, Hispanic origin; sections for Birth file, Mortality file.

Mortality file—The reporting area for an Hispanic-origin item on the death certificate expanded between 1985 and 1997. In 1985, mortality data by Hispanic origin of decedent were based on deaths of residents of the following 17 states and D.C. whose data on the death certificate were at least 90% complete on a place-of-occurrence basis and of comparable format: Arizona, Arkansas, California, Colorado, Georgia, Hawaii, Illinois, Indiana, Kansas, Mississippi, Nebraska, New York, North Dakota, Ohio, Texas, Utah, and Wyoming. In 1986, New Jersey began reporting Hispanic origin of decedent, increasing the number of reporting states to 18 and D.C. in 1986 and 1987. In 1988, Alabama, Kentucky, Maine, Montana, North Carolina, Oregon, Rhode Island, and Washington state were added to the reporting area, increasing the number of states to 26 and D.C. In 1989, an additional 18 states were added, increasing the Hispanic reporting area to 44 states and D.C.; only Connecticut, Louisiana, Maryland, New Hampshire, Oklahoma, and Virginia were not included in the reporting area. Starting with 1990 data in *Health, United States*, the criterion was changed to include states whose data were at least 80% complete. In 1990, Maryland, Virginia, and Connecticut; in 1991 Louisiana; and in 1993 New Hampshire were added, increasing the reporting area for Hispanic origin of decedent to 47 states and D.C. in 1990; 48 states and D.C. in 1991 and 1992; and 49 states and D.C. in 1993–1996. Only Oklahoma did not provide this information in 1993–1996. Starting in 1997, Hispanic origin of decedent was reported by all 50 states and D.C. Based on data from the U.S. Census Bureau, the 1990 reporting area encompassed 99.6% of the U.S. Hispanic population. In 1990, more than 96% of death records included information on Hispanic origin of the decedent.

Starting with 2003 data, some states began using the 2003 revision of the U.S. Standard Certificate of Death, which allows the reporting of more than one race (multiple races) and includes some revisions in the item reporting Hispanic origin. The effect of the 2003 revision of the Hispanic origin item on the reporting of Hispanic origin on death certificates is presumed to be minor. For more information, see Appendix II, Race. Also see the Technical Notes section of the annual series of “Deaths: Final Data” reports, available from: <http://www.cdc.gov/nchs/products/nvsr.htm>; and NCHS procedures for multiple-race and Hispanic origin data: Collection, coding, editing, and transmitting. Hyattsville, MD: NCHS; 2004. Available from: http://www.cdc.gov/nchs/data/dvs/Multiple_race_docu_5-10-04.pdf.

National Health Interview Survey (NHIS) and National Health and Nutrition Examination Survey (NHANES)—Questions on Hispanic origin are self-reported in NHANES III and subsequent years, and since 1976 in NHIS, and precede questions on race. For 1999–2006 data, the NHANES sample was designed to provide estimates specifically for persons of Mexican origin and not for all Hispanic-origin persons in the United States. Persons of Hispanic origin other than Mexican were entered into the sample with different selection probabilities that are not nationally representative of the total U.S. Hispanic population. Starting with 2007–2008 data collection, all Hispanic persons were oversampled, not just persons of Mexican origin. In addition to allowing estimates for the total group of Hispanic persons, the sample size for Hispanic persons of Mexican origin is sufficient to continue to produce reliable estimates for this group. However, the methodology for the oversampling of Hispanic persons did not provide sufficient sample sizes for calculating estimates for other Hispanic subgroups besides Mexican origin. For more information on the NHANES sampling methodology changes, see http://www.cdc.gov/nchs/nhanes/nhanes2007-2008/sampling_0708.htm; and the series of NHANES analytic guidelines available from: http://www.cdc.gov/nchs/nhanes/survey_methods.htm. For more information on race and Hispanic origin in NHIS, see the NHIS Race and Hispanic Origin Information home page. Available from: <http://www.cdc.gov/nchs/nhis/rhoi.htm>.

Surveillance, Epidemiology, and End Results (SEER) Program—SEER data are available from the National Institutes of Health, National Cancer Institute. SEER Hispanic data used in *Health, United States* tables exclude data from Alaska. The North American Association of Central Cancer Registries, Inc. (NAACCR) Hispanic Identification Algorithm was used on a combination of variables to classify incidence cases as Hispanic for analytic purposes. See: NAACCR guideline for enhancing Hispanic–Latino identification. Bethesda, MD: National Cancer Institute; 2003. Available from:

http://seer.cancer.gov/seerstat/variables/seer/yr1973_2004/race_ethnicity/.

Youth Risk Behavior Survey (YRBS)—Prior to 1999, a single question was asked about race and Hispanic origin, with the option of selecting one of the following categories: white not Hispanic, black not Hispanic, Hispanic or Latino, Asian or Other Pacific Islander, American Indian or Alaska Native, or other. Between 1999 and 2003, respondents were asked a single question about race and Hispanic origin with the option of choosing one or more of the following categories: white, black or African American, Hispanic or Latino, Asian, Native Hawaiian or Other Pacific Islander, or American Indian or Alaska Native. Beginning in 2005, respondents were asked a question about Hispanic origin (“Are you Hispanic or Latino?”) and a second separate question about race that included the option of selecting one or more of the following categories: American Indian or Alaska Native, Asian, black or African American, Native Hawaiian or Other Pacific Islander, or white. Because of the differences between questions, the data about race and Hispanic ethnicity for the years prior to 1999 are not strictly comparable with estimates for the subsequent years. However, analyses of data collected between 1991 and 2003 have indicated that the data are comparable across years and can be used to study trends. See Appendix II, Race; and see: Brener ND, Kann L, McManus T. A comparison of two survey questions on race and ethnicity among high school students. *Public Opin Q* 2003;67(2):227–36.

HIV—See Appendix II, Human immunodeficiency virus (HIV) disease.

Home visit—Starting in 1997, the National Health Interview Survey has been collecting information on home visits received during the 12 months prior to interview. Respondents are asked, “During the past 12 months, did you receive care at home from a nurse or other health care professional? What was the total number of home visits received?” These data are combined with data on visits to doctors' offices, clinics, and emergency departments to provide a summary measure of adult health care visits. (Also see Appendix II, Emergency department or emergency room visit; Health care contact.)

Hospital—According to the American Hospital Association (AHA), hospitals are licensed institutions with at least six beds whose primary function is to provide diagnostic and therapeutic patient services for medical conditions; they have an organized physician staff and provide continuous nursing services under the supervision of registered nurses. The World Health Organization (WHO) considers an establishment to be a hospital if it is permanently staffed by at least one physician, can offer inpatient accommodation, and can provide active medical and nursing care. Hospitals may be classified by type of service, ownership, size in terms of number of beds, and length of stay. In the National

Hospital Ambulatory Medical Care Survey, hospitals include all those with an average length of stay for all patients of less than 30 days (short-stay) or hospitals whose specialty is general (medical or surgical) or children's general. Federal hospitals and hospital units of institutions and hospitals with fewer than six beds staffed for patient use are excluded. (Also see Appendix II, Average length of stay; Bed, health facility; Days of care; Emergency department; Inpatient; Outpatient department.)

Community hospital—Community hospitals, based on the AHA definition, include all nonfederal, short-term general and special hospitals whose facilities and services are available to the public. Special hospitals include obstetrics and gynecology; eye, ear, nose, and throat; rehabilitation; orthopedic; and other specialty services. Short-term general and special children's hospitals are also considered to be community hospitals. A hospital may include a nursing-home-type unit and still be classified as short-term, provided the majority of its patients are admitted to units where the average length of stay is less than 30 days. Hospital units of institutions such as prisons and college infirmaries that are not open to the public and are contained within a nonhospital facility are not included in the category of community hospitals. Traditionally, the definition has included all nonfederal short-stay hospitals except facilities for persons with intellectual disabilities (formerly called mentally retarded). In a revised definition, the following additional sites were excluded: hospital units of institutions, and alcoholism and chemical dependency facilities.

Federal hospital—Federal hospitals are those operated by the federal government.

For-profit hospital—For-profit hospitals are operated for profit by individuals, partnerships, or corporations.

General hospital—General hospitals provide diagnostic, treatment, and surgical services for patients with a variety of medical conditions. According to WHO, these hospitals provide medical and nursing care for more than one category of medical discipline (e.g., general medicine, specialized medicine, general surgery, specialized surgery, and obstetrics). Excluded are hospitals, usually in rural areas, that provide a more limited range of care.

Nonprofit hospital—Nonprofit hospitals are those controlled by nonprofit organizations, such as religious organizations and fraternal societies.

Registered hospital—Registered hospitals are those registered with AHA. About 98% of U.S. hospitals are registered.

Short-stay hospital—In the National Health Interview Survey, short-stay hospitals are defined as any hospital or hospital department in which the type of service

provided is general; maternity; eye, ear, nose, and throat; children's; or osteopathic.

Special hospital—Special hospitals are those, such as psychiatric, tuberculosis, chronic disease, rehabilitation, maternity, and alcoholic or narcotic dependency facilities, that provide a particular type of service to the majority of their patients.

Hospital-based physician—See Appendix II, Physician.

Hospital day—See Appendix II, Days of care.

Hospital utilization—Estimates of hospital utilization (such as hospital discharge rate, days of care rate, average length of stay, and percentage of the population with a hospitalization) presented in *Health, United States* are based on data from three sources: Healthcare Cost and Utilization Project, National (Nationwide) Inpatient Sample (HCUP–NIS); National Health Interview Survey (NHIS); and American Hospital Association (AHA). Beginning with the 2012 data year, HCUP–NIS is a 20% sample of discharges (alive or deceased) from all community hospitals participating in HCUP, excluding rehabilitation and long-term acute care hospitals. For prior years, HCUP–NIS estimates are based on hospital stays for persons discharged alive or deceased from about 1,000 hospitals sampled to approximate a 20% stratified sample of U.S. community hospitals, excluding rehabilitation hospitals and long-term acute care hospitals. NHIS hospital utilization data are based on household interviews with a sample of the civilian noninstitutionalized population. NHIS respondents were asked whether they had any hospital stays in the past year. AHA data are from information reported by a census of hospitals. (Also see Appendix II, Average length of stay; Days of care; Discharge; and Appendix I, Healthcare Cost and Utilization Project [HCUP], National [Nationwide] Inpatient Sample; National Health Interview Survey [NHIS].)

Human immunodeficiency virus (HIV) disease—HIV disease is caused by infection with a cytopathic retrovirus, which in turn leads to destruction of parts of the immune system. A surveillance case for HIV requires laboratory-confirmed evidence of infection, including a positive result on a screening test for HIV antibody, followed by a positive result on a confirmatory test, or a positive result or detectable quantity on an HIV virologic test (see, CDC. HIV Surveillance Report, 2014; vol. 26. 2015. Available from: <http://www.cdc.gov/hiv/library/reports/surveillance/>).

Since 1985, many states and U.S. dependent areas have implemented HIV case reporting as part of their comprehensive HIV and AIDS surveillance programs. As of April 2008, all reporting areas (50 states, D.C., and the six U.S. dependent areas of American Samoa, Guam, the Northern Mariana Islands, Puerto Rico, the Republic of Palau, and the U.S. Virgin Islands) had implemented HIV case surveillance using a confidential system for name-based case reporting for both HIV infection and AIDS. To better capture and characterize populations in which HIV infection has been

newly diagnosed, including persons with evidence of recent HIV infection, many states report the prevalence of those living with a diagnosis of HIV infection, including those living with AIDS.

In 2008, changes were made to the case definition for HIV infection. The new case definition combined the two previous case definitions for HIV and AIDS and established a new disease staging classification. The term HIV/AIDS was replaced with the term “diagnosis of HIV infection,” which is defined as diagnosis of HIV infection regardless of the stage of disease (stage 1, 2, 3 [AIDS], or unknown) and refers to all persons with a diagnosis of HIV infection (see MMWR 2008;57[RR-10]:1–8). Mortality and morbidity coding for HIV disease are similar and have evolved over time.

In 2014, the HIV surveillance case definition was revised again to adapt to changes in diagnostic criteria used by laboratories and clinicians. The new case definition recognizes early HIV infection (stage 0); includes the distinction between HIV-1 and HIV-2 infections; consolidates staging systems for adults and children; simplifies surveillance criteria for opportunistic illnesses; and incorporates clinical criteria for reporting diagnoses without laboratory evidence. (See HIV Surveillance Report, 2014; vol. 26. 2015.)

The 2008 case definition was used to classify cases diagnosed from the beginning of the epidemic through 2013 and was used to classify HIV infection among adults and adolescents, using the following HIV infection classification staging system:

- *HIV infection, stage 1*: No AIDS-defining condition and either a CD4 count of 500 cells/μL or more or a CD4 percentage of total lymphocytes of 29% or more.
- *HIV infection, stage 2*: No AIDS-defining condition and either a CD4 count of 200–499 cells/μL or a CD4 percentage of total lymphocytes of 14%–28%.
- *HIV infection, stage 3 (AIDS)*: Documentation of an AIDS-defining condition, or either a CD4 count of less than 200 cells/μL or a CD4 percentage of total lymphocytes of less than 14%. Documentation of an AIDS-defining condition supersedes a CD4 count or percentage that would not by itself be the basis for a stage 3 (AIDS) classification.
- *HIV infection, stage unknown*: No reported information on AIDS-defining conditions and no information available on CD4 count or percentage (see MMWR 2008;57[RR-10]:1–8).

The 2014 case definition was used to classify cases diagnosed beginning in 2014, and is similar to the 2008 case definition except for the following: (a) inclusion of criteria for stage 0, (b) inclusion of CD4 testing criteria for stage 3 in children, and (c) changes in cutoffs for CD4 percentages of total lymphocytes used for classifications of stages 1 and 2 in persons aged 6 years and over. The 2014 case definition classifies HIV infection based on the following stages:

- *HIV infection, stage 0*: First positive HIV test result within 6 months after negative test result. After 6 months, the stage may be reclassified as 1, 2, 3, or unknown.
- *HIV infection, stages 1, 2, and 3*: Documentation of an AIDS-defining condition (excluding stage 0) is stage 3. Otherwise, the stage is determined by the lowest CD4 test result.
- *HIV infection, stage unknown*: No reported information on AIDS-defining conditions and no information available on CD4 count or percentage.

Mortality coding—Starting with 1999 data and the introduction of the 10th revision of the *International Classification of Diseases* (ICD-10), the title for this cause of death was changed from HIV infection to HIV disease, and the ICD codes were changed to B20–B24. Starting with 1987 data, NCHS introduced category numbers *042–*044 for classifying and coding HIV infection as a cause of death in ICD-9. The asterisks before the category numbers indicate that these codes were not part of the original ICD-9. HIV infection was formerly referred to as human T-cell lymphotropic virus-III/lymphadenopathy-associated virus (HTLV-III/LAV) infection. Before 1987, deaths involving HIV infection were classified to Deficiency of cell-mediated immunity (ICD-9, code 279.1) contained in the category All other diseases; to Pneumocystosis (ICD-9, code 136.3) contained in the category All other infectious and parasitic diseases; to Malignant neoplasms, including neoplasms of lymphatic and hematopoietic tissues; and to a number of other causes. Because of these coding changes, death statistics for HIV infection before 1987 are not strictly comparable with data for 1987 and subsequent years and therefore are not shown in *Health, United States*.

(Also see Appendix II, Acquired immunodeficiency syndrome [AIDS]; Cause of death; *International Classification of Diseases* [ICD]; *International Classification of Diseases, 9th Revision, Clinical Modification* [ICD-9-CM]; Table IV.)

Hypercholesterolemia—See Appendix II, Cholesterol.

Hypertension—See Appendix II, Blood pressure, high.

ICD; ICD codes—See Appendix II, Cause of death; *International Classification of Diseases* (ICD).

Illicit drug use—Illicit drug use refers to the use and misuse of illegal and controlled drugs.

Monitoring the Future (MTF) Study—In this school-based survey of secondary school students, information on illicit drug use is collected using self-completed questionnaires. The information is based on the following questions: “On how many occasions (if any) have you used marijuana in the last 30 days?” Similar questions are asked about a range of other drugs, including hallucinogens, inhalants, heroin, and so on. Questions on cocaine use include the following: “On how many occasions (if any) have you taken crack (cocaine in

chunk or rock form) during the last 30 days?” and “On how many occasions (if any) have you taken cocaine in any other form during the last 30 days?” Questions about prescription drugs—tranquilizers, sedatives, narcotic drugs other than heroin, and amphetamines—provide a description of the legitimate uses for those drugs and then ask respondents to include only use “...on your own, that is, without a doctor telling you to take them.”

National Survey on Drug Use & Health (NSDUH)—Information on illicit drug use is collected for survey participants aged 12 and over. Information on any illicit drug use includes any use of marijuana or hashish, cocaine, crack, heroin, hallucinogens, inhalants or methamphetamine, as well as misuse of prescription psychotherapeutic drugs. Current use (within the past month) is based on the question: “How long has it been since you last used [drug name]?” This answer is cross-checked with the following question: “Think specifically about the past 30 days, from [DATE] up to and including today. During the past 30 days, on how many days did you use [drug name]?” Starting in 2013, information about marijuana use that was recommended by a doctor or other health care professional has been collected; however, reported marijuana use is classified as illicit drug use. (Also see Appendix II, Substance use.)

Immunization—See Appendix II, Vaccination.

Incidence—Incidence is the number of cases of disease having their onset during a prescribed period of time. It is often expressed as a rate (e.g., the incidence of measles per 1,000 children aged 5–15 during a specified year). Measuring incidence may be complicated because the population at risk for the disease may change during the period of interest due to births, deaths, or migration, for example. In addition, determining whether a case is new—that is, whether its onset occurred during the prescribed period of time—may be difficult. Because of these difficulties in measuring incidence, many health statistics are instead measured in terms of prevalence. (Also see Appendix II, Prevalence.)

Income—See Appendix II, Family income.

Individual practice association (IPA)—See Appendix II, Health maintenance organization (HMO).

Industry of employment—For the presentation of data in *Health, United States*, industries are classified according to the North American Industry Classification System (NAICS). NAICS groups establishments into industries based on their production or supply function. Establishments using similar raw material inputs, capital equipment, and labor are classified in the same industry. This approach creates homogeneous categories well suited for economic analysis. NAICS uses a two- through six-digit hierarchical coding system to classify all economic activity. The first two digits of

the six-digit code designate the highest level of aggregation, into the government and 20 private industry sectors (Table VIII). Agriculture, forestry, fishing and hunting; mining; construction; and manufacturing are primarily goods-producing sectors, and the remaining 16 sectors are entirely service providing. NAICS allows for the classification of more than 1,000 industries. For more information on NAICS, see: <http://www.census.gov/eos/www/naics>.

Starting in 1997, NAICS replaced the Standard Industrial Classification (SIC) system, which was last updated in 1997. The SIC system focused on the manufacturing sector of the economy and provided significantly less detail for the now-dominant service sector, including newly developed industries in information services, health care delivery, and high-technology manufacturing. Although some titles in SIC and NAICS are similar, there is little comparability between the two systems because industry groupings are defined differently. Estimates classified by NAICS should not be compared with estimates that used SIC.

Infant death—An infant death is the death of a live-born child before his or her first birthday. Age at death may be further classified as neonatal or postneonatal. Neonatal deaths are those that occur before the 28th day of life; postneonatal deaths are those that occur within 28 days to under 1 year of age. (Also see Appendix II, Rate: Death and related rates.)

Injury—The International Classification of External Causes of Injuries (ICECI) Coordination and Maintenance Group defines injury as a (suspected) bodily lesion resulting from acute overexposure to energy (this can be mechanical, thermal, electrical, chemical, or radiant) interacting with the body in amounts or rates that exceed the threshold of physiological tolerance. The time between exposure to the energy and the appearance of an injury is short. In some cases, an injury results from an insufficiency of any of the vital elements (i.e., air, water, or warmth), as in strangulation, drowning, or freezing. Acute poisonings and toxic effects, including overdoses of substances and wrong substances given or taken in error are included, as are adverse effects and complications of therapeutic, surgical, and medical care. Psychological harm is excluded. Injuries can be intentional or unintentional (i.e., accidental). In NCHS data systems, external causes of nonfatal injuries are currently coded to the *International Classification of Diseases, 9th Revision, Clinical Modification, Supplementary Classification of External Causes of Injury and Poisoning*, and the codes are often referred to as E codes. See Table IX for a list of external causes of injury categories and E codes used in *Health, United States*. Also see the NCHS injury website at: <http://www.cdc.gov/nchs/injury.htm>; and see: ICECI Coordination and Maintenance Group. International Classification of External Causes of Injuries (ICECI), ver 1.2. Amsterdam, The Netherlands: Consumer Safety Institute; and Adelaide, Australia: Australian Institute of Health and Welfare National Injury Surveillance Unit, Flinders University;

2004. Available from: <http://www.who.int/classifications/icd/adaptations/iceci/en/index.html>. (Also see Appendix II, Diagnosis; Injury-related visit.)

Injury-related visit—In the National Hospital Ambulatory Medical Care Survey (NHAMCS), an emergency department visit was considered injury-related if the physician diagnosis was injury-related or an external cause-of-injury code (E code) was present (Table IX). Starting with *Health, United States, 2008*, an injury-related visit was redefined as an initial injury visit. In the 2001–2010 NHAMCS, an initial injury visit was the first visit to an emergency department for an injury that was characterized by either the first-listed diagnosis being a valid injury diagnosis or by a valid first-listed E code, regardless of the diagnosis code. Visits for which the first-listed diagnosis or the first-listed E code was for a complication of medical care or for an adverse event were not counted as injury visits. For 2001–2004 and 2007 and subsequent data years, the patient record form had a specific question on whether the episode of care was an initial visit for the problem. In the 2005 and 2006 surveys, this variable was not included, and in its place an imputed variable was constructed that indicated whether the visit was or was not the initial visit for the problem. For an explanation of the methodology used to create the imputed initial visit variable, see: <http://www.cdc.gov/nchs/data/ahcd/initialvisit.pdf>. For more information, see the CDC/NCHS Injury Data and Resources website at: <http://www.cdc.gov/nchs/injury.htm>; and Fingerhut LA. Recommended definition of initial injury visits to emergency departments for use with the NHAMCS–ED data. NCHS. Health E-Stats; 2006. Available from: <http://www.cdc.gov/nchs/data/hestat/injury/injury.htm>. (Also see Appendix II, Emergency department or emergency room visit; External cause of injury; Injury.)

Inpatient—An inpatient is a person who is formally admitted to the inpatient service of a hospital for observation, care, diagnosis, or treatment. (Also see Appendix II, Admission; Average length of stay; Days of care; Discharge; Hospital.)

Inpatient care—See Appendix II, Hospital utilization.

Inpatient day—See Appendix II, Days of care.

Instrumental activities of daily living (IADL)—IADLs are activities related to independent living and include preparing meals, managing money, shopping for groceries or personal items, performing light or heavy housework, and using a telephone. In the National Health Interview Survey, respondents are asked whether they or family members need the help of another person for handling routine IADL needs because of a physical, mental, or emotional problem.

In the Medicare Current Beneficiary Survey, if a sample person had any difficulty performing an activity by him- or herself and without special equipment, or did not perform the activity at all because of health problems, the person

Table VIII. Codes for industries, based on the North American Industry Classification System (NAICS)

<i>Industry</i>	<i>Code</i>
Agriculture, forestry, fishing and hunting	11
Mining, quarrying, and oil and gas extraction	21
Utilities	22
Construction	23
Manufacturing	31–33
Wholesale trade	42
Retail trade	44–45
Transportation and warehousing	48–49
Information	51
Finance and insurance	52
Real estate and rental and leasing	53
Professional, scientific, and technical services	54
Management of companies and enterprises	55
Administrative and support and waste management and remediation services	56
Educational services	61
Health care and social assistance	62
Arts, entertainment, and recreation	71
Accommodation and food services	72
Other services, except public administration	81
Public administration	92

SOURCE: Bureau of Labor Statistics. Available from: <http://www.census.gov/eos/www/naics/>.

was categorized as having a limitation in that activity. The limitation may have been temporary or chronic at the time of interview. Sampled persons in the community answered health status and functioning questions themselves, if able to do so. For sampled persons in a long-term care facility, a proxy such as a nurse answered questions about the sampled person's health status and functioning. (Also see Appendix II, Activities of daily living [ADL]; Complex activity limitation; Limitation of activity.)

Insurance—See Appendix II, Health insurance coverage.

Intermediate care facility—See Appendix II, Nursing home.

International Classification of Diseases (ICD)—The ICD is used to code and classify cause-of-death data. The ICD is developed collaboratively by the World Health Organization and 10 international centers, one of which is housed at NCHS. The purpose of the ICD is to promote international comparability in the collection, classification, processing, and presentation of health statistics. Since 1900, the ICD has been modified about once every 10 years, except for the 20-year interval between the 9th and 10th revisions (ICD–9 and ICD–10) (Table III). The purpose of the revisions is to stay abreast of advances in medical science. New revisions usually introduce major disruptions in time series of mortality statistics (Tables IV and V). For more information, see the NCHS ICD–10 website at: <http://www.cdc.gov/nchs/icd/icd10.htm>. (Also see Appendix II, Cause of death; Comparability ratio; *International Classification of Diseases, 9th Revision, Clinical Modification* [ICD–9–CM].)

Table IX. Codes for external causes of injury, from the *International Classification of Diseases, 9th Revision, Clinical Modification*

<i>External cause of injury category</i>	<i>E code</i>
All injury	E800–E869, E880–E929, E950–E999
Unintentional.	E800–E869, E880–E929
Motor vehicle traffic	E810–E819
Falls	E880–E886, E888
Struck by or against objects or persons	E916–E917
Caused by cutting and piercing instruments or objects.	E920
Intentional (suicide and homicide)	E950–E969, E979, E999.1
Undetermined	E980–E989
Other (includes legal intervention and operations of war).	E970–E978, E990–E999.0

SOURCE: Recommended framework of E code groupings for presenting injury morbidity data. Available from: http://www.cdc.gov/injury/wisqars/ecode_matrix.html, and the *International Classification of Diseases, 9th Revision, Clinical Modification*. Available from: <http://www.cdc.gov/nchs/icd/icd9cm.htm>.

International Classification of Diseases, 9th Revision, Clinical Modification (ICD–9–CM)—ICD–9–CM is based on, and is compatible with, the World Health Organization's ICD–9. The United States used ICD–9–CM to code morbidity diagnoses and inpatient procedures until October 1, 2015, when the *International Classification of Diseases, 10th Revision, Clinical Modification/Procedure Coding System* went into effect. ICD–9–CM consists of three volumes. Volumes 1 and 2 contain the diagnosis tabular list and index; Volume 3 contains the procedure classification (tabular list and index combined).

ICD–9–CM is divided into 17 chapters and two supplemental classifications. The chapters are arranged primarily by body system. In addition, there are chapters for Infectious and parasitic diseases; Neoplasms; Endocrine, nutritional, and metabolic diseases; Mental disorders; Complications of pregnancy, childbirth, and puerperium; Certain conditions originating in the perinatal period; Congenital anomalies; and Symptoms, signs, and ill-defined conditions. The two supplemental classifications are for factors influencing health status and contact with health services (V codes), and for external causes of injury and poisoning (E codes).

In *Health, United States*, morbidity data will be classified using ICD–9–CM until 2016 data are available and then morbidity data will be classified using *International Classification of Diseases, 10th Revision, Clinical Modification/Procedure Coding System* (ICD–10–CM/PCS). ICD–9–CM procedure categories and codes are shown in Table X. For more information about ICD–9–CM, see the NCHS Classification of Diseases, Functioning, and Disability website at: <http://www.cdc.gov/nchs/icd.htm>. (Also see Appendix II, *International Classification of Diseases [ICD]; International Classification of Diseases, 10th Revision, Clinical Modification/Procedure Coding System [ICD–10–CM/PCS]*.)

International Classification of Diseases, 10th Revision, Clinical Modification/Procedure Coding System (ICD–10–CM/PCS)—Use of ICD–10–CM/PCS to report medical diagnoses and inpatient procedures was implemented October 1, 2015. The transition to ICD–10 is

required for everyone covered by the Health Insurance Portability and Accountability Act (HIPAA). This change to ICD–10 does not affect Current Procedural Terminology (CPT) coding for outpatient procedures and physician services. ICD–10–CM/PCS consists of two parts: ICD–10–CM for diagnosis coding, and ICD–10–PCS for inpatient procedure coding. For more information about ICD–10–CM/PCS, see the NCHS Classification of Diseases, Functioning, and Disability website at: <http://www.cdc.gov/nchs/icd.htm> and the Centers for Medicare & Medicaid Services ICD–10 transition website at: <http://www.cms.gov/Medicare/Coding/ICD10/index.html>.

Late fetal death rate—See Appendix II, Rate: Death and related rates.

Leading causes of death—See Appendix II, Cause-of-death ranking.

Length of stay—See Appendix II, Average length of stay.

Life expectancy—Life expectancy is the average number of years of life remaining to a person at a particular age and is based on a given set of age-specific death rates—generally the mortality conditions existing in the period mentioned. Life expectancy may be determined by sex, race and Hispanic origin, or other characteristics, by using age-specific death rates for the population with that characteristic. (Also see Appendix II, Rate: Death and related rates.)

U.S. life tables by Hispanic origin were available starting with 2006 data. Life expectancy data for the Hispanic population were not available before 2006 for three major reasons: (a) coverage of the Hispanic population in the U.S. mortality statistics system was incomplete, (b) misclassification of Hispanic persons on death certificate data underestimated deaths in the Hispanic population, and (c) misstatement of age at the oldest ages in the Hispanic population led to an underestimation of mortality at the oldest ages.

Hispanic origin was added to the U.S. standard death certificate in 1989, but it was not adopted by every state

until 1997. By 1997, all states had reporting rates over 99%. Research on race and Hispanic origin reporting on U.S. death certificates found that misclassification of race and Hispanic origin accounts for a net underestimate of 5% for total Hispanic deaths and 1% for total non-Hispanic black deaths, and a net overestimate of 0.5% for non-Hispanic white deaths. To address the effects of age misstatement at the oldest ages, the probability of death for Hispanic persons over age 80 is estimated as a function of non-Hispanic white mortality with the use of the Brass relational logit model. For more information, see: Arias E. United States life tables by Hispanic origin. NCHS. Vital Health Stat 2010;2(152). Available from: http://www.cdc.gov/nchs/data/series/sr_02/sr02_152.pdf.

In 2000, the life table methodology was revised. The revised methodology is similar to that developed for the 1999–2001 decennial life tables. In 2008, the life table methodology was refined in two important ways. First, a logistic rather than a nonlinear least squares model was used to smooth and extrapolate the Vital and Medicare blended death rates at the older ages. Second, the age at which smoothing is begun was raised from 66 to 85 years or so, depending on the population. Values for 2001 and subsequent data years are based on the latest revision of the life table methodology. As a result, data post-2000 may differ from figures published previously. For a full description of the new life table methodology, see: Arias E. United States life tables, 2008. National vital statistics reports; vol 61 no 3. Hyattsville, MD: NCHS; 2012. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr61/nvsr61_03.pdf.

Limitation of activity—Limitation of activity may be defined in different ways, depending on the conceptual framework. In the National Health Interview Survey, limitation of activity refers to a long-term reduction in a person's capacity to perform the usual kind or amount of activities associated with his or her age group as a result of a chronic condition. Limitation of activity is assessed by asking persons a series of questions about limitations in their or a family member's ability to perform activities usual for their age group because of a physical, mental, or emotional problem. Persons are asked about limitations in activities of daily living, instrumental activities of daily living, play, school, work, difficulty walking or remembering, and any other activity limitations. For reported limitations, the causal health conditions are determined, and persons are considered limited if one or more of these conditions is chronic. Children under age 18 who receive special education or early intervention services are considered to have a limitation of activity. (Also see Appendix II, Activities of daily living [ADL]; Instrumental activities of daily living [IADL].)

Long-term care facility—A long-term care facility is a residence that provides a specific level of personal or medical care or supervision to residents. In the Medicare Current Beneficiary Survey, a residence is considered a long-term care facility if it has three or more long-term care

beds and answers affirmatively to at least one of three questions: “Does this facility (a) provide personal care services to residents, (b) provide continuous supervision of residents, (c) provide any long-term care?” Types of long-term care facilities include licensed nursing homes, skilled nursing homes, intermediate care facilities, retirement homes (that provide services), domiciliary or personal care facilities, distinct long-term care units in a hospital complex, mental health facilities and centers, assisted and foster care homes, and institutions for persons with intellectual disabilities (formerly called mentally retarded) and the developmentally disabled. (Also see Appendix II, Nursing home.)

Low birthweight—See Appendix II, Birthweight.

Mammography—A mammogram is an x-ray image of the breast used to detect irregularities in breast tissue. In the National Health Interview Survey, questions concerning use of mammography are asked on an intermittent schedule, and question content has differed across years. For 2013, women were asked when they had their most recent mammogram, and use of mammography was defined as “percent of women having a mammogram within the past two years.” Survey questions have changed over time as follows.

In 1987 and 1990, women were asked to report when they had their last mammogram. In 1991, women were asked whether they had a mammogram in the past 2 years. In 1993 and 1994, women were asked whether they had a mammogram within the past year, between 1 and 2 years ago, or over 2 years ago. In 1998, women were asked whether they had a mammogram a year ago or less, more than 1 year but not more than 2 years, or more than 2 years ago.

In 1999, women were asked when they had their most recent mammogram, in days, weeks, months, or years. Estimates for 1999 may be slightly overestimated in comparison with previous years due to the inclusion of women who responded “2 years ago” (10% of women), which could have included more than 2 years but less than 3 years.

In 2000 and 2003, women were asked when they had their most recent mammogram (asked to give month and year). Women who did not respond were given a follow-up question that used the 1999 wording, and women who did not respond to the 1999 wording were asked a second follow-up question that used the 1998 wording. Estimates for 2000 and 2003 may be slightly overestimated compared with estimates prior to 1999 due to the inclusion of women who responded “2 years ago” (2% of women), which could have included more than 2 years but less than 3 years.

In 2005, women were asked the same series of mammography questions as in the 2000 and 2003 surveys, but the questionnaire skip pattern was modified so that more women were asked the follow-up question using the

Table X. Codes for procedure categories for Healthcare Cost and Utilization Project data, from the *International Classification of Diseases, 9th Revision, Clinical Modification*

Procedure category	Code
Amputation of lower extremity (amputation of lower limb)	84.10–84.19
Appendectomy	47.0, 47.01, 47.09, 47.1, 47.11, 47.19
Arthroplasty knee (knee replacement)	00.80–00.84, 81.41–81.44, 81.46, 81.47, 81.54, 81.55
Cesarean section	74.0, 74.1, 74.2, 74.4, 74.99
Cholecystectomy (gall bladder removal).	51.21–51.24, 51.41–51.43, 51.49, 51.51, 51.59
Colorectal resection (removal of part of the bowel)	17.31–17.36, 17.39, 45.71–45.76, 45.79, 45.8, 45.81–45.83, 48.40–48.43, 48.49, 48.5, 48.50–48.52, 48.59, 48.61–48.66, 48.69
Coronary artery bypass graft (CABG)	36.10–36.17, 36.19, 36.2, 36.3, 36.31–36.34, 36.39
Enderterectomy (plaque removal from artery lining of brain, head, neck)	38.11, 38.12
Heart valve procedures	35.00–35.14, 35.20–35.28, 35.96, 35.97, 35.99
Hip replacement	00.70–00.77, 00.85–00.87, 81.51–81.53, 81.69
Hysterectomy	68.3, 68.31, 68.39, 68.4, 68.41, 68.49, 68.5, 68.51, 68.59, 68.6, 68.61, 68.69, 68.7, 68.71, 68.79, 68.9
Incision and excision of CNS (brain surgery)	01.01, 01.09, 01.21–01.28, 01.31, 01.32, 01.39, 01.41, 01.42, 01.51–01.53, 01.59
Insertion, revision, replacement, removal of cardiac pacemaker	00.50–00.54, 00.56, 00.57, 17.51, 17.52, 37.70–37.83, 37.85–37.87, 37.89, 37.94–37.98
Laminectomy (spine surgery)	03.02, 03.09, 80.5, 80.50, 80.51, 80.59, 84.59–84.69, 84.80–84.85
Ligation of fallopian tubes (“tying” of fallopian tubes)	66.21, 66.22, 66.29, 66.31, 66.32, 66.39
Oophorectomy (removal of one or both ovaries)	65.3, 65.31, 65.39, 65.4, 65.41, 65.49, 65.51–65.54, 65.61–65.64
Percutaneous coronary angioplasty (PTCA) (balloon angioplasty)	00.66, 17.55, 36.01, 36.02, 36.05
Small bowel resection (removal of part of the small bowel)	45.61–45.63
Spinal fusion.	81.00–81.09, 81.30–81.39, 81.61–81.64, 84.51
Tonsillectomy and/or adenoidectomy	28.2, 28.3, 28.6, 28.7
Treatment, fracture or dislocation of hip and femur	78.55, 78.65, 79.05, 79.15, 79.25, 79.35, 79.45, 79.55, 79.65, 79.75, 79.85, 79.95

NOTES: Procedures were classified by Clinical Classifications Software (CCS). For more information, see: <http://www.hcup-us.ahrq.gov/toolsoftware/ccs/AppendixBSinglePR.txt>.

SOURCE: Agency for Healthcare Research and Quality.

1998 wording. Thus, estimates for 2005 and subsequent years are more precise than estimates for 1999, 2000, and 2003. SAS code to categorize mammography data for 2000 and beyond is available from: http://www.cdc.gov/nchs/nhis/nhis_2005_data_release.htm. In 2008, 2010, and 2013, the mammography questions were identical to those asked in 2005.

Mammography screening recommendations have changed over time and vary in the recommended age to begin screening and the interval for screening. The current recommendation, made by the U.S. Preventive Services Task Force in 2016, is the use of screening mammography for breast cancer every 2 years in women aged 50–74, with additional guidance provided for women aged 40–49. For additional information, see: U.S. Preventive Services Task Force. Breast cancer: Screening. Rockville, MD: Agency for Healthcare Research and Quality; 2016. Available from: <http://www.uspreventiveservicestaskforce.org/Page/Document/UpdateSummaryFinal/breast-cancer-screening1?ds=1&s=breastcancer>; and U.S. Preventive

Services Task Force. Guide to clinical preventive services, 2014. Rockville, MD: Agency for Healthcare Research and Quality; 2014. Available from: <http://www.ahrq.gov/professionals/clinicians-providers/guidelines-recommendations/guide/index.html>.

Managed care—“Managed care” is a term originally used to refer to prepaid health plans (generally, health maintenance organizations, or HMOs) that furnish care through a network of providers under a fixed budget and “manage” costs. Increasingly, the term is also used to include preferred provider organizations (PPOs) and even forms of indemnity insurance coverage (i.e., “fee-for-service” insurance).

Medicare managed care includes a combination of risk- and cost-based plans. Risk-based plans receive a fixed prepayment per beneficiary per month to help pay for the cost of all covered services that a beneficiary may use. Each year, the Centers for Medicare & Medicaid Services (CMS) announces a “benchmark” amount for each county for coverage of Medicare Part A and Part B services. A managed

care plan contracting with Medicare then submits a “bid,” which represents the revenue it needs to cover these services. If the bid is above the benchmark, the difference must be charged in a premium to the enrollees of the plan. If the bid is below the benchmark, then a portion of the difference must be used to provide additional benefits to enrollees, with the Medicare trust funds receiving the remaining share. Starting with 2013 data, the term Medicare Risk Health Maintenance Organization was replaced with Medicare Advantage plan. Medicare Advantage plans include HMOs, PPOs, private fee-for-service plans, special needs plans, and Medicare medical savings account plans.

Cost-based plans are offered by an HMO or a competitive medical plan and are paid for their “reasonable costs” in providing Medicare services to enrollees, based on annual cost reports filed with CMS. For current definitions of the various Medicare managed care plans, see the CMS Medicare managed care manual. Ch 1, section 30, “Types of MA plans,” Baltimore, MD: CMS; 2011. Available from: <https://www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/Internet-Only-Manuals-IOMs-Items/CMS019326.html?DLPage=1&DLEntries=100&DLSort=0&DLSortDir=ascending>.

Medicare enrollees can choose to enroll in a managed care program (if available) or to receive services on a fee-for-service basis.

The two major Medicaid managed care categories are risk-based plans (managed care organizations, or MCOs) and primary care case management (PCCM) arrangements. Risk-based plans (MCOs) are paid a fixed monthly fee per enrollee. MCOs assume some or all of the financial risk for providing the services covered under the contract. PCCM providers are usually physicians, physician group practices, or entities employing or having other arrangements with such physicians, but they can also include nurse practitioners, nurse midwives, or physician assistants. These providers (also called gatekeepers) contract directly with the state to locate, coordinate, and monitor covered primary care (and sometimes additional services). PCCM providers are paid a per-patient case management fee and usually do not assume financial risk for the provision of services. Some states allow Medicaid enrollees to voluntarily enroll in managed care plans; most states require that at least certain categories of Medicaid beneficiaries join such plans. Both MCOs and PCCM arrangements include plans that provide specialized services to certain categories of Medicaid beneficiaries. For more information on state Medicaid managed care plans, see <http://www.medicaid.gov/>.

(Also see Appendix II, Health maintenance organization [HMO]; Medicare; Medicaid; Preferred provider organization [PPO].)

Marital status—Marital status is classified through self-reporting into the categories married and unmarried. The term “married” encompasses all married people, including

those separated from their spouses. “Unmarried” includes those who are single (never married), divorced, or widowed.

Birth file—In 1970, 39 states and D.C., and in 1975, 38 states and D.C., included a direct question about mother's marital status on the birth certificate. Since 1980, national estimates of births to unmarried women have been based on two methods for determining marital status: a direct question in the birth registration process and inferential procedures. In 1980–1996, marital status was reported on the birth certificates of 41–45 states and D.C.; with the addition of California in 1997, 46 states and D.C.; and in 1998–2001, 48 states and D.C. In 1997, all but four states (Connecticut, Michigan, Nevada, and New York), and in 1998, all but two states (Michigan and New York) included a direct question about mother's marital status on their birth certificates. In 1998–2007, marital status was imputed as married on birth records with missing information in the 48 states and D.C. where this information was obtained by a direct question. In 2008–2013 for 49 states and D.C., marital status is reported in the birth registration process.

For states lacking a direct question, marital status was inferred. Before 1980, the incidence of births to unmarried women in states with no direct question on marital status was assumed to be the same as the incidence in reporting states in the same geographic division. Starting in 1980, for states without a direct question, marital status was inferred by comparing the parents' and child's surnames. For 1994–1996, birth certificates in 45 states and D.C. included a question about the mother's marital status. Beginning in 1997, the marital status of women giving birth in California and Nevada has been determined by a direct question in the birth registration process. Beginning in June 15, 1998, Connecticut discontinued inferring the mother's marital status and added a direct question regarding mother's marital status to the state's birth certificate.

In 2005, Michigan added a direct question to the birth registration process but uses inferential procedures to update information collected using the direct question. In both Michigan and New York, a birth is inferred as nonmarital if either of these factors, listed in priority-of-use order, is present: (a) a paternity acknowledgment was received or (b) the father's name is missing. For 2006–2008 data, inferential procedures were used to compile birth statistics by marital status, in full or in part, for New York and Michigan, respectively. For 2009–2014, mother's marital status is inferred for New York.

National Health Interview Survey (NHIS)—In NHIS, marital status is asked of, or about, all persons aged 14 and over. Respondents are asked, “Are you now married, widowed, divorced, separated, never married, or living with a partner?”

Maternal age—See Appendix II, Age.

Medicaid—Medicaid was authorized in 1965 and became Title XIX of the Social Security Act. Medicaid is a jointly funded cooperative venture between the federal and state governments to assist states in the provision of adequate medical care to eligible persons. Within broad federal guidelines, each state establishes its own eligibility standards; determines the type, amount, duration, and scope of services; sets the rate of payment for services; and administers its own program.

Medicaid is the largest program providing medical and health-related services to America's poorest people. Medicaid was originally available only to individuals receiving cash assistance, but over time, Congress has expanded eligibility for children and selected adult groups. Most recently, the Affordable Care Act (ACA) and the Health Care and Education Reconciliation Act (HCERA) initiated significant changes to Medicaid. (Subsequent references to the ACA in this text will include changes enacted by either the ACA or HCERA).

States are mandated by federal law to cover certain population groups (mandatory eligibility groups) but are granted flexibility in covering other groups (optional eligibility groups). In the standard benefit package, states must cover mandatory benefits (e.g., physician services) but may choose to cover optional benefits (e.g., tuberculosis-related services). Prior to the ACA, many states expanded Medicaid coverage above the federal minimums and many states have chosen to continue this additional coverage. The major coverage change introduced by the ACA was to create a new eligibility group—Medicaid Expansion to Low-Income Adults—for those with incomes up to 138% of the federal poverty level (FPL) (133% by statute with an addition 5% income disregard). This is discussed further below.

States set individual eligibility criteria within federal minimum standards. In addition to Medicaid's financial eligibility requirements, individuals must satisfy federal and state requirements regarding residency, immigration status, and documentation of U.S. citizenship. The ACA provided for a standard application—available through the Medicaid program or through the Health Insurance Marketplace—and a standard method for calculating income eligibility for Medicaid, CHIP, and insurance affordability programs offered through the marketplace, primarily based on modified adjusted gross income (MAGI). Effective 2014, MAGI is used to determine Medicaid and CHIP eligibility for most nondisabled children and adults under age 65. By using one set of income eligibility rules across all insurance affordability programs, the ACA made it easier for people to apply for health coverage and enroll in the appropriate program.

Broadly, there are four major eligibility groups covered by most states: Children, Adults with Disabilities, Aged Adults, and Nondisabled Adults. These are discussed in more detail below.

- *Major Eligibility Groups*

Children—Medicaid was originally available only to individuals receiving cash assistance, but Congress has since expanded eligibility for children and other populations, making individuals eligible based on income below a specified percentage of the FPL. The ACA raised the minimum Medicaid eligibility for nondisabled children to 138% FPL (133% by statute with an addition 5% income disregard). Other eligible child groups include: infants born to women covered by Medicaid (known as “deemed newborns”), certain children in foster care or adoption assistance programs, certain children with disabilities, and children who use long-term services and supports. Like disabled adults, most states automatically qualify disabled children in the Supplemental Security Income (SSI) program for Medicaid coverage; eligibility is not determined by the newly introduced MAGI. Some states use more restrictive criteria to determine Medicaid eligibility of children with SSI. These criteria are usually based on income relative to the FPL and assets. Regardless of how they qualify, all children enrolled in Medicaid are entitled to the comprehensive set of health care services known as Early, Periodic Screening, Diagnostic and Treatment (EPSDT). These services include screening for and treatment of any vision or hearing problems, coverage for eyeglasses and hearing aids, and regular preventive dental care and treatment.

Adults with Disabilities—Adults with disabilities from physical conditions, intellectual or development disabilities, serious behavioral disorders, or serious mental illness may be eligible for Medicaid. The Supplemental Security Income (SSI) program pays benefits to disabled adults and children who have limited income and resources. Enrollment in SSI (or the Social Security Disability Insurance program, which provides Medicare to qualified individuals after a 24-month waiting period) automatically qualifies adults with disabilities for Medicaid in most states. However, some states use more restrictive criteria (known as 209(b) of the 1972 amendments to the Social Security Act) to determine Medicaid eligibility. These criteria are often based on income relative to the FPL and assets. As of January 2016, 11 states used more restrictive criteria than enrollment in SSI. Individuals with disabilities who are eligible for Medicaid are entitled to all services that are deemed medically necessary.

All states have the option of covering additional people with low incomes or high medical expenses through other eligibility pathways. These may include covering those at higher income levels; permitting persons with disabilities and high medical expenses to spend down until they are eligible for coverage; setting a special income level to cover institutionalized individuals with incomes up to 300% of the SSI benefit rate; extending coverage to individuals who receive home- and

community-based waiver services as an alternative to institutionalization; permitting working individuals who are severely impaired but whose earnings would otherwise disqualify them from Medicaid to buy into Medicaid; covering adults with disabilities who use long-term services and supports, based on an individual's functional status (known as level-of-care) and use of services (e.g., residence in a nursing facility, intermediate care facility for persons with intellectual disabilities, or mental health facility; or requiring significant home-based services).

Aged Adults—The Supplemental Security Income (SSI) program covers those with disabilities and also people aged 65 and older without disabilities who meet the financial limits. In most states, SSI enrollment automatically qualified those aged 65 and older for Medicaid. However, some states use more restrictive criteria (known as 209(b)) to determine Medicaid eligibility. The more restrictive criteria may consider income and assets, disability, or both. Most Medicaid enrollees aged 65 or over are also Medicare beneficiaries. This group is known as dual-eligible beneficiaries. Dual eligibles are eligible for the same Medicare benefits as other Medicare beneficiaries but have low incomes that make it difficult to afford the premiums and cost sharing required by Medicare, as well as the cost of services not covered by the Medicare program (e.g., long-term services and supports). Dual eligibles may qualify for partial Medicaid benefits (to cover Medicare premium and cost sharing) or full Medicaid benefits, in which case they get coverage for the full range of services offered by their state's Medicaid program.

Like coverage for adults with disabilities, states may extend Medicaid coverage to adults with low incomes or high medical expenses through other eligibility pathways, such as covering those with higher income levels or those with chronic conditions or low functional status requiring institutionalization or significant home-based services. There is considerable variation across states in the optional Medicaid services covered, which results in different benefits for dual-eligible beneficiaries depending on where they live.

Nondisabled Adults—Prior to the enactment of the ACA, most low-income nondisabled adults were not eligible for Medicaid unless they were in special groups (e.g., pregnant women, low-income parents, or other caretaker relatives with dependent children) or in states with demonstration programs that provided expanded coverage.

The major eligibility groups of nondisabled adults are now—Medicaid Expansion Coverage to Low-income Adults (the new adult group), Pregnant Women, Parents and Caretaker Relatives, and Adults Without Dependent Children. These groups and some specialty eligible groups—Breast and Cervical Cancer Prevention and

Treatment Program and Tuberculosis (TB)—are discussed below.

Medicaid Expansion to Low-income Adults—The ACA mandated Medicaid expansion, starting in 2014, to cover most adults under age 65 with incomes below 138% of FPL. This change would have ended the longstanding coverage gap for low-income adults; however, the Supreme Court ruling in June 2012 removed any sanction for states that do not implement the Medicaid expansion. As of January 2016, 31 states and D.C. had chosen to expand their Medicaid programs. They are: Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Hawaii, Illinois, Indiana, Iowa, Kentucky, Louisiana, Maryland, Massachusetts, Michigan, Minnesota, Montana, Nevada, New Jersey, New Hampshire, New Mexico, New York, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, Vermont, Washington, and West Virginia.

Pregnant Women—Since 1989, Congress has required Medicaid to cover pregnant women with low income. Currently, all but four states have extended Medicaid coverage to pregnant women above the currently required level of 138% FPL. Maternity-related services covered by the programs include prenatal care, labor and delivery, and 60 days of postpartum care. In Medicaid-expansion states, women at or below 138% FPL who are pregnant when they apply for Medicaid are not eligible for the new adult group. Medicaid coverage as a pregnant woman ends two months postpartum (after which the individual may be eligible in another Medicaid eligibility group).

Parents and Caretaker Relatives—Parents and caretaker relatives in low-income families with dependent children are eligible for coverage if their income meets the minimum eligibility levels established in 1996 for financial and medical assistance, which averages 41% of poverty. (1996 was the year of enactment for welfare reform, which held in place guaranteed Medicaid eligibility for those receiving cash benefits at that time.) States have the option to be more or less restrictive than the 1996 standards.

Adults without Dependent Children—Prior to the ACA, about one-half of states provide some coverage through Medicaid demonstration projects or state-funded programs for nondisabled adults who have limited incomes but do not otherwise qualify for Medicaid. Currently, 32 states have implemented the ACA Medicaid expansion for adults with incomes at or below 133% of the poverty line (with a 5% income disregard, so effectively 138%).

- *Other Eligibility Groups*

Breast and Cervical Cancer Prevention and Treatment Program—In 2000, Congress passed the Breast and Cervical Cancer Prevention and Treatment Act, which allowed states to offer eligible women who were

diagnosed with cancer through the CDC-funded screening program access to treatment through Medicaid. All states and D.C. have chosen to provide this coverage. For a woman to be eligible under this option, she must be under age 65; been screened through CDC's National Breast and Cervical Cancer Early Detection Program; be diagnosed with either breast or cervical cancer, including precancerous conditions; need treatment for breast or cervical cancer; and be uninsured and otherwise not eligible for Medicaid.

Tuberculosis (TB)—States can choose to provide Medicaid coverage of TB-related services for low-income individuals who are infected with TB. This eligibility group serves individuals who are not otherwise eligible for Medicaid based on the traditional eligibility categories.

Medicaid operates as a vendor payment program. States may pay health care providers directly on a fee-for-service basis, or states may pay for Medicaid services through various prepayment arrangements, such as through health maintenance organizations or other forms of managed care. Within federally imposed upper limits and restrictions, each state generally has broad discretion in determining both the payment method and rate for services. Thus, the Medicaid program varies considerably from state to state, as well as within each state over time. For more information, see: <http://www.medicaid.gov/> and <https://www.macpac.gov/>. (Also see Appendix II, Children's Health Insurance Program (CHIP); Health expenditures, national; Health insurance coverage; Health maintenance organization [HMO]; Managed care; and Appendix I, Medicaid Statistical Information System [MSIS].)

Medicaid payments—Under the Medicaid program, medical vendor payments are payments (expenditures) to medical vendors from the state through a fiscal agent, or to a health insurance plan. Adjustments are made for cost settlements, third-party recoupments, refunds, voided checks, and financial settlements that cannot be related to specific provided claims. Medicaid medical vendor payments, for purposes of this paper do not include payments to providers from other federal programs, or from third party payers, for Medicaid-eligible individuals; payments made from state medical assistance funds that are not federally matchable; disproportionate-share hospital payments, cost sharing, or enrollment fees collected from recipients or a third party; and administration and training costs. Medicaid payment data presented in *Health, United States* are from the Medical Statistical Information System (MSIS), which obtains payment data from electronic Medicaid data submitted to the Centers for Medicare & Medicaid Services by each state. Payment data are based on adjudicated claims for medical services reimbursed with Title XIX funds.

Medical specialty—See Appendix II, Physician specialty.

Medicare—Medicare is a nationwide program providing health insurance coverage to selected groups, regardless of income. The covered groups are (a) most people aged 65 and over; (b) people entitled to Social Security or Railroad Retirement disability benefits for at least 24 months (with the waiting period waived or reduced in certain situations); (c) government employees or spouses with Medicare-only coverage who have been disabled for more than 29 months (with the waiting period waived or reduced in certain situations); (d) most people with end-stage renal disease; and (e) certain people in the Libby, Montana, vicinity who are diagnosed with asbestos-related conditions. The program was enacted on July 30, 1965, as Title XVIII of the Social Security Act, "Health Insurance for the Aged and Disabled," and became effective on July 1, 1966.

From its inception, Medicare has included two separate but coordinated programs: Hospital Insurance (Part A) and Supplementary Medical Insurance (Part B). Part C (Medicare Advantage) was established by the Balanced Budget Act of 1997 as an expanded set of options for the delivery of health care under Medicare. Although all Medicare beneficiaries can receive their benefits through the original fee-for-service program, most beneficiaries enrolled in both Part A and Part B have the option to participate in a Medicare Advantage plan instead.

Organizations that seek to contract as Medicare Advantage plans must meet specific organizational, financial, and other requirements. Although most Medicare Advantage enrollees are in coordinated care plans, such as health maintenance organizations and preferred provider organizations, Medicare Advantage plans also include private fee-for-service plans, provider-sponsored organizations, special needs plans, and medical savings account plans (MSA plans, which provide benefits after a single high deductible is met). Medicare Advantage plans are generally paid on a capitation basis—that is, plans are paid a predetermined amount per member per month, which is adjusted according to the health status of the plans' members—and are required to provide at least those services covered by Parts A and B, except hospice services. Plans may (and in certain situations must) provide extra benefits (such as vision or hearing coverage) or reduce cost sharing or premiums.

The Medicare Prescription Drug, Improvement, and Modernization Act (also called the Medicare Modernization Act, or MMA) was passed on December 8, 2003. The MMA (Pub. L. 108–173) established a voluntary prescription drug benefit for Medicare beneficiaries and created a new Medicare Part D. People eligible for Medicare could begin to enroll in Part D beginning in January 2006. For more information, see: <http://www.medicare.gov/Pubs/pubs/pdf/10050.pdf> and <http://www.cms.gov/Research-Statistics-Data-and-Systems/Statistics-Trends-and-Reports/MedicareMedicaidStatSupp/2013.html>. (Also see Appendix II, Fee-for-service health insurance; Health insurance coverage; Health maintenance organization [HMO];

Managed care; and Appendix I, Medicare Administrative Data.)

Metropolitan statistical area (MSA)—The Office of Management and Budget (OMB) defines MSAs according to published standards that are applied to U.S. Census Bureau data. The standards are revised periodically, generally prior to the decennial census, and are applied to the census data to delineate the statistical areas. Revisions to the areas are implemented between censuses by using updated population estimates. The most recent standards were released in June 2010 (available from: http://www.whitehouse.gov/sites/default/files/omb/assets/fedreg_2010/06282010_metro_standards-Complete.pdf). In July 2015, OMB released a new delineation of the nation's metropolitan and micropolitan statistical areas based on the 2010 standards (available from: <https://www.whitehouse.gov/sites/default/files/omb/bulletins/2015/15-01.pdf>). New MSA delineations are incorporated into individual data systems at different times. In the 2000 and 2010 standards, an MSA is a county, or group of contiguous counties, that contains at least one urbanized area with a population of 50,000 or more. In addition to the county or counties that contain all or part of the urbanized area, an MSA may contain other counties if there are strong social and economic ties with the central county or counties, as measured by commuting. Counties that are not within an MSA are considered to be nonmetropolitan. For more information, see: <http://www.census.gov/population/metro/> and http://www.whitehouse.gov/omb/bulletins_fy05_b05-02. Most data by MSA currently in *Health, United States* are based on the June 2003 OMB definitions (2000 OMB standards applied to 2000 census data). (Also see Appendix II, Urbanization.)

National Health Interview Survey (NHIS)—For respondents to NHIS, designation of place of residence as metropolitan or nonmetropolitan is based on the following MSA definitions: for 2006 and beyond, on the June 2003 OMB definitions (2000 OMB standards applied to 2000 census data); for 1995–2005, on the June 1993 OMB definitions (1990 OMB standards applied to 1990 census data); and for 1985–1994, on the June 1983 OMB definitions (1980 OMB standards applied to 1980 census data). For estimates based on 2006 NHIS data combined with earlier years of NHIS, metropolitan status of residence for all years involved is based on the June 2003 definitions. Introduction of each set of standards may create a discontinuity in trends.

National Immunization Survey (NIS)—Designation of place of residence as metropolitan or nonmetropolitan for respondents to NIS is based on 2000 census data and the MSAs delineated in 2003, as well as the following versions and revisions of MSA definitions: for 2011 and 2012, on the December 2009 definitions; for 2010, on the November 2008 definitions, for New England, the county-based areas were used; for 2009, on the November 2007 definitions, for New England, the

county-based areas were used; for 2008, on the December 2006 definitions, for New England, the county-based areas were used; for quarter 4 of 2007, on the December 2006 definitions; for quarters 1–3 of 2007, on the December 2005 definitions, for New England, the county-based areas were used in 2007; for 2006, on the November 2004 definitions, for New England, the county-based areas were used; for 2005, on the December 2003 definitions, for New England, the county-based areas were used; for quarters 3 and 4 of 2004, on the December 2003 definitions; and for quarters 1 and 2 of 2004 and quarter 4 of 2003, on the June 2003 definitions. For 2003–2004 for New England, the county-based areas were used. For more information, see: <http://www.census.gov/population/metro/>.

Micropolitan statistical area—The Office of Management and Budget (OMB) defines a micropolitan statistical area as a nonmetropolitan county, or group of contiguous nonmetropolitan counties, that contains an urban cluster of 10,000–49,999 persons. A micropolitan statistical area may include surrounding counties that have strong social and economic ties with the central county or counties as measured by commuting. Nonmetropolitan counties that are not classified as part of a micropolitan statistical area are considered noncore.

OMB defines micropolitan statistical areas according to published standards that are applied to U.S. Census Bureau data. The standards are revised periodically, generally prior to the decennial census, and are applied to the census data to delineate the statistical areas. Revisions to the areas are implemented between censuses using updated population estimates. The most recent standards were released in June 2010 (available from: http://www.whitehouse.gov/sites/default/files/omb/assets/fedreg_2010/06282010_metro_standards-Complete.pdf). OMB released a new delineation of the nation's metropolitan and micropolitan statistical areas based on the 2010 standards in July 2015 (available from: <https://www.whitehouse.gov/sites/default/files/omb/bulletins/2015/15-01.pdf>). Data for micropolitan statistical areas currently in *Health, United States* are based on the 2013-based delineation as part of the 2013 NCHS Urban–Rural Classification Scheme for Counties. The micropolitan statistical area data will be updated when the new delineation is incorporated into individual data systems.

For more information about micropolitan statistical areas, see <http://www.census.gov/population/www/metroareas/metroarea.html>. (Also see Appendix II, Metropolitan statistical area [MSA]; Urbanization.)

Multum Lexicon Plus therapeutic class—Starting with 2003 data, NCHS used Lexicon Plus (Cerner Multum, Inc., Denver, CO.), a proprietary database, to assist with data editing and classification of human drugs. Starting with 2005 data, Lexicon Plus has also been used to assist with data collection. Data collected before 2003 were updated by adding a generic drug code from Lexicon Plus.

Lexicon Plus is a comprehensive database of all prescription and some nonprescription drug products available in the U.S. drug market. It uses a three-level nested category system to assign a therapeutic classification to each drug (e.g., for atenolol: cardiovascular agents [level 1]; beta-adrenergic blocking agents [level 2]; cardioselective beta blockers [level 3]). Not all drugs have three classification levels; some may only have two (e.g., for diltiazem: cardiovascular agents [level 1]; calcium channel blocking agents [level 2]). Other drugs may have only one classification level. All drugs in NCHS surveys were assigned into a Lexicon Plus drug category, even those drugs not found in the Lexicon Plus drug database. “Unspecified” drugs were assigned to their respective therapeutic category (e.g., hormones/hormone modifiers—unspecified: category ID = 97, category name = hormones/hormone modifiers).

Data presented in the *Health, United States* Trend Table on prescription drug use by drug class are based on the second level of the Lexicon Plus nested category system (e.g., calcium channel blocking agents). A drug may have up to four drug therapeutic categories; drugs classified into more than one class were counted in each class. For example, if a person reported taking lorazepam, that respondent was classified as taking an anticonvulsant, an antiemetic/antivertigo agent, and an anxiolytic, sedative, hypnotic drug.

The drug information file is updated along with each cycle of prescription medication data release. Some new therapeutic categories could be added, and a few assigned classification levels might be changed (e.g., alendronate now has three classification levels: metabolic agents [level 1], bone resorption inhibitors [level 2], and bisphosphonates [level 3]); under the prior drug information file, alendronate had two classification levels: hormones [level 1] and bisphosphonates [level 2]. Data presented in *Health, United States* used the most recent drug information file for all data years.

For more information, see: http://wwwn.cdc.gov/nchs/nhanes/1999-2000/RXQ_DRUG.htm.

Neonatal mortality rate—See Appendix II, Rate: Death and related rates.

Nonprofit hospital—See Appendix II, Hospital.

North American Industry Classification System (NAICS)—See Appendix II, Industry of employment.

Notifiable disease—A notifiable disease is one that, when diagnosed, health providers are required (usually by law) to report to state or local public health officials. Notifiable diseases are of public interest by reason of their contagiousness, severity, or frequency. For more information, see: http://www.cdc.gov/osels/ph_surveillance/nndss/nndsshis.htm.

Nursing home—In the Quality Improvement Evaluation System (QIES) (formerly the Online Survey Certification and Reporting [OSCAR]) database, a nursing home is a facility that is certified and meets the Centers for Medicare & Medicaid Services' long-term care requirements for Medicare and Medicaid eligibility.

After October 1, 1990, long-term care facilities that met the Omnibus Budget Reconciliation Act of 1987, Pub. L. No. 100–203, 101 Stat. 1330 nursing home reform requirements and were formerly certified under Medicaid as skilled nursing, nursing home, or intermediate care facilities were reclassified as nursing facilities. Medicare continues to certify skilled nursing facilities but not intermediate care facilities. State Medicaid programs can certify intermediate care facilities for individuals with intellectual disabilities (formerly called mentally retarded or developmentally disabled). To be certified for participation in Medicaid, nursing facilities must also be certified to participate in Medicare (except those facilities that have obtained waivers). Thus, most nursing home care is now provided in skilled care facilities.

(Also see Appendix II, Long-term care facility; Nursing home; Resident, health facility.)

Nursing home expenditures—See Appendix II, Health expenditures, national.

Obesity—See Appendix II, Body mass index (BMI).

Occupancy rate—In American Hospital Association statistics, hospital occupancy rate is calculated as the average daily census divided by the number of hospital beds, cribs, and pediatric bassinets set up and staffed on the last day of the reporting period, expressed as a percentage. Average daily census is calculated by dividing the total annual number of inpatients, excluding newborns, by 365 days to derive the number of inpatients receiving care on an average day during the annual reporting period. The occupancy rate for facilities other than hospitals is calculated as the number of residents at the facility reported on the day of interview, divided by the number of reported beds. In the Quality Improvement Evaluation System (QIES) (formerly the Online Survey Certification and Reporting [OSCAR]) database, occupancy is determined as of the day of certification inspection as the total number of residents on that day divided by the total number of beds on that day.

Office-based physician—See Appendix II, Physician.

Office visit—In the National Ambulatory Medical Care Survey, a physician's ambulatory practice (office) can be in any location other than in a hospital, nursing home, other extended care facility, patient's home, industrial clinic, college clinic, or family planning clinic. Offices in health maintenance organizations and private offices in hospitals are included. An office visit is any direct personal exchange between an ambulatory patient and a physician or members

of his or her staff for the purpose of seeking care and rendering health services. (Also see Appendix II, Outpatient visit.)

Operation—See Appendix II, Procedure.

Outpatient department—According to the National Hospital Ambulatory Medical Care Survey (NHAMCS), an outpatient department (OPD) is a hospital facility where nonurgent ambulatory medical care is provided. The following types of OPDs are excluded from NHAMCS: ambulatory surgical centers, chemotherapy, employee health services, renal dialysis, methadone maintenance, and radiology. (Also see Appendix II, Emergency department; Outpatient visit.)

Outpatient surgery—According to the American Hospital Association, outpatient surgery is a surgical operation, whether major or minor, performed on a patient who does not remain in the hospital overnight. Outpatient surgery may be performed in inpatient operating suites, outpatient surgery suites, or procedure rooms within an outpatient care facility. A surgical operation involving more than one surgical procedure is considered one surgical operation. (Also see Appendix II, Procedure.)

Outpatient visit—The American Hospital Association defines outpatient visits as visits for receipt of medical, dental, or other services at a hospital by patients who are not lodged in the hospital. Each appearance by an outpatient to each unit of the hospital is counted individually as an outpatient visit, including all clinic visits, referred visits, observation services, outpatient surgeries, and emergency department visits. In the National Hospital Ambulatory Medical Care Survey, an outpatient department visit is a direct personal exchange between a patient and a physician or other health care provider working under the physician's supervision for the purpose of seeking care and receiving personal health services. (Also see Appendix II, Emergency department or emergency room visit; Outpatient department.)

Overweight—See Appendix II, Body mass index (BMI).

Pap smear—A Pap smear (also known as a Papanicolaou smear or Pap test) is a microscopic examination of cells scraped from the cervix that is used to detect cancerous or precancerous conditions of the cervix or other medical conditions.

In the National Health Interview Survey, questions concerning Pap smear use are asked on an intermittent schedule, and the question content has differed slightly across years. For 2013, women were asked when they had their most recent Pap smear, and use of Pap smears was defined as “percent of women having a Pap smear within the past three years.” Survey questions have changed over time.

In 1987, women were asked to report when they had their most recent Pap smear, in days, weeks, months, or years. Women who did not respond were asked a follow-up question, “Was it 3 years ago or less, between 3 and 5 years, or 5 years or more ago?” In 1990 and 1991, Pap smear data in the past 3 years were not available. In 1993 and 1994, women were asked whether they had a Pap smear within the past year, between 1 and 3 years ago, or more than 3 years ago. In 1998, women were asked whether they had a Pap smear 1 year ago or less, more than 1 year but not more than 2 years ago, more than 2 years but not more than 3 years ago, more than 3 years but not more than 5 years ago, or more than 5 years ago.

In 1999, women were asked when they had their most recent Pap smear, in days, weeks, months, or years. Estimates for 1999 may be slightly overestimated in comparison with estimates for previous years due to the inclusion of women who responded “3 years ago” (4% of women), which could have included more than 3 years but less than 4 years.

In 2000 and 2003, women were asked when they had their most recent Pap smear (give month and year). Women who did not respond were given a follow-up question that used the 1999 wording, and women who did not respond to the follow-up question were asked a second follow-up question that used the 1998 wording. Estimates for 2000 and 2003 may be slightly overestimated in comparison with years prior to 1999 due to the inclusion of women who responded “3 years ago” (less than 1% of women), which could have included more than 3 years but less than 4 years.

In 2005, women were asked the same series of questions about Pap smear use as in the 2000 and 2003 surveys, but the questionnaire skip pattern was modified so that more women were asked the follow-up question using the 1998 wording, and these women were not uniformly coded as having had a Pap smear within the past 3 years. Thus, estimates for 2005 are more precise than estimates for 1999, 2000, and 2003. SAS code to categorize Pap smear data for 2000 and beyond is available from: http://www.cdc.gov/nchs/nhis/nhis_2005_data_release.htm.

In 2008, 2010, and 2013, Pap smear questions were similar to those asked in 2005.

All women aged 18 and over are asked the Pap smear question(s). Women who reported having had a hysterectomy (removal of the uterus, with or without removal of the ovaries and cervix) were still asked the Pap smear questions because a woman who has had a hysterectomy may still have Pap smear testing.

The current general recommendation, made by the U.S. Preventive Services Task Force in 2012, is the use of Pap smears for cervical cancer every 3 years in women aged 21–65, with additional recommendations available for women aged 30–65 who want to lengthen the recommended screening interval. In *Health, United States, 2014*, additional age groups (18–20, 21–24, and 21–44) were added to account for the new recommendation.

For more information on the recommendations, see: <http://www.uspreventiveservicestaskforce.org/Page/Document/RecommendationStatementFinal/cervical-cancer-screening#update-of-previous-uspstf-recommendation>.

The U.S. Preventive Services Task Force recommends against routine Pap smear screening in women who have had a total hysterectomy for benign disease. Therefore, two measures of Pap smear screening are presented in *Health, United States*: one among all women and one among women who did not report having a hysterectomy, although it is not known from NHIS data whether, for women who did report a hysterectomy, if it was for benign disease. Questions about whether the respondent had a hysterectomy were not asked in 2003. For other survey years, questions about hysterectomy in NHIS differed slightly, as follows.

In 1987, women who reported that they had not had a recent Pap smear were asked the most important reason they had not had a Pap smear; one reason women could select was because they had had a hysterectomy. In 1993, 1994, 1998, 1999, and 2013, women were asked, "Have you had a hysterectomy?" In 2000, 2005, 2008, and 2010, two questions were used to determine whether women had had a hysterectomy. Women were asked, "Have you had a hysterectomy?" In addition, women who reported that they had not had a recent Pap smear were asked the most important reason they had not had a Pap smear; one reason women could select was because they had had a hysterectomy. Women indicating in either of these questions that they had had a hysterectomy were excluded from the Pap smear screening estimates.

Pap smear screening recommendations have changed over time and vary in the recommended age to begin and end screening and the interval for screening. For current recommendations, see: U.S. Preventive Services Task Force. Cervical cancer, Screening: Summary of Recommendations and Evidence. Release date, March 2012. Rockville, MD: Agency for Healthcare Research and Quality; 2013. Available from: <http://www.uspreventiveservicestaskforce.org/uspstf/uspscerv.htm>.

Patient—See Appendix II, Inpatient; Office visit; Outpatient visit.

Percent change/percentage change—See Appendix II, Average annual rate of change (percent change).

Perinatal mortality rate; ratio—See Appendix II, Rate: Death and related rates.

Personal care home with or without nursing—See Appendix II, Nursing home.

Personal health care expenditures—See Appendix II, Health expenditures, national.

Physical activity, leisure-time—Starting with *Health, United States, 2010*, estimates on leisure-time physical

activity changed to reflect the federal *2008 Physical Activity Guidelines for Americans* (available from: <http://www.health.gov/PAGuidelines/guidelines/default.aspx>). Adults who met the 2008 guidelines reported at least 150 minutes per week of moderate-intensity or 75 minutes per week of vigorous-intensity aerobic physical activity (or an equivalent combination of moderate- and vigorous-intensity aerobic activity) and muscle-strengthening activities at least twice a week. The estimates for the percentage of Americans who met the 2008 guidelines for aerobic physical activity and muscle strengthening are not comparable with estimates in previous editions of *Health, United States* that showed the percentage of Americans with regular leisure-time physical activity. For more information, see: Carlson SA, Fulton JE, Schoenborn CA, Loustalot F. Trend and prevalence estimates based on the 2008 Physical Activity Guidelines for Americans. *Am J Prev Med* 2010;39(4):305–13.

Starting with 1998 data, leisure-time physical activity has been assessed in the National Health Interview Survey (NHIS) by asking adults a series of questions about how often they do vigorous or light/moderate physical activity of at least 10 minutes duration and about how long these sessions generally last. All questions related to leisure-time physical activity were phrased in terms of current behavior and lack a specific reference period. Vigorous physical activity is described as causing heavy sweating or a large increase in breathing or heart rate and light/moderate as causing light sweating or a slight-to-moderate increase in breathing or heart rate. Adults were also asked about how often they did leisure-time physical activities specifically designed to strengthen their muscles, such as lifting weights or doing calisthenics. The 2008 guidelines recommend any kind of aerobic activity, not just leisure-time aerobic activity, so the leisure-time aerobic activity estimates presented in this report may underestimate the percentage of adults who met the 2008 guidelines for aerobic activity. For more information, see the NHIS Adult Physical Activity Information website at: http://www.cdc.gov/nchs/nhis/physical_activity.htm.

Physician—Data on physician characteristics are obtained through physician self-report from the American Medical Association's (AMA) Physician Masterfile. Although the AMA collects data for both doctors of medicine (MDs) and doctors of osteopathy (DOs), in *Health, United States* data for DOs come from the American Osteopathic Association.

Active (or professionally active) physician—These physicians are currently engaged in patient care or other professional activity for a minimum of 20 hours per week. Other professional activity includes administration, medical teaching, research, and other activities such as employment with insurance carriers, pharmaceutical companies, corporations, voluntary organizations, and medical societies. Physicians who are retired, semiretired, working part-time, or not practicing are classified as inactive and are excluded. Also excluded are physicians with unknown address and physicians who did not

provide information on type of practice or present employment (not classified).

Hospital-based physician—These physicians are employed under contract with hospitals to provide direct patient care and include physicians in residency training (including clinical fellows) and full-time members of the hospital staff.

Office-based physician—These physicians are engaged in seeing patients in solo practice, group practice, two-physician practice, other patient care employment, or in providing inpatient services such as those offered by pathologists and radiologists.

Data for physicians are presented by type of education (doctor of medicine or doctor of osteopathy); place of education (U.S. medical graduates and international medical graduates); activity status (professionally active and inactive); area of specialty; and geographic area. (Also see Appendix II, Physician specialty.)

Physician specialty—A physician specialty is any specific branch of medicine in which a physician may concentrate. Data are based on physician self-reports of their primary area of specialty. Physician data are broadly categorized into two areas of practice: those who provide primary care and those who provide specialty care.

Primary care generalist—These physicians practice in the general fields of family medicine, general practice, internal medicine, obstetrics and gynecology, and pediatrics. Specifically excluded are primary care specialists associated with these generalist fields.

Primary care specialist—These specialists practice in the primary care subspecialties of family medicine, internal medicine, obstetrics and gynecology, and pediatrics. Family medicine subspecialties include geriatric medicine and sports medicine. Internal medicine subspecialties include adolescent medicine, critical care medicine, diabetes, endocrinology, diabetes and metabolism, hematology, hepatology, hematology/oncology, cardiac electrophysiology, infectious diseases, clinical and laboratory immunology, geriatric medicine, sports medicine, nephrology, nutrition, medical oncology, pulmonary critical care medicine, and rheumatology. Obstetrics and gynecology subspecialties include hospice and palliative medicine (obstetrics and gynecology), maternal and fetal medicine, critical care medicine (obstetrics and gynecology), and reproductive endocrinology. Pediatric subspecialties include adolescent medicine, pediatric critical care medicine, pediatrics/internal medicine, neonatal-perinatal medicine, pediatric allergy, pediatric cardiology, pediatric endocrinology, pediatric infectious disease, pediatric pulmonology, medical toxicology (pediatrics), pediatric emergency medicine, pediatric gastroenterology, pediatric hematology/oncology, clinical and laboratory immunology (pediatrics), pediatric nephrology, pediatric rheumatology, and sports medicine (pediatrics).

Specialty care physician—These physicians are sometimes called specialists and include primary care specialists listed above in addition to all other physicians not included in the generalist definition. Specialty fields include allergy and immunology, aerospace medicine, anesthesiology, cardiovascular diseases, child and adolescent psychiatry, colon and rectal surgery, dermatology, diagnostic radiology, forensic pathology, gastroenterology, general surgery, medical genetics, neurology, nuclear medicine, neurological surgery, occupational medicine, ophthalmology, orthopedic surgery, otolaryngology, psychiatry, public health and general preventive medicine, physical medicine and rehabilitation, plastic surgery, anatomic and clinical pathology, pulmonary diseases, radiation oncology, thoracic surgery, urology, addiction medicine, critical care medicine, legal medicine, and clinical pharmacology.

(Also see Appendix II, Physician.)

Population—The U.S. Census Bureau collects and publishes data on populations in the United States according to several different definitions. Various statistical systems then use the appropriate population for calculating rates. (Also see Appendix I, Population Census and Population Estimates.)

Resident population includes persons whose usual place of residence (i.e., the place where one usually lives and sleeps) is in one of the 50 states or D.C. It includes members of the Armed Forces stationed in the United States and their families. It excludes members of the Armed Forces stationed outside the United States and civilian U.S. citizens whose usual place of residence is outside the United States. The resident population is the denominator used to calculate birth and death rates and incidence of disease.

Civilian population is the resident population excluding members of the Armed Forces, although families of members of the Armed Forces are included. The civilian population is the denominator for emergency department visit rates using the National Hospital Ambulatory Medical Care Survey—Emergency Department Component.

Civilian noninstitutionalized population is the civilian population excluding persons residing in institutions (such as nursing homes, prisons, jails, mental hospitals, and juvenile correctional facilities). U.S. Census Bureau estimates of the civilian noninstitutionalized population are used to calculate sample weights for the National Health Interview Survey, the National Health and Nutrition Examination Survey, and the National Survey of Family Growth, and as denominators for rates calculated for the National Ambulatory Medical Care Survey and the National Hospital Ambulatory Medical Care Survey—Outpatient Department Component.

Postneonatal mortality rate—See Appendix II, Rate: Death and related rates.

Poverty—Two related versions of federal poverty measures are shown in *Health, United States*. The first measure—a ratio of family income to federal poverty threshold—is constructed using poverty thresholds from the U.S. Census Bureau. Poverty thresholds are updated annually for inflation by the Census Bureau using the Consumer Price Index for all urban consumers (CPI-U). Poverty thresholds include a set of money income thresholds that vary by family size and composition but do not vary geographically. Families or individuals with income below the appropriate threshold are classified as below poverty. For example, the average poverty threshold for a family of four was \$24,230 in 2014, \$23,834 in 2013, \$22,315 in 2010, \$17,604 in 2000, and \$13,359 in 1990. For more information, see the U.S. Census Bureau's poverty threshold website at: <http://www.census.gov/hhes/www/poverty/poverty.html>.

The second poverty measure used in *Health, United States* is a ratio of family income to the HHS poverty guidelines. Poverty guidelines are derived from the U.S. Census Bureau's poverty thresholds and are issued annually by HHS. These guidelines are often used to determine eligibility in certain federal programs. The HHS poverty guidelines take into account family size and state (coterminous, Alaska, Hawaii), but not family composition. For more information, see HHS, Office of the Assistant Secretary for Planning and Evaluation. Poverty Guidelines, Research, and Measurement website at: <http://aspe.hhs.gov/poverty/index.cfm>.

National Health Interview Survey (NHIS)—For data years prior to 1997, a ratio of family income to U.S. Census Bureau poverty threshold was computed taking into account family income and family size. Starting with 1997 data, the poverty ratio was based on family income, family size, and family composition (number of children in the family, and for families with two or fewer adults the age of the adults in the family). (Also see Appendix II, Consumer Price Index [CPI]; Family income; and Appendix I, Current Population Survey [CPS]; National Health Interview Survey [NHIS].)

National Health and Nutrition Examination Survey (NHANES)—NHANES uses the U.S. Census Bureau's Current Population Survey (CPS) definition of family to group household members into a family unit. A poverty ratio is computed by dividing family income by the HHS poverty guidelines specific to family size, as well as the appropriate guideline year, and state. See: Johnson CL, Paulose-Ram R, Ogden CL, et al. National Health and Nutrition Examination Survey: Analytic guidelines, 1999–2010. NCHS. Vital Health Stat 2(161). 2013. Available from: http://www.cdc.gov/nchs/data/series/sr_02/sr02_161.pdf.

Preferred provider organization (PPO)—A PPO is a type of medical plan in which coverage is provided to participants through a network of selected health care providers, such as

hospitals and physicians. Enrollees may seek care outside the network but pay a greater percentage of the cost of coverage than within the network. (Also see Appendix II, Health maintenance organization [HMO]; Managed care.)

Prevalence—Prevalence is the number of cases of a disease, number of infected persons, or number of persons with some other attribute present during a particular interval of time. It is often expressed as a rate (e.g., the prevalence of diabetes per 1,000 persons during a year). (Also see Appendix II, Incidence.)

Primary care specialty—See Appendix II, Physician specialty.

Private expenditures—See Appendix II, Health expenditures, national.

Procedure—Procedures can include surgical procedures (such as appendectomies), diagnostic procedures (such as spinal taps), and therapeutic treatments (such as infusion of a cancer chemotherapeutic substance) reported on a patient's medical record. In *Health, United States*, procedures are coded according to the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM) until 2016 data are available, and then procedures will be classified using the *International Classification of Diseases, 10th Revision, Clinical Modification/Procedure Coding System* (ICD-10-CM/PCS).

Healthcare Cost and Utilization Project, National (Nationwide) Inpatient Sample (HCUP-NIS)—Currently, up to 15 procedures are coded using ICD-9-CM procedure codes per hospital stay in the HCUP-NIS database. Starting with 2016 data, procedures will be coded according to the *International Classification of Diseases, 10th Revision, Clinical Modification/Procedure Coding System*. For each record, a principal procedure is identified as the first procedure listed. HCUP-NIS procedure data presented in *Health, United States* are limited to operating room procedures that are principal procedures (first-listed). Valid operating room procedures were identified according to diagnosis-related groups (DRGs) software. For DRG development, physician panels classify all ICD-9-CM procedure codes based on whether the procedure would be performed in operating rooms in most hospitals. Clinical Classifications Software (CCS) was used to categorize ICD-9-CM principal operating room procedure codes into 1 of 231 clinically meaningful categories. CCS was developed at the Agency for Healthcare Research and Quality as a tool for clustering patient procedures into a manageable number of clinically meaningful categories. It is periodically updated. For more information on CCS, see: <http://www.hcup-us.ahrq.gov/toolssoftware/ccs/AppendixBSinglePR.txt>. The top-ranking operating room procedure categories by age group, based on the number of discharges and total national costs, are presented in *Health, United States* (Table X).

CCS categories labeled “other” are not presented because these comprise miscellaneous procedures that do not form a homogeneous group.

(Also see Appendix II, Outpatient surgery.)

Proprietary hospital—See Appendix II, Hospital.

Public expenditures—See Appendix II, Health expenditures, national.

Purchasing power parities (PPPs)—PPPs are calculated rates of currency conversion that equalize the purchasing power of different currencies by eliminating the differences in price levels between countries. PPPs show the ratio of prices in national currencies for the same good or service in different countries. PPPs can be used to make intercountry comparisons of the gross domestic product (GDP) and its component expenditures. (Also see Appendix II, Gross domestic product [GDP].)

Race—In 1977, the Office of Management and Budget (OMB) issued “Race and Ethnic Standards for Federal Statistics and Administrative Reporting” (Statistical Policy Directive 15) to promote comparability of data among federal data systems. The 1977 Standards called for the federal government’s data systems to classify individuals into the following four racial groups: American Indian or Alaskan Native, Asian or Pacific Islander, black, and white. Depending on the data source, the classification by race was based on self-classification or on observation by an interviewer or other person filling out the questionnaire.

In 1997, revisions were announced for classification of individuals by race within the federal government’s data systems. (See: Revisions to the standards for the classification of federal data on race and ethnicity. Fed Regist 1997 October 30;62(210):58781–90.) The 1997 Standards specify five racial groups: American Indian or Alaska Native, Asian, black or African American, Native Hawaiian or Other Pacific Islander, and white. These five categories are the minimum set for data on race in federal statistics. The 1997 Standards also offer an opportunity for respondents to select more than one of the five groups, leading to many possible multiple-race categories. As with the single-race groups, data for the multiple-race groups are to be reported when estimates meet agency requirements for reliability and confidentiality. The 1997 Standards allow for observer or proxy identification of race but clearly state a preference for self-classification. The federal government considers race and Hispanic origin to be two separate and distinct concepts. Thus, Hispanic persons may be of any race. Federal data systems were required to comply with the 1997 Standards by 2003.

National Health Interview Survey (NHIS)—Starting with *Health, United States, 2002*, race-specific estimates based on NHIS were tabulated using the 1997 Standards for data year 1999 and beyond and are not strictly comparable with estimates for earlier years. The 1997

Standards specify five single-race categories plus multiple-race categories. Estimates for specific race groups are shown when they meet requirements for statistical reliability and confidentiality. The race categories white only, black or African American only, American Indian or Alaska Native only, Asian only, and Native Hawaiian or Other Pacific Islander only include persons who reported only one racial group; the category 2 or more races includes persons who reported more than one of the five racial groups in the 1997 Standards or one of the five racial groups and “some other race.” Prior to data year 1999, data were tabulated according to the 1977 Standards, with four racial groups, and the Asian only category included Native Hawaiian or Other Pacific Islander. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Differences between estimates tabulated using the two standards for data year 1999 are discussed in the footnotes for each NHIS table in *Health, United States 2002, 2003, and 2004* editions. Available from: <http://www.cdc.gov/nchs/hus/previous.htm#editions>.

Tables XI and XII illustrate NHIS data tabulated by race and Hispanic origin according to the 1997 and 1977 Standards for two health statistics (cigarette smoking and private health insurance coverage). In these examples, three separate tabulations using the 1997 Standards are shown: (a) Race: mutually exclusive race groups, including several multiple-race combinations; (b) Race, any mention: race groups that are not mutually exclusive because each race category includes all persons who mention that race; and (c) Hispanic origin and race: detailed race and Hispanic origin with a multiple-race total category. Where applicable, comparison tabulations by race and Hispanic origin are shown based on the 1977 Standards. Because there are more race groups with the 1997 Standards, the sample size of each race group under the 1997 Standards is slightly smaller than the sample size under the 1977 Standards. Only those few multiple-race groups with sufficient numbers of observations to meet standards of statistical reliability are shown. These tables also illustrate changes in labels and group categories resulting from the 1997 Standards. The race designation black was changed to black or African American, and the ethnicity designation Hispanic was changed to Hispanic or Latino.

Survey data included in *Health, United States*, other than NHIS, the National Survey of Drug Use & Health (NSDUH), and the National Health and Nutrition Examination Survey (NHANES), generally do not permit tabulation of estimates for the detailed race and ethnicity categories shown in Tables XI and XII, either because race data based on the 1997 Standards categories are not yet available or because there are insufficient numbers of observations in certain subpopulation groups to meet statistical reliability or confidentiality requirements.

To improve the quality of data on ethnicity and race in NHIS, hot-deck imputation of selected race and ethnicity variables was done for the first time in the 2000 NHIS and continued to be used for subsequent data years. Starting with 2003 data, records for persons for whom “other race” was the only race response were treated as having missing data on race and were added to the pool of records for which selected race and ethnicity variables were imputed. Prior to the 2000 NHIS, a crude imputation method that assigned a race to persons with missing values for the variable MAINRACE (the respondent's classification of the race he or she most identified with) was used. Under these procedures, if an observed race was recorded by the interviewer, it was used to code a race value. If there was no observed race value, all persons who had a missing value for MAINRACE and were identified as Hispanic on the Hispanic origin question were coded as white. In all other cases, non-Hispanic persons were coded as “other race.” Additional information on the NHIS methodology for imputing race and ethnicity is available from the survey documentation at: http://www.cdc.gov/nchs/nhis/quest_data_related_1997_forward.htm and from the NHIS race and Hispanic origin home page at: <http://www.cdc.gov/nchs/nhis/rhoi.htm>.

National Health and Nutrition Examination Survey (NHANES)—Starting with *Health, United States, 2003*, race-specific estimates based on NHANES were tabulated using the 1997 Standards for data years 1999 and beyond. Prior to data year 1999, the 1977 Standards were used. Because of the differences between the two standards, the race-specific estimates shown in Trend Tables based on NHANES for 1999 and beyond are not strictly comparable with estimates for earlier years. Race in NHANES I and II was determined primarily by interviewer observation; starting with NHANES III, race was self-reported by survey participants.

The NHANES sample for data years 1999–2006 was designed to provide estimates specifically for persons of Mexican origin and not for all Hispanic-origin persons in the United States. Persons of Hispanic origin other than Mexican origin were entered into the sample with different selection probabilities that are not nationally representative of the total U.S. Hispanic population. Starting with 2007–2008 data, all Hispanic persons were oversampled, not just persons of Mexican origin. Oversampling of the black population was continued. Starting in 2011, NHANES oversampled the non-Hispanic Asian population. In *Health, United States*, estimates are shown for non-Hispanic white, non-Hispanic black, and Mexican-origin persons, as well as Hispanic-origin and non-Hispanic Asian persons, where possible. Although data were collected according to the 1997 Standards, there are insufficient numbers of observations during this period to meet statistical reliability or confidentiality requirements for reporting estimates for additional race categories.

National Survey on Drug Use & Health (NSDUH)—Race-specific estimates based on NSDUH are tabulated using the 1997 Standards. Estimates in the NSDUH Trend Table begin with data year 2002. Estimates for specific race groups are shown when they meet requirements for statistical reliability and confidentiality. The race categories white only, black or African American only, American Indian or Alaska Native only, Asian only, and Native Hawaiian or Other Pacific Islander only include persons who reported only one racial group; the category 2 or more races includes persons who reported more than one of the five racial groups in the 1997 Standards or one of the five racial groups and “some other race.”

National Vital Statistics System (NVSS)—Some of the states in the Vital Statistics Cooperative Program are still revising their birth and death records to conform to the 1997 Standards on race and ethnicity. During the transition to full implementation of the 1997 Standards, vital statistics data will continue to be presented for four major race groups (white, black or African American, American Indian or Alaska Native, and Asian or Pacific Islander) in accordance with the 1977 Standards.

Birth file—Information about the race and Hispanic origin of the mother and father is provided by the mother at the time of birth and is recorded on the birth certificate or fetal death record. Since 1980, birth rates, birth characteristics, and death rates for live-born infants and fetal deaths are presented in *Health, United States* according to race of the mother. Before 1980, data were tabulated by race of the newborn and fetus, taking into account the race of both parents. If the parents were of different races and one parent was white, the child was classified according to the race of the other parent. When neither parent was white, the child was classified according to father's race, with one exception: if either parent was Hawaiian, the child was classified Hawaiian. Before 1964, if race was unknown, the birth was classified as white. Starting in 1964, unknown race was classified according to information on the birth record. Starting with the 2000 census, the race and ethnicity data used for denominators (population) to calculate birth and fertility rates have been collected in accordance with the 1997 revised OMB standards for race and ethnicity. However, the numerators (births) will not be compatible with the denominators until all the states revise their birth certificates to reflect the new standards. To compute rates, it is currently necessary to bridge population data for multiple-race persons to single-race categories. (Also see Appendix I, Population Census and Population Estimates, Bridged-race Population Estimates.)

Starting with 2003 data, some states began using the 2003 revision of the U.S. Standard Certificate of Live Birth, which allows the reporting of more than one race (multiple races). For 2014 data, 49 states, D.C., Guam,

Table XI. Current cigarette smoking among adults aged 18 and over, by race and Hispanic origin under the 1997 and 1977 Standards for federal data on race and ethnicity: United States, average annual 1993–1995

1997 Standards	Sample size	Percent	Standard error	1977 Standards	Sample size	Percent	Standard error
White only	46,228	25.2	0.26	White	46,664	25.3	0.26
Black or African American only	7,208	26.6	0.64	Black	7,334	26.5	0.63
American Indian or Alaska Native only	416	32.9	2.53	American Indian or Alaskan Native	480	33.9	2.38
Asian only	1,370	15.0	1.19	Asian or Pacific Islander	1,411	15.5	1.22
2 or more races total	786	34.5	2.00				
Black or African American; white	83	*21.7	6.05				
American Indian or Alaska Native; white	461	40.0	2.58				
Race, any mention							
White, any mention	46,882	25.3	0.26				
Black or African American, any mention	7,382	26.6	0.63				
American Indian or Alaska Native, any mention	965	36.3	1.71				
Asian, any mention	1,458	15.7	1.20				
Native Hawaiian or Other Pacific Islander, any mention	53	*17.5	5.10				
Hispanic origin and race							
Not Hispanic or Latino:				Non-Hispanic:			
White only	42,421	25.8	0.27	White	42,976	25.9	0.27
Black or African American only	7,053	26.7	0.65	Black	7,203	26.7	0.64
American Indian or Alaska Native only	358	33.5	2.69	American Indian or Alaskan Native	407	35.4	2.53
Asian only	1,320	14.8	1.21	Asian or Pacific Islander	1,397	15.3	1.24
2 or more races total	687	35.6	2.15				
Hispanic or Latino	5,175	17.8	0.65	Hispanic	5,175	17.8	0.65

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error of 20%–30%.

NOTES: The Office of Management and Budget's (OMB) *1997 Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* specifies five race groups (white, black or African American, American Indian or Alaska Native, Asian, and Native Hawaiian or Other Pacific Islander) and allows respondents to report one or more race groups. Estimates for single-race and multiple-race groups not shown above do not meet standards for statistical reliability or confidentiality (relative standard error greater than 30%). Race groups under the 1997 Standards were based on the question, "What is the group or groups which represents [person's] race?" For persons who selected multiple groups, race groups under the OMB's 1977 *Race and Ethnic Standards for Federal Statistics and Administrative Reporting* were based on the additional question, "Which of those groups would you say best represents [person's] race?" Race-specific estimates in this table were calculated after excluding respondents of other and unknown race. Other published race-specific estimates are based on files in which such responses have been edited. Estimates are age-adjusted to the year 2000 standard population using five age groups: 18–24, 25–34, 35–44, 45–64, and 65 and over. See Appendix II, Age adjustment.

SOURCE: CDC/NCHS, National Health Interview Survey. See Appendix I, National Health Interview Survey (NHIS).

and Northern Marianas allowed the reporting of multiple-race data. The 49 states and D.C. represented 99% of all U.S. resident births. In 2014, multiple race was reported for slightly more than 2% of mothers in the states that permitted reporting of more than one race. In 2014, data from the vital records of the remaining state, and two territories, followed the 1977 OMB Standards and reported the minimum set of four race categories, compared with the minimum of five race categories for the 1997 Standards. To provide uniformity and comparability of data during the transition to the 2003 revision, before multiple-race

data are available for all reporting areas, the responses of those who reported more than one race are bridged to a single race. For more information on the adoption of the 2003 revision of the U.S. Standard Certificate of Live Birth, see the Technical Notes section of the annual series of "Births: Final Data" reports, available from: <http://www.cdc.gov/nchs/products/nvsr.htm>.

Although the bridging procedure imputes multiple race of mothers to one of the four minimum races stipulated in the 1977 Standards, mothers of a specified Asian or Pacific Islander (API) subgroup (Chinese, Japanese, Hawaiian, or Filipino) in combination with another

American Indian, Asian, or Hispanic on census and survey records may sometimes be reported as white or non-Hispanic on the death certificate, resulting in an underestimation of deaths and death rates for the American Indian, Asian, and Hispanic groups. Bias also results from undercounts of some population groups in the census—particularly young black males, young white males, and elderly persons—resulting in an overestimation of death rates. Race and ethnicity reporting on the death certificate continues to be excellent for the white and black populations. It remains poor for the American Indian or Alaska Native population but is reasonably good for the Hispanic and Asian or Pacific Islander populations. Decedent characteristics such as place of residence and nativity have an important effect on the quality of reporting on the death certificate. The effects of misclassification on mortality estimates were most pronounced for the American Indian or Alaska Native population, where correcting for misclassification reverses a large American Indian or Alaska Native-over-white mortality advantage to a large disadvantage. Among the Hispanic and Asian or Pacific Islander populations, adjustment for death certificate misclassification did not significantly affect minority-majority mortality. For more information, see: Arias E, Schauman WS, Eschbach K, et al. The validity of race and Hispanic origin reporting on death certificates in the United States. *NCHS. Vital Health Stat* 2008;2(148). Available from: http://www.cdc.gov/nchs/data/series/sr_02/sr02_148.pdf.

Denominators for infant mortality rates are based on the number of live births, rather than on population estimates. Race information for the denominator is supplied from the birth certificate. Before 1980, race of child for the denominator took into account the races of both parents. Starting in 1980, race information for the denominator has been based solely on the race of the mother. Race information for the numerator is supplied from the death certificate. For the infant mortality rate, race information for the numerator is race of the deceased child.

Issues affecting the interpretation of vital event rates for the American Indian or Alaska Native population include (a) changes in the classification or self-identification of persons of American Indian or Alaska Native heritage over time, and (b) misclassification of American Indian or Alaska Native persons on death certificates by the funeral director or informant. Vital event rates for the American Indian or Alaska Native population shown in *Health, United States* are based on the total U.S. resident American Indian or Alaska Native population, as enumerated by the U.S. Census Bureau. In contrast, the Indian Health Service calculates vital event rates for this population based on U.S. Census Bureau county data for American Indian or Alaska Native persons who reside on or near reservations. Because of misclassification of American Indian or Alaska Native persons on death

certificates, American Indian or Alaska Native national and state-specific mortality estimates published in *Health, United States* should be interpreted with caution.

Interpretation of trends for the Asian population in the United States should take into account that this population more than doubled between 1980 and 1990, primarily because of immigration. Between 1990 and 2000, the increase in the Asian population was 48% for persons reporting that they were Asian alone and 72% for persons who reported they were either Asian alone or Asian in combination with another race.

For more information on coding race by using vital statistics, see: NCHS. *Vital statistics of the United States, vol I, Natality, and vol II, Mortality, part A, Technical appendix*. Hyattsville, MD; published annually. Available from: <http://www.cdc.gov/nchs/nvss.htm>.

Starting with 2003 data, some states began using the 2003 revision of the U.S. Standard Certificate of Death, which allows the reporting of more than one race (multiple races). This change was implemented to reflect the increasing diversity of the U.S. population and to be consistent with the decennial census. For more information on states' reporting of multiple-race data, see the annual series of "Deaths: Final Data" reports, available from: <http://www.cdc.gov/nchs/products/nvsr.htm>.

To provide uniformity and comparability of data until all states are reporting multiple-race data, it has been necessary to bridge the responses of those for whom more than one race is reported (multiple race) to one single race. For more information, see: NCHS procedures for multiple-race and Hispanic origin data: Collection, coding, editing, and transmitting. Hyattsville, MD: NCHS; 2004. Available from: http://www.cdc.gov/nchs/data/dvs/Multiple_race_docu_5-10-04.pdf; and NCHS. *Vital statistics of the United States, vol I, Natality, and vol II, Mortality, part A, Technical appendix*. Hyattsville, MD; published annually. Available from: <http://www.cdc.gov/nchs/nvss.htm>.

Youth Risk Behavior Survey (YRBS)—Prior to 1999, the 1977 OMB Standards were used. Respondents could select only one of the following categories: white (not Hispanic), black (not Hispanic), Hispanic or Latino, Asian or Pacific Islander, American Indian or Alaska Native, or other. Beginning in 1999, the 1997 OMB Standards were used for race-specific estimates, and respondents were given the option of selecting more than one category to describe their race and ethnicity. Between 1999 and 2003, students were asked a single question about race and Hispanic origin, with the option of choosing more than one of the following responses: white, black or African American, Hispanic or Latino, Asian, Native Hawaiian or Other Pacific Islander, or American Indian or Alaska Native. In 2005, students were asked a question about Hispanic origin ("Are you Hispanic or Latino?") and a second separate question about race that included the

option of selecting more than one of the following categories: American Indian or Alaska Native, Asian, black or African American, Native Hawaiian or Other Pacific Islander, or white. Because of the differences between questions, data about race and Hispanic ethnicity for the years prior to 1999 are not strictly comparable with estimates for the later years. However, analyses of data collected between 1991 and 2003 have indicated that the data are comparable across years and can be used to study trends. See: Brener ND, Kann L, McManus T. A comparison of two survey questions on race and ethnicity among high school students. *Public Opin Q* 2003;67(2):227–36.

(Also see Appendix II, Hispanic origin; and Appendix I, Population Census and Population Estimates.)

Rate—A rate is a measure of some event, disease, or condition in relation to a unit of population, along with some specification of time. (Also see Appendix II, Age adjustment; Population.)

- *Birth and related rates*

Birth rate is calculated by dividing the number of live births in a population in a year by the resident population. For census years, rates are based on unrounded census counts of the resident population as of April 1. For the noncensus years 1981–1989, rates are based on national estimates of the resident population as of July 1, rounded to thousands. Rounded population estimates for 5-year age groups are calculated by summing unrounded population estimates before rounding to thousands. Starting in 1991, rates are based on unrounded national population estimates. Birth rates for 1991–1999 were revised based on the 1990 and 2000 censuses. The rates for 1990, 2000, and 2010 are based on populations from the censuses in those years as of April 1. Birth rates for 2001–2009 were revised based on the 2000 and 2010 censuses. Birth rates for 2011 and subsequent years were computed using 2010-based postcensal estimates. The population estimates have been provided by the U.S. Census Bureau and have been modified to be consistent with OMB racial categories as of 1977 and historical categories for birth data. Beginning in 1997, the birth rate for the maternal age group 45–49 includes data for mothers aged 50–54 in the numerator and is based on the population of women aged 45–49 in the denominator. Birth rates are expressed as the number of live births per 1,000 population. The rate may be restricted to births to women of specific age, race, marital status, or geographic location (specific rate), or it may be related to the entire population (crude rate).

Fertility rate is the total number of live births, regardless of the age of the mother, per 1,000 women of reproductive age (aged 15–44). Beginning in 1997, the birth rate for the maternal age group 45–49 includes data for mothers aged 50–54 in the numerator and is

based on the population of women aged 45–49 in the denominator.

- *Death and related rates*

Death rate is calculated by dividing the number of deaths in a population in a year by the midyear resident population. For census years, rates are based on unrounded census counts of the resident population as of April 1. For the noncensus years 1981–1989, rates are based on national estimates of the resident population as of July 1, rounded to thousands. Rounded population estimates for 10-year age groups are calculated by summing unrounded population estimates before rounding to thousands. Starting in 1991, rates are based on unrounded national population estimates. Rates for the Hispanic and non-Hispanic white populations in each year are based on unrounded state population estimates for states in the Hispanic reporting area. Death rates are expressed as the number of deaths per 100,000 resident population. The rate may be restricted to deaths in specific age, race, sex, or geographic groups or from specific causes of death (specific rate), or it may be related to the entire population (crude rate). (Also see Appendix I, Population Census and Population Estimates.)

Birth cohort infant mortality rates are based on the birth cohort linked birth and infant death files and are computed as the number of deaths under age 1 year to members of the birth cohort, divided by the number of live births, times 1,000. (Also see Appendix II, Birth cohort.)

Fetal mortality rate is the number of fetal deaths with stated or presumed gestation of 20 weeks or more, divided by the sum of live births plus fetal deaths, times 1,000.

Infant mortality rate is based on period files and is calculated by dividing the number of infant deaths during a calendar year by the number of live births reported in the same year. It is expressed as the number of infant deaths per 1,000 live births. Neonatal mortality rate is the number of deaths among infants under age 28 days per 1,000 live births. Postneonatal mortality rate is the number of infant deaths that occur between 28 days to under 1 year after birth, per 1,000 live births. (Also see Appendix II, Infant death.)

Late fetal mortality rate is the number of fetal deaths with stated or presumed gestation of 28 weeks or more, divided by the sum of live births plus late fetal deaths, times 1,000. (Also see Appendix II, Gestation.)

Perinatal mortality rates and ratios relate to the period surrounding the birth event. Rates and ratios are based on events reported in a calendar year. Although several different perinatal mortality definitions exist, the perinatal definition used in *Health, United States* (and used most commonly for international comparisons) is

the sum of late fetal deaths at 28 weeks of gestation or more plus infant deaths within 7 days of birth, divided by the sum of live births plus late fetal deaths, times 1,000. Perinatal mortality ratio is the sum of late fetal deaths plus infant deaths within 7 days of birth, divided by the number of live births, times 1,000.

- **Visit rate**

Visit rate is a basic measure of service utilization for event-based data. Examples of events include physician office visits with drugs provided, or hospital discharges. In the visit rate calculation, the numerator is the number of estimated events, and the denominator is the corresponding U.S. population estimate for those who possibly could have had events during a given period of time. The interpretation is that for every person in the population there were, on average, x events. It does not mean that x persons in the population had events, because some persons in the population had no events while others had multiple events. The only exception is when an event can occur just once for a person (e.g., if an appendectomy is performed during a hospital stay). The visit rate is best used to compare utilization across various subgroups of interest, such as age or race groups or geographic regions.

Region—See Appendix II, Geographic region.

Registered hospital—See Appendix II, Hospital.

Registration area—The United States has separate registration areas for birth, death, marriage, and divorce statistics. In general, registration areas correspond to states and include two separate registration areas for D.C. and New York City. The term “reporting area” may be used interchangeably with the term “registration area.” All registration areas have adopted laws that require registration of births and deaths and the reporting of fetal deaths. It is believed that more than 99% of births and deaths occurring in this country are registered.

The death registration area was established in 1900 with 10 states and D.C., and the birth registration area was established in 1915, also with 10 states and D.C. Beginning in 1933, all states were included in the birth and death registration areas. The specific states added year by year are shown in: Hetzel AM. History and organization of the vital statistics system. Hyattsville, MD: NCHS; 1997. Available from: <http://www.cdc.gov/nchs/data/misc/usvss.pdf>. Currently, Puerto Rico, the U.S. Virgin Islands, and Guam each constitute a separate registration area, although their data are not included in statistical tabulations of U.S. resident data. (Also see Appendix II, Reporting area.)

Relative standard error (RSE)—RSE is a measure of an estimate's reliability. The RSE of an estimate is obtained by dividing the standard error of the estimate, $SE(r)$, by the estimate itself, r . This quantity is expressed as a percentage of the estimate and is calculated as follows:

$$RSE = 100 \times [SE(r)/r]$$

Estimates with large RSEs are considered unreliable. In *Health, United States*, most statistics with large RSEs are preceded by an asterisk or are not presented. The criteria for evaluating RSEs is discussed in the footnotes accompanying each table.

Relative survival rate—The relative survival rate is the ratio of the observed survival rate for the patient group to the expected survival rate for persons in the general population similar to the patient group with respect to age, sex, race, and calendar year of observation. The 5-year relative survival rate estimates the proportion of cancer patients who have survived their cancer 5 years after diagnosis. Because more than one-half of all cancers occur in persons aged 65 and over, many of these individuals die of other causes with no evidence of recurrence of their cancer. However, by adjusting observed survival for the normal life expectancy of the general population of the same age, the relative survival rate gives a more specific estimate of the chance of surviving the effects of cancer alone.

Reporting area—In the National Vital Statistics System, the reporting area for such basic items on the birth and death certificates as age, race, and sex is based on data from residents of all 50 states in the United States, D.C., and New York City. The term “reporting area” may be used interchangeably with the term “registration area.” (Also see Appendix II, Registration area; and Appendix I, National Vital Statistics System [NVSS].)

Resident, health facility—In the Quality Improvement Evaluation System (QIES) (formerly the Online Survey Certification and Reporting [OSCAR]) database, all residents in certified facilities are counted on the day of certification inspection.

Resident population—See Appendix II, Population.

Rural—See Appendix II, Urbanization.

Self-assessment of health—See Appendix II, Health status, respondent-assessed.

Serious psychological distress—The K6 mental health screening instrument is a measure of psychological distress associated with unspecified but potentially diagnosable mental illness that may result in a higher risk for disability and higher utilization of health services. In the National Health Interview Survey, the K6 questions were asked of adults aged 18 and over. The K6 is designed to identify persons with serious psychological distress, using as few questions as possible. The six items included in the K6 are:

During the past 30 days, how often did you feel:

- So sad that nothing could cheer you up?
- Nervous?
- Restless or fidgety?

- Hopeless?
- That everything was an effort?
- Worthless?

Possible answers are “All of the time” (4 points), “Most of the time” (3 points), “Some of the time” (2 points), “A little of the time” (1 point), and “None of the time” (0 points).

To score the K6, the points are added together, yielding a possible total of 0–24 points. A threshold of 13 points or more is used to define serious psychological distress. Persons answering “Some of the time” to all six questions would not reach the threshold for serious psychological distress because they would need to answer “Most of the time” to at least one item to achieve a score of 13. Only respondents who answered all six psychological distress questions would have a computed K6 score for analysis. The version of the K6 used in NHIS provides 1-month prevalence rates because the reference period is the past 30 days. For more information, see: Kessler RC, Barker PR, Colpe LJ, Epstein JF, Gfroerer JC, Hiripi E, et al. Screening for serious mental illness in the general population. *Arch Gen Psychiatry* 2003;60(2):184–9. (Also see Appendix II, Basic actions difficulty.)

Starting in 2013, the K6 questions were moved to the adult selected items section of the Sample Adult questionnaire. Observed differences between the 2012 and earlier estimates and 2013 and later estimates may be partially or fully attributable to this change in question placement within the Sample Adult questionnaire.

Short-stay hospital—See Appendix II, Hospital.

Skilled nursing facility—See Appendix II, Nursing home.

Smoker—See Appendix II, Cigarette smoking.

Special hospital—See Appendix II, Hospital.

Substance use—Substance use refers to the use of selected substances, including alcohol, tobacco products, drugs, inhalants, and other substances that can be consumed, inhaled, injected, or otherwise absorbed into the body with possible dependence and other detrimental effects. (Also see Appendix II, Illicit drug use.)

Monitoring the Future (MTF) Study—MTF collects information on the use of selected substances by using self-completed questionnaires in a school-based survey of secondary school students. MTF has tracked 12th graders' illicit drug use and attitudes toward drugs since 1975. In 1991, 8th and 10th graders were added to the study. The survey includes questions on abuse of substances including (but not limited to) marijuana, inhalants, other illegal drugs, alcohol, cigarettes, and other tobacco products. (Also see Appendix I, Monitoring the Future [MTF] Study.)

National Survey on Drug Use & Health (NSDUH)—NSDUH conducts in-person, computer-assisted interviews of a sample of individuals aged 12 and over at their place of residence. For illicit drug use, alcohol use, and tobacco use, information is collected about use in the lifetime, past year, and past month. However, only estimates of use in the past month are presented in *Health, United States*. For illicit drug use, respondents in NSDUH are asked about use of marijuana/hashish, cocaine (including crack), inhalants, hallucinogens, heroin, and prescription-type psychotherapeutic drugs (pain relievers, tranquilizers, stimulants, and sedatives) used nonmedically. A series of questions is asked about each substance: “Have you ever, even once, used [substance]?” and “How long has it been since you last used [substance]?” Numerous probes and checks are included in the computer-assisted interview system. Nonprescription medications and legitimate use of prescription drugs under a doctor's supervision are not included in the estimates. Starting in 2013, information about marijuana use that was recommended by a doctor or other health care professional has been collected; however, any reported marijuana use is classified as illicit drug use. Summary measures, such as current illicit drug use, are produced. (Also see Appendix II, Alcohol consumption; Cigarette smoking; Illicit drug use; and Appendix I, National Survey on Drug Use & Health [NSDUH].)

Suicidal ideation—Suicidal ideation means having thoughts of suicide or of taking action to end one's own life. Suicidal ideation includes all thoughts of suicide, both when the thoughts include a plan to commit suicide and when they do not include a plan. Suicidal ideation is measured in the Youth Risk Behavior Survey by the following three questions: “During the past 12 months, did you ever seriously consider attempting suicide?”, “During the past 12 months, how many times did you actually attempt suicide?”, and “If you attempted suicide during the past 12 months, did any attempt result in an injury, poisoning, or overdose that had to be treated by a doctor or nurse?” For more information, see: <http://www.cdc.gov/HealthyYouth/yrbs/index.htm>.

Surgery—See Appendix II, Outpatient surgery; Procedure.

Surgical specialty—See Appendix II, Physician specialty.

Tobacco use—See Appendix II, Cigarette smoking.

Uninsured—Broadly, persons are considered uninsured if they do not have coverage under private health insurance, Medicare, Medicaid, public assistance (through 1996), Children's Health Insurance Program (CHIP), a state-sponsored or other government-sponsored plan or program, or a military health plan. Because of differences in methodology, question wording, and recall period, estimates from different sources may vary and are not directly comparable. For more information, see: Health insurance measurement: Differences by data source.

Available from: http://www.census.gov/content/dam/Census/library/infographics/health_insurance_measurement.pdf.

American Community Survey (ACS)—In ACS, persons are considered uninsured if they do not have coverage through private health insurance, Medicare, Medicaid, CHIP, military/TRICARE or veterans coverage, another government program, or other insurance. Persons with only Indian Health Service coverage are considered uninsured. The questions on health insurance are administered throughout the year and ask about current health insurance coverage as of the day of the interview.

National Health Interview Survey (NHIS)—In NHIS, the uninsured are persons who do not have coverage under private health insurance, Medicare, Medicaid, public assistance (through 1996), CHIP, a state-sponsored health plan, other government-sponsored programs, or a military health plan. Persons with only Indian Health Service coverage are considered uninsured. Estimates for the uninsured are shown only for the population under age 65. Estimates of the percentage of persons who are uninsured based on NHIS may differ slightly from those based on the March Current Population Survey or the American Community Survey because of differences in survey questions, recall period, and other aspects of survey methodology.

Survey respondents may be covered by health insurance at the time of interview but may have experienced one or more lapses in coverage during the year prior to interview. Starting with *Health, United States, 2006*, NHIS estimates for people with health insurance coverage for all 12 months prior to interview, for those who were uninsured for any period up to 12 months, and for those who were uninsured for more than 12 months were added as stub variables to selected tables. (Also see Appendix II, Health insurance coverage.)

Urbanization—Urbanization is the degree of urban (city-like) character of a particular geographic area. Urbanization can be measured in a variety of ways. In *Health, United States*, the two measures currently used to categorize counties by urbanization level are the Office of Management and Budget's (OMB) metropolitan and micropolitan statistical area classification and the 2013 NCHS Urban–Rural Classification Scheme for Counties. For more information on the OMB classification of counties, see Appendix II, Metropolitan statistical area (MSA); Micropolitan statistical area.

The 2013 Urban–Rural Classification Scheme is based on the February 2013 OMB delineation of MSAs and micropolitan statistical areas, 2012 postcensal estimates of county and place population, and county-level data on selected settlement density, socioeconomic, and demographic variables from Census 2010. This is an updated version of NCHS' earlier scheme, the 2006 NCHS Urban–Rural Classification Scheme for Counties. The six categories of the NCHS scheme are large central metro (inner-city counties of

MSAs of 1 million or more population), large fringe metro (suburban counties of MSAs of 1 million or more population), medium metro (counties of MSAs of 250,000–999,999 population), small metro (counties of MSAs with less than 250,000 population), nonmetropolitan micropolitan statistical areas, and nonmetropolitan noncore. For more information on the classification scheme, see: http://www.cdc.gov/nchs/data_access/urban_rural.htm.

Usual source of care—Usual source of care was measured in the National Health Interview Survey in 1993 and 1994 by asking the respondent, “Is there a particular person or place that [person] usually goes to when [person] is sick or needs advice about [person's] health?” In the 1995 and 1996 NHIS, the respondent was asked, “Is there one doctor, person, or place that [person] usually goes to when [person] is sick or needs advice about health?” Starting in 1997, the respondent was asked, “Is there a place that [person] usually goes when he/she is sick or you need advice about [his/her] health?” Persons who report the emergency department as their usual source of care are defined in *Health, United States* as having no usual source of care.

Vaccination—Vaccinations, or immunizations, work by stimulating the immune system—the natural disease-fighting system of the body. A healthy immune system is able to recognize invading bacteria and viruses and produce substances (antibodies) to destroy or disable these invaders. Vaccinations prepare the immune system to ward off a disease. In addition to the initial immunization process, the effectiveness of some immunizations can be improved by periodic repeat injections or “boosters.” Vaccines are among the most successful and cost-effective public health tools available for reducing morbidity and mortality from vaccine-preventable diseases. For a comprehensive list of vaccine-preventable diseases, see: <http://www.cdc.gov/vaccines/vpd-vac/vpd-list.htm> and <http://www.cdc.gov/vaccines/spec-grps/default.htm>.

The currently recommended childhood vaccination schedule includes vaccines that prevent infectious diseases including hepatitis A and B, diphtheria, tetanus toxoids, acellular pertussis (whooping cough), measles, mumps, rubella (German measles), polio, varicella (chicken pox), and some forms of meningitis (HIB), influenza, and pneumococcal disease. In February 2006, a rotavirus vaccine (RotaTeq) was licensed for use in U.S. infants.

A vaccine that protects against the four types of human papillomavirus (HPV) that cause most cervical cancers and genital warts began to be marketed in 2006 and is now available for both females and males. The vaccine was recommended for girls aged 11 and 12 and for girls and women aged 13–26 who have not yet been vaccinated or completed the vaccine series. In October 2011, HPV vaccination was recommended for males aged 11 and 12. Further information is available from: <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6050a3.htm>.

Boosters (revaccination) of vaccinations received during childhood or adulthood are necessary for some vaccines. In addition to keeping current with the vaccines listed above, and annual influenza vaccination, some additional vaccinations are recommended for older adults, persons with specific health conditions, or health care workers who are likely to be exposed to infectious persons. Herpes zoster vaccination is recommended for adults aged 60 and over, and pneumococcal vaccination is recommended for adults aged 65 and over.

For a full discussion of recommended vaccination schedules by age and population, see CDC's vaccination and immunization website at: <http://www.cdc.gov/vaccines/schedules/index.html>.

Influenza vaccination—In the National Health Interview Survey, questions concerning influenza vaccination have been slightly different across the survey years. Respondents were asked, “During the past 12 months, have you had a flu shot? A flu shot is usually given in the fall and protects against influenza for the flu season.” Beginning in September 2003, respondents were asked about influenza vaccination by nasal spray (sometimes called by the brand name FluMist) during the past 12 months, in addition to the question regarding the flu shot. Starting with 2005 data, receipt of nasal spray or a flu shot was included in the calculation of influenza vaccination estimates. In 2010, additional questions were asked about the receipt of the H1N1 flu shot and spray, including month and year received. These H1N1 questions, and the original seasonal flu questions, were asked only in quarters 1 and 2 and the first several weeks of quarter 3. Beginning August 11, 2010, revised flu vaccination questions replaced all flu vaccination questions fielded earlier in 2010 and were used in 2011 and beyond. The revised questions reflect the introduction of a new combined flu vaccination that protects against both the seasonal and H1N1 strains. For more information regarding the influenza questions that were introduced in 2010, see: ftp://ftp.cdc.gov/pub/Health_Statistics/NCHS/Dataset_Documentation/NHIS/2010/srvydesc.pdf.

The prevalence of influenza vaccination during the past 12 months may differ from season-specific coverage, and estimates from different data sources may differ (additional estimates are available from: <http://www.cdc.gov/flu/fluview/>). Prevalence of influenza vaccination during the past 12 months is different from season-specific coverage, see: CDC. Surveillance of influenza vaccination coverage—United States, 2007–08 through 2011–12 influenza seasons. MMWR 2013;62(ss04):1–29. Available from: http://www.cdc.gov/mmwr/preview/mmwrhtml/ss6204a1.htm?s_cid=ss6204a1_w.

The recommendations of the Advisory Committee on Immunization Practices regarding who should receive an influenza vaccination have changed over the years, and

changes in coverage estimates may reflect changes in recommendations. An influenza vaccine shortage occurred during the 2004–2005 influenza season. Delays in the availability of influenza shots also occurred in fall 2000 and, to a lesser extent, in fall 2001.

Pneumococcal vaccination—In the National Health Interview Survey, questions concerning pneumococcal vaccination were slightly different across the survey years. Respondents were asked, “Have you EVER had a pneumonia vaccination? This shot is usually given only once in a person's lifetime and is different from the flu shot.” Starting in 1999, respondents were asked, “Have you EVER had a pneumonia vaccination, sometimes called a pneumonia shot? This shot is usually given only once in a person's lifetime and is different from the flu shot.” Starting in 2001, respondents were asked, “Have you EVER had a pneumonia shot? This shot is usually given only once or twice in a person's lifetime and is different from the flu shot. It is also called the pneumococcal vaccine.”

Wages and salaries—See Appendix II, Employer costs for employee compensation.

Years of potential life lost (YPLL)—YPLL is a measure of premature mortality. Starting with *Health, United States, 1996*, YPLL has been presented for persons under age 75 because the average life expectancy in the United States is over 75 years. YPLL–75 is calculated using the following eight age groups: under 1, 1–14, 15–24, 25–34, 35–44, 45–54, 55–64, and 65–74. The number of deaths for each age group is multiplied by years of life lost, calculated as the difference between age 75 years and the midpoint of the age group. For the eight age groups, the midpoints are 0.5, 7.5, 19.5, 29.5, 39.5, 49.5, 59.5, and 69.5 years, respectively. For example, the death of a person aged 15–24 counts as 55.5 years of life lost. YPLL is derived by summing years of life lost over all age groups. In *Health, United States, 1995* and earlier editions, YPLL was presented for persons under age 65. For more information, see: CDC. Premature mortality in the United States: Public health issues in the use of years of potential life lost. MMWR 1986;35(SS-02):1S–11S. Available from: <http://www.cdc.gov/mmwr/preview/mmwrhtml/00001773.htm>.

Index

(Numbers are table and figure numbers)

A	<i>Table/Figure (F)</i>
Abortion	7
Access to care (see also Dental visits; Emergency department visits; Health insurance; Hospital utilization; Injuries; Unmet need for medical care, dental care, prescription drugs)	
Health care visits	65, F14
No recent health care visit, children	64
No usual source of care	61, 62
Accidents, see Motor vehicle-related injuries; Unintentional injuries.	
Activities of daily living (ADL), see Basic actions difficulty; Complex activity limitation; Limitation of activity.	
Adolescents, see Child and adolescent health.	
AIDS, see HIV/AIDS.	
Alcohol consumption	50, 51, 52
Allergy	35
Alzheimer's disease	17, 18, 19, 20, F2
American Indian or Alaska Native population	
Access to care	61, 62, 63, 64, 65
Alcohol consumption	50
Allergy	35
Asthma	35
Attention-deficit/hyperactivity disorder	35
Back pain, low	41
Basic actions difficulty	42
Birth rates	3, 4, F4
Births, preterm	F20
Birthweight, low	5, 6
Breast cancer	26, 36
Cancer incidence rates	36
Cesarean delivery, low-risk	F21
Cancer, respondent-reported	38
Chronic conditions, selected	39
Cigarette smoking	49, 50, F24
Colorectal tests or procedures	72
Complex activity limitation	42
Death rates, all causes	16, 17, 21
Death rates, selected causes	17, 18, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32
Death rates, state and U.S. territory	16
Deaths, leading causes	19
Dental visits	78
Drug poisoning	27
Ear infection	35
Emergency department visits	73, 74
Emotional or behavioral difficulties	35
Expenses, health care	97, 98
Headache, severe or migraine	41
Health care visits	65
Health insurance	102, 103, 104, 105, F26
Health status, respondent-assessed	45
Hearing trouble	44
Heart disease, respondent-reported	38
HIV diagnoses	34

A—Con.	<i>Table/Figure (F)</i>
American Indian or Alaska Native population—Con.	
Hospital utilization, inpatient	81
Illicit drug use	50
Infant mortality	10, 12, F19
Mammography	70
Marijuana use	50
Medicaid	104, 109
Neck pain	41
Occupational injury deaths	32
Out-of-pocket health care expenditures	97, 98
Pap smear (Pap test)	71
Physical activity	57
Population, resident	1
Poverty	2
Serious psychological distress	46
Stroke, respondent-reported	38
Teenage childbearing	3, F4
Unmarried mothers	4
Unmet need	63, F27
Vaccinations	66, 67, 68, 69, F25
Vision trouble	43
Years of potential life lost (YPLL)	18
Asian or Pacific Islander population	
Access to care	61, 62, 63, 64, 65
Alcohol consumption	50
Allergy	35
Asthma	35
Attention-deficit/hyperactivity disorder	35
Back pain, low	41
Basic actions difficulty	42
Birth rates	3, 4, F4
Births, preterm	F20
Birthweight, low	5, 6
Breast cancer	26, 36
Cancer incidence rates	36
Cancer, respondent-reported	38
Cesarean delivery, low-risk	F21
Cholesterol	55
Chronic conditions, selected	39
Cigarette smoking	49, 50, F24
Colorectal tests or procedures	72
Complex activity limitation	42
Death rates, all causes	16, 17, 21
Death rates, selected causes	17, 18, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32
Death rates, state and U.S. territory	16
Deaths, leading causes	19
Dental caries (cavities), untreated	60
Dental visits	78
Diabetes	40
Drug poisoning	27

A—Con.

B—Con.

	<i>Table/Figure (F)</i>		<i>Table/Figure (F)</i>
Asian or Pacific Islander population—Con.		Black or African American population	
Ear infection	35	Access to care	61, 62, 63, 64, 65
Emergency department visits	73, 74	Alcohol consumption	50, 51, 52
Emotional or behavioral difficulties	35	Allergy	35
Expenses, health care	97, 98	Asthma	35
Glycemic control	40	Attention-deficit/hyperactivity disorder	35
Headache, severe or migraine	41	Back pain, low	41
Health care visits	65	Basic actions difficulty	42
Health insurance	102, 103, 104, 105, F26	Birth rates	3, 4, F4
Health status, respondent-assessed	45	Births, preterm	F20
Hearing trouble	44	Birthweight, low	5, 6
Heart disease, respondent-reported	38	Breast cancer	26, 36
HIV diagnoses	34	Breastfeeding	9
Hospital utilization, inpatient	81	Cancer incidence rates	36
Hypertension	54, F23	Cancer, respondent-reported	38
Illicit drug use	50	Cancer survival, 5-year relative	37
Infant mortality	10, 12, F19	Cesarean delivery, low-risk	F21
Mammography	70	Cholesterol	55
Marijuana use	50	Chronic conditions, selected	39
Medicaid	104, 109	Cigarette smoking	47, 48, 49, 50, 51, F24
Neck pain	41	Cocaine use	51
Normal weight	58	Colorectal tests or procedures	72
Obesity	F22	Complex activity limitation	42
Occupational injury deaths	32	Contraception	8
Out-of-pocket health care expenditures	97, 98	Death rates, all causes	16, 17, 21
Overweight and obesity	58, 59	Death rates, selected causes	17, 18, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32
Pap smear (Pap test)	71	Death rates, state and U.S. territory	16
Physical activity	57	Deaths, leading causes	19
Population, resident	1	Dental caries (cavities), untreated	60
Poverty	2	Dental visits	78, 108
Serious psychological distress	46	Diabetes	40
Stroke, respondent-reported	38	Doctor visits	77
Teenage childbearing	3, F4	Drug poisoning	27
Unmarried mothers	4	Drugs, prescription, use in past 30 days	79
Unmet need	63, F27	Ear infection	35
Vaccinations	66, 67, 68, 69, F25	Emergency department visits	73, 74, 76
Vision trouble	43	Emotional or behavioral difficulties	35
Years of potential life lost (YPLL)	18	Expenses, health care	97, 98
Asthma	35	Fetal mortality	10
Attention-deficit/hyperactivity disorder	35	Glycemic control	40
		Headache, severe or migraine	41
		Health care visits	65
		Health insurance	102, 103, 104, 105, 106, F26
		Health status, respondent-assessed	45
		Hearing trouble	44
		Heart disease, respondent-reported	38
		HIV diagnoses	34
		Hospital utilization, inpatient	81, 108
		Hospital utilization, outpatient department	76, 108
		Hypertension	54, F23
		Illicit drug use	50, 51
		Infant mortality	10, 11, 12, F19
		Inhalants	51
		Life expectancy	15, F18
		Limitation of activity	108
		Mammography	70
		Marijuana use	50, 51
B			
Back pain, low	41		
Basic actions difficulty	41, 42, 45, 49, 57, 62, 63, 65, 68, 69, 70, 71, 72, 74, 78, 81, 102, 103, 104, 105		
Bed, health facility	89, 90, 92		
Birth control, see Contraception.			
Births			
Age of mother	3, 4		
Birth rates	3, 4, F4		
Births, number	4		
Births, preterm	F20		
Birthweight, low	5, 6		
Cesarean delivery, low-risk	F21		
Fertility rates	3		
State	6		
Teenage childbearing	F4		
Unmarried mothers	4		

B—Con.

C—Con.

	<i>Table/Figure (F)</i>
Black or African American population—Con.	
Medicaid	98, 104, 106, 109
Medicare	98, 106, 108
Neck pain	41
Normal weight	58
Nursing home expenditures	108
Nursing home utilization	108
Obesity	F22
Occupational injury deaths	32
Out-of-pocket health care expenditures	97, 98
Overweight and obesity	58, 59
Pap smear (Pap test)	71
Physical activity	57
Population, resident	1
Poverty	2
Seatbelt use	52
Serious psychological distress	46
Stroke, respondent-reported	38
Suicidal ideation	52
Teenage childbearing	3, F4
Unmarried mothers	4
Unmet need	63, F27
Vaccinations	66, 67, 68, 69, F25
Violence	52
Vision trouble	43
Years of potential life lost (YPLL)	18
Blood pressure, high, see Hypertension.	
Breastfeeding	9

C

Calories, see Energy and macronutrient intake.	
Cancer (Malignant neoplasms)	
Breast	17, 18, 26, 36, 37
Deaths and death rates	17, 19, 20, 24, 25, 26, F2
Incidence rates	36
Prevalence, respondent-reported	38
Site-specific data	17, 18, 25, 26, 36, 37
Survival, 5-year relative	37
Trachea, bronchus, lung	17, 25, 36, 37
Years of potential life lost (YPLL)	18
Central and South American population, see Hispanic origin subgroups.	
Cerebrovascular disease (stroke)	
Deaths and death rates	17, 19, 20, 23, F2
Prevalence, respondent-reported	38
Years of potential life lost (YPLL)	18
Cesarean section delivery	96, F21
Chancroid, see Diseases, notifiable.	
Child and adolescent health	
Access to care	61, 63, 64, 65
Alcohol consumption	50, 51, 52
Allergy	35
Asthma	35
Attention-deficit/hyperactivity disorder	35
Birthweight, low	5, 6

	<i>Table/Figure (F)</i>
Child and adolescent health—Con.	
Breastfeeding	9
Cigarette smoking	50, 51
Cocaine use	51
Contraception	8
Death rates, all causes	20, 21
Death rates, selected causes	20, 22, 23, 24, 27, 28, 29 30, 31, 32
Deaths, leading causes	20, 21
Dental caries (cavities), untreated	60
Dental visits	78
Doctor visits	77
Drug poisoning	27
Drugs, prescription, use in past 30 days	79, 80
Ear infection	35
Emergency department visits	73, 75, 76
Emotional or behavioral difficulties	35
Expenses, health care	96, 97, 98, 99
Health insurance	102, 103, 104, 105
Health status, respondent-assessed	45
HIV diagnoses	34
Hospital utilization, inpatient	81
Hospital utilization, outpatient department	76
Illicit drug use	50, 51
Infant mortality	10, 11, 12, 13, F19
Inhalants	51
Injury	75
Marijuana use	50, 51
Medicaid	98, 104, 109
Obesity	53, 59, F8, F22
Out-of-pocket health care expenditures	97, 98, 99
Physical activity	52
Population, resident	1
Poverty	2
Seatbelt use	52
Sleep	52
Suicidal ideation	52
Teenage childbearing	3, 4, F4
Vaccinations	66, 67
Violence	52
Chlamydia, see Diseases, notifiable.	
Cholesterol	53, 55
Chronic conditions, selected	39
Chronic liver disease and cirrhosis	17, 18, 19, 20
Chronic lower respiratory diseases	17, 18, 19, 20, F2
Cigarette smoking	47, 48, 49, 50, 51, F7, F24
Cirrhosis, see Chronic liver disease and cirrhosis.	
Cocaine use	51
Colorectal tests or procedures	72
Complex activity limitation	41, 42, 45, 49, 57, 62, 63, 65, 68, 69, 70, 71, 72, 74, 78, 81, 102, 103, 104, 105
Congenital anomalies	19, 20
Contraception	8
Cost, see Employers' costs for health insurance.	
Cuban population, see Hispanic origin subgroups.	

D

Table/Figure (F)

Deaths, death rates (see also Cancer [malignant neoplasms]; Cerebrovascular disease [stroke]; Chronic lower respiratory diseases; Diabetes; Drug poisoning; Firearm-related injuries; Heart disease; Homicide; Infant mortality; Life expectancy; Motor vehicle-related injuries; Occupational injury deaths; Suicide; Years of potential life lost [YPLL])	
All causes	21
Leading causes	19, 20
Selected causes	17, F2
State	16
Dental caries (cavities), untreated	53, 60
Dental services expenditures	94
Dental visits	78, 108
Dentists	86, 88
Schools and students	88
State	86
Diabetes	17, 18, 19, 20, 40, 53, F2
Deaths and death rates	17, 19, 20, F2
Prevalence	40, 53
Years of potential life lost (YPLL)	18
Diphtheria, see Diseases, notifiable; Vaccinations.	
Disability	
Basic actions difficulty	41, 42, 45, 49, 57, 62, 63, 65, 68, 69, 70, 71, 72, 74, 78, 81, 102, 103, 104, 105
Blind and disabled Medicaid expenditures	109
Complex activity limitation	41, 42, 45, 49, 57, 62, 63, 65, 68, 69, 70, 71, 72, 74, 78, 81, 102, 103, 104, 105
Disability measures, selected	F6
Medicaid recipients	110
Medicare beneficiaries	108
Veterans with service-connected disabilities	111
Diseases, notifiable	33, F5
Doctors of Medicine, see Physicians.	
Drug poisoning	27
Drug use, illicit, see Alcohol consumption; Cigarette smoking; Cocaine use; Illicit drug use; Inhalants; Marijuana use.	
Drugs, prescription, use in past 30 days	79, 80
DTP (diphtheria, tetanus, pertussis), see Vaccinations.	

E

Ear infection	35
Education	
Access to care	63
Alcohol consumption	51, 52
Back pain, low	41
Breastfeeding	9
Cancer, respondent-reported	38
Cigarette smoking	48, 49, 51
Cocaine use	51
Colorectal tests or procedures	72
Headache, severe or migraine	41
Hearing trouble	44
Heart disease, respondent-reported	38
Illicit drug use	51
Inhalants	51
Mammography	70
Marijuana use	51
Neck pain	41
Pap smear (Pap test)	71
Physical activity	57

E—Con.

Table/Figure (F)

Education—Con.	
Seatbelt use	52
Stroke, respondent-reported	38
Suicidal ideation	52
Unmet need	63
Violence	52
Vision trouble	43
Elderly population, see Older population aged 65 and over.	
Electronic health record (EHR) components	F13
Emergency department visits	73, 74, 75, F11
Emotional or behavioral difficulties	35
Employed health service personnel	87
Employers' costs for health insurance	101
Energy and macronutrient intake	56
Ethnicity, see Hispanic or Latino population.	
Exercise, see Physical activity.	
Expenditures, national health (see also Consumer Price Index [CPI]; Hospital care expenditures; Medicaid; Medicare; Nursing homes expenditures; Physician services expenditures; Prescription expenditures; Veterans' medical care)	
Amount per capita	93, 95
Percent of Gross Domestic Product	93
Personal health care	93, 94, 95, F15
Source of funds	93, 95
Type of expenditure	94, 95
Type of payer	100
Expenses, health care	97, 98, 99

F

Fertility rates, see Births.	
Fetal mortality	10
Firearm-related injuries, death rates	31
Food intake, see Energy and macronutrient intake.	

G

Geographic region	
Access to care	61, 62, 63, 64, 65
Back pain, low	41
Breastfeeding	9
Cancer, respondent-reported	38
Chronic conditions, selected	39
Colorectal tests or procedures	72
Dental visits	78
Emergency department visits	73, 74
Headache, severe or migraine	41
Health care visits	65
Health insurance	102, 103, 104, 105
Health status, respondent-assessed	45
Hearing trouble	44
Heart disease, respondent-reported	38
Hospital utilization, inpatient	81
Neck pain	41
Physical activity	57
Serious psychological distress	46
Stroke, respondent-reported	38
Unmet need	63
Vaccinations	68, 69
Vision trouble	43

G—Con.

H—Con.

	Table/Figure (F)
Glycemic control	40
Gonorrhea, see Diseases, notifiable.	
Gross Domestic Product (GDP)	93

H

<i>Haemophilus influenza</i> , invasive, see Diseases, notifiable.	
Hawaiian population, see Native Hawaiian or Other Pacific Islander population.	
Headache, severe or migraine	41
Health care expenses, see Expenses, health care.	
Health care utilization	64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, F11
Health expenditures, national, see Expenditures, national health.	
Health insurance (see also Access to care; Emergency department visits; Medicaid; Medicare)	
Basic actions difficulty	102, 103, 104, 105
Complex activity limitation	102, 103, 104, 105
Employer costs	101
Employment related	103
Medicaid	104, F11, F12, F14, F15, F16, F17
Private	102, 103, F11, F12, F14, F15, F16, F17
Race and Hispanic origin	102, 103, 104, 105, 106
65 and over	106
Under age 65	102, 103, 104, 105
Uninsured	105, 114, F11, F12, F14, F15, F16, F17, F26
Health professionals visits, see Visits to health professionals.	
Health status, respondent-assessed	45
Hearing trouble	44
Heart disease	
Deaths and death rates	17, 19, 20, 22, F2
Drugs, prescription, use in past 30 days	80
Ischemic heart disease	17, 18
Prevalence, respondent-reported	38
Years of potential life lost (YPLL)	18
Hepatitis A, see Disease, notifiable.	
Hepatitis B, see Disease, notifiable.	
Heroin poisoning	27
Hib (<i>Haemophilus influenza</i> type b), see Vaccinations.	
Hispanic or Latino population	
Access to care	61, 62, 63, 64, 65
Alcohol consumption	50, 52
Allergy	35
Asthma	35
Attention-deficit/hyperactivity disorder	35
Back pain, low	41
Basic actions difficulty	42
Birth rates	3, 4, F4
Births, preterm	F20
Birthweight, low	5, 6
Breast cancer	26, 36
Breastfeeding	9
Cancer incidence rates	36
Cancer, respondent-reported	38
Cesarean delivery, low-risk	F21
Cholesterol	55
Chronic conditions, selected	39
Cigarette smoking	49, 50, F24
Colorectal tests or procedures	72

Table/Figure (F)

Hispanic or Latino population—Con.	
Complex activity limitation	42
Contraception	8
Death rates, all causes	16, 17, 21
Death rates, selected causes	17, 18, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32
Death rates, state and U.S. territory	16
Deaths, leading causes	19
Dental caries (cavities), untreated	60
Dental visits	78, 108
Diabetes	40
Drug poisoning	27
Drugs, prescription, use in past 30 days	79
Ear infection	35
Emergency department visits	73, 74
Emotional or behavioral difficulties	35
Expenses, health care	97, 98
Fetal mortality	10
Glycemic control	40
Headache, severe or migraine	41
Health care visits	65
Health insurance	102, 103, 104, 105, 106, F26
Health status, respondent-assessed	45
Hearing trouble	44
Heart disease, respondent-reported	38
HIV diagnoses	34
Hospital utilization, inpatient	81, 108
Hospital utilization, outpatient department	108
Hypertension	54, F23
Illicit drug use	50
Infant mortality	10, 12, F19
Life expectancy	15, F18
Limitation of activity	108
Mammography	70
Marijuana use	50
Medicaid	104, 106, 109
Medicare	98, 106, 108
Neck pain	41
Normal weight	58
Nursing home expenditures	108
Nursing home utilization	108
Obesity	F22
Occupational injury deaths	32
Out-of-pocket health care expenditures	97, 98
Overweight and obesity	58, 59
Pap smear (Pap test)	71
Physical activity	57
Population, resident	1
Poverty	2
Seatbelt use	52
Serious psychological distress	46
Stroke, respondent-reported	38
Suicidal ideation	52
Teenage childbearing	3, F4
Unmarried mothers	4
Unmet need	63, F27
Vaccinations	66, 67, 68, 69, F25
Violence	52

H—Con.

I—Con.

	<i>Table/Figure (F)</i>
Hispanic or Latino population—Con.	
Vision trouble	43
Years of potential life lost (YPLL)	18
Hispanic origin subgroups (Central and South American population; Cuban population) (see also Mexican origin population; Puerto Rican population)	
Birth rates	4
Births, preterm	F20
Birthweight, low	5
Cesarean delivery, low-risk	F21
Chronic conditions, selected	39
Health insurance	102, 103, 104, 105
Infant mortality	10, F19
Unmarried mothers	4
HIV/AIDS	
Deaths and death rates	17, 19, 20
HIV diagnoses	34
Years of potential life lost (YPLL)	18
Home health care expenditures	94
Homicide, death rates	17, 18, 19, 20, 29, F3
Hospital care expenditures (see also Consumer Price Index [CPI]; Medicaid; Medicare)	95, 96
Hospital discharges	81
Hospital utilization (see also Access to care; Emergency department visits; Medicaid; Medicare; Veterans' medical care)	
Admissions	82
Average length of stay	82, 112
Cost for procedures or surgeries	96
Outpatient department	76, 82, 108
Race and Hispanic origin	81, 108
Hospitals (see also Mental health; Nursing homes)	
Beds	89, 90
Occupancy rate	89, 91
State	90, 91
Hypercholesterolemia, See Cholesterol.	
Hypertension	53, 54, F23

I

Illicit drug use	27, 30, 50, 51
Immunizations, see Vaccinations.	
Incidence (Cancer)	36
Income, family, see Poverty.	
Infant mortality (see also Fetal mortality)	
Age at death	10, 11
Birth cohort data	10
Cause of death	20
International	13
Race and Hispanic origin	10, 11, 12, F19
State	12
Infectious disease	
Deaths	17, 18, 19, 20
Notifiable diseases	33, 34, F5
Vaccinations	66, 67, 68, 69, F25
Influenza and pneumonia	17, 18, 19, 20
Influenza vaccination, see Vaccinations.	
Inhalants	51
Injuries, see Emergency department visits; Firearm-related injuries; Death rates; Motor vehicle-related injuries; Occupational injuries; Unintentional injuries.	

Table/Figure (F)

Inpatient care, see Hospital utilization; Mental health; Nursing homes, utilization.	
Instrumental activities of daily living (IADL), see Limitation of activity.	
Insurance, see Health insurance.	
International health (see also Expenditures, national health; Infant mortality; Life expectancy)	13, 14, F1
Ischemic heart disease, see Heart disease.	

K

Kidney disease, see Nephritis, nephrotic syndrome and nephrosis.

L

Leading causes of death, see Deaths, leading causes.	
Leisure-time activity, see Physical activity.	
Life expectancy	14, 15, F1, F18
Limitation of activity (see also Basic actions difficulty; Complex activity limitation)	108
Liver disease, see Chronic liver disease and cirrhosis.	
Low birthweight, see Births; Infant mortality.	
Low income, see Poverty.	
Lyme disease, see Diseases, notifiable.	

M

Malignant neoplasms, see Cancer.	
Mammography	70
Marijuana use	50, 51
Marital status	102, 103, 104, 105, 106
Maternal health, see Women's health.	
Measles, see Diseases, notifiable; Vaccinations.	
Medicaid (see also Health insurance)	
Basic actions difficulty	104
Basis of eligibility	109
Complex activity limitation	104
Coverage	104, 106
Expenses, health care	97
Expenditures	95, 100, F15
Payments	109, 110, 113
Race and Hispanic origin	104, 109
State	113
Type of service	110
Medical doctors, see Physicians.	
Medicare (see also Health insurance)	
Age and sex of beneficiaries	106
Coverage	106
Enrollment	107, 108, 112
Expenses, health care	97
Expenditures	95, 100, 107, F15
Hospital utilization	112
Payments	98, 112
Race and Hispanic origin	106, 108
State	112
Type of service	107
Meningococcal disease	20, 33, F5
Men's health	
Access to care	62, 63, 65
Alcohol consumption	50
Back pain, low	41
Basic actions difficulty	42

	<i>Table/Figure (F)</i>
Men's health—Con.	
Cancer incidence rates	36
Cancer, respondent-reported	38
Cancer survival, 5-year relative	37
Cholesterol	55
Chronic conditions, selected	39
Cigarette smoking	47, 48, 49, 50, F7, F24
Colorectal tests or procedures	72
Complex activity limitation	42
Contraception	8
Death rates, all causes	17, 21, F2
Death rates, selected causes	17, 18, 22, 23, 24, 25, 27, 28, 29, 30, 31, 32, 34, F2, F3
Deaths, leading causes	19
Dental caries (cavities), untreated	60
Dental visits	78
Diabetes	40
Disability measures, selected	F6
Doctor visits	77
Drug poisoning	27
Drugs, prescription, use in past 30 days	79, 80
Emergency department visits	74, 75, 76
Energy and macronutrient intake	56
Expenses, health care	97, 98
Glycemic control	40
Headache, severe or migraine	41
Health insurance	102, 103, 104, 105, 106, 108
Health status, respondent-assessed	45
Hearing trouble	44
Heart disease, respondent-reported	38
HIV diagnoses	34
Hospital utilization, inpatient	81
Hospital utilization, outpatient department	76
Hypertension	54, F23
Illicit drug use	50
Injury	75
Life expectancy	14, 15, F1, F18
Marijuana use	50
Neck pain	41
Normal weight	58
Occupational injury deaths	32
Overweight and obesity	58, F9
Physical activity	57
Population, resident	1
Serious psychological distress	46
Stroke, respondent-reported	38
Unmet need	63
Vaccinations	68, 69
Vision trouble	43
Years of potential life lost (YPLL)	18
Mental health (see also Suicide)	
Drugs, prescription, use in past 30 days	80
Emotional or behavioral difficulties, children	35
Expenditures	110
Psychiatrists	84
Serious psychological distress	46

	<i>Table/Figure (F)</i>
Metropolitan/nonmetropolitan data	
Access to care	61, 62, 63, 64, 65, F14
Back pain, low	41
Basic actions difficulty	42
Cancer, respondent-reported	38
Chronic conditions, selected	39
Colorectal tests or procedures	72
Complex activity limitation	42
Dental visits	78
Emergency department visits	73, 74
Headache, severe or migraine	41
Health care visits	65
Health insurance	102, 103, 104, 105
Health status, respondent-assessed	45
Hearing trouble	44
Heart disease, respondent-reported	38
Hospital utilization, inpatient	81
Medicaid	104
Neck pain	41
Physical activity	57
Serious psychological distress	46
Stroke, respondent-reported	38
Unmet need	63
Vaccinations	66, 67, 68, 69
Vision trouble	43
Mexican origin population (see also Hispanic origin subgroups)	
Access to care	62, 63
Back pain, low	41
Birth weight, low	5
Births, preterm	F20
Births, rate	4
Cancer, respondent-reported	38
Cesarean delivery, low-risk	F21
Cholesterol	55
Cigarette smoking	49
Colorectal tests or procedures	72
Dental caries (cavities), untreated	60
Diabetes	40
Drugs, prescription, use in past 30 days	79
Emergency department visits	74
Glycemic control	40
Headache, severe or migraine	41
Health care visits	65
Health insurance	102, 103, 104, 105
Health status, respondent-assessed	45
Hearing trouble	44
Heart disease, respondent-reported	38
Hypertension	54, F23
Infant mortality	10, F19
Medicaid	104
Neck pain	41
Normal weight	58
No usual source of care	62
Overweight and obesity	58, 59
Physical activity	57
Poverty	2
Serious psychological distress	46

M—Con.

O—Con.

	<i>Table/Figure (F)</i>
Mexican origin population (see also Hispanic origin subgroups)—Con.	
Stroke, respondent-reported	38
Unmarried mother	4
Unmet need	63
Vaccinations	68, 69
Vision trouble	43
MMR (measles, mumps, rubella), see Vaccinations.	
Motor vehicle-related injuries	17, 18, 28, 75
Mumps, see Diseases, notifiable; Vaccinations.	

N

National health expenditures, see Expenditures, national health.	
Native Hawaiian or Other Pacific Islander population	
Alcohol consumption	50
Cigarette smoking	50
HIV diagnoses	34
Illicit drugs	50
Occupational injury deaths	32
Vaccinations	66
Neck pain	41
Neonatal mortality, see Infant mortality, age at death.	
Nephritis, nephrotic syndrome and nephrosis	17, 18, 19, 20
Nurses	87
Nursing homes	
Beds, occupancy	92
Expenditures	94, 95, 108
Utilization	92, 108, 111
Nutrition, see Energy and macronutrient intake.	

O

Obesity	53, 58, 59, F8, F9, F22
Occupational injury deaths	32
Occupational therapists	87
Office visits	76, 77
Older population aged 65 and over	
Access to care	63, 65
Back pain, low	41
Basic actions difficulty	42
Cancer, respondent-reported	38
Cholesterol	55
Chronic conditions, selected	39
Cigarette smoking	47, 49, F7
Complex activity limitation	42
Death rates, all causes	21
Death rates, selected causes	22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32
Deaths, leading causes	20
Dental caries (cavities), untreated	60
Dental visits	78, 108
Diabetes	40
Disability measures, selected	F6
Doctor visits	77
Drug poisoning	27
Drugs, prescription, use in past 30 days	79, 80
Emergency department visits	74, 75, 76
Energy and macronutrient intake	56
Expenses, health care	96, 97, 98, 99

Table/Figure (F)

Older population aged 65 and over—Con.	
Glycemic control	40
Headache, severe or migraine	41
Health insurance	106, 108
Health status, respondent-assessed	45
Hearing trouble	44
Heart disease, respondent-reported	38
HIV diagnoses	34
Homicide	F3
Hospital utilization, inpatient	81, 96, 108, 112
Hospital utilization, outpatient department	76, 108
Hypertension	54
Injury	75
Life expectancy	14, 15
Limitation of activity	108
Mammography	70
Medicaid	109
Medicare	98, 106, 107, 108, 112
Neck pain	41
Normal weight	58
Nursing home expenditures	108
Nursing home utilization	92, 108
Occupational injury deaths	32
Out-of-pocket health care expenses	97, 98, 99
Overweight and obesity	58
Pap smear (Pap test)	71, F10
Physical activity	57
Population, resident	1
Serious psychological distress	46
Stroke, respondent-reported	38
Suicide	F3
Unmet need	63
Vaccinations	68, 69, F25
Vision trouble	43
Opioid poisoning	27
Optometry students	88
Osteopaths, see Physicians.	
Out-of-pocket health care expenses	97, 98, 99, 100, F15
Outpatient department, see Hospital utilization, outpatient department.	
Overweight	53, 58, F9

P

Pap smear (Pap test)	71, F10
Perinatal mortality, see Infant mortality, age at death.	
Personal health care expenditures, see Expenditures, national health.	
Pertussis (whooping cough), see Diseases, notifiable; Vaccinations.	
Pharmacists/pharmacy students	87, 88
Physical activity	52, 57
Physician services expenditures (see also Consumer Price Index [CPI]; Medicaid; Medicare)	95
Physician utilization	76, 77
Physicians	
Doctors of osteopathy	88
International medical school graduates	84
Primary care	77, 85
Primary specialty	77, 84, 85
Schools and students	88
State	83

	<i>Table/Figure (F)</i>
Pneumococcal vaccinations, see Vaccinations.	
Podiatry students	88
Poliomyelitis (polio), see Diseases, notifiable; Vaccinations.	
Population, resident	1
Postneonatal mortality, see Infant mortality, age at death.	
Poverty	
Access to care	61, 62, 63, 64, 65
Allergy	35
Asthma	35
Attention-deficit/hyperactivity disorder	35
Back pain, low	41
Basic actions difficulty	42
Cancer, respondent-reported	38
Cholesterol	55
Chronic conditions, selected	39
Cigarette smoking	49
Colorectal tests or procedures	72
Complex activity limitation	42
Dental caries (cavities), untreated	60
Dental visits	78
Diabetes	40
Ear infection	35
Emergency department visits	73, 74
Emotional or behavioral difficulties	35
Glycemic control	40
Headache, severe or migraine	41
Health care visits	65
Health insurance	102, 103, 104, 105, 106
Health status, respondent-assessed	45
Hearing trouble	44
Heart disease, respondent-reported	38
Hospital utilization, inpatient	81
Hypertension	54
Mammography	70
Medicaid	104, 106
Medicare	106
Neck pain	41
Normal weight	58
Overweight and obesity	58, 59
Pap smear (Pap test)	71
Physical activity	57
Population	2
Serious psychological distress	46
Stroke, respondent-reported	38
Unmet need	63
Vaccinations	66, 67, 68, 69
Vision trouble	43
Prescription drug expenditures (see also Medicaid; Medicare)	94, 95, 97
Prescription drug use, see Drugs, prescription, use in past 30 days.	
Primary care physicians, see Physicians.	
Private health insurance, see Health insurance.	
Procedures, see Hospital utilization.	
Public Health, schools of; students	88
Puerto Rican population (see also Hispanic origin subgroups)	
Births	4, 5
Birthweight, low	6
Births, preterm	F20

	<i>Table/Figure (F)</i>
Puerto Rican population (see also Hispanic origin subgroups)—Con.	
Cesarean delivery, low-risk	F21
Death rates, state and U.S. territory	16
Health insurance	102, 103, 104, 105
Infant mortality	10, 12, F19
Poverty	2

R

Race, see specific racial groups.
Rocky Mountain spotted fever, see Diseases, notifiable.
Rubella (German measles), see Diseases, notifiable; Vaccinations.
Rubeola (measles), see Diseases, notifiable; Vaccinations.
Rural data, see Metropolitan/nonmetropolitan data.

S

Salmonellosis, see Diseases, notifiable.	
Self-assessment of health, see Health status, respondent-assessed.	
Septicemia	19, 20
Serious psychological distress (see also Mental health)	46
Shigellosis, see Diseases, notifiable.	
Smoking, see Cigarette smoking.	
Source of funds or payments (see also Expenditures, national health; Health insurance; Medicaid; Medicare)	95, 98, 100
Special Feature on Racial and Ethnic Health Disparities	F18, F19, F20, F21, F22, F23, F24, F25, F26, F27
State and U.S. territory data	
Birthweight, low	6
Death rates	16
Dentists	86
Health insurance, uninsured	114
Hospital beds	90
Hospital occupancy rates	91
Infant mortality	12
Medicaid	113
Medicare	112
Nursing homes, beds, occupancy rates.	92
Physicians	83
Sterilization, see Contraception.	
Stroke, see Cerebrovascular disease (stroke).	
Sudden infant death syndrome, see Infant mortality, cause of death.	
Suicidal ideation	52
Suicide	17, 18, 19, 20, 30, F3
Syphilis, see Diseases, notifiable.	

T

Tetanus, see Diseases, notifiable; Vaccinations.
Tobacco use, see Cigarette smoking.
Tuberculosis, see Diseases, notifiable.

U

Uninsured, health, see Health insurance, uninsured.	
Unintentional injuries	17, 18, 19, 20, 75, F2
Unmet need for medical care, dental care, prescription drugs	63, F12, F27
Urban and rural data, see Metropolitan/nonmetropolitan data.	
U.S. territories, see State and U.S. territory data.	
Usual source of care, see Access to care.	

V

	<i>Table/Figure (F)</i>
Vaccinations	66, 67, 68, 69, F25
Varicella, see Vaccinations.	
Veterans' medical care	111
Vision trouble	43
Visits to health professionals	65

W

Wages and salaries	87, 101
Wages, health care occupations	87
White population	
Access to care	61, 62, 63, 64, 65
Alcohol consumption	50, 51, 52
Allergy	35
Asthma	35
Attention-deficit/hyperactivity disorder	35
Back pain, low	41
Basic actions difficulty	42
Birth rates	3, 4, F4
Births, preterm	F20
Birthweight, low	5, 6
Breast cancer	26, 36
Breastfeeding	9
Cancer incidence rates	36
Cancer, respondent-reported	38
Cancer survival, 5-year relative	37
Cesarean delivery, low-risk	F21
Cholesterol	55
Chronic conditions, selected	39
Cigarette smoking	47, 48, 49, 50, 51, F24
Cocaine use	51
Colorectal tests or procedures	72
Complex activity limitation	42
Contraception	8
Death rates, all causes	16, 17, 21
Death rates, selected causes	17, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32
Death rates, state and U.S. territory	16
Deaths, leading causes	19
Dental caries (cavities), untreated	60
Dental visits	78
Diabetes	40
Doctor visits	77
Drug poisoning	27
Drugs, prescription, use in past 30 days	79
Ear infection	35
Emergency department visits	73, 74, 76
Emotional or behavioral difficulties	35
Expenses, health care	97, 98
Fetal mortality	10
Glycemic control	40
Headache, severe or migraine	41
Health care visits	65
Health insurance	102, 103, 104, 105, 106, F26
Health status, respondent-assessed	45
Hearing trouble	44
Heart disease, respondent-reported	38

W—Con.

	<i>Table/Figure (F)</i>
White population—Con.	
HIV diagnoses	34
Hospital utilization, inpatient	81, 108
Hospital utilization, outpatient department	76, 108
Hypertension	54, F23
Illicit drug use	50, 51
Infant mortality	10, 11, 12, F19
Inhalants	51
Life expectancy	15, F18
Limitation of activity	108
Mammography	70
Marijuana use	50, 51
Medicaid	98, 104, 106, 109
Medicare	98, 106, 108
Neck pain	41
Normal weight	58
Nursing home expenditures	108
Nursing home utilization	108
Obesity	F22
Occupational injury deaths	32
Out-of-pocket health care expenditures	97, 98
Overweight and obesity	58, 59
Pap smear (Pap test)	71
Physical activity	57
Population, resident	1
Poverty	2
Seatbelt use	52
Serious psychological distress	46
Stroke, respondent-reported	38
Suicidal ideation	52
Teenage childbearing	3, F4
Unmarried mothers	4
Unmet need	63, F27
Vaccinations	66, 67, 68, 69, F25
Violence	52
Vision trouble	43
Years of potential life lost (YPLL)	18
Women's health	
Access to care	62, 63, 65
Abortion	7
Alcohol consumption	50
Back pain, low	41
Basic actions difficulty	42
Birth rates, fertility rates	3, 4
Births, number	4
Births, preterm	F20
Breast cancer	26, 36, 37
Breastfeeding	9
Cancer incidence rates	36
Cancer, respondent-reported	38
Cancer survival, 5-year relative	37
Cesarean delivery, low-risk	F21
Cholesterol	55
Chronic conditions, selected	39
Cigarette smoking	47, 48, 49, 50, F7, F24
Colorectal tests or procedures	72
Complex activity limitation	42

	<i>Table/Figure (F)</i>	<i>Table/Figure (F)</i>
Women's health—Con.		
Contraception	8	
Death rates, all causes	17, 21, F2	
Death rates, selected causes	17, 18, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, F2, F3	
Deaths, leading causes	19	
Dental caries (cavities), untreated	60	
Dental visits	78	
Diabetes	40	
Disability measures, selected	F6	
Doctor visits	77	
Drug poisoning	27	
Drugs, prescription, use in past 30 days	79, 80	
Emergency department visits	74, 75, 76	
Energy and macronutrient intake	56	
Expenses, health care	97, 98	
Glycemic control	40	
Headache, severe or migraine	41	
Health insurance	102, 103, 104, 105, 106, 108	
Health status, respondent-assessed	45	
Hearing trouble	44	
Heart disease, respondent-reported	38	
HIV diagnoses	34	
Hospital utilization, inpatient	81	
Hospital utilization, outpatient department	76	
Hypertension	54, F23	
Illicit drug use	50	
Injury	75	
Life expectancy	14, 15, F1, F18	
Mammography	70	
Marijuana use	50	
Neck pain	41	
Normal weight	58	
Occupational injury deaths	32	
Overweight and obesity	58, F9	
Pap smear (Pap test)	71, F10	
Physical activity	57	
Population, resident	1	
Poverty	2	
Serious psychological distress	46	
Stroke, respondent-reported	38	
Teenage childbearing	3, F4	
Unmarried mothers	4	
Unmet need	63	
Vaccinations	68, 69	
Vision trouble	43	
Years of potential life lost (YPLL)	18	
Working-age adults (aged 18–64)	39, 42, 62, 63, 69, 71, 74, 78, 81, 102, 103, 104, 105, F6, F12, F16, F17, F25, F26, F27	
Years of potential life lost (YPLL)		18
Young adults (aged 19–25)		62, 63, 102, 103, 104, 105